

# 1. Kinematics

**Decay rate and cross section** (Note:  $\mathcal{M}$  has a mass dimension of  $4 - N_i - N_f$ .)

$$\text{decay rate (rest frame; } \sqrt{s} = M_0) : \quad d\Gamma = \frac{d\Pi^{N_f}}{2M_0} \left| \mathcal{M}(M_0 \rightarrow \{p_1, p_2, \dots, p_{N_f}\}) \right|^2, \quad (1.1)$$

$$\text{cross section (Lorentz invariant) :} \quad d\sigma = \frac{d\Pi^{N_f}}{2E_A 2E_B v_{\text{Mol}}} \left| \mathcal{M}(p_A, p_B \rightarrow \{p_1, p_2, \dots, p_{N_f}\}) \right|^2, \quad (1.2)$$

where  $d\Pi^n$  is  $n$ -particle Lorentz-invariant phase space with momentum conservation

$$d\Pi^n := d\Pi_1 d\Pi_2 \dots d\Pi_n (2\pi)^4 \delta^{(4)} \left( P_0 - \sum p_n \right); \quad d\Pi := \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{1}{2E_{\mathbf{p}}}. \quad (1.3)$$

At the CM frame, two-body phase-space are characterized by the final momentum  $\|\mathbf{p}\|$  and given by

$$d\Pi^2 = \frac{\|\mathbf{p}\|}{4\pi\sqrt{s}} \frac{d\Omega}{4\pi} = \frac{\|\mathbf{p}\|}{8\pi\sqrt{s}} d\cos\theta = \frac{1}{16\pi} \sqrt{1 - \frac{2(m_1^2 + m_2^2)}{s} + \frac{(m_1^2 - m_2^2)^2}{s^2}} d\cos\theta \quad (1.4)$$

with  $\sqrt{s} = M_0$  or  $E_{\text{CM}}$ ,  $\theta$  is the angle between initial and final motion, and


$$\|\mathbf{p}\| = \frac{\sqrt{s}}{2} \lambda^{1/2} \left( 1; \frac{m_1^2}{s}, \frac{m_2^2}{s} \right), \quad E_1 = \frac{s + m_1^2 - m_2^2}{2\sqrt{s}}, \quad E_2 = \frac{s - m_1^2 + m_2^2}{2\sqrt{s}}, \quad p_1 \cdot p_2 = \frac{s - (m_1^2 + m_2^2)}{2}.$$

**Mandelstam variables** For  $(k_1, k_2) \rightarrow (p_3, p_4)$  collision,

$$s = (k_1 + k_2)^2 = (p_3 + p_4)^2, \quad t = (p_3 - k_1)^2 = (p_4 - k_2)^2, \quad u = (p_3 - k_2)^2 = (p_4 - k_1)^2; \\ s + t + u = m_1^2 + m_2^2 + m_3^2 + m_4^2.$$


If the collision is with the “same mass”  $(m_A, m_A) \rightarrow (m_B, m_B)$ ,

$$t = m_A^2 + m_B^2 - s/2 + 2kp \cos\theta, \quad (k_1 - k_2)^2 = 4m_A^2 - s, \\ u = m_A^2 + m_B^2 - s/2 - 2kp \cos\theta, \quad (p_3 - p_4)^2 = 4m_B^2 - s, \\ k = \frac{\sqrt{s - 4m_A^2}}{2}, \quad k_1 \cdot k_2 = \frac{s}{2} - m_A^2, \quad k_1 \cdot p_3 = k_2 \cdot p_4 = \frac{m_A^2 + m_B^2 - t}{2}, \\ p = \frac{\sqrt{s - 4m_B^2}}{2}, \quad p_3 \cdot p_4 = \frac{s}{2} - m_B^2, \quad k_1 \cdot p_4 = k_2 \cdot p_3 = \frac{m_A^2 + m_B^2 - u}{2}.$$



Instead, if the collision is “initially massless”  $(0, 0) \rightarrow (m_3, m_4)$ ,

$$t = (m_3^2 + m_4^2 - s)/2 + p\sqrt{s} \cos\theta, \\ u = (m_3^2 + m_4^2 - s)/2 - p\sqrt{s} \cos\theta, \\ p = (\sqrt{s}/2) \lambda^{1/2} \left( 1; m_3^2/s, m_4^2/s \right).$$



## 1.1. Fundamentals

Lorentz-invariant phase space:

$$\int d\Pi = \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{1}{2E_{\mathbf{p}}} = \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{1}{2\sqrt{m^2 + \|\mathbf{p}\|^2}} = \int \frac{dp_0 d^3\mathbf{p}}{(2\pi)^4} (2\pi) \delta(p_0^2 - \|\mathbf{p}\|^2 - m^2) \Theta(p_0)$$

Källén function:

$$\begin{aligned} \lambda(x, y, z) &= x^2 + y^2 + z^2 - 2xy - 2yz - 2zx = (x - y - z)^2 - 4yz; \\ \lambda(1; \alpha_1^2, \alpha_2^2) &= (1 - (\alpha_1 + \alpha_2)^2)(1 - (\alpha_1 - \alpha_2)^2) = (1 + \alpha_1 + \alpha_2)(1 - \alpha_1 - \alpha_2)(1 + \alpha_1 - \alpha_2)(1 - \alpha_1 + \alpha_2). \\ \lambda^{1/2}(s; m_1^2, m_2^2) &= s \lambda^{1/2}\left(1; \frac{m_1^2}{s}, \frac{m_2^2}{s}\right); & \lambda^{1/2}\left(1; \frac{m^2}{s}, \frac{m^2}{s}\right) &= \sqrt{1 - \frac{4m^2}{s}}, \\ \lambda^{1/2}\left(1; \frac{m_1^2}{s}, \frac{m_2^2}{s}\right) &= \sqrt{1 - \frac{2(m_1^2 + m_2^2)}{s} + \frac{(m_1^2 - m_2^2)^2}{s^2}}, & \lambda^{1/2}\left(1; \frac{m_1^2}{s}, 0\right) &= \frac{s - m_1^2}{s}. \end{aligned}$$

**Two-body phase space** If  $f(p_1^\mu, p_2^\mu)$  is Lorentz invariant,  $f \equiv f(p_1^2, p_2^2, p_1^\mu p_{2\mu}) \equiv f(p_1, p_2, \cos \theta_{12})$ . Meanwhile,

$$\int d\Pi_1 d\Pi_2 = \int \frac{d^3\mathbf{p}_1}{(2\pi)^3} \frac{d^3\mathbf{p}_2}{(2\pi)^3} \frac{1}{2E_1 2E_2} = \int \frac{(4\pi) dp_1 p_1^2 (2\pi) dp_2 p_2^2 d\cos \theta_{12}}{(2\pi)^3} \frac{1}{2E_1 2E_2} = \int \frac{dE_+ dE_- ds}{128\pi^4}, \quad (1.5)$$

with the replacement of the variables

$$E_{\pm} = E_1 \pm E_2, \quad s = (p_1 + p_2)^2 = m_1^2 + m_2^2 + 2E_1 E_2 - 2\|\mathbf{p}_1\| \|\mathbf{p}_2\| \cos \theta_{12};$$

$$\left| \frac{d(E_+, E_-, s)}{d(p_1, p_2, \cos \theta_{12})} \right| = \frac{4p_1^2 p_2^2}{E_1 E_2}, \quad \left| \frac{d(E_1, E_2, s)}{d(p_1, p_2, \cos \theta_{12})} \right| = \frac{2p_1^2 p_2^2}{E_1 E_2}.$$

Therefore,

$$\int d\Pi_1 d\Pi_2 = \frac{1}{128\pi^4} \int_{(m_1+m_2)^2}^{\infty} ds \int_{\sqrt{s}}^{\infty} dE_+ \int_{\min}^{\max} dE_-, \quad (1.6)$$

where the boundary of  $E_-$  is given by

$$\begin{aligned} \cos \theta_{12} &= \frac{E_+^2 - E_-^2 + 2(m_1^2 + m_2^2 - s)}{\sqrt{(E_+ + E_-)^2 - 4m_1^2} \sqrt{(E_+ - E_-)^2 - 4m_2^2}} \in [-1, 1] \\ \therefore \left| E_- - \frac{m_1^2 - m_2^2}{s} E_+ \right| &\leq \sqrt{E_+^2 - s} \cdot \lambda^{1/2}\left(1; \frac{m_1^2}{s}, \frac{m_2^2}{s}\right) = 2p \sqrt{\frac{E_+^2 - s}{s}}. \end{aligned}$$

**Two-body phase space with momentum conservation** As a general representation in any frame,

$$\frac{d\Pi^2}{16\pi^2} = \frac{dp_1 d\Omega p_1^2}{16\pi^2} \frac{\delta(E_0 - \sqrt{m_1^2 + p_1^2} - \sqrt{m_2^2 + \|\mathbf{P}_0 - \mathbf{p}_1\|^2})}{E_1 E_2} = \frac{1}{8\pi} d\cos \theta_1 \frac{p_1^2}{E_0 p_1 - P_0 E_1 \cos \theta_1}, \quad (1.7)$$

where the momentum  $p_1$  is given by

$$p_1 = \frac{(E_0^2 + m_1^2 - m_2^2 - P_0^2) P_0 \cos \theta_1 + E_0 \sqrt{\lambda(E_0^2, m_1^2, m_2^2) + P_0^4 - 2P_0^2(E_0^2 + m_1^2 - 2m_1^2 \cos^2 \theta_1 - m_2^2)}}{2(E_0^2 - P_0^2 \cos^2 \theta_1)}. \quad (1.8)$$

CM frame result is recovered by setting  $E_0 = \sqrt{s}$  and  $P_0 = 0$ .

## 1.2. Decay rate and Cross section

As  $\langle \text{out} | \text{in} \rangle = (2\pi)^4 \delta^{(4)}(p_i - p_f) i\mathcal{M}$  (for  $\text{in} \neq \text{out}$ ) and  $\langle \mathbf{p} | \mathbf{p} \rangle = 2E_{\mathbf{p}} (2\pi)^3 \delta^{(3)}(\mathbf{0}) = 2E_{\mathbf{p}} V$  for one-particle state,

$$\frac{N_{\text{ev}}}{\prod_{\text{in}} N_{\text{particle}}} = \int d\Pi^{\text{out}} \frac{|\langle \text{out} | \text{in} \rangle|^2}{\langle \text{in} | \text{in} \rangle} = \int d\Pi^{\text{out}} \frac{(2\pi)^8 |\mathcal{M}|^2}{\prod_{\text{in}} (2E)V (2\pi^4)} \delta^{(4)}(p_i - p_f) = VT \int d\Pi^{N_f} \frac{|\mathcal{M}|^2}{\prod_{\text{in}} (2E)V}. \quad (1.9)$$

Therefore, decay rate (at the rest frame) is given by

$$d\Gamma := \frac{1}{T} \frac{dN_{\text{ev}}}{N_{\text{particle}}} = \frac{1}{T} VT d\Pi^{N_f} \frac{|\mathcal{M}|^2}{(2E)V} = \frac{1}{2M_0} d\Pi^{N_f} |\mathcal{M}|^2. \quad (1.10)$$

We also define Lorentz-invariant cross section  $\sigma$  by  $N_{\text{ev}} =: (\rho_A v_{\text{Mø}} T \sigma) N_B = (\rho_A v_{\text{Mø}} T \sigma) (\rho_B V)$ , or

$$d\sigma := \frac{dN_{\text{ev}}}{\rho_A v_{\text{Mø}} T N_B} = \frac{V}{v_{\text{Mø}} T} VT d\Pi^{N_f} \frac{|\mathcal{M}|^2}{2E_A 2E_B V^2} = \frac{1}{2E_A 2E_B v_{\text{Mø}}} d\Pi^{N_f} |\mathcal{M}|^2. \quad (1.11)$$

where the Møller parameter  $v_{\text{Mø}}$  is equal to  $v_{\text{rel}}^{\text{NR}} = \|\mathbf{v}_A - \mathbf{v}_B\|$  if  $\mathbf{v}_A \parallel \mathbf{v}_B$  (cf. Ref. [?]). Generally,

$$v_{\text{Mø}} := \frac{\sqrt{(p_A \cdot p_B)^2 - m_A^2 m_B^2}}{E_A E_B} = \frac{p_A \cdot p_B}{E_A E_B} v_{\text{rel}} = (1 - \mathbf{v}_A \cdot \mathbf{v}_B) v_{\text{rel}}, \quad (1.12)$$

where  $v_{\text{rel}}$  is the actual relative velocity

$$v_{\text{rel}} = \sqrt{1 - \frac{(1 - v_A^2)(1 - v_B^2)}{1 - (\mathbf{v}_A \cdot \mathbf{v}_B)^2}} = \frac{\sqrt{\|\mathbf{v}_A - \mathbf{v}_B\|^2 - \|\mathbf{v}_A \times \mathbf{v}_B\|^2}}{1 - \mathbf{v}_A \cdot \mathbf{v}_B} = \frac{\lambda^{1/2}(s, m_A^2, m_B^2)}{s - (m_A^2 + m_B^2)} \neq v_{\text{rel}}^{\text{NR}}. \quad (1.13)$$

(Note that  $p_A \cdot p_B / E_A E_B = 1$  if  $\mathbf{p}_A = 0$  or  $\mathbf{p}_B = 0$ . Also, Each of  $v_{\text{rel}}$ ,  $VT$ , and  $E_A E_B v_{\text{Mø}}$  is Lorentz invariant.)

## 2. Gauge theory

**SU(2)** Fundamental representation  $\mathbf{2} = (T^a)_{ij}$ , adjoint representation  $\text{adj.} = (\epsilon^a)^{bc}$ .<sup>\*1</sup>

$$T_a = \frac{1}{2}\sigma_a, \quad \text{Tr}(T_a T_b) = \frac{1}{2}\delta_{ab}, \quad [T_a, T_b] = i\epsilon^{abc}T^c, \quad \epsilon^{abc}\epsilon^{ade} = \delta_{bd}\delta_{ce} - \delta_{be}\delta_{cd}$$

Since  $\bar{\mathbf{2}} = -(T^a)^*_{ij}$  has identities  $-\epsilon T^a \epsilon = -T^{a*}$  and  $-\epsilon(-T^{a*})\epsilon = T^a$ , we see that  $\epsilon^{ab}\mathbf{2}^b$  transforms as  $\bar{\mathbf{2}}^a$ :

$$\epsilon^{ab}\mathbf{2}^b \rightarrow \epsilon^{ab}[\exp(i\theta^\alpha T^\alpha)]^{bc}\mathbf{2}^c = \epsilon^{ab}[\exp(i\theta^\alpha T^\alpha)]^{bc}(\epsilon^{-1})^{cd}(\epsilon^{de}\mathbf{2}^e) = [\exp(-i\theta^\alpha T^{\alpha*})]^{ab}(\epsilon^{bc}\mathbf{2}^c). \quad (2.1)$$

**SU(3)** Fundamental representation  $\mathbf{3} = (\tau^a)_{ij}$ ,  $\bar{\mathbf{3}} = -(\tau^a)^*_{ij}$ ; adjoint representation  $\text{adj.} = \mathbf{8} = (f^a)^{bc}$ .  
Gell-Mann matrices:

$$\lambda_{1-8} = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & -i \\ 0 & 0 & 0 \\ i & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix}, \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix}. \quad (2.2)$$

$$\tau_a = \frac{1}{2}\lambda_a, \quad \text{Tr}(\tau_a \tau_b) = \frac{1}{2}\delta_{ab}, \quad [\tau_a, \tau_b] = if^{abc}\tau^c, \quad f^{ade}f^{bcd} + f^{bde}f^{cad} + f^{cde}f^{abd} = 0.$$

$$\begin{aligned} \mathbf{3}: \quad & \phi_a \rightarrow [\exp(i\theta^\alpha \tau^\alpha)]_{ab}\phi_b \simeq \phi_a + i\theta^\alpha \tau_{ab}^\alpha \phi_b & \bar{\mathbf{3}}: \quad & \phi_a \rightarrow [\exp(-i\theta^\alpha \tau^{\alpha*})]_{ab}\phi_b \simeq \phi_a - i\theta^\alpha \tau_{ab}^{\alpha*} \phi_b \\ & \phi_a^* \rightarrow [\exp(-i\theta^\alpha \tau^{\alpha*})]_{ab}\phi_b^* \simeq \phi_a^* - i\theta^\alpha \tau_{ab}^{\alpha*} \phi_b^* & & \phi_a^* \rightarrow [\exp(i\theta^\alpha \tau^\alpha)]_{ab}\phi_b^* \simeq \phi_a^* + i\theta^\alpha \tau_{ab}^\alpha \phi_b^* \\ & = \phi_b^*[\exp(-i\theta^\alpha \tau^\alpha)]_{ba} \simeq \phi_a^* - i\theta^\alpha \phi_b^* \tau_{ba}^\alpha & & \end{aligned}$$

---

<sup>\*1</sup>We do not distinguish sub- and superscripts for gauge indices.

### 3. Spinors

$$(\overline{\psi_1}\psi_2)^* = (\psi_2)^\dagger(\overline{\psi_1})^\dagger = \overline{\psi_2}\psi_1. \quad (3.1)$$

## 4. Supersymmetry with $\eta = \text{diag}(+, -, -, -)$

**Convention** Our convention follows DHM (except for  $D_\mu$ ):

$$\begin{aligned}\eta &= \text{diag}(1, -1, -1, -1); \quad \epsilon^{0123} = -\epsilon_{0123} = 1, \quad \epsilon^{12} = \epsilon_{21} = \epsilon^{\dot{1}\dot{2}} = \epsilon_{\dot{2}\dot{1}} = 1 \quad (\epsilon_{\alpha\beta}\epsilon^{\beta\gamma} = \epsilon^{\alpha\beta}\epsilon_{\beta\gamma} = \delta_\gamma^\alpha), \\ \psi^\alpha &= \epsilon^{\alpha\beta}\psi_\beta, \quad \psi_\alpha = \epsilon_{\alpha\beta}\psi^\beta, \quad \bar{\psi}^{\dot{\alpha}} = \epsilon^{\dot{\alpha}\dot{\beta}}\bar{\psi}_{\dot{\beta}}, \quad \bar{\psi}_{\dot{\alpha}} = \epsilon_{\dot{\alpha}\dot{\beta}}\bar{\psi}^{\dot{\beta}}; \\ \sigma_{\alpha\dot{\alpha}}^\mu &:= (\mathbf{1}, \boldsymbol{\sigma})_{\alpha\dot{\alpha}}, \quad \sigma^{\mu\nu}{}_\alpha{}^\beta := \frac{i}{4}(\sigma^\mu\bar{\sigma}^\nu - \sigma^\nu\bar{\sigma}^\mu)_\alpha{}^\beta,^{*2} \quad (\sigma_{\alpha\dot{\beta}}^\mu = \epsilon_{\alpha\delta}\epsilon_{\dot{\beta}\dot{\gamma}}\bar{\sigma}^{\mu\dot{\gamma}\delta}, \quad \bar{\sigma}^{\mu\dot{\alpha}\beta} = \epsilon^{\dot{\alpha}\dot{\delta}}\epsilon^{\beta\gamma}\sigma_{\gamma\dot{\delta}}^\mu) \\ \bar{\sigma}^{\mu\dot{\alpha}\alpha} &:= (\mathbf{1}, -\boldsymbol{\sigma})^{\dot{\alpha}\alpha}, \quad \bar{\sigma}^{\mu\nu}{}_{\dot{\beta}}{}^{\dot{\alpha}} := \frac{i}{4}(\bar{\sigma}^\mu\sigma^\nu - \bar{\sigma}^\nu\sigma^\mu)_{\dot{\beta}}{}^{\dot{\alpha}},^{*2} \\ (\psi\xi) &:= \psi^\alpha\xi_\alpha, \quad (\bar{\psi}\bar{\chi}) := \bar{\psi}_{\dot{\alpha}}\bar{\chi}^{\dot{\alpha}}; \quad \frac{d}{d\theta^\alpha}(\theta\theta) := \theta_\alpha \quad [\text{left derivative}].\end{aligned}$$

Especially, spinor-index contraction is done as  ${}^\alpha_\alpha$  and  ${}_{\dot{\alpha}}^{\dot{\alpha}}$  except for  $\epsilon_{ab}$  (which always comes from left). Noting that complex conjugate reverses spinor order:  $(\psi^\alpha\xi^\beta)^* := (\xi^\beta)^*(\psi^\alpha)^*$ ,

$$\begin{aligned}\bar{\psi}^{\dot{\alpha}} &:= (\psi^\alpha)^*, \quad \epsilon^{\dot{a}\dot{b}} := (\epsilon^{ab})^*, \quad (\psi\chi)^* = (\bar{\psi}\bar{\chi}), \\ (\sigma_{\alpha\dot{\beta}}^\mu)^* &= \bar{\sigma}^{\mu\dot{\alpha}\beta} = \epsilon_{\beta\delta}\epsilon_{\dot{\alpha}\dot{\gamma}}\bar{\sigma}^{\mu\dot{\gamma}\delta}, \quad (\sigma^{\mu\nu})^\dagger{}_\alpha{}^\beta = \bar{\sigma}^{\mu\nu}{}_{\dot{\beta}}{}^{\dot{\alpha}}, \quad (\sigma^{\mu\nu}{}_\alpha{}^\beta)^* = \bar{\sigma}^{\mu\nu}{}_{\dot{\alpha}}{}^{\dot{\beta}} = \bar{\sigma}^{\mu\nu}{}_{\dot{\alpha}}{}^{\dot{\beta}} = \epsilon_{\dot{\alpha}\dot{\gamma}}\epsilon^{\dot{\beta}\delta}\bar{\sigma}^{\mu\dot{\gamma}\delta}, \\ (\bar{\sigma}^{\mu\dot{\alpha}\beta})^* &= \sigma^{\mu\alpha\dot{\beta}} = \epsilon^{\dot{\beta}\dot{\delta}}\epsilon^{\alpha\gamma}\sigma_{\gamma\dot{\delta}}^\mu, \quad (\bar{\sigma}^{\mu\nu})^\dagger{}_{\dot{\alpha}}{}^{\dot{\beta}} = \sigma^{\mu\nu}{}_\alpha{}^\beta, \quad (\bar{\sigma}^{\mu\nu}{}_{\dot{\beta}}{}^{\dot{\alpha}})^* = \sigma^{\mu\nu}{}_\beta{}^\alpha = \sigma^{\mu\nu}{}_\beta{}^\alpha = \epsilon_{\beta\delta}\epsilon^{\alpha\gamma}\sigma^{\mu\dot{\delta}\gamma}_\gamma.\end{aligned}$$

### Contraction formulae

$$\begin{aligned}\theta^\alpha\theta^\beta &= -\frac{1}{2}(\theta\theta)\epsilon^{\alpha\beta} & \bar{\theta}^{\dot{\alpha}}\bar{\theta}^{\dot{\beta}} &= \frac{1}{2}(\bar{\theta}\bar{\theta})\epsilon^{\dot{\alpha}\dot{\beta}} & (\theta\xi)(\theta\chi) &= -\frac{1}{2}(\theta\theta)(\xi\chi) & (\theta\sigma^\nu\bar{\theta})\theta^\alpha &= \frac{1}{2}(\theta\theta)(\bar{\theta}\bar{\sigma}^\nu)^\alpha \\ \theta_\alpha\theta_\beta &= \frac{1}{2}(\theta\theta)\epsilon_{\alpha\beta} & \bar{\theta}_{\dot{\alpha}}\bar{\theta}_{\dot{\beta}} &= -\frac{1}{2}(\bar{\theta}\bar{\theta})\epsilon_{\dot{\alpha}\dot{\beta}} & (\bar{\theta}\xi)(\bar{\theta}\chi) &= -\frac{1}{2}(\bar{\theta}\bar{\theta})(\bar{\xi}\bar{\chi}) & (\theta\sigma^\nu\bar{\theta})\bar{\theta}_{\dot{\alpha}} &= -\frac{1}{2}(\theta\sigma^\nu)_{\dot{\alpha}}(\bar{\theta}\bar{\theta}) \\ \theta^\alpha\theta_\beta &= \frac{1}{2}(\theta\theta)\delta_\beta^\alpha & \bar{\theta}^{\dot{\alpha}}\bar{\theta}_{\dot{\beta}} &= \frac{1}{2}(\bar{\theta}\bar{\theta})\delta_{\dot{\beta}}^{\dot{\alpha}} & (\theta\sigma^\mu\bar{\theta})(\theta\sigma^\nu\bar{\theta}) &= \frac{1}{2}(\theta\theta)(\bar{\theta}\bar{\theta})\eta^{\mu\nu} \\ (\theta\sigma^\mu\bar{\sigma}^\nu\theta) &= (\theta\theta)\eta^{\mu\nu} & (\bar{\theta}\bar{\sigma}^\mu\sigma^\nu\bar{\theta}) &= (\bar{\theta}\bar{\theta})\eta^{\mu\nu} & (\sigma^\mu\bar{\theta})_\alpha(\theta\sigma^\nu\bar{\theta}) &= \frac{1}{2}(\bar{\theta}\bar{\theta})(\sigma^\mu\bar{\sigma}^\nu\theta)_\alpha\end{aligned}$$

$$\begin{aligned}\sigma^\mu\bar{\sigma}^\nu &= \eta^{\mu\nu} - 2i\sigma^{\mu\nu} & \sigma^\mu\bar{\sigma}^\rho\sigma^\nu + \sigma^\nu\bar{\sigma}^\rho\sigma^\mu &= 2(\sigma^\mu\eta^{\nu\rho} + \sigma^\nu\eta^{\mu\rho} - \sigma^\rho\eta^{\mu\nu}) \\ \bar{\sigma}^\mu\sigma^\nu &= \eta^{\mu\nu} - 2i\bar{\sigma}^{\mu\nu} & \sigma^\mu\bar{\sigma}^\nu\sigma^\rho - \sigma^\rho\bar{\sigma}^\nu\sigma^\mu &= 2i\sigma_\sigma\epsilon^{\mu\nu\rho\sigma} \\ \text{Tr}(\sigma^\mu\bar{\sigma}^\nu) &= \text{Tr}(\bar{\sigma}^\mu\sigma^\nu) = 2\eta^{\mu\nu} & \bar{\sigma}^\mu\sigma^\rho\bar{\sigma}^\nu + \bar{\sigma}^\nu\sigma^\rho\bar{\sigma}^\mu &= 2(\bar{\sigma}^\mu\eta^{\nu\rho} + \bar{\sigma}^\nu\eta^{\mu\rho} - \bar{\sigma}^\rho\eta^{\mu\nu}) \\ \sigma_{\alpha\dot{\alpha}}^\mu\bar{\sigma}_{\dot{\beta}\beta}^\mu &= 2\delta_{\dot{\alpha}}^{\dot{\beta}}\delta_\alpha^\beta & \bar{\sigma}^\mu\sigma^\nu\bar{\sigma}^\rho - \bar{\sigma}^\rho\sigma^\nu\bar{\sigma}^\mu &= -2i\bar{\sigma}_\sigma\epsilon^{\mu\nu\rho\sigma} \\ \sigma_{\mu\alpha\dot{\alpha}}\sigma_{\beta\dot{\beta}}^\mu &= 2\epsilon_{\alpha\beta}\epsilon_{\dot{\alpha}\dot{\beta}} & \epsilon_{\dot{\beta}\dot{\alpha}}\bar{\sigma}^{\mu\dot{\alpha}\alpha} &= \epsilon_{\dot{\beta}\dot{\alpha}}\epsilon^{\dot{\alpha}\dot{\gamma}}\epsilon^{\alpha\gamma}\sigma_{\gamma\dot{\gamma}}^\mu = \epsilon^{\alpha\gamma}\sigma_{\gamma\dot{\beta}}^\mu \\ \bar{\sigma}_{\mu}{}^{\dot{\alpha}\alpha}\bar{\sigma}^{\mu\dot{\beta}\beta} &= 2\epsilon^{\alpha\beta}\epsilon^{\dot{\alpha}\dot{\beta}} & \epsilon_{\beta\alpha}\bar{\sigma}^{\mu\dot{\alpha}\alpha} &= \epsilon_{\beta\alpha}\epsilon^{\dot{\alpha}\dot{\gamma}}\epsilon^{\alpha\gamma}\sigma_{\gamma\dot{\gamma}}^\mu = \epsilon^{\dot{\alpha}\dot{\gamma}}\sigma_{\beta\dot{\gamma}}^\mu \\ \text{Tr}(\sigma^{\mu\nu}) &= \text{Tr}(\bar{\sigma}^{\mu\nu}) = 0 & \text{Tr}(\sigma^{\mu\nu}\sigma^{\rho\sigma}) &= \frac{1}{2}(\eta^{\mu\rho}\eta^{\nu\sigma} - \eta^{\mu\sigma}\eta^{\nu\rho}) - \frac{1}{2}i\epsilon^{\mu\nu\rho\sigma} \\ \bar{\sigma}^{\mu\nu} &= -\bar{\sigma}^{\nu\mu} \quad \sigma^{\mu\nu} = -\sigma^{\nu\mu} & \text{Tr}(\bar{\sigma}^{\mu\nu}\bar{\sigma}^{\rho\sigma}) &= \frac{1}{2}i\epsilon^{\mu\nu\rho\sigma} + \frac{1}{2}(\eta^{\mu\rho}\eta^{\nu\sigma} - \eta^{\mu\sigma}\eta^{\nu\rho}) \\ \sigma^{\mu\nu}{}_\alpha{}^\beta\epsilon_{\beta\gamma} &= \sigma^{\mu\nu}{}_\gamma{}^\beta\epsilon_{\beta\alpha} & \sigma_{\alpha\dot{\alpha}}^\mu\sigma_{\beta\dot{\beta}}^\nu - \sigma_{\alpha\dot{\alpha}}^\nu\sigma_{\beta\dot{\beta}}^\mu &= -2i\epsilon_{\dot{\alpha}\dot{\gamma}}\bar{\sigma}^{\mu\dot{\gamma}\gamma}\epsilon_{\alpha\beta} - 2i\sigma^{\mu\nu}{}_\alpha{}^\gamma\epsilon_{\gamma\beta}\epsilon_{\dot{\alpha}\dot{\beta}} \\ \bar{\sigma}^{\mu\nu}{}_{\dot{\beta}}{}^{\dot{\alpha}}\epsilon^{\dot{\beta}\dot{\gamma}} &= \bar{\sigma}^{\mu\nu}{}_{\dot{\gamma}}{}^{\dot{\alpha}}\epsilon^{\dot{\beta}\dot{\gamma}} & \sigma_{\alpha\dot{\alpha}}^\mu\sigma_{\beta\dot{\beta}}^\nu + \sigma_{\alpha\dot{\alpha}}^\nu\sigma_{\beta\dot{\beta}}^\mu &= 4\sigma^{\rho\mu}{}_\alpha{}^\gamma\epsilon_{\gamma\beta}\epsilon_{\dot{\alpha}\dot{\gamma}}\bar{\sigma}^{\nu\dot{\gamma}\dot{\beta}}\eta_{\rho\sigma} + \epsilon_{\alpha\beta}\epsilon_{\dot{\alpha}\dot{\beta}}\eta^{\mu\nu} \\ \bar{\sigma}_{\rho\sigma}\epsilon^{\mu\nu\rho\sigma} &= -2i\bar{\sigma}^{\mu\nu} & \bar{\sigma}^{\mu\dot{\alpha}\alpha}\bar{\sigma}^{\nu\dot{\beta}\beta} - \bar{\sigma}^{\nu\dot{\alpha}\alpha}\bar{\sigma}^{\mu\dot{\beta}\beta} &= -2i\bar{\sigma}^{\mu\nu}{}_{\dot{\gamma}}{}^{\dot{\alpha}}\epsilon^{\dot{\beta}\dot{\gamma}}\epsilon^{\alpha\beta} - 2i\epsilon^{\alpha\gamma}\sigma^{\mu\nu}{}_\gamma{}^\beta\epsilon_{\dot{\alpha}\dot{\beta}} \\ \sigma_{\rho\sigma}\epsilon^{\mu\nu\rho\sigma} &= 2i\sigma^{\mu\nu} & \bar{\sigma}^{\mu\dot{\alpha}\alpha}\bar{\sigma}^{\nu\dot{\beta}\beta} + \bar{\sigma}^{\nu\dot{\alpha}\alpha}\bar{\sigma}^{\mu\dot{\beta}\beta} &= 4\epsilon^{\alpha\gamma}\sigma^{\sigma\nu}{}_\gamma{}^\beta\bar{\sigma}^{\rho\mu\dot{\alpha}}{}_{\dot{\gamma}}\epsilon^{\dot{\beta}\dot{\gamma}}\eta_{\rho\sigma} + \epsilon^{\alpha\beta}\epsilon^{\dot{\alpha}\dot{\beta}}\eta^{\mu\nu}\end{aligned}$$

$$\begin{aligned}\bar{\xi}\bar{\sigma}^\mu\chi &= -\chi\sigma^\mu\bar{\xi} & \bar{\xi}\bar{\sigma}^\mu\sigma^\nu\bar{\chi} &= \bar{\chi}\bar{\sigma}^\nu\sigma^\mu\bar{\xi} & \xi\sigma^\mu\bar{\sigma}^\nu\chi &= \chi\sigma^\nu\bar{\sigma}^\mu\xi & \bar{\xi}\bar{\sigma}^\mu\sigma^\nu\bar{\sigma}^\rho\chi &= -\chi\sigma^\rho\bar{\sigma}^\nu\sigma^\mu\bar{\xi} \\ (\xi\sigma^\mu\bar{\chi})^* &= \chi\sigma^\mu\bar{\xi} & (\bar{\xi}\bar{\sigma}^\mu\chi)^* &= \bar{\chi}\bar{\sigma}^\mu\xi & (\bar{\chi}\bar{\sigma}^\mu\sigma^\nu\bar{\xi})^* &= \xi\sigma^\nu\bar{\sigma}^\mu\chi & (\xi[\sigma_s]\chi)^* &= \bar{\chi}[\sigma_{\text{sreversed}}]\bar{\xi} \\ (\xi\chi)\psi^\alpha &= -(\psi\xi)\chi^\alpha - (\psi\chi)\xi^\alpha & (\xi\chi)\bar{\psi}_{\dot{\alpha}} &= \frac{1}{2}(\xi\sigma^\mu\bar{\psi})(\chi\sigma_\mu)_{\dot{\alpha}}\end{aligned}$$

### Superfields

<sup>\*2</sup>As the definition of  $\sigma^{\mu\nu}$  and  $\bar{\sigma}^{\mu\nu}$  are not unified in literature, they are not used in this CheatSheet except for this page.

#### 4.1. Lorentz symmetry as $SU(2) \times SU(2)$

#### 4.2. Derivative in superspace

$$y^\mu := x^\mu - i(\theta\sigma^\mu\bar{\theta}) \quad (4.1)$$

$$\phi(y) = \phi(x) - i(\theta\sigma^\mu\bar{\theta})\partial_\mu\phi(x) - \frac{1}{4}\theta^4\partial^2\phi(x) \quad (4.2)$$

#### 4.3. Superfields

##### Chiral superfield

$$\Phi = \phi(y) + \sqrt{2}\theta\psi(y) + \theta^2 F(y) \quad (4.3)$$

$$= \phi(x) + \sqrt{2}\theta\psi(x) - i\partial_\mu\phi(x)(\theta\sigma^\mu\bar{\theta}) + F(x)\theta^2 + \frac{i}{\sqrt{2}}(\partial_\mu\psi(x)\sigma^\mu\bar{\theta})\theta^2 - \frac{1}{4}\partial^2\phi(x)\theta^4 \quad (4.4)$$

$$\Phi^* = \phi^*(x) + \sqrt{2}\bar{\psi}(x)\bar{\theta} + F^*(x)\bar{\theta}^2 + i\partial_\mu\phi^*(x)(\theta\sigma^\mu\bar{\theta}) - \frac{i}{\sqrt{2}}[\theta\sigma^\mu\partial_\mu\bar{\psi}(x)]\bar{\theta}^2 - \frac{1}{4}\partial^2\phi^*(x)\theta^4 \quad (4.5)$$

$$\begin{aligned} \Phi_i^*\Phi_j &= \phi_i^*\phi_j + \sqrt{2}\phi_i^*(\theta\psi_j) + \sqrt{2}(\bar{\psi}_i\bar{\theta})\phi_j + \phi_i^*F_j\theta^2 + 2(\bar{\psi}_i\bar{\theta})(\theta\psi_j) - i(\phi_i^*\partial_\mu\phi_j - \partial_\mu\phi_i^*\phi_j)(\theta\sigma^\mu\bar{\theta}) + F_i^*\phi_j\bar{\theta}^2 \\ &+ \left[ \sqrt{2}\bar{\psi}_i\bar{\theta}F_j - \frac{i(\partial_\mu\phi_i^* \cdot \psi_j\sigma^\mu\bar{\theta} - \phi_i^*\partial_\mu\psi_j\sigma^\mu\bar{\theta})}{\sqrt{2}} \right] \theta^2 + \left[ \sqrt{2}F_i^*\theta\psi_j + \frac{i(\theta\sigma^\mu\bar{\psi}_i\partial_\mu\phi_j - \theta\sigma^\mu\partial_\mu\bar{\psi}_i\phi_j)}{\sqrt{2}} \right] \bar{\theta}^2 \\ &+ \frac{1}{4}(4F_i^*F_j - \phi_i^*\partial^2\phi_j - (\partial^2\phi_i^*)\phi_j + 2(\partial_\mu\phi_i^*)(\partial^\mu\phi_j) + 2i(\psi_j\sigma^\mu\partial_\mu\bar{\psi}_i) - 2i(\partial_\mu\psi_j\sigma^\mu\bar{\psi}_i))\theta^4 \end{aligned} \quad (4.6)$$

$$\begin{aligned} &\equiv \phi_i^*\phi_j + \sqrt{2}\phi_i^*(\theta\psi_j) + \sqrt{2}(\bar{\psi}_i\bar{\theta})\phi_j + \phi_i^*F_j\theta^2 + 2(\bar{\psi}_i\bar{\theta})(\theta\psi_j) - 2i(\phi_i^*\partial_\mu\phi_j)(\theta\sigma^\mu\bar{\theta}) + F_i^*\phi_j\bar{\theta}^2 \\ &+ \sqrt{2}(\bar{\psi}_i\bar{\theta}F_j + i\phi_i^*\partial_\mu\psi_j\sigma^\mu\bar{\theta})\theta^2 + \sqrt{2}(F_i^*\theta\psi_j - i\theta\sigma^\mu\partial_\mu\bar{\psi}_i\phi_j)\bar{\theta}^2 \\ &+ (F_i^*F_j + (\partial_\mu\phi_i^*)(\partial^\mu\phi_j) + i\bar{\psi}_i\sigma^\mu\partial_\mu\psi_j)\theta^4 \end{aligned} \quad (4.7)$$

$$\Phi_i\Phi_j \Big|_{\theta^2} = -\psi_i\psi_j + F_i\phi_j + \phi_iF_j \quad (4.8)$$

$$\Phi_i\Phi_j\Phi_k \Big|_{\theta^2} = -(\psi_i\psi_j)\phi_k - (\psi_k\psi_i)\phi_j - (\psi_j\psi_k)\phi_i + \phi_i\phi_jF_k + \phi_k\phi_iF_j + \phi_j\phi_kF_i \quad (4.9)$$

$$e^{k\Phi} = e^{k\phi} \left[ 1 + \sqrt{2}k\theta\psi + \left( kF - \frac{k^2}{2}\psi\psi \right) \theta^2 - ik\partial_\mu\phi(\theta\sigma^\mu\bar{\theta}) + \frac{ik(\partial_\mu\psi + k\psi\partial_\mu\phi)\sigma^\mu\bar{\theta}\theta^2}{\sqrt{2}} - \frac{k}{4}(\partial^2\phi + k\partial_\mu\phi\partial^\mu\phi)\theta^4 \right] \quad (4.10)$$

## 5. Minimal Supersymmetric Standard Model

Gauge symmetry:  $SU(3)_{\text{color}} \times SU(2)_{\text{weak}} \times U(1)_Y$

Particle content:

(a) Chiral superfields						(b) Vector superfields			
	SU(3)	SU(2)	U(1)	$B$	$L$	scalar/spinor	SU(3)	SU(2)	U(1) ino/boson
$Q_i$	<b>3</b>	<b>2</b>	1/6	1/3		$\tilde{q}_L, q_L \rightarrow (u_L, d_L)$	$g$	adj.	$\tilde{g}, g_\mu$
$L_i$		<b>2</b>	-1/2		1	$\tilde{l}_L, l_L \rightarrow (\nu_L, l_L)$	$W$		$\tilde{w}, W_\mu$
$U_i^c$	$\bar{\mathbf{3}}$		-2/3	-1/3		$\tilde{u}_R^c, u_R^c$	$B$	adj.	$\tilde{b}, B_\mu$
$D_i^c$	$\bar{\mathbf{3}}$		1/3	-1/3		$\tilde{d}_R^c, d_R^c$			
$E_i^c$			1		-1	$\tilde{e}_R^c, e_R^c$			
$H_u$		<b>2</b>	1/2			$h_u, \tilde{h}_u \rightarrow (h_u^+, h_u^0)$			
$H_d$		<b>2</b>	-1/2			$h_d, \tilde{h}_d \rightarrow (h_d^0, h_d^-)$			

Here, each of the column groups shows (from left to right) superfield name, charges for the gauge symmetries, other quantum numbers if relevant, and notation for corresponding fields (and SU(2) decomposition).

**“c”-notation** For scalars,  $\tilde{\phi}_R^c := \phi_R^* = C\phi_R C$  (because the intrinsic phase for  $C$  is +1 for quarks and leptons.)

For matter spinors,  $\psi_R^c := \bar{\psi}_R$  (and  $\psi_R = \bar{\psi}_R^c$ ); Dirac spinors are thus

$$\psi_L = \begin{pmatrix} \psi_L \\ 0 \end{pmatrix}, \quad \bar{\psi}_L = (0 \quad \bar{\psi}_L), \quad \psi_R^c := \begin{pmatrix} \psi_R^c \\ 0 \end{pmatrix} = C \begin{pmatrix} 0 \\ \psi_R \end{pmatrix} = C\psi_R, \quad \bar{\psi}_R^c = (0 \quad \psi_R) = (\bar{\psi}_R \quad 0) C = \bar{\psi}_R C.$$

### Superpotential and SUSY-terms

$$W_{\text{RPC}} = \mu H_u H_d - y_{uij} U_i^c H_u Q_j + y_{dij} D_i^c H_d Q_j + y_{eij} E_i^c H_d L_j, \quad (5.1)$$

$$W_{\text{RPV}} = -\kappa_i L_i H_u + \frac{1}{2} \lambda_{ijk} L_i L_j E_k^c + \lambda'_{ijk} L_i Q_j D_k^c + \frac{1}{2} \lambda''_{ijk} U_i^c D_j^c D_k^c, \quad (5.2)$$

$$\mathcal{L}_{\text{SUSY}} = -\frac{1}{2} \left( M_3 \tilde{g}\tilde{g} + M_2 \tilde{w}\tilde{w} + M_1 \tilde{b}\tilde{b} + \text{H.c.} \right) - V_{\text{SUSY}}, \quad (5.3)$$

$$\begin{aligned} V_{\text{SUSY}}^{\text{RPC}} = & \left( \tilde{q}_L^* m_Q^2 \tilde{q}_L + \tilde{l}_L^* m_L^2 \tilde{l}_L + \tilde{u}_R^* m_{U^c}^2 \tilde{u}_R + \tilde{d}_R^* m_{D^c}^2 \tilde{d}_R + \tilde{e}_R^* m_{E^c}^2 \tilde{e}_R + m_{H_u}^2 |h_u|^2 + m_{H_d}^2 |h_d|^2 \right) \\ & + \left( -\tilde{u}_R^* h_u a_u \tilde{q}_L + \tilde{d}_R^* h_d a_d \tilde{q}_L + \tilde{e}_R^* h_d a_e \tilde{l}_L + b H_u H_d + \text{H.c.} \right) \\ & + \left( -\tilde{u}_R^* h_u^* c_u \tilde{q}_L + \tilde{d}_R^* h_u^* c_d \tilde{q}_L + \tilde{e}_R^* h_u^* c_e \tilde{l}_L + \text{H.c.} \right), \end{aligned} \quad (5.4)$$

$$\begin{aligned} V_{\text{SUSY}}^{\text{RPV}} = & \left( -b_i \tilde{l}_{Li} H_u + \frac{1}{2} T_{ijk} \tilde{l}_{Li} \tilde{l}_{Lj} \tilde{e}_{Rk}^* + T'_{ijk} \tilde{l}_{Li} \tilde{q}_{Lj} \tilde{d}_{Rk}^* + \frac{1}{2} T''_{ijk} \tilde{u}_{Ri}^* \tilde{d}_{Rj}^* \tilde{d}_{Rk}^* + \tilde{l}_{Li}^* M_{Li}^2 H_d + \text{H.c.} \right) \\ & + \left( C_{ijk}^1 \tilde{l}_{Li}^* \tilde{q}_{Lj} \tilde{u}_{Rk}^* + C_i^2 h_u^* h_d \tilde{e}_{Ri}^* + C_{ijk}^3 \tilde{d}_{Ri} \tilde{u}_{Rj}^* \tilde{e}_{Rk}^* + \frac{1}{2} C_{ijk}^4 \tilde{d}_{Ri} \tilde{q}_{Lj} \tilde{q}_{Lk} + \text{H.c.} \right), \end{aligned} \quad (5.5)$$

$$(\lambda_{ijk} = -\lambda_{jik}, \lambda''_{ijk} = -\lambda''_{ikj}, \text{ and } C_{ijk}^4 = C_{ikj}^4.)$$

We follow the notation of DHM [?, PhysRept] and Martin [?, v7] (but note that Martin uses  $(-, +, +, +)$ -metric) for RPC part and SLHA2 convention for RPV part.

## 5.1. Scalar potential

The MSSM scalar potential has contributions from  $F$ -terms and  $D$ -terms:

$$V_{\text{SUSY}} = F_i^* F_i + \frac{1}{2} D^a D^a; \quad F_i = -W_i^* = -\frac{\delta W^*}{\delta \phi_i^*}, \quad D^a = -g(\phi^* T^a \phi), \quad (5.6)$$

where  $T_a$  corresponds to the gauge-symmetry generator relevant for each  $\phi$ . They are given by

$$-F_{h_u^a}^* = \epsilon^{ab} \left( -\tilde{u}_R^{x*} y_u \tilde{q}_L^b + \mu h_d^b + \mu'_i \tilde{l}_{Li}^b \right), \quad (5.7)$$

$$-F_{h_d^a}^* = \epsilon^{ab} \left( \tilde{e}_R^{x*} y_e \tilde{l}_L^b + \tilde{d}_R^{x*} y_d \tilde{q}_L^b - \mu h_u^b \right), \quad (5.8)$$

$$-F_{\tilde{q}_{Li}^{ax}}^* = \epsilon^{ab} \left( -y_{dij} h_d^b \tilde{d}_{Rj}^{x*} + y_{ujj} h_u^b \tilde{u}_{Rj}^{x*} - \lambda'_{kij} \tilde{d}_{Rj}^{x*} \tilde{l}_{Lk}^b \right), \quad (5.9)$$

$$-F_{\tilde{u}_{Ri}^{xx}}^* = -y_{uij} h_u \tilde{q}_{Lj}^x + \frac{1}{2} \epsilon^{xyz} \lambda''_{ijk} \tilde{d}_{Rj}^{y*} \tilde{d}_{Rk}^{z*}, \quad (5.10)$$

$$-F_{\tilde{d}_{Ri}^{xx}}^* = y_{dij} h_d \tilde{q}_{Lj}^x + \lambda'_{jki} \tilde{l}_{Lj} \tilde{q}_{Lk}^x - \lambda''_{jik} \epsilon^{xyz} \tilde{u}_{Rj}^{y*} \tilde{d}_{Rk}^{z*}, \quad (5.11)$$

$$-F_{\tilde{l}_{Li}^a}^* = \epsilon^{ab} \left( -y_{eji} \tilde{e}_{Rj}^* h_d^b - \mu'_i h_u^b + \lambda_{ijk} \tilde{l}_{Lj}^b \tilde{e}_{Rk}^* + \lambda'_{ijk} \tilde{q}_{Lj}^{bx} \tilde{d}_{Rk}^{x*} \right), \quad (5.12)$$

$$-F_{\tilde{e}_{Ri}^*}^* = y_{eij} h_d \tilde{l}_{Lj} + \frac{1}{2} \lambda_{jki} \tilde{l}_{Lj} \tilde{l}_{Lk}. \quad (5.13)$$

$$D_{\text{SU}(3)}^\alpha = -g_3 \sum_{i=1}^3 \left( \sum_{a=1,2} \tilde{q}_{Li}^{a*} \tau^\alpha \tilde{q}_{Li}^a - \tilde{u}_{Ri}^* \tau^\alpha \tilde{u}_{Ri} - \tilde{d}_{Ri}^* \tau^\alpha \tilde{d}_{Ri} \right), \quad (5.14)$$

$$D_{\text{SU}(2)}^\alpha = -g_2 \left[ \sum_{i=1}^3 \left( \sum_{x=1}^3 \tilde{q}_{Li}^{x*} T^\alpha \tilde{q}_{Li}^x + \tilde{l}_{Li}^* T^\alpha \tilde{l}_{Li} \right) + h_u^* T^\alpha h_u + h_d^* T^\alpha h_d \right], \quad (5.15)$$

$$D_{\text{U}(1)} = -g_1 \left( \frac{1}{6} |\tilde{q}_L|^2 - \frac{1}{2} |\tilde{l}_L|^2 - \frac{2}{3} |\tilde{u}_R|^2 + \frac{1}{3} |\tilde{d}_R|^2 + |\tilde{e}_R|^2 + \frac{1}{2} |h_u|^2 - \frac{1}{2} |h_d|^2 \right). \quad (5.16)$$



Combining these, the full SUSY scalar potential is given by

$$\begin{aligned}
V &= V_{\text{SUSY}} + V_{\text{SUSY}}; \tag{5.17} \\
V_{\text{SUSY}} &= |h_u|^2 (|\mu|^2 + |\mu'|^2) + |\mu|^2 |h_d|^2 + \left( \mu_i^* \tilde{\mu}_{Li}^* h_d + \text{H.c.} \right) + \mu_i^* \mu_j' \tilde{\mu}_{Li}^* \tilde{\mu}_{Lj} \\
&+ \left[ -y_{uij} \mu_i^* \tilde{u}_d^* \tilde{u}_{Ri}^* \tilde{q}_{Lj} - y_{uij} \mu_k' \tilde{u}_{Ri}^* \tilde{q}_{Lj} \tilde{t}_{Lk}^* - (y_{dij} \mu_i^* + \lambda_{kji}' \mu_k') h_u^* \tilde{d}_{Ri}^* \tilde{q}_{Lj} \right. \\
&\quad \left. + y_{eij} \mu_j' \tilde{e}_{Ri}^* h_u^* h_d + (\lambda_{jki} \mu_k' - y_{eij} \mu_i^*) h_u^* \tilde{e}_{Ri}^* \tilde{l}_{Lj} + \text{H.c.} \right] \\
&+ \frac{1}{8} (g_2^2 + g_Y^2) |h_d|^4 + \frac{1}{8} (g_2^2 + g_Y^2) |h_u|^4 + \left( -\frac{g_2^2}{4} |h_d|^2 |h_u|^2 - \frac{g_Y^2}{4} |h_d|^2 |h_u|^2 + \frac{g_2^2}{2} |h_d^* h_u|^2 \right) \\
&+ \left( -\frac{g_2^2}{4} |h_u|^2 |\tilde{q}_L|^2 + \frac{g_Y^2}{12} |h_u|^2 |\tilde{q}_L|^2 + \frac{g_2^2}{2} |h_u^b \tilde{q}_{Li}|^2 + \epsilon^{ac} \epsilon^{bd} (y_u^\dagger y_u)_{ji} h_u^a h_u^{b*} \tilde{q}_{Li}^c \tilde{q}_{Lj}^{d*} \right) \\
&+ \left( -\frac{g_2^2}{4} |h_d|^2 |\tilde{q}_L|^2 - \frac{g_Y^2}{12} |h_d|^2 |\tilde{q}_L|^2 + \frac{g_2^2}{2} |h_d^* \tilde{q}_{Li}|^2 + \epsilon^{ac} \epsilon^{bd} (y_d^\dagger y_d)_{ji} h_d^a h_d^{b*} \tilde{q}_{Li}^c \tilde{q}_{Lj}^{d*} \right) \\
&+ \left( -\frac{g_2^2}{4} |h_u|^2 |\tilde{l}_L|^2 - \frac{g_Y^2}{4} |h_u|^2 |\tilde{l}_L|^2 + \frac{g_2^2}{2} |h_u^* \tilde{l}_{Li}|^2 \right) \\
&+ \left( -\frac{g_2^2}{4} |h_d|^2 |\tilde{l}_L|^2 + \frac{g_Y^2}{4} |h_d|^2 |\tilde{l}_L|^2 + \frac{g_2^2}{2} |h_d^* \tilde{l}_{Li}|^2 + \epsilon^{ac} \epsilon^{bd} (y_e^\dagger y_e)_{ji} h_d^a h_d^{b*} \tilde{l}_{Li}^c \tilde{l}_{Lj}^{d*} \right) \\
&+ |h_u|^2 \left( -\frac{g_Y^2}{3} |\tilde{u}_R|^2 + (y_u y_u^\dagger)_{ij} \tilde{u}_{Ri}^* \tilde{u}_{Rj} \right) + \frac{g_Y^2}{3} |h_d|^2 |\tilde{u}_R|^2 \\
&+ \frac{g_Y^2}{6} |h_u|^2 |\tilde{d}_R|^2 + |h_d|^2 \left( -\frac{g_Y^2}{6} |\tilde{d}_R|^2 + (y_d y_d^\dagger)_{ij} \tilde{d}_{Ri}^* \tilde{d}_{Rj} \right) - \left[ (y_u y_d^\dagger)_{ij} \tilde{u}_{Ri}^* \tilde{d}_{Rj} (h_d^* h_u) + \text{H.c.} \right] \\
&+ \frac{g_Y^2}{2} |h_u|^2 |\tilde{e}_R|^2 + |h_d|^2 \left( -\frac{g_Y^2}{2} |\tilde{e}_R|^2 + (y_e y_e^\dagger)_{ij} \tilde{e}_{Ri}^* \tilde{e}_{Rj} \right) \\
&+ \left[ -\frac{1}{2} \epsilon^{ab} \epsilon^{xyz} y_{ulk} \lambda_{lij}'' h_u^a \tilde{d}_{Ri}^x \tilde{q}_{Lj}^y \tilde{q}_{Lk}^{bz} + \epsilon^{ab} \epsilon^{xyz} y_{dlk} \lambda_{ijl}'' h_d^a \tilde{u}_{Ri}^x \tilde{q}_{Lj}^y \tilde{q}_{Lk}^{bz} - y_{uil} \lambda_{klj}' h_u^a \tilde{u}_{Ri}^* \tilde{d}_{Rj} \tilde{l}_{Lk}^{a*} \right. \\
&\quad \left. + y_{dil} \lambda_{klj}' h_d^a \tilde{d}_{Ri}^* \tilde{d}_{Rj} \tilde{l}_{Lk}^{a*} - \epsilon^{ab} \epsilon^{cd} y_{dli} \lambda_{kjl}' h_d^a \tilde{q}_{Li}^b \tilde{q}_{Lj}^{c*} \tilde{l}_{Lk}^{d*} + y_{eil} \lambda_{klj}' h_d^a \tilde{e}_{Ri}^* \tilde{e}_{Rj} \tilde{l}_{Lk}^{a*} \right. \\
&\quad \left. - y_{ejl} \lambda_{kli}' h_d^a \tilde{e}_{Ri}^* \tilde{d}_{Rj} \tilde{q}_{Lk}^{a*} + \frac{1}{2} \epsilon^{ab} \epsilon^{cd} y_{eli} \lambda_{jkl}' h_d^a \tilde{l}_{Li}^b \tilde{l}_{Lj}^{c*} \tilde{l}_{Lk}^{d*} + \text{H.c.} \right] \\
&+ \left[ \left( -\frac{g_3^2}{12} + \frac{g_Y^2}{72} - \frac{g_2^2}{8} \right) |\tilde{q}_L|^4 + \frac{g_2^2}{4} \tilde{q}_{Li}^{ax} \tilde{q}_{Lj}^{by} \tilde{q}_{Li}^{bx*} \tilde{q}_{Lj}^{ay*} + \frac{g_3^2}{4} \tilde{q}_{Li}^{ax} \tilde{q}_{Lj}^{by} \tilde{q}_{Li}^{ay*} \tilde{q}_{Lj}^{bx*} \right] \\
&+ \left[ \left( -\frac{g_3^2}{12} + \frac{2g_Y^2}{9} \right) |\tilde{u}_R|^4 + \frac{g_3^2}{4} \tilde{u}_{Ri}^{x*} \tilde{u}_{Rj}^{y*} \tilde{u}_{Ri}^y \tilde{u}_{Rj}^x \right] \\
&+ \left[ \left( -\frac{g_3^2}{12} + \frac{g_Y^2}{18} \right) |\tilde{d}_R|^4 + \frac{g_3^2}{4} \tilde{d}_{Ri}^{x*} \tilde{d}_{Rj}^{y*} \tilde{d}_{Ri}^y \tilde{d}_{Rj}^x + \frac{1}{2} \lambda_{mij}'' \lambda_{mkl}'' \tilde{d}_{Ri}^{x*} \tilde{d}_{Rj}^{y*} \tilde{d}_{Rk}^x \tilde{d}_{Rl}^y \right] \\
&+ \left[ \left( \frac{g_3^2}{6} - \frac{g_Y^2}{9} \right) |\tilde{u}_R|^2 |\tilde{q}_L|^2 - \frac{g_3^2}{2} \tilde{u}_{Ri}^{x*} \tilde{q}_{Lj}^{ax} \tilde{u}_{Ri}^y \tilde{q}_{Lj}^{ay*} + y_{uik} y_{ujl} \tilde{u}_{Ri}^{x*} \tilde{u}_{Rj}^y \tilde{q}_{Lk}^{ax} \tilde{q}_{Ll}^{ay*} \right] \\
&+ \left[ \left( \frac{g_3^2}{6} + \frac{g_Y^2}{18} \right) |\tilde{d}_R|^2 |\tilde{q}_L|^2 - \frac{g_3^2}{2} \tilde{d}_{Ri}^{x*} \tilde{q}_{Lj}^{ax} \tilde{d}_{Ri}^y \tilde{q}_{Lj}^{ay*} + (y_{dik} y_{djl}^* + \lambda_{mki}' \lambda_{mlj}') \tilde{d}_{Ri}^{x*} \tilde{q}_{Lk}^{ax} \tilde{d}_{Rj}^y \tilde{q}_{Ll}^{ay*} \right] \\
&+ \left[ -\left( \frac{g_3^2}{6} + \frac{2g_Y^2}{9} \right) |\tilde{d}_R|^2 |\tilde{u}_R|^2 + \left( \frac{g_3^2}{2} - \lambda_{ikm}'' \lambda_{ljm}'' \right) \tilde{u}_{Ri}^{x*} \tilde{d}_{Rj}^x \tilde{d}_{Rj}^{y*} \tilde{u}_{Ri}^y + \lambda_{ikm}'' \lambda_{jlm}'' \tilde{u}_{Ri}^{x*} \tilde{u}_{Rj}^x \tilde{d}_{Rk}^{y*} \tilde{d}_{Rl}^y \right] \\
&+ \left[ -\left( \frac{g_2^2}{4} + \frac{g_Y^2}{12} \right) |\tilde{l}_L|^2 |\tilde{q}_L|^2 + \frac{g_2^2}{2} \tilde{q}_{Li}^{ax} \tilde{q}_{Li}^{bx*} \tilde{l}_{Lj}^b \tilde{l}_{Lj}^{a*} + \epsilon^{ac} \epsilon^{bd} \lambda_{kim}' \lambda_{ljm}' \tilde{q}_{Li}^{ax} \tilde{q}_{Lj}^{bx*} \tilde{l}_{Lk}^c \tilde{l}_{Ll}^{d*} \right] + \frac{g_Y^2}{6} |\tilde{e}_R|^2 |\tilde{q}_L|^2 \\
&+ \left( -\frac{g_Y^2}{6} |\tilde{d}_R|^2 |\tilde{l}_L|^2 + \lambda_{lmi}' \lambda_{kmj}' \tilde{d}_{Ri}^* \tilde{d}_{Rj} \tilde{l}_{Lk}^* \tilde{l}_{Ll} \right) + \frac{g_Y^2}{3} |\tilde{u}_R|^2 |\tilde{l}_L|^2 - \frac{2g_Y^2}{3} |\tilde{u}_R|^2 |\tilde{e}_R|^2 + \frac{g_Y^2}{3} |\tilde{d}_R|^2 |\tilde{e}_R|^2 \\
&+ \left[ \left( -\frac{g_2^2}{8} + \frac{g_Y^2}{8} \right) |\tilde{l}_L|^4 + \frac{g_2^2}{4} \tilde{l}_{Li}^a \tilde{l}_{Lj}^b \tilde{l}_{Li}^{b*} \tilde{l}_{Lj}^{a*} + \frac{1}{4} \epsilon^{ab} \epsilon^{cd} \lambda_{ijm} \lambda_{klm}^* \tilde{l}_{Li}^a \tilde{l}_{Lj}^b \tilde{l}_{Lk}^{c*} \tilde{l}_{Ll}^{d*} \right] \\
&+ \frac{g_Y^2}{2} |\tilde{e}_R|^4 + \left[ -\frac{g_Y^2}{2} |\tilde{e}_R|^2 |\tilde{l}_L|^2 + (y_{eik} y_{ejl}^* + \lambda_{kmi} \lambda_{lmj}^*) \tilde{e}_{Ri}^* \tilde{e}_{Rj} \tilde{l}_{Lk}^* \tilde{l}_{Ll} \right] \\
&+ \left[ (y_{dik} y_{ejl}^* - \lambda_{mki}' \lambda_{lmj}') \tilde{d}_{Ri}^{x*} \tilde{q}_{Lk}^{ax} \tilde{e}_{Rj} \tilde{l}_{Ll}^{a*} - \epsilon^{ab} \epsilon^{xyz} \lambda_{lkm}' \lambda_{ijm}'' \tilde{l}_{Li}^b \tilde{q}_{Lk}^{az} \tilde{d}_{Rj}^y \tilde{u}_{Ri}^x + \text{H.c.} \right].
\end{aligned}
\tag{5.18}$$

## 5.2. SLHA convention

The SLHA convention [?] is different from our notation; the reinterpretation rules for the MSSM parameters are given in the right table (**magenta color** for objects in other conventions), while

$\mu, b, m_{Q,L,H_u,H_d}^2$ , RPV-trilinears ( $\lambda$ s and  $T$ s) are in common.

SLHA	our notation	Martin/DHM
$(H_1, H_2)$	$(H_d, H_u)$	
$Y_{u,d,e}$	$(y_{u,d,e})^T$	
$T_{u,d,e}$	$(a_{u,d,e})^T$	
$A_{u,d,e}$	$(A_{u,d,e})^T$	
$m_{U^c, D^c, E^c}^2$	$(m_{U^c, D^c, E^c}^2)^\dagger$	
$M_{1,2,3}$	$-M_{1,2,3}$	
$m_3^2$	$b$	
$m_A^2$	$m_{A_0}^2$ (tree)	
	$\kappa_i$	$= -\mu'_i$ (rarely used)
$D_i$	$b_i$	
$m_{L_i H_1}^2$	$M_{L_i}^2$	

In particular, the chargino/neutralino mass terms in RPC case are given by

$$\mathcal{L} \supset \left[ \frac{1}{2} \mathbf{M}_1 \tilde{b} \tilde{b} + \frac{1}{2} \mathbf{M}_2 \tilde{w} \tilde{w} - \mu \tilde{h}_u \tilde{h}_d - \frac{g_Y}{2\sqrt{2}} \left( h_u^* \tilde{h}_u - h_d^* \tilde{h}_d \right) \tilde{b} - \sqrt{2} g_2 \left( h_u^* T^a \tilde{h}_u + h_d^* T^a \tilde{h}_d \right) \tilde{w} \right] + \text{H.c.} \quad (5.19)$$

$$\rightarrow \frac{1}{2} \begin{pmatrix} \tilde{b} \\ \tilde{w} \\ h_u^0 \\ h_d^0 \end{pmatrix}^T \begin{pmatrix} -M_1 & 0 & -m_{ZC\beta S_w} & m_{ZS\beta S_w} \\ 0 & -M_2 & m_{ZC\beta C_w} & -m_{ZS\beta C_w} \\ -m_{ZC\beta S_w} & m_{ZC\beta C_w} & 0 & -\mu \\ m_{ZS\beta S_w} & -m_{ZS\beta C_w} & -\mu & 0 \end{pmatrix} \begin{pmatrix} \tilde{b} \\ \tilde{w} \\ h_u^0 \\ h_d^0 \end{pmatrix} \quad (5.20)$$

## A. Mathematics

### A.1. Matrix exponential

Excerpted from §2 and §5 of Hall 2015 [?]:

$$e^X := \sum_{m=0}^{\infty} \frac{X^m}{m!} \quad (\text{converges for any } X), \quad \log X := \sum_{m=1}^{\infty} (-1)^{m+1} \frac{(X-I)^m}{m} \quad (\text{conv. if } \|X-I\| < 1). \quad (\text{A.1})$$

$$e^{\log A} = A \quad (\text{if } \|A-I\| < 1), \quad \log e^X = X \text{ and } \|e^X - I\| < 1 \quad (\text{if } \|X\| < \log 2). \quad (\text{A.2})$$

$$\text{Hilbert-Schmidt norm: } \|X\|^2 := \sum_{i,j} |X_{ij}|^2 = \text{Tr } X^\dagger X. \quad (\text{A.3})$$

Properties:

$$e^{(X^T)} = (e^X)^T, \quad e^{(X^*)} = (e^X)^*, \quad (e^X)^{-1} = e^{-X}, \quad e^{YXY^{-1}} = Y e^X Y^{-1},$$

$$\det \exp X = \exp \text{Tr } X, \quad \frac{d}{dt} e^{tX} = X e^{tX} = e^{tX} X \quad e^{(\alpha+\beta)X} = e^{\alpha X} e^{\beta X} \text{ for } \alpha, \beta \in \mathbb{C};$$

Baker-Campbell-Hausdorff:

$$e^X Y e^{-X} = Y + [X, Y] + \frac{1}{2!} [X, [X, Y]] + \frac{1}{3!} [X, [X, [X, Y]]] + \cdots; \quad (\text{A.4})$$

$$\log(e^X e^Y) = X + \int_0^1 dt g(e^{tX}, e^{tY}) Y \quad \left[ g(z) = \frac{\log z}{1-z^{-1}} = 1 + \sum_{n=1}^{\infty} \frac{(-1)^{n+1} (z-1)^n}{n(n+1)} \right] \quad (\text{A.5})$$

$$= X + Y + \frac{1}{2} [X, Y] + \frac{1}{12} [X, [X, Y]] - \frac{1}{12} [Y, [X, Y]] + \cdots \quad (\text{Baker-Campbell-Hausdorff}). \quad (\text{A.6})$$