

ระบบกายภาพบำบัด ผ่านเว็บแคม

เปลี่ยนกล้องมือถือ ให้เป็นนักกายภาพ AI พร้อมส่งผลให้แพทย์



Presented By:
นางสาวพิชญ์สินี มโนมณีรัตน์



PAIN POINTS

01.

1. ทำท่าผิดหรือทำไม่สุระยะ



2. ขาดแรงจูงใจและการติดตามผลระยะยาว



3. ลืมท่ากายภาพบำบัดว่าต้องทำยังไง



คนไข้ยากทำที่บ้าน
แต่ขาดการดูแล

Musculoskeletal Science and Practice 83 (2026) 103547



Contents lists available at ScienceDirect

Musculoskeletal Science and Practice

journal homepage: www.elsevier.com/locate/msksp



Patient expectations and therapist perceptions in low back pain care: A cross-sectional survey

Dylan J. Gibson , John H. Hollman, Stephanie M. Dang, Zach C. Fellows, Jason C. Schad, Ben E. Zuercher, Darren Q. Calley ^{*}

Program in Physical Therapy, Department of Physical Medicine & Rehabilitation, Mayo Clinic, Rochester, MN, USA

ARTICLE INFO

Keywords:

Low back pain
Physical therapists
Patient preference
Health knowledge
Attitudes
Practice
Cross-sectional studies

ABSTRACT

Background: Low back pain is one of the most common conditions managed by physical therapists. Positive patient expectations have been associated with improved clinical outcomes; however, limited research has examined how physical therapists perceive patients' expectations at the initial consultation.

Objective: To explore patients' expectations at their initial physical therapy evaluation and compare them with physical therapists' perceptions of those expectations.

Methods: A cross-sectional survey was conducted at a large medical center in the United States. Patients and physical therapists completed an identical 10-question, 5-point Likert scale assessing expectations for initial care. Patients completed the survey prior to their first visit; therapists responded electronically. Mann-Whitney U tests were used to compare group differences.

Results: 144 patients and 22 physical therapists completed the survey. Patients most frequently rated provision of a home exercise program as "very" or "extremely important" (93%). Physical therapists assigned relatively greater importance to examination, diagnosis, and prognosis. Statistically significant differences (Bonferroni-adjusted $p \leq 0.001$) reflected differing emphases between groups.

Conclusion: Patients and physical therapists assigned varying levels of importance to several components of initial low back pain care. Physical therapists tended to place relatively greater importance on examination-related elements, whereas patients emphasized receiving a home exercise program. These findings suggest that patient priorities and therapist perceptions may not always align during initial consultations. Therapists may consider directly asking patients about their expectations during the initial consultation to clarify what individuals hope to gain from the clinical encounter.

Features

Real-time Pose Feedback

Progress Dashboard

Gamification

Clinical Report ส่งแพทย์

Human-in-the-Loop Safety



หลักการทำงาน

03.



Vision Tracker

MediaPipe ตี Skeleton
คำนวณมุมข้อต่อ Real-time



AI Coach

คิดและเปรียบเทียบ
นำผลลัพธ์ที่ได้ มาเตือนคนไข้



Clinical Reporter

สรุปผล สร้าง PDF รายงาน
และส่งให้แพทย์หรือนักกายภาพ



03.

 **capacitor**


 **tailwindcss**

HTML


JS




MySQL

เป้าหมาย

สิ่งที่อยากเห็น:

- คนใช้ทำกายภาพที่บ้านได้โดยไม่ต้องพึ่งนักกายภาพ ตลอดเวลา
- มี AI คอยบอกแบบเป็นกันเองว่า "งอเข้าอีกนิดนะ" ไม่ใช่แค่จับผิด
- ผู้สูงอายุไม่รู้สึกลำบากตลอดเวลาต้องฝึกคนเดียว
- แพทย์เห็นข้อมูลจริง ไม่ใช่แค่ฟังจากคนไข้





***Thank
You!***

