

POLAR background prediction

S. Nguyen, N. Produit

University of Geneva

stephane.nguyen@etu.unige.ch

nicolas.produit@unige.ch

October 23, 2023

Overview

1 Introduction

- Gamma Ray Bursts
- Related work
- Our work

2 Methodology











- Background model
- Poor predictions
- Cluster and cluster intersections

3 Results & discussion

4 Conclusion











Gamma Ray Bursts (GRBs)

- What are GRBs?

¹<https://www.astro.unige.ch/polar/grb-light-curves>          











Gamma Ray Bursts (GRBs)

- What are GRBs?
 - ▶ Intense, energetic, and extra-galactic explosions

¹<https://www.astro.unige.ch/polar/grb-light-curves>          


Gamma Ray Bursts (GRBs)

- What are GRBs?
 - ▶ Intense, energetic, and extra-galactic explosions
 - ▶ Bursts of high-energy photons

¹<https://www.astro.unige.ch/polar/grb-light-curves>          

Gamma Ray Bursts (GRBs)

- What are GRBs?
 - ▶ Intense, energetic, and extra-galactic explosions
 - ▶ Bursts of high-energy photons
 - ▶ Bursts of gamma rays with an undetermined cause

¹<https://www.astro.unige.ch/polar/grb-light-curves> 

Gamma Ray Bursts (GRBs)

- What are GRBs?
 - ▶ Intense, energetic, and extra-galactic explosions
 - ▶ Bursts of high-energy photons
 - ▶ Bursts of gamma rays with an undetermined cause
- Lead to spikes in photon counts/rates




Figure: A light-curve containing a Gamma Ray Burst¹

¹<https://www.astro.unige.ch/polar/grb-light-curves>

What are astrophysicists interested in?

- Better understand GRBs

What are astrophysicists interested in?

- Better understand GRBs
- For this purpose, there's a need for:
 - ▶ Data taking → POLAR detector




Figure: Picture taken from <https://www.astro.unige.ch/polar/>

What are astrophysicists interested in?

- Better understand GRBs
- For this purpose, we need to:
 - ▶ Capture data → POLAR detector or POLAR-2 detector
 - ▶ Detect

What are astrophysicists interested in?

- Better understand GRBs
- For this purpose, we need to:
 - ▶ Capture data → POLAR detector or POLAR-2 detector
 - ▶ Detect
 - ▶ Localize

What are astrophysicists interested in?

- Better understand GRBs
- For this purpose, we need to:
 - ▶ Capture data → POLAR detector or POLAR-2 detector
 - ▶ Detect
 - ▶ Localize
 - ▶ Further measurements

What are astrophysicists interested in?

- Better understand GRBs
- For this purpose, we need to:
 - ▶ Capture data → POLAR detector or POLAR-2 detector
 - ▶ Detect
 - ▶ Localize
 - ▶ Further measurements
 - ▶ Analysis

Related work

- Koziol's master thesis [1]: HAGRID²

²High Accuracy GRB Rapid Inference with Deep Learning in Space

Related work

- Koziol's master thesis [1]: HAGRID²
 - ▶ Fast real-time GRB detection and localization using ML

²High Accuracy GRB Rapid Inference with Deep Learning in Space

Related work

- Koziol's master thesis [1]: HAGRID²
 - ▶ Fast real-time GRB detection and localization using ML
 - ▶ In particular, for GRB detection, he used:

²High Accuracy GRB Rapid Inference with Deep Learning in Space

Related work

- Koziol's master thesis [1]: HAGRID²
 - ▶ Fast real-time GRB detection and localization using ML
 - ▶ In particular, for GRB detection, he used:
 - ★ Labeled simulated data with simulated GRBs

²High Accuracy GRB Rapid Inference with Deep Learning in Space

- Koziol's master thesis [1]: HAGRID²
 - ▶ Fast real-time GRB detection and localization using ML
 - ▶ In particular, for GRB detection, he used:
 - ★ Labeled simulated data with simulated GRBs
 - ★ From past 60 [s] simulated light-curves + energy info.

²High Accuracy GRB Rapid Inference with Deep Learning in Space

Related work

- Koziol's master thesis [1]: HAGRID²
 - ▶ Fast real-time GRB detection and localization using ML
 - ▶ In particular, for GRB detection, he used:
 - ★ Labeled simulated data with simulated GRBs
 - ★ From past 60 [s] simulated light-curves + energy info.
 - ★ Predict the occurrence of a GRB at time t

²High Accuracy GRB Rapid Inference with Deep Learning in Space

Related work

- Koziol's master thesis [1]: HAGRID³
 - ▶ Possible data distribution mismatch

³High Accuracy GRB Rapid Inference with Deep Learning in Space

Related work

- Koziol's master thesis [1]: HAGRID³
 - ▶ Possible data distribution mismatch
 - ▶ But low miss-detection: correctly detected all real GRBs within 10 days of POLAR data.

³High Accuracy GRB Rapid Inference with Deep Learning in Space

Related work

- Koziol's master thesis [1]: HAGRID³
 - ▶ Possible data distribution mismatch
 - ▶ But low miss-detection: correctly detected all real GRBs within 10 days of POLAR data.
 - ▶ Does not account for other measurements

³High Accuracy GRB Rapid Inference with Deep Learning in Space

Our work

Would like to

- Train our model from real POLAR data

Our work

Would like to

- Train our model from real POLAR data
 - ▶ No longer have access to labels

Our work

Would like to

- Train our model from real POLAR data
 - ▶ No longer have access to labels
 - ▶ → Background model

Our work

Would like to

- Train our model from real POLAR data
 - ▶ No longer have access to labels
 - ▶ → Background model
- Explicitly account for other measurements

Our work

Would like to

- Train our model from real POLAR data
 - ▶ No longer have access to labels
 - ▶ → Background model
- Explicitly account for other measurements
 - ▶ Predict photon rates from diverse measurements

Our work

Would like to

- Train our model from real POLAR data
 - ▶ No longer have access to labels
 - ▶ → Background model
- Explicitly account for other measurements
 - ▶ Predict photon rates from diverse measurements
- Reduce complexity barrier to model interpretability

Our work

Would like to

- Train our model from real POLAR data
 - ▶ No longer have access to labels
 - ▶ → Background model
- Explicitly account for other measurements
 - ▶ Predict photon rates from diverse measurements
- Reduce complexity barrier to model interpretability
 - ▶ → Replace the sequential model for a Multi-Layer Perceptron (MLP)

Our work

Would like to

- Train our model from real POLAR data
 - ▶ No longer have access to labels
 - ▶ → Background model
- Explicitly account for other measurements
 - ▶ Predict photon rates from diverse measurements
- Reduce complexity barrier to model interpretability
 - ▶ → Replace the sequential model for a Multi-Layer Perceptron (MLP)
- Try to interpret our trained model

Our work

Would like to

- Train our model from real POLAR data
 - ▶ No longer have access to labels
 - ▶ → Background model
- Explicitly account for other measurements
 - ▶ Predict photon rates from diverse measurements
- Reduce complexity barrier to model interpretability
 - ▶ → Replace the sequential model for a Multi-Layer Perceptron (MLP)
- Try to interpret our trained model
 - ▶ → Gradients w.r.t input variables

Methodology

- Train background model **without** 25 known GRBs⁴.

⁴25 of 55 known GRBs [2] happening within our data

Methodology

- Train background model **without** 25 known GRBs⁴.
- Apply our model to everything

⁴25 of 55 known GRBs [2] happening within our data

Methodology

- Train background model **without** 25 known GRBs⁴.
- Apply our model to everything
- Extract time intervals based on the residuals

⁴25 of 55 known GRBs [2] happening within our data


Methodology

- Train background model **without** 25 known GRBs⁴.
- Apply our model to everything
- Extract time intervals based on the residuals
- Peek at model interpretability using Jacobian matrices

⁴25 of 55 known GRBs [2] happening within our data

Methodology



- Train background model **without** 25 known GRBs⁴.
- Apply our model to everything
- Extract time intervals based on the residuals
- Peek at model interpretability using Jacobian matrices
- A few technologies:



⁴25 of 55 known GRBs [2] happening within our data

Background model

- Multi-Layer Perceptron (MLP) background model⁵





- (a) Multi-Layer Perceptron high-level
- (b) Graph of operators describing my Multi-Layer Perceptron

⁵First picture taken from <https://medium.com/unpackai/from-anns-artificial-neural-networks-to-rnns-recurrent-neural-networks-93b638772fd1>

Background model

- Multi-Layer Perceptron (MLP) background model⁵



- (a) Multi-Layer Perceptron high-level
- (b) Graph of operators describing my Multi-Layer Perceptron

- Trained to minimize a weighted MSE

⁵First picture taken from <https://medium.com/unpackai/from-anns-artificial-neural-networks-to-rnns-recurrent-neural-networks-93b638772fd1>

Poor predictions




Figure: Normalized pulls for rate[0]

Cluster and cluster intersections

- Aggregating extracted data-points \rightarrow clusters

Cluster and cluster intersections

- Aggregating extracted data-points \rightarrow clusters
- GRBs should be visible in different energy bins

Cluster and cluster intersections

- Aggregating extracted data-points \rightarrow clusters
- GRBs should be visible in different energy bins
- Cluster of data points extracted using different energy bins:
 - ▶ Each from the same set of energy bins
 - ▶ Or each satisfying same conditions (e.g. number of energy bins)

Results & discussion: Known GRBs




Figure: Energy bin 0, time windows of ± 100 [s] around 5 known GRB trigger times [2], based on pulls with $k = 5$. Predictions are in black.

Results & discussion: Known GRBs






Figure: Energy bin 0, time windows of ± 100 [s] around 5 other known GRB trigger times [2], based on pulls with $k = 5$.


Results & discussion: Cluster intersections



(a) Cluster intersection with more the 3 energy bins; Solar flare confirmed by Prof. Nicolas Produit




(b) Undetermined cluster intersection with all energy bins and with a 30-second discard window.




(c) Undetermined cluster intersection with all energy bins and with a 30-second discard window.

Results & discussion: Model interpretability




(a) Histogram of $\frac{\partial \text{rate}[0]}{\partial \text{crabarf}}$ (un-normalized features)




(b) Avg slope μ greater (lesser) than its (negative) standard error $\frac{\sigma}{\sqrt{N}}$?

Results & discussion: Model interpretability



(a) Histogram of $\frac{\partial \text{rate}[0]}{\partial \text{crabarf}}$ (un-normalized features)



(b) Avg slope μ greater (lesser) than its (negative) standard error $\frac{\sigma}{\sqrt{N}}$?

- Need for further analysis

Results & discussion

Table 6: Mean \pm standard deviation (over 10 different seeds) number of positive/negative cluster intersections for different sets of energy bins bs or conditions²⁹. This time, there's a discard window of 30 seconds.

bs or condition	negative	positive	bs or condition	negative	positive	bs or condition	negative	positive
# inter > 0	5762.1 \pm 965.708	6487.6 \pm 826.131	0,2	6.9 \pm 2.726	10.7 \pm 4.111	1,3	15.6 \pm 6.398	51.2 \pm 42.695
# inter > 1	4802.1 \pm 822.34	5450.6 \pm 752.954	0,2,3	6.0 \pm 3.432	7.1 \pm 1.37	1,3,4	1.0*	–
# inter > 2	3493.8 \pm 610.247	4083.5 \pm 564.85	0,2,3,4	8.0*	–	1,3,4,5	1.333 \pm 0.577	1.5 \pm 0.577
# inter > 3	3238.4 \pm 602.769	3821.5 \pm 538.957	0,2,3,4,5	212.2 \pm 42.158	242.7 \pm 65.109	1,3,5	–	5.5 \pm 6.364
# inter > 4	1150.2 \pm 232.379	1595.0 \pm 202.499	0,2,3,5	1.0 \pm 0.0	3.5 \pm 3.536	1,4	1.0*	3.0*
# inter > 5	791.3 \pm 165.14	1135.2 \pm 149.199	0,2,4	1.5 \pm 0.707	1.0*	1,4,5	2.125 \pm 1.356	5.5 \pm 3.742
0	23.8 \pm 8.23	71.6 \pm 18.368	0,2,4,5	5.8 \pm 4.367	15.8 \pm 5.095	1,5	–	2.0 \pm 0.0
0,1	16.6 \pm 6.059	62.3 \pm 45.631	0,2,5	1.0 \pm 0.0	2.5 \pm 2.38	2	734.8 \pm 169.59	939.8 \pm 235.236
0,1,2	4.0 \pm 2.0	15.2 \pm 9.739	0,3	1.0*	3.0 \pm 1.414	2,3	2459.9 \pm 493.897	3265.5 \pm 779.499
0,1,2,3	12.4 \pm 3.806	30.5 \pm 17.219	0,3,4,5	1.667 \pm 1.155	1.75 \pm 0.957	2,3,4	114.8 \pm 335.669	11.75 \pm 12.395
0,1,2,3,4	23.0*	2.5 \pm 0.707	0,4	2.0*	–	2,3,4,5	3156.9 \pm 657.884	4049.1 \pm 676.033
0,1,2,3,4,5	791.3 \pm 165.14	1135.2 \pm 149.199	0,4,5	2.25 \pm 1.753	11.6 \pm 3.062	2,3,5	60.444 \pm 11.479	225.0 \pm 392.74
0,1,2,3,5	1.0 \pm 0.0	3.667 \pm 4.899	0,5	1.0*	1.75 \pm 1.5	2,4	18.333 \pm 38.563	2.333 \pm 1.751
0,1,2,4	1.0*	3.0*	1	52.6 \pm 15.16	104.0 \pm 118.135	2,4,5	212.3 \pm 59.913	244.9 \pm 73.942
0,1,2,4,5	2.8 \pm 1.751	11.2 \pm 7.315	1,2	3.444 \pm 1.424	8.5 \pm 4.143	2,5	9.8 \pm 6.356	55.5 \pm 127.411
0,1,2,5	1.0*	2.667 \pm 2.082	1,2,3	56.1 \pm 19.221	115.1 \pm 41.391	3	1206.3 \pm 282.876	1633.0 \pm 282.032
0,1,3	1.25 \pm 0.5	4.875 \pm 7.2	1,2,3,4	14.75 \pm 27.5	1.4 \pm 0.548	3,4	6.25 \pm 9.215	1.0 \pm 0.0
0,1,3,4,5	2.5 \pm 2.811	2.25 \pm 0.5	1,2,3,4,5	413.5 \pm 141.535	714.4 \pm 132.802	3,4,5	19.7 \pm 8.166	16.5 \pm 4.503
0,1,3,5	–	1.0*	1,2,3,5	2.167 \pm 0.753	9.667 \pm 19.268	3,5	1.889 \pm 0.782	21.0 \pm 58.877
0,1,4	1.0*	6.0*	1,2,4	–	4.0*	4	9.333 \pm 20.174	3.833 \pm 4.167
0,1,4,5	2.875 \pm 1.959	5.9 \pm 4.202	1,2,4,5	3.143 \pm 2.673	4.889 \pm 3.333	4,5	123.5 \pm 30.395	124.0 \pm 31.383
0,1,5	2.0*	2.0*	1,2,5	–	3.0*	5	7.3 \pm 3.302	88.1 \pm 234.779

Results & discussion

Table 6: Mean \pm standard deviation (over 10 different seeds) number of positive/negative cluster intersections for different sets of energy bins bs or conditions²⁹. This time, there's a discard window of 30 seconds.

bs or condition	negative	positive	bs or condition	negative	positive	bs or condition	negative	positive
# inter > 0	5762.1 \pm 965.708	6487.6 \pm 826.131	0,2	6.9 \pm 2.726	10.7 \pm 4.111	1,3	15.6 \pm 6.398	51.2 \pm 42.695
# inter > 1	4802.8 \pm 822.34	5450.6 \pm 752.954	0,2,3	6.0 \pm 3.432	7.1 \pm 1.37	1,3,4	1.0*	–
# inter > 2	3493.8 \pm 610.247	4083.5 \pm 564.85	0,2,3,4	8.0*	–	1,3,4,5	1.333 \pm 0.577	1.5 \pm 0.577
# inter > 3	3238.4 \pm 602.769	3821.5 \pm 538.957	0,2,3,4,5	212.2 \pm 42.158	242.7 \pm 65.109	1,3,5	–	5.5 \pm 6.364
# inter > 4	1150.2 \pm 232.379	1595.0 \pm 202.499	0,2,3,5	1.0 \pm 0.0	3.5 \pm 3.536	1,4	1.0*	3.0*
# inter > 5	791.3 \pm 165.14	1135.2 \pm 149.199	0,2,4	1.5 \pm 0.707	1.0*	1,4,5	2.125 \pm 1.356	5.5 \pm 3.742
0	23.8 \pm 8.23	71.6 \pm 18.368	0,2,4,5	5.8 \pm 4.367	15.8 \pm 5.095	1,5	–	2.0 \pm 0.0
0,1	16.6 \pm 6.059	62.3 \pm 45.631	0,2,5	1.0 \pm 0.0	2.5 \pm 2.38	2	734.8 \pm 169.59	939.8 \pm 235.236
0,1,2	4.0 \pm 2.0	15.2 \pm 9.739	0,3	1.0*	3.0 \pm 1.414	2,3	2459.9 \pm 493.897	3265.5 \pm 779.499
0,1,2,3	12.4 \pm 3.806	30.5 \pm 17.219	0,3,4,5	1.667 \pm 1.155	1.75 \pm 0.957	2,3,4	114.8 \pm 335.669	11.75 \pm 12.395
0,1,2,3,4	23.0*	2.5 \pm 0.707	0,4	2.0*	–	2,3,4,5	3156.9 \pm 657.884	4049.1 \pm 676.033
0,1,2,3,4,5	791.3 \pm 165.14	1135.2 \pm 149.199	0,4,5	2.25 \pm 1.753	11.6 \pm 3.062	2,3,5	60.444 \pm 11.479	225.0 \pm 392.74
0,1,2,3,5	1.0 \pm 0.0	3.667 \pm 4.899	0,5	1.0*	1.75 \pm 1.5	2,4	18.333 \pm 38.563	2.333 \pm 1.751
0,1,2,4	1.0*	3.0*	1	52.6 \pm 15.16	104.0 \pm 118.135	2,4,5	212.3 \pm 59.913	244.9 \pm 73.942
0,1,2,4,5	2.8 \pm 1.751	11.2 \pm 7.315	1,2	3.444 \pm 1.424	8.5 \pm 4.143	2,5	9.8 \pm 6.356	55.5 \pm 127.411
0,1,2,5	1.0*	2.667 \pm 2.082	1,2,3	56.1 \pm 19.221	115.1 \pm 41.391	3	1206.3 \pm 282.876	1633.0 \pm 282.032
0,1,3	1.25 \pm 0.5	4.875 \pm 7.2	1,2,3,4	14.75 \pm 27.5	1.4 \pm 0.548	3,4	6.25 \pm 9.215	1.0 \pm 0.0
0,1,3,4,5	2.5 \pm 2.811	2.25 \pm 0.5	1,2,3,4,5	413.5 \pm 141.535	714.4 \pm 132.802	3,4,5	19.7 \pm 8.166	16.5 \pm 4.503
0,1,3,5	–	1.0*	1,2,3,5	2.167 \pm 0.753	9.667 \pm 19.268	3,5	1.889 \pm 0.782	21.0 \pm 58.877
0,1,4	1.0*	6.0*	1,2,4	–	4.0*	4	9.333 \pm 20.174	3.833 \pm 4.167
0,1,4,5	2.875 \pm 1.959	5.9 \pm 4.202	1,2,4,5	3.143 \pm 2.673	4.889 \pm 3.333	4,5	123.5 \pm 30.395	124.0 \pm 31.383
0,1,5	2.0*	2.0*	1,2,5	–	3.0*	5	7.3 \pm 3.302	88.1 \pm 234.779

- Need an analysis of clusters' stability due to initial seeds

Conclusion: Summary

- Built a model of the background (light curve) using data collected from the POLAR detector

Conclusion: Summary

- Built a model of the background (light curve) using data collected from the POLAR detector
- Used poor predictions to extract time intervals

Conclusion: Summary

- Built a model of the background (light curve) using data collected from the POLAR detector
- Used poor predictions to extract time intervals
- Brief peek at model interpretability using partial derivatives of output w.r.t. input

Conclusion: Possible improvements

- Future analysis of our extracted time intervals is required
 - ▶ Categorize them into solar events, GRBs and so on

Conclusion: Possible improvements

- Future analysis of our extracted time intervals is required
 - ▶ Categorize them into solar events, GRBs and so on
 - ▶ Assess the clusters' stabilities due to initial seeds

Conclusion: Possible improvements

- Future analysis of our extracted time intervals is required
 - ▶ Categorize them into solar events, GRBs and so on
 - ▶ Assess the clusters' stabilities due to initial seeds
- Manual inspection of our clusters by experts can give crucial information telling us what to do next.

Conclusion: Possible improvements

- Future analysis of our extracted time intervals is required
 - ▶ Categorize them into solar events, GRBs and so on
 - ▶ Assess the clusters' stabilities due to initial seeds
- Manual inspection of our clusters by experts can give crucial information telling us what to do next.
- Other possible improvements:
 - ▶ Dive deeper into model interpretability

Conclusion: Possible improvements

- Future analysis of our extracted time intervals is required
 - ▶ Categorize them into solar events, GRBs and so on
 - ▶ Assess the clusters' stabilities due to initial seeds
- Manual inspection of our clusters by experts can give crucial information telling us what to do next.
- Other possible improvements:
 - ▶ Dive deeper into model interpretability
 - ▶ Change the methodology, e.g. use sequential models

Conclusion: Possible improvements

- Future analysis of our extracted time intervals is required
 - ▶ Categorize them into solar events, GRBs and so on
 - ▶ Assess the clusters' stabilities due to initial seeds
- Manual inspection of our clusters by experts can give crucial information telling us what to do next.
- Other possible improvements:
 - ▶ Dive deeper into model interpretability
 - ▶ Change the methodology, e.g. use sequential models
 - ▶ Improve code quality, GPU usage and decrease memory footprint.

The End

References

Figures without references come from author of the slides.



[1] Gilles Koziol, (2023)

HAGRID in Space. *Master's thesis*, University of Geneva.

<https://cernbox.cern.ch/s/X10vZ4iY19vewcV>. Accessed: 2023-09-22.



[2] François Fleuret, (2023)

Deep Learning Course.

<https://fleuret.org/dlc/>. Accessed: 2023-09-22



[2] Shaolin Xiong, Yuanhao Wang, Zhengheng Li, Jianchao Sun, Yi Zhao, Hancheng Li, and Yue Huang, (2017)

Overview of the GRB observation by POLAR.

ICRC2017. page 640