

05: Eigenbasis Example & Diagonalization

Chapter Goal: To demonstrate with a concrete example how to use [Diagonalization](#) to compute powers of a matrix, and to verify that it gives the same result as the "brute force" method.

1. Problem Setup

- **Transformation** T : A 2x2 matrix.

$$T = \begin{bmatrix} 1 & 1 \\ 0 & 2 \end{bmatrix}$$

- **Geometric Interpretation:** This is a combination of a vertical scaling (by a factor of 2) and a horizontal shear.
- **Initial Vector** \vec{v} :

$$\vec{v} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

- **Goal:** Calculate $T^2\vec{v}$. (Where will \vec{v} land after being transformed by T twice?).
-

2. Method #1: "Brute Force" (Direct Calculation)

This method does not use diagonalization, only standard matrix multiplication.

Option A: Apply T to the Vector Twice

- **First Step:**

$$\vec{v}_1 = T\vec{v} = \begin{bmatrix} 1 & 1 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} (-1 + 1) \\ (0 + 2) \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \end{bmatrix}$$

- **Second Step:**

$$\vec{v}_2 = T\vec{v}_1 = \begin{bmatrix} 1 & 1 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 0 \\ 2 \end{bmatrix} = \begin{bmatrix} (0 + 2) \\ (0 + 4) \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$$

- **Final Result:** $\begin{bmatrix} 2 \\ 4 \end{bmatrix}$.

Option B: Calculate T^2 First, then Multiply by Vector

- **Calculate** T^2 :

$$T^2 = T \cdot T = \begin{bmatrix} 1 & 1 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 2 \end{bmatrix} = \begin{bmatrix} (1+0) & (1+2) \\ (0+0) & (0+4) \end{bmatrix} = \begin{bmatrix} 1 & 3 \\ 0 & 4 \end{bmatrix}$$

- **Multiply by \vec{v} :**

$$T^2 \vec{v} = \begin{bmatrix} 1 & 3 \\ 0 & 4 \end{bmatrix} \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} (-1+3) \\ (0+4) \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$$

- **Final Result:** $\begin{bmatrix} 2 \\ 4 \end{bmatrix}$. Both ways give the same result, as expected.

3. Method #2: "The Elegant Way" (Using Diagonalization)

Now we will use the recipe $T^2 = CD^2C^{-1}$.

Step 1: Find the "Ingredients" (λ , C , D , C^{-1})

- **Find the Eigenvalues and Eigenvectors of T :**

(This is skipped in the video, but we can calculate it quickly:

$$\det(T - \lambda I) = (1 - \lambda)(2 - \lambda) = 0)$$

- $\lambda_1 = 1$, with corresponding eigenvector $\vec{v}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$.
- $\lambda_2 = 2$, with corresponding eigenvector $\vec{v}_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$.

- **Form Matrices C and D :**

- **C (Eigenvector Matrix):** Its columns are the eigenvectors.

$$C = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$

- **D (Diagonal Matrix):** Its diagonal entries are the eigenvalues, in the same order as the columns in C .

$$D = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}$$

- **Find C^{-1} (The Inverse of C):**

- $\det(C) = (1 \cdot 1) - (1 \cdot 0) = 1$.

- $$C^{-1} = \frac{1}{1} \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}$$

Step 2: Calculate T^2 with the Diagonalization Formula

$$T^2 = CD^2C^{-1}$$

$$T^2 = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1^2 & 0 \\ 0 & 2^2 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 4 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}$$

- **Calculate D^2C^{-1} first:**

$$\begin{bmatrix} 1 & 0 \\ 0 & 4 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} (1+0) & (-1+0) \\ (0+0) & (0+4) \end{bmatrix} = \begin{bmatrix} 1 & -1 \\ 0 & 4 \end{bmatrix}$$

- **Multiply C by the result:**

$$\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 0 & 4 \end{bmatrix} = \begin{bmatrix} (1+0) & (-1+4) \\ (0+0) & (0+4) \end{bmatrix} = \begin{bmatrix} 1 & 3 \\ 0 & 4 \end{bmatrix}$$

- **Result for T^2 :** $\begin{bmatrix} 1 & 3 \\ 0 & 4 \end{bmatrix}$. This is **EXACTLY the same** as the T^2 we calculated with the brute force method!

Step 3: Multiply by the Initial Vector

$$T^2 \vec{v} = \begin{bmatrix} 1 & 3 \\ 0 & 4 \end{bmatrix} \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} (-1+3) \\ (0+4) \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$$

- **Final Result:** $\begin{bmatrix} 2 \\ 4 \end{bmatrix}$. Once again, the result is identical.

4. Conclusion

- This example provides concrete proof that both methods (brute force and diagonalization) yield the identical result.
- While the difference in effort is not dramatic for a 2x2 matrix raised to the power of 2, imagine if we had to calculate T^{100} . The diagonalization method would be vastly more efficient because calculating D^{100} is instantaneous.
- This serves as a validation that the theory of $T^n = CD^nC^{-1}$ truly works in practice.

Tags: #mml-specialization #linear-algebra #eigenbasis #diagonalization #worked-example