
Assignment 2, COMP 540

Chen Zeng(cz39), Zhihui Xie(zx18)

January 24, 2018

1 Gradient and Hessian of $NLL(\theta)$ for logistic regression

Answer1 Because of:

$$g(z) = \frac{1}{(1 + e^{-z})}$$

We can get:

$$g(z)(1 + e^{-z}) = 1$$

Take the derivation on both side of the z, then we can get:

$$g'(z)(1 + e^{-z}) - g(z)e^{-z} = 0$$

Finally, after some basic transformation, we can get:

$$\frac{\partial g(z)}{\partial z} = g(z)(1 - g(z))$$

Therefore, the conclusion is proved.

Answer2 As we know:

$$NLL(\theta) = \frac{1}{2} \sum_{i=1}^m \left(y^{(i)} - \theta^T x^{(i)} \right)^2$$

Take the derivation on it of the parameter θ_j , we can get:

$$\frac{\partial NLL(\theta)}{\partial \theta_j} = x_j^{(i)} \sum_{i=1}^m \left(\sum_{j=1}^d \left(x_j^{(i)} \theta_j \right) - y^{(i)} \right)$$

Therefore, take the derivation on $NLL(\theta)$ of vector θ , we can get:

$$\frac{\partial NLL(\theta)}{\partial \theta} = \sum_{i=1}^m \left(\theta^T x^{(i)} - y^{(i)} \right) x^{(i)}$$

It's the same as another format:

$$\frac{\partial NLL(\theta)}{\partial \theta} = \sum_{i=1}^m \left(h_{\theta}(x^{(i)}) - y^{(i)} \right) x^{(i)}$$

Therefore, the conclusion is proved.

Answer3 Assume z is a nonzero vector with the size $n \times 1$, X is a $m \times n$ full rank matrix, $S_{i,i}$ is the i th element in the diagonal of S and it is strictly positive.

$$\begin{aligned} z^T H z &= z^T X^T S X z \\ &= (X z)^T H (X z) \\ &= \sum_{i=1}^m S_{i,i} \sum_{j=1}^k (x_j z_j)^2 \\ &\geq 0 \end{aligned}$$

So H is positive definite.

2 Properties of L2 regularized logistic regression

Answer1 False. $J(\theta)$ is a convex function and has a global minimum, so it does not have multiple locally optimal solutions.

Answer2 False. Ridge regularization drives θ components to zero but not to exactly zero while Lasso regularization drives many θ components to exactly zero especially when the regularization parameter λ is large.

Answer3 True. For example, if the data points of two classes are linear separable in 2-dimension plane and a single line L crossing the origin can separate them into two classes, the line L can be represented as $ax + by = 0$ and the coefficients a and b can be infinite.

Answer4 False. The λ term determines the relative importance of the first term and the penalty term on θ . Thus, the first term of $J(\theta)$ always decrease as we increase λ .

3 Implementing a k-nearest-neighbor classifier

Please help to refer to the knn folder.




Figure 4.1: Overfitting




Figure 4.2: under-fitting

4 Implementing logistic regression

Problem 3, Part B: Varying λ

- When $\lambda = 0.00001$, the model is overfitting on this data set. Figure 4.1 shows the decision boundary for overfitting.
- When $\lambda = 100$, the model is under-fitting on this data set. Figure 4.2 shows the decision boundary for under-fitting.

Problem 3 Part C: Fitting regularized logistic regression models (L2 and L1)

- L2 regularization and log transformation is the best combination for logistic regression model on for this data set for the reason that it has the highest accuracy 0.94140625 on

the test set with the best lambda 0.1.