

Analysis of Algorithms

Homework 2

Thomas Schollenberger (tss2344)

September 29, 2022

Problem #1

Problem #2

The function `fibItHelper` implemented the recurrence $f(n; a, b)$. What is the time complexity of `fibItHelper`? Write down a recurrence relation $T_f(n)$ that characterizes the time complexity in terms of the number of additions (plusses) performed; then solve the recurrence exactly using iteration.

$$T_f(0) = 0 \quad (1a)$$

$$T_f(1) = 0 \quad (1b)$$

$$T_f(n+1) = 1 + T_f(n) \quad (1c)$$

Which makes $T_f(n) = n - 1$

Problem #3

Let $L : \mathbb{N}^2 \rightarrow \mathbb{N}^2$ be defined as $L(a, b) = (b, a + b)$. Then $f(n; a, b)$ can be understood as $(L^n(a, b))_1$. Prove that $n \in \mathbb{N}, L^n(a, b) = (f(n; a, b), f(n+1; a, b))$. Base case:

$$L^0(a, b) = 1(a, b)_1 = a = f(0; a, b) \quad (2a)$$

Assume that $L^k(a, b) = (f(k; a, b), f(k+1; a, b))$ is true, where $0 \leq k < n$, then:

$$L^{k+1}(a, b) = L_{k+1}(L_k(\dots(L_2(L_1(a, b))))) \quad (2b)$$

$$L^{k+1}(a, b) = L_{k+1}(f(k; a, b), f(k+1; a, b)) \quad (2c)$$

$$L^{k+1}(a, b) = (f(k+1; a, b), f(k+1; a, b) + f(k; a, b)) \quad (2d)$$

$$L^{k+1}(a, b) = (f(k+1; a, b), f(k+2; a, b)) \quad (2e)$$

Problem #5

- (a) An algorithm runs in pseudo-polynomial time if the runtime is some polynomial in the numeric value of the input, rather than in the number of bits required to represent it
- (b) Yes. It is pseudo-polynomial because it scales exponentially in steps whenever you increase n by 1. This explains why it slows drastically by $n = 30$.
- (c) No. It is a polynomial time algorithm, as it linearly scales ($\text{bits} * n$), which is what allows it to be extremely fast.

Problem #6

Solve the following recurrences exactly using the iteration method. In all cases, $T(0) = 0$. Answers should be expressed in terms of $T(n)$

a. $T(n + 1) = T(n) + 5$

$$T(n + 1) = T(n) + 5 \quad (3a)$$

$$= (T(n - 1) + 5) + 5 \quad (3b)$$

$$= ((T(n - 2) + 5) + 5) + 5 \quad (3c)$$

$$T(n) = T(n - k) + 5k \quad (3d)$$

$$T(n) = 5n \quad (3e)$$

b. $T(n + 1) = n + T(n)$

$$T(n + 1) = n + T(n) \quad (4a)$$

$$= n(n + T(n - 1)) \quad (4b)$$

$$= (n + 1)(n((n - 1) + T(n - 2))) \quad (4c)$$

$$T(n) = T(n - k) + \sum_{i=0}^k (n - i) \quad (4d)$$

$$T(n) = n! \quad (4e)$$

Problem #7

Solve the following recurrences exactly using the iteration method. In all cases, $T(0) = 1$. Answers should be expressed in terms of $T(n)$

a. $T(n+1) = 2T(n)$

$$T(n+1) = 2T(n) \quad (5a)$$

$$= 2(2T(n-1)) \quad (5b)$$

$$= 2(2(2T(n-2))) \quad (5c)$$

$$T(n) = 2kT(n-k) \quad (5d)$$

$$T(n) = 2n \quad (5e)$$

b. $T(n+1) = 2^{n+1} + T(n)$

$$T(n+1) = 2^{n+1} + T(n) \quad (6a)$$

$$= 2^{n+1} + (2^n + T(n-1)) \quad (6b)$$

$$= 2^{n+1} + (2^n(2^{n-1} + T(n-2))) \quad (6c)$$

$$T(n) = \sum_{i=0}^k (2^{n-i}) + T(n-k)T(n) = 2^{n!} \quad (6d)$$

Problem #8

Solve the following recurrences exactly using the iteration method. In all cases, $T(1) = 1$. Answers should be expressed in terms of $T(n)$

a. $T(n) = n + T(n/2)$ (**Assume n has the form $n = 2^m$.**)

$$T(n) = n + T(n/2) \quad (7a)$$

$$= n + (n/2 + T((n/2)/2)) \quad (7b)$$

$$= n + (n/2 + (n/4 + T((n/4)/2))) \quad (7c)$$

$$m = \lg(n) \quad (7d)$$

$$T(n) = n + n(\lg(n)) \quad (7e)$$

b. $T(n) = 1 + T(n/3)$ (**Assume n has the form $n = 3^m$.**)

$$T(n) = 1 + T(n/3) \quad (8a)$$

$$= 1 + (1 + T((n/3)/3)) \quad (8b)$$

$$= 1 + (1 + (1 + T(((n/3)/3)/3))) \quad (8c)$$

$$m = \log_3(n) \quad (8d)$$

$$T(n) = 1 + \log_3(n) \quad (8e)$$