

Part IA — Probability

Based on lectures by Dr D. Yeo

Lent 2022

Contents

0	Introduction	3
1	Formal Setup	4
1.1	Examples of Probability Spaces	6
2	Combinatorial Analysis	8
2.1	Subsets	8
2.2	Random Walks	8
2.3	Stirling's Formula	9
2.4	(Ordered) compositions	10
3	Properties of Probability measures	11
3.1	Countable sub-additivity	11
3.2	Continuity	12
3.3	Inclusion-Exclusion Principle	13
3.4	Bonferroni Inequalities	15
3.5	Counting with IEP	16
3.6	Independence	18
3.6.1	Properties	20
3.7	Conditional Probability	20
3.7.1	Properties	21
3.7.2	Law of Total Probability and Bayes' Formula	23
4	Discrete Random Variables	27
4.1	Discrete Probability Distributions	28
4.1.1	Bernoulli distribution - "(biased) coin toss"	28
4.1.2	Binomial distribution	29
4.1.3	More than one RV	29
4.1.4	Geometric distribution ("Waiting for success")	30
4.1.5	Poisson Distribution	30

4.2	Expectation	31
4.2.1	Properties of Expectation	34
4.3	Variance	37
4.3.1	Sums of RVs	40
4.3.2	Chebyshev's Inequality	43
4.4	Conditional Expectation	44
4.5	Random Walks	46
4.5.1	Simple Random Walk (SRW) on \mathbb{Z} - Main example in our course .	46

§0 Introduction

Text in blue is usually less important.

Example 0.1

Dice: outcomes $1, 2, \dots, 6$.

- $\mathbb{P}(2) = \frac{1}{6}$.
- $\mathbb{P}(\text{multiple of } 3) = \frac{2}{6} = \frac{1}{3}$.
- $\mathbb{P}(\text{not a multiple of } 3) = \frac{2}{3}$
- $\mathbb{P}(\text{prime}) = \frac{1}{2}$.
-

$$\begin{aligned}\mathbb{P}(\text{prime or multiple of } 3) &= \frac{1}{3} + \frac{1}{2} - \frac{5}{6} \\ &= \frac{4}{6} - \frac{2}{3}.\end{aligned}$$

$$\mathbb{P}(\text{prime or multiple of } 3) = \frac{1}{3} + \frac{1}{2} - \frac{1}{6} = \frac{2}{3}$$

§1 Formal Setup

Definition 1.1 (Sample Space)

The **sample space** Ω is a set of outcomes.

Definition 1.2 (σ -algebra)

- Let \mathcal{F} a collection of subsets of Ω (called *events*).
- \mathcal{F} is a **σ -algebra** if

F1. $\Omega \in \mathcal{F}$.

F2. $A \in \mathcal{F}$ then $A^c = \Omega \setminus A \in \mathcal{F}$.

F3. \forall countable collections $(A_n)_{n \in \mathbb{N}} \in \mathcal{F}$, the union $\bigcup_{n \in \mathbb{N}} A_n \in \mathcal{F}$ also.

Remark 1. The motivation for F2 is so that $\mathbb{P}(A^c) = 1 - \mathbb{P}(A)$ (the probability of not A is defined as expected).

Definition 1.3 (Probability Measure)

Given σ -algebra \mathcal{F} on Ω , function $\mathbb{P} : \mathcal{F} \rightarrow [0, 1]^a$ is a **probability measure** if

P2. $\mathbb{P}(\Omega) = 1$.

P3. \forall countable collections $(A_n)_{n \in \mathbb{N}}$ of *disjoint* events in \mathcal{F} :

$$\mathbb{P}\left(\bigcup_{n \in \mathbb{N}} A_n\right) = \sum_{n \in \mathbb{N}} \mathbb{P}(A_n).$$

Then $(\Omega, \mathcal{F}, \mathbb{P})$ is a *probability space*.

^aP1. $\mathbb{P}(A) \geq 0$

Example 1.1

Coming back to Example 0.1. $\Omega = \{1, 2, \dots, 6\}$ so

$\mathbb{P}(\Omega) = \mathbb{P}(1 \text{ or } 2 \text{ or } 3 \text{ or } 4 \text{ or } 5 \text{ or } 6) = 1$ and \mathcal{F} is all subsets of Ω .

Question

Why $\mathbb{P} : \mathcal{F} \rightarrow [0, 1]$

and not $\mathbb{P} : \Omega \rightarrow [0, 1]$?

If Ω is countable:

- In general: \mathcal{F} = all subsets of Ω , i.e. $\mathcal{P}(\Omega)$ (the power set).
- $\mathbb{P}(2)$ is shorthand for $\mathbb{P}(\{2\})$.
- \mathbb{P} is determined by $(\mathbb{P}(\{w\}), \forall w \in \Omega)$ (e.g. unfair dice).

If Ω is uncountable:

- E.g. $\Omega = [0, 1]$. Want to choose a real number, all equally likely.
- If $\mathbb{P}(\{0\}) = \alpha > 0$ then $\mathbb{P}\left(\left\{0, 1, \frac{1}{2}, \dots, \frac{1}{n}\right\}\right) = (n+1)\alpha \nless \text{if } n \text{ large as } \mathbb{P} > 1$.
- So $\mathbb{P}(\{0\}) = 0$, or $\mathbb{P}(\{0\})$ is undefined.
- What about $\mathbb{P}\left(\left\{x : x \leq \frac{1}{3}\right\}\right)$?
 - ? “Add up” all $\mathbb{P}(\{x\})$ for $x \leq \frac{1}{3}$. However this range is uncountable and we can’t take a sum of uncountably many terms.

Aside

Question

Can we choose uniformly from an infinite countable set? (E.g. $\Omega = \mathbb{N}$ or $\Omega = \mathbb{Q} \cap [0, 1]$)

Answer

No it is not possible but that’s ok there \exists lots of interesting probability measures of \mathbb{N} !

Proof. Suppose possible

- $\mathbb{P}(\{0\}) = \alpha > 0 \quad \forall \omega \in \Omega$. Then $\mathbb{P}(\Omega) = \sum_{\omega \in \Omega} \mathbb{P}(\{\omega\}) = \sum_{\omega \in \Omega} \alpha = \infty$. \nless of $\mathbb{P}2 : \mathbb{P}(\Omega) = 1$.
- $\mathbb{P}(\{0\}) = 0 \quad \forall \omega \in \Omega$. Then $\mathbb{P}(\Omega) = \sum_{\omega \in \Omega} \mathbb{P}(\{\omega\}) = \sum_{\omega \in \Omega} 0 = 0$.

□

Proposition 1.1 (From the axioms)

- $\mathbb{P}(A^c) = 1 - \mathbb{P}(A)$

Proof. A, A^c are disjoint. $A \cup A^c = \Omega$.
 $\implies \mathbb{P}(A) + \mathbb{P}(A^c) \stackrel{P_3}{=} \mathbb{P}(\Omega) \stackrel{P_2}{=} 1$

□

- $\mathbb{P}(\emptyset) = 0$
- If $A \subseteq B$ then $\mathbb{P}(A) \leq \mathbb{P}(B)$
- $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$

§1.1 Examples of Probability Spaces

Example 1.2 (Uniform Choice)

Ω finite, $\Omega = \{\omega_1, \dots, \omega_n\}$, \mathcal{F} = all subsets. *uniform* choice (equally likely)

$$\mathbb{P} : \mathcal{F} \rightarrow [0, 1], \quad \mathbb{P}(A) = \frac{|A|}{|\Omega|}.$$

In particular: $\mathbb{P}(\{\omega\}) = \frac{1}{|\Omega|} \quad \forall \omega \in \Omega$.

Example 1.3 (Choosing without replacement)

n indistinguishable marbles labelled $\{1, \dots, n\}$. Pick $k \leq n$ marbles uniformly at random. Here: $\Omega = \{A \subseteq \{1, \dots, n\}, |A| = k\}$ $|\Omega| = \binom{n}{k}$

Example 1.4 (Well-shuffled deck of cards)

Uniformly chosen *permutation* of 52 cards.

$$\Omega = \{\text{all permutations of 52 cards}\}$$

$$|\Omega| = 52!$$

$$\mathbb{P}(\text{first three cards have the same suit}) = \frac{52 \times 12 \times 11 \times 49!}{52!} = \frac{22}{425}$$

$$\text{Note: } = \frac{12}{51} \times \frac{11}{50}$$

Example 1.5 (Coincident Birthdays)

There are n people; what is the probability that at least two of them share a birthday?

Assumptions:

- No leap years! (365 days)
- All birthdays are equally likely

Let $\Omega = \{1, \dots, 365\}^n$ and $\mathcal{F} = \mathcal{P}(\Omega)$.

Let $A = \{\text{at least two people share the same birthday}\}$ and so $A^c = \{\text{all } n \text{ birthdays are different}\}$.

$$\mathbb{P}(A^c) = \frac{|A^c|}{|\Omega|} = \frac{365 \times 364 \cdots \times (365 - n + 1)}{365^n}$$
$$\mathbb{P}(A) = 1 - \mathbb{P}(A^c)$$

Note that at $n = 22$, $\mathbb{P}(A) \approx 0.476$ and at $n = 23$, $\mathbb{P}(A) \approx 0.507$. So when there are at least 23 people in a room, the probability that two of them share a birthday is around 50%.

KEY IDEA: Calculating $\mathbb{P}(A^c)$ is easier than $\mathbb{P}(A)$.

$$\mathbb{P}(X_n = n) = \frac{1}{2^n}$$

$$\mathbb{P}(X_n = 0) = 0 \text{ if } n \text{ is odd}$$

What about $\mathbb{P}(X_n = 0)$ when n is even

Idea - Choose $\frac{n}{2}$ k s for $X_k = X_{k-1} + 1$ and the rest $X_k = X_{k-1} - 1$ (i.e. go up half the time and down the other half).

$$\begin{aligned}\mathbb{P}(X_n = 0) &= 2^{-n} \binom{n}{\frac{n}{2}} \\ &= \frac{n!}{2^n \left(\frac{n}{2}!\right)^2}\end{aligned}$$

Question

What happens when n is large?

§2.3 Stirling's Formula

Notation. Let $(a_n), (b_n)$ be two sequences. Say $a_n \sim b_n$ as $n \rightarrow \infty$ if $\frac{a_n}{b_n} \rightarrow 1$ as $n \rightarrow \infty$.

Example 2.1

$$n^2 + 5n + \frac{6}{n} \sim n^2$$

Example 2.2 (Non-Example)

$$\exp\left(n^2 + 5n + \frac{6}{n}\right) \not\sim \exp(n^2)$$


Theorem 2.1 (Stirling)

$$n! \sim \sqrt{2\pi n} n^{n+\frac{1}{2}} e^{-n} \text{ as } n \rightarrow \infty.$$

Theorem 2.2 (Weaker Version)

$$\log n! \sim n \log n.$$

Proof. $\log(n!) = \log 2 + \cdots + \log n.$



$$\begin{array}{c}
 \text{"Upper Integral"} \\
 \int_1^n \log x \, dx \leq \log n! \leq \int_1^{n+1} \log x \, dx \\
 \underbrace{n \log n - n + 1}_{\sim n \log n} \leq \log n! \leq \underbrace{(n+1) \log(n+1) - n}_{\sim n \log n}
 \end{array}$$

Key idea: Sandwiching between lower/upper integrals. It was useful that

- $\log x$ is increasing
- $\log x$ has a nice integral!

□

§2.4 (Ordered) compositions

Definition 2.1 (Composition)

A **composition** of m with k parts is a sequence (m_1, \dots, m_k) of non-negative integers with $m_1 + \dots + m_k = m$.

Example 2.3

$$\begin{array}{c}
 3 + 0 + 1 + 2 = 6 \neq 1 + 2 + 0 + 3 = 6 \\
 \star \star \star || \star | \star \star
 \end{array}$$

There is a bijection between compositions *and* sequences of m stars and $(k-1)$ dividers. So the number of compositions is $\binom{m+k-1}{m}$.

Comment: Easy to mistake k with $k-1$ in no. of dividers.

§3 Properties of Probability measures

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and $\mathbb{P} : \mathcal{F} \rightarrow [0, 1]$.

Definition 3.1 (Countable additivity)

P3 : $\mathbb{P}(\bigcup_{n \in \mathbb{N}} A_n) = \sum_{n \in \mathbb{N}} \mathbb{P}(A_n)$ for $(A_n)_{n \in \mathbb{N}}$ disjoint.

Question

What if the sets are not disjoint?

§3.1 Countable sub-additivity


Proposition 3.1 (Countable sub-additivity)

Let $(A_n)_{n \in \mathbb{N}}$ be a sequence of events in \mathcal{F} . Then

$$\mathbb{P}\left(\bigcup_{n \in \mathbb{N}} A_n\right) \leq \sum_{n \in \mathbb{N}} \mathbb{P}(A_n).$$


May also be called a *union bound*.

Intuition:



$\sum_{n \in \mathbb{N}} \mathbb{P}(A_n)$ “double counts” some sub-events.

Proof. Idea: Rewrite $\bigcup_{n \in \mathbb{N}} A_n$ as a *disjoint* union. Define $B_1 = A_1$ and $B_n = \underbrace{A_n \setminus (A_1 \cup \dots \cup A_{n-1})}_{\in \mathcal{F} \text{ (by Sheet 1)}} \quad \forall n \geq 2$.



So

- $\bigcup_{n \in \mathbb{N}} B_n = \bigcup_{n \in \mathbb{N}} A_n$.
- $(B_n)_{n \in \mathbb{N}}$ is disjoint (by construction).
- $B_n \subseteq A_n \implies \mathbb{P}(B_n) \leq \mathbb{P}(A_n)$
Q4, Sheet 1

$$\mathbb{P}\left(\bigcup_{n \in \mathbb{N}} A_n\right) = \mathbb{P}\left(\bigcup_{n \in \mathbb{N}} B_n\right) \stackrel{P3 \text{ on } (B_n)}{=} \sum_{n \in \mathbb{N}} \mathbb{P}(B_n) \leq \sum_{n \in \mathbb{N}} \mathbb{P}(A_n)$$

□

§3.2 Continuity

Proposition 3.2 (Continuity)

Let $(A_n)_{n \in \mathbb{N}}$ be an increasing sequence of events in \mathcal{F} , i.e. $A_n \subseteq A_{n+1} \quad \forall n$. Then $\mathbb{P}(A_n) \leq \mathbb{P}(A_{n+1})$. So $\mathbb{P}(A_n)$ converges as $n \rightarrow \infty$.^a


In fact: $\lim_{n \rightarrow \infty} \mathbb{P}(A_n) = \mathbb{P}(\bigcup_{n \in \mathbb{N}} A_n)$.

^aAs probabilities are bounded above by 1 and increasing.

For motivation try Q6, Sheet 1.

Proof. Let us reuse the B_n s from the previous subsection.

- $\bigcup_{k=1}^n B_k = A_n$ (disjoint union).
- $\bigcup_{n \in \mathbb{N}} B_n = \bigcup_{n \in \mathbb{N}} A_n$



$$\mathbb{P}(A_n) = \sum_{k=1}^n \mathbb{P}(B_k) \xrightarrow{n \rightarrow \infty} \sum_{k \geq 1} \mathbb{P}(B_k) = \mathbb{P}\left(\bigcup_{n \in \mathbb{N}} B_n\right) = \mathbb{P}\left(\bigcup_{n \in \mathbb{N}} A_n\right)$$

□

§3.3 Inclusion-Exclusion Principle

Background: $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$.

Similarly: $A, B, C \in \mathcal{F}$

$$\mathbb{P}(A \cup B \cup C) = \mathbb{P}(A) + \mathbb{P}(B) + \mathbb{P}(C) - \mathbb{P}(A \cup B) - \mathbb{P}(B \cup C) - \mathbb{P}(C \cup A) + \mathbb{P}(A \cap B \cap C).$$

Proposition 3.3 (Inclusion-Exclusion Principle)

Let $A_1, \dots, A_n \in \mathcal{F}$, then:

$$\begin{aligned} \mathbb{P}\left(\bigcup_{i=1}^n A_i\right) &= \sum_{i=1}^n \mathbb{P}(A_i) - \sum_{1 \leq i_1 < i_2 \leq n} \mathbb{P}(A_{i_1} \cap A_{i_2}) + \sum_{1 \leq i_1 < i_2 < i_3 \leq n} \mathbb{P}(A_{i_1} \cap A_{i_2} \cap A_{i_3}) - \dots \\ &\quad + (-1)^{n+1} \mathbb{P}(A_1 \cap \dots \cap A_n) \\ &= \sum_{\substack{I \subset \{1, \dots, n\} \\ I \neq \emptyset}} (-1)^{|I|+1} \mathbb{P}\left(\bigcap_{i \in I} A_i\right) \end{aligned}$$

Note: $\sum_{1 \leq i_1 < i_2 \leq n}$ is the sum of all triples that are distinct and unordered.

Proof. By induction. For $n = 2$ it holds (Q4e, Sheet 1).

$$\mathbb{P}\left(\bigcup_{i=1}^n A_i\right) = \mathbb{P}\left(\left(\bigcup_{i=1}^{n-1} A_i\right) \cup A_n\right)$$

Using $n = 2$ case we get:

$$= \mathbb{P} \left(\bigcup_{i=1}^{n-1} A_i \right) + \mathbb{P}(A_n) - \mathbb{P} \left(\left(\bigcup_{i=1}^{n-1} A_i \right) \cap A_n \right)$$

We want to break down the final element on the RHS

$$\text{Idea: } \left(\bigcup_{i=1}^{n-1} A_i \right) \cap A_n = \bigcup_{i=1}^{n-1} (A_i \cap A_n)$$

If we apply IEP to $\bigcup_{i=1}^{n-1} (A_i \cap A_n)$ we need to calculate $\bigcap_{i \in J} (A_i \cap A_n)$

$$\bigcap_{i \in J} (A_i \cap A_n) = \bigcap_{i \in J \cup \{n\}} A_i, \quad J \subset \{1, \dots, n-1\}$$

$$\begin{aligned} \mathbb{P} \left(\bigcup_{i=1}^n A_i \right) &= \underbrace{\sum_{\substack{J \subset \{1, \dots, n-1\} \\ J \neq \emptyset}} (-1)^{|J|+1} \mathbb{P} \left(\bigcap_{i \in J} A_i \right)}_{n-1 \text{ case}} + \mathbb{P}(A_n) \\ &\quad - \underbrace{\sum_{\substack{J \subset \{1, \dots, n-1\} \\ J \neq \emptyset}} (-1)^{|J|+1} \mathbb{P} \left(\bigcap_{i \in J \cup \{n\}} A_i \right)}_{n-1 \text{ case on } (A_i \cap A_n)} \end{aligned}$$

$$J \cup \{n\} \mapsto I.$$

$$-(-1)^{|J|+1} \mapsto (-1)^{|I|+1}$$

$$= \underbrace{\sum_{\substack{I \subset \{1, \dots, n-1\} \\ I \neq \emptyset}} (-1)^{|I|+1} \mathbb{P} \left(\bigcap_{i \in I} A_i \right)}_{\text{Just changed the labels}} + \mathbb{P}(A_n)$$

$$+ \sum_{\substack{I \subset \{1, \dots, n\} \\ n \in I, |I| > 2}} (-1)^{|I|+1} \mathbb{P} \left(\bigcap_{i \in I} A_i \right)$$

$$= \sum_{\substack{I \subset \{1, \dots, n\} \\ I \neq \emptyset}} (-1)^{|I|+1} \mathbb{P} \left(\bigcap_{i \in I} A_i \right).$$

Let us check that we have indeed counted all subsets I .

- $\sum_{\substack{I \subset \{1, \dots, n-1\} \\ I \neq \emptyset}} (-1)^{|I|+1} \mathbb{P} \left(\bigcap_{i \in I} A_i \right)$ accounts for all subsets where $n \notin I$.
- $\mathbb{P}(A_n)$ accounts for $\{n\}$
- $\sum_{\substack{I \subset \{1, \dots, n\} \\ n \in I, |I| > 2}} (-1)^{|I|+1} \mathbb{P} \left(\bigcap_{i \in I} A_i \right)$ accounts for all subsets where $n \in I$ and $I \neq \{n\}$.

□

§3.4 Bonferroni Inequalities

Question

What if you *truncate* IEP (Inclusion-Exclusion Principle)?

Proposition 3.4 (Bonferroni Inequality)

Recall: Countable sub-additivity - $\mathbb{P}(\cup A_i) \leq \sum \mathbb{P}(A_i)$.

$$\begin{aligned} \mathbb{P} \left(\bigcup_{i=1}^n A_i \right) &\leq \sum_{k=1}^r (-1)^{k+1} \sum_{i_1 < i_2 < \dots < i_k} \mathbb{P}(A_{i_1} \cap \dots \cap A_{i_k}) \quad \text{if } r \text{ is odd} \\ \mathbb{P} \left(\bigcup_{i=1}^n A_i \right) &\geq \sum_{k=1}^r (-1)^{k+1} \sum_{i_1 < i_2 < \dots < i_k} \mathbb{P}(A_{i_1} \cap \dots \cap A_{i_k}) \quad \text{if } r \text{ is even} \end{aligned}$$

Proof. By induction on r and n . Let r be odd

$$\begin{aligned} \mathbb{P} \left(\bigcup_{i=1}^n A_i \right) &= \mathbb{P} \left(\bigcup_{i=1}^{n-1} A_i \right) + \mathbb{P}(A_n) - \mathbb{P} \left(\bigcup_{i=1}^{n-1} (A_i \cap A_n) \right) \\ \mathbb{P} \left(\bigcup_{i=1}^{n-1} A_i \right) &\leq \underbrace{\sum_{\substack{J \subset \{1, \dots, n-1\} \\ 1 \leq |J| \leq r}} (-1)^{|J|+1} \mathbb{P} \left(\bigcap_{i \in J} A_i \right)}_{n-1 \text{ case.}} \\ \mathbb{P} \left(\bigcup_{i=1}^{n-1} (A_i \cap A_n) \right) &\geq \underbrace{\sum_{\substack{J \subset \{1, \dots, n-1\} \\ 1 \leq |J| \leq r-1^a}} (-1)^{|J|+1} \mathbb{P} \left(\bigcap_{i \in J \cup \{n\}} A_i \right)}_{r-1 \text{ case on } (A_i \cap A_n)} \end{aligned}$$

Using the same rearranging as in the proof of [Inclusion-Exclusion Principle](#)

$$\mathbb{P}\left(\bigcup_{i=1}^n A_i\right) \leq \sum_{\substack{I \subseteq \{1, \dots, n\} \\ 1 \leq |I| \leq r}} (-1)^{|I|+1} \mathbb{P}\left(\bigcap_{i \in I} A_i\right).$$

The case of r being even is similar, simply note all three inequalities are reversed. \square

^a J doesn't include n and we only want r elements in the intersection

^a J doesn't include n and we only want r elements in the intersection

Question

When is it good to truncate at e.g. $r = 2$?

§3.5 Counting with IEP

Uniform probability measure on Ω , $|\Omega| < \infty$. $\mathbb{P}(A) = \frac{|A|}{|\Omega|} \quad \forall A \subseteq \Omega$. Then $\forall A_1, \dots, A_n \subseteq \Omega$

$$|A_1 \cup \dots \cup A_n| = \sum_{k=1}^n (-1)^{k+1} \sum_{i_1 < \dots < i_k} |A_{i_1} \cap \dots \cap A_{i_k}|$$

(and similarly for Bonferroni Inequalities).

Example 3.1 (Surjections)

What is the probability that a function $f : \{1, \dots, n\} \rightarrow \{1, \dots, m\}$, $n \geq m$ is a surjection? Let $\Omega = \{f : \{1, \dots, n\} \rightarrow \{1, \dots, m\}\}$ and $A = \{f \in \Omega : \text{Image}(f) = \{1, \dots, m\}\}$.

$\forall i \in \{1, \dots, m\}$ define $B_i = \{f \in \Omega : i \notin \text{Image}(f)\}$.

Key observations:

-

$$\begin{aligned} A &= B_1^c \cap \dots \cap B_m^c \\ &= (B_1 \cup \dots \cup B_m)^c \end{aligned}$$

- $|B_{i_1} \cap \dots \cap B_{i_k}|$ is nice to calculate.

$$\begin{aligned} |B_{i_1} \cap \dots \cap B_{i_k}| &= |\{f \in \Omega : i_1, \dots, i_k \notin \text{Image}(f)\}| \\ &= (m - k)^n \end{aligned}$$

$$\begin{aligned}
IEP \rightarrow |B_1 \cup \dots \cup B_m| &= \sum_{k=1}^m (-1)^{k+1} \sum_{\substack{i_1 < \dots < i_k \\ \text{same for all } i_1, \dots, i_k}} |B_{i_1} \cap \dots \cap B_{i_k}| \\
&= \sum_{k=1}^m (-1)^{k+1} \binom{m}{k} (m-k)^n \\
|A| &= m^n - |B_{i_1} \cap \dots \cap B_{i_k}| \\
&= \sum_{k=0}^m (-1)^k \binom{m}{k} (m-k)^n
\end{aligned}$$

Example 3.2 (Derangements)

What is the probability that a permutation has no fixed points? Derangements can be useful in a Secret Santa.

$\Omega = \{\text{permutations of } \{1, \dots, n\}\}$ and the derangements, D , are $\{\sigma \in \Omega : \sigma(i) \neq i \ \forall i = 1, \dots, n\}$.

Question

Is $\mathbb{P}(D) = \frac{|D|}{|\Omega|}$ large or small (e.g. when $n \rightarrow \infty$)?

$\forall i \in \{1, \dots, n\} : A_i = \{\sigma \in \Omega : \sigma(i) = i\}$.

Key observations:

- $D = A_1^c \cap \dots \cap A_n^c = (\bigcup_{i=1}^n A_i)^c$.
-

$$\begin{aligned}
\mathbb{P}(A_{i_1} \cap \dots \cap A_{i_k}) &= \frac{(n-k)!}{n!} \\
IEP \rightarrow \mathbb{P}\left(\bigcup_{i=1}^n A_i\right) &= \sum_{k=1}^n (-1)^{k+1} \sum_{i_1 < \dots < i_k} \mathbb{P}(A_{i_1} \cap \dots \cap A_{i_k}) \\
&= \sum_{k=1}^n (-1)^{k+1} \binom{n}{k} \frac{(n-k)!}{n!} \\
&= \sum_{k=1}^n (-1)^{k+1} \frac{1}{k!} \\
\mathbb{P}(D) &= 1 - \mathbb{P}\left(\bigcup_{i=1}^n A_i\right) \\
&= 1 - \sum_{k=1}^n \frac{(-1)^{k+1}}{k!}
\end{aligned}$$

$$\begin{aligned}
&= \sum_{k=0}^n \frac{(-1)^k}{k!} \\
\lim_{n \rightarrow \infty} \mathbb{P}(D) &= \sum_{k=0}^{\infty} \frac{(-1)^{k+1}}{k!} \\
&= e^{-1} \approx 0.37.
\end{aligned}$$

Remark 2.

- What if instead $\Omega' = \{\text{all functions } f : \{1, \dots, n\} \text{ to itself}\}$?

$$\begin{aligned}
D &= \{f \in \Omega' : f(i) \neq i \quad \forall i = 1, \dots, n\}. \\
\mathbb{P}(D) &= \frac{(n-1)^n}{n^n} \\
&= \left(1 - \frac{1}{n}\right)^n \\
\lim_{n \rightarrow \infty} \mathbb{P}(D) &= e^{-1}.
\end{aligned}$$

- We would liked to have calculated $\mathbb{P}(D)$ by doing $\left(\frac{n-1}{n}\right)^n$ as we have n choices each with probability $\frac{n-1}{n}$. We will be allowed to do this soon! See Example 3.6
- $f(i)$ is a random quantity associated to Ω . We will be allowed to study $f(i)$ as a *random variable* soon.
- We are allowed to toss a fair coin n times, $\Omega = \{H, T\}^n$. But we have not yet studied tossing an unfair coin n times.

§3.6 Independence

$(\Omega, \mathcal{F}, \mathbb{P})$ as before.

Definition 3.2 (Independence)

Events $A, B \in \mathcal{F}$ are **independent** ($A \perp\!\!\!\perp B$) if

$$\mathbb{P}(A \cap B) = \mathbb{P}(A)\mathbb{P}(B).$$

A countable^a collection of events (A_n) are **independent** if \forall distinct i_1, \dots, i_k ^b we have:

$$\mathbb{P}(A_{i_1} \cap \dots \cap A_{i_k}) = \prod_{j=1}^k \mathbb{P}(A_{i_j}).$$

^aincluding finite
^b k is finite

Remark 3 (Caution). “Pairwise independence” does not imply independence.

Example 3.3

$$\begin{aligned}\Omega &= \{(H, H), (H, T), (T, H), (T, T)\} \\ \mathbb{P}(\{\omega\}) &= \frac{1}{4} \quad \forall \omega \in \Omega. \\ A &= \text{first coin is } H = \{(H, H), (H, T)\}. \\ B &= \text{second coin is } H = \{(T, H), (H, H)\}. \\ C &= \text{both coins have the same outcome} = \{(T, T), (H, H)\}. \\ \mathbb{P}(A) &= \mathbb{P}(B) = \mathbb{P}(C) = \frac{1}{2}. \\ A \cap B &= A \cap C = B \cap C = \{(H, H)\}. \\ \mathbb{P}(A \cap B) &= \mathbb{P}(A \cap C) = \mathbb{P}(B \cap C) = \frac{1}{4}. \quad \text{Pairwise independence } \checkmark \\ \mathbb{P}(A \cap B \cap C) &= \frac{1}{4} \neq \mathbb{P}(A)\mathbb{P}(B)\mathbb{P}(C). \quad \text{Independence } \times\end{aligned}$$

Example 3.4 (Independence)

•

$$\begin{aligned}\Omega' &= \{\text{all functions } f : \{1, \dots, n\} \text{ to itself}\} \\ A_i &= \{f \in \Omega' : f(i) = i\}. \\ \mathbb{P}(A_i) &= \frac{n^{n-1}}{n^n} = \frac{1}{n} \\ \mathbb{P}(A_{i_1} \cap \dots \cap A_{i_k}) &= \frac{n^{n-k}}{n^n} \\ &= \frac{1}{n^k} \\ &= \prod_{j=1}^k \mathbb{P}(A_{i_j})\end{aligned}$$

Here: (A_i) are independent events.

•

$$\begin{aligned}\Omega &= \{\sigma : \text{permutation of } \{1, \dots, n\}\} \\ A_i &= \{\sigma \in \Omega : \sigma(i) = i\} \\ \mathbb{P}(A_i) &= \frac{(n-1)!}{n!} = \frac{1}{n}. \\ i \neq j \quad \mathbb{P}(A_i \cap A_j) &= \frac{(n-2)!}{n!} \\ &= \frac{1}{n(n-1)} \\ &\neq \mathbb{P}(A_i)\mathbb{P}(A_j)\end{aligned}$$

Here: (A_i) are not independent events.

§3.6.1 Properties

Claim 3.1

If A is independent of B , then A is also independent of B^c .

Proof.

$$\begin{aligned}\mathbb{P}(A \cap B^c) &= \mathbb{P}(A) - \mathbb{P}(A \cap B) \\ &= \mathbb{P}(A) - \mathbb{P}(A)\mathbb{P}(B) \\ &= \mathbb{P}(A)[1 - \mathbb{P}(B)] \\ &= \mathbb{P}(A)\mathbb{P}(B^c)\end{aligned}$$

□

Claim 3.2

A is independent of $B = \Omega$ and of $C = \emptyset$

Proof. $\mathbb{P}(A \cap \Omega) = P(A) = \mathbb{P}(A) \underbrace{\mathbb{P}(\Omega)}_{=1}$ and so $A \perp\!\!\!\perp \emptyset$ by Claim 1.

□

§3.7 Conditional Probability

$(\Omega, \mathcal{F}, \mathbb{P})$ as before.

Consider $B \in \mathcal{F}$ with $\mathbb{P}(B) > 0$, $A \in \mathcal{F}$

Definition 3.3 (Conditional Probability)

The **conditional probability** of A given B is $\mathbb{P}(A \mid B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}$.

“The probability of A is we know B happened”. (e.g. revealing information in succession)

Example 3.5

A, B independent.

$$\mathbb{P}(A \mid B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)} = \frac{\mathbb{P}(A)\mathbb{P}(B)}{\mathbb{P}(B)} = \mathbb{P}(A)$$

“Knowing whether B happened doesn’t affect the probability of A ”.

§3.7.1 Properties

- P1 - $\mathbb{P}(A \mid B) \geq 0$
- P2 - $\mathbb{P}(B \mid B) = 1 = \mathbb{P}(\Omega \mid B)$
- P3 - (A_n) disjoint events $\in \mathcal{F}$:

Claim 3.3

$$\mathbb{P}\left(\bigcup_{n \in \mathbb{N}} A_n \mid B\right) = \sum_{n \in \mathbb{N}} \mathbb{P}(A_n \mid B)$$

Proof.

$$\begin{aligned} \mathbb{P}\left(\bigcup_{n \in \mathbb{N}} A_n \mid B\right) &= \frac{\mathbb{P}((\bigcup_{n \in \mathbb{N}} A_n) \cap B)}{\mathbb{P}(B)} \\ &= \frac{\mathbb{P}(\bigcup_{n \in \mathbb{N}} (A_n \cap B))}{\mathbb{P}(B)}, \quad \bigcup_{n \in \mathbb{N}} (A_n \cap B) \text{ is a disjoint union} \\ &= \frac{\sum_{n \in \mathbb{N}} \mathbb{P}(A_n \cap B)}{\mathbb{P}(B)} \\ &= \sum_{n \in \mathbb{N}} \mathbb{P}(A_n \mid B) \end{aligned}$$

Summary: Use definition and apply P1, P2, P3 to the numerator. □

$\mathbb{P}(\bullet \mid B)$ is a function from $\mathcal{F} \rightarrow [0, 1]$ that satisfies the rules to be a probability measure on Ω .

Aside?

Consider $\Omega' = B$ (especially in finite or countable setting). Let $\mathcal{F}' = \mathcal{P}(B)$. Then $(\Omega', \mathcal{F}', \mathbb{P}(\bullet \mid B))$ also satisfies rules to be a probability measure on Ω' .

$$\begin{aligned}\mathbb{P}(A \cap B) &= \mathbb{P}(A)\mathbb{P}(B \mid A) \\ \mathbb{P}(A_1 \cap A_2 \cap \dots \cap A_n) &= \mathbb{P}(A_1)\mathbb{P}(A_2 \mid A_1)\mathbb{P}(A_3 \mid A_1 \cap A_2) \dots \mathbb{P}(A_n \mid A_1 \cap \dots \cap A_{n-1})\end{aligned}\tag{1}$$

Example 3.6

Uniform choice of a permutation $(\sigma(1), \sigma(2), \dots, \sigma(n)) \in \Sigma_n$.

Claim 3.4

$$\begin{aligned}\mathbb{P}(\sigma(k) = i_k \mid \sigma(1) = i_1, \dots, \sigma(k-1) = i_{k-1}), \quad i_1, \dots, i_{k-1} \text{ distinct.} \\ = \begin{cases} 0 & \text{if } i_k \in \{i_1, \dots, i_{k-1}\}. \\ \frac{1}{n-k+1} & \text{else} \end{cases}\end{aligned}$$

^aThis is an example of [\(Ordered\) compositions](#)

^aThis is an example of [\(Ordered\) compositions](#)

Proof.

$$\begin{aligned}\mathbb{P}(\sigma(k) = i_k \mid \sigma(1) = i_1, \dots, \sigma(k-1) = i_{k-1}) &= \frac{\mathbb{P}(\sigma(1) = i_1, \dots, \sigma(k) = i_k)}{\mathbb{P}(\sigma(1) = i_1, \dots, \sigma(k-1) = i_{k-1})} \\ &= \frac{\frac{(n-k)!}{n!}}{\frac{(n-k+1)!}{n!}} \\ &= \frac{(n-k)!}{(n-k+1)!} \\ &= \frac{1}{n-k+1} \\ \mathbb{P}(\sigma(1) = i_1, \dots, \sigma(k) = i_k) &= 0 \quad \text{if } i_k \in \{i_1, \dots, i_{k-1}\}.\end{aligned}$$

□

§3.7.2 Law of Total Probability and Bayes' Formula

Definition 3.4 (Partition)

$(B_1, B_2, \dots)^a \in \Omega$ is a **partition** of Ω if:

- $\Omega = \bigcup_n B_n$
- (B_n) are disjoint

^afinite or countable

Theorem 3.1 (Law of Total Probability)

(B_n) a finite or countable partition of Ω with $B_n \in \mathcal{F} \forall n$ s.t. $\mathbb{P}(B_n) > 0$. Then $\forall A \in \mathcal{F}$:

$$\mathbb{P}(A) = \sum_n \mathbb{P}(A \mid B_n) \mathbb{P}(B_n).$$

Also known as “Partition Theorem”.

Proof. Note that $\bigcup_n (A \cap B_n) = A$.

$$\begin{aligned} \mathbb{P}(A) &= \sum_{n \in \mathbb{N}} \mathbb{P}(A \cap B_n) \\ &= \sum_n \mathbb{P}(A \mid B_n) \mathbb{P}(B_n) \quad \text{by Equation (1)} \end{aligned}$$

□

Theorem 3.2 (Bayes' Formula)

Same setup as above

$$\begin{aligned} \mathbb{P}(B_n \mid A) &= \frac{\mathbb{P}(A \cap B_n)}{\mathbb{P}(A)} \\ &= \frac{\mathbb{P}(A \mid B_n) \mathbb{P}(B_n)}{\sum_m \mathbb{P}(A \mid B_m) \mathbb{P}(B_m)}. \end{aligned}$$

Let $n = 2$: $\mathbb{P}(B \mid A) \mathbb{P}(A) = \mathbb{P}(A \mid B) \mathbb{P}(B) = \mathbb{P}(A \cap B)$

Example 3.7 (Lecture course)

Consider a Lecture course which has 2/3 of the lectures on weekdays and 1/3 on

weekends. Let

$$\begin{aligned}\mathbb{P}(\text{forget notes} \mid \text{weekday}) &= \frac{1}{8} \\ \mathbb{P}(\text{forget notes} \mid \text{weekend}) &= \frac{1}{2}\end{aligned}$$

What is $\mathbb{P}(\text{weekend} \mid \text{forget notes})$? Let $B_1 = \{\text{weekday}\}$, $B_2 = \{\text{weekend}\}$ and $A = \{\text{forget notes}\}$.

By LTP ([Law of Total Probability](#)): $\mathbb{P}(A) = \frac{2}{3} \times \frac{1}{8} + \frac{1}{3} \times \frac{1}{2} = \frac{1}{4}$.

By [Bayes' Formula](#): $\mathbb{P}(B_2 \mid A) = \frac{1}{3} \times \frac{1/2}{1/4} = \frac{2}{3}$.

Example 3.8 (Disease Testing)

Suppose p are infected and $(1 - p)$ are not. $\mathbb{P}(\text{tests positive} \mid \text{infected}) = 1 - \alpha$ and $\mathbb{P}(\text{tests positive} \mid \text{not infected}) = \beta$ where $\alpha, \beta \in (0, 1)$.

We want to work out $\mathbb{P}(\text{infected} \mid \text{test positive})$.

By LTP: $\mathbb{P}(\text{test positive}) = p(1 - \alpha) + (1 - p)\beta$.

By [Bayes' Formula](#): $\mathbb{P}(\text{infected} \mid \text{test positive}) = \frac{p(1-\alpha)}{p(1-\alpha) + (1-p)\beta}$.

Suppose $p \ll \beta$ then $p(1 - \alpha) \ll (1 - p)\beta$ so $\mathbb{P}(\text{infected} \mid \text{test positive}) \sim \frac{p(1-\alpha)}{(1-p)\beta} \sim \frac{p}{\beta}$ which is small.

Example 3.9 (Simpson's Paradox)

Scientists ask: do jelly beans make you tongue change colour?

Oxford	Change	No change	% change	$\Delta = 3\%$
Blue	15	22	41%	
Green	5	8	38%	

Cambridge	Change	No change	% change	$\Delta = 15\%$
Blue	10	3	77%	
Green	23	14	62%	

Total	Change	No change	% change	$\Delta = -6\%$
Blue	25	25	50%	
Green	28	22	56%	

The conclusion from this example should be that the Cambridge methodology is different to the Oxford one rather than anything about blue/ green jelly beans.^a

Let $A = \{\text{change colour}\}$, $B = \{\text{blue}\}$, $B^c = \{\text{green}\}$, $C = \{\text{Cambridge}\}$, $C^c = \{\text{Oxford}\}$.

$$\begin{aligned}\mathbb{P}(A \mid B \cap C) &> \mathbb{P}(A \mid B^c \cap C) \\ \mathbb{P}(A \mid B \cap C^c) &> \mathbb{P}(A \mid B^c \cap C^c) \\ \not\Rightarrow \mathbb{P}(A \mid B) &> \mathbb{P}(A \mid B^c)\end{aligned}$$

^aObviously this is a frivolous example however if we changed Oxford to November 2021, Cambridge to January 2021 and we were measuring vaccine efficacy for different vaccines we would get similar results. And it would be reasonable to conclude that the main underlying factor was a change in the viral landscape rather than waning efficacy.

Theorem 3.3 (Law of Total Probability for Conditional Probabilities)

Suppose C_1, C_2, \dots a partition of B .

$$\mathbb{P}(A \mid B) = \sum_n \mathbb{P}(A \mid C_n) \mathbb{P}(C_n \mid B)$$

Proof.

$$\begin{aligned}\mathbb{P}(A \mid B) &= \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)} \\ &= \frac{\mathbb{P}(A \cap (\bigcup_n C_n))}{\mathbb{P}(B)} \\ &= \frac{\mathbb{P}(\bigcup_n (A \cap C_n))}{\mathbb{P}(B)} \\ &= \frac{\sum_n \mathbb{P}(A \cap C_n)}{\mathbb{P}(B)} \\ &= \frac{\sum_n \mathbb{P}(A \mid C_n) \mathbb{P}(C_n)}{\mathbb{P}(B)} \\ &= \sum_n \mathbb{P}(A \mid C_n) \frac{\mathbb{P}(B \cap C_n)}{\mathbb{P}(B)} \quad C_n \subset B \implies B \cap C_n = C_n \\ &= \sum_n \mathbb{P}(A \mid C_n) \mathbb{P}(C_n \mid B)\end{aligned}$$

□

Non Examinable

Special Case:

- If all $\mathbb{P}(C_n)$ are equal then so are $\mathbb{P}(C_n \mid B)$. Note $\sum_n \mathbb{P}(C_n \mid B) = 1$.
- If $\mathbb{P}(A \mid C_n)$ are all equal.

Then $\mathbb{P}(A \mid B) = \mathbb{P}(A \mid C_n)$.

Example 3.10 (Well-shuffled deck of cards)

Uniformly chosen *permutation*, $\sigma \in \Sigma_{52}$, of 52 cards. $\{1, 2, 3, 4\}$ are *aces*. Let $A = \{\sigma(1), \sigma(2) \text{ are aces}\}$, $B = \{\sigma(1) \text{ is an ace}\} = \{\sigma(1) \leq 4\}$, $C_1 = \{\sigma(1) = 1\} \dots C_4 = \{\sigma(1) = 4\}$.

Note:

-

$$\begin{aligned} \mathbb{P}(A \mid C_i) &= \mathbb{P}(\sigma(2) \in \{1, 2, 3, 4\} \mid \sigma(1) = i) \quad i \leq 4 \\ &= \frac{3}{51} \text{ by Example 3.6} \end{aligned}$$

- $\mathbb{P}(C_1) = \dots = \mathbb{P}(C_4) = \frac{1}{52}$.

So $\mathbb{P}(A \mid B) = \frac{3}{51}$ and $\mathbb{P}(A) = \mathbb{P}(B)\mathbb{P}(A \mid B) = \frac{4}{52} \times \frac{3}{51}$

§4 Discrete Random Variables

Motivation: Roll two dice. $\Omega = \{1, \dots, 6\}^2 = \{(i, j) : 1 \leq i, j \leq 6\}$. If we restrict our attention to:

- the first dice e.g. $\{(i, j) : i = 3\}$.
- the sum of the dice e.g. $\{(i, j) : i + j = 8\}$.
- the max of the dice e.g. $\{(i, j) : i, j \leq 4, i \text{ or } j = 4\}$.

This is annoying and we want to move on from sets.

Goal: “Random real-valued measurements”, we want the value of the first dice to be X and the sum to be $X + Y$...


Definition 4.1 (Discrete Random Variable)

A **discrete random variable** X on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ is a function $X : \Omega \rightarrow \mathbb{R}$ s.t.

- $\{\omega \in \Omega : X(\omega) = x\} \in \mathcal{F}$
- $\text{Image}(X)$ is finite or countable (subset of \mathbb{R}).
- We abbreviate $\{\omega \in \Omega : X(\omega) = x\}$ as $\{X = x\}$. So $\mathbb{P}(X = x)$ is valid.
- Often $\text{Image}(X) = \mathbb{Z}$ or \mathbb{N}_0 or $\{0, 1\}$ etc. *not* $\{\text{Heads or Tails}\}$.

If Ω is finite or countable and $\mathcal{F} = \mathcal{P}(\Omega)$ both blue bullet points hold automatically.

Example 4.1 (Part II Applied Probability)



“random arrival process”. Let $\Omega = \{\text{countable subsets } (a_1, a_2, \dots) \text{ of } (0, \infty)\}$ and $N_t = \text{number of arrivals by time } t = |\{a_i : a_i \leq t\}| \in \mathbb{N}_0$ is a discrete RV (random variable) for each time t .

Definition 4.2 (Probability Mass Function)

The **probability mass function** of discrete RV X is the function $p_X : \mathbb{R} \rightarrow [0, 1]$ given by $p_X(x) = \mathbb{P}(X = x) \quad \forall x \in \mathbb{R}$.

Note.

- if $x \notin \text{Image}(X)$ then $p_X(x) = \mathbb{P}(\{\omega \in \Omega : X(\omega) = x\}) = \mathbb{P}(\emptyset) = 0$.
-

$$\begin{aligned}
 \sum_{x \in \text{Im}(X)} p_X(x) &= \sum_{x \in \text{Im}(X)} \underbrace{\mathbb{P}(\{\omega \in \Omega : X(\omega) = x\})}_{\text{disjoint}} \\
 &= \mathbb{P}\left(\bigcup_{x \in \text{Im}(X)} \{\omega \in \Omega : X(\omega) = x\}\right) \\
 &= \mathbb{P}(\Omega) \\
 &= 1
 \end{aligned}$$

Example 4.2 (Indicator function)

For event $A \in \mathcal{F}$, define $1_A : \Omega \rightarrow \mathbb{R}$ by

$$1_A(\omega) = \begin{cases} 1 & \text{if } \omega \in A \\ 0 & \text{else} \end{cases}$$

1_A is a discrete RV with $\text{Image} = \{0, 1\}$. $p_{1_A}(1) = \mathbb{P}(1_A = 1) = \mathbb{P}(A)$, $p_{1_A}(0) = \mathbb{P}(1_A = 0) = \mathbb{P}(A^c)$ and $p_{1_A}(x) = 0 \quad \forall x \notin \{0, 1\}$.

This encodes “did A happen” as a real number.

Remark 4. Given a pmf p_X (probability mass function), we can always construct a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and a RV defined on it with this pmf.

- $\Omega = \text{Im}(X)$ i.e. $\{x \in \mathbb{R} : p_X(x) > 0\}$
- $\mathcal{F} = \mathcal{P}(\Omega)$
- $\mathbb{P}(\{x\}) = p_X(x)$ and extend to all $A \in \mathcal{F}$

§4.1 Discrete Probability Distributions

Ω is finite.

§4.1.1 Bernoulli distribution - “(biased) coin toss”

If $X \sim \text{Bern}(p)$ where $p \in [0, 1]$ then $\text{Im}(X) = \{0, 1\}$ and $p_X(1) = \mathbb{P}(X = 1) = p$ and $p_X(0) = 1 - p$.

Example 4.3

$1_A \sim \text{Bern}(p)$ with $p = \mathbb{P}(A)$.

§4.1.2 Binomial distribution

This can be used to model the number of heads when a coin is tossed n times.

If $X \sim \text{Bin}(n, p)$ where $n \in \mathbb{Z}^+$ and $p \in [0, 1]$ then $\text{Im}(X) = \{0, 1, \dots, n\}$, $p_X(k) = \mathbb{P}(X = k) = \binom{n}{k} p^k (1-p)^{n-k}$. $\sum_{k=0}^n p_X(k) = (p + (1-p))^n = 1$.

§4.1.3 More than one RV

Motivation: Roll a dice with outcome $X \in \{1, 2, \dots, 6\}$. Events: $A = \{1 \text{ or } 2\}$, $B = \{1 \text{ or } 2 \text{ or } 3\}$, $C = \{1 \text{ or } 3 \text{ or } 5\}$. $1_A \sim \text{Bern}\left(\frac{1}{3}\right)$, $1_B \sim \text{Bern}\left(\frac{1}{2}\right)$, $1_C \sim \text{Bern}\left(\frac{1}{2}\right)$.

Note: $1_A \leq 1_B$ for all outcomes
but $1_A \leq 1_C$ for all outcomes *is false*.

Definition 4.3 (Independent RVs)

Let X_1, \dots, X_n be discrete RVs. We say X_1, \dots, X_n are *independent* if:

$$\mathbb{P}(X_1 = x_1, \dots, X_n = x_n) = \mathbb{P}(X_1 = x_1) \dots \mathbb{P}(X_n = x_n) \quad \forall x_1, \dots, x_n \in \mathbb{R}.$$

(suffices to check $\forall x_i \in \text{Im}(X_i)$)

Example 4.4

X_1, \dots, X_n independent RVs each with the Bernoulli(p) distribution. Study $S_n = X_1 + \dots + X_n$. Then

$$\begin{aligned}
\mathbb{P}(S_n = k) &= \sum_{\substack{x_1 + \dots + x_n = k \\ x_i \in \{0, 1\}}} \mathbb{P}(X_1 = x_1, \dots, X_n = x_n) \\
&= \sum_{x_1 + \dots + x_n = k} \mathbb{P}(X_1 = x_1) \dots \mathbb{P}(X_n = x_n) \\
&= \sum_{x_1 + \dots + x_n = k} p^{|\{i: x_i = 1\}|} (1-p)^{|\{i: x_i = 0\}|} \\
&= \sum_{x_1 + \dots + x_n = k} p^k (1-p)^{n-k} \\
&= \binom{n}{k} p^k (1-p)^{n-k}.
\end{aligned}$$

So $S_n \sim \text{Bin}(n, p)$.

Example 4.5 (Non-example)

$(\sigma(1), \sigma(2), \dots, \sigma(n))$ a uniform permutation.

Claim 4.1

$\sigma(1)$ and $\sigma(2)$ are *not* independent.

Suffices to find i_1, i_2 s.t. $\mathbb{P}(\sigma(1) = i_1, \sigma(2) = i_2) \neq \mathbb{P}(\sigma(1) = i_1)\mathbb{P}(\sigma(2) = i_2)$. E.g.
 $\mathbb{P}(\sigma(1) = 1, \sigma(2) = 1) = 0 \neq \underbrace{\mathbb{P}(\sigma(1) = 1)\mathbb{P}(\sigma(2) = 1)}_{=1/n \times 1/n}$

Consequence of definition

Let X_1, \dots, X_n be independent. Then $\mathbb{P}(X_1 \in A_1, \dots, X_n \in A_n) = \mathbb{P}(X \in A_1) \dots \mathbb{P}(X_n \in A_n) \quad \forall A_1, \dots, A_n \subset \mathbb{R} \text{ countable.}$

Let $\Omega = \mathbb{N}$, “Ways of choosing a random integer”

§4.1.4 Geometric distribution (“Waiting for success”)

This can be used to model the number of coin tosses until we get a head.

If $X \sim \text{Geo}(p)$ where $p \in (0, 1)$. $\text{Im}(X) = \{1, 2, \dots\}$,
 $p_X(k) = \mathbb{P}((k-1) \text{ failures, then success on the } k\text{th trial}) = (1-p)^{k-1}p$. Check: $\sum_{k \geq 1} (1-p)^{k-1}p = p \sum_{t \geq 0} (1-p)^t = \frac{p}{1-(1-p)} = 1$.

Alternatively: “Count how many failures before a success”


$\text{Im}(Y) = \{0, 1, 2, \dots\}$, $p_Y(k) = \mathbb{P}(k \text{ failures, then success on the } (k+1)\text{th trial})$. Check: $\sum_{k \geq 0} (1-p)^k p = 1$.

§4.1.5 Poisson Distribution

If $X \sim \text{Po}(\lambda)$ (or $\text{Poi}(\lambda)$) with $\lambda \in (0, \infty)$. $\text{Im}(X) = \{0, 1, 2, \dots\}$ and $\mathbb{P}(X = k) = e^{-\lambda} \lambda^k / k! \quad \forall k \geq 0$. Check: $\sum_{k \geq 0} \mathbb{P}(X = k) = e^{-\lambda} \sum_{k \geq 0} \frac{\lambda^k}{k!} = e^{-\lambda} e^{\lambda} = 1$.

Motivation: Consider $X_n \sim \text{Bin}(n, \frac{\lambda}{n})$

Example 4.6 (“Arrival process”)



Split time interval $[0, \lambda]$ into n small intervals.

- Probability of an arrival in each interval is p , independently across intervals.
- Total no. of arrivals is X_n .

$$\mathbb{P}(X_n = k) = \binom{n}{k} \left(\frac{\lambda}{n}\right)^k \left(1 - \frac{\lambda}{n}\right)^{n-k}$$

Fix k and let $n \rightarrow \infty$

$$\begin{aligned} &= \frac{n!}{n^k(n-k)!} \times \underbrace{\frac{\lambda^k}{k!}}_{\text{no } n} \times \underbrace{\left(1 - \frac{\lambda}{n}\right)^n}_{\rightarrow e^{-\lambda}} \times \underbrace{\left(1 - \frac{1}{n}\right)^{-k}}_{\rightarrow 1} \\ \frac{n!}{n^k(n-k)!} &= \frac{n(n-1)\dots(n-k+1)}{n^k} \\ &= 1 \times \left(1 - \frac{1}{n}\right) \times \left(1 - \frac{2}{n}\right) \times \dots \times \left(1 - \frac{k-1}{n}\right) \\ &\rightarrow 1 \quad \text{There are a fixed number of terms all converging to 1} \\ \mathbb{P}(X_n = k) &\xrightarrow{n \rightarrow \infty} e^{-\lambda} \frac{\lambda^k}{k!}. \end{aligned}$$

We might want to say $\text{Bin}(n, \frac{\lambda}{n})$ converges to $\text{Po}(\lambda)$, but what does convergence of random variables mean?

§4.2 Expectation

$(\Omega, \mathcal{F}, \mathbb{P})$ and X a discrete RV. For now: X only takes non-negative values. “ $X \geq 0$ ”

Definition 4.4 (Expectation)

The **expectation of X** (or **expected value** or **mean**)

$$\begin{aligned} \mathbb{E}[X] &= \sum_{x \in \text{Im}(X)} x \mathbb{P}(X = x) \\ &= \sum_{\omega \in \Omega} X(\omega) \mathbb{P}(\{\omega\}) \end{aligned}$$

“Average of values taken by X , weighted by p_X ”.

Example 4.7 (Uniform Dice)

X uniform on $\{1, 2, \dots, 6\}$

$$\begin{aligned}\mathbb{E}[X] &= \frac{1}{6} \cdot 1 + \frac{1}{6} \cdot 2 + \dots + \frac{1}{6} \cdot 6 \\ &= 3.5\end{aligned}$$

Note. $\mathbb{E}[X]$ need not be in $\text{Im}(X)$.

Example 4.8 (Binomial Distribution)

Let $X \sim \text{Binomial}(n, p)$

$$\begin{aligned}\mathbb{E}[X] &= \sum_{k=0}^n k \mathbb{P}(X = k) \\ &= \sum_{k=0}^n k \binom{n}{k} p^k (1-p)^{n-k} \\ \text{Trick: } k \binom{n}{k} &= \frac{k \times n!}{k! \times (n-k)!} \\ &= \frac{n!}{(k-1)!(n-k)!} \\ &= n \binom{n-1}{k-1} \\ \mathbb{E}[X] &= n \sum_{k=1}^n \binom{n-1}{k-1} p^k (1-p)^{n-k} \\ &= np \sum_{k=1}^n \binom{n-1}{k-1} p^{k-1} (1-p)^{n-k} \\ &= np \sum_{l=0}^{n-1} \binom{n-1}{l} p^l (1-p)^{(n-1)-l} \\ &\quad \underbrace{\hspace{10em} \text{pmf of Bin}(n-1, p)} \\ &= np(p + (1-p))^{n-1} \\ &= np.\end{aligned}$$

Note. We would like to say:

$$\mathbb{E}[\text{Bin}(n, p)] = \mathbb{E}[\text{Bern}(p)] + \dots + \mathbb{E}[\text{Bern}(p)]$$

We will be able to do this soon.

Example 4.9 (Poisson Distribution)

Let $X \sim \text{Poisson}(\lambda)$

$$\begin{aligned}
 \mathbb{E}[X] &= \sum_{k \geq 0} k \mathbb{P}(X = k) \\
 &= \sum_{k \geq 0} k \cdot e^{-\lambda} \frac{\lambda^k}{k!} \\
 &= \sum_{k \geq 1} e^{-\lambda} \frac{\lambda^k}{(k-1)!} \\
 &= \lambda \sum_{k \geq 1} e^{-\lambda} \frac{\lambda^{k-1}}{(k-1)!} \quad \text{pmf of Poisson}(\lambda) \\
 &= \lambda.
 \end{aligned}$$

Note. We would like to say

$$\mathbb{E}[\text{Poisson}(\lambda)] \approx \mathbb{E}\left[\text{Bin}\left(n, \frac{\lambda}{n}\right)\right] = \lambda$$

It is not true in general that $\mathbb{P}(X_n = k) \approx \mathbb{P}(X = k) \implies \mathbb{E}[X_n] \approx \mathbb{E}[X]$

Not important

If X can take on any real value (not necessarily $X \geq 0$)

$$\mathbb{E}[X] = \sum_{x \in \text{Im}(X)} x \mathbb{P}(X = x)$$

unless: A, $\sum_{\substack{x > 0 \\ x \in \text{Im}(X)}} x \mathbb{P}(X = x) = +\infty$ and B, $\sum_{\substack{x < 0 \\ x \in \text{Im}(X)}} x \mathbb{P}(X = x) = +\infty$.

Then we say $\mathbb{E}[X]$ is not defined. Do we really want to study $\infty + \frac{2}{3}(-\infty)$

Summary:

- A and B, $\mathbb{E}[X]$ is not defined.
- A but not B, $\mathbb{E}[X] = +\infty$.¹
- B but not A, $\mathbb{E}[X] = -\infty$.
- neither A nor B, X is then integrable i.e. $\mathbb{E}[X]$ absolutely converges.

¹Some people say not defined instead of letting $\mathbb{E}[X] = \pm\infty$

Example 4.10

Most examples in the course are integrable *except*:

- $\mathbb{P}(X = n) = \frac{6}{\pi^2} \frac{1}{n^2}$ for $n \geq 1$. Note that $\sum \mathbb{P}(X = n) = 1$.
Then $\mathbb{E}[X] = \sum \frac{6}{\pi^2} \frac{1}{n} = +\infty$.
- $\mathbb{P}(X = n) = \frac{3}{\pi^2} \frac{1}{n^2}$ for $n \in \mathbb{Z} \setminus \{0\}$. Then $\mathbb{E}[X]$ is not defined. “It’s symmetric so $\mathbb{E}[X] = 0$ ”, we have decided that this is wrong to prevent many things going wrong in second and third year courses in probability.

Example 4.11 (Indicator Function)

$$\mathbb{E}[1_A] = \mathbb{P}(A).$$

§4.2.1 Properties of Expectation

Proposition 4.1

If $X \geq 0$, then $\mathbb{E}[X] \geq 0$ with equality iff $\mathbb{P}(X = 0) = 1$.

Proof. $\mathbb{E}[X] = \sum_{\substack{x \in \text{Im}(X) \\ x \neq 0}} x \mathbb{P}(X = x)$ □

Proposition 4.2 (Linearity of expectation)

Given random variables X, Y (both integrable) on same probability space $\forall \lambda, \mu \in \mathbb{R}$

$$\mathbb{E}[\lambda X + \mu Y] = \lambda \mathbb{E}[X] + \mu \mathbb{E}[Y]$$

$$\text{Similarly } \mathbb{E}[\lambda_1 X_1 + \dots + \lambda_n X_n] = \lambda_1 \mathbb{E}[X_1] + \dots + \lambda_n \mathbb{E}[X_n]^a$$

^aholds for countably infinite collection though proof is omitted until more analysis experience.

^aholds for countably infinite collection though proof is omitted until more analysis experience.

Note. Independence is NOT a condition.

Proof. If Ω is countable:

$$\begin{aligned} \mathbb{E}[\lambda X + \mu Y] &= \sum_{z \in \text{Im}(\lambda X + \mu Y)} z \mathbb{P}(\lambda X + \mu Y = z) \quad \text{awkward} \\ &= \sum_{\omega \in \Omega} (\lambda X(\omega) + \mu Y(\omega)) \mathbb{P}(\{\omega\}) \end{aligned}$$

$$\begin{aligned}
&= \lambda \sum_{\omega \in \Omega} X(\omega) \mathbb{P}(\{\omega\}) + \mu \sum_{\omega \in \Omega} Y(\omega) \mathbb{P}(\{\omega\}) \\
&= \lambda \mathbb{E}[X] + \mu \mathbb{E}[Y].
\end{aligned}$$

□

Aside - Special Cases

1. If $\lambda, c \in \mathbb{R}$ then:

a) $\mathbb{E}[X + c] = \mathbb{E}[X] + c$

b) $\mathbb{E}[\lambda X] = \lambda \mathbb{E}[X]$

2. a) X, Y random variables (both integrable) on same probability space. $\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$.

b) in fact $\lambda, \mu \in \mathbb{R}$ $\mathbb{E}[\lambda X + \mu Y] = \lambda \mathbb{E}[X] + \mu \mathbb{E}[Y]$ (similarly $\mathbb{E}[\lambda_1 X_1 + \cdots + \lambda_n X_n] = \lambda_1 \mathbb{E}[X_1] + \cdots + \lambda_n \mathbb{E}[X_n]$)

Corollary 4.1

$X \geq Y^a$ then $\mathbb{E}[X] \geq \mathbb{E}[Y]$.

$$^a X(\omega) > Y(\omega) \quad \forall \omega \in \Omega$$

Proof.

$$X = (X - Y) + Y$$

$$\mathbb{E}[X] = \mathbb{E}[X - Y] + \mathbb{E}[Y]$$

$$X - Y \geq 0 \implies \mathbb{E}[X - Y] \geq 0 \quad \text{by Proposition 4.1.}$$

□

Example 4.12 (Counting Problems)

$(\sigma(1), \dots, \sigma(n))$ uniform on Σ_n . $Z = |\{i : \sigma(i) = i\}|$ = number of fixed points. Let $A_i = \{\sigma(i) = i\}$, recall from Example 3.4 A_i s not independent.

Key step:

$$Z = 1_{A_1} + \cdots + 1_{A_n}$$

$$\mathbb{E}[Z] = \mathbb{E}[1_{A_1} + \cdots + 1_{A_n}]$$

$$\begin{aligned}
&= \mathbb{E}[1_{A_1}] + \cdots + \mathbb{E}[1_{A_n}] \quad \text{by Linearity of expectation} \\
&= \mathbb{P}(A_1) + \cdots + \mathbb{P}(A_n) \quad \text{by Example 4.11} \\
&= \frac{1}{n}n = 1.
\end{aligned}$$

Note. Same answer as $\text{Bin}(n, \frac{1}{n})$

Proposition 4.3

If X takes values in $\{0, 1, 2, \dots\}$ then

$$\mathbb{E}[X] = \sum_{k \geq 0} \mathbb{P}(X \geq k)$$

Proof. One can carefully re-arrange the summands which is left as an exercise to the reader. \square

Alternative. Write $X = \sum_{k \geq 1} 1_{X \geq k}$ ^a then take $\mathbb{E}[X]$:

$$\begin{aligned}
\mathbb{E}[X] &= \mathbb{E}\left[\sum 1_{X \geq k}\right] \\
&= \sum \mathbb{E}[1_{X \geq k}] \\
&= \sum \mathbb{P}(X \geq k)
\end{aligned}$$

\square

^aSanity check: let $X = 7$, $1_{X \geq 1} = \cdots = 1_{X \geq 7} = 1$ whilst $1_{X \geq 8} = 1_{X \geq 9} = \cdots = 0$.

Claim 4.2 (Markov's Inequality)

Let $X \geq 0$ be a random variable. Then $\forall a > 0$:

$$\mathbb{P}(X \geq a) \leq \frac{\mathbb{E}[X]}{a}$$

The LHS is interesting, e.g. if we want to bound the probability of an extreme outcome, whilst the RHS is easy to study.

Note. Is $a = \frac{\mathbb{E}[X]}{2}$ useful? No, we already know probabilities are less than 2. If a is large it might be useful

Proof. Observe $X \geq a 1_{X \geq a}$ ^a and take \mathbb{E}

$$\mathbb{E}[X] \geq a \mathbb{E}[1_{X \geq a}]$$

$$= a\mathbb{P}(X \geq a)$$

□

^aCheck: If $X \in [0, a)$ then RHS = 0 else RHS = a .

Note. Markov's Inequality is also true for continuous RVs.

Studying $\mathbb{E}[f(X)]$ Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function. Then $f(x)$ is also a *random variable*².

Claim 4.3

$$\mathbb{E}[f(x)] = \sum_{x \in \text{Im}(x)} f(x)\mathbb{P}(X = x)^a.$$

^aif it exists

^aif it exists

Proof. Let $A = \text{Im}(f(x)) = \{f(x) : x \in \text{Im}(X)\}$. Starting with RHS

$$\begin{aligned} \sum_{x \in \text{Im}(X)} f(x)\mathbb{P}(X = x) &= \sum_{y \in A} \sum_{\substack{x \in \text{Im}(X) \\ f(x)=y}} f(x)\mathbb{P}(X = x) \\ &= \sum_{y \in A} y \sum_{\substack{x \in \text{Im}(X) \\ f(x)=y}} \mathbb{P}(X = x) \\ &= \sum_{y \in A} y\mathbb{P}(f(X) = y) \quad \text{by additivity} \\ &= \mathbb{E}[f(X)] \end{aligned}$$

□

§4.3 Variance

Motivation

$$U_n \sim \text{Uniform}(\{-n, -n+1, \dots, n\})$$

$$V_n \sim \text{Uniform}(\{-n, n\})$$

$$Z_n = 0$$

$$S_n = \text{random walk for } n \text{ steps}$$

$$\sim n - 2 \text{Bin}\left(n, \frac{1}{2}\right)$$

² $X : \Omega \rightarrow \mathbb{R}$ so $f(X) : \Omega \rightarrow \mathbb{R}$.

All of these have $\mathbb{E} = 0$.

Variance is a way to “measure how concentrated a RV is around its mean”.

Definition 4.5

The **variance** of X is:

$$\text{Var}(X) = \mathbb{E} \left[(X - \mathbb{E}[X])^2 \right]$$

Proposition 4.4

$\text{Var}(X) \geq 0$ with equality $\iff \mathbb{P}(X = \mathbb{E}[X]) = 1$ (as $(X - \mathbb{E}[X])^2$ so by Proposition 4.1).

Definition 4.6 (Alternative characterisation)

$$\text{Var}(X) = \mathbb{E} \left[X^2 \right] - (\mathbb{E}[X])^2 \quad (\geq 0)$$

Proof. Write $\mu = \mathbb{E}[X]$

$$\begin{aligned} \text{Var}(X) &= \mathbb{E} \left[(X - \mu)^2 \right] \\ &= \mathbb{E} \left[X^2 - 2\mu X + \mu^2 \right] \\ &= \mathbb{E}[X^2] - 2\mu \underbrace{\mathbb{E}[X]}_{\mu} + \mu^2 \\ &= \mathbb{E}[X^2] - \mu^2 \end{aligned}$$

□

Proposition 4.5 (Properties)

If $\lambda, c \in \mathbb{R}$

- $\text{Var}(\lambda X) = \lambda^2 \text{Var}(X)$.
- $\text{Var}(X + c) = \text{Var}(X)$

Proof.

$$\begin{aligned} \mathbb{E}[X + c] &= \mu + c \\ \text{Var}(X + c) &= \mathbb{E} \left[(X + c - (\mu + c))^2 \right] \end{aligned}$$

$$\begin{aligned}
&= \mathbb{E}[(X - \mu)^2] \\
&= \text{Var}(X).
\end{aligned}$$

□

Example 4.13 (Poisson Distribution)

Let $X \sim \text{Poisson}(\lambda)$

$$\text{Var}(X) = \mathbb{E}[X^2] - \lambda^2$$

“Falling factorial trick”: sometimes easier to calculate $\mathbb{E}[X(X-1)]$ than $\mathbb{E}[X^2]$

$$\begin{aligned}
\mathbb{E}[X(X-1)] &= \sum_{k \geq 2} \underbrace{k(k-1)}_{\text{function}} \underbrace{e^{-\lambda} \frac{\lambda^k}{k!}}_{\text{PMF}} \\
&= \lambda^2 e^{-\lambda} \sum_{k \geq 2} \underbrace{\frac{\lambda^{k-1}}{(k-2)!}}_{e^\lambda} \\
&= \lambda^2 \\
\mathbb{E}[X^2] &= \mathbb{E}[X(X-1)] + \mathbb{E}[X] \\
&= \lambda^2 + \lambda \\
\text{Var}(X) &= \lambda
\end{aligned}$$

Example 4.14 (Geometric Distribution)

Let $Y \sim \text{Geom}(p)$ where $Y \in \mathbb{N}$.

$$\mathbb{E}[Y] = \frac{1}{p}, \quad \text{Var}(Y) = \frac{1-p}{p^2}.$$

Proof left as an exercise.

Note. λ large: $\text{Var}(x) = \mathbb{E}[X]$, more concentrated
 p small: $\text{Var}(Y) \approx \frac{1}{p^2} = (\mathbb{E}[X])^2$.

Example 4.15 (Bernouli Distribution)

Let $X \sim \text{Bern}(p)$.

$$\mathbb{E}[X] = 1 \times p = p$$

$$\begin{aligned}\mathbb{E}[X^2] &= 1^2 \times p = p \\ \text{Var}(X) &= p - p^2 \\ &= p(1 - p)\end{aligned}$$

Example 4.16 (Binomial Distribution)

Let $X \sim \text{Bin}(n, p)$

$$\begin{aligned}\mathbb{E}[X] &= np \\ \mathbb{E}[X^2] &= \text{ugly}\end{aligned}$$

§4.3.1 Sums of RVs

Goal: Study $\text{Var}(X_1 + \cdots + X_n)$. Do the X_i s need to be independent.

Proposition 4.6 (Preliminary: Expectation of Product of RVs)

If X, Y are **independent** RVs and f, g are functions $\mathbb{R} \rightarrow \mathbb{R}$.

Then: $\mathbb{E}[f(X)g(Y)] = \mathbb{E}[f(X)]\mathbb{E}[g(Y)]$ “Splits as a product” .

Example 4.17

$$\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$$

Example 4.18

Let $f(x) = g(x) = z^x$ (or e^{tx}).

Proof. (X, Y discrete)

$$\begin{aligned}\text{LHS} &= \sum_{x,y \in \text{Im}} f(x)g(y)\mathbb{P}(X=x, Y=y) \\ &= \sum_{x,y \in \text{Im}} f(x)g(y)\mathbb{P}(X=x)\mathbb{P}(Y=y) \\ &= \left[\sum_{x \in \text{Im}} f(x)\mathbb{P}(X=x) \right] \left[\sum_{y \in \text{Im}} g(y)\mathbb{P}(Y=y) \right] \\ &= \mathbb{E}[f(X)]\mathbb{E}[g(Y)]\end{aligned}$$

□

Proposition 4.7 (Sums of independent RVs)

Let X_1, \dots, X_n be independent. Then

$$\text{Var}(X_1 + \dots + X_n) = \text{Var}(X_1) + \dots + \text{Var}(X_n)$$

Proof. (Suffices to Prove (STP) $n = 2$). Say $\mathbb{E}[X] = \mu$, $\mathbb{E}[Y] = \nu$

$$\begin{aligned} \text{Var}(X + Y) &= \mathbb{E}[(X + Y - \mu - \nu)^2] \\ &= \mathbb{E}[(X - \mu)^2] + \mathbb{E}[(Y - \nu)^2] + 2\mathbb{E}[(X - \mu)(Y - \nu)]. \\ &= \text{Var}(X) + \text{Var}(Y) + 2\mathbb{E}[X - \mu]\mathbb{E}[Y - \nu] \\ &= \text{Var}(X) + \text{Var}(Y). \end{aligned}$$

□

Example 4.19 (Binomial)

Going back to [Binomial Distribution](#), $\text{Var}(\text{Bin}(n, p)) = np(1 - p)$.

Goal: Study $\text{Var}(X + Y)$ when X, Y not independent.

Definition 4.7 (Covariance)

Let X, Y be two RVs. Their **covariance** is

$$\text{Cov}(X, Y) = \mathbb{E}[(X - \mathbb{E}[X])(Y - \mathbb{E}[Y])]$$

“Measures how dependent X, Y are and in which direction” (X large $\implies Y$ larger if $\text{Cov} > 0$ else Y smaller).

Proposition 4.8 (Properties)

$$\text{Cov}(X, Y) = \text{Cov}(Y, X)$$

$$\text{Cov}(X, X) = \text{Var}(X)$$

Definition 4.8 (Alternative Characterisation)

$$\text{Cov}(X, Y) = \mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y]$$

Proof.

$$\begin{aligned}
 \text{Cov}(X, Y) &= \mathbb{E}[(X - \mu)(Y - \nu)] \\
 &= \mathbb{E}[XY] - \underbrace{\mu}_{\nu} \underbrace{\mathbb{E}[Y]}_{\mu} - \nu \underbrace{\mathbb{E}[X]}_{\mu} + \mu\nu \\
 &= \mathbb{E}[XY] - \mu\nu.
 \end{aligned}$$

□

Proposition 4.9 (More Properties)

Let $\lambda \in \mathbb{R}$

$$\begin{aligned}
 \text{Cov}(\lambda, X) &= 0 \\
 \text{Cov}(X + \lambda, Y) &= \text{Cov}(X, Y) \\
 \text{Cov}(\lambda X, Y) &= \lambda \text{Cov}(X, Y) \\
 \text{Var}(X + Y) &= \text{Var}(X) + \text{Var}(Y) + 2 \text{Cov}(X, Y)
 \end{aligned}$$

Covariance is linear in each argument i.e. $\text{Cov}(\sum \lambda_i X_i, Y) = \sum \lambda_i \text{Cov}(X_i, Y)$ and $\text{Cov}(\sum \lambda_i X_i, \sum \mu_j Y_j) = \sum_{i=1}^n \sum_{j=1}^m \lambda_i \mu_j \text{Cov}(X_i, Y_j)$.

Proposition 4.10 (Sums of RVs)

$$\begin{aligned}
 \text{Var}\left(\sum_{i=1}^n X_i\right) &= \text{Cov}\left(\sum_{i=1}^n X_i, \sum_{i=1}^n X_i\right) \\
 &= \sum_{i=1}^n \text{Var}(X_i) + \sum_{i \neq j} \text{Cov}(X_i, X_j)^a
 \end{aligned}$$

^aor $2 \sum_{i < j} \text{Cov}(X_i, X_j)$

^aor $2 \sum_{i < j} \text{Cov}(X_i, X_j)$

Proposition 4.11

X, Y independent $\implies \text{Cov}(X, Y) = 0$, **the converse is false.**

Example 4.20

Let $Y = -X$, $\text{Var}(Y) = \text{Var}(X)$. So $\text{Var}(X + Y) = \text{Var}(0) = 0 \neq \text{Var}(X) + \text{Var}(Y)$.

Example 4.21 (Uniform Permutation)

Let $(\sigma(1), \dots, \sigma(n))$ be uniformly chosen on Σ_n . Let $A_i = \{\sigma(i) = i\}$ and $N = 1_{A_1} + \dots + 1_{A_n}$ (N is the number of fixed points).

$$\mathbb{E}[N] = n \times \frac{1}{n} = 1$$

A_i and A_j are *not* independent

$$\begin{aligned} \text{Var}(1_{A_1}) &= \frac{1}{n} \left(1 - \frac{1}{n}\right) \\ \text{Cov}(1_{A_i}, 1_{A_j}) &= \mathbb{E}[1_{A_i} 1_{A_j}] - \mathbb{E}[1_{A_i}] \mathbb{E}[1_{A_j}] \\ &= \mathbb{E}[1_{A_i \cap A_j}] - \mathbb{E}[1_{A_i}] \mathbb{E}[1_{A_j}] \\ &= \mathbb{P}(A_i \cap A_j) - \mathbb{P}(A_i) \mathbb{P}(A_j) \\ &= \frac{1}{n(n-1)} - \frac{1}{n} \times \frac{1}{n} \\ &= \frac{1}{n^2(n-1)} > 0 \end{aligned}$$

Note: Cov doesn't depend on i, j

$$\begin{aligned} \text{Var}(N) &= \sum_{i=1}^n \text{Var}(1_{A_i}) + \sum_{i \neq j} \text{Cov}(1_{A_i}, 1_{A_j}) \\ &= n \times n \left(1 - \frac{1}{n}\right) + n(n-1) \times \frac{1}{n^2(n-1)} \\ &= 1 - \frac{1}{n} + \frac{1}{n} \\ &= 1. \end{aligned}$$

Compare with $\text{Bin}\left(n, \frac{1}{n}\right)$: $\mathbb{E} = 1$, $\text{Var} = n \times \frac{1}{n} \left(1 - \frac{1}{n}\right) = 1 - \frac{1}{n}$.

§4.3.2 Chebyshev's Inequality**Proposition 4.12 (Chebyshev's Inequality)**

Let X be a RV, $\mathbb{E}[X] = \mu$, $\text{Var}(X) = \sigma^2 < \infty$.

$$\mathbb{P}(|X - \mu| \geq \lambda) \leq \frac{\text{Var}(X)}{\lambda^2}$$

Remember the proof, not the statement.

Proof. Idea: Apply [Markov's Inequality](#) to $(X - \mu)^2$.

$$\begin{aligned}\mathbb{P}\left((X - \mu)^2 \geq \lambda^2\right) &\leq \frac{\mathbb{E}[(X - \mu)^2]}{\lambda^2} \\ &= \frac{\text{Var}(X)}{\lambda^2}\end{aligned}$$

□

Danger: Applying [Markov's Inequality](#) to $|X - \mu|$, $\mathbb{E}[|X - \mu|]$ is less nice than $\mathbb{E}[(X - \mu)^2]$.

Comments

- Chebyshev's Inequality gives better bounds than Markov's Inequality (decays with λ^2 instead of λ).
- We can apply it to all RVs, not just those ≥ 0 .
- Caveat: We need $\text{Var}(X) < \infty$ which is a stronger condition than $\mathbb{E}[X] < \infty$.

Definition 4.9 (Standard Deviation)

$\sqrt{\text{Var}(X)}$ is the **standard deviation**, σ , of X .

It has the same “units” as X but not many nice properties so Var is generally preferred.

We can rewrite Chebyshev as $\mathbb{P}(|X - \mu| \geq k\sigma) \leq 1/k^2$

§4.4 Conditional Expectation

Setting: $(\Omega, \mathcal{F}, \mathbb{P})$.

Recall the definition of [Conditional Probability](#).

Definition 4.10 (Conditional Expectation)

$B \in \mathcal{F}$ with $\mathbb{P}(B) > 0$, X a RV. The **conditional expectation** is

$$\mathbb{E}[X \mid B] = \frac{\mathbb{E}[X1_B]}{\mathbb{P}(B)}$$

Example 4.22 (Uniform Dice)

Let X be a dice, uniform on $\{1, \dots, 6\}$.

$$\begin{aligned}\mathbb{E}[X \mid X \text{ prime}] &= \frac{\frac{1}{6}[0 + 2 + 3 + 0 + 5 + 0]}{\frac{1}{2}} \\ &= \frac{1}{3}(2 + 3 + 5) \\ &= \frac{10}{3}.\end{aligned}$$

Definition 4.11 (Alternative Characterisation)

$$\mathbb{E}[X \mid B] = \sum_{x \in \text{Im } X} x \mathbb{P}(X = x \mid B).$$

Proof.

$$\begin{aligned}\text{RHS} &= \sum \frac{x \mathbb{P}(\{X = x\} \cap B)}{\mathbb{P}(B)} \\ &= \sum_{\substack{x \neq 0 \\ x \in \text{Im } X}} \frac{x \mathbb{P}(X 1_B = x)}{\mathbb{P}(B)} \\ \text{Note: } \mathbb{E}[X 1_B] &= \sum_{\substack{x \neq 0 \\ x \in \text{Im } X}} x \mathbb{P}(X 1_B = x)\end{aligned}$$

□

Proposition 4.13 (Law of Total Expectation)

Let (B_1, B_2, \dots) be a finite or countably-infinite partition of Ω with $B_n \in \mathcal{F} \quad \forall n$ s.t. $\mathbb{P}(B_n) > 0$. X a RV.

$$\mathbb{E}[X] = \sum_n \mathbb{E}[X \mid B_n] \mathbb{P}(B_n).$$

Example 4.23

Let $X = 1_A$ we recover the [Law of Total Probability](#).

Proof.

$$\text{RHS} = \sum_n \mathbb{E}[X 1_{B_n}]$$

$$\begin{aligned}
&= \mathbb{E} \left[\sum X \cdot (1_{B_1} + \cdots + 1_{B_n}) \right] \quad \text{by Linearity of expectation} \\
&= \mathbb{E}[X \cdot 1] \\
&= \mathbb{E}[X].
\end{aligned}$$

□

Application: Two stage randomness where (B_n) describes what happens in stage 1.

Example 4.24 (Sums of random number of terms)

Let $(X_n)_{n \geq 1}$ be IID and $N \in \{0, 1, 2, \dots\}$ be a random index independent of (X_n) . $S_n = X_1 + \cdots + X_n$ with $\mathbb{E}[X_n] = \mu$ so $\mathbb{E}[S_n] = n\mu$. Then

$$\begin{aligned}
\mathbb{E}[S_N] &= \sum_{n \geq 0} \mathbb{E}[S_N \mid N = n] \mathbb{P}(N = n) \\
&= \sum \mathbb{E}[S_n] \mathbb{P}(N = n)^a \\
&= \sum_{n \geq 0} n\mu \mathbb{P}(N = n) \\
&= \mu \mathbb{E}[N].
\end{aligned}$$

^aKEY STEP $\mathbb{E}[S_N \mid N = n] = \mathbb{E}[S_n \mid N = n] = \mathbb{E}[S_n]$ last step follows as S_n and $\{N = n\}$ are independent.

^aKEY STEP $\mathbb{E}[S_N \mid N = n] = \mathbb{E}[S_n \mid N = n] = \mathbb{E}[S_n]$ last step follows as S_n and $\{N = n\}$ are independent.

§4.5 Random Walks

Definition 4.12 (Random Walk)

Let $(X_n)_{n \geq 1}$ be IID RVs then $S_n = x_0 + X_1 + \cdots + X_n$. (S_0, S_1, S_2, \dots) is a random process called a **Random Walk** started from x_0 .

§4.5.1 Simple Random Walk (SRW) on \mathbb{Z} - Main example in our course

$P(X_i = +1) = p, \mathbb{P}(X = -1) = q = 1 - p.$

$x_0 \in \mathbb{Z}$ (often $x_0 = 0$).

Special case: $p = q = \frac{1}{2}$ is “symmetric”.

$$\mathbb{P}(S_2 = x_0) = pq + qp = 2pq$$

Useful interpretation: A gamble repeatedly plays a game where he wins £1 with $\mathbb{P} = p$, loses £1 with $\mathbb{P} = q$.

Often: stops at £0.

Question

Suppose the gambler starts with $\mathcal{L}x$ at time 0. What is the probability he reaches $\mathcal{L}a$ before $\mathcal{L}0$. ($0 < x < a$)

Notation. $\mathbb{P}_x(\cdot) = \mathbb{P}(\cdot \mid x_0 = x)$ “measure of RW started from x_0 .”

Answer

Key idea: Conditional on $S_1 = z$, (S_1, S_2, \dots) is a random walk started from z .
Apply [Law of Total Probability](#)

$$\begin{aligned}\mathbb{P}_x(S \text{ hits } a \text{ before } 0) &= \sum_z \mathbb{P}_x(S \text{ hits } a \text{ before } 0 \mid S_1 = z) \mathbb{P}_x(S_1 = z) \\ &= \sum_z \mathbb{P}_z(S \text{ hits } a \text{ before } 0) \mathbb{P}_x(S_1 = z)\end{aligned}$$

let $h_x = \mathbb{P}_x(S \text{ hits } a \text{ before } 0)$.

$$S_1 = x \pm 1$$

$$h_x = ph_{x+1} + qh_{x-1}.$$

Important to specify boundary conditions: $h_0 = 0$, $h_a = 1$

Solving Linear Recurrence Equations

$ph_{x+1} - h_x + qh_{x-1} = 0$ is a homogenous equation whose solutions form a vector space. We want to find two LI solutions and we guess $h_x = \lambda^x$ So

$$\begin{aligned}p\lambda^{x+1} - \lambda^x + q\lambda^{x-1} &= 0 \\ p\lambda^2 - \lambda + q &= 0 \\ \lambda &= 1, \frac{q}{p}\end{aligned}$$

Case $q \neq p$

$$h_x = A + B \left(\frac{q}{p}\right)^x$$

Use BCs to find A,B:

$$x = 0 : h_0 = 0 = A + B$$

$$x = a : h_a = 1 = A + \left(\frac{q}{p}\right)^a$$

$$h_x = \frac{\left(\frac{q}{p}\right)^x - 1}{\left(\frac{q}{p}\right)^a - 1}.$$

Case $p = q = \frac{1}{2}$: Note $h_x = x$ “ x is the average of $x + 1$ and $x - 1$ ” General solution:
 $h_x = A + Bx$ BCs

$$\begin{aligned} h_0 &= 0 = A \\ h_a &= 1 = Ba \\ h_x &= \frac{x}{a} \end{aligned}$$

^aSubscript: $z \in \text{Im}(S_1)$ but we will bother with that any more

^aSubscript: $z \in \text{Im}(S_1)$ but we will bother with that any more

Question

Suppose the gambler starts with $\mathcal{L}x$ at time 0. What is the expected absorption time, $T = \min\{n \geq 0 : S_n = 0 \text{ or } S_n = a\}$. “first time S hits $\{0, a\}$ ”

Answer

Apply Law of Total Expectation

We want $\mathbb{E}_x[T]$ ($\mathbb{E}[T]$ when we start from x) which we label as τ_x

$$\begin{aligned} \tau_x &= \mathbb{E}_x[T] \\ &= p\mathbb{E}_x[T \mid S_1 = x + 1] + q\mathbb{E}_x[T \mid S_1 = x - 1] \\ &= p\mathbb{E}_{x+1}[T + 1] + q\mathbb{E}_{x-1}[T - 1] \\ &= p(1 + \mathbb{E}_{x+1}[T]) + q(1 + \mathbb{E}_{x-1}[T]) \\ &= 1 + p\tau_{x+1} + q\tau_{x-1} \end{aligned}$$

Boundary conditions: $\tau_0 = \tau_a = 0$ “We’re already there” .

We already solved the homogenous case of this equation previously. Inhomogeneous case: $ph_{x+1} - h_x + qh_{x-1} = -1$. Find a *particular solution* Guess: “one level more complicated than general solution” Add on general solution. Solve for BCs.

$p \neq q$: Guess: $h_x = \frac{x}{q-p}$ works as a particular solution

$p = q = \frac{1}{2}$: Guess $h_x = Cx^2$ might work

Sub in: $\frac{C}{2}(x+1)^2 - Cx^2 + \frac{C}{2}(x-1)^2 = -1$ $C = -1$ so $h_x = A + Bx - x^2$ (find A, B with BCs) roots are 0 and a so $h_x = x(a-x)$ and $h_x \geq 0$

Probability sanity check: $p = q = \frac{1}{2}$. “Fair game” Study: Expected profit if you start from $\mathcal{L}x$ and play until time T . $\mathbb{E}_x[S_T] = a\mathbb{P}_x(S_T = a) + 0 \times \mathbb{P}_x(S_T = 0) = ah_x = x$. Fits our intuition for fair games. ✓