1 Hyper-Detailed Pedagogical Blueprint Real Analysis: Change of Variables & Rigorous Foundations

Director's Cut & AI Extraction Protocol
Lecture Segment: 00:00 – 15:00

Insight: The "Wall" of Multivariable Calculus

This segment serves as the bridge between computational calculus and rigorous Real Analysis. The professor is actively shifting the students' mindset away from "memorizing integration formulas" and towards "understanding the geometric distortions of space via linear algebra (Jacobians)." Notice how carefully he handles domains and inverse mappings—this rigor is the core of the discipline.

2 Reminder: The Setup for the Change of Variables

Cleaned Spoken Protocol [Time: 00:00:11 - 00:01:29]

I hope you all had a great holiday and have recovered from the first half of the semester so we can continue with our work! Let me start by reminding you of what we did last time. In our previous lecture, we stated and began to prove what we playfully call the "final boss of integration": the Change of Variables formula. Let us review the setup.
As always, we begin with our domain $U \subset \mathbb{R}^n$. We apply a diffeomorphism—which we call Φ—that maps U to another domain V.

Board State & Physical Actions: Board State 1: The Diffeomorphism

The professor writes the title "Change of vars." and begins defining the mathematical environment on the far left board. He writes the mapping $U \to V$ and explicitly writes "diffeo" in orange chalk beneath the arrow.

Mathematical Setup (Stroke-by-Stroke): Let $U, V \subset \mathbb{R}^n$. We define the mapping between these spaces:

$$U \xrightarrow[\text{diffeo}]{\Phi} V$$

Redundant Explanation of Step: We establish a transformation Φ mapping a domain U to a codomain V. The orange annotation "diffeo" indicates this mapping is a *diffeomorphism*. A diffeomorphism is a heavily constrained, "well-behaved" mapping: it must be a bijection (one-to-one and onto), it must be differentiable everywhere, and its inverse Φ^{-1} must also be differentiable everywhere. This ensures the space is smoothly warped without tearing or folding.



Cleaned Spoken Protocol [Time: 00:01:29 - 00:02:45]

We then consider a Jordan measurable set $A \subset \mathbb{R}^n$, with the strict condition that its closure, $\overline{A}$, must be entirely contained within the domain U of our diffeomorphism. Next, we need a continuous function to integrate. We define a function f mapping from the closure $\overline{A}$ to the real numbers $\mathbb{R}$.

Board State & Physical Actions: Board State 2: Domain Correction

The professor writes $A \subset \mathbb{R}^n$ J. m. st. $\overline{A} \subset U$. When defining the function f, he initially says "from U or from A to $\mathbb{R}$", but immediately catches his imprecision and writes $\overline{A} \to \mathbb{R}$ continuous.

Mathematical Setup (Continued): We restrict our attention to a specific subset of U:

$$A \subset \mathbb{R}^n \text{ is Jordan measurable, such that } \overline{A} \subset U$$

We define our integrand function:

$$f : \overline{A} \to \mathbb{R} \quad \text{(continuous)}$$

Redundant Explanation of Step: Why the closure $\overline{A}$? By requiring the *closure* of A to be strictly contained within the open set U, we ensure that the boundary of A behaves nicely under the mapping Φ, and we avoid pathological singularities that could exist on the absolute edge of the domain U. We then ensure f is continuous over this closed, bounded region, guaranteeing Riemann integrability.

3 The Clunky Formulation of the Theorem

Cleaned Spoken Protocol [Time: 00:02:45 - 00:03:50]

With this setup, the theorem states that the integral of $f(x)$ over the region A is equivalent to the integral evaluated over the mapped image, $\Phi(A)$. However, to calculate this, we must evaluate our function f at the pre-image, $\Phi^{-1}(y)$. Furthermore, the differential element dy must be corrected by the distortion of the volume, which is exactly the absolute value of the determinant of the Jacobian of Φ, evaluated at $\Phi^{-1}(y)$.

Board State & Physical Actions: Board State 3: The Clunky Formula

The professor writes out the full Change of Variables formula exactly as it was derived in the previous lecture. It is notationally heavy due to the inverse mappings inside the arguments.

The Initial Theorem (Inline Annotated):

$$\underbrace{\int_A f(x)\,dx}_{\text{Integral in Original Space}} = \int_{\Phi(A)} f(\underbrace{\Phi^{-1}(y)}_{\text{Pre-image under } \Phi}) \underbrace{\frac{1}{|\det J\Phi(\Phi^{-1}(y))|}}_{\text{Correction for Volume Distortion}} dy \tag{1}$$

Redundant Explanation of Step: To evaluate the integral in the distorted space V (where the variable is y), every x in the original function must be replaced with the mapping that takes y back to x, which is $\Phi^{-1}(y)$. Because the transformation Φ stretches and squishes space, the resulting "volume pixels" (dy) are distorted. To preserve the total "mass" of the integral, we must divide by the scaling factor of that distortion. The scaling factor of a linear transformation

is the determinant of its matrix—hence, we divide by the determinant of the Jacobian matrix of Φ.

> ***Cleaned Spoken Protocol [Time: 00:03:50 - 00:06:50]***
>
> *I previously explained the geometric intuition behind why this theorem is true, but we didn't have time to complete the formal proof. I propose that we leave the proof for now. Instead, we will look at some concrete examples to see how we apply this theorem in actual calculations. Once you see it in action, you will understand why it is worth the effort to prove it. We will revisit the formal proof in two weeks during our revision buffer week.*
> **(The professor then spends a few minutes discussing an upcoming interactive teaching session on Thursday with guest coaches from ETH. He asks students to attend in person for the interactive exercises.)**

4 Rewriting the Theorem via the Inverse Map

> ***Cleaned Spoken Protocol [Time: 00:06:50 - 00:08:20]***
>
> *Let us rewrite this theorem into a format that is much easier to use in practice. Instead of constantly writing Φ^{-1} and dragging it through our equations, let us define a new mapping Ψ to be exactly equal to the inverse, Φ^{-1}.*
> *If we substitute this into our theorem, the integration region $\Phi(A)$ remains, but the function evaluation $f(\Phi^{-1}(y))$ simply becomes $f(\Psi(y))$. But what happens to that clunky determinant in the denominator?*

Board State & Physical Actions: Board State 4: Defining Psi

The professor moves to the center board to clean up the notation. He defines Ψ and prepares to algebraically manipulate the Jacobian fraction.

Simplifying the Notation: Let us define the inverse map explicitly:

$$\Psi = \Phi^{-1}$$

Substituting this directly into Equation 1, we get an intermediate step:

$$\int_A f(x)\,dx = \int_{\Phi(A)} f(\Psi(y)) \frac{1}{|\det J\Phi(\Psi(y))|}\,dy$$

> ***Cleaned Spoken Protocol [Time: 00:08:20 - 00:10:45]***
>
> *Remember the lemma we proved regarding the differential of an inverse function. If we compose Φ with its inverse Ψ, we get the identity map.*
> *By applying the chain rule, the differential of this composition evaluated at a point y is the differential of Φ multiplied by the differential of Ψ. Because the differential of the identity map is just the identity matrix, this means that the Jacobian matrix of Φ multiplied by the Jacobian matrix of Ψ equals the unit (identity) matrix.*
> *Because the determinant is a multiplicative property for matrices, the determinant of one matrix must be the exact inverse (reciprocal) of the other. Therefore, instead of dividing by the determinant of $J\Phi$, we can simply multiply by the determinant of $J\Psi$!*

Board State & Physical Actions: Board State 5: Chain Rule Derivation

The professor formally derives the relationship between the determinants on the board, line by line, mapping the abstract function composition to concrete matrix algebra.

Derivation of the Jacobian Reciprocal Relation:

$$\underbrace{\Phi \circ \Psi = \mathrm{id}}_{\text{Function Composition}} \implies \underbrace{D\Phi_{\Psi(y)} \circ D\Psi_y = \mathrm{id}}_{\text{Taking the Differential (Chain Rule)}}$$
$$\implies \underbrace{J\Phi(\Psi(y))}_{\text{Matrix 1}} \cdot \underbrace{J\Psi(y)}_{\text{Matrix 2}} = \underbrace{I_n}_{\text{Identity Matrix}}$$

Redundant Explanation of Step: The chain rule in multivariable calculus dictates that the derivative of a composition of functions is the matrix product of their Jacobian matrices. Since the composition yields the identity function ($y \mapsto y$), the product of their Jacobians must yield the identity matrix I_n.

Taking the determinant of both sides, and using the property that $\det(AB) = \det(A)\det(B)$ and $\det(I_n) = 1$:

$$\det\Big(J\Phi(\Psi(y)) \cdot J\Psi(y)\Big) = \det(I_n)$$
$$\det(J\Phi(\Psi(y))) \cdot \det(J\Psi(y)) = 1$$
$$\implies \underbrace{\frac{1}{\det J\Phi(\Psi(y))}}_{\text{Clunky Denominator}} = \underbrace{\det J\Psi(y)}_{\text{Clean Numerator}}$$

Insight: The Power of Linear Algebra

The professor takes a moment here to show how beautifully linear algebra serves calculus. The geometric reality that "stretching space forward by factor k, then pulling it backward, returns it to normal" is flawlessly encoded in the algebraic fact that the determinants of inverse matrices multiply to 1.

Cleaned Spoken Protocol [Time: 00:10:45 - 00:12:15]

This gives us a highly practical, informal way to remember the substitution rule. Just like in Analysis 1, when you substitute $x = \Psi(y)$, you must replace the differential. But instead of replacing dx with the 1D derivative $\Psi'(y)\,dy$, you replace it with the absolute determinant of the Jacobian matrix, $|\det J\Psi(y)|\,dy$.

Board State & Physical Actions: Board State 6: The Orange Rule

To emphasize its importance for practical computation, the professor writes this finalized substitution rule on the board in bright orange chalk.

The Practical Substitution Rule

When substituting $x = \Psi(y)$, the differential transforms as:

$$\underbrace{dx}_{\text{Original Volume}} = \underbrace{|\det J\Psi(y)|}_{\text{Scaling Factor}} \underbrace{dy}_{\text{New Volume}}$$

Yielding the clean theorem:

$$\int_A f(x)\,dx = \int_{\Phi(A)} f(\Psi(y))\,|\det J\Psi(y)|\,dy$$

5 Example: Area of a Unit Disk & The Definition of π

Cleaned Spoken Protocol [Time: 00:12:15 - 00:15:00]

Let us compute a concrete example. But before we do, I must ask you: how was the constant π officially defined in your Analysis 1 course?
(A student answers correctly: "It is defined as the unique number in the open interval $(0, 4)$ such that the sine of π is zero.")
Exactly! You know many things about π from geometry, but your official, rigorous definition is that π is the first positive root of the sine function.
The sine and cosine functions themselves are defined purely analytically (usually via power series), and their defining properties are that the derivative of sine is cosine, the derivative of cosine is negative sine, $\sin(0) = 0$, and $\cos(0) = 1$. Those relations are definitive.

Board State & Physical Actions: Board State 7: Defining Pi rigorously

The professor steps to the far right board to lay down the absolute axioms of Analysis 1 before computing the area of a circle. He wants to prove the area is π *without* using circular geometric logic.

Rigorous Axioms from Analysis 1: The trigonometric functions are defined purely by their analytical properties (not triangles):

$$\sin'(x) = \cos(x) \tag{2}$$
$$\cos'(x) = -\sin(x) \tag{3}$$
$$\sin^2(x) + \cos^2(x) = 1 \tag{4}$$

The constant π is defined analytically as the unique root:

$$\sin(\pi) = 0 \quad \text{(for the first positive root)}$$

Redundant Explanation of Step: By establishing these purely algebraic definitions, the professor is ensuring that when we compute the integral of a disk and get the symbol π, we are arriving at it through rigorous calculus and limits, not by relying on a high-school geometric formula ($A = \pi r^2$) which assumes the conclusion we are trying to prove.

Insight: Avoiding Circular Logic

This is a brilliant teaching moment. It is very easy for students to take geometry for granted. By forcing them to recall that π is just "the root of a specific power series",

the professor elevates the upcoming Change of Variables computation from a "boring high-school geometry problem" into a "profound demonstration of analytical power."

Note: The video segment ends right as the professor finishes writing these trigonometric axioms on the board. In the next segment, he will define the polar coordinate mapping $\Psi(y_1, y_2)$ and use the Orange Substitution Rule to calculate the Jacobian determinant and prove the area of the unit disk is exactly π.

6 Hyper-Detailed Pedagogical Blueprint Real Analysis: Change of Variables & Rigorous Foundations (Part 2)

Director's Cut & AI Extraction Protocol
Lecture Segment: 15:00 – 32:00

4 Computing the Area of a Unit Disk

Cleaned Spoken Protocol [Time: 00:15:00 - 00:18:37]

Now that we have established the rigorous analytical definition of π, we will prove—perhaps for the first time in your mathematical lives—that the area of the unit disk is exactly π.
In our rigorous language, we are asking for the 2-dimensional Jordan measure of the set B_1, which is the set of all points (x_1, x_2) such that $x_1^2 + x_2^2 \leq 1$. To compute this measure, we will use polar coordinates.
Let the first parameter y_1 represent the radius, and the second parameter y_2 represent the angle. We define our coordinate mapping Ψ such that $x_1 = y_1 \cos(y_2)$ and $x_2 = y_1 \sin(y_2)$. Our parameter domain D will be the open rectangle where the radius y_1 goes from 0 to 1, and the angle y_2 goes from 0 to 2π.

Board State & Physical Actions: Board State 8: Setting up the Transformation
The professor writes down the set definition for the disk and formally defines the polar mapping Ψ alongside its parameter domain D.

Mathematical Setup (Stroke-by-Stroke): The Unit Disk in the original coordinates:

$$B_1 = \{(x_1, x_2) \in \mathbb{R}^2 \mid x_1^2 + x_2^2 \leq 1\}$$

The Polar Coordinate Mapping Ψ:

$$\begin{aligned} x_1 &= y_1 \cos(y_2) \\ x_2 &= y_1 \sin(y_2) \end{aligned}$$

The Parameter Domain D:

$$D = \underbrace{(0,1)}_{\text{Radius } y_1} \times \underbrace{(0,2\pi)}_{\text{Angle } y_2} \implies \Psi : D \to B_1$$

Redundant Explanation of Step: We are transitioning from Cartesian coordinates (x_1, x_2) to polar coordinates (y_1, y_2). The domain D is strictly an *open* rectangle. This strict open boundary guarantees that Ψ remains a proper diffeomorphism (bijective and smooth), as including $y_1 = 0$ (the origin) or $y_2 = 0, 2\pi$ (the overlapping seam) would break the bijection.

Cleaned Spoken Protocol [Time: 00:18:37 - 00:20:17]

Our first crucial observation is that the image of our parameter domain, $\Psi(D)$, is not entirely equal to the closed disk B_1. Because we are using the open interval $(0,1)$ for the radius and $(0,2\pi)$ for the angle, our mapping completely misses the origin and the line segment lying on the positive x-axis.
However, we can easily prove that this missing 1-dimensional line segment has a 2-dimensional

Jordan measure of zero. If you cover it with small squares of side length ϵ, the total area goes to zero. Therefore, even though the sets are not identical, their measures are perfectly equal: $\mu_2(B_1) = \mu_2(\Psi(D))$.

Insight: The "Measure Zero" Safety Net

This is a critical Real Analysis concept. Students often panic when a mapping isn't perfectly surjective over the boundaries. The professor is teaching them the "cheat code" of integration: integrals do not "see" sets of measure zero. You can freely delete boundaries, seams, and single points without changing the volume.

Board State & Physical Actions: TikZ: The Measure Zero Boundary Issue

The professor draws the mapping, actively highlighting the "missing" seam on the right side of the circle.



Cleaned Spoken Protocol [Time: 00:20:17 - 00:22:55]

Because the measures are equal, we can apply our Change of Variables substitution rule. The area of the disk is the integral of the constant function 1 over our domain D, multiplied by the determinant of the Jacobian matrix of Ψ.
Let us compute this Jacobian matrix. Taking the partial derivatives with respect to y_1 and y_2, we construct our 2×2 matrix. When we compute the determinant, we get $y_1 \cos^2(y_2) + y_1 \sin^2(y_2)$. Factoring out the y_1, we use our analytical axiom that $\cos^2 + \sin^2 = 1$. The determinant is simply y_1. Because y_1 is strictly positive on our domain, its absolute value is just y_1.

Board State & Physical Actions: Board State 9: The Jacobian Calculation

The professor explicitly writes out the matrix and its determinant. He jokes about making arithmetic mistakes, but executes it flawlessly.

Applying the Substitution Rule:

$$\mu_2(B_1) = \int_{\Psi(D)} 1\, dx = \int_D 1 \cdot \underbrace{|\det J\Psi(y_1, y_2)|}_{\text{Distortion Factor}}\, dy$$

Computing the Jacobian Matrix:

$$J\Psi(y_1, y_2) = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \frac{\partial x_1}{\partial y_2} \\ \frac{\partial x_2}{\partial y_1} & \frac{\partial x_2}{\partial y_2} \end{pmatrix} = \begin{pmatrix} \cos(y_2) & -y_1 \sin(y_2) \\ \sin(y_2) & y_1 \cos(y_2) \end{pmatrix}$$

Evaluating the Determinant:

$$\begin{aligned}\det J\Psi &= \big(\cos(y_2)\cdot y_1\cos(y_2)\big) - \big(-y_1\sin(y_2)\cdot\sin(y_2)\big)\\ &= y_1\cos^2(y_2) + y_1\sin^2(y_2)\\ &= y_1\underbrace{\big(\cos^2(y_2)+\sin^2(y_2)\big)}_{=1\text{ (Axiom)}}\\ &= y_1\end{aligned}$$

Redundant Explanation of Step: The Jacobian determinant tells us exactly how much a tiny square of parameter space dy_1dy_2 is stretched when it is mapped into the disk. Because the determinant is y_1, it means the "stretching" scales linearly with the radius. Grids near the outer edge of the circle are stretched much larger than grids near the origin.

> ***Cleaned Spoken Protocol [Time: 00:22:55 - 00:26:33]***
>
> *Substituting this back in, we must compute the double integral of y_1 over our rectangular domain D. But we have a problem: we haven't learned how to systematically compute multivariable integrals yet!*
> *We can solve this using a geometric trick. The integral of a positive function over a 2D domain represents the 3-dimensional volume of its hypograph—the physical region trapped underneath the function's surface.*
> *Let's add a vertical coordinate, x_3. We want the volume where $x_3 \leq y_1$, over our base rectangle $(0,1)\times(0,2\pi)$. Consider a full 3D bounding box with dimensions $1\times 2\pi\times 1$. The total volume is 2π. Our function, $x_3 = y_1$, forms a slanted plane that cuts this box perfectly in half along its diagonal. Therefore, the volume of our hypograph is exactly half the volume of the box, which gives us π.*

Board State & Physical Actions: Board State 10: The Hypograph Wedge

The professor completes the proof not with calculus, but by drawing a 3D bounding box and showing how the function $f(y_1,y_2) = y_1$ cuts it precisely in half.

The Final Integration via Geometry:

$$\mu_2(B_1) = \int_{(0,1)\times(0,2\pi)} \underbrace{y_1}_{\text{Height } x_3}\ \underbrace{dy_1\,dy_2}_{\text{Base Area}} \tag{5}$$



Redundant Explanation of Step: By interpreting the integral physically, we see the function y_1 creates a wedge. Because the wedge is a perfect diagonal bisection of a rectangular prism, we can compute its volume using simple geometry instead of advanced integration rules.

$$\mu_2(B_1) = \frac{1}{2}\,\text{Vol}(\text{Bounding Box}) = \frac{1}{2}(1\cdot 2\pi\cdot 1) = \pi$$

5 Example: Volume of the Unit Ball in $\mathbb{R}^3$

Cleaned Spoken Protocol [Time: 00:26:33 - 00:32:00]

That trick is wonderful, but let us move to a harder question. Let's compute the volume of the unit ball in $\mathbb{R}^3$. We know from geometry that the answer should be $\frac{4}{3}\pi$, but we have not proved it analytically yet.
We will try the exact same approach using 3D polar coordinates—spherical coordinates. We like spherical coordinates because they flawlessly transform rectangular boxes into spheres. Let our mapping be $\Psi(y_1, y_2, y_3)$.
The first coordinate, y_1, is always the radius. The second coordinate, y_2, is the longitude—the deviation from a fixed meridian. The third coordinate, y_3, represents the elevation. Let me write the formulas down...

Board State & Physical Actions: Board State 11: The Famous Mistake

The professor attempts to write the spherical coordinates. He intends for y_3 to be the "elevation" (latitude, measured from the equator). However, muscle memory kicks in, and he writes the standard physics convention where the angle is measured from the North Pole (colatitude). This is mathematically incorrect for his stated domain.

The Initial (Incorrect) Board State:

$$\begin{aligned} x_1 &= y_1 \cos(y_2) \sin(y_3) \\ x_2 &= y_1 \sin(y_2) \sin(y_3) \\ x_3 &= y_1 \cos(y_3) \end{aligned} \tag{6}$$

Insight: The Coordinate Trap

There are two competing standards for spherical coordinates. Mathematics often uses latitude (angle from the equator, meaning $z = r \sin \phi$). Physics usually uses colatitude (angle from the North Pole, meaning $z = r \cos \phi$). The professor mixed the words of one with the math of the other. This happens to the best of us, and it is a phenomenal active-learning moment for the room.

Cleaned Spoken Protocol [Time: 00:32:00 - 00:32:53]

Wait, looking at this... did I make a mistake?
Student: "If we would like y_3 to represent the elevation from the equator, shouldn't we switch the sine and cosine in the formula for the polar coordinates?"
Ah! The y_3 is the elevation, right? Yes, you are completely right. If we use the cosine for the x_3 coordinate, we are measuring the deviation from the North Pole. If we want the elevation from the equator, the vertical x_3 coordinate must be governed by the sine function. Let me fix this on the board. The x_3 gets the sine, and the others get the cosine. Excellent catch.

Board State & Physical Actions: Board State 12: The Correction

The professor erases the sines and cosines and formally swaps them, bringing the math into perfect alignment with his intended domain for elevation.

The Corrected Board State (Stroke-by-Stroke): The mapping $\Psi : D \to \mathbb{R}^3$ for elevation-

based spherical coordinates:

$$
\begin{aligned}
x_1 &= y_1 \cos(y_2) \underbrace{\cos(y_3)}_{\text{Corrected}} \\
x_2 &= y_1 \sin(y_2) \underbrace{\cos(y_3)}_{\text{Corrected}} \\
x_3 &= y_1 \underbrace{\sin(y_3)}_{\text{Corrected}}
\end{aligned}
\tag{7}
$$

Redundant Explanation of Step: By setting $x_3 = y_1 \sin(y_3)$, we ensure that when the elevation angle $y_3 = 0$, our height x_3 is 0 (putting us on the equator). When $y_3 = \pi/2$, $\sin(\pi/2) = 1$, putting us exactly at the North Pole $x_3 = y_1$.

The parameter domain D is formally defined as:

$$
D = \underbrace{(0,1)}_{y_1(\text{Radius})} \times \underbrace{(0,2\pi)}_{y_2(\text{Longitude})} \times \underbrace{\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)}_{y_3(\text{Elevation})}
$$

Board State & Physical Actions: TikZ: Spherical Coordinates Visualized

Visual representation of the corrected coordinate system, explicitly showing y_3 sweeping up from the equatorial plane.



Board State & Physical Actions: End of Second Segment

The lecture protocol currently ends here. The next mathematical step will be computing the 3×3 Jacobian determinant of this corrected system, leading directly into Cavalieri's Principle to evaluate the resulting 4-dimensional integral.

6 Hyper-Detailed Pedagogical Blueprint Real Analysis: Change of Variables & Rigorous Foundations (Part 3)

Director's Cut & AI Extraction Protocol
Lecture Segment: 32:00 – End

5 Completing the 3D Unit Ball Setup

Cleaned Spoken Protocol [Time: 00:32:53 - 00:36:10]

Now that we have the correct spherical coordinates defined, we need the Jacobian determinant. I am not going to spend seven minutes on the board computing this 3×3 matrix, but I highly encourage you to do it at home. Because of the trigonometric identity $\sin^2(\theta) + \cos^2(\theta) = 1$, many terms simplify beautifully.
In the end, the determinant of the Jacobian evaluates to $y_1^2 \cos(y_3)$. Notice that because our elevation angle y_3 goes strictly from $-\pi/2$ to $\pi/2$, the cosine function is always positive. Therefore, the absolute value of the determinant is simply $y_1^2 \cos(y_3)$ itself.
Using our Change of Variables formula, the volume of the unit ball in $\mathbb{R}^3$ is equal to the integral over our rectangular parameter domain D.

Board State & Physical Actions: Board State 13: Finalizing the 3D Integral
The professor skips the brute-force matrix algebra, writes the final Jacobian determinant, and constructs the triple integral on the board.

Mathematical Setup (Stroke-by-Stroke): The Jacobian determinant simplifies to:

$$\det J\Psi = y_1^2 \cos(y_3)$$

Because $y_3 \in (-\pi/2, \pi/2)$, $\cos(y_3) > 0$. Applying the substitution rule:

$$\mu_3(B_1) = \int_D \underbrace{y_1^2 \cos(y_3)}_{\text{Distortion Factor}} \, dy = \int_0^1 \int_0^{2\pi} \int_{-\pi/2}^{\pi/2} y_1^2 \cos(y_3) \, dy_3 \, dy_2 \, dy_1 \tag{8}$$

Cleaned Spoken Protocol [Time: 00:36:10 - 00:40:07]

Now, you might ask me: "How do we do a geometric trick to compute this, like we did with the 2D wedge?" The answer is, we don't. I know of no simple geometric trick to evaluate this 4-dimensional hyper-volume.
While we could try to invent more sophisticated coordinate changes, we will instead approach this integral systematically. To compute integrals of this type effortlessly, we must introduce the slicing idea. We will establish Cavalieri's Principle, which leads directly to Fubini's Theorem.

6 The Slicing Idea: Cavalieri's Principle

Cleaned Spoken Protocol [Time: 00:40:07 - 00:42:00]

Let me motivate the slicing idea informally. The Cavalieri Principle states that if you have a 3D body—which for us will be a Jordan measurable set—you can compute its total volume by slicing it.
Imagine a potato. If you take a potato and cut a very thin vertical slice, that slice is almost perfectly uniform from its front face to its back face. You can approximate the volume of that single slice by multiplying its cross-sectional area by its tiny thickness. If you sum the volumes of all these slices, the total volume of the potato is perfectly preserved.

Board State & Physical Actions: TikZ: The Potato Analogy

The professor draws a 3D blob (a potato) and highlights a single, thin vertical slice to illustrate the concept of $V \approx \sum A(x)\Delta x$.



Cleaned Spoken Protocol [Time: 00:42:00 - 00:46:30]

How do we do this systematically? We state it in terms of Jordan measures. We split our n-dimensional space, $\mathbb{R}^n$, into a product space: $\mathbb{R}^k \times \mathbb{R}^{n-k}$. Let the base coordinates be x and the vertical coordinates be y.
Let $A \subset \mathbb{R}^n$ be a Jordan measurable set. For a specific base point x, we define the slice A_x as the set of all vertical points y that lie inside A.
*Here is the technical hurdle: Just because A is Jordan measurable in $\mathbb{R}^n$ does not guarantee that the slice A_x is Jordan measurable in the lower-dimensional space $\mathbb{R}^{n-k}$. But we don't care! The outer measure and inner measure are *always* well-defined for any bounded set. So, we define the upper marginal density $\overline{\Phi}(x)$ using the outer measure, and the lower marginal density $\underline{\Phi}(x)$ using the inner measure.*

Board State & Physical Actions: Board State 14: Formalizing the Slices

The professor writes down the formal set notation for slices and explicitly addresses the "non-measurable slice" edge case using inner/outer measures.

Mathematical Setup (Stroke-by-Stroke): Split the space: $\mathbb{R}^n = \mathbb{R}^k \times \mathbb{R}^{n-k}$. A point is (x, y). Let $A \subset \mathbb{R}^n$ be Jordan measurable.

Definition 1 (Slices)**.** For a fixed base coordinate $x \in \mathbb{R}^k$, the slice A_x is defined as:

$$A_x = \{y \in \mathbb{R}^{n-k} \mid (x, y) \in A\}$$

To bypass slices that are not Jordan measurable, we define the **marginal densities**:

$$\overline{\Phi}(x) = \mu^{\text{out}}_{n-k}(A_x) \quad \text{(Upper Marginal)}$$
$$\underline{\Phi}(x) = \mu^{\text{in}}_{n-k}(A_x) \quad \text{(Lower Marginal)}$$

Cavalieri's Principle

Both marginal functions are Riemann integrable over x, and their integrals yield the total volume of A:

$$\int \overline{\Phi}(x)\, dx = \int \underline{\Phi}(x)\, dx = \mu_n(A)$$

Insight: The "I Don't Care" Philosophy

When the professor says "I don't care" about the slice not being Jordan measurable, he is imparting profound mathematical wisdom. In integration, sets of measure zero or pathological boundary anomalies disappear when you sum over them. The upper and lower integrals act as a squeeze theorem—even if individual slices are "fuzzy," integrating over the entire domain squeezes that fuzziness down to zero.

7 Proof by Counting Pixels

Cleaned Spoken Protocol [Time: 00:46:30 - 00:52:00]

How do we prove this? Because A is Jordan measurable, we can approximate it using a very small grid of dyadic cubes—like pixels on a screen.
For any $\epsilon > 0$, we can find an inner approximation F (made entirely of yellow pixels inside A) and an outer approximation G (made of red pixels covering A), such that the volume difference between G and F is less than ϵ.
Because these sets are built from perfect cubes aligned to a grid, what happens when we slice them? As long as our base point x stays within the same column of the grid, the slice of F or G cannot change! Therefore, the measures of these slices, $\mu(F_x)$ and $\mu(G_x)$, are just dyadic step functions.

Board State & Physical Actions: TikZ: The Dyadic Pixel Proof

The professor illustrates the inner (F) and outer (G) approximations. He notes that integrating a dyadic step function is just combinatorics—counting the total number of pixels.



Slice Column x

Proof Logic (Redundant Mathematical Breakdown): 1. A is sandwiched: $F \subset A \subset G$. 2. This forces the slices to be sandwiched: $F_x \subset A_x \subset G_x$. 3. Taking the measure of the slices:

$$\mu(F_x) \leq \underline{\Phi}(x) \leq \overline{\Phi}(x) \leq \mu(G_x)$$

4. Integrating the step functions $\mu(F_x)$ and $\mu(G_x)$ over the base dx simply counts the total number of pixels in F and G. Thus:

$$\text{Vol}(F) \leq \int \underline{\Phi}(x)dx \leq \int \overline{\Phi}(x)dx \leq \text{Vol}(G)$$

5. Since we chose our grid such that $\text{Vol}(G) - \text{Vol}(F) < \epsilon$, as $\epsilon \to 0$, the outer and inner pixel volumes squeeze the marginal integrals perfectly to $\text{Vol}(A)$.

8 Fubini's Theorem: Cavalieri for Functions

Cleaned Spoken Protocol [Time: End of Lecture Segment]

Fubini's theorem is the immediate consequence of Cavalieri's Principle applied to functions instead of just shapes.
If you have a function $f(x, y)$ that is Riemann integrable over A, you can compute the total multi-dimensional integral by integrating over the slices first. You integrate over y for a fixed x, and then you sum those results over all x.
The proof is identical. In fact, remember that the integral of a function is just the volume of its hypograph! Fubini's Theorem is quite literally just Cavalieri's Principle applied to the $n + 1$ dimensional hypograph wedge of the function.

Fubini's Theorem

$$\int_A f(x, y)\, d(x, y) = \int_{\mathbb{R}^k} \left(\int_{\mathbb{R}^{n-k}} \tilde{f}_x(y)\, dy \right) dx$$

Insight: The Grand Finale

The entire lecture culminates here. The 2D disk trick (where the function $x_3 = y_1$ created a wedge) was a manual, geometric application of Fubini's theorem! The professor used a visual trick to solve the integral before proving the theorem that justifies it, completing the pedagogical loop.