



UNIFICATION MODULO BOOLEAN RINGS

Florian Starke

Dresden, 2015/1/24

Structure

Equational Unification

Boolean Rings

Equational Unification

Unification Problem

$$S := \left\{ f(a, x) \stackrel{?}{\approx} x \right\}$$

Equational Unification

Unification Problem

$$I := \{f(x, x) \approx x\}$$

$$S := \left\{ f(a, x) \stackrel{?}{\approx} x \right\}$$

Equational Unification

Unification Problem

$$I := \{f(x, x) \approx x\}$$

$$S := \left\{ f(a, x) \stackrel{?}{\approx} x \right\} \quad \Rightarrow \quad S := \left\{ f(a, x) \stackrel{?}{\approx}_I x \right\}$$

Equational Unification

Unification Classes

Let S be an I-unification problem ...

$$\text{elementary: } S := \left\{ f(y, x) \stackrel{?}{\approx}_I x \right\}$$

$$\text{with constants: } S := \left\{ f(a, x) \stackrel{?}{\approx}_I x \right\}$$

$$\text{general: } S := \left\{ f(g(a), x) \stackrel{?}{\approx}_I x \right\}$$

Equational Unification

MGU?

$$C := \{f(x, y) \approx f(y, x)\}$$

$$S := \left\{ f(x, y) \stackrel{?}{\approx}_C f(a, b) \right\}$$

Equational Unification

MGU?

$$C := \{f(x, y) \approx f(y, x)\}$$

$$S := \left\{ f(x, y) \stackrel{?}{\approx}_C f(a, b) \right\}$$

$$\sigma_1 := \{x \mapsto a, y \mapsto b\}$$

$$\sigma_2 := \{x \mapsto b, y \mapsto a\}$$

Equational Unification

More General

$$\begin{aligned} \sigma &< \sigma' \\ \text{iff} \\ \exists \delta : \delta(\sigma) &= \sigma' \end{aligned}$$

Equational Unification

More General

$$\begin{array}{ccc} \sigma < \sigma' & & \sigma \lesssim_E^X \sigma' \\ \text{iff} & \Rightarrow & \text{iff} \\ \exists \delta : \delta(\sigma) = \sigma' & & \exists \delta : \delta(\sigma(x)) \approx_E \sigma'(x) \text{ for all } x \in X \end{array}$$

Equational Unification

Minimal Complete Sets

Let S be an E-unification problem. A minimal complete set of S is a set of substitutions \mathcal{M} that satisfy the following properties.

- each $\sigma \in \mathcal{M}$ is an E-unifier of S
- for all E-unifiers θ of S there exists a $\sigma \in \mathcal{M}$ such that $\sigma \lesssim_E^X \theta$
- for all $\sigma, \sigma' \in \mathcal{M}$, $\sigma \lesssim_E^X \sigma'$ implies $\sigma = \sigma'$.

Equational Unification

Unification Type of \approx_E

unitary iff for all E-unification problems S there exists a minimal complete set of cardinality ≤ 1 .

Equational Unification

Unification Type of \approx_E

- unitary iff for all E-unification problems S there exists a minimal complete set of cardinality ≤ 1 .
- finitary iff for all E-unification problems S there exists a minimal complete set with finite cardinality.

Equational Unification

Unification Type of \approx_E

- unitary iff for all E-unification problems S there exists a minimal complete set of cardinality ≤ 1 .
- finitary iff for all E-unification problems S there exists a minimal complete set with finite cardinality.
- infinitary iff for all E-unification problems S there exists a minimal complete set, and there exists an E-unification problem for which this set is infinite.

Equational Unification

Unification Type of \approx_E

- unitary iff for all E-unification problems S there exists a minimal complete set of cardinality ≤ 1 .
- finitary iff for all E-unification problems S there exists a minimal complete set with finite cardinality.
- infinitary iff for all E-unification problems S there exists a minimal complete set, and there exists an E-unification problem for which this set is infinite.
- zero iff there exists an E-unification problem that does not have a minimal complete set.

Boolean Rings

$$B := \left\{ \begin{array}{ll} x + y \approx y + x, & x * y \approx y * x, \\ (x + y) + z \approx x + (y + z), & (x * y) * z \approx x * (y * z), \\ x + x \approx 0, & x * x \approx x, \\ 0 + x \approx x, & 0 * x \approx 0, \\ x * (y + z) \approx (x * y) + (x * z), & 1 * x \approx x \end{array} \right\}$$

Boolean Rings

Interpretation

$$\mathcal{B}_2 := (\Delta^{\mathcal{B}_2}, \cdot^{\mathcal{B}_2})$$

$$\Delta^{\mathcal{B}_2} := \{\perp, \top\}$$

Boolean Rings

Interpretation

$$\mathcal{B}_2 := (\Delta^{\mathcal{B}_2}, \cdot^{\mathcal{B}_2})$$

$$\Delta^{\mathcal{B}_2} := \{\perp, \top\}$$

$$(x + y)^{\mathcal{B}_2} := \left(x^{\mathcal{B}_2} \wedge \neg y^{\mathcal{B}_2} \right) \vee \left(\neg x^{\mathcal{B}_2} \wedge y^{\mathcal{B}_2} \right)$$

$$(x * y)^{\mathcal{B}_2} := x^{\mathcal{B}_2} \wedge y^{\mathcal{B}_2}$$

$$0^{\mathcal{B}_2} := \perp$$

$$1^{\mathcal{B}_2} := \top$$

Boolean Rings

Example

$$\begin{aligned}(1 + 0)^{\mathcal{B}_2} &= \left(1^{\mathcal{B}_2} \wedge \neg 0^{\mathcal{B}_2}\right) \vee \left(\neg 1^{\mathcal{B}_2} \wedge 0^{\mathcal{B}_2}\right) \\&= (\top \wedge \neg \perp) \vee (\neg \top \wedge \perp) \\&= \top \vee \perp \\&= \top\end{aligned}$$

Boolean Rings

Polynomial Form

asd