



TECHNISCHE
UNIVERSITÄT
DRESDEN

SMOOTH DIGRAPHS MODULO PP-CONSTRUCTABILITY

Florian Starke

Definitions

Let \mathfrak{A} be a structure.

A structure \mathfrak{B} is in $H(\mathfrak{A})$ if there are homomorphisms $f: \mathfrak{A} \rightarrow \mathfrak{B}$ and $g: \mathfrak{B} \rightarrow \mathfrak{A}$.

Definitions

Let \mathfrak{A} be a structure.

A structure \mathfrak{B} is in $H(\mathfrak{A})$ if there are homomorphisms $f: \mathfrak{A} \rightarrow \mathfrak{B}$ and $g: \mathfrak{B} \rightarrow \mathfrak{A}$.

$$\mathfrak{A}^2, \mathfrak{A} \cup \mathfrak{A} \in H(\mathfrak{A})$$

Definitions

Let \mathfrak{A} be a structure.

A structure \mathfrak{B} is in $H(\mathfrak{A})$ if there are homomorphisms $f: \mathfrak{A} \rightarrow \mathfrak{B}$ and $g: \mathfrak{B} \rightarrow \mathfrak{A}$.

$$\mathfrak{A}^2, \mathfrak{A} \cup \mathfrak{A} \in H(\mathfrak{A})$$

A structure \mathfrak{B} is in $PP(\mathfrak{A})$ if there is a d such that $B = A^d$ and every k -ary relation in \mathfrak{B} is, as a $k \cdot d$ -ary relation in \mathfrak{A} , pp-definable.

Definitions

Let \mathfrak{A} be a structure.

A structure \mathfrak{B} is in $H(\mathfrak{A})$ if there are homomorphisms $f: \mathfrak{A} \rightarrow \mathfrak{B}$ and $g: \mathfrak{B} \rightarrow \mathfrak{A}$.

$$\mathfrak{A}^2, \mathfrak{A} \cup \mathfrak{A} \in H(\mathfrak{A})$$

A structure \mathfrak{B} is in $PP(\mathfrak{A})$ if there is a d such that $B = A^d$ and every k -ary relation in \mathfrak{B} is, as a $k \cdot d$ -ary relation in \mathfrak{A} , pp-definable.

$$\begin{array}{c} \hookrightarrow \hookrightarrow \\ \hookrightarrow \hookrightarrow \end{array} \in PP(\mathfrak{A})$$
$$\Phi_E(x, y) = "x = y"$$

Definitions

Let \mathfrak{A} be a structure.

A structure \mathfrak{B} is in $H(\mathfrak{A})$ if there are homomorphisms $f: \mathfrak{A} \rightarrow \mathfrak{B}$ and $g: \mathfrak{B} \rightarrow \mathfrak{A}$.

$$\mathfrak{A}^2, \mathfrak{A} \cup \mathfrak{A} \in H(\mathfrak{A})$$

A structure \mathfrak{B} is in $PP(\mathfrak{A})$ if there is a d such that $B = A^d$ and every k -ary relation in \mathfrak{B} is, as a $k \cdot d$ -ary relation in \mathfrak{A} , pp-definable.

$$\begin{array}{c} \circlearrowleft \\ \circlearrowright \end{array} \in PP(\mathfrak{A})$$
$$\Phi_E(x, y) = "x = y"$$

A structure \mathfrak{B} is **pp-constructable** from \mathfrak{A} if $\mathfrak{B} \in H(PP(\mathfrak{A}))$. In this case we say $\mathfrak{B} \geq \mathfrak{A}$.

Definitions

Let \mathfrak{A} be a structure.

A structure \mathfrak{B} is in $H(\mathfrak{A})$ if there are homomorphisms $f: \mathfrak{A} \rightarrow \mathfrak{B}$ and $g: \mathfrak{B} \rightarrow \mathfrak{A}$.

$$\mathfrak{A}^2, \mathfrak{A} \cup \mathfrak{A} \in H(\mathfrak{A})$$

A structure \mathfrak{B} is in $PP(\mathfrak{A})$ if there is a d such that $B = A^d$ and every k -ary relation in \mathfrak{B} is, as a $k \cdot d$ -ary relation in \mathfrak{A} , pp-definable.

$$\begin{array}{c} \circlearrowleft \\ \circlearrowright \end{array} \in PP(\mathfrak{A})$$

$$\Phi_E(x, y) = "x = y"$$

A structure \mathfrak{B} is **pp-constructable** from \mathfrak{A} if $\mathfrak{B} \in H(PP(\mathfrak{A}))$. In this case we say $\mathfrak{B} \geq \mathfrak{A}$.


$$\textcircled{1} \in H(PP(\mathfrak{A}))$$

Definitions

A directed graph G is a **smooth digraphs** if every vertex has in-degree and out-degree at least 1.

Definitions


A directed graph G is a **smooth digraphs** if every vertex has in-degree and out-degree at least 1.



Definitions

A directed graph G is a **smooth digraphs** if every vertex has in-degree and out-degree at least 1.

Let \mathfrak{S} be the poset induced by the quasi order \geq on all finite smooth digraphs.



Dividing the graphs

$$[G] \in \mathfrak{G}$$

Dividing the graphs

G can pp-construct
every finite structure



G has 4-ary Siggers
and its core is a disjoint
union of cycles

$[G] \in \mathfrak{G}$

Dividing the graphs

G can pp-construct
every finite structure




$[G]$ contains a disjoint
union of cycles



G has 4-ary Siggers
and its core is a dis-
joint union of cycles

$[G] \in \mathfrak{G}$


Dividing the graphs



Poset first glance



Poset first glance



Multiples




Multiples



Multiples



Multiples



Division




Division



Division

$$\Phi_E(x, y) = x \stackrel{2}{\rightarrow} y$$



Division

$$\Phi_E(x, y) = x \stackrel{2}{\rightarrow} y$$



Division

$$\Phi_E(x, y) = x \xrightarrow{k} y$$




$$a \div k = \frac{a}{\gcd(a, k)}$$

Multiplication



Multiplication




Multiplication

$$\Phi_E \begin{pmatrix} x_1, x_2, x_3, \\ y_1, y_2, y_3 \end{pmatrix} = x_1 \rightarrow y_3$$

$$\wedge x_2 = y_1$$

$$\wedge x_3 = y_2$$




Multiplication

$$\Phi_E \begin{pmatrix} x_1, x_2, x_3, \\ y_1, y_2, y_3 \end{pmatrix} = x_1 \rightarrow y_3$$

$$\wedge x_2 = y_1$$

$$\wedge x_3 = y_2$$




Multiplication

$$\Phi_E \begin{pmatrix} x_1, x_2, x_3, \\ y_1, y_2, y_3 \end{pmatrix} = x_1 \rightarrow y_3$$

$$\wedge x_2 = y_1$$

$$\wedge x_3 = y_2$$




Multiplication

$$\Phi_E \left(\begin{matrix} x_1, x_2, x_3, \\ y_1, y_2, y_3 \end{matrix} \right) = x_1 \rightarrow y_3$$

$$\wedge x_2 = y_1$$

$$\wedge x_3 = y_2$$



Multiplication


$$\Phi_E \begin{pmatrix} x_1, x_2, \\ y_1, y_2 \end{pmatrix} = x_1 \rightarrow y_2$$
$$\wedge x_2 = y_1$$



Multiplication

$$\Phi_E \begin{pmatrix} x_1, x_2, \\ y_1, y_2 \end{pmatrix} = x_1 \rightarrow y_2$$


$$\wedge x_2 = y_1$$



Multiplication


$$\Phi_E \begin{pmatrix} x_1, x_2, \\ y_1, y_2 \end{pmatrix} = x_1 \rightarrow y_2$$

$$\wedge x_2 = y_1$$



Multiplication

$$\Phi_E \begin{pmatrix} x_1, x_2, \\ y_1, y_2 \end{pmatrix} = x_1 \rightarrow y_2 \\ \wedge x_2 = y_1$$



Multiplication

$$\begin{aligned}\Phi_E \left(\begin{array}{c} x_1, \dots, x_k, \\ y_1, \dots, y_k \end{array} \right) &= x_1 \rightarrow y_k \\ &\wedge x_2 = y_1 \\ &\quad \vdots \\ &\wedge x_k = y_{k-1}\end{aligned}$$



Multiplication

$$\begin{aligned}\Phi_E \left(\begin{matrix} x_1, \dots, x_k, \\ y_1, \dots, y_k \end{matrix} \right) &= x_1 \rightarrow y_k \\ &\wedge x_2 = y_1 \\ &\quad \vdots \\ &\wedge x_k = y_{k-1}\end{aligned}$$



Multiplication

$$\begin{aligned}\Phi_E \left(\begin{matrix} x_1, \dots, x_k, \\ y_1, \dots, y_k \end{matrix} \right) &= x_1 \rightarrow y_k \\ &\wedge x_2 = y_1 \\ &\quad \vdots \\ &\wedge x_k = y_{k-1}\end{aligned}$$



$$a \times k = \prod_{\alpha_j \neq 0} p_i^{\alpha_j + \kappa_j}$$

Multiplication

$$\begin{aligned}\Phi_E \left(\begin{matrix} x_1, \dots, x_k, \\ y_1, \dots, y_k \end{matrix} \right) &= x_1 \rightarrow y_k \\ &\wedge x_2 = y_1 \\ &\quad \vdots \\ &\wedge x_k = y_{k-1}\end{aligned}$$



$$a \ltimes k = \prod_{\alpha_j \neq 0} p_i^{\alpha_j + \kappa_j}$$


$$3 \ltimes 3 = 9$$

$$3 \ltimes 2 = 3$$


$$3 \ltimes 6 = 9$$

$$2 \ltimes 12 = 8$$

Normal form




Normal form



G is in **normal form** if


Normal form



G is in **normal form** if

- for all $a, a' \in G$ we have $a \mid a'$ implies $a = a'$ and


Normal form



G is in **normal form** if

- for all $a, a' \in G$ we have $a \mid a'$ implies $a = a'$ and
- if for an $a \in G$ we have $p \mid a$, then there is an $a' \in G$ with $p \mid a'$ but $p^2 \nmid a'$.

Poset second glance




Combination



Combination

$$\Phi_E \left(\begin{matrix} x_1, x_2, \\ y_1, y_2 \end{matrix} \right) = x_1 \rightarrow y_1 \\ \wedge x_2 \rightarrow y_2$$




Combination

$$\Phi_E \left(\begin{matrix} x_1, x_2, \\ y_1, y_2 \end{matrix} \right) = x_1 \rightarrow y_1$$

$$\wedge x_2 \rightarrow y_2$$

$$\wedge x_1 \xrightarrow{2} x_1$$

$$\wedge x_2 \xrightarrow{3} x_2$$




Combination

$$\Phi_E \begin{pmatrix} x_1, x_2, \\ y_1, y_2 \end{pmatrix} = x_1 \rightarrow y_1$$

$$\wedge x_2 \rightarrow y_2$$

$$\wedge x_1 \xrightarrow{2} x_1$$

$$\wedge x_2 \xrightarrow{3} x_2$$



Combination

$$\Phi_E \left(\begin{matrix} x_1, x_2, \\ y_1, y_2 \end{matrix} \right) = x_1 \rightarrow y_1$$

$$\wedge x_2 \rightarrow y_2$$


$$\wedge x_1 \xrightarrow{a} x_1$$

$$\wedge x_2 \xrightarrow{b} x_2$$



$$\begin{aligned} a \nmid b, b \nmid a \\ a \vee b = \text{lcm}(a, b) \end{aligned}$$

Poset final glance



Poset final glance



Summary

$$a \vee b = \text{lcm}(a, b) \qquad a \dot{-} k = \frac{a}{\text{gcd}(a, k)} \qquad a \ltimes k = \prod_{\alpha_j \neq 0} p_j^{\alpha_j + \kappa_j}$$

Summary

$$a \vee b = \text{lcm}(a, b) \qquad a \dot{\div} k = \frac{a}{\text{gcd}(a, k)} \qquad a \ltimes k = \prod_{\alpha_j \neq 0} p_j^{\alpha_j + \kappa_j}$$

$$G \dot{\div} (k_1, \dots, k_d) = \{(a_1 \dot{\div} k_1) \vee \dots \vee (a_d \dot{\div} k_d) \mid a_1, \dots, a_d \in G\}$$

$$G \ltimes (k_1, \dots, k_d) = \{(a_1 \ltimes k_1) \vee \dots \vee (a_d \ltimes k_d) \mid a_1, \dots, a_d \in G\}$$

Summary

$$a \vee b = \text{lcm}(a, b) \qquad a \dot{\div} k = \frac{a}{\text{gcd}(a, k)} \qquad a \ltimes k = \prod_{\alpha_j \neq 0} p_j^{\alpha_j + \kappa_j}$$

$$G \dot{\div} (k_1, \dots, k_d) = \{(a_1 \dot{\div} k_1) \vee \dots \vee (a_d \dot{\div} k_d) \mid a_1, \dots, a_d \in G\}$$

$$G \ltimes (k_1, \dots, k_d) = \{(a_1 \ltimes k_1) \vee \dots \vee (a_d \ltimes k_d) \mid a_1, \dots, a_d \in G\}$$

$$2 \cdot 3, 5 \cdot 7 \dot{\div} (2 \cdot 5, 3 \cdot 7)$$

$$\begin{array}{c} | \\ 2 \cdot 3, 5 \cdot 7 \end{array}$$

Summary

$$a \vee b = \text{lcm}(a, b) \qquad a \dot{\div} k = \frac{a}{\text{gcd}(a, k)} \qquad a \ltimes k = \prod_{\alpha_j \neq 0} p_j^{\alpha_j + \kappa_j}$$

$$G \dot{\div} (k_1, \dots, k_d) = \{(a_1 \dot{\div} k_1) \vee \dots \vee (a_d \dot{\div} k_d) \mid a_1, \dots, a_d \in G\}$$

$$G \ltimes (k_1, \dots, k_d) = \{(a_1 \ltimes k_1) \vee \dots \vee (a_d \ltimes k_d) \mid a_1, \dots, a_d \in G\}$$

$$\begin{array}{c} 2 \cdot 3, 5 \cdot 7, 2 \cdot 7, 3 \cdot 5 \\ \parallel \\ 2 \cdot 3, 5 \cdot 7 \dot{\div} (2 \cdot 5, 3 \cdot 7) \\ \mid \\ 2 \cdot 3, 5 \cdot 7 \end{array}$$

Summary

$$a \vee b = \text{lcm}(a, b) \qquad a \dot{\div} k = \frac{a}{\text{gcd}(a, k)} \qquad a \ltimes k = \prod_{\alpha_j \neq 0} p_j^{\alpha_j + \kappa_j}$$

$$G \dot{\div} (k_1, \dots, k_d) = \{(a_1 \dot{\div} k_1) \vee \dots \vee (a_d \dot{\div} k_d) \mid a_1, \dots, a_d \in G\}$$

$$G \ltimes (k_1, \dots, k_d) = \{(a_1 \ltimes k_1) \vee \dots \vee (a_d \ltimes k_d) \mid a_1, \dots, a_d \in G\}$$

$$\begin{array}{ccc}
 2 \cdot 3, 5 \cdot 7, 2 \cdot 7, 3 \cdot 5 & & \\
 \parallel & & \\
 2 \cdot 3, 5 \cdot 7 \dot{\div} (2 \cdot 5, 3 \cdot 7) & & 2, 3 \ltimes (4, 9) \\
 | & & | \\
 2 \cdot 3, 5 \cdot 7 & & 2, 3
 \end{array}$$

Summary

$$a \vee b = \text{lcm}(a, b) \qquad a \dot{\div} k = \frac{a}{\text{gcd}(a, k)} \qquad a \ltimes k = \prod_{\alpha_j \neq 0} p_j^{\alpha_j + \kappa_j}$$

$$G \dot{\div} (k_1, \dots, k_d) = \{(a_1 \dot{\div} k_1) \vee \dots \vee (a_d \dot{\div} k_d) \mid a_1, \dots, a_d \in G\}$$

$$G \ltimes (k_1, \dots, k_d) = \{(a_1 \ltimes k_1) \vee \dots \vee (a_d \ltimes k_d) \mid a_1, \dots, a_d \in G\}$$

$$\begin{array}{cc}
 2 \cdot 3, 5 \cdot 7, 2 \cdot 7, 3 \cdot 5 & 2^3, 2 \cdot 3, 3^3 \\
 \parallel & \parallel \\
 2 \cdot 3, 5 \cdot 7 \dot{\div} (2 \cdot 5, 3 \cdot 7) & 2, 3 \ltimes (4, 9) \\
 | & | \\
 2 \cdot 3, 5 \cdot 7 & 2, 3
 \end{array}$$

Summary

$$a \vee b = \text{lcm}(a, b) \qquad a \dot{\div} k = \frac{a}{\text{gcd}(a, k)} \qquad a \ltimes k = \prod_{\alpha_j \neq 0} p_j^{\alpha_j + \kappa_j}$$

$$G \dot{\div} (k_1, \dots, k_d) = \{(a_1 \dot{\div} k_1) \vee \dots \vee (a_d \dot{\div} k_d) \mid a_1, \dots, a_d \in G\}$$

$$G \ltimes (k_1, \dots, k_d) = \{(a_1 \ltimes k_1) \vee \dots \vee (a_d \ltimes k_d) \mid a_1, \dots, a_d \in G\}$$

$$Gf(t_1, \dots, t_d), \text{ where } f \in \{\dot{\div}, \ltimes\}^n \text{ and } t_i \in \{1, 2, \dots\}^n$$

$$\begin{array}{ccc} 2 \cdot 3, 5 \cdot 7, 2 \cdot 7, 3 \cdot 5 & & 2^3, 2 \cdot 3, 3^3 \\ \parallel & & \parallel \\ 2 \cdot 3, 5 \cdot 7 \dot{\div} (2 \cdot 5, 3 \cdot 7) & & 2, 3 \ltimes (4, 9) \\ | & & | \\ 2 \cdot 3, 5 \cdot 7 & & 2, 3 \end{array}$$

Summary

$$a \vee b = \text{lcm}(a, b)$$

$$a \dot{-} k = \frac{a}{\text{gcd}(a, k)}$$

$$a \ltimes k = \prod_{\alpha_j \neq 0} p_j^{\alpha_j + \kappa_j}$$

$$G \dot{-} (k_1, \dots, k_d) = \{(a_1 \dot{-} k_1) \vee \dots \vee (a_d \dot{-} k_d) \mid a_1, \dots, a_d \in G\}$$

$$G \ltimes (k_1, \dots, k_d) = \{(a_1 \ltimes k_1) \vee \dots \vee (a_d \ltimes k_d) \mid a_1, \dots, a_d \in G\}$$

$$G f(t_1, \dots, t_d), \text{ where } f \in \{\dot{-}, \ltimes\}^n \text{ and } t_i \in \{1, 2, \dots\}^n$$

$$(G_1 f_1 T_1) \vee \dots \vee (G_n f_n T_n)$$

|
G

For all $a \in G \setminus G_i$ we have $a \nmid \text{lcm}(G_i)$.

$$\begin{array}{c} 2 \cdot 3, 5 \cdot 7, 2 \cdot 7, 3 \cdot 5 \\ \parallel \\ 2 \cdot 3, 5 \cdot 7 \dot{-} (2 \cdot 5, 3 \cdot 7) \end{array}$$

|
 $2 \cdot 3, 5 \cdot 7$

$$\begin{array}{c} 2^3, 2 \cdot 3, 3^3 \\ \parallel \\ 2, 3 \ltimes (4, 9) \end{array}$$

|
 $2, 3$

Summary

$$a \vee b = \text{lcm}(a, b) \quad a \dot{\div} k = \frac{a}{\gcd(a, k)} \quad a \ltimes k = \prod_{\alpha_j \neq 0} p_j^{\alpha_j + \kappa_j}$$

$$G \dot{\div} (k_1, \dots, k_d) = \{(a_1 \dot{\div} k_1) \vee \dots \vee (a_d \dot{\div} k_d) \mid a_1, \dots, a_d \in G\}$$

$$G \ltimes (k_1, \dots, k_d) = \{(a_1 \ltimes k_1) \vee \dots \vee (a_d \ltimes k_d) \mid a_1, \dots, a_d \in G\}$$

$$G f(t_1, \dots, t_d), \text{ where } f \in \{\dot{\div}, \ltimes\}^n \text{ and } t_i \in \{1, 2, \dots\}^n$$

$$(G_1 f_1 T_1) \vee \dots \vee (G_n f_n T_n)$$

$$\begin{array}{c} 2 \cdot 3, 5 \cdot 7, 2 \cdot 7, 3 \cdot 5 \\ \parallel \\ 2 \cdot 3, 5 \cdot 7 \dot{\div} (2 \cdot 5, 3 \cdot 7) \end{array}$$

$$\begin{array}{c} 2^3, 2 \cdot 3, 3^3 \\ \parallel \\ 2, 3 \ltimes (4, 9) \end{array}$$

For all $a \in G \setminus G_i$ we have $a \nmid \text{lcm}(G_i)$.

$$2 \cdot 3, 5 \cdot 7$$

$$2, 3$$