

Combinatorics I, List 1

GUILHERME ZEUS DANTAS E MOURA

zeusdanmou@gmail.com

IMPA, Summer 2021

Problem 1

By defining, and calculating the expectation of, a suitable random variable, show that every graph G has a bipartite subgraph with at least $e(G)/2$ edges.

Solution. Let's pick a random bipartition (A, B) of $V(G)$, with probability $1/2$ of $v \in A$, chosen independently.

Define

$$X = \#(ab \in E(G) : a \in A \text{ and } b \in B).$$

Then,

$$\begin{aligned} \mathbb{E}[X] &= \sum_{uv \in E(G)} \mathbb{1}[\text{one of } u, v \text{ is in } A \text{ and the other is in } B] \\ &= \sum_{uv \in E(G)} \frac{1}{2} \\ &= \frac{e(G)}{2}. \end{aligned}$$

Therefore, there exists a bipartition (A, B) such that the subgraph $G' \subset G$ that keeps the edges going through A to B has at least $e(G)/2$ edges.

Problem 2

Show that $R(3,4) \leq 9$, $R(4,4) \leq 18$ and $R(3,3,3) \leq 17$.

Solution.

Problem 3

Show that every graph of average degree d contains a subgraph of minimum degree at least $d/2$. Deduce that $\text{ex}(n, T) \leq (k-1)n$ for every tree T with k vertices.

Solution of the first part. While there are vertices with degree smaller than $\frac{d}{2}$, throw them away.

Suppose we threw away all vertices. At each step, we threw away at most $\frac{d}{2}$ edges. Since we threw away all edges, this means $n \cdot \frac{d}{2} < e(G) = n \cdot \frac{d}{2}$; a contradiction.

Therefore, we stopped before throwing away all vertices, so we're done.

Solution of the second part. Suppose $e(G) \geq (k-1)n \implies \bar{d}(G) = 2(k-1)$. By the first part of this problem, there exists a subgraph $G' \subset G$ such that $\delta(G') \geq k-1$. By the first part of Problem 4, $T \subset G' \subset G$.

Thus, $\text{ex}(n, T) \leq (k-1)n$.

Problem 4

Show that if T is a tree with k vertices and G is a graph with minimum degree $k - 1$, then $T \subset G$. Deduce that $r(K_3, T) = 2k - 1$.

Solution of the first part. We'll use induction on k . If $k = 1$, we're done!

Pick a leaf v of T . Its unique edge connects it to u . Let T' be the tree without v . By induction, there is a copy $C_{T'}$ of T' in G . Let c_u be the copy of u in $C_{T'}$. Since $\deg(c_u) \leq k - 2$ in $C_{T'}$, but $\deg(c_u) \geq k - 1$ in G , there is some vertex that is connected to u outside of $C_{T'}$, say c_v . Thus, let C_T be $C_{T'}$, adding c_v . C_T is a copy of T inside G .

Solution of the second part. For the lower bound, pick two complete graphs with $k - 1$ vertices and paint them blue. All other edges are colored red.

For the upper bound, pick a vertex v . Suppose $d_R(v) \geq k$. If there is a red edge among the red neighborhood of v , we find a red triangle; on the other hand, if all edges are blue, then there is a blue-colored complete graph with k vertices, which surely contains a blue copy of the tree T .

Therefore, $d_R \leq k - 1$, and consequently, $d_B(v) \geq k - 1$, for all vertices v . The first part implies that there is a blue copy of the tree T .

Problem 5

Let T_1, \dots, T_k be subtrees of a tree T , any two of which have at least one vertex in common. Prove that there is a vertex in all the T_i .

Sketch. Induction on k . For $k = 1$, we're good!

Suppose there is no common vertex to all T_i . Induction hypothesis implies that, for each i , there is a vertex v_i that is in all the T_j , for all $j \neq i$. This implies that $v_i \notin V(T_i)$,

There is a path from v_1 to v_2 that does not go through v_3 inside T_3 (and also inside T).

There is a path from v_1 to v_3 that does not go through v_2 inside T_1 (and also inside T).

There is a path from v_2 to v_3 that does not go through v_1 inside T_2 (and also inside T).

Then, there must exist a cycle in T , which is a contradiction. See fig. 1.




Figure 1: Finding a cycle

Problem 6

Let $R_r(3)$ denote the r -colour Ramsey number of a triangle. Show that

$$2^r \leq R_r(3) \leq 3 \cdot r!.$$

Show moreover that $R_r(3) \leq 5^{r/2}$.

Solution for the lower bound.

Solution for the first upper bound. Let G be a graph with $R_r(3) - 1$ vertices. There must exist a coloring of the edges that does not produce a monochromatic triangle.

Pick a vertex v . For each color, look to the neighborhood of v for that color. There can't be a edge of this color inside this neighborhood, neither a triangle of any other color. Thus, the size of this neighborhood is at most $R_{r-1}(3) - 1$. Repeating this for every color, we have

$$R_r(3) - 1 \leq r(R_{r-1}(3) - 1) + 1,$$

which implies, for $r \geq 2$,

$$R_r(3) \leq r \cdot R_{r-1}(3)$$

Telescoping this recursion, we get

$$R_r(3) \geq r! \cdot R_1(3) = 3 \cdot r!$$

Solution for the second upper bound.

Problem 7

Let $g(n)$ be the largest integer such that there exists a graph with the following properties: $|V(G)| = n$, $e(G) = m$, and it is possible to red-blue colour the edges of G without creating a monochromatic triangle.

Show that $g(n)/\binom{n}{2}$ converges, and find c such that $g(n)/\binom{n}{2} \rightarrow c$ as $n \rightarrow \infty$.

Solution. For the lower bound, pick Turán's graph $T_5(n)$. Color its edges as in fig. 2.

For the upper bound, if there are more edges in G than $t_5(n)$, by Turán, 1941, there must be a K_6 in the graph. Therefore, there is a monochromatic triangle, since $R(3,3) = 6$.

Therefore, $g(n) = t_5(n) \approx \left(\frac{1}{5}\right) \binom{n}{2}$, which implies that $g(n)/\binom{n}{2} \rightarrow \frac{1}{5}$ as $n \rightarrow \infty$.




Figure 2: Construction for the lower bound

Problem 8

Recall that $\alpha(G)$ denotes the size of the largest independent set in G . Show that, for every graph G ,

$$\alpha(G) \geq \sum_{v \in V(G)} \frac{1}{d(v) + 1}.$$

Sketch. Let $V(G) = [n]$. Pick a random permutation $\pi: [n] \rightarrow [n]$. For each $v \in [n]$, define

$$A_v := \{u \in V(G) : u \sim v \text{ and } \pi(v) < \pi(u)\}.$$

Problem 9

Let $C(s)$ be the smallest n such that every connected graph on n vertices has, as an *induced* subgraph, either a complete K_s , a star $K_{1,s}$ or a path P_s of length s .

Show that $C(s) \leq R(s)^s$, where $R(S)$ is the Ramsey number of s .

Solution.

Problem 10

Prove that $R(3, k) \geq k^{1+c}$ for some $c > 0$.

Sketch.

$$X_k = \#(K_k \subset G(n, p))$$

$$X_3 = \#(K_3 \subset \overline{G(n, p)})$$

Problem 11

Show that there is an infinite set S of positive integers such that the sum of any two distinct elements of S has an even number of distinct prime factors.

Solution. Let $S = \{n : n \equiv 1 \pmod{3}\}$. Note that $3 \nmid x + y$ for any $x, y \in S$.

Let $x \sim y$ be blue iff $x + y$ has a even number of distinct prime factors. Ramsey's Theorem implies that there is an infinite monochromatic set. If this set is blue, we are done! If this set is red, then multiply every number by 3 (a new prime factor!) — the scaled set has the property we want.

Problem 12

Suppose we are given n points and n lines in the plane. Show that there are at most $O(n^{3/2})$ point-line incidences.

Solution.