

Assignment

Barath surya M
EE22BTECH11014

Consider the experiment of throwing a die. If a multiple of 3 comes up, throw the die again. If any other number comes up, toss a coin. Find the conditional probability of the event the coin shows a tail, given that at least one die shows a 3.

Solution: Let, the states S_0 and S_1 describe the

S_0	$\Sigma(Y = k); k \in (3, 6)$
S_1	$\Sigma(Y = k); k \in (1, 2, 4, 5)$
S_2	Outcome of coin toss is heads
S_3	Outcome of coin toss is tails

outcomes of dice throws.

S_2 and S_3 describe the outcomes of coin toss. Conditional Probability is that "The coin shows tails" given that "at least one die shows a 3". Since, a Markov chain does not depend on the past outcomes,

$$p_{S_3|S_0} = \Pr(X_n = S_3 | X_1 = S_0) \quad (1)$$

Transition Probability Matrix is given as,

$$\mathbf{P} = \begin{pmatrix} \frac{1}{3} & \frac{2}{3} & 0 & 0 \\ 0 & 0 & \frac{1}{2} & \frac{1}{2} \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad (2)$$

The given condition is "3 occurs at least once", and we let occurrence of 3 as the initial state, Since Markov chain does not depend on past outcomes. So, $\Pr(X = S_0) = 1$ and 0 for all other states.

And State vector is ,

$$\mathbf{Q}_0 = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} \quad (3)$$

The long term Probability that system will be in each state is called stationary state, and stationary state probability is given as,

$$\mathbf{P}\mathbf{x} = \mathbf{x} \quad (4)$$

where \mathbf{x} is steady state probability vector and \mathbf{P} is the transition probability matrix.

So, after long time,

$$\mathbf{Q}_1 = \mathbf{P}\mathbf{Q}_0 \quad (5)$$

$$\mathbf{Q}_2 = \mathbf{P}\mathbf{Q}_1 \quad (6)$$

$$\vdots \quad (7)$$

$$\mathbf{Q}_n = \mathbf{P}\mathbf{Q}_{n-1} \quad (8)$$

So substituting the state vectors we get,

$$\mathbf{Q}_n = \mathbf{P}^n \mathbf{Q}_0 \quad (9)$$

$$(10)$$

applying limits to find the stationary probability vector,

$$\lim_{n \rightarrow \infty} \mathbf{Q}_n = \mathbf{P}^n \mathbf{Q}_0 \quad (11)$$

By substituting the values of \mathbf{Q}_0 and \mathbf{P} in the above equation, We get the steady state probability vector as,

$$\mathbf{x} = \begin{pmatrix} 0 \\ 0 \\ \frac{1}{2} \\ \frac{1}{2} \end{pmatrix} \quad (12)$$

So, the probability that "coin shows tails" given that "Die shows at least one 3" is,

$$p_{3|1} = \frac{1}{2} \quad (13)$$




Fig. 1: State diagram generated using LatexTikZ