



STATISTICS AND DATA ANALYSIS, EXAM-PLÉ

Department of statistics

Edgar Bueno

2024-11-15

Approved aid: Hand-held calculator with no stored text, data or formulas

Provided aid: Formula Sheet and Probability Distribution Tables, returned after the exam.

Problems 1 — 12: Multiple choice questions (max 60 points):

- A total of 12 multiple choice questions with five alternative answers per question one of which is the correct answer. Mark your answers on the attached **answer form**.
- Marking more than one alternative will result in zero points for that question.
- Each correct answer is worth 5 points.
- Written solutions should not be submitted; only your answers on the answer form will be considered in the assessment and final grading.

Problems 13 — 14: Complete written solutions (max 40 points):

- Use only the provided answer sheets when submitting your solutions and answers.
- For full marks, clear, comprehensive and well-motivated solutions are required. Unclear and unexplained solutions will result in point deductions even if the final answer is correct.
- Check your calculations and solutions before submitting. Careless mistakes will result in unnecessary point deductions.

The maximum total number of points is $60 + 40 = 100$. At least 50 points are required to pass (grades A-E). The grading scale is as follows:

| Points | 0—39 | 40—49 | 50—59 | 60—69 | 70—79 | 80—89 | 90—100 |
|--------|------|-------|-------|-------|-------|-------|--------|
| Grade | F | Fx | E | D | C | B | A |

NOTE: Fx and F are failing grades that require re-examination. Students who receive the grade Fx or F cannot supplement extra assignments for a higher grade.

Part one. Multiple choice


1. A salesperson has classified each of her potential customers regarding how likely they are to buy her product. The categories are: high, medium and low. She wants to summarize the data in an adequate chart. Which of the following is a type of chart that is adequate for this situation?
 - (a) Bar chart;
 - (b) Dotplot;
 - (c) Histogram;
 - (d) Scatter plot;
 - (e) Box-and-whisker plot.
2. Which of the following charts describes the information of **only one** variable?
 - (a) Histogram;
 - (b) Stacked bar chart;
 - (c) Mosaic plot;
 - (d) Scatter plot;
 - (e) All of the above.

3. Table 1 summarizes the scores of 170 students in an exam of statistics:

| Points | [0 , 40) | [40 , 50) | [50 , 60) | [60 , 70) | [70 , 80) | [80 , 90) | [90 , 100) |
|-----------|----------|-----------|-----------|-----------|-----------|-----------|------------|
| Frequency | 51 | 17 | 22 | 34 | 21 | 17 | 8 |

Table 1: Scores of 170 students in an exam of statistics

Which of the following is a histogram that correctly represents the scores in Table 1?




4. A researcher in survey methodology is studying the effect of incentives on item nonresponse. To this end she has selected a sample of ten individuals, offered them different amounts of money and submitted them to a long questionnaire. Then she has measured how many questions they answer before they get tired and decide to stop. Table 2 shows the results.

| Incentive (in SEK) | 0 | 100 | 200 | 300 | 400 | 500 | 600 | 700 | 800 | 900 |
|--------------------|---|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Questions answered | 0 | 7 | 100 | 25 | 57 | 33 | 70 | 50 | 100 | 100 |

Table 2: Incentives offered to and number of questions answered by a sample of ten respondents

Which of the following is a scatter plot that adequately represents the measurements in Table 2?



All of
the above

5. Which of the following is **correct** as an interpretation of the mean, \bar{x} :
- (a) It is the most frequently occurring value;
 - (b) It is the middle observation;
 - (c) It is the most likely value;
 - (d) It is the center of gravity of the observations;
 - (e) None of the above.

6. A researcher has asked the thirteen married men in a small community about the brideprice they had to pay to the bride's family when they got married. The brideprice values (in USD) are

20000 3000 10000 20000 13000 0 31000 20000 63000 8000 3000 12000 4000

What is the **interquartile range** of the brideprice?

- (a) -12000;
 - (b) 11000;
 - (c) 12000;
 - (d) 16000;
 - (e) 63000.
7. Which of the following sentences is **correct** regarding the coefficient of determination R^2 :
- (a) In *simple* linear regression, if there is a perfect negative linear association between the independent variable x and the dependent variable y , R^2 is equal to -1 (minus one);
 - (b) In *simple* linear regression, R^2 is equal to the coefficient of correlation between the independent variable x and the dependent variable y , that is, $R^2 = r_{xy}$;
 - (c) In *multiple* linear regression, R^2 should always be preferred over the adjusted coefficient of determination \bar{R}^2 ;
 - (d) R^2 may decrease when more variables are added to a model;
 - (e) R^2 is equal to the square coefficient of correlation between the predictions \hat{y} and the dependent variable y , that is, $R^2 = r_{\hat{y}y}^2$.
8. In the context of simple linear regression, which of the following is **not** correct?
- (a) the least squares regression is the one that minimizes the *sum of squares error*;
 - (b) the intercept b_0 indicates the expected value of the dependent variable y when the independent variable x equals zero;
 - (c) the slope b_1 indicates the expected increment in the dependent variable y associated to a one unit increment in the independent variable x ;
 - (d) the coefficient of determination R^2 indicates the proportion of variability of the dependent variable y that is explained by the independent variable x ;
 - (e) the coefficient of determination R^2 is equal to the coefficient of correlation between the independent variable x and the dependent variable y .

9. A real estate agent has estimated the regression that explains the closing price (variable *price*, in SEK) of the housing units in a region of interest with respect to their size (variable *size*, in m^2). The fitted regression line is

$$\widehat{price} = 2\,500\,000 + 5000\,size.$$

Which of the following is **not correct**:

- (a) if housing unit A has one more square meter than housing unit B, we expect the closing price of A to be around 5000 SEK higher than the closing price of B;
 - (b) the closing price of a housing unit of size $100m^2$ is expected to be around 3 000 000;
 - (c) the mean closing price of the housing units in the region of interest is 2 500 000;
 - (d) the intercept of the fitted regression is 2 500 000;
 - (e) the slope of the fitted regression is 5000.
10. The teacher of a course in statistics wants to explain the score of students in the final exam (variable *exam*) in terms of the score in a previous home assignment (variable *assignment*) through a linear regression of the form:

$$exam = \beta_0 + \beta_1 assignment + \epsilon$$

The following table shows the scores of the eight students in the course:

| <i>Assignment</i> | 42 | 48 | 50 | 50 | 51 | 55 | 59 | 67 |
|-------------------|----|----|----|----|----|----|----|----|
| <i>Exam</i> | 38 | 43 | 57 | 33 | 81 | 50 | 48 | 84 |

The estimated intercept of the regression line of interest is:

- (a) -548.7;
 - (b) - 25.2;
 - (c) 0.6;
 - (d) 1.5;
 - (e) 11.4.
11. Fitting a regression that explains the score of students in the final exam of a course in statistics (variable *exam*) in terms of the score in a previous home assignment (variable *assignment*), yields an intercept $b_0 = -25.2$ and a slope $b_1 = 1.5$. The following table shows the scores of the eight students in the course:

| <i>Assignment</i> | 42 | 48 | 50 | 50 | 51 | 55 | 59 | 67 |
|-------------------|----|----|----|----|----|----|----|----|
| <i>Exam</i> | 38 | 43 | 57 | 33 | 81 | 50 | 48 | 84 |

The *sum of squares error* —SSE— is:

- (a) 0;
- (b) 656;
- (c) 1594;
- (d) 1881;
- (e) 2508;

12. Fitting a regression that explains the score of students in the final exam of a course in statistics in terms of the score in a previous home assignment, yields an intercept $b_0 = -25.2$ and a slope $b_1 = 1.5$. The predicted score in the final exam for a student with 50 points in the home assignment is:
- (a) -1258.5;
 - (b) -1185.0;
 - (c) 12.2;
 - (d) 49.8;
 - (e) 75.0.

Part two. Complete solution

13. On September 2024 a company that offers audio streaming services released a new logo. A random sample of six users was drawn and their time using the service before and after the release was measured. Let x_i = “time (in minutes) spent by the i th user using the service one week *before* the new logo was released” and y_i = “time (in minutes) spent by the i th user using the service one week *after* the new logo was released” ($i = 1, \dots, 6$). The measurements are shown in the table below:

| User | 1 | 2 | 3 | 4 | 5 | 6 |
|------|----|----|-----|-----|-----|-----|
| x | 40 | 53 | 123 | 139 | 205 | 243 |
| y | 64 | 92 | 124 | 90 | 171 | 212 |

- (a) Calculate the following parameters: **i.** the mean of x , \bar{x} ; **ii.** the median of x , \check{x} ; **iii.** the standard deviation of x , S_x ; and **iv.** the interquartile range of x , IQR_x . (10p.)
- (b) It is known that the mean and variance of y are, respectively, $\bar{y} = 125.5$ and $S_y^2 = 3144$. It is also known that the correlation between x and y is $r_{xy} = 0.93$. Using this and your results from (a) find the intercept and the slope of a regression that explains the time using the service *after* the release of the new logo in terms of the time using the service *before* the release of the new logo. (**Note:** If you did not solve part (a) use $\bar{x} = 125.5$ and $S_x^2 = 3144$) (5p.)
- (c) Using the fitted regression in (b), predict the time (in minutes) that a person using the service 100 minutes *before* the release of the new logo will spend using the service *after* the release of the new logo. (5p.)
- (d) Using the fitted regression in (b), find the six residuals e_i . (10p.)
- (e) Find the coefficient of determination of the fitted regression. (10p.)