

Introduction to Machine Learning

Lecture 6: Stochastic Gradient Descent

Hao Wang

Email: wanghao1@shanghaitech.edu.cn

School of Information Science and Technology
ShanghaiTech University, Shanghai

March 9, 2023

Outline

Stochastic gradient descent (SGD)

Convergence Analysis

Noise Reduction

Stochastic gradient descent (SGD)

- ▶ Empirical loss:

$$J(w) = \frac{1}{2n} \sum_j^n J_j(w)$$

e.g. MSE: $J_j(w) = (\mathbf{x}^j{}^T w - y^j)^2$

- ▶ Batch gradient of empirical loss:

$$\nabla J(w) = \frac{1}{n} \sum_j^n \nabla J_j(w)$$

e.g. $\nabla J_j(w) = (\mathbf{x}^j{}^T w - y^j) \cdot \mathbf{x}^j$

- ▶ Stochastic (or “online”) gradient descent:

$$w^{k+1} \leftarrow w^k - \alpha^k \nabla J_j(w^k)$$

- ▶ Use updates based on individual datum j , chosen at random


Stochastic gradient descent

- ▶ Batch GD is a monotone (for what?) algorithm (why?).
- ▶ SGD is not a monotone algorithm (why?).


Definition



Each set of n consecutive accesses is called an **epoch**.



- ▶ The batch method performs only one step per epoch.
- ▶ SG performs n steps per epoch.







SGD and LBFGS


























Stochastic vs deterministic




Fixed learning rate

$$J(\theta) = \frac{1}{m} \sum_{i=1}^m J_i(\theta)$$




Fixed learning rate

$$J(\theta) = \frac{1}{m} \sum_{i=1}^m J_i(\theta)$$




Fixed learning rate

$$J(\theta) = \frac{1}{m} \sum_{i=1}^m J_i(\theta)$$




Fixed learning rate

$$J(\theta) = \frac{1}{m} \sum_{i=1}^m J_i(\theta)$$




Fixed learning rate

$$J(\theta) = \frac{1}{m} \sum_{i=1}^m J_i(\theta)$$




Diminishing learning rate

$$J(\theta) = \frac{1}{m} \sum_{i=1}^m J_i(\theta)$$




Diminishing learning rate

$$J(\theta) = \frac{1}{m} \sum_{i=1}^m J_i(\theta)$$




Diminishing learning rate

$$J(\theta) = \frac{1}{m} \sum_{i=1}^m J_i(\theta)$$



Diminishing learning rate

$$J(\theta) = \frac{1}{m} \sum_{i=1}^m J_i(\theta)$$



Is this what we want?

Mini-Batch stochastic gradient

- ▶ Empirical loss:

$$J(w) = \frac{1}{2n} \sum_j^n J_j(w)$$

- ▶ Batch gradient of empirical loss:

$$\nabla J(w) = \frac{1}{n} \sum_j^n \nabla J_j(w)$$

- ▶ Stochastic (or “online”) gradient descent: $\mathcal{S}_k \subset \{1, \dots, n\}$

- $w^{k+1} \leftarrow w^k - \alpha^k \frac{1}{|\mathcal{S}_k|} \sum_{j \in \mathcal{S}_k} \nabla J_j(w^k)$

- $|\mathcal{S}_k|$ may also vary

Outline

Stochastic gradient descent (SGD)

Convergence Analysis

Noise Reduction

Risk minimization

Minimizing the loss:

$$\min_w F(w) = \begin{cases} R(w) = \mathbb{E}[f(w; \xi)] & \text{expected risk} \\ \text{or} \\ R_n(w) = \frac{1}{n} \sum_{i=1}^n f_i(w) & \text{empirical risk} \end{cases}$$

For empirical risk: (每个样本都是随机变量的一次采样)

$$R_n(w) = \frac{1}{n} \sum_{i=1}^n f_i(w) = \frac{1}{n} \sum_{i=1}^n f(w; \xi_i)$$
$$f_i(w) = f(w; \xi_i)$$

Stochastic Gradient

The stochastic gradient is then defined as $g(w_k, \xi_k)$:

$$g(w_k, \xi_k) = \begin{cases} \nabla f(w_k; \xi_k), & \text{or} \\ \frac{1}{n_k} \sum_{i=1}^{n_k} \nabla f(w_k; \xi_{k,i}) \end{cases}$$

- ▶ ξ_k is a seed for generating a stochastic direction; e.g., a realization of it may represent the choice of a **single** training sample as in the simple SG method, or may represent a **set of samples** as in the minibatch SG method.
- ▶ $g(w_k, \xi_k)$ could represent a stochastic gradient—i.e., an **unbiased** estimator of $\nabla F(w_k)$

Algorithm

Algorithm 2.1 Stochastic Gradient (SG) Method

- 1: Choose an initial iterate w_1 .
 - 2: **for** $k = 1, 2, \dots$ **do**
 - 3: Generate a realization of the random variable ξ_k
 - 4: Compute a stochastic vector $g(w_k, \xi_k)$
 - 5: Choose a stepwise $\alpha_k > 0$
 - 6: Set the new iterate as $w_{k+1} \leftarrow w_k - \alpha_k g(w_k, \xi_k)$
 - 7: **end for**
-

Convergence

Assumption

(Lipschitz-continuous objective gradients). The objective function $F: \mathbb{R}^d \rightarrow \mathbb{R}$ is continuously differentiable and the gradient function of F , namely, $\nabla F: \mathbb{R}^d \rightarrow \mathbb{R}^d$, is Lipschitz continuous with Lipschitz constant $L > 0$, i.e.,

$$\|\nabla F(w) - \nabla F(\bar{w})\|_2 \leq L\|w - \bar{w}\|_2 \quad \text{for all } \{w, \bar{w}\} \subset \mathbb{R}^d.$$

This means

$$F(w) \leq F(\bar{w}) + \nabla F(\bar{w})^T(w - \bar{w}) + \frac{1}{2}L\|w - \bar{w}\|_2^2 \quad \text{for all } \{w, \bar{w}\} \subset \mathbb{R}^d.$$

Convergence

Lemma

The iterates of SG, satisfy the following inequality for all $k \in \mathbb{N}$:

$$\mathbb{E}_{\xi_k}[F(w_{k+1})] - F(w_k) \leq -\alpha_k \nabla F(w_k)^T \mathbb{E}_{\xi_k}[g(w_k, \xi_k)] + \frac{1}{2} \alpha_k^2 L \mathbb{E}_{\xi_k}[\|g(w_k, \xi_k)\|_2^2].$$

Therefore, if $g(w_k, \xi_k)$ is an unbiased estimate of $\nabla F(w_k)$, then it follows from Lemma 4.2 that

$$\mathbb{E}_{\xi_k}[F(w_{k+1})] - F(w_k) \leq -\alpha_k \|\nabla F(w_k)\|_2^2 + \frac{1}{2} \alpha_k^2 L \mathbb{E}_{\xi_k}[\|g(w_k, \xi_k)\|_2^2]$$

In order to limit the second-order term of α , we need to restrict the variance of $g(w_k, \xi_k)$, i.e.,

$$\mathbb{V}_{\xi_k}[g(w_k, \xi_k)] := \mathbb{E}_{\xi_k}[\|g(w_k, \xi_k)\|_2^2] - \|\mathbb{E}_{\xi_k}[g(w_k, \xi_k)]\|_2^2$$

Convergence

Assumption

(first and second moment limits). The objective function and SG satisfy the following conditions:

1. The sequence of iterates $\{w_k\}$ is contained in an open set over which F is bounded below by a scalar F_{\inf} .
2. There exists scalar $\mu_G \geq \mu > 0$ such that, for all $k \in \mathbb{N}$,

$$\begin{aligned}\nabla F(w_k)^T \mathbb{E}_{\xi_k}[g(w_k; \xi_k)] &\geq \mu \|\nabla F(w_k)\|_2^2 \quad \text{and} \\ \|\mathbb{E}_{\xi_k}[g(w_k, \xi_k)]\|_2 &\leq \mu_G \|\nabla F(w_k)\|_2\end{aligned}$$

3. There exist scalars $M \geq 0$ and $M_V \geq 0$ such that, for all $k \in \mathbb{N}$

$$\mathbb{V}_{\xi_k}[g(w_k, \xi_k)] \leq M + M_V \|\nabla F(w_k)\|_2^2$$

Convergence

From the above assumption, we have that

$$\mathbb{E}_{\xi_k}[\|g(w_k, \xi_k)\|_2^2] \leq M + M_G \|\nabla F(w_k)\|_2^2 \quad \text{with} \quad M_G := M_V + \mu_G^2 \geq \mu^2 > 0.$$

Lemma

The iterates of SG satisfy the following inequalities for all $k \in \mathbb{N}$:

$$\begin{aligned} \mathbb{E}_{\xi_k}[F(w_{k+1})] - F(w_k) &\leq -\mu\alpha_k \|\nabla F(w_k)\|_2^2 + \frac{1}{2}\alpha_k^2 L \mathbb{E}_{\xi_k}[\|g(w_k, \xi_k)\|_2^2] \\ &\leq -(\mu - \frac{1}{2}\alpha_k L M_G)\alpha_k \|\nabla F(w_k)\|_2^2 + \frac{1}{2}\alpha_k^2 L M. \end{aligned}$$

Convergence

Assumption

(strong convexity). The objective function $F: \mathbb{R}^d \rightarrow \mathbb{R}$ is strongly convex in that there exists a constant $c > 0$ such that

$$F(\bar{w}) \geq F(w) + \nabla F(w)^T(\bar{w} - w) + \frac{1}{2}c\|\bar{w} - w\|_2^2 \quad \forall (\bar{w}, w) \in \mathbb{R}^d \times \mathbb{R}^d.$$

Hence, F has a unique minimizer, denoted as $w_* \in \mathbb{R}^d$ with $F_* := F(w_*)$.

$$\implies F(w) - F(w_*) \leq \frac{1}{2c}\|\nabla F(w)\|_2^2 \quad \forall w \in \mathbb{R}^d$$

Since w_k is determined by the realization of the independent random variables $\{\xi_1, \xi_2, \dots, \xi_{k-1}\}$, the *total expectation* of $F(w_k)$ for any k can be taken as

$$\mathbb{E}[F(w_k)] = \mathbb{E}_{\xi_1} \mathbb{E}_{\xi_2} \dots \mathbb{E}_{\xi_{k-1}} [F(w_k)]$$

Convergence (strongly convex objective, fixed stepsize)

Theorem


Suppose that the SG method is run with a fixed stepsize, $\alpha_k = \bar{\alpha}$, satisfying

$$0 < \bar{\alpha} \leq \frac{\mu}{LM_G}.$$

Then the expected optimality gap satisfies the following inequality for all k

$$\begin{aligned} \mathbb{E}[F(w_k) - F_*] &\leq \frac{\bar{\alpha}LM}{2c\mu} + (1 - \bar{\alpha}\mu)^{k-1} \left(F(w_1) - F_* - \frac{\bar{\alpha}LM}{2c\mu} \right) \\ &\xrightarrow{k \rightarrow \infty} \frac{\bar{\alpha}LM}{2c\mu} \end{aligned}$$

Convergence (fixed learning rate)



Convergence (strongly convex objective, diminishing stepsizes)

Theorem

Suppose that the SG method is run with a fixed stepsize,

$$\alpha_k = \frac{\beta}{\gamma + k} \text{ for some } \beta > \frac{1}{c\mu} \text{ and } \gamma > 0 \text{ such that } \alpha_1 \leq \frac{\mu}{LM_G}.$$

Then, for all $k \in \mathbb{N}$, the expected optimality gap satisfies

$$\mathbb{E}[F(w_k) - F_*] \leq \frac{\nu}{\gamma + k}$$

where

$$\nu := \max\left\{\frac{\beta^2 LM}{2(\beta c\mu - 1)}, (\gamma + 1)(F(w_1) - F_*)\right\}$$

Convergence (nonconvex objective, fixed stepsize)

Now suppose F is not necessarily convex.

Theorem

Suppose that SG is run with a fixed stepsize $\alpha_k = \bar{\alpha}$ for all k , satisfying

$$0 < \bar{\alpha} \leq \frac{\mu}{LM_G}.$$

Then, the expected sum of squares and average-squared gradient of F corresponding to the SG iterates satisfy the following inequalities for all $K \in \mathbb{N}$:

$$\mathbb{E} \left[\sum_{k=1}^K \|\nabla F(w_k)\|_2^2 \right] \leq \frac{K\bar{\alpha}LM}{\mu} + \frac{2(F(w_1) - F_{\inf})}{\mu\bar{\alpha}},$$

so that

$$\frac{1}{K} \mathbb{E} \left[\sum_{k=1}^K \|\nabla F(w_k)\|_2^2 \right] \leq \frac{K\bar{\alpha}LM}{\mu} + \frac{2(F(w_1) - F_{\inf})}{K\mu\bar{\alpha}} \xrightarrow{K \rightarrow \infty} \frac{\bar{\alpha}LM}{\mu}.$$

Convergence (nonconvex objective, diminishing stepsize)

Theorem

Suppose that the SG method is run with a stepsize sequence satisfying. Then

$$\sum_{k=1}^{\infty} \alpha_k = \infty \quad \text{and} \quad \sum_{k=1}^{\infty} \alpha_k^2 < \infty.$$

More precisely, let $A_K = \sum_{k=1}^K \alpha_k$, then

$$\mathbb{E} \left[\sum_{k=1}^K \alpha_k \|\nabla F(w_k)\|_2^2 \right] < \infty,$$

so that

$$\mathbb{E} \left[\frac{1}{A_K} \sum_{k=1}^K \|\nabla F(w_k)\|_2^2 \right] \xrightarrow{K \rightarrow \infty} 0.$$

Convergence (nonconvex objective, diminishing stepsize)

Corollary

For any K , let $k(K) \in \{1, \dots, K\}$ represents a random index chosen with probabilities proportional to $\{\alpha_k\}_{k=1}^K$. Then, $\|\nabla F(w_{k(K)})\|_2 \xrightarrow{K \rightarrow \infty} 0$ in probability.


Corollary

If F is twice differentiable, and that the mapping $w \rightarrow \|\nabla F(w)\|_2^2$ has Lipschitz-continuous derivatives, then

$$\lim_{k \rightarrow \infty} \mathbb{E}[\|\nabla F(w_k)\|_2^2] = 0.$$

Batch or Stochastic? Early termination


Empirical loss contour



Expected loss contour

Early termination

Empirical loss contour



Expected loss contour

Batch or Stochastic? Work complexity for large-scale learning

- ▶ In a **big data** scenario, let's compare GD and SGD.
- ▶ Suppose that both the expected risk R and the empirical risk R_n attain their minima with parameter vectors

$$w_* \in \arg \min R(w) \quad \text{and} \quad w_n \in \arg \min R_n(w).$$

- ▶ Let \tilde{w}_n be the approximate empirical risk minimizer returned by a given optimization algorithm when the time budget \mathcal{T}_{\max} is exhausted.
- ▶ w_n is the minimizer of $R_n(w)$
- ▶ Let $\epsilon := \mathbb{E}[R_n(\tilde{w}_n) - R_n(w_n)]$ you end up with your optimization tool, within time \mathcal{T}_{\max} .

Work complexity for large-scale learning

- The total error

$$\mathbb{E}[R(\tilde{w}_n)] = \underbrace{R(w_*)}_{\mathcal{E}_{app}(\mathcal{H})} + \underbrace{\mathbb{E}[R(w_n) - R(w_*)]}_{\mathcal{E}_{est}(\mathcal{H}, n)} + \underbrace{\mathbb{E}[R(\tilde{w}_n) - R(w_n)]}_{\mathcal{E}_{opt}(\mathcal{H}, n, \epsilon)}.$$

- The “quality” of your learning

$$\min_{n, \epsilon} \mathcal{E}(n, \epsilon) = \mathbb{E}[R(\tilde{w}_n) - R(w_*)] \text{ s.t. } \mathcal{T}(n, \epsilon) \leq \mathcal{T}_{\max}.$$

- For the error function, a direct application of the uniform laws of large numbers yields:

$$\begin{aligned} \mathcal{E}(n, \epsilon) = \mathbb{E}[R(\tilde{w}_n) - R(w_*)] &= \underbrace{\mathbb{E}[R(\tilde{w}_n) - R_n(\tilde{w}_n)]}_{= \mathcal{O}\left(\sqrt{\log(n)/n}\right)} + \underbrace{\mathbb{E}[R_n(\tilde{w}_n) - R_n(w_n)]}_{= \epsilon} \\ &\quad + \underbrace{\mathbb{E}[R_n(w_n) - R_n(w_*)]}_{\leq 0} + \underbrace{\mathbb{E}[R_n(w_*) - R(w_*)]}_{= \mathcal{O}\left(\sqrt{\log(n)/n}\right)}, \end{aligned}$$

Work complexity for large-scale learning

- ▶ We have the upper bound

$$\mathcal{E}(n, \epsilon) = \mathcal{O} \left(\sqrt{\frac{\log(n)}{n}} + \epsilon \right).$$

- ▶ For cases where loss function is strongly convex, or the data distribution satisfies certain assumptions, it is possible to show that

$$\mathcal{E}(n, \epsilon) = \mathcal{O} \left(\frac{\log(n)}{n} + \epsilon \right).$$

- ▶ To simplify further, let us work with the asymptotic equivalence (for large n , big data)

$$\mathcal{E}(n, \epsilon) \sim \frac{1}{n} + \epsilon$$

Work complexity for large-scale learning

$$\mathcal{E}(n, \epsilon) \sim \frac{1}{n} + \epsilon$$

- For SGD, achieve ϵ -optimality with a computing time of $\mathcal{T}_{stoch} \sim 1/\epsilon$.
- Within the time budget \mathcal{T}_{\max} , the accuracy achieved is proportional to $1/\mathcal{T}_{\max}$, **regardless of n** .
- To minimize the error $\mathcal{E}(n, \epsilon)$, simply choose n as large as possible.
- Since the max number of examples that can be processed by SG is proportional to \mathcal{T}_{\max} , so the optimal error is proportional $1/\mathcal{T}_{\max}$
- For GD, achieve ϵ -optimality with a computing time of $\mathcal{T}_{batch} \sim n \log(1/\epsilon)$.
- Within the time budget \mathcal{T}_{\max} , to achieve ϵ -accuracy, need to process $n \sim \mathcal{T}_{\max} / \log(1/\epsilon)$ examples.
- Optimal error is not necessarily achieved by choosing n as large as possible. But rather by choosing ϵ to minimize the $\mathcal{E}(n, \epsilon) = \log(1/\epsilon) / \mathcal{T}_{\max} + \epsilon$.
- Optimal $\epsilon \sim 1/\mathcal{T}_{\max}$, so that optimal error is

$$\log(\mathcal{T}_{\max}) / \mathcal{T}_{\max} + 1/\mathcal{T}_{\max}$$

Batch or Stochastic?

	Batch	Stochastic
$\mathcal{T}(n, \epsilon)$	$\sim n \log \left(\frac{1}{\epsilon} \right)$	$\frac{1}{\epsilon}$
\mathcal{E}^*	$\sim \frac{\log(\mathcal{T}_{\max})}{\mathcal{T}_{\max}} + \frac{1}{\mathcal{T}_{\max}}$	$\frac{1}{\mathcal{T}_{\max}}$

Comments

- ▶ Fragility of the asymptotic performance of SG.
- ▶ SG and ill-conditioning.
- ▶ Opportunities for distributed computing.
- ▶ Alternatives with faster convergence.

Outline

Stochastic gradient descent (SGD)

Convergence Analysis

Noise Reduction

Noise Reduction Methods I (optional)

What if choosing $\alpha^k = \alpha$, must reduce the noise in sampled gradient at a geometric rate

Dynamic Sample Size Methods

- $w^{k+1} \leftarrow w^k - \alpha \frac{1}{|\mathcal{S}_k|} \sum_{j \in \mathcal{S}_k} \nabla J_j(w^k)$
- $|\mathcal{S}_k| = \lceil \tau^{k-1} \rceil$ with $\tau > 1$.

Noise Reduction Methods II (optional)

Gradient Aggregation

► SVRG

$$\begin{aligned}\nabla J_j(\tilde{w}^k) &\leftarrow \nabla J_j(\tilde{w}^k) - [\nabla J_j(w^k) - \nabla J(w^k)] \\ \tilde{w}^{k+1} &\leftarrow \tilde{w}^k - \alpha \nabla J_j(\tilde{w}^k)\end{aligned}$$

► SAGA

t chosen randomly $\in \{k - n, k - n + 1, \dots, k\}$

$$\begin{aligned}\nabla J_j(w^k) &\leftarrow \nabla J_j(w^k) - \nabla J_j(w^{[t]}) + \frac{1}{n} \sum_{i=1}^n \nabla J_j(w^{[i]}) \\ w^{k+1} &\leftarrow w^k - \alpha \nabla J_j(w^k)\end{aligned}$$

Noise Reduction Methods III (optional)

Iterate Averaging Methods

$$w^{k+1} \leftarrow w^k - \alpha^k \nabla J_j(w^k)$$

$$\tilde{w}^{k+1} \leftarrow \frac{1}{k+1} \sum_{i=1}^{k+1} w^i$$

- $\alpha^k \sim O(1/k)$ or slower
- \tilde{w}^k is *not* used for iterate update

Learning Algorithms

