arXiv:2611.05334v1 [math.OC] 3 Nov 2026

A simple adaptive accelerated optimistic dual averaging method for unconstrained convex optimization in p-norm geometry

Bowen Li* Danqing Zhou* Chuan He*

Abstract

Efficient preconditioning plays an important role in large-scale first-order optimization. In this paper, we propose an adaptive accelerated optimistic dual averaging (A^2ODA) method for unconstrained convex optimization that adopts low-cost preconditioning based on p-norm geometry for $p \in [2, \infty]$. The method is line-search-free and employs a new stepsize selection technique based on an aggregated affine lower model. Its update structure is simple and unified across $p \in [2, \infty]$, with the case $p > 2$ requiring no more elaborate updates than the Euclidean case $p = 2$. Under a generalized local smoothness condition, we derive a convergence bound that separates the effects of constant and gap-dependent curvature, with the latter potentially yielding faster-decaying terms. In the special case of local smoothness, this bound recovers a nearly optimal convergence rate.

Keywords: Convex optimization; accelerated first-order methods; adaptive methods; p-norm geometry; optimistic dual averaging; generalized local smoothness.

Mathematics Subject Classification: 90C25; 90C30; 90C06; 68Q25.

1 Introduction

In this paper, we consider the unconstrained convex optimization problem

$$\min_{x \in \mathbb{R}^n} f(x), \tag{1}$$

where $f : \mathbb{R}^n \to \mathbb{R}$ is a continuously differentiable convex function. We assume that problem (1) has at least one optimal solution. Since the seminal work on accelerated gradient methods [27], the development of fast first-order methods has remained a central focus in the study of smooth convex optimization. In practice, the performance of first-order methods is influenced by many factors, including the acceleration mechanism, the parameter selection, and the underlying geometry. The goal of this paper is to develop an accelerated first-order method in p-norm geometry that adapts its parameters to the optimization trajectory and achieves a unified convergence bound for all $p \in [2, \infty]$.

In recent years, there has been fruitful research on first-order methods for smooth convex optimization [1, 28, 29]. In particular, many recent efforts have focused on parameter selection in first-order methods, which is often guided by estimates of the Lipschitz constant of the objective gradient. Instead of fixing the parameters based on a global Lipschitz constant, adapting them to local curvature typically leads to much faster practical progress. In this direction, there has recently been growing interest in first-order methods that adapt their parameters based on local curvature without requiring costly line searches. The use of local curvature to select stepsizes has a long history, as exemplified by the well-known Barzilai-Borwein

*Department of Mathematics, Linköping University, Sweden (Email: bowen.a.li@liu.se, danqing.zhou@liu.se, and chuan.he@liu.se). All authors contributed equally to this work.

(BB) stepsize [2]. However, directly applying the BB stepsize without line searches does not generally yield convergence guarantees for smooth convex problems. Recently, several studies have proposed novel stepsize schemes for line-search-free gradient methods and established provable convergence guarantees for solving problem (1); see, e.g., [17, 24, 36]. In addition, notable adaptive accelerated gradient methods, including AC-FGM [18] and AdaNAG [33], achieve optimal convergence rates for smooth convex optimization without requiring line searches.

Beyond parameter selection, the choice of geometry is another important aspect of first-order method design. For many constrained problems, replacing the Euclidean squared distance with a suitable Bregman divergence in first-order method design allows the resulting updates to reflect the intrinsic geometry of the feasible region. Well-known examples include the Kullback-Leibler divergence on the probability simplex [4] and the Itakura-Saito divergence on the positive orthant [3]. These examples can be unified under the general mirror descent framework [4, 26], whose update takes the form:

$$x_{k+1} \in \arg\min_{x \in X} \left\{ \langle \nabla f(x_k), x - x_k \rangle + \gamma_k D_h(x, x_k) \right\},$$

where $\gamma_k > 0$, X denotes the feasible region, and $D_h(x, y) := h(x) - h(y) - \langle \nabla h(y), x - y \rangle$ is the Bregman divergence generated by h. When $h(x) = \|x\|_2^2/2$, the update reduces to projected gradient descent. More generally, suitable choices of h allow mirror descent to exploit non-Euclidean geometry. Under the relative smoothness condition

$$D_f(x, y) \le L D_h(x, y) \qquad \forall x, y \in \Omega$$

for some $L > 0$, convergence bounds of mirror descent methods can be established in terms of D_h, where Ω is a suitable convex set containing the iterates and relevant comparison points [4, 20]. Mirror descent and relative smoothness extend gradient descent and Lipschitz smoothness in the Euclidean norm, respectively, to non-Euclidean settings. Nonetheless, extending accelerated gradient methods to non-Euclidean mirror descent is nontrivial. There have been many recent studies on acceleration in mirror descent, including both specific accelerated variants [12, 16, 25] and lower bounds on achievable convergence rates [11]. In addition, adaptive mirror descent methods have also been developed [30].

Another line of research explores first-order methods that directly exploit norm geometry to shape search directions. In particular, replacing the squared Euclidean distance with the squared distance induced by a general norm yields the dual preconditioned gradient update [15]:

$$x_{k+1} = x_k - \frac{1}{\gamma_k} J_*(\nabla f(x_k)) \in \arg\min_{x \in \mathbb{R}^n} \left\{ \langle \nabla f(x_k), x - x_k \rangle + \frac{\gamma_k}{2} \|x - x_k\|^2 \right\},$$

where $\gamma_k > 0$, $\|\cdot\|$ and $\|\cdot\|_*$ denote a general norm and its dual, respectively, and $J_*(g) \in \partial(\|\cdot\|_*^2/2)(g)$ for all $g \in \mathbb{R}^n$. For convex problems, this preconditioned method achieves an $\mathcal{O}(T^{-1})$ convergence rate in the objective gap under a Lipschitz gradient condition defined using the general norm and its dual, together with a level-boundedness assumption [15], where T denotes the number of iterations. This bound is analogous to that of classical gradient descent. Moreover, this preconditioned gradient descent has also been analyzed under a dual relative smoothness condition [23]. This condition serves as a dual counterpart to the primal relative smoothness condition used to analyze mirror descent [20]. In addition, accelerated variants of preconditioned gradient descent in general norm geometries have been developed in [6, 8]. In particular, when the primal and dual norms are the p- and q-norms, respectively, with $1/p + 1/q = 1$ and $2 \le p < \infty$, these methods achieve an $\mathcal{O}(T^{-(1+2/p)})$ convergence rate in the objective gap, matching the lower bounds up to logarithmic factors [7]. From a technical standpoint, these methods rely on the uniform convexity of the proximal regularizer to control linearization errors in their accelerated updates [6, 8]. For the case $p > 2$, the resulting pth-power proximal terms interact with the quadratic terms arising

from Lipschitz smoothness, making parameter selection and the convergence analysis more involved than in the case $p = 2$ (see Remark 1). It is also noteworthy that p-norm geometry has attracted significant recent interest in the design of deep learning optimizers, particularly in the endpoint case $p = \infty$, which leads to sign-based update directions; see, e.g., [5, 10, 13, 22, 31]. These theoretical advances and growing practical interest motivate the following natural question:

Can we design a line-search-free accelerated first-order method for smooth convex optimization that exploits p-norm geometry for preconditioning throughout the range $p \in [2, \infty]$?

This paper provides an affirmative answer to this question. Specifically, our proposed method employs an acceleration mechanism that combines optimistic dual averaging [32] in p-norm geometry with weighted iterate averaging [14]. To adapt the regularization parameters, we aggregate past linearizations into an affine model and use this model to derive a new parameter selection rule that directly controls a Lyapunov potential. This construction differs fundamentally from the parameter selection rules used by existing line-search-free accelerated gradient methods in the case $p = 2$ [18, 33].

Moreover, our analysis relies on the relaxed dual-potential expansion inequality in Lemma 1, which avoids the need for a uniform convexity inequality for the proximal regularizer. Consequently, our method directly uses the squared p-norm as the proximal regularizer for $p > 2$, in contrast to existing accelerated methods that rely on p-uniformly convex regularizers [6, 8] (see Remark 1). On the theoretical side, we establish an accelerated convergence rate for the proposed method under a generalized local smoothness condition (see Assumption 1(b)). This condition incorporates gap-dependent curvature into the local smoothness condition and allows us to distinguish the contributions of the constant and gap-dependent curvature terms to the convergence bound. Our main contributions are summarized below.

- We develop an adaptive accelerated optimistic dual averaging (A^2ODA) method under p-norm geometry for solving problem (1). To the best of our knowledge, this is the first line-search-free first-order method that combines adaptive acceleration with squared p-norm regularization in a unified framework covering all $p \in [2, \infty]$.
- Under the generalized local smoothness condition (Assumption 1(b)), we establish a unified convergence bound for Algorithm 1 across all $p \in [2, \infty]$ that explicitly separates the contributions of constant and gap-dependent curvature (see Theorem 2). In the special case of local smoothness, this bound recovers a nearly optimal convergence rate (see Section 3.4).

The rest of the paper is organized as follows. In Section 2, we introduce the notation and preliminary results. In Section 3, we present the proposed algorithm and state its convergence guarantees. The proof of our main convergence result is presented in Section 4. Additional lemmas and proofs are deferred to the appendix.

2 Notation and preliminaries

Throughout this paper, we use $\mathbb{R}^n$ and $\mathbb{R}_+$ to denote the n-dimensional Euclidean space and the set of nonnegative real numbers, respectively, and $\langle \cdot, \cdot \rangle$ to denote the standard inner product. We adopt the convention $0^0 = 1$ and $1/\infty = 0$. For a positive integer N, we define $[N] := \{1, 2, \dots, N\}$. For any $p \in [1, \infty)$ and $z \in \mathbb{R}^n$, we define $\|z\|_p := (\sum_{i=1}^n |z_i|^p)^{1/p}$ and $\|z\|_\infty := \max_{1 \le i \le n} |z_i|$. For any $A \in \mathbb{R}^{n \times n}$ and any $p, q \in [1, \infty]$, the induced p-to-q operator norm is defined by $\|A\|_{p \to q} := \sup_{\|x\|_p \le 1} \|Ax\|_q$. For any convex function $h : \mathbb{R}^n \to \mathbb{R}$, its Fenchel conjugate is defined by $h^*(u) := \sup_{x \in \mathbb{R}^n} \{\langle x, u \rangle - h(x)\}$ for

every $u \in \mathbb{R}^n$. Let $\partial h(x)$ denote the convex subdifferential of h at x. The Fenchel-Young equality holds if and only if $u \in \partial h(x)$:

$$h(x) + h^*(u) = \langle x, u \rangle. \tag{2}$$

For $x, y \in \mathbb{R}^n$ and $g_y \in \partial h(y)$, define the generalized Bregman divergence by $D_h^{g_y}(x, y) := h(x) - h(y) - \langle g_y, x - y \rangle$. When h is differentiable at y, we omit the superscript and denote the Bregman divergence by $D_h(x, y)$. In addition, we use $\mathcal{O}(\cdot)$ and $\Theta(\cdot)$ to denote asymptotic upper and two-sided bounds, respectively.

For conjugate exponents $p \in [2, \infty]$ and $q \in [1, 2]$ satisfying $1/p + 1/q = 1$, define the primal and dual regularizers by

$$d_p(x) := \frac{1}{2}\|x\|_p^2, \qquad V_q(u) := d_p^*(u) = \frac{1}{2}\|u\|_q^2 \qquad \forall x, u \in \mathbb{R}^n. \tag{3}$$

For any $q \in [1, 2]$ and $u \in \mathbb{R}^n$, define

$$J_q(u) := \begin{cases} \nabla V_q(u), & q \in (1, 2], \\ \|u\|_1 \operatorname{sgn}_0(u), & q = 1, \end{cases} \qquad (\operatorname{sgn}_0(u))_i := \begin{cases} 1, & u_i > 0, \\ 0, & u_i = 0, \\ -1, & u_i < 0, \end{cases} \quad i = 1, \ldots, n. \tag{4}$$

Then, for any pair of conjugate exponents $p \in [2, \infty]$ and $q \in [1, 2]$, one can verify that

$$J_q(u) \in \partial V_q(u), \qquad \|J_q(u)\|_p = \|u\|_q, \qquad \langle J_q(u), u \rangle = \|u\|_q^2. \tag{5}$$

For any $q \in [1, 2]$, we also introduce an exponent that plays an important role in our analysis:

$$r_q := \begin{cases} \frac{q}{2-q}, & q \in [1, 2), \\ \infty, & q = 2. \end{cases} \tag{6}$$

The next lemma establishes an expansion inequality for the squared ℓ_q-norm, including the nonsmooth case $q = 1$. It provides a counterpart of the well-known Euclidean identity

$$V_2(u + v) = V_2(u) + \langle J_2(u), v \rangle + V_2(v). \tag{7}$$

Lemma 1 (Relaxed dual-potential expansion)**.** *Let $q \in [1, 2)$, and let V_q and J_q be defined in (3) and (4), respectively. Then,*

$$c\, V_q(u + v) \le V_q(u) + \langle J_q(u), v \rangle + \left(1 + \frac{2^{2/r_q} c}{(1 - c^{r_q})^{1/r_q}}\right) V_q(v) \tag{8}$$

holds for all $u, v \in \mathbb{R}^n, c \in (0, 1)$, where r_q is defined in (6).

Remark 1. *Existing accelerated methods in p-norm geometry typically rely on p-uniformly convex proximal regularizers for $p > 2$ [6, 8]. The resulting pth-power proximal terms cannot be directly balanced against the quadratic errors arising from Lipschitz smoothness, requiring additional approximation arguments and more involved parameter choices.*

In contrast, our method directly uses the squared p-norm as the proximal regularizer, without requiring a uniform convexity lower bound. The key technical ingredient in our analysis is the relaxed dual-potential expansion in Lemma 1, which controls perturbations through a quadratic term in the dual norm. Combined with our adaptive parameter-selection rule, this inequality provides an alternative approach to acceleration in p-norm geometry and simplifies the convergence analysis.

We make the following assumptions throughout the paper.

Assumption 1. *(a) The function f is convex and attains its minimum, with optimal value denoted by $f_* := \min_{x\in\mathbb{R}^n} f(x)$.*

(b) Let $p \in [2,\infty]$ and $q \in [1,2]$ be conjugate exponents with $1/p + 1/q = 1$, and let $\nu \in [0,1)$. For every compact convex set $\Omega \subset \mathbb{R}^n$, there exist constants $L_p(\Omega), H_p(\Omega) \geq 0$ such that

$$\|\nabla f(x) - \nabla f(y)\|_q^2 \leq 2\left(L_p(\Omega) + H_p(\Omega)\max\{\Delta(x),\Delta(y)\}^\nu\right) D_f(x,y) \tag{9}$$

holds for all $x, y \in \Omega$, where $\Delta(x) := f(x) - f_*$.

Remark 2. *Assumption 1(b) is a cocoercivity-type generalized local smoothness condition. This condition is related to local smoothness [21] and gap-dependent generalized smoothness [19]. Applying (9) to both (x,y) and (y,x), and then using Hölder's inequality, yields the following Lipschitz-type generalized local smoothness condition:*

$$\|\nabla f(x) - \nabla f(y)\|_q \leq (L_p(\Omega) + H_p(\Omega)\max\{\Delta(x),\Delta(y)\}^\nu)\|x-y\|_p \qquad \forall x, y \in \Omega. \tag{10}$$

Conversely, for any given Ω containing x_*, (9) holds with constants $L_p(\Omega), H_p(\Omega) \geq 0$ provided that

$$\|\nabla f(x) - \nabla f(y)\|_q \leq (L_p(\Omega) + 2^{-\nu}H_p(\Omega)\max\{\Delta(x),\Delta(y)\}^\nu)\|x-y\|_p \qquad \forall x, y \in \bar{\Omega},$$

where $\bar{\Omega}$ is a suitably enlarged compact convex set containing Ω. A precise choice of $\bar{\Omega}$ and a proof of this implication are provided in Appendix B. Thus, the cocoercivity-type and Lipschitz-type generalized local smoothness conditions are equivalent, up to an enlargement of the domain and a change in constants.

When $\nu = 0$, Assumption 1(b) reduces to the usual local smoothness condition. For $\nu > 0$, it separates the curvature bound into a constant part $L_p(\Omega)$ and a gap-dependent part $H_p(\Omega)\max\{\Delta(x),\Delta(y)\}^\nu$, which vanishes as the objective values approach the minimum. This decomposition allows the leading term of the convergence bound to depend on $L_p(\Omega)$, which can be substantially smaller than a uniform curvature bound over Ω, while the gap-dependent curvature contributes terms that may decay faster.

3 An adaptive accelerated optimistic dual averaging method

In this section, we propose an adaptive accelerated optimistic dual averaging (A^2ODA) method in p-norm geometry for $p \in [2,\infty]$ and establish its convergence properties under the generalized local smoothness condition stated in Assumption 1(b). Specifically, we first describe the main components of the method in Sections 3.1-3.3 and then present the complete algorithm and its convergence results in Section 3.4.

Before proceeding, we let $\{\lambda_k\}$ be the positive weights for averaging iterates and gradients, and define the cumulative weights by $A_k = \sum_{i=1}^k \lambda_i$ for all $k \geq 1$. The A^2ODA method generates two sequences: the optimistic dual-averaging iterates $\{y_k\}$ and the weighted-average iterates $\{x_k\}$. For convenience, we denote the sequences of function values and gradients evaluated at $\{x_k\}$ by $\{f_k\}$ and $\{g_k\}$, respectively, where $f_k = f(x_k)$ and $g_k = \nabla f(x_k)$ for all $k \geq 1$.

3.1 Optimistic dual averaging and acceleration

The proposed method combines optimistic dual averaging with weighted primal averaging to achieve acceleration. It mainly consists of three parts: dual averaging to accumulate past gradients, optimism to predict the next gradient contribution, and weighted primal averaging to construct the iterates at which gradients are evaluated.

Following the dual-averaging strategy [35], we maintain the weighted gradient sum

$$s_0 = 0, \qquad s_k = \sum_{i=1}^{k} \lambda_i g_i, \qquad \forall k \geq 1.$$

At the beginning of iteration k, the new gradient g_k is not yet available. We therefore use the most recent gradient g_{k-1} to predict the contribution of g_k to s_k, forming

$$\tilde{s}_k = s_{k-1} + \lambda_k g_{k-1} \qquad \forall k \geq 1.$$

The term $\lambda_k g_{k-1}$ constitutes the optimistic part: it adds a prediction of the next weighted gradient before that gradient is observed, as in optimistic follow-the-regularized-leader methods [32].

The optimistic dual-averaging iterate y_k is then obtained by minimizing the sum of the predicted linear model and a regularization term:

$$y_k \in \arg\min_{y \in \mathbb{R}^n} \left\{ \langle \tilde{s}_k, y - x_0 \rangle + \frac{\gamma_k}{2} \|y - x_0\|_p^2 \right\},$$

where $\gamma_k > 0$ controls the regularization strength. Our novel adaptive strategy for updating γ_k will be introduced later.

To achieve acceleration, a key ingredient is the coupling of optimistic updates with gradient evaluations at running weighted averages [14]. Specifically, we form the weighted primal average

$$x_k = \frac{A_{k-1}}{A_k} x_{k-1} + \frac{\lambda_k}{A_k} y_k = \frac{1}{A_k} \sum_{i=1}^{k} \lambda_i y_i,$$

and evaluate the new gradient at the averaged iterate x_k, rather than at y_k, setting $g_k = \nabla f(x_k)$.

3.2 Aggregated affine lower model for adaptive regularization

We now motivate our adaptive strategy for updating the regularization parameters based on an accumulated affine lower model. Following the estimate-sequence and approximate-duality-gap techniques [9, 34], we aggregate supporting affine functions and add a regularization term.

For any $x \in \mathbb{R}^n$, we define the aggregated affine lower model φ_k by

$$\varphi_k(x) := c_k + \langle s_k, x - x_0 \rangle = \sum_{i=1}^{k} \lambda_i (f_i + \langle g_i, x - x_i \rangle) \leq A_k f(x) \tag{11}$$

$$\text{with} \quad c_k := \sum_{i=1}^{k} \lambda_i (f_i + \langle g_i, x_0 - x_i \rangle),$$

where the inequality is due to the convexity of f. This model collects the first-order information obtained along the optimization trajectory. It requires only the vector s_k and the scalar c_k, both of which can be updated recursively.

Directly minimizing φ_k over $\mathbb{R}^n$ does not generally yield a useful comparison value. We therefore define the regularized estimating function for any $\gamma > 0$ and its minimum value by

$$\psi_k(x; \gamma) := \varphi_k(x) + \frac{\gamma}{2} \|x - x_0\|_p^2, \qquad m_k(\gamma) := \min_{x \in \mathbb{R}^n} \psi_k(x; \gamma) = c_k - \frac{\|s_k\|_q^2}{2\gamma}, \tag{12}$$

where the last equality follows from Fenchel conjugacy. For any minimizer x_* of f, (11) then implies

$$m_k(\gamma) \le \psi_k(x_*;\gamma) \le A_k f_* + \frac{\gamma}{2}\|x_* - x_0\|_p^2. \tag{13}$$

Although $m_k(\gamma)$ is not necessarily a lower bound on $A_k f_*$, this inequality provides the comparison needed to control the objective gap.

Motivated by (13), we seek a regularization parameter satisfying

$$A_k f_k \le m_k(\gamma) + \rho_k^2 \gamma, \tag{14}$$

where $\rho_k > 0$ is a scale to be specified later. The additional term $\rho_k^2\gamma$ is chosen to scale linearly with γ, matching the regularization term in (13). The condition (14) is computable without knowing f_*. Combining it with (13) yields

$$f_k - f_* \le \frac{\gamma}{A_k}\Big(\rho_k^2 + \frac{1}{2}\|x_* - x_0\|_p^2\Big). \tag{15}$$

For a fixed ρ_k, the resulting bound is proportional to γ.

Given $\gamma_k > 0$ and a safeguard parameter $\theta \in (0,1)$, we select the smallest $\gamma \ge \theta\gamma_k$ satisfying (14):

$$\gamma_{k+1} := \min\left\{\gamma \ge \theta\gamma_k : A_k f_k \le m_k(\gamma) + \rho_k^2\gamma\right\}. \tag{16}$$

The safeguard limits the decrease in the regularization parameter, while the estimating-function comparison (14) determines whether a value larger than $\theta\gamma_k$ is needed. Denote $\chi_k := A_k f_k - c_k$. Using the expression for $m_k(\gamma)$ in (12), we can rewrite the update (16) explicitly as

$$\gamma_{k+1} = \max\left\{\theta\gamma_k,\ \frac{\chi_k + \sqrt{\chi_k^2 + 2\rho_k^2\|s_k\|_q^2}}{2\rho_k^2}\right\}.$$

Since (14) holds with $\gamma = \gamma_{k+1}$, it follows from (15) that

$$f_k - f_* \le \frac{\gamma_{k+1}}{A_k}\Big(\rho_k^2 + \frac{1}{2}\|x_* - x_0\|_p^2\Big).$$

3.3 Bounded Lyapunov-potential certificate

We now use the estimating-function comparison in Section 3.2 to construct a natural potential sequence $\{\mathcal{P}_k\}$ for the convergence analysis. Specifically, for all $k \ge 1$, define

$$\mathcal{P}_k := A_k f_k - m_k(\gamma_{k+1}) = \chi_k + \frac{\|s_k\|_q^2}{2\gamma_{k+1}},$$

where the last equality follows from the definition of χ_k and (12). The connection between this potential and the objective gap follows from (13), which gives

$$\mathcal{P}_k \ge A_k(f_k - f_*) - \frac{\gamma_{k+1}}{2}\|x_* - x_0\|_p^2.$$

Thus, controlling $\mathcal{P}_k$ relative to γ_{k+1} provides a bound on the objective gap. In addition, the condition (14), with $\gamma = \gamma_{k+1}$, ensures that

$$\mathcal{P}_k = \chi_k + \frac{\|s_k\|_q^2}{2\gamma_{k+1}} \le \rho_k^2\gamma_{k+1}.$$

This bounded potential condition also provides a basis for establishing the boundedness of the iterates $\{x_k\}$ and $\{y_k\}$. A detailed statement will be given in Theorem 1.

3.4 The A^{2}ODA method and its convergence rate

We now summarize the A^{2}ODA method in Algorithm 1, which combines the updates described above.

Algorithm 1 A^{2}ODA: An **A**daptive **A**ccelerated **O**ptimistic **D**ual **A**veraging method

Input: Initial iterate $x_0 \in \mathbb{R}^n$ satisfying $\nabla f(x_0) \neq 0$, a conjugate pair (p, q) with $q \in [1, 2]$ and $1/p + 1/q = 1$, averaging scale $m \geq 2$, initial regularization parameter $\gamma_1 > 0$, and scaling parameters α, θ satisfying $\theta \in (0, 1)$ and $\alpha > 2(1 + e + e/\theta)$.
Set $f_0 = f(x_0)$, $g_0 = \nabla f(x_0)$, $s_0 = 0$, $c_0 = 0$, $A_0 = 0$, and $\rho_0 = 0$.
for $k = 1, 2, \ldots$ **do**

Compute the averaging weights:

$$\lambda_k = \left(1 + \frac{k-1}{m-1}\right)^{m-1}, \qquad A_k = A_{k-1} + \lambda_k. \tag{17}$$

Perform an optimistic dual averaging step:

$$\tilde{s}_k = s_{k-1} + \lambda_k\, g_{k-1}, \qquad y_k = x_0 + \frac{1}{\gamma_k}\, J_q(-\tilde{s}_k), \qquad x_k = \frac{A_{k-1}}{A_k}\, x_{k-1} + \frac{\lambda_k}{A_k}\, y_k. \tag{18}$$

Evaluate $f_k = f(x_k)$, $g_k = \nabla f(x_k)$, and update the safeguard radius:

$$\rho_k = \max\left\{\rho_{k-1}, \frac{\|y_k - x_0\|_p}{\alpha}\right\}. \tag{19}$$

Update the accumulated gradient and affine model:

$$s_k = s_{k-1} + \lambda_k\, g_k, \qquad c_k = c_{k-1} + \lambda_k\, (f_k + \langle g_k, x_0 - x_k\rangle), \qquad \chi_k = A_k\, f_k - c_k. \tag{20}$$

Update the regularization parameter:

$$\gamma_{k+1} = \max\left\{\theta\gamma_k, \frac{\chi_k + \sqrt{\chi_k^2 + 2(\rho_k \|s_k\|_q)^2}}{2\rho_k^2}\right\}. \tag{21}$$

end for

We now make a few remarks about Algorithm 1.

- The parameter $m \geq 2$ controls the growth of the averaging weights: for fixed m, $\lambda_k = \Theta(k^{m-1})$ and $A_k = \Theta(k^m)$. In particular, $m = 2$ gives $\lambda_k = k$ and $A_k = k(k+1)/2$, recovering the linearly increasing weights used in accelerated optimistic dual averaging [14]. Allowing $m > 2$ provides flexibility to exploit the diminishing gap-dependent curvature under generalized local smoothness, potentially yielding faster decay of the corresponding terms in the convergence bound.

  The benefit of increasing m depends on its interaction with the adaptive regularization. Indeed, when the sequence of radius scales $\{\rho_k\}$ is bounded, the objective-gap bound derived above is governed by γ_{k+1}/A_k. Thus, faster growth of A_k can translate into faster convergence only if it is not offset by the growth of γ_{k+1}. The convergence analysis below quantifies this interaction.

- The scale ρ_k controls the tolerance in the estimating-function comparison (see (14)) and therefore affects the update of the regularization parameter γ_{k+1}. We choose

$$\rho_k = \frac{1}{\alpha} \max_{1 \leq i \leq k} \{\|y_i - x_0\|_p\}$$

so that this scale adapts to the observed size of the optimization trajectory, without requiring prior knowledge of the distance from x_0 to a minimizer.

This choice balances two effects: a larger ρ_k relaxes the estimating-function comparison and may allow a smaller regularization parameter, but it also increases the radius-dependent term in the objective-gap bound. This adaptive update is designed to achieve a favorable dependence on problem constants while using only quantities already available from the iterates. A simpler alternative is to use a fixed scale $\rho_k \equiv \rho > 0$.

- The initial regularization parameter γ_1 can be selected through an optional backtracking line search, similar to that in [18]. We present its procedure as follows:

Calibration 1 (Selection of γ_1): Given $g_0 \neq 0$ and an initial $\gamma > 0$, define $x_1(\gamma) = x_0 + \gamma^{-1} J_q(-g_0)$. A trial γ is accepted if

$$f(x_1(\gamma)) \leq f(x_0) - \frac{\|g_0\|_q^2}{2\gamma}.$$

Test the initial γ and proceed as follows:

1. *Initial trial rejected.* Repeatedly set $\gamma \leftarrow 2\gamma$ and test the new γ until it is accepted.
2. *Initial trial accepted.* Repeatedly test $\gamma/2$. If accepted, set $\gamma \leftarrow \gamma/2$; otherwise, stop.

Return $\gamma_1 = \gamma$ and $x_1 = x_1(\gamma_1)$.

This procedure terminates after finitely many trials: differentiability of f at x_0 ensures acceptance for all sufficiently large γ, while $f \geq f_*$ and $g_0 \neq 0$ ensure rejection for all sufficiently small γ. Let L_p and H_p be the corresponding curvature moduli on a compact convex set $\mathcal{S}_p$ containing the iterates and a minimizer x_*, where $\mathcal{S}_p$ will be specified in Theorem 1 below. Under Assumption 1, the calibrated parameter γ_1 satisfies

$$\frac{\|g_0\|_q^2}{2\Delta(x_0)} \leq \gamma_1 < 2\big(L_p + H_p \Delta(x_0)^\nu\big). \tag{22}$$

A proof of this bound is provided in Appendix C. The calibration requires no prior knowledge of the smoothness constants and improves the dependence of the convergence rate on L_p and H_p, as detailed in Remark 5. Since the line search is performed only during initialization, subsequent iterations remain line-search-free.

We conclude this section by presenting the convergence rate of the A^{2}ODA method under the generalized local smoothness condition in Assumption 1(b). A formal statement is provided in Theorem 2 below. For any fixed $m \geq 2$ and $\nu \in [0, 1)$, with a calibration of γ_1, we establish

$$f(x_T) - f_* = \mathcal{O}\left(\frac{L_p \|x_0 - x_*\|_p^2 + (H_p \|x_0 - x_*\|_p^2)^{\frac{1}{1-\nu}}}{T^m} + \frac{L_p \|x_0 - x_*\|_p^2}{T^{1+2/p}} + \frac{(H_p \|x_0 - x_*\|_p^2)^{\frac{1}{1-\nu}}}{T^{\min\left\{m, \frac{1+2/p}{1-\nu}\right\}}} \right), \tag{23}$$

with an additional logarithmic factor $(\log T)^{(1-2/p)/(1-\nu)}$ in the last term when $m = (1 + 2/p)/(1 - \nu)$. The three terms reflect the effects of initialization, constant curvature, and gap-dependent curvature, respectively. In particular, when $\nu > 0$, the gap-dependent curvature term can decay faster than the constant-curvature term. For $L_p > 0$, the overall rate remains $\mathcal{O}(T^{-(1+2/p)})$, which matches the lower bound on the convergence-rate exponent in [7], up to dimension-dependent factors in the high-dimensional oracle regime. When $L_p = 0$ and $\nu > 0$, a suitable choice of m yields a faster convergence rate. These results highlight the advantages of exploiting generalized local smoothness through adaptive regularization and weighted averaging.

4 Convergence analysis

In this section, we establish convergence rate guarantees for the A^{2}ODA method by analyzing the potential sequence

$$\mathcal{P}_0 = 0, \qquad \mathcal{P}_k := \chi_k + \frac{1}{\gamma_{k+1}} V_q(-s_k) \qquad \forall k \geq 1, \tag{24}$$

where $\{\chi_k\}$, $\{s_k\}$, and $\{\gamma_k\}$ are generated by the algorithm.

The following theorem establishes upper bounds on $\{\mathcal{P}_k\}$ and the boundedness of the iterates generated by Algorithm 1.

Theorem 1. *Suppose that Assumption 1(a) holds. Let $\{(x_k, y_k, \gamma_k, \rho_k)\}$ be generated by Algorithm 1 and let $\{\mathcal{P}_k\}$ be defined in (24). Fix any minimizer x_* of f, and denote*

$$C_\theta = 2(1 + e + e/\theta), \quad \text{and} \quad \bar{\rho} = \max\left\{ \|g_0\|_q/(\alpha\gamma_1),\, C_\theta \|x_0 - x_*\|_p / \sqrt{\alpha^2 - C_\theta^2} \right\}. \tag{25}$$

Then, $0 < \rho_k \leq \bar{\rho}$ for $k \geq 1$, and for any $k \geq 0$,

$$\mathcal{P}_k \leq \rho_k^2 \gamma_{k+1}. \tag{26}$$

Moreover, it holds that $\|x_k - x_0\|_p, \|y_k - x_0\|_p \leq \alpha\rho_k$ for any $k \geq 1$, and $\|x_* - x_0\|_p \leq \alpha\bar{\rho}$. Consequently, for all $k \geq 1$,

$$x_k, y_k, x_* \in \mathcal{S}_p := \{x \in \mathbb{R}^n : \|x - x_0\|_p \leq \alpha\bar{\rho}\}. \tag{27}$$

Proof. We first prove (26). For $k = 0$, it follows from $\mathcal{P}_0 = 0$ and $\gamma_1 > 0$. Using the update rule of ρ_k given in (19), we have $\rho_1 = \|g_0\|_q/(\alpha\gamma_1) > 0$ and $\{\rho_k\}_{k\geq 1}$ is a nondecreasing sequence, which implies that $\rho_k > 0$ for all $k \geq 1$. Fix any $k \geq 1$, by the definitions of V_q and $\mathcal{P}_k$ in (3) and (24), one has

$$\mathcal{P}_k = \chi_k + \frac{1}{\gamma_{k+1}} V_q(-s_k) = \chi_k + \frac{\|s_k\|_q^2}{2\gamma_{k+1}}. \tag{28}$$

Observe from the update of γ_{k+1} in (21) that

$$\gamma_{k+1} \geq \frac{\chi_k + \sqrt{\chi_k^2 + 2\rho_k^2 \|s_k\|_q^2}}{2\rho_k^2} \qquad \Longrightarrow \qquad \rho_k^2 \gamma_{k+1}^2 - \chi_k \gamma_{k+1} - \frac{1}{2}\|s_k\|_q^2 \geq 0.$$

Dividing this inequality by $\gamma_{k+1} > 0$ and using (28), we obtain

$$\mathcal{P}_k = \chi_k + \frac{\|s_k\|_q^2}{2\gamma_{k+1}} \leq \rho_k^2 \gamma_{k+1}.$$

This proves (26). We next prove the boundedness of the iterates. Recall from (20) and (17) that

$$s_k = \sum_{i=1}^k \lambda_i g_i, \qquad c_k = \sum_{i=1}^k \lambda_i \left(f_i + \langle g_i, x_0 - x_i \rangle\right), \qquad A_k = \sum_{i=1}^k \lambda_i, \tag{29}$$

hold for any $k \geq 1$. Using these identities, we obtain

$$\mathcal{P}_k \overset{(28)}{=} \chi_k + \frac{\|s_k\|_q^2}{2\gamma_{k+1}} \overset{(20),(17)}{=} A_k f_k - c_k + \frac{\|s_k\|_q^2}{2\gamma_{k+1}} = A_k(f_k - f_*) + \sum_{i=1}^k \lambda_i D_f(x_*, x_i) + \langle s_k, x_* - x_0 \rangle + \frac{\|s_k\|_q^2}{2\gamma_{k+1}}$$

$$\geq \frac{\|s_k\|_q^2}{2\gamma_{k+1}} + \langle s_k, x_* - x_0\rangle \geq \frac{\|s_k\|_q^2}{2\gamma_{k+1}} - \|s_k\|_q\|x_* - x_0\|_p \geq \frac{\|s_k\|_q^2}{4\gamma_{k+1}} - \gamma_{k+1}\|x_* - x_0\|_p^2,$$

where the third equality follows from (29) and the definition of D_f, the first inequality is due to $f_k \geq f_*$ and the convexity of f, and the last two inequalities follow from Hölder's and Young's inequalities, respectively. Combining this inequality with $\mathcal{P}_k \leq \rho_k^2\gamma_{k+1}$ yields

$$\frac{\|s_k\|_q}{\gamma_{k+1}} \leq 2\sqrt{\|x_* - x_0\|_p^2 + \rho_k^2}. \tag{30}$$

This also holds for $k = 0$ because $s_0 = 0$. By (5) and the update of y_k in (18), one has for any $k \geq 1$ that

$$\|y_k - x_0\|_p = \frac{1}{\gamma_k}\|J_q(-\tilde{s}_k)\|_p = \frac{\|\tilde{s}_k\|_q}{\gamma_k}. \tag{31}$$

For $k = 1$, the update rule gives $\|y_1 - x_0\|_p = \|g_0\|_q/\gamma_1 = \alpha\rho_1$. For $k \geq 2$, combining $s_{k-1} - s_{k-2} = \lambda_{k-1}g_{k-1}$ and $\tilde{s}_k - s_{k-1} = \lambda_k g_{k-1}$ gives

$$\tilde{s}_k = \Big(1 + \frac{\lambda_k}{\lambda_{k-1}}\Big)s_{k-1} - \frac{\lambda_k}{\lambda_{k-1}}s_{k-2}. \tag{32}$$

By Lemma 8, we have $\lambda_k/\lambda_{k-1} \leq e$. Combining with $\gamma_k \geq \theta\gamma_{k-1}$, and $\rho_{k-2} \leq \rho_{k-1}$ gives

$$\begin{aligned}
\|y_k - x_0\|_p &\overset{(31)}{=} \frac{\|\tilde{s}_k\|_q}{\gamma_k} \leq \Big(1 + \frac{\lambda_k}{\lambda_{k-1}}\Big)\frac{\|s_{k-1}\|_q}{\gamma_k} + \frac{\lambda_k}{\lambda_{k-1}}\frac{\|s_{k-2}\|_q}{\gamma_k} \leq (1+e)\frac{\|s_{k-1}\|_q}{\gamma_k} + \frac{e}{\theta}\frac{\|s_{k-2}\|_q}{\gamma_{k-1}} \\
&\overset{(30)}{\leq} 2(1+e)\sqrt{\|x_0 - x_*\|_p^2 + \rho_{k-1}^2} + \frac{2e}{\theta}\sqrt{\|x_0 - x_*\|_p^2 + \rho_{k-2}^2} \\
&\leq 2\Big(1 + e + \frac{e}{\theta}\Big)\sqrt{\|x_0 - x_*\|_p^2 + \rho_{k-1}^2} \overset{(25)}{=} C_\theta\sqrt{\|x_0 - x_*\|_p^2 + \rho_{k-1}^2}.
\end{aligned}$$

Consequently, by (19) and the preceding bound on $\|y_k - x_0\|_p$, for every $k \geq 2$, we have

$$\rho_k \leq \max\left\{\rho_{k-1}, \frac{C_\theta}{\alpha}\sqrt{\|x_0 - x_*\|_p^2 + \rho_{k-1}^2}\right\}.$$

Since in Algorithm 1, we set $\alpha > C_\theta$, the definition of $\bar{\rho}$ in (25) gives $\rho_1 = \|g_0\|_q/(\alpha\gamma_1) \leq \bar{\rho}$ and

$$C_\theta^2\|x_0 - x_*\|_p^2 \leq (\alpha^2 - C_\theta^2)\bar{\rho}^2 \qquad \Longrightarrow \qquad \frac{C_\theta}{\alpha}\sqrt{\|x_0 - x_*\|_p^2 + \bar{\rho}^2} \leq \bar{\rho}.$$

We now prove $\rho_k \leq \bar{\rho}$ by induction. The case $k = 1$ has already been established. Suppose that $\rho_{k-1} \leq \bar{\rho}$ for some $k \geq 2$, we have

$$\rho_k \leq \max\left\{\bar{\rho}, \frac{C_\theta}{\alpha}\sqrt{\|x_0 - x_*\|_p^2 + \bar{\rho}^2}\right\} = \bar{\rho}.$$

Thus, we have $\rho_k \leq \bar{\rho}$ for all $k \geq 1$. It remains to bound the iterates. By (19), $\|y_k - x_0\|_p \leq \alpha\rho_k$ for every $k \geq 1$. To obtain the corresponding bound for x_k, note first that $\|x_0 - x_0\|_p = 0 = \alpha\rho_0$. Suppose that $\|x_{k-1} - x_0\|_p \leq \alpha\rho_{k-1}$. Using the update of x_k in (18), $A_k = A_{k-1} + \lambda_k$, and $\rho_{k-1} \leq \rho_k$, we obtain

$$\begin{aligned}
\|x_k - x_0\|_p &= \left\|\frac{A_{k-1}}{A_k}(x_{k-1} - x_0) + \frac{\lambda_k}{A_k}(y_k - x_0)\right\|_p \leq \frac{A_{k-1}}{A_k}\|x_{k-1} - x_0\|_p + \frac{\lambda_k}{A_k}\|y_k - x_0\|_p \\
&\leq \alpha\left(\frac{A_{k-1}}{A_k}\rho_{k-1} + \frac{\lambda_k}{A_k}\rho_k\right) \leq \alpha\frac{A_{k-1} + \lambda_k}{A_k}\rho_k = \alpha\rho_k.
\end{aligned}$$

Combining these bounds with $\rho_k \le \bar{\rho}$ yields $x_k, y_k \in \mathcal{S}_p$. Moreover, since $\alpha > C_\theta > 1$, the definition of $\bar{\rho}$ in (25) gives

$$\alpha\bar{\rho} \ge \frac{\alpha C_\theta}{\sqrt{\alpha^2 - C_\theta^2}}\|x_0 - x_*\|_p > \|x_0 - x_*\|_p,$$

which also implies $x_* \in \mathcal{S}_p$. This completes the proof. □

Remark 3 (Upper bounds on $\bar{\rho}$)**.** *We now derive upper bounds on $\bar{\rho}$ for the cases with and without calibration of γ_1. The calibration procedure is described in Calibration 1.*

(i) Without calibration of γ_1. Since $\nabla f(x_*) = 0$, (10) gives $\|g_0\|_q \le (L_p + H_p\Delta(x_0)^\nu)\|x_0 - x_*\|_p$. Hence, the definition of $\bar{\rho}$ yields

$$\bar{\rho} \le \max\left\{\frac{L_p + H_p\Delta(x_0)^\nu}{\alpha\gamma_1}, \frac{C_\theta}{\sqrt{\alpha^2 - C_\theta^2}}\right\}\|x_0 - x_*\|_p.$$

(ii) With calibration of γ_1. Since $f(x_1) \ge f_*$, the sufficient-decrease condition and convexity imply

$$\frac{\|g_0\|_q^2}{2\gamma_1} \le \Delta(x_0) \le \langle g_0, x_0 - x_*\rangle \le \|g_0\|_q\|x_0 - x_*\|_p.$$

Consequently, $\|g_0\|_q/(\alpha\gamma_1) \le (2/\alpha)\|x_0 - x_*\|_p$. Hence, the definition of $\bar{\rho}$ yields

$$\bar{\rho} \le \max\left\{\frac{2}{\alpha}, \frac{C_\theta}{\sqrt{\alpha^2 - C_\theta^2}}\right\}\|x_0 - x_*\|_p.$$

Therefore, calibrating γ_1 eliminates the explicit dependence on L_p and H_p in the upper bound for $\bar{\rho}$, leading to improved dependence on these constants in the convergence rate.

Using Assumption 1(b) and the compactness and convexity of $\mathcal{S}_p$, there exist constants $L_p(\mathcal{S}_p), H_p(\mathcal{S}_p) \ge 0$, denoted simply by L_p, H_p, such that

$$\|\nabla f(y) - \nabla f(x)\|_q^2 \le 2(L_p + H_p \max\{\Delta(x), \Delta(y)\}^\nu)D_f(x, y) \qquad \forall x, y \in \mathcal{S}_p. \tag{33}$$

For each $k \ge 2$, define the auxiliary curvature estimate by

$$\ell_k := \begin{cases} h_k\dfrac{\|g_k - g_{k-1}\|_q^2}{2D_f(x_{k-1}, x_k)}, & D_f(x_{k-1}, x_k) > 0, \\ 0, & D_f(x_{k-1}, x_k) = 0, \end{cases} \qquad h_k := \frac{\lambda_k^2}{2A_{k-1}}. \tag{34}$$

For each $k \ge 1$, define the local curvature bound

$$M_k := L_p + H_p \max\{\Delta(x_{k-1}), \Delta(x_k)\}^\nu. \tag{35}$$

Since $x_{k-1}, x_k \in \mathcal{S}_p$, applying (33) yields $\|g_k - g_{k-1}\|_q^2 \le 2M_k D_f(x_{k-1}, x_k)$. Consequently,

$$\ell_k \le h_k M_k \qquad \forall k \ge 2. \tag{36}$$

Our convergence analysis relies on comparisons with reference potentials. Specifically, we define the reference regularization scales $\{\eta_k\}$ and the reference potentials $\{\mathcal{Q}_k\}$ by

$$\eta_{k+1} := \|[\gamma_k, r_a\ell_k]\|_{r_q}, \qquad \mathcal{Q}_k := \chi_k + \frac{1}{\eta_{k+1}}V_q(-s_k) \qquad \forall k \ge 2, \tag{37}$$

where $r_a > 8$ is fixed and r_q is defined in (6). For notational convenience, we also define the auxiliary sequences denoting displacements from x_0:

$$z_k := \frac{1}{\gamma_k} J_q(-\tilde{s}_k), \qquad z_k^- := \frac{1}{\gamma_k} J_q(-s_{k-1}) \qquad \forall k \geq 1. \tag{38}$$

Lemma 2. *Suppose that Assumption 1 holds. Let $\{(x_k, s_k, \tilde{s}_k, \rho_k, \gamma_k)\}$ be generated by Algorithm 1, let $\{\mathcal{P}_k\}$ be defined in (24), let $\{\eta_k\}$ and $\{\mathcal{Q}_k\}$ be defined in (37), and let $\{(z_k, z_k^-)\}$ be defined in (38). Then, for any $k \geq 2$,*

$$\begin{aligned}\mathcal{Q}_k - \mathcal{P}_{k-1} = &-\frac{1}{\gamma_k} V_q(-\tilde{s}_k) + \langle z_k, s_k - \tilde{s}_k \rangle + \frac{1}{\eta_{k+1}} V_q(-s_k) \\ &- A_{k-1} D_f(x_{k-1}, x_k) - \gamma_k D_{d_p}^{-s_{k-1}/\gamma_k}(z_k, z_k^-).\end{aligned} \tag{39}$$

Proof. Fix any $k \geq 2$. By (18), (17), and (38), one has that $A_{k-1}(x_k - x_{k-1}) = \lambda_k(y_k - x_k)$ and $z_k = y_k - x_0$. Using these and the updates of A_k, c_k, and χ_k in (20) and (17), we have

$$\begin{aligned}\chi_k - \chi_{k-1} &= A_k f_k - A_{k-1} f_{k-1} - \lambda_k(f_k + \langle g_k, x_0 - x_k \rangle) = A_{k-1}(f_k - f_{k-1}) + \lambda_k \langle g_k, x_k - x_0 \rangle \\ &= A_{k-1}\langle g_k, x_k - x_{k-1} \rangle + \lambda_k \langle g_k, x_k - x_0 \rangle - A_{k-1} D_f(x_{k-1}, x_k) \\ &= \lambda_k \langle g_k, y_k - x_0 \rangle - A_{k-1} D_f(x_{k-1}, x_k) = \lambda_k \langle g_k, z_k \rangle - A_{k-1} D_f(x_{k-1}, x_k),\end{aligned}$$

where the third equality is due to the definition of D_f. By this and the definition of $\mathcal{P}_{k-1}$ in (24), one has

$$\chi_k - \mathcal{P}_{k-1} = \lambda_k \langle \nabla f(x_k), z_k \rangle - A_{k-1} D_f(x_{k-1}, x_k) - \frac{1}{\gamma_k} V_q(-s_{k-1}). \tag{40}$$

Since $V_q = d_p^*$, it follows from (4), (38), and conjugacy that $-\tilde{s}_k/\gamma_k \in \partial d_p(z_k)$ and $-s_{k-1}/\gamma_k \in \partial d_p(z_k^-)$. By Fenchel-Young equality in (2), one has

$$\gamma_k d_p(z_k) = -\langle \tilde{s}_k, z_k \rangle - \frac{1}{\gamma_k} V_q(-\tilde{s}_k), \qquad \gamma_k d_p(z_k^-) = -\langle s_{k-1}, z_k^- \rangle - \frac{1}{\gamma_k} V_q(-s_{k-1}). \tag{41}$$

Consequently,

$$\begin{aligned}\gamma_k D_{d_p}^{-s_{k-1}/\gamma_k}(z_k, z_k^-) &= \gamma_k d_p(z_k) - \gamma_k d_p(z_k^-) + \langle s_{k-1}, z_k - z_k^- \rangle \\ &\overset{(41)}{=} -\langle \tilde{s}_k, z_k \rangle - \frac{1}{\gamma_k} V_q(-\tilde{s}_k) + \langle s_{k-1}, z_k^- \rangle + \frac{1}{\gamma_k} V_q(-s_{k-1}) + \langle s_{k-1}, z_k - z_k^- \rangle \\ &= \frac{1}{\gamma_k}\big(V_q(-s_{k-1}) - V_q(-\tilde{s}_k)\big) - \langle \tilde{s}_k - s_{k-1}, z_k \rangle \\ &= \frac{1}{\gamma_k}\big(V_q(-s_{k-1}) - V_q(-\tilde{s}_k)\big) - \lambda_k \langle g_{k-1}, z_k \rangle,\end{aligned}$$

where the last equality follows from $\tilde{s}_k - s_{k-1} = \lambda_k g_{k-1}$. Substituting this into (40) and using $s_k - \tilde{s}_k = \lambda_k(g_k - g_{k-1})$, we obtain

$$\begin{aligned}\chi_k - \mathcal{P}_{k-1} &= -\frac{1}{\gamma_k} V_q(-\tilde{s}_k) + \lambda_k \langle g_k - g_{k-1}, z_k \rangle - A_{k-1} D_f(x_{k-1}, x_k) - \gamma_k D_{d_p}^{-s_{k-1}/\gamma_k}(z_k, z_k^-) \\ &= -\frac{1}{\gamma_k} V_q(-\tilde{s}_k) + \langle z_k, s_k - \tilde{s}_k \rangle - A_{k-1} D_f(x_{k-1}, x_k) - \gamma_k D_{d_p}^{-s_{k-1}/\gamma_k}(z_k, z_k^-).\end{aligned}$$

Combining this with the definition of $\mathcal{Q}_k$ in (37) completes the proof. □

We define the threshold

$$a_q := \frac{1}{2} - \frac{2^{2/r_q}}{r_a} > 0, \tag{42}$$

where r_q is defined in (6). We partition the iteration indices $k \geq 2$ as follows:

$$\mathcal{A} := \{k \geq 2 : \ell_k \leq a_q \gamma_k\}, \qquad \mathcal{B} := \{k \geq 2 : \ell_k > a_q \gamma_k\}. \tag{43}$$

This partition is based on a comparison between the regularization parameters $\{\gamma_k\}$ used by Algorithm 1 and the local curvature estimates $\{\ell_k\}$.

The next lemma shows that, for $k \in \mathcal{A}$, the next regularization parameter satisfies $\gamma_{k+1} \leq \eta_{k+1}$, which indicates that Algorithm 1 takes a larger step than the reference step.

Lemma 3. *Suppose that Assumption 1 holds. Let $\{(\rho_k, \gamma_k)\}$ be generated by Algorithm 1, and let $\{\mathcal{P}_k\}$, $\{\eta_k\}$, $\{\mathcal{Q}_k\}$, and $\mathcal{A}$ be defined in (24), (37), and (43), respectively. Then, we have $\mathcal{Q}_k \leq \mathcal{P}_{k-1} \leq \rho_k^2 \eta_{k+1}$ and $\gamma_{k+1} \leq \eta_{k+1}$ for all $k \in \mathcal{A}$.*

Proof. Fix any $k \in \mathcal{A}$. If $\ell_k = 0$, it follows from (34) and the cocoercivity of f that $g_k = g_{k-1}$. Then, one has $s_k = \tilde{s}_k$ and $\eta_{k+1} = \gamma_k$. Using these and (39), we obtain that

$$\begin{aligned}\mathcal{Q}_k - \mathcal{P}_{k-1} &= \frac{1}{\eta_{k+1}} V_q(-s_k) - \frac{1}{\gamma_k} V_q(-\tilde{s}_k) + \langle z_k, s_k - \tilde{s}_k \rangle - A_{k-1} D_f(x_{k-1}, x_k) - \gamma_k D_{d_p}^{-s_{k-1}/\gamma_k}(z_k, z_k^-) \\ &= -A_{k-1} D_f(x_{k-1}, x_k) - \gamma_k D_{d_p}^{-s_{k-1}/\gamma_k}(z_k, z_k^-) \leq 0.\end{aligned}$$

This together with $\eta_{k+1} = \gamma_k$ proves the conclusion.

Now suppose $\ell_k > 0$. We next prove $\mathcal{Q}_k \leq \mathcal{P}_{k-1}$ by considering two separate cases.

(i) $q = 2$. By (38) and (6), one has that $z_k = -\tilde{s}_k/\gamma_k$ and $r_q = \infty$. It then follows from (37) that $\eta_{k+1} = \max\{\gamma_k, r_a \ell_k\} \geq \gamma_k$. Then, the Euclidean identity (7) gives

$$\begin{aligned}&\frac{1}{\eta_{k+1}} V_2(-s_k) - \frac{1}{\gamma_k} V_2(-\tilde{s}_k) + \langle z_k, s_k - \tilde{s}_k \rangle \\ &= \frac{\|s_k\|_2^2}{2\eta_{k+1}} - \frac{\|\tilde{s}_k\|_2^2}{2\gamma_k} - \frac{\langle \tilde{s}_k, s_k - \tilde{s}_k \rangle}{\gamma_k} = \frac{\|s_k - \tilde{s}_k\|_2^2}{2\gamma_k} - \frac{1}{2}\Big(\frac{1}{\gamma_k} - \frac{1}{\eta_{k+1}}\Big)\|s_k\|_2^2 \leq \frac{\|s_k - \tilde{s}_k\|_2^2}{2\gamma_k},\end{aligned}$$

where the last inequality is due to $\eta_{k+1} \geq \gamma_k$. The definition of ℓ_k in (34), together with $s_k - \tilde{s}_k = \lambda_k(g_k - g_{k-1})$ (see (18) and (20)), yields

$$A_{k-1} D_f(x_{k-1}, x_k) = \frac{\|s_k - \tilde{s}_k\|_2^2}{4\ell_k}.$$

Substituting the above two relations into the potential identity (39) gives

$$\mathcal{Q}_k - \mathcal{P}_{k-1} \leq \Big(\frac{1}{2\gamma_k} - \frac{1}{4\ell_k}\Big)\|s_k - \tilde{s}_k\|_2^2 - \gamma_k D_{d_2}(z_k, z_k^-) \leq -\gamma_k D_{d_2}(z_k, z_k^-) \leq 0,$$

where the second inequality follows from $k \in \mathcal{A}$ and $\ell_k \leq a_2 \gamma_k \leq \gamma_k/2$, and the last inequality follows from the convexity of d_2.

(ii) $q \in [1, 2)$. Since $\ell_k > 0$, the definition of η_{k+1} in (37) gives $\eta_{k+1} > \gamma_k$ and $(\eta_{k+1}^{r_q} - \gamma_k^{r_q})^{1/r_q} = r_a \ell_k$. Applying (8) with $u = -\tilde{s}_k$, $v = \tilde{s}_k - s_k$, and $c = \gamma_k/\eta_{k+1} \in (0, 1)$, and using $z_k = J_q(-\tilde{s}_k)/\gamma_k$, we obtain

$$\frac{1}{\eta_{k+1}} V_q(-s_k) - \frac{1}{\gamma_k} V_q(-\tilde{s}_k) + \langle z_k, s_k - \tilde{s}_k \rangle = \frac{1}{\gamma_k}\left(\frac{\gamma_k}{\eta_{k+1}} V_q(-s_k) - V_q(-\tilde{s}_k) + \langle J_q(-\tilde{s}_k), s_k - \tilde{s}_k \rangle\right)$$

$$\leq \frac{1}{\gamma_k}\left(1+\frac{2^{2/r_q}\gamma_k}{(\eta_{k+1}^{r_q}-\gamma_k^{r_q})^{1/r_q}}\right)V_q(s_k-\tilde{s}_k)=\left(\frac{1}{2\gamma_k}+\frac{2^{2/r_q}}{2r_a\ell_k}\right)\|s_k-\tilde{s}_k\|_q^2. \tag{44}$$

The definition of ℓ_k in (34), together with $s_k-\tilde{s}_k=\lambda_k(g_k-g_{k-1})$, yields

$$A_{k-1}D_f(x_{k-1},x_k)=\frac{\|s_k-\tilde{s}_k\|_q^2}{4\ell_k}. \tag{45}$$

Substituting (44) and (45) into the potential identity (39) gives

$$\begin{aligned}\mathcal{Q}_k-\mathcal{P}_{k-1}&\leq\Big(\frac{1}{2\gamma_k}+\frac{2^{2/r_q}}{2r_a\ell_k}-\frac{1}{4\ell_k}\Big)\|s_k-\tilde{s}_k\|_q^2-\gamma_k D_{d_p}^{-s_{k-1}/\gamma_k}(z_k,z_k^-)\\&=\frac{1}{2}\Big(\frac{1}{\gamma_k}-\frac{a_q}{\ell_k}\Big)\|s_k-\tilde{s}_k\|_q^2-\gamma_k D_{d_p}^{-s_{k-1}/\gamma_k}(z_k,z_k^-)\leq-\gamma_k D_{d_p}^{-s_{k-1}/\gamma_k}(z_k,z_k^-)\leq 0,\end{aligned}$$

where the equality uses (42), the second inequality is due to $k\in\mathcal{A}$ and $\ell_k\leq a_q\gamma_k$, and the last inequality follows from the convexity of d_p.

Further, by the definition of η_{k+1} in (37), one has $\eta_{k+1}\geq\gamma_k$. Then, using (26) and the definition of $\mathcal{Q}_k$ in (37), we have

$$\chi_k+\frac{\|s_k\|_q^2}{2\eta_{k+1}}=\mathcal{Q}_k\leq\mathcal{P}_{k-1}\leq\rho_{k-1}^2\gamma_k\leq\rho_k^2\eta_{k+1}\qquad\Longrightarrow\qquad\eta_{k+1}\geq\frac{\chi_k+\sqrt{\chi_k^2+2\rho_k^2\|s_k\|_q^2}}{2\rho_k^2}.$$

This along with (21) and $\eta_{k+1}\geq\gamma_k\geq\theta\gamma_k$ implies $\eta_{k+1}\geq\gamma_{k+1}$, which completes the proof. □

The next lemma bounds γ_{k+1} by the scaled local curvature estimate, with a multiplicative constant independent of the iteration index.

Lemma 4. *Suppose that Assumption 1 holds. Let $\{(\rho_k,\gamma_k)\}$ be generated by Algorithm 1, let L_p and H_p satisfy (33), and let a_q be defined in (42), and recall that $\alpha>2(1+e+e/\theta)$, where $\theta\in(0,1)$ is the scaling parameter specified in Algorithm 1. Define*

$$\Gamma_{\mathcal{B}}:=\Big(4+\frac{1}{a_q}\Big)(1+\alpha)^2. \tag{46}$$

Then, $\gamma_{k+1}\leq\Gamma_{\mathcal{B}}h_kM_k$ for all $k\in\mathcal{B}$, where $\{h_k\}$ and $\{M_k\}$ are defined in (34) and (35), respectively.

Proof. Fix any $k\in\mathcal{B}$. Recall from (36) that $\ell_k\leq h_kM_k$. By this and the definition of $\mathcal{B}$, one has

$$\gamma_k<\frac{\ell_k}{a_q}\leq\frac{h_kM_k}{a_q}. \tag{47}$$

Recall that the update of γ_{k+1} has the form

$$\gamma_{k+1}=\max\left\{\theta\gamma_k,\ \frac{\chi_k+\sqrt{\chi_k^2+2(\rho_k\|s_k\|_q)^2}}{2{\rho_k}^2}\right\},$$

it remains to derive upper bounds on χ_k and $\|s_k\|_q$ to bound γ_{k+1}. By the updates of x_k, $\tilde{s}_k$, and s_k in (18) and (20), the bounds $\|y_k-x_0\|_p,\|x_k-x_0\|_p,\|x_{k-1}-x_0\|_p\leq\alpha\rho_k$ from Theorem 1, and (10), one has

$$\|s_k-\tilde{s}_k\|_q=\lambda_k\|g_k-g_{k-1}\|_q\leq\lambda_kM_k\|x_k-x_{k-1}\|_p=\frac{\lambda_k^2M_k}{A_k}\|y_k-x_{k-1}\|_p\leq 4h_kM_k\alpha\rho_k. \tag{48}$$

In addition, using the definition of z_k in (38) and the updates of y_k, we have

$$\|z_k\|_p = \|y_k - x_0\|_p \le \alpha\rho_k, \qquad \|\tilde{s}_k\|_q \le \gamma_k\|z_k\|_p \le \gamma_k\alpha\rho_k. \tag{49}$$

Combining the identity (39) with the definition of $\mathcal{Q}_k$ in (37), we obtain

$$\begin{aligned}\chi_k &= \mathcal{P}_{k-1} - \frac{1}{\gamma_k}V_q(-\tilde{s}_k) + \langle z_k, s_k - \tilde{s}_k\rangle - A_{k-1}D_f(x_{k-1}, x_k) - \gamma_k D_{d_p}^{-s_{k-1}/\gamma_k}(z_k, z_k^-)\\ &\le {\rho_k}^2\gamma_k + \langle z_k, s_k - \tilde{s}_k\rangle \le {\rho_k}^2\gamma_k + \|z_k\|_p\|s_k - \tilde{s}_k\|_q \overset{(47)(48)(49)}{\le} \frac{{\rho_k}^2 h_k M_k}{a_q} + 4h_k M_k\alpha^2\rho_k^2,\end{aligned}$$

where the first inequality is due to (26), $\rho_{k-1} \le \rho_k$, $V_q(-\tilde{s}_k) \ge 0$, and the convexity of f and d_p. Moreover, it follows from (47), (48), and (49) that

$$\|s_k\|_q \le \|\tilde{s}_k\|_q + \|s_k - \tilde{s}_k\|_q \le \gamma_k\alpha\rho_k + 4h_k M_k\alpha\rho_k \le \frac{h_k M_k\alpha\rho_k}{a_q} + 4h_k M_k\alpha\rho_k.$$

Combining the upper bounds for χ_k and $\|s_k\|_q$, we obtain

$$\begin{aligned}\frac{\chi_k + \sqrt{\chi_k^2 + 2(\rho_k\|s_k\|_q)^2}}{2{\rho_k}^2} &\le \frac{\chi_k + |\chi_k|}{2{\rho_k}^2} + \frac{\|s_k\|_q}{\sqrt{2}\rho_k} \le \left(a_q^{-1} + 4\alpha^2 + \frac{\alpha}{\sqrt{2}}(a_q^{-1} + 4)\right) h_k M_k\\ &\le \Gamma_{\mathcal{B}} h_k M_k,\end{aligned}$$

where the first inequality uses $\sqrt{a^2 + b^2} \le |a| + |b|$ for all $a, b \in \mathbb{R}$, and the last inequality follows from the direct calculation. This inequality, together with (46), (47), and (21) implies $\gamma_{k+1} \le \Gamma_{\mathcal{B}} h_k M_k$. Hence, the conclusion of this lemma holds. □

Next, we introduce the quantity

$$\Gamma_0 = \max\{\gamma_1, \gamma_2\} \tag{50}$$

to separate the initialization from the subsequent curvature estimates, while retaining $\gamma_1 > 0$ as an arbitrary input.

Lemma 5. *Suppose that Assumption 1 holds. Let $\{\gamma_k\}$ be generated by Algorithm 1, and let Γ_0 and M_1 be defined in (50) and (35), respectively. Then, $\Gamma_0 \le \alpha^2 M_1 + \alpha(\gamma_1 + M_1)/\sqrt{2}$.*

Proof. Since $\lambda_1 = A_1 = 1$, the first iteration gives $x_1 = y_1$, $s_1 = g_1$, and $\chi_1 = \langle g_1, x_1 - x_0\rangle$. Moreover, it is easy to verify that $\|x_1 - x_0\|_p = \alpha\rho_1$, $\|g_0\|_q = \gamma_1\alpha\rho_1$, and $\langle g_0, x_1 - x_0\rangle = -\gamma_1\alpha^2\rho_1^2$. Let L_p, H_p satisfy (33). Applying (10) yields

$$\begin{aligned}\chi_1 &= \langle g_0, x_1 - x_0\rangle + \langle g_1 - g_0, x_1 - x_0\rangle \le -\gamma_1\alpha^2\rho_1^2 + \|g_1 - g_0\|_q\|x_1 - x_0\|_p\\ &\le -\gamma_1\alpha^2\rho_1^2 + M_1\|x_1 - x_0\|_p^2 = (M_1 - \gamma_1)\alpha^2\rho_1^2 \le M_1\alpha^2\rho_1^2, \quad \text{and}\\ \|s_1\|_q &= \|g_1\|_q \le \|g_0\|_q + \|g_1 - g_0\|_q \le \alpha\rho_1(\gamma_1 + M_1).\end{aligned}$$

Using these estimates in (21), we obtain

$$\begin{aligned}\gamma_2 &= \max\left\{\theta\gamma_1, \frac{\chi_1 + \sqrt{\chi_1^2 + 2\rho_1^2\|s_1\|_q^2}}{2\rho_1^2}\right\} \le \max\left\{\theta\gamma_1, \frac{\max\{\chi_1, 0\}}{\rho_1^2} + \frac{\|s_1\|_q}{\sqrt{2}\rho_1}\right\}\\ &\le \max\left\{\theta\gamma_1, \alpha^2 M_1 + \frac{\alpha(\gamma_1 + M_1)}{\sqrt{2}}\right\} \le \alpha^2 M_1 + \frac{\alpha(\gamma_1 + M_1)}{\sqrt{2}},\end{aligned}$$

where the first inequality follows from $\sqrt{a^2 + b^2} \le |a| + |b|$, the last follows from $\alpha > \sqrt{2}$, and $\theta \in (0, 1)$. Since the second entry is also at least γ_1, we obtain $\Gamma_0 \le \alpha^2 M_1 + \alpha(\gamma_1 + M_1)/\sqrt{2}$. □

The next lemma provides upper bounds for $\{\gamma_k\}_{k\geq 2}$ generated by Algorithm 1.

Lemma 6. *Suppose that Assumption 1 holds. Let $\{\gamma_k\}$ be generated by Algorithm 1, and let r_q, $\{\ell_k\}$,$\{M_k\}$, $\Gamma_{\mathcal{B}}$, and Γ_0 be defined in (6), (34), (35), (46), and (50) respectively. Define $\Gamma_1 := \max\{r_a, \Gamma_{\mathcal{B}}\}$. Then, for all $k \geq 1$,*

$$\gamma_{k+1} \leq G_k, \qquad G_k := \begin{cases} \Gamma_0, & k = 1, \\ \Gamma_0 + \Gamma_1 \|(h_2 M_2, \ldots, h_k M_k)\|_{r_q}, & k \geq 2. \end{cases} \tag{51}$$

Proof. We prove $\gamma_{k+1} \leq G_k$ for all $k \geq 1$ by induction, starting from the established bound $\gamma_2 \leq \Gamma_0$. Fix $k \geq 2$ and suppose that $\gamma_k \leq G_{k-1}$. Consider the following two cases.

(i) Case $k \in \mathcal{A}$. By invoking the bound $\gamma_{k+1} \leq \eta_{k+1}$ established in Lemma 3, together with the definition of η_{k+1}, the estimate $\ell_k \leq h_k M_k$ in (36), the inequality $r_a \leq \Gamma_1$ implied by the definition of Γ_1, and the induction hypothesis $\gamma_k \leq G_{k-1}$, it follows that

$$\begin{aligned} \gamma_{k+1} \leq \eta_{k+1} &= \|(\gamma_k, r_a \ell_k)\|_{r_q} \leq \left\|\left(\Gamma_0 + \Gamma_1 \|(h_2 M_2, \ldots, h_{k-1} M_{k-1})\|_{r_q}, r_a \ell_k\right)\right\|_{r_q} \\ &\leq \left\|\left(\Gamma_0 + \Gamma_1 \|(h_2 M_2, \ldots, h_{k-1} M_{k-1})\|_{r_q}, r_a h_k M_k\right)\right\|_{r_q} \leq \Gamma_0 + \Gamma_1 \|(h_2 M_2, \ldots, h_k M_k)\|_{r_q} = G_k. \end{aligned}$$

(ii) Case $k \in \mathcal{B}$. By invoking the bound $\gamma_{k+1} \leq \Gamma_{\mathcal{B}} h_k M_k$ established in Lemma 4, together with the inequality $\Gamma_{\mathcal{B}} \leq \Gamma_1$ implied by the definition of Γ_1, we obtain

$$\gamma_{k+1} \leq \Gamma_{\mathcal{B}} h_k M_k \leq \Gamma_0 + \Gamma_1 \|(h_2 M_2, \ldots, h_k M_k)\|_{r_q} = G_k.$$

Combining these two cases completes the proof. □

Next, we provide our main theorem which establishes the convergence rate of Algorithm 1.

Theorem 2. *Suppose that Assumption 1 holds. Let $\{x_k\}$ be generated by Algorithm 1, let r_q, $\{\ell_k\}$, $\bar{\rho}$, $\Gamma_{\mathcal{B}}$, Γ_0, and $\{G_k\}$ be defined in (6), (34), (25), (46), (50) and (51), respectively, and let Γ_1 be defined in Lemma 6. Then, for any integer $T \geq 1$,*

$$f(x_T) - f_* \leq \frac{B_p G_T}{A_T}, \qquad B_p = \|x_0 - x_*\|_p^2/2 + \bar{\rho}^2. \tag{52}$$

Moreover, for $T \geq 2$, we further have

$$f(x_T) - f_* \leq \frac{2m^m \Gamma_0 B_p}{T^m} + \frac{2^m m^{2m} \Gamma_1 L_p B_p}{T^{3-2/q}} + \left(2^{m(1+\nu)} m^{2m} \Gamma_1 H_p C B_p\right)^{\frac{1}{1-\nu}} r_{m,\nu,q}(T), \tag{53}$$

where $C := C(m, \nu, q) > 0$ is independent of T and is given by

$$C = \begin{cases} 1, & q = 2 \text{ or } m(1-\nu) = 3 - \frac{2}{q}, \\ \left[\max\left\{1, \frac{2/q-1}{|m(1-\nu)-3+2/q|}\right\}\right]^{2/q-1}, & \text{otherwise}, \end{cases}$$

$$\text{and } \; r_{m,\nu,q}(T) = \begin{cases} T^{-m}, & m < \frac{3-2/q}{1-\nu}, \\ T^{-\frac{3-2/q}{1-\nu}} (\log T)^{\frac{2/q-1}{1-\nu}}, & m = \frac{3-2/q}{1-\nu}, \\ T^{-\frac{3-2/q}{1-\nu}}, & m > \frac{3-2/q}{1-\nu}. \end{cases} \tag{54}$$

Proof. For $T \geq 1$, by the definitions of $\mathcal{P}_T$, s_T, and c_T in (24) and (20), we have

$$\mathcal{P}_T = A_T f_T - c_T + \frac{\|s_T\|_q^2}{2\gamma_{T+1}} = A_T(f_T - f_*) + \sum_{k=1}^{T} \lambda_k D_f(x_*, x_k) + \langle s_T, x_* - x_0 \rangle + \frac{\|s_T\|_q^2}{2\gamma_{T+1}}.$$

It then follows from the convexity of f, Hölder's and Young's inequalities that

$$\mathcal{P}_T \geq A_T(f_T - f_*) - \|s_T\|_q \|x_0 - x_*\|_p + \frac{\|s_T\|_q^2}{2\gamma_{T+1}} \geq A_T(f_T - f_*) - \frac{\gamma_{T+1}}{2}\|x_0 - x_*\|_p^2.$$

Combining this with $\mathcal{P}_T \leq \rho_T^2 \gamma_{T+1}$ in (26), $\gamma_{k+1} \leq G_k$ for all $k \geq 1$ in (51), we obtain

$$A_T(f_T - f_*) \leq \mathcal{P}_T + \frac{\gamma_{T+1}}{2}\|x_0 - x_*\|_p^2 \overset{(26)}{\leq} \frac{\gamma_{T+1}}{2}(\|x_0 - x_*\|_p^2 + 2\rho_T^2) \leq B_p \gamma_{T+1} \overset{(51)}{\leq} B_p G_T,$$

where we use $\rho_T \leq \bar{\rho}$ and the definition of B_p. Dividing both sides by A_T yields (52). We now assume $T \geq 2$ and estimate G_T. For simplicity, we denote

$$V_T := \|(h_k)_{k=2}^T\|_{r_q}, \qquad U_T := \|(h_k A_{k-1}^{-\nu})_{k=2}^T\|_{r_q}. \tag{55}$$

First, for $2 \leq k \leq T$, applying (52) with T replaced by $k-1$ and k, respectively, and using $G_{k-1} \leq G_k \leq G_T$, we have $\Delta(x_{k-1}) \leq B_p G_T / A_{k-1}$ and $\Delta(x_k) \leq B_p G_T / A_k$. Moreover, by the definition of M_k and the monotonicity of $\{A_k\}$, we have

$$M_k = L_p + H_p \max\{\Delta(x_{k-1}), \Delta(x_k)\}^\nu \leq L_p + H_p B_p^\nu G_T^\nu A_{k-1}^{-\nu}.$$

This together with the definition of G_T gives

$$\begin{aligned} G_T &= \Gamma_0 + \Gamma_1 \|(h_2 M_2, \ldots, h_T M_T)\|_{r_q} \leq \Gamma_0 + \Gamma_1 L_p \|(h_2, \ldots, h_T)\|_{r_q} + \Gamma_1 H_p B_p^\nu G_T^\nu \|(h_2 A_1^{-\nu}, \ldots, h_T A_{T-1}^{-\nu})\|_{r_q} \\ &\overset{(55)}{=} \Gamma_0 + \Gamma_1 L_p V_T + \Gamma_1 H_p B_p^\nu U_T G_T^\nu \leq 2\Gamma_0 + 2\Gamma_1 L_p V_T + \left(2\Gamma_1 H_p B_p^\nu U_T\right)^{\frac{1}{1-\nu}}, \end{aligned} \tag{56}$$

where the first inequality uses Minkowski's inequality, and the last inequality follows from Lemma 10 with $a = G_T$, $b = \Gamma_0 + \Gamma_1 L_p V_T$, and $c = \Gamma_1 H_p B_p^\nu U_T$. We then estimate V_T and U_T. Using Lemma 8, we have $A_k \geq c_A k^m$ and $h_k \leq C_h k^{m-2}$, where $c_A = (m(m-1)^{m-1})^{-1}$ and $C_h = 2^{m-1}/c_A$ are the same constants as those defined in Lemma 8. It follows that

$$V_T = \|(h_k)_{k=2}^T\|_{r_q} \leq C_h \|(k^{m-2})_{k=2}^T\|_{r_q} \leq C_h T^{m-2+1/r_q} = C_h T^{m-3+2/q}, \tag{57}$$

$$U_T = \|(h_k A_{k-1}^{-\nu})_{k=2}^T\|_{r_q} \leq C_h \left(\frac{2^m}{c_A}\right)^\nu \|(k^{m(1-\nu)-2})_{k=2}^T\|_{r_q}, \tag{58}$$

where we use $1/r_q = 2/q - 1$, with $1/r_q = 0$ when $q = 2$. If $r_q < \infty$ (equivalently, $q \in [1, 2)$), define $\beta = (m(1-\nu) - 2)r_q$. The standard power-sum estimate yields

$$\sum_{k=2}^{T} k^\beta \leq \begin{cases} C_\beta, & \beta < -1, \\ C_\beta \log T, & \beta = -1, \\ C_\beta T^{\beta+1}, & \beta > -1, \end{cases} \qquad C_\beta = \begin{cases} -1/(\beta+1), & \beta < -1, \\ 1, & \beta = -1, \\ \max\{1, 1/(\beta+1)\}, & \beta > -1. \end{cases} \tag{59}$$

Taking the r_q-th root of (59) and using (58), we obtain

$$U_T \leq C_h C_\beta^{1/r_q} \left(\frac{2^m}{c_A}\right)^\nu \tilde{r}_{m,\nu,q}(T), \qquad \tilde{r}_{m,\nu,q}(T) = \begin{cases} 1, & m < \frac{3-2/q}{1-\nu}, \\ (\log T)^{2/q-1}, & m = \frac{3-2/q}{1-\nu}, \\ T^{m(1-\nu)-3+2/q} & m > \frac{3-2/q}{1-\nu}. \end{cases} \tag{60}$$

If $r_q = \infty$ (equivalently, $q = 2$), we instead take the maximum of $k^{m(1-\nu)-2}$ in (58). Consequently,

$$U_T \le C_h \left(\frac{2^m}{c_A}\right)^\nu \tilde{r}_{m,\nu,2}(T), \qquad \tilde{r}_{m,\nu,2}(T) = \begin{cases} 1, & m \le \frac{2}{1-\nu}, \\ T^{m(1-\nu)-2}, & m > \frac{2}{1-\nu}. \end{cases} \tag{61}$$

Combining the two cases in (60) and (61), for each fixed $q \in [1,2]$ with C defined in (54), we have $U_T \le C_h C\,(2^m/c_A)^\nu\, \tilde{r}_{m,\nu,q}(T)$. Substituting the bounds on V_T and U_T into the estimate of G_T, we obtain

$$\begin{aligned} G_T &\overset{(56)}{\le} 2\Gamma_0 + 2\Gamma_1 L_p V_T + \left(2\Gamma_1 H_p B_p^\nu U_T\right)^{\frac{1}{1-\nu}} \\ &\overset{(57),(60),(61)}{\le} 2\Gamma_0 + 2\Gamma_1 L_p C_h T^{m-3+2/q} + \left(2\Gamma_1 H_p B_p^\nu C_h C\,(2^m/c_A)^\nu\right)^{\frac{1}{1-\nu}}\, \tilde{r}_{m,\nu,q}^{\frac{1}{1-\nu}}(T). \end{aligned}$$

Finally, using (52) and $A_T \ge c_A T^m$, we obtain

$$f(x_T) - f_* \le \frac{B_p G_T}{A_T} \le \frac{B_p}{c_A}\left[2\Gamma_0 T^{-m} + 2\Gamma_1 L_p C_h T^{-(3-2/q)} + \left(2\Gamma_1 H_p B_p^\nu C_h C\,(2^m/c_A)^\nu\right)^{\frac{1}{1-\nu}}\, T^{-m}\, \tilde{r}_{m,\nu,q}^{\frac{1}{1-\nu}}(T)\right].$$

Since $r_{m,\nu,q}(T) = T^{-m}\tilde{r}_{m,\nu,q}(T)^{1/(1-\nu)}$, rearranging the constants and using $m - 1 \le m$ yields (53), which completes the proof. □

Remark 4 (Upper bounds on $\Gamma_0 B_p$)**.** *We now derive upper bounds on $\Gamma_0 B_p$ for the cases with and without calibration of γ_1. The calibration procedure is described in Calibration 1.*

(i) Without calibration of γ_1. By convexity, $\|g_0\|_q = \alpha\gamma_1\rho_1$, and the definitions of Γ_0 in (50) and B_p in (52), we have $\Delta(x_0) \le \alpha\gamma_1 B_p \le \alpha\Gamma_0 B_p$. Moreover, (52) with $T = 1$ gives $\Delta(x_1) \le \Gamma_0 B_p$, since $A_1 = 1$ and $G_1 = \Gamma_0$. Since $\alpha > 1$, the quantity M_1 defined in (35) satisfies

$$M_1 = L_p + H_p \max\{\Delta(x_0), \Delta(x_1)\}^\nu \le L_p + H_p(\alpha\Gamma_0 B_p)^\nu.$$

The bound $\Gamma_0 \le \alpha^2 M_1 + \alpha(\gamma_1 + M_1)/\sqrt{2}$ in Lemma 5 yields

$$\begin{aligned} \Gamma_0 B_p &\le (\tfrac{\alpha}{\sqrt{2}}\gamma_1 + (\alpha^2 + \tfrac{\alpha}{\sqrt{2}})L_p)B_p + \alpha^\nu(\alpha^2 + \tfrac{\alpha}{\sqrt{2}})H_p B_p(\Gamma_0 B_p)^\nu \\ &\le 2\big(\tfrac{\alpha}{\sqrt{2}}\gamma_1 + (\alpha^2 + \tfrac{\alpha}{\sqrt{2}})L_p\big)B_p + \left(2\alpha^\nu(\alpha^2 + \tfrac{\alpha}{\sqrt{2}})H_p B_p\right)^{\frac{1}{1-\nu}}. \end{aligned}$$

where the second inequality uses Lemma 10. For arbitrary $\gamma_1 > 0$, combining this estimate with Remark 3 gives

$$\begin{aligned} &\Gamma_0 B_p \le 2\kappa_0\left(\tfrac{\alpha}{\sqrt{2}}\gamma_1 + (\alpha^2 + \tfrac{\alpha}{\sqrt{2}})L_p\right)\|x_0 - x_*\|_p^2 + \left(2\alpha^\nu(\alpha^2 + \tfrac{\alpha}{\sqrt{2}})H_p\kappa_0\|x_0 - x_*\|_p^2\right)^{\frac{1}{1-\nu}} \\ &= \mathcal{O}\left((\gamma_1 + L_p)\left[1 + \frac{(L_p + H_p\Delta(x_0)^\nu)^2}{\gamma_1^2}\right]\|x_0 - x_*\|_p^2 + \left\{H_p\left[1 + \frac{(L_p + H_p\Delta(x_0)^\nu)^2}{\gamma_1^2}\right]\|x_0 - x_*\|_p^2\right\}^{\frac{1}{1-\nu}}\right) \\ &\text{with } \kappa_0 = \tfrac{1}{2} + \max\left\{\tfrac{(L_p + H_p\Delta(x_0)^\nu)^2}{\alpha^2\gamma_1^2}, \tfrac{C_\theta^2}{\alpha^2 - C_\theta^2}\right\}. \end{aligned} \tag{62}$$

This inequality follows from the estimate given in Remark 3 with

$$\begin{aligned} B_p &= \frac{1}{2}\|x_0 - x_*\|_p^2 + \max\left\{\frac{\|g_0\|_q^2}{\alpha^2\gamma_1^2}, \frac{C_\theta^2\|x_0 - x_*\|_p^2}{\alpha^2 - C_\theta^2}\right\} \\ &\le \left[\frac{1}{2} + \max\left\{\frac{(L_p + H_p\Delta(x_0)^\nu)^2}{\alpha^2\gamma_1^2}, \frac{C_\theta^2}{\alpha^2 - C_\theta^2}\right\}\right]\|x_0 - x_*\|_p^2 = \kappa_0\|x_0 - x_*\|_p^2. \end{aligned}$$

(ii) *With calibration of γ_1.* *If γ_1 is selected by Calibration 1, then $\|g_0\|_q^2/(2\Delta(x_0)) \le \gamma_1 < 2(L_p + H_p\Delta(x_0)^\nu)$ in (22) and $\Delta(x_1) \le \Delta(x_0)$ hold. Hence, the quantity M_1 defined in (35) satisfies $M_1 = L_p + H_p\Delta(x_0)^\nu$, and applying Lemma 5 together with $\gamma_1 < 2M_1$ yields $\Gamma_0 \le (\alpha^2 + 3\alpha/\sqrt{2})M_1$. Furthermore, the convexity of f and (10) imply $\Delta(x_0) \le M_1\|x_0 - x_*\|_p^2$. Substituting this into M_1 yields*

$$\begin{aligned} M_1\|x_0 - x_*\|_p^2 &\le L_p\|x_0 - x_*\|_p^2 + H_p\|x_0 - x_*\|_p^2 \left(M_1\|x_0 - x_*\|_p^2\right)^\nu \\ &\le 2L_p\|x_0 - x_*\|_p^2 + \left(2H_p\|x_0 - x_*\|_p^2\right)^{\frac{1}{1-\nu}}, \end{aligned}$$

where the second inequality uses Lemma 10. We then obtain

$$\begin{aligned} \Gamma_0 B_p &\le \left(\alpha^2 + \frac{3\alpha}{\sqrt{2}}\right) M_1 B_p = \left(\alpha^2 + \frac{3\alpha}{\sqrt{2}}\right)(L_p + H_p\Delta(x_0)^\nu)B_p \\ &\le \left(\alpha^2 + \frac{3\alpha}{\sqrt{2}}\right)\left[\frac{1}{2} + \max\left\{\frac{4}{\alpha^2}, \frac{C_\theta^2}{\alpha^2 - C_\theta^2}\right\}\right]\left[2L_p\|x_0 - x_*\|_p^2 + \left(2H_p\|x_0 - x_*\|_p^2\right)^{\frac{1}{1-\nu}}\right] \\ &= \mathcal{O}\left(L_p\|x_0 - x_*\|_p^2 + \left(H_p\|x_0 - x_*\|_p^2\right)^{\frac{1}{1-\nu}}\right), \end{aligned} \tag{63}$$

where the last inequality follows from the estimate given in Remark 3 with

$$B_p \le \left[\frac{1}{2} + \max\left\{\frac{4}{\alpha^2}, \frac{C_\theta^2}{\alpha^2 - C_\theta^2}\right\}\right]\|x_0 - x_*\|_p^2.$$

Estimate (62) holds for arbitrary $\gamma_1 > 0$ and retains explicit dependence on γ_1 and $\Delta(x_0)$. Calibration yields (63), which removes this dependence and the additional factors involving $(L_p + H_p\Delta(x_0)^\nu)^2/\gamma_1^2$, thereby improving the dependence on L_p and H_p in the upper bound on $\Gamma_0 B_p$.

Remark 5 (Convergence rate under Calibration 1)**.** *This remark aims to confirm the convergence rate stated earlier in (23). Under the assumptions of Theorem 2, with $0 \le \nu < 1$, choose γ_1 using Calibration 1. Remarks 3 and 4 then bound B_p and $\Gamma_0 B_p$, respectively. Substituting these estimates into (53) yields the following explicit convergence bound for every integer $T \ge 2$:*

$$\begin{aligned} f(x_T) - f_* \le{}& \frac{2m^m}{T^m}\left(\alpha^2 + \frac{3\alpha}{\sqrt{2}}\right)\left[\frac{1}{2} + \max\left\{\frac{4}{\alpha^2}, \frac{C_\theta^2}{\alpha^2 - C_\theta^2}\right\}\right]\left[2L_p\|x_0 - x_*\|_p^2 + \left(2H_p\|x_0 - x_*\|_p^2\right)^{\frac{1}{1-\nu}}\right] \\ &+ \frac{2^m m^{2m}\Gamma_1}{T^{3-2/q}}\left[\frac{1}{2} + \max\left\{\frac{4}{\alpha^2}, \frac{C_\theta^2}{\alpha^2 - C_\theta^2}\right\}\right] L_p\|x_0 - x_*\|_p^2 \\ &+ \left\{2^{m(1+\nu)} m^{2m}\Gamma_1 C\left[\frac{1}{2} + \max\left\{\frac{4}{\alpha^2}, \frac{C_\theta^2}{\alpha^2 - C_\theta^2}\right\}\right]\right\}^{\frac{1}{1-\nu}} \left(H_p\|x_0 - x_*\|_p^2\right)^{\frac{1}{1-\nu}} r_{m,\nu,q}(T), \end{aligned}$$

where C and $r_{m,\nu,q}$ are defined in (54). In addition, recall that $\Gamma_1 = \max\{r_a, (4 + a_q^{-1})(1 + \alpha)^2\}$, defined in Lemma 6, is independent of γ_1. This calibration sharpens the upper bounds on both $\Gamma_0 B_p$ and B_p, and consequently improves the dependence of the convergence bound on L_p and H_p.

References

[1] Z. Allen-Zhu and L. Orecchia. Linear coupling: An ultimate unification of gradient and mirror descent. In *8th Innovations in Theoretical Computer Science Conference*, 2017.

[2] J. Barzilai and J. M. Borwein. Two-point step size gradient methods. *IMA J. Numer. Anal.*, 8(1):141–148, 1988.

[3] H. H. Bauschke and J. M. Borwein. Legendre functions and the method of random Bregman projections. *Journal of Convex Analysis*, 4(1):27–67, 1997.

[4] A. Beck and M. Teboulle. Mirror descent and nonlinear projected subgradient methods for convex optimization. *Operations Research Letters*, 31(3):167–175, 2003.

[5] J. Bernstein, Y.-X. Wang, K. Azizzadenesheli, and A. Anandkumar. signSGD: Compressed optimisation for non-convex problems. In *Proceedings of the 35th International Conference on Machine Learning*, volume 80 of *Proceedings of Machine Learning Research*, pages 560–569. PMLR, 2018.

[6] A. d'Aspremont, C. Guzman, and M. Jaggi. Optimal affine-invariant smooth minimization algorithms. *SIAM Journal on Optimization*, 28(3):2384–2405, 2018.

[7] J. Diakonikolas and C. Guzmán. Lower bounds for parallel and randomized convex optimization. *Journal of Machine Learning Research*, 21(5):1–31, 2020.

[8] J. Diakonikolas and C. Guzmán. Complementary composite minimization, small gradients in general norms, and applications. *Mathematical Programming*, 208(1–2):319–363, 2024.

[9] J. Diakonikolas and L. Orecchia. The approximate duality gap technique: A unified theory of first-order methods. *SIAM Journal on Optimization*, 29(1):660–689, 2019.

[10] Y. Dong and W. Sawin. Muon^p: Muon with fractional spectral powers. arXiv:2606.13867, 2026.

[11] R.-A. Dragomir, A. B. Taylor, A. d'Aspremont, and J. Bolte. Optimal complexity and certification of Bregman first-order methods. *Mathematical Programming*, 194:41–83, 2022.

[12] F. Hanzely, P. Richtárik, and L. Xiao. Accelerated Bregman proximal gradient methods for relatively smooth convex optimization. *Computational Optimization and Applications*, 79(2):405–440, 2021.

[13] K. Jordan, Y. Jin, V. Boza, J. You, F. Cesista, L. Newhouse, and J. Bernstein. Muon: An optimizer for hidden layers in neural networks, 2024. Technical write-up, December 8, 2024.

[14] P. Joulani, A. Raj, A. György, and C. Szepesvári. A simpler approach to accelerated optimization: Iterative averaging meets optimism. In *Proceedings of the 37th International Conference on Machine Learning*, volume 119 of *Proceedings of Machine Learning Research*, pages 4984–4993, 2020.

[15] J. A. Kelner, Y. T. Lee, L. Orecchia, and A. Sidford. An almost-linear-time algorithm for approximate max flow in undirected graphs, and its multicommodity generalizations. In *Proceedings of the Twenty-Fifth Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 217–226. SIAM, 2014.

[16] W. Krichene, A. M. Bayen, and P. L. Bartlett. Accelerated mirror descent in continuous and discrete time. In *Advances in Neural Information Processing Systems*, volume 28, 2015.

[17] P. Latafat, A. Themelis, and P. Patrinos. On the convergence of adaptive first order methods: proximal gradient and alternating minimization algorithms. In *Proceedings of the 6th Annual Learning for Dynamics & Control Conference*, volume 242 of *Proceedings of Machine Learning Research*, pages 197–208. PMLR, 2024.

[18] T. Li and G. Lan. A simple uniformly optimal method without line search for convex optimization. *Mathematical Programming*, 219:219–256, 2026.

[19] A. Lobanov. A few accelerated algorithms for convex optimization under (H_0, H_1)-smoothness. arXiv:2608.04884, 2026.

[20] H. Lu, R. M. Freund, and Y. Nesterov. Relatively smooth convex optimization by first-order methods, and applications. *SIAM Journal on Optimization*, 28(1):333–354, 2018.

[21] Z. Lu and S. Mei. Accelerated first-order methods for convex optimization with locally Lipschitz continuous gradient. *SIAM Journal on Optimization*, 33(3):2275–2310, 2023.

[22] B. Ma, J. Zhang, and C. He. Schattor: Schatten-family methods for deep learning optimization. arXiv:2606.15702, 2026.

[23] C. J. Maddison, D. Paulin, Y. W. Teh, and A. Doucet. Dual space preconditioning for gradient descent. *SIAM Journal on Optimization*, 31(1):991–1016, 2021.

[24] Y. Malitsky and K. Mishchenko. Adaptive gradient descent without descent. In *Proceedings of the 37th International Conference on Machine Learning*, volume 119 of *Proceedings of Machine Learning Research*, pages 6702–6712. PMLR, 2020.

[25] R. Merchav, S. Sabach, and M. Teboulle. The adaptive integrated mean: A novel perspective for acceleration and its extension to Bregman first-order methods. *Mathematical Programming*, 2026.

[26] A. Nemirovski and D. B. Yudin. *Problem Complexity and Method Efficiency in Optimization*. Wiley, New York, 1983.

[27] Y. Nesterov. A method for solving the convex programming problem with convergence rate $O(1/k^2)$. *Soviet Mathematics Doklady*, 27:372–376, 1983.

[28] Y. Nesterov. *Introductory Lectures on Convex Optimization: A Basic Course*, volume 87 of *Applied Optimization*. Kluwer Academic Publishers, 2004.

[29] Y. Nesterov. Universal gradient methods for convex optimization problems. *Math. Program.*, 152(1-2):381–404, 2015.

[30] H. Ou, P. Latafat, and A. Themelis. Linesearch-free adaptive Bregman proximal gradient for convex minimization under local relative smoothness. *Journal of Optimization Theory and Applications*, 210(3), 2026. Article 68.

[31] T. Pethick. When to use what Schatten-p norm in deep learning? arXiv:2606.15268, 2026.

[32] A. Rakhlin and K. Sridharan. Online learning with predictable sequences. In *Proceedings of the 26th Annual Conference on Learning Theory*, volume 30, pages 993–1019, 2013.

[33] J. J. Suh and S. Ma. An adaptive and parameter-free Nesterov's accelerated gradient method for convex optimization. arXiv:2505.11670, 2025.

[34] A. Wibisono and A. C. Wilson. On accelerated methods in optimization. arXiv:1509.03616, 2015.

[35] L. Xiao. Dual averaging methods for regularized stochastic learning and online optimization. *Journal of Machine Learning Research*, 11:2543–2596, 2010.

[36] D. Zhou, S. Ma, and J. Yang. AdaBB: Adaptive Barzilai–Borwein method for convex optimization. *Mathematics of Operations Research*, 51(1):715–745, 2026.

Appendix

A Proof of the q-norm expansion inequality

Proof of Lemma 1. When $u = 0$, (8) holds clearly. Suppose that $\|u\|_q > 0$. Lemma 9 gives

$$\|u+v\|_q^q \le \|u\|_q^q + q\sum_{i=1}^n \omega_q(u_i)v_i + 2^{2-q}\|v\|_q^q = \|u\|_q^q + \frac{q}{\|u\|_q^{2-q}}\langle J_q(u), v\rangle + 2^{2-q}\|v\|_q^q, \tag{64}$$

where the last equality uses the definitions of ω_q and J_q. Recall from (5) that $\|J_q(u)\|_p = \|u\|_q$. By this and Hölder's inequality, one has $|\langle J_q(u), v\rangle| \le \|J_q(u)\|_p\|v\|_q = \|u\|_q\|v\|_q$. We now derive a nonnegative upper bound for the first two terms on the right-hand side of the above inequality:

$$\begin{aligned}\|u\|_q^q + \frac{q}{\|u\|_q^{2-q}}\langle J_q(u), v\rangle &\le \left|\|u\|_q + \frac{\langle J_q(u), v\rangle}{\|u\|_q}\right|^q = \left(\|u\|_q^2 + 2\langle J_q(u), v\rangle + \frac{\langle J_q(u), v\rangle^2}{\|u\|_q^2}\right)^{q/2} \\ &\le \left(\|u\|_q^2 + 2\langle J_q(u), v\rangle + \|v\|_q^2\right)^{q/2},\end{aligned}$$

where the first inequality is due to the convexity of $t \mapsto |t|^q$. Combining this with (64) yields

$$\|u+v\|_q^q \le (\|u\|_q^2 + 2\langle J_q(u), v\rangle + \|v\|_q^2)^{q/2} + 2^{2-q}\|v\|_q^q. \tag{65}$$

By Hölder's inequality with conjugate exponents $2/q$ and $2/(2-q)$, one has that $a_1b_1 + a_2b_2 \le (a_1^{2/q} + a_2^{2/q})^{q/2}(b_1^{2/(2-q)} + b_2^{2/(2-q)})^{(2-q)/2}$ for $a_1, b_1, a_2, b_2 \ge 0$. Fix any $c \in (0,1)$. By (65) and applying this Hölder's inequality with

$$a_1 = (\|u\|_q^2 + 2\langle J_q(u), v\rangle + \|v\|_q^2)^{q/2}, \quad a_2 = \frac{2^{2-q}c^{q/2}\|v\|_q^q}{(1-c^{r_q})^{(2-q)/2}}, \quad b_1 = c^{q/2}, \quad b_2 = (1-c^{r_q})^{(2-q)/2},$$

we obtain

$$\begin{aligned}c^{q/2}\|u+v\|_q^q &\le c^{q/2}(\|u\|_q^2 + 2\langle J_q(u), v\rangle + \|v\|_q^2)^{q/2} + (1-c^{r_q})^{(2-q)/2}\frac{2^{2-q}c^{q/2}\|v\|_q^q}{(1-c^{r_q})^{(2-q)/2}} \\ &\le \left(\|u\|_q^2 + 2\langle J_q(u), v\rangle + \|v\|_q^2 + \left(\frac{2^{2-q}c^{q/2}\|v\|_q^q}{(1-c^{r_q})^{(2-q)/2}}\right)^{2/q}\right)^{q/2} \\ &= \left(\|u\|_q^2 + 2\langle J_q(u), v\rangle + \left(1 + \frac{2^{2/r_q}c}{(1-c^{r_q})^{1/r_q}}\right)\|v\|_q^2\right)^{q/2},\end{aligned}$$

where we use the definition $r_q = q/(2-q)$. Raising both sides to the power $2/q$ and dividing by 2 proves (8) as desired. □

B Generalized local cocoercivity

Lemma 7. *Suppose that $f : \mathbb{R}^n \to \mathbb{R}$ is convex and continuously differentiable and attains its minimum value f_*. Define $\Delta(x) := f(x) - f_*$, and fix an arbitrary optimal solution x_*. Let $p \in [2, \infty]$ and $q \in [1, 2]$ satisfy $1/p + 1/q = 1$, and let $x_c \in \mathbb{R}^n$ be arbitrary. Fix any $R > 0$ satisfying $R \ge \|x_* - x_c\|_p$, and define*

$$\mathcal{C}_p := \{x \in \mathbb{R}^n : \|x - x_c\|_p \le R\}, \qquad \bar{\mathcal{C}}_p := \{x \in \mathbb{R}^n : \|x - x_c\|_p \le 3R\}.$$

Then, for any $L_p, H_p \ge 0$ and $\nu \in [0, 1)$, the Lipschitz-type condition

$$\|\nabla f(u) - \nabla f(v)\|_q \le \left(L_p + H_p \max\{\Delta(u), \Delta(v)\}^\nu\right)\|u - v\|_p \qquad \forall u, v \in \bar{\mathcal{C}}_p \tag{66}$$

implies the cocoercivity-type condition

$$\|\nabla f(x) - \nabla f(y)\|_q^2 \le 2\,(L_p + 2^\nu H_p \max\{\Delta(x), \Delta(y)\}^\nu)\, D_f(x,y) \qquad \forall x, y \in \mathcal{C}_p. \tag{67}$$

Proof. Suppose that (66) holds. We now prove (67). Fix any $x, y \in \mathcal{C}_p$. For convenience, denote

$$g_x := \nabla f(x), \quad g_y := \nabla f(y), \quad \Phi_{x,y} := \max\{\Delta(x), \Delta(y)\},$$
$$M_{x,y} := L_p + H_p\, \Phi_{x,y}^\nu, \quad \widetilde{M}_{x,y} := L_p + 2^\nu H_p\, \Phi_{x,y}^\nu.$$

The conclusion is immediate if $g_x = g_y$. Henceforth, suppose that $g_x \neq g_y$, which implies $\Phi_{x,y} > 0$. For any $z \in \mathcal{C}_p$ and any $v \in \mathbb{R}^n$ with $\|v\|_p = 1$, we define $\varphi_{z,v}(s) := \Delta(z - sv)$ for all $s \ge 0$. Taking the derivative, we have $\varphi'_{z,v}(s) = -\langle \nabla f(z - sv), v\rangle$. By this, Hölder's inequality, and (66), one has

$$|\varphi'_{z,v}(s) - \varphi'_{z,v}(r)| \le \|\nabla f(z - sv) - \nabla f(z - rv)\|_q \le (L_p + H_p \max\{\varphi_{z,v}(s), \varphi_{z,v}(r)\}^\nu)|s - r| \tag{68}$$

for all $s, r \ge 0$ such that $z - sv, z - rv \in \bar{\mathcal{C}}_p$. This one-dimensional Lipschitz-type condition will be used repeatedly below.

We first bound $\|g_y\|_q$ in terms of the objective gap. Claim that

$$\|g_y\|_q^2 \le 2M_{x,y}\Phi_{x,y}. \tag{69}$$

This claim holds clearly if $\|g_y\|_q = 0$. Otherwise, let $w \in \mathbb{R}^n$ be such that $\|w\|_p = 1$ and $\langle g_y, w\rangle = \|g_y\|_q > 0$. Since $\nabla f(x_*) = 0$ and $x_* \in \mathcal{C}_p$, it follows from (66) that

$$\|g_y\|_q \le M_{x,y}\|y - x_*\|_p \le 2M_{x,y}R.$$

Thus, setting $s_{x,y} := \|g_y\|_q / M_{x,y}$, we have $\{y - sw : 0 \le s \le s_{x,y}\} \subset \bar{\mathcal{C}}_p$, since

$$\|y - sw - x_c\|_p \le \|y - x_c\|_p + s\|w\|_p \le R + s_{x,y} \le 3R \qquad \forall s \in [0, s_{x,y}].$$

Then, applying (68) with $(z, v) = (y, w)$ and $r = 0$ and using $\varphi_{y,w}(0) = \Delta(y) \le \Phi_{x,y}$, we obtain

$$\varphi'_{y,w}(s) \le \varphi'_{y,w}(0) + (L_p + H_p \max\{\varphi_{y,w}(s), \Phi_{x,y}\}^\nu)s \qquad \forall s \in [0, s_{x,y}]. \tag{70}$$

We next prove $\varphi_{y,w}(s) \le \Phi_{x,y}$ for all $s \in [0, s_{x,y}]$ and relate the coefficient of s in (70) to $M_{x,y}$. Notice from the definition of $\varphi_{y,w}$ and the choice of w that $\varphi'_{y,w}(0) = -\|g_y\|_q < 0$. We show by contradiction that $\varphi_{y,w}(s) < \varphi_{y,w}(0)$ for all $s \in (0, s_{x,y}]$. Otherwise, there exists an $s_0 \in (0, s_{x,y}]$ such that $\varphi_{y,w}(s_0) = \varphi_{y,w}(0)$ and $\varphi_{y,w}(s) < \varphi_{y,w}(0)$ for all $s \in (0, s_0)$. Therefore, $\varphi_{y,w}(s) \le \varphi_{y,w}(0) \le \Phi_{x,y}$ for all $s \in [0, s_0]$. By this and (70), one has

$$\varphi'_{y,w}(s) \le \varphi'_{y,w}(0) + (L_p + H_p\Phi_{x,y}^\nu)s = \varphi'_{y,w}(0) + M_{x,y}s = -\|g_y\|_q + M_{x,y}s \qquad \forall s \in [0, s_0].$$

Integrating this inequality over $s \in [0, s_0]$, we obtain $\varphi_{y,w}(s_0) \le \varphi_{y,w}(0) - \|g_y\|_q\, s_0 + M_{x,y}s_0^2/2 < \varphi_{y,w}(0)$, where the strict inequality follows from $s_0 \in (0, s_{x,y}]$ and $s_{x,y} = \|g_y\|_q / M_{x,y}$. This contradicts $\varphi_{y,w}(s_0) = \varphi_{y,w}(0)$. Hence, $\varphi_{y,w}(s) \le \varphi_{y,w}(0) \le \Phi_{x,y}$ for all $s \in [0, s_{x,y}]$. Combining this with (70), we obtain

$$\varphi'_{y,w}(s) \le \varphi'_{y,w}(0) + (L_p + H_p\Phi_{x,y}^\nu)s = -\|g_y\|_q + M_{x,y}s \qquad \forall s \in [0, s_{x,y}].$$

Integrating this inequality over $s \in [0, s_{x,y}]$, we obtain

$$\varphi_{y,w}(s_{x,y}) \le \varphi_{y,w}(0) - \|g_y\|_q\, s_{x,y} + \frac{M_{x,y}}{2}s_{x,y}^2 = \varphi_{y,w}(0) - \frac{\|g_y\|_q^2}{2M_{x,y}} \le \Phi_{x,y} - \frac{\|g_y\|_q^2}{2M_{x,y}}.$$

Together with $\varphi_{y,w}(s_{x,y}) \geq 0$, which follows from the definition of $\varphi_{y,w}$, this yields the claim (69).

We are now ready to prove (67). Let $u \in \mathbb{R}^n$ be such that $\|u\|_p = 1$ and $\langle g_x - g_y, u\rangle = \|g_x - g_y\|_q$. By $x, y \in \mathcal{C}_p \subset \bar{\mathcal{C}}_p$, one has from (66) that

$$\|g_x - g_y\|_q \leq M_{x,y}\|x - y\|_p \leq 2M_{x,y}R. \tag{71}$$

Since $g_x \neq g_y$, it then follows that $M_{x,y} > 0$, and hence $\widetilde{M}_{x,y} \geq M_{x,y} > 0$. Set $t_{x,y} := \|g_x - g_y\|_q/\widetilde{M}_{x,y}$. By (71), we have $t_{x,y} \leq 2R$. Thus, $\{x - su : 0 \leq s \leq t_{x,y}\} \subset \bar{\mathcal{C}}_p$, since

$$\|x - su - x_c\|_p \leq \|x - x_c\|_p + s\|u\|_p \leq R + t_{x,y} \leq 3R \qquad \forall s \in [0, t_{x,y}].$$

Then, applying (68) with $(z, v) = (x, u)$ and $r = 0$, and using $\varphi_{x,u}(0) = \Delta(x) \leq \Phi_{x,y}$, we obtain

$$\varphi'_{x,u}(s) \leq \varphi'_{x,u}(0) + (L_p + H_p \max\{\varphi_{x,u}(s), \Phi_{x,y}\}^\nu)s \qquad \forall s \in [0, t_{x,y}]. \tag{72}$$

Claim that

$$\varphi_{x,u}(s) \leq 2\Phi_{x,y} \qquad \forall s \in [0, t_{x,y}]. \tag{73}$$

Suppose for contradiction that this bound fails. Since $\varphi_{x,u}(0) \leq \Phi_{x,y} < 2\Phi_{x,y}$, continuity implies that there exists an $s_0 \in (0, t_{x,y})$ such that $\varphi_{x,u}(s_0) = 2\Phi_{x,y}$ and $\varphi_{x,u}(s) \leq 2\Phi_{x,y}$ for all $s \in [0, s_0]$. Combining this with (72) yields $\varphi'_{x,u}(s) \leq \varphi'_{x,u}(0) + \widetilde{M}_{x,y}s$ for all $s \in [0, s_0]$. By the choice of u and Hölder's inequality, one has

$$\varphi'_{x,u}(0) = -\langle g_x, u\rangle = -\|g_x - g_y\|_q - \langle g_y, u\rangle \leq \|g_y\|_q - \|g_x - g_y\|_q.$$

For convenience, denote

$$U(t) := \Phi_{x,y} + (\|g_y\|_q - \|g_x - g_y\|_q)\, t + \frac{\widetilde{M}_{x,y}}{2}\, t^2 \qquad \forall t \in [0, t_{x,y}].$$

Then, by $\varphi_{x,u}(s) \leq 2\Phi_{x,y}$ for all $s \in [0, s_0]$ and integrating (72) over $s \in [0, s_0]$, we obtain

$$\varphi_{x,u}(s_0) \leq \varphi_{x,u}(0) + \varphi'_{x,u}(0)\, s_0 + (L_p + 2^\nu H_p \Phi^\nu_{x,y})\, \frac{s_0^2}{2} \leq U(s_0). \tag{74}$$

On the other hand, by the definitions of U and $t_{x,y}$, we have

$$U(0) = \Phi_{x,y}, \qquad U(t_{x,y}) = \Phi_{x,y} + \frac{\|g_y\|_q\, \|g_x - g_y\|_q}{\widetilde{M}_{x,y}} - \frac{\|g_x - g_y\|_q^2}{2\widetilde{M}_{x,y}} \leq \Phi_{x,y} + \frac{\|g_y\|_q^2}{2\widetilde{M}_{x,y}} \leq 2\Phi_{x,y},$$

where the first inequality follows by completing the square, and the second inequality follows from (69) and $M_{x,y} \leq \widetilde{M}_{x,y}$. By convexity of U and the fact that $s_0 \in [0, t_{x,y})$, one has

$$U(s_0) \leq \left(1 - \frac{s_0}{t_{x,y}}\right) U(0) + \frac{s_0}{t_{x,y}} U(t_{x,y}) \leq \left(1 + \frac{s_0}{t_{x,y}}\right) \Phi_{x,y} < 2\Phi_{x,y},$$

which along with (74) implies $\varphi_{x,u}(s_0) \leq U(s_0) < 2\Phi_{x,y}$. This contradicts $\varphi_{x,u}(s_0) = 2\Phi_{x,y}$. Hence, the claim (73) holds.

Combining (73) with (72), we obtain

$$\varphi'_{x,u}(s) \leq \varphi'_{x,u}(0) + \left(L_p + 2^\nu H_p \Phi^\nu_{x,y}\right) s = \varphi'_{x,u}(0) + \widetilde{M}_{x,y}s \qquad \forall s \in [0, t_{x,y}]. \tag{75}$$

Define $h(z) := f(z) - \langle g_y, z\rangle$ for all $z \in \mathbb{R}^n$. Since h is convex and $\nabla h(y) = 0$, y is a global minimizer of h. Moreover, by the definitions of h and $\varphi_{x,u}$, one has

$$h(x) - h(y) = D_f(x,y), \quad \nabla h(x) = g_x - g_y, \quad h(x - su) - h(x) = \varphi_{x,u}(s) - \varphi_{x,u}(0) + s\langle g_y, u\rangle. \tag{76}$$

Integrating (75) over $s \in [0, t_{x,y}]$ and using $\varphi'_{x,u}(0) = -\langle g_x, u\rangle$, we obtain

$$h(x - t_{x,y}u) - h(x) \le -t_{x,y}\langle g_x - g_y, u\rangle + \frac{\widetilde{M}_{x,y}}{2} t_{x,y}^2 = -\|g_x - g_y\|_q \, t_{x,y} + \frac{\widetilde{M}_{x,y}}{2} t_{x,y}^2.$$

Since y is a global minimizer of h, we have $h(y) \le h(x - t_{x,y}u)$. Therefore,

$$D_f(x,y) = h(x) - h(y) \ge \|g_x - g_y\|_q \, t_{x,y} - \frac{\widetilde{M}_{x,y}}{2} t_{x,y}^2 = \frac{\|g_x - g_y\|_q^2}{2\widetilde{M}_{x,y}}.$$

Multiplying both sides by $2\widetilde{M}_{x,y}$ proves (67) as desired. □

C Proof of the calibration bound

Proof of (22). Since γ_1 is accepted and $f(x_1) \ge f_*$, the sufficient-decrease condition gives $\frac{\|g_0\|_q^2}{2\gamma_1} \le \Delta(x_0)$, which proves the lower bound.

For the upper bound, we first show that the rejected trial $x_1(\gamma_1/2)$ belongs to $\mathcal{S}_p$ defined in (27). By convexity, for any minimizer x_*, we have $\Delta(x_0) \le \langle g_0, x_0 - x_*\rangle \le \|g_0\|_q \|x_0 - x_*\|_p$. Combining (5) with the lower bound above yields

$$\|x_1(\gamma_1/2) - x_0\|_p = \frac{2\|g_0\|_q}{\gamma_1} \le \frac{4\Delta(x_0)}{\|g_0\|_q} \le 4\|x_0 - x_*\|_p.$$

On the other hand, (25) and $\alpha > C_\theta > 4$ imply

$$\alpha\bar{\rho} \ge \frac{\alpha C_\theta}{\sqrt{\alpha^2 - C_\theta^2}} \|x_0 - x_*\|_p > 4\|x_0 - x_*\|_p.$$

Hence $x_1(\gamma_1/2) \in \mathcal{S}_p$. Second, we show that every trial $\gamma \ge \max\{\gamma_1/2, L_p + H_p\Delta(x_0)^\nu\}$ satisfies the sufficient-decrease condition. Define

$$z(t) = x_0 + \frac{t}{\gamma} J_q(-g_0), \qquad t \in [0,1].$$

Since $\mathcal{S}_p$ is convex and contains x_0 and $x_1(\gamma_1/2)$, we have $z(t) \in \mathcal{S}_p$ for all $t \in [0,1]$. Suppose that $f(z(t)) \ge f(x_0)$ for some $t \in (0,1]$. Since

$$\frac{\mathrm{d}}{\mathrm{d}t} f(z(t))\Big|_{t=0} = -\frac{\|g_0\|_q^2}{\gamma} < 0,$$

there exists a first $t_0 \in (0,1]$ such that $f(z(t_0)) = f(x_0)$ and $f(z(t)) < f(x_0)$ for all $t \in (0, t_0)$. Thus, for $t \in [0, t_0]$, $\Delta(z(t)) \le \Delta(x_0)$. By (10), we have

$$\|\nabla f(z(t)) - g_0\|_q \le (L_p + H_p\Delta(x_0)^\nu)\|z(t) - x_0\|_p = \frac{(L_p + H_p\Delta(x_0)^\nu)t}{\gamma}\|g_0\|_q.$$

Integrating along the segment and using (5), we obtain

$$f(z(t_0)) - f(x_0) \le -\frac{t_0}{\gamma}\|g_0\|_q^2 + \frac{(L_p + H_p\Delta(x_0)^\nu)t_0^2}{2\gamma^2}\|g_0\|_q^2 \le -\frac{t_0}{2\gamma}\|g_0\|_q^2 < 0,$$

where the last inequality uses $\gamma \geq (L_p + H_p \Delta(x_0)^\nu)$ and $t_0 \leq 1$. This is a contradiction. Consequently, $f(z(t)) \leq f(x_0)$ throughout $[0, 1]$. Applying the same estimate at $t = 1$ gives

$$f(x_1(\gamma)) \leq f(x_0) - \frac{\|g_0\|_q^2}{\gamma} + \frac{(L_p + H_p \Delta(x_0)^\nu)\|g_0\|_q^2}{2\gamma^2} \leq f(x_0) - \frac{\|g_0\|_q^2}{2\gamma}.$$

Thus, every $\gamma \geq \max\{\gamma_1/2, L_p + H_p \Delta(x_0)^\nu\}$ is accepted. By the calibration procedure, the trial $\gamma_1/2$ is rejected. Therefore, we obtain $\gamma_1 < 2(L_p + H_p \Delta(x_0)^\nu)$. This completes the proof. □

D Technical lemmas

Lemma 8. *Let $c_A = (m(m-1)^{m-1})^{-1}$, and $C_h = 2^{m-1}/c_A$. For $m \geq 2$ and $k \geq 2$, the weights λ_k and A_k in Algorithm 1 satisfy*

$$1 \leq \frac{\lambda_k}{\lambda_{k-1}} \leq e, \qquad c_A k^m \leq A_k \leq k^m, \qquad \frac{k^{m-2}}{2(m-1)^{2m-2}} \leq h_k := \frac{\lambda_k^2}{2A_{k-1}} \leq C_h k^{m-2}. \tag{77}$$

Proof. We prove the results given in (77) one by one. First, for $k \geq 2$, using the update of λ_k in (17), the first inequality can be verified directly by

$$1 \leq \frac{\lambda_k}{\lambda_{k-1}} = \left(\frac{k+m-2}{k+m-3}\right)^{m-1} \leq \left(1 + \frac{1}{m-1}\right)^{m-1} \leq e.$$

Second, for every $k \geq 1$, we have $\frac{k}{m-1} \leq 1 + \frac{k-1}{m-1} = \lambda_k^{1/(m-1)} \leq k$. Hence, taking the $m-1$ power, summing λ_i from $i = 1$ to k, and using the update form of A_k in (17), we obtain

$$\frac{k^{m-1}}{(m-1)^{m-1}} \leq \lambda_k \leq k^{m-1}, \qquad \frac{1}{(m-1)^{m-1}} \sum_{i=1}^{k} i^{m-1} \leq A_k \leq \sum_{i=1}^{k} i^{m-1}.$$

Using the elementary inequality $k^m/m \leq \sum_{i=1}^{k} i^{m-1} \leq k^m$ for $m \geq 1$, we directly have $c_A k^m \leq A_k \leq k^m$ by using the definition $c_A = (m(m-1)^{m-1})^{-1}$. Finally, for $k \geq 2$, the inequality $k - 1 \geq k/2$ gives

$$h_k = \frac{\lambda_k^2}{2A_{k-1}} \leq \frac{k^{2m-2}}{2c_A(k-1)^m} \leq \frac{2^{m-1}}{c_A} k^{m-2} = C_h k^{m-2}.$$

For the reverse order estimate, using $\lambda_k \geq (k/(m-1))^{m-1}$ and $A_{k-1} \leq (k-1)^m$, we obtain

$$h_k = \frac{\lambda_k^2}{2A_{k-1}} \geq \frac{k^{2m-2}}{2(m-1)^{2m-2}(k-1)^m} \geq \frac{k^{m-2}}{2(m-1)^{2m-2}}.$$

This completes the proof. □

Lemma 9. *Let $q \in [1, 2)$, and define $\omega_q(t) := \operatorname{sgn}_0(t)|t|^{q-1}$ for all $t \in \mathbb{R}$. Then,*

$$\|u + v\|_q^q \leq \|u\|_q^q + q \sum_{i=1}^{n} \omega_q(u_i) v_i + 2^{2-q} \|v\|_q^q \tag{78}$$

holds for all $u, v \in \mathbb{R}^n$.

Proof. Define $\omega_q(t) := \operatorname{sgn}_0(t)|t|^{q-1}$ for all $t \in \mathbb{R}$. We first show that, for all $q \in [1, 2)$,

$$|\omega_q(a) - \omega_q(b)| \leq 2^{2-q}|a - b|^{q-1} \qquad \forall a, b \in \mathbb{R}. \tag{79}$$

To prove the claim, we consider the following three cases.

(i) If $ab > 0$, the subadditivity of $t \mapsto t^{q-1}$ on $\mathbb{R}_+$ gives $|\omega_q(a) - \omega_q(b)| = \big||a|^{q-1} - |b|^{q-1}\big| \le \big||a| - |b|\big|^{q-1} = |a-b|^{q-1}$, which further implies (79) using $q \in [1,2)$;

(ii) If $ab < 0$, the concavity of $t \mapsto t^{q-1}$ on $\mathbb{R}_+$ gives $|\omega_q(a) - \omega_q(b)| = |a|^{q-1} + |b|^{q-1} \le 2^{2-q}(|a|+|b|)^{q-1} = 2^{2-q}|a-b|^{q-1}$;

(iii) If $ab = 0$, then $|\omega_q(a) - \omega_q(b)| \le |a-b|^{q-1}$, which further implies (79).

Combining all these cases establishes (79). Moreover, for any $q \in [1,2)$, the function $t \mapsto |t|^q$ is locally absolutely continuous, and its derivative equals $q\omega_q(t)$ almost everywhere. Thus, for all $a, b \in \mathbb{R}$,

$$\begin{aligned} |a+b|^q - |a|^q - q\omega_q(a)b &= q\int_0^1 (\omega_q(a+tb) - \omega_q(a))b \,\mathrm{d}t \le q\int_0^1 |\omega_q(a+tb) - \omega_q(a)|\,|b| \,\mathrm{d}t \\ &\overset{(79)}{\le} q\,2^{2-q}|b|^q \int_0^1 t^{q-1}\,\mathrm{d}t = 2^{2-q}|b|^q. \end{aligned}$$

Applying it with $a = u_i$ and $b = v_i$ and summing over $i = 1, \dots, n$ yields the desired result (78). □

Lemma 10. *Let $a, b, c \ge 0$, $\nu \in [0,1)$. If $a \le b + ca^\nu$, then $a \le 2b + (2c)^{\frac{1}{1-\nu}}$.*

Proof. By the weighted Young's inequality, we have

$$ca^\nu = \frac{1}{2}\big[(2c)^{1/(1-\nu)}\big]^{1-\nu} a^\nu \le \frac{1}{2}\Big((1-\nu)(2c)^{\frac{1}{1-\nu}} + \nu a\Big) \le \frac{1}{2}\Big((2c)^{\frac{1}{1-\nu}} + a\Big).$$

Substituting this into the inequality $a \le b + ca^\nu$ and rearranging gives the claim. □