

Beyond Report Imitation: Clinically Aware Multi-Image Ultrasound Report Generation from Visible Evidence

Yuchen Yang¹

3319733548@bupt.edu.cn

Wang Xin²

wangxin21@pumch.cn

Lufan Wang¹

wlf@bupt.edu.cn

Yinghong Pan¹

panyinghong@bupt.edu.cn

Yujuan Feng³

yujuanf15@bjut.edu.cn

Yuying Yang^{1,*}

yangyuying@bupt.edu.cn

¹ State Key Laboratory of Networking and Switching Technology, Beijing University of Posts and Telecommunications, Beijing, China

² Department of Ultrasound, State Key Laboratory of Complex Severe and Rare Diseases, Peking Union Medical College Hospital, Chinese Academy of Medical Sciences & Peking Union Medical College, Beijing, China

³ College of Computer Science, Beijing University of Technology, Beijing, China

Abstract

Generating ultrasound reports from multiple images requires aggregating clinical evidence across views, yet archived key frames capture only part of the dynamic examination. Raw-report imitation is therefore misaligned with visual supervision: content that is clinically valid for the full examination may be unverifiable from the images available to a model. This gap creates a clinical behavior alignment problem. A model must preserve visible findings, avoid diagnostic reversals and unsupported completion, and not collapse into conservative templates. We propose **CAMEO**, a **Clinically Aware Multi-image Evidence-grounded Orchestration** framework for ultrasound report generation. Stage I learns ultrasound visual-language primitives; Stage II performs Cross-View Evidence Grounding by distilling trusted visible report points into multi-image QA and report-style supervision; and Stage III performs Clinically Aware Preference Alignment using clinical-error-oriented preference pairs. From USReport, we construct USREPORT-DISTILLED with 17,670 evidence-grounded paired-image training instances and USREPORT-PREF with 21,869 preference pairs; we additionally use 25,631 PUBMEDVISION-US ultrasound instruction samples for domain adaptation and multi-image instruction tuning. On the primary USREPORT-DISTILLED benchmark, CAMEO improves over EchoVLM from 0.25 to 0.40 BLEU-1, 0.28 to 0.45 ROUGE-1, and 0.27 to 0.43 METEOR, while raising ClinicalScore from 55.02 to 74.20. These results underscore the value of evidence-grounded supervision, clinically aware alignment, and clinically structured evaluation for reliable ultrasound report generation. Code and project resources: <https://github.com/NiHaoWoJiaoYYC/CAMEO>.




Figure 1: Motivation. Archived ultrasound key frames only partially observe the dynamic examination. Raw report imitation can therefore introduce unsupported supervision, and reference-based metrics can reward non-grounded completion. CAMEO uses trusted visible evidence for cross-view training, preference alignment, and clinical error evaluation.

1 Introduction

Ultrasound is safe, relatively inexpensive, real-time, and widely used in routine care. Its interpretation is inherently multi-view: clinicians integrate scanning planes, probe-dependent appearances, and complementary key frames before writing a report. Multi-image ultrasound report generation is therefore not a collection of independent frame captions, but a problem of aggregating evidence across views and expressing it in clinically bounded language. A useful model must determine which observations are visible, how they relate across views, and how strongly they support report-level statements.

Recent MLLMs make this task increasingly feasible. Medical instruction tuning and large-scale biomedical image-text data improve domain coverage [3, 8, 17], while multi-image MLLMs support comparison, coreference, and cross-image aggregation [6, 10, 9]. However, input capacity and domain coverage alone do not guarantee reliable ultrasound reports. A model may produce fluent report text from modality priors or common templates while failing to preserve the specific evidence visible in the archived images.

A central source of this reliability problem is partial visual observability. A clinical ultrasound report may incorporate dynamic scanning, unarchived views, measurements, and contextual decisions, whereas a model usually receives only archived key frames. Some report statements are visible in these images; others may be clinically valid for the full examination but unverifiable from the available image set. Direct raw-report imitation therefore turns unobserved examination content into apparent image-grounded supervision.

This mismatch creates the twofold misalignment illustrated in Fig. 1. During training, unverifiable report points can become positive visual targets, encouraging completion from language priors. During evaluation, lexical metrics may reward plausible but unsupported completion while penalizing evidence-faithful outputs that omit unverifiable reference text. This is not ordinary label noise: a reference report may be clinically correct for the full examination yet still be an inappropriate image-grounded target for the archived image pair.

The central challenge is not merely to remove unverifiable report content, but to define the desired model behavior under incomplete visual evidence. A reliable model should report

visible abnormalities, preserve view-specific attribution, avoid unsupported diagnostic escalation, qualify claims when evidence is insufficient, and avoid conservative templates that omit visible findings. These requirements make report generation a clinically aware alignment problem rather than a data-cleaning exercise: the model must be informative when evidence is present and restrained when it is absent.

We formulate the task as *clinically aware evidence-grounded generation under partial visual observability*. CAMEO distills trusted visible evidence from raw reports, converts it into cross-view SFT data and clinical-error-oriented preference pairs, and evaluates generations against the same evidence structure. Training proceeds through Stage I Ultrasound Domain Adaptation, Stage II Cross-View Evidence Grounding, and Stage III Clinically Aware Preference Alignment.

Our contributions are:

1. We formulate multi-image ultrasound report generation as clinically aware evidence-grounded generation under partial visual observability.
2. We introduce visual-evidence distillation, which decomposes raw reports into atomic points and retains visually supportable findings as positive image-grounded supervision.
3. We propose CAMEO, a stage-wise clinically aware alignment framework that combines cross-view SFT with clinical-error-oriented preference alignment to balance evidence attribution, informative reporting, and clinically bounded generation.
4. We introduce an evidence-grounded, LLM-assisted Clinical Error Evaluation protocol that measures whether generated reports are informative, diagnostically consistent, and bounded by trusted visible evidence.

2 Related Work

2.1 Radiology Report Generation and Grounded Medical Reporting

Radiology report generation commonly uses encoder-decoder models, memory modules, knowledge bases, or LLM adapters, as in R2Gen [4] and R2GenGPT [19]. These methods improve fluency and report-style generation, but typically treat the paired reference report as direct image-grounded supervision. This assumption is reasonable when most reference content is observable in the input images, but becomes fragile for archived ultrasound, where the report may summarize information from unobserved portions of the scan.

Grounded medical reporting further links report phrases to local visual evidence [16]. This direction is complementary to ours, but typically begins with report content assumed to be visually relevant. We ask an earlier question: is each raw-report statement supportable from the archived ultrasound image pair at all? Only statements that pass this verification are used for positive supervision, preference construction, or evidence-grounded evaluation.

2.2 Medical MLLMs and Ultrasound Intelligence

Medical MLLMs such as LLaVA-Med [8] and HuatuoGPT-Vision [9] improve biomedical visual instruction following, while ultrasound-specific models such as EchoVLM [17] address sonographic domain shift. These models provide useful priors for ultrasound language and image understanding, but domain adaptation alone does not specify how a model

should respond when the report contains information that cannot be verified from the available views.

KMVE [10] is the closest prior work on ultrasound report generation. It introduces US-Report and uses knowledge-guided alignment to reduce the image–report feature discrepancy. We use USReport as source data but pursue a different objective: rather than aligning images directly with raw reports, we reconstruct visually supportable evidence, build clinical-error-oriented preferences, and evaluate generations against trusted visible evidence.

2.3 Multi-image and Cross-view Reasoning

MANTIS [6], LLaVA-OneVision [2], and LLaVA-NeXT-Interleave [9] improve comparison and aggregation across images. MMIU [12] further demonstrates that multi-image understanding cannot be reduced to single-image VQA. Ultrasound reporting, however, requires more than generic comparison. A finding may be visible in one view, corroborated by both views, or unsupported by the available pair despite appearing in the raw report.

Cross-view reasoning therefore becomes a problem of clinical evidence attribution. The model must determine whether a statement is supported by the first image, the second image, both images, or neither, and express the result in appropriate report language. Stage II targets this requirement rather than generic multi-image conversation.

2.4 Clinical Evaluation and Preference Alignment

Clinical metrics such as CheXbert [13], RadGraph [8], RadCliQ [21], and GREEN [14] move beyond lexical overlap, but generally compare generated reports with reference reports. In our setting, those references may contain unverifiable statements, so even clinical reference matching can reward unsupported completion. We instead evaluate against trusted visible evidence and separate omissions, incorrect statements, and unsupported additions.

Preference optimization provides a direct mechanism for behavior alignment. DPO [15] avoids explicit reward modeling, and MMedPO [23] demonstrates the value of clinically meaningful negative examples. Our preferences target ultrasound-specific omissions, polarity errors, view mismatches, diagnostic overreach, unsupported templates, and conservative shortcuts, using the same clinical error axes as the evaluation.

3 Method

3.1 From Partial Observability to Clinically Aware Evidence Grounding

Let $X = \{x_1, \dots, x_N\}$ denote the archived ultrasound images and R the raw report. In our primary benchmark, $N = 2$. Unlike conventional report generation, which learns $f_\theta : X \rightarrow R$, we do not assume that every statement in R is visually verifiable from X . An unsupported statement may still be clinically valid for the full examination, but it should not serve as positive image-grounded supervision. This distinction frames partial observability as an evidence-allocation problem rather than an assertion that the original clinical report is incorrect.

We decompose R into atomic report points,

$$P = \{p_1, p_2, \dots, p_M\}, \quad (1)$$




Figure 2: Overview of CAMEO. Raw reports are decomposed into atomic clinical points and filtered into trusted visible evidence. The retained evidence supports grounded QA, report-style SFT, clinical preference pairs, and LLM-assisted Clinical Error Evaluation, aligning training and assessment within a shared evidence space.

where each p_i expresses one factual clinical statement. For each point, we estimate a four-way visibility label

$$v_i \in \{\text{visible}, \text{partial}, \text{unsupported}, \text{excluded}\}, \quad (2)$$

where partial denotes a partially visible point. We define the scorable set and its fully and partially visible subsets as

$$\begin{aligned} P_{\text{scorable}} &= \{p_i \in P \mid v_i \neq \text{excluded}\}, \\ E_{\text{vis}} &= \{p_i \in P_{\text{scorable}} \mid v_i = \text{visible}\}, \\ E_{\text{partial}} &= \{p_i \in P_{\text{scorable}} \mid v_i = \text{partial}\}. \end{aligned} \quad (3)$$

Each point also has a support source $s_i \in \{\text{image 1}, \text{image 2}, \text{both}, \text{none}\}$. Visible and partial points must identify at least one supporting view; $s_i = \text{none}$ is reserved for unsupported or excluded points. The trusted positive image-grounded evidence is E_{vis} , and coverage is

$$\text{coverage}(X, R) = \frac{|E_{\text{vis}}|}{|P_{\text{scorable}}|}, \quad (4)$$

with coverage set to zero when P_{scorable} is empty. Partial points are included in the denominator but not the numerator; they support qualified supervision rather than fully positive report targets. Unsupported points remain scorable and motivate refusal or negative supervision, whereas excluded points enter neither the denominator nor downstream supervision. Coverage is a diagnostic signal rather than the task objective: low coverage indicates that report-level imitation is visually underdetermined, not that the case is uninformative. The reported training configuration does not discard cases solely because of low coverage. Coverage is retained as metadata for auditing this supervision; the complete policy is given in the supplementary material.

3.2 Multi-source Ultrasound Data Organization

Each data source has a distinct role. PUBMEDVISION-US [9] provides broad ultrasound instruction data: its single-image subset supports Stage I, and its multi-image subset sup-

ports Stage II. USReport, released with KMVE [14], provides paired-image reports for evidence distillation and preference construction. From USReport, we construct USREPORT-DISTILLED with 5,890 training cases, 11,780 images, 17,670 evidence-grounded instances, and 1,474 test cases. We also construct USREPORT-PREF with 21,869 training and 1,354 validation preference pairs. Thus, PUBMEDVISION-US supplies broad ultrasound instruction coverage, while the two derived datasets provide CAMEO’s evidence and behavior supervision.

3.3 Trustworthy Visual Evidence Distillation

Atomic report point decomposition. We split each raw report into atomic points, each expressing one clinical fact, such as lesion presence, location, size tendency, echogenicity, boundary, wall appearance, polarity, or an assessable negative finding. Multi-organ, multi-lesion, and multi-attribute sentences are separated before verification so that a visible clause is not discarded merely because it appears alongside an unverifiable one. Negative findings are retained only when they are visually assessable from the archived views; otherwise, absence statements are treated as unsupported rather than as negative visual evidence.

Debate-style visibility verification. Two independent verifiers assign each point a visibility label, supporting view, visual cue, reason for non-support, and source span. The labels are *visible*, *partially visible*, *unsupported*, and *excluded*; support sources are image 1, image 2, both, or neither. An arbiter resolves disagreements over visibility, support source, and clinical interpretation. Only visible points become positive image-grounded supervision, whereas partially visible points support qualified responses. This procedure is scalable weak-supervision purification, not radiologist annotation.

Evidence reconstruction. Visible and partial points are reconstructed into a trusted evidence template containing key findings, optional visible findings, qualified partial findings, unsupported or excluded statements, support-image metadata, and coverage. The same template is used to generate SFT targets, construct preference contrasts, and evaluate model outputs. This shared representation keeps training and clinical evaluation anchored to a consistent definition of the available evidence.

Radiologist audit of evidence selection. LLM-generated supervision can fail silently: a fluent rationale need not reflect what is actually visible. We therefore assign distinct roles to the teacher models. Disagreements are resolved by the arbiter, after which schema validation and aggregation are deterministic. The resulting inclusion decisions are then subjected to blinded radiologist review. The reader saw only archived images and anonymized atomic points, without teacher labels or evidence templates. Across 301 annotations, the binary teacher labels achieved 78.9% accuracy and 84.8% F1 when the reader’s “supported” and “partially supported” judgments were treated as positive. The reader also marked 25.2% of scorable raw-report points as unsupported, directly quantifying the supervision noise introduced by raw-report imitation. The complete protocol and an independent output evaluation are provided in the supplementary material.

USR-D QA type	Question / target form	Stage-II purpose
Evidence QA	Is the report point visible in the paired images?	Filter out non-verifiable raw-report content.
View Attribution QA	Which view supports the finding: first view, second view, both, or neither?	Learn view-conditioned attribution and reduce cross-view confusion.
Cross-view QA	How do the two views jointly support or limit the finding?	Train reasoning over complementary views rather than independent single-image captioning.
Refusal QA	Should an unsupported claim be answered, qualified, or refused?	Teach conservative responses when the image pair is insufficient.
Grounded Report QA	Produce a concise report-style summary from trusted visible evidence.	Convert verified evidence into clinically readable report supervision.

Table 1: SFT supervision reconstructed from USREPORT-DISTILLED. Each QA type targets a distinct cross-view reporting behavior.

3.4 Dense Evidence-grounded Supervision Construction

We convert E_{vis} and E_{partial} into two complementary forms of supervision: fully visible points provide positive targets, while partial points provide explicitly qualified targets. SFT teaches multi-image reasoning, view attribution, and report-style expression, while preference data penalizes clinically plausible but unsafe behavior that likelihood training does not directly discourage. The model must learn both what the visible evidence supports and when fluent report completion would exceed that evidence.

SFT supervision from visible evidence. The SFT branch uses multi-image supervision from USREPORT-DISTILLED rather than imitating raw reports. The QA types in Table 1 target verification, attribution, cross-view synthesis, bounded refusal, and report-style generation. Verification separates learnable evidence from unverifiable text; attribution reduces view mismatch; synthesis teaches aggregation across complementary views; refusal teaches bounded generation when evidence is insufficient; and grounded-report QA converts trusted evidence into readable clinical language.

Preference data from clinical failure modes. The preference branch constructs (y^+, y^-) pairs from the same evidence template. Rejected responses represent clinically plausible failure modes rather than generic low-quality text. After SFT, fluent normal templates or common diagnostic phrases may improve lexical overlap while omitting visible evidence, reversing attributes, or conflating views. Conversely, excessive refusal may reduce hallucination at the cost of under-reporting visible findings. The preferences therefore favor responses that are both informative and evidence-bounded, rather than merely shorter, more cautious, or more template-like.

Chosen responses cover visible findings, preserve attribution, and qualify incomplete support. Rejected responses instantiate organ-specific errors such as echogenicity reversal, boundary or wall-status confusion, benign-versus-suspicious wording, incorrect view attribution, unsupported normal templates, and conservative shortcuts. We group these errors under the same report-level axes used in evaluation: missed diagnosis, misdiagnosis, and overdiagnosis (Table 2). The goal is evidence-bounded clinical informativeness, not shorter or more hesitant text.

Preference contrast	Clinical axis	Rejected behavior	Desired alignment
Omission	Missed diagnosis	A visible lesion, abnormality, or key descriptor is dropped.	Preserve clinically important visible evidence.
Unsupported content	Overdiagnosis	Template-like normal findings or organ status are added without visual support.	Suppress hallucinated and template-driven content.
Clinical attribute confusion	Misdiagnosis	Echogenicity, wall status, lesion boundary, or diagnostic tendency is reversed or confused.	Penalize fluent but clinically wrong statements.
View mismatch	Misdiagnosis	Findings are assigned to the wrong view or mixed across views.	Strengthen view-specific attribution and cross-view consistency.
Diagnostic overreach	Overdiagnosis	The conclusion is stronger than the paired images justify.	Encourage conservative language under partial observability.
Template shortcut	Missed diagnosis / Overdiagnosis	The response relies on fixed benign phrases or empty caution without reporting visible evidence.	Maintain clinical informativeness while staying evidence-grounded.

Table 2: Preference data from USREPORT-PREF. Rejected responses instantiate clinically meaningful failure modes; chosen responses remain informative and evidence-bounded.

3.5 Stage-wise Model Alignment

Reliable multi-image ultrasound reporting requires domain adaptation, cross-view evidence grounding, and clinical risk calibration. We learn these capabilities sequentially because they address distinct sources of error: visual-language domain mismatch, multi-view evidence attribution, and generation behavior under partial evidence. Following LLaVA-Med [8], we keep the visual encoder fixed. Stage I begins with projector-only alignment and then instruction-tunes the projector and LLM; Stages II and III continue from the resulting check-point and update both the projector and LLM.

Stage I: Ultrasound Domain Adaptation. Stage I establishes ultrasound visual-language primitives. Projector-only alignment first maps ultrasound visual features into the language space while keeping the visual encoder and LLM fixed; single-image ultrasound instruction tuning then updates both the projector and LLM. This stage exposes the model to sonographic textures, acoustic artifacts, echogenicity descriptors, and organ-specific terminology. It reduces domain mismatch but does not address multi-image reporting, because view attribution and partial observability are not yet modeled.

Stage II: Cross-View Evidence Grounding. Stage II moves from ultrasound recognition to multi-image evidence use. It teaches which view supports a statement, whether the views jointly support it, and when insufficient evidence requires qualification. Multi-image ultrasound instructions provide cross-view practice, while USREPORT-DISTILLED supplies report-style targets derived from trusted visible evidence. At this stage, CAMEO learns to preserve visible findings and descriptors rather than defaulting to normal templates. We update the projector and LLM and optimize

$$\mathcal{L}_{\text{SFT}} = - \sum_t \log \pi_{\theta}(y_t | X, q, y_{<t}). \quad (5)$$

Stage III: Clinically Aware Preference Alignment. Stage III calibrates clinical behavior beyond likelihood training. USREPORT-PREF contrasts preferred responses with alternatives containing unsupported completion, attribute or polarity errors, view mismatch, diagnostic overreach, and fixed-template shortcuts. This stage preserves the informativeness learned in Stage II while penalizing clinically plausible failures; it is behavior calibration rather than generic stylistic refinement. The aim is to reduce hallucinated or overly specific completion without driving the model toward empty refusals when visible evidence is present. We update the projector and LLM while keeping the reference model fixed. Given input (X, q) , preferred response y^+ , rejected response y^- , and reference model π_{ref} , we optimize DPO [15]

$$\mathcal{L}_{\text{DPO}} = -\log \sigma \left(\beta \left[\log \frac{\pi_{\theta}(y^+ | X, q)}{\pi_{\text{ref}}(y^+ | X, q)} - \log \frac{\pi_{\theta}(y^- | X, q)}{\pi_{\text{ref}}(y^- | X, q)} \right] \right). \quad (6)$$

This increases the relative likelihood of evidence-bounded responses over clinically plausible failure cases.

3.6 LLM-assisted Clinical Error Evaluation

Conventional text metrics do not distinguish omissions from incorrect statements or unsupported additions. We therefore compare generated reports with the trusted evidence template. Here, *missed diagnosis*, *misdiagnosis*, and *overdiagnosis* denote report-level textual errors rather than patient-level outcomes. The LLM judge emits structured semantic labels, from which all case-level scores are computed deterministically.

Let G denote trusted gold units, P_{clin} non-style predicted units, and $W_G = \sum_{g \in G} w_g$. Gold priorities use $w_g \in \{3, 2, 1\}$ for key, visible, and optional findings. The judge assigns $\ell_g \in \{0, 0.5, 1\}$ for matched, partially matched, and missed units. Wrong and unsupported-extra predictions receive subtype weights $w_{\text{wrong}}(p)$ and $w_{\text{extra}}(p)$ in $\{1, 2, 3\}$; the complete mapping is provided in the supplementary material. The implemented losses are

$$L_{\text{miss}} = \frac{\sum_{g \in G} w_g \ell_g}{W_G}, \quad L_{\text{miss}} = 0 \text{ if } W_G = 0, \quad (7)$$

$$L_{\text{mis}} = \frac{\sum_{p \in P_{\text{clin}}} w_{\text{wrong}}(p)/3}{\max(|P_{\text{clin}}|, 1)}, \quad L_{\text{over}} = \frac{\sum_{p \in P_{\text{clin}}} w_{\text{extra}}(p)/3}{\max(|P_{\text{clin}}|, 1)}. \quad (8)$$

Only units labeled wrong contribute to L_{mis} , and only units labeled extra contribute to L_{over} . If a unit is both wrong and unsupported, we count it primarily under L_{mis} . We report

$$\text{MissScore} = 100(1 - L_{\text{miss}}), \quad \text{MisdiagScore} = 100(1 - L_{\text{mis}}), \quad (9)$$

$$\text{OverdiagScore} = 100(1 - L_{\text{over}}), \quad (10)$$

$$\text{ClinicalScore} = 0.40 \text{MissScore} + 0.35 \text{MisdiagScore} + 0.25 \text{OverdiagScore}. \quad (11)$$

The aggregate weights prioritize omissions, followed by incorrect statements and unsupported completion. ClinicalScore is an expert-audited auxiliary proxy, not a clinical utility function; we report all three sub-scores to expose the underlying trade-offs.




Figure 3: Dataset coverage and supervision flow. PUBMEDVISION-US [8] provides broad ultrasound instruction data, while USReport [10] is distilled into USREPORT-DISTILLED and USREPORT-PREF for cross-view grounding, preference alignment, and clinical error evaluation.

4 Experiments

4.1 Experimental Setup

Datasets. Fig. 3 summarizes the role of each dataset. The PUBMEDVISION-US single- and multi-image test sets each contain 500 samples. USREPORT-DISTILLED serves as the primary evidence-grounded benchmark, with 1,474 test cases. Its training split covers Breast, Thyroid, and Liver and contains 5,890 cases and 11,780 images. USREPORT-PREF is used only for preference alignment and validation, never as a source of test references. This separation distinguishes broad ultrasound instruction coverage from the multi-image evidence and preference supervision used to train CAMEO.

Baselines and variants. We evaluate four external baselines. These are LLaVA-Med [8], Qwen3-VL-32B [10], Lingshu-32B [20], and EchoVLM [10]. Together, they cover medical instruction tuning, model scale, medical generalist pretraining, and ultrasound-specific modeling. The component analysis then isolates our evidence-grounded alignment using branches continued from Stage I, including Raw Report SFT and Distilled Report SFT. Dashes mark unavailable metrics, such as clinical scores for PUBMEDVISION-US Multi.

All systems receive the same case-level multi-image inputs whenever their interfaces support them, together with a semantically consistent report-generation instruction. English outputs are scored directly; non-English outputs are translated only for text metrics, with clinical content preserved. Clinical metrics are computed by structured evidence-unit matching under the same rubric for every system. For PUBMEDVISION-US Multi, EchoVLM was evaluated on the same 500 cases using matched English and Chinese prompts. Table 3 reports the 500 English-prompt cases, and the supplementary material provides the paired Chinese-prompt control.

Implementation details. CAMEO is initialized from LLaVA-Med and trained in three stages on two modified NVIDIA RTX 4090 GPUs, each with 48GB of memory. The vi-

Model	Type	Params	BLEU-1	R-1	R-L	METEOR	BERT	Miss	Misdiag	Overdiag	Clinical
<i>PubMedVision-US Multi</i>											
LLaVA-Med [8]	medical	7B	0.19	0.30	0.20	0.23	0.19	–	–	–	–
Qwen3-VL-32B [9]	general	32B	0.32	0.38	0.23	0.34	0.17	–	–	–	–
Lingshu-32B [10]	medical	32B	0.33	0.42	0.25	0.35	0.25	–	–	–	–
EchoVLM [11]	ultrasound	12B	0.12	0.19	0.13	0.14	0.08	–	–	–	–
CAMEO	proposed	7B	0.38	0.44	0.33	0.43	0.30	–	–	–	–
<i>USReport-Distilled</i>											
LLaVA-Med [8]	medical	7B	0.17	0.21	0.16	0.16	0.18	22.35	65.28	69.64	49.19
Qwen3-VL-32B [9]	general	32B	0.26	0.33	0.24	0.36	0.28	25.25	70.39	73.26	53.05
Lingshu-32B [10]	medical	32B	0.25	0.30	0.21	0.30	0.25	26.31	69.35	75.48	53.66
EchoVLM [11]	ultrasound	12B	0.25	0.28	0.21	0.27	0.20	29.28	77.41	64.87	55.02
CAMEO	proposed	7B	0.40	0.45	0.34	0.43	0.42	56.36	82.90	90.53	74.20

Table 3: Main results on PUBMEDVISION-US Multi [8] and USREPORT-DISTILLED, constructed from USReport [11]. R-1, R-L, and BERT denote ROUGE-1, ROUGE-L, and BERTScore. Clinical metrics are evaluated only on USREPORT-DISTILLED; the EchoVLM PUBMEDVISION-US row uses all 500 English-prompt cases.

sual encoder remains frozen. Stage I performs projector-only alignment followed by joint projector–LLM instruction tuning; Stages II and III then update the projector and LLM using SFT and DPO, respectively. The teacher/debate pipeline operates offline and uses images plus raw reports solely to construct supervision. At inference, the 7B student receives only the images. Detailed model roles and settings are provided in the supplementary material.

Metrics. We report BLEU-1 [14], ROUGE-1/L [15], METEOR [9], and BERTScore [12]. On USREPORT-DISTILLED, the references are distilled evidence-grounded targets rather than raw reports; these metrics therefore quantify agreement with the target style, not independent clinical utility. We additionally report the MissScore, MisdiagScore, OverdiagScore, and ClinicalScore defined in Section 3.6. Higher values are better for every metric.

4.2 Main Results on Multi-Image Ultrasound Report Generation

Table 3 evaluates two settings. PUBMEDVISION-US Multi measures transferable multi-image ultrasound generation using text metrics only, while USREPORT-DISTILLED is the primary evidence-grounded benchmark because it additionally supports clinical error metrics. On PUBMEDVISION-US Multi, CAMEO reaches 0.38 BLEU-1, 0.44 ROUGE-1, and 0.43 METEOR, outperforming the external baselines under this protocol. This auxiliary result suggests that task-specific cross-view training benefits multi-image ultrasound instruction following. It is not, however, a general claim about ultrasound model quality, because the benchmark does not support clinical error evaluation.

On USREPORT-DISTILLED, Qwen3-VL-32B and Lingshu-32B achieve competitive lexical scores among the external baselines, yet their ClinicalScores are 53.05 and 53.66, respectively, showing that lexical plausibility and clinical evidence grounding can diverge. The ultrasound-specific EchoVLM attains a higher ClinicalScore than the larger general baselines but still trails CAMEO in this report-generation setting. Relative to EchoVLM, CAMEO improves BLEU-1 from 0.25 to 0.40, ROUGE-1 from 0.28 to 0.45, METEOR from 0.27 to 0.43, and ClinicalScore from 55.02 to 74.20. Ultrasound-specific modeling alone cannot replace task-specific cross-view evidence supervision and clinical alignment.

Variant	Evidence	Cross-view QA	Preference	BLEU-1	ROUGE-1	Clinical	Main purpose
LLaVA-Med [8]	no	no	no	0.17	0.21	49.19	base model
Stage I	no	no	no	0.18	0.22	55.71	domain adaptation
Stage I + Raw Report SFT	no	partial	no	0.26	0.32	61.28	raw imitation
Stage I + Distilled Report SFT	yes	limited	no	0.36	0.41	63.01	purification
Stage I + Stage II	yes	yes	no	0.39	0.45	71.44	cross-view reasoning
CAMEO	yes	yes	yes	0.40	0.45	74.20	clinical alignment

Table 4: Component ablation study. Rows after Stage I continue from the Stage I checkpoint. Complete component metrics are provided in the supplementary material.

Independent radiologist evaluation. Automatic gains may reflect conformity to the evidence rubric rather than clinically meaningful report quality. We therefore asked one radiologist to blindly score four anonymized, randomly ordered outputs for each of 30 cases using only the images and reports. On a 1–5 scale, CAMEO scored 3.57 ± 1.41 , compared with 1.73 ± 0.98 for EchoVLM, 1.90 ± 0.99 for Lingshu-32B, and 2.73 ± 1.11 for Raw Report SFT. The corresponding CAMEO win/tie/loss counts were 23/5/2, 23/7/0, and 22/3/5. ClinicalScore correlated with the radiologist’s ratings of CAMEO (Spearman $\rho = 0.698$, permutation $p < 0.001$). In a separate 20-case preference audit, the radiologist selected the constructed chosen response in 17 cases (85%; exact binomial $p < 0.001$ against a 1/3 chance rate). Together, these audits show that gains persist without access to the evidence schema, while residual losses expose remaining LLM failures. Full protocols for all three expert audits are provided in the supplementary material.

4.3 Component Ablation Study

Table 4 isolates the role of each component. Stage I reduces ultrasound domain mismatch but remains insufficient without cross-view supervision. Raw Report SFT improves lexical overlap while retaining unverifiable content, reflecting the mismatch introduced by raw-report supervision. Distilled Report SFT improves target quality but lacks explicit paired-view attribution and consequently underperforms Stage I + Stage II. Stage II provides the largest gain by teaching cross-view evidence grounding. Stage III largely preserves text metrics while improving ClinicalScore, indicating that preference alignment primarily calibrates clinical behavior rather than lexical overlap.

4.4 Clinical Error Analysis


Table 5 breaks down the clinical scores by organ. Stage III raises the overall ClinicalScore by 2.76 points, with improvements along all three axes. Liver and thyroid improve consistently, suggesting that preference alignment reduces both incorrect statements and omissions when relevant structures are well represented in the archived views. Breast improves mainly in MisdiagScore; its essentially unchanged MissScore instead points to a remaining visual-recall bottleneck for subtle mammary findings. We therefore interpret the organ-level variation as an evidence-recognition limitation that preference alignment alone cannot resolve.

4.5 Qualitative Analysis and Limitations

Fig. 4 illustrates the gap between report fluency and evidence grounding. EchoVLM produces a normal-style report and Stage I + Stage II remains conservative, both omitting the

Organ	Miss			Misdiagnosis			Overdiagnosis			Clinical		
	SFT	DPO	Δ	SFT	DPO	Δ	SFT	DPO	Δ	SFT	DPO	Δ
Liver	61.52	66.11	+4.59	91.04	91.81	+0.77	92.44	92.58	+0.14	79.58	81.72	+2.14
Breast	58.58	58.55	-0.03	81.20	84.86	+3.66	88.26	88.27	+0.01	73.92	75.19	+1.27
Thyroid	37.68	44.43	+6.75	66.44	72.04	+5.60	89.97	90.75	+0.78	60.82	65.68	+4.86
All	52.59	56.36	+3.77	79.56	82.90	+3.34	90.22	90.53	+0.31	71.44	74.20	+2.76

Table 5: Organ-wise clinical error evaluation. SFT denotes Stage II and DPO denotes CAMEO. Breast corresponds to the original Mammary label.



View 1

View 2



Ultrasound demonstrates disordered glandular structure with uneven echoes. A well-defined, oval hypoechoic nodule containing a visible lymphatic hilum is identified in the left axilla.

EchoVLM:

Bilateral breast tissue shows coarse heterogeneous echogenicity, uniformly dilated ducts, and normal vascularity, with no focal lesion. No abnormally enlarged lymph nodes are seen in the bilateral axillae

Reports coarse breast tissue and duct dilation, but concludes that there is no focal lesion and no abnormal axillary lymph node.

Misses the visible axillary finding.

Stage I+II :

The images show bilateral breast tissue with slightly disordered glandular structure. No definite space-occupying lesions are identified in the visualized fields.

Captures disordered breast tissue, but states that no definite space-occupying lesion is identified in the visualized fields.

Conservative template shortcut.

CAMEO :

In the provided images, the left axilla (labeled L-AX) demonstrates a hypoechoic nodule with clear boundaries, regular morphology, and a visible central echogenic hilum. The surrounding glandular tissue appears disordered with uneven echoes."

Attributes the finding to the left axilla (L-AX), describes the hypoechoic nodule, clear boundary, regular morphology, and central echogenic hilum, while retaining the surrounding disordered tissue.

Aligned with visible evidence.

Figure 4: Qualitative comparison on a breast/axillary case. EchoVLM and Stage I + Stage II omit the visible left-axillary nodule through normal or conservative templates, whereas CAMEO attributes the finding to the L-AX view and preserves the key descriptors.

visible left-axillary nodule. CAMEO preserves the L-AX attribution and key descriptors without adding unsupported content. The example reflects the intended effect of Stage III: suppressing template shortcuts while preserving visible findings. It also remains consistent with Table 5, where subtle mammary findings are still recall-limited.

This case also clarifies what CAMEO does not optimize for: the target is neither shorter output nor universal caution. A normal template may reduce unsupported abnormal claims while omitting visible evidence; an over-complete report may be informative but contain unsupported statements. The desired behavior is to preserve report-level informativeness when it can be anchored to view-specific evidence. Accordingly, our preference construction penalizes both unsupported completion and conservative shortcuts, and the clinical evaluation reports omissions and over-completion separately.

The quantitative results support the same interpretation. Stage I improves the base model's ultrasound language prior, but its gains are limited because single-image instruction tuning does not teach how evidence is distributed across views. Stage II delivers the largest report-generation improvement by converting trusted visible evidence into cross-view supervision. Stage III then reshapes the error profile rather than simply increasing text overlap: ClinicalScore improves while the lexical metrics remain similar, indicating that preference

alignment primarily affects what the model includes, omits, or qualifies. This distinction matters in ultrasound reporting, where a small change in wording can alter clinical meaning through polarity, attribution, or diagnostic strength.

The baseline comparison further motivates evidence-grounded alignment. Strong general or medical VLMs can generate plausible report-like text, and ultrasound-specific pretraining supplies useful modality priors, but neither directly addresses the mismatch between raw reports and archived evidence. CAMEO makes this mismatch explicit throughout the pipeline: report points are filtered by visible support, SFT targets are tied to view-specific evidence, preference pairs encode clinically meaningful failures, and evaluation tests whether generated statements remain within the trusted evidence. Data construction, alignment, and evaluation are therefore treated as coupled design choices rather than independent components.

The radiologist analyses characterize, rather than eliminate, the limitations of current LLMs. They use a single reader and validate binary evidence inclusion, not four-class visibility, support-view attribution, or inter-reader reliability. Even CAMEO averages only 3.57 ± 1.41 on the 1–5 scale and loses to Raw Report SFT in 5 of 30 cases, revealing substantial case-level variation despite the aggregate improvement. The inputs are archived key frames rather than full dynamic examinations, and ClinicalScore remains an auxiliary proxy despite its correlation with reader ratings on the audited subset. The benchmark covers two-view report generation across three common organ categories, but not complete ultrasound protocols, dynamic scanning decisions, or rare findings. Our claims are therefore restricted to evidence-grounded report generation from archived images and do not extend to clinical deployment.

5 Conclusion

We introduced CAMEO, an evidence-grounded framework for multi-image ultrasound report generation under partial visual observability. It aligns visible evidence, supervision, generation, and evaluation through atomic-point distillation, cross-view tuning, and clinical-error-oriented preferences. Cross-View Evidence Grounding yields the largest gain, while preference alignment improves the clinical error profile.

The blinded radiologist studies support this design and expose its limits: raw reports contain image-unverifiable content, constructed preferences generally agree with expert judgment, and CAMEO improves report quality but retains case-specific failures. With one reader, archived key frames, three organ categories, and an auxiliary LLM-assisted metric, these results do not establish clinical utility.

Acknowledgements

This research was supported by the National Natural Science Foundation of China (grants 62203060 and 62403492).

References

- [1] Shuai Bai, Yuxuan Cai, Ruizhe Chen, Keqin Chen, Xionghui Chen, Zesen Cheng, Lianghao Deng, Wei Ding, Chang Gao, Chunjiang Ge, Wenbin Ge, Zhifang Guo, Qidong Huang, Jie Huang, Fei Huang, Binyuan Hui, Shutong Jiang, Zhaohai Li,

- Mingsheng Li, Mei Li, Kaixin Li, Zicheng Lin, Junyang Lin, Xuejing Liu, Jiawei Liu, Chenglong Liu, Yang Liu, Dayiheng Liu, Shixuan Liu, Dunjie Lu, Ruilin Luo, Chenxu Lv, Rui Men, Lingchen Meng, Xuancheng Ren, Xingzhang Ren, Sibao Song, Yuchong Sun, Jun Tang, Jianhong Tu, Jianqiang Wan, Peng Wang, Pengfei Wang, Qiuyue Wang, Yuxuan Wang, Tianbao Xie, Yiheng Xu, Haiyang Xu, Jin Xu, Zhibo Yang, Mingkun Yang, Jianxin Yang, An Yang, Bowen Yu, Fei Zhang, Hang Zhang, Xi Zhang, Bo Zheng, Humen Zhong, Jingren Zhou, Fan Zhou, Jing Zhou, Yuanzhi Zhu, and Ke Zhu. Qwen3-VL technical report. *arXiv preprint arXiv:2511.21631*, 2025. URL <https://arxiv.org/abs/2511.21631>.
- [2] Satanjeev Banerjee and Alon Lavie. METEOR: An automatic metric for MT evaluation with improved correlation with human judgments. In *Proceedings of the ACL Workshop on Intrinsic and Extrinsic Evaluation Measures for Machine Translation and/or Summarization*, pages 65–72, 2005. URL <https://aclanthology.org/W05-0909/>.
- [3] Junying Chen, Chi Gui, Ruyi Ouyang, Anningzhe Gao, Shunian Chen, Guiming Hardy Chen, Xidong Wang, Zhenyang Cai, Ke Ji, Xiang Wan, and Benyou Wang. HuatuoGPT-Vision, towards injecting medical visual knowledge into multi-modal LLMs at scale. In *Proceedings of the 2024 Conference on Empirical Methods in Natural Language Processing*, pages 7346–7370. Association for Computational Linguistics, 2024. doi: 10.18653/v1/2024.emnlp-main.418. URL <https://aclanthology.org/2024.emnlp-main.418/>.
- [4] Zhihong Chen, Yan Song, Tsung-Hui Chang, and Xiang Wan. Generating radiology reports via memory-driven transformer. In *Proceedings of the 2020 Conference on Empirical Methods in Natural Language Processing*, pages 1439–1449. Association for Computational Linguistics, 2020. doi: 10.18653/v1/2020.emnlp-main.112. URL <https://aclanthology.org/2020.emnlp-main.112/>.
- [5] Saahil Jain, Ashwin Agrawal, Adriel Saporta, Steven QH Truong, Du Nguyen Duong, Tan Bui, Pierre Chambon, Yuhao Zhang, Matthew P. Lungren, Andrew Y. Ng, Curtis P. Langlotz, and Pranav Rajpurkar. RadGraph: Extracting clinical entities and relations from radiology reports. In *Advances in Neural Information Processing Systems, Datasets and Benchmarks Track*, 2021. URL <https://arxiv.org/abs/2106.14463>.
- [6] Dongfu Jiang, Xuan He, Huaye Zeng, Cong Wei, Max Ku, Qian Liu, and Wenhua Chen. MANTIS: Interleaved multi-image instruction tuning. *Transactions on Machine Learning Research*, 2024. URL <https://openreview.net/forum?id=skLtdUVaJa>.
- [7] Bo Li, Yuanhan Zhang, Dong Guo, Renrui Zhang, Feng Li, Hao Zhang, Kaichen Zhang, Peiyuan Zhang, Yanwei Li, Ziwei Liu, and Chunyuan Li. LLaVA-OneVision: Easy visual task transfer. *arXiv preprint arXiv:2408.03326*, 2024. URL <https://arxiv.org/abs/2408.03326>.
- [8] Chunyuan Li, Cliff Wong, Sheng Zhang, Naoto Usuyama, Haotian Liu, Jianwei Yang, Tristan Naumann, Hoifung Poon, and Jianfeng Gao. LLaVA-Med: Training a large language-and-vision assistant for biomedicine in one day. In *Advances in*

- Neural Information Processing Systems*, volume 36, pages 28541–28564, 2023. URL https://proceedings.neurips.cc/paper_files/paper/2023/hash/5abcdf8ecdacba028c6662789194572-Abstract-Datasets_and_Benchmarks.html.
- [9] Feng Li, Renrui Zhang, Hao Zhang, Yuanhan Zhang, Bo Li, Wei Li, Zejun Ma, and Chunyuan Li. LLaVA-Interleave: Tackling multi-image, video, and 3d in large multi-modal models. In *International Conference on Learning Representations*, 2025. URL https://proceedings.iclr.cc/paper_files/paper/2025/hash/c9f95e9ec39fa5ad3d0a562b993b92aa-Abstract-Conference.html.
- [10] Jun Li, Tongkun Su, Baoliang Zhao, Faqin Lv, Qiong Wang, Nassir Navab, Ying Hu, and Zhongliang Jiang. Ultrasound report generation with cross-modality feature alignment via unsupervised guidance. *IEEE Transactions on Medical Imaging*, 44(1):19–30, 2024. doi: 10.1109/TMI.2024.3424978. URL <https://doi.org/10.1109/TMI.2024.3424978>.
- [11] Chin-Yew Lin. ROUGE: A package for automatic evaluation of summaries. In *Text Summarization Branches Out*, pages 74–81, 2004. URL <https://aclanthology.org/W04-1013/>.
- [12] Fanqing Meng, Jin Wang, Chuanhao Li, Quanfeng Lu, Hao Tian, Tianshuo Yang, Jiaqi Liao, Xizhou Zhu, Jifeng Dai, Yu Qiao, Ping Luo, Kaipeng Zhang, and Wenqi Shao. MMIU: Multimodal multi-image understanding for evaluating large vision-language models. In *International Conference on Learning Representations*, 2025. URL https://proceedings.iclr.cc/paper_files/paper/2025/hash/5f92032278b1c70946e0b753068de51e-Abstract-Conference.html.
- [13] Sophie Ostmeier, Justin Xu, Zhihong Chen, Maya Varma, Louis Blankemeier, Christian Bluethgen, Arne Edward Michalson, Michael Moseley, Curtis Langlotz, Akshay S. Chaudhari, and Jean-Benoit Delbrouck. GREEN: Generative radiology report evaluation and error notation. In *Findings of the Association for Computational Linguistics: EMNLP 2024*, pages 374–390. Association for Computational Linguistics, 2024. doi: 10.18653/v1/2024.findings-emnlp.21. URL <https://aclanthology.org/2024.findings-emnlp.21/>.
- [14] Kishore Papineni, Salim Roukos, Todd Ward, and Wei-Jing Zhu. BLEU: A method for automatic evaluation of machine translation. In *Proceedings of the 40th Annual Meeting of the Association for Computational Linguistics*, pages 311–318, 2002. doi: 10.3115/1073083.1073135. URL <https://aclanthology.org/P02-1040/>.
- [15] Rafael Rafailov, Archit Sharma, Eric Mitchell, Christopher D. Manning, Stefano Ermon, and Chelsea Finn. Direct preference optimization: Your language model is secretly a reward model. In *Advances in Neural Information Processing Systems*, volume 36, 2023. URL https://proceedings.neurips.cc/paper_files/paper/2023/hash/a85b405ed65c6477a4fe8302b5e06ce7-Abstract-Conference.html.
- [16] Sergio Sanchez Santisteban, Muhammad Awais, Yi-Zhe Song, and Josef Kittler. Enhancing radiology report generation: The impact of locally grounded vision and language training. In *35th British Machine Vision Conference 2024, BMVC 2024*. BMVA, 2024. URL <https://bmvc2024.org/proceedings/857/>.

- [17] Chaoyin She, Ruifang Lu, Lida Chen, Wei Wang, and Qinghua Huang. EchoVLM: Dynamic mixture-of-experts vision-language model for universal ultrasound intelligence. In *Proceedings of the 64th Annual Meeting of the Association for Computational Linguistics (Volume 1: Long Papers)*, pages 10800–10822. Association for Computational Linguistics, 2026. doi: 10.18653/v1/2026.acl-long.494. URL <https://aclanthology.org/2026.acl-long.494/>.
- [18] Akshay Smit, Saahil Jain, Pranav Rajpurkar, Anuj Pareek, Andrew Y. Ng, and Matthew P. Lungren. CheXbert: Combining automatic labelers and expert annotations for accurate radiology report labeling using BERT. In *Proceedings of the 2020 Conference on Empirical Methods in Natural Language Processing*, pages 1500–1519. Association for Computational Linguistics, 2020. doi: 10.18653/v1/2020.emnlp-main.117. URL <https://aclanthology.org/2020.emnlp-main.117/>.
- [19] Zhanyu Wang, Lingqiao Liu, Lei Wang, and Luping Zhou. R2GenGPT: Radiology report generation with frozen LLMs. *Meta-Radiology*, 1(3):100033, 2023. doi: 10.1016/j.metrad.2023.100033. URL <https://doi.org/10.1016/j.metrad.2023.100033>.
- [20] Weiwen Xu, Hou Pong Chan, Long Li, Mahani Aljunied, Ruifeng Yuan, Jianyu Wang, Chenghao Xiao, Guizhen Chen, Chaoqun Liu, Zhaodonghui Li, Yu Sun, Junao Shen, Chaojun Wang, Jie Tan, Deli Zhao, Tingyang Xu, Hao Zhang, and Yu Rong. Lingshu: A generalist foundation model for unified multimodal medical understanding and reasoning. *arXiv preprint arXiv:2506.07044*, 2025. URL <https://arxiv.org/abs/2506.07044>.
- [21] Feiyang Yu, Mark Endo, Rayan Krishnan, Ian Pan, Andy Tsai, Eduardo Pontes Reis, Eduardo Kaiser Ururahy Nunes Fonseca, Henrique Min Ho Lee, Zahra Shakeri Hossein Abad, Andrew Y. Ng, Curtis P. Langlotz, Vasantha Kumar Venugopal, and Pranav Rajpurkar. Evaluating progress in automatic chest x-ray radiology report generation. *Patterns*, 4(9):100802, 2023. doi: 10.1016/j.patter.2023.100802. URL <https://doi.org/10.1016/j.patter.2023.100802>.
- [22] Tianyi Zhang, Varsha Kishore, Felix Wu, Kilian Q. Weinberger, and Yoav Artzi. BERTScore: Evaluating text generation with BERT. *arXiv preprint arXiv:1904.09675*, 2019. URL <https://arxiv.org/abs/1904.09675>.
- [23] Kangyu Zhu, Peng Xia, Yun Li, Hongtu Zhu, Sheng Wang, and Huaxiu Yao. MMedPO: Aligning medical vision-language models with clinical-aware multimodal preference optimization. In *Proceedings of the 42nd International Conference on Machine Learning*, volume 267 of *Proceedings of Machine Learning Research*, pages 80207–80222. PMLR, 2025. URL <https://proceedings.mlr.press/v267/zhu25v.html>.