

Magnon-induced phononic Chern insulator

Rui-Chang Shen,¹ Yihao Yang,^{2,3,4,*} and Haoran Xue^{1,5,†}

¹*Department of Physics, The Chinese University of Hong Kong, Shatin, Hong Kong SAR, China*

²*State Key Laboratory of Extreme Photonics and Instrumentation, ZJU-Hangzhou Global Scientific and Technological Innovation Center, Zhejiang University, Hangzhou 310027, China.*

³*The Electromagnetics Academy at Zhejiang University, College of Information Science and Electronic Engineering, Zhejiang University, Hangzhou 310027, China.*

⁴*Zhejiang Key Lab. of Intelligent Electromagnetic Control and Advanced Electronic Integration, Jinhua Institute of Zhejiang University, Zhejiang University, Jinhua 321099, China.*

⁵*State Key Laboratory of Quantum Information Technologies and Materials, The Chinese University of Hong Kong, Shatin, Hong Kong SAR, China*

High-frequency artificial phononic crystals offer a low-loss platform compatible with on-chip integration, yet realizing Chern phononic phases at GHz frequencies remains challenging. Here, we propose a magnon-induced phononic Chern insulator in a honeycomb phononic crystal hybridized with ferromagnetic islands at the hexagon centers. A circularly polarized Kittel mode couples to the surrounding phonons with a phase winding, which breaks time-reversal symmetry and opens a full Chern gap. In the large-detuning regime, this mechanism leads to an effective Haldane-type phononic model with magnon-induced complex hopping. By tuning the magnon-phonon interaction, the full hybrid system accesses Chern phases with tunable Chern numbers $|C| = 1$ and $|C| = 2$. The predicted gaps can exceed realistic phonon and magnon linewidths, enabling their observation in GHz acoustic devices. Our work establishes chiral magnon-phonon hybridization as a route to magnetically reconfigurable topological phononics.

Introduction— Phonons, the quantized excitations of lattice vibrations, are fundamental carriers of thermal energy, mechanical signals, and coherent information [1]. The introduction of topological band theory into acoustic and elastic systems has given rise to topological phononics, enabling robust boundary transport and unconventional control of sound and mechanical waves [2–14]. Most experimental realizations, however, operate at kilohertz-to-megahertz frequencies, where the comparatively long acoustic wavelengths limit miniaturization and integration. Gigahertz (GHz) phonons, by contrast, combine compact wavelengths and low dissipation with compatibility with integrated electromechanical architectures [15, 16]. They can furthermore interface coherently with superconducting qubits, spin excitations, microwave-frequency resonators, and ferromagnetic dynamics [17–21], making them promising information carriers for hybrid quantum and classical devices. To date, GHz topological-phononics experiments have primarily realized valley-Hall, crystalline, or pumping-based phases [22–24]. Although such phases can suppress scattering under appropriate symmetry conditions, they cannot provide the strictly one-way boundary transport of a Chern insulator and thus remain vulnerable to symmetry-breaking disorders [25].

Realizing a phononic Chern phase requires breaking the time-reversal symmetry and opening a complete bulk gap. Existing approaches have employed gyroscopic motion [2, 8], circulating fluids [4–6, 26, 27], Floquet modulation [28, 29], and charge-doped piezomagnetic platforms [30]. These schemes have established the basic principles of acoustic Chern transport, but their reliance on moving media, continuous external driving,

or material-specific responses presents substantial challenges for miniaturized high-frequency devices.

Magnons, the collective excitations of spin precession in ordered magnets, provide a natural microscopic source of time-reversal symmetry breaking. Their intrinsic GHz-frequency dynamics and magnetic tunability have made them important carriers of coherent information [31–33]. Magnons couple to lattice vibrations through effects like magnetoelastic and magneto-rotation interactions [34, 35], and suitable symmetry breaking can convert these interactions into nonreciprocal acoustic responses [35–38].

Here, we propose a magnon-induced phononic Chern insulator in a honeycomb phononic crystal containing ferromagnetic islands. Unlike previous studies of topological magnon-phonon polaritons [39–43], our scheme incorporates magnon-phonon coupling into an artificial phononic-crystal platform, where magnons not merely serve as a constituent of a hybrid excitation, but as virtual mediators that imprint a synthetic gauge flux on a lattice and thereby produce a globally gapped phononic Chern phase. Specifically, due to the discrete rotational symmetry, the magnon couples to the surrounding phonons with a phase winding. This induces virtual complex phonon hoppings and an effective Haldane-type flux, opening a complete Chern gap with a tunable Chern number up to two. We also present the phase diagram of the model and analyse the gap size using realistic parameters, further paving the way for possible experiments.

Human-AI discovery workflow— The theoretical framework was developed through an AI scientific-discovery workflow centered on the Qiushi Discovery Engine [44], as illustrated in Fig. 1(a). Starting from




FIG. 1. (a) Schematic of Human-Agent hybrid workflow. (b) A honeycomb phononic crystal with one phonon mode on each A and B sublattice site. A magnetic island at the plaquette center supports a right-handed circular Kittel mode. The six surrounding phononic sites are labeled in their physical counterclockwise order as $A_1, B_1, A_2, B_2, A_3, B_3$. Their radial vectors are $\mathbf{d}_{\alpha,i} = a_0(\cos \theta_{\alpha,i}, \sin \theta_{\alpha,i})$, with $\theta_{A,i} = 2\pi(i-1)/3$, $\theta_{B,i} = (2i-1)\pi/3$, and a_0 is the distance from the plaquette center to each surrounding phononic site. (c) Momentum-space intensity of the magnon–phonon terms $|G_A(\mathbf{k})|^2$ and $|G_B(\mathbf{k})|^2$, showing that the magnon couples selectively to sublattice A at K and to sublattice B at K'. (d) Schematic of the magnon-induced effective phonon hopping, showing $H_{\text{eff}} - H_{\text{ph}}$, where only the magnon-induced coupling between phonons is depicted. The coupling phase is illustrated in the right panel. Each NN bond receives virtual-magnon contributions from the center magnons of its two adjacent plaquettes.

the open-ended objective of using magnons to realize a phononic Chern insulator, the engine autonomously constructed the initial lattice model, derived its effective Hamiltonian, explored candidate topological mechanisms, and performed closed-loop consistency checks and self-verification. Human researchers subsequently examined the physical assumptions, independently verified the analytical and numerical results, and formulated the final interpretation. This human–AI workflow therefore combined autonomous theoretical exploration with human scientific validation.

Phonon-magnon model— We consider a honeycomb phononic crystal with one localized phonon mode on each A and B sublattice [Fig. 1(b)]. A ferromagnetic island at each plaquette center supports a uniform Kittel mode [45–47]. The system is governed by the Hamiltonian

$$H = H_{\text{ph}} + H_{\text{m}} + H_{\text{int}}. \quad (1)$$

Here, H_{ph} is the phonon Hamiltonian given by

$$H_{\text{ph}} = \sum_{r_A} \omega_{\text{ph}} a_{r_A}^\dagger a_{r_A} + \sum_{r_B} \omega_{\text{ph}} b_{r_B}^\dagger b_{r_B} + \sum_{\langle r_A, r_B \rangle} t (a_{r_A}^\dagger b_{r_B} + \text{H.c.}), \quad (2)$$

where $a_{r_{A/B}}^\dagger$ ($a_{r_{A/B}}$) and $b_{r_{A/B}}^\dagger$ ($b_{r_{A/B}}$) create (annihilate) a phonon mode at position r_A and r_B , respectively, ω_{ph}

is the phonon frequency, t is the nearest-neighbor phonon hopping, and H.c. stands for the Hermitian conjugate. Due to the large spatial separation between the ferromagnetic islands, the magnon Hamiltonian H_{m} only consists of on-site resonances:

$$H_{\text{m}} = \sum_R \omega_{\text{m}} m_R^\dagger m_R, \quad (3)$$

where $m_R^\dagger$ (m_R) are magnon creation (annihilation) operators at position R and ω_{m} is the magnon frequency. Without phonon-magnon interaction, the phononic crystal behaves like a phononic graphene with two Dirac points at the Brillouin corners (i.e., K and K' valleys), whereas the magnon dispersion is a flatband. This setup has the advantage that, once the Dirac points are properly gapped, a clean and full Chern gap can be easily obtained without further parameter optimization, and a low-energy description can directly capture the underlying mechanism.

The interaction term takes the form:

$$H_{\text{int}} = \sum_R \sum_{i=1}^3 [g \xi^{2(i-1)} a_{R+\mathbf{d}_{A,i}}^\dagger m_R + g \xi^{2i-1} b_{R+\mathbf{d}_{B,i}}^\dagger m_R + \text{H.c.}]. \quad (4)$$

Here, g is the magnitude of the magnon–phonon coupling and $\xi = e^{-i\pi/3}$ is the phase increment between




FIG. 2. (a) Bulk band structures in the absence (gray dashed curves) and presence (colored solid curves) of the magnon-phonon coupling, which explicitly opens a gap at the Dirac points. The lower, middle and upper bands have the Chern numbers $C = (-1, +1, 0)$, respectively. (b) The corresponding strip spectrum for $g = 0.4$, with periodic boundary conditions applied along $\mathbf{a}_1 = a_0(3, \sqrt{3})/2$. (c) Spatial distribution of the edge states, calculated as the sum of the probability of all localized modes within the bandgap. Parameters are set as $t = 1$ and $\delta_\omega = 3.2$.

adjacent radial directions. This phase factor originates from projecting the right-circular Kittel mode onto the radial effective fields associated with the six surrounding phononic sites (see Supplemental Material (SM), Sec. S1 for more details on the model construction [48]). The A and B sites are labeled in their physical counterclockwise order, with $\theta_{A,i} = 2\pi(i-1)/3$ and $\theta_{B,i} = (2i-1)\pi/3$. For a fixed magnetization direction, these complex coupling phases generate the synthetic flux responsible for the Chern gap.

In the following, we consider a generic large-detuning scenario where $|\delta_\omega| \equiv |\omega_m - \omega_{ph}| \gg g, t$. In such a case, the magnon serves as an auxiliary degree of freedom to manipulate the phonon band topology, and the resulting topological transport remains phonon-dominated, as desired for phononic applications. In the Bloch basis $\Psi_{\mathbf{k}} = (a_{\mathbf{k}A}, a_{\mathbf{k}B}, m_{\mathbf{k}})^T$, the model Hamiltonian is [48]

$$H(\mathbf{k}) = \begin{pmatrix} 0 & tf(\mathbf{k}) & G_A(\mathbf{k}) \\ tf^*(\mathbf{k}) & 0 & G_B(\mathbf{k}) \\ G_A^*(\mathbf{k}) & G_B^*(\mathbf{k}) & \delta_\omega \end{pmatrix}, \quad (5)$$

where $\mathbf{k} = (k_x, k_y)$ is the momentum and $f(\mathbf{k}) = \sum_{i=1}^3 e^{i\mathbf{k} \cdot \mathbf{d}_{A,i}}$. The magnon-phonon coupling terms are given by

$$G_A(\mathbf{k}) = g \sum_{i=1}^3 \xi^{2(i-1)} e^{-i\mathbf{k} \cdot \mathbf{d}_{A,i}}, \quad (6)$$

$$G_B(\mathbf{k}) = g \sum_{i=1}^3 \xi^{2i-1} e^{-i\mathbf{k} \cdot \mathbf{d}_{B,i}}.$$

At the two Dirac points, the phases sum constructively on one sublattice and destructively on the other: $|G_A(\mathbf{K})|^2 = 9g^2$, $|G_B(\mathbf{K})|^2 = 0$; and $|G_A(\mathbf{K}')|^2 = 0$, $|G_B(\mathbf{K}')|^2 = 9g^2$. Consequently, the magnon couples exclusively to sublattice A at K and to sublattice B at K', as plotted in Fig. 1(c). This valley-sublattice locking generates opposite Dirac masses at the two val-

leys ($m_{K/K'} = \mp 9g^2/2\delta_\omega$), ensuring that their Berry-curvature contributions constructively add up to open a topological Chern gap (see SM Sec. S2 [48]). Interestingly, this mechanism closely mimics the gap-opening process in off-resonantly driven graphene, where photon-assisted transitions similarly generate a valley-opposite Dirac mass [49–52].

Effective Haldane-like model— To gain a deeper insight into the role of the central magnon on the phononic crystal, we apply the Schrieffer-Wolff transformation to derive the effective phonon Hamiltonian as [53, 54]:

$$H_{\text{eff}} = H_{\text{ph}} - \sum_R \frac{V_R^\dagger V_R}{\delta_\omega}, \quad (7)$$

where $V_R = \sum_{i=1}^3 [g\xi^{-2(i-1)}a_{R+\mathbf{d}_{A,i}} + g\xi^{-(2i-1)}b_{R+\mathbf{d}_{B,i}}]$. The magnon-phonon interaction generates four types of effective terms [see Fig. 1(d)]: an on-site energy renormalization, nearest-neighbor (NN) hoppings, next-nearest-neighbor (NNN) hoppings, and third-nearest-neighbor (NNNN) hoppings. Specifically, the effective hopping amplitudes between two sites of the same sublattice around the same magnon center are

$$t_{A_i \leftarrow A_j}^{\text{eff}} = -\frac{g^2}{\delta_\omega} \xi^{2(i-j)}, \text{ and } t_{B_i \leftarrow B_j}^{\text{eff}} = -\frac{g^2}{\delta_\omega} \xi^{2(i-j)}, \quad (8)$$

where i, j label phonon sites surrounding the plaquette. For $i = j$, these expressions reduce to real on-site energy shifts $-g^2/\delta_\omega$; for $i \neq j$, they yield complex NNN hoppings on the two sublattices. This is precisely the Haldane-type flux pattern that generates a nonzero Chern number without an external magnetic field [55].

In addition, a magnon-mediated hopping from a B_j site to an A_i site carries the phase associated with the difference between their physical radial angles. It is given by

$$t_{A_i \leftarrow B_j}^{\text{eff}} = -\frac{g^2}{\delta_\omega} \xi^{2(i-j)-1}. \quad (9)$$




FIG. 3. (a) and (b) Chern-number phase diagrams of the lowest band and the middle band, respectively, as functions of the magnon detuning δ_ω and the magnon–phonon coupling strength g , with $t = 1$. The dashed curve $4g^2 = t^2 + t\delta_\omega$, marks the gap-closing condition at the M point, while the vertical dashed line in (b) indicates the Γ -point transition at $\delta_\omega = 3t$. (c) and (d) Plots of bulk bandgaps Δ_1 (gap between the lower and middle bands) and Δ_2 (gap between the middle and upper bands), respectively, as functions of δ_ω and g .

This cross-sublattice term includes both NN and NNNN contributions, depending on the relative positions of the two sites.

Topological bandgap and boundary modes— To elucidate the emergence of the topological phase, we investigate the bulk band structure of the full coupled system. At $g = 0$ the two phononic bands host gapless Dirac points at the K and K' valleys, while the magnon band is decoupled [Fig. 2(a), dashed curves]. The chiral magnon–phonon coupling hybridizes these modes and opens a bandgap at the phonon Dirac points. Through a numerical calculation of the Chern number [56], we find $C = \mp 1$ for the two phonon-dominated bands and $C = 0$ for the magnon-dominated bands [Fig. 2(a), solid curves], consistent with previous analysis.

The nontrivial bulk topology implies the existence of chiral edge states via the bulk–boundary correspondence [57]. For a strip geometry, Fig. 2(b) shows the edge modes traversing the bulk gap, strictly localized at opposite boundaries and propagating unidirectionally. Figure 2(c) further visualizes the boundary localization through the spatial intensity obtained by summing the probability of all in-gap states on a finite lattice. Reversing the magnetization direction ($+z \rightarrow -z$) complex-conjugates the local coupling phases ($\xi \rightarrow \xi^*$), reversing the effective synthetic flux and changing the lowest-band Chern number from -1 to $+1$. Accordingly, the chiral edge states reverse their propagation direction.

Phase diagram and gap visibility— Finally, we work out the full phase diagram of the model by going beyond the large detuning limit and also allowing for strong magnon–phonon interaction. Figures 3(a) and (b) show the evolution of the lower- and middle-band Chern numbers as functions of δ_ω and g , with the dashed line denoting analytic phase boundaries. For the lower band, the Chern number changes from -1 to $+2$ at $4g^2 = t^2 + t\delta_\omega$, suggesting a large Chern number phase with multiple chiral edge states can be obtained via increasing the

magnon–phonon hopping. In the large detuning limit, this phase transition can be inferred from the effective NNNN hopping [Fig. 1(d)]. For the middle band, an additional phase transition occurs at $\delta_\omega = 3t$, making the phase diagram much richer than the lowest band. Notably, when the sum of the Chern numbers of the lowest two bands is nonzero, the second bandgap will also host a chiral edge state [48].

Apart from the Chern numbers, gap size is also a crucial quantity that determines the performance of the system. Under strong detuning and weak coupling, the first bandgap Δ_1 can be enlarged by reducing the frequency detuning δ_ω or increasing the coupling g [Fig. 3(c)], consistent with the low-energy theory (i.e., $\Delta_1 \propto g^2/\delta_\omega$). Away from this regime, the gap size is jointly determined by valley and M point modes, thus showing a nonmonotonic behavior. For the second bandgap, its size Δ_2 vanishes along $\delta_\omega = 3t$, which corresponds to the band touching between the middle and upper bands. The gap size generally grows away from the closing points.

A realistic GHz phononic-crystal platform can be implemented using SAW devices or suspended elastic crystals on LiNbO₃, AlN, or Si, with nearest-neighbor hopping rates of $t/2\pi \sim 10$ – 100 MHz and phonon linewidths approaching $\gamma_{\text{ph}}/2\pi \sim 1$ MHz [22, 58–62]. A CoFeB or YIG island placed at each plaquette center provides a Kittel mode whose frequency can be tuned by an external magnetic field, thereby enabling control of the magnon–phonon detuning δ_ω , while magnetoelastic and magnetorotational interactions generate the required coupling [34, 35, 37, 63]. CoFeB can offer relatively strong coupling, $g/2\pi \sim 10$ – 30 MHz [64], but its larger Gilbert damping can lead to magnon linewidths of $\gamma_{\text{m}}/2\pi \sim 10$ – 100 MHz [65]. YIG provides a favorable low-loss alternative, with $g/2\pi \sim 5$ – 10 MHz and magnon linewidths near $\gamma_{\text{m}}/2\pi \sim 1$ MHz [66]. For representative parameters $t/2\pi = 30$ MHz, $\delta_\omega = 3.2t$, and $g/2\pi = 7$ MHz, the estimated topological gap is approximately 5 MHz, exceed-

ing the expected phonon and YIG magnon linewidths and therefore allowing clear spectral resolution of the gap, which can be further enhanced by reducing the frequency detuning.

Conclusion— We have proposed a magnon-induced phononic Chern insulator in a graphene-type phononic crystal system. The key to the design is the placement of the phonon and magnon sites, which generates favored time-reversal-breaking hoppings and opens a full topological gap. The model is studied analytically by analyzing the full and reduced phonon Hamiltonians, which clearly reveal the core function of the magnon-phonon interaction. We also investigate the model numerically, showing the existence of the chiral edge states and the fruitful phase diagram. More broadly, this work exemplifies an emerging human-AI discovery workflow in which the Qishu Discovery Engine constructed the initial model and carried out the subsequent theoretical exploration and numerical checks, the authors then examined its physical assumptions, verified the results, and formulated the final interpretation. These results demonstrate the power of using discrete lattices to engineer magnon-phonon interactions and establish magnons as tunable mediators of synthetic gauge fields for reconfigurable GHz phonon transport.

Acknowledgements— This work was supported by the National Natural Science Foundation of China under Grant Nos. 62401491 (H.X.) 62175215 (Y.Y.), and U25D8017 (Y.Y.), the National Key Research and Development Program of China under Grant Nos. 2022YFA1405200 (Y.Y.), and 2022YFA1404900 (Y.Y.), the Class D for Young Scientific Research Peak Creation No. K20250230 (Y.Y.), the Fundamental Research Funds for the Zhejiang Provincial Universities No. 226-2025-00231 (Y.Y.), the Science Challenge Project NO. TZ2025015 (Y.Y.), the Research Grants Council of the Hong Kong SAR, China, under Grant No. 24304825 (H.X.), the Guangdong Provincial Quantum Science Strategic Initiative under Grant No. GDZX2501012 (H.X.), and the Chinese University of Hong Kong under Grant Nos. 4053729 (H.X.) and 4053794 (H.X.).

* yangyihao@zju.edu.cn

† haoranxue@cuhk.edu.hk

- [1] M. Maldovan, Sound and heat revolutions in phononics, *Nature* **503**, 209 (2013).
- [2] P. Wang, L. Lu, and K. Bertoldi, Topological phononic crystals with one-way elastic edge waves, *Phys. Rev. Lett.* **115**, 104302 (2015).
- [3] S. H. Mousavi, A. B. Khanikaev, and Z. Wang, Topologically protected elastic waves in phononic metamaterials, *Nat. Commun.* **6**, 8682 (2015).
- [4] Z. Yang, F. Gao, X. Shi, X. Lin, Z. Gao, Y. Chong, and B. Zhang, Topological acoustics, *Phys. Rev. Lett.* **114**, 114301 (2015).
- [5] A. B. Khanikaev, R. Fleury, S. H. Mousavi, and A. Alu, Topologically robust sound propagation in an angular-momentum-biased graphene-like resonator lattice, *Nat. Commun.* **6**, 8260 (2015).
- [6] X. Ni, C. He, X.-C. Sun, X.-p. Liu, M.-H. Lu, L. Feng, and Y.-F. Chen, Topologically protected one-way edge mode in networks of acoustic resonators with circulating air flow, *New J. Phys.* **17**, 053016 (2015).
- [7] R. Süssstrunk and S. D. Huber, Observation of phononic helical edge states in a mechanical topological insulator, *Science* **349**, 47 (2015).
- [8] L. M. Nash, D. Kleckner, A. Read, V. Vitelli, A. M. Turner, and W. T. Irvine, Topological mechanics of gyroscopic metamaterials, *Proc. Natl. Acad. Sci. USA* **112**, 14495 (2015).
- [9] C. L. Kane and T. C. Lubensky, Topological boundary modes in isostatic lattices, *Nat. Phys.* **10**, 39 (2014).
- [10] M. Xiao, G. Ma, Z. Yang, P. Sheng, Z. Zhang, and C. T. Chan, Geometric phase and band inversion in periodic acoustic systems, *Nat. Phys.* **11**, 240 (2015).
- [11] C. He, X. Ni, H. Ge, X.-C. Sun, Y.-B. Chen, M.-H. Lu, X.-P. Liu, and Y.-F. Chen, Acoustic topological insulator and robust one-way sound transport, *Nat. Phys.* **12**, 1124 (2016).
- [12] M. Yan, J. Lu, F. Li, W. Deng, X. Huang, J. Ma, and Z. Liu, On-chip valley topological materials for elastic wave manipulation, *Nat. Mater.* **17**, 993 (2018).
- [13] G. Ma, M. Xiao, and C. T. Chan, Topological phases in acoustic and mechanical systems, *Nat. Rev. Phys.* **1**, 281 (2019).
- [14] H. Xue, Y. Yang, and B. Zhang, Topological acoustics, *Nat. Rev. Mater.* **7**, 974 (2022).
- [15] D. Hatanaka, I. Mahboob, K. Onomitsu, and H. Yamaguchi, Phonon waveguides for electromechanical circuits, *Nat. Nanotechnol.* **9**, 520 (2014).
- [16] P. Delsing, A. N. Cleland, M. J. Schuetz, J. Knörzer, G. Giedke, J. I. Cirac, K. Srinivasan, M. Wu, K. C. Balram, C. Bäuerle, *et al.*, The 2019 surface acoustic waves roadmap, *J. Phys. D: Appl. Phys.* **52**, 353001 (2019).
- [17] M. V. Gustafsson, T. Aref, A. F. Kockum, M. K. Ekström, G. Johansson, and P. Delsing, Propagating phonons coupled to an artificial atom, *Science* **346**, 207 (2014).
- [18] Y. Chu, P. Kharel, W. H. Renninger, L. D. Burkhardt, L. Frunzio, and R. J. Schoelkopf, Quantum acoustics with superconducting qubits, *Science* **358**, 199 (2017).
- [19] K. J. Satzinger, Y. Zhong, H.-S. Chang, G. A. Peairs, A. Bienfait, M.-H. Chou, A. Y. Cleland, C. R. Conner, É. Dumur, J. Grebel, I. Gutierrez, B. H. November, R. G. Povey, S. J. Whiteley, D. D. Awschalom, D. I. Schuster, and A. N. Cleland, Quantum control of surface acoustic-wave phonons, *Nature* **563**, 661 (2018).
- [20] A. H. Safavi-Naeini, D. Van Thourhout, R. Baets, and R. Van Laer, Controlling phonons and photons at the wavelength scale: integrated photonics meets integrated phononics, *Optica* **6**, 213 (2019).
- [21] M. Weiler, H. Huebl, F. S. Goerg, F. D. Czeschka, R. Gross, and S. T. B. Goennenwein, Elastically driven ferromagnetic resonance in nickel thin films, *Phys. Rev. Lett.* **106**, 117601 (2011).
- [22] Q. Zhang, D. Lee, L. Zheng, X. Ma, S. I. Meyer, L. He, H. Ye, Z. Gong, B. Zhen, K. Lai, and A. T. C. Johnson, Gigahertz topological valley hall effect in nanoelectromechanical phononic crystals, *Nat. Electron.* **5**, 157 (2022).

- [23] Z.-D. Zhang, S.-Y. Yu, M.-H. Lu, and Y.-F. Chen, Near ghz lithium niobate higher-order topological nanomechanical metamaterials, *Nano Lett.* **24**, 15421 (2024).
- [24] X.-B. Xu, M. Oudich, Y. Zeng, J.-Z. Zhang, Y.-H. Yang, J.-Q. Wang, W. Wang, L. Sun, G.-C. Guo, Y. Jing, and C.-L. Zou, Gigahertz topological phononic circuits based on micrometre-scale unsuspended waveguide arrays, *Nat. Electron.* **8**, 689 (2025).
- [25] D. Leykam, H. Xue, B. Zhang, and Y. Chong, Limitations and possibilities of topological photonics, *Nat. Rev. Phys.* **8**, 55 (2026).
- [26] A. Souslov, B. C. Van Zuiden, D. Bartolo, and V. Vitelli, Topological sound in active-liquid metamaterials, *Nat. Phys.* **13**, 1091 (2017).
- [27] Y. Ding, Y. Peng, Y. Zhu, X. Fan, J. Yang, B. Liang, X. Zhu, X. Wan, and J. Cheng, Experimental demonstration of acoustic chern insulators, *Phys. Rev. Lett.* **122**, 014302 (2019).
- [28] R. Fleury, A. B. Khanikaev, and A. Alu, Floquet topological insulators for sound, *Nat. Commun.* **7**, 11744 (2016).
- [29] A. Darabi, X. Ni, M. Leamy, and A. Alù, Reconfigurable floquet elastodynamic topological insulator based on synthetic angular momentum bias, *Sci. Adv.* **6**, eaba8656 (2020).
- [30] Q. Zhang, L. He, E. J. Mele, B. Zhen, and A. C. Johnson, General duality and magnet-free passive phononic chern insulators, *Nat. Commun.* **14**, 916 (2023).
- [31] P. Pirro, V. I. Vasyuchka, A. A. Serga, and B. Hillebrands, Advances in coherent magnonics, *Nat. Rev. Mater.* **6**, 1114 (2021).
- [32] B. Z. Rameshti, S. V. Kusminskiy, J. A. Haigh, K. Usami, D. Lachance-Quirion, Y. Nakamura, C.-M. Hu, H. X. Tang, G. E. Bauer, and Y. M. Blanter, Cavity magnonics, *Phys. Rep.* **979**, 1 (2022).
- [33] H. Yuan, Y. Cao, A. Kamra, R. A. Duine, and P. Yan, Quantum magnonics: When magnon spintronics meets quantum information science, *Phys. Rep.* **965**, 1 (2022).
- [34] C. Kittel, Interaction of spin waves and ultrasonic waves in ferromagnetic crystals, *Phys. Rev.* **110**, 836 (1958).
- [35] M. Xu, K. Yamamoto, J. Puebla, K. Baumgaertl, B. Rana, K. Miura, H. Takahashi, D. Grundler, S. Maekawa, and Y. Otani, Nonreciprocal surface acoustic wave propagation via magneto-rotation coupling, *Sci. Adv.* **6**, eabb1724 (2020).
- [36] P. J. Shah, D. A. Bas, I. Lisenkov, A. Matyushov, N. X. Sun, and M. R. Page, Giant nonreciprocity of surface acoustic waves enabled by the magnetoelastic interaction, *Sci. Adv.* **6**, eabc5648 (2020).
- [37] L. Liao, J. Puebla, K. Yamamoto, J. Kim, S. Maekawa, Y. Hwang, Y. Ba, and Y. Otani, Valley-selective phonon-magnon scattering in magnetoelastic superlattices, *Phys. Rev. Lett.* **131**, 176701 (2023).
- [38] L. Liao, J. Liu, J. Puebla, Q. Shao, and Y. Otani, Hybrid magnon-phonon crystals, *npj Spintronics* **2**, 47 (2024).
- [39] R. Takahashi and N. Nagaosa, Berry curvature in magnon-phonon hybrid systems, *Phys. Rev. Lett.* **117**, 217205 (2016).
- [40] G. Go, S. K. Kim, and K.-J. Lee, Topological magnon-phonon hybrid excitations in two-dimensional ferromagnets with tunable chern numbers, *Phys. Rev. Lett.* **123**, 237207 (2019).
- [41] E. Thingstad, A. Kamra, A. Brataas, and A. Sudbø, Chiral phonon transport induced by topological magnons, *Phys. Rev. Lett.* **122**, 107201 (2019).
- [42] X. Zhang, Y. Zhang, S. Okamoto, and D. Xiao, Thermal hall effect induced by magnon-phonon interactions, *Phys. Rev. Lett.* **123**, 167202 (2019).
- [43] S. Zhang, G. Go, K.-J. Lee, and S. K. Kim, Su (3) topology of magnon-phonon hybridization in 2d antiferromagnets, *Phys. Rev. Lett.* **124**, 147204 (2020).
- [44] S. Yang, F. Chen, R. Zhao, J. Wu, Y. Wang, H. Luo, N. Han, Q. Chen, Y. Hu, W. Li, M. Li, H. Chen, and Y. Yang, End-to-end autonomous scientific discovery on a real optical platform, *arXiv:2604.27092* 10.48550/arXiv.2604.27092 (2026).
- [45] C. Kittel, On the theory of ferromagnetic resonance absorption, *Phys. Rev.* **73**, 155 (1948).
- [46] T. Holstein and H. Primakoff, Field dependence of the intrinsic domain magnetization of a ferromagnet, *Phys. Rev.* **58**, 1098 (1940).
- [47] X. Zhang, C.-L. Zou, L. Jiang, and H. X. Tang, Cavity magnomechanics, *Sci. Adv.* **2**, e1501286 (2016).
- [48] See Supplemental Material for additional details on the theoretical model, Dirac mass and topological phase transitions.
- [49] T. Oka and H. Aoki, Photovoltaic hall effect in graphene, *Phys. Rev. B* **79**, 081406(R) (2009).
- [50] T. Kitagawa, T. Oka, A. Brataas, L. Fu, and E. Demler, Transport properties of nonequilibrium systems under the application of light: Photoinduced quantum hall insulators without landau levels, *Phys. Rev. B* **84**, 235108 (2011).
- [51] G. Usaj, P. M. Perez-Piskunow, L. E. F. Foa Torres, and C. A. Balseiro, Irradiated graphene as a tunable floquet topological insulator, *Phys. Rev. B* **90**, 115423 (2014).
- [52] M. S. Rudner and N. H. Lindner, Band structure engineering and non-equilibrium dynamics in floquet topological insulators, *Nat. Rev. Phys.* **2**, 229 (2020).
- [53] J. R. Schrieffer and P. A. Wolff, Relation between the Anderson model and the Kondo problem, *Phys. Rev.* **149**, 491 (1966).
- [54] S. Bravyi, D. P. DiVincenzo, and D. Loss, Schrieffer-wolff transformation for quantum many-body systems, *Ann. Phys.* **326**, 2793 (2011).
- [55] F. D. M. Haldane, Model for a quantum hall effect without landau levels: Condensed-matter realization of the “parity anomaly”, *Phys. Rev. Lett.* **61**, 2015 (1988).
- [56] T. Fukui, Y. Hatsugai, and H. Suzuki, Chern numbers in discretized Brillouin zone: Efficient method of computing (spin) Hall conductances, *J. Phys. Soc. Jpn.* **74**, 1674 (2005).
- [57] D. J. Thouless, M. Kohmoto, M. P. Nightingale, and M. den Nijs, Quantized Hall conductance in a two-dimensional periodic potential, *Phys. Rev. Lett.* **49**, 405 (1982).
- [58] L. Shao, S. Maity, L. Zheng, L. Wu, A. Shams-Ansari, Y.-I. Sohn, E. Puma, M. N. Gadalla, M. Zhang, C. Wang, E. L. Hu, K. Lai, and M. Loncar, Phononic band structure engineering for high- q gigahertz surface acoustic wave resonators on lithium niobate, *Phys. Rev. Appl.* **12**, 014022 (2019).
- [59] D. Hatanaka, M. Asano, H. Okamoto, and H. Yamaguchi, Phononic crystal cavity magnomechanics, *Phys. Rev. Appl.* **19**, 054071 (2023).
- [60] Z.-D. Zhang, S.-Y. Yu, M.-H. Lu, and Y.-F. Chen, Gigahertz surface acoustic wave topological rainbow in nanoscale phononic crystals, *Phys. Rev. Lett.* **133**, 267001 (2024).

- [61] M. Shen, J. Xie, C.-L. Zou, Y. Xu, W. Fu, and H. X. Tang, High frequency lithium niobate film-thickness-mode optomechanical resonator, [Appl. Phys. Lett. **117**, 131104 \(2020\)](#).
- [62] Z. Zheng, H. Feng, A. T. Işık, P. J. M. van der Slot, C. Wang, and D. Marpaung, Gigahertz thermoelastic acousto-optic modulation in lithium niobate integrated photonic device, [Nanophotonics **14**, 4683 \(2025\)](#).
- [63] L. Dreher, M. Weiler, M. Pernpeintner, H. Huebl, R. Gross, M. S. Brandt, and S. T. B. Goennenwein, Surface acoustic wave driven ferromagnetic resonance in nickel thin films: Theory and experiment, [Phys. Rev. B **86**, 134415 \(2012\)](#).
- [64] Y. Hwang, J. Puebla, K. Kondou, C. Gonzalez-Ballester, H. Isshiki, C. Sánchez Muñoz, L. Liao, F. Chen, W. Luo, S. Maekawa, and Y. Otani, Strongly coupled spin waves and surface acoustic waves at room temperature, [Phys. Rev. Lett. **132**, 056704 \(2024\)](#).
- [65] H. Matsumoto, I. Yasuda, M. Asano, Y. Todaka, T. Kawada, M. Kawaguchi, D. Hatanaka, and M. Hayashi, Magnon-phonon coupling of synthetic anti-ferromagnets in a surface acoustic wave cavity resonator, [Nano Lett. **24**, 5683 \(2024\)](#).
- [66] K. Künstle, Y. Kunz, T. Moussa, K. Lasinger, K. Yamamoto, P. Pirro, J. F. Gregg, A. Kamra, and M. Weiler, Magnon-polaron control in a surface magnetoacoustic wave resonator, [Nat. Commun. **16**, 10116 \(2025\)](#).