arXiv:2610.18494v1 [eess.SP] 13 Oct 2026

Task-evoked eye movements as candidate digital biomarkers of depression severity in adolescents and young adults

Youxiang Sun[1,2,†], Yujie Nie[1,2,†], Zhuoru Wu[4,†], Mengyuan Yu[3], Wenkong Wang[1,2], Xiang Li[1,2], Yumei Wang[3,*], Xin Ma[1,2,*], and Lizhou Fan[4,*]

[1]School of Control Science and Engineering, Shandong University, Jinan, Shandong, China

[2]Engineering Research Center of Intelligent Unmanned System, Ministry of Education, China

[3]Department of Psychology, Shandong Provincial Hospital Affiliated to Shandong First Medical University, Jinan, Shandong, China

[4]Department of Psychiatry, The Chinese University of Hong Kong, Shatin, Hong Kong SAR, China

[†]These authors contributed equally: Youxiang Sun, Yujie Nie, and Zhuoru Wu.

[*]Correspondence should be addressed to Yumei Wang (wangym@bjmu.edu.cn), Xin Ma (maxin@sdu.edu.cn), or Lizhou Fan (leofan@cuhk.edu.hk)

Abstract:

Objective digital measures of depression severity could complement conventional symptom assessments and facilitate more timely clinical evaluation. Here, we investigated whether task-evoked eye movements provide interpretable behavioral markers of depression severity in adolescents and young adults. In this single-center, cross-sectional study, 155 participants aged 14–20 years completed emotional-spatial free-viewing and memory-guided recall tasks and were stratified by 17-item Hamilton Depression Rating Scale scores. We examined 19 prespecified eye-movement features across six task-informed domains using covariate-adjusted ordinal-trend models. Greater depression severity was associated with lower gaze-allocation entropy (standardized $\beta = -0.599$), fewer inter-area switches ($\beta = -0.526$), longer first target-entry latency ($\beta = 0.572$), and lower target dwell proportion ($\beta = -0.678$; all false-discovery-rate-adjusted $q < 0.001$). Emotion-specific attentional-bias measures were not associated with severity, whereas associations involving fixation dispersion weakened after adjustment for anxiety symptoms. In a complementary machine-learning analysis incorporating 44 candidate eye-movement features and demographic covariates, a three-stage cascade classifier achieved the highest four-class stratification performance (accuracy, 0.665; macro-F1, 0.670; area under the receiver operating characteristic curve, 0.869), outperforming single-stage and two-stage approaches. Together, these findings identify altered visual exploration and memory-guided gaze control as candidate digital biomarkers of depression severity, rather than generalized negative attentional bias. These results support further prospective evaluation of task-evoked eye tracking as a complementary approach for timely severity assessment and clinical decision support.

Keywords: adolescent depression, eye tracking, digital biomarker, visual attention, spatial memory, psychomotor behavior

Introduction

Depression during adolescence is common, disabling, and associated with recurrent illness, impaired social and educational functioning, self-harm, and suicide risk[1–6]. This developmental period also involves substantial maturation of emotion regulation, executive control, memory, and social cognition[7,8]. Clinical assessment therefore needs to capture not only affective symptoms but also the cognitive and behavioral changes through which depression may interfere with daily functioning.

Depression has been linked to alterations in cognitive control, attention, memory, and processing of socially salient emotional information[9–11]. Eye tracking provides temporally resolved measurement of overt visual attention and oculomotor behavior and can distinguish initial orienting, sustained allocation, scanpath organization, and fixation-related processing[12–14]. These properties make eye movement a candidate task-based digital behavioral measure, although individual metrics are not specific to a single latent cognitive process and require interpretation within the demands of the task that generated them.

Most depression eye-tracking research has focused on emotional attentional bias. Meta-analytic evidence suggests that clinical depression is more consistently associated with prolonged maintenance on dysphoric material and reduced attention to positive material than with altered initial orienting[15,16]. Studies using emotional faces and other affective stimuli have also reported mood-congruent gaze patterns and associations with particular symptom dimensions[17–20]. However, findings vary across stimulus sets, viewing durations, and task instructions, indicating that emotional valence alone may not account for depression-related gaze behavior.

Broader alterations have been reported in visual exploration, fixation control, saccadic behavior, and memory-related processing[21–26]. Eye-tracking studies involving memory tasks further suggest inefficient cue detection, impaired recognition of learned face–scene associations, and reduced allocation to remembered targets in depression[27,28]. In adolescents, eye-movement characteristics have also been associated with cognitive functioning in major depressive disorder[29]. Together, these findings motivate paradigms that jointly examine how participants distribute gaze while encoding information and how they use that information to guide subsequent behavior.

Three limitations remain. First, case–control comparisons cannot establish whether a feature changes progressively with symptom burden. Second, single-task studies and isolated metrics provide limited information about whether observed effects arise during visual sampling, scanpath organization, memory-

guided orienting, or maintenance after target acquisition. Third, statistical associations are difficult to interpret when eye-movement features are not prespecified according to task demands and plausible behavioral processes[14,30].

We therefore combined emotional-spatial free-viewing with memory-guided recall and organized 19 primary eye-movement features into six prespecified domains: visual scanning coverage, emotion-related attentional bias, gaze transition dynamics, memory-guided target acquisition, target-area gaze maintenance, and fixation stability. We tested ordered associations across four HAMD-17 symptom-severity strata and evaluated dimensional consistency using continuous HAMD-17 scores. We hypothesized that greater symptom severity would be associated with less comprehensive and less organized visual sampling, greater negative attentional maintenance, delayed target acquisition, weaker target maintenance, and poorer fixation stability. Beyond these dimensional tests, we used the task-informed feature set as input to an exploratory machine-learning analysis of whether the candidate features jointly separate the four severity strata and whether hierarchical (cascade) decision rules improve on single-stage classification. The study was designed to characterize task-related behavioral correlates of symptom severity, not to develop or validate a diagnostic test.

Results

Cohort characteristics and data quality

Of the 203 recruited participants, 48 were excluded because their proportion of valid eye-tracking data was below the prespecified threshold of 80%, leaving 155 participants for analysis. The analytic cohort comprised 32 participants with HAMD-17 scores of 0–7, 39 with scores of 8–16, 49 with scores of 17–23, and 35 with scores of 24 or higher. Mean age and years of education differed across severity groups, whereas the proportion of male participants did not (Table 1).

The observed severity gradient must be interpreted in relation to the recruitment design. Participants without depression were recruited from the community, whereas participants with major depressive disorder were recruited from a clinical service. Consequently, comparisons that span the no-depression and symptomatic groups may reflect both symptom severity and recruitment-source or diagnostic-status differences.

Table 1. Demographic and clinical characteristics across HAMD-17 symptom-severity groups.

	No depression (n = 32)	Mild (n = 39)	Moderate (n = 49)	Severe (n = 35)	Statistic	p
Age, years	17.22 (2.56)	17.23 (2.24)	16.06 (2.67)	15.60 (2.63)	H = 10.84	0.013
Male sex, no. (%)	14 (43.8)	16 (41.0)	17 (34.7)	10 (28.6)	χ^2 = 2.08	0.557
Education, years	11.34 (2.35)	11.18 (2.04)	10.14 (2.37)	9.69 (2.27)	H = 12.94	0.005
HAMD-17 score	4.28 (2.44)	12.51 (3.09)	19.16 (2.22)	27.06 (3.87)	H = 144.02	< 0.001
HAMA score	5.09 (5.01)	9.31 (5.05)	14.10 (7.23)	18.37 (8.11)	H = 53.47	< 0.001

Continuous variables are mean (SD). Group comparisons used Kruskal–Wallis tests for continuous variables and a chi-square test for sex. HAMA, Hamilton Anxiety Rating Scale; HAMD-17, 17-item Hamilton Depression Rating Scale.

Ordered-severity associations in visual scanning and gaze transition dynamics

A total of 44 eye-movement features were extracted to assess their associations with ordered HAMD-17 symptom severity. Operational definitions and calculation procedures for all features are provided in Table S1. Table 2 presents the prespecified primary features, whereas results for the remaining supplementary features are reported in Table S2.

Severity-related associations during free-viewing were concentrated in visual scanning coverage and gaze transition dynamics (Table 2). All three primary visual-scanning features were associated with the ordered severity variable after false-discovery-rate correction. Each one-level increase in severity was associated with a 0.305-SD longer time to full area-of-interest (AOI) coverage (95% CI, 0.165–0.445), a 0.447-SD lower first-pass AOI coverage ratio (95% CI, −0.604 to −0.291), and a 0.599-SD lower AOI allocation entropy (95% CI, −0.733 to −0.466; all $q_{\mathrm{FDR}} < 0.001$). These estimates indicate slower, less complete, and less balanced sampling of the three emotional face locations as symptom severity increased (Fig. 1).

Gaze transitions showed a similar ordered pattern. Greater severity was associated with a lower inter-AOI switch rate (β = −0.526, 95% CI, −0.674 to −0.379), lower transition diversity (β = −0.567, 95% CI, −0.705 to −0.429), and greater transition tortuosity (β = 0.334, 95% CI, 0.186–0.482; all $q_{\mathrm{FDR}} < 0.001$). Within-AOI gaze dispersion was not significant after correction. The convergence of lower switch frequency, lower transition diversity, and greater tortuosity supports an association with less efficient scanpath organization rather than a simple reduction in total looking.



Fig. 1. Representative free-viewing eye-movement features across HAMD-17 symptom-severity groups. Violin plots with embedded boxplots show unadjusted participant-level distributions for selected features from visual scanning coverage, emotion-related attentional bias, and gaze transition dynamics. Displayed p values were obtained from ordered-severity regression models adjusted for age, sex, and education, with HC3 standard errors. The displayed q values were adjusted across the 19 prespecified primary features.

Continuous HAMD-17 associations and sensitivity analyses

Analyses using continuous HAMD-17 scores showed generally similar directions to the ordered-severity results for the F1, F3, R1, and R2 domains, whereas associations in the F2 domain remained limited. The primary-feature results are summarized in Table 3, whereas complete results for supplementary features are provided in Table S3.

In the F1 domain, higher continuous HAMD-17 scores were associated with longer time to full AOI coverage (β = 0.299, 95% CI, 0.157–0.441), lower first-pass AOI coverage ratio (β = −0.435, 95% CI, −0.586 to −0.283), and lower AOI allocation entropy (β = −0.549, 95% CI, −0.682 to −0.416; all $q_{\mathrm{FDR}} < 0.001$). Spearman correlations showed the same directions (ρ_s = 0.278, −0.405, and −0.651, respectively; all $q_{\mathrm{FDR}} < 0.001$).

In the F3 domain, higher HAMD-17 scores were associated with a lower inter-AOI switch rate (β = −0.500, 95% CI, −0.652 to −0.347), lower AOI transition diversity (β = −0.506, 95% CI, −0.644 to −0.369), and greater transition tortuosity (β = 0.270, 95% CI, 0.128–0.412; all $q_{\mathrm{FDR}} < 0.001$). The

Table 2. Associations between ordered HAMD-17 symptom severity and prespecified eye-movement features.

Feature	Domain	Standardized β (95% CI)	p	q_{FDR}
Time to full AOI coverage	F1	0.305 (0.165, 0.445)	< 0.001	< 0.001
First-pass AOI coverage ratio	F1	−0.447 (−0.604, −0.291)	< 0.001	< 0.001
AOI allocation entropy	F1	−0.599 (−0.733, −0.466)	< 0.001	< 0.001
Negative–positive gaze allocation bias	F2	0.124 (−0.019, 0.266)	0.090	0.114
Negative AOI visit duration	F2	−0.065 (−0.238, 0.109)	0.466	0.492
Negative stickiness after first hit	F2	−0.020 (−0.184, 0.145)	0.813	0.813
Inter-AOI switch rate	F3	−0.526 (−0.674, −0.379)	< 0.001	< 0.001
AOI transition diversity	F3	−0.567 (−0.705, −0.429)	< 0.001	< 0.001
Transition tortuosity	F3	0.334 (0.186, 0.482)	< 0.001	< 0.001
Within-AOI gaze dispersion	F3	0.131 (−0.013, 0.275)	0.075	0.102
First target-entry latency	R1	0.572 (0.442, 0.702)	< 0.001	< 0.001
Initial target-entry accuracy	R1	−0.600 (−0.706, −0.493)	< 0.001	< 0.001
First target-fixation latency	R1	0.580 (0.454, 0.706)	< 0.001	< 0.001
Target dwell proportion	R2	−0.678 (−0.778, −0.578)	< 0.001	< 0.001
Target lock-on ratio	R2	−0.437 (−0.575, −0.299)	< 0.001	< 0.001
Post-hit off-target ratio	R2	0.224 (0.103, 0.344)	< 0.001	< 0.001
Target-area fixation dispersion	R3	0.179 (0.022, 0.336)	0.025	0.037
Within-target fixation step size	R3	0.097 (−0.058, 0.251)	0.219	0.245
Target-area stability ellipse area	R3	0.125 (−0.047, 0.297)	0.153	0.182

Each feature was standardized before fitting a separate linear regression model. Ordered severity was coded 0–3 and models adjusted for age, sex, and education. HC3 standard errors were used. The coefficient is the SD difference in the eye-movement feature associated with a one-level increase in severity. FDR correction was applied across all 19 primary features. AOI, area of interest; CI, confidence interval; F1, visual scanning coverage; F2, emotion-related attentional bias; F3, gaze transition dynamics; R1, memory-guided target acquisition; R2, target-area gaze maintenance; R3, fixation stability.

corresponding Spearman correlations showed the same directions (ρ_s = −0.465, −0.516, and 0.326, respectively; all $q_{\mathrm{FDR}} < 0.001$). Within-AOI gaze dispersion was not significant in either analysis.

In the R1 domain, higher HAMD-17 scores were associated with longer first target-entry latency (β = 0.515, 95% CI, 0.351–0.679), lower initial target-entry accuracy (β = −0.562, 95% CI, −0.682 to −0.441), and longer first target-fixation latency (β = 0.515, 95% CI, 0.350–0.681; all $q_{\mathrm{FDR}} < 0.001$). In the R2 domain, higher HAMD-17 scores were associated with lower target dwell proportion (β = −0.624, 95% CI, −0.750 to −0.498), lower target lock-on ratio (β = −0.369, 95% CI, −0.539 to −0.199), and higher post-hit off-target ratio (β = 0.199, 95% CI, 0.076–0.323; q_{FDR} = 0.003 for post-hit off-target ratio and < 0.001 for the other two). Spearman correlations for these six R1 and R2 features were also significant and showed the same directions.

Associations in the F2 domain were weaker. Negative–positive gaze allocation bias was not significant in the Spearman analysis (ρ_s = 0.114, q_{FDR} = 0.206), although it reached the FDR-corrected threshold in the adjusted regression model (β = 0.152, 95% CI, 0.011–0.293; q_{FDR} = 0.049). Negative AOI visit duration

Table 3. Associations between continuous HAMD-17 scores and prespecified eye-movement features.

Feature	Domain	Spearman correlation			Adjusted regression model		
		ρ_s	p	q_{FDR}	Standardized β (95% CI)	p	q_{FDR}
Time to full AOI coverage	F1	0.278	< 0.001	< 0.001	0.299 (0.157, 0.441)	< 0.001	< 0.001
First-pass AOI coverage ratio	F1	−0.405	< 0.001	< 0.001	−0.435 (−0.586, −0.283)	< 0.001	< 0.001
AOI allocation entropy	F1	−0.651	< 0.001	< 0.001	−0.549 (−0.682, −0.416)	< 0.001	< 0.001
Negative–positive gaze allocation bias	F2	0.114	0.158	0.206	0.152 (0.011, 0.293)	0.035	0.049
Negative AOI visit duration	F2	−0.095	0.242	0.273	0.033 (−0.163, 0.230)	0.740	0.776
Negative stickiness after first hit	F2	0.021	0.798	0.817	0.045 (−0.122, 0.212)	0.598	0.675
Inter-AOI switch rate	F3	−0.465	< 0.001	< 0.001	−0.500 (−0.652, −0.347)	< 0.001	< 0.001
AOI transition diversity	F3	−0.516	< 0.001	< 0.001	−0.506 (−0.644, −0.369)	< 0.001	< 0.001
Transition tortuosity	F3	0.326	< 0.001	< 0.001	0.270 (0.128, 0.412)	< 0.001	< 0.001
Within-AOI gaze dispersion	F3	0.103	0.202	0.233	0.054 (−0.099, 0.208)	0.489	0.566
First target-entry latency	R1	0.586	< 0.001	< 0.001	0.515 (0.351, 0.679)	< 0.001	< 0.001
Initial target-entry accuracy	R1	−0.597	< 0.001	< 0.001	−0.562 (−0.682, −0.441)	< 0.001	< 0.001
First target-fixation latency	R1	0.597	< 0.001	< 0.001	0.515 (0.350, 0.681)	< 0.001	< 0.001
Target dwell proportion	R2	−0.666	< 0.001	< 0.001	−0.624 (−0.750, −0.498)	< 0.001	< 0.001
Target lock-on ratio	R2	−0.482	< 0.001	< 0.001	−0.369 (−0.539, −0.199)	< 0.001	< 0.001
Post-hit off-target ratio	R2	0.290	< 0.001	< 0.001	0.199 (0.076, 0.323)	0.002	0.003
Target-area fixation dispersion	R3	0.225	0.005	0.008	0.221 (0.069, 0.374)	0.004	0.007
Within-target fixation step size	R3	0.114	0.159	0.206	0.131 (−0.028, 0.291)	0.106	0.141
Target-area stability ellipse area	R3	0.135	0.096	0.133	0.164 (−0.022, 0.351)	0.084	0.116

ρ_s denotes the Spearman rank correlation coefficient between continuous HAMD-17 total scores and eye-movement features. Adjusted β values were estimated using separate linear regression models with standardized HAMD-17 total score as the predictor and each standardized eye-movement feature as the outcome, adjusting for age, sex, and years of education. Robust standard errors were used. The β coefficient represents the SD difference in the eye-movement feature associated with a one-SD increase in HAMD-17 total score. FDR correction was applied separately to the Spearman correlation tests and the adjusted regression tests across all 44 eye-movement features, including 19 primary and 25 supplementary features. AOI, area of interest; CI, confidence interval; FDR, false discovery rate; F1, visual scanning coverage; F2, emotion-related attentional bias; F3, gaze transition dynamics; R1, memory-guided target acquisition; R2, target-area gaze maintenance; R3, fixation stability.

and negative stickiness after first hit were not significant in either analysis. In the R3 domain, target-area fixation dispersion was positively associated with continuous HAMD-17 scores ($\rho_s = 0.225$, $q_{FDR} = 0.008$; $\beta = 0.221$, 95% CI, 0.069–0.374; $q_{FDR} = 0.007$), whereas within-target fixation step size and target-area stability ellipse area were not significant after correction.

Additional adjustment for HAMA scores in Table S4 generally preserved the F1, F3, R1, and R2 associations, with most estimates remaining similar in direction and magnitude. In contrast, the association for target-area fixation dispersion was attenuated and was no longer significant after FDR correction. Excluding feature-specific extreme observations ($|z| > 3$) yielded a similar overall pattern, with the principal F1, F3, R1, and R2 associations remaining significant.

Emotion-related attentional bias across symptom severity

None of the three primary emotion-related attentional-bias features remained significant after correction. The negative–positive gaze allocation bias had $\beta = 0.124$ (95% CI, −0.019 to 0.266; $q_{FDR} = 0.114$),



Fig. 2. Representative memory-guided recall features across HAMD-17 symptom-severity groups. Violin plots with embedded boxplots show unadjusted participant-level distributions for selected features from target acquisition, target-area gaze maintenance, and fixation stability. Displayed p values were obtained from ordered-severity regression models adjusted for age, sex, and education, with HC3 standard errors. The displayed q values were adjusted across the 19 prespecified primary features.

negative-AOI visit duration had $\beta = -0.065$ (95% CI, −0.238 to 0.109; $q_{\text{FDR}} = 0.492$), and negative stickiness after first hit had $\beta = -0.020$ (95% CI, −0.184 to 0.145; $q_{\text{FDR}} = 0.813$). Thus, the broad severity gradients in coverage and transition organization were not accompanied by evidence of a generalized shift toward negative faces in this memory-relevant viewing context.

Ordered-severity associations in memory-guided target acquisition and gaze maintenance

Overall recall performance decreased across the four groups ($H = 71.97$, $p < 0.001$). Group-specific numbers of correct recall trials are provided in Table S5.

All three target-acquisition features were strongly associated with ordered severity (Fig. 2). Each severity step was associated with a 0.572-SD longer first target-entry latency (95% CI, 0.442–0.702), a 0.600-SD lower initial target-entry accuracy (95% CI, −0.706 to −0.493), and a 0.580-SD longer first target-fixation latency (95% CI, 0.454–0.706; all $q_{\text{FDR}} < 0.001$). These findings indicate that participants with greater symptom severity were slower and less accurate when orienting to the remembered target location.

Target maintenance after acquisition showed the largest standardized association. Greater severity was associated with a lower target dwell proportion (β = −0.678, 95% CI, −0.778 to −0.578), lower target lock-on ratio (β = −0.437, 95% CI, −0.575 to −0.299), and higher post-hit off-target ratio (β = 0.224, 95% CI, 0.103–0.344; all $q_{\mathrm{FDR}} < 0.001$). By contrast, evidence for local fixation stability was limited. Target-area fixation dispersion was associated with severity in the primary model (β = 0.179, 95% CI, 0.022–0.336; $q_{\mathrm{FDR}} = 0.037$), whereas within-target fixation step size and stability ellipse area were not.

Machine-learning classification of the four severity strata

As an exploratory analysis, we asked whether the candidate eye-movement features jointly separate the four severity strata. Model input comprised the 44 candidate features (25 free-viewing and 19 recall features, of which the 19 prespecified primary features are a subset) together with age, sex, and education, giving 47 inputs per participant. We evaluated 21 classifiers (three baseline models, the TabPFN transformer, fourteen AutoGluon configurations, and three simple-average ensembles; Methods) under four protocols: single-stage four-class classification; a two-stage cascade that first separates no/mild from moderate/severe strata and then splits each branch (2+2); a two-stage cascade that first separates no depression from the symptomatic strata and then classifies mild, moderate, and severe (1+3); and a three-stage cascade of successive binary decisions (2+2+2).

Performance was the mean over five folds of stratified cross-validation. Configurations were ranked by the sum of their ranks across accuracy, macro F1, and one-vs-rest macro AUC (rank 1 = best per metric), and Table 4 lists the top five configurations per protocol (grouped bar plots in Fig. 3); each listed configuration ranks within the top five for at least two of the three metrics. The three-stage cascade occupied the top of the composite ranking, with TabPFN to support vector machine (RBF) to TabPFN first (accuracy 0.658, macro F1 0.664, AUC 0.869). The highest individual metric value across all protocols was accuracy 0.665, macro F1 0.670, and AUC 0.851 (AutoGluon ExtraTreesGini to support vector machine (RBF) to TabPFN). Both two-stage cascades and single-stage classification were lower (best accuracy 0.632, 0.632, and 0.619, respectively). In an exploratory two-class task that asked only whether participants fell in the no/mild or the moderate/severe strata, the best classifier (logistic regression) reached accuracy 0.839 (macro F1 0.835) and AUC 0.929 (Fig. S1). The coarse no/mild versus moderate/severe boundary is therefore well supported by these features, whereas fine-grained separation, in particular between the moderate and severe strata, remains difficult. Because best-of-many configurations were selected from 441 or 9261 candidates, these

Table 4. Top-5 machine-learning configurations per protocol by composite ACC/F1/AUC rank.

Protocol	Model	Accuracy	Macro F1	AUC
Single-stage four-class	TabPFN	0.619 (0.062)	0.627 (0.064)	0.865 (0.027)
Single-stage four-class	Avg(R8)	0.613 (0.032)	0.618 (0.035)	0.843 (0.014)
Single-stage four-class	Avg(13)	0.613 (0.051)	0.614 (0.047)	0.838 (0.014)
Single-stage four-class	SVM	0.594 (0.029)	0.594 (0.029)	0.843 (0.019)
Single-stage four-class	AG:NN	0.600 (0.093)	0.605 (0.094)	0.837 (0.027)
Two-stage cascade 2+2	LR → TabPFN	0.626 (0.018)	0.631 (0.022)	0.859 (0.025)
Two-stage cascade 2+2	LR → SVM	0.626 (0.049)	0.634 (0.054)	0.855 (0.022)
Two-stage cascade 2+2	AG:LR → SVM	0.632 (0.071)	0.640 (0.073)	0.853 (0.036)
Two-stage cascade 2+2	LR → LR	0.632 (0.043)	0.641 (0.045)	0.851 (0.018)
Two-stage cascade 2+2	SVM → SVM	0.626 (0.059)	0.638 (0.061)	0.851 (0.022)
Two-stage cascade 1+3	TabPFN → Avg(R8)	0.619 (0.053)	0.626 (0.060)	0.851 (0.011)
Two-stage cascade 1+3	AG:ETG → Avg(T7)	0.626 (0.054)	0.635 (0.056)	0.843 (0.015)
Two-stage cascade 1+3	AG:ETG → Avg(13)	0.632 (0.067)	0.637 (0.075)	0.838 (0.015)
Two-stage cascade 1+3	AG:ETG → Avg(R8)	0.626 (0.063)	0.631 (0.072)	0.838 (0.014)
Two-stage cascade 1+3	AG:RFE → Avg(13)	0.626 (0.067)	0.631 (0.075)	0.832 (0.017)
Three-stage cascade 2+2+2	TabPFN → SVM → TabPFN	0.658 (0.063)	0.664 (0.066)	0.869 (0.031)
Three-stage cascade 2+2+2	AG:ETG → SVM → RF	0.645 (0.079)	0.651 (0.085)	0.854 (0.026)
Three-stage cascade 2+2+2	**AG:ETG → SVM → TabPFN**	**0.665 (0.081)**	**0.670 (0.085)**	**0.851 (0.033)**
Three-stage cascade 2+2+2	Avg(T7) → SVM → TabPFN	0.652 (0.053)	0.657 (0.056)	0.845 (0.036)
Three-stage cascade 2+2+2	AG:ETG → SVM → LR	0.645 (0.091)	0.653 (0.095)	0.840 (0.038)

Values are mean (SD) over five folds of stratified cross-validation. Configurations are ordered by the sum of their ranks across accuracy, macro F1, and AUC (rank 1 = best per metric); each listed configuration ranks within the top five for at least two of the three metrics. AG, AutoGluon; NN, NeuralNetTorch; ETE, ExtraTreesEntr; ETG, ExtraTreesGini; RFE, RandomForestEntr; LR, logistic regression; AG:LR, AutoGluon LinearModel; RF, random forest; SVM, support vector machine (RBF); TabPFN, TabPFN transformer; Avg(13), simple-average ensemble of 13 base AutoGluon models; Avg(T7), ensemble of the seven tree models; Avg(R8), ensemble of one representative per model family.

values are an upper-bound characterization of what the feature set can support rather than the performance of a single prespecified model.

Discussion

In this cross-sectional eye-tracking study, depression symptom severity was associated with broad changes in how young participants sampled, organized, retrieved, and maintained visual information. The most consistent associations were lower gaze-allocation entropy, fewer and less diverse transitions between emotional faces, slower and less accurate orienting to a remembered location, and weaker gaze maintenance after target acquisition. These effects were large in standardized units and remained significant after correction across the prespecified primary feature set. In contrast, none of the emotion-specific attentional-



Fig. 3. Top-5 machine-learning configurations for the four classification protocols. **a**, single-stage four-class (21 models); **b**, two-stage cascade 2+2 (441 configurations); **c**, two-stage cascade 1+3 (441 configurations); **d**, three-stage cascade 2+2+2 (9261 configurations). For each protocol, configurations are ordered by the sum of their ranks across accuracy, macro F1, and AUC (top 5 shown, left to right); grouped bars show the three metric values of each configuration. Values are means over five folds of stratified cross-validation. AG, AutoGluon; NN, NeuralNetTorch; ETE, ExtraTreesEntr; ETG, ExtraTreesGini; RFE, RandomForestEntr; LR, logistic regression; AG:LR, AutoGluon LinearModel; RF, random forest; SVM, support vector machine (RBF); TabPFN, TabPFN transformer; Avg(13), simple-average ensemble of 13 base AutoGluon models; Avg(T7), ensemble of the seven tree models; Avg(R8), ensemble of one representative per model family.

bias features was associated with severity, and fixation-stability findings were limited. The pattern therefore points more strongly to distributed visual exploration and memory-guided control than to a generalized negative gaze bias.

The free-viewing findings extend prior reports of altered scanpath length, saccade amplitude, fixation behavior, and visual exploration in depression[23–26]. Lower first-pass coverage and entropy indicate that participants with greater symptoms sampled the available facial information less comprehensively and less evenly. Lower inter-AOI switch rate and transition diversity, together with greater tortuosity, further indicate altered organization of the scanpath. These measures should not be interpreted as direct readouts of a single cognitive mechanism. They may jointly reflect differences in task engagement, processing speed, spatial encoding, executive allocation, or psychomotor behavior, and the present design cannot distinguish among these explanations.

The absence of a significant F2 signal is clinically and theoretically informative. Meta-analyses of passive or sustained viewing have identified greater maintenance on negative material and reduced attention to positive material in depression[15,16]. Here, however, every emotional face was potentially relevant because the target category was revealed only after encoding. This task demand may have reduced spontaneous valence-driven allocation by encouraging participants to encode all locations. The null results do not exclude context-dependent affective bias. Rather, they show that, under a memory-relevant viewing requirement, symptom severity was expressed more consistently in coverage and transition organization than in negative-versus-positive allocation.

The recall findings suggest that the largest severity-related differences emerged when encoded information had to guide a goal-directed gaze response. Longer target-entry and target-fixation latencies, lower initial target-entry accuracy, and reduced dwell after target acquisition are consistent with less efficient memory-guided orienting and maintenance. Prior studies have linked depression to cue-detection deficits and impaired memory-guided allocation[27,28], and adolescent work has associated eye-movement characteristics with cognitive functioning[29]. Nevertheless, the current task combines spatial memory, response selection, visual search, and oculomotor stabilization. The results should therefore be described as task-related behavioral correlates rather than evidence of isolated memory impairment or a specific neurocognitive pathway.

From a digital-medicine perspective, a brief, standardized eye-tracking task could complement clinical interviews by providing repeatable behavioral measurements that are difficult to obtain from symptom reports

alone. The domain framework also provides a more interpretable alternative to treating a large collection of eye-movement variables as independent biomarkers. Potential applications could include longitudinal phenotyping, treatment-response research, or stratification in mechanistic studies. Such uses remain prospective. The study did not evaluate diagnostic discrimination, test–retest reliability, responsiveness to clinical change, patient acceptability, implementation burden, or added value beyond routine assessment. Specialized hardware, calibration requirements, stimulus licensing, and intersite standardization also limit immediate scalability.

The exploratory machine-learning results extend the statistical framework to a discriminative question. Among the four protocols, three-stage cascades outperformed two-stage and single-stage classification (best four-class accuracy 0.665, macro F1 0.670, AUC 0.869), suggesting that severity discrimination benefits from progressively finer hierarchical decisions rather than a single four-way decision. The exploratory two-class task (no/mild versus moderate/severe) was substantially easier (best accuracy 0.839, AUC 0.929), indicating that the coarse symptomatic boundary is well supported by these features, while the residual four-class errors concentrated near the moderate–severe boundary. This mirrors clinical practice, in which the moderate–severe distinction rests on threshold items of the HAMD-17 that are intrinsically less well defined than the distinction between no depression and symptomatic illness. Refining moderate-versus-severe discrimination, for example with richer severity ratings or longitudinal anchors, is therefore a key direction for future work.

Several design features constrain the severity interpretation. The no-depression group was recruited from the community, whereas symptomatic participants were recruited from a clinical service, so the ordered gradient may partly encode differences in diagnostic status, referral pathway, medication exposure, illness chronicity, or testing context. Age and education also differed across groups. Although both were included as covariates, residual confounding remains possible, and adjusting simultaneously for highly correlated developmental variables may not fully recover comparability. A within-MDD continuous HAMD-17 analysis, propensity or overlap weighting, flexible age adjustment, and demographic-only benchmark models would strengthen the claim that the eye-movement features track symptom burden.

The HAMD-17 grouping also warrants caution. Thresholds create clinically recognizable strata but discard within-group information and impose equal spacing between categories. Moreover, HAMD-17 includes psychomotor items, raising the possibility that some eye-movement associations reflect overlap with clinician-rated motor behavior. Continuous HAMD-17 models should be emphasized, and sensitivity

analyses should exclude psychomotor items or examine symptom dimensions. Anxiety was addressed only by covariate adjustment. Because depression and anxiety symptoms overlap, HAMA adjustment may either reduce confounding or remove clinically shared variance, and both interpretations should be considered.

Other limitations include the cross-sectional and single-center design, modest group sizes, exclusion of 48 participants on eye-tracking quality criteria, and incomplete reporting of medication, treatment, illness duration, comorbidity, sleep, and visual characteristics. The sample age range of 14–20 years extends beyond conventional definitions of adolescence and should be described as adolescents and young adults unless a prespecified alternative definition is justified. The fixed trial sequence and progressively shorter display duration may introduce order, fatigue, learning, or target-emotion confounding. The requirement for valid data in all 12 trials may also select participants with better task adherence. The machine-learning analysis was exploratory, used a single five-fold split of a small sample, and reported best-of-many configurations (441 two-stage or 9261 three-stage), so its best values are optimistic upper bounds on generalizable performance. External validation, trial-level mixed-effects analyses, calibration and reliability studies, and prospective longitudinal evaluation are required.

In conclusion, the combined encoding and recall paradigm identified a coherent severity-related pattern across visual sampling, scanpath organization, memory-guided acquisition, and target maintenance, while providing little evidence for generalized negative gaze bias; an exploratory hierarchical classifier reached the best four-class separation (accuracy 0.665, macro F1 0.670, AUC 0.869). These findings support eye tracking as a candidate research tool for task-based behavioral phenotyping of depression symptom severity. They do not establish a depression-specific mechanism, diagnostic biomarker, or clinically deployable assessment.

Methods

Study design and reporting

This was a single-center, cross-sectional observational eye-tracking study. Patients with major depressive disorder were recruited from the Department of Psychology, Shandong Provincial Hospital Affiliated to Shandong First Medical University, and participants without depression were recruited from local communities.

Participants

Eligible participants were aged 14–20 years, had completed at least primary education, and had normal or corrected-to-normal vision. Patients were required to meet DSM-5 criteria for major depressive disorder in a clinical interview conducted by an experienced psychiatrist and to have a HAMD-17 score above the normal range. Community participants were required to have a HAMD-17 score within the normal range and no personal or family history of depression.

Exclusion criteria were a history of neurological disease, traumatic brain injury, epilepsy, or another organic brain disorder; a current or previous major psychiatric disorder other than major depressive disorder, including schizophrenia, bipolar disorder, or substance-use disorder; intellectual disability or cognitive impairment sufficient to interfere with task completion; color blindness or another ocular condition affecting eye tracking; and missing key clinical or behavioral data or inadequate eye-tracking quality.

Clinical assessment and symptom-severity strata

Depressive symptoms were rated using the HAMD-17, and anxiety symptoms were rated using the HAMA[31]. Both scales were administered before eye tracking by trained clinical assessors. The HAMD-17 was used as the principal measure of depression symptom severity[32–34].

Participants were grouped using HAMD-17 thresholds of 0–7, 8–16, 17–23, and 24 or higher, corresponding to no, mild, moderate, and severe symptom strata[35–37]. These labels describe HAMD-17 symptom strata and should not be interpreted as independent diagnostic categories.

Ethics and consent

The study complied with the Declaration of Helsinki and was approved by the Ethics Committee for Biomedical Research Involving Human Subjects, Shandong Provincial Hospital Affiliated to Shandong First Medical University (R202411120385). Written informed consent was obtained from adult participants. For participants younger than 18 years, written permission was obtained from a parent or legal guardian.

Eye-tracking apparatus

Eye movements were recorded using a Tobii Pro Fusion system (Tobii Technology, Stockholm, Sweden) at 250 Hz and exported using Tobii Pro Lab version 25.23. Participants completed a nine-point calibration before the task. Fixations were identified using the Tobii Pro Lab fixation filter with a minimum duration of 100 ms.

Stimuli

The task used 36 facial images from the Chinese Facial Affective Picture System, with equal numbers of male and female identities and 12 images in each of the positive, neutral, and negative categories[38]. Normative mean (SD) valence and arousal ratings were 2.75 (0.46) and 5.97 (0.78) for negative faces, 4.42 (0.32) and 3.97 (0.69) for neutral faces, and 6.56 (0.59) and 5.28 (0.71) for positive faces. Images were converted to grayscale and standardized for size and resolution. Each image appeared three times, with pseudo-randomized screen position across displays.

Emotional-spatial free-viewing and memory-guided recall paradigm

The paradigm paired emotional-spatial free-viewing with a subsequent memory-guided recall response (Fig. 4). During free-viewing, each display contained one positive, one neutral, and one negative face at three locations in an upright or inverted triangular arrangement. Participants knew that a subsequent cue could request any emotional category, but the target was not revealed during viewing. The task therefore required simultaneous sampling of emotional faces and encoding of their spatial locations rather than unconstrained passive viewing.

After the free-viewing displays in a trial, a cue indicated the emotional category to be recalled from the final display. A probe screen then showed six fixation points corresponding to all possible face locations. Participants were instructed to direct gaze to the remembered target location and maintain fixation until the next trial. The paradigm included 12 trials, with each emotion cued four times. Trials contained two, three, or four free-viewing displays in a repeated cycle, yielding 36 displays. Display duration decreased from 4.5 s to 3.4 s in 0.1-s steps across trials. The cue was presented for 4.0 s and the probe for 3.5 s. The target-emotion sequence was pseudo-randomized but identical across participants. Stimuli were displayed on a 24-inch, 1920 $\times$ 1080 monitor at a viewing distance of approximately 60–80 cm.



Fig. 4. Emotional-spatial free-viewing and memory-guided recall paradigm. Each trial comprised two to four displays containing positive, neutral, and negative faces, followed by an emotional-category cue and a six-location recall probe. The cue referred to the final face display immediately preceding it.

The identical sequence and progressively changing display duration standardized exposure but also coupled trial order with task difficulty.

Gaze preprocessing and quality control

Raw gaze samples were restricted to valid samples and segmented according to task phase and trial. Gaze coordinates were mapped to prespecified AOIs. Free-viewing features were aggregated from display to trial and then participant level. Recall features were computed by trial relative to the target AOI and then aggregated to participant level. Participants were included only if their overall valid-sample ratio was at least 80% and all 12 trials were usable.

A recall trial was classified as correct when gaze entered and remained within the target AOI according to a prespecified offline rule.

Table 5. Prespecified eye-movement feature domains.

Task	Domain	Task-related interpretation	Primary features
Free-viewing	F1 Visual scanning coverage	Completeness and balance of visual sampling across the three emotional face AOIs	Time to full AOI coverage; first-pass AOI coverage ratio; AOI allocation entropy
Free-viewing	F2 Emotion-related attentional bias	Valence-related allocation and maintenance, particularly negative maintenance and delayed disengagement	Negative–positive gaze allocation bias; negative-AOI visit duration; negative stickiness after first hit
Free-viewing	F3 Gaze transition dynamics	Spatial and temporal organization of gaze across and within emotional AOIs	Inter-AOI switch rate; AOI transition diversity; transition tortuosity; within-AOI gaze dispersion
Recall	R1 Memory-guided target acquisition	Efficiency and accuracy of orienting to the cued location using encoded spatial information	First target-entry latency; initial target-entry accuracy; first target-fixation latency
Recall	R2 Target-area gaze maintenance	Preferential and sustained gaze on the recalled target after acquisition	Target dwell proportion; target lock-on ratio; post-hit off-target ratio
Recall	R3 Fixation stability	Local spatial stability of gaze within the target area	Target-area fixation dispersion; within-target fixation step size; target-area stability ellipse area

AOI, area of interest.

Task-informed eye-movement feature framework

Features were prespecified according to the task phase and the aspect of gaze behavior they were intended to characterize. Free-viewing features were grouped into visual scanning coverage (F1), emotion-related attentional bias (F2), and gaze transition dynamics (F3). Recall features were grouped into memory-guided target acquisition (R1), target-area gaze maintenance (R2), and fixation stability (R3) (Table 5). These domains are task-informed interpretations and not direct measurements of isolated cognitive mechanisms. Operational definitions and calculation procedures for all features are provided in Table S1.

Statistical analysis

Analyses were conducted at participant level. Continuous demographic and clinical variables were compared across the four HAMD-17 groups using Kruskal–Wallis tests, and sex was compared using a chi-square test. Each primary eye-movement feature was standardized before model fitting. For feature Y_j, the primary

model was

$$Y_{ij}^{(z)} = \beta_0 + \beta_1 G_i + \beta_2 \text{Age}_i + \beta_3 \text{Sex}_i + \beta_4 \text{Education}_i + \epsilon_i, \quad (1)$$

where G_i was coded 0, 1, 2, or 3 for the ordered HAMD-17 strata. HC3 heteroscedasticity-consistent standard errors were used. Thus, β_1 represents the SD change in the feature associated with a one-level increase in symptom-severity stratum.

The primary family contained 19 prespecified features across six domains. Benjamini–Hochberg false-discovery-rate correction was applied across these 19 tests, and $q_{\text{FDR}} < 0.05$ was considered statistically significant. All tests were two-sided. The 25 supplementary features were analyzed as supportive evidence. The Benjamini–Hochberg false-discovery-rate correction was applied across the 25 supplementary features as a separate correction family.

Dimensional analyses examined Spearman correlations and linear regressions using continuous HAMD-17 scores, adjusted for age, sex, and education. Sensitivity analyses additionally adjusted for HAMA score and repeated the primary models after values with feature-specific $|z| > 3$ were set to missing.

Machine-learning classification

The machine-learning analysis was exploratory and used the same 155 participants and HAMD-17 severity strata as the statistical analysis. Inputs were the 44 candidate eye-movement features (25 free-viewing and 19 recall features, including the 19 prespecified primary features) plus age, sex, and education (47 inputs). The severity label, raw HAMD-17 and HAMA scores, and participant identifiers were excluded from all model inputs. The single missing value of one recall feature in three participants was imputed with the training-fold mean.

We evaluated 21 classifiers: logistic regression (L2 penalty, maximum 5000 iterations), random forest (500 trees), radial-basis-function support vector machine with probability calibration, the TabPFN transformer in its default configuration, fourteen AutoGluon 1.1.1 configurations (LightGBM, LightGBM extra-trees, large LightGBM, XGBoost, CatBoost, two random-forest, two extra-trees, two linear, two k-nearest-neighbour, and one neural-network model, plus a weighted ensemble fitted with a 20% holdout of each training fold, refit on the full fold, without stacking or bagging), and three simple-average ensembles of the out-of-fold class probabilities of the 13 base AutoGluon models, of the seven tree models, or of one representative per model family (eight models).

Four protocols were tested. Single-stage: direct four-class classification of the severity strata. Two-stage 2+2: a first binary classifier separated no/mild from moderate/severe strata, and two second-stage binary classifiers separately split no from mild and moderate from severe. Two-stage 1+3: a first binary classifier separated no depression from the symptomatic strata, followed by a three-class classifier for mild, moderate, and severe. Three-stage 2+2+2: the 2+2 first stage followed by both second-stage binary classifiers. In cascade protocols, second- and third-stage models were trained, within each fold, on the true-branch training subsets, and test-fold participants were routed by the previous stage's predicted probabilities; the four stratum probabilities were the products of the stage probabilities along each path of the cascade tree (for the three-stage protocol, $(1-p_1)(1-p_2)$, $(1-p_1)p_2$, $p_1(1-p_2)(1-p_3)$, and $p_1(1-p_2)p_3$, where p_1, p_2, and p_3 are the first-stage moderate/severe, second-stage mild, and third-stage severe probabilities), and the predicted stratum was the argmax. All protocols used the same five-fold cross-validation (stratified on the four severity labels, shuffled, fixed seed), with 124 participants for training and 31 for testing in each fold; the exploratory two-class task was stratified on its two labels. Imputation and, for the three baseline classifiers, standardization, were refit within each training fold. Accuracy, macro-averaged F1, and one-vs-rest macro-averaged AUC were reported as the mean (sample SD) over folds. Analyses used Python 3.8.20, scikit-learn 1.3.2, AutoGluon 1.1.1, and TabPFN 8.4.0 on an NVIDIA RTX 5880 Ada Generation GPU.

Data availability

Data supporting the findings of this study are not publicly available because they contain sensitive behavioral and clinical information from adolescent participants. Access to these data is restricted by ethical and institutional requirements.

Code availability

The custom code used for eye tracking data preprocessing, model training, and evaluation is not publicly available but may be made available to qualified researchers upon reasonable request to the corresponding authors, subject to institutional approval.

Acknowledgements

This work was supported in part by the National Key Research and Development Program of China under Grant 2023YFB4706100, the National Natural Science Foundation of China under Grant 62573271, the Major Basic Research Project of Shandong Provincial Natural Science Foundation under Grant ZR2025ZD24, and the Chinese University of Hong Kong Vice-Chancellor Early Career Professorship Scheme under Grant 441.

Author contributions

Youxiang Sun: Writing – original draft, Writing – review and editing, Conceptualization, Data curation. Yujie Nie: Writing – original draft, Writing – review and editing. Zhuoru Wu: Writing – original draft, Writing – review and editing, Formal analysis, Visualization. Mengyuan Yu: Writing – original draft, Writing – review and editing, Investigation, Data curation. Wenkong Wang: Writing – original draft, Writing – review and editing. Xiang Li: Writing – original draft, Writing – review and editing. Yumei Wang: Writing – original draft, Writing – review and editing, Supervision, Data curation. Xin Ma: Writing – original draft, Writing – review and editing, Supervision, Funding acquisition. Lizhou Fan: Writing – original draft, Writing – review and editing, Conceptualization, Supervision, Funding acquisition.

Competing interests

The authors declare no competing interests.

References

[1] Shorey, S., Ng, E.D., Wong, C.H.J., 2022. Global prevalence of depression and elevated depressive symptoms among adolescents: A systematic review and meta-analysis. British Journal of Clinical Psychology 61, 287–305. doi:10.1111/bjc.12333.

[2] Friedrich, M.J., 2017. Depression is the leading cause of disability around the world. JAMA 317, 1517. doi:10.1001/jama.2017.3826.

[3] Clayborne, Z.M., Varin, M., Colman, I., 2019. Systematic review and meta-analysis: Adolescent depression and long-term psychosocial outcomes. Journal of the American Academy of Child & Adolescent Psychiatry 58, 72–79. doi:10.1016/j.jaac.2018.07.896.

[4] Johnson, D., Dupuis, G., Piche, J., Clayborne, Z., Colman, I., 2018. Adult mental health outcomes of adolescent depression: A systematic review. Depression and Anxiety 35, 700–716. doi:10.1002/da.22777.

[5] Grossberg, A., Rice, T., 2023. Depression and suicidal behavior in adolescents. Medical Clinics of North America 107, 169–182. doi:10.1016/j.mcna.2022.04.005.

[6] Dwyer, J.B., Stringaris, A., Brent, D.A., Bloch, M.H., 2020. Annual research review: Defining and treating pediatric treatment-resistant depression. Journal of Child Psychology and Psychiatry 61, 312–332. doi:10.1111/jcpp.13202.

[7] Fitzpatrick, C., Lemieux, A., Smith, J., West, G.L., Bohbot, V., Asbridge, M., 2023. Is adolescent internet use a risk factor for the development of depression symptoms or vice-versa? Psychological Medicine 53, 6773–6779. doi:10.1017/S0033291723000284.

[8] Rice, F., Riglin, L., Lomax, T., Souter, E., Potter, R., Smith, D.J., Thapar, A.K., Thapar, A., 2019. Adolescent and adult differences in major depression symptom profiles. Journal of Affective Disorders 243, 175–181. doi:10.1016/j.jad.2018.09.015.

[9] Beck, A.T., 2008. The evolution of the cognitive model of depression and its neurobiological correlates. American Journal of Psychiatry 165, 969–977. doi:10.1176/appi.ajp.2008.08050721.

[10] LeMoult, J., Gotlib, I.H., 2019. Depression: A cognitive perspective. Clinical Psychology Review 69, 51–66. doi:10.1016/j.cpr.2018.06.008.

[11] Everaert, J., Grahek, I., Koster, E.H.W., 2017. Individual differences in cognitive control over emotional material modulate cognitive biases linked to depressive symptoms. Cognition & Emotion 31, 736–746. doi:10.1080/02699931.2016.1144562.

[12] Carter, B.T., Luke, S.G., 2020. Best practices in eye tracking research. International Journal of Psychophysiology 155, 49–62. doi:10.1016/j.ijpsycho.2020.05.010.

[13] Skaramagkas, V., Giannakakis, G., Ktistakis, E., Manousos, D., Karatzanis, I., Tachos, N.S., Tripoliti, E., Marias, K., Fotiadis, D.I., Tsiknakis, M., 2023. Review of eye tracking metrics involved in emotional and cognitive processes. IEEE Reviews in Biomedical Engineering 16, 260–277. doi:10.1109/RBME.2021.3066072.

[14] Hessels, R.S., Nuthmann, A., Nyström, M., Andersson, R., Niehorster, D.C., Hooge, I.T.C., 2025. The fundamentals of eye tracking part 1: The link between theory and research question. Behavior Research Methods 57, 16. doi:10.3758/s13428-024-02544-8.

[15] Armstrong, T., Olatunji, B.O., 2012. Eye tracking of attention in the affective disorders: A meta-analytic review and synthesis. Clinical Psychology Review 32, 704–723. doi:10.1016/j.cpr.2012.09.004.

[16] Suslow, T., Hußlack, A., Kersting, A., Bodenschatz, C.M., 2020. Attentional biases to emotional information in clinical depression: A systematic and meta-analytic review of eye tracking findings. Journal of Affective Disorders 274, 632–642. doi:10.1016/j.jad.2020.05.140.

[17] Akram, U., Ellis, J.G., Cau, G., Hershaw, F., Rajenthran, A., Lowe, M., Trommelen, C., Drabble, J., 2021. Eye tracking and attentional bias for depressive internet memes in depression. Experimental Brain Research 239, 575–581. doi:10.1007/s00221-020-06001-8.

[18] Klawohn, J., Bruchnak, A., Burani, K., Meyer, A., Lazarov, A., Bar-Haim, Y., Hajcak, G., 2020. Aberrant attentional bias to sad faces in depression and the role of stressful life events: Evidence from an eye-tracking paradigm. Behaviour Research and Therapy 135, 103762. doi:10.1016/j.brat.2020.103762.

[19] Sun, R., Fietz, J., Erhart, M., et al., 2024. Free-viewing gaze patterns reveal a mood-congruency bias in MDD during an affective fMRI/eye-tracking task. European Archives of Psychiatry and Clinical Neuroscience 274, 559–571. doi:10.1007/s00406-023-01608-8.

[20] Liu, X., Li, Y., Chen, Y., Xue, C., Fan, J., Zhang, J., Zhong, D., Dong, Q., Zheng, Z., Li, J., Jin, R., 2025. Attentional bias toward threatening stimuli in major depressive disorder: A free-viewing eye-tracking study. Journal of Affective Disorders 378, 352–359. doi:10.1016/j.jad.2025.02.039.

[21] Carvalho, N., Laurent, E., Noiret, N., Chopard, G., Haffen, E., Bennabi, D., Vandel, P., 2015. Eye movement in unipolar and bipolar depression: A systematic review of the literature. Frontiers in Psychology 6, 1809. doi:10.3389/fpsyg.2015.01809.

[22] Toh, W.L., Rossell, S.L., Castle, D.J., 2011. Current visual scanpath research: A review of investigations into the psychotic, anxiety, and mood disorders. Comprehensive Psychiatry 52, 567–579. doi:10.1016/j.comppsych.2010.12.005.

[23] Takahashi, J., Hirano, Y., Miura, K., Morita, K., Fujimoto, M., Yamamori, H., Yasuda, Y., Kudo, N., Shishido, E., Okazaki, K., Shiino, T., Nakao, T., Kasai, K., Hashimoto, R., Onitsuka, T., 2021. Eye movement abnormalities in major depressive disorder. Frontiers in Psychiatry 12, 673443. doi:10.3389/fpsyt.2021.673443.

[24] Wang, Y., Lyu, H.L., Tian, X.H., Lang, B., Wang, X.Y., St Clair, D., Wu, R., Zhao, J., 2022. The similar eye movement dysfunction between major depressive disorder, bipolar depression and bipolar mania. The World Journal of Biological Psychiatry 23, 689–702. doi:10.1080/15622975.2022.2025616.

[25] Gao, M., Xin, R., Wang, Q., Gao, D., Wang, J., Yu, Y., 2023. Abnormal eye movement features in patients with depression: Preliminary findings based on eye tracking technology. General Hospital Psychiatry 84, 25–30. doi:10.1016/j.genhosppsych.2023.04.010.

[26] Zhang, D., Liu, X., Xu, L., Li, Y., Xu, Y., Xia, M., Qian, Z., Tang, Y., Liu, Z., Chen, T., Liu, H., Zhang, T., Wang, J., 2022. Effective differentiation between depressed patients and controls using discriminative eye movement features. Journal of Affective Disorders 307, 237–243. doi:10.1016/j.jad.2022.03.077.

[27] Chen, S., Zhou, R., Cui, H., Chen, X., 2013. Deficits in cue detection underlie event-based prospective memory impairment in major depression: An eye tracking study. Psychiatry Research 209, 453–458. doi:10.1016/j.psychres.2013.01.015.

[28] Nemeth, V.L., Csete, G., Drotos, G., Greminger, N., Janka, Z., Vecsei, L., Must, A., 2016. The effect of emotion and reward contingencies on relational memory in major depression: An eye-movement study with follow-up. Frontiers in Psychology 7, 1849. doi:10.3389/fpsyg.2016.01849.

[29] Lu, Y., Song, L., Huang, C., Fan, T., Huang, J., Zhang, L., Luo, X., Li, Y., Shen, Y., 2024. The association between eye movement characteristics and cognitive function in adolescents with major depressive disorder. Psychiatry Research: Neuroimaging 345, 111914. doi:10.1016/j.pscychresns.2024.111914.

[30] Smith, K.G., Schmidt, J., Wang, B., Henderson, J.M., Fridriksson, J., 2018. Task-related differences in eye movements in individuals with aphasia. Frontiers in Psychology 9, 2430. doi:10.3389/fpsyg.2018.02430.

[31] Shear, M.K., Vander Bilt, J., Rucci, P., Endicott, J., Lydiard, B., Otto, M.W., Pollack, M.H., Chandler, L., Williams, J., Ali, A., Frank, D.M., 2001. Reliability and validity of a structured interview guide for the hamilton anxiety rating scale (SIGH-A). Depression and Anxiety 13, 166–178. doi:10.1002/da.1033.

[32] Khan, A., Bhat, A., Kolts, R., Thase, M.E., Brown, W., 2010. Why has the antidepressant–placebo difference in antidepressant clinical trials diminished over the past three decades? CNS Neuroscience & Therapeutics 16, 217–226. doi:10.1111/j.1755-5949.2010.00151.x.

[33] Shafer, A.B., 2006. Meta-analysis of the factor structures of four depression questionnaires: Beck, CES-D, Hamilton, and Zung. Journal of Clinical Psychology 62, 123–146. doi:10.1002/jclp.20213.

[34] Seemüller, F., Schennach, R., Musil, R., Obermeier, M., Adli, M., Bauer, M., Brieger, P., Laux, G., Gaebel, W., Falkai, P., Riedel, M., Möller, H.J., 2023. A factor analytic comparison of three commonly used depression scales (HAMD, MADRS, BDI) in a large sample of depressed inpatients. BMC Psychiatry 23, 548. doi:10.1186/s12888-023-05038-7.

[35] Zimmerman, M., Martinez, J.H., Young, D., Chelminski, I., Dalrymple, K., 2013. Severity classification on the hamilton depression rating scale. Journal of Affective Disorders 150, 384–388. doi:10.1016/j.jad.2013.04.028.

[36] Zimmerman, M., Clark, H.L., Multach, M.D., Walsh, E., Rosenstein, L.K., Gazarian, D., 2016. Symptom severity and the generalizability of antidepressant efficacy trials: Changes during the past 20 years. Journal of Clinical Psychopharmacology 36, 153–156. doi:10.1097/JCP.0000000000000466.

[37] Lecrubier, Y., Bech, P., 2007. The Ham D 6 is more homogenous and as sensitive as the Ham D 17. European Psychiatry 22, 252–255. doi:10.1016/j.eurpsy.2007.01.1218.

[38] Gong, X., Huang, Y.X., Wang, Y., Luo, Y.J., 2011. Revision of the chinese facial affective picture system. Chinese Mental Health Journal 25, 40–46. doi:10.3969/j.issn.1000-6729.2011.01.011.

Supplementary information



Fig. S1. Top-5 classifiers for the exploratory two-class task (no/mild versus moderate/severe strata). Models are ordered by the sum of their ranks across accuracy, macro F1, and AUC (top 5 shown, left to right); grouped bars show the three metric values of each model. Values are means over five folds of stratified cross-validation. Abbreviations as in Fig. 3.

Table S1. Operational definitions and interpretations of extracted eye-movement features.

Domain	Feature	Operational calculation	Interpretation
F1	Complete AOI coverage proportion	For each free-viewing display, code whether gaze entered all three emotional face AOIs at least once; average this binary value across displays.	Completeness of emotional AOI sampling during encoding; higher values indicate more complete coverage.
F1	AOI allocation entropy	Compute Shannon entropy of dwell-time proportions across negative, neutral, and positive AOIs for each display; average across displays.	Balance of gaze allocation across emotional AOIs; higher values indicate more even sampling.
F1	AOI allocation imbalance	Compute the L1 distance between the observed dwell-time distribution across the three emotional AOIs and a uniform distribution of 1/3, 1/3, 1/3; average across displays.	Deviation from balanced AOI allocation; higher values indicate more uneven sampling.
F1	First-pass AOI coverage	During the first pass through AOIs before the first revisit, count the number of unique emotional AOIs visited and divide by three; average across displays.	Breadth of initial visual sampling; higher values indicate broader first-pass coverage.
F1	Time to full AOI coverage	Compute the latency from display onset until all three emotional AOIs have been entered at least once; average across displays with full coverage.	Efficiency of completing emotional AOI coverage; higher values indicate slower coverage.
F2	Negative-AOI disengagement latency	After the first entry into the negative AOI, compute the duration until gaze first leaves the negative AOI; average across displays with a negative-AOI hit.	Disengagement from negative information; higher values indicate slower disengagement from the negative AOI.
F2	High-low load change in negative-positive gaze allocation bias	Compute negative-positive gaze allocation bias separately for high-load displays with four stimuli and low-load displays with two stimuli, then subtract low-load from high-load values.	Load-related change in negative versus positive gaze allocation; positive values indicate stronger negative bias under higher encoding load.
F2	Negative AOI visit duration	Identify continuous visits to the negative AOI and compute the mean duration of these visits; average across displays.	Maintenance of gaze on negative emotional information; higher values indicate longer negative-AOI visits.
F2	Negative-neutral gaze allocation bias	Compute the difference between the negative AOI dwell proportion and the neutral AOI dwell proportion for each display; average across displays.	Relative allocation of gaze to negative versus neutral stimuli; positive values indicate greater negative than neutral allocation.
F2	Negative-positive gaze allocation bias	Compute the difference between the negative AOI dwell proportion and the positive AOI dwell proportion for each display; average across displays.	Relative allocation of gaze to negative versus positive stimuli; positive values indicate greater negative than positive allocation.
F2	Negative stickiness after first hit	After the first hit on the negative AOI, compute the proportion of the remaining viewing time spent on the negative AOI; average across displays with a negative-AOI hit.	Post-engagement maintenance on negative information; higher values indicate stronger negative stickiness after first hit.
F2	Late-early change in negative-positive gaze allocation bias	Compute negative-positive gaze allocation bias separately for late and early task phases, then subtract early-phase from late-phase values.	Task-phase change in negative versus positive gaze allocation; positive values indicate increasing negative bias over time.
F2	Late-early change in negative stickiness	Compute negative stickiness after first hit separately for late and early task phases, then subtract early-phase from late-phase values.	Task-phase change in negative attention stickiness; positive values indicate increasing negative stickiness over time.
F2	Negative AOI revisit count	Count repeated entries into the negative AOI after the initial negative-AOI visit; average across displays.	Recurrent return to negative information; higher values indicate more negative-AOI revisits.
F3	Inter-AOI switch rate	Count transitions between different emotional AOIs and divide by valid free-viewing time in seconds; average across displays.	Frequency of switching among emotional AOIs; higher values indicate more frequent inter-AOI exploration.
F3	AOI transition entropy	Compute Shannon entropy over observed AOI-to-AOI transition categories for each display; average across displays.	Diversity of inter-AOI transition patterns; higher values indicate more varied transition sequences.

Continued on next page

Supplementary Table S1 (continued)

Domain	Feature	Operational calculation	Interpretation
F3	Normalized gaze convex-hull area	Compute the convex-hull area covering valid gaze samples during free viewing and normalize it by the fixed stimulus/display coordinate scale; average across displays.	Global spatial spread of visual exploration; higher values indicate a wider exploration field.
F3	Mean saccade amplitude	Compute the mean amplitude of saccadic gaze shifts during free viewing and normalize by the fixed stimulus/display coordinate scale; average across displays.	Spatial extent of saccadic exploration; higher values indicate larger gaze shifts.
F3	Normalized scanpath length	Sum consecutive gaze-step distances during free viewing and normalize by the fixed stimulus/display coordinate scale; average across displays.	Overall amount of gaze movement during encoding; higher values indicate longer scanpaths.
F3	AOI transition duration	For inter-AOI transitions, compute the elapsed time from leaving one AOI to entering the next AOI; average across transitions and displays.	Temporal cost of moving between AOIs; higher values indicate slower inter-AOI transitions.
F3	Outside-AOI transition pause duration	During inter-AOI transitions, compute time spent outside all emotional AOIs before the next AOI is entered; average across transitions and displays.	Off-AOI pause during transitions; higher values indicate longer non-AOI dwell while shifting between emotional faces.
F3	Transition tortuosity	For inter-AOI transitions, divide the actual gaze path length by the straight-line distance between the source and destination AOI centers; average across transitions.	Geometric efficiency of inter-AOI transitions; higher values indicate more curved or indirect transition trajectories.
F3	Within-AOI gaze dispersion	Within AOI visits, compute the root-mean-square distance of gaze samples from the local AOI-visit centroid and normalize by AOI/display scale; average across visits and displays.	Local spatial dispersion of gaze within AOIs; higher values indicate less concentrated within-AOI viewing.
F3	Within-AOI gaze step size	Within AOI visits, compute the mean distance between consecutive gaze samples and normalize by AOI/display scale; average across visits and displays.	Local within-AOI gaze wandering; higher values indicate greater moment-to-moment movement within AOIs.
R1	Target-hit proportion	Code whether gaze ever entered the cued target AOI on each recall trial and average across trials.	Whether the recalled target was successfully reached; higher values indicate more frequent target hits.
R1	Initial target-entry accuracy	For each recall trial, identify the first candidate AOI entered and code whether it was the cued target; average across trials.	Accuracy of initial memory-guided orienting; higher values indicate more accurate first target entry.
R1	Pre-target non-target AOI count	Before first target entry, count entries into non-target candidate AOIs; average across recall trials.	Pre-target search errors or distractor sampling; higher values indicate more non-target exploration before target acquisition.
R1	Pre-target search path length	Before first target entry, sum consecutive gaze-step distances and normalize by the fixed stimulus/display coordinate scale; average across trials.	Spatial search effort before acquiring the target; higher values indicate longer pre-target search paths.
R1	Strict target-fixation latency	Compute the latency from recall onset to the first strict fixation event within the target AOI; average across trials with strict target fixation.	Time required to form stable fixation on the recalled target; higher values indicate delayed stable target fixation.
R1	Target-entry latency	Compute the latency from recall onset to the first entry into the cued target AOI; average across trials with target entry.	Speed of memory-guided target acquisition; higher values indicate delayed target entry.
R2	Longest continuous target run	For each recall trial, compute the longest continuous dwell episode within the target AOI; average across trials.	Sustained continuous gaze maintenance on the recalled target; higher values indicate longer uninterrupted target holding.
R2	Off-target dwell proportion	Compute the proportion of recall viewing time spent in non-target candidate AOIs; average across trials.	Allocation to competing non-target locations; higher values indicate stronger off-target allocation.
R2	Post-hit off-target ratio	After first target entry, compute the proportion of the remaining recall time spent in non-target AOIs; average across trials with target entry.	Off-target drift after target acquisition; higher values indicate weaker maintenance of the recalled target.

Continued on next page

Supplementary Table S1 (continued)

Domain	Feature	Operational calculation	Interpretation
R2	Target dwell proportion	Compute the proportion of recall viewing time spent within the cued target AOI; average across trials.	Preferential allocation to the recalled target; higher values indicate stronger target prioritization.
R2	Target lock-on ratio	After first target entry, compute the proportion of the remaining recall time spent within the target AOI; average across trials with target entry.	Maintenance of gaze on the target after acquisition; higher values indicate stronger target lock-on.
R2	Target re-entry count after first hit	After first target entry, count subsequent re-entries into the target AOI following exits; average across trials.	Return behavior after leaving the target; higher values indicate more repeated target reacquisition.
R3	Candidate target-area fixation dispersion	Within any candidate AOI during recall, compute the root-mean-square distance of strict fixation samples from their local centroid and normalize by AOI/display scale; average across candidate AOIs/trials.	General fixation dispersion within candidate target areas, including target and non-target AOIs; higher values indicate lower local stability.
R3	Candidate target-area fixation step size	Within any candidate AOI during recall, compute the mean consecutive step size of strict fixation samples and normalize by AOI/display scale; average across candidate AOIs/trials.	General fixation wandering within candidate target areas; higher values indicate less stable local fixation.
R3	Target-area stability ellipse area	For strict fixation samples within the target AOI, compute a fixation-stability ellipse area from the horizontal and vertical gaze dispersion and normalize by AOI/display scale; average across trials.	Spatial spread of target-area fixation; higher values indicate lower fixation stability.
R3	Strict target-fixation count	Count strict fixation events within the target AOI on each recall trial; average across trials.	Frequency of stable target fixations; higher values indicate more target fixation episodes.
R3	Total strict target-fixation duration	Sum the duration of strict fixation events within the target AOI on each recall trial; average across trials.	Total stable fixation time on the recalled target; higher values indicate more sustained stable target viewing.
R3	Target-area fixation dispersion	For strict fixation samples within the target AOI, compute the root-mean-square distance from the local fixation centroid and normalize by AOI/display scale; average across trials.	Local fixation dispersion within the target area; higher values indicate lower fixation stability.
R3	Within-target fixation step size	For strict fixation samples within the target AOI, compute the mean distance between consecutive samples and normalize by AOI/display scale; average across trials.	Moment-to-moment instability within target fixation; higher values indicate greater local fixation wandering.
R3	Total strict target-fixation duration	Sum the duration of strict fixation events within the target AOI on each recall trial; average across trials.	Total stable fixation time on the recalled target; higher values indicate more sustained stable target viewing.

Table S2. Supplementary results from ordered-severity models.

Domain	Supplementary feature	Standardized β (95% CI)	p	q_{FDR}
F1	Complete AOI coverage proportion	−0.538 (−0.679, −0.398)	< 0.001	< 0.001
F1	AOI allocation imbalance	0.603 (0.477, 0.729)	< 0.001	< 0.001
F2	Negative-neutral gaze allocation bias	0.118 (−0.031, 0.267)	0.120	0.157
F2	Negative AOI revisit count	−0.215 (−0.368, −0.062)	0.006	0.009
F2	Negative-AOI disengagement latency	−0.072 (−0.238, 0.093)	0.392	0.466
F2	Late-early change in negative-positive gaze allocation bias	−0.129 (−0.295, 0.036)	0.126	0.157
F2	Late-early change in negative stickiness	−0.006 (−0.157, 0.144)	0.933	0.933
F2	High-low load change in negative-positive gaze allocation bias	0.019 (−0.143, 0.181)	0.822	0.857
F2	High-low load change in negative stickiness	−0.061 (−0.218, 0.097)	0.450	0.512
F3	Normalized scanpath length	0.326 (0.182, 0.471)	< 0.001	< 0.001
F3	Normalized gaze convex-hull area	0.146 (−0.015, 0.306)	0.075	0.104
F3	AOI transition duration	0.409 (0.255, 0.563)	< 0.001	< 0.001
F3	Outside-AOI transition pause duration	0.288 (0.159, 0.416)	< 0.001	< 0.001
F3	Within-AOI gaze step size	0.308 (0.168, 0.449)	< 0.001	< 0.001
F3	Mean saccade amplitude	−0.348 (−0.493, −0.204)	< 0.001	< 0.001
R1	Target-hit proportion	−0.577 (−0.690, −0.463)	< 0.001	< 0.001
R1	Pre-target non-target AOI count	0.442 (0.318, 0.566)	< 0.001	< 0.001
R1	Pre-target search path length	0.550 (0.416, 0.684)	< 0.001	< 0.001
R2	Off-target dwell proportion	0.390 (0.269, 0.511)	< 0.001	< 0.001
R2	Longest continuous target run	−0.516 (−0.646, −0.386)	< 0.001	< 0.001
R2	Target re-entry count after first hit	−0.050 (−0.190, 0.090)	0.483	0.525
R3	Strict target-fixation count	−0.215 (−0.360, −0.070)	0.004	0.006
R3	Total strict target-fixation duration	−0.683 (−0.785, −0.580)	< 0.001	< 0.001
R3	Candidate target-area fixation dispersion	0.251 (0.100, 0.402)	0.001	0.002
R3	Candidate target-area fixation step size	0.143 (−0.008, 0.294)	0.063	0.093

Standardized coefficients were obtained from ordered-severity regression models adjusted for age, gender, and years of education, using heteroscedasticity-consistent robust standard errors. FDR correction was applied across all 25 features. Supplementary features were used to evaluate convergent patterns within the same mechanism-informed domain. FDR, false discovery rate; CI, confidence interval.

Table S3. Associations between continuous HAMD-17 scores and supplementary eye-movement features.

Feature domain	Supplementary feature	Spearman correlation			Adjusted regression model		
		ρ_s	p	q_{FDR}	β (95% CI)	p	q_{FDR}
F1	Complete AOI coverage proportion	−0.600	< 0.001	< 0.001	−0.490 (−0.627, −0.354)	< 0.001	< 0.001
F1	AOI allocation imbalance	0.614	< 0.001	< 0.001	0.550 (0.418, 0.682)	< 0.001	< 0.001
F2	Negative–neutral gaze allocation bias	0.151	0.060	0.088	0.113 (−0.034, 0.261)	0.132	0.171
F2	Negative AOI revisit count	−0.229	0.004	0.007	−0.230 (−0.375, −0.085)	0.002	0.003
F2	Negative-AOI disengagement latency	−0.107	0.187	0.222	0.037 (−0.152, 0.226)	0.701	0.752
F2	Late–early change in negative–positive gaze allocation bias	−0.109	0.176	0.221	−0.106 (−0.275, 0.064)	0.222	0.271
F2	Late–early change in negative stickiness	0.029	0.717	0.752	0.001 (−0.159, 0.161)	0.991	0.991
F2	High–low load change in negative–positive gaze allocation bias	0.018	0.820	0.820	−0.011 (−0.168, 0.146)	0.890	0.910
F2	High–low load change in negative stickiness	−0.060	0.461	0.507	−0.088 (−0.258, 0.082)	0.312	0.371
F3	Normalized scanpath length	0.213	0.008	0.012	0.253 (0.096, 0.411)	0.002	0.003
F3	Normalized gaze convex-hull area	0.107	0.183	0.222	0.121 (−0.052, 0.293)	0.170	0.214
F3	AOI transition duration	0.343	< 0.001	< 0.001	0.344 (0.183, 0.506)	< 0.001	< 0.001
F3	Outside-AOI transition pause duration	0.312	< 0.001	< 0.001	0.243 (0.121, 0.364)	< 0.001	< 0.001
F3	Within-AOI gaze step size	0.225	0.005	0.008	0.242 (0.090, 0.394)	0.002	0.003
F3	Mean saccade amplitude	−0.303	< 0.001	< 0.001	−0.302 (−0.454, −0.150)	< 0.001	< 0.001
R1	Target-hit proportion	−0.571	< 0.001	< 0.001	−0.547 (−0.671, −0.422)	< 0.001	< 0.001
R1	Pre-target non-target AOI count	0.533	< 0.001	< 0.001	0.423 (0.299, 0.547)	< 0.001	< 0.001
R1	Pre-target search path length	0.618	< 0.001	< 0.001	0.485 (0.339, 0.632)	< 0.001	< 0.001
R2	Off-target dwell proportion	0.415	< 0.001	< 0.001	0.382 (0.252, 0.511)	< 0.001	< 0.001
R2	Longest continuous target run	−0.525	< 0.001	< 0.001	−0.495 (−0.651, −0.339)	< 0.001	< 0.001
R2	Target re-entry count after first hit	−0.031	0.698	0.749	−0.033 (−0.182, 0.116)	0.665	0.732
R3	Strict target-fixation count	−0.185	0.021	0.032	−0.169 (−0.324, −0.014)	0.033	0.048
R3	Total strict target-fixation duration	−0.661	< 0.001	< 0.001	−0.622 (−0.755, −0.489)	< 0.001	< 0.001
R3	Candidate target-area fixation dispersion	0.273	< 0.001	0.001	0.280 (0.130, 0.430)	< 0.001	< 0.001
R3	Candidate target-area fixation step size	0.150	0.063	0.090	0.173 (0.014, 0.332)	0.033	0.048

ρ_s denotes Spearman correlation with continuous HAMD-17 total score. Adjusted β values were estimated using separate linear regression models with standardized HAMD-17 total score as the predictor and each standardized eye-movement feature as the outcome, adjusting for age, sex, and years of education. Robust standard errors were used. The β coefficient represents the SD difference in the eye-movement feature associated with a one-SD increase in HAMD-17 total score. FDR correction was applied separately to the Spearman correlation tests and the adjusted regression tests across all 44 eye-movement features, including 19 primary and 25 supplementary features. HAMD-17, the 17-item Hamilton Depression Rating Scale; FDR, false discovery rate; CI, confidence interval.

Table S4. Sensitivity analyses for primary eye-movement features.

Features domain	Features	Unadjusted model		HAMA-adjusted model		Extreme-excluded	
		β (95% CI)	q_{FDR}	β (95% CI)	q_{FDR}	β (95% CI)	q_{FDR}
F1	Time to full AOI coverage	0.284 (0.143, 0.425)	< 0.001	0.265 (0.084, 0.446)	0.007	0.302 (0.159, 0.444)	< 0.001
F1	First-pass AOI coverage ratio	−0.446 (−0.596, −0.296)	< 0.001	−0.530 (−0.744, −0.316)	< 0.001	−0.442 (−0.603, −0.281)	< 0.001
F1	AOI allocation entropy	−0.600 (−0.732, −0.468)	< 0.001	−0.747 (−0.930, −0.565)	< 0.001	−0.618 (−0.750, −0.486)	< 0.001
F2	Negative-positive gaze allocation bias	0.142 (−0.029, 0.313)	0.130	0.140 (−0.080, 0.359)	0.269	0.071 (−0.070, 0.211)	0.341
F2	Negative AOI visit duration	−0.077 (−0.237, 0.083)	0.364	−0.165 (−0.368, 0.038)	0.163	−0.107 (−0.273, 0.059)	0.247
F2	Negative stickiness after first hit	−0.008 (−0.179, 0.162)	0.923	−0.055 (−0.281, 0.171)	0.668	−0.049 (−0.219, 0.122)	0.575
F3	Inter-AOI switch rate	−0.508 (−0.653, −0.363)	< 0.001	−0.545 (−0.721, −0.369)	< 0.001	−0.527 (−0.674, −0.380)	< 0.001
F3	AOI transition diversity	−0.548 (−0.678, −0.418)	< 0.001	−0.643 (−0.829, −0.457)	< 0.001	−0.560 (−0.696, −0.425)	< 0.001
F3	Transition tortuosity	0.315 (0.186, 0.445)	< 0.001	0.344 (0.168, 0.521)	< 0.001	0.366 (0.209, 0.523)	< 0.001
F3	Within-AOI gaze dispersion	0.158 (0.014, 0.302)	0.043	0.132 (−0.064, 0.328)	0.253	0.143 (−0.005, 0.291)	0.078
R1	First target-entry latency	0.574 (0.435, 0.713)	< 0.001	0.628 (0.477, 0.779)	< 0.001	0.605 (0.468, 0.742)	< 0.001
R1	Initial target-entry accuracy	−0.599 (−0.709, −0.490)	< 0.001	−0.718 (−0.863, −0.574)	< 0.001	−0.600 (−0.706, −0.493)	< 0.001
R1	First target-fixation latency	0.594 (0.461, 0.728)	< 0.001	0.666 (0.522, 0.810)	< 0.001	0.609 (0.487, 0.731)	< 0.001
R2	Target dwell proportion	−0.699 (−0.794, −0.604)	< 0.001	−0.753 (−0.884, −0.623)	< 0.001	−0.678 (−0.778, −0.578)	< 0.001
R2	Target lock-on ratio	−0.474 (−0.607, −0.340)	< 0.001	−0.450 (−0.615, −0.285)	< 0.001	−0.409 (−0.538, −0.280)	< 0.001
R2	Post-hit off-target ratio	0.233 (0.125, 0.340)	< 0.001	0.245 (0.029, 0.461)	0.041	0.257 (0.108, 0.406)	0.001
R3	Target-area fixation dispersion	0.185 (0.033, 0.337)	0.025	0.083 (−0.121, 0.286)	0.506	0.176 (0.015, 0.337)	0.047
R3	Within-target fixation step size	0.088 (−0.065, 0.242)	0.289	0.029 (−0.161, 0.219)	0.767	0.109 (−0.052, 0.269)	0.233
R3	Target-area stability ellipse area	0.138 (−0.032, 0.308)	0.133	0.082 (−0.154, 0.318)	0.555	0.083 (−0.075, 0.240)	0.339

Sensitivity analyses were conducted for prespecified primary features. Unadjusted models included ordered severity as the only predictor. HAMA-adjusted models additionally included HAMA total scores alongside age, gender, and years of education. Extreme-excluded models repeated the primary analyses after excluding extreme eye-movement values. HAMA, Hamilton Anxiety Rating Scale; FDR, false discovery rate; CI, confidence interval.

Table S5. Recall-task performance across depression severity groups.

	No (n = 32)	Mild (n = 39)	Moderate (n = 49)	Severe (n = 35)	H	p
Correct recall trials out of 12, mean (SD)	11.09 (1.06)	8.95 (2.05)	6.98 (2.84)	5.03 (2.99)	71.97	< 0.001

Correct recall trials were defined as trials in which the participant oriented gaze to the cued target location during the recall phase. Group comparisons were performed using the Kruskal–Wallis test. H denotes the Kruskal–Wallis test statistic.