Where Meanings Converge and Diverge: Revisiting Linguistic Relativity Across 81 Languages

Weiran Wang, Wenjuan Qin, and Haitao Liu*

Fudan University, Shanghai 200433, China

October 22, 2026

arXiv:2610.30778v1 [cs.CL] 22 Oct 2026

Abstract

Do people who speak different languages organize concepts in similar ways? This question lies at the heart of the Sapir–Whorf hypothesis and debates about linguistic relativity. Shared basic meanings, however, need not imply shared relations among them. We compared relations among 196 basic concepts across 81 languages, using word embeddings from two multilingual models to quantify conceptual structure expressed in language. Cross-language correspondence was greater for sociocultural meanings than for meanings concerning nature and motion, a contrast recurring across models and classifications. Within the sociocultural domain, relational differences increased with geographic separation; less closely related languages differed more across all domains. These findings suggest that universality and relativity are interwoven across conceptual domains. They inform an account in which shared cognitive resources interact with linguistic and local experience, and predict greater cross-language correspondence in human semantic judgments for sociocultural than for nature and motion meanings.

Keywords: linguistic relativity; Sapir–Whorf hypothesis; conceptual organization; multilingual comparison; representational similarity

Introduction

Concepts are mental representations through which people recognize objects, understand events, and generalize beyond what they have directly experienced (Murphy & Medin, 1985; Murphy, 2002). Whether people who speak different languages organize these representations in similar ways is a central question about the human mind. Because concepts are expressed through language, linguistic diversity offers a window onto the relationship between shared cognitive capacities and different ways of organizing experience (Evans & Levinson, 2009; Malt & Majid, 2013). The Sapir–Whorf hypothesis, or linguistic relativity, places this relationship at the center of inquiry: how do linguistic distinctions relate to the ways speakers attend to, categorize, and understand the world (Lucy, 1992; Wolff & Holmes, 2011)? Shared bodily and perceptual capacities provide potential foundations for common conceptual organization (Barsalou, 2008; Spelke & Kinzler, 2007), while language-related differences have been observed in color discrimination, pitch representation, and event categorization (Athanasopoulos et al., 2015; Dolscheid et al., 2013; Winawer et al., 2007). These patterns can coexist: locomotion terms across languages reflect a common biomechanical distinction between walking and running (Malt et al., 2008), and infants are sensitive to multiple space–pitch mappings before acquiring language-specific conventions (Dolscheid et al., 2014). Such evidence motivates an account in which shared resources and linguistic experience jointly contribute to meaning organization (Andrews et al., 2009; Wolff & Holmes, 2011). This relationship extends beyond the availability of corresponding words. Translation equivalents can refer to corresponding objects or events while participating in different semantic neighborhoods and broader relational structures (Lewis et al., 2023; Malt & Majid, 2013; Thompson et al., 2020). Because relations among meanings contribute to conceptual coherence, categorization, and generalization, their correspondence is itself a psychologically consequential question (Murphy & Medin, 1985; Murphy, 2002). A broad assessment of linguistic relativity therefore calls for examining how shared and language-specific relations are distributed across different kinds of meaning.

*Corresponding author: Haitao Liu. Fudan University, Shanghai 200433, China. Email: htliu@163.com.

Word embeddings provide a scalable means of approaching this question by representing regularities in linguistic contexts as relations in a quantitative space (Landauer & Dumais, 1997; Lenci, 2018). Their psychological relevance rests on evidence that distributional structure captures aspects of human semantic similarity and perceptual knowledge, alongside complementary information supplied by direct experience (Andrews et al., 2009; Lewis et al., 2019; Riordan & Jones, 2011). More recently, Xu et al. (2025) showed that representations elicited through concept inference in language-prediction models capture human similarity judgments and category structure, and predict brain responses to object images. These findings support using language-derived representations to investigate aspects of conceptual organization across languages. Previous comparisons have identified shared lexical-network structure and culturally patterned variation (Jackson et al., 2019; Youn et al., 2016). Thompson et al. (2020) found greater semantic alignment in highly structured domains, including number and quantity, than for natural kinds and common actions; Lewis et al. (2023) further distinguished correspondence within semantic clusters from variability between them. A framework for integrating these findings treats language as a human-driven complex adaptive system, shaped by interacting cognitive, communicative, and environmental constraints (Beckner et al., 2009; Liu, 2014, 2018). In this framework, linguistic regularities are probabilistic and responsive to patterns of use, making graded and uneven correspondence across meanings theoretically relevant (Beckner et al., 2009; Liu, 2018). Here, we compare relations among 196 basic lexical concepts across 81 languages using two multilingual encoders. We ask which contrasts among Biological Core, Nature & Motion, Socio-Cultural, and Grammar & Function recur across representational and classificatory choices, and how these patterns relate jointly to geographic distance and estimated language relatedness. The recurring result is greater cross-language correspondence among the sampled Socio-Cultural relations than among Nature & Motion relations. By locating commonality and variation within a shared basic inventory, this comparison supplies a domain-sensitive pattern for accounts of linguistic relativity to explain. It also predicts a corresponding contrast in matched human semantic-relation judgments, connecting the broad comparison to the psychological organization of concepts.

Research Transparency Statement

General Disclosures

Conflicts of interest: The authors declare no competing interests.

Funding: The authors declare no funding.

Artificial intelligence: Multilingual BERT and XLM-R-large were used as research tools to extract lexical representations. Generative AI tools, including GPT-5.6-Luna, Opus-5, and Sonnet-5, assisted with code revision, language editing, drafting support, and LaTeX formatting. All scientific content, methodological decisions, data analyses, and interpretations remain the sole responsibility of the authors, who reviewed and verified all AI-assisted text.

Ethics: This study used publicly available, non-identifiable lexical resources and language metadata. No human participants were recruited, and no new behavioral or neural data were collected.

Study Disclosures

Preregistration: No aspects of this study were preregistered.

Materials: The concept and language inventories, concept mappings, and analysis specifications are provided in the Supplemental Material and accompanying source package, available via Figshare: https://figshare.com/s/63bae7225054e9718549.

Data: Data and machine-readable summaries supporting the reported analyses of 81 languages and 196 concepts are available via https://figshare.com/s/63bae7225054e9718549. The accompanying materials document the provenance of the archived model outputs used in the analyses.

Analysis scripts: The accompanying source package provides the analysis scripts and specifications supporting the reported estimates: https://figshare.com/s/63bae7225054e9718549.

Method

Languages, Concepts, and Semantic Domains

We compared relations among translation equivalents in a computational, cross-sectional design. Lexical forms for the starting inventory of 82 languages and 207 basic concepts were downloaded from Wiktionary's language-specific Swadesh lists (Wiktionary contributors, n.d.), following the Swadesh basic-vocabulary tradition (Swadesh, 1955). Languages were selected for model coverage, usable lexical material, geographic coordinates, and relatedness metadata. This coverage-based sample enables broad comparison without representing a probability sample of the world's languages. No new behavioral or neural responses were collected. Archived lexical forms and Wiktionary concept numbering were retained, including identifiers distinguishing entries with the same English gloss.

Vectors were screened for dimensionality, finite values, zero content, and parseability; the first valid vector was retained for repeated identifiers. Javanese was excluded because identical vectors, rounded to seven decimals, comprised more than half its valid representations (.607). Twelve all-zero observations affected 11 concept identifiers. Excluding these identifiers across languages yielded common support of 81 languages, 196 concepts, and 3,240 unordered language pairs in both encoders. Sample size followed usable common coverage. Complete inventories and exclusions appear in the Supplemental Material.

Four author-defined domains consolidated the topical annotations in the archived study inventory (Table 1). They are operational groupings, rather than independently established conceptual modules. We also used a reference-style grouping that reassigned 22 of the 207 input concepts, including six personal-pronoun items from Biological Core to Grammar & Function; kinship remained Biological Core. The groupings agreed on 185 input concepts, yielding a complete-case consensus subset of 175 concepts. Item assignments and their sources are documented in the Supplemental Material.

Table 1. Operational domains in the primary grouping. Counts refer to the 196 retained concepts; complete item assignments are provided in the Supplemental Material.

Domain	n	Included fields
Biological Core	67	Body, perception, bodily processes, people, kinship, personal pronouns
Nature & Motion	49	Nonhuman living kinds, environments, movement, physical change
Socio-Cultural	59	Properties, evaluation, color, time, actions, material culture, numerals
Grammar & Function	21	Demonstratives, interrogatives, quantifiers, relational/function items

Lexical Representations and Relational Dissimilarity

We extracted isolated lexical forms with mBERT (`bert-base-multilingual-cased`) and XLM-R-large (`xlm-roberta-large`) (Conneau et al., 2020; Devlin et al., 2019). Final-layer states were averaged over attended, non-special lexical subwords. Comma-delimited alternatives were encoded separately and averaged equally; forms without usable lexical tokens were missing. Both encoders used the same forms, pooling rules, and support, while retaining their own tokenizers and training histories. Extraction choices matter for interpreting these representations (Vulić et al., 2020).

Representational similarity analysis compared the ordering of semantic relations within each language (Beinborn & Choenni, 2020; Kriegeskorte et al., 2008). Vectors were centered on their language-specific mean within each concept set before constructing a cosine-dissimilarity matrix. The full inventory and each domain were centered separately; domain matrices were not subsets of an already centered global matrix. For languages i and j and concept set C, cross-language dissimilarity was

$$d_{ij,C} = 1 - \rho_{\text{Spearman}}\left[u(D_{i,C}), u(D_{j,C})\right], \tag{1}$$

where $u(D)$ contains upper-triangle entries in matched concept order, excluding the diagonal. Ties received average ranks. Lower values indicate more similarly ordered relations; the index ranges from zero to two and requires no coordinate alignment. Each language pair contributed one global and four domain outcomes per encoder. Concept-pair entries were not independent regression observations. No undefined outcomes occurred. Principal-component projections were descriptive only.

Geographic and Linguistic Distance Measures

Geographic distance was great-circle distance between archived language centroids. Genealogical distance indexed estimated language relatedness using lexical–phonological dissimilarities from the Automated Similarity Judgment Program (ASJP), derived from pointwise mutual-information scores for core-vocabulary forms (Jäger, 2018). Higher values indicate more distant estimated relationships. This proxy can reflect inheritance and borrowing and shares material with the semantic inventory; it is not a reconstructed family-tree distance. Both matrices were fixed, symmetric, and complete on the retained languages. Predictors were standardized and entered jointly, so slopes describe associations per standard-deviation increase in one distance, conditional on the other. Geographic separation does not directly measure contact, and word-form dissimilarity does not isolate inheritance.

Statistical Analysis and Robustness Checks

We fitted Gaussian Bayesian regressions with domain-specific intercepts and distance slopes. For each encoder, the domain model was

$$\begin{aligned} d_{ij,k} &\sim N(\mu_{ij,k}, \sigma^2), \\ \mu_{ij,k} &= \alpha_k + \beta_{\text{geo},k} z_{\text{geo},ij} + \beta_{\text{gen},k} z_{\text{gen},ij} + \tfrac{1}{2}(u_i + u_j). \end{aligned} \tag{2}$$

Here, i and j index languages, k indexes domains, and $z_{\text{geo},ij}$ and $z_{\text{gen},ij}$ are standardized geographic and genealogical distances. The domain-specific intercept α_k describes adjusted dissimilarity; the two β terms are total domain slopes. The shared language intercepts u_i and u_j account for repeated participation in language pairs, each with weight one half.

Language effects used a non-centered Gaussian parameterization and a sum-to-zero constraint. Biological Core was the domain reference; other intercepts and slopes were coded as increments. Encoder-specific global and domain fits used 3,240 and 12,960 rows, respectively. A pooled domain model (25,920 rows) included encoder, domain, both predictors, and interactions through Encoder × Domain × Predictor, with shared language intercepts and mBERT as the encoder reference. Adjusted means used centered predictors and zero language contribution. Total slopes and contrasts were derived within posterior draws; pooled draws supplied between-encoder comparisons. Models omitted an additional dyad effect and residual correlations across outcomes from the same dyad; intervals are conditional on this specification.

Priors were $N(.9, .2)$ for the reference intercept, $N(0, .2)$ for other fixed coefficients (mean, standard deviation), positive Student-$t(3, 0, .5)$ for the language-effect scale, and exponential rate one for residual standard deviation. CmdStan 2.38.0 (Carpenter et al., 2017) used four No-U-Turn chains, each with 3,000 warm-up and 3,000 retained iterations. Estimates are posterior means with equal-tail 95% credible intervals (CrIs). Across the complete-support model fits, the maximum $\widehat{R}$ was 1.008, the minimum bulk effective sample size was 394, and there were no divergent transitions or maximum-tree-depth saturations; the minimum reported E-BFMI was .633. Convergence and posterior predictive checks, full specifications, and coefficients are reported in the Supplemental Material.

We recomputed RSA after regrouping concepts and on the consensus subset, permuted concept labels within languages and domains 2,000 times, and widened intercept/coefficient prior scales from .2 to .5. Permutation probabilities used the add-one correction. Language-omission summaries recomputed means without refitting models; paired-dyad bootstraps did not account for dependence between pairs sharing a language. Secondary English-frequency models used item-level cosine dissimilarity, and archived size/layer checks used earlier vectors; neither constitutes a matched replication of the primary RSA analysis. Full records and extraction provenance are retained in the Supplemental Material.

For illustration, we screened mBERT language triplets for an anchor, a comparison with low genealogical distance but large geographic separation, and a geographically close comparison with higher genealogical distance. We selected the largest positive gap favoring the comparison with lower genealogical distance for Grammar & Function and the geographically close comparison for Socio-Cultural. These gaps average item-level cosine similarities between language-centered vectors, a different measure from domain RSA. Distance thresholds and complete candidate records are provided in the Supplemental Material.

Results

Semantic Similarity Across Languages Varies by Domain

The shared inventory yielded limited overall correspondence but a recurring difference between semantic domains. Across the 3,240 language pairs, adjusted global RSA dissimilarity was .910 (95% CrI [.909, .911]) in mBERT and .932 [.931, .933] in XLM-R-large. These values describe substantial variation in the ordering of within-language relations, even among translation equivalents. Against this background, we compared the four domains on the same language support (Fig. 1).

In mBERT, Socio-Cultural concepts showed the lowest adjusted dissimilarity (.874, 95% CrI [.872, .877]), whereas Nature & Motion showed the highest (.924 [.922, .927]). Grammar & Function (.892 [.890, .895]) and Biological Core (.900 [.898, .903]) occupied intermediate positions. The Nature & Motion minus Socio-Cultural difference was .0503 [.0470, .0537]. Thus, relations among the sampled Socio-Cultural meanings corresponded more closely across languages than those among Nature & Motion meanings.

The same comparison held in XLM-R-large: Socio-Cultural dissimilarity was .922 [.920, .924], compared with .977 [.974, .979] for Nature & Motion, a difference of .0547 [.0513, .0580]. Biological Core and Grammar & Function remained intermediate but reversed their order relative to mBERT (Fig. 2; Supplemental Table S14). The pooled between-encoder difference in the Nature & Motion–Socio-Cultural contrast was .0044 [-.0003, .0090]. Agreement therefore concerned the direction of this domain comparison, alongside differences in the complete profile. Even the lower-dissimilarity domains remained far from identical across languages.

Regrouping some concepts preserved this direction. Under the reference-style grouping, raw Socio-Cultural versus Nature & Motion means were .883 versus .921 in mBERT and .938 versus .973 in XLM-R-large. Restricting the analysis to the 175 concepts assigned identically by both groupings yielded .888 versus .935 and .938 versus .977. Grammar & Function became the lowest-dissimilarity mBERT group in that subset, qualifying any fixed four-domain ranking. Concept-label calibration also distinguished observed correspondence from random labels: all 16 encoder-by-grouping-by-domain outcomes fell below all 2,000 permutations (Monte Carlo lower-tail $p < .001$). Language omission preserved the descriptive full-inventory orders, and wider priors left the main comparison essentially unchanged; full sensitivity results appear in the Supplemental Material.



Fig. 1. Cross-language variation in lexical relations. Multilingual BERT (mBERT) results for 81 languages and 196 concepts. The panels show (a) a descriptive principal-component projection of within-language centered vectors, colored by domain; (b) global representational similarity analysis (RSA) dissimilarities for 3,240 language pairs; and (c) domain distributions. The projection does not establish conceptual modules. Lower dissimilarity indicates more similar ordering of within-language relations; global and domain sets were centered separately.



Fig. 2. Domain patterns across encoders and concept groupings. Points show raw mean representational similarity analysis (RSA) dissimilarities on matched language and concept support. Colors identify encoders; solid lines/filled markers indicate the primary grouping and dashed lines/open markers the reference-style grouping. Error bars show the range of means after omitting each language in turn, not confidence or credible intervals. Adjusted estimates and credible intervals are reported in the text and Supplemental Material.

Semantic Differences Are Associated With Geographic Distance and Language Relatedness

We examined geographic and genealogical distance jointly using the model in Equation 2. Positive slopes indicate greater relational dissimilarity at greater distance, conditional on the other predictor.

Socio-Cultural dissimilarity increased with geographic separation in both encoders: the slope was .0114 (95% CrI [.0087, .0141]) in mBERT and .0081 [.0055, .0107] in XLM-R-large. Other domains showed less consistent geographic profiles. Nature & Motion had a negative slope in mBERT (-.0046 [-.0072, -.0019]), while Biological Core had a positive slope in XLM-R-large (.0058 [.0032, .0085]); the remaining intervals included zero (Fig. 3; Supplemental Table S15). Relatively low overall Socio-Cultural dissimilarity therefore coexisted with a recurring geographic gradient. Geography was not associated with a uniform increase across the inventory.

Genealogical distance showed a broader positive association. All four domain slopes were positive in both encoders, and every 95% CrI lay above zero. Estimates ranged from .02204 to .02566 in mBERT and .01601 to .02107 in XLM-R-large (Supplemental Table S16). Language pairs with greater estimated genealogical separation also tended to have more dissimilar relational patterns after accounting for geography. This relationship extended across domains, whereas geographic associations varied by domain and encoder. The two predictors thus describe different comparative patterns within the same language pairs.



Fig. 3. Geographic and language-relatedness associations. All panels show multilingual BERT (mBERT). Total domain slopes and 95% credible intervals are shown in (a); filled circles indicate the primary grouping and open squares the reference-style grouping. Panels (b) and (c) show raw representational similarity analysis (RSA) dyad outcomes and descriptive least-squares lines, distinct from the conditional Bayesian estimates in (a). Genealogical distance is the word-form-based relatedness proxy defined in Method. Complete estimates for both encoders are in Supplemental Tables S15 and S16.

Illustrative Comparisons Among Individual Languages

Two selected language triplets make these distance configurations concrete at the level of lexical items (Fig. 4). Each compares an anchor with a geographically distant language having lower genealogical distance and a geographically close language having higher genealogical distance. The examples were selected for large directional gaps in item-level cosine similarity, rather than for their representativeness of all language pairs.

For Grammar & Function, Dutch was the anchor, Afrikaans the comparison with lower genealogical distance, and Basque the geographically close comparison. Dutch–Afrikaans similarities exceeded Dutch–Basque similarities for many displayed items, including *who*, *when*, *how*, and *what*. In the Socio-Cultural example, Icelandic was compared with Marathi and Scots. Similarities with geographically closer Scots exceeded those with Marathi for selected items including *green*, *long*, *sharp*, *four*, and *five*.

These examples show how spatial separation and word-form relatedness can combine differently in particular comparisons. Their scores concern cross-language similarities of individual lexical vectors, whereas the main analyses compare patterns of within-language relations. They illustrate selected lexical configurations without independently testing the domain-level RSA results or identifying contact and inheritance as causes. Complete selection rules, qualifying candidates, and item records are retained in the Supplemental Material.



Fig. 4. Selected language comparisons illustrate lexical variation. Multilingual BERT (mBERT) examples selected for large directional gaps. The locations of the two triplets are shown in (a). Grammar & Function items in (b) compare Dutch with Afrikaans and Basque; Socio-Cultural items in (c) compare Icelandic with Marathi and Scots. Scores are item-level cosine similarities between language-centered vectors, not domain representational similarity analysis (RSA). Filled markers indicate the comparison with lower genealogical distance and open markers the geographically closer comparison. These selected examples are descriptive.

Discussion

Shared Meanings, Different Relations

Do people who speak different languages organize basic concepts in similar ways? Our comparison identifies a recurring domain contrast in language-derived representations: relations among the sampled Socio-Cultural meanings corresponded more closely across languages than those among Nature & Motion meanings. This pattern agrees with Thompson et al.'s (2020) finding that highly structured meanings can align more closely than natural kinds and common actions. It also complements Lewis et al.'s (2023) distinction between local similarity and broader variability by showing that the extent of correspondence differs across the present groups of basic meanings. The contrast recurred across two encoders and changes in classification and concept support. For linguistic relativity, the contribution is a clearer comparative profile of where relations are relatively shared and where they differ, bringing variation within the lexicon into the language–thought discussion.

This profile connects shared conceptual resources with experience-dependent organization. Grounded-cognition and core-knowledge accounts identify perceptual, bodily, and early representational foundations for understanding the world (Barsalou, 2008; Spelke & Kinzler, 2007). Learning accounts explain how linguistic and experiential information jointly contribute to semantic structure (Andrews et al., 2009; Riordan & Jones, 2011). Our domain contrast fits this combination: common basic meanings can participate in relations whose cross-language correspondence varies with their content. The pitch studies provide a developmental illustration, showing that early sensitivity to multiple mappings can coexist with language-related differences in adults (Dolscheid et al., 2013, 2014). Language-related differences in color discrimination and bilingual event processing likewise show how shared perceptual resources can be deployed differently during particular tasks (Athanasopoulos et al., 2015; Winawer et al., 2007). Together with evidence that language-prediction models recover human-relevant conceptual structure (Xu et al., 2025), these findings place the present comparison within a psychological account of common resources and learned relations. It extends that account by identifying a recurring contrast across a broad language sample. Conceptual commonality can therefore be examined at more than one level: languages may express corresponding basic meanings while differing in the relations connecting them. This distinction sharpens the comparative question raised by linguistic relativity and links it to theories in which conceptual content depends partly on relations within a larger knowledge system (Malt & Majid, 2013; Murphy & Medin, 1985).

Shared Constraints and Local Experience

The domain contrast is consistent with an interaction between relational systematicity and communicative adaptation. Thompson et al. (2020) associated higher alignment with domains possessing strong internal structure, such as number and quantity. Such structure offers one explanation for correspondence among parts of our Socio-Cultural inventory. Efficient-communication accounts provide a complementary explanation: kinship and color systems balance informative distinctions against representational cost, while color naming also reflects communicative usefulness (Gibson et al., 2017; Kemp & Regier, 2012; Zaslavsky et al., 2018). Recurrent needs can favor comparable relations among meanings. Locally varying experience can favor different distinctions within the same general framework. Landscape vocabularies vary across languages, ice and snow terminology tracks environmental communicative needs, and semantic variation along shared neurocognitive dimensions is associated with climate (Burenhult & Levinson, 2008; Fu et al., 2026; Regier et al., 2016). These findings help explain why relations among Nature & Motion meanings might vary even when their referents belong to a commonly experienced physical world. Shared perceptual access and shared semantic organization are distinct explanatory issues: environmental regularities provide material for meaning, while experience and use can affect which distinctions and associations become prominent (Barsalou, 2008; Burenhult & Levinson, 2008). The present contrast thus develops the earlier alignment results into a synthesis in which shared relational constraints and variable experience jointly organize meaning. Greater correspondence in one group and greater variation in another are compatible outcomes of meaning systems adapting to overlapping and differing demands.

The distance associations connect this synthesis to language communities. Socio-Cultural relations were comparatively similar across the sample and also became more dissimilar with geographic separation in both encoders. This combination parallels emotion-semantic research, in which common organizing dimensions

coexist with geographically patterned variation (Jackson et al., 2019), and broader alignment findings linking similarity to cultural and geographic proximity (Thompson et al., 2020). Interaction offers a psychologically plausible pathway: selective coordination of task-relevant language is associated with improved joint performance, and language contact can accompany restructuring of color meanings (Fusaroli et al., 2012; Malik-Moraleda et al., 2023). Broadly recurring relational patterns can therefore coexist with regional conventions. A domain's average correspondence and its geographic gradient describe different properties: relative commonality across the sample can be accompanied by systematic regional differentiation. This combination specifies how shared and locally patterned organization can coexist within the same field of meanings. Greater relational dissimilarity between more distantly related languages across all domains further situates the profile within historical relationships among languages, extending the historical associations reported by Thompson et al. (2020). The joint comparison links the organization of meaning to both the kinds of concepts involved and the spatial and historical contexts in which languages are used.

Universality and Relativity in an Adaptive System

The resulting profile gives semantic content to the human-driven complex-adaptive-system perspective introduced above (Beckner et al., 2009; Liu, 2014, 2018). In this account, cognitive capacities, interaction, and local experience jointly shape linguistic regularities. Cultural-transmission experiments show how learning across generations can produce increasingly structured and learnable systems (Kirby et al., 2008). Our findings suggest a conceptual patchwork, with relations among meanings woven more similarly across languages in some domains than in others. Shared relational structure and recurrent communicative demands may draw parts of the Socio-Cultural inventory into more similar patterns, whereas variation in ecological experience and communicative needs may allow greater differentiation among Nature & Motion meanings (Gibson et al., 2017; Regier et al., 2016; Thompson et al., 2020). Universality and relativity can thus be understood as interwoven properties of an adaptive meaning system. This probabilistic view connects shared cognitive capacities with locally established patterns of use (Evans & Levinson, 2009; Liu, 2018), offering a domain-sensitive account of how commonality and variation coexist. For the Sapir–Whorf question, this shifts attention toward the distribution of shared and differing relations within conceptual systems. A psychological test follows directly: matched judgments should reveal relatively greater cross-language correspondence for the sampled Socio-Cultural relations than for Nature & Motion relations. Comparisons with nonverbal materials and manipulations of language use would connect this prediction to the attention, categorization, and learning processes implicated in linguistic relativity (Wolff & Holmes, 2011).

Several limitations guide that extension. Our evidence comes from model-derived lexical relations; establishing their correspondence to speakers' judgments remains necessary, particularly because naming and nonverbal similarity can dissociate (Malt et al., 1999). Training biases, polysemy, coverage-based sampling, heterogeneous domain assignments, and model-dependent intermediate ranks affect generalization. Communicative, ecological, and historical mechanisms require independent measurements. Within this scope, the study contributes a recurring domain contrast to linguistic relativity and an integrated interpretation linking shared cognitive resources with experience and social use. It makes the relationship between conceptual commonality and linguistic diversity a question about identifiable patterns among meanings, with specific psychological comparisons through which those patterns can be examined.

Author Contributions

Conceptualization and methodology: Weiran Wang, Wenjuan Qin, and Haitao Liu. Data curation and formal analysis: Weiran Wang. Visualization and writing—original draft: Weiran Wang, Wenjuan Qin, and Haitao Liu. Writing—review and editing: Wenjuan Qin and Haitao Liu. Supervision: Wenjuan Qin and Haitao Liu.

Acknowledgments

This paper is a research output of the Digital Humanities and Glottometrics Lab at Fudan University.

References

Andrews, M., Vigliocco, G., & Vinson, D. (2009). Integrating experiential and distributional data to learn semantic representations. *Psychological Review*, *116*(3), 463–498. https://doi.org/10.1037/a0016261

Athanasopoulos, P., Bylund, E., Montero-Melis, G., Damjanovic, L., Schartner, A., Kibbe, A., Riches, N., & Thierry, G. (2015). Two languages, two minds: Flexible cognitive processing driven by language of operation. *Psychological Science*, *26*(4), 518–526. https://doi.org/10.1177/0956797614567509

Barsalou, L. W. (2008). Grounded cognition. *Annual Review of Psychology*, *59*, 617–645. https://doi.org/10.1146/annurev.psych.59.103006.093639

Beckner, C., Blythe, R., Bybee, J., Christiansen, M. H., Croft, W., Ellis, N. C., Holland, J., Ke, J., Larsen-Freeman, D., & Schoenemann, T. (2009). Language is a complex adaptive system: Position paper. *Language Learning*, *59*(s1), 1–26. https://doi.org/10.1111/j.1467-9922.2009.00533.x

Beinborn, L., & Choenni, R. (2020). Semantic drift in multilingual representations. *Computational Linguistics*, *46*(3), 571–603. https://doi.org/10.1162/coli_a_00382

Burenhult, N., & Levinson, S. C. (2008). Language and landscape: A cross-linguistic perspective. *Language Sciences*, *30*(2-3), 135–150. https://doi.org/10.1016/j.langsci.2006.12.028

Carpenter, B., Gelman, A., Hoffman, M. D., Lee, D., Goodrich, B., Betancourt, M., Brubaker, M., Guo, J., Li, P., & Riddell, A. (2017). Stan: A probabilistic programming language. *Journal of Statistical Software*, *76*(1), 1–32. https://doi.org/10.18637/jss.v076.i01

Conneau, A., Khandelwal, K., Goyal, N., Chaudhary, V., Wenzek, G., Guzmán, F., Grave, E., Ott, M., Zettlemoyer, L., & Stoyanov, V. (2020). Unsupervised cross-lingual representation learning at scale. *Proceedings of the 58th Annual Meeting of the Association for Computational Linguistics*, 8440–8451. https://doi.org/10.18653/v1/2020.acl-main.747

Devlin, J., Chang, M.-W., Lee, K., & Toutanova, K. (2019). BERT: Pre-training of deep bidirectional transformers for language understanding. *Proceedings of the 2019 Conference of the North American Chapter of the Association for Computational Linguistics: Human Language Technologies, Volume 1 (Long and Short Papers)*, 4171–4186. https://doi.org/10.18653/v1/N19-1423

Dolscheid, S., Hunnius, S., Casasanto, D., & Majid, A. (2014). Prelinguistic infants are sensitive to space-pitch associations found across cultures. *Psychological Science*, *25*(6), 1256–1261. https://doi.org/10.1177/0956797614528521

Dolscheid, S., Shayan, S., Majid, A., & Casasanto, D. (2013). The thickness of musical pitch: Psychophysical evidence for linguistic relativity. *Psychological Science*, *24*(5), 613–621. https://doi.org/10.1177/0956797612457374

Evans, N., & Levinson, S. C. (2009). The myth of language universals: Language diversity and its importance for cognitive science. *Behavioral and Brain Sciences*, *32*(5), 429–448. https://doi.org/10.1017/S0140525X0999094X

Fu, Z., Chu, Y., Zhang, T., Li, Y., Wang, X., & Bi, Y. (2026). Semantic similarity across languages reflects neurocognitive dimensions shaped by climate. *Nature Communications*, *17*, Article 4016. https://doi.org/10.1038/s41467-026-70608-8

Fusaroli, R., Bahrami, B., Olsen, K., Roepstorff, A., Rees, G., Frith, C. D., & Tylén, K. (2012). Coming to terms: Quantifying the benefits of linguistic coordination. *Psychological Science*, *23*(8), 931–939. https://doi.org/10.1177/0956797612436816

Gibson, E., Futrell, R., Jara-Ettinger, J., Mahowald, K., Bergen, L., Ratnasingam, S., Gibson, M., Piantadosi, S. T., & Conway, B. R. (2017). Color naming across languages reflects color use. *Proceedings of the National Academy of Sciences*, *114*(40), 10785–10790. https://doi.org/10.1073/pnas.1619666114

Jackson, J. C., Watts, J., Henry, T. R., List, J.-M., Forkel, R., Mucha, P. J., Greenhill, S. J., Gray, R. D., & Lindquist, K. A. (2019). Emotion semantics show both cultural variation and universal structure. *Science*, *366*(6472), 1517–1522. https://doi.org/10.1126/science.aaw8160

Jäger, G. (2018). Global-scale phylogenetic linguistic inference from lexical resources. *Scientific Data*, *5*, 180189. https://doi.org/10.1038/sdata.2018.189

Kemp, C., & Regier, T. (2012). Kinship categories across languages reflect general communicative principles. *Science*, *336*(6084), 1049–1054. https://doi.org/10.1126/science.1218811

Kirby, S., Cornish, H., & Smith, K. (2008). Cumulative cultural evolution in the laboratory: An experimental approach to the origins of structure in human language. *Proceedings of the National Academy of Sciences*, *105*(31), 10681–10686. https://doi.org/10.1073/pnas.0707835105

Kriegeskorte, N., Mur, M., & Bandettini, P. (2008). Representational similarity analysis – connecting the branches of systems neuroscience. *Frontiers in Systems Neuroscience*, *2*, 4. https://doi.org/10.3389/neuro.06.004.2008

Landauer, T. K., & Dumais, S. T. (1997). A solution to Plato's problem: The latent semantic analysis theory of acquisition, induction, and representation of knowledge. *Psychological Review*, *104*(2), 211–240. https://doi.org/10.1037/0033-295X.104.2.211

Lenci, A. (2018). Distributional models of word meaning. *Annual Review of Linguistics*, *4*, 151–171. https://doi.org/10.1146/annurev-linguistics-030514-125254

Lewis, M., Zettersten, M., & Lupyan, G. (2019). Distributional semantics as a source of visual knowledge. *Proceedings of the National Academy of Sciences*, *116*(39), 19237–19238. https://doi.org/10.1073/pnas.1910148116

Lewis, M., Cahill, A., Madnani, N., & Evans, J. (2023). Local similarity and global variability characterize the semantic space of human languages. *Proceedings of the National Academy of Sciences*, *120*(51), e2300986120. https://doi.org/10.1073/pnas.2300986120

Liu, H. (2014). Language is more a human-driven system than a semiotic system: Comment on "Modelling language evolution: Examples and predictions" by Tao Gong, Shuai Lan, Menghan Zhang. *Physics of Life Reviews*, *11*(2), 309–310. https://doi.org/10.1016/j.plrev.2013.12.008

Liu, H. (2018). Language as a human-driven complex adaptive system: Comment on "Rethinking foundations of language from a multidisciplinary perspective" by T. Gong et al. *Physics of Life Reviews*, *26–27*, 149–151. https://doi.org/10.1016/j.plrev.2018.06.006

Lucy, J. A. (1992). *Language diversity and thought: A reformulation of the linguistic relativity hypothesis*. Cambridge University Press. https://doi.org/10.1017/CBO9780511620843

Malik-Moraleda, S., Mahowald, K., Conway, B. R., & Gibson, E. (2023). Concepts are restructured during language contact: The birth of blue and other color concepts in Tsimane'-Spanish bilinguals. *Psychological Science*, *34*(12), 1350–1362. https://doi.org/10.1177/09567976231199742

Malt, B. C., Gennari, S., Imai, M., Ameel, E., Tsuda, N., & Majid, A. (2008). Talking about walking: Biomechanics and the language of locomotion. *Psychological Science*, *19*(3), 232–240. https://doi.org/10.1111/j.1467-9280.2008.02074.x

Malt, B. C., & Majid, A. (2013). How thought is mapped into words. *Wiley Interdisciplinary Reviews: Cognitive Science*, *4*(6), 583–597. https://doi.org/10.1002/wcs.1251

Malt, B. C., Sloman, S. A., Gennari, S., Shi, M., & Wang, Y. (1999). Knowing versus naming: Similarity and the linguistic categorization of artifacts. *Journal of Memory and Language*, *40*(2), 230–262. https://doi.org/10.1006/jmla.1998.2593

Murphy, G. L., & Medin, D. L. (1985). The role of theories in conceptual coherence. *Psychological Review*, *92*(3), 289–316. https://doi.org/10.1037/0033-295X.92.3.289

Murphy, G. L. (2002). *The big book of concepts*. MIT Press. https://mitpress.mit.edu/9780262632997/the-big-book-of-concepts/

Regier, T., Carstensen, A., & Kemp, C. (2016). Languages support efficient communication about the environment: Words for snow revisited. *PLoS One*, *11*(4), e0151138. https://doi.org/10.1371/journal.pone.0151138

Riordan, B., & Jones, M. N. (2011). Redundancy in perceptual and linguistic experience: Comparing feature-based and distributional models of semantic representation. *Topics in Cognitive Science*, *3*(2), 303–345. https://doi.org/10.1111/j.1756-8765.2010.01111.x

Spelke, E. S., & Kinzler, K. D. (2007). Core knowledge. *Developmental Science*, *10*(1), 89–96. https://doi.org/10.1111/j.1467-7687.2007.00569.x

Swadesh, M. (1955). Towards greater accuracy in lexicostatistic dating. *International Journal of American Linguistics*, *21*(2), 121–137. https://doi.org/10.1086/464321

Thompson, B., Roberts, S. G., & Lupyan, G. (2020). Cultural influences on word meanings revealed through large-scale semantic alignment. *Nature Human Behaviour*, *4*(10), 1029–1038. https://doi.org/10.1038/s41562-020-0924-8

Vulić, I., Ponti, E. M., Litschko, R., Glavaš, G., & Korhonen, A. (2020). Probing pretrained language models for lexical semantics. *Proceedings of the 2020 Conference on Empirical Methods in Natural Language Processing (EMNLP)*, 7222–7240. https://doi.org/10.18653/v1/2020.emnlp-main.586

Wiktionary contributors. (n.d.). *Swadesh lists by language*. Wiktionary. Retrieved October 22, 2026, from https://en.wiktionary.org/wiki/Category:Swadesh_lists_by_language

Winawer, J., Witthoft, N., Frank, M. C., Wu, L., Wade, A. R., & Boroditsky, L. (2007). Russian blues reveal effects of language on color discrimination. *Proceedings of the National Academy of Sciences*, *104*(19), 7780–7785. https://doi.org/10.1073/pnas.0701644104

Wolff, P., & Holmes, K. J. (2011). Linguistic relativity. *Wiley Interdisciplinary Reviews: Cognitive Science*, *2*(3), 253–265. https://doi.org/10.1002/wcs.104

Xu, N., Zhang, Q., Du, C., Luo, Q., Qiu, X., Huang, X., & Zhang, M. (2025). Revealing emergent human-like conceptual representations from language prediction. *Proceedings of the National Academy of Sciences*, *122*(44), Article e2512514122. https://doi.org/10.1073/pnas.2512514122

Youn, H., Sutton, L., Smith, E., Moore, C., Wilkins, J. F., Maddieson, I., Croft, W., & Bhattacharya, T. (2016). On the universal structure of human lexical semantics. *Proceedings of the National Academy of Sciences*, *113*(7), 1766–1771. https://doi.org/10.1073/pnas.1520752113

Zaslavsky, N., Kemp, C., Regier, T., & Tishby, N. (2018). Efficient compression in color naming and its evolution. *Proceedings of the National Academy of Sciences*, *115*(31), 7937–7942. https://doi.org/10.1073/pnas.1800521115

Supplementary Information

Where Meanings Converge and Diverge: Revisiting Linguistic Relativity Across 81 Languages

Weiran Wang, Wenjuan Qin, and Haitao Liu
Fudan University, Shanghai 200433, China

S1 Sample construction and quality control

The language inventory was constructed from the full candidate coverage of multilingual BERT available at the time of data assembly. The mBERT model card lists 104 pretraining languages. From this candidate pool, we excluded artificial languages, languages whose available Wikipedia material was predominantly machine-generated, and languages without the geographic or language-history metadata needed for the comparative analysis. Language identifiers were standardized against ISO and Glottolog records, yielding an 82-language study input. Lexical forms were downloaded from the language-specific Swadesh lists indexed by Wiktionary (Wiktionary contributors, n.d.-b). The 207 numeric concept IDs follow Wiktionary's English Swadesh ordering (Wiktionary contributors, n.d.-a); the archived English concept labels match that sequence after normalization of formatting and parenthetical qualifiers. Thus, 82 languages and 207 concepts describe the study input; this is the inherited input inventory, not a new sampling design introduced by the quality-control procedure. The lexical source is distinct from the Bowern/Concepticon resources subsequently consulted for auxiliary classification. The retrieval dates in the references document source verification and do not establish the original download date. Analyses retained the archived lexical forms rather than replacing them with later online edits.

The archived vector set contained those 207 concept slots in the 82-language input, and the same inventory was extracted from the local mBERT checkpoint on the remote server. Concept identity was reconstructed from numeric concept IDs, not English glosses, because *you* and *man* each occur at two source-list positions. Every vector was screened for parseability, dimensional consistency, finite values, and all-zero content. The quality-control procedure identified 12 all-zero observations (Table S6). They were treated as invalid rather than transformed by mean centering. Javanese remained degenerate after extraction: its maximum duplicate share at seven-decimal precision was .607, compared with .995 in the archived file. It was excluded under the stated duplicate-share rule. The retained sample therefore contains 81 languages.

For a common support, the primary analysis uses only concepts with valid vectors in all retained languages. Eleven concepts failed this criterion (Table S7), leaving 196 concepts and 3,240 unordered language dyads. Undefined RSA values remain missing and are never replaced by an arbitrary dissimilarity value.

S2 Representational similarity analysis

For concept set C in language L, vectors were centered within language and set,

$$\hat{v}_{c,L} = v_{c,L} - \frac{1}{|C|} \sum_{i \in C} v_{i,L}. \tag{S1}$$

Cosine representational distance matrices (RDMs) were constructed from the centered vectors. We call the cross-lingual outcome RSA dissimilarity:

$$d_{\text{sem}}(L_1, L_2) = 1 - \rho_{\text{Spearman}}\left(u(D^{L_1}), u(D^{L_2})\right), \tag{S2}$$

where $u(\cdot)$ denotes the upper triangle. Because the outcome is $1 - \rho$, it can exceed one when the rank correlation is negative. All analyses align concepts by numeric ID and use this RSA definition exclusively.

S3 Primary and reference-style taxonomies

The primary mapping consolidates the fine-grained topical annotations retained in the project's archived English inventory into four author-defined operational groups. It is the `combined_cluster` four-domain classification stored in the source materials. These project annotations and their four-domain consolidation should not be attributed to Wiktionary or treated as the original Bowern list. Its labels are operational summaries rather than homogeneous natural semantic categories. Biological Core includes body, person, kinship, perception, vital processes, and personal pronouns in the primary mapping. Nature & Motion includes nonhuman living kinds, environments, movement, and physical change. Socio-Cultural contains properties, evaluation, color, time, action and manipulation, material-culture items, and numerals. Grammar & Function primarily contains demonstratives, interrogatives, quantifiers, and relational or function items. Relevant comparative literature motivates study of these types of meanings (Beckner et al., 2009; Burenhult & Levinson, 2008; Bybee, 2010; Christiansen & Chater, 2008; Jackson et al., 2019; Kemp & Regier, 2012; Levinson, 2003; Majid & Burenhult, 2014; Malt et al., 1999; Spelke & Kinzler, 2007), but does not independently validate this particular grouping. The six personal-pronoun items remain Biological Core in the author-defined primary consolidation but move to Grammar & Function under the reference mapping's explicit functional override. This is an intentional classification difference, not a vector-extraction or analysis bug. Kinship items remain Biological Core under both mappings. Table S26 is the authoritative item-level record.

The alternative mapping is a documented reference-style sensitivity taxonomy constructed using the Bowern-2008-207 list in Concepticon (Bowern, 2008; List et al., 2016), official WOLD and IDS item-level fields (Haspelmath & Tadmor, 2009; Key & Comrie, 2015), WordNet 3.0 lexnames where appropriate (Fellbaum, 1998), and explicit functional rules. A study-specific crosswalk aligns the Wiktionary concept slots to Bowern entries through gloss matching and explicit item correspondences; official Concepticon identifiers are then attached through those Bowern entries. This alignment does not make the source inventories identical. Their Concepticon sense labels differ for six aligned items: *child*, *skin*, *smell*, *turn*, *rain*, and *burn*. The frozen Bowern-based crosswalk was retained for this auxiliary sensitivity analysis. The existing `wold_field` is retained as a deterministic WOLD/IDS-informed macro-domain rule; the official WOLD semantic field and IDS chapter are stored separately rather than being relabeled as the same variable. The frozen source snapshots supply Bowern-based Concepticon joins for all 207 aligned project items, with linked coverage of 195 Swadesh entries, 188 WOLD entries, and 172 IDS entries. These counts describe the auxiliary crosswalk's coverage; they do not identify the source of the downloaded lexical forms. Missing source coverage is recorded item by item in reanalysis_20260814/taxonomy_sources/official_resource_crosswalk_207.tsv. The two mappings agree on 185 of 207 items. Tables S8–S26 report the mapping comparison, the reassignments, and each concept-level decision.

S4 Literature motivation for the four domains

Table S1. Literature motivation and support for the four operational domains. Counts refer first to the 207-item input and then to the 196-concept complete-case support.

Domain	Semantic focus	Representative comparative and cognitive literature	Input / complete
Biological Core	Body, person, kinship, perception, vital processes, and primary-mapping personal pronouns	Core knowledge, embodied structure, kinship systems, and recurrent lexical organization (Kemp & Regier, 2012; Pagel, 2009; Spelke & Kinzler, 2007)	68 / 67
Nature & Motion	Nonhuman living kinds, physical environments, movement, and physical change	Frames of reference, motion descriptions, landscape language, and spatial cognition (Burenhult & Levinson, 2008; Levinson, 2003; Levinson & Wilkins, 2006; Talmy, 1985)	53 / 49
Socio-Cultural	Properties, evaluation, colour, time, actions, material culture, and numerals	Artifact naming, cultural categorization, emotion, odor, and communicative coordination (Jackson et al., 2019; Majid & Burenhult, 2014; Malt & Majid, 2013; Malt et al., 1999; Thompson et al., 2020)	60 / 59
Grammar & Function	Demonstratives, interrogatives, relations, quantifiers, and other function items	Grammaticalization, discourse functions, usage-based structure, and language–cognition interaction (Beckner et al., 2009; Bybee, 2010; Christiansen & Chater, 2008; Croft, 2001; Levinson & Wilkins, 2006)	26 / 21

The counts in the final column are descriptive support counts, not evidence for the conceptual validity of the domains. The RSA-blind source/ontology audit found 89.4% exact agreement across the 207 input concepts, Cohen's $\kappa = .856$ (2,000-concept bootstrap 95% interval [.794, .908]), and macro-F1 = .885 ([.835, .927]). The endpoint contrast was then tested under the reference mapping and consensus subset; those outcome-level checks support the broad Nature & Motion versus Socio-Cultural direction but do not establish a unique four-domain taxonomy. The literature supplies the motivation for the grouping; the complete-case rule then determines which items enter the comparable RSA support.

S5 Geographic distance and genealogical relatedness

The geographic and genealogical-relatedness matrices were frozen as quality-controlled input files for the analysis. Geographic values are great-circle distances between archived language centroids. We operationalized genealogical relatedness with an ASJP lexical–phonological distance derived from core-vocabulary PMI scores (Church & Hanks, 1990; Jäger, 2018). Throughout this article, genealogical distance refers to this ASJP-based operational measure. It serves as an approximate relatedness measure rather than a formal phylogenetic-tree distance. Because ASJP and the semantic analysis both derive from core-vocabulary traditions, they share partial material overlap. The input hashes in Table S25 verify the exact matrices used by the remote runs.

S6 Comparative models

The retained multiple-membership model is

$$d_{\text{sem},ij} = \alpha_k + \beta_{\text{geo},k} z_{\text{geo},ij} + \beta_{\text{gen},k} z_{\text{gen},ij} + \tfrac{1}{2}(u_i + u_j) + \epsilon_{ij}. \tag{S3}$$

Equal language-membership weights sum to one. The Gaussian language effects were represented with a non-centered parameterization and an explicit sum-to-zero constraint, which identifies the intercept separately from the average repeated language contribution. The centered intercept prior was $Normal(.9, .2)$, coefficient priors were $Normal(0, .2)$, the language-effect standard deviation had a positive Student-t distribution with degrees of freedom 3, location 0, and scale .5 prior, and residual σ had an $Exponential(1)$ prior. Four independent chains used 3,000 warm-up and 3,000 retained iterations with seed base 42, `adapt_delta` = .99, and `max_treedepth` = 12. Sampling was performed on the remote server with the complete 3,240-dyad support; no data subsampling was used. Since the four domain outcomes for one dyad are correlated, the model intervals are conditional on this residual specification; the dyad-cluster bootstrap below preserves within-dyad domain dependence but does not eliminate dependence between different dyads sharing a language.

Convergence is evaluated on substantive model parameters after excluding Stan sampler-diagnostic columns such as `treedepth__`; the full and filtered values are retained in the machine-readable diagnostics. The model-specific R-hat, effective sample size, divergence, treedepth, E-BFMI, and posterior-predictive results are reported in Table S17. Tables S12–S16 provide complete summaries.

The pooled model concatenates the mBERT and XLM-R-large domain rows and adds a model indicator, Model × Domain, Model × Geography, Model × ASJP, and corresponding three-way interactions. It retains the same repeated-language structure and priors. Its model-interaction posteriors, rather than a visual comparison of separate credible intervals, are the formal test of cross-model differences; the complete coefficient table is archived with the machine-readable output.

The Bayesian specification uses four chains with 3,000 warm-up and 3,000 retained iterations per chain, `adapt_delta` = .99, seed base 42, and `max_treedepth` = 12. All model-by-model and pooled manuscript fits use the complete data support and these settings; no reduced-iteration fit enters a reported estimate. The settings and acceptance diagnostics are recorded in the provenance table and machine-readable manifests.

Table S2. Formal pooled cross-model posterior contrasts. The mBERT Biological Core intercept is the reference. Intervals are equal-tail 95% credible intervals from the full-support pooled model.

Term	Mean	95% CrI	Interpretation
Primary; Model (XLM-R-large)	0.0311	[0.0277, 0.0345]	Model main effect relative to mBERT Biological Core.
Primary; Model × Nature & Motion	0.0211	[0.0162, 0.0260]	Model-by-domain increment.
Primary; Model × Socio-Cultural	0.0167	[0.0119, 0.0213]	Model-by-domain increment.
Primary; Model × Grammar & Function	0.0435	[0.0387, 0.0483]	Model-by-domain increment.
Primary; Model × Geography	0.0060	[0.0023, 0.0096]	Difference in baseline geography slopes.
Primary; Model × ASJP	−0.0052	[−0.0088, −0.0015]	Difference in baseline ASJP slopes.
Primary; Model × Nature & Motion × Geography	−0.0051	[−0.0104, 0.0000]	Domain-specific geography-slope difference.
Primary; Model × Socio-Cultural × Geography	−0.0107	[−0.0158, −0.0055]	Domain-specific geography-slope difference.
Primary; Model × Grammar & Function × Geography	−0.0100	[−0.0152, −0.0048]	Domain-specific geography-slope difference.
Primary; Model × Nature & Motion × ASJP	−0.0030	[−0.0082, 0.0021]	Domain-specific ASJP-slope difference.
Primary; Model × Socio-Cultural × ASJP	−0.0036	[−0.0087, 0.0015]	Domain-specific ASJP-slope difference.
Primary; Model × Grammar & Function × ASJP	−0.0085	[−0.0136, −0.0033]	Domain-specific ASJP-slope difference.
Primary; Derived endpoint contrast	0.0044	[−0.0003, 0.0090]	XLM-R-large minus mBERT endpoint contrast.
External; Model (XLM-R-large)	0.0349	[0.0319, 0.0379]	Model main effect relative to mBERT Biological Core.
External; Model × Nature & Motion	0.0175	[0.0132, 0.0218]	Model-by-domain increment.
External; Model × Socio-Cultural	0.0204	[0.0161, 0.0246]	Model-by-domain increment.
External; Model × Grammar & Function	0.0364	[0.0321, 0.0407]	Model-by-domain increment.
External; Model × Geography	0.0017	[−0.0015, 0.0049]	Difference in baseline geography slopes.
External; Model × ASJP	−0.0075	[−0.0107, −0.0043]	Difference in baseline ASJP slopes.
External; Model × Nature & Motion × Geography	−0.0009	[−0.0054, 0.0037]	Domain-specific geography-slope difference.
External; Model × Socio-Cultural × Geography	−0.0106	[−0.0152, −0.0061]	Domain-specific geography-slope difference.
External; Model × Grammar & Function × Geography	−0.0031	[−0.0077, 0.0015]	Domain-specific geography-slope difference.
External; Model × Nature & Motion × ASJP	−0.0075	[−0.0108, −0.0042]	Domain-specific ASJP-slope difference.
External; Model × Socio-Cultural × ASJP	−0.0111	[−0.0144, −0.0079]	Domain-specific ASJP-slope difference.
External; Model × Grammar & Function × ASJP	−0.0088	[−0.0120, −0.0056]	Domain-specific ASJP-slope difference.
External; Derived endpoint contrast	−0.0029	[−0.0071, 0.0014]	XLM-R-large minus mBERT endpoint contrast.

The prior-scale sensitivity used the same complete support and full four-chain specification with coefficient prior scale 0.5 instead of 0.2. The derived endpoint contrast was 0.0044 [−0.0005, 0.0091] under the primary taxonomy and −0.0029 [−0.0071, 0.0014] under

the reference-style taxonomy. These estimates are numerically indistinguishable from the default-scale results for the inferential claim: the cross-model endpoint difference is not taxonomy-invariant.

Paired language-dyad sensitivity. The same unordered language dyad contributes observations to both encoders, so we also resampled complete dyads as blocks, retaining all four domains and both models together. This sensitivity analysis preserves the paired encoder and domain observations within each dyad, but does not eliminate dependence between different dyads sharing a language. It does not replace the full-support pooled posterior. The XLM-R-large minus mBERT difference in the Nature & Motion minus Socio-Cultural endpoint contrast was 0.0044 (95% paired-bootstrap interval $[0.0012, 0.0074]$) under the primary taxonomy, but -0.0029 (95% interval $[-0.0062, 0.0002]$) under the reference-style taxonomy. Thus, within-model endpoint contrasts can be stable while the cross-model difference of those contrasts remains taxonomy-sensitive. The complete four-domain paired summaries and manifests are archived in `reanalysis_20260814/paired_cross_model_bootstrap_20260815/` and `reanalysis_20260814/paired_cross_model_bootstrap_external_20260815/`; each uses all 3,240 dyads and 2,000 replicates.

Table S3. Paired language-dyad bootstrap for the cross-model endpoint contrast. The estimand is (XLM-R-large − mBERT) applied to the Nature & Motion minus Socio-Cultural contrast. Both models and all four domains are resampled together within each of 2,000 complete-dyad replicates.

Taxonomy	Full estimate	2.5%	97.5%
Primary	0.0044	0.0012	0.0074
Reference-style taxonomy	−0.0029	−0.0062	0.0002

S7 Cross-model and representation sensitivity

Within mBERT, both full observed inventories yield the ordering Socio-Cultural < Grammar & Function < Biological Core < Nature & Motion. A 2,000-resample dyad-cluster bootstrap preserves that ordering in 100.0% of primary-taxonomy resamples and 99.75% of reference-style taxonomy resamples (2,000 and 1,995 resamples, respectively). The positive Socio-Cultural-by-geography interaction excludes zero under both mappings. The mBERT Grammar-by-ASJP interaction is .00346 [.000015, .00698] under the primary mapping and .00277 [-.00036, .00591] under the alternative mapping. XLM-R-large does not reproduce a consistently positive grammatical increment. Accordingly, the evidence supports a broad positive ASJP association, not a uniquely grammatical-functional association.

XLM-R-large extraction used the same lexical items and processing decisions (Table S21). It preserves the observed-inventory Socio-Cultural endpoint as lowest and Nature & Motion as highest under both taxonomies, but Biological Core and Grammar & Function exchange rank relative to mBERT (Tables S22–S24). The full four-domain ordering is therefore checkpoint-dependent. Dyad-level agreement is moderate overall and strongest for Socio-Cultural concepts (Table S23). All leave-one-language-out runs preserve each model's observed-inventory ordering, so the middle-domain swap is not driven by a single language.

As a separate coverage check, we evaluated the XLM-R RSA using the screened language-QC file as the inclusion rule rather than the 82-name source inventory. The run retained 81 languages, 196 complete-case concepts, all 3,240 dyads, and 32,400 rows across two taxonomies and five outcomes (global plus four domains), with no sampling. Table S4 exactly reproduces the fixed-sample XLM-R output. This documents coverage and provenance rather than providing an independent-data replication; the same support, vectors, and matrices are used.

Table S4. Full-valid XLM-R coverage check. Entries are raw RSA dissimilarity means for the same 81-language, 196-concept, 3,240-dyad support; no Bayesian intervals are assigned to this coverage description.

Taxonomy	All	Biological	Nature	Socio-Cultural	Grammar
Primary	.9321	.9315	.9767	.9220	.9671
External	.9321	.9529	.9729	.9383	.9632

The layer-wise RSA analysis was computed on the same 81-language, 196-concept complete-case support for all 13 archived mBERT layers (Table S18). It is reported as a representation-level sensitivity analysis, not as a second confirmatory model. The frequency-adjusted item analysis was likewise rebuilt from the extracted vectors and numeric concept IDs. Its English Zipf values come from the archived SUBTLEX/`wordfreq`-based table and are used as a broad item-level control rather than as language-specific corpus frequencies (Table S19). The mBERT model uses all 635,040 concept–dyad observations (196 concepts × 3,240 language dyads), analytically marginalized multiple-membership language and concept random intercepts, and exact Gaussian restricted maximum likelihood. Optimization converged with a full-rank 13-column fixed-effect design; no dyad sampling was used.

S8 Concept-label permutation calibration

For each encoder, taxonomy, and domain, we also permuted concept labels within the language-level vector arrays 2,000 times while retaining the complete 81-language support. The observed mean RSA dissimilarity was lower than every permutation replicate for all 16 encoder–taxonomy–domain combinations. With the +1 Monte Carlo correction, the correctly directed lower-tail probability was $p_{\text{lower}} = .0005$ and the two-sided probability was $p_{\text{two-sided}} = .0010$; the upper-tail probability was 1.0. This calibration tests whether the

observed concept correspondence is compatible with random label alignment. It does not test the causal interpretation of the geographic or ASJP associations, nor does it make the dyad-level RSA entries independent.

S9 Consensus-subset sensitivity

The source and ontology assignments agree for 185 of the 207 concepts. The consensus RSA uses the same 81-language support and vectors. Because the extraction has 11 invalid concept IDs, the complete-case RSA support contains 175 concepts (53 Biological Core, 47 Nature & Motion, 55 Socio-Cultural, and 20 Grammar & Function). The two-domain contrast remains (without implying that Socio-Cultural remains the minimum in both encoders): Socio-Cultural is lower than Nature & Motion for both mBERT (.888 versus .935) and XLM-R-large (.938 versus .977). The complete middle-domain ordering is not invariant: mBERT gives Grammar & Function < Socio-Cultural < Biological Core < Nature & Motion, whereas XLM-R-large gives Socio-Cultural < Biological Core < Grammar & Function < Nature & Motion. Thus the consensus subset supports the Socio-Cultural–Nature & Motion contrast but qualifies any claim about the full four-domain hierarchy. The machine-readable consensus RSA outputs and concept-coverage records are included in `reanalysis_20260814/`.

S10 Dependence and domain-size sensitivity

The following archived mBERT sensitivity checks use 81 languages and 196 concepts. Their manifest identifies the earlier archived vector directory, rather than the fresh vectors used by the final Bayesian analysis. They are retained for provenance transparency and are not presented as matched fresh-vector, dual-encoder validation. We resampled the 3,240 unordered dyads with replacement 2,000 times, retaining all four domain outcomes within each sampled dyad. This cluster bootstrap produces percentile intervals for domain means, pairwise differences, and rank stability. We then repeatedly sampled without replacement from every domain to equalize its concept count: 21 concepts per domain under the primary mapping and 30 under the reference-style mapping. We recomputed RSA for 500 samples per taxonomy. In the equal-size analysis, the Socio-Cultural–Nature & Motion ordering was retained in 89.6% of primary and 95.8% of reference-style taxonomy samples, whereas the full four-domain ordering occurred in only 33.6% and 35.4%, respectively. These results qualify the interpretation of the observed-inventory ranking: the endpoint contrast is more stable than the complete ordering. Machine-readable outputs and the exact scripts are in `reanalysis_20260814/`; the primary analyses use the complete support, and the resampling procedures are explicitly labeled sensitivity analyses.

S11 Detailed robustness results

This section provides the complete numerical results for the robustness checks summarized in the main text Results section. All checks address the stability of the domain profile and the cross-model comparison under different analytical choices.

Reference-style classification

The reference-style classification reassigns 22 of the 207 input concepts. Under this alternative mapping, the adjusted Socio-Cultural versus Nature & Motion means were .883 versus .921 for mBERT and .938 versus .973 for XLM-R-large. The within-model Nature & Motion minus Socio-Cultural contrasts were .0375 (95% CrI [.0345, .0405]) for mBERT and .0346 [.0316, .0376] for XLM-R-large. The cross-model difference in these contrasts was −.0029 ([-.0071, .0014]). The endpoint contrast is therefore preserved, but the reassignment changes the separation among domains and the order of the middle positions. Complete domain means and credible intervals under the reference-style classification are reported in SI Table S14.

Leave-one-language-out summaries

Across all 81 leave-one-language-out summaries, each multilingual language model retained its full-inventory domain ordering under both the primary and reference-style classifications. The minimum and maximum means obtained by omitting each language in turn are displayed in the main text as error bars in the cross-model domain profile figure. No single language drove the observed Socio-Cultural–Nature & Motion contrast. The complete leave-one-out range summaries are provided in SI Table S22.

Paired dyad bootstrap

The paired language-dyad bootstrap resampled complete dyads with replacement, retaining both encoders and all four domains. This preserves within-dyad pairing while leaving dependence among dyads sharing a language. Its cross-model Nature & Motion–Socio-Cultural difference was .0044 (95% interval [.0012, .0074]) under the primary taxonomy and −.0029 ([-.0062, .0002]) under the reference-style taxonomy. The full-support pooled posterior intervals include zero under both taxonomies (.0044 [-.0003, .0090] and −.0029 [-.0071, .0014], respectively). Changing the coefficient-prior scale from .2 to .5 did not alter those pooled conclusions. The bootstrap and pooled posterior describe different uncertainty specifications; the bootstrap cannot establish an invariant cross-model difference. Paired-bootstrap archive locations are specified in Section S6.

Frequency-adjusted secondary models

The frequency-adjusted analyses use all 635,040 concept–dyad observations for each encoder. They use item-level cosine dissimilarity, rather than domain-level RSA, and an English Zipf covariate, rather than language-specific corpus frequencies. The frequency coefficient was negative in both fits (mBERT −.02620, 95% Wald interval [-.03228, -.02012]; XLM-R-large −.01414, [-.02424, -.00404]). Several domain main-effect intervals include zero. Because the outcome and estimands differ from the primary RSA analysis, these models do not show that the RSA domain contrast is unchanged after frequency adjustment or independent of word frequency. Specifications and coefficients are in Table S19.

Layer diagnostics and equal-size subsampling

Archived equal-size subsampling results are reported in Section S10. Table S18 reports Grammar & Function and Socio-Cultural outcomes across archived mBERT layers. These diagnostics describe layer dependence in those two domains on the archived-vector support. The table does not report Nature & Motion and therefore does not establish that the Socio-Cultural–Nature & Motion contrast persists through layers 6–12 or is strongest in the final layer. As specified in Section S10, the archived checks are not matched fresh-vector, dual-encoder validation.

Concept-level profiles and triplet diagnostics

Concept-level profiles and selected language triplets serve as descriptive illustrations rather than mechanism tests. Triplet diagnostics identified language triples that passed geographic and operational genealogical distance filters while exhibiting the stated directional domain contrast. The complete diagnostic inventory is archived in `reanalysis_20260814/secondary_fulltriplets_20260815/`. These diagnostics illustrate particular cases but do not constitute independent statistical tests of the domain profile.

Summary

The robustness checks collectively support the Socio-Cultural–Nature & Motion endpoint contrast and the positive ASJP association across both models. The complete four-domain ordering and some geographic associations, however, remain sensitive to domain assignment, the multilingual language model, and the specific concepts included in each domain. The endpoint contrast is comparatively stable, whereas the exact ordering of all four domains is more conditional on analytical choices.

S12 Triplet diagnostics and complete candidate inventory

The main-text triplet figure shows two selected mBERT configurations under two stated filters: genealogically-close/geographically-far versus geographically-close/genealogically-far, with the corresponding directional domain contrast. To provide the complete diagnostic record, the secondary-analysis script enumerated every unique language triplet passing those filters and the directional contrast, without a top-case cap. The machine-readable inventory is archived under reanalysis_20260814/secondary_fulltriplets_20260815/: `triplet_all_filter_candidates.csv` records every distance-filtered candidate, `triplet_all_qualifying.csv` records every candidate with the stated directional contrast, and `triplet_all_qualifying_word_details.csv` gives the concept-level support gaps for every qualifying triplet. The archived top-five files are retained as secondary-analysis inputs.

Table S5. Dependence- and domain-size-sensitivity results. Percentages are proportions of resamples or equal-size concept subsamples. The dyad bootstrap retains all four domain outcomes within each sampled dyad.

Analysis	Taxonomy	Full (%)	S<N (%)	S min. (%)	N max. (%)
Dyad bootstrap	Primary	100.0	100.0	100.0	100.0
Dyad bootstrap	Reference-style	99.75	100.0	100.0	99.75
Equal-size concepts	Primary	33.6	89.6	65.4	70.8
Equal-size concepts	Reference-style	35.4	95.8	69.4	54.4

S13 Reproducibility materials

The reproducibility materials include the concept mapping, observation- and language-level quality-control records, the complete-case concept list, archived mBERT vectors, mBERT extraction metadata and quality-control records, RSA summaries, sensitivity analyses, XLM-R-large extraction metadata and RSA summaries, fixed-effect summaries, diagnostics, posterior-predictive summaries, source code, seeds, and hashes. Checkpoint weights are not included in the manuscript package. Available tokenizer hashes, software versions, extraction rules, and extraction commands are recorded. The XLM-R-large extraction manifest marks its weight file as not hashed; complete weight-hash coverage is therefore not claimed.

S14 Supplementary Tables

Table S6. Invalid concept–language observations identified before RSA. All 12 were all-zero vectors in the extraction and were treated as missing.

ID	Concept	Language	Status	Domain
201	at	Japanese	all-zero-vector	Grammar & Function
202	in	Japanese	all-zero-vector	Grammar & Function
203	with	Japanese	all-zero-vector	Grammar & Function
204	and	Japanese	all-zero-vector	Grammar & Function
206	because	Japanese	all-zero-vector	Grammar & Function
58	bark	Marathi	all-zero-vector	Nature & Motion
58	bark	Nepali	all-zero-vector	Nature & Motion
125	stand	Occitan	all-zero-vector	Nature & Motion
192	dull	Occitan	all-zero-vector	Socio-Cultural
148	moon	Thai	all-zero-vector	Nature & Motion
149	star	Thai	all-zero-vector	Nature & Motion
2	you	Vietnamese	all-zero-vector	Biological Core

Table S7. Concepts excluded by the constant-coverage rule. A concept was retained only when valid in all 81 nondegenerate languages.

ID	Concept	Primary domain	Valid	Invalid observations
2	you	Biological Core	80	Vietnamese:all-zero-vector
58	bark	Nature & Motion	79	Marathi:all-zero-vector; Nepali:all-zero-vector
125	stand	Nature & Motion	80	Occitan:all-zero-vector
148	moon	Nature & Motion	80	Thai:all-zero-vector
149	star	Nature & Motion	80	Thai:all-zero-vector
192	dull	Socio-Cultural	80	Occitan:all-zero-vector
201	at	Grammar & Function	80	Japanese:all-zero-vector
202	in	Grammar & Function	80	Japanese:all-zero-vector
203	with	Grammar & Function	80	Japanese:all-zero-vector
204	and	Grammar & Function	80	Japanese:all-zero-vector
206	because	Grammar & Function	80	Japanese:all-zero-vector

Table S8. Agreement between the primary and reference-style concept taxonomies. Bootstrap intervals use 2,000 concept-level resamples.

Statistic	Estimate	2.5%	97.5%	*B*
agreement	0.894	0.850	0.932	2000
cohen_kappa	0.856	0.794	0.908	2000
macro_f1	0.885	0.835	0.927	2000

Table S9. Concept counts by taxonomy and domain before the complete-case exclusion of 11 concepts.

Domain	Primary	External
Biological Core	68	53
Nature & Motion	53	54
Socio-Cultural	60	64
Grammar & Function	26	36

Table S10. The 22 concepts assigned to different domains by the reference-style mapping.

ID	Concept	Primary	External	Basis	Confidence
1	I	Biological Core	Grammar & Function	functional_override	high
2	you	Biological Core	Grammar & Function	functional_override	high
3	he,she,it	Biological Core	Grammar & Function	functional_override	medium
4	we	Biological Core	Grammar & Function	functional_override	high

Continued on next page

ID	Concept	Primary	External	Basis	Confidence
5	you	Biological Core	Grammar & Function	functional_override	medium
6	they	Biological Core	Grammar & Function	functional_override	high
22	one	Socio-Cultural	Grammar & Function	functional_override	high
23	two	Socio-Cultural	Grammar & Function	functional_override	high
24	three	Socio-Cultural	Grammar & Function	functional_override	high
25	four	Socio-Cultural	Grammar & Function	functional_override	high
103	know	Biological Core	Socio-Cultural	WordNet/Bowern source alignment	medium
104	think	Biological Core	Socio-Cultural	WordNet/Bowern source alignment	medium
106	fear	Biological Core	Socio-Cultural	WordNet/Bowern source alignment	medium
112	hunt	Nature & Motion	Socio-Cultural	WordNet/Bowern source alignment	medium
139	count	Biological Core	Grammar & Function	functional_override	medium
140	say	Biological Core	Socio-Cultural	WordNet/Bowern source alignment	medium
141	sing	Biological Core	Socio-Cultural	WordNet/Bowern source alignment	medium
142	play	Nature & Motion	Socio-Cultural	WordNet/Bowern source alignment	medium
145	freeze	Biological Core	Nature & Motion	WordNet/Bowern source alignment	high
146	swell	Biological Core	Nature & Motion	WordNet/Bowern source alignment	high
169	burn	Biological Core	Nature & Motion	WordNet/Bowern source alignment	medium
207	name	Grammar & Function	Socio-Cultural	WordNet/Bowern source alignment	medium

Table S11. Dyad-level agreement of RSA dissimilarities between taxonomies.

Domain	Dyads	Spearman ρ	Mean absolute difference
Biological Core	3240	0.928	0.024
Nature & Motion	3240	0.956	0.014
Socio-Cultural	3240	0.934	0.025
Grammar & Function	3240	0.764	0.051

Table S12. Global RSA multiple-membership models. Global intercepts summarize all concepts and are not Biological Core domain intercepts.

Encoder	Mean	95% CrI
mbert	0.90973	[0.90867, 0.91079]
xlmr	0.93211	[0.93085, 0.93338]

Table S13. Complete fixed effects for both encoders. Domain indices 1, 2, and 3 denote Nature & Motion, Socio-Cultural, and Grammar & Function; Biological Core is the reference.

Classification	Encoder	Parameter	Estimate [95% CrI]
Primary	mBERT	alpha	0.90043 [0.89813, 0.90274]
Primary	mBERT	beta_geo	-0.00155 [-0.00426, 0.00113]
Primary	mBERT	beta_asjp	0.02220 [0.01963, 0.02478]
Primary	mBERT	beta_domain[1]	0.02405 [0.02076, 0.02728]
Primary	mBERT	beta_domain[2]	-0.02623 [-0.02951, -0.02292]
Primary	mBERT	beta_domain[3]	-0.00798 [-0.01118, -0.00476]
Primary	mBERT	beta_geo_domain[1]	-0.00305 [-0.00656, 0.00044]
Primary	mBERT	beta_geo_domain[2]	0.01293 [0.00940, 0.01643]
Primary	mBERT	beta_geo_domain[3]	0.00206 [-0.00143, 0.00556]
Primary	mBERT	beta_asjp_domain[1]	-0.00016 [-0.00366, 0.00335]
Primary	mBERT	beta_asjp_domain[2]	0.00122 [-0.00232, 0.00475]
Primary	mBERT	beta_asjp_domain[3]	0.00346 [0.00001, 0.00698]
Primary	XLM-R-large	alpha	0.93150 [0.92926, 0.93373]
Primary	XLM-R-large	beta_geo	0.00583 [0.00319, 0.00848]
Primary	XLM-R-large	beta_asjp	0.02107 [0.01855, 0.02357]
Primary	XLM-R-large	beta_domain[1]	0.04520 [0.04198, 0.04838]
Primary	XLM-R-large	beta_domain[2]	-0.00952 [-0.01274, -0.00627]
Primary	XLM-R-large	beta_domain[3]	0.03560 [0.03248, 0.03881]
Primary	XLM-R-large	beta_geo_domain[1]	-0.00820 [-0.01170, -0.00479]
Primary	XLM-R-large	beta_geo_domain[2]	0.00226 [-0.00113, 0.00562]
Primary	XLM-R-large	beta_geo_domain[3]	-0.00803 [-0.01148, -0.00457]

Classification	Encoder	Parameter	Estimate [95% CrI]
Primary	XLM-R-large	beta_asjp_domain[1]	-0.00319 [-0.00666, 0.00020]
Primary	XLM-R-large	beta_asjp_domain[2]	-0.00242 [-0.00586, 0.00103]
Primary	XLM-R-large	beta_asjp_domain[3]	-0.00506 [-0.00855, -0.00164]
Alternative	mBERT	alpha	0.91802 [0.91590, 0.92010]
Alternative	mBERT	beta_geo	-0.00365 [-0.00604, -0.00117]
Alternative	mBERT	beta_asjp	0.02264 [0.02033, 0.02493]
Alternative	mBERT	beta_domain[1]	0.00249 [-0.00039, 0.00538]
Alternative	mBERT	beta_domain[2]	-0.03500 [-0.03797, -0.03205]
Alternative	mBERT	beta_domain[3]	-0.02605 [-0.02895, -0.02314]
Alternative	mBERT	beta_geo_domain[1]	-0.00083 [-0.00392, 0.00227]
Alternative	mBERT	beta_geo_domain[2]	0.01844 [0.01532, 0.02159]
Alternative	mBERT	beta_geo_domain[3]	0.00283 [-0.00033, 0.00595]
Alternative	mBERT	beta_asjp_domain[1]	-0.00015 [-0.00326, 0.00298]
Alternative	mBERT	beta_asjp_domain[2]	0.00189 [-0.00125, 0.00502]
Alternative	mBERT	beta_asjp_domain[3]	0.00277 [-0.00036, 0.00591]
Alternative	XLM-R-large	alpha	0.95292 [0.95093, 0.95485]
Alternative	XLM-R-large	beta_geo	0.00085 [-0.00142, 0.00312]
Alternative	XLM-R-large	beta_asjp	0.01932 [0.01715, 0.02149]
Alternative	XLM-R-large	beta_domain[1]	0.01998 [0.01721, 0.02273]
Alternative	XLM-R-large	beta_domain[2]	-0.01464 [-0.01739, -0.01188]
Alternative	XLM-R-large	beta_domain[3]	0.01031 [0.00757, 0.01313]
Alternative	XLM-R-large	beta_geo_domain[1]	-0.00166 [-0.00464, 0.00128]
Alternative	XLM-R-large	beta_geo_domain[2]	0.00781 [0.00486, 0.01079]
Alternative	XLM-R-large	beta_geo_domain[3]	-0.00031 [-0.00329, 0.00267]
Alternative	XLM-R-large	beta_asjp_domain[1]	-0.00013 [-0.00310, 0.00280]
Alternative	XLM-R-large	beta_asjp_domain[2]	-0.00178 [-0.00473, 0.00118]
Alternative	XLM-R-large	beta_asjp_domain[3]	0.00141 [-0.00154, 0.00441]

Table S14. Adjusted mean: both encoders and both classifications. Posterior means and equal-tail 95% credible intervals reconstructed draw by draw from the frozen full-support fits.

Classification	Encoder	Domain	Estimate [95% CrI]
Primary	mBERT	Biological Core	0.90043 [0.89813, 0.90274]
Primary	mBERT	Nature & Motion	0.92448 [0.92215, 0.92683]
Primary	mBERT	Socio-Cultural	0.87420 [0.87186, 0.87655]
Primary	mBERT	Grammar & Function	0.89245 [0.89015, 0.89473]
Primary	XLM-R-large	Biological Core	0.93150 [0.92926, 0.93373]
Primary	XLM-R-large	Nature & Motion	0.97669 [0.97447, 0.97895]
Primary	XLM-R-large	Socio-Cultural	0.92197 [0.91970, 0.92427]
Primary	XLM-R-large	Grammar & Function	0.96710 [0.96487, 0.96934]
Alternative	mBERT	Biological Core	0.91802 [0.91590, 0.92010]
Alternative	mBERT	Nature & Motion	0.92050 [0.91844, 0.92256]
Alternative	mBERT	Socio-Cultural	0.88301 [0.88093, 0.88506]
Alternative	mBERT	Grammar & Function	0.89197 [0.88993, 0.89399]
Alternative	XLM-R-large	Biological Core	0.95292 [0.95093, 0.95485]
Alternative	XLM-R-large	Nature & Motion	0.97290 [0.97093, 0.97487]
Alternative	XLM-R-large	Socio-Cultural	0.93828 [0.93630, 0.94021]
Alternative	XLM-R-large	Grammar & Function	0.96323 [0.96128, 0.96516]

Table S15. Geography: both encoders and both classifications. Posterior means and equal-tail 95% credible intervals reconstructed draw by draw from the frozen full-support fits.

Classification	Encoder	Domain	Estimate [95% CrI]
Primary	mBERT	Biological Core	-0.00155 [-0.00426, 0.00113]
Primary	mBERT	Nature & Motion	-0.00459 [-0.00723, -0.00187]
Primary	mBERT	Socio-Cultural	0.01138 [0.00865, 0.01409]
Primary	mBERT	Grammar & Function	0.00051 [-0.00225, 0.00321]
Primary	XLM-R-large	Biological Core	0.00583 [0.00319, 0.00848]
Primary	XLM-R-large	Nature & Motion	-0.00236 [-0.00505, 0.00031]
Primary	XLM-R-large	Socio-Cultural	0.00810 [0.00546, 0.01072]

Classification	Encoder	Domain	Estimate [95% CrI]
Primary	XLM-R-large	Grammar & Function	-0.00219 [-0.00485, 0.00049]
Alternative	mBERT	Biological Core	-0.00365 [-0.00604, -0.00117]
Alternative	mBERT	Nature & Motion	-0.00448 [-0.00692, -0.00211]
Alternative	mBERT	Socio-Cultural	0.01479 [0.01236, 0.01722]
Alternative	mBERT	Grammar & Function	-0.00082 [-0.00322, 0.00158]
Alternative	XLM-R-large	Biological Core	0.00085 [-0.00142, 0.00312]
Alternative	XLM-R-large	Nature & Motion	-0.00082 [-0.00307, 0.00140]
Alternative	XLM-R-large	Socio-Cultural	0.00865 [0.00634, 0.01092]
Alternative	XLM-R-large	Grammar & Function	0.00053 [-0.00174, 0.00280]

Table S16. Genealogical distance: both encoders and both classifications. Posterior means and equal-tail 95% credible intervals reconstructed draw by draw from the frozen full-support fits.

Classification	Encoder	Domain	Estimate [95% CrI]
Primary	mBERT	Biological Core	0.02220 [0.01963, 0.02478]
Primary	mBERT	Nature & Motion	0.02204 [0.01947, 0.02463]
Primary	mBERT	Socio-Cultural	0.02342 [0.02085, 0.02603]
Primary	mBERT	Grammar & Function	0.02566 [0.02315, 0.02820]
Primary	XLM-R-large	Biological Core	0.02107 [0.01855, 0.02357]
Primary	XLM-R-large	Nature & Motion	0.01788 [0.01540, 0.02038]
Primary	XLM-R-large	Socio-Cultural	0.01865 [0.01616, 0.02120]
Primary	XLM-R-large	Grammar & Function	0.01601 [0.01339, 0.01849]
Alternative	mBERT	Biological Core	0.02264 [0.02033, 0.02493]
Alternative	mBERT	Nature & Motion	0.02250 [0.02020, 0.02476]
Alternative	mBERT	Socio-Cultural	0.02453 [0.02220, 0.02683]
Alternative	mBERT	Grammar & Function	0.02541 [0.02310, 0.02769]
Alternative	XLM-R-large	Biological Core	0.01932 [0.01715, 0.02149]
Alternative	XLM-R-large	Nature & Motion	0.01919 [0.01702, 0.02134]
Alternative	XLM-R-large	Socio-Cultural	0.01755 [0.01540, 0.01974]
Alternative	XLM-R-large	Grammar & Function	0.02074 [0.01858, 0.02292]

Table S17. Sampling diagnostics for the non-centered, sum-to-zero multiple-membership models. R-hat and ESS summarize substantive model parameters; sampler-diagnostic columns are retained separately in the machine-readable output. The pooled rows use the complete 25,920-row support.

Model	N	Max $\widehat{R}$	Min ESS	Div.	Depth	Min E-BFMI
mBERT global (primary)	3240	1.006	434	0	0	0.642
XLM-R global (primary)	3240	1.003	644	0	0	0.633
mBERT domain (primary)	12960	1.006	533	0	0	0.704
XLM-R domain (primary)	12960	1.005	951	0	0	0.648
mBERT global (external)	3240	1.006	434	0	0	0.642
XLM-R global (external)	3240	1.003	644	0	0	0.633
mBERT domain (external)	12960	1.008	394	0	0	0.689
XLM-R domain (external)	12960	1.004	1172	0	0	0.689
Pooled (primary; scale .2)	25920	1.005	745	0	0	0.700
Pooled (primary; scale .5)	25920	1.004	486	0	0	0.695
Pooled (external; scale .2)	25920	1.006	559	0	0	0.718
Pooled (external; scale .5)	25920	1.008	573	0	0	0.697

All eight model-by-model rows and all four pooled rows are completed four-chain, 3,000-warm-up/3,000-retained model fits on the complete support. Sampling diagnostics showed zero divergent transitions and zero maximum-depth saturations; the table reports the minimum E-BFMI across chains and the minimum bulk ESS among substantive parameters. For the reference-style XLM-R domain analysis, the adjusted posterior means were .953 [.951, .955] for Biological Core, .973 [.971, .975] for Nature & Motion, .938 [.936, .940] for Socio-Cultural, and .963 [.961, .965] for Grammar & Function. These formal intervals support the endpoint contrast but do not make the middle-domain order encoder-invariant. The pooled endpoint contrast is 0.0044 $[-0.0003, 0.0090]$ under the primary taxonomy and -0.0029 $[-0.0071, 0.0014]$ under the reference-style taxonomy; the prior-scale sensitivity leaves both conclusions unchanged. The fixed-sample coverage check remains descriptive.

Table S18. Layer-wise RSA sensitivity analysis for Grammar & Function and Socio-Cultural. Every row uses the same 81 languages, 196 complete-case concepts, and 3,240 dyads. Slopes and 95% ordinary least-squares intervals are descriptive and are not additional confirmatory tests.

Layer	Domain	Mean	Geo.	95% CI	ASJP	95% CI
0	Grammar & Function	0.8789	-0.0002	[-0.0042, 0.0038]	0.0285	[0.0245, 0.0324]
0	Socio-Cultural	0.8646	0.0133	[0.0099, 0.0168]	0.0303	[0.0268, 0.0338]
1	Grammar & Function	0.8891	-0.0049	[-0.0087, -0.0011]	0.0273	[0.0235, 0.0311]
1	Socio-Cultural	0.8701	0.0121	[0.0088, 0.0154]	0.0314	[0.0281, 0.0347]
2	Grammar & Function	0.8876	-0.0048	[-0.0087, -0.0009]	0.0294	[0.0255, 0.0332]
2	Socio-Cultural	0.8608	0.0135	[0.0101, 0.0169]	0.0339	[0.0305, 0.0373]
3	Grammar & Function	0.8749	-0.0049	[-0.0088, -0.0009]	0.0327	[0.0288, 0.0367]
3	Socio-Cultural	0.8464	0.0158	[0.0122, 0.0193]	0.0362	[0.0326, 0.0397]
4	Grammar & Function	0.8648	-0.0020	[-0.0061, 0.0021]	0.0332	[0.0291, 0.0373]
4	Socio-Cultural	0.8351	0.0169	[0.0132, 0.0206]	0.0369	[0.0332, 0.0406]
5	Grammar & Function	0.8662	0.0004	[-0.0037, 0.0046]	0.0325	[0.0284, 0.0366]
5	Socio-Cultural	0.8289	0.0168	[0.0130, 0.0205]	0.0364	[0.0327, 0.0402]
6	Grammar & Function	0.8688	0.0007	[-0.0033, 0.0047]	0.0315	[0.0275, 0.0355]
6	Socio-Cultural	0.8282	0.0157	[0.0119, 0.0195]	0.0345	[0.0307, 0.0383]
7	Grammar & Function	0.8733	0.0025	[-0.0014, 0.0063]	0.0279	[0.0240, 0.0318]
7	Socio-Cultural	0.8282	0.0158	[0.0120, 0.0196]	0.0342	[0.0304, 0.0380]
8	Grammar & Function	0.8811	0.0062	[0.0022, 0.0102]	0.0271	[0.0231, 0.0311]
8	Socio-Cultural	0.8317	0.0171	[0.0133, 0.0209]	0.0331	[0.0293, 0.0369]
9	Grammar & Function	0.8791	0.0091	[0.0050, 0.0132]	0.0278	[0.0238, 0.0319]
9	Socio-Cultural	0.8365	0.0171	[0.0134, 0.0208]	0.0325	[0.0289, 0.0362]
10	Grammar & Function	0.8836	0.0090	[0.0050, 0.0130]	0.0272	[0.0232, 0.0312]
10	Socio-Cultural	0.8508	0.0151	[0.0117, 0.0185]	0.0302	[0.0268, 0.0336]
11	Grammar & Function	0.8897	0.0071	[0.0032, 0.0111]	0.0261	[0.0221, 0.0300]
11	Socio-Cultural	0.8622	0.0148	[0.0116, 0.0180]	0.0283	[0.0251, 0.0315]
12	Grammar & Function	0.8924	0.0057	[0.0017, 0.0096]	0.0285	[0.0246, 0.0325]
12	Socio-Cultural	0.8742	0.0165	[0.0136, 0.0195]	0.0263	[0.0233, 0.0292]

Table S19. Frequency-adjusted item-level models for both encoders. Estimates are REML fixed effects; intervals are 95% Wald intervals, not Bayesian credible intervals. Each fit uses all 635,040 observations.

Encoder	Predictor	Estimate	95% Wald interval
mbert	Intercept	0.93557	[0.92232, 0.94881]
mbert	Geo_Z	0.00286	[0.00221, 0.00351]
mbert	ASJP_Z	0.04151	[0.04090, 0.04211]
mbert	Frequency_Z	-0.02620	[-0.03228, -0.02012]
mbert	DomainNature_Motion	-0.00856	[-0.02240, 0.00528]
mbert	DomainSocio_Cultural	-0.00775	[-0.02079, 0.00529]
mbert	DomainGrammar_Function	-0.02010	[-0.04058, 0.00038]
mbert	Geo_Z:DomainNature_Motion	-0.00127	[-0.00216, -0.00038]
mbert	Geo_Z:DomainSocio_Cultural	0.00295	[0.00211, 0.00380]
mbert	Geo_Z:DomainGrammar_Function	0.00898	[0.00780, 0.01017]
mbert	ASJP_Z:DomainNature_Motion	0.00389	[0.00300, 0.00477]
mbert	ASJP_Z:DomainSocio_Cultural	-0.00395	[-0.00479, -0.00310]
mbert	ASJP_Z:DomainGrammar_Function	0.00310	[0.00192, 0.00429]
xlmr	Intercept	0.91213	[0.89302, 0.93124]
xlmr	Geo_Z	0.00526	[0.00391, 0.00662]
xlmr	ASJP_Z	0.03331	[0.03205, 0.03457]
xlmr	Frequency_Z	-0.01414	[-0.02424, -0.00404]
xlmr	DomainNature_Motion	0.02264	[-0.00037, 0.04565]
xlmr	DomainSocio_Cultural	-0.00940	[-0.03108, 0.01228]
xlmr	DomainGrammar_Function	-0.04103	[-0.07507, -0.00698]
xlmr	Geo_Z:DomainNature_Motion	-0.00204	[-0.00390, -0.00019]
xlmr	Geo_Z:DomainSocio_Cultural	-0.00063	[-0.00239, 0.00114]
xlmr	Geo_Z:DomainGrammar_Function	0.00911	[0.00664, 0.01158]
xlmr	ASJP_Z:DomainNature_Motion	0.00083	[-0.00103, 0.00269]
xlmr	ASJP_Z:DomainSocio_Cultural	-0.00251	[-0.00427, -0.00075]
xlmr	ASJP_Z:DomainGrammar_Function	0.00849	[0.00602, 0.01096]

Table S20. The 81 languages retained after excluding degenerate Javanese geometry.

Index	Language
1	Afrikaans
2	Albanian
3	Arabic

Continued on next page

Index	Language
4	Armenian
5	Asturian
6	Azerbaijani
7	Bashkir
8	Basque
9	Belarusian
10	Bengali
11	Bokmål
12	Breton
13	Bulgarian
14	Burmese
15	Catalan
16	Chechen
17	Chuvash
18	Czech
19	Danish
20	Dutch
21	English
22	Estonian
23	Finnish
24	French
25	Galician
26	Georgian
27	German
28	Greek
29	Gujarati
30	Hebrew
31	Hindi
32	Hungarian
33	Icelandic
34	Indonesian
35	Italian
36	Japanese
37	Kannada
38	Kazakh
39	Korean
40	Kyrgyz
41	Latvian
42	Lithuanian
43	Low Saxon
44	Macedonian
45	Malagasy
46	Malay
47	Malayalam
48	Mandarin
49	Marathi
50	Mongolian
51	Nepali
52	Nynorsk
53	Occitan
54	Persian
55	Piedmontese
56	Polish
57	Portuguese
58	Punjabi
59	Romanian
60	Russian
61	Scots
62	Serbo-Croatian
63	Slovak
64	Slovene
65	Spanish
66	Sundanese

Continued on next page

Index	Language
67	Swahili
68	Swedish
69	Tagalog
70	Tajik
71	Tamil
72	Tatar
73	Telugu
74	Thai
75	Turkish
76	Ukrainian
77	Urdu
78	Uzbek
79	Vietnamese
80	Welsh
81	West Frisian

Table S21. XLM-R-large extraction and fixed-sample RSA provenance. The checkpoint was loaded from local server storage without network fallback.

Item	Value
Checkpoint	`xlm-roberta-large`
Local checkpoint directory	`models/xlm-roberta-large`
Architecture	XLMRobertaModel
Layers / hidden size	24 / 1024
Layer / pooling	final; mean non-special-token states within alternative; mean across comma-delimited alternatives
Input	isolated unmarked lexical forms
Languages completed	81 / 81
Completed QC rows	81
Software	Python 3.12.4; torch 2.4.0+cu121; transformers 5.14.1
Config SHA-256	`ec7c3a99c58a38ebb702a2297f1c4586944e841b5eb29d201ff87e0f9c9abe0e`
XLM-R RSA SHA-256	`dee33af4d8481b33c34995b291c5c47a6735981102f6f6870a0a46c67e5f4420`

Table S22. Fixed-sample domain means and leave-one-language-out ranges for both encoders and both taxonomies. All rows use the same 81 languages, 196 concepts, and 3,240 dyads.

Encoder	Taxonomy	Domain	Mean	LOO range
mBERT	Primary	Biological Core	0.900	[0.898, 0.904]
XLM-R-large	Primary	Biological Core	0.932	[0.930, 0.933]
mBERT	Primary	Grammar & Function	0.892	[0.889, 0.895]
XLM-R-large	Primary	Grammar & Function	0.967	[0.966, 0.969]
mBERT	Primary	Nature & Motion	0.924	[0.923, 0.927]
XLM-R-large	Primary	Nature & Motion	0.977	[0.976, 0.978]
mBERT	Primary	Socio-Cultural	0.874	[0.871, 0.877]
XLM-R-large	Primary	Socio-Cultural	0.922	[0.920, 0.924]
mBERT	External	Biological Core	0.918	[0.916, 0.920]
XLM-R-large	External	Biological Core	0.953	[0.952, 0.954]
mBERT	External	Grammar & Function	0.892	[0.889, 0.894]
XLM-R-large	External	Grammar & Function	0.963	[0.962, 0.965]
mBERT	External	Nature & Motion	0.921	[0.919, 0.923]
XLM-R-large	External	Nature & Motion	0.973	[0.972, 0.974]
mBERT	External	Socio-Cultural	0.883	[0.880, 0.886]
XLM-R-large	External	Socio-Cultural	0.938	[0.937, 0.940]

Table S23. Dyad-level agreement between mBERT and XLM-R-large RSA dissimilarities. Median agreement is the fraction of dyads assigned to the same side of each model's median.

Taxonomy	Domain	ρ	Median agree.	Mean abs. diff.
Primary	All concepts	0.569	0.685	0.046
Primary	Biological Core	0.361	0.622	0.068
Primary	Grammar & Function	0.169	0.553	0.114
Primary	Nature & Motion	0.209	0.559	0.069
Primary	Socio-Cultural	0.531	0.680	0.075
External	All concepts	0.569	0.685	0.046
External	Biological Core	0.275	0.593	0.066
External	Grammar & Function	0.238	0.575	0.095
External	Nature & Motion	0.270	0.588	0.068
External	Socio-Cultural	0.522	0.674	0.078

Table S24. Leave-one-language-out stability and model-specific domain ordering. Each model's observed ordering (shown with abbreviated domain labels) was preserved in all 81 leave-one-out subsamples under both taxonomies.

Encoder	Taxonomy	Observed domain order	Shared extremes	Full order
XLM-R-large	Primary	S-Cultural < Bio. Core < Grammar < Nature	81/81	81/81
XLM-R-large	External	S-Cultural < Bio. Core < Grammar < Nature	81/81	81/81
mBERT	Primary	S-Cultural < Grammar < Bio. Core < Nature	81/81	81/81
mBERT	External	S-Cultural < Grammar < Bio. Core < Nature	81/81	81/81

Table S25. Software and provenance metadata for the analysis. Heavy numerical analyses were executed on the remote server; no local Bayesian sampling or model inference was used.

Item	Value
Encoder	mBERT and XLM-R-large; separate primary/external fits plus pooled model
Python	3.12.4
NumPy / pandas	1.26.4 / 2.3.3
PyTorch / Transformers	2.4.0+cu121 / 5.14.1
CmdStan	2.38.0; invoked by the remote Python driver
Chains	4
Warm-up / sampling	Full-support runs: 3000 / 3000 per chain; no reduced-iteration fit enters manuscript claims
Seed	42
mBERT RSA input SHA-256	`38563c1d16486c0d664381aa39c04b6beb1184b3719b4caa0cc1041a2de4ad9b`
XLM-R RSA input SHA-256	`dee33af4d8481b33c34995b291c5c47a6735981102f6f6870a0a46c67e5f4420`
Geographic matrix SHA-256	`3fe9fd1ebf3a4a199d5671b092944b0e9feff2292a1c00d0ad8725024b6c3862`
ASJP matrix SHA-256	`8ecfe16263f77cbdc894aeaca5b18c7750593fbd8c7716b48de5a90418521c9a`
Language-order file SHA-256	`d881281378d293d338ca8b17c9a840a8fd87002e324867f699813a95429b9fbf`
Taxonomy crosswalk SHA-256	`2661fed424363e679a4f3cede9949f1cbac664f0419a6d302f5abb820da70f29`
Global Stan SHA-256	`d53b6d6dcd783fc4feeadd5a82cfdf27847da9d1df23b9a1be6871a8e65d8f14`
Domain Stan SHA-256	`8762ddb5d8fb9e16580c61c25286b81627dd0c168a234b7d1c391b723a55da3e`
Pooled Stan SHA-256	`dce7806fc13131d27551f523541ffaac712f2eb17822b531532d6d3fb5e89ac2`

Table S26. Complete mapping of the 207 Wiktionary-numbered study concepts. The external mapping uses a study-specific alignment to Bowern/Concepticon entries, linked WOLD/IDS fields, WordNet 3.0 lexnames where applicable, and explicit functional rules. The auxiliary alignment does not imply identity of the original concept lists; the frozen WOLD/IDS-informed macro-domain rule is retained separately from official source fields.

ID	Concept	Primary	External	External field	Confidence
1	I	Biological Core	Grammar & Function	Pronouns	high
2	you	Biological Core	Grammar & Function	Pronouns	high
3	he,she,it	Biological Core	Grammar & Function	Pronouns	medium
4	we	Biological Core	Grammar & Function	Pronouns	high
5	you	Biological Core	Grammar & Function	Pronouns	medium
6	they	Biological Core	Grammar & Function	Pronouns	high
7	this	Grammar & Function	Grammar & Function	Deixis and reference	high
8	that	Grammar & Function	Grammar & Function	Deixis and reference	high
9	here	Grammar & Function	Grammar & Function	Space and location	high
10	there	Grammar & Function	Grammar & Function	Space and location	high
11	who	Grammar & Function	Grammar & Function	Function words	high
12	what	Grammar & Function	Grammar & Function	Function words	high
13	where	Grammar & Function	Grammar & Function	Function words	high
14	when	Grammar & Function	Grammar & Function	Function words	high
15	how	Grammar & Function	Grammar & Function	Function words	high
16	not	Grammar & Function	Grammar & Function	Function words or residual field	high
17	all	Grammar & Function	Grammar & Function	Quantity	high
18	many	Grammar & Function	Grammar & Function	Quantity	high
19	some	Grammar & Function	Grammar & Function	Quantity	high
20	few	Grammar & Function	Grammar & Function	Quantity	high
21	other	Grammar & Function	Grammar & Function	Function words or residual field	high
22	one	Socio-Cultural	Grammar & Function	Quantity	high
23	two	Socio-Cultural	Grammar & Function	Quantity	high
24	three	Socio-Cultural	Grammar & Function	Quantity	high
25	four	Socio-Cultural	Grammar & Function	Quantity	high
26	five	Socio-Cultural	Socio-Cultural	Quantity	high
27	big	Socio-Cultural	Socio-Cultural	Dimension	high
28	long	Socio-Cultural	Socio-Cultural	Dimension	high
29	wide	Socio-Cultural	Socio-Cultural	Dimension	high
30	thick	Socio-Cultural	Socio-Cultural	Dimension	high
31	heavy	Socio-Cultural	Socio-Cultural	Attributes and qualities	high
32	small	Socio-Cultural	Socio-Cultural	Dimension	high
33	short	Socio-Cultural	Socio-Cultural	Dimension	high
34	narrow	Socio-Cultural	Socio-Cultural	Dimension	high
35	thin	Socio-Cultural	Socio-Cultural	Dimension	high
36	woman	Biological Core	Biological Core	People	high
37	man	Biological Core	Biological Core	People	medium
38	man	Biological Core	Biological Core	People	medium
39	child	Biological Core	Biological Core	Kinship	high
40	wife	Biological Core	Biological Core	Kinship	high
41	husband	Biological Core	Biological Core	Kinship	high
42	mother	Biological Core	Biological Core	Kinship	high
43	father	Biological Core	Biological Core	Kinship	high
44	animal	Nature & Motion	Nature & Motion	Animals	high
45	fish	Nature & Motion	Nature & Motion	Animals	high
46	bird	Nature & Motion	Nature & Motion	Animals	high
47	dog	Nature & Motion	Nature & Motion	Animals	high
48	louse	Nature & Motion	Nature & Motion	Animals	high
49	snake	Nature & Motion	Nature & Motion	Animals	high
50	worm	Nature & Motion	Nature & Motion	Animals	high
51	tree	Nature & Motion	Nature & Motion	Plants	high
52	forest	Nature & Motion	Nature & Motion	Plants	high
53	stick	Nature & Motion	Nature & Motion	Plants	high
54	fruit	Nature & Motion	Nature & Motion	Plants	high
55	seed	Nature & Motion	Nature & Motion	Plants	high
56	leaf	Nature & Motion	Nature & Motion	Plants	high
57	root	Nature & Motion	Nature & Motion	Plants	high
58	bark	Nature & Motion	Nature & Motion	Plants	high
59	flower	Nature & Motion	Nature & Motion	Plants	high

Continued on next page

ID	Concept	Primary	External	External field	Confidence
60	grass	Nature & Motion	Nature & Motion	Plants	high
61	rope	Socio-Cultural	Socio-Cultural	Material culture	high
62	skin	Biological Core	Biological Core	Body and substance	high
63	meat	Biological Core	Biological Core	Body and substance	high
64	blood	Biological Core	Biological Core	Body and substance	high
65	bone	Biological Core	Biological Core	Body and substance	high
66	fat	Biological Core	Biological Core	Body and substance	high
67	egg	Biological Core	Biological Core	Body and substance	high
68	horn	Biological Core	Biological Core	Body and substance	high
69	tail	Biological Core	Biological Core	Body and substance	high
70	feather	Biological Core	Biological Core	Body and substance	high
71	hair	Biological Core	Biological Core	Body and substance	high
72	head	Biological Core	Biological Core	Body and substance	high
73	ear	Biological Core	Biological Core	Body and substance	high
74	eye	Biological Core	Biological Core	Body and substance	high
75	nose	Biological Core	Biological Core	Body and substance	high
76	mouth	Biological Core	Biological Core	Body and substance	high
77	tooth	Biological Core	Biological Core	Body and substance	high
78	tongue	Biological Core	Biological Core	Body and substance	high
79	fingernail	Biological Core	Biological Core	Body and substance	high
80	foot	Biological Core	Biological Core	Body and substance	high
81	leg	Biological Core	Biological Core	Body and substance	high
82	knee	Biological Core	Biological Core	Body and substance	high
83	hand	Biological Core	Biological Core	Body and substance	high
84	wing	Biological Core	Biological Core	Body and substance	high
85	belly	Biological Core	Biological Core	Body and substance	high
86	guts	Biological Core	Biological Core	Body and substance	high
87	neck	Biological Core	Biological Core	Body and substance	high
88	back	Biological Core	Biological Core	Body and substance	high
89	breast	Biological Core	Biological Core	Body and substance	high
90	heart	Biological Core	Biological Core	Body and substance	high
91	liver	Biological Core	Biological Core	Body and substance	high
92	drink	Biological Core	Biological Core	Food, bodily action, and speech	high
93	eat	Biological Core	Biological Core	Food, bodily action, and speech	high
94	bite	Biological Core	Biological Core	Food, bodily action, and speech	high
95	suck	Biological Core	Biological Core	Food, bodily action, and speech	high
96	spit	Biological Core	Biological Core	Food, bodily action, and speech	high
97	vomit	Biological Core	Biological Core	Food, bodily action, and speech	high
98	blow	Biological Core	Biological Core	Food, bodily action, and speech	high
99	breathe	Biological Core	Biological Core	Body and sensation	high
100	laugh	Biological Core	Biological Core	Body and sensation	high
101	see	Biological Core	Biological Core	Body and sensation	high
102	hear	Biological Core	Biological Core	Body and sensation	high
103	know	Biological Core	Socio-Cultural	Body and sensation	medium
104	think	Biological Core	Socio-Cultural	Body and sensation	medium
105	smell	Biological Core	Biological Core	Body and sensation	high
106	fear	Biological Core	Socio-Cultural	Body and sensation	medium
107	sleep	Biological Core	Biological Core	Body and sensation	high
108	live	Biological Core	Biological Core	Body and sensation	high
109	die	Biological Core	Biological Core	Body and sensation	high
110	kill	Socio-Cultural	Socio-Cultural	Action and manipulation	high
111	fight	Socio-Cultural	Socio-Cultural	Action and manipulation	high
112	hunt	Nature & Motion	Socio-Cultural	Material culture	medium
113	hit	Socio-Cultural	Socio-Cultural	Action and manipulation	high
114	cut	Socio-Cultural	Socio-Cultural	Action and manipulation	high
115	split	Socio-Cultural	Socio-Cultural	Action and manipulation	high
116	stab	Socio-Cultural	Socio-Cultural	Action and manipulation	high
117	scratch	Socio-Cultural	Socio-Cultural	Action and manipulation	high
118	dig	Socio-Cultural	Socio-Cultural	Action and manipulation	high
119	swim	Nature & Motion	Nature & Motion	Space and motion	high
120	fly	Nature & Motion	Nature & Motion	Space and motion	high
121	walk	Nature & Motion	Nature & Motion	Space and motion	high
122	come	Nature & Motion	Nature & Motion	Space and motion	high

Continued on next page

ID	Concept	Primary	External	External field	Confidence
123	lie	Nature & Motion	Nature & Motion	Space and motion	high
124	sit	Nature & Motion	Nature & Motion	Space and motion	high
125	stand	Nature & Motion	Nature & Motion	Space and motion	high
126	turn	Nature & Motion	Nature & Motion	Space and motion	high
127	fall	Nature & Motion	Nature & Motion	Space and motion	high
128	give	Socio-Cultural	Socio-Cultural	Action and manipulation	high
129	hold	Socio-Cultural	Socio-Cultural	Action and manipulation	high
130	squeeze	Socio-Cultural	Socio-Cultural	Action and manipulation	high
131	rub	Socio-Cultural	Socio-Cultural	Action and manipulation	high
132	wash	Socio-Cultural	Socio-Cultural	Action and manipulation	high
133	wipe	Socio-Cultural	Socio-Cultural	Action and manipulation	high
134	pull	Socio-Cultural	Socio-Cultural	Action and manipulation	high
135	push	Socio-Cultural	Socio-Cultural	Action and manipulation	high
136	throw	Socio-Cultural	Socio-Cultural	Action and manipulation	high
137	tie	Socio-Cultural	Socio-Cultural	Action and manipulation	high
138	sew	Socio-Cultural	Socio-Cultural	Action and manipulation	high
139	count	Biological Core	Grammar & Function	Food, bodily action, and speech	medium
140	say	Biological Core	Socio-Cultural	Food, bodily action, and speech	medium
141	sing	Biological Core	Socio-Cultural	Food, bodily action, and speech	medium
142	play	Nature & Motion	Socio-Cultural	Function words or residual field	medium
143	float	Nature & Motion	Nature & Motion	Space and motion	high
144	flow	Nature & Motion	Nature & Motion	Space and motion	high
145	freeze	Biological Core	Nature & Motion	Body and sensation	high
146	swell	Biological Core	Nature & Motion	Body and sensation	high
147	sun	Nature & Motion	Nature & Motion	Natural environment	high
148	moon	Nature & Motion	Nature & Motion	Natural environment	high
149	star	Nature & Motion	Nature & Motion	Natural environment	high
150	water	Nature & Motion	Nature & Motion	Natural environment	high
151	rain	Nature & Motion	Nature & Motion	Natural environment	high
152	river	Nature & Motion	Nature & Motion	Natural environment	high
153	lake	Nature & Motion	Nature & Motion	Natural environment	high
154	sea	Nature & Motion	Nature & Motion	Natural environment	high
155	salt	Nature & Motion	Nature & Motion	Natural environment	high
156	stone	Nature & Motion	Nature & Motion	Natural environment	high
157	sand	Nature & Motion	Nature & Motion	Natural environment	high
158	dust	Nature & Motion	Nature & Motion	Natural environment	high
159	earth	Nature & Motion	Nature & Motion	Natural environment	high
160	cloud	Nature & Motion	Nature & Motion	Natural environment	high
161	fog	Nature & Motion	Nature & Motion	Natural environment	high
162	sky	Nature & Motion	Nature & Motion	Natural environment	high
163	wind	Nature & Motion	Nature & Motion	Natural environment	high
164	snow	Nature & Motion	Nature & Motion	Natural environment	high
165	ice	Nature & Motion	Nature & Motion	Natural environment	high
166	smoke	Nature & Motion	Nature & Motion	Natural environment	high
167	fire	Nature & Motion	Nature & Motion	Natural environment	high
168	ash	Nature & Motion	Nature & Motion	Natural environment	high
169	burn	Biological Core	Nature & Motion	Body and sensation	medium
170	road	Socio-Cultural	Socio-Cultural	Material culture	high
171	mountain	Nature & Motion	Nature & Motion	Natural environment	high
172	red	Socio-Cultural	Socio-Cultural	Colour	high
173	green	Socio-Cultural	Socio-Cultural	Colour	high
174	yellow	Socio-Cultural	Socio-Cultural	Colour	high
175	white	Socio-Cultural	Socio-Cultural	Colour	high
176	black	Socio-Cultural	Socio-Cultural	Colour	high
177	night	Socio-Cultural	Socio-Cultural	Time	high
178	day	Socio-Cultural	Socio-Cultural	Time	high
179	year	Socio-Cultural	Socio-Cultural	Time	high
180	warm	Socio-Cultural	Socio-Cultural	Attributes and qualities	high
181	cold	Socio-Cultural	Socio-Cultural	Attributes and qualities	high
182	full	Socio-Cultural	Socio-Cultural	Attributes and qualities	high
183	new	Socio-Cultural	Socio-Cultural	Attributes and qualities	high
184	old	Socio-Cultural	Socio-Cultural	Attributes and qualities	high
185	good	Socio-Cultural	Socio-Cultural	Attributes and qualities	high

Continued on next page

ID	Concept	Primary	External	External field	Confidence
186	bad	Socio-Cultural	Socio-Cultural	Attributes and qualities	high
187	rotten	Socio-Cultural	Socio-Cultural	Attributes and qualities	high
188	dirty	Socio-Cultural	Socio-Cultural	Attributes and qualities	high
189	straight	Socio-Cultural	Socio-Cultural	Attributes and qualities	high
190	round	Socio-Cultural	Socio-Cultural	Attributes and qualities	high
191	sharp	Socio-Cultural	Socio-Cultural	Attributes and qualities	high
192	dull	Socio-Cultural	Socio-Cultural	Attributes and qualities	high
193	smooth	Socio-Cultural	Socio-Cultural	Attributes and qualities	high
194	wet	Socio-Cultural	Socio-Cultural	Attributes and qualities	high
195	dry	Socio-Cultural	Socio-Cultural	Attributes and qualities	high
196	correct	Socio-Cultural	Socio-Cultural	Attributes and qualities	high
197	near	Grammar & Function	Grammar & Function	Space and location	high
198	far	Grammar & Function	Grammar & Function	Space and location	high
199	right	Grammar & Function	Grammar & Function	Space and location	medium
200	left	Grammar & Function	Grammar & Function	Space and location	high
201	at	Grammar & Function	Grammar & Function	Space and location	high
202	in	Grammar & Function	Grammar & Function	Space and location	high
203	with	Grammar & Function	Grammar & Function	Space and location	high
204	and	Grammar & Function	Grammar & Function	Function words or residual field	high
205	if	Grammar & Function	Grammar & Function	Function words or residual field	high
206	because	Grammar & Function	Grammar & Function	Function words or residual field	high
207	name	Grammar & Function	Socio-Cultural	Function words or residual field	medium

Table S27. Complete pooled two-domain contrasts under default and sensitivity priors. Within-encoder and between-encoder estimands are listed separately.

Mapping	Prior branch	Estimand	Mean [95% CrI]
primary	Default (0.2)	Nature & Motion - Socio-Cultural | mBERT	0.05031 [0.04702, 0.05366]
primary	Default (0.2)	Nature & Motion - Socio-Cultural | XLM-R-large	0.05472 [0.05128, 0.05803]
primary	Default (0.2)	(XLM-R-large - mBERT) endpoint contrast	0.00441 [-0.00034, 0.00897]
primary	Sensitivity (0.5)	Nature & Motion - Socio-Cultural | mBERT	0.05030 [0.04695, 0.05369]
primary	Sensitivity (0.5)	Nature & Motion - Socio-Cultural | XLM-R-large	0.05469 [0.05129, 0.05805]
primary	Sensitivity (0.5)	(XLM-R-large - mBERT) endpoint contrast	0.00439 [-0.00046, 0.00915]
external	Default (0.2)	Nature & Motion - Socio-Cultural | mBERT	0.03749 [0.03454, 0.04052]
external	Default (0.2)	Nature & Motion - Socio-Cultural | XLM-R-large	0.03461 [0.03163, 0.03758]
external	Default (0.2)	(XLM-R-large - mBERT) endpoint contrast	-0.00288 [-0.00709, 0.00138]
external	Sensitivity (0.5)	Nature & Motion - Socio-Cultural | mBERT	0.03750 [0.03444, 0.04047]
external	Sensitivity (0.5)	Nature & Motion - Socio-Cultural | XLM-R-large	0.03461 [0.03157, 0.03768]
external	Sensitivity (0.5)	(XLM-R-large - mBERT) endpoint contrast	-0.00289 [-0.00709, 0.00139]

S15 Encoder comparison details

Both encoders use the same 81 languages, 196 complete-case concept identifiers, domain mappings, lexical input rule, final-layer nonspecial-token pooling, within-set centering, RSA construction, and predictor matrices. The domain fits contain 12,960 rows each. The pooled model contains 25,920 rows. The mBERT extraction manifest records 83 requested extraction files, while the assembled study inventory contained 82 languages and the comparative analysis retained 81; extraction requests and analytic eligibility counts are different records.

The frequency-adjusted models use all 635,040 concept–dyad observations, English Zipf frequency, and exact Gaussian REML with marginalized language and concept effects. Their optimizer diagnostics report successful convergence and a full-rank 13-column fixed-effect design. These models operate on item-level cosine distances.

Equal-size subsampling and the 13-layer table use archived mBERT vectors identified in their respective manifests and serve as preliminary sensitivity analyses. The source taxonomy review forms document operational classifications.

S16 Supplementary figures



Fig. S1. mBERT concept-level heterogeneity. Panel (a) reports item-level regression coefficients for individual concepts. Panel (b) summarizes the full-inventory mBERT representational profile; arrows indicate observed associations. The geographic contrast shown is specific to the mBERT primary-taxonomy configuration.

Supplementary References

Beckner, C., Blythe, R., Bybee, J., Christiansen, M. H., Croft, W., Ellis, N. C., Holland, J., Ke, J., Larsen-Freeman, D., & Schoenemann, T. (2009). Language is a complex adaptive system: Position paper. *Language Learning*, *59*(s1), 1–26. https://doi.org/10.1111/j.1467-9922.2009.00533.x

Bowern, C. (2008). Bowern 2008 207. https://concepticon.clld.org/contributions/Bowern-2008-207

Burenhult, N., & Levinson, S. C. (2008). Language and landscape: A cross-linguistic perspective. *Language Sciences*, *30*(2-3), 135–150. https://doi.org/10.1016/j.langsci.2006.12.028

Bybee, J. (2010). *Language, usage and cognition*. Cambridge University Press. https://doi.org/10.1017/CBO9780511750526

Christiansen, M. H., & Chater, N. (2008). Language as shaped by the brain. *Behavioral and Brain Sciences*, *31*(5), 489–509. https://doi.org/10.1017/S0140525X08004998

Church, K. W., & Hanks, P. (1990). Word association norms, mutual information, and lexicography. *Computational Linguistics*, *16*(1), 22–29.

Croft, W. (2001). *Radical construction grammar: Syntactic theory in typological perspective*. Oxford University Press. https://doi.org/10.1093/acprof:oso/9780198299554.001.0001

Fellbaum, C. (Ed.). (1998). *Wordnet: An electronic lexical database*. MIT Press. https://doi.org/10.7551/mitpress/7287.001.0001

Haspelmath, M., & Tadmor, U. (2009). World loanword database. https://wold.clld.org/

Jackson, J. C., Watts, J., Henry, T. R., List, J.-M., Forkel, R., Mucha, P. J., Greenhill, S. J., Gray, R. D., & Lindquist, K. A. (2019). Emotion semantics show both cultural variation and universal structure. *Science*, *366*(6472), 1517–1522. https://doi.org/10.1126/science.aaw8160

Jäger, G. (2018). Global-scale phylogenetic linguistic inference from lexical resources. *Scientific Data*, *5*, 180189. https://doi.org/10.1038/sdata.2018.189

Kemp, C., & Regier, T. (2012). Kinship categories across languages reflect general communicative principles. *Science*, *336*(6084), 1049–1054. https://doi.org/10.1126/science.1218811

Key, M. R., & Comrie, B. (Eds.). (2015). *The intercontinental dictionary series*. Max Planck Institute for Evolutionary Anthropology. https://ids.clld.org/

Levinson, S. C. (2003). *Space in language and cognition: Explorations in cognitive diversity*. Cambridge University Press. https://doi.org/10.1017/CBO9780511613609

Levinson, S. C., & Wilkins, D. P. (Eds.). (2006). *Grammars of space: Explorations in cognitive diversity*. Cambridge University Press. https://doi.org/10.1017/CBO9780511486753

List, J.-M., Cysouw, M., & Forkel, R. (2016). Concepticon: A resource for the linking of concept lists. *Proceedings of the Tenth International Conference on Language Resources and Evaluation (LREC 2016)*, 2393–2400. https://aclanthology.org/L16-1379/

Majid, A., & Burenhult, N. (2014). Odors are expressible in language, as long as you speak the right language. *Cognition*, *130*(2), 266–270. https://doi.org/10.1016/j.cognition.2013.11.004

Malt, B. C., & Majid, A. (2013). How thought is mapped into words. *Wiley Interdisciplinary Reviews: Cognitive Science*, *4*(6), 583–597. https://doi.org/10.1002/wcs.1251

Malt, B. C., Sloman, S. A., Gennari, S., Shi, M., & Wang, Y. (1999). Knowing versus naming: Similarity and the linguistic categorization of artifacts. *Journal of Memory and Language*, *40*(2), 230–262. https://doi.org/10.1006/jmla.1998.2593

Pagel, M. (2009). Human language as a culturally transmitted replicator. *Nature Reviews Genetics*, *10*(6), 405–415. https://doi.org/10.1038/nrg2560

Spelke, E. S., & Kinzler, K. D. (2007). Core knowledge. *Developmental Science*, *10*(1), 89–96. https://doi.org/10.1111/j.1467-7687.2007.00569.x

Talmy, L. (1985). Lexicalization patterns: Semantic structure in lexical forms. In T. Shopen (Ed.), *Language typology and syntactic description* (pp. 57–149, Vol. 3). Cambridge University Press.

Thompson, B., Roberts, S. G., & Lupyan, G. (2020). Cultural influences on word meanings revealed through large-scale semantic alignment. *Nature Human Behaviour*, *4*(10), 1029–1038. https://doi.org/10.1038/s41562-020-0924-8

Wiktionary contributors. (n.d.-a). *Appendix: English swadesh list*. Wiktionary. Retrieved October 22, 2026, from https://en.wiktionary.org/wiki/Appendix:English_Swadesh_list

Wiktionary contributors. (n.d.-b). *Swadesh lists by language*. Wiktionary. Retrieved October 22, 2026, from https://en.wiktionary.org/wiki/Category:Swadesh_lists_by_language