arXiv:2611.06471v1 [cs.CV] 4 Nov 2026

A 3D foundation model unifying CT interpretation and segmentation

Yulu Xie[1,2], **Zilong Li**[2], **Shen Xu**[3,2], **Yifan Bai**[2], **Liuhong Shi**[4], **Jiao Chen**[5], **Xianglei Kong**[5], **Wenchao Zhu**[5], **Na Shen**[6], **Xianghua Ye**[7], **Yang Yan**[8], **Jiaju Zheng**[8], **Yunsong Peng**[9], **Yanzhou Su**[2], **Chaohui Yu**[2], **Hao Luo**[2,10,11,†], **Fan Wang**[2,†]

[1]Department of Statistics and Data Science, Tsinghua University, Beijing, China [2]Alibaba DAMO Academy, Hangzhou, China [3]School of Psychological and Cognitive Sciences and Beijing Key Laboratory of Behavior and Mental Health, Peking University, Beijing, China [4]Department of Head and Neck Surgery, Sir Run Run Shaw Hospital, Zhejiang University School of Medicine, Hangzhou, China [5]Department of Radiology, Sir Run Run Shaw Hospital, Zhejiang University School of Medicine, Hangzhou, China [6]Department of Otolaryngology-Head & Neck Surgery, Zhongshan Hospital, Fudan University, Shanghai, China [7]Department of Radiation Oncology, The First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou, China [8]Department of Radiology, The Affiliated Yangming Hospital of Ningbo University, Yuyao, China [9]Guizhou Province International Science and Technology Cooperation Base for Precision Imaging Diagnosis and Treatment, Key Laboratory of Advanced Medical Imaging and Intelligent Computing of Guizhou Province, Department of Radiology, Guizhou Provincial People's Hospital, Guiyang, China [10]College of Computer Science and Technology, Zhejiang University, Hangzhou, China [11]Hupan Laboratory, Hangzhou, China
[†]Corresponding authors

Radiologists interpret computed tomography (CT) scans by jointly determining what a finding represents and where it lies in three-dimensional (3D) anatomy. However, deep learning models for CT have largely treated CT interpretation and 3D segmentation as separate tasks, leaving textual findings spatially unverifiable and overlooking potential synergy between clinical understanding and spatial perception. Here we present HoloCT, a generalist 3D foundation model unifying CT interpretation and voxel-level segmentation by jointly modelling whole-volume context and fine-grained anatomical detail, performing diverse clinical tasks while providing 3D masks as evidence. To develop and evaluate HoloCT at scale, we curated HoloDATA, comprising 195,921 CT scans from 73 public datasets and six clinical cohorts, with 6.2 million instruction dialogues and 2.8 million CT–mask pairs. Across 346 tasks spanning segmentation, lesion screening, visual question answering, and report generation on 61 internal and 12 external datasets, HoloCT outperformed both specialist and foundation models. In multicentre validation across 11,455 clinical CT scans from five hospitals, HoloCT outperformed competing models in lesion segmentation and report generation by 18.2 and 15.3 points. It also achieved 85.1% mean accuracy in screening thyroid tumours and colon lesions, exceeding all four participating radiologists in a retrospective reader study. Further analyses revealed bidirectional semantic–spatial synergy: language-guided segmentation delineated unseen anatomical structures and lesions as well as diffuse free-text findings, outperforming the leading approach by 22.1 points, whereas prompting the model to generate corresponding 3D masks alongside its textual outputs significantly improved interpretation performance. In summary, HoloCT shows that clinical understanding and spatial perception can be learned together within a single foundation model and offers a path towards more accurate and reliable CT analysis.

Code: https://github.com/alibaba-damo-academy/HoloCT

1 Introduction

Computed tomography (CT) provides rapid, high-resolution volumetric imaging for clinical screening, diagnosis, and treatment planning[1]. When interpreting a scan, radiologists continuously link *what* a finding represents to *where* it lies in three-dimensional (3D) anatomy. Reliable CT interpretation requires both: textual descriptions without anatomical localization are difficult to verify across hundreds of slices, and segmented regions without clinical characterization carry limited diagnostic meaning.

Existing deep learning systems have largely developed these two capabilities in isolation. Segmentation networks delineate organs and abnormalities at the voxel level yet do not interpret their clinical significance[2–5]. Conversely, CT vision–language models (VLMs) generate diagnostic text without providing voxel-level localization, making the anatomical basis of their predictions difficult to inspect[6–8]. Recent grounded medical models connect language and segmentation at the output level[9–14]; however, for 3D scans, existing models route high-resolution spatial features only to the mask decoder rather than the language model, limiting the use of fine-grained spatial information during text generation[12–14]. Moreover, despite advances in voxel-level visual encoders[15], many of these models still rely on 2D slice encoders or coarsely downsampled 3D inputs at the expense of cross-slice continuity or fine spatial detail[12;14]. Whether clinical understanding and spatial perception can be learned jointly within a single foundation model and mutually reinforce one another therefore remains an open question.

Here we introduce HoloCT, a generalist 3D foundation model that unifies CT interpretation and 3D segmentation by jointly modelling whole-volume context and fine-grained anatomical detail within a unified vision–language representation (Fig. 1). Specifically, HoloCT couples a dual-scale visual encoding module, a unified multimodal large language model, and a mask decoder. First, each scan is encoded through two parallel branches: a global encoder for whole-volume context and a detailed encoder that preserves fine-grained anatomical detail at standardized isotropic spacing. Second, global tokens and fine-grained spatial tokens from the two encoders are supplied directly to the language model, enabling clinical reasoning to draw on fine-grained spatial features. Third, segmentation tokens emitted by the language model guide the mask decoder for voxel-level delineation. Accordingly, a single HoloCT model performs language-guided segmentation, lesion screening, visual question answering (VQA), and radiology report generation. For grounded interpretation, HoloCT can pair its textual outputs with 3D masks of the corresponding anatomical regions or findings, providing spatial evidence that readers can inspect.

To develop and evaluate HoloCT at scale, we curated HoloDATA by converting segmentation annotations, question–answer pairs, and radiology reports across datasets into a unified instruction–response format, with 3D masks incorporated where available. HoloDATA comprises 195,921 CT scans totalling more than 61 million slices from 73 public datasets and six clinical cohorts at five hospitals, and contains 6.2 million instruction dialogues and 2.8 million CT–mask pairs across 246 anatomical structures and 50 lesion categories.

Evaluated on 346 tasks across 11 task formulations on 61 internal and 12 external datasets, a single HoloCT-8B model outperformed 82 dataset-specific nnU-Net models on internal segmentation benchmarks and surpassed segmentation foundation models in zero-shot external transfer across both seen and unseen categories of anatomical structures and lesions. HoloCT also exceeded task-specific classifiers as well as medical and general-purpose VLMs in lesion screening, VQA, and report generation. In multicentre clinical validation across 11,455 CT scans from six cohorts at five hospitals, HoloCT surpassed competing models in the segmentation of five lesion categories, including laryngeal lesions, lung cancer and thyroid tumours, as well as in zero-shot abdominal and chest CT report generation. In retrospective reader studies of histopathology-confirmed thyroid tumours and colon lesions, HoloCT-8B achieved higher sensitivity and accuracy than all four participating radiologists, including two senior readers.

Beyond benchmark performance, our analyses revealed bidirectional synergy between clinical understanding and spatial perception. During training, adding text-only dialogues improved 3D segmentation accuracy, whereas incorporating mask-paired dialogues improved lesion screening, VQA, and report generation. At inference, language-guided segmentation delineated previously unseen anatomical structures and lesions as well as diffuse free-text findings, outperforming the leading approach by 22.1 points; conversely, prompting HoloCT to generate corresponding 3D masks alongside its textual outputs significantly improved performance across these interpretation tasks while maintaining mask quality. Together, these results demonstrate that unifying clinical understanding and spatial perception within a single foundation model strengthens both capabilities, aligning AI with how radiologists link diagnostic characterization to 3D anatomy when interpreting CT scans.

2 Results

2.1 Overview of HoloCT and HoloDATA

HoloCT is a generalist 3D foundation model developed and evaluated on the HoloDATA corpus to unify CT interpretation and voxel-level segmentation across volumetric CT tasks (Fig. 1). HoloDATA comprises 195,921 volumetric CT scans with more than 61 million slices, integrating 184,466 scans from 73 public datasets with 11,455 scans from six clinical cohorts at five hospitals (Fig. 1a). Of the 73 public datasets, 61 internal datasets supported model development and held-out testing, whereas 12 external datasets were held out entirely for zero-shot and fine-tuned transfer. By standardizing segmentation, question–answer, and radiology report annotations across datasets into a unified dialogue format with 3D masks without manual relabelling, HoloDATA provides 6.2 million instruction dialogues and 2.8 million CT–mask pairs from non-contrast and contrast-enhanced scans, covering 246 anatomical structures and 50 lesion categories across six major anatomical systems (Fig. 1b).

Architecturally, HoloCT encodes each CT scan through two parallel branches (Fig. 1c): a global encoder for whole-volume context and a detailed encoder that preserves fine-grained anatomical detail at standardized isotropic spacing. Features from both encoders are projected into whole-volume global tokens and fine-grained spatial tokens and supplied directly to a unified multimodal large language model instantiated as HoloCT-4B and HoloCT-8B, integrating whole-volume and fine-grained features within a shared vision–language representation. When spatial localization is requested, segmentation tokens emitted by the language model interact with multiscale skip features from the detailed encoder in a mask decoder, pairing diagnostic text with 3D masks.

2.2 Evaluation settings

We evaluated HoloCT on 346 tasks spanning four task families and 11 formulations (Fig. 1d): referring segmentation (target-level and finding-level), lesion screening (single-target and multi-target), VQA (closed-ended, open-ended and grounded) and report generation (narrative, structured, anatomy-grounded and finding-grounded). Evaluations were organized into three tiers without scan overlap: held-out test splits of 61 internal public datasets, 12 external public datasets excluded from training, covering seen and unseen categories as well as VQA and report generation, and six clinical cohorts for lesion segmentation (centres 1–3), histopathology-confirmed lesion screening (centres 3 and 4) and report generation (centre 5). More details can be found in Supplementary Tables 1–5.

A single pretrained HoloCT model was evaluated without task-specific fine-tuning on all internal benchmarks and in zero-shot external and clinical evaluations. For supervised comparisons in external lesion screening and segmentation, we additionally fine-tuned all unified models per task on the pooled training splits.

HoloCT was compared with three families of state-of-the-art models: specialist networks trained per benchmark, including nnU-Net[2] for segmentation and 3D ResNet-18[16] for screening; segmentation foundation models VISTA3D[3], SAT-Pro[4] and VoxTell[5]; and vision–language models Qwen3-VL-32B[17], MedGemma 1.5-4B[18], RadFM-14B[8], HuluMed-32B[19] and M3D-Phi3-4B (M3D)[12]. Details of these methods can be found in Supplementary Methods. As few existing models support finding-grounded report generation, we additionally constructed a two-stage pipeline using HuluMed-32B and VoxTell for comparison. For fair comparison, pretrained baselines were evaluated using released weights, whereas specialist and fine-tuned models used the corresponding training splits. In clinical screening, HoloCT was also compared with four radiologists in a blinded reader study.

Segmentation was assessed using Dice and HD95[20], with hit rates at Dice thresholds of 0.05 and 0.10 ($HIT_{5\%}$ and $HIT_{10\%}$) for finding-level referring segmentation. Screening was assessed using F1, sensitivity, specificity and accuracy, together with conditional Dice on true-positive detections; closed-ended VQA was assessed by accuracy, and grounded outputs additionally by match rate and Dice. Free-text outputs were scored with lexical metrics (BLEU-4[21], ROUGE-L[22] and METEOR[23]) and clinical metrics (LlamaScore, evaluated using Llama-3.1-70B-Instruct[24], plus RaTEScore[25] and GREEN[26] for reports). Building on prior work emphasizing both linguistic and clinical measures of report quality[27;28], we summarized free-text quality using the Language–Clinical Score (LCS), the equal-weight mean of linguistic similarity and clinical agreement (Methods).

2.3 Open-vocabulary 3D segmentation

We first evaluated HoloCT's foundational spatial perception through language-guided target-level referring segmentation across 272 internal and 51 external anatomical and lesion categories (Fig. 2, Supplementary Table 8). On the 128 internal categories supported by the segmentation foundation model VISTA3D, HoloCT-8B increased mean Dice from 0.837 to 0.870 ($P < 0.001$; Fig. 2a). Across all 272 internal categories, a single HoloCT-8B model achieved a mean Dice of 0.816, outperforming the collection of 82 dedicated specialist nnU-Net models trained on individual datasets (0.799, $P < 0.001$; Fig. 2b).

To assess zero-shot generalization beyond the training distribution, we next evaluated HoloCT without fine-tuning across 51 external categories comprising 37 seen and 14 unseen structures and lesions, where it consistently outperformed VISTA3D, SAT-Pro, M3D, and VoxTell (Fig. 2c–f). Specifically, on the 37 seen categories, HoloCT-8B reached a mean Dice of 0.658, surpassing specialist nnU-Net models by 0.339 (0.319, $P < 0.001$; Fig. 2b) and the strongest text-prompted baseline, VoxTell, by 0.108 ($P < 0.001$). It led five of the six anatomical and lesion groups, with VoxTell slightly higher on vascular structures (0.436 compared with 0.423; Fig. 2c,d). On the 14 unseen categories, HoloCT-8B increased mean Dice over VoxTell from 0.340 to 0.414 ($P < 0.001$) and led four of the five groups, with VISTA3D slightly higher on musculoskeletal bone lesions (0.165 compared with 0.154; Fig. 2e,f). In summary, a single HoloCT surpassed both specialist and foundation models on in-distribution benchmarks and demonstrated strong generalization performance to unseen anatomy and lesions.

2.4 Single- and multi-target grounded clinical screening

Beyond delineating known targets, clinical screening requires jointly determining whether an abnormality is present and localizing detected lesions on scans (Fig. 3a). We evaluated single-target screening against ResNet-18, nnU-Net, and M3D across internal testing, external zero-shot transfer, and external fine-tuning (Fig. 3b,c), and multi-target screening against ResNet-18 and nnU-Net across internal testing and external fine-tuning (Fig. 3d,e). In single-target screening, HoloCT-8B achieved the highest macro F1 in all three settings: it exceeded nnU-Net by 19.4 percentage points internally (0.651 and 0.458, respectively) and, when fine-tuned as a single pooled model across external screening datasets, surpassed dedicated specialist nnU-Net models by 9.2 percentage points (0.859 compared with 0.767, $P < 0.001$; Fig. 3b,c). Across nine external lesion types evaluated without fine-tuning (three seen and six unseen), HoloCT-8B reached a macro F1 of 0.397, substantially outperforming text-prompted M3D (0.037) as well as ResNet-18 (0.050) and nnU-Net (0.036), which are restricted to predefined label sets (Fig. 3c).

To reflect clinical workflows in which multiple abnormalities are assessed simultaneously, we next evaluated multi-target screening by querying and segmenting multiple lesion types in a single pass without per-lesion prompting (Fig. 3d). HoloCT-8B raised macro F1 over nnU-Net from 0.368 to 0.718 internally and from 0.647 to 0.776 after external fine-tuning ($P < 0.001$; Fig. 3e). On detected positive scans, HoloCT's conditional Dice exceeded that of nnU-Net internally (0.679 compared with 0.337) and was lower after external fine-tuning (0.360 compared with 0.519), consistent with HoloCT returning positive masks on a larger and more difficult set of detected cases. Across the 13 lesion types shared between single- and multi-target benchmarks, querying multiple targets jointly improved overall macro F1 over separate single-target queries for both HoloCT-8B (0.740 versus 0.727) and HoloCT-4B (0.729 versus 0.716; both $P < 0.001$), although single-target queries remained higher on the externally fine-tuned subset (Supplementary Tables 9 and 10). By contrast, ResNet-18 and nnU-Net performed worse under multi-target prediction than with separate single-target models (Fig. 3f).

To test whether generating 3D masks aided lesion-presence classification, we removed the grounding request from the prompt and retained only the question (Fig. 3g). Across 18 paired internal and external zero-shot lesion types, omitting mask generation reduced mean macro F1 from 0.562 to 0.505 for HoloCT-8B and from 0.531 to 0.504 for HoloCT-4B (both $P < 0.001$; Fig. 3h). Meanwhile, conditional Dice under grounded screening remained on par with standalone segmentation (Extended Data Fig. 1a).

2.5 Visual question answering and anatomy-grounded report generation

In addition to clinical screening, we evaluated HoloCT's broader clinical reasoning on closed-ended, open-ended, and grounded VQA, together with narrative, structured, and anatomy-grounded report generation (Fig. 4). Across internal closed- and open-ended VQA benchmarks and external BIMCV-VQA [29], HoloCT

outperformed competing general-purpose and medical VLMs (Fig. 4a,e). In closed-ended VQA for CT-RATE, HoloCT-8B achieved an accuracy of 93.4% (95% confidence interval (CI), 92.9%–94.0%) compared with 84.7% (95% CI, 83.9%–85.5%) for the strongest baseline, HuluMed-32B (Supplementary Table 11). On external BIMCV-VQA, HoloCT-8B reached an LCS of 0.401, compared with 0.390 for HuluMed-32B (Supplementary Table 12). In grounded VQA across AMOS-MM-VQA[30] and DeepTumorVQA[31], generating supporting 3D masks alongside answers significantly improved mean accuracy from 75.9% to 77.1% for HoloCT-8B and from 68.0% to 76.0% for HoloCT-4B (both $P < 0.001$; Fig. 4b–d); both models localized 94.0% of reference target objects at Dice levels comparable to standalone segmentation (Extended Data Fig. 1a, Supplementary Tables 15 and 16).

Extending from targeted question answering to comprehensive scan interpretation, HoloCT also led all seven internal narrative and structured report-generation benchmarks as well as external BIMCV-RG (Fig. 4f). On RadGenome and external BIMCV-RG, HoloCT-8B achieved LCS values of 0.595 and 0.268, exceeding HuluMed-32B (0.487 and 0.256, respectively). In anatomy-grounded report generation on RadGenome, both HoloCT-4B and HoloCT-8B matched all reference thoracic regions (100%) and raised report LCS to 0.600, outperforming text-only generation (0.587 and 0.595, respectively; both $P < 0.001$) and HuluMed-32B (0.487; Fig. 4g, Supplementary Tables 17 and 18). In qualitative comparisons, pairing regional descriptions with 3D anatomical masks corrected a left-sided mislocalization of middle-lobe atelectasis and recovered a sliding hiatal hernia and mild bone degeneration missed by text-only report generation (Fig. 4h).

2.6 Referring segmentation and finding-grounded report generation

Whereas anatomy-grounded reporting links text to predefined organs, clinical radiology reports describe diverse findings in free text, ranging from focal lesions such as pulmonary nodules or masses to diffuse, non-focal abnormalities such as bilateral ground-glass opacities (Fig. 5a,b). To evaluate fine-grained localization from clinical descriptions, we first benchmarked referring segmentation on ReXGroundingCT[32], where HoloCT-8B outperformed VoxTell and SAT-Pro across focal Dice, non-focal Dice, and spatial hit rates: overall mean Dice reached 0.291 (95% CI, 0.277–0.304), compared with 0.259 (95% CI, 0.240–0.274) for VoxTell and 0.168 (95% CI, 0.152–0.185) for SAT-Pro, and non-focal Dice reached 0.252 (95% CI, 0.224–0.277), compared with 0.149 (95% CI, 0.129–0.183) for VoxTell and 0.098 (95% CI, 0.081–0.124) for SAT-Pro (Fig. 5c, Supplementary Table 19). HoloCT thus localized both focal lesions and diffuse, non-focal chest CT findings more accurately than standalone text-prompted segmentation models from free-text clinical descriptions.

To unify free-text report generation with voxel-level finding localization, we next evaluated finding-grounded report generation, in which HoloCT autonomously identified abnormal findings across the chest CT scan and generated a dedicated 3D mask for each reported finding without pre-specified target prompts (Fig. 5b). Compared with a two-stage pipeline combining HuluMed-32B for report generation with VoxTell for post-hoc finding segmentation, HoloCT-8B raised finding-detection F1 from 0.239 (95% CI, 0.213–0.278) to 0.278 (95% CI, 0.248–0.314) and mean conditional Dice from 0.186 (95% CI, 0.165–0.213) to 0.234 (95% CI, 0.209–0.258; Fig. 5d, Supplementary Table 20). Removing the grounding request lowered all six linguistic and clinical report scores for HoloCT-8B (Fig. 5e); in qualitative review, grounded generation detected and localized bilateral consolidation missed by text-only report generation (Fig. 5f).

2.7 Multicentre clinical validation and reader study

Routine hospital CT workflows span diverse anatomical sites, scanner protocols, and contrast phases, requiring models that generalize across centres without per-cohort re-engineering. To evaluate clinical applicability across real-world diagnostic settings, we tested HoloCT on 11,455 CT scans from six retrospective cohorts at five hospitals, covering lesion delineation across five targets—laryngeal lesions on contrast-enhanced CT (centre 1), lung cancer on non-contrast CT (centre 2), and benign, malignant and combined thyroid tumours on non-contrast CT (centre 3)—as well as clinical screening of histopathology-confirmed thyroid tumours (centre 3) and colon lesions (centre 4), and high-volume non-contrast chest and abdominal report generation (centre 5; Fig. 6). In zero-shot segmentation across the five clinical lesion categories at centres 1–3, HoloCT-4B and HoloCT-8B substantially outperformed M3D, SAT-Pro, and VoxTell; HoloCT-8B increased mean Dice over VoxTell from 0.107 (95% CI, 0.066–0.150) to 0.488 (95% CI, 0.441–0.531) on laryngeal lesions and from 0.543 (95% CI, 0.506–0.579) to 0.715 (95% CI, 0.687–0.741)

on lung cancer (Fig. 6a, Supplementary Table 8). Fine-tuned as a single model on the pooled training splits of all five clinical lesion categories, HoloCT-8B achieved a mean Dice of 0.696 compared with 0.681 across five dedicated specialist nnU-Net models, leading four of the five categories (Fig. 6b). Specifically, HoloCT-8B achieved a Dice of 0.684 (95% CI, 0.627–0.720) compared with 0.655 (95% CI, 0.602–0.707) on laryngeal lesions and 0.786 (95% CI, 0.762–0.809) compared with 0.776 (95% CI, 0.742–0.790) on lung cancer, while matching nnU-Net on combined thyroid tumours at 0.819 (95% CI, 0.774–0.864) compared with 0.819 (95% CI, 0.759–0.853). Averaged over the five clinical lesion categories and both evaluation settings (zero-shot versus VoxTell; fine-tuned versus nnU-Net), HoloCT-8B improved Dice by 18.2 points (Methods).

We next evaluated diagnostic performance in two clinical scenarios prone to missed findings in routine CT interpretation: thyroid tumours on non-contrast CT (centre 3), where subtle low-attenuation lesions within iodine-rich thyroid parenchyma are difficult to discern without ultrasound or intravenous contrast, and colon lesions on routine contrast-enhanced CT (centre 4), where focal wall thickening or intraluminal masses can mimic peristalsis or luminal contents. A single HoloCT model was fine-tuned on the merged training splits of both histopathology-confirmed cohorts, pairing classification with 3D tumour segmentation for thyroid scans and text-only classification for colon scans, and evaluated on held-out cases in the matching task format ($n = 141$ thyroid and $n = 166$ colon scans; Fig. 6c,d, Extended Data Fig. 1b). In blinded reader studies against two junior and two senior radiologists per cohort, HoloCT-8B surpassed every reader in sensitivity and overall accuracy: on thyroid tumours, it achieved 81.6% accuracy (radiologists, 72.3%–78.0%), with 86.2% sensitivity (95% CI, 79.1%–92.4%), 65.6% specificity (95% CI, 50.0%–82.4%), and an F1 score of 0.879 (95% CI, 0.829–0.917; Supplementary Table 9); on colon lesions, it achieved 88.6% accuracy (radiologists, 75.3%–83.1%), with 92.6% sensitivity (95% CI, 86.9%–97.1%), 83.1% specificity (95% CI, 73.7%–90.7%), and an F1 score of 0.903 (95% CI, 0.853–0.939). Compared case by case with the most accurate senior radiologist in each cohort, HoloCT-8B correctly identified 11 thyroid scans and 15 colon scans missed by the reader while misclassifying six scans per cohort that the reader judged correctly, increasing total correct diagnoses from 110 to 115 of 141 thyroid cases and from 138 to 147 of 166 colon cases (Fig. 6d).

Finally, to assess automated report drafting across high-volume routine workloads without site-specific adaptation, we evaluated zero-shot anatomy-grounded report generation across 5,307 non-contrast abdominal and 2,974 non-contrast chest CT examinations from centre 5. HoloCT-8B achieved LCS values of 0.443 (95% CI, 0.434–0.451) on abdominal CT and 0.384 (95% CI, 0.376–0.392) on chest CT, substantially outperforming HuluMed-32B at 0.236 (95% CI, 0.228–0.243) and 0.285 (95% CI, 0.275–0.296), respectively (Fig. 6e). In qualitative review, HoloCT captured subtle multi-organ abnormalities—including small pericardial effusion, reduced bone density with end-plate changes, and tiny thyroid calcifications—that HuluMed-32B omitted or misdescribed, while linking each regional statement to a verifiable 3D anatomical mask (Fig. 6f).

2.8 Bidirectional synergy between clinical understanding and spatial perception

Having demonstrated strong performance across public benchmarks and clinical cohorts, we next investigated the underlying mechanism of semantic–spatial synergy during training by comparing six configurations varying task supervision and the integration of whole-volume and fine-grained spatial features (Fig. 7a–c, Extended Data Fig. 1c). With only fine-grained tokens supplied to the language model, adding language supervision to segmentation-only training improved Dice from 0.601 to 0.613, whereas adding segmentation supervision to language-only training increased screening macro F1 from 0.256 to 0.351, alongside gains in VQA and report generation (Fig. 7c). Adding the global encoder further improved all four task families, while directly supplying both whole-volume and fine-grained tokens to the language model outperformed separate pathways and early token fusion (Fig. 7b,c).

To further determine at inference whether clinical interpretation relied directly on fine-grained 3D spatial features, we masked language-model attention to fine-grained tokens in HoloCT-8B while leaving skip connections from the detailed encoder to the mask decoder intact (Fig. 7d). Blocking this attention significantly degraded performance across all four tasks ($P < 0.001$; Fig. 7e): segmentation Dice fell from 0.840 to 0.811, screening macro F1 dropped from 0.678 to 0.061, VQA accuracy declined from 77.1% to 70.9%, and report LCS decreased from 0.507 to 0.429. Crossing spatial-token masking with retention or removal of the grounding request further showed a significant interaction ($P < 0.001$; Fig. 7f,g): masking fine-grained tokens reduced the gain from mask generation from 1.47 to 0.53 percentage points in screening F1, from 1.24 to 1.06 percentage points in VQA accuracy, and from 1.19 to 0.24 points in report LCS

(0–100 scale). These results indicate that grounded interpretation depends on the fine-grained spatial tokens shared with segmentation, although masking constitutes an out-of-distribution intervention.

3 Discussion

HoloCT integrates whole-volume and fine-grained 3D features within a unified multimodal large language model, uniting CT interpretation and voxel-level segmentation across tasks largely addressed separately by segmentation models[2–5] and medical VLMs[6–8], or connected only at the decoder output level[12–14]. Built upon the large-scale HoloDATA corpus, HoloCT outperformed specialist and foundation segmentation models, task-specific classifiers, and VLMs across 346 benchmark tasks and six clinical cohorts at five hospitals.

A central finding of this work is the bidirectional synergy between clinical understanding and 3D spatial perception in volumetric CT. Prior studies on 2D radiographs aligned report text with global and local image features[33–35], and recent medical visual tokenizers encode anatomy and semantics within unified representations[36;37]. Extending this joint representation design to volumetric CT, HoloCT supplies whole-volume global tokens and fine-grained spatial tokens directly to a unified language–segmentation model. During training, adding text-only dialogues improved 3D segmentation accuracy, whereas incorporating mask-paired dialogues improved lesion screening, VQA, and radiology report generation. At inference, this synergy operated in both directions: semantic representations enabled language-guided segmentation to generalize zero-shot to unseen anatomy, novel lesion categories, and diffuse free-text findings, while retaining the grounding request to generate 3D masks alongside text significantly improved screening, VQA, and report generation without compromising mask quality. Crucially, masking language-model attention to fine-grained spatial tokens markedly attenuated the gains from mask generation, indicating that grounded clinical reasoning draws directly on the fine-grained spatial features shared with voxel-level segmentation. Moreover, although HoloCT-8B achieved stronger overall performance, explicit grounding requests in VQA and report generation yielded larger gains for HoloCT-4B, suggesting that larger models may partially internalize spatial–semantic alignment, whereas smaller models benefit more from explicit spatial grounding during clinical interpretation.

Beyond single-task performance, unifying semantic reasoning and segmentation consolidates multi-step clinical workflows within a single model[38]. Whereas interactive and text-prompted segmentation models require users to specify each target in advance[39], HoloCT screened multiple candidate lesions in a single pass, improving overall detection accuracy over separate single-target queries while producing verifiable 3D masks. In addition, in finding-grounded report generation, HoloCT detected and localized chest CT abnormalities simultaneously, outperforming a two-stage pipeline combining HuluMed-32B and VoxTell.

In clinical practice, pairing diagnostic text with voxel-level masks directly addresses the verifiability bottleneck of medical VLMs. Although recent VLMs achieve strong aggregate scores in report generation[40], fine-grained perceptual errors and anatomical mislocalizations often remain hidden in free text[41] and are labour-intensive to audit across CT volumes spanning hundreds of slices. Whereas concept-based explanations support auditing only at the whole-image level[42;43], grounding classifications, question answers, and report statements with 3D masks allows clinicians to inspect the volumetric evidence underlying each textual output. Alongside spatial verifiability, multicentre clinical validation demonstrated strong real-world applicability: HoloCT-8B surpassed human radiologists in sensitivity and overall accuracy on histopathology-confirmed thyroid tumours and colon lesions and generalized zero-shot to high-volume chest and abdominal report generation, highlighting its potential for subtle-lesion triage and automated report drafting.

Several dataset and modelling limitations warrant future investigation. First, although HoloDATA spans six major anatomical systems for segmentation, grounded report generation remains concentrated in thoracic and abdominal CT, with finding-level 3D masks currently limited to chest CT; extending fine-grained finding annotations to abdominal, head-and-neck, neuroimaging, and musculoskeletal CT will broaden grounded reporting. Second, despite substantial gains over existing foundation models, segmentation accuracy on musculoskeletal bone lesions and diffuse, non-focal chest findings remains limited; adaptive high-resolution cropping and iterative mask refinement may further improve delineation of these challenging targets. Third, HoloCT currently interprets single CT volumes independently, whereas routine radiological workflows rely on longitudinal comparisons with prior scans, multi-phase contrast studies, and integration with electronic health records or complementary modalities such as MRI and

PET–CT; extending the unified architecture to these longitudinal and multimodal settings is an important next step.

From a clinical translation perspective, our hospital evaluation was retrospective and conducted within a single country, motivating prospective validation across international cohorts and diverse scanner protocols. Further reader-in-the-loop trials are needed to quantify how interactive 3D mask verification affects diagnostic accuracy, reading efficiency, and clinician trust in routine workflows[44;45]. Finally, safe clinical deployment will require calibrated uncertainty estimation and automated quality control alongside expert radiological oversight.

In conclusion, HoloCT demonstrates that clinical understanding and 3D spatial perception can be learned synergistically within a single generalist foundation model, offering a unified path towards more accurate, generalizable, and spatially verifiable CT interpretation.

4 Methods

4.1 Data curation

HoloDATA was assembled from 73 public datasets, including 62 segmentation datasets and 11 with language annotations, alongside six clinical cohorts from five hospitals. Dataset sources, annotations and data splits are listed in Supplementary Tables 1–5. Predefined splits were retained; for public datasets without predefined splits, 20% of scans were randomly held out for testing. For entries marked as aligned, scans shared across datasets and their associated annotations were assigned to the same split. We verified that none of the 12 external datasets, the clinical cohorts or the internal test scans were included in the training data of VISTA3D or Med3DVLM, according to their published data descriptions. The 6.2 million instruction dialogues were counted as single-turn dialogues, with each turn of a multi-turn dialogue counted separately. The 2.8 million 3D segmentation masks were counted as unique CT–mask pairs, excluding repeated uses of the same mask across tasks.

We extended the SAT label vocabulary[4] into a shared target vocabulary of 246 anatomical structures and 50 lesion categories (Supplementary Table 7). For target-level referring segmentation, prompts specified target names; for finding-level referring segmentation, prompts contained free-text findings from ReXGroundingCT[32]. Single-target and multi-target lesion screening required presence-or-absence decisions and masks for positive targets. Missing annotations were not treated as negative labels. Closed-ended VQA required binary or multiple-choice answers, whereas open-ended VQA required free-text answers. Grounded VQA paired closed-ended answers with supporting segmentation masks. Reference reports comprised findings or impressions for narrative report generation and region-specific findings for structured report generation. Each dataset's original reporting format was preserved. Anatomy-grounded report generation paired regional descriptions with anatomical masks, whereas finding-grounded report generation paired individual findings with their corresponding masks.

GPT-4o was used to generate report-question templates and to match questions from DeepTumorVQA[31] and AMOS-MM-VQA[30] with supporting masks. It was also used to clean translated BIMCV reports[29] and to generate open-ended questions from them. BIMCV image–text pairs were manually reviewed before external evaluation. Dataset-specific processing is described in Supplementary Methods, and prompts are provided in Supplementary Tables 24–29.

4.2 Model architecture

Each CT scan was converted into a global input and a detailed input. Scans were first reoriented to superior–right–anterior axes using MONAI[46]. The global input was resized to $128 \times 256 \times 256$ voxels by linear interpolation. The detailed input was cropped to the bounding box of voxels above -1000 Hounsfield units (HU), linearly resampled to $1.5 \times 1.5 \times 1.5$ mm and symmetrically padded with -1000 HU so that each dimension was divisible by 32. Intensities in both inputs were clipped to $[-1000, 1000]$ HU and scaled to $[0, 1]$.

HoloCT combined a Med3DVLM-derived global encoder[47], a VISTA3D-derived detailed encoder[3] and a Qwen3-VL language backbone[17]. HoloCT-4B and HoloCT-8B used Qwen3-VL-4B and Qwen3-VL-8B, respectively. A global projector produced 256 whole-volume global tokens from the global input, and a

detailed projector produced a fine-grained spatial token grid with one token per 32^3 voxels of the detailed input. Whole-volume tokens preceded fine-grained tokens in the language-model input, so that fine-grained tokens could attend to whole-volume tokens through causal self-attention.

Qwen3-VL's multimodal rotary position embedding has three positional components (Extended Data Fig. 2f). Each whole-volume or text token received the same sequential index for all three components. The fine-grained token at row i and column j of depth plane k received indices $(b_k, b_k + i, b_k + j)$, where b_k is the starting index of plane k. A text depth marker for each plane encoded the position of the plane centre on the training grid:

$$r_k = \left(32k + \frac{31}{2}\right) \times 1.5 \text{ mm}, \tag{1}$$

where k is the zero-based plane index, 32 is the voxel stride and 1.5 mm is the resampled spacing.

Each target description was followed by a <SEG> token, whose final hidden state was projected to a 48-dimensional target representation q_k. A mask decoder with multiscale skip connections from the detailed encoder produced shared 48-channel voxel features $F(u)$ at each voxel u. Target-specific logits and probabilities were obtained from their inner product:

$$\begin{aligned} z_k(u) &= q_k^{\mathsf{T}} F(u), \\ p_k(u) &= \text{sigmoid}(z_k(u)) \,. \end{aligned} \tag{2}$$

Mask channels followed the order of the generated <SEG> tokens. Predictions were restored to the original scan geometry and binarized at 0.5. Training augmentation, layer specifications and implementation details are provided in Supplementary Methods.

4.3 Training and ablations

Training comprised three stages: vision–language alignment (Stage 1), multitask pre-training (Stage 2) and high-quality multitask tuning (Stage 3) (Supplementary Table 6). The encoders and language backbone were initialized from their pretrained weights. Stage 1 trained only the global and detailed projectors on approximately 0.68 million report-generation samples.

Stage 2 used approximately 2.70 million language-supervised and 3.50 million segmentation-supervised samples, including single-region report-generation samples in which each region of a structured report formed a separate sample. The language backbone was trained with low-rank adaptation (LoRA) [48], and all other modules were fully trainable. Stage 3 used approximately 1.69 million language-supervised and 1.15 million segmentation-supervised samples with the same trainable modules. Stage 3 excluded single-region report-generation samples, so that all report-generation tasks used complete reports rather than individual regional descriptions. Stage 3 excluded RadGenome VQA, because of its limited question diversity and disproportionate sample size, and PASTA-Gen-TotalSeg, because of its low-quality TotalSegmentator pseudo-labels. In Stages 2 and 3, segmentation-supervised and language-supervised updates alternated.

Language supervision used next-token cross-entropy over assistant-response tokens:

$$\mathcal{L}_{\text{text}} = -\frac{1}{|\mathcal{A}|} \sum_{t \in \mathcal{A}} \log P_\theta(y_t \mid X, y_{<t}) \,, \tag{3}$$

where θ denotes the model parameters, $\mathcal{A}$ the set of supervised response positions, y_t the token at position t and X the visual and conversational context. For segmentation-supervised samples, <SEG> hidden states were obtained by teacher forcing with the reference response, and each predicted mask was supervised with binary cross-entropy (BCE) and Dice losses:

$$\mathcal{L}_{\text{BCE},k} = -\frac{1}{V_k} \sum_{u=1}^{V_k} \left[m_k(u) \log p_k(u) + (1 - m_k(u)) \log(1 - p_k(u)) \right], \tag{4}$$

$$\mathcal{L}_{\text{Dice},k} = 1 - \frac{2 \sum_u p_k(u) m_k(u) + \epsilon_{\text{n}}}{\sum_u p_k(u)^2 + \sum_u m_k(u)^2 + \epsilon_{\text{d}}}, \tag{5}$$

where $p_k(u)$ is the predicted foreground probability of target k at voxel u, $m_k(u)$ is the binary reference label and V_k is the number of voxels. Both smoothing constants, ϵ_{n} and ϵ_{d}, were 10^{-5}. The segmentation

loss was averaged over the K supervised masks in each minibatch:

$$\mathcal{L}_{\text{seg}} = \frac{1}{K}\sum_{k=1}^{K}\left(0.3\mathcal{L}_{\text{BCE},k} + 0.7\mathcal{L}_{\text{Dice},k}\right). \tag{6}$$

The total loss was $\mathcal{L}_{\text{text}} + \mathcal{L}_{\text{seg}}$ for segmentation-supervised samples and $\mathcal{L}_{\text{text}}$ otherwise.

Both model sizes were trained with an effective batch size of 64 in bfloat16 precision. AdamW[49] used learning rates of 5×10^{-4} in Stage 1 and 5×10^{-5} in Stages 2 and 3 with cosine decay, $(\beta_1, \beta_2) = (0.9, 0.999)$, $\epsilon = 10^{-8}$, zero weight decay and gradient clipping at a norm of 1.0. LoRA used rank $r = 128$, scaling factor $\alpha = 256$ and dropout 0.05.

Downstream fine-tuning started from the Stage 3 checkpoints, ran for five epochs and otherwise used Stage 3 settings. For external lesion screening, each HoloCT model was fine-tuned once on the pooled training data for single-target and multi-target lesion screening. For clinical segmentation, training data for all five targets from centres 1–3 were pooled. For clinical screening, thyroid and colon training cases from centres 3 and 4 were pooled; thyroid cases received screening and segmentation supervision, and colon cases received screening supervision only. Evaluation prompts requested the outputs of each cohort's training task.

All ablations used HoloCT-8B. Training ablations S1–S6 were initialized from the same pretrained weights as Stage 1 and trained for five epochs with Stage 3 settings on a random 10% of Stage 3 data plus the training splits of the ablation benchmarks. S1–S3 supplied only fine-grained tokens to the language model and differed in supervision: S1 used language supervision, S2 used segmentation supervision with text and mask losses, and S3 combined both. S4–S6 combined both forms of supervision and differed in how whole-volume and fine-grained tokens were integrated. S4 supplied only whole-volume tokens to the language model and used fine-grained features solely for mask decoding. S5 fused projected whole-volume and fine-grained tokens before the language model using two-way attention inspired by the Segment Anything Model (SAM)[50], supplying the updated whole-volume tokens to the language model and passing the updated fine-grained tokens to the mask decoder. S6 supplied both token sets directly to the language model and corresponded to the HoloCT architecture. Inference ablations evaluated a single HoloCT-8B checkpoint on the training-ablation benchmarks in a 2×2 design: fine-grained tokens were either visible or masked in causal attention, and grounding requests were either retained in or removed from the prompts. Masking preserved the position indices of fine-grained tokens, and the mask decoder still received fine-grained features. Initialization of the remaining modules, ablation architectures and benchmarks are detailed in Supplementary Methods, and results are shown in Fig. 7.

4.4 Compared methods

Segmentation baselines were nnU-Net v2, VISTA3D, SAT-Pro, VoxTell and M3D-Phi3-4B (hereafter M3D)[2–5;12], and lesion-screening baselines additionally included 3D ResNet-18[16]. Text-generation baselines were M3D, HuluMed-32B, RadFM-14B, MedGemma 1.5-4B and Qwen3-VL-32B[8;17–19]. Released weights were used unless stated otherwise.

We trained nnU-Net from scratch on each dataset. For single-target lesion screening, we trained a separate ResNet-18 for each dataset and lesion category; for multi-target lesion screening, ResNet-18 used a separate binary head for each target. We fine-tuned M3D on pooled lesion-screening training data and HuluMed-32B only on ReXGroundingCT training data, using LoRA for the language backbone with all other modules fully trainable. We fully fine-tuned SAT-Pro on ReXGroundingCT for finding-level referring segmentation. Training configurations are detailed in Supplementary Methods.

The finding-grounded report generation baseline was a pipeline in which released VoxTell segmented each finding generated by fine-tuned HuluMed-32B. VoxTell's training annotations overlap with the ReXGroundingCT test set, so its standalone and pipeline results are affected by label leakage.

4.5 Benchmark evaluation

Public benchmarks were used to assess performance on internal test splits, zero-shot performance on external datasets and performance after task-specific fine-tuning. Test scans were excluded from training and checkpoint selection.

Target-level referring segmentation was evaluated using Dice and the 95th-percentile Hausdorff distance (HD95, in millimetres)[20]:

$$D(P,G) = \frac{2|P \cap G|}{|P| + |G|}, \tag{7}$$

$$\mathrm{HD95} = \max\left\{Q^A_{0.95}(\mathcal{D}_{P,G}), Q^A_{0.95}(\mathcal{D}_{G,P})\right\}, \tag{8}$$

where P and G are the predicted and reference foreground voxel sets, $\mathcal{D}_{P,G}$ and $\mathcal{D}_{G,P}$ contain directed nearest-surface distances in physical space, and $Q^A_{0.95}$ is the surface-area-weighted 95th percentile.

Lesion screening was evaluated from counts of true positives (TP), true negatives (TN), false positives (FP) and false negatives (FN):

$$\begin{aligned} \mathrm{Accuracy} &= \frac{\mathrm{TP}+\mathrm{TN}}{\mathrm{TP}+\mathrm{TN}+\mathrm{FP}+\mathrm{FN}}, \\ \mathrm{F1} &= \frac{2\mathrm{TP}}{2\mathrm{TP}+\mathrm{FP}+\mathrm{FN}}, \\ \mathrm{Sensitivity} &= \frac{\mathrm{TP}}{\mathrm{TP}+\mathrm{FN}}, \\ \mathrm{Specificity} &= \frac{\mathrm{TN}}{\mathrm{TN}+\mathrm{FP}}. \end{aligned} \tag{9}$$

Responses from HoloCT and generative baselines were classified as positive or negative by case-insensitive string matching, ignoring surrounding whitespace. When prompts included grounding requests, a positive prediction additionally required a nonempty target mask. An nnU-Net prediction was positive whenever its target mask contained foreground, irrespective of overlap with the reference. Conditional Dice was the mean Dice score over true-positive CT–target pairs $\mathcal{T}$:

$$D_{\mathrm{cond}} = \frac{1}{|\mathcal{T}|}\sum_{i\in\mathcal{T}} D(P_i, G_i). \tag{10}$$

Closed-ended VQA answers (yes or no, or option labels A–D) were parsed using the same string-matching rule, and invalid or incorrect answers counted as errors. DeepTumorVQA and AMOS-MM-VQA accuracies were also reported by question category (Supplementary Table 11). For grounded VQA, predicted and reference objects were matched by category; for anatomy-grounded report generation, regions were matched by anatomical name (Supplementary Tables 16 and 18). For both tasks, match rate was the number of matched references divided by the total number of references, pooled across scans and categories or regions. Dice and HD95 were computed for matched objects or regions only.

Open-ended VQA and report generation were evaluated using BLEU-4[21], ROUGE-L[22], METEOR[23] and LlamaScore, an integer clinical-agreement score from 0 to 10 assigned by Llama-3.1-70B-Instruct[24] as a large language model (LLM) judge. Report generation was additionally evaluated using RaTEScore[25] and GREEN[26]. We defined the Language–Clinical Score (LCS) as the equally weighted mean of linguistic similarity N and clinical agreement C, each on a 0–1 scale:

$$\begin{aligned} N &= \frac{\mathrm{BLEU\text{-}4}+\mathrm{ROUGE\text{-}L}+\mathrm{METEOR}}{3}, \\ C_{\mathrm{VQA}} &= \mathrm{LlamaScore}/10, \\ C_{\mathrm{RG}} &= \frac{\mathrm{RaTEScore}+\mathrm{LlamaScore}/10+\mathrm{GREEN}}{3}, \end{aligned} \tag{11}$$

$$\mathrm{LCS}_{\mathrm{VQA}} = \frac{N + C_{\mathrm{VQA}}}{2}, \qquad \mathrm{LCS}_{\mathrm{RG}} = \frac{N + C_{\mathrm{RG}}}{2}. \tag{12}$$

For finding-level referring segmentation, each ReXGroundingCT[32] finding was supplied as a prompt, and the predicted mask was compared with its reference mask using Dice and hit rates. For M findings with Dice scores D_j, hit rates were calculated as

$$\mathrm{HIT}_\tau = \frac{1}{M}\sum_{j=1}^{M} \mathbf{1}[D_j > \tau], \qquad \tau \in \{0.05, 0.10\}, \tag{13}$$

and denoted $\mathrm{HIT}_{5\%}$ and $\mathrm{HIT}_{10\%}$, respectively. For finding-grounded report generation, the LLM judge assessed whether each generated finding matched a reference finding from the same scan, requiring agreement in pathology, morphology and location without contradictory clinical attributes. Each reference finding could be matched to at most one prediction, and only such matches counted as TP; when several predictions matched one reference, the prediction with the highest Dice was retained. Finding detection was evaluated using F1, with

$$\mathrm{FP} = n_{\mathrm{pred}} - \mathrm{TP}, \qquad \mathrm{FN} = n_{\mathrm{ref}} - \mathrm{TP}, \tag{14}$$

where n_{pred} and n_{ref} are the numbers of predicted and reference findings. Conditional Dice and hit rates were computed over true-positive findings. For both finding-level tasks, localization metrics were reported overall and separately for focal and non-focal findings. Evaluation implementation is detailed in Supplementary Methods.

4.6 Clinical evaluation

Clinical evaluation used six retrospective cohorts from five centres, with one CT scan per patient. The research ethics committees of all five participating centres (Zhongshan Hospital, Fudan University; the First Affiliated Hospital, Zhejiang University School of Medicine; Sir Run Run Shaw Hospital, Zhejiang University School of Medicine; the Affiliated Yangming Hospital of Ningbo University; and Guizhou Provincial People's Hospital) approved the study and waived informed consent; images and records were de-identified. Sample sizes were determined by the number of eligible cases, without an a priori power calculation. Scans from centres 1 and 4 were contrast-enhanced, and the remaining scans were non-contrast. Centres 1 and 2 included only lesion-positive cases, whereas centres 3 and 4 included both positive and negative cases. Four physicians delineated lesion masks for centres 1–3 using clinical and pathological findings. Histopathology provided reference diagnoses for thyroid (centre 3) and colon (centre 4) lesion screening; colon cases had no reference masks. The cohort design is shown in Fig. 1a.

Lesion segmentation was evaluated at centres 1–3, and lesion screening at centres 3 and 4. Each cohort was split at the patient level into 80% training and 20% testing. Models were evaluated zero-shot and after fine-tuning on the training split, using the same test patients. Two junior radiologists (3 and 5 years of experience) and two senior radiologists (13 and 17 years of experience) independently reviewed the original CT volumes of the test patients in shuffled order, without access to reference diagnoses or model outputs. Screening performance of models and readers was measured by F1, sensitivity, specificity and accuracy. Diagnostic correctness of each model was compared with that of the highest-accuracy senior reader in each cohort. Reader comparisons are shown in Fig. 6c,d.

Abdominal and chest reporting cohorts from centre 5 were reserved for external zero-shot evaluation. GPT-4o cleaned the Chinese reports, translated them into English and organized them by the Merlin abdominal and RadGenome thoracic region schemes. HoloCT performed anatomy-grounded report generation, and its report LCS was compared with that of HuluMed-32B. Report processing is detailed in Supplementary Table 29.

4.7 Statistical analysis

Segmentation scores were averaged across cases within each dataset and target, then with equal weight across datasets for each target:

$$\bar{s}_\ell = \frac{1}{|\mathcal{D}_\ell|} \sum_{d \in \mathcal{D}_\ell} \left(\frac{1}{n_{d\ell}} \sum_{i=1}^{n_{d\ell}} s_{id\ell} \right). \tag{15}$$

Here, $s_{id\ell}$ is the score for case i, dataset d and target ℓ; $n_{d\ell}$ is the number of evaluated cases, and $\mathcal{D}_\ell$ is the set of datasets containing target ℓ. Category and overall summaries weighted target means equally. Lesion-screening F1 was computed from counts pooled within each dataset and target, followed by the same hierarchical averaging. The external zero-shot screening comparison used the same nine targets for every model, assigning a score of zero to the six targets not supported by nnU-Net and 3D ResNet-18. VQA scores were averaged over questions and report LCS over CT scans; finding-level F1 was computed from counts pooled across scans. Figure-specific aggregations and error-bar definitions are provided in Supplementary Methods.

Confidence intervals were estimated from 1,000 percentile bootstrap resamples of CT scans within datasets (seed 42), keeping all observations from each scan together. Paired conditions used identical resamples, and all aggregation steps were repeated for each resample. The 2.5th and 97.5th percentiles defined 95% confidence intervals.

All hypothesis tests were two-sided, and P values were not adjusted for multiple comparisons. Segmentation comparisons used paired t-tests of target-level scores $\bar{s}_\ell$. Screening, VQA and report comparisons, including model–reader comparisons, used paired permutation tests with 10,000 permutations, in which the two conditions' complete predictions for each CT scan were exchanged with probability 0.5 before the metric was recomputed. Score drops after grounding-request removal were compared between intact and masked attention using paired t-tests, with pairing by target for screening and by CT scan for VQA and report generation.

Abstract summary values for HoloCT-8B were computed as follows. (1) The 18.2-point clinical segmentation improvement is the macro average over the five clinical lesion categories of, per category, the equally weighted mean of two Dice differences: zero-shot HoloCT-8B minus zero-shot VoxTell, and fine-tuned HoloCT-8B minus specialist nnU-Net. (2) The 15.3-point report-generation improvement is the equally weighted mean, over the abdominal and chest cohorts of centre 5, of the LCS difference between grounded HoloCT-8B and HuluMed-32B. (3) The 85.1% screening accuracy is the equally weighted mean of per-cohort accuracies (115/141 thyroid; 147/166 colon). (4) The 22.1-point open-vocabulary margin is the difference in macro Dice between HoloCT-8B and VoxTell over 16 equally weighted items: 10 external unseen targets (PANORAMA targets were excluded because they were used in VoxTell training), the five clinical categories evaluated zero-shot, and overall finding-level Dice on ReXGroundingCT.

4.8 Computational hardware and software

All models were trained on NVIDIA H100 GPUs. HoloCT-4B and HoloCT-8B each used 16 GPUs across two nodes. nnU-Net and 3D ResNet-18 each used two GPUs, whereas M3D, HuluMed-32B and SAT-Pro each used eight. HoloCT training used gradient checkpointing and DeepSpeed ZeRO-3, with optimizer states offloaded to CPU memory[51]. Training required approximately 3,500 GPU hours for HoloCT-4B and 4,500 GPU hours for HoloCT-8B.

The HoloCT software environment used Python 3.10.16, PyTorch 2.6.0 and CUDA runtime 12.4.127. Core libraries included Transformers 4.57.0, MONAI 1.5.0, DeepSpeed 0.17.5, PEFT 0.17.1 and Accelerate 1.10.0. Image processing and evaluation used NiBabel 5.2.1, SimpleITK 2.5.2, NumPy 1.26.4, SciPy 1.14.0, scikit-learn 1.7.1 and surface-distance 0.1.

Acknowledgements

We thank the researchers and institutions that made the public CT datasets used in this study available. This work was supported by Alibaba DAMO Academy.

References

1. Smith-Bindman, R. et al. Trends in use of medical imaging in US health care systems and in Ontario, Canada, 2000-2016. *JAMA* **322**, 843–856 (2019).

2. Isensee, F., Jaeger, P. F., Kohl, S. A. A., Petersen, J. & Maier-Hein, K. H. nnU-Net: a self-configuring method for deep learning-based biomedical image segmentation. *Nat. Methods* **18**, 203–211 (2021).

3. He, Y. et al. VISTA3D: a unified segmentation foundation model for 3D medical imaging. In *Proc. IEEE/CVF Conference on Computer Vision and Pattern Recognition*, pp. 20863–20873 (IEEE, 2025).

4. Zhao, Z. et al. Large-vocabulary segmentation for medical images with text prompts. *npj Digit. Med.* **8**, 566 (2025).

5. Rokuss, M. et al. VoxTell: free-text promptable universal 3D medical image segmentation. In *Proc. IEEE/CVF Conference on Computer Vision and Pattern Recognition*, pp. 37538–37557 (IEEE, 2026).

6. Blankemeier, L. et al. Merlin: a computed tomography vision–language foundation model and dataset. *Nature* **652**, 1318–1328 (2026).

7. Hamamci, I. E. et al. Generalist foundation models from a multimodal dataset for 3D computed tomography. *Nat. Biomed. Eng.* **10**, 1610–1628 (2026).

8. Wu, C., Zhang, X., Zhang, Y., Hui, H., Wang, Y. & Xie, W. Towards generalist foundation model for radiology by leveraging web-scale 2D&3D medical data. *Nat. Commun.* **16**, 7866 (2025).

9. Zhao, T. et al. A foundation model for joint segmentation, detection and recognition of biomedical objects across nine modalities. *Nat. Methods* **22**, 166–176 (2025).

10. Zhou, H.-Y., Acosta, J. N., Adithan, S., Datta, S., Topol, E. J. & Rajpurkar, P. MedVersa: a generalist foundation model for diverse medical imaging tasks. *NEJM AI* **3**, AIoa2500595 (2026).

11. Shen, L. et al. From image to pixels: towards fine-grained medical vision-language models. *IEEE Trans. Pattern Anal. Mach. Intell.* **48**, 9982–9996 (2026).

12. Bai, F., Du, Y., Huang, T., Meng, M. Q.-H. & Zhao, B. M3D: advancing 3D medical image analysis with multi-modal large language models. Preprint at https://doi.org/10.48550/arXiv.2404.00578 (2024).

13. Luo, L., Tang, B., Chen, X., Han, R. & Chen, T. VividMed: vision language model with versatile visual grounding for medicine. In *Proc. 2025 Conference of the Nations of the Americas Chapter of the Association for Computational Linguistics: Human Language Technologies (Volume 1: Long Papers)*, pp. 1800–1821 (ACL, 2025).

14. Wu, L. et al. A universal foundation model for grounded biomedical image interpretation. *Nat. Commun.* **17**, 7173 (2026).

15. Varma, M. et al. MedVAE: efficient automated interpretation of medical images with large-scale generalizable autoencoders. In *Proc. International Conference on Medical Imaging with Deep Learning*, vol. 301, pp. 1597–1626 (PMLR, 2026).

16. He, K., Zhang, X., Ren, S. & Sun, J. Deep residual learning for image recognition. In *Proc. IEEE Conference on Computer Vision and Pattern Recognition*, pp. 770–778 (IEEE, 2016).

17. Bai, S. et al. Qwen3-VL technical report. Preprint at https://doi.org/10.48550/arXiv.2511.21631 (2025).

18. Sellergren, A. et al. MedGemma 1.5 technical report. Preprint at https://doi.org/10.48550/arXiv.2604.05081 (2026).

19. Jiang, S. et al. Hulu-Med: a transparent generalist model towards holistic medical vision-language understanding. Preprint at https://doi.org/10.48550/arXiv.2510.08668 (2025).

20. Taha, A. A. & Hanbury, A. Metrics for evaluating 3D medical image segmentation: analysis, selection, and tool. *BMC Med. Imaging* **15**, 29 (2015).

21. Papineni, K., Roukos, S., Ward, T. & Zhu, W.-J. BLEU: a method for automatic evaluation of machine translation. In *Proc. 40th Annual Meeting of the Association for Computational Linguistics*, pp. 311–318 (ACL, 2002).

22. Lin, C.-Y. ROUGE: a package for automatic evaluation of summaries. In *Proc. Text Summarization Branches Out*, pp. 74–81 (ACL, 2004).

23. Banerjee, S. & Lavie, A. METEOR: an automatic metric for MT evaluation with improved correlation with human judgments. In *Proc. ACL Workshop on Intrinsic and Extrinsic Evaluation Measures for Machine Translation and/or Summarization*, pp. 65–72 (ACL, 2005).

24. Grattafiori, A. et al. The Llama 3 herd of models. Preprint at https://doi.org/10.48550/arXiv.2407.21783 (2024).

25. Zhao, W., Wu, C., Zhang, X., Zhang, Y., Wang, Y. & Xie, W. RaTEScore: a metric for radiology report generation. In *Proc. 2024 Conference on Empirical Methods in Natural Language Processing*, pp. 15004–15019 (ACL, 2024).

26. Ostmeier, S. et al. GREEN: generative radiology report evaluation and error notation. In *Findings of the Association for Computational Linguistics: EMNLP 2024*, pp. 374–390 (ACL, 2024).

27. Yu, F. et al. Evaluating progress in automatic chest X-ray radiology report generation. *Patterns* **4**, 100802 (2023).

28. Li, C.-Y. et al. Towards a holistic framework for multimodal LLM in 3D brain CT radiology report generation. *Nat. Commun.* **16**, 2258 (2025).

29. de la Iglesia Vayá, M. et al. BIMCV COVID-19+: a large annotated dataset of RX and CT images from COVID-19 patients. Preprint at https://doi.org/10.48550/arXiv.2006.01174 (2020).

30. Ji, Y., Ge, C., Li, R. & Luo, P. AMOS-MM: abdominal multimodal analysis challenge: structured description of the challenge design. Zenodo https://doi.org/10.5281/zenodo.10992155 (2024).

31. Chen, Y. et al. Are vision language models ready for clinical diagnosis? A 3D medical benchmark for tumor-centric visual question answering. Preprint at https://doi.org/10.48550/arXiv.2505.18915 (2025).

32. Baharoon, M. et al. State-of-the-art text-prompted medical segmentation models struggle to ground chest CT findings. In *Proc. Machine Learning for Healthcare Conference*, vol. 298, pp. 1–30 (PMLR, 2025).

33. Huang, S.-C., Shen, L., Lungren, M. P. & Yeung, S. GLoRIA: a multimodal global-local representation learning framework for label-efficient medical image recognition. In *Proc. IEEE/CVF International Conference on Computer Vision*, pp. 3942–3951 (IEEE, 2021).

34. Tiu, E., Talius, E., Patel, P., Langlotz, C. P., Ng, A. Y. & Rajpurkar, P. Expert-level detection of pathologies from unannotated chest X-ray images via self-supervised learning. *Nat. Biomed. Eng.* **6**, 1399–1406 (2022).

35. Zhang, X., Wu, C., Zhang, Y., Xie, W. & Wang, Y. Knowledge-enhanced visual-language pre-training on chest radiology images. *Nat. Commun.* **14**, 4542 (2023).

36. Ma, C. et al. Unified medical image tokenizer for autoregressive synthesis and understanding. Preprint at https://doi.org/10.48550/arXiv.2505.19225 (2025).

37. Ma, C. et al. MedVTok: a general-purpose medical visual tokenizer. In *Proc. Advances in Neural Information Processing Systems* (2026). In press.

38. Moor, M. et al. Foundation models for generalist medical artificial intelligence. *Nature* **616**, 259–265 (2023).

39. Ma, J., He, Y., Li, F., Han, L., You, C. & Wang, B. Segment anything in medical images. *Nat. Commun.* **15**, 654 (2024).

40. Zambrano Chaves, J. M. et al. A clinically accessible small multimodal radiology model and evaluation metric for chest X-ray findings. *Nat. Commun.* **16**, 3108 (2025).

41. D'Cunha, R. et al. MMBU: a massive multi-modal biomedical understanding benchmark to probe the perception capabilities of vision-language models. In *Proc. European Conference on Computer Vision*, pp. 588–604 (ECCV, 2026).

42. Kim, C. et al. Transparent medical image AI via an image–text foundation model grounded in medical literature. *Nat. Med.* **30**, 1154–1165 (2024).

43. Nie, Y. et al. An explainable biomedical foundation model via large-scale concept-enhanced vision–language pretraining. *Nat. Biomed. Eng.* https://doi.org/10.1038/s41551-026-01764-x (2026).

44. Tanno, R. et al. Collaboration between clinicians and vision–language models in radiology report generation. *Nat. Med.* **31**, 599–608 (2025).

45. Yu, F., Moehring, A., Banerjee, O., Salz, T., Agarwal, N. & Rajpurkar, P. Heterogeneity and predictors of the effects of AI assistance on radiologists. *Nat. Med.* **30**, 837–849 (2024).

46. Cardoso, M. J. et al. MONAI: an open-source framework for deep learning in healthcare. Preprint at https://doi.org/10.48550/arXiv.2211.02701 (2022).

47. Xin, Y., Ates, G. C., Gong, K. & Shao, W. Med3DVLM: an efficient vision-language model for 3D medical image analysis. *IEEE J. Biomed. Health Inform.* **30**, 2524–2536 (2026).

48. Hu, E. J. et al. LoRA: low-rank adaptation of large language models. In *Proc. International Conference on Learning Representations* (ICLR, 2022).

49. Loshchilov, I. & Hutter, F. Decoupled weight decay regularization. In *Proc. International Conference on Learning Representations* (ICLR, 2019).

50. Kirillov, A. et al. Segment anything. In *Proc. IEEE/CVF International Conference on Computer Vision*, pp. 4015–4026 (IEEE, 2023).

51. Rajbhandari, S., Rasley, J., Ruwase, O. & He, Y. ZeRO: memory optimizations toward training trillion parameter models. In *Proc. International Conference for High Performance Computing, Networking, Storage and Analysis*, pp. 1–16 (IEEE, 2020).



Figure 1. HoloDATA and HoloCT for unified CT interpretation and segmentation. **a**, Public datasets and multicentre cohorts comprising 195,921 CT scans, grouped by source and contrast status. **b**, Language and segmentation annotations across datasets and anatomical systems. **c**, Dual-scale encoding combines global context and high-resolution spatial tokens in a multimodal language model for text generation and 3D mask decoding. **d**, Representative dialogues and masks for simultaneous multi-target lesion screening, grounded visual question answering (VQA) and finding-level grounded report generation.



Figure 2. Language-guided segmentation across internal and external CT datasets. **a**, Internal comparison with VISTA3D across 128 shared targets. **b**, Comparison with specialist nnU-Net models across 272 internal and 37 seen external targets. **c,e**, Internal reference examples alongside external-seen and external-unseen targets, respectively. **d,f**, External zero-shot Dice by anatomical and lesion group for 37 seen and 14 unseen targets, respectively. In **a,b**, bars show means across targets; whiskers span target minima–maxima. Brackets indicate unadjusted $P < 0.001$ from two-sided paired t-tests across targets. Radar axes in **d,f** are normalized to HoloCT-8B, with native Dice values on labelled rings.



Figure 3. Single-target and multi-target lesion screening. **a**, Screening workflow with lesion classification and 3D masks. **b**,**d**, Single- and multi-target examples, respectively. **c**,**e**, Macro F1 and conditional Dice on true-positive cases for single- and multi-target screening, respectively. **f**, Single- versus multi-target macro F1 across 13 matched public lesions. **g**, Prompts with and without mask requests. **h**, Grounding comparison averaged across 18 paired internal and external zero-shot lesions. In **c**,**e**,**f**, bars show lesion means, circles denote lesions, and whiskers span minima–maxima. Filled and open bars in **h** show means with and without grounding. Brackets indicate unadjusted $P < 0.001$ from two-sided scan-level paired permutation tests.



Figure 4. Visual question answering (VQA) and anatomy-grounded report generation. **a**, Closed-ended VQA accuracy. **b**, VQA prompts with and without mask requests. **c**, Mean overall accuracy across DeepTumorVQA and AMOS-MM-VQA. **d**, Question–answer examples with supporting masks. **e**,**f**, Open-ended VQA and report-generation Language–Clinical Score (LCS), respectively, including external BIMCV. **g**, RadGenome report LCS (%) with and without grounding versus HuluMed-32B. **h**, Example of anatomy-grounded report generation; green and red text indicate agreement and discrepancy with the reference. In **a**,**e**,**f**, radar axes are normalized to HoloCT-8B, with native scores on labelled rings. In **c**,**g**, filled and open bars show means with and without grounding; brackets indicate unadjusted $P < 0.001$ from two-sided scan-level paired permutation tests.



Figure 5. Referring segmentation and finding-grounded report generation. **a**,**b**, Task illustrations for finding-level referring segmentation and finding-grounded report generation, respectively. **c**, ReXGroundingCT referring-segmentation Dice and spatial hit rates; $HIT_{5\%}$ and $HIT_{10\%}$ denote the fractions of findings with Dice above 0.05 and 0.10. **d**, Finding-detection F1 and conditional localization compared with a two-stage pipeline pairing HuluMed-32B with VoxTell. **e**, Report-quality metrics with and without 3D mask requests; heatmap values show raw means and colours indicate within-metric standardized scores. **f**, Representative grounded and text-only reports; green and red text indicate agreement and discrepancy with the reference.



Figure 6. Multicentre clinical evaluation of segmentation, screening, and report generation. **a**,**b**, Zero-shot and fine-tuned segmentation across five clinical tumour categories at centres 1–3; bars show mean Dice with scan-level bootstrap 95% confidence intervals, and red outlines and yellow overlays indicate reference and predicted masks. **c**, Sensitivity and specificity of fine-tuned models (squares) and two junior and two senior radiologists (triangles) on held-out thyroid ($n = 141$) and colon ($n = 166$) scans. **d**, Case-level classification comparison between HoloCT-8B and the most accurate senior radiologist on the same cohorts; green and red indicate correct and incorrect diagnoses, respectively. **e**, Zero-shot report Language–Clinical Score (LCS, %) across 5,307 abdominal and 2,974 chest CT scans. Filled and open bars denote inference with and without grounding; whiskers show scan-level bootstrap 95% confidence intervals. **f**, Report excerpts and anatomical masks compared with HuluMed-32B and reference reports.

Figure 7. Training, dual-scale fusion, and spatial-attention ablations. **a**, Three-stage training curriculum and the 10% Stage 3 subset combined with the complete training splits of the evaluation datasets for S1–S6. **b**, Ablation settings S1–S6; S5 exchanges projected tokens prior to the language model, and S6 feeds both visual streams directly into it. **c**, Task scores (%), with colours indicating within-task standardized scores; segmentation is not applicable (N/A) to the text-only configuration S1. **d**, Masking language-model attention to high-resolution spatial tokens with skip connections from the encoder to the mask decoder kept intact. **e**, Performance under intact or masked spatial attention with mask requests retained (filled bars) or removed (open bars). **f**, Design for removing the 3D mask request. **g**, Score reductions (in percentage points) upon removing the mask request under intact versus masked spatial attention. In **e**,**g**, bars show means across the two evaluation datasets per task and whiskers span the two dataset-level values. Brackets indicate unadjusted $P < 0.001$: two-sided paired t-tests for segmentation in **e** and score reductions in **g**, and scan-level paired permutation tests for other tasks in **e**.





Extended Data Fig. 1. Segmentation quality, clinical screening, and ablation configurations. **a**, Grounded versus segmentation-only requests: conditional Dice across 15 screening lesions and Dice across 12 AMOS-MM-VQA and 20 DeepTumorVQA categories. **b**, Fine-tuned single- versus multi-target thyroid screening ($n = 141$ scans), with benign and malignant tumours shown separately. In **a**,**b**, bars show category means, circles denote categories, and whiskers span minima–maxima. **c**, S1–S3 share a high-resolution encoder across different training tasks; S4 separates global language and high-resolution mask pathways; S5 adds bidirectional feature exchange after projection; S6 supplies both projected visual streams to the language model. High-resolution modules are labelled 'Detailed' in **c**.



Token	D	H	W
g0	0	0	0
g1	1	1	1
g2	2	2	2
g3	3	3	3
<23.25mm>	4 … b0−1	4 … b0−1	4 … b0−1
v0	b0	b0	b0
v1	b0	b0	b0+1
v2	b0	b0+1	b0
v3	b0	b0+1	b0+1
<71.25mm>	b0+2 … b1−1	b0+2 … b1−1	b0+2 … b1−1
v4	b1	b1	b1
v5	b1	b1	b1+1
v6	b1	b1+1	b1
v7	b1	b1+1	b1+1
T0	b1+2	b1+2	b1+2
T1	b1+3	b1+3	b1+3

Extended Data Fig. 2. HoloCT encoder, projection, and position-encoding components. **a**, DCFormer global encoder. **b**, Global projector with parallel Mixer streams. **c**, VISTA3D high-resolution encoder and mask decoder with multiscale skip connections. **d**, High-resolution projector with depth pooling, feature projection, patch grouping, and visual merging. **e**, Bidirectional attention after visual projection in ablation configuration S5. **f**, Three-axis position-index example for global, high-resolution, and language tokens. High-resolution modules are labelled ‘Detailed’ in **c**–**e**.

Supplementary Information

Table of contents

Supplementary Methods **26**

Supplementary Table 1 | Dataset details for segmentation 30

Supplementary Table 2 | Dataset details for grounded lesion screening 33

Supplementary Table 3 | Dataset details for visual question answering 34

Supplementary Table 4 | Dataset details for report generation 34

Supplementary Table 5 | Dataset details for grounded closed-ended visual question answering and grounded report generation 35

Supplementary Table 6 | Dataset composition across training stages 36

Supplementary Table 7 | Segmentation label hierarchy and counts 37

Supplementary Table 8 | Detailed Dice and HD95 scores across segmentation benchmarks 38

Supplementary Table 9 | Single-target grounded lesion screening 77

Supplementary Table 10 | Multiple-target grounded lesion screening 82

Supplementary Table 11 | Detailed closed-ended VQA scores across benchmarks 85

Supplementary Table 12 | Detailed open-ended VQA scores across benchmarks 86

Supplementary Table 13 | Detailed narrative report generation scores across benchmarks 87

Supplementary Table 14 | Detailed structured report generation scores across benchmarks 88

Supplementary Table 15 | Effect of grounding on AMOS-MM-VQA and DeepTumorVQA 89

Supplementary Table 16 | Segmentation performance on AMOS-MM-VQA and DeepTumorVQA . 90

Supplementary Table 17 | Anatomy-grounded and finding-grounded report generation: text quality 92

Supplementary Table 18 | RadGenome segmentation performance for anatomy-grounded report generation 93

Supplementary Table 19 | Finding-level referring segmentation 94

Supplementary Table 20 | Finding-grounded report generation 95

Supplementary Table 21 | Zero-shot conditional Dice across classification targets 96

Supplementary Table 22 | Training-strategy ablations S1–S6 across tasks 97

Supplementary Table 23 | Masking and grounding-request ablations across tasks 98

Supplementary Table 24 | Prompts for report cleaning, question construction and evidence-mask selection. 103

Supplementary Table 25 | Segmentation, lesion-screening and finding-grounded report generation templates. 105

Supplementary Table 26 | Question-answering and VQA-grounding templates. 106

Supplementary Table 27 | Dataset-specific report templates for five source datasets. 109

Supplementary Table 28 | INSPECT, BIMCV and anatomy-grounded report generation templates. 111

Supplementary Table 29 | Clinical report cleaning, translation and anatomical structuring. 114

Supplementary Methods

Data processing and task construction

For training, each multi-turn dialogue was used as a single sample. Only the CT-RATE[1] long- and short-answer data were organized as multi-turn dialogues; all other instruction data were single-turn dialogues. Stage-specific inventories count training samples, including samples reused across stages, and task-specific mask inventories include repeated uses of shared masks. Target-level referring segmentation used positive cases only, whereas lesion screening used explicitly annotated positive and negative cases. Dataset and stage inventories are listed in Supplementary Tables 1–6.

Report-question templates were derived from each source dataset's description and 50 sample reports. Grounding requests were added to the prompts for grounded tasks. For anatomy-grounded report generation, thoracic reports were organized into ten RadGenome[2] regions, and abdominal reports were organized according to the Merlin[3] abdominal region scheme. Task prompts are listed in Supplementary Tables 25–28, and data-construction prompts in Supplementary Table 24.

CT-RATE-Option preprocessing. Image markers and task-specific tags were removed. We excluded questions without recognizable answer options and responses that did not contain exactly one answer label (A–D). The remaining questions, options and answers were converted into a standardized format.

Grounded VQA construction. For DeepTumorVQA[4] and AMOS-MM-VQA[5], masks were selected according to anatomical specificity, laterality and lesion relevance. Questions about organ measurements required organ masks, and questions about lesion presence, count, size or volume required lesion masks rather than parent-organ masks. Multiple masks were retained only for bilateral findings or multi-object comparisons. DeepTumorVQA questions assigned three or more masks were reviewed to retain only the masks needed to support the answer. Questions without suitable masks received no grounding annotation.

BIMCV report and VQA construction.[6] We excluded empty reports, reports shorter than 10 or longer than 512 words, and duplicate reports. Cleaning of the Spanish-to-English translations retained current imaging findings and removed non-imaging content and references to prior studies, without adding unsupported facts. Cleaned narratives formed BIMCV-RG, and BIMCV-VQA contained up to five open-ended single-turn dialogues per CT scan. Only non-redundant questions supported by the cleaned report were retained.

Model implementation

CT preprocessing and augmentation. Masks underwent the same cropping, resampling and padding as the detailed input, using nearest-neighbour interpolation and zero padding. Spatial augmentation applied identical random transformations to the global input, the detailed input and the masks. Each spatial axis was flipped independently with probability 0.2, and rotations of 90°, 180° or 270° in the plane of the last two spatial axes were applied with probability 0.2. For intensity augmentation, scaled intensities were multiplied by a factor sampled uniformly from $[0.9, 1.1]$ with probability 0.1 and shifted by an offset in $[-0.1, 0.1]$ with probability 0.5.

Network architecture. The global encoder was adapted from the Med3DVLM DCFormer[7], with five stages of 64, 96, 192, 384 and 768 channels containing 1, 2, 3, 6 and 2 blocks, respectively. Its decomposed 3D convolutions summed the outputs of three axis-aligned branches. The penultimate feature map had 256 spatial locations and 384 channels, and the final map had 32 locations and 768 channels. The global projector passed each map through two Mixer layers, each using two-layer GELU multilayer perceptrons (MLPs) for token and channel mixing, and concatenated the outputs into 256 whole-volume tokens.

The detailed encoder had five resolution levels with 48, 96, 192, 384 and 768 channels containing 1, 2, 2, 4 and 4 residual blocks, respectively. Four strided operations downsampled the input by a factor of 16 along each axis. The detailed projector applied average pooling with a kernel and stride of 2 along the depth axis and 1 along the other axes, followed by a GELU MLP and the Qwen3-VL visual merger[8], which combined each 2×2 in-plane neighbourhood into one fine-grained token of the language model's hidden dimension. These operations gave an overall stride of 32 along each axis. The encoders and projectors are shown in Extended Data Fig. 2a–d.

Visual tokens and position embedding. Whole-volume tokens followed <global_feature>, and each plane of fine-grained tokens followed <local_feature>. The chat template combined these blocks with the query using the Qwen3-VL vision-start and vision-end delimiters. Position indices advanced sequentially through text and delimiters, and each visual block started after the largest preceding index, which defined b_k. Depth markers were tokenized as ordinary text and could span multiple positions.

Target representation and mask decoding. Each target description was enclosed by <s> and </s>. These two tokens, <SEG>, <global_feature> and <local_feature> were added to the tokenizer vocabulary, and the embedding matrix was expanded accordingly. The projection from the <SEG> hidden state to q_k was a GELU MLP. The mask decoder upsampled the bottleneck features of the detailed encoder through four stages with skip connections, followed by two residual 3D convolutional blocks. Output restoration removed padding and reversed resampling, cropping and reorientation, preserving the original affine transformation.

Alternative interaction pathways. In variants S4 and S5, the mask decoder received fine-grained features from the detailed encoder and <SEG> hidden states from the language model. S5 additionally applied two successive two-way attention blocks[9] to the projected whole-volume and fine-grained tokens before the language model. Each block had 16 attention heads and residual connections and applied, in order, self-attention among whole-volume tokens, cross-attention from whole-volume queries to fine-grained keys and values, an MLP, and reverse cross-attention from fine-grained queries to whole-volume keys and values. Queries and keys used 3D rotary position embeddings. After both blocks, each token stream was added to its projected input features. The updated whole-volume tokens entered the language model, and the updated fine-grained tokens were passed to the mask decoder. These pathways are shown in Extended Data Fig. 1c, and the two-way attention block in Extended Data Fig. 2e.

Training implementation and ablation benchmarks

Initialization. The visual merger in the detailed projector inherited Qwen3-VL weights, and mask-decoder layers retained from VISTA3D inherited VISTA3D weights[10]. The remaining projector layers, the <SEG> projection, the embeddings of newly added tokens and the S5 two-way attention blocks were randomly initialized.

Sampling and loss computation. Language-supervised and segmentation-supervised samples were drawn from separately shuffled pools, and samples were repeated as needed to complete each distributed sampling cycle. During training and validation, each GPU processed one example per minibatch, and gradients were accumulated over four minibatches per optimizer update. Supervised response positions included the response-end marker and excluded system and user tokens, visual placeholders and padding. Mask losses were computed in float32, and BCE was computed from logits for numerical stability.

Ablation benchmarks. Training and inference ablations used the same benchmarks, preprocessing and prompts. Target-level referring segmentation was evaluated on TotalSegmentatorV2[11] and AbdomenAtlas3.0[12] using macro Dice, and single-target lesion screening was evaluated on AbdomenAtlas3.0 and LiMT[13] using macro F1. Grounded VQA was evaluated on AMOS-MM-VQA and DeepTumorVQA using accuracy. Anatomy-grounded and finding-grounded report generation were evaluated on RadGenome and ReXGroundingCT[14], respectively, using report LCS. Prompts for textual outputs included grounding requests, except in the inference settings in which these requests were removed. Segmentation was not evaluated for S1, which received no segmentation supervision. Results are shown in Fig. 7 and Extended Data Fig. 1c, and scores are listed in Supplementary Tables 22 and 23.

Baseline training

nnU-Net. nnU-Net[15] used its automatically planned 3D full-resolution configuration and batch size. Following SAT[16], TotalSegmentator labels were divided into six non-overlapping subsets, each assigned a separate model, whereas a single model was trained for each other dataset. For datasets with overlapping annotations, labels were converted to mutually exclusive classes for training only, with higher-index labels taking precedence. Models were trained for 1,000 epochs of 250 iterations

each, using SGD with an initial learning rate of 0.01 to minimize the sum of Dice and cross-entropy losses.

3D ResNet-18.[17] Inputs were resized to 128^3 voxels, with intensities clipped to $[-1000, 1000]$ HU. Models were trained for 100 epochs with a batch size of 100 in bfloat16 precision, using AdamW[18] with a fixed learning rate of 10^{-4} and weight decay of 10^{-4} to minimize class-weighted cross-entropy loss.

M3D and HuluMed-32B. M3D[19] inputs were min–max normalized and resized to $32 \times 256 \times 256$ voxels. HuluMed-32B[20] processed 180 axial slices with a spatial merge size of 2 and token compression. Both models were trained for three epochs with a batch size of 64, using AdamW with a learning rate of 5×10^{-5}, 3% warmup followed by cosine decay, and weight decay of 0.01; HuluMed-32B was trained in bfloat16 precision. LoRA[21] used $r = 16$, $\alpha = 32$ and dropout 0.05 for M3D, and $r = 64$, $\alpha = 128$ and dropout 0.05 for HuluMed-32B.

SAT-Pro. The image encoder, BioLORD-based text encoder and projector were updated. Inputs were reoriented to RAS, resampled to $1 \times 1 \times 3$ mm spacing, clipped to $[-500, 1000]$ HU and z-score normalized per scan. Training crops contained $288 \times 288 \times 64$ voxels, with a patch size of 32^3 voxels. The model was trained for 25 epochs with a batch size of 16 in bfloat16 precision, using AdamW with a learning rate of 10^{-4} and 2,000 warmup steps.

Evaluation implementation

Evaluation prompts and target-name mappings are listed in Supplementary Tables 25–28 and 7, and benchmark scores in Supplementary Tables 8–21.

Segmentation and lesion screening. Predicted probability maps were binarized at 0.5, and predicted and reference masks were scored on the reference voxel grid without further resampling. Foreground comprised nonzero voxels in binary masks or the target label in multiclass masks; reference labels came from the original annotations, and each target was scored independently. Pairs in which both masks were empty were excluded, and pairs with one empty mask received Dice 0. HD95[22] was computed using the DeepMind surface-distance implementation. When HoloCT generated a mask for the same target more than once, only the first mask channel was evaluated.

Segmentation baselines. VISTA3D was prompted with class labels only, without points or boxes, and masks of multiple classes mapped to one target were merged by union. Text-prompted baselines received each model's own target names when available and HoloDATA target names otherwise. VoxTell[23] predictions were restored from the cropped region to native space. Targets unsupported by nnU-Net or VISTA3D received Dice 0.

Closed-ended and grounded VQA. Questions with unparseable reference answers were excluded. Visual grounding was scored independently of answer accuracy. Object-set exact match required the predicted object set to equal the reference set, with no unknown or duplicate objects.

Open-ended VQA and report generation. BLEU-4[24] was computed with smoothing and ROUGE-L[25] with stemming. When several references were available, the first was used; empty predictions received a score of zero. The LLM judge[26] used temperature 0 throughout, with separate evaluation prompts for VQA and report generation. For VQA, the judge received the question, question type, reference and prediction, and assessed accuracy, relevance, completeness and clinical soundness. RaTEScore[27] used RaTE-NER-DeBERTa for entity extraction and BioLORD-2023-C for semantic comparison.

Anatomy-grounded report generation. Grounded and ungrounded report generation were compared on the same scans, with the same reference reports and evaluation prompts. Before text scoring, grounding tokens were removed from generated reports, leaving clinical wording, anatomical labels and their order unchanged.

Finding-grounded report generation. Predicted findings were extracted from finding tags in HoloCT reports and, for the pipeline baseline, by splitting HuluMed-32B reports at commas; both sets were then evaluated identically. Contradictory clinical attributes assessed by the LLM judge included laterality, negation, distribution and spatial relationships. Ties between highest-Dice predictions were resolved in favour of the earliest prediction.

Metric aggregation and figure summaries

Lesion screening. For each of the 18 targets in Fig. 3h, scores were averaged over all dataset–task settings, covering single-target and multi-target lesion screening in internal and external zero-shot evaluations. Each model's summary score was the unweighted mean of its 18 target scores.
VQA. In Fig. 4a, AMOS-MM-VQA and DeepTumorVQA scores were unweighted means of question-category accuracies, whereas Fig. 4c shows the mean of the two datasets' question-level accuracies.
Report generation. Segmentation scores for anatomy-grounded report generation were averaged with equal weight across anatomical regions.
Ablation. In Fig. 7c,e, ablation scores were averaged within each dataset and then with equal weight across datasets. The score drop after grounding-request removal was the score with grounding requests minus the score without them, computed from the values in Fig. 7e.
Radar plots. Radar plots in Fig. 2d,f and Fig. 4a,e,f normalized each axis to the HoloCT-8B score for visualization only; statistical tests used unnormalized scores.
Error bars. Whiskers in Fig. 6a,b,e show CT-level bootstrap 95% confidence intervals for clinical lesion segmentation and report LCS. Whiskers in Fig. 2a,b, Fig. 3c,e,f and Extended Data Fig. 1a,b span the minimum and maximum target scores. In Fig. 7e, whiskers span the two dataset-level scores, and in Fig. 7g, the two dataset-level score drops.

Supplementary Table 1 | Dataset details for segmentation. Dataset Type: Internal, External or Private. Here, aligned means that a dataset follows the target dataset's split to prevent label leakage.

ID	Dataset Name	3D CTs	Classes	Masks	Body Parts	Split Rule	Dataset Type	Download Links
001	AbdomenCT-1k [28]	957	4	3,606	Abdomen, Genitourinary	VISTA3D	Internal	Download
002	AMOS22 [29]	588	15	5,167	Abdomen, Thorax	VISTA3D	Internal	Download
003	MSD-colon [30]	190	1	126	Abdomen	VISTA3D	Internal	Download
004	MSD-hepatic-vessel [30]	443	2	606	Abdomen, Vascular	VISTA3D	Internal	Download
005	MSD-liver [30]	201	2	249	Abdomen	VISTA3D	Internal	Download
006	MSD-lung [30]	95	1	63	Thorax	VISTA3D	Internal	Download
007	MSD-pancreas [30]	420	2	562	Abdomen	VISTA3D	Internal	Download
008	MSD-spleen [30]	61	1	41	Abdomen	VISTA3D	Internal	Download
009	AeroPath [31]	27	2	54	Thorax	VISTA3D	Internal	Download
010	VerSe [32]	374	25	4,543	Musculoskeletal	VISTA3D	Internal	Download
011	Total-SegmentatorV2 [11]	1,228	117	86,557	Whole body	VISTA3D	Internal	Download
012	Pancreas-CT [33]	80	1	80	Abdomen	VISTA3D	Internal	Download
013	BTCV-Abdomen [34]	50	15	388	Abdomen, Thorax	VISTA3D	Internal	Download
014	BTCV-Cervix [34]	50	4	118	Abdomen, Genitourinary	VISTA3D	Internal	Download
015	CRLM-CT [35]	197	4	788	Abdomen, Vascular	VISTA3D	Internal	Download
016	HaN-Seg [36]	42	30	1,259	Head & Neck, Vascular	SAT	Internal	Download
017	KiTS23 [37]	489	3	1,226	Genitourinary	SAT	Internal	Download
018	KiPA22 [38]	100	4	280	Genitourinary, Vascular	SAT	Internal	Download
019	CHAOS [39]	40	1	20	Abdomen	SAT	Internal	Download
020	Flare22 [40]	50	13	650	Abdomen, Thorax	VISTA3D	Internal	Download
021	Flare23 [41]	2,550	14	14,506	Abdomen, Thorax	Flare22-aligned	Internal	Download
022	CT-ORG [42]	112	6	539	Whole body	VISTA3D	Internal	Download
023	LIDC-IDRI [43]	703	1	703	Thorax	random	Internal	Download
024	SLIVER07 [44]	20	1	20	Abdomen	SAT	Internal	Download
025	LUNA16 [45]	888	3	2,663	Thorax	SAT	Internal	Download
026	COVID-19 [46]	20	3	60	Thorax	SAT	Internal	Download
027	CTPelvic1K-CLINIC [47]	178	4	468	Musculoskeletal	SAT	Internal	Download
028	CTPelvic1K-COLONOG [47]	713	4	2,851	Musculoskeletal	SAT	Internal	Download
029	CTPelvic1K-KiTS [47]	44	4	175	Musculoskeletal	KiTS23-aligned	Internal	Download
030	CTPelvic1K-MSD-colon [47]	155	4	620	Musculoskeletal	MSD-colon-aligned	Internal	Download
031	CTPelvic1K-BTCV-Cervix [47]	41	4	164	Musculoskeletal	BTCV-Cervix-aligned	Internal	Download
032	CTPelvic1K-BTCV-Abdomen [47]	35	4	140	Musculoskeletal	BTCV-Abdomen-aligned	Internal	Download
033	Couinaud-Liver (MSD-hepaticvessel) [48]	193	8	1,530	Abdomen	MSD-hepatic-vessel-aligned	Internal	Download
034	DAP-Atlas (AutoPETIII) [49]	533	142	73,971	Whole body	AutoPETIII-aligned	Internal	Download
035	FUMPE [50]	33	1	33	Thorax	SAT	Internal	Download
036	LNDb [51]	229	1	229	Thorax	SAT	Internal	Download
037	MM-WHS-CT [52]	20	7	140	Thorax, Vascular	SAT	Internal	Download
038	PDDCA [53]	48	9	405	Head & Neck	SAT	Internal	Download
039	Parse2022 [54]	100	1	100	Vascular	SAT	Internal	Download

Supplementary Table 1 (continued) | Dataset details for segmentation. Dataset Type: Internal, External or Private. Here, aligned means that a dataset follows the target dataset's split to prevent label leakage.

ID	Dataset Name	3D CTs	Classes	Masks	Body Parts	Split Rule	Dataset Type	Download Links
040	SEGA [55]	56	1	56	Vascular	SAT	Internal	Download
041	SegRap2023-Task1 [56]	120	45	5,400	Head & Neck, Thorax	SAT	Internal	Download
042	SegRap2023-Task2 [56]	120	2	240	Head & Neck	SAT	Internal	Download
043	SegTHOR [57]	40	4	160	Thorax, Vascular	SAT	Internal	Download
044	Seg-CQ500 [58]	51	2	102	Head & Neck	random	Internal	Download
045	NSCLC-Radiomics [59]	422	6	1,938	Head & Neck, Thorax	SAT	Internal	Download
046	PleThora (NSCLC-Radiomics) [60]	402	2	479	Thorax	NSCLC-Radiomics-aligned	Internal	Download
047	WORD [61]	150	16	1,920	Abdomen, Thorax	SAT	Internal	Download
048	RAOS [62]	317	19	5,748	Abdomen, Thorax	official	Internal	Download
049	CT-ICH [63]	75	1	36	Head & Neck	random	Internal	Download
050	Mediastinal-Lymph-Node-SEG [64]	513	1	513	Thorax	random	Internal	Download
051	3D-ircadb-01 [65]	20	32	182	Abdomen, Thorax	random	Internal	Download
052	NSCLC-Radiogenomics [66]	295	3	542	Thorax	SAT	Internal	Download
053	PanTS [67]	9,901	28	227,102	Abdomen, Thorax	random	Internal	Download
054	PASTA-Gen-30K-Bladder cancer [68]	2,000	1	2,000	Genitourinary	random	Internal	Download
055	PASTA-Gen-30K-Bone metastasis [68]	2,000	1	2,000	Musculoskeletal	random	Internal	Download
056	PASTA-Gen-30K-Colorectal cancer [68]	2,000	1	2,000	Abdomen	random	Internal	Download
057	PASTA-Gen-30K-Esophageal cancer [68]	2,000	1	2,000	Thorax	random	Internal	Download
058	PASTA-Gen-30K-Gallbladder cancer [68]	2,000	1	2,000	Abdomen	random	Internal	Download
059	PASTA-Gen-30K-Gallstone [68]	2,000	1	2,000	Abdomen	random	Internal	Download
060	PASTA-Gen-30K-Gastric cancer [68]	2,000	1	2,000	Abdomen	random	Internal	Download
061	PASTA-Gen-30K-Kidney cyst [68]	2,000	1	2,000	Genitourinary	random	Internal	Download
062	PASTA-Gen-30K-Kidney stone [68]	2,000	1	2,000	Genitourinary	random	Internal	Download
063	PASTA-Gen-30K-Kidney tumor [68]	2,000	1	2,000	Genitourinary	random	Internal	Download
064	PASTA-Gen-30K-Liver cyst [68]	2,000	1	2,000	Abdomen	random	Internal	Download
065	PASTA-Gen-30K-Liver tumor [68]	2,000	1	2,000	Abdomen	random	Internal	Download
066	PASTA-Gen-30K-Lung tumor [68]	2,000	1	2,000	Thorax	random	Internal	Download
067	PASTA-Gen-30K-Pancreatic cyst [68]	2,000	1	2,000	Abdomen	random	Internal	Download
068	PASTA-Gen-30K-Pancreatic tumor [68]	2,000	1	2,000	Abdomen	random	Internal	Download
069	PASTA-Gen-30K-TotalSeg [68]	30,000	104	1,691,982	Whole body	Total-SegmentatorV2-aligned	Internal	Download
070	AbdomenAtlas 3.0 [12]	9,262	44	361,562	Abdomen, Thorax	Several-dataset-aligned	Internal	Download
071	LiMT [13]	150	5	290	Abdomen	random	Internal	Download
072	ULS23-KiTS23 [69]	332	1	332	Genitourinary	KiTS23-aligned	Internal	Download
073	ULS23-LiTS [69]	832	1	832	Abdomen	MSD-liver-aligned	Internal	Download

Supplementary Table 1 (continued) | Dataset details for segmentation. Dataset Type: Internal, External or Private. Here, aligned means that a dataset follows the target dataset's split to prevent label leakage.

ID	Dataset Name	3D CTs	Classes	Masks	Body Parts	Split Rule	Dataset Type	Download Links
074	ULS23-LIDC-IDRI [69]	2,236	1	2,236	Thorax	LIDC-IDRI-aligned	Internal	Download
075	ULS23-MSD-lung [69]	70	1	70	Thorax	MSD-lung-aligned	Internal	Download
076	ULS23-MSD-colon [69]	131	1	131	Abdomen	MSD-colon-aligned	Internal	Download
077	ULS23-MSD-pancreas [69]	283	1	283	Abdomen	MSD-pancreas-aligned	Internal	Download
078	LiMT-oneclass [13]	150	2	250	Abdomen	LiMT-aligned	Internal	Download
079	ULS23-NIG-LN-ABD [69]	557	1	557	Abdomen	CT-Lymph-Nodes-aligned	External	Download
080	ULS23-NIG-LN-MED [69]	375	1	375	Thorax	CT-Lymph-Nodes-aligned	External	Download
081	ULS23-Radboudumc-Bone [69]	697	1	697	Musculoskeletal	random	External	Download
082	ULS23-Radboudumc-Pancreas [69]	120	1	120	Abdomen	random	External	Download
083	ULS23-DeepLesion3D [69]	743	1	743	Abdomen, Thorax	random	External	Download
084	CT-Lymph-Nodes [70]	176	2	176	Thorax, Abdomen	random	External	Download
085	HCC-TACE-Seg [71]	337	4	724	Abdomen	random	External	Download
086	QUBIQ2021 [72]	58	2	90	Abdomen	random	External	Download
087	PANORAMA [73]	2,203	5	9,116	Abdomen, Vascular	random	External	Download
088	TotalSeg-subtask-liver-lesion [74]	842	1	815	Abdomen	Total-SegmentatorV2-aligned	External	Download
089	MHB-SEG-multiclass [75]	240	5	594	Head & Neck	MHB-SEG-aligned	External	Download
090	MHB-SEG-oneclass [75]	315	1	270	Head & Neck	MHB-SEG-aligned	External	Download
091	Pediatric-CT-SEG [76]	359	29	8,880	Abdomen, Thorax	random	External	Download
092	VEELA2025 [77]	20	3	60	Abdomen, Vascular	random	External	Download
093	Pengwin [78]	100	3	300	Musculoskeletal	random	External	Download
094	Adrenal-ACC-Ki67 [79]	41	1	41	Genitourinary	random	External	Download
095	Private01(laryngeal cancer)	439	1	439	Head & Neck	random	Private	Not public
096	Private02(lung cancer)	1,191	1	1,191	Thorax	random	Private	Not public
097	Private03(thyroid cancer-multi)	712	2	625	Head & Neck	Private03-aligned	Private	Not public
098	Private03(thyroid cancer-one)	712	1	549	Head & Neck	Private03-aligned	Private	Not public
099	Private04(colon lesion)	832	1	0	Abdomen	random	Private	Not public
TOTAL	**All included datasets**	**109,294**	**873**	**2,564,448**				

Supplementary Table 2 | Dataset details for grounded lesion screening. Dataset Type: Internal, External or Private. Here, aligned means that a dataset follows the target dataset's split to prevent label leakage. Masks equal the number of positive CTs for mask-annotated labels; Private04 has classification labels only and no masks. Positive and Negative are full-cohort label-specific counts; NR denotes not reported. The Private04 held-out test set contains 166 CTs (95 positive, 71 negative); these test counts are not full-cohort counts. Total CTs count each listed dataset once across its labels; overlapping dataset variants are not deduplicated.

ID	Dataset Name	3D CTs	Label Name	Masks	Positive	Negative	Split Rule	Dataset Type	Download Links
001	Total-SegmentatorV2 [11]	1,228	left kidney cyst	30	30	1,198	VISTA3D	Internal	Download
002			right kidney cyst	25	25	1,203			
003	KiTS23 [37]	489	kidney cyst	248	248	241	SAT	Internal	Download
004	CT-ICH [63]	75	intracranial hemorrhage	36	36	39	random	Internal	Download
005	PanTS [67]	9,901	pancreas lesion	1,032	1,032	8,869	random	Internal	Download
006	AbdomenAtlas 3.0 [12]	9,262	kidney lesion	2,222	2,222	7,040	Several-dataset-aligned	Internal	Download
007			liver lesion	1,508	1,508	7,754			
008			pancreas lesion	1,113	1,113	8,149			
009			colon lesion	183	183	9,079			
010	LiMT [13]	150	liver cyst	50	50	100	random	Internal	Download
011			liver hemangioma	37	37	113			
012			metastatic liver cancer	22	22	128			
013			primary liver cancer	31	31	119			
014	LiMT-oneclass [13]	150	liver lesion	100	100	50	LiMT-aligned	Internal	Download
015	QUBIQ2021 [72]	58	pancreas lesion	32	32	26	random	External	Download
016	PANORAMA [73]	2,203	pancreatic ductal adenocarcinoma	662	662	1,541	random	External	Download
017	TotalSeg-subtask-liver-lesion [74]	842	liver lesion	815	815	27	Total-SegmentatorV2-aligned	External	Download
018	MHB-SEG-multiclass [75]	240	epidural hemorrhage	32	32	208	MHB-SEG-aligned	External	Download
019			subdural hemorrhage	158	158	82			
020			subarachnoid hemorrhage	142	142	98			
021			intraparenchymal hemorrhage	142	142	98			
022			intraventricular hemorrhage	120	120	120			
023	MHB-SEG-oneclass [75]	315	intracranial hemorrhage	270	270	45	MHB-SEG-aligned	External	Download
024	Private03(thyroid cancer-multi)	712	malignant thyroid tumor	193	193	519	Private03-aligned	Private	Not public
025			benign thyroid tumor	432	432	280			
026	Private03(thyroid cancer-one)	712	thyroid tumor	549	549	163	Private03-aligned	Private	Not public
027	Private04(colon lesion)	832	colon lesion	0	NR	NR	Patient-level 80/20	Private	Not public
TOTAL		**27,169**		**10,184**					

Supplementary Table 3 | Dataset details for visual question answering. Dataset Type: Internal, External or Private. Here, aligned means that a dataset follows the target dataset's split to prevent label leakage.

Dataset Name	3D CTs	Single-turn QAs	Body Parts	Split Rule	Dataset Type	Download Links
CT-RATE-option [1]	46,609	135,384	Thorax, Vascular, Musculoskeletal	official	Internal	Download
CT-RATE-long&short [1]	50,175	2,472,480 (within 669,798 multi-turn QAs)	Thorax, Vascular, Musculoskeletal	official	Internal	Download
3D-RAD [80]	16,188	170,105	Thorax, Vascular, Musculoskeletal	official	Internal	Download
RadGenome-VQA [2]	25,692	481,752	Thorax, Vascular, Musculoskeletal	official	Internal	Download
AMOS-MM-VQA [5]	1,687	16,538	Abdomen, Genitourinary, Vascular, Musculoskeletal	AMOS-22-aligned	Internal	Download
FLARE-2025-VQA [81]	3,917	37,795	Abdomen, Genitourinary, Vascular, Musculoskeletal	AMOS-MM-aligned	Internal	Download
DeepTumorVQA [4]	9,255	439,973	Abdomen, Genitourinary, Vascular, Musculoskeletal	AbdomenAtlas3.0-aligned	Internal	Download
BIMCV-VQA [6]	6,991	26,434	Thorax, Vascular, Musculoskeletal	random	External	Download
TOTAL	**160,514**	**3,780,461**				

Supplementary Table 4 | Dataset details for report generation. Dataset Type: Internal, External or Private. Here, aligned means that a dataset follows the target dataset's split to prevent label leakage.

Dataset Name	3D CTs	Single-turn QAs	Body Parts	Split Rule	Dataset Type	Download Links
CT-RATE-RG [1]	50,175	50,175	Thorax, Vascular, Musculoskeletal	official	Internal	Download
RadGenome [2]	25,692	223,643	Thorax, Vascular, Musculoskeletal	official	Internal	Download
INSPECT [82]	23,226	23,226	Thorax, Vascular, Musculoskeletal	random	Internal	Download
Merlin [3]	25,372	328,318	Abdomen, Genitourinary, Vascular, Musculoskeletal	random	Internal	Download
AMOS-MM-RG [5]	1,687	5,350	Abdomen, Genitourinary, Vascular, Musculoskeletal	AMOS-22-aligned	Internal	Download
FLARE-2025-RG [81]	3,917	35,295	Abdomen, Genitourinary, Vascular, Musculoskeletal	AMOS-MM-aligned	Internal	Download
AbdomenAtlas3.0 [12]	9,262	9,262	Abdomen, Genitourinary, Vascular, Musculoskeletal	Several-dataset-aligned	Internal	Download
BIMCV-RG [6]	6,991	6,991	Thorax, Vascular, Musculoskeletal	random	External	Download
Private05(Abdomen Report)	5,307	5,307	Abdomen, Genitourinary, Vascular, Musculoskeletal	Zero-shot test	Private	Not public
Private06(Chest Report)	2,974	2,974	Thorax, Vascular, Musculoskeletal	Zero-shot test	Private	Not public
TOTAL	**154,603**	**690,541**				

Supplementary Table 5 | Dataset details for grounded closed-ended visual question answering and grounded report generation. Dataset Type: Internal, External or Private. Here, aligned means that a dataset follows the target dataset's split to prevent label leakage. This table lists public grounding-supervision sources. Private report-evaluation cohorts are listed in Supplementary Table 4; their text results appear in Supplementary Table 17. RadGenome-grounded contains 192,242 + 25,690 = 217,932 single-turn QAs.

Dataset Name	3D CTs	Masks	Single-turn QAs	Body Parts	Split Rule	Dataset Type	Download Links
AMOS-MM-VQA-grounded [5]	207	1,642	1,438	Abdomen, Genitourinary, Vascular, Musculoskeletal	AMOS-22-aligned	Internal	Download
DeepTumorVQA-grounded [4]	9,169	329,929	218,226	Abdomen, Genitourinary, Vascular, Musculoskeletal	AbdomenAtlas3.0-aligned	Internal	Download
RadGenome-VQA-grounded [2]	25,687	162,480	468,643	Thorax, Vascular, Musculoskeletal	CT-RATE-aligned	Internal	Download
RadGenome-grounded [2]	25,690	192,242	217,932	Thorax, Vascular, Musculoskeletal	CT-RATE-aligned	Internal	Download
ReXGroundingCT [14]	3,042	7,802	10,844	Thorax, Vascular, Musculoskeletal	CT-RATE-aligned	Internal	Download
TOTAL	**63,795**	**694,095**	**917,083**				

Supplementary Table 6 | Dataset composition across training stages. Each dialogue represents one training sample: a single-turn dialogue and a multi-turn dialogue each count as one sample. Counts are stage-specific and expressed in millions (M); they are not sums of the single-turn QA counts in the source inventories.

Training stage	Supervision stream	Source data	Dialogue types included	Stage-specific selection	No. of dialogues
Stage 1: 3D CT vision–language alignment	Language	Supplementary Table 4	Report generation	High-quality report-generation dialogues	**0.68M**
Stage 2: 3D CT multitask pre-training	Language	Supplementary Tables 2–4	Single-target and multi-target lesion screening without grounding; closed-ended and open-ended VQA; narrative and structured report generation	All eligible training dialogues from the corresponding training splits	**2.70M**
	Segmentation	Supplementary Tables 1, 2 and 5	Target-level and finding-level referring segmentation; single-target and multi-target lesion screening with grounding; grounded VQA; anatomy-grounded and finding-grounded report generation	All eligible training dialogues from the corresponding training splits	**3.50M**
Stage 3: 3D CT high-quality multitask tuning	Language	Supplementary Tables 2–4	Single-target and multi-target lesion screening without grounding; closed-ended and open-ended VQA; narrative and structured report generation	Single-region report-generation dialogues were excluded; RadGenome-VQA dialogues were excluded	**1.69M**
	Segmentation	Supplementary Tables 1, 2 and 5	Target-level and finding-level referring segmentation; single-target and multi-target lesion screening with grounding; grounded VQA; anatomy-grounded and finding-grounded report generation	Low-quality segmentation data including PASTA-Gen-TotalSeg were excluded; RadGenome grounded-VQA dialogues were excluded; single-region grounded-report dialogues were excluded	**1.15M**

Supplementary Table 7 | Segmentation label hierarchy and counts.

Primary Category	Secondary Category	Standard Terms and Counts	
		ID	OOD
Head & Neck	Brain & CNS	brain(962), brainstem(210), optic chiasm(210), pituitary gland(162), left temporal lobe(120), right temporal lobe(120), left hippocampus(120), right hippocampus(120), intracranial hemorrhage(357)	epidural hemorrhage(32), subdural hemorrhage(158), subarachnoid hemorrhage(142), intraparenchymal hemorrhage(142), intraventricular hemorrhage(120)
	Sensory Organs	left cochlea(162), right cochlea(162), left anterior eyeball(42), right anterior eyeball(42), left posterior eyeball(42), right posterior eyeball(42), left lacrimal gland(42), right lacrimal gland(42), left optic nerve(210), right optic nerve(210), left eyeball(652), right eyeball(648), left lens(120), right lens(120), left middle ear(120), right middle ear(120), left internal auditory canal(120), right internal auditory canal(120), left tympanic cavity(120), right tympanic cavity(120), left vestibule semicircular canal(120), right vestibule semicircular canal(120), left eustachian tube bone(120), right eustachian tube bone(120), left mastoid process(120), right mastoid process(120)	–
	Upper Aerodigestive Tract	arytenoid(42), buccal mucosa(42), oral cavity(162), cricopharyngeal inlet(42), cervical esophagus(42), larynx glottis(162), larynx supraglottis(162), lips(42), nasal cavity(533), larynx(120), pharynx constrictor muscle(120), nasopharyngeal tumor(120)	laryngeal cancer(439)
	Neck Glands & Maxillofacial	thyroid gland(704), skull(1,006), mandible(82), left submandibular gland(203), right submandibular gland(198), thyroid(162), left parotid gland(210), right parotid gland(210), left thyroid(533), right thyroid(532), left cheek(468), right cheek(451), left mandible(120), right mandible(120), left temporomandibular joint(120), right temporomandibular joint(120), nasopharyngeal lymph node(120), face(259)	malignant thyroid tumor(193), benign thyroid tumor(432), thyroid tumor(549)
Thorax	Lung & Pleura	lung tumor(2,664), lung(435), left lung upper lobe(21,740), left lung lower lobe(27,320), right lung upper lobe(6,386), right lung middle lobe(20,878), right lung lower lobe(27,582), pulmonary nodule(3,306), left lung(20,152), right lung(19,936), covid-19 infection(20), lung effusion(78)	lung cancer(1,191)
	Heart & Great Airways	trachea(5,148), heart(1,980), left atrial appendage of heart(673), left heart atrium(10,595), heart tissue(533), right heart atrium(13,503), myocardium(16,330), left heart ventricle(14,750), right heart ventricle(16,339), bronchie(533)	–
	Mediastinum & Esophagus	esophagus(34,566), mediastinal tissue(533), thymus(462), mediastinal lymph node(978), esophageal cancer(2,000)	–
	Chest Wall & Breast	left breast(593), right breast(597)	–
	Thoracic Cavity	thoracic cavity(401)	–
Abdomen	Liver	liver(53,664), liver tumor(3,465), caudate lobe(9,398), left lateral superior segment of liver(9,398), left lateral inferior segment of liver(9,383), left medial segment of liver(9,407), right anterior inferior segment of liver(9,381), right posterior inferior segment of liver(9,383), right posterior superior segment of liver(9,397), right anterior superior segment of liver(9,398), liver cyst(2,053), liver lesion(2,423), liver hemangioma(37), metastatic liver cancer(22), primary liver cancer(31)	hepatocellular carcinoma(337)
	Pancreatobiliary System	pancreas(52,082), gallbladder(47,561), pancreas tumor(2,564), biliary system(1), head of pancreas(18,382), body of pancreas(18,320), tail of pancreas(18,109), common bile duct(17,597), pancreatic duct(8,801), pancreas lesion(2,297), gallbladder cancer(2,000), gallstone(2,000), pancreatic cyst(2,000)	pancreatic ductal adenocarcinoma(662), pancreatic parenchyma(2,203)
	Spleen	spleen(50,753)	–
	Gastrointestinal Tract	stomach(49,473), duodenum(50,005), colon cancer(257), small bowel(30,790), colon(50,453), rectum(7,233), intestine(9,627), gastric cancer(2,000), colorectal cancer(2,000), colon lesion(183)	–
	Retroperitoneum & Abdominal Soft Tissue	abdominal tissue(533)	abdominal lymph node(643)
Genitourinary	Adrenal & Kidney	kidney(1,574), right kidney(50,972), left kidney(50,777), right adrenal gland(48,494), left adrenal gland(48,150), left kidney cyst(30), right kidney cyst(25), adrenal gland(121), kidney tumor(2,559), kidney cyst(2,248), left adrenal gland tumor(1), right adrenal gland tumor(1), kidney stone(2,000), kidney lesion(2,554)	adrenocortical carcinoma(41)
	Urinary Bladder	urinary bladder(23,396), bladder cancer(2,000)	–
	Reproductive Organs	prostate or uterus(291), prostate(12,191), uterus(265), gonad(399), uterocervix(342), seminal vesicles(181)	–
Musculoskeletal	Spine & Spinal Canal	cervical vertebrae 1 (c1)(859), cervical vertebrae 2 (c2)(862), cervical vertebrae 3 (c3)(842), cervical vertebrae 4 (c4)(866), cervical vertebrae 5 (c5)(989), cervical vertebrae 6 (c6)(1,362), cervical vertebrae 7 (c7)(2,988), thoracic vertebrae 1 (t1)(3,489), thoracic vertebrae 2 (t2)(3,603), thoracic vertebrae 3 (t3)(3,718), thoracic vertebrae 4 (t4)(3,784), thoracic vertebrae 5 (t5)(4,165), thoracic vertebrae 6 (t6)(5,526), thoracic vertebrae 7 (t7)(8,570), thoracic vertebrae 8 (t8)(13,085), thoracic vertebrae 9 (t9)(18,635), thoracic vertebrae 10 (t10)(23,345), thoracic vertebrae 11 (t11)(26,616), thoracic vertebrae 12 (t12)(28,443), lumbar vertebrae 1 (l1)(28,706), lumbar vertebrae 2 (l2)(28,705), lumbar vertebrae 3 (l3)(28,951), lumbar vertebrae 4 (l4)(28,871), lumbar vertebrae 5 (l5)(28,140), lumbar vertebrae 6 (l6)(50), sacrum(27,216), sacral vertebrae 1 (s1)(597), spinal cord(1,755), lumbar vertebrae(1,103), spinal canal(883)	–
	Thoracic Cage & Shoulder Girdle	left scapula(4,102), right scapula(4,819), left clavicle(3,443), right clavicle(3,453), right rib 4(4,584), right rib 3(3,702), left rib 1(3,280), left rib 2(3,405), left rib 3(3,603), left rib 4(4,594), left rib 5(9,649), left rib 6(16,099), left rib 7(19,073), left rib 8(20,612), left rib 9(23,008), left rib 10(25,761), left rib 11(27,512), left rib 12(27,442), right rib 1(3,370), right rib 2(3,480), right rib 5(9,270), right rib 6(17,062), right rib 7(22,915), right rib 8(24,992), right rib 9(26,115), right rib 10(27,494), right rib 11(28,326), right rib 12(27,538), sternum(1,491), costal cartilage(1,017), rib cartilage(533), manubrium of sternum(533)	–
	Pelvis & Extremities	left humerus(2,159), right humerus(2,806), left femur(17,803), right femur(17,331), left hip(29,131), right hip(29,105), head of left femur(437), head of right femur(437)	–
	Muscles & Other Musculoskeletal Structures	left gluteus maximus(23,602), right gluteus maximus(24,535), left gluteus medius(26,457), right gluteus medius(27,066), left gluteus minimus(16,618), right gluteus minimus(19,221), left autochthon(31,167), right autochthon(31,189), left iliopsoas(30,096), right iliopsoas(30,212), bone(479), muscle(533), bone metastasis(2,000)	bone lesion(697)
Vascular	Arteries	aorta(51,628), brachiocephalic trunk(1,264), right subclavian artery(1,264), left subclavian artery(1,284), right common carotid artery(1,254), left common carotid artery(1,286), left iliac artery(27,922), right iliac artery(27,885), left carotid artery(42), right carotid artery(42), renal artery(70), celiac trunk(9,756), pulmonary artery(4,313), right internal carotid artery(533), left internal carotid artery(533), pulmonary embolism(33), heart ascending aorta(20), superior mesenteric artery(18,176), celiac artery(17,242)	abdominal aorta(25), pancreatic arteries(2,203)
	Veins	inferior vena cava(51,328), hepatic vein(9,761), pulmonary vein(793), left brachiocephalic vein(1,263), right brachiocephalic vein(1,268), superior vena cava(777), portal vein and splenic vein(37,856), left iliac vena(27,374), right iliac vena(27,416), portal vein(217), renal vein(70), right internal jugular vein(533), left internal jugular vein(533), veins(18,644)	pancreatic veins(2,203)
Other	Other	tumor(1,086), fat(533), skin(901), stone(1)	lesion(743)

Supplementary Table 8 | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell. PASTA-Gen-30K-TotalSeg was not included in evaluation. Internal results cover 77 inventory subdatasets and 272 distinct targets.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	AbdomenCT-1k	liver	Dice	0.9618 (0.9605–0.9629)	0.8959 (0.8914–0.9002)	0.9722 (0.9706–0.9735)	0.9755 (0.9739–0.9771)	0.9684 (0.9633–0.9752)	0.9879 (0.9858–0.9898)	0.9750 (0.9734–0.9764)
			HD95	5.026 (4.052–6.588)	9.311 (8.485–10.33)	3.224 (2.941–3.589)	3.938 (3.142–4.817)	4.261 (3.221–4.743)	3.237 (2.633–3.922)	3.438 (2.922–4.085)
		kidney	Dice	0.9123 (0.9042–0.9194)	0.7248 (0.6859–0.7648)	0.9554 (0.9529–0.9577)	0.9624 (0.9594–0.9649)	0.9419 (0.9411–0.9504)	0.9617 (0.9557–0.9667)	0.9532 (0.9472–0.9577)
			HD95	7.731 (6.851–8.702)	9.910 (8.828–11.08)	2.638 (2.451–2.852)	4.799 (3.227–6.799)	4.140 (1.988–4.163)	4.694 (3.493–6.348)	4.093 (3.555–4.726)
		spleen	Dice	0.9530 (0.9508–0.9551)	0.8423 (0.8326–0.8521)	0.9656 (0.9628–0.9679)	0.9739 (0.9707–0.9764)	0.9420 (0.9222–0.9443)	0.9866 (0.9847–0.9883)	0.9752 (0.9735–0.9769)
			HD95	3.394 (2.847–4.172)	7.503 (6.953–8.156)	1.929 (1.713–2.207)	2.158 (1.496–2.990)	2.142 (1.927–2.842)	0.8630 (0.6660–1.178)	1.491 (1.193–1.865)
		pancreas	Dice	0.8531 (0.8459–0.8599)	0.5979 (0.5820–0.6136)	0.8801 (0.8741–0.8858)	0.8550 (0.8465–0.8626)	0.8506 (0.8439–0.8523)	0.8740 (0.8647–0.8828)	0.8739 (0.8665–0.8809)
			HD95	6.900 (6.023–7.961)	14.99 (13.36–16.82)	3.836 (3.461–4.314)	6.722 (5.822–7.778)	4.011 (3.562–4.280)	4.859 (4.194–5.717)	5.254 (4.517–6.079)
		spleen	Dice	0.9222 (0.9072–0.9330)	0.7122 (0.6707–0.7459)	0.9502 (0.9444–0.9554)	0.9364 (0.9227–0.9481)	0.9379 (0.9189–0.9505)	0.9738 (0.9694–0.9776)	0.9552 (0.9503–0.9599)
			HD95	7.332 (4.318–11.60)	17.31 (11.70–26.98)	2.788 (2.120–3.600)	6.354 (3.789–9.422)	3.920 (3.532–5.522)	1.966 (1.287–2.759)	2.686 (2.100–3.345)
		right kidney	Dice	0.9315 (0.9243–0.9373)	0.7018 (0.6565–0.7413)	0.9513 (0.9443–0.9570)	0.9477 (0.9398–0.9544)	0.9448 (0.9331–0.9538)	0.9646 (0.9571–0.9704)	0.9503 (0.9408–0.9576)
			HD95	4.480 (3.783–5.261)	33.48 (22.22–46.98)	2.092 (1.760–2.489)	3.309 (2.553–4.246)	2.684 (1.978–3.406)	2.411 (1.392–4.028)	2.848 (1.933–4.127)
		left kidney	Dice	0.9183 (0.8858–0.9378)	0.6538 (0.6020–0.7016)	0.9509 (0.9456–0.9555)	0.9461 (0.9367–0.9545)	0.9459 (0.9272–0.9521)	0.9668 (0.9593–0.9727)	0.9491 (0.9390–0.9570)
			HD95	4.827 (3.448–7.265)	49.20 (35.35–64.36)	2.157 (1.807–2.626)	6.699 (2.685–13.91)	4.305 (3.451–6.695)	2.010 (1.212–3.234)	2.502 (1.753–3.619)
	AMOS22	gallbladder	Dice	0.8275 (0.7967–0.8560)	0.2908 (0.2269–0.3523)	0.8700 (0.8445–0.8903)	0.7982 (0.7492–0.8408)	0.8366 (0.8034–0.8546)	0.8668 (0.8298–0.8975)	0.8783 (0.8585–0.8972)
			HD95	7.525 (5.984–9.327)	35.56 (22.63–51.06)	4.093 (3.303–5.000)	9.833 (6.999–13.03)	9.396 (7.871–10.45)	6.728 (4.611–9.292)	6.026 (4.265–7.950)
		esophagus	Dice	0.7618 (0.7342–0.7883)	0.3826 (0.3438–0.4268)	0.7955 (0.7699–0.8184)	0.7798 (0.7529–0.8061)	0.7954 (0.7891–0.8161)	0.8307 (0.8056–0.8544)	0.8264 (0.8052–0.8458)
			HD95	51.87 (30.83–76.71)	17.52 (12.58–25.23)	9.354 (4.839–15.49)	11.57 (6.910–18.50)	8.821 (5.067–13.79)	7.676 (4.118–12.24)	8.182 (3.515–14.63)
		liver	Dice	0.9459 (0.9384–0.9506)	0.8206 (0.7867–0.8429)	0.9617 (0.9530–0.9669)	0.9576 (0.9475–0.9643)	0.9712 (0.9657–0.9871)	0.9788 (0.9712–0.9840)	0.9669 (0.9625–0.9705)
			HD95	5.540 (4.589–6.882)	20.12 (13.07–31.50)	3.703 (3.050–4.631)	6.137 (4.450–8.366)	5.968 (4.322–8.316)	3.529 (2.405–4.924)	3.924 (3.266–4.796)
		stomach	Dice	0.8614 (0.8198–0.8941)	0.5750 (0.5264–0.6213)	0.8780 (0.8339–0.9126)	0.8243 (0.7756–0.8644)	0.8563 (0.8157–0.8781)	0.9201 (0.8814–0.9465)	0.9078 (0.8692–0.9341)
			HD95	16.85 (9.919–25.54)	26.73 (19.20–38.88)	20.68 (6.859–38.10)	24.30 (14.77–38.37)	12.75 (8.997–17.21)	6.857 (4.310–10.83)	8.632 (5.319–12.71)
		aorta	Dice	0.9032 (0.8941–0.9110)	0.7327 (0.7077–0.7556)	0.9113 (0.9013–0.9200)	0.9236 (0.9130–0.9325)	0.9480 (0.9304–0.9619)	0.9397 (0.9309–0.9473)	0.9326 (0.9262–0.9386)
			HD95	14.86 (10.17–20.03)	15.70 (11.54–20.62)	12.84 (7.656–18.25)	9.546 (5.394–15.02)	9.920 (6.321–16.52)	7.591 (3.545–12.13)	5.250 (2.893–8.914)
		inferior vena cava	Dice	0.8443 (0.8249–0.8629)	0.6286 (0.5920–0.6618)	0.8834 (0.8693–0.8960)	0.8658 (0.8460–0.8845)	0.8653 (0.8483–0.8706)	0.8642 (0.8435–0.8822)	0.8527 (0.8344–0.8694)
			HD95	7.634 (6.560–8.909)	17.03 (12.11–23.34)	3.885 (3.532–4.281)	5.266 (4.016–7.208)	5.487 (4.166–5.964)	5.709 (4.320–7.583)	5.681 (4.517–7.278)
		pancreas	Dice	0.7877 (0.7437–0.8228)	0.4633 (0.4212–0.5015)	0.8238 (0.7854–0.8522)	0.7745 (0.7334–0.8112)	0.8382 (0.8039–0.8402)	0.8441 (0.8051–0.8740)	0.8401 (0.8038–0.8659)
			HD95	10.24 (7.381–13.91)	25.69 (17.33–37.21)	6.421 (4.395–9.042)	11.08 (8.635–14.28)	8.071 (6.318–12.57)	6.185 (4.188–9.088)	6.226 (4.412–9.015)
		right adrenal gland	Dice	0.6934 (0.6580–0.7261)	0.2357 (0.2006–0.2721)	0.7568 (0.7310–0.7800)	0.7379 (0.7153–0.7601)	0.7019 (0.6861–0.7187)	0.7409 (0.7170–0.7628)	0.7676 (0.7483–0.7863)
			HD95	6.486 (5.530–7.501)	24.52 (14.99–36.95)	3.402 (2.772–4.087)	5.089 (4.334–5.954)	10.98 (9.162–11.19)	6.092 (3.958–8.669)	3.687 (3.114–4.299)
		left adrenal gland	Dice	0.6905 (0.6468–0.7265)	0.2094 (0.1730–0.2451)	0.7517 (0.7072–0.7855)	0.7362 (0.7014–0.7668)	0.7232 (0.7118–0.7546)	0.7547 (0.7167–0.7883)	0.7584 (0.7283–0.7850)
			HD95	6.943 (5.746–8.554)	35.73 (20.90–53.08)	3.355 (2.594–4.531)	5.550 (4.400–6.889)	7.885 (3.908–8.232)	5.244 (3.732–7.169)	5.928 (4.374–7.723)
		duodenum	Dice	0.6869 (0.6441–0.7275)	0.3833 (0.3394–0.4198)	0.7734 (0.7342–0.8041)	0.7035 (0.6654–0.7363)	0.7463 (0.6935–0.7923)	0.7662 (0.7233–0.8004)	0.7716 (0.7304–0.8059)
			HD95	15.08 (12.26–18.52)	30.45 (21.73–40.58)	8.060 (6.321–10.25)	14.87 (12.31–17.59)	9.371 (8.042–12.93)	11.49 (8.873–14.49)	9.211 (7.034–11.92)
		urinary bladder	Dice	0.8094 (0.7659–0.8480)	0.4055 (0.3434–0.4703)	0.8466 (0.8078–0.8783)	0.8079 (0.7704–0.8447)	0.8088 (0.7657–0.8553)	0.8884 (0.8499–0.9176)	0.8687 (0.8314–0.8974)
			HD95	15.03 (8.262–25.44)	44.00 (27.84–63.25)	7.687 (5.239–10.73)	12.55 (9.624–16.14)	8.819 (6.488–9.952)	6.858 (4.620–9.520)	7.230 (5.115–9.884)
		prostate or uterus	Dice	0.7680 (0.7425–0.7899)	0.3878 (0.3283–0.4445)	0.0000 (0.0000–0.0000)	0.7808 (0.7500–0.8070)	0.8247 (0.8109–0.8286)	0.8782 (0.8542–0.8986)	0.8260 (0.7996–0.8527)
			HD95	15.93 (12.60–18.52)	31.77 (22.81–40.86)	239.3 (175.5–310.9)	14.31 (11.43–17.62)	14.32 (9.235–20.42)	9.235 (5.354–15.61)	45.97 (27.17–67.76)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	MSD-colon	colon cancer	Dice	0.4005 (0.2856–0.5077)	0.02744 (0.007287–0.05207)	0.6740 (0.5450–0.7863)	0.6852 (0.6052–0.7550)	0.4700 (0.4376–0.5049)	0.5877 (0.4793–0.6891)	0.6159 (0.5046–0.7023)
			HD95	165.8 (130.6–200.2)	159.5 (132.5–184.9)	18.35 (4.606–38.52)	50.57 (25.94–77.81)	50.57 (35.86–69.49)	45.92 (21.01–80.09)	39.64 (16.25–71.21)
	MSD-hepatic-vessel	hepatic vein	Dice	0.5389 (0.5072–0.5713)	0.2274 (0.1928–0.2622)	0.6835 (0.6572–0.7063)	0.3526 (0.3163–0.3904)	0.5466 (0.5051–0.5754)	0.5575 (0.5313–0.5861)	0.5431 (0.5141–0.5733)
			HD95	25.02 (21.58–28.42)	40.57 (35.26–46.66)	11.10 (9.039–13.33)	33.97 (30.81–37.54)	12.67 (11.13–14.95)	23.64 (20.75–27.14)	24.68 (21.68–28.71)
		liver tumor	Dice	0.6686 (0.6051–0.7261)	0.4081 (0.3255–0.4983)	0.8317 (0.7966–0.8588)	0.7227 (0.6660–0.7752)	0.6561 (0.6394–0.7082)	0.7244 (0.6608–0.7792)	0.7550 (0.7148–0.7929)
			HD95	25.97 (17.43–35.69)	52.85 (39.88–65.89)	14.05 (7.233–23.11)	28.49 (20.02–37.63)	46.78 (37.52–56.03)	27.97 (19.16–38.22)	32.13 (22.91–43.21)
	MSD-liver	liver	Dice	0.9322 (0.9153–0.9468)	0.8293 (0.8088–0.8503)	0.9413 (0.9234–0.9556)	0.9441 (0.9274–0.9597)	0.9359 (0.9359–0.9551)	0.9614 (0.9458–0.9751)	0.9452 (0.9282–0.9599)
			HD95	14.75 (8.405–23.80)	15.03 (13.05–17.05)	6.676 (4.726–8.900)	13.53 (7.388–21.05)	8.406 (7.344–9.245)	7.553 (5.417–10.00)	8.063 (6.008–10.57)
		liver tumor	Dice	0.5706 (0.4435–0.6824)	0.1419 (0.05291–0.2286)	0.7759 (0.7414–0.8088)	0.7015 (0.6234–0.7741)	0.6154 (0.5858–0.6505)	0.6953 (0.6381–0.7538)	0.7238 (0.6599–0.7827)
			HD95	64.06 (29.87–101.9)	104.1 (78.86–133.4)	15.17 (7.883–23.64)	41.16 (23.29–60.48)	26.58 (20.21–34.43)	24.32 (13.93–37.01)	29.35 (15.05–46.90)
	MSD-lung	lung tumor	Dice	0.5982 (0.4332–0.7361)	0.2832 (0.1272–0.4450)	0.7533 (0.6173–0.8433)	0.8410 (0.7961–0.8816)	0.5558 (0.5023–0.6008)	0.6743 (0.5162–0.7923)	0.6819 (0.5300–0.7988)
			HD95	72.63 (19.53–149.0)	87.14 (44.64–132.1)	10.76 (2.730–25.65)	19.73 (3.047–38.36)	52.84 (35.25–78.39)	28.67 (9.081–55.22)	11.80 (4.696–23.45)
	MSD-pancreas	pancreas	Dice	0.8028 (0.7900–0.8156)	0.6045 (0.5823–0.6276)	0.8444 (0.8336–0.8545)	0.8119 (0.7984–0.8253)	0.8039 (0.7900–0.8122)	0.8232 (0.8112–0.8364)	0.8174 (0.8050–0.8303)
			HD95	7.667 (6.807–8.631)	10.30 (9.576–11.11)	4.161 (3.842–4.489)	6.579 (5.675–7.642)	4.739 (4.116–5.713)	5.890 (5.108–6.790)	7.376 (5.848–9.335)
		pancreas tumor	Dice	0.3072 (0.2335–0.3874)	0.2065 (0.1529–0.2669)	0.7656 (0.6933–0.8271)	0.6981 (0.6480–0.7422)	0.4819 (0.4381–0.5140)	0.5931 (0.5269–0.6444)	0.6346 (0.5683–0.6856)
			HD95	67.51 (45.61–93.08)	43.86 (33.98–54.64)	4.453 (2.965–6.489)	16.29 (9.225–24.80)	45.81 (14.42–78.70)	21.10 (15.39–27.45)	25.74 (18.24–34.65)
	MSD-spleen	spleen	Dice	0.9385 (0.9289–0.9470)	0.7589 (0.7100–0.8104)	0.9478 (0.9286–0.9642)	0.9514 (0.9396–0.9617)	0.9441 (0.9437–0.9482)	0.9672 (0.9594–0.9749)	0.9576 (0.9511–0.9638)
			HD95	3.186 (2.783–3.640)	9.828 (8.470–11.17)	2.165 (1.491–3.151)	2.316 (1.708–3.167)	2.605 (2.512–3.128)	1.521 (1.181–1.947)	1.880 (1.601–2.185)
	AeroPath	lung	Dice	0.9456 (0.9254–0.9580)	0.9214 (0.9067–0.9351)	0.9663 (0.9508–0.9768)	0.9710 (0.9652–0.9769)	0.9679 (0.9565–0.9772)	0.9894 (0.9818–0.9950)	0.9799 (0.9757–0.9840)
			HD95	18.77 (15.19–22.29)	16.16 (11.22–21.90)	13.86 (7.278–21.13)	13.62 (10.38–17.51)	10.04 (6.125–14.85)	6.497 (2.352–10.64)	6.263 (2.703–9.824)
		trachea	Dice	0.7231 (0.6776–0.7673)	0.6700 (0.6259–0.7140)	0.7718 (0.7370–0.8038)	0.8129 (0.7474–0.8788)	0.8954 (0.8679–0.9310)	0.8403 (0.7765–0.8994)	0.8043 (0.7434–0.8570)
			HD95	74.29 (67.40–83.76)	47.20 (39.14–57.74)	40.55 (30.65–51.73)	39.40 (28.77–50.20)	34.57 (28.66–42.83)	34.76 (21.59–44.43)	46.46 (36.11–55.08)
	VerSe	cervical vertebrae 1 (c1)	Dice	0.8468 (0.8289–0.8636)	0.1527 (0.06277–0.2515)	0.8426 (0.8266–0.8574)	0.8881 (0.8612–0.9114)	0.8599 (0.8486–0.9011)	0.8623 (0.8395–0.8860)	0.8821 (0.8682–0.8963)
			HD95	3.696 (2.858–4.615)	110.3 (23.98–250.4)	2.131 (1.732–2.671)	3.649 (2.165–5.280)	3.950 (3.679–5.088)	5.063 (3.034–7.517)	1.688 (1.420–1.991)
		cervical vertebrae 2 (c2)	Dice	0.8700 (0.8483–0.8892)	0.1498 (0.04401–0.2614)	0.8884 (0.8800–0.8969)	0.9252 (0.9136–0.9357)	0.8569 (0.8458–0.8619)	0.8980 (0.8713–0.9159)	0.9099 (0.9018–0.9181)
			HD95	14.72 (2.685–37.57)	106.3 (23.47–252.4)	1.608 (1.417–1.833)	2.541 (1.503–3.918)	7.315 (5.970–11.67)	15.96 (1.422–43.54)	1.679 (1.381–2.058)
		cervical vertebrae 3 (c3)	Dice	0.8195 (0.7598–0.8664)	0.08709 (0.03173–0.1618)	0.8315 (0.8107–0.8506)	0.8989 (0.8668–0.9224)	0.8475 (0.8299–0.8569)	0.8274 (0.7911–0.8584)	0.8721 (0.8527–0.8891)
			HD95	3.971 (2.595–5.545)	95.47 (26.83–221.7)	23.98 (2.033–67.53)	1.776 (1.165–2.689)	4.077 (3.052–6.408)	4.955 (3.272–6.939)	2.226 (1.589–3.122)
		cervical vertebrae 4 (c4)	Dice	0.8446 (0.8187–0.8647)	0.1827 (0.08219–0.2875)	0.8146 (0.7767–0.8473)	0.8905 (0.8565–0.9168)	0.8695 (0.8447–0.8877)	0.5642 (0.5470–0.5817)	0.8648 (0.8469–0.8806)
			HD95	21.13 (2.539–57.23)	82.17 (22.58–189.6)	24.92 (1.997–70.20)	2.247 (1.314–3.485)	6.338 (4.069–8.156)	72.95 (14.13–151.9)	2.055 (1.697–2.413)
		cervical vertebrae 5 (c5)	Dice	0.8399 (0.8209–0.8580)	0.1405 (0.06912–0.2270)	0.8192 (0.7831–0.8459)	0.8915 (0.8656–0.9122)	0.8573 (0.8495–0.8897)	0.4128 (0.3164–0.5123)	0.8583 (0.8414–0.8743)
			HD95	4.731 (0.2658–12.78)	133.3 (67.00–217.0)	2.358 (1.907–2.972)	2.888 (1.794–3.988)	6.781 (6.073–10.34)	8.910 (6.957–10.56)	2.837 (1.964–3.955)
		cervical vertebrae 6 (c6)	Dice	0.8575 (0.8483–0.8664)	0.04104 (0.009703–0.07959)	0.8371 (0.8228–0.8502)	0.8789 (0.8536–0.9009)	0.8186 (0.8082–0.8266)	0.8555 (0.8374–0.8711)	0.8531 (0.8352–0.8709)
			HD95	2.686 (2.464–2.959)	161.6 (101.2–236.2)	2.186 (1.893–2.528)	9.161 (2.019–22.28)	3.583 (3.552–6.216)	2.186 (1.785–2.679)	2.743 (2.193–3.441)
		cervical vertebrae 7 (c7)	Dice	0.8497 (0.8136–0.8729)	0.006017 (0.0001674–0.01519)	0.8579 (0.8469–0.8680)	0.8817 (0.8535–0.9036)	0.8247 (0.7978–0.8322)	0.8883 (0.8793–0.8968)	0.8869 (0.8774–0.8955)
			HD95	3.746 (2.675–5.184)	172.7 (130.4–227.5)	2.357 (1.930–2.875)	4.047 (2.819–5.444)	6.017 (5.307–9.112)	3.415 (2.624–4.274)	2.756 (2.200–3.424)
		thoracic vertebrae 1 (t1)	Dice	0.8354 (0.8056–0.8597)	0.09463 (0.05570–0.1328)	0.8784 (0.8690–0.8864)	0.9056 (0.8880–0.9208)	0.8035 (0.7923–0.8118)	0.9109 (0.9038–0.9178)	0.9074 (0.9029–0.9121)
			HD95	27.29 (20.06–35.09)	133.1 (76.78–199.4)	2.555 (1.743–3.650)	17.74 (4.209–43.51)	7.411 (5.121–11.52)	1.969 (1.454–2.593)	1.939 (1.517–2.629)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	VerSe	thoracic vertebrae 2 (t2)	Dice	0.8617 (0.8251–0.8890)	0.1196 (0.07368–0.1690)	0.8786 (0.8646–0.8893)	0.8730 (0.8373–0.9037)	0.8367 (0.8271–0.8380)	0.8744 (0.8431–0.8949)	0.9051 (0.8861–0.9163)
			HD95	14.36 (3.476–35.35)	125.0 (75.95–178.8)	2.919 (2.010–3.974)	17.91 (5.456–40.78)	5.217 (4.842–6.189)	2.887 (2.116–3.852)	1.847 (1.507–2.347)
		thoracic vertebrae 3 (t3)	Dice	0.8411 (0.7943–0.8775)	0.2320 (0.1647–0.3043)	0.8538 (0.8158–0.8820)	0.8635 (0.8246–0.8976)	0.8417 (0.8078–0.8764)	0.8648 (0.8212–0.8916)	0.8923 (0.8603–0.9092)
			HD95	6.925 (4.805–9.191)	111.4 (59.10–170.5)	3.415 (2.031–5.133)	10.90 (6.874–15.44)	4.667 (2.294–6.346)	8.124 (6.242–10.22)	3.038 (1.734–5.154)
		thoracic vertebrae 4 (t4)	Dice	0.8465 (0.8049–0.8768)	0.2668 (0.1859–0.3450)	0.8326 (0.7698–0.8804)	0.8842 (0.8434–0.9140)	0.7800 (0.7352–0.8093)	0.7039 (0.6738–0.7342)	0.8960 (0.8824–0.9067)
			HD95	6.357 (4.233–8.729)	105.5 (53.64–162.7)	32.51 (2.937–77.49)	5.874 (3.226–9.008)	9.235 (6.495–12.27)	45.52 (27.57–66.34)	2.612 (1.867–3.557)
		thoracic vertebrae 5 (t5)	Dice	0.8307 (0.7784–0.8728)	0.2880 (0.2106–0.3669)	0.8052 (0.7185–0.8741)	0.8822 (0.8378–0.9152)	0.7954 (0.7593–0.8337)	0.5814 (0.4934–0.6747)	0.9015 (0.8888–0.9118)
			HD95	18.11 (4.653–42.24)	101.1 (51.75–157.7)	4.197 (2.398–6.371)	5.758 (3.277–9.028)	10.34 (6.452–12.78)	8.838 (6.613–10.93)	2.359 (1.727–3.342)
		thoracic vertebrae 6 (t6)	Dice	0.8298 (0.7593–0.8815)	0.2813 (0.2040–0.3603)	0.7896 (0.6966–0.8640)	0.8697 (0.8082–0.9150)	0.7830 (0.7399–0.7974)	0.8806 (0.8577–0.8982)	0.8981 (0.8807–0.9101)
			HD95	5.911 (3.796–8.354)	86.21 (42.69–133.0)	4.865 (2.730–7.392)	6.409 (3.694–9.568)	13.37 (10.51–16.65)	3.182 (2.557–3.824)	2.443 (1.752–3.518)
		thoracic vertebrae 7 (t7)	Dice	0.8324 (0.7669–0.8829)	0.2895 (0.2084–0.3664)	0.7844 (0.6869–0.8657)	0.8768 (0.8277–0.9148)	0.8891 (0.8690–0.9356)	0.9177 (0.9021–0.9282)	0.9133 (0.9065–0.9197)
			HD95	6.512 (3.999–9.233)	72.59 (40.87–111.0)	4.996 (2.924–7.291)	7.826 (5.149–10.69)	4.881 (3.955–7.530)	3.242 (1.769–5.197)	3.176 (2.397–4.106)
		thoracic vertebrae 8 (t8)	Dice	0.8025 (0.7210–0.8730)	0.2390 (0.1617–0.3251)	0.7822 (0.6868–0.8687)	0.8681 (0.8150–0.9099)	0.8433 (0.7904–0.8812)	0.8936 (0.8791–0.9056)	0.9134 (0.9054–0.9204)
			HD95	16.07 (5.108–35.83)	91.69 (56.01–128.7)	5.960 (3.441–8.595)	9.761 (6.781–13.07)	6.391 (4.264–7.508)	3.453 (2.756–4.286)	5.142 (3.227–7.179)
		thoracic vertebrae 9 (t9)	Dice	0.8048 (0.7238–0.8771)	0.2446 (0.1892–0.3000)	0.7991 (0.6966–0.8827)	0.8562 (0.7798–0.9152)	0.8221 (0.7723–0.8704)	0.9270 (0.9169–0.9343)	0.9181 (0.9073–0.9258)
			HD95	13.45 (5.108–27.42)	75.94 (48.41–105.8)	5.358 (3.098–7.733)	8.198 (5.391–11.16)	5.165 (2.669–6.414)	5.091 (2.856–7.617)	2.821 (1.920–4.085)
		thoracic vertebrae 10 (t10)	Dice	0.7927 (0.7085–0.8617)	0.2990 (0.2427–0.3527)	0.8202 (0.7366–0.8892)	0.8601 (0.7967–0.9110)	0.8534 (0.8095–0.8867)	0.8884 (0.8676–0.9068)	0.9189 (0.9096–0.9270)
			HD95	6.741 (4.750–9.258)	74.76 (49.98–104.2)	4.585 (2.740–6.643)	8.732 (5.985–11.96)	5.913 (5.138–8.030)	9.530 (7.580–11.57)	3.539 (2.322–4.990)
		thoracic vertebrae 11 (t11)	Dice	0.8162 (0.7406–0.8785)	0.2560 (0.2064–0.3071)	0.8385 (0.7671–0.8974)	0.8796 (0.8167–0.9236)	0.8322 (0.7896–0.8690)	0.8852 (0.8478–0.9149)	0.9148 (0.8849–0.9342)
			HD95	7.269 (4.836–10.05)	62.47 (47.60–78.31)	4.523 (2.687–6.648)	7.459 (4.451–10.24)	7.065 (2.731–8.076)	8.917 (3.837–16.54)	2.724 (1.803–3.819)
		thoracic vertebrae 12 (t12)	Dice	0.8597 (0.8056–0.9019)	0.09522 (0.05695–0.1374)	0.8930 (0.8500–0.9194)	0.9194 (0.8883–0.9444)	0.7955 (0.7721–0.8127)	0.9113 (0.8876–0.9289)	0.9241 (0.9009–0.9388)
			HD95	7.798 (5.044–10.99)	83.32 (67.01–102.8)	18.10 (3.406–40.09)	4.843 (2.572–7.825)	16.04 (9.603–23.31)	6.373 (4.590–8.481)	2.906 (1.622–4.798)
		lumbar vertebrae 1 (l1)	Dice	0.8593 (0.8110–0.8950)	0.1364 (0.09955–0.1758)	0.8828 (0.8461–0.9110)	0.9295 (0.9076–0.9464)	0.8725 (0.8526–0.8905)	0.8938 (0.8587–0.9235)	0.9326 (0.9194–0.9419)
			HD95	9.023 (6.015–12.24)	69.96 (60.72–80.93)	17.58 (4.458–35.97)	6.275 (3.540–9.527)	10.35 (7.565–13.58)	13.75 (6.306–23.10)	3.263 (1.944–4.841)
		lumbar vertebrae 2 (l2)	Dice	0.8143 (0.7532–0.8681)	0.2809 (0.2307–0.3334)	0.8230 (0.7543–0.8771)	0.9197 (0.8935–0.9402)	0.8036 (0.7748–0.8171)	0.8598 (0.8081–0.8954)	0.9318 (0.9198–0.9400)
			HD95	10.11 (6.875–13.33)	51.11 (42.19–62.19)	19.94 (5.825–39.06)	7.195 (4.066–11.92)	10.47 (10.42–15.27)	9.990 (7.908–12.41)	6.684 (2.035–15.71)
		lumbar vertebrae 3 (l3)	Dice	0.7412 (0.6574–0.8137)	0.3597 (0.2983–0.4175)	0.7539 (0.6777–0.8299)	0.8838 (0.8380–0.9221)	0.8802 (0.8304–0.9163)	0.8818 (0.8525–0.9064)	0.9239 (0.9132–0.9325)
			HD95	12.13 (8.842–15.98)	48.05 (38.38–58.63)	31.31 (11.51–57.54)	10.82 (6.012–18.49)	8.910 (6.393–11.83)	14.70 (12.06–17.31)	5.772 (4.068–7.568)
		lumbar vertebrae 4 (l4)	Dice	0.6976 (0.6109–0.7812)	0.3387 (0.2783–0.3921)	0.6898 (0.5936–0.7707)	0.8494 (0.7950–0.8973)	0.7697 (0.7269–0.8140)	0.7975 (0.7503–0.8393)	0.9169 (0.9006–0.9298)
			HD95	149.6 (101.8–196.6)	56.24 (43.52–70.70)	40.16 (15.84–72.41)	13.83 (8.509–21.71)	16.19 (13.79–17.36)	47.61 (24.76–77.16)	5.735 (4.079–7.662)
		lumbar vertebrae 5 (l5)	Dice	0.7304 (0.6466–0.8073)	0.2154 (0.1603–0.2751)	0.6830 (0.5935–0.7738)	0.8191 (0.7727–0.8596)	0.7591 (0.7004–0.8000)	0.8089 (0.7501–0.8666)	0.8581 (0.8293–0.8843)
			HD95	12.68 (9.305–15.90)	67.52 (51.32–83.28)	11.21 (7.845–14.88)	17.06 (12.02–24.12)	15.45 (11.68–22.11)	12.89 (9.221–17.24)	10.16 (7.465–13.05)
		lumbar vertebrae 6 (l6)	Dice	Not supported	0.02595 (0.0000–0.06496)	0.5369 (0.4201–0.6494)	0.07041 (0.0000–0.1781)	0.4742 (0.4334–0.5082)	0.8005 (0.7014–0.8828)	0.8047 (0.7585–0.8385)
			HD95		220.2 (174.1–264.6)	19.36 (16.67–21.89)	19.10 (18.56–19.63)	41.88 (33.45–49.79)	12.47 (8.074–17.32)	21.45 (18.71–23.77)
	Total-SegmentatorV2	spleen	Dice	0.9526 (0.9470–0.9577)	0.7209 (0.6897–0.7494)	0.9369 (0.9172–0.9527)	0.9649 (0.9578–0.9702)	0.9643 (0.9642–0.9772)	0.9691 (0.9593–0.9779)	0.9657 (0.9596–0.9710)
			HD95	20.33 (9.383–33.11)	23.89 (17.68–32.07)	2.988 (2.186–3.893)	3.014 (2.161–4.126)	4.934 (4.846–5.274)	3.500 (1.738–6.533)	2.501 (1.941–3.220)
		right kidney	Dice	0.9187 (0.8958–0.9370)	0.6254 (0.5895–0.6588)	0.9088 (0.8860–0.9300)	0.9356 (0.9117–0.9554)	0.9430 (0.9226–0.9668)	0.9352 (0.9098–0.9581)	0.9201 (0.8963–0.9414)
			HD95	11.13 (5.069–19.07)	48.52 (39.08–58.62)	2.366 (1.977–2.774)	2.583 (1.982–3.318)	3.493 (2.839–3.970)	5.282 (2.173–10.27)	3.957 (3.089–4.900)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	Total-SegmentatorV2	left kidney	Dice	0.9063 (0.8749–0.9335)	0.6382 (0.6000–0.6727)	0.9168 (0.8909–0.9368)	0.9185 (0.8847–0.9474)	0.9211 (0.8985–0.9327)	0.9281 (0.8957–0.9549)	0.9324 (0.9090–0.9525)
			HD95	4.359 (3.384–5.442)	46.40 (36.99–56.72)	2.442 (1.836–3.414)	3.221 (2.020–4.724)	2.859 (0.4540–3.014)	9.600 (4.172–16.28)	2.998 (2.228–3.827)
		gallbladder	Dice	0.8171 (0.7730–0.8575)	0.3347 (0.2908–0.3806)	0.8209 (0.7848–0.8560)	0.8042 (0.7566–0.8477)	0.8406 (0.7956–0.8817)	0.8113 (0.7616–0.8539)	0.8131 (0.7653–0.8530)
			HD95	8.566 (5.904–12.67)	40.53 (30.70–51.73)	7.975 (3.897–15.76)	5.802 (4.418–7.529)	6.723 (3.888–9.440)	9.292 (5.804–14.58)	6.153 (4.756–7.936)
		liver	Dice	0.9503 (0.9341–0.9626)	0.8167 (0.7877–0.8394)	0.9535 (0.9370–0.9655)	0.9568 (0.9371–0.9711)	0.9618 (0.9235–0.9632)	0.9619 (0.9428–0.9791)	0.9539 (0.9367–0.9689)
			HD95	14.37 (5.884–24.77)	16.94 (13.96–20.39)	3.326 (2.698–4.383)	3.539 (3.164–3.971)	4.306 (1.958–6.752)	5.986 (3.033–10.58)	4.018 (3.101–5.252)
		stomach	Dice	0.9173 (0.8972–0.9345)	0.6204 (0.5871–0.6514)	0.8805 (0.8565–0.9029)	0.9190 (0.8977–0.9363)	0.9372 (0.9287–0.9646)	0.9319 (0.9110–0.9490)	0.9262 (0.9091–0.9402)
			HD95	9.378 (6.074–13.42)	26.21 (21.18–31.87)	41.35 (23.07–61.45)	6.047 (5.005–7.271)	3.029 (2.718–3.568)	6.361 (4.796–8.532)	13.70 (7.073–21.46)
		pancreas	Dice	0.8460 (0.8216–0.8675)	0.4827 (0.4510–0.5130)	0.8412 (0.8171–0.8615)	0.8395 (0.8136–0.8637)	0.8608 (0.8304–0.9013)	0.8535 (0.8277–0.8770)	0.8587 (0.8378–0.8784)
			HD95	6.937 (5.505–8.669)	23.37 (18.65–28.76)	3.959 (3.327–4.752)	5.938 (5.006–7.076)	4.553 (1.444–4.613)	4.918 (3.757–6.341)	5.570 (4.056–7.724)
		right adrenal gland	Dice	0.8503 (0.8290–0.8691)	0.2793 (0.2579–0.3021)	0.7947 (0.7745–0.8105)	0.8186 (0.7957–0.8375)	0.8747 (0.8594–0.8877)	0.8089 (0.7876–0.8288)	0.8231 (0.7982–0.8424)
			HD95	3.409 (2.992–3.980)	22.64 (16.34–29.55)	2.093 (1.863–2.382)	3.114 (2.583–3.729)	4.608 (3.277–5.360)	10.92 (3.328–20.61)	3.443 (2.537–4.523)
		left adrenal gland	Dice	0.8603 (0.8397–0.8788)	0.2709 (0.2462–0.2930)	0.8054 (0.7869–0.8221)	0.8278 (0.8064–0.8468)	0.8650 (0.8467–0.8918)	0.8345 (0.8134–0.8537)	0.8290 (0.8077–0.8482)
			HD95	3.493 (3.067–4.006)	30.63 (22.70–39.61)	2.186 (1.901–2.598)	3.010 (2.612–3.441)	2.184 (1.826–2.352)	6.072 (2.203–13.64)	5.112 (3.465–7.134)
		left lung upper lobe	Dice	0.9396 (0.9293–0.9498)	0.6772 (0.6416–0.7147)	0.9204 (0.9070–0.9322)	0.9306 (0.9180–0.9431)	0.9387 (0.9315–0.9586)	0.9163 (0.9023–0.9291)	0.9148 (0.9013–0.9267)
			HD95	8.872 (6.606–11.73)	27.11 (22.96–31.53)	7.215 (5.484–9.370)	9.149 (7.246–11.80)	10.49 (7.197–12.85)	24.51 (21.43–27.73)	9.551 (7.918–11.59)
		left lung lower lobe	Dice	0.9261 (0.9074–0.9428)	0.7189 (0.6870–0.7484)	0.9102 (0.8907–0.9276)	0.9160 (0.8952–0.9340)	0.9504 (0.9302–0.9557)	0.8837 (0.8594–0.9063)	0.9028 (0.8817–0.9223)
			HD95	6.129 (5.054–7.231)	30.15 (25.44–35.55)	5.131 (4.369–5.995)	7.252 (6.103–8.605)	3.846 (2.143–5.269)	24.01 (20.34–28.03)	14.07 (9.883–19.30)
		right lung upper lobe	Dice	0.8703 (0.8337–0.8997)	0.6295 (0.5833–0.6744)	0.8485 (0.8095–0.8832)	0.8492 (0.8088–0.8903)	0.9158 (0.8830–0.9362)	0.8290 (0.7874–0.8709)	0.8297 (0.7884–0.8703)
			HD95	14.71 (9.192–21.21)	38.67 (32.10–46.07)	6.893 (5.135–9.001)	8.079 (6.242–10.52)	4.145 (2.326–5.878)	20.26 (15.20–26.26)	21.97 (15.86–28.74)
		right lung middle lobe	Dice	0.9037 (0.8802–0.9232)	0.5085 (0.4650–0.5499)	0.8648 (0.8353–0.8906)	0.8839 (0.8581–0.9049)	0.9423 (0.9115–0.9709)	0.8320 (0.8025–0.8578)	0.8364 (0.8075–0.8610)
			HD95	16.06 (7.463–30.13)	33.14 (27.46–39.37)	7.163 (5.844–8.795)	8.752 (7.440–10.46)	4.376 (2.266–6.330)	14.59 (12.22–17.51)	13.97 (11.33–17.55)
		right lung lower lobe	Dice	0.9293 (0.9097–0.9464)	0.7127 (0.6785–0.7443)	0.9050 (0.8811–0.9257)	0.9164 (0.8908–0.9367)	0.9547 (0.9353–0.9744)	0.8932 (0.8684–0.9141)	0.8932 (0.8694–0.9136)
			HD95	13.94 (5.646–28.32)	28.31 (23.68–33.07)	5.976 (5.049–6.978)	6.604 (5.380–7.986)	6.800 (4.668–9.140)	25.04 (22.59–27.54)	13.96 (11.86–16.27)
		esophagus	Dice	0.8906 (0.8756–0.9022)	0.5364 (0.5143–0.5556)	0.8684 (0.8543–0.8791)	0.8650 (0.8532–0.8754)	0.9172 (0.8927–0.9256)	0.8738 (0.8605–0.8852)	0.8730 (0.8592–0.8840)
			HD95	15.36 (8.887–23.35)	10.89 (9.337–12.77)	2.292 (2.090–2.540)	2.707 (2.405–3.060)	3.253 (1.603–4.305)	3.327 (2.442–4.686)	2.635 (2.373–2.939)
		trachea	Dice	0.9275 (0.9200–0.9342)	0.6449 (0.6188–0.6711)	0.9048 (0.8938–0.9138)	0.9395 (0.9305–0.9466)	0.9434 (0.9333–0.9472)	0.9135 (0.9017–0.9233)	0.9096 (0.8999–0.9178)
			HD95	4.196 (3.270–5.272)	12.07 (10.52–13.95)	4.798 (3.238–6.581)	4.647 (2.915–6.988)	4.848 (4.054–5.435)	10.22 (7.917–12.51)	7.672 (5.998–9.512)
		thyroid gland	Dice	0.8486 (0.8265–0.8667)	0.0000 (0.0000–0.0000)	0.8315 (0.8152–0.8465)	0.8524 (0.8328–0.8679)	0.8817 (0.8497–0.8948)	0.8555 (0.8338–0.8730)	0.8436 (0.8243–0.8606)
			HD95	4.314 (3.850–4.876)	215.6 (201.2–229.6)	3.028 (2.506–3.784)	2.916 (2.584–3.240)	4.421 (4.421–5.321)	2.647 (2.246–3.131)	5.460 (3.077–9.717)
		small bowel	Dice	0.8176 (0.7922–0.8417)	0.5136 (0.4828–0.5463)	0.8294 (0.8061–0.8501)	0.8030 (0.7739–0.8320)	0.8649 (0.8367–0.9015)	0.8168 (0.7859–0.8412)	0.8181 (0.7891–0.8416)
			HD95	16.68 (11.24–22.78)	29.45 (23.56–37.83)	8.517 (7.119–10.11)	12.66 (10.47–15.48)	7.870 (6.470–9.740)	20.48 (13.54–28.48)	15.99 (9.938–23.66)
		duodenum	Dice	0.8087 (0.7854–0.8304)	0.4202 (0.3898–0.4499)	0.8003 (0.7762–0.8211)	0.7696 (0.7389–0.7964)	0.8076 (0.7998–0.8081)	0.7698 (0.7439–0.7939)	0.7797 (0.7559–0.8027)
			HD95	9.475 (7.470–12.11)	30.06 (24.28–36.83)	9.347 (5.518–14.95)	10.93 (8.443–13.77)	10.27 (7.463–14.29)	10.17 (8.252–12.75)	9.418 (7.678–11.86)
		colon	Dice	0.8827 (0.8696–0.8932)	0.5392 (0.5115–0.5653)	0.8624 (0.8510–0.8726)	0.8482 (0.8359–0.8597)	0.8770 (0.8727–0.9047)	0.8386 (0.8211–0.8544)	0.8562 (0.8447–0.8665)
			HD95	15.74 (11.10–22.29)	33.11 (27.21–40.25)	14.61 (7.962–26.42)	18.67 (14.87–24.06)	14.30 (9.280–21.50)	28.44 (19.64–38.74)	23.79 (15.37–35.11)
		urinary bladder	Dice	0.8851 (0.8568–0.9071)	0.5290 (0.4812–0.5779)	0.8839 (0.8590–0.9061)	0.8838 (0.8506–0.9070)	0.9019 (0.8731–0.9371)	0.8739 (0.8437–0.8974)	0.8900 (0.8621–0.9104)
			HD95	49.39 (28.46–74.91)	35.81 (26.00–46.56)	8.932 (4.121–17.69)	6.262 (5.064–7.712)	7.030 (5.279–9.140)	9.550 (6.288–13.43)	7.924 (5.331–11.28)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	Total-SegmentatorV2	prostate	Dice	0.7170 (0.6548–0.7757)	0.1789 (0.1333–0.2293)	0.7621 (0.6851–0.8319)	0.7330 (0.6600–0.7933)	0.7486 (0.6963–0.8186)	0.7427 (0.6752–0.8011)	0.7255 (0.6623–0.7831)
			HD95	22.65 (18.30–27.44)	50.83 (41.34–60.90)	5.539 (3.876–7.526)	5.944 (4.733–7.368)	8.427 (7.057–9.650)	20.06 (9.117–34.89)	71.86 (46.64–100.8)
		left kidney cyst	Dice	0.6743 (0.4239–0.8632)	0.1251 (0.01252–0.2704)	0.8678 (0.8259–0.8992)	0.0000 (0.0000–0.0000)	0.8544 (0.8083–0.8807)	0.9079 (0.8712–0.9380)	0.8763 (0.8345–0.9168)
			HD95	56.57 (5.137–152.9)	71.05 (47.56–90.74)	68.36 (18.89–122.3)	– (– –)	5.811 (3.895–7.440)	41.05 (2.765–96.98)	21.44 (4.177–52.10)
		right kidney cyst	Dice	0.5917 (0.3232–0.7966)	0.1737 (0.0000–0.3921)	0.8106 (0.7687–0.8515)	0.0000 (0.0000–0.0000)	0.8449 (0.8440–0.8484)	0.7168 (0.4366–0.8904)	0.7062 (0.4024–0.8871)
			HD95	48.20 (7.517–126.9)	58.37 (27.66–94.05)	4.709 (3.132–6.527)	– (– –)	4.541 (2.121–6.424)	11.97 (5.290–21.72)	4.819 (3.371–6.919)
		sacrum	Dice	0.8684 (0.8401–0.8901)	0.5202 (0.4930–0.5450)	0.8049 (0.7858–0.8192)	0.8985 (0.8787–0.9133)	0.9345 (0.9324–0.9516)	0.8756 (0.8451–0.8992)	0.8927 (0.8710–0.9084)
			HD95	21.91 (17.93–25.66)	16.04 (13.25–19.39)	18.88 (7.670–37.37)	6.703 (5.313–8.410)	3.533 (1.258–6.300)	81.49 (51.69–110.1)	14.30 (5.737–30.27)
		sacral vertebrae 1 (s1)	Dice	0.8861 (0.8585–0.9092)	0.01010 (0.002950–0.01869)	0.8742 (0.8446–0.8974)	0.8548 (0.8279–0.8756)	0.9076 (0.8774–0.9197)	0.8280 (0.7950–0.8552)	0.8540 (0.8221–0.8806)
			HD95	66.00 (38.72–98.64)	61.98 (56.12–70.77)	9.882 (2.528–21.13)	6.009 (5.073–7.164)	6.168 (4.305–7.560)	72.40 (46.20–99.45)	9.667 (4.957–18.44)
		lumbar vertebrae 5 (l5)	Dice	0.9017 (0.8668–0.9304)	0.1649 (0.1275–0.2005)	0.8837 (0.8516–0.9100)	0.9142 (0.8821–0.9408)	0.9341 (0.9132–0.9660)	0.9096 (0.8788–0.9348)	0.9097 (0.8795–0.9332)
			HD95	16.81 (4.933–35.50)	37.64 (34.33–40.83)	22.79 (9.069–40.23)	4.696 (3.315–6.248)	2.850 (1.119–4.014)	12.93 (6.979–20.51)	4.655 (3.465–6.098)
		lumbar vertebrae 4 (l4)	Dice	0.8570 (0.8161–0.8918)	0.3136 (0.2731–0.3571)	0.8746 (0.8410–0.9032)	0.9053 (0.8665–0.9380)	0.9328 (0.8959–0.9800)	0.8487 (0.8129–0.8802)	0.9011 (0.8684–0.9292)
			HD95	19.39 (16.04–22.81)	31.03 (25.41–38.56)	38.13 (21.16–60.28)	4.183 (2.950–5.630)	2.594 (1.176–3.897)	47.39 (29.38–68.16)	4.785 (3.288–6.810)
		lumbar vertebrae 3 (l3)	Dice	0.9124 (0.8848–0.9353)	0.3852 (0.3450–0.4255)	0.8955 (0.8711–0.9147)	0.9193 (0.8898–0.9447)	0.9297 (0.9173–0.9570)	0.8855 (0.8522–0.9149)	0.9058 (0.8766–0.9291)
			HD95	18.08 (6.349–35.85)	29.26 (21.89–38.65)	15.55 (4.717–28.72)	2.899 (2.219–3.803)	3.219 (3.007–4.017)	20.86 (13.03–29.76)	7.629 (5.062–11.33)
		lumbar vertebrae 2 (l2)	Dice	0.9256 (0.9056–0.9425)	0.4017 (0.3715–0.4352)	0.9122 (0.9027–0.9202)	0.9427 (0.9280–0.9548)	0.9297 (0.9106–0.9554)	0.8699 (0.8455–0.8892)	0.9362 (0.9247–0.9461)
			HD95	21.63 (6.402–40.86)	24.65 (20.70–30.59)	2.082 (1.758–2.438)	3.447 (2.552–4.633)	2.215 (2.186–2.587)	10.84 (9.580–12.35)	4.079 (3.026–5.240)
		lumbar vertebrae 1 (l1)	Dice	0.9127 (0.8865–0.9376)	0.1864 (0.1552–0.2161)	0.9061 (0.8932–0.9174)	0.9255 (0.8993–0.9476)	0.9124 (0.8950–0.9340)	0.8953 (0.8667–0.9190)	0.9258 (0.9025–0.9437)
			HD95	4.818 (3.769–5.910)	36.97 (32.43–42.33)	4.661 (1.858–9.745)	3.171 (2.309–4.191)	5.824 (5.493–6.007)	27.06 (19.02–35.31)	7.480 (3.440–12.65)
		thoracic vertebrae 12 (t12)	Dice	0.8918 (0.8593–0.9201)	0.08564 (0.06386–0.1105)	0.8934 (0.8703–0.9131)	0.9166 (0.8863–0.9420)	0.9108 (0.8860–0.9306)	0.8933 (0.8686–0.9132)	0.9137 (0.8860–0.9373)
			HD95	25.77 (15.53–36.85)	43.82 (39.77–48.06)	12.88 (4.505–22.98)	3.429 (2.501–4.478)	4.629 (4.603–5.073)	9.650 (7.169–13.39)	4.522 (2.597–8.145)
		thoracic vertebrae 11 (t11)	Dice	0.8896 (0.8597–0.9177)	0.2960 (0.2605–0.3308)	0.8991 (0.8829–0.9131)	0.9238 (0.8997–0.9428)	0.9209 (0.8847–0.9390)	0.8907 (0.8626–0.9155)	0.9241 (0.9069–0.9379)
			HD95	7.373 (4.878–11.64)	30.35 (27.38–33.61)	2.360 (1.932–2.817)	3.148 (2.421–3.933)	2.379 (2.253–3.302)	6.331 (4.517–8.720)	3.653 (2.798–4.582)
		thoracic vertebrae 10 (t10)	Dice	0.8898 (0.8607–0.9162)	0.4046 (0.3680–0.4375)	0.8905 (0.8733–0.9060)	0.9229 (0.8996–0.9436)	0.9166 (0.9039–0.9211)	0.8634 (0.8392–0.8842)	0.9055 (0.8864–0.9231)
			HD95	5.460 (4.485–6.663)	27.43 (23.45–32.07)	2.313 (1.892–2.786)	3.120 (2.467–3.891)	4.172 (2.796–5.120)	12.74 (10.99–15.04)	6.774 (5.110–9.103)
		thoracic vertebrae 9 (t9)	Dice	0.8856 (0.8545–0.9132)	0.2593 (0.2284–0.2892)	0.8765 (0.8577–0.8924)	0.9189 (0.8917–0.9418)	0.9160 (0.8920–0.9329)	0.9055 (0.8889–0.9205)	0.9080 (0.8882–0.9247)
			HD95	10.23 (6.109–15.69)	35.17 (31.02–40.14)	2.817 (2.192–3.485)	3.198 (2.520–3.980)	1.981 (0.5620–2.236)	7.350 (5.874–8.921)	5.678 (4.354–7.152)
		thoracic vertebrae 8 (t8)	Dice	0.8687 (0.8325–0.8987)	0.3180 (0.2843–0.3527)	0.8504 (0.8240–0.8755)	0.8924 (0.8591–0.9164)	0.8977 (0.8968–0.9432)	0.7506 (0.7005–0.7984)	0.8797 (0.8514–0.9043)
			HD95	21.68 (10.36–36.99)	50.74 (43.02–59.24)	2.885 (2.344–3.546)	3.870 (2.941–5.028)	4.166 (3.285–7.583)	8.120 (6.468–9.970)	7.874 (6.643–9.155)
		thoracic vertebrae 7 (t7)	Dice	0.8440 (0.8099–0.8763)	0.3718 (0.3310–0.4113)	0.8215 (0.7876–0.8510)	0.8420 (0.8014–0.8815)	0.8908 (0.8665–0.9242)	0.7980 (0.7522–0.8395)	0.8427 (0.8069–0.8773)
			HD95	6.841 (5.543–8.243)	32.16 (25.79–38.78)	3.890 (2.761–5.129)	4.317 (3.329–5.525)	3.552 (2.035–4.949)	7.998 (5.360–12.06)	6.014 (4.862–7.371)
		thoracic vertebrae 6 (t6)	Dice	0.8751 (0.8448–0.9051)	0.4224 (0.3891–0.4551)	0.8438 (0.8175–0.8671)	0.8852 (0.8528–0.9120)	0.8975 (0.8837–0.9103)	0.8336 (0.8011–0.8642)	0.8729 (0.8497–0.8950)
			HD95	5.367 (4.561–6.349)	24.56 (19.69–30.17)	2.766 (2.272–3.333)	3.621 (2.884–4.401)	4.317 (3.995–6.186)	5.229 (4.116–6.511)	5.078 (3.880–6.405)
		thoracic vertebrae 5 (t5)	Dice	0.8807 (0.8499–0.9068)	0.3973 (0.3634–0.4311)	0.8428 (0.8143–0.8679)	0.8918 (0.8601–0.9160)	0.9272 (0.9060–0.9322)	0.6285 (0.5780–0.6729)	0.8840 (0.8570–0.9074)
			HD95	8.177 (4.774–14.26)	17.35 (14.89–20.57)	3.457 (2.599–4.545)	3.439 (2.758–4.177)	3.864 (2.921–4.332)	7.866 (6.884–8.860)	3.877 (3.256–4.624)
		thoracic vertebrae 4 (t4)	Dice	0.8743 (0.8390–0.9037)	0.4247 (0.3952–0.4539)	0.8604 (0.8357–0.8802)	0.8982 (0.8674–0.9240)	0.9135 (0.9006–0.9155)	0.7150 (0.6919–0.7370)	0.8950 (0.8719–0.9129)
			HD95	5.581 (4.578–6.719)	13.78 (12.81–14.81)	3.297 (2.412–4.467)	3.305 (2.629–4.180)	5.004 (2.390–6.671)	54.52 (45.94–62.56)	3.726 (2.987–4.528)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	Total-SegmentatorV2	thoracic vertebrae 3 (t3)	Dice	0.8828 (0.8501–0.9143)	0.3902 (0.3644–0.4184)	0.8526 (0.8257–0.8763)	0.8973 (0.8662–0.9248)	0.9278 (0.9145–0.9617)	0.8738 (0.8498–0.8951)	0.9045 (0.8824–0.9235)
			HD95	5.508 (4.435–6.844)	15.51 (14.16–16.95)	2.545 (2.071–3.086)	3.007 (2.394–3.694)	4.991 (4.373–5.252)	12.78 (9.550–16.84)	4.168 (2.888–6.235)
		thoracic vertebrae 2 (t2)	Dice	0.9087 (0.8797–0.9315)	0.2012 (0.1733–0.2302)	0.8694 (0.8520–0.8861)	0.9052 (0.8746–0.9334)	0.9176 (0.8927–0.9360)	0.8744 (0.8482–0.8955)	0.9215 (0.9089–0.9328)
			HD95	21.08 (10.39–34.75)	22.01 (20.01–24.56)	2.232 (1.848–2.735)	2.415 (1.912–3.033)	4.266 (3.180–4.435)	3.338 (2.635–4.215)	2.547 (2.091–3.055)
		thoracic vertebrae 1 (t1)	Dice	0.9060 (0.8852–0.9249)	0.2260 (0.1997–0.2535)	0.8711 (0.8513–0.8863)	0.9254 (0.9047–0.9429)	0.9455 (0.9337–0.9542)	0.9170 (0.8970–0.9312)	0.9236 (0.9125–0.9323)
			HD95	16.02 (12.53–19.54)	23.63 (20.72–26.82)	2.144 (1.834–2.498)	2.099 (1.688–2.624)	2.185 (0.8640–3.492)	2.365 (1.823–2.946)	2.277 (1.918–2.734)
		cervical vertebrae 7 (c7)	Dice	0.9039 (0.8795–0.9226)	0.0000 (0.0000–0.0000)	0.8404 (0.8195–0.8581)	0.8948 (0.8690–0.9177)	0.9466 (0.9396–0.9834)	0.8722 (0.8451–0.8938)	0.8963 (0.8731–0.9152)
			HD95	3.346 (2.878–4.001)	157.6 (150.9–164.3)	2.354 (2.023–2.835)	2.514 (2.051–3.143)	1.624 (1.548–2.010)	7.425 (4.791–11.81)	3.837 (2.644–5.489)
		cervical vertebrae 6 (c6)	Dice	0.8796 (0.8526–0.9021)	0.02665 (0.01215–0.04564)	0.7637 (0.7338–0.7907)	0.8124 (0.7675–0.8553)	0.8432 (0.8130–0.8816)	0.8168 (0.7832–0.8490)	0.8336 (0.8007–0.8626)
			HD95	4.042 (3.208–5.004)	127.2 (117.9–136.5)	2.839 (2.313–3.630)	3.529 (2.633–4.577)	2.708 (1.588–3.102)	4.358 (3.343–5.515)	6.453 (2.940–12.69)
		cervical vertebrae 5 (c5)	Dice	0.8064 (0.7467–0.8568)	0.05967 (0.03120–0.09090)	0.7435 (0.6931–0.7904)	0.7841 (0.7166–0.8456)	0.8496 (0.8151–0.8707)	0.4599 (0.4001–0.5181)	0.8142 (0.7626–0.8564)
			HD95	10.01 (5.684–14.33)	62.68 (53.37–73.14)	2.986 (2.227–4.416)	2.218 (1.861–2.694)	2.542 (2.376–5.631)	7.871 (6.975–8.897)	3.146 (2.752–3.592)
		cervical vertebrae 4 (c4)	Dice	0.8866 (0.8432–0.9170)	0.1273 (0.08531–0.1720)	0.8180 (0.7798–0.8468)	0.8967 (0.8609–0.9223)	0.8613 (0.8346–0.8795)	0.5396 (0.5073–0.5679)	0.8642 (0.8224–0.8976)
			HD95	29.37 (11.62–50.15)	32.37 (25.87–38.83)	2.454 (1.992–3.170)	2.031 (1.746–2.387)	3.803 (3.439–4.239)	100.8 (65.56–135.8)	2.597 (2.128–3.162)
		cervical vertebrae 3 (c3)	Dice	0.8976 (0.8618–0.9244)	0.03431 (0.006584–0.06950)	0.8405 (0.8039–0.8717)	0.8997 (0.8584–0.9320)	0.9371 (0.9119–0.9532)	0.8354 (0.7884–0.8739)	0.8857 (0.8387–0.9174)
			HD95	30.29 (14.23–50.38)	39.94 (34.26–46.31)	2.037 (1.725–2.486)	2.086 (1.691–2.566)	2.637 (1.891–3.906)	5.589 (4.380–6.847)	2.472 (1.937–3.130)
		cervical vertebrae 2 (c2)	Dice	0.8974 (0.8608–0.9261)	0.1426 (0.09709–0.1910)	0.8550 (0.8095–0.8919)	0.9273 (0.8929–0.9494)	0.9541 (0.9127–0.9792)	0.8960 (0.8662–0.9232)	0.8664 (0.8097–0.9139)
			HD95	21.44 (9.527–34.46)	31.33 (26.71–36.68)	3.363 (1.595–6.245)	2.130 (1.450–3.031)	2.673 (2.020–3.843)	8.551 (1.966–20.96)	7.246 (2.489–15.69)
		cervical vertebrae 1 (c1)	Dice	0.8614 (0.8092–0.9022)	0.3072 (0.2601–0.3544)	0.8145 (0.7621–0.8546)	0.8862 (0.8314–0.9281)	0.8572 (0.8238–0.8943)	0.8463 (0.7849–0.8922)	0.8567 (0.7904–0.9044)
			HD95	31.12 (9.915–60.06)	17.58 (15.33–20.10)	10.48 (1.730–35.74)	2.253 (1.624–3.201)	13.67 (10.65–17.13)	25.31 (6.757–51.48)	19.17 (2.219–45.49)
		heart	Dice	0.8963 (0.8728–0.9159)	0.2719 (0.2516–0.2916)	0.9016 (0.8783–0.9208)	0.9077 (0.8828–0.9275)	0.9264 (0.9098–0.9455)	0.9102 (0.8874–0.9302)	0.9050 (0.8837–0.9234)
			HD95	11.39 (9.217–13.59)	47.98 (46.01–50.00)	11.42 (6.743–17.69)	4.550 (4.021–5.232)	5.745 (2.767–8.090)	18.79 (13.47–24.64)	7.490 (5.150–10.40)
		aorta	Dice	0.9419 (0.9305–0.9503)	0.7028 (0.6822–0.7205)	0.9174 (0.9002–0.9314)	0.9545 (0.9439–0.9625)	0.9568 (0.9365–0.9609)	0.9493 (0.9356–0.9592)	0.9416 (0.9297–0.9512)
			HD95	3.260 (2.936–3.634)	15.45 (13.37–17.74)	3.837 (2.688–5.495)	2.637 (2.056–3.434)	2.124 (1.590–2.890)	7.521 (4.748–11.48)	5.631 (3.773–7.862)
		pulmonary vein	Dice	0.6835 (0.6445–0.7245)	0.005716 (0.003492–0.008508)	0.7795 (0.7512–0.8041)	0.8276 (0.8016–0.8528)	0.8223 (0.8119–0.8508)	0.7772 (0.7523–0.8019)	0.8154 (0.7889–0.8413)
			HD95	184.9 (158.9–212.2)	92.10 (86.63–97.59)	5.382 (3.814–7.358)	7.469 (5.164–10.02)	7.522 (5.663–10.25)	12.45 (9.479–16.18)	9.880 (7.530–12.52)
		brachiocephalic trunk	Dice	0.8840 (0.8683–0.8968)	0.009225 (0.005542–0.01349)	0.8568 (0.8371–0.8710)	0.8804 (0.8621–0.8951)	0.8962 (0.8920–0.9368)	0.8691 (0.8505–0.8856)	0.8799 (0.8642–0.8922)
			HD95	7.689 (5.078–10.47)	100.7 (89.96–111.8)	5.574 (2.210–12.10)	2.774 (2.475–3.131)	4.090 (3.936–4.683)	4.711 (3.962–5.534)	5.749 (3.170–9.979)
		right subclavian artery	Dice	0.8463 (0.8267–0.8642)	0.001032 (0.0001986–0.002105)	0.8133 (0.8015–0.8236)	0.8584 (0.8382–0.8758)	0.8671 (0.8641–0.8727)	0.8341 (0.8173–0.8494)	0.8490 (0.8288–0.8649)
			HD95	27.34 (13.44–43.13)	124.2 (107.8–142.7)	4.528 (2.314–8.721)	3.350 (2.779–3.954)	2.759 (2.653–4.715)	15.49 (8.998–23.59)	4.987 (3.956–6.116)
		left subclavian artery	Dice	0.8565 (0.8404–0.8713)	0.006123 (0.001615–0.01262)	0.8346 (0.8205–0.8450)	0.8804 (0.8714–0.8894)	0.9045 (0.9003–0.9086)	0.8439 (0.8274–0.8580)	0.8596 (0.8460–0.8708)
			HD95	7.682 (4.926–10.51)	184.4 (156.7–214.9)	5.744 (2.008–13.03)	3.372 (2.567–4.613)	2.702 (1.734–3.505)	7.552 (5.809–9.590)	6.327 (4.657–8.604)
		right common carotid artery	Dice	0.7975 (0.7736–0.8169)	0.00003876 (0.00003283–0.00008297)	0.7922 (0.7726–0.8086)	0.8043 (0.7829–0.8243)	0.8547 (0.8529–0.8583)	0.7855 (0.7645–0.8061)	0.8163 (0.7944–0.8354)
			HD95	9.831 (5.803–16.79)	80.31 (71.79–89.57)	7.681 (2.705–15.10)	4.833 (3.959–5.788)	5.897 (5.399–6.303)	6.763 (5.014–8.964)	7.510 (4.134–12.35)
		left common carotid artery	Dice	0.8541 (0.8351–0.8704)	0.005155 (0.001001–0.01390)	0.8416 (0.8305–0.8516)	0.8766 (0.8651–0.8872)	0.8845 (0.8623–0.9053)	0.8274 (0.8073–0.8464)	0.8772 (0.8626–0.8893)
			HD95	9.919 (4.973–17.52)	117.0 (99.84–135.3)	10.96 (4.452–20.61)	4.133 (3.317–5.111)	2.727 (1.785–5.124)	5.346 (4.410–6.486)	15.74 (8.905–24.13)
		left brachiocephalic vein	Dice	0.7650 (0.7340–0.7944)	0.0006165 (0.00006731–0.001503)	0.8547 (0.8412–0.8669)	0.9008 (0.8881–0.9133)	0.9228 (0.9197–0.9366)	0.8808 (0.8657–0.8940)	0.8895 (0.8792–0.8990)
			HD95	17.51 (13.70–21.07)	235.3 (210.1–262.4)	55.44 (35.78–78.44)	3.417 (2.359–4.876)	3.332 (2.850–5.045)	6.915 (4.538–10.00)	3.607 (2.719–4.759)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	Total-SegmentatorV2	right brachiocephalic vein	Dice	0.8706 (0.8563–0.8833)	0.0006354 (0.0001379–0.001390)	0.8471 (0.8282–0.8625)	0.8801 (0.8644–0.8939)	0.9121 (0.9001–0.9124)	0.8474 (0.8337–0.8598)	0.8731 (0.8559–0.8864)
			HD95	4.469 (4.038–5.055)	249.5 (223.9–273.4)	24.30 (10.85–39.31)	3.384 (2.868–3.991)	2.358 (0.8580–2.753)	7.001 (5.154–10.28)	2.816 (2.554–3.086)
		left atrial appendage of heart	Dice	0.8819 (0.8655–0.8949)	0.0004524 (0.0000–0.001224)	0.0000 (0.0000–0.0000)	0.8727 (0.8551–0.8862)	0.8961 (0.8908–0.9130)	0.6563 (0.6257–0.6856)	0.8670 (0.8481–0.8815)
			HD95	12.50 (3.378–24.59)	132.2 (127.0–138.0)	70.90 (60.28–88.34)	3.321 (2.753–3.931)	5.325 (4.323–6.645)	4.926 (4.458–5.432)	4.276 (2.911–6.042)
		superior vena cava	Dice	0.7630 (0.7270–0.7963)	0.003658 (0.0002010–0.009604)	0.8840 (0.8653–0.9011)	0.9106 (0.8907–0.9266)	0.8991 (0.8660–0.9052)	0.8756 (0.8498–0.8982)	0.8936 (0.8686–0.9135)
			HD95	16.64 (13.48–19.77)	157.4 (145.7–168.4)	2.683 (2.345–3.086)	2.572 (2.290–2.895)	2.357 (1.579–2.645)	15.28 (8.722–22.51)	3.817 (3.119–4.787)
		inferior vena cava	Dice	0.8880 (0.8680–0.9047)	0.6637 (0.6357–0.6885)	0.8990 (0.8833–0.9119)	0.8898 (0.8734–0.9044)	0.9020 (0.8838–0.9140)	0.8815 (0.8610–0.8997)	0.8801 (0.8604–0.8962)
			HD95	5.022 (4.071–6.449)	13.28 (10.29–16.62)	3.041 (2.316–4.353)	4.272 (3.324–5.655)	6.089 (6.046–7.927)	4.037 (3.195–5.289)	4.486 (3.565–5.601)
		portal vein and splenic vein	Dice	0.7943 (0.7659–0.8194)	0.2655 (0.2417–0.2893)	0.7647 (0.7397–0.7887)	0.7627 (0.7299–0.7944)	0.8338 (0.8072–0.8653)	0.7656 (0.7339–0.7914)	0.7848 (0.7563–0.8096)
			HD95	6.945 (5.706–8.486)	49.46 (42.83–56.27)	6.035 (4.658–7.789)	12.45 (9.619–16.02)	7.266 (5.731–8.780)	10.69 (8.402–13.39)	10.08 (7.806–12.56)
		left iliac artery	Dice	0.8423 (0.8139–0.8653)	0.3313 (0.3052–0.3564)	0.8622 (0.8468–0.8750)	0.8571 (0.8289–0.8810)	0.9008 (0.8732–0.9347)	0.8223 (0.7893–0.8519)	0.8144 (0.7901–0.8378)
			HD95	16.41 (13.29–19.78)	21.44 (17.78–25.63)	14.62 (4.953–27.57)	4.410 (3.606–5.226)	3.824 (3.158–6.674)	13.91 (7.857–23.77)	14.70 (8.518–24.77)
		right iliac artery	Dice	0.7870 (0.7563–0.8160)	0.2815 (0.2574–0.3048)	0.8552 (0.8329–0.8730)	0.8501 (0.8190–0.8780)	0.8751 (0.8543–0.9004)	0.8199 (0.7835–0.8498)	0.8423 (0.8150–0.8653)
			HD95	16.21 (13.50–18.78)	24.30 (19.53–29.54)	10.71 (2.049–21.96)	4.328 (3.434–5.371)	4.889 (3.042–6.145)	92.35 (65.90–119.9)	9.788 (5.352–17.40)
		left iliac vena	Dice	0.9044 (0.8911–0.9147)	0.3923 (0.3626–0.4205)	0.8900 (0.8780–0.9007)	0.8797 (0.8626–0.8936)	0.8978 (0.8885–0.8999)	0.8654 (0.8441–0.8838)	0.8718 (0.8549–0.8861)
			HD95	6.022 (3.119–10.76)	20.81 (15.99–26.34)	5.233 (1.843–14.72)	3.042 (2.561–3.581)	5.074 (2.677–6.051)	8.522 (4.534–14.67)	5.861 (4.443–7.790)
		right iliac vena	Dice	0.8540 (0.8248–0.8771)	0.3742 (0.3470–0.4012)	0.8722 (0.8468–0.8903)	0.8559 (0.8277–0.8794)	0.8879 (0.8623–0.9021)	0.8401 (0.8154–0.8633)	0.8534 (0.8273–0.8767)
			HD95	13.84 (11.08–16.52)	20.55 (15.75–25.87)	4.935 (1.894–10.84)	3.430 (2.801–4.160)	4.193 (4.108–5.165)	12.27 (5.145–22.25)	6.681 (3.100–12.96)
		left humerus	Dice	0.6027 (0.5586–0.6420)	0.5589 (0.5103–0.6047)	0.8388 (0.7951–0.8799)	0.8814 (0.8346–0.9191)	0.8689 (0.8474–0.9057)	0.8742 (0.8285–0.9150)	0.8732 (0.8325–0.9121)
			HD95	41.64 (38.16–45.04)	22.75 (13.73–36.17)	28.85 (12.68–48.41)	4.345 (2.535–6.367)	6.993 (0.8770–15.50)	15.93 (8.006–26.61)	8.073 (4.804–12.98)
		right humerus	Dice	0.7016 (0.6587–0.7386)	0.5677 (0.5231–0.6125)	0.8251 (0.7762–0.8714)	0.8600 (0.8048–0.9025)	0.7946 (0.7540–0.8351)	0.8412 (0.7851–0.8896)	0.8689 (0.8204–0.9107)
			HD95	39.23 (35.87–42.35)	31.57 (18.21–48.06)	20.11 (8.357–34.00)	4.522 (2.543–7.316)	8.303 (6.117–11.68)	11.20 (5.826–18.78)	9.540 (3.507–18.80)
		left scapula	Dice	0.8984 (0.8807–0.9145)	0.5227 (0.4888–0.5551)	0.8517 (0.8253–0.8775)	0.9040 (0.8809–0.9215)	0.9487 (0.9370–0.9570)	0.8935 (0.8728–0.9087)	0.9026 (0.8845–0.9177)
			HD95	6.170 (3.730–10.09)	13.97 (9.929–19.31)	3.502 (2.517–4.733)	2.907 (2.340–3.557)	2.349 (0.1230–2.893)	14.67 (9.478–20.56)	4.723 (2.739–7.878)
		right scapula	Dice	0.9021 (0.8832–0.9178)	0.5181 (0.4783–0.5500)	0.8801 (0.8610–0.8964)	0.9166 (0.8956–0.9335)	0.9236 (0.9032–0.9502)	0.8886 (0.8702–0.9029)	0.9078 (0.8896–0.9225)
			HD95	41.58 (27.13–59.38)	13.31 (9.605–18.11)	4.617 (1.675–10.29)	1.977 (1.686–2.343)	2.770 (1.884–2.863)	14.41 (7.021–23.38)	3.660 (2.110–6.434)
		left clavicle	Dice	0.9383 (0.9209–0.9495)	0.4012 (0.3696–0.4298)	0.8882 (0.8644–0.9049)	0.9542 (0.9382–0.9641)	0.9192 (0.9020–0.9370)	0.9416 (0.9204–0.9572)	0.9261 (0.9022–0.9444)
			HD95	17.51 (6.668–29.83)	22.58 (18.69–26.77)	13.07 (1.642–32.76)	1.458 (1.079–2.044)	4.132 (3.566–5.770)	17.21 (9.680–26.64)	6.038 (1.824–14.08)
		right clavicle	Dice	0.9331 (0.9208–0.9414)	0.4364 (0.4067–0.4652)	0.8982 (0.8881–0.9062)	0.9453 (0.9343–0.9523)	0.9375 (0.9280–0.9483)	0.9389 (0.9242–0.9507)	0.9380 (0.9283–0.9467)
			HD95	14.36 (5.300–26.21)	19.94 (16.45–23.97)	3.982 (1.628–8.570)	1.763 (1.561–1.973)	1.427 (1.165–1.629)	7.426 (4.416–10.97)	4.557 (1.789–11.38)
		left femur	Dice	0.9102 (0.8832–0.9325)	0.7449 (0.7084–0.7773)	0.8957 (0.8667–0.9217)	0.9555 (0.9286–0.9759)	0.8843 (0.8565–0.8903)	0.9449 (0.9143–0.9703)	0.9135 (0.8852–0.9369)
			HD95	35.43 (32.96–37.94)	18.40 (11.77–25.75)	130.4 (88.51–172.3)	2.851 (1.682–4.909)	8.004 (6.995–12.05)	81.21 (50.79–113.2)	63.62 (46.15–81.86)
		right femur	Dice	0.8858 (0.8552–0.9107)	0.7436 (0.7111–0.7725)	0.8888 (0.8647–0.9108)	0.9682 (0.9468–0.9805)	0.9130 (0.8828–0.9245)	0.9588 (0.9296–0.9799)	0.9619 (0.9435–0.9749)
			HD95	28.76 (23.82–33.30)	14.20 (9.334–20.12)	196.9 (153.4–246.0)	1.799 (1.556–2.099)	11.84 (8.827–14.44)	16.20 (5.888–28.11)	18.97 (5.073–38.01)
		left hip	Dice	0.9262 (0.9024–0.9429)	0.6507 (0.6274–0.6730)	0.9214 (0.8999–0.9380)	0.9452 (0.9218–0.9633)	0.9334 (0.9091–0.9504)	0.9342 (0.9106–0.9537)	0.9360 (0.9160–0.9516)
			HD95	51.92 (30.18–75.20)	16.50 (11.02–24.38)	10.02 (2.128–22.07)	1.965 (1.605–2.524)	11.61 (6.224–19.56)	72.35 (53.77–92.73)	47.67 (26.47–74.19)
		right hip	Dice	0.8979 (0.8703–0.9213)	0.6754 (0.6398–0.7051)	0.9275 (0.9031–0.9429)	0.9379 (0.9092–0.9592)	0.9321 (0.9250–0.9340)	0.9391 (0.9093–0.9601)	0.9357 (0.9098–0.9539)
			HD95	21.60 (17.71–25.35)	27.27 (18.98–36.90)	18.17 (3.874–36.68)	2.703 (1.553–4.792)	3.953 (2.551–7.664)	32.57 (18.58–49.50)	19.05 (6.468–35.19)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	Total-SegmentatorV2	spinal cord	Dice	0.9205 (0.9157–0.9251)	0.006804 (0.004890–0.009071)	0.9068 (0.8980–0.9141)	0.9227 (0.9185–0.9273)	0.9546 (0.9332–0.9662)	0.8497 (0.8382–0.8596)	0.8993 (0.8923–0.9054)
			HD95	2.844 (2.669–3.084)	177.7 (164.2–191.5)	1.968 (1.724–2.333)	1.871 (1.676–2.127)	1.512 (0.7330–3.169)	20.34 (17.52–23.65)	9.701 (6.697–13.13)
		left gluteus maximus	Dice	0.9257 (0.9098–0.9380)	0.8166 (0.7887–0.8406)	0.9333 (0.9137–0.9473)	0.9311 (0.9089–0.9459)	0.9294 (0.9177–0.9299)	0.9156 (0.8936–0.9311)	0.9115 (0.8873–0.9267)
			HD95	20.14 (10.19–33.14)	9.173 (7.843–11.06)	2.818 (2.440–3.330)	2.912 (2.497–3.501)	5.131 (4.930–5.903)	38.15 (26.51–50.48)	9.690 (5.374–16.50)
		right gluteus maximus	Dice	0.9228 (0.9024–0.9402)	0.8226 (0.7897–0.8521)	0.9362 (0.9138–0.9514)	0.9298 (0.9026–0.9487)	0.9342 (0.9125–0.9551)	0.9086 (0.8757–0.9349)	0.9109 (0.8839–0.9310)
			HD95	14.24 (5.442–26.36)	18.64 (12.36–26.47)	2.533 (2.309–2.778)	2.880 (2.403–3.414)	4.192 (3.764–4.595)	21.42 (10.14–35.20)	14.72 (4.741–27.10)
		left gluteus medius	Dice	0.8969 (0.8680–0.9200)	0.7628 (0.7316–0.7923)	0.9103 (0.8801–0.9349)	0.9107 (0.8908–0.9252)	0.9254 (0.9083–0.9423)	0.8849 (0.8579–0.9062)	0.8882 (0.8674–0.9058)
			HD95	12.47 (8.787–16.15)	14.11 (9.604–20.55)	5.868 (2.604–12.26)	3.330 (3.107–3.589)	4.332 (4.130–4.998)	24.61 (12.40–37.91)	5.669 (4.153–8.288)
		right gluteus medius	Dice	0.9143 (0.8978–0.9264)	0.7600 (0.7308–0.7863)	0.9283 (0.9199–0.9354)	0.9319 (0.9273–0.9361)	0.9319 (0.9294–0.9321)	0.8945 (0.8849–0.9034)	0.9091 (0.9005–0.9164)
			HD95	14.36 (11.22–17.50)	21.26 (12.83–31.76)	2.901 (2.712–3.117)	2.841 (2.650–3.038)	3.180 (2.137–5.510)	15.43 (7.517–26.23)	4.710 (4.122–5.440)
		left gluteus minimus	Dice	0.9008 (0.8952–0.9063)	0.6467 (0.6185–0.6717)	0.8921 (0.8858–0.8970)	0.8866 (0.8810–0.8920)	0.9210 (0.9085–0.9248)	0.8485 (0.8382–0.8573)	0.8564 (0.8475–0.8640)
			HD95	3.373 (3.201–3.572)	13.96 (9.674–19.44)	10.69 (2.500–23.25)	2.820 (2.668–2.975)	3.672 (3.352–5.210)	39.87 (26.87–54.60)	4.259 (3.493–5.612)
		right gluteus minimus	Dice	0.9010 (0.8822–0.9121)	0.6194 (0.5844–0.6526)	0.8974 (0.8916–0.9024)	0.8914 (0.8719–0.9031)	0.9232 (0.9212–0.9372)	0.8609 (0.8401–0.8729)	0.8330 (0.8174–0.8438)
			HD95	7.164 (3.046–15.30)	38.25 (25.95–51.36)	6.921 (2.462–15.69)	2.634 (2.489–2.791)	1.858 (0.9770–2.397)	3.489 (3.213–3.801)	6.009 (5.197–7.023)
		left autochthon	Dice	0.9373 (0.9334–0.9408)	0.8024 (0.7848–0.8198)	0.9396 (0.9335–0.9446)	0.9360 (0.9306–0.9406)	0.9315 (0.9228–0.9466)	0.8968 (0.8848–0.9060)	0.8969 (0.8846–0.9063)
			HD95	5.171 (2.978–8.747)	7.971 (6.790–9.273)	2.058 (1.955–2.169)	2.276 (2.101–2.454)	2.411 (2.184–3.361)	15.93 (12.16–20.91)	6.414 (5.158–7.894)
		right autochthon	Dice	0.9282 (0.9144–0.9388)	0.7973 (0.7756–0.8141)	0.9398 (0.9343–0.9439)	0.9292 (0.9161–0.9403)	0.9306 (0.9144–0.9479)	0.8907 (0.8748–0.9036)	0.8983 (0.8851–0.9090)
			HD95	4.925 (2.909–9.425)	8.120 (6.875–9.651)	3.593 (1.929–8.351)	3.694 (1.984–6.994)	3.254 (1.651–4.451)	11.59 (8.356–15.48)	5.766 (4.321–7.699)
		left iliopsoas	Dice	0.9085 (0.8957–0.9189)	0.6271 (0.5945–0.6586)	0.8980 (0.8816–0.9111)	0.8868 (0.8632–0.9058)	0.9057 (0.8706–0.9118)	0.8558 (0.8307–0.8781)	0.8527 (0.8308–0.8715)
			HD95	3.469 (2.965–4.352)	19.77 (12.85–27.97)	8.272 (2.358–16.70)	2.982 (2.632–3.381)	3.120 (2.380–4.301)	20.99 (11.58–32.82)	14.61 (8.849–22.11)
		right iliopsoas	Dice	0.8941 (0.8700–0.9144)	0.5868 (0.5478–0.6225)	0.8949 (0.8730–0.9131)	0.8857 (0.8605–0.9080)	0.9129 (0.8888–0.9251)	0.8535 (0.8258–0.8776)	0.8087 (0.7801–0.8369)
			HD95	8.183 (3.407–14.97)	27.33 (20.73–35.77)	2.195 (2.015–2.425)	2.655 (2.258–3.123)	2.695 (1.898–5.079)	23.61 (12.68–35.91)	26.64 (18.84–36.68)
		brain	Dice	0.8690 (0.8197–0.9134)	0.6046 (0.5225–0.6758)	0.8650 (0.8167–0.9080)	0.8801 (0.8309–0.9220)	0.9276 (0.8727–0.9907)	0.8661 (0.8064–0.9100)	0.8717 (0.8146–0.9135)
			HD95	21.11 (9.915–36.00)	25.43 (18.29–33.98)	6.622 (4.652–8.810)	6.631 (4.526–9.174)	4.507 (0.7870–4.952)	53.79 (22.81–90.11)	21.21 (7.750–38.59)
		skull	Dice	0.7819 (0.7223–0.8331)	0.03119 (0.02192–0.04115)	0.7638 (0.7146–0.8107)	0.8195 (0.7595–0.8714)	0.8529 (0.8518–0.8633)	0.7912 (0.7434–0.8342)	0.7950 (0.7474–0.8400)
			HD95	11.86 (7.900–16.58)	108.1 (98.57–117.5)	17.09 (9.511–26.47)	4.548 (2.492–7.213)	3.128 (2.535–6.792)	110.9 (87.08–138.1)	29.89 (14.60–47.55)
		right rib 4	Dice	0.8913 (0.8653–0.9134)	0.1689 (0.1502–0.1886)	0.8313 (0.8111–0.8483)	0.9115 (0.8879–0.9315)	0.9421 (0.9249–0.9555)	0.6920 (0.6632–0.7219)	0.8930 (0.8667–0.9159)
			HD95	6.637 (4.586–9.178)	25.33 (21.58–30.08)	4.806 (1.918–8.383)	3.937 (1.990–6.433)	3.302 (1.159–5.125)	30.69 (25.93–36.04)	7.006 (4.311–10.66)
		right rib 3	Dice	0.8835 (0.8591–0.9044)	0.1436 (0.1269–0.1606)	0.7976 (0.7738–0.8181)	0.8925 (0.8673–0.9156)	0.9367 (0.9070–0.9693)	0.6803 (0.6455–0.7143)	0.8570 (0.8253–0.8863)
			HD95	5.956 (3.771–9.639)	44.79 (36.87–53.18)	5.181 (1.840–9.844)	4.139 (1.816–8.162)	4.206 (2.094–7.801)	24.38 (21.67–27.69)	8.339 (4.995–13.14)
		left rib 1	Dice	0.8908 (0.8698–0.9057)	0.2101 (0.1908–0.2310)	0.8014 (0.7857–0.8138)	0.8952 (0.8787–0.9084)	0.9385 (0.9325–0.9580)	0.8704 (0.8458–0.8897)	0.8823 (0.8619–0.8970)
			HD95	3.265 (2.990–3.608)	15.57 (13.27–18.28)	28.19 (16.67–40.82)	2.188 (1.968–2.436)	1.898 (1.879–3.600)	17.24 (13.25–21.64)	4.183 (3.320–5.162)
		left rib 2	Dice	0.9181 (0.9005–0.9318)	0.2124 (0.1921–0.2321)	0.8225 (0.8084–0.8334)	0.9293 (0.9216–0.9365)	0.9298 (0.9223–0.9500)	0.8617 (0.8429–0.8791)	0.9159 (0.9044–0.9250)
			HD95	6.685 (3.045–13.05)	19.01 (16.43–22.00)	10.73 (4.346–19.27)	1.957 (1.250–3.167)	3.350 (2.858–7.031)	12.80 (10.12–15.75)	3.791 (2.429–5.585)
		left rib 3	Dice	0.8141 (0.7853–0.8422)	0.2142 (0.1932–0.2369)	0.8114 (0.7879–0.8303)	0.9103 (0.8906–0.9258)	0.9308 (0.9294–0.9401)	0.7546 (0.7313–0.7765)	0.8707 (0.8492–0.8889)
			HD95	16.17 (13.26–19.21)	24.00 (20.82–27.40)	4.588 (2.412–7.920)	2.752 (1.890–3.717)	2.421 (0.4370–5.045)	23.74 (19.70–30.03)	8.192 (6.684–9.722)
		left rib 4	Dice	0.8865 (0.8585–0.9123)	0.2364 (0.2125–0.2600)	0.8231 (0.7978–0.8447)	0.9038 (0.8797–0.9259)	0.9358 (0.9241–0.9606)	0.6584 (0.6371–0.6801)	0.8737 (0.8481–0.8946)
			HD95	6.645 (5.191–8.341)	20.84 (18.71–23.16)	3.987 (2.313–6.419)	3.217 (2.096–4.544)	2.632 (2.459–4.368)	98.73 (88.94–108.7)	9.200 (7.387–11.14)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	Total-SegmentatorV2	left rib 5	Dice	0.8706 (0.8446–0.8963)	0.2265 (0.2018–0.2540)	0.8263 (0.8057–0.8435)	0.8809 (0.8512–0.9064)	0.8972 (0.8888–0.9005)	0.7741 (0.7433–0.8011)	0.8428 (0.8149–0.8671)
			HD95	8.756 (5.338–14.31)	23.95 (20.78–28.11)	3.809 (2.862–4.988)	4.101 (3.036–5.231)	3.217 (2.803–3.879)	27.78 (21.23–34.42)	5.496 (4.135–7.114)
		left rib 6	Dice	0.8563 (0.8259–0.8840)	0.2535 (0.2286–0.2784)	0.8229 (0.7970–0.8469)	0.8760 (0.8468–0.9037)	0.9078 (0.8978–0.9184)	0.8310 (0.8048–0.8534)	0.8479 (0.8215–0.8707)
			HD95	7.988 (6.143–10.16)	27.76 (23.23–33.31)	3.372 (2.565–4.285)	6.129 (4.683–7.602)	5.056 (3.764–6.348)	19.13 (15.61–23.42)	5.920 (4.451–7.858)
		left rib 7	Dice	0.8694 (0.8444–0.8922)	0.2819 (0.2563–0.3076)	0.8508 (0.8301–0.8697)	0.8815 (0.8536–0.9051)	0.9022 (0.8880–0.9182)	0.8940 (0.8721–0.9120)	0.9007 (0.8839–0.9162)
			HD95	7.803 (6.262–9.687)	25.70 (22.00–30.04)	3.525 (2.513–4.760)	9.493 (7.060–12.47)	3.938 (3.223–5.417)	14.05 (10.80–17.82)	6.631 (5.240–8.094)
		left rib 8	Dice	0.8788 (0.8541–0.8997)	0.2856 (0.2587–0.3101)	0.8478 (0.8257–0.8677)	0.9067 (0.8885–0.9217)	0.8924 (0.8813–0.8938)	0.9058 (0.8875–0.9206)	0.9185 (0.9073–0.9281)
			HD95	8.642 (7.200–10.25)	26.20 (22.18–30.69)	3.597 (2.523–4.865)	5.646 (4.354–7.131)	4.863 (3.854–7.132)	6.351 (4.401–8.733)	4.440 (3.344–5.662)
		left rib 9	Dice	0.8856 (0.8587–0.9082)	0.2709 (0.2454–0.2952)	0.8499 (0.8282–0.8693)	0.9003 (0.8781–0.9175)	0.9002 (0.8980–0.9268)	0.8958 (0.8809–0.9092)	0.9144 (0.9010–0.9259)
			HD95	8.355 (6.245–11.35)	29.41 (25.05–34.26)	4.865 (2.512–7.967)	7.087 (5.142–9.239)	6.396 (4.774–7.760)	19.45 (15.80–23.61)	5.921 (4.490–7.351)
		left rib 10	Dice	0.8808 (0.8525–0.9068)	0.2551 (0.2333–0.2782)	0.8556 (0.8344–0.8733)	0.8972 (0.8738–0.9166)	0.8817 (0.8543–0.8967)	0.8394 (0.8201–0.8570)	0.9108 (0.8927–0.9252)
			HD95	9.876 (7.504–12.70)	28.24 (23.49–33.47)	5.692 (3.192–9.352)	8.802 (6.136–11.90)	5.867 (4.339–8.097)	65.85 (56.31–75.93)	6.729 (5.206–8.463)
		left rib 11	Dice	0.8929 (0.8684–0.9136)	0.2634 (0.2408–0.2875)	0.8506 (0.8314–0.8646)	0.8932 (0.8743–0.9109)	0.8913 (0.8783–0.9122)	0.7983 (0.7696–0.8266)	0.9088 (0.8916–0.9234)
			HD95	6.830 (5.198–8.740)	23.54 (19.68–28.03)	5.530 (2.463–9.357)	7.459 (5.193–10.13)	6.223 (6.171–9.228)	42.07 (30.45–53.84)	3.975 (2.589–5.561)
		left rib 12	Dice	0.8759 (0.8504–0.9002)	0.1967 (0.1723–0.2189)	0.8102 (0.7896–0.8295)	0.8667 (0.8357–0.8941)	0.8716 (0.8613–0.9009)	0.8500 (0.8219–0.8762)	0.8107 (0.7752–0.8419)
			HD95	6.053 (4.099–8.407)	25.06 (20.72–30.51)	11.34 (4.729–18.86)	5.280 (3.598–7.335)	5.887 (5.525–7.483)	36.85 (26.15–49.40)	19.24 (15.55–23.64)
		right rib 1	Dice	0.8911 (0.8722–0.9053)	0.1657 (0.1491–0.1840)	0.7913 (0.7763–0.8048)	0.8978 (0.8791–0.9131)	0.9371 (0.9249–0.9709)	0.8616 (0.8407–0.8820)	0.8923 (0.8717–0.9091)
			HD95	8.505 (3.562–14.80)	17.12 (14.64–20.05)	37.74 (24.34–52.42)	2.134 (1.801–2.550)	1.354 (0.6460–1.928)	14.90 (10.62–20.60)	2.772 (2.156–3.606)
		right rib 2	Dice	0.9096 (0.8945–0.9226)	0.1562 (0.1382–0.1729)	0.8074 (0.7942–0.8175)	0.9211 (0.9035–0.9329)	0.9481 (0.9339–0.9739)	0.8497 (0.8326–0.8644)	0.9091 (0.8937–0.9198)
			HD95	4.638 (2.820–8.246)	22.92 (19.21–27.13)	8.587 (2.858–15.99)	1.503 (1.180–1.894)	2.849 (1.371–4.467)	13.63 (11.79–15.43)	2.731 (1.848–3.880)
		right rib 5	Dice	0.8574 (0.8298–0.8844)	0.1779 (0.1555–0.2003)	0.8060 (0.7789–0.8292)	0.8570 (0.8176–0.8899)	0.8977 (0.8693–0.9254)	0.6700 (0.6333–0.7094)	0.8730 (0.8460–0.8968)
			HD95	15.20 (6.169–30.30)	34.49 (27.02–43.97)	4.887 (2.613–7.675)	4.538 (2.728–7.287)	3.433 (0.5940–6.004)	23.23 (18.45–28.03)	5.944 (3.648–9.037)
		right rib 6	Dice	0.8702 (0.8458–0.8962)	0.1726 (0.1506–0.1964)	0.8388 (0.8212–0.8562)	0.8718 (0.8437–0.8963)	0.9311 (0.9155–0.9567)	0.8506 (0.8282–0.8712)	0.8952 (0.8780–0.9110)
			HD95	8.293 (6.751–9.878)	90.16 (77.60–104.4)	4.387 (2.543–7.452)	9.197 (6.653–12.36)	3.547 (0.9560–4.877)	20.38 (16.14–25.67)	6.314 (3.702–9.700)
		right rib 7	Dice	0.8722 (0.8426–0.8973)	0.2241 (0.2027–0.2462)	0.8655 (0.8522–0.8774)	0.8974 (0.8730–0.9178)	0.9167 (0.9144–0.9329)	0.8640 (0.8492–0.8771)	0.8888 (0.8689–0.9053)
			HD95	9.559 (7.457–12.17)	27.94 (23.17–33.30)	4.112 (2.582–6.362)	8.399 (6.165–11.03)	5.448 (3.359–9.480)	28.45 (22.88–34.32)	11.47 (8.975–14.11)
		right rib 8	Dice	0.8730 (0.8447–0.8977)	0.2378 (0.2166–0.2611)	0.8625 (0.8506–0.8737)	0.8954 (0.8736–0.9146)	0.9272 (0.9141–0.9514)	0.8898 (0.8733–0.9045)	0.8928 (0.8809–0.9040)
			HD95	13.62 (8.421–20.43)	30.25 (25.81–35.25)	4.998 (2.936–7.500)	7.458 (5.281–10.23)	3.385 (1.281–7.351)	14.56 (11.40–18.28)	11.75 (9.435–14.33)
		right rib 9	Dice	0.8720 (0.8423–0.8980)	0.2691 (0.2478–0.2895)	0.8676 (0.8524–0.8798)	0.8960 (0.8739–0.9147)	0.9232 (0.9033–0.9426)	0.8439 (0.8258–0.8615)	0.8973 (0.8816–0.9106)
			HD95	14.30 (8.937–22.34)	30.94 (25.91–36.90)	4.152 (2.465–6.432)	7.299 (5.443–9.486)	6.475 (5.652–8.120)	28.28 (25.38–31.41)	12.66 (10.64–14.81)
		right rib 10	Dice	0.8871 (0.8618–0.9082)	0.2549 (0.2331–0.2780)	0.8711 (0.8603–0.8795)	0.8966 (0.8724–0.9169)	0.9227 (0.8744–0.9341)	0.7202 (0.6911–0.7481)	0.8156 (0.7885–0.8426)
			HD95	35.12 (21.54–51.64)	45.95 (38.02–54.56)	4.713 (2.780–7.575)	7.605 (5.511–9.682)	3.581 (2.937–5.813)	26.76 (24.82–28.74)	8.484 (6.642–10.41)
		right rib 11	Dice	0.8790 (0.8484–0.9040)	0.2291 (0.2077–0.2503)	0.8407 (0.8199–0.8575)	0.8715 (0.8437–0.8955)	0.9183 (0.8982–0.9308)	0.6799 (0.6369–0.7193)	0.8891 (0.8629–0.9140)
			HD95	7.688 (5.780–9.855)	37.96 (31.48–45.05)	4.217 (2.271–7.623)	8.350 (5.853–10.99)	4.356 (0.4170–4.400)	20.94 (15.93–26.59)	3.456 (2.386–4.726)
		right rib 12	Dice	0.8701 (0.8410–0.8953)	0.1270 (0.1112–0.1431)	0.8076 (0.7877–0.8256)	0.8615 (0.8307–0.8897)	0.9088 (0.9010–0.9399)	0.8521 (0.8229–0.8787)	0.8854 (0.8604–0.9065)
			HD95	9.898 (5.946–14.86)	35.99 (29.24–43.45)	9.969 (3.906–17.81)	6.963 (4.587–9.889)	4.602 (1.213–4.969)	31.12 (20.25–42.57)	8.049 (4.722–12.49)
		sternum	Dice	0.8878 (0.8672–0.9058)	0.0002803 (0.0001041–0.0005131)	0.8530 (0.8261–0.8791)	0.8923 (0.8686–0.9131)	0.9031 (0.8827–0.9161)	0.8842 (0.8608–0.9049)	0.8658 (0.8366–0.8924)
			HD95	4.628 (3.461–6.173)	146.8 (141.4–152.3)	13.71 (3.583–29.32)	3.837 (2.679–5.293)	3.394 (1.041–3.407)	7.751 (5.016–11.23)	3.400 (2.433–4.728)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	Total-SegmentatorV2	costal cartilage	Dice	0.8416 (0.8217–0.8590)	0.003181 (0.002157–0.004239)	0.7986 (0.7800–0.8155)	0.8186 (0.7952–0.8384)	0.8630 (0.8490–0.8843)	0.8144 (0.7938–0.8327)	0.8223 (0.8033–0.8392)
			HD95	8.004 (5.274–11.20)	192.4 (182.4–203.0)	8.492 (4.494–13.81)	5.949 (4.279–7.978)	5.372 (4.728–7.700)	9.422 (6.533–12.97)	7.822 (5.481–10.77)
	Pancreas-CT	pancreas	Dice	0.8410 (0.8269–0.8530)	0.5967 (0.5461–0.6355)	0.8636 (0.8472–0.8779)	0.8581 (0.8440–0.8712)	0.8562 (0.8432–0.8676)	0.8634 (0.8495–0.8780)	0.8527 (0.8391–0.8670)
			HD95	5.249 (4.469–6.225)	13.80 (9.282–18.85)	3.188 (2.522–3.985)	3.607 (3.139–4.142)	4.044 (3.235–4.075)	3.884 (3.102–4.999)	5.350 (3.750–7.595)
		spleen	Dice	0.9315 (0.9071–0.9473)	0.8031 (0.7775–0.8267)	0.9585 (0.9520–0.9649)	0.9606 (0.9524–0.9671)	0.9583 (0.9580–0.9684)	0.9759 (0.9659–0.9845)	0.9639 (0.9560–0.9699)
			HD95	7.124 (3.387–14.20)	9.811 (8.487–11.21)	2.017 (1.722–2.333)	1.946 (1.565–2.388)	4.213 (2.787–4.812)	1.293 (0.8280–1.713)	1.631 (1.411–1.928)
		right kidney	Dice	0.9297 (0.9219–0.9373)	0.7291 (0.6499–0.7857)	0.9452 (0.9387–0.9508)	0.9449 (0.9354–0.9520)	0.9497 (0.9411–0.9599)	0.9637 (0.9547–0.9705)	0.9507 (0.9432–0.9577)
			HD95	4.527 (3.363–5.800)	71.81 (34.37–107.3)	2.642 (2.103–3.166)	3.219 (2.187–4.361)	3.211 (2.502–4.534)	1.713 (0.9450–2.737)	2.413 (1.278–3.771)
		left kidney	Dice	0.9284 (0.9181–0.9385)	0.6127 (0.5298–0.7008)	0.9504 (0.9422–0.9572)	0.9501 (0.9422–0.9571)	0.9332 (0.9205–0.9423)	0.9665 (0.9610–0.9710)	0.9571 (0.9523–0.9603)
			HD95	6.735 (3.023–13.04)	98.78 (62.57–123.9)	2.351 (1.732–3.079)	2.348 (1.598–3.025)	4.079 (3.068–6.058)	1.917 (0.8350–3.132)	1.473 (1.137–1.805)
		gallbladder	Dice	0.7718 (0.5755–0.8902)	0.3024 (0.07825–0.5348)	0.7902 (0.5760–0.9149)	0.7085 (0.4070–0.8995)	0.7731 (0.5719–0.8910)	0.8988 (0.8471–0.9360)	0.8997 (0.8698–0.9260)
			HD95	7.718 (4.473–12.89)	40.39 (8.823–97.51)	6.483 (2.324–13.75)	15.92 (3.669–37.58)	6.960 (3.066–10.81)	2.349 (1.477–3.439)	2.723 (2.148–3.383)
	BTCV-Abdomen	esophagus	Dice	0.7672 (0.7559–0.7809)	0.5313 (0.4868–0.5817)	0.7962 (0.7832–0.8113)	0.7978 (0.7624–0.8324)	0.7400 (0.7303–0.7621)	0.8763 (0.8518–0.8983)	0.8692 (0.8459–0.8954)
			HD95	11.82 (4.270–19.88)	9.610 (6.890–13.98)	4.437 (3.731–5.036)	6.018 (4.259–7.901)	6.910 (5.091–7.447)	2.400 (1.625–3.453)	2.360 (1.858–2.807)
		liver	Dice	0.9546 (0.9498–0.9580)	0.8678 (0.8544–0.8802)	0.9668 (0.9626–0.9702)	0.9660 (0.9619–0.9702)	0.9659 (0.9588–0.9824)	0.9840 (0.9802–0.9872)	0.9706 (0.9673–0.9735)
			HD95	6.730 (4.729–8.876)	11.57 (9.813–13.75)	4.011 (3.247–4.874)	6.011 (4.380–7.866)	2.920 (1.042–4.767)	3.102 (2.596–3.783)	4.049 (3.330–4.855)
		stomach	Dice	0.9315 (0.9221–0.9393)	0.7742 (0.7543–0.7968)	0.8463 (0.6740–0.9399)	0.9254 (0.9097–0.9408)	0.7927 (0.7891–0.8026)	0.9628 (0.9540–0.9707)	0.9519 (0.9452–0.9575)
			HD95	5.419 (4.527–6.430)	12.64 (11.39–13.98)	57.95 (3.510–165.7)	10.37 (4.882–18.47)	19.33 (18.84–20.42)	3.372 (2.631–4.108)	3.596 (3.034–4.174)
		aorta	Dice	0.9169 (0.9070–0.9271)	0.8156 (0.7929–0.8403)	0.9140 (0.8851–0.9324)	0.9380 (0.9284–0.9462)	0.9059 (0.8968–0.9173)	0.9535 (0.9454–0.9611)	0.9388 (0.9290–0.9479)
			HD95	3.071 (2.315–4.295)	5.147 (3.658–6.742)	27.74 (2.038–77.35)	1.283 (1.058–1.498)	2.035 (1.724–2.555)	0.6480 (0.3822–0.9160)	1.260 (1.054–1.467)
		inferior vena cava	Dice	0.8267 (0.7541–0.8840)	0.6952 (0.5514–0.8062)	0.8684 (0.8223–0.9094)	0.8603 (0.8056–0.9042)	0.8256 (0.7690–0.8642)	0.8901 (0.8497–0.9274)	0.8889 (0.8571–0.9194)
			HD95	19.67 (13.42–26.18)	8.720 (6.027–11.70)	5.979 (4.415–7.855)	6.302 (3.749–9.362)	8.475 (6.942–11.29)	3.400 (1.922–5.087)	3.463 (2.339–4.579)
		portal vein and splenic vein	Dice	0.7689 (0.7208–0.8116)	0.2380 (0.1280–0.3294)	0.7772 (0.7416–0.8112)	0.6829 (0.5884–0.7675)	0.7480 (0.6960–0.7807)	0.8339 (0.7968–0.8755)	0.8326 (0.8027–0.8676)
			HD95	7.122 (5.254–9.389)	38.39 (29.44–48.19)	6.256 (3.726–8.802)	22.56 (9.725–38.85)	7.913 (7.224–11.53)	3.857 (2.361–5.606)	4.635 (3.280–5.897)
		pancreas	Dice	0.8263 (0.7991–0.8492)	0.5561 (0.5151–0.6010)	0.8494 (0.8376–0.8614)	0.8130 (0.7640–0.8485)	0.8250 (0.8138–0.8376)	0.8769 (0.8571–0.8955)	0.8510 (0.8042–0.8822)
			HD95	11.28 (5.723–20.17)	15.73 (11.51–20.70)	4.466 (3.874–4.936)	12.58 (5.076–26.22)	7.612 (5.376–8.179)	4.532 (3.258–6.250)	10.13 (3.966–20.56)
		right adrenal gland	Dice	0.6703 (0.6271–0.7159)	0.3156 (0.2546–0.3767)	0.7467 (0.7155–0.7742)	0.7151 (0.6759–0.7547)	0.6616 (0.6341–0.6851)	0.7598 (0.7131–0.7967)	0.8074 (0.7904–0.8255)
			HD95	6.527 (4.039–9.228)	9.723 (8.302–11.07)	3.454 (2.371–4.575)	4.591 (2.190–7.142)	7.318 (6.779–7.522)	2.873 (1.694–4.241)	1.658 (1.408–1.990)
		left adrenal gland	Dice	0.6417 (0.4345–0.7760)	0.2156 (0.1050–0.3244)	0.7827 (0.7472–0.8307)	0.7394 (0.6197–0.8212)	0.6395 (0.6029–0.6872)	0.8098 (0.7530–0.8548)	0.8223 (0.7911–0.8547)
			HD95	9.681 (4.296–18.81)	24.01 (13.07–35.09)	2.824 (1.892–3.705)	5.621 (3.435–8.278)	5.767 (5.287–7.035)	2.207 (1.494–3.192)	2.139 (1.422–3.334)
		urinary bladder	Dice	0.4910 (0.2551–0.7404)	0.3014 (0.04782–0.5618)	0.8934 (0.8665–0.9238)	0.8993 (0.8710–0.9267)	0.7559 (0.7346–0.7719)	0.9071 (0.8774–0.9405)	0.9188 (0.9023–0.9388)
			HD95	65.95 (20.37–145.2)	35.84 (21.71–49.89)	3.643 (3.061–4.319)	3.986 (2.724–5.684)	10.62 (10.37–11.87)	3.142 (2.322–4.409)	2.718 (2.587–2.849)
	BTCV-Cervix	uterus	Dice	0.4706 (0.2592–0.7134)	0.2136 (0.1118–0.3222)	0.7954 (0.7460–0.8506)	0.7969 (0.7352–0.8602)	0.6703 (0.6268–0.7298)	0.8277 (0.7752–0.8808)	0.8200 (0.7715–0.8681)
			HD95	74.66 (47.90–112.2)	55.42 (40.94–71.19)	12.81 (8.631–17.50)	17.65 (10.34–25.28)	33.27 (30.47–38.61)	10.19 (6.806–13.59)	36.03 (8.990–88.31)
		rectum	Dice	Not supported	0.1099 (0.01854–0.2124)	0.8379 (0.7930–0.8780)	0.8662 (0.8459–0.8866)	0.6899 (0.6678–0.7188)	0.8373 (0.7783–0.8886)	0.8731 (0.8512–0.8933)
			HD95		52.36 (30.36–78.00)	6.321 (3.955–8.383)	5.734 (3.871–8.025)	12.87 (11.00–15.52)	7.075 (4.262–10.16)	19.81 (4.399–40.14)
		small bowel	Dice	0.3293 (0.2089–0.4368)	0.2758 (0.1549–0.4249)	0.6778 (0.5450–0.7653)	0.6588 (0.4877–0.7776)	0.5251 (0.3808–0.6363)	0.7680 (0.7192–0.8169)	0.7677 (0.7057–0.8204)
			HD95	107.7 (63.57–141.8)	105.9 (76.76–131.2)	51.13 (11.93–90.43)	48.38 (19.75–84.00)	64.56 (35.58–90.66)	50.38 (21.50–92.12)	43.32 (13.22–85.87)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	CRLM-CT	liver	Dice	0.9313 (0.9264–0.9357)	0.8676 (0.8338–0.8890)	0.9420 (0.9372–0.9461)	0.9446 (0.9392–0.9492)	0.9404 (0.9274–0.9482)	0.9610 (0.9576–0.9641)	0.9456 (0.9417–0.9494)
			HD95	22.62 (21.40–23.81)	18.60 (15.97–22.75)	17.51 (16.18–18.88)	24.41 (22.85–26.37)	19.89 (18.75–21.21)	19.45 (18.53–20.36)	22.11 (21.19–22.96)
		hepatic vein	Dice	0.4671 (0.4135–0.5113)	0.1001 (0.07545–0.1270)	0.4195 (0.3761–0.4622)	0.1494 (0.1155–0.1865)	0.4918 (0.4586–0.5188)	0.5073 (0.4607–0.5503)	0.5260 (0.4789–0.5717)
			HD95	28.01 (22.75–34.02)	81.33 (71.86–90.28)	27.13 (22.28–32.62)	39.03 (34.93–44.54)	29.65 (24.57–34.89)	24.53 (19.76–30.99)	25.74 (20.46–33.09)
		portal vein	Dice	0.4619 (0.4322–0.4938)	0.08952 (0.06067–0.1220)	0.2715 (0.2321–0.3132)	0.4883 (0.4627–0.5129)	0.5301 (0.5113–0.6482)	0.5934 (0.5635–0.6204)	0.6508 (0.6231–0.6729)
			HD95	97.68 (88.51–106.2)	91.06 (81.59–101.1)	91.46 (83.49–99.73)	80.52 (70.32–92.03)	37.60 (31.88–43.50)	29.80 (24.15–37.04)	27.82 (22.93–34.63)
		liver tumor	Dice	0.6206 (0.5362–0.7031)	0.08934 (0.02796–0.1611)	0.6307 (0.5362–0.7203)	0.6161 (0.5261–0.6978)	0.6270 (0.5519–0.6862)	0.6875 (0.6099–0.7583)	0.6898 (0.6202–0.7546)
			HD95	43.58 (25.26–64.56)	125.8 (106.2–145.6)	21.65 (10.65–37.00)	47.54 (27.73–66.35)	44.21 (29.58–60.99)	36.05 (20.86–53.45)	40.84 (25.58–59.00)
		left carotid artery	Dice	Not supported	0.1697 (0.1209–0.2185)	0.7552 (0.7069–0.8011)	0.7870 (0.7644–0.8138)	0.7673 (0.7451–0.7896)	0.7096 (0.6631–0.7530)	0.8218 (0.8003–0.8413)
			HD95		39.53 (29.49–50.55)	10.53 (7.754–13.12)	16.72 (14.91–18.28)	8.325 (6.872–10.38)	24.98 (19.95–32.43)	2.388 (2.024–2.776)
		right carotid artery	Dice	Not supported	0.1760 (0.1327–0.2183)	0.7716 (0.7350–0.8001)	0.7911 (0.7706–0.8132)	0.7755 (0.7648–0.7967)	0.7586 (0.7298–0.7877)	0.8086 (0.7848–0.8331)
			HD95		30.61 (22.03–40.08)	10.58 (7.730–13.05)	18.13 (14.31–20.94)	18.51 (14.47–20.59)	20.54 (18.21–22.72)	2.260 (1.738–2.782)
		arytenoid	Dice	Not supported	0.05639 (0.01929–0.09761)	0.4791 (0.3834–0.5704)	0.5628 (0.4592–0.6517)	0.5436 (0.5042–0.5970)	0.5992 (0.5287–0.6584)	0.6959 (0.6442–0.7502)
			HD95		27.05 (14.16–47.72)	3.164 (2.621–3.684)	3.120 (2.399–3.818)	4.733 (3.546–5.792)	2.783 (1.981–3.840)	1.907 (1.551–2.327)
		mandible	Dice	Not supported	0.5040 (0.3718–0.5916)	0.9211 (0.9131–0.9291)	0.9417 (0.9316–0.9490)	0.9103 (0.8955–0.9325)	0.9151 (0.8989–0.9266)	0.9301 (0.9198–0.9394)
			HD95		16.24 (11.12–23.02)	1.941 (1.560–2.319)	1.922 (1.265–2.697)	2.064 (1.627–2.810)	10.29 (2.369–24.04)	2.258 (1.481–3.092)
		brainstem	Dice	Not supported	0.4967 (0.3413–0.6424)	0.8125 (0.7847–0.8372)	0.7820 (0.7405–0.8194)	0.8165 (0.7805–0.8423)	0.8302 (0.8000–0.8568)	0.8748 (0.8686–0.8815)
			HD95		18.47 (9.270–33.67)	5.125 (4.031–6.435)	5.988 (4.268–8.059)	6.864 (4.840–8.766)	5.417 (3.677–7.378)	2.955 (2.732–3.174)
		buccal mucosa	Dice	Not supported	0.1970 (0.1306–0.2690)	0.5573 (0.4876–0.6167)	0.5857 (0.5310–0.6456)	0.5487 (0.4798–0.6160)	0.6096 (0.5583–0.6639)	0.7467 (0.7120–0.7802)
			HD95		29.35 (23.83–34.14)	6.513 (4.708–8.963)	7.630 (5.645–9.779)	7.171 (3.824–9.181)	15.21 (8.625–23.82)	3.292 (3.060–3.532)
		oral cavity	Dice	Not supported	0.7066 (0.6718–0.7398)	0.8645 (0.8359–0.8912)	0.8551 (0.8209–0.8851)	0.8623 (0.8411–0.8972)	0.8803 (0.8613–0.8983)	0.9070 (0.8963–0.9184)
			HD95		15.12 (12.74–17.77)	6.065 (4.433–8.073)	8.052 (5.516–11.24)	8.020 (5.730–8.381)	6.795 (4.542–9.966)	3.933 (3.256–4.701)
	HaN-Seg	left cochlea	Dice	Not supported	0.0000 (0.0000–0.0000)	0.5713 (0.5075–0.6408)	0.6478 (0.5868–0.6997)	0.6312 (0.6255–0.6346)	0.7068 (0.6320–0.7802)	0.7716 (0.7225–0.8257)
			HD95		97.34 (84.64–110.9)	2.340 (1.946–2.661)	1.790 (1.413–2.195)	3.194 (2.161–4.702)	1.699 (1.221–2.276)	1.118 (0.6804–1.804)
		right cochlea	Dice	Not supported	0.0000 (0.0000–0.0000)	0.5778 (0.5211–0.6316)	0.6486 (0.5666–0.7316)	0.5686 (0.5362–0.5887)	0.7442 (0.6725–0.8068)	0.8038 (0.7798–0.8296)
			HD95		113.5 (77.09–132.8)	2.228 (1.847–2.625)	1.834 (1.275–2.553)	2.754 (1.922–4.005)	1.674 (1.120–2.191)	0.7156 (0.6286–0.8110)
		cricopharyngeal inlet	Dice	Not supported	0.1526 (0.07340–0.2367)	0.6526 (0.5859–0.7043)	0.7162 (0.6705–0.7547)	0.6663 (0.6209–0.7063)	0.6955 (0.6648–0.7228)	0.8182 (0.7852–0.8444)
			HD95		37.56 (17.49–68.89)	5.173 (3.589–7.383)	3.899 (3.033–4.890)	3.930 (2.925–6.293)	4.057 (3.294–4.904)	2.076 (1.652–2.494)
		cervical esophagus	Dice	Not supported	0.1777 (0.1276–0.2359)	0.6533 (0.6049–0.6996)	0.7388 (0.6985–0.7785)	0.6974 (0.6544–0.7539)	0.7995 (0.7632–0.8255)	0.8681 (0.8443–0.8944)
			HD95		29.99 (20.76–41.77)	5.342 (4.466–6.139)	4.286 (3.228–5.330)	5.864 (5.453–6.539)	3.427 (2.625–4.402)	1.788 (1.370–2.225)
		left anterior eyeball	Dice	Not supported	0.1063 (0.04647–0.1806)	0.6971 (0.6575–0.7395)	0.7514 (0.7086–0.7933)	0.7210 (0.6818–0.7753)	0.8339 (0.7848–0.8702)	0.8428 (0.8164–0.8718)
			HD95		14.18 (11.82–16.64)	3.139 (2.262–4.130)	2.595 (2.196–2.957)	3.810 (2.860–4.034)	1.778 (1.307–2.280)	1.676 (1.298–2.050)
		right anterior eyeball	Dice	Not supported	0.06729 (0.03001–0.1004)	0.7071 (0.6539–0.7580)	0.7682 (0.7241–0.8094)	0.7667 (0.7475–0.8131)	0.8136 (0.7846–0.8426)	0.8290 (0.7996–0.8574)
			HD95		16.92 (14.73–19.39)	3.234 (2.476–4.190)	2.652 (1.869–3.653)	4.737 (3.634–5.184)	1.758 (1.413–2.183)	1.477 (1.135–1.852)
		left posterior eyeball	Dice	Not supported	0.4885 (0.4366–0.5566)	0.9040 (0.8928–0.9137)	0.8988 (0.8911–0.9075)	0.8944 (0.8923–0.9042)	0.9202 (0.9026–0.9330)	0.9316 (0.9211–0.9398)
			HD95		10.82 (8.974–12.58)	1.994 (1.693–2.323)	2.321 (2.036–2.599)	4.022 (3.169–4.122)	1.569 (1.257–2.009)	1.283 (1.100–1.476)
		right posterior eyeball	Dice	Not supported	0.5039 (0.4184–0.5784)	0.9062 (0.9005–0.9134)	0.8955 (0.8823–0.9103)	0.8852 (0.8702–0.8969)	0.9122 (0.9033–0.9218)	0.9293 (0.9211–0.9362)
			HD95		10.71 (8.487–12.79)	1.885 (1.657–2.034)	2.078 (1.798–2.421)	3.890 (3.140–4.349)	1.810 (1.505–2.204)	1.445 (1.151–1.774)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	HaN-Seg	left lacrimal gland	Dice	Not supported	0.2007 (0.09178–0.3300)	0.5300 (0.3870–0.6326)	0.0000 (0.0000–0.0000)	0.5280 (0.5161–0.5298)	0.3817 (0.2499–0.4961)	0.7802 (0.7454–0.8059)
			HD95		33.46 (10.02–61.37)	4.404 (2.728–6.754)	– (– –)	5.584 (4.995–6.750)	4.392 (3.381–5.333)	1.807 (1.238–2.494)
		right lacrimal gland	Dice	Not supported	0.1025 (0.05153–0.1520)	0.4334 (0.3186–0.5382)	0.0000 (0.0000–0.0000)	0.5158 (0.4654–0.5681)	0.6667 (0.5507–0.7585)	0.6795 (0.5950–0.7390)
			HD95		17.85 (13.86–22.33)	4.933 (3.582–6.427)	– (– –)	5.791 (4.841–6.983)	2.914 (1.744–4.816)	2.028 (1.464–2.718)
		left submandibular gland	Dice	Not supported	0.4114 (0.3025–0.5113)	0.8340 (0.8048–0.8642)	0.8565 (0.8417–0.8716)	0.8553 (0.8245–0.8789)	0.8450 (0.8148–0.8702)	0.8882 (0.8660–0.9034)
			HD95		33.71 (16.03–55.84)	3.894 (2.635–5.343)	3.691 (2.548–5.214)	3.988 (3.395–4.017)	2.820 (2.174–3.506)	2.038 (1.522–2.569)
		right submandibular gland	Dice	Not supported	0.3547 (0.2342–0.4755)	0.7913 (0.7188–0.8501)	0.8498 (0.8309–0.8698)	0.8398 (0.8274–0.8845)	0.8528 (0.8178–0.8851)	0.8566 (0.8311–0.8770)
			HD95		30.94 (15.38–45.70)	4.851 (2.932–7.223)	3.178 (2.511–4.018)	5.004 (4.535–7.257)	2.883 (2.133–3.719)	2.432 (2.006–2.900)
		thyroid	Dice	0.8137 (0.7950–0.8325)	0.0000 (0.0000–0.0000)	0.8479 (0.8367–0.8613)	0.8677 (0.8532–0.8816)	0.8253 (0.7967–0.8263)	0.8703 (0.8588–0.8852)	0.8716 (0.8543–0.8910)
			HD95	3.905 (3.128–4.753)	110.7 (99.82–118.6)	2.293 (1.992–2.771)	2.147 (1.619–2.775)	3.296 (3.193–3.456)	1.992 (1.641–2.400)	2.021 (1.617–2.428)
		larynx glottis	Dice	Not supported	0.1268 (0.05605–0.1989)	0.6792 (0.6251–0.7346)	0.7068 (0.6351–0.7692)	0.6764 (0.6255–0.7295)	0.7462 (0.6932–0.7965)	0.7859 (0.7703–0.8006)
			HD95		15.92 (13.03–19.25)	3.798 (2.746–4.798)	3.852 (2.524–5.109)	5.479 (4.040–5.657)	2.570 (2.246–2.901)	2.663 (2.252–3.113)
		larynx supraglottis	Dice	Not supported	0.3091 (0.2042–0.4112)	0.7602 (0.7392–0.7794)	0.7934 (0.7563–0.8271)	0.7573 (0.7337–0.7868)	0.8200 (0.8075–0.8336)	0.8336 (0.8135–0.8513)
			HD95		34.45 (18.97–52.07)	3.880 (3.358–4.530)	3.872 (2.877–5.020)	6.034 (4.601–6.892)	3.452 (2.883–4.099)	2.756 (2.490–2.988)
		lips	Dice	Not supported	0.4118 (0.3514–0.4768)	0.6170 (0.5683–0.6715)	0.6486 (0.5858–0.7207)	0.6411 (0.5798–0.6915)	0.7135 (0.6789–0.7486)	0.7840 (0.7431–0.8269)
			HD95		12.61 (10.76–14.29)	6.915 (5.774–8.462)	8.342 (5.769–10.95)	6.332 (5.583–8.446)	4.667 (4.034–5.315)	3.112 (2.742–3.458)
		optic chiasm	Dice	Not supported	0.0000 (0.0000–0.0000)	0.4159 (0.3603–0.4634)	0.5223 (0.4716–0.5662)	0.3865 (0.3505–0.4273)	0.5704 (0.5148–0.6292)	0.6780 (0.6170–0.7286)
			HD95		156.7 (142.2–168.2)	3.677 (3.080–4.360)	3.721 (2.971–4.548)	4.678 (3.971–6.244)	3.097 (2.618–3.556)	2.658 (2.216–3.122)
		left optic nerve	Dice	Not supported	0.0001222 (0.0000–0.0003666)	0.5747 (0.4451–0.6662)	0.6153 (0.5424–0.6899)	0.5652 (0.5223–0.6177)	0.7131 (0.6553–0.7583)	0.7903 (0.7339–0.8368)
			HD95		123.9 (95.42–140.3)	3.003 (2.443–3.588)	5.691 (3.519–8.045)	4.792 (3.726–7.302)	12.02 (2.525–23.16)	1.936 (1.434–2.435)
		right optic nerve	Dice	Not supported	0.004694 (0.0000–0.01408)	0.6576 (0.6025–0.7092)	0.6609 (0.6082–0.6971)	0.6433 (0.5937–0.6743)	0.7405 (0.7170–0.7655)	0.7218 (0.6509–0.7756)
			HD95		77.57 (41.38–117.6)	3.037 (2.404–3.991)	4.108 (2.687–5.596)	4.330 (2.523–5.427)	3.571 (2.653–4.564)	2.478 (1.812–3.136)
		left parotid gland	Dice	Not supported	0.5415 (0.4404–0.6314)	0.8186 (0.7915–0.8418)	0.8267 (0.7975–0.8537)	0.8070 (0.7661–0.8416)	0.8446 (0.8274–0.8582)	0.8567 (0.8365–0.8737)
			HD95		24.11 (10.43–44.59)	6.398 (4.891–8.804)	7.011 (5.452–9.054)	8.537 (6.659–10.82)	5.482 (3.694–7.709)	5.381 (3.799–7.526)
		right parotid gland	Dice	Not supported	0.5631 (0.4671–0.6505)	0.7916 (0.7402–0.8383)	0.7985 (0.7503–0.8393)	0.7861 (0.7333–0.8297)	0.8312 (0.8099–0.8499)	0.8581 (0.8388–0.8763)
			HD95		21.11 (8.910–44.17)	5.459 (4.519–6.602)	6.442 (5.288–7.806)	9.373 (8.275–11.24)	4.875 (3.784–6.161)	3.221 (2.911–3.542)
		pituitary gland	Dice	Not supported	0.0000 (0.0000–0.0000)	0.6126 (0.5044–0.7052)	0.7403 (0.6793–0.8064)	0.6551 (0.5798–0.7039)	0.7674 (0.6220–0.8487)	0.7953 (0.7589–0.8254)
			HD95		147.0 (138.8–154.7)	2.815 (2.138–3.414)	1.965 (1.469–2.500)	4.164 (2.061–4.501)	1.393 (1.007–1.889)	1.462 (1.234–1.686)
		spinal cord	Dice	0.3798 (0.3482–0.4187)	0.3246 (0.2266–0.4031)	0.7291 (0.6480–0.7843)	0.4867 (0.4522–0.5249)	0.7224 (0.6868–0.7596)	0.8082 (0.7866–0.8327)	0.7892 (0.7642–0.8125)
			HD95	80.45 (74.34–87.15)	47.27 (32.51–63.18)	10.21 (2.433–25.42)	78.12 (71.77–84.23)	12.13 (7.890–17.83)	2.127 (1.824–2.453)	8.460 (2.442–20.12)
		kidney	Dice	0.8548 (0.8311–0.8749)	0.7126 (0.6868–0.7349)	0.8560 (0.8271–0.8850)	0.8589 (0.8298–0.8867)	0.9031 (0.8762–0.9199)	0.9231 (0.9032–0.9374)	0.9196 (0.8977–0.9352)
			HD95	14.67 (10.48–19.51)	18.87 (14.81–24.01)	9.862 (6.794–13.77)	11.49 (7.874–15.84)	13.89 (10.11–17.81)	11.01 (7.595–15.52)	9.792 (6.622–13.92)
	KiTS23	kidney tumor	Dice	Not supported	0.2293 (0.1747–0.2853)	0.6648 (0.6083–0.7217)	0.8045 (0.7578–0.8429)	0.7980 (0.7797–0.8258)	0.7273 (0.6728–0.7784)	0.7423 (0.6918–0.7921)
			HD95		102.9 (88.85–116.2)	21.43 (13.85–30.37)	24.21 (15.95–33.43)	21.88 (14.33–30.05)	39.78 (28.49–52.17)	30.95 (21.79–41.64)
		kidney cyst	Dice	0.2168 (0.1476–0.2892)	0.04249 (0.01424–0.07457)	0.4887 (0.4021–0.5744)	0.0000 (0.0000–0.0000)	0.5587 (0.5342–0.6164)	0.5008 (0.4198–0.5839)	0.6101 (0.5298–0.6828)
			HD95	100.9 (84.82–118.7)	111.2 (93.01–129.0)	46.93 (32.80–60.68)	– (– –)	46.42 (32.77–60.41)	52.41 (38.00–66.49)	58.91 (45.17–75.34)
	KiPA22	renal vein	Dice	Not supported	0.6385 (0.6065–0.6700)	0.6772 (0.6098–0.7348)	0.8394 (0.8168–0.8607)	0.7543 (0.7246–0.7997)	0.7422 (0.6977–0.7823)	0.7976 (0.7622–0.8286)
			HD95		6.152 (5.076–7.427)	6.641 (4.734–9.093)	3.773 (2.805–4.862)	4.370 (1.470–7.038)	8.281 (6.031–11.29)	5.026 (3.795–6.334)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	KiPA22	kidney	Dice	0.8311 (0.6946–0.9087)	0.9259 (0.9194–0.9311)	0.8802 (0.8415–0.9135)	0.8849 (0.8324–0.9271)	0.9133 (0.9106–0.9198)	0.9377 (0.9277–0.9463)	0.9475 (0.9429–0.9517)
			HD95	10.85 (6.635–16.81)	3.969 (3.537–4.473)	11.22 (6.606–17.01)	13.44 (8.381–20.07)	4.564 (2.791–5.287)	5.803 (4.671–7.620)	3.989 (3.538–4.513)
		renal artery	Dice	Not supported	0.4047 (0.3813–0.4295)	0.6205 (0.5985–0.6397)	0.8634 (0.8353–0.8878)	0.8552 (0.8412–0.8781)	0.6887 (0.6540–0.7210)	0.6579 (0.6157–0.6935)
			HD95		16.05 (13.97–17.82)	10.48 (3.746–23.20)	2.765 (2.252–3.333)	14.08 (13.66–15.20)	12.18 (9.391–16.66)	14.27 (12.63–16.09)
		kidney tumor	Dice	Not supported	0.8362 (0.8121–0.8623)	0.5844 (0.3893–0.7493)	0.9332 (0.9230–0.9436)	0.8347 (0.8127–0.8627)	0.8031 (0.6565–0.9043)	0.8585 (0.8113–0.8995)
			HD95		4.719 (3.396–6.790)	7.071 (4.251–10.02)	1.877 (1.458–2.322)	4.924 (4.365–5.946)	12.79 (3.201–26.65)	3.950 (2.630–5.652)
	CHAOS	liver	Dice	0.8941 (0.8801–0.9133)	0.6998 (0.2360–0.9467)	0.8343 (0.8050–0.8637)	0.9393 (0.9117–0.9596)	0.9431 (0.9416–0.9511)	0.9687 (0.9632–0.9747)	0.9705 (0.9647–0.9763)
			HD95	25.69 (20.36–31.01)	42.36 (7.634–109.9)	34.21 (27.13–46.20)	24.71 (11.78–43.68)	7.890 (4.861–8.858)	5.325 (4.077–6.754)	4.235 (3.273–5.420)
		liver	Dice	0.9641 (0.9615–0.9668)	0.9058 (0.8919–0.9155)	0.9767 (0.9745–0.9788)	0.9804 (0.9760–0.9843)	0.9405 (0.9313–0.9570)	0.9933 (0.9897–0.9964)	0.9800 (0.9761–0.9837)
			HD95	3.763 (3.414–4.343)	7.101 (6.296–8.215)	2.737 (2.500–3.210)	2.174 (1.535–3.056)	3.819 (2.350–3.892)	1.681 (1.344–2.058)	1.978 (1.633–2.366)
		right kidney	Dice	0.8977 (0.8219–0.9402)	0.7498 (0.5355–0.8737)	0.9682 (0.9642–0.9720)	0.9718 (0.9669–0.9767)	0.9332 (0.9270–0.9352)	0.9826 (0.9777–0.9873)	0.9786 (0.9736–0.9828)
			HD95	11.48 (7.652–15.56)	34.20 (7.088–67.31)	3.524 (2.288–5.153)	5.591 (3.348–7.891)	5.163 (4.174–7.799)	6.805 (3.274–10.63)	2.718 (1.608–4.440)
	Flare22	spleen	Dice	0.9586 (0.9533–0.9629)	0.8574 (0.8214–0.8872)	0.9716 (0.9688–0.9747)	0.9842 (0.9789–0.9884)	0.9608 (0.9526–0.9680)	0.9957 (0.9920–0.9993)	0.9829 (0.9783–0.9868)
			HD95	2.681 (2.355–3.028)	5.692 (5.103–6.631)	1.897 (1.557–2.229)	0.7127 (0.4451–0.9165)	2.821 (1.739–4.213)	0.3951 (0.3166–0.4873)	0.8911 (0.8289–0.9557)
		pancreas	Dice	0.8610 (0.8419–0.8789)	0.6625 (0.6255–0.7008)	0.8756 (0.8417–0.8999)	0.8849 (0.8738–0.8976)	0.8796 (0.8791–0.8865)	0.9067 (0.8948–0.9199)	0.8954 (0.8828–0.9078)
			HD95	6.081 (4.531–8.499)	9.191 (7.637–10.66)	4.809 (3.029–7.594)	4.420 (3.218–5.906)	5.145 (3.820–5.795)	2.876 (2.330–3.532)	3.282 (2.854–3.748)
		aorta	Dice	0.9089 (0.8716–0.9357)	0.8390 (0.8211–0.8558)	0.9467 (0.9385–0.9538)	0.9490 (0.9297–0.9647)	0.9345 (0.9193–0.9471)	0.9740 (0.9669–0.9800)	0.9580 (0.9523–0.9633)
			HD95	19.71 (2.426–44.01)	4.260 (3.779–4.775)	7.923 (2.052–18.81)	17.30 (0.9245–40.96)	2.735 (1.477–3.997)	1.568 (0.3805–3.808)	2.132 (1.001–3.986)
		inferior vena cava	Dice	0.8892 (0.8712–0.9048)	0.7685 (0.7402–0.7943)	0.9176 (0.8962–0.9350)	0.9202 (0.9051–0.9342)	0.9197 (0.9076–0.9372)	0.9201 (0.9060–0.9320)	0.9323 (0.9197–0.9444)
			HD95	10.37 (7.105–13.70)	11.01 (7.534–15.36)	4.232 (2.713–6.623)	5.181 (3.222–7.485)	4.828 (3.311–6.897)	4.898 (2.498–9.024)	3.027 (2.193–3.995)
		right adrenal gland	Dice	0.7829 (0.7191–0.8333)	0.3339 (0.2752–0.3872)	0.8380 (0.8111–0.8601)	0.8405 (0.7783–0.8900)	0.8165 (0.7935–0.8579)	0.8623 (0.8305–0.8900)	0.8651 (0.8312–0.8934)
			HD95	4.653 (3.217–6.362)	9.848 (8.252–11.48)	1.873 (1.497–2.245)	2.657 (1.046–4.546)	5.358 (5.015–6.450)	1.375 (0.8850–1.825)	1.439 (0.8488–2.343)
		left adrenal gland	Dice	0.8070 (0.7741–0.8386)	0.3795 (0.3159–0.4437)	0.8379 (0.8198–0.8530)	0.8697 (0.8427–0.8933)	0.8312 (0.8061–0.8391)	0.8899 (0.8614–0.9124)	0.8824 (0.8639–0.8997)
			HD95	3.593 (2.985–4.223)	10.93 (8.756–13.26)	1.986 (1.741–2.268)	1.737 (1.210–2.288)	3.195 (2.459–4.858)	0.8690 (0.4833–1.403)	1.061 (0.9029–1.249)
		gallbladder	Dice	0.8971 (0.8781–0.9158)	0.5550 (0.3989–0.6810)	0.9145 (0.9001–0.9289)	0.9075 (0.8820–0.9307)	0.8999 (0.8932–0.9212)	0.9241 (0.8906–0.9546)	0.9264 (0.9067–0.9453)
			HD95	4.139 (3.308–5.492)	40.35 (14.06–81.01)	2.462 (2.147–2.803)	3.951 (2.295–6.022)	3.575 (3.009–4.955)	3.003 (1.487–4.684)	1.999 (1.489–2.491)
		esophagus	Dice	0.8154 (0.7851–0.8493)	0.6599 (0.6163–0.7015)	0.8747 (0.8584–0.8922)	0.8750 (0.8587–0.8918)	0.8678 (0.8554–0.8823)	0.9101 (0.8975–0.9245)	0.8956 (0.8779–0.9112)
			HD95	8.587 (6.280–11.48)	7.227 (5.508–9.251)	3.604 (2.553–5.137)	4.395 (2.854–6.410)	4.042 (2.782–6.122)	1.980 (1.608–2.418)	2.908 (2.324–3.628)
		stomach	Dice	0.9319 (0.9201–0.9423)	0.7885 (0.7380–0.8325)	0.9468 (0.9360–0.9547)	0.9426 (0.9222–0.9582)	0.9448 (0.9330–0.9638)	0.9745 (0.9695–0.9792)	0.9594 (0.9536–0.9644)
			HD95	5.138 (3.852–6.645)	8.958 (7.334–10.78)	2.772 (2.500–3.272)	11.32 (3.590–21.90)	4.460 (2.930–5.059)	2.056 (1.538–2.725)	2.498 (2.037–3.089)
		duodenum	Dice	0.7722 (0.7503–0.7922)	0.5796 (0.5136–0.6356)	0.8608 (0.8458–0.8768)	0.8318 (0.7925–0.8659)	0.8152 (0.7988–0.8438)	0.8767 (0.8590–0.8904)	0.8798 (0.8554–0.8996)
			HD95	18.59 (14.21–23.43)	15.03 (11.08–19.01)	5.419 (3.973–7.062)	10.13 (6.702–14.07)	8.918 (8.783–10.30)	8.105 (5.321–11.00)	7.327 (3.877–11.48)
	Flare23	left kidney	Dice	0.9306 (0.9220–0.9389)	0.7600 (0.5819–0.8691)	0.9643 (0.9601–0.9690)	0.9686 (0.9636–0.9733)	0.9599 (0.9522–0.9658)	0.9848 (0.9791–0.9899)	0.9724 (0.9680–0.9762)
			HD95	7.479 (5.235–10.17)	44.69 (7.154–83.34)	3.351 (2.272–5.016)	3.732 (2.504–5.516)	6.401 (4.397–6.936)	2.885 (1.222–5.654)	3.526 (1.912–6.077)
		liver	Dice	0.9691 (0.9646–0.9726)	0.6001 (0.5393–0.6586)	0.6568 (0.5946–0.7160)	0.6922 (0.6358–0.7502)	0.9611 (0.9511–0.9768)	0.9704 (0.9666–0.9735)	0.9691 (0.9653–0.9725)
			HD95	5.875 (4.074–8.386)	10.30 (9.352–11.49)	4.424 (3.477–5.855)	43.43 (35.94–51.26)	7.335 (6.506–8.890)	4.920 (4.004–5.988)	5.082 (3.973–6.568)
		right kidney	Dice	0.9007 (0.8823–0.9168)	0.4992 (0.4439–0.5478)	0.6187 (0.5595–0.6822)	0.6264 (0.5668–0.6880)	0.9141 (0.8885–0.9270)	0.9247 (0.9112–0.9369)	0.9402 (0.9291–0.9496)
			HD95	7.150 (5.863–9.209)	72.96 (62.02–83.64)	4.794 (4.075–5.582)	37.92 (30.98–45.05)	6.318 (5.618–8.355)	5.826 (4.816–7.288)	5.035 (4.036–6.458)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	Flare23	spleen	Dice	0.9660 (0.9632–0.9682)	0.5599 (0.5020–0.6130)	0.6536 (0.5897–0.7226)	0.6585 (0.5920–0.7232)	0.9358 (0.9349–0.9430)	0.9697 (0.9632–0.9742)	0.9710 (0.9644–0.9754)
			HD95	1.884 (1.690–2.127)	8.940 (7.697–10.48)	2.126 (1.719–2.682)	64.13 (51.45–77.15)	3.541 (2.754–5.397)	1.758 (1.289–2.371)	1.960 (1.378–2.712)
		pancreas	Dice	0.8412 (0.8253–0.8541)	0.3905 (0.3491–0.4333)	0.5791 (0.5182–0.6365)	0.5590 (0.5066–0.6136)	0.7998 (0.7802–0.8217)	0.8350 (0.8148–0.8536)	0.8413 (0.8233–0.8581)
			HD95	7.216 (5.620–8.998)	14.97 (13.02–17.07)	5.589 (4.236–7.698)	48.91 (40.23–58.21)	9.322 (6.771–13.14)	6.854 (5.338–9.014)	7.735 (5.902–9.948)
		left kidney	Dice	0.9177 (0.9064–0.9263)	0.5120 (0.4585–0.5625)	0.6223 (0.5664–0.6812)	0.6337 (0.5720–0.7006)	0.9320 (0.9183–0.9415)	0.9397 (0.9300–0.9481)	0.9429 (0.9377–0.9481)
			HD95	6.414 (5.542–7.527)	69.17 (58.26–80.57)	6.087 (4.263–8.764)	59.48 (47.84–70.57)	5.511 (5.217–5.902)	4.932 (3.995–6.139)	4.902 (4.095–5.903)
		tumor	Dice	Not supported	0.2622 (0.2207–0.3041)	0.003614 (0.0000–0.009096)	0.6045 (0.5621–0.6437)	0.5997 (0.5651–0.6359)	0.5915 (0.5384–0.6407)	0.6312 (0.5844–0.6753)
			HD95		89.03 (79.72–98.30)	182.6 (164.4–204.9)	83.64 (72.94–95.86)	42.56 (31.95–50.35)	48.99 (37.26–61.30)	48.20 (38.37–59.08)
		liver	Dice	0.9316 (0.9010–0.9556)	0.7965 (0.6988–0.8623)	0.9641 (0.9589–0.9690)	0.9605 (0.9485–0.9701)	0.9581 (0.9395–0.9640)	0.9749 (0.9666–0.9825)	0.9672 (0.9620–0.9721)
			HD95	27.59 (9.466–50.69)	30.93 (12.89–56.54)	3.290 (2.778–3.865)	23.18 (8.058–45.09)	9.563 (6.468–14.16)	6.221 (3.516–9.520)	4.011 (2.988–5.222)
		urinary bladder	Dice	0.8502 (0.7703–0.9115)	0.4785 (0.3376–0.6073)	0.9024 (0.8682–0.9321)	0.8595 (0.7708–0.9223)	0.8609 (0.8363–0.8913)	0.8749 (0.7966–0.9352)	0.8841 (0.8386–0.9257)
			HD95	6.424 (0.8233–14.46)	82.24 (23.46–148.7)	4.880 (3.572–6.444)	13.63 (6.651–21.27)	8.232 (7.286–10.79)	43.63 (7.129–91.16)	9.856 (5.227–17.64)
	CT-ORG	lung	Dice	0.9133 (0.8774–0.9397)	0.7051 (0.5866–0.8075)	0.9522 (0.9335–0.9644)	0.9545 (0.9289–0.9691)	0.9456 (0.9212–0.9644)	0.9773 (0.9537–0.9906)	0.9643 (0.9394–0.9788)
			HD95	31.53 (13.14–56.49)	41.02 (19.10–65.02)	16.85 (9.673–28.95)	25.57 (9.520–48.71)	17.32 (9.853–29.56)	22.12 (6.698–43.45)	20.29 (8.203–36.06)
		kidney	Dice	0.9106 (0.8971–0.9236)	0.6838 (0.5773–0.7704)	0.9318 (0.9226–0.9415)	0.9203 (0.9038–0.9369)	0.9035 (0.8704–0.9303)	0.9344 (0.9179–0.9487)	0.9265 (0.9118–0.9392)
			HD95	8.231 (6.746–9.961)	35.71 (12.53–65.52)	4.578 (3.622–5.602)	6.229 (5.015–7.709)	5.628 (5.247–6.297)	5.371 (4.331–6.530)	6.195 (5.112–7.321)
		bone	Dice	Not supported	0.6677 (0.5772–0.7437)	0.7172 (0.5945–0.8202)	0.8050 (0.7554–0.8400)	0.7921 (0.7578–0.8000)	0.8355 (0.7877–0.8683)	0.8362 (0.7889–0.8682)
			HD95		51.95 (32.43–75.51)	59.52 (34.62–87.62)	47.32 (33.95–63.23)	49.05 (35.84–64.77)	50.54 (34.01–66.62)	45.97 (30.15–62.23)
	LIDC-IDRI	pulmonary nodule	Dice	0.1598 (0.1188–0.2042)	0.01719 (0.004692–0.03352)	0.3663 (0.3045–0.4234)	0.5179 (0.4611–0.5771)	0.5425 (0.5271–0.5876)	0.6134 (0.5650–0.6601)	0.6614 (0.6189–0.7052)
			HD95	184.2 (175.7–193.1)	168.1 (158.5–178.5)	98.44 (84.44–113.4)	93.66 (79.00–108.9)	44.70 (27.31–52.22)	73.98 (60.32–88.28)	52.53 (40.93–64.77)
	SLIVER07	liver	Dice	0.9721 (0.9689–0.9761)	0.9190 (0.8940–0.9389)	0.9726 (0.9682–0.9755)	0.9750 (0.9730–0.9766)	0.9693 (0.9470–0.9726)	0.9747 (0.9714–0.9775)	0.9750 (0.9715–0.9777)
			HD95	4.239 (2.740–6.107)	8.254 (5.282–13.34)	3.323 (2.427–4.366)	4.555 (2.713–6.397)	3.822 (2.051–4.774)	4.581 (3.647–5.514)	4.650 (2.500–7.717)
	LUNA16	left lung	Dice	0.9715 (0.9601–0.9775)	0.8927 (0.8689–0.9134)	0.9771 (0.9656–0.9831)	0.9618 (0.9472–0.9699)	0.9599 (0.9526–0.9627)	0.9761 (0.9647–0.9821)	0.9776 (0.9662–0.9836)
			HD95	9.887 (8.527–11.54)	12.94 (9.781–17.60)	3.116 (1.802–5.193)	16.13 (13.92–18.31)	3.371 (2.861–4.843)	3.297 (2.306–5.138)	3.502 (2.526–5.543)
		right lung	Dice	0.9697 (0.9585–0.9760)	0.8924 (0.8692–0.9112)	0.9786 (0.9671–0.9844)	0.9695 (0.9568–0.9768)	0.9709 (0.9574–0.9718)	0.9767 (0.9654–0.9828)	0.9790 (0.9676–0.9850)
			HD95	19.75 (18.52–21.43)	20.69 (16.25–25.75)	2.747 (1.796–4.356)	16.06 (13.85–18.56)	7.386 (7.111–7.888)	10.49 (8.829–12.43)	4.958 (3.684–6.637)
		trachea	Dice	0.8383 (0.8327–0.8432)	0.6779 (0.6590–0.6912)	0.8947 (0.8905–0.8984)	0.8827 (0.8781–0.8871)	0.8760 (0.8616–0.8781)	0.8917 (0.8872–0.8957)	0.8867 (0.8819–0.8908)
			HD95	26.20 (24.61–27.70)	14.83 (12.97–17.35)	4.225 (3.455–5.332)	16.52 (15.19–18.00)	4.549 (1.616–6.051)	7.353 (6.351–8.514)	7.772 (6.780–8.957)
	COVID-19	left lung	Dice	0.4581 (0.0000–0.9162)	0.9024 (0.8516–0.9364)	0.8499 (0.7594–0.9349)	0.8199 (0.7280–0.9348)	0.9601 (0.9232–0.9764)	0.9515 (0.9115–0.9763)	0.9643 (0.9404–0.9802)
			HD95	13.09 (11.64–14.54)	10.91 (8.718–12.87)	17.46 (7.221–27.70)	49.84 (13.84–85.84)	6.010 (3.455–9.142)	7.263 (4.123–10.54)	5.421 (3.152–8.301)
		right lung	Dice	0.4656 (0.0000–0.9312)	0.8864 (0.8363–0.9365)	0.8934 (0.8336–0.9552)	0.7656 (0.5692–0.9375)	0.9322 (0.8986–0.9406)	0.9588 (0.9403–0.9773)	0.9614 (0.9409–0.9812)
			HD95	197.8 (13.38–382.2)	16.60 (7.227–29.82)	45.13 (3.688–86.57)	54.55 (12.80–96.30)	5.748 (4.957–6.569)	7.403 (3.519–13.74)	5.415 (2.494–8.288)
	CTPelvic1K-CLINIC	covid-19 infection	Dice	Not supported	0.2189 (0.02432–0.4707)	0.4256 (0.2012–0.6500)	0.6341 (0.4630–0.8052)	0.7247 (0.6373–0.7767)	0.5948 (0.4876–0.6919)	0.7218 (0.6176–0.7959)
			HD95		103.3 (39.29–167.4)	82.85 (25.23–164.6)	97.60 (38.15–157.1)	18.73 (15.90–24.00)	22.19 (14.24–30.14)	18.95 (9.096–29.27)
		sacrum	Dice	0.9256 (0.9225–0.9283)	0.5029 (0.3841–0.6059)	0.9384 (0.9360–0.9405)	0.9586 (0.9472–0.9655)	0.9360 (0.9310–0.9441)	0.9370 (0.9271–0.9460)	0.9511 (0.9465–0.9548)
			HD95	2.561 (2.381–2.770)	49.60 (27.72–74.25)	1.298 (1.227–1.376)	1.395 (1.069–1.837)	3.424 (2.945–3.439)	2.729 (2.028–3.490)	1.565 (1.215–2.023)
		left hip	Dice	0.9402 (0.9369–0.9428)	0.5004 (0.3707–0.6192)	0.9565 (0.9543–0.9586)	0.9782 (0.9745–0.9816)	0.9418 (0.9363–0.9463)	0.9747 (0.9700–0.9791)	0.9701 (0.9663–0.9735)
			HD95	2.218 (2.118–2.336)	73.94 (46.72–104.6)	1.060 (0.9921–1.140)	2.297 (0.8213–5.165)	2.155 (1.959–3.059)	19.88 (9.540–31.90)	1.465 (0.9573–2.127)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	CTPelvic1K-CLINIC	right hip	Dice	0.9375 (0.9311–0.9418)	0.5015 (0.3707–0.6242)	0.9542 (0.9487–0.9578)	0.9736 (0.9660–0.9798)	0.9351 (0.9342–0.9385)	0.9768 (0.9687–0.9829)	0.9659 (0.9576–0.9722)
			HD95	2.378 (2.152–2.743)	83.44 (59.90–107.8)	1.249 (1.118–1.468)	2.782 (0.8894–6.253)	1.810 (1.422–3.800)	1.184 (0.5340–1.983)	1.642 (0.9241–2.449)
		lumbar vertebrae	Dice	0.9270 (0.9242–0.9297)	0.4387 (0.3573–0.5134)	0.9345 (0.9307–0.9379)	0.9567 (0.9370–0.9698)	0.9307 (0.9277–0.9321)	0.9685 (0.9650–0.9715)	0.9586 (0.9557–0.9612)
			HD95	2.241 (2.172–2.317)	60.96 (32.64–91.73)	1.238 (1.172–1.317)	3.053 (0.8704–6.514)	2.100 (1.898–3.499)	7.106 (1.050–17.57)	1.762 (1.110–2.606)
		sacrum	Dice	0.9167 (0.9071–0.9252)	0.4816 (0.4426–0.5208)	0.9264 (0.9148–0.9337)	0.9728 (0.9703–0.9750)	0.9534 (0.9396–0.9638)	0.9229 (0.9171–0.9282)	0.9499 (0.9469–0.9527)
			HD95	4.375 (2.930–6.452)	37.55 (29.66–45.36)	1.956 (1.552–2.511)	1.888 (1.196–2.904)	4.080 (1.425–5.627)	3.865 (3.332–4.458)	1.860 (1.567–2.237)
	CTPelvic1K-COLONOG	left hip	Dice	0.9365 (0.9156–0.9505)	0.6547 (0.6241–0.6819)	0.9259 (0.9062–0.9403)	0.9778 (0.9706–0.9827)	0.9437 (0.9305–0.9530)	0.9672 (0.9646–0.9692)	0.9672 (0.9647–0.9691)
			HD95	4.987 (1.691–9.549)	22.32 (17.58–27.97)	4.406 (1.686–8.144)	6.495 (2.837–11.00)	4.124 (3.563–5.920)	14.65 (7.572–23.26)	2.178 (0.8370–4.386)
		right hip	Dice	0.9354 (0.9149–0.9493)	0.5441 (0.5089–0.5786)	0.9204 (0.8994–0.9344)	0.9786 (0.9711–0.9838)	0.9640 (0.9524–0.9735)	0.9699 (0.9678–0.9716)	0.9688 (0.9666–0.9706)
			HD95	6.173 (1.356–11.96)	78.93 (66.81–90.86)	4.478 (1.846–8.194)	4.655 (1.458–8.643)	2.250 (2.000–2.495)	1.284 (0.8303–2.107)	1.165 (0.7772–1.921)
		lumbar vertebrae	Dice	0.6543 (0.6368–0.6713)	0.4506 (0.4263–0.4763)	0.6484 (0.6305–0.6656)	0.6807 (0.6622–0.6982)	0.8248 (0.8189–0.8414)	0.8492 (0.8376–0.8605)	0.8789 (0.8676–0.8895)
			HD95	71.52 (67.80–75.50)	42.72 (37.14–48.64)	71.27 (65.67–76.88)	70.55 (66.77–74.30)	13.84 (11.67–15.80)	16.23 (12.62–21.00)	18.69 (14.14–24.03)
	CTPelvic1K-KiTS	sacrum	Dice	0.8322 (0.8197–0.8435)	0.04111 (0.007349–0.08688)	0.9261 (0.9093–0.9373)	0.9551 (0.9395–0.9679)	0.9488 (0.9252–0.9754)	0.8771 (0.8501–0.9027)	0.8931 (0.8613–0.9218)
			HD95	13.02 (4.269–25.44)	176.3 (119.1–262.9)	2.667 (1.646–4.170)	3.031 (1.005–5.375)	3.463 (2.920–4.026)	7.240 (4.790–9.987)	4.972 (2.990–7.384)
		left hip	Dice	0.9415 (0.9245–0.9548)	0.3451 (0.2494–0.4227)	0.9348 (0.9176–0.9481)	0.9645 (0.9469–0.9784)	0.9096 (0.9017–0.9297)	0.9521 (0.9344–0.9673)	0.9492 (0.9334–0.9635)
			HD95	5.851 (2.026–10.81)	127.2 (86.23–164.7)	4.291 (1.971–7.867)	5.406 (1.581–10.87)	6.387 (3.327–9.970)	8.481 (3.362–15.01)	5.727 (1.729–10.87)
		right hip	Dice	0.9263 (0.9063–0.9437)	0.2479 (0.1970–0.3053)	0.9275 (0.9084–0.9413)	0.9640 (0.9433–0.9791)	0.9218 (0.9012–0.9313)	0.9518 (0.9336–0.9672)	0.9463 (0.9281–0.9623)
			HD95	12.11 (0.8379–23.78)	163.3 (149.9–176.0)	4.959 (2.083–9.008)	5.240 (0.8015–11.77)	5.478 (3.100–9.060)	6.196 (1.211–11.92)	5.491 (1.114–11.13)
		lumbar vertebrae	Dice	0.4156 (0.2920–0.5208)	0.2165 (0.1326–0.2947)	0.4105 (0.2856–0.5123)	0.4302 (0.3150–0.5321)	0.6397 (0.5184–0.7467)	0.7022 (0.5160–0.8399)	0.7034 (0.5083–0.8530)
			HD95	177.6 (105.0–275.3)	120.2 (98.67–143.7)	115.1 (94.13–138.2)	116.8 (93.39–140.4)	31.97 (18.81–45.16)	79.73 (18.67–183.5)	85.11 (29.82–157.5)
		sacrum	Dice	0.8123 (0.8038–0.8204)	0.5830 (0.5012–0.6306)	0.9299 (0.9219–0.9369)	0.9133 (0.8967–0.9271)	0.8991 (0.8981–0.9109)	0.8963 (0.8753–0.9153)	0.9262 (0.9121–0.9385)
			HD95	9.761 (3.524–18.13)	11.09 (9.948–12.43)	1.977 (1.676–2.319)	5.347 (3.558–7.420)	5.670 (5.397–6.162)	5.000 (3.683–6.763)	2.742 (1.923–3.721)
	CTPelvic1K-MSD-colon	left hip	Dice	0.9388 (0.9344–0.9432)	0.6900 (0.6739–0.7079)	0.9467 (0.9428–0.9501)	0.9606 (0.9553–0.9660)	0.9365 (0.9284–0.9376)	0.9559 (0.9505–0.9609)	0.9564 (0.9506–0.9621)
			HD95	2.780 (2.444–3.165)	10.23 (9.631–10.86)	1.866 (1.673–2.045)	1.927 (1.295–2.754)	2.556 (1.848–3.487)	12.07 (3.318–24.79)	13.23 (1.152–36.98)
		right hip	Dice	0.9351 (0.9288–0.9406)	0.5504 (0.4703–0.6315)	0.9451 (0.9408–0.9485)	0.9614 (0.9566–0.9656)	0.9607 (0.9408–0.9680)	0.9582 (0.9514–0.9640)	0.9574 (0.9511–0.9629)
			HD95	7.501 (1.880–13.97)	89.88 (55.56–125.0)	2.533 (2.149–2.945)	1.470 (1.105–1.875)	1.996 (1.885–2.415)	1.483 (1.065–1.979)	1.307 (0.9499–1.722)
		lumbar vertebrae	Dice	0.6423 (0.6073–0.6792)	0.4567 (0.4032–0.5113)	0.6598 (0.6252–0.6975)	0.6547 (0.6184–0.6935)	0.8135 (0.7884–0.8406)	0.8361 (0.8059–0.8649)	0.8621 (0.8409–0.8818)
			HD95	131.2 (87.25–182.2)	36.53 (30.39–41.88)	74.85 (60.45–92.93)	74.60 (65.73–83.01)	28.42 (22.75–32.50)	40.53 (12.76–82.62)	60.81 (22.05–107.9)
		sacrum	Dice	0.4370 (0.2470–0.6753)	0.5539 (0.4622–0.6372)	0.3867 (0.2177–0.6611)	0.9445 (0.9389–0.9503)	0.8933 (0.8853–0.9094)	0.9034 (0.8796–0.9255)	0.9362 (0.9263–0.9461)
			HD95	46.58 (28.10–61.85)	25.90 (9.754–55.15)	23.82 (12.55–32.32)	1.963 (1.300–2.666)	5.003 (4.326–5.702)	3.747 (2.353–5.142)	2.013 (1.477–2.464)
		left hip	Dice	0.2018 (0.007974–0.5748)	0.6768 (0.6454–0.7075)	0.1916 (0.0000–0.5744)	0.6785 (0.5911–0.8250)	0.9227 (0.9009–0.9369)	0.9613 (0.9570–0.9656)	0.9634 (0.9588–0.9662)
			HD95	138.7 (70.13–174.7)	36.93 (8.907–91.85)	136.8 (68.21–175.2)	128.8 (64.06–163.9)	8.413 (5.008–13.63)	1.299 (0.9766–1.782)	0.9766 (0.9766–0.9766)
	CTPelvic1K-BTCV-Cervix	right hip	Dice	0.1917 (0.0005502–0.5721)	0.6283 (0.4701–0.7265)	0.1913 (0.0000–0.5738)	0.6867 (0.5979–0.8326)	0.9437 (0.9250–0.9499)	0.9627 (0.9591–0.9664)	0.9651 (0.9600–0.9687)
			HD95	140.2 (71.21–178.5)	39.70 (8.653–95.46)	139.1 (68.41–177.6)	131.3 (65.72–167.0)	2.374 (1.948–2.895)	0.9766 (0.9766–0.9766)	0.9766 (0.9766–0.9766)
		lumbar vertebrae	Dice	0.5000 (0.4508–0.5457)	0.3137 (0.1782–0.4369)	0.2146 (0.1487–0.3195)	0.4517 (0.4214–0.4925)	0.8175 (0.7714–0.8341)	0.8160 (0.7863–0.8453)	0.8381 (0.8097–0.8721)
			HD95	96.09 (81.57–111.2)	44.58 (23.63–77.29)	119.6 (96.00–145.8)	116.7 (96.66–138.3)	12.94 (8.876–17.44)	20.64 (13.15–28.54)	9.199 (4.623–14.39)
	CTPelvic1K-BTCV-Abdomen	sacrum	Dice	0.8126 (0.8035–0.8223)	0.6179 (0.5610–0.6676)	0.9303 (0.9201–0.9427)	0.9303 (0.9190–0.9433)	0.9166 (0.8984–0.9402)	0.8944 (0.8712–0.9156)	0.9267 (0.9187–0.9366)
			HD95	13.71 (11.86–15.51)	10.82 (8.072–15.71)	1.838 (1.330–2.281)	2.132 (1.593–2.638)	1.724 (1.570–2.418)	3.757 (2.663–4.834)	2.600 (1.822–3.518)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	CTPelvic1K-BTCV-Abdomen	right hip	Dice	0.9434 (0.9406–0.9465)	0.5621 (0.4270–0.6891)	0.9445 (0.9398–0.9490)	0.9635 (0.9615–0.9660)	0.9444 (0.9325–0.9526)	0.9633 (0.9593–0.9673)	0.9618 (0.9585–0.9656)
			HD95	2.030 (1.566–2.608)	77.01 (24.33–133.3)	2.506 (2.152–2.860)	1.178 (0.8530–1.578)	3.053 (2.728–3.892)	0.9573 (0.7570–1.196)	0.7812 (0.6963–0.8934)
		left hip	Dice	0.9433 (0.9392–0.9480)	0.7105 (0.6835–0.7331)	0.9494 (0.9428–0.9556)	0.9665 (0.9615–0.9721)	0.9628 (0.9467–0.9664)	0.9609 (0.9564–0.9639)	0.9628 (0.9588–0.9665)
			HD95	1.837 (1.398–2.340)	25.20 (7.329–59.87)	1.860 (1.376–2.374)	1.174 (0.7265–1.740)	1.977 (1.678–3.409)	13.83 (1.662–35.05)	0.8535 (0.6921–1.119)
		lumbar vertebrae	Dice	0.6483 (0.5893–0.6862)	0.5031 (0.3993–0.5843)	0.6577 (0.5978–0.6961)	0.6505 (0.5820–0.6922)	0.8457 (0.8338–0.8692)	0.8485 (0.8155–0.8791)	0.8646 (0.8387–0.8850)
			HD95	74.49 (65.26–87.57)	26.58 (20.62–32.10)	67.51 (57.34–81.17)	72.12 (63.41–85.30)	9.361 (6.437–11.02)	12.14 (4.684–19.73)	13.15 (7.635–19.32)
		caudate lobe	Dice	Not supported	0.05244 (0.03391–0.07375)	0.8621 (0.8449–0.8742)	0.5513 (0.5232–0.5778)	0.7882 (0.7738–0.8122)	0.8315 (0.8183–0.8451)	0.8458 (0.8356–0.8550)
			HD95		66.12 (52.78–79.67)	4.882 (4.492–5.397)	25.19 (23.76–26.63)	11.07 (9.365–12.63)	6.885 (5.930–7.957)	5.606 (5.166–6.105)
		left lateral superior segment of liver	Dice	Not supported	0.0000 (0.0000–0.0000)	0.8844 (0.8526–0.9082)	0.8186 (0.7894–0.8446)	0.8431 (0.8172–0.8786)	0.8855 (0.8612–0.9054)	0.9081 (0.8965–0.9182)
			HD95		130.8 (119.4–140.7)	6.506 (5.203–8.467)	13.35 (10.41–16.79)	11.34 (11.15–12.84)	8.853 (6.911–11.06)	6.609 (5.605–7.926)
		left lateral inferior segment of liver	Dice	Not supported	0.0000 (0.0000–0.0000)	0.8633 (0.8306–0.8897)	0.7637 (0.7110–0.8091)	0.8613 (0.8499–0.8662)	0.8689 (0.8181–0.9059)	0.8752 (0.8227–0.9103)
			HD95		160.1 (147.1–172.3)	7.674 (5.527–10.47)	16.61 (12.18–22.01)	11.69 (6.601–16.43)	8.620 (5.486–12.78)	8.003 (4.927–12.22)
	Couinaud-Liver (MSD-hepaticvessel)	left medial segment of liver	Dice	Not supported	0.0000 (0.0000–0.0000)	0.8793 (0.8449–0.9006)	0.7888 (0.7694–0.8056)	0.8166 (0.7923–0.8276)	0.8623 (0.8479–0.8755)	0.8699 (0.8588–0.8812)
			HD95		158.9 (149.3–169.0)	5.829 (4.613–8.033)	18.53 (14.25–23.90)	10.69 (9.193–13.24)	9.761 (7.534–12.38)	6.615 (5.937–7.351)
		right anterior inferior segment of liver	Dice	Not supported	0.008415 (0.0006640–0.02088)	0.8437 (0.7886–0.8820)	0.7630 (0.7210–0.7992)	0.7886 (0.7575–0.8248)	0.8365 (0.8013–0.8647)	0.8519 (0.8172–0.8793)
			HD95		76.84 (70.87–83.20)	8.190 (6.597–10.21)	18.42 (15.94–20.89)	16.44 (13.75–18.89)	14.78 (10.61–19.93)	11.00 (9.245–13.00)
		right posterior inferior segment of liver	Dice	Not supported	0.04553 (0.009687–0.1063)	0.8694 (0.8427–0.8931)	0.7636 (0.7207–0.8006)	0.8639 (0.8374–0.8948)	0.8424 (0.7983–0.8801)	0.8725 (0.8466–0.8944)
			HD95		70.60 (60.32–82.29)	8.832 (6.639–11.83)	20.14 (16.09–24.23)	12.44 (10.95–15.26)	13.63 (10.23–18.03)	20.26 (9.812–33.12)
		right posterior superior segment of liver	Dice	Not supported	0.006829 (0.002929–0.01113)	0.9041 (0.8844–0.9180)	0.8505 (0.8293–0.8686)	0.8922 (0.8832–0.9034)	0.8911 (0.8715–0.9072)	0.9040 (0.8872–0.9177)
			HD95		89.51 (81.71–98.33)	10.77 (6.270–18.13)	17.83 (14.81–21.00)	10.90 (8.603–13.88)	13.42 (8.863–20.29)	8.250 (6.491–10.23)
		right anterior superior segment of liver	Dice	Not supported	0.004849 (0.0009141–0.01040)	0.8746 (0.8429–0.8976)	0.8204 (0.8017–0.8392)	0.8556 (0.8379–0.8692)	0.8687 (0.8398–0.8931)	0.8845 (0.8617–0.9026)
			HD95		103.7 (95.49–111.1)	10.57 (8.080–13.17)	19.97 (16.64–23.79)	15.29 (13.74–18.74)	11.37 (8.349–14.73)	9.916 (7.405–13.54)
	DAP-Atlas (AutoPETIII)	muscle	Dice	Not supported	0.009012 (0.007278–0.01067)	0.9442 (0.9393–0.9479)	0.6639 (0.6493–0.6773)	0.8536 (0.8337–0.8682)	0.8676 (0.8601–0.8745)	0.8651 (0.8570–0.8718)
			HD95		466.8 (446.4–488.7)	2.627 (2.507–2.831)	25.29 (24.41–26.18)	15.16 (14.71–17.95)	17.72 (16.62–19.07)	13.29 (12.13–14.67)
		fat	Dice	Not supported	0.000003231 (0.0000001422–0.00000861)	0.9530 (0.9472–0.9581)	0.9329 (0.9251–0.9398)	0.9176 (0.9172–0.9253)	0.8737 (0.8616–0.8870)	0.9019 (0.8913–0.9126)
			HD95		424.1 (411.2–435.9)	2.352 (2.065–2.770)	6.006 (4.827–7.329)	3.452 (2.171–4.842)	33.98 (30.35–37.47)	12.01 (10.72–13.58)
		abdominal tissue	Dice	Not supported	0.1121 (0.1023–0.1223)	0.9130 (0.9000–0.9252)	0.9089 (0.8924–0.9226)	0.9082 (0.9062–0.9346)	0.8677 (0.8498–0.8839)	0.8838 (0.8647–0.9009)
			HD95		93.79 (87.51–100.8)	2.463 (2.414–2.501)	4.481 (3.528–5.696)	2.562 (2.358–2.799)	61.29 (52.29–70.11)	7.783 (5.778–10.78)
		mediastinal tissue	Dice	Not supported	0.01913 (0.01386–0.02495)	0.8406 (0.8240–0.8550)	0.8088 (0.7902–0.8271)	0.8281 (0.8243–0.8390)	0.7647 (0.7364–0.7870)	0.7541 (0.7291–0.7754)
			HD95		115.5 (104.8–125.9)	2.454 (2.398–2.507)	4.358 (3.688–5.157)	4.664 (4.238–5.142)	10.64 (8.659–13.02)	9.545 (8.093–11.34)
		esophagus	Dice	0.8334 (0.8064–0.8561)	0.3318 (0.3007–0.3631)	0.8882 (0.8724–0.9004)	0.8666 (0.8387–0.8902)	0.8867 (0.8833–0.9150)	0.8658 (0.8520–0.8792)	0.8718 (0.8541–0.8845)
			HD95	7.785 (4.249–13.11)	20.51 (17.41–25.60)	4.763 (2.058–9.639)	5.840 (2.535–10.72)	8.665 (5.849–13.99)	5.533 (2.574–10.46)	7.644 (2.317–14.78)
		stomach	Dice	0.9558 (0.9508–0.9599)	0.5807 (0.5450–0.6148)	0.1405 (0.1128–0.1707)	0.9612 (0.9568–0.9652)	0.9418 (0.9209–0.9448)	0.9587 (0.9542–0.9628)	0.9592 (0.9552–0.9626)
			HD95	3.451 (2.725–4.602)	23.37 (21.81–24.92)	442.4 (431.1–453.4)	12.69 (2.489–32.64)	5.174 (3.756–5.865)	3.178 (2.596–4.019)	3.019 (2.577–3.584)
		small bowel	Dice	0.8767 (0.8629–0.8892)	0.5020 (0.4746–0.5284)	0.9317 (0.9264–0.9364)	0.9037 (0.8927–0.9143)	0.8993 (0.8836–0.9117)	0.9012 (0.8903–0.9112)	0.8912 (0.8794–0.9030)
			HD95	8.799 (6.805–11.01)	31.53 (26.31–39.00)	3.068 (2.762–3.406)	13.91 (8.500–24.46)	6.194 (4.839–7.272)	7.498 (6.103–9.162)	34.70 (14.57–64.18)
		duodenum	Dice	0.8508 (0.8252–0.8758)	0.3621 (0.3387–0.3880)	0.8825 (0.8629–0.9013)	0.8378 (0.8016–0.8694)	0.8638 (0.8492–0.8826)	0.8685 (0.8398–0.8929)	0.8737 (0.8511–0.8936)
			HD95	7.366 (5.589–9.131)	28.14 (25.37–31.42)	3.741 (3.022–4.630)	13.48 (6.915–24.70)	4.752 (3.560–4.821)	5.309 (3.678–7.542)	5.392 (3.692–7.559)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	DAP-Atlas (AutoPETIII)	colon	Dice	0.8806 (0.8703–0.8906)	0.4457 (0.4170–0.4758)	0.9305 (0.9241–0.9361)	0.8980 (0.8870–0.9091)	0.8988 (0.8910–0.9105)	0.8595 (0.8480–0.8700)	0.8968 (0.8865–0.9068)
			HD95	14.56 (10.51–19.00)	35.16 (31.71–39.08)	3.751 (3.077–4.553)	12.62 (10.43–14.88)	5.098 (3.280–5.356)	25.80 (12.54–44.31)	11.13 (9.050–13.28)
		gallbladder	Dice	0.7116 (0.6030–0.8096)	0.2240 (0.1644–0.2843)	0.7513 (0.6483–0.8458)	0.7276 (0.6180–0.8221)	0.7216 (0.6354–0.7908)	0.7644 (0.6627–0.8548)	0.7963 (0.6946–0.8794)
			HD95	52.06 (19.69–91.94)	45.72 (30.71–62.92)	19.45 (7.457–33.90)	20.94 (8.252–37.47)	19.13 (6.383–35.00)	18.53 (5.304–34.10)	47.14 (15.98–84.82)
		liver	Dice	0.9533 (0.9497–0.9564)	0.7686 (0.7593–0.7778)	0.9743 (0.9731–0.9755)	0.9731 (0.9720–0.9741)	0.9499 (0.9453–0.9786)	0.9754 (0.9742–0.9766)	0.9766 (0.9753–0.9778)
			HD95	36.18 (33.48–38.45)	22.49 (21.61–23.34)	2.630 (2.521–2.772)	2.685 (2.534–2.883)	3.681 (3.553–3.994)	2.784 (2.530–3.078)	2.841 (2.570–3.172)
		pancreas	Dice	0.8783 (0.8414–0.9024)	0.3736 (0.3276–0.4214)	0.9049 (0.8885–0.9168)	0.9076 (0.8739–0.9272)	0.8815 (0.8702–0.9068)	0.9042 (0.8762–0.9230)	0.9094 (0.8857–0.9252)
			HD95	18.26 (3.672–40.21)	27.17 (22.38–33.01)	3.297 (2.511–4.762)	4.415 (2.587–7.261)	3.244 (1.737–5.426)	3.941 (2.447–6.417)	3.664 (2.413–5.915)
		left kidney	Dice	0.8996 (0.8241–0.9482)	0.6216 (0.5649–0.6664)	0.9170 (0.8417–0.9669)	0.9106 (0.8286–0.9666)	0.9017 (0.8391–0.9522)	0.9235 (0.8437–0.9764)	0.9206 (0.8390–0.9753)
			HD95	14.40 (4.297–31.94)	64.59 (47.99–81.97)	2.729 (1.932–3.832)	3.872 (3.090–4.937)	3.554 (2.913–3.976)	2.856 (1.718–4.516)	2.579 (1.991–3.551)
		right kidney	Dice	0.9523 (0.9486–0.9554)	0.6754 (0.6641–0.6864)	0.9600 (0.9578–0.9619)	0.9621 (0.9591–0.9651)	0.9518 (0.9410–0.9584)	0.9742 (0.9694–0.9778)	0.9744 (0.9713–0.9767)
			HD95	4.462 (4.022–4.982)	19.74 (16.64–25.03)	2.233 (2.109–2.385)	3.537 (3.000–4.014)	3.614 (2.605–4.989)	1.817 (1.454–2.278)	2.057 (1.675–2.531)
		urinary bladder	Dice	0.8749 (0.8455–0.8997)	0.2285 (0.1778–0.2836)	0.9122 (0.8972–0.9259)	0.8725 (0.8457–0.8974)	0.9087 (0.8810–0.9135)	0.9169 (0.9014–0.9313)	0.9252 (0.9103–0.9382)
			HD95	9.717 (4.621–15.97)	39.48 (34.13–45.19)	77.13 (25.85–136.6)	5.728 (4.047–7.749)	8.880 (7.783–9.757)	15.00 (4.148–30.97)	4.147 (3.354–5.022)
		gonad	Dice	Not supported	0.0000 (0.0000–0.0000)	0.4151 (0.2881–0.5385)	0.0000 (0.0000–0.0000)	0.08021 (0.07733–0.09382)	0.08758 (0.04011–0.1419)	0.2624 (0.1761–0.3503)
			HD95		662.0 (546.4–766.3)	60.44 (14.06–137.5)	126.7 (116.4–136.6)	62.12 (51.61–72.65)	270.4 (158.5–392.1)	99.54 (24.87–200.8)
		prostate	Dice	0.5521 (0.5009–0.6028)	0.05424 (0.02490–0.09018)	0.8669 (0.8369–0.8920)	0.6782 (0.6143–0.7346)	0.8266 (0.7815–0.8470)	0.8233 (0.8008–0.8445)	0.7919 (0.7585–0.8231)
			HD95	38.65 (35.88–40.70)	107.4 (50.62–187.2)	3.131 (2.783–3.517)	8.583 (6.780–10.48)	4.435 (3.701–5.808)	4.100 (3.547–4.693)	8.998 (4.130–17.98)
		uterocervix	Dice	Not supported	0.03835 (0.0006883–0.09554)	0.5232 (0.2860–0.7446)	0.3348 (0.1187–0.5857)	0.4204 (0.2071–0.6275)	0.4032 (0.1875–0.6289)	0.4939 (0.2622–0.6914)
			HD95		87.91 (60.61–116.0)	13.31 (5.590–23.16)	28.83 (13.92–45.34)	13.64 (7.953–19.65)	10.13 (5.014–15.69)	30.20 (9.509–62.70)
		uterus	Dice	Not supported	0.03145 (0.01007–0.05463)	0.7764 (0.7290–0.8254)	0.5552 (0.3767–0.6999)	0.6871 (0.6111–0.7341)	0.6952 (0.6120–0.7699)	0.6916 (0.6188–0.7614)
			HD95		83.43 (63.24–103.3)	6.711 (4.551–9.036)	19.73 (13.89–27.05)	17.16 (11.97–23.77)	70.74 (8.865–185.0)	160.7 (64.95–273.2)
		left breast	Dice	Not supported	0.000007628 (0.0000–0.00002248)	0.6123 (0.4964–0.7146)	0.3997 (0.2744–0.5219)	0.4434 (0.3568–0.5258)	0.5491 (0.4311–0.6567)	0.5137 (0.3931–0.6339)
			HD95		234.2 (184.0–291.2)	74.99 (23.99–142.6)	238.1 (168.4–308.6)	76.49 (24.82–134.2)	84.46 (27.78–151.1)	95.46 (28.55–166.3)
		right breast	Dice	Not supported	0.00001726 (0.0000–0.00005052)	0.5869 (0.4758–0.6875)	0.4356 (0.3264–0.5410)	0.5751 (0.5716–0.5755)	0.5866 (0.4781–0.6790)	0.5813 (0.4722–0.6745)
			HD95		225.7 (184.4–279.7)	63.03 (15.06–123.3)	133.3 (77.95–197.3)	73.43 (28.16–129.5)	91.85 (34.08–150.2)	82.47 (31.89–137.1)
		spinal canal	Dice	Not supported	0.003224 (0.001240–0.005448)	0.9183 (0.9096–0.9257)	0.9117 (0.9025–0.9211)	0.9088 (0.8975–0.9179)	0.8165 (0.8012–0.8312)	0.8859 (0.8767–0.8949)
			HD95		239.9 (200.4–284.4)	4.027 (2.050–6.332)	19.81 (12.45–28.53)	4.881 (2.757–6.233)	83.49 (74.82–92.87)	34.16 (24.18–44.81)
		brain	Dice	0.9400 (0.9283–0.9506)	0.6182 (0.5582–0.6695)	0.9676 (0.9623–0.9722)	0.9737 (0.9706–0.9765)	0.9694 (0.9648–0.9705)	0.9609 (0.9556–0.9662)	0.9643 (0.9575–0.9700)
			HD95	16.89 (6.629–32.19)	31.85 (28.54–35.59)	3.190 (2.786–3.623)	2.507 (2.309–2.747)	3.713 (3.205–4.631)	196.0 (115.2–279.2)	12.58 (3.227–39.19)
		spleen	Dice	0.9499 (0.9339–0.9613)	0.6603 (0.6223–0.6907)	0.9418 (0.9306–0.9510)	0.9718 (0.9640–0.9773)	0.9442 (0.9223–0.9583)	0.9692 (0.9600–0.9751)	0.9691 (0.9579–0.9772)
			HD95	21.75 (16.10–27.39)	21.85 (18.92–26.96)	256.0 (195.7–307.9)	1.332 (0.9155–2.103)	4.887 (4.337–6.546)	2.408 (1.113–4.338)	1.910 (1.040–3.494)
		left adrenal gland	Dice	0.8327 (0.8116–0.8518)	0.1565 (0.1167–0.1958)	0.8206 (0.7986–0.8381)	0.8887 (0.8674–0.9071)	0.8685 (0.8336–0.8875)	0.8702 (0.8517–0.8859)	0.8766 (0.8611–0.8905)
			HD95	3.606 (2.626–4.746)	24.79 (21.18–29.10)	25.73 (2.279–60.62)	2.669 (1.674–3.914)	3.688 (3.646–4.524)	2.196 (1.684–2.849)	1.970 (1.358–2.897)
		right adrenal gland	Dice	0.7753 (0.7159–0.8191)	0.1912 (0.1523–0.2289)	0.7763 (0.7201–0.8211)	0.8503 (0.8003–0.8921)	0.8419 (0.8086–0.8763)	0.8194 (0.7646–0.8626)	0.8461 (0.7950–0.8816)
			HD95	9.532 (2.948–22.14)	20.92 (18.48–23.75)	2.998 (2.077–4.106)	2.036 (1.403–2.885)	6.361 (6.070–7.654)	4.137 (2.287–6.342)	6.709 (2.144–12.68)
		left thyroid	Dice	Not supported	0.0000 (0.0000–0.0000)	0.6797 (0.5567–0.7816)	0.6112 (0.5027–0.7127)	0.6586 (0.5705–0.7366)	0.6879 (0.5831–0.7820)	0.7009 (0.5917–0.7938)
			HD95		292.4 (267.2–317.5)	76.53 (18.33–155.6)	15.09 (6.298–27.55)	13.83 (5.222–24.32)	12.40 (3.720–24.98)	12.10 (3.305–25.10)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	DAP-Atlas (AutoPETIII)	right thyroid	Dice	Not supported	0.0000 (0.0000–0.0000)	0.7590 (0.6559–0.8451)	0.7041 (0.6002–0.7992)	0.7506 (0.7306–0.7707)	0.7456 (0.6431–0.8304)	0.7217 (0.6104–0.8212)
			HD95		281.9 (271.3–294.9)	17.40 (6.194–31.19)	11.56 (4.541–21.11)	11.57 (5.551–20.82)	7.794 (3.159–15.42)	10.30 (2.984–20.29)
		thymus	Dice	Not supported	0.0000 (0.0000–0.0000)	0.4088 (0.2787–0.5439)	0.0000 (0.0000–0.0000)	0.4017 (0.3930–0.4019)	0.3794 (0.2324–0.5277)	0.3311 (0.1994–0.4529)
			HD95		180.3 (163.7–194.6)	79.16 (4.618–191.4)	– (– –)	14.74 (8.039–25.18)	10.80 (3.860–21.04)	23.49 (15.73–31.70)
		left gluteus maximus	Dice	0.9250 (0.9198–0.9295)	0.7879 (0.7744–0.8016)	0.9609 (0.9585–0.9630)	0.9508 (0.9471–0.9540)	0.9466 (0.9355–0.9490)	0.9371 (0.9325–0.9412)	0.9459 (0.9427–0.9490)
			HD95	52.42 (19.13–94.62)	15.73 (13.63–18.54)	2.321 (2.137–2.524)	3.109 (2.695–3.639)	2.885 (2.551–3.353)	28.83 (16.33–43.83)	3.266 (2.992–3.539)
		right gluteus maximus	Dice	0.9264 (0.9200–0.9324)	0.7849 (0.7730–0.7969)	0.9625 (0.9583–0.9659)	0.9420 (0.9370–0.9471)	0.9555 (0.9533–0.9578)	0.9376 (0.9318–0.9423)	0.9485 (0.9425–0.9533)
			HD95	38.56 (11.48–72.16)	17.48 (14.84–21.63)	2.969 (2.368–3.863)	3.866 (3.265–4.639)	3.120 (2.475–3.998)	5.334 (4.277–6.863)	3.443 (2.771–4.511)
		left gluteus medius	Dice	0.8486 (0.8331–0.8635)	0.7093 (0.6893–0.7272)	0.9603 (0.9584–0.9620)	0.9423 (0.9393–0.9450)	0.9588 (0.9562–0.9734)	0.9059 (0.9029–0.9089)	0.9358 (0.9329–0.9386)
			HD95	63.75 (62.57–64.81)	22.34 (17.00–29.16)	19.92 (2.062–55.53)	2.761 (2.573–2.964)	6.383 (6.137–6.401)	4.688 (4.360–5.045)	3.162 (2.953–3.388)
		right gluteus medius	Dice	0.9104 (0.9068–0.9141)	0.7320 (0.7197–0.7438)	0.9612 (0.9593–0.9631)	0.9379 (0.9351–0.9406)	0.9368 (0.9342–0.9491)	0.9171 (0.9139–0.9206)	0.9345 (0.9303–0.9382)
			HD95	29.17 (23.93–34.43)	14.71 (13.63–16.25)	1.985 (1.869–2.118)	3.244 (3.005–3.515)	3.275 (1.918–5.110)	4.203 (3.943–4.504)	3.197 (2.923–3.515)
		left gluteus minimus	Dice	0.8492 (0.8425–0.8564)	0.4525 (0.3697–0.5208)	0.8883 (0.8786–0.8983)	0.9076 (0.9023–0.9125)	0.8840 (0.8699–0.8871)	0.8862 (0.8771–0.8948)	0.8969 (0.8905–0.9031)
			HD95	27.89 (5.032–61.98)	50.43 (28.62–75.72)	455.1 (399.0–496.5)	2.900 (2.587–3.284)	4.596 (3.597–6.038)	3.707 (3.355–4.079)	2.777 (2.581–2.975)
		right gluteus minimus	Dice	0.8741 (0.8671–0.8807)	0.4653 (0.3990–0.5292)	0.8674 (0.8573–0.8787)	0.9070 (0.8997–0.9141)	0.8920 (0.8786–0.9092)	0.8850 (0.8765–0.8930)	0.8626 (0.8516–0.8731)
			HD95	5.227 (4.863–5.620)	44.08 (24.80–68.70)	482.4 (439.9–516.5)	2.466 (2.285–2.645)	4.538 (3.697–5.966)	3.244 (2.914–3.667)	4.257 (3.828–4.724)
		left iliopsoas	Dice	0.9377 (0.9340–0.9411)	0.5615 (0.5022–0.6153)	0.9429 (0.9383–0.9471)	0.9575 (0.9538–0.9607)	0.9404 (0.9368–0.9452)	0.9434 (0.9393–0.9471)	0.9319 (0.9282–0.9356)
			HD95	3.045 (2.478–4.041)	31.13 (22.53–40.47)	2.107 (1.977–2.267)	1.792 (1.559–2.140)	4.784 (4.415–7.069)	4.925 (2.706–8.040)	9.488 (5.217–16.23)
		right iliopsoas	Dice	0.9400 (0.9362–0.9437)	0.5845 (0.5356–0.6259)	0.9518 (0.9489–0.9546)	0.9599 (0.9571–0.9625)	0.9393 (0.9377–0.9401)	0.9395 (0.9354–0.9433)	0.9065 (0.9003–0.9124)
			HD95	2.362 (2.227–2.490)	24.42 (17.04–33.16)	1.790 (1.745–1.831)	1.440 (1.309–1.579)	2.165 (1.401–2.407)	9.407 (3.621–18.70)	35.96 (31.24–40.83)
		left autochthon	Dice	0.9462 (0.9416–0.9501)	0.7395 (0.7269–0.7505)	0.9627 (0.9603–0.9649)	0.9506 (0.9464–0.9544)	0.9620 (0.9573–0.9683)	0.9162 (0.9107–0.9212)	0.9249 (0.9190–0.9301)
			HD95	2.527 (2.229–2.947)	14.38 (11.99–18.19)	1.737 (1.630–1.849)	2.013 (1.755–2.380)	2.891 (2.638–3.574)	22.35 (15.80–29.65)	4.286 (3.172–5.876)
		right autochthon	Dice	0.9399 (0.9356–0.9434)	0.7396 (0.7244–0.7529)	0.9613 (0.9587–0.9636)	0.9453 (0.9405–0.9497)	0.9585 (0.9420–0.9596)	0.9118 (0.9049–0.9170)	0.9266 (0.9213–0.9309)
			HD95	2.554 (2.358–2.817)	15.77 (12.52–20.05)	1.677 (1.579–1.784)	2.220 (1.958–2.541)	4.044 (2.946–5.117)	9.458 (4.806–15.76)	3.207 (2.936–3.571)
		skin	Dice	Not supported	0.0006846 (0.0004049–0.001021)	0.7606 (0.7547–0.7663)	0.7798 (0.7721–0.7865)	0.7761 (0.7620–0.7883)	0.7347 (0.7286–0.7399)	0.7466 (0.7411–0.7516)
			HD95		115.9 (106.4–128.3)	1.336 (1.211–1.478)	21.35 (13.20–30.45)	2.245 (1.912–3.152)	3.549 (3.092–4.016)	2.566 (2.264–2.850)
		cervical vertebrae 1 (c1)	Dice	0.8652 (0.8582–0.8711)	0.1337 (0.1009–0.1672)	0.8663 (0.8552–0.8754)	0.9089 (0.9027–0.9146)	0.8360 (0.8262–0.8381)	0.8881 (0.8796–0.8959)	0.9033 (0.8968–0.9088)
			HD95	55.47 (24.99–93.20)	39.34 (34.85–43.70)	42.37 (11.94–82.21)	1.997 (1.762–2.193)	1.974 (1.704–2.682)	2.466 (2.194–2.760)	1.947 (1.716–2.168)
		cervical vertebrae 2 (c2)	Dice	0.8886 (0.8814–0.8950)	0.09402 (0.05907–0.1298)	0.6908 (0.6411–0.7415)	0.9217 (0.9170–0.9263)	0.8593 (0.8378–0.8639)	0.9057 (0.8969–0.9124)	0.9203 (0.9153–0.9251)
			HD95	10.44 (5.533–16.24)	45.17 (41.15–49.21)	354.5 (309.0–395.3)	1.863 (1.610–2.144)	2.482 (2.225–2.846)	2.180 (1.952–2.411)	1.741 (1.509–1.954)
		cervical vertebrae 3 (c3)	Dice	0.8555 (0.8437–0.8667)	0.03676 (0.01260–0.06253)	0.3834 (0.3495–0.4221)	0.9014 (0.8935–0.9089)	0.8853 (0.8699–0.8951)	0.8936 (0.8822–0.9035)	0.9052 (0.8993–0.9102)
			HD95	8.191 (4.535–12.42)	56.13 (51.42–61.33)	408.3 (399.3–416.2)	2.122 (1.841–2.421)	2.825 (2.054–3.317)	2.199 (1.973–2.429)	1.757 (1.543–1.964)
		cervical vertebrae 4 (c4)	Dice	0.8670 (0.8575–0.8757)	0.03874 (0.01351–0.06714)	0.2733 (0.2628–0.2857)	0.9068 (0.8978–0.9146)	0.9009 (0.8711–0.9301)	0.6374 (0.6239–0.6518)	0.8865 (0.8750–0.8978)
			HD95	41.55 (16.23–76.42)	63.03 (55.77–70.54)	424.1 (414.5–432.7)	1.751 (1.541–1.989)	3.487 (3.479–5.099)	119.2 (70.16–175.0)	2.296 (2.026–2.575)
		cervical vertebrae 5 (c5)	Dice	0.8166 (0.7853–0.8404)	0.000009553 (0.0000–0.00002866)	0.3229 (0.3043–0.3452)	0.8904 (0.8787–0.8998)	0.7931 (0.7865–0.8011)	0.7119 (0.6687–0.7530)	0.8590 (0.8457–0.8714)
			HD95	26.35 (18.56–35.17)	128.2 (121.5–134.9)	437.8 (428.7–445.7)	2.017 (1.817–2.223)	3.457 (2.050–5.522)	4.911 (4.282–5.621)	3.161 (2.769–3.581)
		cervical vertebrae 6 (c6)	Dice	0.8553 (0.8437–0.8658)	0.0000 (0.0000–0.0000)	0.8489 (0.8259–0.8667)	0.8944 (0.8830–0.9036)	0.8704 (0.8570–0.8958)	0.8681 (0.8582–0.8774)	0.8788 (0.8705–0.8872)
			HD95	21.56 (2.727–46.19)	160.6 (152.7–169.1)	58.84 (14.32–107.1)	2.034 (1.813–2.301)	3.319 (2.285–3.629)	2.556 (2.359–2.787)	2.285 (2.068–2.512)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	DAP-Atlas (AutoPETIII)	cervical vertebrae 7 (c7)	Dice	0.8850 (0.8728–0.8946)	0.0000 (0.0000–0.0000)	0.6335 (0.5790–0.6878)	0.9074 (0.8985–0.9160)	0.9063 (0.8871–0.9205)	0.8681 (0.8566–0.8789)	0.8875 (0.8792–0.8952)
			HD95	2.509 (2.295–2.816)	173.9 (164.8–183.4)	397.9 (353.3–436.3)	1.871 (1.647–2.100)	3.805 (2.607–4.658)	4.378 (3.598–5.280)	2.540 (2.418–2.667)
		thoracic vertebrae 1 (t1)	Dice	0.8697 (0.8471–0.8867)	0.04665 (0.02127–0.07478)	0.8186 (0.7643–0.8576)	0.9201 (0.9087–0.9301)	0.9180 (0.9045–0.9223)	0.9076 (0.8936–0.9195)	0.9096 (0.9026–0.9165)
			HD95	32.50 (24.64–39.14)	46.20 (41.81–50.65)	111.4 (52.15–180.7)	1.922 (1.560–2.375)	2.937 (1.751–3.865)	2.336 (1.849–2.981)	2.263 (1.934–2.721)
		thoracic vertebrae 2 (t2)	Dice	0.8815 (0.8438–0.9054)	0.1498 (0.1061–0.2014)	0.8767 (0.8560–0.8925)	0.9108 (0.8853–0.9279)	0.8865 (0.8661–0.8962)	0.8261 (0.7971–0.8504)	0.9135 (0.9019–0.9234)
			HD95	36.55 (4.912–72.80)	33.72 (29.27–38.18)	2.870 (2.526–3.331)	3.464 (2.185–5.117)	3.609 (2.909–4.689)	5.902 (4.428–7.690)	2.252 (1.600–3.183)
		thoracic vertebrae 3 (t3)	Dice	0.8630 (0.8031–0.9019)	0.2691 (0.2137–0.3261)	0.8712 (0.8444–0.8912)	0.8887 (0.8461–0.9220)	0.8723 (0.8524–0.8765)	0.8593 (0.8389–0.8777)	0.9043 (0.8818–0.9199)
			HD95	3.716 (2.388–5.554)	24.15 (20.58–28.46)	3.214 (2.467–4.246)	3.885 (2.489–5.784)	3.495 (2.876–5.008)	8.934 (5.834–13.76)	2.556 (1.734–3.748)
		thoracic vertebrae 4 (t4)	Dice	0.8282 (0.7574–0.8892)	0.2990 (0.2578–0.3399)	0.8445 (0.8110–0.8719)	0.8711 (0.8129–0.9137)	0.8383 (0.8183–0.8484)	0.6945 (0.6582–0.7287)	0.8896 (0.8621–0.9107)
			HD95	4.875 (3.039–7.248)	22.35 (20.21–24.49)	71.35 (18.82–136.4)	3.742 (2.196–5.729)	4.342 (3.084–7.294)	85.24 (70.53–98.28)	2.702 (1.905–3.778)
		thoracic vertebrae 5 (t5)	Dice	0.8483 (0.8056–0.8858)	0.3722 (0.3318–0.4075)	0.8729 (0.8561–0.8877)	0.8806 (0.8353–0.9140)	0.8797 (0.8727–0.8956)	0.5884 (0.5166–0.6625)	0.8967 (0.8814–0.9103)
			HD95	5.424 (3.849–7.185)	20.76 (18.59–23.27)	3.337 (2.706–4.252)	3.930 (2.611–5.468)	3.557 (1.653–4.966)	10.68 (8.760–12.75)	3.020 (2.225–3.944)
		thoracic vertebrae 6 (t6)	Dice	0.8619 (0.8218–0.8941)	0.4088 (0.3583–0.4548)	0.8698 (0.8562–0.8818)	0.8860 (0.8479–0.9154)	0.8656 (0.8624–0.8824)	0.8245 (0.7874–0.8564)	0.8851 (0.8678–0.9001)
			HD95	4.294 (3.120–5.873)	22.86 (16.81–32.56)	2.925 (2.652–3.215)	3.775 (2.638–5.161)	5.833 (5.390–7.790)	5.538 (4.277–6.869)	3.511 (2.723–4.377)
		thoracic vertebrae 7 (t7)	Dice	0.8672 (0.8285–0.8972)	0.2734 (0.2048–0.3403)	0.8790 (0.8646–0.8917)	0.8951 (0.8560–0.9242)	0.8653 (0.8617–0.8718)	0.8986 (0.8818–0.9125)	0.9004 (0.8816–0.9146)
			HD95	4.591 (3.186–6.308)	43.84 (32.07–58.44)	3.198 (2.663–3.871)	3.704 (2.619–4.901)	4.950 (3.863–5.624)	3.632 (2.696–4.826)	3.338 (2.439–4.458)
		thoracic vertebrae 8 (t8)	Dice	0.8723 (0.8526–0.8910)	0.2173 (0.1549–0.2820)	0.8763 (0.8493–0.8953)	0.9106 (0.8826–0.9332)	0.8846 (0.8506–0.9006)	0.8770 (0.8532–0.8967)	0.9005 (0.8840–0.9143)
			HD95	25.00 (19.53–30.23)	63.35 (46.77–81.52)	2.911 (2.533–3.417)	3.142 (2.204–4.367)	4.126 (3.255–6.915)	4.716 (3.737–5.830)	3.723 (2.742–4.859)
		thoracic vertebrae 9 (t9)	Dice	0.9013 (0.8854–0.9144)	0.3543 (0.2906–0.4167)	0.8779 (0.8463–0.9005)	0.9187 (0.8913–0.9396)	0.9042 (0.8845–0.9374)	0.9050 (0.8864–0.9195)	0.9147 (0.8993–0.9271)
			HD95	52.22 (23.84–85.27)	28.54 (22.08–36.96)	3.529 (2.626–4.682)	2.805 (2.015–3.754)	3.816 (3.062–4.779)	3.615 (2.798–4.661)	2.749 (2.173–3.475)
		thoracic vertebrae 10 (t10)	Dice	0.9029 (0.8829–0.9210)	0.2824 (0.2272–0.3424)	0.8786 (0.8472–0.9001)	0.9211 (0.8952–0.9402)	0.8920 (0.8890–0.9238)	0.9058 (0.8916–0.9188)	0.9177 (0.9078–0.9273)
			HD95	4.485 (2.869–6.493)	31.01 (25.11–38.28)	3.460 (2.583–4.614)	3.215 (2.197–4.486)	4.769 (3.730–6.316)	4.726 (3.619–5.963)	3.015 (2.333–3.755)
		thoracic vertebrae 11 (t11)	Dice	0.8898 (0.8243–0.9339)	0.2793 (0.2252–0.3450)	0.8774 (0.8379–0.9091)	0.9202 (0.8799–0.9473)	0.8932 (0.8771–0.9089)	0.9232 (0.8944–0.9404)	0.9273 (0.9125–0.9394)
			HD95	8.889 (2.415–19.73)	37.60 (30.27–45.60)	4.209 (2.464–6.993)	3.864 (1.942–6.546)	4.005 (2.536–7.482)	3.276 (1.866–5.597)	2.549 (1.798–3.459)
		thoracic vertebrae 12 (t12)	Dice	0.8943 (0.8417–0.9304)	0.02524 (0.003933–0.05160)	0.8199 (0.7724–0.8504)	0.9251 (0.8859–0.9511)	0.9249 (0.9100–0.9262)	0.9093 (0.8853–0.9284)	0.9315 (0.9151–0.9439)
			HD95	11.14 (6.518–15.85)	78.20 (66.52–90.16)	358.5 (350.2–366.3)	2.972 (1.611–4.916)	4.693 (4.353–8.157)	4.560 (3.385–5.985)	2.822 (1.720–4.311)
		lumbar vertebrae 1 (l1)	Dice	0.9096 (0.8711–0.9357)	0.2764 (0.2254–0.3282)	0.8643 (0.8414–0.8841)	0.9377 (0.9101–0.9575)	0.9061 (0.8791–0.9172)	0.9220 (0.9085–0.9336)	0.9364 (0.9225–0.9467)
			HD95	4.364 (2.086–7.420)	36.08 (31.66–41.02)	279.7 (225.2–326.7)	3.727 (1.710–6.899)	4.285 (3.695–6.641)	4.597 (2.795–7.668)	3.210 (1.609–5.340)
		lumbar vertebrae 2 (l2)	Dice	0.9119 (0.8814–0.9325)	0.4744 (0.4355–0.5082)	0.8980 (0.8834–0.9108)	0.9436 (0.9261–0.9570)	0.9262 (0.9059–0.9592)	0.8923 (0.8659–0.9107)	0.9329 (0.9218–0.9422)
			HD95	4.261 (2.290–6.578)	20.42 (18.47–22.58)	62.43 (22.21–113.7)	2.826 (1.542–4.758)	2.961 (1.873–4.046)	5.972 (4.503–7.804)	3.944 (2.264–6.202)
		lumbar vertebrae 3 (l3)	Dice	0.9096 (0.8826–0.9307)	0.2596 (0.2123–0.3080)	0.9056 (0.8923–0.9161)	0.9377 (0.9163–0.9543)	0.9356 (0.9278–0.9451)	0.9271 (0.9167–0.9362)	0.9314 (0.9201–0.9409)
			HD95	4.911 (0.4736–10.32)	34.62 (30.98–37.97)	3.170 (2.417–4.462)	3.520 (1.935–5.893)	4.244 (3.220–4.391)	3.442 (2.431–4.941)	3.192 (1.913–5.088)
		lumbar vertebrae 4 (l4)	Dice	0.7764 (0.7577–0.7937)	0.1351 (0.09130–0.1807)	0.8552 (0.8441–0.8658)	0.9343 (0.9148–0.9515)	0.9236 (0.9195–0.9324)	0.8996 (0.8842–0.9137)	0.9328 (0.9228–0.9422)
			HD95	48.94 (47.00–50.61)	41.84 (38.45–45.26)	380.3 (340.3–414.3)	3.815 (2.051–6.043)	4.141 (3.206–7.114)	29.72 (8.124–59.82)	2.585 (2.044–3.405)
		lumbar vertebrae 5 (l5)	Dice	0.8936 (0.8458–0.9252)	0.01211 (0.003800–0.02174)	0.8119 (0.7890–0.8299)	0.9066 (0.8702–0.9386)	0.8761 (0.8527–0.8906)	0.9097 (0.8811–0.9309)	0.9152 (0.8900–0.9344)
			HD95	4.553 (2.660–7.054)	61.21 (57.77–64.84)	410.6 (381.5–429.6)	5.295 (2.972–8.298)	4.991 (2.918–5.515)	3.901 (2.546–5.621)	2.861 (2.117–3.973)
		left rib 1	Dice	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.7729 (0.6902–0.8384)	0.0000 (0.0000–0.0000)	0.7162 (0.7055–0.7314)	0.6772 (0.5914–0.7467)	0.8080 (0.7169–0.8774)
			HD95	293.8 (281.6–308.4)	299.4 (291.1–308.2)	39.71 (15.24–74.58)	294.2 (286.5–303.1)	16.31 (7.073–26.27)	15.64 (6.509–27.33)	17.54 (8.816–27.97)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	DAP-Atlas (AutoPETIII)	right rib 1	Dice	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.7755 (0.6977–0.8295)	0.0000 (0.0000–0.0000)	0.7675 (0.7536–0.7806)	0.8306 (0.7547–0.8872)	0.8588 (0.7941–0.9031)
			HD95	287.4 (275.8–300.3)	296.8 (290.1–303.5)	50.91 (22.26–85.79)	293.0 (285.8–300.4)	9.462 (2.332–17.00)	10.10 (1.929–21.87)	7.922 (1.645–18.58)
		left rib 2	Dice	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.7736 (0.6890–0.8472)	0.0000 (0.0000–0.0000)	0.7522 (0.7478–0.7524)	0.6456 (0.5473–0.7278)	0.8224 (0.7352–0.8967)
			HD95	265.4 (253.2–278.8)	265.5 (259.9–271.3)	14.36 (4.206–29.27)	– (– –)	8.816 (3.597–15.62)	21.70 (13.04–31.70)	6.982 (1.399–13.93)
		right rib 2	Dice	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.7712 (0.6881–0.8461)	0.0000 (0.0000–0.0000)	0.7581 (0.7574–0.7635)	0.7247 (0.6201–0.8117)	0.8306 (0.7435–0.9051)
			HD95	246.8 (236.2–256.8)	268.4 (261.4–275.9)	15.04 (4.525–29.01)	– (– –)	15.07 (4.737–26.54)	18.03 (7.981–31.17)	11.27 (2.336–23.72)
		left rib 3	Dice	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.8032 (0.7306–0.8623)	0.0000 (0.0000–0.0000)	0.7801 (0.7613–0.7812)	0.7716 (0.6998–0.8332)	0.8746 (0.8037–0.9214)
			HD95	227.5 (218.1–237.4)	216.6 (208.5–225.6)	11.97 (3.101–28.67)	298.9 (298.9–298.9)	13.60 (12.53–14.31)	34.20 (24.59–47.82)	8.379 (0.9367–21.99)
		right rib 3	Dice	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.8061 (0.7329–0.8642)	0.0000 (0.0000–0.0000)	0.7979 (0.7927–0.8071)	0.7313 (0.6718–0.7791)	0.8875 (0.8325–0.9185)
			HD95	213.5 (208.5–219.0)	220.2 (215.3–225.3)	6.308 (2.856–11.17)	279.8 (254.0–307.7)	6.677 (4.253–12.01)	28.36 (24.42–32.26)	3.284 (1.104–7.394)
		left rib 4	Dice	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.7814 (0.6971–0.8553)	0.0000 (0.0000–0.0000)	0.7715 (0.7602–0.7726)	0.6545 (0.5981–0.6924)	0.8582 (0.7786–0.9194)
			HD95	159.7 (154.1–165.0)	164.3 (158.9–169.7)	7.470 (4.416–11.11)	189.9 (179.8–200.3)	7.044 (3.020–11.94)	178.8 (167.9–189.1)	4.229 (1.593–7.544)
		right rib 4	Dice	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.8036 (0.7327–0.8625)	0.0000 (0.0000–0.0000)	0.7809 (0.7769–0.7813)	0.7743 (0.7175–0.8135)	0.9019 (0.8464–0.9311)
			HD95	160.5 (156.3–165.0)	170.1 (164.5–176.2)	5.985 (2.412–10.65)	211.3 (198.0–224.5)	8.714 (3.735–13.39)	36.68 (30.74–43.92)	3.432 (0.8995–7.819)
		left rib 5	Dice	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.7616 (0.6606–0.8390)	0.0000 (0.0000–0.0000)	0.7439 (0.7365–0.7447)	0.8175 (0.7319–0.8862)	0.8469 (0.7685–0.9110)
			HD95	108.5 (101.8–117.4)	115.7 (107.7–126.2)	16.57 (5.943–33.02)	166.8 (150.5–184.5)	20.16 (12.66–29.90)	16.99 (6.043–32.45)	13.15 (2.246–28.49)
		right rib 5	Dice	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.7945 (0.7215–0.8514)	0.0000 (0.0000–0.0000)	0.7794 (0.7718–0.7855)	0.8766 (0.8192–0.9099)	0.8869 (0.8311–0.9185)
			HD95	103.7 (100.1–107.2)	116.9 (112.0–121.6)	5.993 (2.725–10.70)	158.7 (145.3–173.5)	9.059 (6.756–13.63)	5.986 (2.038–10.31)	4.832 (0.9457–9.613)
		left rib 6	Dice	0.03498 (0.006075–0.07448)	0.01384 (0.007653–0.02123)	0.8008 (0.7258–0.8626)	0.8408 (0.7673–0.9056)	0.8353 (0.8168–0.8561)	0.6925 (0.6299–0.7575)	0.8828 (0.8360–0.9195)
			HD95	38.75 (35.25–42.84)	61.33 (56.42–66.46)	7.162 (3.859–11.49)	9.775 (5.861–13.59)	7.435 (3.558–10.66)	20.93 (16.82–24.54)	5.378 (1.913–9.863)
		right rib 6	Dice	0.04803 (0.01547–0.09125)	0.008577 (0.002090–0.01766)	0.7928 (0.7256–0.8428)	0.8795 (0.8132–0.9249)	0.8528 (0.8449–0.8635)	0.7825 (0.7190–0.8305)	0.8913 (0.8332–0.9247)
			HD95	48.16 (35.06–69.09)	94.67 (77.44–113.3)	8.075 (3.956–13.77)	10.31 (5.466–15.71)	7.632 (3.311–11.70)	23.06 (19.16–26.80)	5.215 (0.9992–11.09)
		left rib 7	Dice	0.05600 (0.006283–0.1203)	0.1169 (0.09759–0.1367)	0.7754 (0.6882–0.8498)	0.8414 (0.7519–0.9124)	0.8179 (0.7930–0.8311)	0.6368 (0.5791–0.6923)	0.8661 (0.8345–0.8923)
			HD95	36.14 (33.79–38.77)	40.27 (36.22–44.47)	7.266 (3.834–11.57)	9.182 (5.219–12.99)	8.917 (7.668–10.43)	24.14 (22.24–25.92)	7.022 (3.074–12.17)
		right rib 7	Dice	0.01763 (0.001467–0.04477)	0.1298 (0.1072–0.1524)	0.7912 (0.7118–0.8455)	0.8693 (0.7863–0.9288)	0.7878 (0.7589–0.8255)	0.6777 (0.6125–0.7335)	0.8814 (0.8253–0.9138)
			HD95	38.03 (35.63–40.97)	40.43 (33.90–47.43)	6.101 (3.036–11.17)	10.83 (6.243–16.10)	10.10 (7.682–12.52)	24.89 (22.14–28.34)	5.009 (1.683–10.74)
		left rib 8	Dice	0.0000 (0.0000–0.0000)	0.001722 (0.0000–0.005053)	0.7498 (0.6549–0.8283)	0.0000 (0.0000–0.0000)	0.7123 (0.6958–0.7165)	0.3965 (0.3275–0.4670)	0.8754 (0.8457–0.8988)
			HD95	104.7 (99.97–109.3)	81.06 (76.53–85.52)	9.550 (4.742–15.71)	173.1 (139.0–205.9)	10.43 (6.927–14.70)	21.92 (19.80–23.93)	5.318 (2.148–9.859)
		right rib 8	Dice	0.0004303 (0.0000–0.001291)	0.006221 (0.00001330–0.01751)	0.7834 (0.7100–0.8365)	0.0000 (0.0000–0.0000)	0.7785 (0.7732–0.7816)	0.3387 (0.2788–0.4004)	0.8886 (0.8334–0.9202)
			HD95	127.6 (106.8–152.4)	76.60 (71.65–81.08)	5.766 (2.967–9.718)	176.9 (142.0–212.1)	6.514 (3.465–12.39)	24.76 (22.78–27.23)	3.792 (1.543–7.700)
		left rib 9	Dice	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.7535 (0.6645–0.8213)	0.8421 (0.7421–0.9207)	0.8154 (0.8015–0.8206)	0.5507 (0.4867–0.5983)	0.8766 (0.8408–0.9049)
			HD95	164.6 (158.9–171.2)	135.8 (129.8–141.7)	11.28 (4.818–21.08)	5.300 (2.402–8.808)	8.459 (4.534–13.17)	30.78 (24.92–39.37)	10.94 (4.196–20.82)
		right rib 9	Dice	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.7678 (0.6983–0.8197)	0.8798 (0.7975–0.9350)	0.7603 (0.7482–0.7684)	0.5612 (0.5077–0.6102)	0.8923 (0.8374–0.9227)
			HD95	186.8 (168.8–207.5)	132.5 (125.1–139.2)	5.421 (2.584–9.582)	4.869 (1.460–9.581)	6.220 (4.330–12.33)	24.74 (23.04–26.78)	3.293 (1.006–7.497)
		left rib 10	Dice	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.7451 (0.6666–0.8074)	0.8350 (0.7417–0.9106)	0.7343 (0.7195–0.7368)	0.6687 (0.6071–0.7237)	0.8501 (0.8024–0.8920)
			HD95	222.7 (216.8–229.3)	189.3 (183.0–195.5)	18.42 (4.726–37.01)	6.436 (2.736–10.58)	7.472 (2.577–11.44)	29.29 (23.04–39.76)	8.170 (4.556–12.18)
		right rib 10	Dice	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.7573 (0.7008–0.7970)	0.8833 (0.8136–0.9294)	0.8469 (0.8435–0.8587)	0.8015 (0.7522–0.8394)	0.8829 (0.8486–0.9075)
			HD95	36.42 (31.15–41.83)	186.2 (178.2–193.0)	5.291 (2.500–8.643)	9.410 (1.007–24.49)	6.317 (5.008–11.28)	16.01 (12.74–19.50)	4.370 (1.947–7.824)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	DAP-Atlas (AutoPETIII)	left rib 11	Dice	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.6567 (0.5751–0.7249)	0.8215 (0.7182–0.9033)	0.8162 (0.8081–0.8215)	0.7618 (0.6705–0.8301)	0.8394 (0.7657–0.8949)
			HD95	275.8 (270.0–281.6)	236.5 (230.5–242.8)	158.7 (113.9–200.2)	5.350 (1.948–9.408)	5.920 (2.209–10.12)	33.87 (11.98–60.50)	7.218 (3.470–11.64)
		right rib 11	Dice	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.6795 (0.5988–0.7405)	0.8431 (0.7598–0.9228)	0.6684 (0.6572–0.6761)	0.8112 (0.7345–0.8761)	0.8699 (0.8228–0.9066)
			HD95	269.5 (264.8–274.0)	231.7 (225.5–238.5)	133.5 (91.90–174.9)	3.825 (1.289–6.959)	5.081 (2.618–8.333)	8.034 (3.905–12.68)	4.726 (2.096–8.500)
		left rib 12	Dice	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.6567 (0.5732–0.7254)	0.8071 (0.7029–0.8883)	0.8000 (0.7981–0.8044)	0.7527 (0.6694–0.8239)	0.8147 (0.7255–0.8889)
			HD95	293.0 (285.1–299.5)	266.5 (258.8–276.1)	160.2 (115.6–204.7)	10.59 (2.381–20.61)	14.22 (7.281–21.41)	187.5 (148.8–224.2)	10.69 (2.719–20.55)
		right rib 12	Dice	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.6617 (0.5933–0.7222)	0.8167 (0.7267–0.8955)	0.7891 (0.7802–0.7979)	0.7979 (0.7378–0.8493)	0.8572 (0.7990–0.8979)
			HD95	290.5 (284.0–297.7)	260.1 (251.3–269.5)	158.6 (115.6–200.6)	9.132 (3.362–15.68)	11.69 (4.299–16.96)	33.50 (16.34–56.27)	7.081 (2.707–12.59)
		rib cartilage	Dice	0.7651 (0.7524–0.7776)	0.004292 (0.001973–0.006857)	0.8707 (0.8614–0.8791)	0.8665 (0.8527–0.8779)	0.8487 (0.8341–0.8704)	0.8162 (0.8026–0.8276)	0.8314 (0.8172–0.8438)
			HD95	39.58 (26.14–54.30)	262.7 (239.3–291.8)	4.480 (2.825–6.827)	51.17 (29.20–77.69)	5.076 (3.219–7.223)	60.84 (38.93–89.34)	58.09 (36.45–86.83)
		sternum	Dice	0.6086 (0.5766–0.6299)	0.0000 (0.0000–0.0000)	0.8617 (0.8035–0.8962)	0.6398 (0.5998–0.6640)	0.8090 (0.7799–0.8267)	0.8767 (0.8142–0.9136)	0.8783 (0.8135–0.9151)
			HD95	83.57 (59.88–111.4)	209.1 (188.2–233.9)	38.07 (13.26–68.86)	87.02 (60.05–117.0)	42.71 (22.34–67.87)	50.43 (20.30–89.72)	49.92 (19.79–89.78)
		left clavicle	Dice	0.9119 (0.9015–0.9208)	0.1040 (0.07896–0.1290)	0.9000 (0.8934–0.9067)	0.9497 (0.9464–0.9525)	0.9430 (0.9346–0.9472)	0.9515 (0.9440–0.9577)	0.9505 (0.9456–0.9545)
			HD95	29.38 (4.073–62.78)	66.41 (52.36–80.13)	2.408 (2.338–2.468)	0.9654 (0.8845–1.077)	2.854 (1.913–3.932)	11.02 (1.064–28.48)	1.012 (0.9015–1.151)
		right clavicle	Dice	0.9150 (0.9101–0.9192)	0.1167 (0.08780–0.1434)	0.8997 (0.8924–0.9067)	0.9508 (0.9484–0.9531)	0.9361 (0.9337–0.9435)	0.9536 (0.9506–0.9570)	0.9508 (0.9475–0.9539)
			HD95	43.17 (4.440–97.51)	74.00 (61.34–86.97)	48.91 (2.467–109.7)	0.9665 (0.8961–1.045)	4.045 (3.026–4.748)	2.528 (1.339–4.255)	1.101 (0.9699–1.238)
		left scapula	Dice	0.8681 (0.8550–0.8797)	0.4403 (0.4199–0.4584)	0.9127 (0.9080–0.9164)	0.9290 (0.9263–0.9311)	0.9020 (0.8858–0.9066)	0.9299 (0.9221–0.9355)	0.9362 (0.9319–0.9393)
			HD95	11.21 (6.958–15.63)	25.78 (19.56–35.64)	1.859 (1.790–1.937)	0.9944 (0.9189–1.100)	2.518 (2.008–4.047)	1.858 (0.9838–3.114)	0.9449 (0.8768–1.035)
		right scapula	Dice	0.8925 (0.8821–0.9010)	0.4064 (0.3834–0.4258)	0.9126 (0.9049–0.9181)	0.9278 (0.9218–0.9322)	0.9032 (0.8891–0.9069)	0.9212 (0.9154–0.9268)	0.9370 (0.9330–0.9406)
			HD95	11.68 (8.108–15.05)	36.79 (25.11–51.11)	2.218 (1.823–2.919)	1.718 (0.9676–3.054)	1.870 (1.610–3.240)	7.234 (2.445–14.64)	1.188 (0.9483–1.529)
		left humerus	Dice	0.4440 (0.4258–0.4631)	0.3937 (0.3594–0.4244)	0.9260 (0.9118–0.9373)	0.9145 (0.8986–0.9284)	0.9212 (0.9131–0.9284)	0.9318 (0.9190–0.9424)	0.9473 (0.9383–0.9556)
			HD95	71.47 (69.86–72.97)	38.73 (28.62–51.20)	15.29 (2.654–36.41)	27.70 (10.50–48.53)	15.96 (5.705–29.08)	20.51 (8.737–34.91)	5.814 (2.690–9.590)
		right humerus	Dice	0.6257 (0.6024–0.6481)	0.3731 (0.3273–0.4182)	0.9296 (0.9182–0.9383)	0.9224 (0.9123–0.9324)	0.9190 (0.8905–0.9285)	0.9370 (0.9267–0.9458)	0.9385 (0.9282–0.9470)
			HD95	66.45 (64.31–68.27)	47.88 (29.26–79.01)	19.24 (4.726–39.86)	19.30 (7.555–32.98)	12.86 (6.103–23.54)	26.81 (10.41–47.14)	11.37 (5.179–19.08)
		skull	Dice	0.7588 (0.7518–0.7668)	0.1191 (0.09238–0.1434)	0.1822 (0.1355–0.2501)	0.8450 (0.8393–0.8509)	0.8154 (0.8124–0.8346)	0.8704 (0.8648–0.8764)	0.8768 (0.8711–0.8836)
			HD95	34.42 (18.34–51.90)	142.5 (93.98–200.6)	736.1 (666.7–786.1)	5.132 (3.126–8.688)	6.176 (1.867–9.256)	10.92 (4.050–19.63)	60.67 (7.712–125.7)
		left hip	Dice	0.9385 (0.9329–0.9431)	0.5903 (0.5805–0.5999)	0.8820 (0.8732–0.8903)	0.9659 (0.9621–0.9687)	0.9583 (0.9542–0.9596)	0.9551 (0.9494–0.9588)	0.9587 (0.9552–0.9618)
			HD95	31.81 (25.12–38.95)	21.90 (19.60–24.73)	401.8 (392.7–410.0)	1.363 (1.023–1.757)	3.547 (3.133–4.707)	15.98 (7.405–27.34)	1.692 (1.346–2.145)
		right hip	Dice	0.9062 (0.8974–0.9133)	0.5697 (0.5329–0.5957)	0.9242 (0.9202–0.9281)	0.9658 (0.9613–0.9692)	0.9632 (0.9537–0.9749)	0.9576 (0.9520–0.9614)	0.9572 (0.9540–0.9601)
			HD95	45.00 (42.48–47.58)	47.64 (23.40–77.97)	363.9 (318.1–397.9)	1.689 (1.009–2.737)	2.386 (2.113–3.142)	1.806 (1.277–2.468)	1.691 (1.262–2.266)
		sacrum	Dice	0.7911 (0.7831–0.7992)	0.4763 (0.4492–0.4989)	0.8692 (0.8661–0.8724)	0.9402 (0.9329–0.9474)	0.8754 (0.8640–0.8792)	0.9147 (0.9068–0.9222)	0.9284 (0.9212–0.9348)
			HD95	41.46 (38.39–43.39)	21.23 (16.68–29.87)	418.8 (409.4–427.1)	1.956 (1.592–2.427)	3.733 (2.605–4.513)	3.274 (2.856–3.697)	2.674 (2.267–3.093)
		left femur	Dice	0.8730 (0.8624–0.8835)	0.6658 (0.6516–0.6800)	0.8119 (0.7978–0.8255)	0.9629 (0.9595–0.9659)	0.9283 (0.9240–0.9383)	0.9717 (0.9680–0.9743)	0.9746 (0.9719–0.9766)
			HD95	55.97 (52.36–59.55)	17.25 (15.93–18.55)	606.3 (589.0–622.5)	1.579 (1.266–1.999)	2.888 (2.195–3.226)	1.496 (1.214–1.858)	1.109 (0.9289–1.319)
		right femur	Dice	0.8147 (0.8017–0.8272)	0.6831 (0.6730–0.6930)	0.7965 (0.7796–0.8102)	0.9618 (0.9595–0.9638)	0.9349 (0.9244–0.9530)	0.9718 (0.9655–0.9757)	0.9742 (0.9714–0.9766)
			HD95	66.08 (63.69–68.35)	19.64 (14.50–29.31)	622.9 (602.5–647.4)	1.554 (1.381–1.754)	2.752 (2.642–3.579)	1.695 (1.247–2.435)	1.171 (0.9767–1.461)
		heart	Dice	0.04151 (0.03750–0.04599)	0.05205 (0.04385–0.06060)	0.8007 (0.7689–0.8181)	0.6634 (0.6348–0.6894)	0.7794 (0.7784–0.7971)	0.7407 (0.7326–0.7487)	0.7233 (0.7133–0.7319)
			HD95	234.6 (180.9–284.0)	58.91 (55.79–62.27)	2.908 (2.543–3.604)	8.978 (7.583–10.71)	5.532 (4.187–5.804)	8.180 (7.393–9.052)	6.264 (5.707–6.988)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	DAP-Atlas (AutoPETIII)	left heart atrium	Dice	Not supported	0.4086 (0.3667–0.4459)	0.9000 (0.8874–0.9120)	0.9572 (0.9541–0.9602)	0.9459 (0.9378–0.9659)	0.9143 (0.9088–0.9196)	0.9460 (0.9420–0.9498)
			HD95		32.78 (28.32–38.37)	309.3 (252.8–360.9)	1.867 (1.702–2.034)	3.953 (3.455–6.030)	3.973 (3.676–4.290)	8.327 (3.777–13.28)
		heart tissue	Dice	Not supported	0.05063 (0.03829–0.06230)	0.7843 (0.7603–0.8061)	0.7367 (0.7039–0.7634)	0.7710 (0.7583–0.7923)	0.7164 (0.6850–0.7417)	0.7113 (0.6820–0.7351)
			HD95		58.52 (54.73–62.63)	2.599 (2.472–2.795)	6.358 (5.330–7.444)	7.237 (5.886–8.624)	5.995 (4.292–8.004)	8.919 (7.472–10.61)
		right heart atrium	Dice	Not supported	0.2813 (0.2133–0.3488)	0.9449 (0.9378–0.9496)	0.9340 (0.9273–0.9393)	0.9216 (0.9135–0.9329)	0.9160 (0.9081–0.9229)	0.9141 (0.9040–0.9212)
			HD95		41.05 (33.88–49.12)	2.484 (2.322–2.702)	2.924 (2.735–3.159)	5.011 (5.002–6.248)	5.133 (3.474–8.087)	3.797 (3.356–4.372)
		myocardium	Dice	Not supported	0.2702 (0.2313–0.3028)	0.9061 (0.9023–0.9096)	0.8890 (0.8812–0.8957)	0.8751 (0.8388–0.8822)	0.7880 (0.7765–0.7978)	0.8903 (0.8829–0.8971)
			HD95		38.70 (28.07–55.32)	2.505 (2.500–2.516)	2.757 (2.602–2.937)	3.162 (2.927–3.990)	3.767 (3.501–4.107)	2.492 (2.402–2.574)
		left heart ventricle	Dice	Not supported	0.3833 (0.3306–0.4331)	0.9479 (0.9433–0.9515)	0.7039 (0.6985–0.7098)	0.9109 (0.8894–0.9217)	0.9053 (0.8982–0.9114)	0.8937 (0.8860–0.9010)
			HD95		34.21 (28.17–41.15)	21.71 (2.496–51.06)	9.343 (9.060–9.642)	4.996 (4.590–6.756)	3.321 (3.045–3.684)	3.773 (3.504–4.057)
		right heart ventricle	Dice	Not supported	0.4086 (0.3400–0.4727)	0.9515 (0.9493–0.9536)	0.9482 (0.9444–0.9513)	0.9095 (0.9045–0.9179)	0.9291 (0.9246–0.9333)	0.9312 (0.9277–0.9345)
			HD95		32.60 (25.03–41.70)	2.497 (2.448–2.531)	2.630 (2.477–2.808)	3.587 (3.432–4.547)	3.865 (3.516–4.223)	3.230 (2.997–3.497)
		left iliac artery	Dice	0.7955 (0.7760–0.8139)	0.1794 (0.1518–0.2043)	0.7190 (0.6996–0.7388)	0.8889 (0.8732–0.9024)	0.8853 (0.8670–0.8974)	0.8490 (0.8287–0.8670)	0.8620 (0.8500–0.8742)
			HD95	34.38 (30.62–37.63)	37.28 (28.74–48.63)	374.1 (365.5–382.3)	12.71 (4.340–24.04)	13.37 (6.081–21.96)	12.68 (4.496–24.90)	13.19 (5.144–24.80)
		right iliac artery	Dice	0.6567 (0.6356–0.6774)	0.1349 (0.1080–0.1621)	0.6620 (0.6387–0.6875)	0.8887 (0.8748–0.9018)	0.8695 (0.8570–0.8824)	0.8699 (0.8523–0.8853)	0.8887 (0.8772–0.8990)
			HD95	32.74 (31.95–33.55)	44.59 (33.94–57.55)	369.2 (358.8–378.5)	21.81 (8.067–37.18)	22.30 (15.86–32.64)	33.29 (11.02–63.89)	21.54 (7.319–38.00)
		aorta	Dice	0.9444 (0.9420–0.9468)	0.5825 (0.5550–0.6068)	0.9539 (0.9518–0.9561)	0.9467 (0.9434–0.9494)	0.9355 (0.9228–0.9578)	0.9575 (0.9545–0.9604)	0.9546 (0.9504–0.9582)
			HD95	2.553 (2.299–2.930)	24.05 (21.94–26.29)	1.974 (1.877–2.074)	1.476 (1.282–1.716)	1.811 (1.461–1.855)	2.158 (1.506–2.981)	1.972 (1.560–2.577)
		left iliac vena	Dice	0.8826 (0.8668–0.8966)	0.1877 (0.1446–0.2320)	0.8555 (0.8428–0.8677)	0.9016 (0.8838–0.9173)	0.8702 (0.8409–0.8871)	0.8763 (0.8602–0.8902)	0.8986 (0.8853–0.9102)
			HD95	11.12 (3.599–21.04)	58.08 (38.93–79.16)	319.5 (271.5–360.8)	9.915 (2.566–19.44)	10.05 (3.134–19.23)	10.96 (4.076–20.61)	9.247 (2.298–19.15)
		right iliac vena	Dice	0.8070 (0.7831–0.8299)	0.2097 (0.1729–0.2475)	0.8318 (0.8151–0.8475)	0.8854 (0.8615–0.9053)	0.8647 (0.8518–0.8759)	0.8635 (0.8412–0.8843)	0.8867 (0.8683–0.9018)
			HD95	30.13 (28.07–31.69)	47.02 (32.97–62.99)	311.1 (254.6–360.9)	15.45 (5.299–27.17)	15.95 (5.687–26.32)	14.89 (4.671–26.90)	17.04 (5.911–29.27)
		inferior vena cava	Dice	0.9140 (0.9080–0.9198)	0.6443 (0.6211–0.6660)	0.9400 (0.9363–0.9435)	0.9362 (0.9316–0.9408)	0.9010 (0.8988–0.9079)	0.9110 (0.9028–0.9190)	0.9116 (0.9022–0.9208)
			HD95	4.336 (3.412–5.380)	17.21 (14.72–20.08)	2.134 (1.963–2.312)	2.698 (2.132–3.622)	2.674 (2.126–3.375)	3.747 (3.252–4.309)	3.557 (2.779–4.754)
		portal vein and splenic vein	Dice	0.8498 (0.8399–0.8596)	0.2180 (0.1927–0.2445)	0.8641 (0.8592–0.8685)	0.9029 (0.8959–0.9084)	0.8818 (0.8757–0.8840)	0.8990 (0.8930–0.9040)	0.9012 (0.8964–0.9057)
			HD95	7.703 (5.828–9.654)	54.82 (47.05–63.06)	2.476 (2.394–2.576)	4.566 (3.290–6.183)	4.813 (4.630–5.936)	2.538 (2.234–2.895)	3.283 (2.827–3.792)
		celiac trunk	Dice	Not supported	0.001992 (0.0005469–0.003963)	0.7930 (0.7688–0.8161)	0.8351 (0.8100–0.8590)	0.7977 (0.7864–0.8066)	0.8265 (0.8017–0.8498)	0.8565 (0.8304–0.8771)
			HD95		158.9 (144.5–173.4)	5.895 (3.326–9.285)	5.753 (3.524–8.671)	6.086 (5.102–9.636)	12.34 (4.206–26.21)	3.847 (2.377–5.845)
		left lung lower lobe	Dice	0.9508 (0.9374–0.9590)	0.6236 (0.5901–0.6596)	0.9484 (0.9395–0.9544)	0.9564 (0.9463–0.9629)	0.9360 (0.9194–0.9517)	0.9013 (0.8857–0.9156)	0.9240 (0.9118–0.9356)
			HD95	9.940 (6.569–13.52)	38.53 (31.96–45.58)	4.307 (3.692–4.996)	11.21 (9.895–12.55)	11.99 (10.26–15.00)	11.66 (10.03–13.48)	20.26 (10.32–38.39)
		left lung upper lobe	Dice	0.9633 (0.9525–0.9716)	0.6563 (0.6309–0.6816)	0.9566 (0.9477–0.9628)	0.9641 (0.9516–0.9724)	0.9338 (0.9303–0.9574)	0.9335 (0.9233–0.9424)	0.9406 (0.9297–0.9500)
			HD95	10.52 (8.006–13.70)	26.88 (23.30–31.73)	3.915 (3.242–4.827)	7.646 (6.400–9.193)	4.211 (3.496–5.825)	11.48 (10.00–13.09)	9.471 (8.084–10.83)
		right lung lower lobe	Dice	0.9490 (0.9221–0.9635)	0.6257 (0.6050–0.6473)	0.9511 (0.9466–0.9546)	0.9564 (0.9464–0.9634)	0.9372 (0.9224–0.9390)	0.9189 (0.9074–0.9301)	0.9318 (0.9232–0.9400)
			HD95	13.54 (12.07–15.45)	30.87 (27.94–34.20)	4.041 (3.563–4.560)	11.37 (10.13–12.58)	12.41 (11.37–14.68)	11.75 (10.44–13.25)	12.40 (10.95–14.09)
		right lung middle lobe	Dice	0.9091 (0.8707–0.9344)	0.3246 (0.2741–0.3787)	0.9088 (0.8883–0.9249)	0.9057 (0.8795–0.9254)	0.9011 (0.8825–0.9282)	0.8539 (0.8257–0.8768)	0.8673 (0.8364–0.8869)
			HD95	22.06 (6.496–48.50)	41.85 (35.62–48.42)	5.373 (4.296–6.692)	8.951 (7.637–10.35)	9.358 (9.156–11.30)	11.47 (9.731–13.48)	9.371 (8.233–10.69)
		right lung upper lobe	Dice	0.9583 (0.9544–0.9619)	0.6734 (0.6592–0.6883)	0.9470 (0.9436–0.9503)	0.9572 (0.9520–0.9618)	0.9480 (0.9264–0.9530)	0.9271 (0.9205–0.9329)	0.9393 (0.9325–0.9449)
			HD95	9.265 (7.795–11.31)	24.05 (21.59–26.79)	5.028 (4.496–5.597)	8.283 (7.113–9.525)	8.711 (6.287–10.53)	10.06 (8.625–11.63)	8.901 (7.688–10.17)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	DAP-Atlas (AutoPETIII)	bronchie	Dice	0.1935 (0.1825–0.2044)	0.001621 (0.001136–0.002129)	0.7220 (0.7096–0.7339)	0.7774 (0.7628–0.7916)	0.7678 (0.7522–0.7752)	0.6934 (0.6777–0.7091)	0.7332 (0.7193–0.7476)
			HD95	124.0 (107.5–149.6)	85.37 (82.10–88.64)	3.114 (2.815–3.448)	16.53 (14.57–18.60)	3.853 (1.889–4.030)	13.07 (11.48–14.95)	19.42 (17.43–21.23)
		trachea	Dice	0.8897 (0.8810–0.8967)	0.4201 (0.3873–0.4509)	0.9256 (0.9201–0.9309)	0.9363 (0.9278–0.9432)	0.9306 (0.9236–0.9393)	0.9204 (0.9140–0.9264)	0.9264 (0.9196–0.9318)
			HD95	7.150 (6.071–8.367)	29.79 (26.01–34.29)	1.900 (1.681–2.222)	2.537 (1.774–3.372)	2.046 (1.409–2.090)	3.180 (2.614–3.805)	2.513 (2.072–3.097)
		pulmonary artery	Dice	Not supported	0.2201 (0.1846–0.2580)	0.8639 (0.8577–0.8698)	0.8976 (0.8911–0.9033)	0.8901 (0.8847–0.9075)	0.8419 (0.8305–0.8524)	0.8536 (0.8414–0.8643)
			HD95		69.19 (63.97–74.71)	3.180 (2.792–3.835)	10.87 (10.23–11.53)	3.746 (3.675–5.235)	35.03 (32.74–37.39)	26.28 (24.63–28.04)
		left cheek	Dice	Not supported	0.0000 (0.0000–0.0000)	0.6021 (0.5242–0.6763)	0.8524 (0.8215–0.8764)	0.7623 (0.7339–0.8048)	0.7582 (0.7128–0.7959)	0.8461 (0.8073–0.8750)
			HD95		617.6 (572.1–651.2)	589.8 (544.1–627.4)	2.719 (2.376–3.079)	5.421 (4.088–7.167)	4.765 (4.167–5.377)	5.016 (2.368–10.03)
		right cheek	Dice	Not supported	0.00004926 (0.0000–0.0001478)	0.6020 (0.5057–0.7009)	0.7784 (0.6956–0.8442)	0.7763 (0.7716–0.7820)	0.7641 (0.6811–0.8281)	0.7839 (0.6969–0.8455)
			HD95		210.6 (159.4–266.9)	505.9 (429.4–566.4)	3.300 (2.710–3.950)	5.423 (3.645–7.751)	4.264 (3.125–5.701)	23.46 (2.557–65.00)
		left eyeball	Dice	Not supported	0.0000 (0.0000–0.0000)	0.8342 (0.8085–0.8557)	0.7543 (0.7138–0.7904)	0.8340 (0.8231–0.8509)	0.8617 (0.8430–0.8787)	0.8062 (0.7699–0.8376)
			HD95		631.6 (596.0–665.5)	2.338 (2.151–2.552)	3.411 (2.933–3.937)	3.995 (3.988–5.619)	2.126 (1.935–2.323)	2.396 (2.124–2.680)
		right eyeball	Dice	Not supported	0.0000 (0.0000–0.0000)	0.8133 (0.7784–0.8390)	0.7375 (0.6815–0.7808)	0.7959 (0.7956–0.8021)	0.8080 (0.7742–0.8344)	0.8079 (0.7531–0.8474)
			HD95		531.9 (489.4–576.0)	2.521 (2.277–2.872)	3.568 (2.972–4.309)	3.020 (2.825–4.115)	2.307 (2.018–2.688)	2.366 (2.167–2.575)
		nasal cavity	Dice	Not supported	0.000004220 (0.0000–0.00001266)	0.4648 (0.4356–0.4934)	0.8291 (0.8180–0.8398)	0.8246 (0.8152–0.8414)	0.8135 (0.8033–0.8229)	0.7790 (0.7673–0.7914)
			HD95		224.5 (213.8–236.3)	603.7 (580.3–626.9)	8.195 (7.300–9.164)	9.806 (7.515–12.58)	8.999 (6.358–13.41)	9.875 (8.829–11.26)
		right common carotid artery	Dice	0.7044 (0.6733–0.7359)	0.001229 (0.000007105–0.003369)	0.8131 (0.7902–0.8333)	0.8180 (0.7957–0.8372)	0.8075 (0.8046–0.8113)	0.7222 (0.6977–0.7451)	0.8342 (0.8216–0.8452)
			HD95	17.20 (7.349–35.21)	262.9 (217.1–306.9)	81.85 (27.35–137.8)	13.30 (7.653–20.13)	7.149 (4.790–11.66)	5.964 (4.302–7.799)	3.790 (2.327–5.654)
		left common carotid artery	Dice	0.7553 (0.7315–0.7749)	0.004649 (0.002764–0.007214)	0.7765 (0.7503–0.7997)	0.8694 (0.8574–0.8801)	0.7726 (0.7517–0.7926)	0.7841 (0.7680–0.7995)	0.8585 (0.8469–0.8687)
			HD95	8.814 (6.060–11.89)	297.9 (275.7–315.6)	264.2 (194.9–329.3)	5.067 (3.351–7.494)	6.471 (4.017–8.771)	4.304 (3.068–5.635)	2.077 (1.619–2.618)
		manubrium of sternum	Dice	Not supported	0.0000 (0.0000–0.0000)	0.8706 (0.8140–0.9006)	0.9008 (0.8438–0.9342)	0.8574 (0.8039–0.8841)	0.8864 (0.8294–0.9202)	0.9003 (0.8434–0.9305)
			HD95		132.2 (125.7–138.5)	36.96 (4.096–73.63)	4.173 (2.400–7.237)	4.296 (3.655–8.133)	5.473 (3.499–8.517)	3.647 (2.072–6.526)
		right internal carotid artery	Dice	Not supported	0.0000 (0.0000–0.0000)	0.7431 (0.7190–0.7682)	0.6977 (0.6567–0.7328)	0.6838 (0.6551–0.7024)	0.5546 (0.5166–0.5878)	0.7460 (0.7259–0.7651)
			HD95		489.4 (454.5–527.1)	3.881 (2.250–5.760)	7.902 (6.019–10.14)	9.794 (8.409–10.97)	8.210 (6.336–10.61)	3.496 (2.780–4.543)
		left internal carotid artery	Dice	Not supported	0.0000 (0.0000–0.0000)	0.7193 (0.6836–0.7522)	0.6930 (0.6427–0.7351)	0.7161 (0.7097–0.7216)	0.5012 (0.4571–0.5462)	0.6857 (0.6545–0.7116)
			HD95		493.7 (453.6–530.8)	4.096 (2.985–5.364)	6.939 (4.845–9.418)	8.430 (8.410–9.471)	11.66 (8.410–15.79)	22.00 (4.136–57.20)
		right internal jugular vein	Dice	Not supported	0.0000 (0.0000–0.0000)	0.8405 (0.7933–0.8723)	0.8869 (0.8575–0.9125)	0.8828 (0.8801–0.8910)	0.7248 (0.6761–0.7660)	0.8436 (0.8001–0.8785)
			HD95		349.8 (329.4–372.3)	185.9 (111.0–261.8)	7.732 (2.358–15.38)	8.191 (6.077–9.213)	9.242 (5.016–16.55)	7.338 (2.438–15.82)
		left internal jugular vein	Dice	Not supported	0.0000 (0.0000–0.0000)	0.7837 (0.7625–0.8034)	0.8833 (0.8665–0.8973)	0.8359 (0.8248–0.8512)	0.7659 (0.7405–0.7902)	0.8475 (0.8316–0.8620)
			HD95		460.1 (431.3–488.8)	429.2 (381.9–468.7)	3.949 (2.398–6.378)	7.319 (6.881–10.65)	10.56 (6.761–15.48)	9.436 (6.057–13.91)
		brachiocephalic trunk	Dice	0.7131 (0.6837–0.7389)	0.01764 (0.01304–0.02257)	0.7839 (0.7574–0.8070)	0.8492 (0.8293–0.8649)	0.8279 (0.8179–0.8340)	0.8519 (0.8364–0.8654)	0.8727 (0.8617–0.8828)
			HD95	31.67 (24.45–38.38)	259.5 (235.6–278.8)	398.6 (322.2–462.7)	5.000 (4.036–6.013)	4.498 (4.292–5.115)	5.242 (3.861–6.642)	2.994 (2.442–3.763)
		right brachiocephalic vein	Dice	0.5967 (0.5697–0.6251)	0.0000 (0.0000–0.0000)	0.8423 (0.8200–0.8608)	0.8532 (0.8285–0.8733)	0.8531 (0.8399–0.8551)	0.8335 (0.8155–0.8512)	0.8505 (0.8384–0.8638)
			HD95	23.48 (21.11–25.50)	361.0 (334.0–387.3)	4.802 (3.228–7.091)	5.831 (4.203–8.042)	5.676 (3.909–7.417)	9.650 (6.274–14.09)	5.927 (4.254–8.131)
		left brachiocephalic vein	Dice	0.2992 (0.2731–0.3256)	0.0000 (0.0000–0.0000)	0.7839 (0.7537–0.8095)	0.8582 (0.8366–0.8780)	0.8483 (0.8375–0.8755)	0.8408 (0.8151–0.8631)	0.8738 (0.8624–0.8842)
			HD95	49.32 (48.30–50.25)	390.4 (360.0–419.9)	173.2 (104.7–243.6)	5.495 (3.887–7.681)	6.837 (6.031–8.795)	16.37 (9.353–24.34)	3.426 (2.648–4.439)
		right subclavian artery	Dice	0.6952 (0.6765–0.7120)	0.0002523 (0.0000–0.0007574)	0.7656 (0.7467–0.7828)	0.8016 (0.7854–0.8166)	0.7878 (0.7744–0.7922)	0.7804 (0.7583–0.8012)	0.8090 (0.7945–0.8241)
			HD95	45.73 (10.83–84.18)	272.3 (230.3–317.7)	30.23 (4.168–67.93)	8.730 (6.845–10.88)	10.96 (8.097–14.42)	7.036 (5.164–9.132)	6.104 (4.345–8.447)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	DAP-Atlas (AutoPETIII)	left subclavian artery	Dice	0.4970 (0.4317–0.5579)	0.0006889 (0.0000–0.002056)	0.7630 (0.7402–0.7839)	0.8524 (0.8378–0.8658)	0.7489 (0.7314–0.7903)	0.8163 (0.7978–0.8340)	0.8282 (0.8087–0.8451)
			HD95	61.48 (60.25–62.77)	338.6 (290.2–387.5)	245.6 (173.9–314.6)	5.119 (3.683–6.605)	6.036 (4.774–8.516)	7.034 (3.805–12.04)	5.676 (4.345–7.089)
	FUMPE	pulmonary embolism	Dice	Not supported	0.01693 (0.0005908–0.04238)	0.2523 (0.08824–0.4299)	0.6329 (0.3842–0.7930)	0.6576 (0.6373–0.6712)	0.5593 (0.3316–0.7671)	0.6613 (0.4730–0.8039)
			HD95		71.29 (60.01–83.89)	29.91 (16.67–43.34)	29.49 (7.115–61.80)	16.96 (5.256–29.15)	21.02 (8.213–37.26)	18.68 (9.391–28.36)
	LNDb	pulmonary nodule	Dice	0.07889 (0.02078–0.1486)	0.02781 (0.00002581–0.06910)	0.2075 (0.1288–0.2919)	0.3976 (0.2999–0.4837)	0.4286 (0.3901–0.4768)	0.2866 (0.2116–0.3738)	0.2431 (0.1748–0.3177)
			HD95	143.6 (120.9–165.6)	133.8 (119.7–148.9)	109.4 (84.78–137.0)	111.0 (87.85–133.4)	73.17 (58.55–89.63)	117.6 (97.83–138.5)	101.9 (80.54–120.9)
		myocardium	Dice	Not supported	0.6081 (0.5764–0.6324)	0.8590 (0.8444–0.8735)	0.5957 (0.5483–0.6717)	0.8313 (0.7728–0.8543)	0.9054 (0.8950–0.9249)	0.9276 (0.9203–0.9393)
			HD95		28.93 (24.38–32.67)	3.614 (3.039–4.306)	48.65 (36.67–59.17)	15.38 (11.66–19.83)	2.587 (2.317–3.006)	1.993 (1.764–2.223)
		left heart atrium	Dice	Not supported	0.6115 (0.3514–0.7949)	0.9207 (0.8762–0.9583)	0.8315 (0.8184–0.8446)	0.9171 (0.9085–0.9397)	0.9351 (0.9146–0.9555)	0.9602 (0.9478–0.9726)
			HD95		27.07 (12.00–38.92)	4.429 (2.025–6.832)	39.38 (15.60–64.78)	5.050 (2.439–7.131)	5.219 (3.677–8.122)	2.027 (1.550–2.748)
		left heart ventricle	Dice	Not supported	0.6943 (0.4306–0.8603)	0.8864 (0.8572–0.9157)	0.5935 (0.5135–0.6734)	0.8828 (0.8584–0.8948)	0.9441 (0.9257–0.9626)	0.9442 (0.9253–0.9631)
			HD95		19.52 (7.927–36.93)	4.674 (3.499–5.439)	25.21 (13.34–40.02)	6.664 (6.587–7.614)	2.360 (1.775–3.189)	2.265 (1.820–2.709)
	MM-WHS-CT	right heart atrium	Dice	Not supported	0.2231 (0.005519–0.4406)	0.8987 (0.8805–0.9169)	0.6648 (0.5830–0.7465)	0.8788 (0.8592–0.9031)	0.9248 (0.9120–0.9323)	0.9318 (0.9231–0.9383)
			HD95		43.99 (34.19–54.00)	4.969 (3.809–6.923)	57.93 (47.03–68.45)	9.052 (5.461–10.95)	3.451 (3.159–3.714)	2.908 (2.815–3.029)
		right heart ventricle	Dice	Not supported	0.3194 (0.09066–0.5370)	0.8801 (0.8400–0.9247)	0.7211 (0.5942–0.8481)	0.8574 (0.8231–0.9139)	0.9106 (0.8794–0.9278)	0.9412 (0.9311–0.9540)
			HD95		42.50 (34.95–51.98)	6.131 (3.849–8.351)	18.52 (9.775–27.26)	7.463 (5.694–8.146)	3.989 (3.255–4.967)	2.444 (2.000–2.750)
		heart ascending aorta	Dice	Not supported	0.1904 (0.0000–0.5455)	0.9573 (0.9509–0.9625)	0.9511 (0.9417–0.9593)	0.9397 (0.9263–0.9497)	0.9702 (0.9600–0.9777)	0.9747 (0.9718–0.9771)
			HD95		83.21 (37.65–111.2)	1.397 (1.385–1.414)	38.77 (15.74–61.81)	2.480 (2.303–2.935)	1.529 (0.8414–2.347)	0.9220 (0.7687–1.042)
		pulmonary artery	Dice	Not supported	0.5160 (0.4455–0.5702)	0.8511 (0.8214–0.8899)	0.6634 (0.5639–0.7630)	0.8473 (0.8358–0.8645)	0.9289 (0.9135–0.9426)	0.9409 (0.9326–0.9498)
			HD95		26.11 (11.01–49.00)	12.62 (5.990–16.83)	60.64 (49.40–78.45)	12.46 (6.757–16.83)	6.162 (2.967–9.357)	4.429 (2.413–6.446)
		brainstem	Dice	Not supported	0.6153 (0.5603–0.6702)	0.8625 (0.8477–0.8749)	0.8420 (0.8162–0.8658)	0.8273 (0.8071–0.8351)	0.8754 (0.8597–0.8892)	0.8819 (0.8675–0.8937)
			HD95		9.847 (7.874–12.45)	3.884 (3.238–4.656)	6.226 (4.205–8.293)	5.746 (4.387–6.782)	3.726 (3.100–4.454)	3.211 (2.724–3.688)
	PDDCA	optic chiasm	Dice	Not supported	0.0000 (0.0000–0.0000)	0.5630 (0.4955–0.6301)	0.4431 (0.3986–0.4980)	0.4965 (0.4574–0.5488)	0.6481 (0.6062–0.6896)	0.7737 (0.7439–0.7999)
			HD95		175.9 (171.7–181.1)	3.822 (3.291–4.444)	5.338 (4.598–6.106)	5.315 (3.458–5.822)	3.282 (2.928–3.815)	2.741 (1.802–3.896)
		mandible	Dice	Not supported	0.5184 (0.4733–0.5575)	0.8954 (0.8873–0.9033)	0.8949 (0.8826–0.9076)	0.8638 (0.8474–0.8830)	0.8673 (0.8469–0.8848)	0.9125 (0.9052–0.9193)
			HD95		15.28 (12.02–18.88)	2.492 (2.183–2.762)	2.156 (1.724–2.576)	3.863 (3.050–3.866)	7.679 (3.631–12.67)	1.938 (1.500–2.398)
		left optic nerve	Dice	Not supported	0.0000 (0.0000–0.0000)	0.7070 (0.6452–0.7620)	0.6626 (0.6272–0.6959)	0.6577 (0.6156–0.7051)	0.7765 (0.7505–0.8053)	0.8098 (0.7812–0.8354)
			HD95		106.5 (77.89–133.2)	2.792 (1.774–3.960)	11.85 (9.834–13.80)	3.332 (2.372–4.161)	10.58 (2.947–19.88)	1.654 (1.200–2.174)
		right optic nerve	Dice	Not supported	0.02323 (0.006107–0.04765)	0.6973 (0.6530–0.7342)	0.6758 (0.6194–0.7272)	0.6794 (0.6733–0.7025)	0.7632 (0.7186–0.8023)	0.7601 (0.7186–0.7974)
			HD95		68.86 (39.27–103.1)	2.606 (2.055–3.097)	8.878 (7.019–10.64)	5.081 (3.547–8.034)	5.623 (2.312–10.04)	1.983 (1.344–2.747)
		left parotid gland	Dice	Not supported	0.5095 (0.3683–0.6314)	0.8597 (0.8442–0.8744)	0.8560 (0.8368–0.8737)	0.8591 (0.8443–0.8683)	0.8626 (0.8442–0.8784)	0.8806 (0.8696–0.8898)
			HD95		24.98 (11.64–43.96)	4.398 (3.305–5.606)	6.265 (4.040–9.234)	6.543 (4.460–6.710)	3.965 (3.183–5.232)	3.348 (2.822–4.028)
		right parotid gland	Dice	Not supported	0.4319 (0.3026–0.5364)	0.8115 (0.7407–0.8584)	0.8049 (0.7109–0.8639)	0.7940 (0.7560–0.8142)	0.8499 (0.8137–0.8795)	0.8803 (0.8674–0.8915)
			HD95		39.31 (21.77–62.28)	5.413 (3.928–7.804)	9.726 (5.397–13.99)	7.985 (6.031–9.407)	5.249 (3.556–7.514)	3.204 (2.621–4.069)
		left submandibular gland	Dice	Not supported	0.2459 (0.1449–0.3622)	0.7976 (0.7579–0.8336)	0.8127 (0.7682–0.8477)	0.7917 (0.7705–0.8159)	0.8431 (0.8226–0.8623)	0.8555 (0.8369–0.8771)
			HD95		63.69 (40.89–84.26)	4.417 (3.253–5.760)	4.063 (3.114–5.333)	6.889 (6.157–8.636)	3.208 (2.399–4.338)	2.354 (1.970–2.694)
		right submandibular gland	Dice	Not supported	0.2038 (0.1526–0.2576)	0.7851 (0.7607–0.8089)	0.7615 (0.7223–0.7995)	0.7455 (0.7187–0.7795)	0.8442 (0.8220–0.8634)	0.8433 (0.8210–0.8637)
			HD95		42.13 (24.70–60.87)	4.741 (3.549–6.021)	5.828 (4.385–7.428)	5.632 (3.594–7.057)	2.553 (2.305–2.847)	2.601 (2.379–2.873)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	Parse2022	pulmonary artery	Dice	Not supported	0.4241 (0.4018–0.4473)	0.7171 (0.6921–0.7391)	0.7607 (0.7380–0.7802)	0.7608 (0.7444–0.7797)	0.6592 (0.6368–0.6813)	0.6863 (0.6633–0.7088)
			HD95		91.97 (88.05–95.73)	12.71 (9.875–16.87)	10.51 (10.04–10.98)	15.15 (14.31–16.02)	37.47 (35.63–39.20)	17.88 (16.90–19.02)
	SEGA	aorta	Dice	0.7740 (0.6601–0.8384)	0.5003 (0.3442–0.6359)	0.8306 (0.7755–0.8700)	0.8418 (0.8217–0.8637)	0.8664 (0.8567–0.8931)	0.8479 (0.8235–0.8706)	0.8711 (0.8524–0.8904)
			HD95	13.70 (9.765–17.13)	109.0 (77.64–139.2)	41.59 (20.42–66.65)	133.3 (89.68–175.8)	16.95 (10.61–22.16)	82.98 (45.52–126.3)	44.09 (23.92–80.65)
		brain	Dice	0.9660 (0.9634–0.9679)	0.1132 (0.05332–0.1867)	0.9655 (0.9640–0.9669)	0.9783 (0.9775–0.9791)	0.9673 (0.9599–0.9791)	0.9692 (0.9677–0.9706)	0.9755 (0.9747–0.9763)
			HD95	33.44 (3.443–73.77)	78.50 (70.50–85.79)	21.30 (14.64–27.55)	2.783 (2.657–2.894)	5.422 (4.875–6.679)	15.54 (10.17–21.44)	25.93 (3.705–60.01)
		brainstem	Dice	Not supported	0.4554 (0.3789–0.5264)	0.9065 (0.8986–0.9140)	0.8634 (0.8476–0.8785)	0.8930 (0.8798–0.8990)	0.8884 (0.8836–0.8940)	0.8853 (0.8777–0.8924)
			HD95		27.82 (19.52–37.81)	3.224 (2.956–3.552)	4.249 (3.554–5.067)	4.565 (4.262–5.158)	3.266 (3.052–3.559)	3.147 (2.936–3.413)
		optic chiasm	Dice	Not supported	0.0000 (0.0000–0.0000)	0.6811 (0.6291–0.7325)	0.5421 (0.4857–0.5945)	0.6743 (0.6635–0.6854)	0.6744 (0.6394–0.7064)	0.7272 (0.6940–0.7570)
			HD95		144.2 (126.9–157.2)	13.89 (2.634–36.01)	4.145 (3.810–4.527)	5.145 (4.137–7.064)	3.276 (3.100–3.490)	2.714 (2.430–2.986)
		left temporal lobe	Dice	Not supported	0.0005553 (0.0000–0.001666)	0.9352 (0.9273–0.9413)	0.9117 (0.9032–0.9193)	0.9339 (0.9300–0.9388)	0.8920 (0.8825–0.8998)	0.9183 (0.9103–0.9248)
			HD95		74.62 (67.20–86.61)	35.01 (2.894–83.21)	4.066 (3.545–4.633)	5.290 (4.008–5.398)	4.539 (4.079–5.163)	17.46 (3.903–44.21)
		right temporal lobe	Dice	Not supported	0.0000 (0.0000–0.0000)	0.9380 (0.9335–0.9424)	0.9169 (0.9089–0.9241)	0.9239 (0.9214–0.9298)	0.9067 (0.8996–0.9132)	0.8993 (0.8883–0.9085)
			HD95		75.77 (72.17–80.32)	3.025 (2.903–3.159)	4.164 (3.634–4.730)	3.767 (2.429–5.315)	4.333 (3.892–4.881)	4.173 (3.787–4.560)
		left hippocampus	Dice	Not supported	0.0000 (0.0000–0.0000)	0.7873 (0.7573–0.8163)	0.7328 (0.6903–0.7691)	0.7801 (0.7795–0.7856)	0.7372 (0.7076–0.7640)	0.7952 (0.7728–0.8166)
			HD95		113.9 (99.57–132.9)	2.803 (2.541–3.078)	18.74 (3.289–49.20)	3.809 (2.731–4.990)	3.821 (3.393–4.332)	2.938 (2.613–3.380)
		right hippocampus	Dice	Not supported	0.0000 (0.0000–0.0000)	0.7719 (0.7319–0.8033)	0.7289 (0.6923–0.7629)	0.7565 (0.7495–0.7736)	0.7405 (0.7113–0.7651)	0.7559 (0.7269–0.7866)
			HD95		127.3 (117.7–142.7)	2.823 (2.546–3.165)	3.707 (3.217–4.227)	5.969 (5.365–6.873)	9.871 (5.584–15.69)	3.796 (2.854–5.174)
		left eyeball	Dice	Not supported	0.3173 (0.2483–0.3906)	0.9342 (0.9266–0.9404)	0.8972 (0.8738–0.9129)	0.8747 (0.8502–0.8843)	0.9198 (0.9133–0.9262)	0.9368 (0.9309–0.9428)
			HD95		30.06 (13.69–55.30)	1.623 (1.355–1.932)	2.054 (1.783–2.346)	3.295 (3.281–4.255)	2.019 (1.732–2.271)	14.05 (1.278–39.38)
		right eyeball	Dice	Not supported	0.3824 (0.3106–0.4545)	0.9232 (0.9099–0.9335)	0.8977 (0.8828–0.9107)	0.8816 (0.8675–0.8871)	0.9141 (0.9025–0.9243)	0.9287 (0.9199–0.9365)
			HD95		17.24 (11.09–24.63)	1.753 (1.497–2.038)	2.343 (2.068–2.632)	2.920 (2.178–3.799)	4.567 (1.791–9.789)	1.577 (1.313–1.875)
	SegRap2023-Task1	left lens	Dice	Not supported	0.04166 (0.02331–0.06401)	0.7540 (0.7287–0.7778)	0.7133 (0.6823–0.7454)	0.7263 (0.7120–0.7416)	0.7640 (0.7416–0.7864)	0.8307 (0.8036–0.8555)
			HD95		62.95 (42.04–85.42)	2.128 (1.757–2.487)	2.438 (2.053–2.757)	3.600 (2.501–3.757)	1.821 (1.519–2.153)	1.130 (0.8656–1.469)
		right lens	Dice	Not supported	0.03684 (0.01618–0.06324)	0.7605 (0.7235–0.7948)	0.7313 (0.6976–0.7669)	0.7467 (0.7234–0.7789)	0.7547 (0.7262–0.7824)	0.8195 (0.8018–0.8374)
			HD95		63.51 (44.47–83.19)	1.711 (1.411–2.037)	2.128 (1.771–2.488)	2.483 (1.624–2.490)	2.071 (1.647–2.476)	8.196 (1.322–15.33)
		left optic nerve	Dice	Not supported	0.0009220 (0.0000–0.002766)	0.7755 (0.7482–0.8006)	0.7598 (0.7413–0.7773)	0.7495 (0.7327–0.7567)	0.7995 (0.7769–0.8188)	0.8215 (0.8018–0.8385)
			HD95		103.7 (83.60–121.3)	2.035 (1.680–2.424)	2.774 (2.420–3.156)	3.546 (1.610–3.894)	22.74 (14.96–30.25)	17.84 (2.496–47.29)
		right optic nerve	Dice	Not supported	0.01034 (0.001394–0.02128)	0.8024 (0.7733–0.8275)	0.7717 (0.7498–0.7938)	0.7902 (0.7797–0.8266)	0.8175 (0.7963–0.8375)	0.8423 (0.8283–0.8563)
			HD95		83.99 (62.35–103.5)	2.061 (1.511–2.678)	3.075 (2.714–3.488)	3.725 (3.323–4.670)	2.234 (1.844–2.579)	17.63 (1.674–49.11)
		left middle ear	Dice	Not supported	0.0000 (0.0000–0.0000)	0.8427 (0.7932–0.8794)	0.6232 (0.5797–0.6611)	0.8193 (0.8128–0.8275)	0.8025 (0.7626–0.8323)	0.8228 (0.7924–0.8507)
			HD95		101.4 (99.47–103.4)	1.573 (1.104–2.223)	8.420 (7.658–9.161)	5.287 (5.028–6.574)	4.505 (2.365–8.399)	2.265 (1.918–2.628)
		right middle ear	Dice	Not supported	0.0000 (0.0000–0.0000)	0.8382 (0.7939–0.8684)	0.5335 (0.4938–0.5701)	0.8198 (0.8179–0.8260)	0.7367 (0.6920–0.7738)	0.8236 (0.7975–0.8486)
			HD95		100.8 (96.03–104.6)	1.765 (1.343–2.267)	10.75 (10.20–11.29)	2.284 (2.064–5.129)	2.827 (2.534–3.217)	2.323 (1.999–2.658)
		left internal auditory canal	Dice	Not supported	0.0000 (0.0000–0.0000)	0.8296 (0.8025–0.8549)	0.7951 (0.7654–0.8206)	0.8013 (0.7912–0.8242)	0.7869 (0.7336–0.8285)	0.8231 (0.7931–0.8504)
			HD95		97.83 (92.50–102.6)	1.024 (0.7869–1.283)	1.396 (1.084–1.735)	3.227 (1.484–4.340)	7.363 (1.711–14.49)	1.286 (0.9647–1.630)
		right internal auditory canal	Dice	Not supported	0.0000 (0.0000–0.0000)	0.8229 (0.7985–0.8445)	0.8197 (0.7914–0.8453)	0.7710 (0.7673–0.7793)	0.8139 (0.7904–0.8332)	0.8406 (0.8083–0.8680)
			HD95		105.9 (100.4–111.2)	1.096 (0.8031–1.463)	1.524 (1.118–1.918)	2.125 (1.995–3.533)	1.281 (1.021–1.568)	1.055 (0.7603–1.383)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	SegRap2023-Task1	left tympanic cavity	Dice	Not supported	0.0000 (0.0000–0.0000)	0.7740 (0.7468–0.8000)	0.7242 (0.6853–0.7621)	0.7736 (0.7720–0.7801)	0.7040 (0.6653–0.7399)	0.7873 (0.7634–0.8118)
			HD95		99.75 (95.55–103.6)	2.007 (1.702–2.327)	2.356 (2.025–2.652)	6.398 (6.051–7.854)	3.161 (2.389–4.442)	1.726 (1.503–2.000)
		right tympanic cavity	Dice	Not supported	0.0000 (0.0000–0.0000)	0.7953 (0.7622–0.8243)	0.7564 (0.7189–0.7892)	0.7323 (0.6928–0.7506)	0.7465 (0.7035–0.7938)	0.7339 (0.6984–0.7718)
			HD95		98.81 (93.21–103.7)	1.868 (1.547–2.207)	2.309 (2.003–2.619)	3.297 (2.943–4.379)	1.946 (1.631–2.276)	1.866 (1.600–2.128)
		left vestibule semicircular canal	Dice	Not supported	0.0000 (0.0000–0.0000)	0.8257 (0.8007–0.8481)	0.7883 (0.7602–0.8140)	0.8093 (0.8042–0.8258)	0.7808 (0.7615–0.8007)	0.7726 (0.7391–0.7982)
			HD95		90.92 (87.15–94.57)	1.197 (0.8641–1.579)	1.900 (1.559–2.283)	2.205 (1.846–3.174)	16.76 (5.152–28.92)	1.383 (1.098–1.694)
		right vestibule semicircular canal	Dice	Not supported	0.0000 (0.0000–0.0000)	0.8026 (0.7791–0.8234)	0.7946 (0.7728–0.8157)	0.7967 (0.7597–0.8263)	0.7144 (0.6839–0.7419)	0.7915 (0.7500–0.8258)
			HD95		96.76 (91.51–102.7)	1.160 (0.8867–1.475)	1.540 (1.208–1.941)	1.787 (1.457–2.393)	2.581 (2.272–2.907)	1.494 (1.148–1.856)
		left cochlea	Dice	Not supported	0.0000 (0.0000–0.0000)	0.8313 (0.8083–0.8536)	0.8022 (0.7797–0.8240)	0.7953 (0.7670–0.8135)	0.8057 (0.7686–0.8340)	0.8193 (0.7936–0.8420)
			HD95		106.9 (88.20–135.2)	1.064 (0.7804–1.412)	1.402 (1.063–1.799)	1.752 (1.541–2.636)	1.630 (1.196–2.037)	0.9620 (0.7490–1.236)
		right cochlea	Dice	Not supported	0.0000 (0.0000–0.0000)	0.8188 (0.7821–0.8483)	0.8190 (0.7921–0.8410)	0.8051 (0.8026–0.8075)	0.8136 (0.7916–0.8350)	0.8175 (0.7987–0.8358)
			HD95		101.0 (87.69–113.1)	1.060 (0.8283–1.359)	1.140 (0.8660–1.471)	2.035 (1.469–3.151)	1.758 (1.316–2.173)	1.161 (0.8825–1.487)
		left eustachian tube bone	Dice	Not supported	0.0000 (0.0000–0.0000)	0.7966 (0.7725–0.8175)	0.7415 (0.7181–0.7709)	0.7215 (0.6943–0.7564)	0.7354 (0.7145–0.7575)	0.7144 (0.6772–0.7521)
			HD95		78.80 (77.06–80.60)	1.523 (1.169–1.907)	2.024 (1.673–2.383)	2.978 (2.408–3.375)	1.962 (1.658–2.243)	1.778 (1.515–2.072)
		right eustachian tube bone	Dice	Not supported	0.0000 (0.0000–0.0000)	0.7901 (0.7505–0.8217)	0.7267 (0.6954–0.7560)	0.7725 (0.7643–0.7782)	0.7646 (0.7303–0.7954)	0.7876 (0.7619–0.8111)
			HD95		80.10 (76.35–84.03)	1.430 (1.094–1.782)	2.147 (1.850–2.485)	2.605 (2.398–3.855)	1.632 (1.356–1.924)	1.653 (1.322–1.975)
		pituitary gland	Dice	Not supported	0.0000 (0.0000–0.0000)	0.7516 (0.7059–0.7943)	0.6432 (0.5977–0.6878)	0.6317 (0.5931–0.6589)	0.6987 (0.6580–0.7374)	0.7089 (0.6176–0.7863)
			HD95		141.6 (136.4–147.2)	1.840 (1.403–2.292)	2.848 (2.533–3.192)	2.909 (1.857–3.028)	2.621 (2.317–2.886)	2.272 (1.899–2.643)
		oral cavity	Dice	Not supported	0.5814 (0.5512–0.6106)	0.9180 (0.9076–0.9271)	0.8707 (0.8613–0.8797)	0.9082 (0.9031–0.9202)	0.9003 (0.8910–0.9075)	0.9165 (0.9106–0.9221)
			HD95		17.08 (15.36–19.15)	3.781 (3.430–4.247)	6.309 (5.498–7.258)	4.176 (3.530–4.416)	14.20 (3.987–34.50)	4.723 (3.831–5.858)
		left mandible	Dice	Not supported	0.4324 (0.4092–0.4519)	0.9357 (0.9299–0.9404)	0.9187 (0.9141–0.9228)	0.9286 (0.9260–0.9372)	0.9274 (0.9212–0.9320)	0.9258 (0.9205–0.9304)
			HD95		74.77 (71.26–78.22)	1.788 (1.574–2.032)	2.412 (2.147–2.638)	2.663 (1.826–3.067)	7.625 (1.731–19.24)	6.087 (2.503–12.97)
		right mandible	Dice	Not supported	0.4186 (0.4033–0.4344)	0.9388 (0.9327–0.9435)	0.9198 (0.9158–0.9240)	0.9191 (0.8919–0.9322)	0.9177 (0.9108–0.9235)	0.9352 (0.9298–0.9396)
			HD95		78.61 (76.11–81.26)	1.715 (1.490–1.952)	2.190 (1.952–2.416)	2.079 (1.732–4.093)	21.63 (2.594–51.25)	1.611 (1.372–1.865)
		left submandibular gland	Dice	Not supported	0.2330 (0.1479–0.3233)	0.8765 (0.8527–0.8965)	0.8372 (0.8232–0.8495)	0.8220 (0.8161–0.8279)	0.8415 (0.8276–0.8571)	0.8844 (0.8663–0.9000)
			HD95		53.27 (40.93–65.07)	2.481 (2.063–2.980)	2.981 (2.687–3.286)	5.078 (4.921–5.476)	3.088 (2.811–3.386)	2.429 (1.884–3.248)
		right submandibular gland	Dice	Not supported	0.2871 (0.2123–0.3590)	0.8723 (0.8514–0.8904)	0.8442 (0.8282–0.8602)	0.8638 (0.8498–0.8840)	0.8490 (0.8302–0.8662)	0.8767 (0.8642–0.8888)
			HD95		45.84 (34.18–57.92)	2.492 (2.103–2.948)	2.793 (2.545–3.122)	3.005 (1.478–3.328)	14.93 (2.636–39.11)	2.266 (1.964–2.507)
		left parotid gland	Dice	Not supported	0.4154 (0.3494–0.4743)	0.9170 (0.9113–0.9228)	0.8788 (0.8675–0.8887)	0.8688 (0.8658–0.8738)	0.8954 (0.8895–0.9007)	0.9085 (0.9022–0.9139)
			HD95		34.94 (22.96–47.82)	2.396 (2.153–2.634)	3.442 (3.034–3.997)	3.517 (2.488–3.691)	3.325 (3.057–3.647)	2.725 (2.423–3.059)
		right parotid gland	Dice	Not supported	0.4737 (0.3963–0.5456)	0.9050 (0.8918–0.9149)	0.8764 (0.8681–0.8847)	0.8998 (0.8946–0.9068)	0.8740 (0.8636–0.8838)	0.8916 (0.8853–0.8987)
			HD95		29.54 (17.23–42.58)	2.881 (2.608–3.137)	4.190 (3.613–4.811)	4.127 (3.867–5.500)	5.468 (4.635–6.460)	4.190 (3.343–5.439)
		left mastoid process	Dice	Not supported	0.0000 (0.0000–0.0000)	0.9001 (0.8912–0.9084)	0.8625 (0.8505–0.8737)	0.8583 (0.8519–0.8801)	0.8804 (0.8674–0.8926)	0.8896 (0.8799–0.8993)
			HD95		88.12 (78.53–103.5)	2.132 (1.899–2.388)	3.638 (3.181–4.263)	4.105 (2.892–4.359)	4.731 (2.613–8.690)	2.622 (2.339–2.905)
		right mastoid process	Dice	Not supported	0.0000 (0.0000–0.0000)	0.9016 (0.8836–0.9153)	0.8576 (0.8386–0.8731)	0.8836 (0.8820–0.8944)	0.8717 (0.8497–0.8882)	0.8858 (0.8717–0.8970)
			HD95		88.99 (84.59–93.08)	2.460 (1.910–3.495)	3.515 (2.942–4.307)	3.660 (2.745–5.490)	2.979 (2.666–3.341)	2.964 (2.529–3.510)
		left temporomandibular joint	Dice	Not supported	0.002152 (0.00004084–0.004916)	0.8665 (0.8439–0.8876)	0.8346 (0.8170–0.8510)	0.8516 (0.8462–0.8533)	0.8147 (0.7916–0.8355)	0.8676 (0.8518–0.8832)
			HD95		101.2 (99.04–103.3)	2.156 (1.802–2.527)	2.785 (2.585–2.939)	3.614 (2.983–3.750)	24.30 (10.24–38.55)	2.385 (2.072–2.670)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	SegRap2023-Task1	right temporomandibular joint	Dice	Not supported	0.009331 (0.002716–0.01814)	0.8894 (0.8763–0.9004)	0.8466 (0.8321–0.8601)	0.8785 (0.8555–0.8929)	0.7384 (0.7136–0.7646)	0.8687 (0.8535–0.8824)
			HD95		93.57 (84.93–100.5)	2.022 (1.711–2.372)	2.740 (2.548–2.912)	5.153 (4.454–5.202)	19.16 (3.782–49.41)	2.330 (2.057–2.621)
		spinal cord	Dice	0.7975 (0.7839–0.8107)	0.1869 (0.1496–0.2226)	0.8709 (0.8525–0.8862)	0.8239 (0.7806–0.8604)	0.8319 (0.8033–0.8507)	0.8396 (0.8222–0.8547)	0.8623 (0.8448–0.8775)
			HD95	7.998 (4.392–12.72)	110.4 (99.98–120.9)	5.125 (2.021–9.731)	43.10 (7.395–82.88)	5.727 (2.054–10.91)	7.202 (3.840–11.83)	25.43 (11.31–41.74)
		esophagus	Dice	0.6289 (0.6077–0.6512)	0.09060 (0.05086–0.1324)	0.7896 (0.7726–0.8043)	0.7504 (0.7336–0.7663)	0.7850 (0.7714–0.7871)	0.8123 (0.7943–0.8293)	0.8198 (0.8038–0.8360)
			HD95	12.22 (10.78–13.81)	96.59 (80.59–114.1)	3.381 (2.816–4.080)	4.376 (3.837–4.921)	4.496 (3.445–5.751)	3.294 (2.871–3.873)	3.510 (2.976–4.112)
		larynx	Dice	Not supported	0.02382 (0.006618–0.05056)	0.8597 (0.8258–0.8803)	0.7720 (0.7524–0.7914)	0.8651 (0.8496–0.8683)	0.9003 (0.8898–0.9087)	0.9029 (0.8955–0.9099)
			HD95		71.01 (64.22–78.57)	30.49 (8.881–62.24)	21.37 (10.13–42.58)	7.188 (5.649–10.95)	24.15 (4.508–54.72)	3.837 (3.228–4.595)
		larynx glottis	Dice	Not supported	0.1967 (0.1373–0.2578)	0.7457 (0.7108–0.7799)	0.6913 (0.6665–0.7126)	0.7078 (0.6653–0.7273)	0.7432 (0.7107–0.7746)	0.7731 (0.7452–0.8003)
			HD95		16.26 (13.47–19.24)	14.75 (3.112–37.68)	3.873 (3.534–4.288)	6.623 (5.703–7.933)	3.567 (3.289–3.897)	2.992 (2.827–3.179)
		larynx supraglottis	Dice	Not supported	0.2230 (0.1535–0.2908)	0.8223 (0.7946–0.8490)	0.7856 (0.7660–0.8056)	0.7797 (0.7510–0.7838)	0.8235 (0.8035–0.8416)	0.8269 (0.8133–0.8403)
			HD95		44.00 (33.76–53.85)	3.013 (2.614–3.517)	4.192 (3.678–4.703)	3.982 (3.792–4.973)	3.269 (3.030–3.557)	3.135 (3.005–3.295)
		pharynx constrictor muscle	Dice	Not supported	0.03987 (0.02739–0.05410)	0.7754 (0.7549–0.7938)	0.7443 (0.7206–0.7646)	0.7726 (0.7623–0.7801)	0.7229 (0.7106–0.7369)	0.7880 (0.7767–0.7989)
			HD95		29.24 (26.77–32.01)	14.99 (2.933–38.85)	3.632 (3.085–4.306)	5.987 (3.404–6.582)	15.16 (3.236–38.74)	2.394 (2.205–2.624)
	SegRap2023-Task2	thyroid	Dice	0.7331 (0.7136–0.7501)	0.0000 (0.0000–0.0000)	0.8539 (0.8384–0.8666)	0.8239 (0.8074–0.8400)	0.7876 (0.7789–0.7981)	0.8320 (0.8192–0.8443)	0.8466 (0.8339–0.8587)
			HD95	6.385 (5.551–7.396)	110.8 (104.9–116.3)	3.018 (2.842–3.226)	3.268 (2.959–3.752)	3.315 (3.009–3.725)	2.907 (2.697–3.091)	3.002 (2.774–3.245)
		trachea	Dice	0.5316 (0.5136–0.5516)	0.009948 (0.0000–0.02355)	0.8612 (0.8376–0.8831)	0.8221 (0.7862–0.8502)	0.8424 (0.8394–0.8568)	0.8784 (0.8633–0.8932)	0.8993 (0.8839–0.9139)
			HD95	36.33 (31.81–40.35)	166.1 (145.6–184.9)	11.87 (7.287–17.43)	10.83 (7.197–15.84)	11.52 (7.207–15.70)	5.006 (3.575–6.777)	5.039 (3.185–7.666)
		nasopharyngeal tumor	Dice	Not supported	0.002187 (0.0001856–0.005513)	0.7788 (0.7398–0.8097)	0.3064 (0.2417–0.3724)	0.8296 (0.8294–0.8526)	0.6328 (0.5843–0.6755)	0.7244 (0.6923–0.7484)
			HD95		71.72 (65.00–78.35)	19.98 (7.535–41.61)	38.52 (21.02–64.17)	20.53 (8.774–41.37)	14.47 (11.22–18.62)	29.87 (14.93–56.91)
		nasopharyngeal lymph node	Dice	Not supported	0.001640 (0.0000–0.004416)	0.6381 (0.5694–0.6997)	0.004323 (0.0000–0.01297)	0.7123 (0.6978–0.7196)	0.3367 (0.2793–0.3920)	0.5250 (0.4580–0.5865)
			HD95		64.11 (59.90–67.79)	30.79 (15.00–59.02)	108.0 (84.78–131.3)	23.40 (20.04–27.76)	30.33 (24.93–36.53)	33.31 (24.50–43.14)
	SegTHOR	esophagus	Dice	0.7929 (0.7621–0.8268)	0.3342 (0.2458–0.4129)	0.7966 (0.7598–0.8337)	0.8287 (0.8036–0.8532)	0.8099 (0.8012–0.8177)	0.8294 (0.8077–0.8500)	0.8339 (0.8143–0.8561)
			HD95	8.426 (4.842–12.72)	21.88 (13.53–35.45)	6.908 (4.097–11.00)	3.744 (3.096–4.567)	7.661 (6.495–8.199)	3.664 (3.053–4.280)	3.812 (2.825–4.952)
		heart	Dice	0.7648 (0.7395–0.7896)	0.2601 (0.1695–0.3392)	0.9186 (0.8740–0.9524)	0.9120 (0.8794–0.9349)	0.8955 (0.8808–0.9100)	0.9350 (0.9256–0.9438)	0.9305 (0.9194–0.9409)
			HD95	29.31 (27.25–31.12)	63.70 (43.22–90.88)	8.279 (5.200–11.80)	11.78 (7.189–16.80)	12.81 (10.46–13.11)	47.16 (20.97–82.00)	31.47 (6.373–79.55)
		trachea	Dice	0.8959 (0.8813–0.9106)	0.5922 (0.5182–0.6514)	0.8885 (0.8750–0.9026)	0.9172 (0.9071–0.9281)	0.9016 (0.8936–0.9099)	0.8959 (0.8908–0.9015)	0.9003 (0.8924–0.9077)
			HD95	4.159 (2.820–5.655)	26.44 (14.78–40.81)	4.447 (3.442–5.533)	5.246 (2.574–8.599)	6.156 (5.129–7.758)	5.775 (4.437–7.170)	4.669 (3.653–5.682)
		aorta	Dice	0.7894 (0.7438–0.8260)	0.6177 (0.5032–0.6998)	0.8495 (0.7910–0.8990)	0.8296 (0.7816–0.8696)	0.8257 (0.7921–0.8571)	0.8597 (0.8252–0.8894)	0.8551 (0.8073–0.8912)
			HD95	52.74 (34.71–76.24)	40.05 (23.19–58.40)	30.12 (10.38–54.99)	45.19 (24.96–69.20)	41.08 (21.89–64.22)	43.83 (21.16–73.13)	44.52 (22.48–73.98)
		brain	Dice	0.9487 (0.9454–0.9521)	0.1908 (0.1360–0.2567)	0.9492 (0.9440–0.9542)	0.8216 (0.7494–0.8873)	0.9286 (0.9194–0.9423)	0.9686 (0.9653–0.9716)	0.9780 (0.9761–0.9796)
			HD95	4.111 (3.708–4.630)	26.39 (23.04–29.58)	5.958 (4.384–8.392)	22.43 (15.53–28.61)	3.987 (2.675–4.132)	14.41 (8.480–20.76)	5.168 (4.422–5.894)
	Seg-CQ500	intracranial hemorrhage	Dice	Not supported	0.02812 (0.002430–0.07246)	0.4952 (0.2779–0.7230)	0.1473 (0.01200–0.2897)	0.8216 (0.7790–0.8381)	0.6198 (0.4365–0.7955)	0.5941 (0.3719–0.8015)
			HD95		84.83 (76.67–94.50)	23.52 (8.838–41.17)	76.04 (57.66–94.57)	5.583 (4.114–11.80)	17.52 (6.278–31.80)	35.20 (11.78–64.26)
	NSCLC-Radiomics	esophagus	Dice	0.7441 (0.7082–0.7751)	0.3488 (0.2735–0.4136)	0.7510 (0.7183–0.7785)	0.7314 (0.6948–0.7654)	0.7287 (0.7044–0.7469)	0.7622 (0.7383–0.7843)	0.7689 (0.7436–0.7913)
			HD95	14.45 (10.15–19.22)	26.51 (18.60–36.25)	10.72 (7.604–14.35)	12.07 (8.671–16.01)	15.39 (12.46–20.01)	11.11 (7.256–15.19)	10.60 (7.008–14.54)
		lung tumor	Dice	0.3292 (0.1941–0.4679)	0.3666 (0.2287–0.5092)	0.3165 (0.1727–0.4573)	0.7002 (0.5498–0.8220)	0.6031 (0.4662–0.7297)	0.5078 (0.3461–0.6692)	0.6038 (0.4493–0.7512)
			HD95	120.8 (81.65–171.3)	76.72 (39.93–120.0)	69.50 (42.54–98.61)	48.63 (21.44–80.43)	49.43 (24.79–77.47)	66.03 (39.77–93.92)	45.00 (23.72–72.16)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	NSCLC-Radiomics	heart	Dice	0.8111 (0.7760–0.8462)	0.2579 (0.1326–0.3645)	0.8787 (0.8572–0.9042)	0.8588 (0.8325–0.8830)	0.8843 (0.8501–0.8955)	0.8946 (0.8705–0.9224)	0.8947 (0.8757–0.9175)
			HD95	32.06 (30.08–35.56)	56.88 (37.64–80.16)	13.52 (10.13–16.90)	22.37 (16.20–29.69)	15.02 (14.76–15.79)	26.80 (10.04–52.09)	15.33 (12.09–20.51)
		left lung	Dice	0.9452 (0.9157–0.9669)	0.8312 (0.7301–0.8980)	0.9415 (0.9036–0.9650)	0.9616 (0.9458–0.9734)	0.9557 (0.9400–0.9741)	0.9711 (0.9616–0.9773)	0.9678 (0.9591–0.9737)
			HD95	12.56 (5.295–22.32)	20.99 (10.55–39.87)	6.996 (4.826–9.815)	5.922 (3.167–9.832)	3.761 (3.369–5.570)	2.970 (2.524–3.424)	4.224 (3.211–5.400)
		right lung	Dice	0.9527 (0.9432–0.9619)	0.8164 (0.6881–0.9053)	0.9633 (0.9523–0.9714)	0.9652 (0.9583–0.9713)	0.9296 (0.9213–0.9411)	0.9702 (0.9635–0.9754)	0.9721 (0.9671–0.9757)
			HD95	15.28 (12.32–18.42)	24.90 (10.56–50.77)	5.912 (4.355–7.527)	6.316 (3.701–10.59)	6.890 (3.147–9.680)	8.072 (5.297–11.02)	5.759 (3.182–9.861)
		spinal cord	Dice	0.7842 (0.7344–0.8251)	0.006259 (0.0003417–0.01737)	0.3921 (0.3035–0.4863)	0.8107 (0.7638–0.8503)	0.7653 (0.7064–0.8074)	0.7847 (0.7308–0.8309)	0.8163 (0.7625–0.8645)
			HD95	34.38 (12.20–61.09)	249.5 (217.2–277.6)	60.58 (38.68–83.81)	29.46 (10.71–53.04)	32.95 (13.76–55.11)	46.43 (27.37–73.98)	30.19 (9.560–56.22)
	PleThora (NSCLC-Radiomics)	lung effusion	Dice	Not supported	0.04945 (0.007386–0.1178)	0.5973 (0.5033–0.6830)	0.6690 (0.5858–0.7394)	0.5744 (0.4791–0.6476)	0.5300 (0.4047–0.6480)	0.5465 (0.4292–0.6523)
			HD95		123.9 (100.5–146.9)	41.69 (26.59–59.37)	53.55 (35.96–73.33)	51.98 (35.84–71.85)	65.09 (38.85–101.3)	48.50 (34.29–64.61)
		thoracic cavity	Dice	Not supported	0.7464 (0.6227–0.8425)	0.9540 (0.9455–0.9607)	0.2585 (0.1935–0.3326)	0.9407 (0.9200–0.9543)	0.9476 (0.9375–0.9544)	0.9468 (0.9393–0.9524)
			HD95		32.50 (19.49–53.79)	5.425 (3.570–8.044)	32.32 (28.08–37.18)	7.518 (6.408–10.30)	7.958 (5.830–11.17)	9.108 (6.889–11.58)
	WORD	liver	Dice	0.9561 (0.9504–0.9614)	0.8224 (0.8126–0.8315)	0.9625 (0.9587–0.9662)	0.9627 (0.9590–0.9661)	0.9276 (0.9261–0.9276)	0.9621 (0.9585–0.9655)	0.9627 (0.9595–0.9660)
			HD95	5.376 (3.831–8.065)	16.24 (14.54–18.00)	3.539 (3.175–3.974)	4.218 (3.429–5.193)	5.881 (5.434–6.735)	4.207 (3.348–5.537)	4.916 (3.508–7.178)
		spleen	Dice	0.9437 (0.9375–0.9490)	0.6545 (0.5700–0.7203)	0.9528 (0.9485–0.9573)	0.9560 (0.9520–0.9600)	0.9422 (0.9379–0.9472)	0.9554 (0.9508–0.9595)	0.9519 (0.9474–0.9565)
			HD95	2.760 (2.437–3.189)	25.74 (14.54–41.17)	2.331 (2.174–2.561)	2.091 (1.739–2.608)	2.308 (2.269–2.728)	2.038 (1.839–2.233)	2.189 (1.897–2.556)
		left kidney	Dice	0.9402 (0.9339–0.9459)	0.6298 (0.5064–0.7208)	0.9498 (0.9439–0.9554)	0.9518 (0.9458–0.9574)	0.9374 (0.9341–0.9379)	0.9505 (0.9460–0.9550)	0.9466 (0.9404–0.9524)
			HD95	2.947 (2.651–3.360)	49.39 (25.18–80.52)	2.750 (2.492–3.036)	2.342 (2.003–2.726)	3.689 (2.955–4.706)	2.578 (2.351–2.806)	3.569 (2.741–4.480)
		right kidney	Dice	0.9425 (0.9344–0.9490)	0.6971 (0.6089–0.7622)	0.9522 (0.9455–0.9580)	0.9529 (0.9448–0.9591)	0.9374 (0.9367–0.9603)	0.9486 (0.9416–0.9546)	0.9504 (0.9446–0.9556)
			HD95	3.003 (2.686–3.420)	29.79 (12.19–57.95)	2.479 (2.226–2.697)	2.161 (1.863–2.484)	2.594 (1.848–3.269)	2.721 (2.140–3.507)	2.570 (2.163–3.040)
		stomach	Dice	0.9248 (0.9162–0.9333)	0.6049 (0.5649–0.6431)	0.8931 (0.8224–0.9344)	0.9230 (0.9073–0.9349)	0.9040 (0.8936–0.9091)	0.9306 (0.9228–0.9384)	0.9310 (0.9222–0.9406)
			HD95	8.752 (5.570–12.66)	34.81 (24.50–46.19)	44.23 (4.850–101.9)	9.741 (6.021–14.58)	10.55 (5.212–14.93)	7.678 (4.563–11.80)	7.250 (3.783–11.83)
		gallbladder	Dice	0.6820 (0.5374–0.8048)	0.1990 (0.1243–0.2705)	0.7758 (0.6790–0.8486)	0.6624 (0.5235–0.7896)	0.7604 (0.6827–0.8236)	0.7916 (0.6927–0.8637)	0.8013 (0.7334–0.8602)
			HD95	11.69 (5.726–18.98)	36.07 (26.23–47.81)	5.879 (3.664–8.850)	11.81 (6.575–18.22)	8.354 (6.296–9.125)	6.723 (4.002–10.28)	5.564 (4.099–7.152)
		esophagus	Dice	0.7668 (0.7381–0.7922)	0.2922 (0.2220–0.3561)	0.7973 (0.7758–0.8199)	0.7887 (0.7672–0.8076)	0.7910 (0.7595–0.8096)	0.7953 (0.7757–0.8149)	0.8077 (0.7886–0.8267)
			HD95	6.856 (2.874–12.02)	18.86 (15.61–22.73)	5.172 (3.924–6.774)	5.484 (4.476–6.650)	6.299 (5.175–6.738)	4.816 (3.920–5.804)	4.419 (3.588–5.442)
		pancreas	Dice	0.8051 (0.7719–0.8312)	0.3685 (0.3015–0.4333)	0.8349 (0.8063–0.8593)	0.8217 (0.7879–0.8483)	0.8201 (0.7803–0.8477)	0.8415 (0.8207–0.8583)	0.8442 (0.8192–0.8638)
			HD95	21.78 (5.518–52.62)	28.37 (20.44–37.13)	5.426 (3.865–7.636)	7.738 (5.342–11.00)	5.755 (4.015–8.534)	6.037 (4.696–7.833)	5.675 (4.309–7.397)
		duodenum	Dice	0.6498 (0.5802–0.7098)	0.3182 (0.2619–0.3680)	0.7024 (0.6356–0.7593)	0.6791 (0.6088–0.7356)	0.6900 (0.6358–0.7355)	0.7156 (0.6618–0.7573)	0.7491 (0.6938–0.7943)
			HD95	18.97 (13.37–25.85)	36.53 (28.25–45.74)	15.10 (9.578–21.58)	19.33 (13.68–25.90)	17.69 (11.89–24.65)	18.18 (13.85–23.63)	13.61 (8.403–19.70)
		colon	Dice	0.7646 (0.7145–0.8037)	0.3529 (0.2545–0.4412)	0.8537 (0.8273–0.8746)	0.7982 (0.7603–0.8291)	0.7949 (0.7523–0.8196)	0.8205 (0.7846–0.8509)	0.8439 (0.8226–0.8636)
			HD95	33.62 (24.69–44.81)	63.04 (45.15–81.67)	13.31 (6.995–23.67)	39.11 (23.16–56.89)	23.60 (15.07–33.76)	17.79 (10.44–28.35)	20.11 (13.12–29.16)
		intestine	Dice	Not supported	0.4437 (0.3432–0.5353)	0.8718 (0.8613–0.8831)	0.7057 (0.6793–0.7331)	0.8536 (0.8445–0.8619)	0.8427 (0.8270–0.8571)	0.8516 (0.8328–0.8672)
			HD95		42.23 (28.14–59.90)	6.439 (5.231–7.722)	40.65 (35.26–45.63)	7.284 (5.424–8.345)	16.31 (9.763–26.94)	9.200 (6.981–11.50)
		adrenal gland	Dice	0.6666 (0.6256–0.7069)	0.1799 (0.1239–0.2339)	0.7212 (0.6967–0.7446)	0.7110 (0.6752–0.7423)	0.6981 (0.6830–0.7173)	0.7223 (0.6895–0.7550)	0.7493 (0.7253–0.7734)
			HD95	7.641 (5.610–10.48)	31.66 (22.67–42.24)	5.437 (4.164–7.441)	6.760 (4.863–9.333)	5.890 (3.245–6.450)	7.579 (5.000–10.79)	5.193 (3.919–6.912)
		rectum	Dice	Not supported	0.3615 (0.2513–0.4570)	0.8143 (0.7940–0.8369)	0.7670 (0.7200–0.8008)	0.7936 (0.7813–0.8252)	0.7852 (0.7314–0.8252)	0.7967 (0.7655–0.8247)
			HD95		55.30 (23.55–95.91)	8.083 (6.297–10.10)	13.12 (10.46–15.77)	10.46 (7.858–12.74)	12.41 (8.604–16.50)	11.46 (8.807–14.43)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	WORD	urinary bladder	Dice	0.9302 (0.9120–0.9462)	0.5440 (0.4430–0.6387)	0.9386 (0.9188–0.9526)	0.9392 (0.9205–0.9538)	0.9259 (0.9196–0.9583)	0.9390 (0.9175–0.9547)	0.9387 (0.9185–0.9542)
			HD95	3.797 (3.219–4.564)	37.30 (20.22–63.40)	3.199 (2.939–3.555)	3.194 (2.831–3.783)	6.697 (5.355–6.890)	5.799 (2.865–11.05)	3.649 (2.875–4.758)
		head of left femur	Dice	Not supported	0.6506 (0.6189–0.6827)	0.9211 (0.9047–0.9350)	0.9208 (0.9039–0.9364)	0.9205 (0.9185–0.9278)	0.9140 (0.9001–0.9263)	0.9246 (0.9113–0.9351)
			HD95		28.61 (17.34–41.61)	91.48 (23.69–167.1)	7.549 (4.657–11.45)	8.945 (6.646–12.70)	10.51 (5.966–15.77)	28.48 (9.954–51.01)
		head of right femur	Dice	Not supported	0.6252 (0.5480–0.6835)	0.9234 (0.9062–0.9359)	0.9203 (0.9016–0.9358)	0.9059 (0.8786–0.9314)	0.9245 (0.9105–0.9366)	0.9263 (0.9146–0.9361)
			HD95		32.73 (17.56–55.25)	46.12 (5.689–100.4)	7.149 (4.382–10.57)	9.459 (4.065–12.59)	5.970 (3.900–9.030)	4.915 (3.471–7.136)
		liver	Dice	0.9545 (0.9493–0.9587)	0.8250 (0.8112–0.8370)	0.9596 (0.9562–0.9628)	0.9599 (0.9556–0.9637)	0.9524 (0.9401–0.9584)	0.9614 (0.9592–0.9634)	0.9614 (0.9586–0.9636)
			HD95	4.994 (4.014–6.185)	14.54 (13.62–15.52)	3.597 (3.274–3.967)	4.561 (3.546–5.934)	6.038 (4.464–6.394)	4.126 (3.525–4.922)	4.382 (3.571–5.383)
		spleen	Dice	0.9374 (0.9306–0.9443)	0.7319 (0.7068–0.7536)	0.9418 (0.9362–0.9472)	0.9481 (0.9418–0.9539)	0.9362 (0.9351–0.9368)	0.9524 (0.9490–0.9555)	0.9515 (0.9484–0.9543)
			HD95	3.617 (2.749–4.782)	13.28 (11.68–15.27)	2.765 (2.497–3.053)	3.931 (2.181–6.172)	5.082 (4.815–5.305)	2.204 (2.048–2.366)	2.238 (2.081–2.391)
		left kidney	Dice	0.9365 (0.9320–0.9409)	0.7612 (0.7403–0.7803)	0.9317 (0.9242–0.9383)	0.9464 (0.9411–0.9511)	0.9357 (0.9313–0.9534)	0.9454 (0.9396–0.9494)	0.9429 (0.9378–0.9467)
			HD95	3.354 (2.971–3.880)	11.46 (10.39–12.60)	3.449 (3.014–4.047)	2.765 (2.332–3.223)	4.628 (2.939–5.962)	3.480 (2.912–4.247)	4.980 (3.242–8.035)
		right kidney	Dice	0.9382 (0.9313–0.9432)	0.7887 (0.7714–0.8026)	0.9365 (0.9304–0.9425)	0.9466 (0.9384–0.9524)	0.9218 (0.9127–0.9369)	0.9438 (0.9393–0.9480)	0.9455 (0.9420–0.9488)
			HD95	3.316 (2.936–3.992)	10.79 (9.995–11.53)	3.147 (2.857–3.554)	2.650 (2.223–3.359)	3.450 (3.359–3.919)	3.410 (2.927–4.003)	7.064 (4.369–10.71)
		stomach	Dice	0.8676 (0.8018–0.9095)	0.6473 (0.5819–0.6971)	0.8765 (0.8121–0.9134)	0.8746 (0.8148–0.9134)	0.9041 (0.8781–0.9612)	0.9184 (0.9081–0.9272)	0.9217 (0.9118–0.9306)
			HD95	14.78 (10.02–20.06)	22.80 (17.77–29.24)	10.36 (6.315–16.00)	16.39 (11.43–22.44)	8.450 (4.787–14.38)	10.16 (7.629–13.21)	9.883 (7.418–12.80)
		gallbladder	Dice	0.7358 (0.6713–0.7868)	0.3236 (0.2514–0.3946)	0.7673 (0.7296–0.8033)	0.7393 (0.6812–0.7864)	0.8033 (0.7574–0.8409)	0.8169 (0.7926–0.8393)	0.8240 (0.7975–0.8451)
			HD95	10.92 (7.612–14.62)	20.63 (17.99–23.60)	10.25 (6.568–15.45)	11.35 (7.865–15.29)	7.185 (6.599–9.051)	6.754 (5.153–8.691)	5.886 (4.651–7.428)
		esophagus	Dice	0.7685 (0.7353–0.7943)	0.4198 (0.3874–0.4515)	0.7760 (0.7424–0.8036)	0.7673 (0.7385–0.7944)	0.8336 (0.7989–0.8482)	0.8053 (0.7914–0.8180)	0.8099 (0.7964–0.8236)
			HD95	30.57 (8.348–59.75)	12.02 (10.51–13.56)	5.966 (4.438–8.250)	6.512 (4.966–8.513)	5.853 (4.826–6.825)	4.492 (4.014–5.007)	4.515 (3.987–5.036)
		pancreas	Dice	0.7892 (0.7649–0.8094)	0.5131 (0.4806–0.5411)	0.8053 (0.7818–0.8257)	0.8016 (0.7780–0.8242)	0.8129 (0.7849–0.8556)	0.8196 (0.8013–0.8368)	0.8241 (0.8053–0.8410)
			HD95	10.32 (6.822–14.76)	15.70 (13.43–18.54)	7.053 (5.038–9.524)	9.073 (6.444–12.20)	10.90 (8.258–13.88)	9.211 (7.126–11.75)	9.012 (6.840–11.76)
	RAOS	duodenum	Dice	0.6143 (0.5403–0.6784)	0.4098 (0.3613–0.4537)	0.6310 (0.5575–0.7017)	0.6450 (0.5709–0.7078)	0.7201 (0.6682–0.7556)	0.7041 (0.6704–0.7333)	0.7288 (0.6905–0.7636)
			HD95	22.14 (17.21–28.21)	25.75 (21.40–31.05)	19.52 (14.35–25.32)	21.12 (15.94–26.93)	18.45 (16.18–22.84)	25.33 (22.00–28.73)	18.34 (14.96–21.87)
		colon	Dice	0.7469 (0.7073–0.7825)	0.3845 (0.3066–0.4612)	0.7715 (0.7316–0.8062)	0.7669 (0.7236–0.8037)	0.8754 (0.8502–0.8972)	0.8163 (0.7990–0.8317)	0.8351 (0.8176–0.8495)
			HD95	29.23 (25.11–33.38)	43.23 (34.60–53.47)	17.42 (13.96–21.16)	27.79 (23.78–32.04)	14.72 (10.72–18.49)	22.12 (17.97–27.24)	21.77 (17.44–26.85)
		intestine	Dice	Not supported	0.3515 (0.2814–0.4179)	0.8458 (0.8276–0.8625)	0.6839 (0.6589–0.7069)	0.8630 (0.8469–0.8925)	0.8388 (0.8275–0.8503)	0.8505 (0.8383–0.8626)
			HD95		58.43 (43.30–75.63)	9.352 (7.254–11.87)	41.62 (36.79–47.07)	11.09 (10.32–13.61)	11.89 (9.730–14.54)	9.696 (7.925–12.25)
		left adrenal gland	Dice	0.01525 (0.0000–0.04466)	0.002400 (0.0000–0.007201)	0.01603 (0.0000–0.04624)	0.01824 (0.0000–0.05232)	0.7215 (0.7087–0.7407)	0.6964 (0.6529–0.7305)	0.7053 (0.6625–0.7392)
			HD95	71.57 (68.15–74.92)	70.72 (67.71–73.82)	70.32 (66.90–73.90)	71.78 (67.77–75.30)	8.610 (6.440–12.11)	8.615 (6.024–11.74)	9.519 (6.009–13.69)
		right adrenal gland	Dice	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.7193 (0.7187–0.7233)	0.7152 (0.6723–0.7527)	0.7072 (0.6689–0.7409)
			HD95	70.44 (66.85–74.02)	70.94 (65.53–78.46)	70.01 (66.82–73.32)	70.49 (66.92–73.99)	11.78 (9.431–14.44)	7.578 (5.391–10.34)	42.40 (35.89–48.67)
		rectum	Dice	Not supported	0.1497 (0.08999–0.2126)	0.7783 (0.7403–0.8119)	0.7365 (0.6982–0.7702)	0.8329 (0.7934–0.8718)	0.8072 (0.7896–0.8219)	0.8087 (0.7921–0.8247)
			HD95		55.81 (37.81–79.58)	9.956 (8.103–12.00)	16.37 (12.93–20.08)	10.50 (8.550–12.87)	16.14 (10.09–26.96)	13.09 (10.72–15.79)
		urinary bladder	Dice	0.8954 (0.8245–0.9367)	0.5695 (0.4928–0.6453)	0.9042 (0.8380–0.9451)	0.8928 (0.8215–0.9368)	0.9004 (0.8574–0.9432)	0.9356 (0.9184–0.9484)	0.9369 (0.9186–0.9497)
			HD95	5.114 (3.745–7.050)	23.86 (17.65–32.51)	4.041 (3.185–5.571)	5.695 (3.991–7.951)	7.878 (6.553–11.25)	5.819 (3.046–9.718)	3.461 (2.895–4.411)
		head of left femur	Dice	Not supported	0.6079 (0.5587–0.6512)	0.9151 (0.8953–0.9317)	0.9136 (0.8943–0.9292)	0.8304 (0.8046–0.8421)	0.9106 (0.8969–0.9213)	0.9129 (0.9005–0.9239)
			HD95		62.66 (46.11–82.92)	46.23 (11.48–85.58)	9.496 (6.699–13.27)	13.99 (12.78–14.97)	13.84 (7.138–25.02)	20.75 (11.86–33.51)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	RAOS	head of right femur	Dice	Not supported	0.5986 (0.5664–0.6284)	0.9209 (0.9040–0.9342)	0.9136 (0.8945–0.9282)	0.7018 (0.6888–0.7365)	0.9118 (0.8998–0.9233)	0.9150 (0.9043–0.9245)
			HD95		74.17 (61.63–88.30)	6.826 (4.622–9.580)	8.983 (6.080–12.69)	17.04 (15.10–19.35)	9.930 (7.096–13.17)	13.10 (6.734–24.08)
		prostate	Dice	0.7673 (0.7177–0.8065)	0.2085 (0.1081–0.3163)	0.7353 (0.6681–0.7862)	0.5378 (0.4298–0.6341)	0.9290 (0.8990–0.9703)	0.8348 (0.8146–0.8524)	0.8129 (0.7770–0.8421)
			HD95	7.388 (2.587–13.25)	29.07 (17.47–41.76)	6.835 (5.780–8.072)	11.71 (9.720–13.83)	6.866 (5.423–8.651)	5.673 (4.951–6.464)	34.36 (18.55–55.45)
		seminal vesicles	Dice	Not supported	0.02716 (0.002789–0.06138)	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.8400 (0.8374–0.8475)	0.7599 (0.7392–0.7789)	0.7818 (0.7598–0.8013)
			HD95		189.8 (85.70–296.9)	237.8 (210.1–269.0)	– (– –)	5.419 (4.215–7.770)	6.569 (5.599–7.540)	8.075 (5.282–12.24)
	CT-ICH	intracranial hemorrhage	Dice	Not supported	0.0000 (0.0000–0.0000)	0.08367 (0.0000–0.1979)	0.5030 (0.3032–0.6733)	0.8201 (0.8127–0.8340)	0.3163 (0.1163–0.4908)	0.5493 (0.3747–0.7037)
			HD95		122.8 (100.2–145.4)	82.02 (54.01–106.7)	33.98 (13.57–54.43)	3.460 (2.367–4.882)	47.93 (27.41–69.88)	46.60 (17.99–75.98)
	Mediastinal-Lymph-Node-SEG	mediastinal lymph node	Dice	Not supported	0.01460 (0.005448–0.02726)	0.0000 (0.0000–0.0000)	0.2783 (0.2171–0.3352)	0.7820 (0.7806–0.7914)	0.5012 (0.4340–0.5682)	0.4822 (0.4106–0.5526)
			HD95		125.9 (112.0–140.2)	158.7 (157.0–160.3)	62.42 (54.47–70.47)	11.67 (9.630–15.04)	48.21 (40.38–57.15)	35.92 (30.22–41.89)
		bone	Dice	Not supported	0.5845 (0.4665–0.6773)	0.0000 (0.0000–0.0000)	0.9024 (0.8538–0.9378)	0.8171 (0.7605–0.8559)	0.8811 (0.8263–0.9214)	0.8795 (0.8248–0.9217)
			HD95		112.8 (68.89–156.8)	132.9 (120.0–141.3)	11.05 (1.767–22.89)	8.409 (2.383–13.47)	8.229 (2.000–14.52)	7.145 (2.000–12.29)
	3D-ircadb-01	liver	Dice	0.9678 (0.9628–0.9727)	0.9222 (0.9191–0.9251)	0.9650 (0.9598–0.9685)	0.9721 (0.9681–0.9756)	0.9677 (0.9569–0.9792)	0.9723 (0.9688–0.9761)	0.9719 (0.9679–0.9761)
			HD95	3.142 (2.362–3.896)	7.052 (5.940–8.419)	3.091 (2.711–3.498)	2.740 (1.902–3.886)	5.842 (3.723–7.285)	2.468 (1.957–2.978)	2.692 (2.119–3.195)
		portal vein and splenic vein	Dice	0.4911 (0.4250–0.5473)	0.2635 (0.2221–0.2957)	0.4503 (0.4146–0.4810)	0.5759 (0.5259–0.6252)	0.4812 (0.4343–0.5168)	0.5875 (0.5317–0.6463)	0.6472 (0.6025–0.6919)
			HD95	51.78 (49.72–53.91)	42.12 (35.64–48.35)	53.32 (51.96–55.27)	43.72 (37.14–51.50)	42.92 (41.33–47.20)	33.90 (26.05–45.64)	23.68 (18.46–28.89)
		aorta	Dice	0.7878 (0.7393–0.8332)	0.7307 (0.6808–0.7732)	0.7872 (0.7315–0.8250)	0.7945 (0.7518–0.8408)	0.7863 (0.7468–0.8170)	0.8036 (0.7584–0.8318)	0.8038 (0.7523–0.8299)
			HD95	53.17 (48.02–60.93)	46.47 (42.87–53.08)	47.76 (43.01–56.13)	52.60 (48.06–59.85)	47.15 (44.49–54.07)	51.87 (46.14–59.70)	51.79 (46.03–59.57)
		liver tumor	Dice	0.5198 (0.2507–0.8050)	0.07503 (0.0000–0.2251)	0.4454 (0.02872–0.7456)	0.8311 (0.7883–0.8941)	0.6983 (0.6132–0.8362)	0.8194 (0.7534–0.9027)	0.8457 (0.8176–0.8923)
			HD95	51.34 (42.91–66.24)	97.45 (66.08–113.3)	20.68 (3.490–40.96)	43.37 (1.600–93.57)	8.199 (2.845–18.58)	2.973 (1.600–4.644)	2.584 (1.600–3.892)
		veins	Dice	Not supported	0.5324 (0.4392–0.5965)	0.0000 (0.0000–0.0000)	0.7299 (0.6719–0.7918)	0.7281 (0.7268–0.7286)	0.6620 (0.6524–0.6717)	0.6669 (0.6051–0.7344)
			HD95		72.67 (70.04–75.74)	– (– –)	29.36 (12.11–57.09)	29.77 (28.59–34.18)	49.66 (26.73–65.78)	37.42 (20.01–59.97)
		inferior vena cava	Dice	0.5794 (0.5794–0.5794)	0.5353 (0.5353–0.5353)	0.5705 (0.5705–0.5705)	0.6000 (0.6000–0.6000)	0.5644 (0.5575–0.5684)	0.5829 (0.5829–0.5829)	0.5872 (0.5872–0.5872)
			HD95	76.57 (76.57–76.57)	80.83 (80.83–80.83)	83.67 (83.67–83.67)	83.29 (83.29–83.29)	78.05 (78.05–78.83)	83.34 (83.34–83.34)	84.85 (84.85–84.85)
		gallbladder	Dice	0.8815 (0.8815–0.8815)	0.6900 (0.6900–0.6900)	0.8776 (0.8776–0.8776)	0.8965 (0.8965–0.8965)	0.8789 (0.8730–0.8968)	0.8867 (0.8867–0.8867)	0.8943 (0.8943–0.8943)
			HD95	19.75 (19.75–19.75)	14.16 (14.16–14.16)	9.857 (9.857–9.857)	24.64 (24.64–24.64)	14.57 (14.08–15.08)	25.25 (25.25–25.25)	25.01 (25.01–25.01)
		left kidney	Dice	0.9628 (0.9628–0.9628)	0.9042 (0.9042–0.9042)	0.9499 (0.9499–0.9499)	0.9683 (0.9683–0.9683)	0.9502 (0.9488–0.9511)	0.9592 (0.9592–0.9592)	0.9683 (0.9683–0.9683)
			HD95	1.600 (1.600–1.600)	3.927 (3.927–3.927)	1.755 (1.755–1.755)	1.440 (1.440–1.440)	2.050 (1.601–3.018)	1.610 (1.610–1.610)	1.610 (1.610–1.610)
		right kidney	Dice	0.9601 (0.9601–0.9601)	0.8177 (0.8177–0.8177)	0.9509 (0.9509–0.9509)	0.9705 (0.9705–0.9705)	0.9538 (0.9385–0.9619)	0.9672 (0.9672–0.9672)	0.9674 (0.9674–0.9674)
			HD95	1.610 (1.610–1.610)	117.9 (117.9–117.9)	1.755 (1.755–1.755)	1.600 (1.600–1.600)	2.339 (1.620–2.775)	3.200 (3.200–3.200)	1.897 (1.897–1.897)
		left adrenal gland	Dice	0.8121 (0.8121–0.8121)	0.5377 (0.5377–0.5377)	0.8138 (0.8138–0.8138)	0.8982 (0.8982–0.8982)	0.7860 (0.7815–0.7896)	0.8941 (0.8941–0.8941)	0.9046 (0.9046–0.9046)
			HD95	1.755 (1.755–1.755)	11.06 (11.06–11.06)	1.610 (1.610–1.610)	0.7200 (0.7200–0.7200)	3.289 (3.089–3.314)	1.440 (1.440–1.440)	0.7200 (0.7200–0.7200)
	NSCLC-Radiogenomics	right adrenal gland	Dice	0.8104 (0.8104–0.8104)	0.4224 (0.4224–0.4224)	0.8088 (0.8088–0.8088)	0.8991 (0.8991–0.8991)	0.8751 (0.8606–0.8787)	0.8440 (0.8440–0.8440)	0.8821 (0.8821–0.8821)
			HD95	2.270 (2.270–2.270)	14.57 (14.57–14.57)	1.610 (1.610–1.610)	0.7200 (0.7200–0.7200)	3.140 (1.665–3.158)	2.270 (2.270–2.270)	1.018 (1.018–1.018)
		lung	Dice	0.9472 (0.9369–0.9566)	0.6152 (0.5479–0.6779)	0.9538 (0.9436–0.9619)	0.9546 (0.9443–0.9643)	0.9392 (0.9296–0.9462)	0.9543 (0.9440–0.9634)	0.9556 (0.9453–0.9647)
			HD95	49.72 (30.58–71.65)	51.88 (38.96–64.97)	5.690 (3.152–9.277)	6.630 (4.998–8.923)	5.714 (3.814–7.182)	5.979 (3.594–9.650)	4.318 (3.693–4.981)
		pulmonary nodule	Dice	0.4320 (0.2977–0.5661)	0.02529 (0.0000–0.07588)	0.3483 (0.2092–0.4943)	0.5581 (0.4313–0.6751)	0.6351 (0.6127–0.6894)	0.5554 (0.4137–0.6900)	0.5672 (0.4249–0.6971)
			HD95	93.74 (66.38–121.2)	155.6 (131.4–180.8)	99.94 (62.40–135.3)	112.2 (77.67–147.0)	94.37 (66.66–121.9)	105.8 (74.06–140.6)	104.2 (68.58–139.1)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	NSCLC-Radiogenomics	lung tumor	Dice	0.3638 (0.2777–0.4394)	0.02776 (0.0000–0.07949)	0.3917 (0.2764–0.5044)	0.5069 (0.4106–0.5937)	0.6019 (0.5772–0.6640)	0.5040 (0.3922–0.6015)	0.5293 (0.4267–0.6226)
			HD95	34.78 (31.30–38.29)	205.8 (158.6–256.4)	127.2 (77.62–181.4)	92.41 (58.31–134.8)	45.15 (28.48–64.70)	61.82 (32.35–98.13)	48.41 (27.87–71.93)
	PanTS	aorta	Dice	0.4541 (0.4300–0.4766)	0.7624 (0.7530–0.7707)	0.4356 (0.4122–0.4581)	0.6878 (0.6647–0.7086)	0.8252 (0.8185–0.8280)	0.8672 (0.8607–0.8740)	0.8734 (0.8673–0.8800)
			HD95	39.06 (36.21–42.56)	11.29 (10.18–12.39)	42.40 (38.88–46.10)	24.16 (21.97–26.45)	10.55 (9.339–12.25)	12.45 (11.18–13.63)	10.92 (9.794–12.05)
		inferior vena cava	Dice	0.3369 (0.3133–0.3618)	0.7009 (0.6881–0.7124)	0.3534 (0.3311–0.3762)	0.4062 (0.3802–0.4316)	0.7823 (0.7732–0.7851)	0.7711 (0.7603–0.7809)	0.7961 (0.7867–0.8050)
			HD95	44.18 (40.76–47.49)	15.35 (13.83–17.09)	43.68 (39.94–47.21)	36.38 (34.11–38.92)	15.07 (14.41–17.10)	16.15 (14.47–17.86)	15.16 (13.53–16.94)
		liver	Dice	0.6077 (0.5863–0.6305)	0.8984 (0.8939–0.9029)	0.6115 (0.5912–0.6320)	0.7486 (0.7267–0.7695)	0.9346 (0.9340–0.9387)	0.9487 (0.9445–0.9527)	0.9505 (0.9469–0.9538)
			HD95	55.10 (52.10–58.02)	4.960 (4.622–5.374)	51.67 (48.93–54.40)	47.70 (44.42–50.80)	6.835 (6.480–6.978)	6.905 (6.224–7.635)	6.792 (6.112–7.579)
		stomach	Dice	0.3325 (0.3088–0.3542)	0.7145 (0.6999–0.7290)	0.2082 (0.1887–0.2285)	0.6270 (0.6031–0.6500)	0.8181 (0.8083–0.8185)	0.7996 (0.7863–0.8115)	0.8022 (0.7907–0.8141)
			HD95	67.03 (63.01–71.02)	15.68 (13.78–17.67)	95.60 (91.47–99.96)	43.62 (40.19–47.28)	16.83 (15.88–17.49)	15.18 (13.86–16.64)	15.81 (14.38–17.34)
		colon	Dice	0.5691 (0.5525–0.5863)	0.2119 (0.1953–0.2293)	0.4808 (0.4648–0.4979)	0.6438 (0.6261–0.6615)	0.7847 (0.7743–0.7968)	0.7601 (0.7502–0.7698)	0.7866 (0.7779–0.7952)
			HD95	37.37 (35.34–39.56)	67.36 (64.38–70.47)	36.46 (34.39–38.51)	36.61 (34.12–39.06)	18.36 (17.68–20.25)	18.40 (16.85–20.04)	16.58 (15.23–18.03)
		right lung	Dice	0.3535 (0.3274–0.3793)	0.7908 (0.7746–0.8069)	0.3660 (0.3407–0.3921)	0.4576 (0.4257–0.4888)	0.8913 (0.8658–0.8949)	0.9276 (0.9168–0.9374)	0.9306 (0.9207–0.9398)
			HD95	71.01 (66.95–74.93)	20.09 (17.71–22.62)	73.88 (70.05–77.68)	70.62 (65.90–75.58)	10.03 (9.038–11.11)	11.60 (9.775–13.55)	10.90 (9.086–12.76)
		left lung	Dice	0.3674 (0.3403–0.3951)	0.7780 (0.7612–0.7935)	0.3600 (0.3352–0.3870)	0.4539 (0.4228–0.4830)	0.8742 (0.8643–0.8856)	0.8983 (0.8862–0.9103)	0.8996 (0.8876–0.9111)
			HD95	84.98 (80.72–89.18)	23.14 (20.44–25.63)	88.28 (84.45–92.56)	78.85 (73.80–83.99)	15.52 (14.19–17.68)	17.18 (14.96–19.70)	13.45 (11.06–15.90)
		pancreas	Dice	0.3120 (0.2916–0.3341)	0.6155 (0.6051–0.6258)	0.2678 (0.2488–0.2873)	0.6739 (0.6565–0.6903)	0.7548 (0.7394–0.7604)	0.7709 (0.7628–0.7802)	0.7892 (0.7814–0.7976)
			HD95	50.85 (47.33–54.15)	11.94 (10.63–13.38)	52.76 (49.66–56.11)	18.11 (16.57–19.72)	8.055 (8.010–8.980)	9.050 (8.101–10.10)	8.815 (7.881–9.832)
		veins	Dice	Not supported	0.008733 (0.006887–0.01088)	0.2638 (0.2334–0.2947)	0.3386 (0.3039–0.3726)	0.6376 (0.6281–0.6465)	0.6529 (0.6425–0.6633)	0.6825 (0.6723–0.6923)
			HD95		62.92 (61.64–64.21)	76.27 (58.92–92.89)	54.28 (52.44–56.14)	18.35 (17.32–19.29)	18.26 (17.21–19.31)	15.13 (14.14–16.10)
		right adrenal gland	Dice	0.1957 (0.1782–0.2141)	0.3739 (0.3607–0.3864)	0.1447 (0.1283–0.1617)	0.3238 (0.2989–0.3461)	0.5824 (0.5626–0.5921)	0.6659 (0.6543–0.6768)	0.7012 (0.6920–0.7095)
			HD95	28.82 (26.04–31.51)	15.39 (13.83–17.10)	30.33 (26.52–33.99)	32.12 (29.50–34.83)	5.974 (5.225–6.720)	10.81 (9.769–11.98)	4.553 (4.201–4.928)
		duodenum	Dice	0.2378 (0.2205–0.2551)	0.4751 (0.4628–0.4867)	0.1855 (0.1695–0.2010)	0.4623 (0.4403–0.4820)	0.7457 (0.7449–0.7597)	0.6729 (0.6620–0.6831)	0.7310 (0.7221–0.7398)
			HD95	45.45 (42.49–48.54)	18.66 (17.49–19.84)	55.11 (51.06–58.75)	28.09 (26.25–30.01)	10.83 (9.180–12.07)	13.19 (12.50–14.07)	10.50 (9.851–11.26)
		head of pancreas	Dice	Not supported	0.3991 (0.3864–0.4129)	0.004113 (0.002059–0.006583)	0.3186 (0.3054–0.3323)	0.7616 (0.7581–0.7805)	0.7294 (0.7210–0.7382)	0.7756 (0.7673–0.7839)
			HD95		60.14 (58.31–61.85)	90.92 (78.69–104.2)	58.14 (56.32–59.85)	6.441 (5.380–9.631)	11.00 (10.37–11.76)	7.414 (6.752–8.345)
		right kidney	Dice	0.3233 (0.2969–0.3525)	0.7752 (0.7659–0.7835)	0.3198 (0.2917–0.3488)	0.4395 (0.4065–0.4697)	0.9013 (0.9008–0.9089)	0.9292 (0.9236–0.9339)	0.9330 (0.9285–0.9372)
			HD95	74.58 (70.16–78.80)	51.16 (47.35–54.93)	69.94 (65.56–74.55)	68.24 (63.49–72.90)	4.831 (4.690–5.364)	5.436 (4.592–6.389)	3.957 (3.573–4.345)
		spleen	Dice	0.3821 (0.3514–0.4126)	0.8462 (0.8390–0.8536)	0.3611 (0.3337–0.3878)	0.4702 (0.4361–0.5024)	0.9031 (0.8931–0.9206)	0.9225 (0.9143–0.9302)	0.9275 (0.9190–0.9349)
			HD95	96.24 (90.27–101.9)	4.753 (4.196–5.447)	98.91 (93.31–104.5)	93.12 (87.54–99.38)	4.329 (2.904–5.500)	5.706 (4.860–6.622)	4.516 (3.821–5.276)
		body of pancreas	Dice	Not supported	0.2426 (0.2356–0.2498)	0.0001518 (0.0000–0.0004402)	0.3031 (0.2938–0.3126)	0.6637 (0.6489–0.6666)	0.6061 (0.5937–0.6190)	0.6361 (0.6236–0.6478)
			HD95		47.47 (46.35–48.66)	112.2 (96.20–128.6)	49.27 (48.05–50.51)	10.81 (9.444–11.64)	18.49 (17.78–19.25)	11.95 (11.36–12.56)
		superior mesenteric artery	Dice	Not supported	0.001502 (0.001262–0.001810)	0.0000 (0.0000–0.0000)	0.3106 (0.2928–0.3284)	0.5539 (0.5474–0.5633)	0.5634 (0.5498–0.5783)	0.6331 (0.6204–0.6454)
			HD95		73.68 (71.21–76.31)	95.09 (77.14–115.3)	26.68 (24.40–28.88)	11.01 (10.02–11.95)	10.63 (9.923–11.48)	7.956 (7.392–8.559)
		gallbladder	Dice	0.2265 (0.2043–0.2490)	0.3669 (0.3469–0.3868)	0.1866 (0.1668–0.2063)	0.3553 (0.3249–0.3846)	0.6950 (0.6900–0.7104)	0.6831 (0.6662–0.6996)	0.7001 (0.6833–0.7178)
			HD95	66.96 (61.75–71.99)	42.18 (38.88–45.51)	60.70 (55.72–65.77)	34.26 (29.12–39.68)	11.05 (9.813–12.97)	12.01 (11.04–13.00)	11.65 (10.45–12.90)
		left adrenal gland	Dice	0.1753 (0.1560–0.1937)	0.3308 (0.3183–0.3433)	0.1263 (0.1092–0.1437)	0.3644 (0.3404–0.3902)	0.6066 (0.6063–0.6217)	0.6579 (0.6451–0.6703)	0.6926 (0.6819–0.7022)
			HD95	32.11 (28.30–36.38)	17.12 (15.75–18.82)	36.33 (32.52–40.04)	34.44 (31.72–37.10)	5.939 (4.865–7.699)	8.514 (7.701–9.484)	6.871 (6.287–7.419)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	PanTS	tail of pancreas	Dice	Not supported	0.1961 (0.1880–0.2048)	0.0002480 (0.0000–0.0007440)	0.2034 (0.1911–0.2166)	0.6605 (0.6466–0.6652)	0.6700 (0.6553–0.6849)	0.6791 (0.6636–0.6935)
			HD95		72.24 (70.34–74.02)	109.8 (98.81–121.0)	66.05 (64.55–67.68)	11.21 (10.87–11.93)	11.46 (10.63–12.32)	11.25 (10.43–12.14)
		left kidney	Dice	0.3374 (0.3110–0.3632)	0.7646 (0.7530–0.7757)	0.3329 (0.3060–0.3588)	0.4582 (0.4281–0.4891)	0.8613 (0.8424–0.8699)	0.9055 (0.8970–0.9136)	0.9140 (0.9052–0.9220)
			HD95	78.55 (73.63–83.39)	44.43 (40.84–48.05)	74.00 (69.42–78.65)	69.12 (64.32–73.76)	5.424 (3.630–5.886)	9.096 (7.847–10.44)	4.878 (4.277–5.549)
		common bile duct	Dice	Not supported	0.007194 (0.006046–0.008456)	0.00002753 (0.0000–0.0001101)	0.07140 (0.06078–0.08253)	0.4212 (0.4127–0.4320)	0.3648 (0.3511–0.3783)	0.4235 (0.4089–0.4370)
			HD95		61.56 (59.49–63.54)	90.41 (83.14–98.09)	19.47 (17.40–21.91)	8.625 (7.538–8.836)	9.908 (9.421–10.37)	9.677 (9.090–10.39)
		celiac artery	Dice	Not supported	0.02258 (0.02044–0.02468)	0.04085 (0.03455–0.04803)	0.2139 (0.2022–0.2247)	0.3640 (0.3432–0.3847)	0.4466 (0.4347–0.4587)	0.4465 (0.4340–0.4593)
			HD95		46.82 (45.10–48.70)	64.97 (59.38–71.26)	50.10 (48.76–51.36)	18.57 (16.65–20.39)	18.01 (17.08–19.08)	18.19 (17.34–19.11)
		pancreatic duct	Dice	Not supported	0.009559 (0.008147–0.01087)	0.0007438 (0.0004090–0.001198)	0.07934 (0.06683–0.09165)	0.08184 (0.08122–0.1003)	0.09234 (0.08333–0.1016)	0.1093 (0.09973–0.1195)
			HD95		67.08 (64.85–69.52)	61.66 (53.77–70.42)	41.54 (38.83–44.37)	32.57 (32.19–34.00)	30.38 (28.63–32.10)	27.79 (26.30–29.43)
		urinary bladder	Dice	0.04774 (0.03465–0.06204)	0.1568 (0.1348–0.1784)	0.02217 (0.01417–0.03083)	0.2718 (0.2394–0.3025)	0.2682 (0.2617–0.2841)	0.2815 (0.2511–0.3122)	0.2795 (0.2489–0.3096)
			HD95	126.1 (117.2–135.4)	98.23 (89.90–106.3)	110.3 (103.0–118.0)	36.91 (27.96–46.14)	73.04 (64.27–80.63)	78.51 (71.14–86.93)	92.78 (85.73–101.0)
		prostate	Dice	0.02414 (0.01472–0.03374)	0.1537 (0.1316–0.1765)	0.002018 (0.00004899–0.005162)	0.09034 (0.06799–0.1115)	0.3782 (0.3703–0.3944)	0.3281 (0.3009–0.3537)	0.3260 (0.3014–0.3497)
			HD95	114.5 (108.0–121.4)	58.18 (52.87–63.82)	105.0 (96.28–114.0)	25.53 (20.33–31.36)	26.05 (25.27–28.63)	29.71 (26.60–32.61)	42.48 (39.67–45.65)
		left femur	Dice	0.09359 (0.06801–0.1187)	0.6168 (0.5817–0.6522)	0.05557 (0.03669–0.07624)	0.3236 (0.2647–0.3829)	0.7649 (0.7563–0.7783)	0.7668 (0.7303–0.8071)	0.7817 (0.7426–0.8198)
			HD95	181.0 (173.2–188.8)	23.78 (17.87–29.95)	161.1 (146.9–173.8)	128.7 (113.0–143.3)	26.00 (18.98–34.56)	48.65 (38.41–60.34)	41.71 (31.96–51.47)
		right femur	Dice	0.1608 (0.1331–0.1888)	0.6020 (0.5690–0.6328)	0.09125 (0.06707–0.1178)	0.3914 (0.3389–0.4481)	0.7943 (0.7604–0.8202)	0.7919 (0.7636–0.8211)	0.8161 (0.7892–0.8424)
			HD95	170.7 (162.0–180.0)	61.41 (50.63–71.63)	150.7 (139.1–163.0)	116.4 (103.1–130.2)	10.74 (9.998–14.10)	16.05 (12.06–20.58)	23.12 (16.90–29.99)
		pancreas lesion	Dice	0.06855 (0.04504–0.09540)	0.1678 (0.1399–0.1958)	0.005854 (0.0003434–0.01550)	0.3028 (0.2503–0.3549)	0.4828 (0.4786–0.4957)	0.3491 (0.3049–0.3945)	0.3888 (0.3437–0.4361)
			HD95	62.26 (50.25–74.92)	66.19 (62.26–70.07)	87.82 (73.00–101.2)	37.21 (29.14–45.85)	26.08 (22.17–29.24)	35.63 (30.88–40.20)	32.46 (27.85–37.71)
		lung tumor	Dice	0.4279 (0.3847–0.4730)	0.01395 (0.009369–0.01918)	0.006545 (0.004327–0.009109)	0.5506 (0.4988–0.5993)	0.7934 (0.7689–0.8216)	0.6637 (0.6261–0.6958)	0.7201 (0.6936–0.7437)
			HD95	101.5 (83.38–126.0)	189.5 (177.4–201.6)	167.1 (159.0–175.3)	96.30 (76.90–121.4)	22.45 (17.12–29.84)	44.44 (36.29–52.47)	37.14 (29.83–44.56)
	PASTA-Gen-30K-Lesions	liver tumor	Dice	0.4881 (0.4394–0.5376)	0.03574 (0.02355–0.04969)	0.02772 (0.01968–0.03687)	0.5305 (0.4719–0.5863)	0.6957 (0.6581–0.7368)	0.6716 (0.6364–0.7065)	0.6913 (0.6596–0.7221)
			HD95	39.94 (33.23–48.10)	117.2 (110.1–124.1)	95.29 (90.06–100.8)	37.48 (29.42–45.84)	26.89 (19.73–36.41)	22.73 (18.93–26.96)	26.73 (20.62–33.56)
		gallbladder cancer	Dice	Not supported	0.4430 (0.3823–0.5005)	0.0004982 (0.0000–0.001441)	0.1540 (0.1148–0.1985)	0.8062 (0.7848–0.8385)	0.7456 (0.7043–0.7881)	0.7763 (0.7367–0.8117)
			HD95		45.06 (34.06–57.59)	131.0 (102.2–157.0)	35.93 (31.44–41.89)	16.27 (12.44–22.91)	7.748 (5.708–10.40)	12.45 (7.990–17.78)
		pancreas tumor	Dice	0.01734 (0.002756–0.03950)	0.3699 (0.3314–0.4104)	0.003484 (0.0008306–0.007482)	0.01990 (0.006567–0.03696)	0.6022 (0.5486–0.6739)	0.3708 (0.3132–0.4311)	0.3767 (0.3221–0.4324)
			HD95	146.4 (121.9–168.9)	30.77 (24.47–37.25)	100.1 (90.48–110.4)	74.93 (60.82–90.05)	26.52 (18.81–38.27)	31.13 (26.32–36.00)	28.02 (22.70–33.80)
		esophageal cancer	Dice	Not supported	0.006366 (0.0007561–0.01462)	0.2960 (0.2506–0.3421)	0.09959 (0.07436–0.1272)	0.8244 (0.8095–0.8478)	0.8083 (0.7784–0.8332)	0.8506 (0.8377–0.8615)
			HD95		150.9 (141.8–159.8)	13.78 (11.09–17.00)	45.59 (37.20–55.10)	17.58 (13.27–24.66)	54.55 (42.78–67.33)	31.27 (24.93–38.35)
		gastric cancer	Dice	Not supported	0.2462 (0.1820–0.3161)	0.0000 (0.0000–0.0000)	0.08088 (0.07075–0.09160)	0.6613 (0.6154–0.7218)	0.7031 (0.6769–0.7264)	0.7764 (0.7565–0.7939)
			HD95		86.61 (76.22–98.79)	270.3 (263.2–276.9)	86.32 (80.36–91.68)	22.40 (15.92–31.77)	19.13 (13.11–26.75)	53.48 (47.46–60.57)
		colorectal cancer	Dice	0.1073 (0.07417–0.1438)	0.002069 (0.0009012–0.003467)	0.001519 (0.0004687–0.002943)	0.0001242 (0.0000–0.0003301)	0.6411 (0.5862–0.7084)	0.6195 (0.5787–0.6594)	0.6655 (0.6316–0.6985)
			HD95	88.28 (74.84–102.0)	143.7 (135.6–152.3)	190.3 (180.7–200.1)	134.1 (120.4–149.3)	37.01 (26.35–51.92)	34.62 (25.25–44.54)	35.49 (26.28–46.02)
		kidney tumor	Dice	Not supported	0.3110 (0.2776–0.3425)	0.2623 (0.2332–0.2889)	0.5417 (0.4866–0.6001)	0.7915 (0.7658–0.8197)	0.8265 (0.7977–0.8499)	0.8641 (0.8349–0.8853)
			HD95		89.77 (80.50–98.65)	67.42 (60.52–74.62)	50.37 (41.08–59.57)	19.60 (14.21–27.83)	14.91 (10.34–20.13)	15.36 (10.48–21.63)
		bladder cancer	Dice	Not supported	0.4614 (0.3896–0.5333)	0.00003449 (0.000004922–0.00007862)	0.07388 (0.06005–0.08933)	0.7379 (0.7015–0.7814)	0.7979 (0.7626–0.8288)	0.8236 (0.7932–0.8480)
			HD95		68.22 (55.58–81.90)	105.2 (95.22–115.3)	62.66 (59.45–66.08)	14.40 (10.82–20.47)	11.99 (8.791–15.45)	16.04 (10.12–23.14)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	PASTA-Gen-30K-Lesions	bone metastasis	Dice	0.01197 (0.008088–0.01644)	0.4279 (0.3890–0.4665)	0.01553 (0.0000001250–0.03821)	0.0000 (0.0000–0.0000)	0.7108 (0.6684–0.7693)	0.2995 (0.2623–0.3387)	0.3649 (0.3278–0.4015)
			HD95	131.9 (124.1–140.1)	103.4 (92.78–113.9)	168.0 (71.42–262.8)	215.0 (187.1–241.7)	34.27 (24.11–48.62)	91.15 (83.77–98.51)	98.35 (90.91–105.1)
		liver cyst	Dice	Not supported	0.006115 (0.003762–0.009278)	0.0003101 (0.00001329–0.0009211)	0.01456 (0.005277–0.02461)	0.6381 (0.5794–0.7098)	0.8138 (0.7956–0.8307)	0.8255 (0.8101–0.8392)
			HD95		120.3 (113.5–128.4)	176.3 (167.8–184.7)	119.5 (109.7–130.0)	25.03 (17.66–36.84)	14.85 (11.42–18.52)	13.27 (9.722–16.73)
		gallstone	Dice	Not supported	0.4721 (0.4139–0.5271)	0.4089 (0.3366–0.4879)	0.0000 (0.0000–0.0000)	0.6763 (0.6325–0.7411)	0.7688 (0.7371–0.7984)	0.7761 (0.7336–0.8113)
			HD95		14.78 (10.93–20.20)	17.94 (10.05–27.94)	30.37 (24.47–36.59)	18.83 (13.42–27.56)	8.742 (4.750–13.41)	4.607 (2.787–6.945)
		pancreatic cyst	Dice	Not supported	0.5222 (0.4887–0.5526)	0.005648 (0.002598–0.009657)	0.2784 (0.2087–0.3438)	0.7108 (0.6659–0.7724)	0.7639 (0.7166–0.8089)	0.8040 (0.7586–0.8439)
			HD95		16.05 (13.56–18.98)	95.03 (86.87–103.6)	29.16 (17.67–42.84)	18.92 (13.11–28.73)	17.73 (12.08–23.86)	9.400 (4.893–15.48)
		kidney cyst	Dice	0.3112 (0.2537–0.3720)	0.05148 (0.03418–0.07201)	0.07921 (0.06116–0.09887)	0.0000 (0.0000–0.0000)	0.7190 (0.6718–0.7796)	0.8236 (0.7919–0.8527)	0.8643 (0.8420–0.8847)
			HD95	90.18 (78.70–102.9)	98.92 (89.19–108.4)	73.51 (67.42–79.96)	– (– –)	25.64 (17.95–37.42)	26.68 (17.73–36.61)	20.05 (13.54–27.38)
		kidney stone	Dice	Not supported	0.1707 (0.1419–0.2009)	0.01014 (0.005443–0.01647)	0.0000 (0.0000–0.0000)	0.7153 (0.6691–0.7818)	0.5778 (0.5225–0.6277)	0.5190 (0.4672–0.5714)
			HD95		84.94 (72.41–97.52)	106.4 (96.56–116.0)	– (– –)	28.40 (19.86–41.57)	49.32 (38.38–60.23)	57.87 (46.78–70.08)
	AbdomenAtlas 3.0	left adrenal gland	Dice	0.7650 (0.7571–0.7730)	0.2788 (0.2680–0.2894)	0.7483 (0.7393–0.7555)	0.7462 (0.7368–0.7550)	0.7993 (0.7761–0.8004)	0.8295 (0.8233–0.8349)	0.8267 (0.8212–0.8318)
			HD95	4.087 (3.670–4.587)	21.72 (20.41–23.06)	4.808 (3.624–6.391)	5.076 (4.558–5.606)	6.599 (6.083–8.839)	3.642 (3.137–4.235)	4.604 (3.963–5.334)
		right adrenal gland	Dice	0.7105 (0.7035–0.7166)	0.3388 (0.3294–0.3486)	0.6825 (0.6746–0.6907)	0.6876 (0.6782–0.6957)	0.7390 (0.7271–0.7430)	0.8195 (0.8135–0.8254)	0.8124 (0.8069–0.8180)
			HD95	4.278 (3.856–4.798)	17.89 (16.54–19.43)	3.779 (3.382–4.210)	4.883 (4.434–5.440)	3.219 (2.501–3.661)	4.699 (3.786–5.929)	3.804 (3.287–4.341)
		aorta	Dice	0.8651 (0.8589–0.8713)	0.7653 (0.7587–0.7726)	0.8516 (0.8449–0.8580)	0.8823 (0.8768–0.8876)	0.8718 (0.8620–0.8775)	0.9017 (0.8961–0.9070)	0.9015 (0.8960–0.9067)
			HD95	29.72 (27.84–31.94)	13.62 (12.20–15.17)	34.58 (31.94–37.23)	24.86 (22.74–27.03)	15.05 (13.92–16.54)	13.19 (11.60–14.79)	13.35 (11.82–14.96)
		urinary bladder	Dice	0.7610 (0.7341–0.7882)	0.3943 (0.3735–0.4181)	0.6965 (0.6685–0.7243)	0.6662 (0.6391–0.6928)	0.7558 (0.7239–0.7862)	0.6983 (0.6707–0.7253)	0.7063 (0.6777–0.7337)
			HD95	16.54 (13.33–20.07)	86.65 (78.95–93.84)	14.47 (10.61–18.84)	25.47 (20.17–31.23)	17.17 (11.41–19.18)	37.18 (30.38–44.05)	51.67 (44.15–60.22)
		celiac artery	Dice	Not supported	0.02728 (0.02520–0.02949)	0.2691 (0.2611–0.2769)	0.2524 (0.2424–0.2619)	0.5306 (0.5146–0.5448)	0.5901 (0.5807–0.5989)	0.6144 (0.6045–0.6234)
			HD95		51.61 (49.89–53.24)	52.48 (51.06–54.13)	52.43 (51.53–53.43)	15.57 (12.99–15.69)	14.50 (13.30–15.80)	12.27 (11.32–13.28)
		celiac trunk	Dice	Not supported	0.003588 (0.002881–0.004415)	0.5584 (0.5413–0.5736)	0.5017 (0.4830–0.5200)	0.5682 (0.5319–0.5734)	0.6290 (0.6119–0.6460)	0.6711 (0.6540–0.6875)
			HD95		77.53 (74.52–80.92)	19.38 (15.76–23.01)	20.76 (17.92–23.65)	12.63 (10.89–16.54)	23.91 (20.38–27.48)	14.30 (11.55–17.32)
		colon	Dice	0.8091 (0.8013–0.8166)	0.2006 (0.1870–0.2139)	0.7939 (0.7858–0.8014)	0.6895 (0.6765–0.7019)	0.8666 (0.8558–0.8764)	0.8660 (0.8603–0.8716)	0.8778 (0.8721–0.8834)
			HD95	40.78 (37.68–44.58)	87.81 (84.22–91.37)	26.99 (25.22–28.74)	63.05 (58.16–68.24)	30.13 (27.88–33.51)	18.07 (16.24–20.34)	19.95 (17.01–23.28)
		common bile duct	Dice	Not supported	0.009287 (0.007782–0.01110)	0.0000 (0.0000–0.0000)	0.1886 (0.1734–0.2035)	0.5641 (0.5492–0.5682)	0.5712 (0.5594–0.5824)	0.6131 (0.6015–0.6246)
			HD95		77.27 (75.55–79.26)	135.2 (117.3–157.0)	19.30 (18.11–20.76)	9.119 (8.284–9.133)	9.101 (8.674–9.525)	8.856 (7.804–10.09)
		duodenum	Dice	0.7783 (0.7719–0.7847)	0.4695 (0.4601–0.4785)	0.7657 (0.7576–0.7729)	0.7390 (0.7290–0.7482)	0.8393 (0.8312–0.8533)	0.8258 (0.8192–0.8322)	0.8461 (0.8400–0.8515)
			HD95	13.76 (12.52–15.42)	22.02 (20.89–23.35)	10.76 (9.866–11.74)	16.45 (15.54–17.27)	8.450 (8.357–8.499)	10.82 (9.553–12.30)	8.857 (7.713–10.47)
		esophagus	Dice	0.7541 (0.7454–0.7624)	0.4656 (0.4535–0.4767)	0.7561 (0.7468–0.7645)	0.7425 (0.7327–0.7515)	0.7543 (0.7499–0.7647)	0.8015 (0.7936–0.8094)	0.7946 (0.7868–0.8026)
			HD95	35.49 (31.49–39.72)	20.98 (19.36–22.74)	17.22 (15.08–19.16)	16.31 (14.68–18.02)	15.28 (12.55–17.77)	17.16 (15.30–19.12)	17.19 (15.31–19.20)
		left femur	Dice	0.6586 (0.6237–0.6938)	0.4990 (0.4716–0.5238)	0.6362 (0.6021–0.6696)	0.6750 (0.6425–0.7063)	0.8216 (0.7872–0.8446)	0.7515 (0.7242–0.7750)	0.7624 (0.7383–0.7864)
			HD95	34.07 (33.27–34.86)	82.03 (74.45–90.22)	154.0 (138.8–169.4)	72.90 (63.18–81.70)	49.50 (43.22–58.01)	117.7 (107.5–128.6)	107.4 (98.92–115.6)
		right femur	Dice	0.6249 (0.5891–0.6603)	0.4810 (0.4567–0.5055)	0.6058 (0.5707–0.6368)	0.6638 (0.6278–0.6966)	0.7936 (0.7850–0.8074)	0.7210 (0.6921–0.7473)	0.7384 (0.7078–0.7665)
			HD95	229.5 (211.1–247.8)	127.6 (119.1–135.6)	175.7 (159.1–194.7)	62.44 (53.87–71.29)	45.53 (39.63–52.37)	113.0 (103.1–122.8)	85.94 (77.80–94.24)
		gallbladder	Dice	0.7897 (0.7716–0.8078)	0.4004 (0.3830–0.4195)	0.7723 (0.7546–0.7894)	0.7636 (0.7430–0.7814)	0.8220 (0.8051–0.8239)	0.8358 (0.8212–0.8498)	0.8347 (0.8192–0.8486)
			HD95	10.98 (9.099–13.27)	37.03 (34.13–40.14)	7.786 (6.727–8.931)	11.61 (10.18–13.20)	7.270 (6.360–7.973)	7.792 (6.896–8.740)	9.688 (7.594–12.14)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	AbdomenAtlas 3.0	hepatic vein	Dice	0.4706 (0.4561–0.4864)	0.06012 (0.05445–0.06569)	0.4884 (0.4726–0.5028)	0.2513 (0.2407–0.2617)	0.4980 (0.4807–0.5089)	0.5590 (0.5449–0.5721)	0.5906 (0.5765–0.6037)
			HD95	30.83 (29.36–32.34)	62.55 (60.43–64.79)	28.35 (26.35–30.74)	55.53 (54.00–57.09)	29.01 (26.59–32.53)	26.03 (24.53–27.86)	22.64 (21.22–24.23)
		intestine	Dice	Not supported	0.2127 (0.1994–0.2272)	0.8005 (0.7935–0.8077)	0.6354 (0.6270–0.6444)	0.8575 (0.8558–0.8868)	0.8630 (0.8573–0.8678)	0.8664 (0.8610–0.8712)
			HD95		81.55 (78.74–84.60)	13.93 (12.97–15.11)	55.41 (53.01–57.93)	13.52 (11.08–15.26)	11.48 (10.61–12.40)	11.35 (10.46–12.37)
		left kidney	Dice	0.9243 (0.9173–0.9308)	0.7681 (0.7605–0.7762)	0.9164 (0.9094–0.9231)	0.9316 (0.9253–0.9380)	0.9190 (0.8993–0.9303)	0.9446 (0.9378–0.9507)	0.9432 (0.9358–0.9495)
			HD95	6.562 (5.666–7.738)	44.13 (40.48–47.37)	6.370 (5.282–7.583)	5.649 (4.906–6.495)	6.265 (5.196–7.706)	4.750 (3.807–5.770)	4.517 (3.747–5.320)
		kidney lesion	Dice	Not supported	0.2352 (0.1965–0.2734)	0.00000001668 (0.0000–0.00000005005)	0.3382 (0.2926–0.3862)	0.6902 (0.6877–0.6956)	0.5990 (0.5575–0.6357)	0.6667 (0.6323–0.6999)
			HD95		103.6 (94.81–112.5)	282.4 (242.9–316.0)	100.7 (86.55–116.1)	30.88 (24.86–41.08)	39.23 (31.93–46.49)	39.48 (33.32–46.10)
		right kidney	Dice	0.9269 (0.9205–0.9328)	0.7719 (0.7640–0.7794)	0.9241 (0.9169–0.9304)	0.9299 (0.9227–0.9366)	0.9158 (0.9075–0.9211)	0.9547 (0.9501–0.9585)	0.9559 (0.9513–0.9599)
			HD95	7.516 (6.441–8.757)	48.09 (44.58–51.57)	5.636 (4.764–6.727)	6.810 (5.900–7.838)	6.163 (4.900–7.583)	4.418 (3.370–5.707)	3.300 (2.817–3.830)
		liver	Dice	0.9699 (0.9680–0.9718)	0.8725 (0.8689–0.8762)	0.9666 (0.9651–0.9681)	0.9691 (0.9673–0.9707)	0.9621 (0.9609–0.9714)	0.9736 (0.9719–0.9750)	0.9724 (0.9708–0.9738)
			HD95	18.24 (14.07–22.60)	11.79 (11.34–12.30)	3.915 (3.525–4.366)	8.304 (6.995–9.789)	8.300 (5.619–9.131)	4.308 (3.831–4.797)	4.297 (3.806–4.819)
		liver lesion	Dice	0.5379 (0.4548–0.6162)	0.1881 (0.1427–0.2382)	0.3760 (0.3319–0.4223)	0.5086 (0.4621–0.5528)	0.6286 (0.5803–0.6713)	0.6848 (0.6557–0.7116)	0.6800 (0.6482–0.7099)
			HD95	75.95 (47.65–109.8)	95.82 (86.04–104.9)	52.29 (39.01–66.55)	61.50 (51.60–72.13)	32.20 (27.03–40.36)	26.20 (21.70–31.36)	42.32 (36.23–48.52)
		caudate lobe	Dice	Not supported	0.02273 (0.01939–0.02635)	0.6012 (0.5887–0.6144)	0.4472 (0.4367–0.4578)	0.7899 (0.7881–0.7907)	0.8236 (0.8172–0.8297)	0.8207 (0.8141–0.8273)
			HD95		154.4 (148.8–159.9)	124.9 (116.1–134.3)	27.17 (26.50–28.00)	8.128 (7.078–8.463)	7.140 (6.782–7.495)	6.837 (6.534–7.155)
		left lateral superior segment of liver	Dice	Not supported	0.0000 (0.0000–0.0000)	0.7362 (0.7241–0.7468)	0.7302 (0.7188–0.7407)	0.8985 (0.8876–0.9059)	0.9114 (0.9062–0.9157)	0.9055 (0.9003–0.9100)
			HD95		133.2 (131.5–135.1)	29.90 (27.37–32.69)	19.53 (18.64–20.59)	5.728 (4.197–7.049)	6.452 (5.981–6.946)	5.936 (5.548–6.424)
		left lateral inferior segment of liver	Dice	Not supported	0.0000 (0.0000–0.0000)	0.6724 (0.6583–0.6875)	0.6725 (0.6615–0.6824)	0.8418 (0.8291–0.8463)	0.8879 (0.8810–0.8945)	0.8828 (0.8755–0.8894)
			HD95		146.2 (144.1–148.2)	20.01 (17.78–22.68)	26.83 (25.64–28.11)	7.668 (6.665–9.803)	6.121 (5.636–6.742)	6.640 (6.014–7.277)
		left medial segment of liver	Dice	Not supported	0.0001119 (0.0000–0.0003358)	0.4737 (0.4557–0.4914)	0.6394 (0.6259–0.6537)	0.8833 (0.8819–0.8911)	0.8733 (0.8668–0.8795)	0.8844 (0.8777–0.8904)
			HD95		161.3 (159.5–163.2)	42.59 (39.92–45.35)	31.31 (30.11–32.55)	9.313 (8.210–10.93)	11.70 (10.72–12.89)	7.633 (6.970–8.428)
		right anterior inferior segment of liver	Dice	Not supported	0.004746 (0.003353–0.006413)	0.4646 (0.4470–0.4824)	0.6710 (0.6602–0.6820)	0.8840 (0.8708–0.8921)	0.8849 (0.8775–0.8925)	0.8789 (0.8707–0.8863)
			HD95		88.80 (86.67–90.97)	37.44 (34.01–41.35)	26.70 (24.71–28.97)	9.850 (8.409–12.29)	10.29 (8.802–11.96)	11.09 (9.585–12.69)
		right posterior inferior segment of liver	Dice	Not supported	0.01492 (0.01234–0.01768)	0.4959 (0.4810–0.5105)	0.6315 (0.6221–0.6413)	0.8588 (0.8374–0.8664)	0.8733 (0.8654–0.8803)	0.8780 (0.8702–0.8849)
			HD95		78.61 (75.09–82.56)	30.55 (28.03–33.41)	38.66 (35.24–42.40)	20.98 (16.32–23.63)	14.45 (11.59–17.30)	15.29 (12.40–18.61)
		right posterior superior segment of liver	Dice	Not supported	0.006630 (0.005476–0.008073)	0.5426 (0.5263–0.5577)	0.6981 (0.6883–0.7068)	0.8481 (0.8434–0.8601)	0.8980 (0.8917–0.9037)	0.9092 (0.9032–0.9146)
			HD95		106.3 (102.0–111.2)	48.56 (45.00–52.39)	43.86 (39.45–48.60)	22.79 (19.18–27.13)	14.12 (11.16–17.79)	13.67 (10.61–17.47)
		right anterior superior segment of liver	Dice	Not supported	0.0002546 (0.00007141–0.0004958)	0.3987 (0.3795–0.4180)	0.6655 (0.6557–0.6751)	0.8822 (0.8736–0.8944)	0.8838 (0.8779–0.8896)	0.8957 (0.8900–0.9012)
			HD95		117.3 (114.3–120.5)	47.12 (44.97–49.37)	40.47 (37.21–43.93)	17.12 (15.83–20.93)	12.25 (9.878–15.14)	11.67 (9.352–14.37)
		left lung	Dice	0.9262 (0.9173–0.9346)	0.7406 (0.7270–0.7536)	0.9254 (0.9175–0.9327)	0.9231 (0.9139–0.9319)	0.9367 (0.9312–0.9540)	0.9497 (0.9427–0.9558)	0.9507 (0.9438–0.9568)
			HD95	13.33 (11.55–15.38)	30.46 (27.84–33.43)	14.37 (12.30–16.50)	18.41 (15.81–21.23)	14.59 (12.15–15.47)	12.86 (10.94–14.83)	10.49 (8.562–12.35)
		right lung	Dice	0.9281 (0.9195–0.9364)	0.7441 (0.7285–0.7573)	0.9151 (0.9067–0.9221)	0.9316 (0.9217–0.9409)	0.9351 (0.9195–0.9511)	0.9568 (0.9499–0.9632)	0.9578 (0.9514–0.9640)
			HD95	15.14 (13.03–17.34)	34.13 (31.64–36.66)	18.82 (16.56–21.21)	17.56 (15.39–20.10)	13.98 (9.792–15.31)	10.46 (8.985–12.01)	16.70 (13.80–19.76)
		pancreas	Dice	0.8632 (0.8563–0.8700)	0.5804 (0.5705–0.5894)	0.8569 (0.8487–0.8636)	0.8382 (0.8298–0.8460)	0.8500 (0.8498–0.8504)	0.8780 (0.8719–0.8840)	0.8804 (0.8737–0.8865)
			HD95	9.028 (7.748–10.64)	15.12 (14.29–16.09)	6.515 (5.665–7.556)	10.19 (9.023–11.45)	9.791 (8.389–10.31)	6.216 (5.412–7.156)	7.195 (5.962–8.863)
		body of pancreas	Dice	Not supported	0.2245 (0.2183–0.2303)	0.0000 (0.0000–0.0000)	0.3327 (0.3267–0.3382)	0.7523 (0.7443–0.7636)	0.7367 (0.7272–0.7461)	0.7519 (0.7418–0.7610)
			HD95		54.14 (53.36–54.96)	360.5 (331.2–382.1)	59.01 (58.18–59.85)	12.07 (9.701–13.74)	18.02 (17.16–18.90)	12.83 (12.17–13.56)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	AbdomenAtlas 3.0	head of pancreas	Dice	Not supported	0.4275 (0.4175–0.4367)	0.0006126 (0.0001487–0.001251)	0.4152 (0.4052–0.4247)	0.8342 (0.8219–0.8395)	0.8053 (0.7950–0.8138)	0.8352 (0.8247–0.8443)
			HD95		67.64 (66.20–69.01)	257.2 (212.3–302.8)	77.32 (76.08–78.61)	8.377 (6.042–11.32)	13.99 (13.03–15.03)	10.06 (8.082–12.23)
		tail of pancreas	Dice	Not supported	0.1686 (0.1609–0.1765)	0.0000 (0.0000–0.0000)	0.3645 (0.3551–0.3735)	0.8170 (0.8114–0.8202)	0.7965 (0.7857–0.8068)	0.8057 (0.7947–0.8163)
			HD95		78.96 (77.73–80.26)	341.9 (307.8–375.1)	72.06 (71.19–73.02)	11.69 (11.05–12.44)	11.35 (10.35–12.44)	9.428 (8.599–10.44)
		pancreas lesion	Dice	0.1120 (0.05198–0.1821)	0.1324 (0.1039–0.1610)	0.1563 (0.1094–0.2077)	0.3278 (0.2762–0.3808)	0.4751 (0.4361–0.5097)	0.4138 (0.3700–0.4594)	0.4684 (0.4245–0.5148)
			HD95	147.8 (110.6–184.3)	79.15 (71.15–89.01)	56.93 (34.52–82.53)	46.23 (35.45–57.72)	35.23 (27.13–43.13)	36.19 (31.16–41.88)	34.26 (29.28–39.53)
		portal vein and splenic vein	Dice	0.5962 (0.5831–0.6090)	0.2105 (0.2024–0.2189)	0.5868 (0.5738–0.5995)	0.5693 (0.5570–0.5822)	0.6163 (0.5958–0.6431)	0.6534 (0.6413–0.6665)	0.6635 (0.6509–0.6767)
			HD95	41.44 (37.95–45.04)	51.12 (48.52–54.03)	27.58 (24.90–30.31)	53.61 (49.93–57.30)	29.22 (26.14–33.21)	42.25 (38.95–45.64)	43.56 (39.97–47.06)
		inferior vena cava	Dice	0.8189 (0.8091–0.8285)	0.6919 (0.6825–0.7010)	0.8221 (0.8121–0.8316)	0.8153 (0.8059–0.8253)	0.8035 (0.7892–0.8158)	0.8345 (0.8254–0.8429)	0.8371 (0.8285–0.8451)
			HD95	25.72 (23.41–27.99)	23.05 (21.05–25.14)	21.93 (19.56–24.42)	23.56 (21.17–26.27)	22.08 (19.59–23.71)	21.63 (19.60–23.97)	21.17 (19.20–23.43)
		prostate	Dice	0.4815 (0.4529–0.5108)	0.2388 (0.2200–0.2583)	0.3241 (0.2971–0.3530)	0.2307 (0.2082–0.2537)	0.6270 (0.5852–0.6459)	0.5800 (0.5544–0.6069)	0.5470 (0.5244–0.5707)
			HD95	155.9 (144.2–167.9)	68.28 (61.72–75.28)	34.55 (31.58–37.62)	38.64 (34.77–43.06)	39.24 (33.81–43.73)	37.00 (31.16–44.16)	74.84 (69.05–80.70)
		rectum	Dice	Not supported	0.02811 (0.02303–0.03378)	0.5733 (0.5495–0.5960)	0.5293 (0.5069–0.5519)	0.7238 (0.7093–0.7362)	0.6933 (0.6740–0.7121)	0.7071 (0.6895–0.7247)
			HD95		75.69 (70.43–80.91)	29.15 (24.97–35.03)	27.81 (25.72–29.99)	13.64 (12.63–15.31)	46.08 (36.21–57.06)	30.73 (24.22–37.58)
		spleen	Dice	0.9445 (0.9374–0.9516)	0.7915 (0.7843–0.7984)	0.9425 (0.9367–0.9479)	0.9423 (0.9353–0.9482)	0.9225 (0.9182–0.9380)	0.9563 (0.9507–0.9609)	0.9560 (0.9506–0.9607)
			HD95	13.08 (9.766–16.63)	11.25 (10.49–12.10)	10.15 (7.131–13.69)	8.321 (7.027–9.793)	5.294 (3.950–6.147)	2.727 (2.220–3.352)	2.726 (2.366–3.109)
		stomach	Dice	0.8853 (0.8756–0.8954)	0.6867 (0.6758–0.6981)	0.7114 (0.6891–0.7338)	0.8495 (0.8384–0.8604)	0.8811 (0.8744–0.8905)	0.9018 (0.8934–0.9101)	0.8980 (0.8894–0.9062)
			HD95	15.92 (14.09–17.80)	18.40 (17.28–19.72)	89.39 (80.73–98.49)	21.06 (19.22–23.08)	13.06 (11.54–15.51)	12.16 (10.80–13.51)	13.65 (12.10–15.37)
		superior mesenteric artery	Dice	Not supported	0.001816 (0.001635–0.002024)	0.001417 (0.0004707–0.002726)	0.4793 (0.4660–0.4938)	0.7078 (0.7009–0.7167)	0.7758 (0.7682–0.7829)	0.8231 (0.8185–0.8277)
			HD95		121.8 (118.9–124.9)	102.1 (95.29–109.1)	34.19 (32.06–36.41)	9.399 (8.015–12.35)	7.994 (7.363–8.600)	5.130 (4.742–5.562)
		veins	Dice	Not supported	0.004697 (0.003654–0.005945)	0.1213 (0.1039–0.1426)	0.2205 (0.2075–0.2328)	0.7868 (0.7805–0.7894)	0.8061 (0.7992–0.8128)	0.8175 (0.8109–0.8238)
			HD95		78.25 (77.35–79.14)	85.69 (79.34–93.03)	67.01 (66.06–67.49)	12.00 (8.766–13.33)	9.079 (8.488–9.803)	7.477 (6.890–8.078)
	LiMT	colon lesion	Dice	Not supported	0.03393 (0.0001163–0.08354)	0.0000 (0.0000–0.0000)	0.6030 (0.4972–0.7061)	0.6854 (0.6839–0.6893)	0.6212 (0.5086–0.7165)	0.6518 (0.5500–0.7353)
			HD95		186.9 (157.3–217.6)	– (––)	70.07 (39.90–104.8)	34.54 (11.62–66.17)	28.56 (14.76–46.14)	41.25 (17.06–72.64)
		liver	Dice	0.8720 (0.8141–0.9192)	0.08766 (0.002237–0.2003)	0.7628 (0.6613–0.8643)	0.6592 (0.5397–0.7613)	0.9381 (0.9266–0.9466)	0.9429 (0.9254–0.9578)	0.9495 (0.9384–0.9592)
			HD95	13.45 (9.159–18.01)	133.9 (108.8–157.0)	19.03 (12.44–26.55)	102.9 (69.36–137.4)	8.283 (6.750–11.90)	7.824 (4.504–12.57)	6.202 (4.424–8.453)
		liver cyst	Dice	Not supported	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.6831 (0.6823–0.7010)	0.5513 (0.3939–0.6726)	0.5525 (0.3805–0.6922)
			HD95		– (––)	400.7 (385.3–416.2)	– (––)	13.29 (6.723–23.73)	26.29 (6.734–49.53)	37.64 (23.66–53.32)
		liver hemangioma	Dice	Not supported	0.0000 (0.0000–0.0000)	0.3428 (0.1047–0.5815)	0.2417 (0.0000–0.5089)	0.7385 (0.7292–0.7385)	0.7789 (0.7022–0.8441)	0.7105 (0.5357–0.8288)
			HD95		– (––)	87.16 (49.97–125.0)	71.59 (5.750–153.7)	38.57 (14.58–68.21)	19.56 (8.792–31.53)	40.47 (11.82–69.13)
		metastatic liver cancer	Dice	0.3684 (0.09556–0.6286)	0.0000 (0.0000–0.0000)	0.02375 (0.0000–0.06959)	0.1174 (0.0000–0.3203)	0.4385 (0.4141–0.4834)	0.3798 (0.1154–0.5868)	0.4520 (0.1643–0.6442)
			HD95	50.23 (26.19–76.95)	177.9 (177.9–177.9)	136.7 (94.46–177.7)	83.57 (46.14–126.0)	50.87 (28.55–75.34)	40.42 (19.15–74.88)	40.53 (18.05–75.66)
		primary liver cancer	Dice	0.4323 (0.1632–0.6778)	0.000001252 (0.0000–0.000003756)	0.3477 (0.1408–0.5614)	0.3747 (0.1390–0.6476)	0.7234 (0.6444–0.7951)	0.7394 (0.6181–0.8488)	0.6927 (0.4554–0.8568)
			HD95	62.16 (36.32–87.10)	168.5 (141.0–212.4)	46.39 (25.89–69.49)	43.99 (16.66–79.89)	22.60 (11.26–35.78)	40.33 (11.73–75.55)	31.25 (15.05–54.17)
	ULS23-KiTS23	kidney lesion	Dice	Not supported	0.3935 (0.3192–0.4668)	0.0001014 (0.0000–0.0003043)	0.7737 (0.7098–0.8330)	0.7771 (0.7251–0.7936)	0.7338 (0.6647–0.7955)	0.7705 (0.7142–0.8215)
			HD95		49.15 (38.58–60.27)	111.4 (97.94–127.8)	12.70 (8.181–17.29)	15.65 (9.966–20.31)	21.57 (13.32–32.14)	15.11 (10.34–20.05)
	ULS23-LiTS	liver tumor	Dice	0.04586 (0.01075–0.08097)	0.04802 (0.02934–0.07236)	0.1157 (0.08589–0.1483)	0.1277 (0.09877–0.1616)	0.6518 (0.6444–0.6867)	0.3608 (0.3183–0.4032)	0.3647 (0.3243–0.4019)
			HD95	111.3 (106.4–116.3)	87.46 (83.62–91.14)	70.78 (66.76–74.79)	85.12 (80.88–89.30)	10.10 (9.703–12.23)	34.33 (30.76–38.01)	43.67 (40.11–47.27)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
Internal	ULS23-LIDC-IDRI	pulmonary nodule	Dice	Not supported	0.02544 (0.01489–0.03840)	0.1377 (0.1156–0.1602)	0.3140 (0.2770–0.3518)	0.6392 (0.6039–0.6539)	0.5756 (0.5466–0.6013)	0.6145 (0.5862–0.6386)
			HD95		103.3 (99.73–106.3)	64.86 (60.92–68.74)	44.16 (39.76–48.79)	7.559 (4.777–9.595)	11.96 (9.934–14.08)	11.44 (9.205–13.68)
	ULS23-MSD-lung	lung tumor	Dice	Not supported	0.2841 (0.1097–0.4727)	0.5587 (0.3853–0.7002)	0.7853 (0.6504–0.8695)	0.7826 (0.7051–0.8457)	0.7133 (0.5900–0.8122)	0.7365 (0.6088–0.8291)
			HD95		61.46 (36.31–89.15)	20.63 (6.943–38.43)	9.120 (2.656–20.83)	5.581 (3.685–6.837)	19.10 (3.972–39.28)	15.96 (4.259–36.35)
	ULS23-MSD-colon	colon cancer	Dice	Not supported	0.02641 (0.0003254–0.08548)	0.6397 (0.5143–0.7500)	0.7137 (0.6537–0.7674)	0.7484 (0.7180–0.7704)	0.6619 (0.5737–0.7377)	0.6531 (0.5453–0.7330)
			HD95		130.8 (105.4–156.7)	16.26 (4.970–31.90)	11.09 (7.324–15.67)	11.76 (7.867–15.60)	12.15 (8.331–17.15)	20.72 (10.12–33.79)
	ULS23-MSD-pancreas	pancreas tumor	Dice	Not supported	0.2248 (0.1756–0.2756)	0.7307 (0.6583–0.7969)	0.6651 (0.5948–0.7287)	0.7768 (0.7636–0.7950)	0.6121 (0.5635–0.6602)	0.6432 (0.5875–0.6960)
			HD95		40.33 (34.17–47.17)	4.266 (3.455–5.159)	14.25 (8.106–21.50)	7.176 (7.052–7.233)	16.54 (11.97–21.90)	19.27 (13.61–25.63)
	LiMT-oneclass	liver	Dice	0.8720 (0.8037–0.9206)	0.08766 (0.002036–0.1985)	0.7628 (0.6507–0.8616)	0.6592 (0.5415–0.7704)	0.9266 (0.9169–0.9400)	0.9429 (0.9245–0.9575)	0.9495 (0.9379–0.9596)
			HD95	13.45 (9.350–17.99)	133.9 (111.2–156.2)	19.03 (12.33–27.20)	102.9 (69.03–139.5)	8.071 (5.867–13.19)	7.824 (4.552–12.56)	6.202 (4.428–8.336)
		liver lesion	Dice	0.07319 (0.006662–0.1552)	0.0000 (0.0000–0.0000)	0.03527 (0.0000–0.1054)	0.04505 (0.01061–0.09095)	0.3532 (0.3108–0.3904)	0.3203 (0.1757–0.4760)	0.2896 (0.1500–0.4240)
			HD95	94.28 (58.91–148.4)	172.4 (129.4–210.7)	160.6 (84.52–252.0)	117.9 (94.67–141.2)	22.56 (9.291–38.33)	25.61 (9.964–42.83)	41.04 (24.79–58.20)
External-seen (zero-shot)	ULS23-NIG-LN-MED	mediastinal lymph node	Dice	Not supported	0.02758 (0.003452–0.05970)	0.0000 (0.0000–0.0000)	0.1791 (0.1326–0.2312)	0.05950 (0.05100–0.06620)	0.2901 (0.2012–0.3765)	0.3488 (0.2669–0.4303)
			HD95		63.52 (56.97–70.40)	– (––)	70.65 (62.13–78.95)	79.22 (65.03–97.30)	36.90 (28.71–46.15)	37.37 (26.98–48.03)
	ULS23-Radboudumc-Pancreas	pancreas lesion	Dice	Not supported	0.1669 (0.09012–0.2573)	0.05259 (0.0004245–0.1337)	0.4539 (0.2718–0.6191)	0.09135 (0.05473–0.1321)	0.5139 (0.4165–0.5923)	0.6181 (0.5015–0.7177)
			HD95		55.56 (41.74–67.40)	49.52 (16.31–85.55)	17.75 (9.539–26.41)	58.02 (45.67–83.12)	16.92 (11.31–23.42)	24.09 (14.17–34.74)
	CT-Lymph-Nodes	mediastinal lymph node	Dice	Not supported	0.05771 (0.01808–0.1121)	0.0000 (0.0000–0.0000)	0.1986 (0.06696–0.3376)	0.06686 (0.05627–0.1174)	0.3697 (0.1488–0.5788)	0.3884 (0.1903–0.5772)
			HD95		54.71 (46.11–64.05)	– (––)	125.6 (87.19–184.3)	55.36 (30.69–86.93)	61.99 (32.46–95.41)	40.74 (22.61–59.04)
		left adrenal gland	Dice	0.3328 (0.2779–0.3837)	0.1581 (0.1303–0.1876)	0.4150 (0.3695–0.4605)	0.4033 (0.3490–0.4580)	0.1807 (0.04050–0.3250)	0.4617 (0.4182–0.5073)	0.5014 (0.4577–0.5423)
			HD95	16.67 (14.52–18.93)	20.35 (18.46–22.45)	14.27 (12.13–16.51)	18.96 (16.24–21.76)	37.36 (23.33–56.47)	14.43 (12.08–17.26)	12.42 (10.44–15.09)
		right adrenal gland	Dice	0.4021 (0.3543–0.4468)	0.1484 (0.1218–0.1727)	0.4416 (0.3947–0.4879)	0.4707 (0.4147–0.5235)	0.1658 (0.02130–0.3196)	0.4782 (0.4331–0.5189)	0.4828 (0.4391–0.5237)
			HD95	11.11 (9.458–13.07)	19.89 (17.32–22.68)	8.961 (7.374–11.13)	13.88 (11.12–16.78)	36.48 (22.42–56.55)	8.851 (7.061–10.96)	9.356 (7.295–11.70)
		urinary bladder	Dice	0.7850 (0.7274–0.8385)	0.5710 (0.5129–0.6316)	0.7739 (0.7079–0.8301)	0.8063 (0.7571–0.8493)	0.5555 (0.4462–0.6745)	0.8540 (0.8070–0.8931)	0.8585 (0.8122–0.8968)
			HD95	16.89 (8.583–27.64)	24.07 (17.71–32.67)	18.26 (9.680–29.43)	14.15 (10.15–19.43)	28.02 (17.23–43.70)	7.773 (5.396–10.75)	13.09 (7.888–22.13)
		bone	Dice	Not supported	0.5998 (0.5799–0.6175)	0.1139 (0.05941–0.1722)	0.9359 (0.9306–0.9411)	0.6198 (0.5293–0.7197)	0.9031 (0.8973–0.9089)	0.9131 (0.9083–0.9177)
			HD95		36.86 (31.22–42.81)	163.7 (147.7–181.5)	4.560 (2.063–7.881)	16.87 (9.919–26.87)	15.49 (9.998–22.17)	18.17 (11.16–26.16)
	Pediatric-CT-SEG	left breast	Dice	Not supported	0.00008134 (0.0000–0.0002440)	0.3291 (0.1942–0.4713)	0.4196 (0.3188–0.5381)	0.07670 (0.001400–0.1872)	0.3270 (0.2171–0.4452)	0.3445 (0.2232–0.4686)
			HD95		143.4 (118.4–174.1)	19.92 (12.37–29.10)	83.28 (49.66–121.3)	108.3 (77.81–149.4)	26.16 (14.11–40.20)	61.41 (23.78–105.6)
		right breast	Dice	Not supported	0.0001054 (0.0000–0.0003140)	0.4546 (0.3291–0.5887)	0.3916 (0.2732–0.5177)	0.07520 (0.004600–0.1924)	0.4374 (0.3272–0.5378)	0.4916 (0.3671–0.5993)
			HD95		131.0 (119.0–144.6)	14.65 (9.509–20.76)	67.96 (39.73–98.16)	132.7 (92.33–178.9)	65.59 (44.00–94.37)	25.16 (12.23–46.33)
		duodenum	Dice	0.4969 (0.4485–0.5446)	0.3453 (0.3117–0.3772)	0.5253 (0.4778–0.5714)	0.5726 (0.5348–0.6073)	0.2540 (0.1025–0.3893)	0.5661 (0.5231–0.6074)	0.6202 (0.5858–0.6554)
			HD95	16.76 (14.66–18.83)	21.28 (18.76–24.09)	15.47 (13.13–18.41)	20.18 (16.95–24.18)	44.83 (29.71–63.11)	16.15 (14.21–18.36)	12.61 (11.23–14.33)
		esophagus	Dice	0.6488 (0.6183–0.6783)	0.4391 (0.4106–0.4666)	0.6782 (0.6439–0.7079)	0.6707 (0.6411–0.6980)	0.3341 (0.2136–0.4616)	0.6450 (0.6118–0.6759)	0.6817 (0.6536–0.7090)
			HD95	58.76 (39.62–79.61)	10.71 (9.638–11.86)	6.654 (5.687–7.805)	7.941 (6.580–9.616)	30.92 (19.80–47.95)	10.23 (7.951–13.00)	9.630 (7.519–12.10)
		left hip	Dice	0.01970 (0.006673–0.03838)	0.07471 (0.04781–0.1060)	0.003148 (0.001675–0.005009)	0.01868 (0.01431–0.02310)	0.08640 (0.007300–0.2088)	0.4617 (0.3738–0.5424)	0.6304 (0.5653–0.6907)
			HD95	82.75 (77.75–87.59)	88.41 (82.83–94.57)	120.6 (101.3–141.2)	87.30 (78.71–97.08)	166.5 (116.9–215.4)	104.1 (79.58–127.7)	246.9 (206.7–289.4)
		right hip	Dice	0.01821 (0.009888–0.02719)	0.08068 (0.06607–0.09556)	0.02296 (0.009777–0.04006)	0.01134 (0.007850–0.01556)	0.02440 (0.007300–0.09330)	0.5554 (0.4570–0.6480)	0.5766 (0.5024–0.6476)
			HD95	274.0 (236.5–313.8)	148.6 (131.1–167.5)	110.3 (93.66–130.3)	81.66 (76.29–87.42)	152.4 (106.9–201.8)	63.76 (45.51–85.28)	83.77 (64.61–109.1)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
External-seen (zero-shot)	Pediatric-CT-SEG	gallbladder	Dice	0.8224 (0.7810–0.8557)	0.4515 (0.3976–0.4994)	0.8146 (0.7670–0.8530)	0.8281 (0.7894–0.8613)	0.5037 (0.3970–0.6212)	0.8585 (0.8331–0.8802)	0.8505 (0.8166–0.8827)
			HD95	7.141 (4.600–11.12)	16.76 (14.19–19.48)	4.967 (3.629–6.823)	6.141 (4.496–8.171)	29.85 (18.21–43.38)	4.343 (3.349–5.654)	5.367 (3.891–6.904)
		gonad	Dice	Not supported	0.0002080 (0.000004660–0.0004483)	0.003774 (0.0000–0.01036)	0.00001475 (0.0000–0.00004425)	0.001921 (0.0006837–0.03588)	0.04580 (0.02509–0.06976)	0.04204 (0.01917–0.07182)
			HD95		148.4 (132.1–166.0)	97.08 (33.91–215.2)	67.28 (35.99–92.56)	115.3 (79.87–158.7)	99.22 (82.50–119.8)	145.5 (97.65–200.1)
		heart	Dice	0.8519 (0.8417–0.8620)	0.2640 (0.2335–0.2926)	0.8782 (0.8643–0.8889)	0.5514 (0.5044–0.6005)	0.5544 (0.4506–0.6653)	0.9110 (0.9015–0.9197)	0.8977 (0.8880–0.9072)
			HD95	64.07 (37.07–91.61)	34.06 (31.41–37.30)	26.12 (16.30–38.27)	21.66 (17.31–26.64)	49.24 (32.56–69.05)	13.18 (10.56–16.48)	12.92 (11.18–14.74)
		left kidney	Dice	0.9236 (0.8841–0.9495)	0.7649 (0.7334–0.7896)	0.9318 (0.8941–0.9565)	0.9431 (0.9168–0.9623)	0.6795 (0.6088–0.7591)	0.9573 (0.9442–0.9666)	0.9550 (0.9402–0.9648)
			HD95	4.392 (3.384–5.890)	33.47 (25.15–41.47)	3.944 (2.382–6.439)	5.512 (3.138–9.007)	17.05 (9.360–26.92)	3.527 (2.273–5.153)	4.072 (3.305–4.959)
		right kidney	Dice	0.9259 (0.8916–0.9457)	0.7739 (0.7514–0.7946)	0.9273 (0.8890–0.9520)	0.9315 (0.8995–0.9567)	0.6911 (0.6206–0.7853)	0.9391 (0.9087–0.9622)	0.9439 (0.9200–0.9620)
			HD95	4.932 (3.698–6.638)	24.81 (18.19–31.65)	4.341 (2.803–6.436)	6.816 (4.287–10.70)	16.13 (9.237–25.28)	4.544 (2.954–6.534)	4.275 (2.811–6.262)
		colon	Dice	0.6503 (0.6102–0.6834)	0.3658 (0.3231–0.4103)	0.6670 (0.6254–0.7063)	0.6280 (0.5867–0.6666)	0.4146 (0.3097–0.5308)	0.6978 (0.6618–0.7327)	0.7040 (0.6640–0.7417)
			HD95	31.42 (28.32–34.82)	38.11 (33.84–44.21)	25.12 (21.46–29.73)	43.60 (35.90–52.28)	51.61 (35.68–74.29)	36.64 (28.66–46.83)	34.63 (28.51–40.92)
		liver	Dice	0.9533 (0.9500–0.9564)	0.8538 (0.8464–0.8610)	0.9565 (0.9535–0.9596)	0.9582 (0.9525–0.9624)	0.7122 (0.6480–0.7959)	0.9626 (0.9600–0.9647)	0.9642 (0.9615–0.9665)
			HD95	4.808 (4.185–5.498)	12.65 (11.41–14.38)	3.920 (3.375–4.598)	7.901 (5.509–10.96)	18.71 (10.62–29.93)	4.668 (3.858–5.729)	3.874 (3.343–4.530)
		left lung	Dice	0.9170 (0.8959–0.9339)	0.7449 (0.7066–0.7767)	0.9044 (0.8831–0.9225)	0.9268 (0.9072–0.9438)	0.5874 (0.4965–0.6812)	0.9312 (0.9110–0.9470)	0.9343 (0.9136–0.9512)
			HD95	14.00 (11.02–17.43)	16.99 (13.98–20.56)	27.10 (17.90–37.23)	15.17 (11.25–20.38)	51.17 (34.02–73.85)	17.20 (12.39–22.45)	9.662 (6.744–13.16)
		right lung	Dice	0.9253 (0.9101–0.9382)	0.6514 (0.5884–0.7069)	0.9188 (0.9005–0.9351)	0.9402 (0.9248–0.9541)	0.5866 (0.5003–0.6913)	0.9417 (0.9289–0.9527)	0.9435 (0.9307–0.9545)
			HD95	12.81 (10.28–16.77)	27.81 (22.21–34.07)	16.96 (12.17–22.53)	13.59 (10.04–17.74)	46.58 (30.01–66.73)	16.08 (11.97–20.88)	11.46 (9.065–14.22)
		pancreas	Dice	0.7301 (0.6896–0.7613)	0.4640 (0.4333–0.4936)	0.7338 (0.6957–0.7650)	0.7385 (0.7051–0.7686)	0.5101 (0.4057–0.6291)	0.7732 (0.7506–0.7948)	0.7746 (0.7483–0.7985)
			HD95	9.082 (7.610–10.75)	14.95 (12.80–17.32)	7.807 (6.306–9.509)	13.65 (10.64–17.31)	30.77 (19.86–47.47)	7.570 (6.497–8.769)	8.190 (6.732–9.881)
		prostate	Dice	0.1408 (0.07076–0.2272)	0.06738 (0.02323–0.1193)	0.2763 (0.1836–0.3761)	0.1114 (0.04114–0.1957)	0.05250 (0.002500–0.1658)	0.3505 (0.2618–0.4414)	0.2341 (0.1592–0.3167)
			HD95	147.8 (94.27–203.7)	50.54 (38.96–64.15)	8.190 (6.210–10.56)	22.06 (11.18–39.70)	32.49 (19.55–51.89)	21.13 (15.00–28.83)	118.9 (89.17–149.1)
		rectum	Dice	Not supported	0.08611 (0.05616–0.1171)	0.6553 (0.6072–0.6984)	0.4765 (0.4177–0.5325)	0.007900 (0.005100–0.09110)	0.6589 (0.6098–0.7050)	0.6587 (0.6087–0.7051)
			HD95		39.90 (35.84–44.07)	29.61 (18.71–41.90)	23.35 (19.66–27.23)	122.2 (85.49–167.6)	59.34 (33.38–87.80)	32.70 (20.98–48.44)
		skin	Dice	Not supported	0.00008226 (0.00004011–0.0001389)	0.6575 (0.6405–0.6721)	0.5786 (0.5505–0.6057)	0.1684 (0.01960–0.3218)	0.6201 (0.5986–0.6390)	0.6621 (0.6439–0.6789)
			HD95		78.94 (73.08–85.01)	11.70 (9.560–14.04)	26.80 (23.23–30.48)	46.66 (30.59–65.87)	15.55 (12.63–18.96)	11.71 (9.145–14.63)
		small bowel	Dice	0.6352 (0.5899–0.6782)	0.3886 (0.3400–0.4382)	0.6871 (0.6460–0.7221)	0.6624 (0.6188–0.7013)	0.4384 (0.3298–0.5518)	0.7709 (0.7383–0.7942)	0.7641 (0.7290–0.7914)
			HD95	25.98 (20.84–32.57)	39.92 (32.65–48.25)	19.77 (16.03–26.53)	29.32 (24.19–35.65)	48.19 (30.73–70.83)	20.44 (16.47–26.14)	19.18 (15.49–24.80)
		spinal canal	Dice	Not supported	0.002467 (0.0007330–0.004736)	0.7677 (0.7528–0.7841)	0.8668 (0.8464–0.8816)	0.4504 (0.3487–0.5662)	0.7455 (0.7298–0.7610)	0.8563 (0.8438–0.8671)
			HD95		191.9 (166.0–214.5)	54.81 (51.12–58.85)	3.736 (3.031–4.421)	31.64 (19.39–48.13)	77.40 (72.54–82.94)	37.04 (29.31–44.87)
		spleen	Dice	0.9306 (0.9219–0.9389)	0.7554 (0.7367–0.7749)	0.9260 (0.9126–0.9386)	0.9239 (0.9076–0.9374)	0.6689 (0.5956–0.7633)	0.9390 (0.9290–0.9480)	0.9453 (0.9341–0.9534)
			HD95	4.204 (3.275–5.541)	10.97 (9.858–12.15)	3.972 (2.858–5.481)	11.52 (8.538–14.69)	19.13 (11.64–30.04)	3.791 (2.803–4.837)	3.096 (2.324–4.184)
		stomach	Dice	0.8568 (0.8213–0.8832)	0.5876 (0.5447–0.6264)	0.7226 (0.6561–0.7825)	0.7722 (0.7340–0.8116)	0.6390 (0.5595–0.7466)	0.8794 (0.8513–0.9047)	0.8754 (0.8466–0.9028)
			HD95	9.943 (7.632–12.59)	19.23 (16.97–21.48)	55.23 (36.57–75.27)	24.25 (19.66–29.16)	26.92 (16.75–40.13)	8.683 (6.514–11.06)	10.62 (7.803–13.93)
		thymus	Dice	Not supported	0.007088 (0.001957–0.01311)	0.1952 (0.1233–0.2787)	0.0000 (0.0000–0.0000)	0.1080 (0.007500–0.2358)	0.4862 (0.4123–0.5558)	0.5471 (0.4676–0.6157)
			HD95		130.4 (117.1–143.5)	22.29 (17.19–27.32)	– (– –)	54.13 (35.64–79.24)	16.80 (13.37–20.24)	22.44 (12.98–38.40)
		uterocervix	Dice	Not supported	0.1226 (0.07594–0.1710)	0.009672 (0.001133–0.02196)	0.1036 (0.04398–0.1802)	0.06510 (0.002300–0.1836)	0.3242 (0.2396–0.4005)	0.3626 (0.2763–0.4457)
			HD95		47.65 (40.80–54.78)	156.6 (113.9–194.7)	42.95 (31.55–54.10)	75.33 (51.78–107.1)	41.42 (28.67–57.11)	34.81 (24.69–46.68)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
External-seen (zero-shot)	VEELA2025	hepatic vein	Dice	0.3136 (0.2600–0.3601)	0.2238 (0.1629–0.2842)	0.1086 (0.06328–0.1660)	0.4117 (0.3754–0.4468)	0.1793 (0.1651–0.2041)	0.3050 (0.2590–0.3497)	0.3460 (0.3044–0.3822)
			HD95	27.74 (25.06–30.48)	39.88 (34.06–47.81)	41.34 (33.04–49.92)	21.22 (18.39–24.76)	41.49 (31.19–45.82)	24.66 (22.24–27.06)	26.64 (24.44–28.67)
		portal vein	Dice	Not supported	0.1366 (0.08519–0.1931)	0.1034 (0.06409–0.1489)	0.4604 (0.4388–0.4831)	0.1030 (0.06272–0.1246)	0.4382 (0.4020–0.4752)	0.4995 (0.4658–0.5350)
			HD95		67.87 (56.94–79.32)	69.33 (57.23–80.76)	78.94 (71.22–87.59)	68.92 (62.11–75.02)	32.31 (30.15–34.55)	34.69 (28.50–42.27)
		liver	Dice	0.9693 (0.9665–0.9720)	0.7977 (0.6556–0.9350)	0.9731 (0.9707–0.9750)	0.9717 (0.9682–0.9748)	0.7143 (0.6828–0.7392)	0.9702 (0.9681–0.9726)	0.9728 (0.9700–0.9757)
			HD95	3.713 (3.166–4.416)	5.482 (5.057–5.917)	2.487 (2.276–2.702)	4.327 (3.436–5.328)	15.87 (14.64–17.95)	4.886 (4.166–5.696)	3.291 (2.738–3.940)
		sacrum	Dice	0.9347 (0.9296–0.9393)	0.3551 (0.2878–0.4223)	0.9361 (0.9330–0.9388)	0.8502 (0.7926–0.8976)	0.5583 (0.4634–0.6784)	0.8892 (0.8793–0.8989)	0.9284 (0.9204–0.9353)
			HD95	2.290 (1.712–3.019)	52.06 (38.07–66.36)	1.572 (1.345–1.875)	7.080 (4.229–10.40)	16.76 (9.575–27.19)	5.716 (4.902–6.497)	2.844 (2.285–3.447)
	Pengwin	left hip	Dice	0.9529 (0.9515–0.9542)	0.3445 (0.2760–0.4149)	0.9570 (0.9557–0.9582)	0.8881 (0.8350–0.9299)	0.5793 (0.4899–0.6908)	0.9562 (0.9498–0.9607)	0.9631 (0.9545–0.9687)
			HD95	1.215 (1.175–1.265)	76.26 (61.04–92.17)	1.067 (1.038–1.099)	17.68 (11.13–24.76)	18.21 (9.790–29.56)	38.90 (30.96–47.71)	3.594 (1.534–6.123)
		right hip	Dice	0.9522 (0.9505–0.9539)	0.3648 (0.2951–0.4290)	0.9559 (0.9540–0.9575)	0.8800 (0.8322–0.9234)	0.5812 (0.4980–0.6929)	0.9628 (0.9553–0.9677)	0.9633 (0.9575–0.9680)
			HD95	2.352 (1.236–4.520)	76.50 (61.79–91.34)	1.318 (1.080–1.773)	13.99 (8.368–20.46)	18.10 (10.47–30.63)	5.434 (1.465–10.81)	4.167 (1.105–8.428)
	MHB-SEG-oneclass	intracranial hemorrhage	Dice	Not supported	0.0006353 (0.00002192–0.001675)	0.2819 (0.1972–0.3694)	0.2154 (0.1446–0.2881)	0.1873 (0.09772–0.2877)	0.3219 (0.2393–0.4067)	0.4069 (0.3181–0.4835)
			HD95		87.13 (80.73–93.08)	49.39 (40.36–58.28)	61.83 (53.43–70.32)	79.92 (70.62–89.30)	56.13 (46.17–65.28)	54.52 (45.81–63.56)
	QUBIQ2021	pancreas	Dice	0.2222 (0.05556–0.4444)	0.05491 (0.008992–0.1157)	0.0001425 (0.0000–0.0004275)	0.04201 (0.0000–0.1035)	0.1389 (0.1169–0.3456)	0.7761 (0.7302–0.7931)	0.7841 (0.7405–0.8041)
			HD95	53.47 (0.0000–126.3)	67.65 (51.01–83.90)	116.3 (85.79–147.8)	78.52 (44.38–121.6)	64.11 (40.59–100.1)	8.600 (7.439–9.381)	7.953 (6.330–9.587)
		pancreas lesion	Dice	0.0000 (0.0000–0.0000)	0.06874 (0.01625–0.1300)	0.0000 (0.0000–0.0000)	0.1035 (0.02533–0.1865)	0.03778 (0.01692–0.08354)	0.2608 (0.1629–0.3595)	0.3342 (0.2669–0.4065)
			HD95	169.0 (140.0–199.6)	106.0 (95.29–119.8)	67.62 (46.26–91.31)	83.94 (32.54–140.0)	111.3 (97.42–183.6)	39.97 (32.72–48.42)	28.69 (24.11–33.59)
	TotalSeg-subtask-liver-lesion	liver lesion	Dice	0.5862 (0.5510–0.6225)	0.1780 (0.1482–0.2067)	0.2491 (0.2132–0.2837)	0.5485 (0.5103–0.5844)	0.1338 (0.1127–0.1462)	0.6493 (0.6166–0.6803)	0.6936 (0.6650–0.7204)
			HD95	41.44 (36.53–47.30)	83.83 (78.70–89.34)	45.47 (39.86–51.45)	48.04 (42.68–53.40)	84.91 (81.97–90.66)	30.12 (26.09–34.36)	32.69 (28.54–37.04)
	HCC-TACE-Seg	liver	Dice	0.4004 (0.3189–0.4733)	0.7549 (0.6756–0.8280)	0.5683 (0.5069–0.6276)	0.5929 (0.4990–0.6750)	0.3983 (0.2868–0.4596)	0.9241 (0.9022–0.9420)	0.9166 (0.8905–0.9375)
			HD95	93.19 (80.65–104.8)	25.62 (16.60–37.10)	82.96 (70.19–95.35)	95.30 (82.89–107.6)	87.83 (84.58–111.0)	24.22 (15.64–33.88)	12.84 (10.20–15.45)
	PANORAMA	pancreatic duct	Dice	Not supported	0.01906 (0.01440–0.02426)	0.001345 (0.0002719–0.003120)	0.3549▲ (0.3089–0.3980)	0.07302 (0.06472–0.07823)	0.1910 (0.1607–0.2208)	0.2647 (0.2272–0.2997)
			HD95		62.18 (56.79–67.75)	70.79 (50.44–94.79)	30.57▲ (26.85–34.50)	59.85 (53.11–65.39)	43.84 (40.58–47.13)	34.43 (31.08–38.12)
External-unseen (zero-shot)	ULS23-NIG-LN-ABD	abdominal lymph node	Dice	Not supported	0.02013 (0.006038–0.04199)	0.0000 (0.0000–0.0000)	0.1529 (0.09906–0.2146)	Not supported	0.1170 (0.07211–0.1660)	0.2109 (0.1568–0.2659)
			HD95		82.61 (76.86–88.16)	38.31 (38.31–38.31)	63.36 (58.03–68.74)		43.04 (36.58–49.25)	39.13 (33.36–45.06)
	ULS23-Radboudumc-Bone	bone lesion	Dice	0.1648 (0.0000–0.3696)	0.008280 (0.0003008–0.02242)	0.002161 (0.0001841–0.005225)	0.01918 (0.0008726–0.05219)	Not supported	0.08578 (0.05759–0.1178)	0.1538 (0.1192–0.1906)
			HD95	32.85 (27.73–38.85)	89.56 (81.95–98.03)	93.70 (84.75–102.7)	76.63 (66.01–87.73)		37.75 (31.07–46.35)	32.35 (28.07–36.28)
	ULS23-DeepLesion3D	lesion	Dice	Not supported	0.004582 (0.001279–0.009446)	0.0004724 (0.0000–0.001265)	0.06145 (0.02352–0.1064)	Not supported	0.1722 (0.1136–0.2407)	0.2102 (0.1435–0.2798)
			HD95		98.13 (90.15–107.2)	94.79 (87.87–101.8)	54.29 (44.39–64.53)		40.27 (31.97–49.39)	37.66 (29.42–46.09)
	CT-Lymph-Nodes	abdominal lymph node	Dice	Not supported	0.01018 (0.001135–0.02459)	0.0000 (0.0000–0.0000)	0.03879 (0.01030–0.06972)	Not supported	0.1404 (0.03606–0.2741)	0.2723 (0.08376–0.4746)
			HD95		164.5 (104.1–222.7)	237.0 (196.5–280.0)	283.3 (227.3–354.6)		163.4 (108.9–210.5)	141.8 (84.89–201.7)
	MHB-SEG-multiclass	epidural hemorrhage	Dice	Not supported	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	Not supported	0.02401 (0.0000–0.05538)	0.1481 (0.03280–0.2635)
			HD95		99.04 (85.50–112.6)	63.25 (18.36–98.73)	100.3 (100.3–100.3)		60.53 (29.79–84.06)	65.61 (43.02–87.97)
		subdural hemorrhage	Dice	Not supported	0.001488 (0.0000–0.004252)	0.3481 (0.2215–0.4899)	0.2614 (0.1448–0.3892)	Not supported	0.4107 (0.2589–0.5586)	0.4532 (0.3026–0.6053)
			HD95		84.68 (76.94–93.21)	37.66 (25.78–49.66)	69.98 (57.18–82.05)		42.69 (29.71–58.29)	46.94 (31.14–64.33)
		subarachnoid hemorrhage	Dice	Not supported	0.00005765 (0.0000–0.0001729)	0.2185 (0.1089–0.3331)	0.1549 (0.05791–0.2796)	Not supported	0.1458 (0.06387–0.2362)	0.2698 (0.1478–0.3947)
			HD95		81.28 (73.69–89.61)	54.63 (41.70–67.04)	67.43 (53.18–84.89)		50.17 (41.31–60.45)	44.32 (34.48–54.15)

Supplementary Table 8 (continued) | Detailed Dice and HD95 scores of HoloCT and competing methods on internal, External-seen (zero-shot), External-unseen (zero-shot), and private zero-shot/fine-tuning segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. ▲ marks VoxTell results on PANORAMA; because PANORAMA was used to train VoxTell, these results do not constitute external validation for VoxTell.

Dataset Type	Dataset Name	Label Name	Metric	VISTA3D	M3D-Phi3-4B	SAT-Pro	VoxTell	nnU-Net	HoloCT-4B	HoloCT-8B
External-unseen (zero-shot)	MHB-SEG-multiclass	intraparenchymal hemorrhage	Dice	Not supported	0.003667 (0.0000–0.009758)	0.09978 (0.04260–0.1712)	0.07165 (0.01379–0.1510)	Not supported	0.1014 (0.05298–0.1557)	0.1073 (0.04726–0.1846)
			HD95		83.40 (73.95–92.38)	54.88 (43.18–67.93)	78.71 (63.92–94.72)		60.88 (52.72–69.58)	62.58 (51.77–73.73)
		intraventricular hemorrhage	Dice	Not supported	0.001244 (0.0000–0.003699)	0.02984 (0.01136–0.05626)	0.08302 (0.03221–0.1446)	Not supported	0.2030 (0.1175–0.2972)	0.3504 (0.2483–0.4575)
			HD95		86.53 (78.41–94.72)	75.45 (66.81–84.17)	66.66 (56.81–77.41)		64.31 (52.04–75.83)	60.31 (47.25–73.60)
	Adrenal-ACC-Ki67	adrenocortical carcinoma	Dice	Not supported	0.4918 (0.3794–0.5986)	0.4771 (0.4562–0.4989)	0.6579 (0.5546–0.7651)	Not supported	0.8370 (0.7055–0.9170)	0.8923 (0.8164–0.9342)
			HD95		49.52 (36.34–63.01)	54.61 (52.81–56.40)	23.37 (14.90–32.49)		34.18 (10.40–63.09)	18.36 (4.355–44.44)
	PANORAMA	pancreatic ductal adenocarcinoma	Dice	Not supported	0.06358 (0.03703–0.09149)	0.00003823 (0.0000–0.0001147)	0.5403▲ (0.4713–0.6060)	Not supported	0.4040 (0.3465–0.4634)	0.4382 (0.3665–0.5032)
			HD95		92.86 (81.70–104.5)	227.4 (195.0–260.0)	21.06▲ (14.52–27.61)		27.52 (22.95–32.52)	28.36 (22.99–34.05)
		pancreatic veins	Dice	Not supported	0.09721 (0.08661–0.1081)	0.3073 (0.2752–0.3387)	0.8379▲ (0.8216–0.8529)	Not supported	0.7008 (0.6835–0.7177)	0.7661 (0.7498–0.7798)
			HD95		39.71 (37.06–43.06)	57.15 (49.05–66.57)	15.99▲ (13.59–18.85)		18.00 (16.31–19.86)	25.97 (24.44–27.81)
		pancreatic arteries	Dice	Not supported	0.02302 (0.02053–0.02560)	0.002528 (0.000004579–0.007547)	0.8251▲ (0.8146–0.8356)	Not supported	0.4146 (0.3956–0.4339)	0.2630 (0.2521–0.2734)
			HD95		44.20 (41.60–47.22)	171.9 (101.8–255.7)	15.90▲ (14.50–17.34)		96.76 (88.29–104.8)	86.21 (73.86–98.08)
		pancreatic parenchyma	Dice	Not supported	0.4686 (0.4439–0.4942)	0.0005745 (0.00001441–0.001338)	0.8422▲ (0.8247–0.8587)	Not supported	0.8331 (0.8152–0.8506)	0.8521 (0.8359–0.8686)
			HD95		24.27 (20.89–27.80)	223.1 (187.8–262.0)	11.70▲ (9.504–14.37)		6.966 (5.901–8.070)	7.511 (6.211–8.870)
	HCC-TACE-Seg	hepatocellular carcinoma	Dice	Not supported	0.2264 (0.1120–0.3515)	0.2471 (0.1647–0.3415)	0.3140 (0.2036–0.4299)	Not supported	0.5944 (0.4785–0.7013)	0.6428 (0.5403–0.7401)
			HD95		96.72 (74.08–119.9)	128.8 (103.2–153.9)	133.6 (109.9–158.4)		28.84 (19.96–38.95)	21.47 (15.03–29.33)
Private (zero-shot)	Private01 (laryngeal cancer)	laryngeal cancer	Dice	Not supported	0.009046 (0.002355–0.01793)	0.004320 (0.0003728–0.01074)	0.1070 (0.06581–0.1499)	Not supported	0.3713 (0.3313–0.4143)	0.4880 (0.4413–0.5308)
			HD95		122.0 (112.4–129.6)	44.68 (28.75–63.27)	40.84 (28.62–54.97)		16.32 (12.75–21.19)	14.62 (10.83–20.13)
	Private02 (lung cancer)	lung cancer	Dice	Not supported	0.2409 (0.2038–0.2777)	0.4417 (0.4008–0.4831)	0.5428 (0.5057–0.5794)	Not supported	0.6962 (0.6681–0.7257)	0.7152 (0.6872–0.7412)
			HD95		112.8 (101.8–123.5)	36.70 (29.74–44.33)	93.66 (85.68–102.1)		33.61 (27.75–40.91)	30.01 (24.21–36.39)
	Private03 (thyroid cancer-multi)	malignant thyroid tumor	Dice	Not supported	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	Not supported	0.1116 (0.04548–0.1923)	0.1461 (0.06687–0.2438)
			HD95		143.7 (122.3–163.2)	127.2 (115.5–138.2)	34.60 (30.97–38.22)		35.27 (30.47–40.12)	34.83 (29.40–39.71)
		benign thyroid tumor	Dice	Not supported	0.01230 (0.0002859–0.02863)	0.0003445 (0.0000–0.001017)	0.01297 (0.001348–0.02792)	Not supported	0.5331 (0.4659–0.5910)	0.6232 (0.5625–0.6787)
			HD95		93.72 (87.13–100.4)	127.9 (123.2–132.7)	27.58 (23.65–31.57)		24.41 (21.38–27.49)	19.29 (16.25–22.16)
	Private03 (thyroid cancer-one)	thyroid tumor	Dice	Not supported	0.002521 (0.0006734–0.004839)	0.004120 (0.001315–0.007779)	0.1804 (0.1340–0.2260)	Not supported	0.5442 (0.4878–0.6007)	0.6183 (0.5610–0.6719)
			HD95		103.1 (94.03–111.9)	81.93 (63.88–103.4)	25.88 (23.33–28.39)		21.12 (18.00–24.71)	18.74 (15.26–22.80)
Private (fine-tuning)	Private01 (laryngeal cancer)	laryngeal cancer	Dice	0.6157 (0.5559–0.6720)	0.3903 (0.3372–0.4595)	0.4822 (0.4312–0.5418)	Not supported	0.6545 (0.6017–0.7070)	0.6684 (0.6213–0.7147)	0.6839 (0.6270–0.7202)
			HD95	17.08 (12.91–23.86)	27.04 (19.56–35.33)	19.29 (13.08–25.48)		15.34 (10.69–20.47)	13.18 (9.160–18.31)	12.21 (9.160–18.21)
	Private02 (lung cancer)	lung cancer	Dice	0.7440 (0.7098–0.7685)	0.6559 (0.6163–0.6970)	0.7263 (0.6950–0.7503)	Not supported	0.7758 (0.7415–0.7901)	0.7791 (0.7549–0.8024)	0.7859 (0.7616–0.8090)
			HD95	35.99 (28.97–42.18)	46.50 (37.89–54.87)	37.46 (29.60–43.27)		31.70 (25.94–37.91)	29.37 (23.91–35.84)	31.94 (25.44–38.08)
	Private03 (thyroid cancer-multi)	malignant thyroid tumor	Dice	0.3231 (0.2288–0.4239)	0.2269 (0.1479–0.3112)	0.3020 (0.2126–0.4095)	Not supported	0.3546 (0.2603–0.4668)	0.3746 (0.2811–0.4902)	0.3809 (0.2874–0.4967)
			HD95	25.84 (18.55–32.77)	36.32 (28.19–45.22)	28.41 (20.72–35.03)		23.26 (16.42–29.92)	21.79 (15.63–29.31)	20.84 (14.81–27.19)
		benign thyroid tumor	Dice	0.7521 (0.6819–0.8033)	0.6457 (0.5895–0.7023)	0.7339 (0.6792–0.7943)	Not supported	0.7993 (0.7345–0.8487)	0.7941 (0.7368–0.8472)	0.8084 (0.7512–0.8613)
			HD95	10.44 (7.080–12.11)	16.46 (12.73–20.72)	11.67 (8.230–14.50)		8.320 (6.024–10.54)	8.388 (6.424–9.930)	8.126 (6.453–9.171)
	Private03 (thyroid cancer-one)	thyroid tumor	Dice	0.7677 (0.7141–0.8235)	0.6758 (0.6222–0.7390)	0.7577 (0.7057–0.8077)	Not supported	0.8194 (0.7592–0.8533)	0.8049 (0.7604–0.8501)	0.8187 (0.7742–0.8643)
			HD95	10.39 (8.140–12.22)	15.46 (11.14–19.96)	11.45 (8.070–14.25)		8.528 (6.322–10.89)	8.370 (6.395–9.747)	8.341 (6.292–9.511)

Supplementary Table 9 | Detailed sensitivity, specificity, F1, and conditional Dice scores of HoloCT and competing methods on internal, external-seen zero-shot/fine-tuning, external-unseen zero-shot/fine-tuning, and private zero-shot/fine-tuning classification-segmentation benchmarks.
Each score is reported as mean with 95% confidence interval. Cond. Dice denotes conditional Dice. For the external zero-shot Macro F1 summary in Fig. 3c, unsupported ResNet and nnU-Net targets are scored as zero, so every model is averaged across the same nine targets.

Dataset Type	Dataset Name	Label Name	Metric	M3D-Phi3-4B	nnU-Net	ResNet	HoloCT-4B w/o grounding	HoloCT-4B w/ grounding	HoloCT-8B w/o grounding	HoloCT-8B w/ grounding
Internal	Total-SegmentatorV2	left kidney cyst	Sens.	0.5000 (0.0000–1.000)	1.000 (1.000–1.000)	0.8333 (0.5000–1.000)	0.6667 (0.2500–1.000)	0.6667 (0.2500–1.000)	1.000 (1.000–1.000)	0.8333 (0.4611–1.000)
			Spec.	0.6529 (0.5950–0.7137)	0.7810 (0.7265–0.8340)	0.7769 (0.7231–0.8306)	0.9752 (0.9535–0.9918)	0.9669 (0.9419–0.9876)	0.9711 (0.9463–0.9877)	0.9752 (0.9542–0.9918)
			F1	0.06452 (0.0000–0.1379)	0.1846 (0.06667–0.3158)	0.1538 (0.03704–0.2759)	0.5000 (0.1536–0.7692)	0.4444 (0.1248–0.7059)	0.6316 (0.3071–0.8464)	0.5882 (0.2500–0.8235)
			Cond. Dice	0.03792 (0.0000–0.07468)	0.7320 (0.5424–0.8979)	Not supported	Not supported	0.9016 (0.8845–0.9240)	Not supported	0.8872 (0.8380–0.9342)
		right kidney cyst	Sens.	0.5000 (0.0000–1.000)	0.5000 (0.0000–1.000)	0.6667 (0.2500–1.000)	0.5000 (0.0000–1.000)	0.5000 (0.0000–1.000)	0.3333 (0.0000–0.7537)	0.5000 (0.0000–1.000)
			Spec.	0.7769 (0.7231–0.8299)	0.8017 (0.7520–0.8506)	0.8347 (0.7860–0.8776)	0.9959 (0.9874–1.000)	0.9917 (0.9793–1.000)	0.9959 (0.9874–1.000)	0.9917 (0.9793–1.000)
			F1	0.09524 (0.0000–0.1967)	0.1053 (0.0000–0.2105)	0.1600 (0.03844–0.2963)	0.6000 (0.0000–0.9098)	0.5455 (0.0000–0.8750)	0.4444 (0.0000–0.8240)	0.5455 (0.0000–0.8750)
			Cond. Dice	0.2163 (0.0000–0.6490)	0.6539 (0.5931–0.7558)	Not supported	Not supported	0.8536 (0.7821–0.9343)	Not supported	0.8563 (0.7787–0.9328)
	KiTS23	kidney cyst	Sens.	0.0000 (0.0000–0.0000)	0.8909 (0.8043–0.9643)	0.6727 (0.5500–0.7917)	0.5636 (0.4259–0.6897)	0.6545 (0.5245–0.7778)	0.6909 (0.5660–0.8085)	0.6909 (0.5671–0.8077)
			Spec.	1.000 (1.000–1.000)	0.0000 (0.0000–0.0000)	0.4884 (0.3333–0.6410)	0.9535 (0.8837–1.000)	0.9070 (0.8181–0.9778)	0.8372 (0.7273–0.9459)	0.8605 (0.7561–0.9545)
			F1	0.0000 (0.0000–0.0000)	0.6667 (0.5693–0.7516)	0.6491 (0.5361–0.7458)	0.7045 (0.5823–0.8036)	0.7579 (0.6506–0.8462)	0.7600 (0.6591–0.8462)	0.7677 (0.6667–0.8519)
			Cond. Dice	-	0.2649 (0.1930–0.3366)	Not supported	Not supported	0.5781 (0.4930–0.6636)	Not supported	0.7046 (0.6268–0.7816)
	CT-ICH	intracranial hemorrhage	Sens.	1.000 (1.000–1.000)	1.000 (1.000–1.000)	1.000 (1.000–1.000)	0.2857 (0.0000–0.6667)	0.4286 (0.0000–0.8333)	0.4286 (0.0000–0.8333)	0.4286 (0.0000–0.8333)
			Spec.	0.1250 (0.0000–0.4000)	0.8750 (0.5714–1.000)	0.5000 (0.1424–0.8571)	0.7500 (0.4286–1.000)	0.7500 (0.4000–1.000)	0.7500 (0.4000–1.000)	0.7500 (0.4000–1.000)
			F1	0.6667 (0.3529–0.8696)	0.9333 (0.7273–1.000)	0.7778 (0.5000–0.9524)	0.3636 (0.0000–0.7143)	0.5000 (0.0000–0.8000)	0.5000 (0.0000–0.8000)	0.5000 (0.0000–0.8000)
			Cond. Dice	0.0000 (0.0000–0.0000)	0.8793 (0.7911–0.9419)	Not supported	Not supported	0.5557 (0.5180–0.6021)	Not supported	0.6853 (0.5247–0.8460)
	PanTS	pancreas lesion	Sens.	0.5894 (0.5102–0.6690)	0.4636 (0.3831–0.5411)	0.5232 (0.4429–0.6015)	0.3046 (0.2313–0.3784)	0.3974 (0.3200–0.4694)	0.4305 (0.3551–0.5172)	0.4834 (0.4057–0.5690)
			Spec.	0.3320 (0.2972–0.3644)	0.8867 (0.8623–0.9073)	0.9107 (0.8899–0.9302)	0.9960 (0.9908–1.000)	0.9907 (0.9827–0.9973)	0.9787 (0.9674–0.9879)	0.9760 (0.9647–0.9863)
			F1	0.2402 (0.1983–0.2804)	0.4575 (0.3852–0.5228)	0.5320 (0.4590–0.5970)	0.4600 (0.3696–0.5405)	0.5505 (0.4660–0.6224)	0.5603 (0.4830–0.6360)	0.6033 (0.5315–0.6767)
			Cond. Dice	0.2034 (0.1575–0.2502)	0.3146 (0.2930–0.3796)	Not supported	Not supported	0.4920 (0.4242–0.5574)	Not supported	0.5412 (0.4820–0.6016)
	AbdomenAtlas 3.0	kidney lesion	Sens.	0.7290 (0.6779–0.7807)	0.7214 (0.6641–0.7744)	0.8588 (0.8162–0.9011)	0.3702 (0.3113–0.4286)	0.5153 (0.4527–0.5772)	0.2939 (0.2422–0.3494)	0.3740 (0.3162–0.4317)
			Spec.	0.2319 (0.1997–0.2633)	0.09337 (0.07217–0.1162)	0.6717 (0.6344–0.7067)	0.9714 (0.9587–0.9826)	0.9458 (0.9286–0.9629)	0.9940 (0.9867–0.9985)	0.9804 (0.9691–0.9898)
			F1	0.3967 (0.3563–0.4353)	0.3590 (0.3224–0.3944)	0.6383 (0.5975–0.6771)	0.5132 (0.4485–0.5714)	0.6236 (0.5670–0.6742)	0.4490 (0.3834–0.5121)	0.5255 (0.4615–0.5829)
			Cond. Dice	0.2381 (0.1936–0.2842)	0.7081 (0.6614–0.7303)	Not supported	Not supported	0.7344 (0.6924–0.7729)	Not supported	0.8221 (0.7941–0.8475)
		liver lesion	Sens.	0.9776 (0.9563–0.9954)	0.9058 (0.8643–0.9420)	0.4529 (0.3883–0.5156)	0.6502 (0.5855–0.7129)	0.7668 (0.7110–0.8217)	0.8027 (0.7489–0.8543)	0.7130 (0.6529–0.7718)
			Spec.	0.03698 (0.02446–0.05234)	0.0000 (0.0000–0.0000)	0.8009 (0.7717–0.8280)	0.9602 (0.9459–0.9739)	0.9388 (0.9207–0.9558)	0.8947 (0.8712–0.9177)	0.9388 (0.9214–0.9563)
			F1	0.3900 (0.3539–0.4271)	0.3582 (0.3225–0.3924)	0.4353 (0.3795–0.4907)	0.7323 (0.6808–0.7775)	0.7826 (0.7387–0.8227)	0.7521 (0.7072–0.7936)	0.7482 (0.7005–0.7933)
			Cond. Dice	0.1827 (0.1332–0.2372)	0.6494 (0.5935–0.6831)	Not supported	Not supported	0.7248 (0.6988–0.7506)	Not supported	0.7102 (0.6796–0.7402)
		pancreas lesion	Sens.	0.6277 (0.5500–0.7045)	0.6204 (0.5378–0.7035)	0.4818 (0.3968–0.5630)	0.4745 (0.3931–0.5573)	0.5693 (0.4857–0.6462)	0.6277 (0.5454–0.7054)	0.6131 (0.5349–0.6957)
			Spec.	0.4385 (0.4038–0.4736)	0.3118 (0.2809–0.3442)	0.7275 (0.6975–0.7584)	0.9886 (0.9806–0.9950)	0.9797 (0.9698–0.9887)	0.9341 (0.9160–0.9499)	0.9569 (0.9427–0.9706)
			F1	0.2583 (0.2140–0.3022)	0.2222 (0.1820–0.2627)	0.3158 (0.2571–0.3723)	0.6161 (0.5384–0.6854)	0.6753 (0.6021–0.7369)	0.6255 (0.5574–0.6869)	0.6588 (0.5892–0.7207)
			Cond. Dice	0.1387 (0.09945–0.1816)	0.4728 (0.4217–0.5494)	Not supported	Not supported	0.5480 (0.4906–0.6055)	Not supported	0.5712 (0.5176–0.6266)
		colon lesion	Sens.	1.000 (1.000–1.000)	1.000 (1.000–1.000)	0.8000 (0.6250–0.9545)	0.8800 (0.7391–1.000)	0.9200 (0.7931–1.000)	0.5600 (0.3529–0.7587)	0.7200 (0.5332–0.8889)
			Spec.	0.05771 (0.04319–0.07325)	0.02775 (0.01774–0.03859)	0.03219 (0.02111–0.04373)	0.9589 (0.9457–0.9711)	0.9345 (0.9183–0.9501)	0.9922 (0.9865–0.9978)	0.9834 (0.9735–0.9911)
			F1	0.05562 (0.03778–0.07827)	0.05400 (0.03486–0.07463)	0.04362 (0.02623–0.06243)	0.5238 (0.3846–0.6429)	0.4299 (0.3010–0.5391)	0.6087 (0.4241–0.7619)	0.6207 (0.4641–0.7568)
			Cond. Dice	0.05536 (0.001067–0.1158)	0.6813 (0.5886–0.8194)	Not supported	Not supported	0.6678 (0.5685–0.7530)	Not supported	0.7231 (0.6598–0.7816)

Supplementary Table 9 (continued) | Detailed sensitivity, specificity, F1, and conditional Dice scores of HoloCT and competing methods on internal, external-seen zero-shot/fine-tuning, external-unseen zero-shot/fine-tuning, and private zero-shot/fine-tuning classification-segmentation benchmarks. Each score is reported as mean with 95% confidence interval. Cond. Dice denotes conditional Dice. For the external zero-shot Macro F1 summary in Fig. 3c, unsupported ResNet and nnU-Net targets are scored as zero, so every model is averaged across the same nine targets.

Dataset Type	Dataset Name	Label Name	Metric	M3D-Phi3-4B	nnU-Net	ResNet	HoloCT-4B w/o grounding	HoloCT-4B w/ grounding	HoloCT-8B w/o grounding	HoloCT-8B w/ grounding
Internal	LiMT	liver cyst	Sens.	0.0000 (0.0000–0.0000)	0.8000 (0.5000–1.000)	0.0000 (0.0000–0.0000)	0.6000 (0.2854–0.8892)	0.6000 (0.2857–0.9000)	0.7000 (0.3747–1.000)	0.7000 (0.3747–1.000)
			Spec.	1.000 (1.000–1.000)	0.7000 (0.5000–0.9000)	0.9500 (0.8333–1.000)	0.9500 (0.8333–1.000)	0.9000 (0.7500–1.000)	0.9000 (0.7500–1.000)	0.9000 (0.7500–1.000)
			F1	0.0000 (0.0000–0.0000)	0.6667 (0.4000–0.8667)	0.0000 (0.0000–0.0000)	0.7059 (0.4000–0.9167)	0.6667 (0.3636–0.8750)	0.7368 (0.4444–0.9231)	0.7368 (0.4444–0.9231)
			Cond. Dice	-	0.6562 (0.5824–0.7276)	Not supported	Not supported	0.6003 (0.4689–0.7237)	Not supported	0.6448 (0.5193–0.7476)
		liver hemangioma	Sens.	0.0000 (0.0000–0.0000)	0.8333 (0.5000–1.000)	0.6667 (0.2000–1.000)	0.6667 (0.2500–1.000)	0.6667 (0.2500–1.000)	1.000 (1.000–1.000)	1.000 (1.000–1.000)
			Spec.	1.000 (1.000–1.000)	0.7917 (0.6190–0.9524)	0.7500 (0.5652–0.9200)	1.000 (1.000–1.000)	1.000 (1.000–1.000)	0.8750 (0.7199–1.000)	0.8750 (0.7199–1.000)
			F1	0.0000 (0.0000–0.0000)	0.6250 (0.2500–0.8462)	0.5000 (0.1429–0.7619)	0.8000 (0.4000–1.000)	0.8000 (0.4000–1.000)	0.8000 (0.5000–1.000)	0.8000 (0.5000–1.000)
			Cond. Dice	-	0.6200 (0.2938–0.8864)	Not supported	Not supported	0.8313 (0.7636–0.8765)	Not supported	0.7106 (0.4914–0.8313)
		metastatic liver cancer	Sens.	0.0000 (0.0000–0.0000)	0.7500 (0.0000–1.000)	0.5000 (0.0000–1.000)	0.2500 (0.0000–0.7625)	0.2500 (0.0000–0.7625)	0.2500 (0.0000–1.000)	0.2500 (0.0000–1.000)
			Spec.	1.000 (1.000–1.000)	0.7692 (0.5998–0.9231)	0.7308 (0.5600–0.8846)	1.000 (1.000–1.000)	1.000 (1.000–1.000)	0.9615 (0.8696–1.000)	0.9615 (0.8696–1.000)
			F1	0.0000 (0.0000–0.0000)	0.4615 (0.0000–0.7505)	0.3077 (0.0000–0.6250)	0.4000 (0.0000–0.8651)	0.4000 (0.0000–0.8651)	0.3333 (0.0000–0.8000)	0.3333 (0.0000–0.8000)
			Cond. Dice	-	0.4623 (0.2354–0.6347)	Not supported	Not supported	0.6313 (0.6313–0.6313)	Not supported	0.6605 (0.6605–0.6605)
	LiMT-oneclass	primary liver cancer	Sens.	0.0000 (0.0000–0.0000)	0.8571 (0.5000–1.000)	0.8571 (0.5000–1.000)	1.000 (1.000–1.000)	1.000 (1.000–1.000)	1.000 (1.000–1.000)	1.000 (1.000–1.000)
			Spec.	1.000 (1.000–1.000)	0.8261 (0.6538–0.9583)	0.7391 (0.5455–0.9092)	1.000 (1.000–1.000)	1.000 (1.000–1.000)	1.000 (1.000–1.000)	1.000 (1.000–1.000)
			F1	0.0000 (0.0000–0.0000)	0.7059 (0.4000–0.9091)	0.6316 (0.3077–0.8462)	1.000 (1.000–1.000)	1.000 (1.000–1.000)	1.000 (1.000–1.000)	1.000 (1.000–1.000)
			Cond. Dice	-	0.6500 (0.3433–0.8813)	Not supported	Not supported	0.7407 (0.5913–0.8614)	Not supported	0.6904 (0.4280–0.8608)
		liver lesion	Sens.	0.0000 (0.0000–0.0000)	0.4000 (0.1000–0.7278)	0.5000 (0.1667–0.8333)	0.7000 (0.3750–1.000)	0.7000 (0.3750–1.000)	0.9000 (0.6667–1.000)	0.9000 (0.6667–1.000)
			Spec.	1.000 (1.000–1.000)	0.7500 (0.5383–0.9202)	0.7500 (0.5500–0.9375)	0.8500 (0.6667–1.000)	0.8500 (0.6667–1.000)	0.8000 (0.6000–0.9501)	0.8000 (0.6000–0.9501)
			F1	0.0000 (0.0000–0.0000)	0.4211 (0.1176–0.6667)	0.5000 (0.1818–0.7410)	0.7000 (0.4211–0.8889)	0.7000 (0.4211–0.8889)	0.7826 (0.5455–0.9375)	0.7826 (0.5455–0.9375)
			Cond. Dice	-	0.2835 (0.2158–0.3212)	Not supported	Not supported	0.6100 (0.5105–0.7276)	Not supported	0.5606 (0.4261–0.6810)
External-seen (zero-shot)	QUBIQ2021	pancreas lesion	Sens.	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.1000 (0.0000–0.3333)	0.1000 (0.0000–0.3333)	0.1000 (0.0000–0.3333)	0.2000 (0.0000–0.5000)
			Spec.	1.000 (1.000–1.000)	1.000 (1.000–1.000)	0.2500 (0.0000–0.5714)	1.000 (1.000–1.000)	1.000 (1.000–1.000)	1.000 (1.000–1.000)	1.000 (1.000–1.000)
			F1	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.1818 (0.0000–0.5000)	0.1818 (0.0000–0.5000)	0.1818 (0.0000–0.5000)	0.3333 (0.0000–0.6667)
			Cond. Dice	-	-	Not supported	Not supported	0.3361 (0.3361–0.3361)	Not supported	0.3867 (0.3384–0.4350)
	TotalSeg-subtask-liver-lesion	liver lesion	Sens.	0.0000 (0.0000–0.0000)	0.09191 (0.05861–0.1301)	0.1287 (0.08855–0.1715)	0.8088 (0.7593–0.8535)	0.8676 (0.8261–0.9074)	0.8750 (0.8327–0.9135)	0.7978 (0.7491–0.8456)
			Spec.	1.000 (1.000–1.000)	0.1111 (0.0000–0.3750)	0.1111 (0.0000–0.3750)	0.8889 (0.6248–1.000)	0.7778 (0.4989–1.000)	1.000 (1.000–1.000)	1.000 (1.000–1.000)
			F1	0.0000 (0.0000–0.0000)	0.1639 (0.1081–0.2222)	0.2222 (0.1579–0.2866)	0.8925 (0.8613–0.9206)	0.9255 (0.9006–0.9482)	0.9333 (0.9087–0.9548)	0.8875 (0.8565–0.9163)
			Cond. Dice	-	0.07529 (0.03029–0.1113)	Not supported	Not supported	0.7096 (0.6827–0.7361)	Not supported	0.7456 (0.7185–0.7684)
	MHB-SEG-oneclass	intracranial hemorrhage	Sens.	0.0000 (0.0000–0.0000)	0.09524 (0.02273–0.2000)	0.1429 (0.04651–0.2564)	0.5714 (0.4146–0.7251)	0.6429 (0.4878–0.7750)	0.5238 (0.3684–0.6818)	0.4762 (0.3191–0.6364)
			Spec.	1.000 (1.000–1.000)	0.1667 (0.0000–0.6000)	0.1667 (0.0000–0.5014)	1.000 (1.000–1.000)	1.000 (1.000–1.000)	1.000 (1.000–1.000)	1.000 (1.000–1.000)
			F1	0.0000 (0.0000–0.0000)	0.1569 (0.04082–0.2963)	0.2264 (0.08159–0.3730)	0.7273 (0.5862–0.8406)	0.7826 (0.6557–0.8732)	0.6875 (0.5384–0.8108)	0.6452 (0.4838–0.7778)
			Cond. Dice	-	0.1263 (0.07529–0.1755)	Not supported	Not supported	0.4056 (0.2862–0.5260)	Not supported	0.5143 (0.3883–0.6315)
External-seen (fine-tuning)	QUBIQ2021	pancreas lesion	Sens.	0.2000 (0.0000–0.5000)	0.9000 (0.6667–1.000)	0.9000 (0.6667–1.000)	Not supported	1.000 (1.000–1.000)	Not supported	0.9000 (0.7000–1.000)
			Spec.	0.8750 (0.6000–1.000)	0.7500 (0.4000–1.000)	0.2500 (0.0000–0.6250)		0.6250 (0.2500–1.000)		1.000 (1.000–1.000)
			F1	0.3077 (0.0000–0.6154)	0.8571 (0.6250–1.000)	0.7200 (0.4545–0.8966)		0.8696 (0.6667–1.000)		0.9474 (0.8235–1.000)
			Cond. Dice	0.1346 (0.08605–0.1754)	0.4318 (0.2274–0.6469)	Not supported		0.7173 (0.6478–0.7964)		0.7808 (0.7108–0.8577)

Supplementary Table 9 (continued) | Detailed sensitivity, specificity, F1, and conditional Dice scores of HoloCT and competing methods on internal, external-seen zero-shot/fine-tuning, external-unseen zero-shot/fine-tuning, and private zero-shot/fine-tuning classification-segmentation benchmarks. Each score is reported as mean with 95% confidence interval. Cond. Dice denotes conditional Dice. For the external zero-shot Macro F1 summary in Fig. 3c, unsupported ResNet and nnU-Net targets are scored as zero, so every model is averaged across the same nine targets.

Dataset Type	Dataset Name	Label Name	Metric	M3D-Phi3-4B	nnU-Net	ResNet	HoloCT-4B w/o grounding	HoloCT-4B w/ grounding	HoloCT-8B w/o grounding	HoloCT-8B w/ grounding
External-seen (fine-tuning)	TotalSeg-subtask-liver-lesion	liver lesion	Sens.	0.9265 (0.8926–0.9554)	0.9706 (0.9485–0.9889)	1.000 (1.000–1.000)	Not supported	0.9890 (0.9743–1.000)	Not supported	0.9926 (0.9814–1.000)
			Spec.	0.3333 (0.0000–0.7146)	0.2222 (0.0000–0.5556)	0.1111 (0.0000–0.3750)		0.7778 (0.4444–1.000)		0.8889 (0.6000–1.000)
			F1	0.9509 (0.9305–0.9669)	0.9724 (0.9568–0.9837)	0.9855 (0.9744–0.9946)		0.9908 (0.9814–0.9982)		0.9945 (0.9870–1.000)
			Cond. Dice	0.2444 (0.2064–0.2807)	0.7723 (0.7427–0.8009)	Not supported		0.7171 (0.6954–0.7444)		0.7665 (0.7443–0.7985)
	MHB-SEG-oneclass	intracranial hemorrhage	Sens.	0.9286 (0.8421–1.000)	1.000 (1.000–1.000)	0.9762 (0.9231–1.000)	Not supported	1.000 (1.000–1.000)	Not supported	1.000 (1.000–1.000)
			Spec.	0.3333 (0.0000–0.8000)	0.3333 (0.0000–0.7500)	0.0000 (0.0000–0.0000)		0.6667 (0.1415–1.000)		0.8333 (0.4286–1.000)
			F1	0.9176 (0.8500–0.9680)	0.9545 (0.9048–0.9895)	0.9213 (0.8571–0.9787)		0.9767 (0.9383–1.000)		0.9882 (0.9620–1.000)
			Cond. Dice	0.1528 (0.07214–0.1854)	0.6995 (0.6429–0.7484)	Not supported		0.5750 (0.5116–0.6475)		0.6220 (0.5651–0.6912)
External-unseen (zero-shot)	PANORAMA	pancreatic ductal adenocarcinoma	Sens.	0.07576 (0.01639–0.1429)	Not supported	Not supported	0.7846 (0.6849–0.8814)	0.8462 (0.7535–0.9311)	0.4923 (0.3709–0.6112)	0.8462 (0.7581–0.9333)
			Spec.	0.1234 (0.07333–0.1768)			0.6883 (0.6207–0.7639)	0.7403 (0.6730–0.8077)	0.9481 (0.9125–0.9805)	0.8442 (0.7871–0.8976)
			F1	0.04854 (0.009950–0.09391)			0.6220 (0.5393–0.7097)	0.6875 (0.6000–0.7683)	0.6095 (0.4898–0.7091)	0.7639 (0.6880–0.8382)
			Cond. Dice	0.1064 (0.07157–0.1459)			Not supported	0.4112 (0.3469–0.4737)	Not supported	0.4468 (0.3702–0.5143)
		epidural hemorrhage	Sens.	1.000 (1.000–1.000)	Not supported	Not supported	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)
			Spec.	0.0000 (0.0000–0.0000)			1.000 (1.000–1.000)	1.000 (1.000–1.000)	1.000 (1.000–1.000)	1.000 (1.000–1.000)
			F1	0.1887 (0.04082–0.3448)			0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)
			Cond. Dice	0.0000 (0.0000–0.0000)			Not supported	-	Not supported	-
		subdural hemorrhage	Sens.	0.05000 (0.0000–0.1667)	Not supported	Not supported	0.4000 (0.1905–0.6190)	0.3500 (0.1429–0.5789)	0.3500 (0.1429–0.5789)	0.3500 (0.1429–0.5789)
			Spec.	1.000 (1.000–1.000)			0.9643 (0.8787–1.000)	0.9643 (0.8787–1.000)	0.9643 (0.8787–1.000)	0.9643 (0.8787–1.000)
			F1	0.09524 (0.0000–0.2857)			0.5517 (0.2999–0.7500)	0.5000 (0.2353–0.7097)	0.5000 (0.2353–0.7097)	0.5000 (0.2353–0.7097)
			Cond. Dice	0.002020 (0.002020–0.002020)			Not supported	0.6676 (0.4534–0.8512)	Not supported	0.6724 (0.4538–0.8441)
	MHB-SEG-multiclass	subarachnoid hemorrhage	Sens.	0.0000 (0.0000–0.0000)	Not supported	Not supported	0.3000 (0.1153–0.5238)	0.2000 (0.04545–0.4000)	0.1000 (0.0000–0.2500)	0.3000 (0.1111–0.5238)
			Spec.	1.000 (1.000–1.000)			0.9286 (0.8182–1.000)	0.9643 (0.8750–1.000)	1.000 (1.000–1.000)	0.9643 (0.8800–1.000)
			F1	0.0000 (0.0000–0.0000)			0.4286 (0.1818–0.6452)	0.3200 (0.08333–0.5386)	0.1818 (0.0000–0.4000)	0.4444 (0.1905–0.6667)
			Cond. Dice	-			Not supported	0.1777 (0.0000–0.5956)	Not supported	0.5016 (0.2761–0.7039)
		intraparenchymal hemorrhage	Sens.	0.0000 (0.0000–0.0000)	Not supported	Not supported	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)
			Spec.	1.000 (1.000–1.000)			1.000 (1.000–1.000)	1.000 (1.000–1.000)	1.000 (1.000–1.000)	1.000 (1.000–1.000)
			F1	0.0000 (0.0000–0.0000)			0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)
			Cond. Dice	-			Not supported	-	Not supported	-
		intraventricular hemorrhage	Sens.	0.0000 (0.0000–0.0000)	Not supported	Not supported	0.04762 (0.0000–0.1667)	0.04762 (0.0000–0.1579)	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)
			Spec.	1.000 (1.000–1.000)			1.000 (1.000–1.000)	0.9630 (0.8749–1.000)	0.9630 (0.8667–1.000)	1.000 (1.000–1.000)
			F1	0.0000 (0.0000–0.0000)			0.09091 (0.0000–0.2857)	0.08696 (0.0000–0.2727)	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)
			Cond. Dice	-			Not supported	0.6548 (0.6548–0.6548)	Not supported	-
External-unseen (fine-tuning)	PANORAMA	pancreatic ductal adenocarcinoma	Sens.	0.8333 (0.7376–0.9143)	0.6667 (0.5556–0.7742)	0.6970 (0.5857–0.7937)	Not supported	0.8485 (0.7586–0.9298)	Not supported	0.8636 (0.7750–0.9444)
			Spec.	0.1883 (0.1250–0.2483)	0.0000 (0.0000–0.0000)	0.2208 (0.1554–0.2875)		0.7532 (0.6818–0.8258)		0.8442 (0.7785–0.8968)
			F1	0.4472 (0.3739–0.5152)	0.3333 (0.2609–0.4000)	0.3966 (0.3196–0.4706)		0.7000 (0.6182–0.7791)		0.7755 (0.6880–0.8400)
			Cond. Dice	0.2283 (0.1588–0.3044)	0.4000 (0.3244–0.4844)	Not supported		0.5376 (0.4840–0.6082)		0.5177 (0.4665–0.5843)

Supplementary Table 9 (continued) | Detailed sensitivity, specificity, F1, and conditional Dice scores of HoloCT and competing methods on internal, external-seen zero-shot/fine-tuning, external-unseen zero-shot/fine-tuning, and private zero-shot/fine-tuning classification-segmentation benchmarks. Each score is reported as mean with 95% confidence interval. Cond. Dice denotes conditional Dice. For the external zero-shot Macro F1 summary in Fig. 3c, unsupported ResNet and nnU-Net targets are scored as zero, so every model is averaged across the same nine targets.

Dataset Type	Dataset Name	Label Name	Metric	M3D-Phi3-4B	nnU-Net	ResNet	HoloCT-4B w/o grounding	HoloCT-4B w/ grounding	HoloCT-8B w/o grounding	HoloCT-8B w/ grounding
External-unseen (fine-tuning)	MHB-SEG-multiclass	epidural hemorrhage	Sens.	0.8000 (0.3333–1.000)	0.8000 (0.3333–1.000)	0.2000 (0.0000–0.6667)	Not supported	0.4000 (0.0000–0.8857)	Not supported	0.6000 (0.0000–1.000)
			Spec.	0.5116 (0.3571–0.6591)	0.7442 (0.6000–0.8667)	0.9302 (0.8409–1.000)		0.9535 (0.8810–1.000)		0.9070 (0.8140–0.9773)
			F1	0.2667 (0.06897–0.4707)	0.4000 (0.1111–0.6401)	0.2222 (0.0000–0.6000)		0.4444 (0.0000–0.8000)		0.5000 (0.0000–0.8235)
			Cond. Dice	0.08540 (0.02290–0.1635)	0.5202 (0.1516–0.8571)	Not supported		0.2678 (0.1545–0.3922)		0.3043 (0.1908–0.4269)
		subdural hemorrhage	Sens.	0.9000 (0.7368–1.000)	1.000 (1.000–1.000)	0.7500 (0.5293–0.9412)	Not supported	0.9000 (0.7497–1.000)	Not supported	0.9500 (0.8235–1.000)
			Spec.	0.1429 (0.03125–0.2800)	0.8214 (0.6774–0.9600)	0.1786 (0.03996–0.3333)		0.9286 (0.8148–1.000)		0.8929 (0.7692–1.000)
			F1	0.5806 (0.4209–0.7213)	0.8889 (0.7726–0.9756)	0.5172 (0.3333–0.6667)		0.9000 (0.7742–0.9778)		0.9048 (0.7878–0.9804)
			Cond. Dice	0.1231 (0.06690–0.2047)	0.7027 (0.5860–0.8007)	Not supported		0.5851 (0.4909–0.6855)		0.6043 (0.5220–0.6809)
		subarachnoid hemorrhage	Sens.	0.9000 (0.7368–1.000)	0.9500 (0.8333–1.000)	0.9000 (0.7368–1.000)	Not supported	0.9500 (0.8400–1.000)	Not supported	0.9500 (0.8421–1.000)
			Spec.	0.1429 (0.03125–0.2800)	0.8929 (0.7500–1.000)	0.2143 (0.06897–0.3667)		0.9286 (0.8213–1.000)		0.9643 (0.8750–1.000)
			F1	0.5806 (0.4209–0.7213)	0.9048 (0.7826–0.9804)	0.6000 (0.4313–0.7385)		0.9268 (0.8333–1.000)		0.9500 (0.8649–1.000)
			Cond. Dice	0.1073 (0.05880–0.1898)	0.6375 (0.5301–0.7286)	Not supported		0.3712 (0.2888–0.4706)		0.4785 (0.3952–0.5697)
		intraparenchymal hemorrhage	Sens.	0.6500 (0.4118–0.8637)	0.8500 (0.6814–1.000)	0.6500 (0.4118–0.8637)	Not supported	0.8500 (0.6842–1.000)	Not supported	0.9000 (0.7500–1.000)
			Spec.	0.3571 (0.1786–0.5358)	0.6786 (0.5000–0.8463)	0.6071 (0.4074–0.8000)		0.7500 (0.5713–0.9000)		0.7143 (0.5357–0.8800)
			F1	0.5098 (0.3254–0.6667)	0.7391 (0.5714–0.8772)	0.5909 (0.4000–0.7500)		0.7727 (0.6109–0.9020)		0.7826 (0.6285–0.8980)
			Cond. Dice	0.08500 (0.03620–0.1527)	0.4724 (0.3697–0.5586)	Not supported		0.2283 (0.1485–0.3382)		0.2608 (0.1773–0.3676)
		intraventricular hemorrhage	Sens.	0.9048 (0.7500–1.000)	0.9524 (0.8333–1.000)	0.7143 (0.4814–0.9048)	Not supported	0.9048 (0.7500–1.000)	Not supported	0.9524 (0.8333–1.000)
			Spec.	0.1481 (0.03226–0.2917)	0.7778 (0.6154–0.9201)	0.2593 (0.1034–0.4287)		0.8889 (0.7500–1.000)		0.8519 (0.7083–0.9655)
			F1	0.6032 (0.4444–0.7324)	0.8511 (0.7142–0.9434)	0.5357 (0.3555–0.6875)		0.8837 (0.7618–0.9730)		0.8889 (0.7778–0.9767)
			Cond. Dice	0.1151 (0.05780–0.1962)	0.4493 (0.3364–0.5572)	Not supported		0.4145 (0.3327–0.5063)		0.5169 (0.4371–0.6078)
Private (zero-shot)	Private03 (thyroid cancer-multi)	malignant thyroid tumor	Sens.	Not supported	Not supported	Not supported	Not supported	0.5263 (0.3823–0.6924)	Not supported	0.5526 (0.4000–0.7106)
			Spec.					0.5922 (0.4949–0.6789)		0.6019 (0.5052–0.6931)
			F1					0.4000 (0.2782–0.5099)		0.4200 (0.2970–0.5358)
			Cond. Dice					0.1403 (0.05336–0.2679)		0.1422 (0.03349–0.2738)
		benign thyroid tumor	Sens.	Not supported	Not supported	Not supported	Not supported	0.3838 (0.2941–0.4763)	Not supported	0.4040 (0.3157–0.4952)
			Spec.					0.6429 (0.4998–0.7778)		0.6429 (0.5000–0.7839)
			F1					0.5000 (0.4028–0.5898)		0.5195 (0.4246–0.6072)
			Cond. Dice					0.6429 (0.5881–0.6942)		0.6397 (0.5802–0.6991)
	Private03 (thyroid cancer-one)	thyroid tumor	Sens.	Not supported	Not supported	Not supported	Not supported	0.3578 (0.2752–0.4415)	Not supported	0.3761 (0.2909–0.4602)
			Spec.					0.5938 (0.4194–0.7667)		0.5938 (0.4286–0.7586)
			F1					0.4845 (0.3875–0.5715)		0.5031 (0.4103–0.5883)
			Cond. Dice					0.6475 (0.5948–0.7016)		0.6671 (0.6101–0.7195)
	Private04(colon lesion)	colon lesion	Sens.	Not supported	Not supported	Not supported	Not supported	0.4211 (0.3182–0.5196)	Not supported	0.4421 (0.3370–0.5422)
			Spec.					0.3803 (0.2666–0.5000)		0.3944 (0.2881–0.5132)
			F1					0.4469 (0.3432–0.5326)		0.4667 (0.3678–0.5510)
			Cond. Dice					Not supported		Not supported

Supplementary Table 9 (continued) | Detailed sensitivity, specificity, F1, and conditional Dice scores of HoloCT and competing methods on internal, external-seen zero-shot/fine-tuning, external-unseen zero-shot/fine-tuning, and private zero-shot/fine-tuning classification-segmentation benchmarks. Each score is reported as mean with 95% confidence interval. Cond. Dice denotes conditional Dice. For the external zero-shot Macro F1 summary in Fig. 3c, unsupported ResNet and nnU-Net targets are scored as zero, so every model is averaged across the same nine targets.

Dataset Type	Dataset Name	Label Name	Metric	M3D-Phi3-4B	nnU-Net	ResNet	HoloCT-4B w/o grounding	HoloCT-4B w/ grounding	HoloCT-8B w/o grounding	HoloCT-8B w/ grounding
Private (fine-tuning)	Private03 (thyroid cancer-multi)	malignant thyroid tumor	Sens.	Not supported	0.5789 (0.4250–0.7353)	0.6316 (0.4736–0.7813)	Not supported	0.6842 (0.5313–0.8205)	Not supported	0.7105 (0.5581–0.8571)
			Spec.		0.6117 (0.5200–0.7113)	0.6602 (0.5714–0.7522)		0.7087 (0.6176–0.7925)		0.7184 (0.6277–0.8036)
			F1		0.4400 (0.3146–0.5631)	0.4948 (0.3636–0.6111)		0.5532 (0.4186–0.6729)		0.5745 (0.4565–0.6838)
			Cond. Dice		0.2338 (0.1930–0.2796)	Not supported		0.2137 (0.1270–0.3335)		0.2009 (0.1283–0.3145)
		benign thyroid tumor	Sens.	Not supported	0.4141 (0.3178–0.5104)	0.4545 (0.3516–0.5521)	Not supported	0.5354 (0.4343–0.6346)	Not supported	0.5455 (0.4454–0.6429)
			Spec.		0.6667 (0.5143–0.8096)	0.7143 (0.5713–0.8445)		0.6429 (0.5000–0.7857)		0.6667 (0.5143–0.8095)
			F1		0.5325 (0.4354–0.6216)	0.5769 (0.4755–0.6667)		0.6347 (0.5432–0.7135)		0.6467 (0.5584–0.7232)
			Cond. Dice		0.6012 (0.5573–0.6621)	Not supported		0.6518 (0.5906–0.7297)		0.7152 (0.6650–0.7737)
	Private03 (thyroid cancer-one)	thyroid tumor	Sens.	Not supported	0.3853 (0.2952–0.4775)	0.4220 (0.3229–0.5176)	Not supported	0.8440 (0.7706–0.9107)	Not supported	0.8624 (0.7909–0.9238)
			Spec.		0.6250 (0.4500–0.7931)	0.6875 (0.5000–0.8485)		0.6875 (0.5294–0.8378)		0.6563 (0.5000–0.8236)
			F1		0.5153 (0.4161–0.6057)	0.5576 (0.4557–0.6437)		0.8720 (0.8200–0.9182)		0.8785 (0.8293–0.9174)
			Cond. Dice		0.6114 (0.5725–0.6887)	Not supported		0.6592 (0.6044–0.7289)		0.7197 (0.6718–0.7792)
	Private04(colon lesion)	colon lesion	Sens.	Not supported	0.4526 (0.3535–0.5474)	0.5263 (0.4285–0.6250)	0.9053 (0.8421–0.9588)	Not supported	0.9263 (0.8687–0.9714)	Not supported
			Spec.		0.4085 (0.2957–0.5352)	0.4789 (0.3662–0.6057)	0.8310 (0.7424–0.9167)		0.8310 (0.7368–0.9067)	
			F1		0.4778 (0.3820–0.5672)	0.5495 (0.4578–0.6334)	0.8912 (0.8414–0.9340)		0.9026 (0.8528–0.9394)	
			Cond. Dice		Not supported	Not supported	Not supported		Not supported	

Supplementary Table 10 | Detailed sensitivity, specificity, F1, and conditional Dice scores of HoloCT and competing methods on internal, external-unseen, and private classification-segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. Cond. Dice denotes conditional Dice.

Dataset Type	Dataset Name	Label Name	Metric	nnU-Net	ResNet	HoloCT-4B w/o grounding	HoloCT-4B w/ grounding	HoloCT-8B w/o grounding	HoloCT-8B w/ grounding
Internal	AbdomenAtlas 3.0	kidney lesion	Sens.	0.6450 (0.5893–0.7020)	0.7786 (0.7210–0.8296)	0.4809 (0.4196–0.5451)	0.6374 (0.5792–0.6935)	0.6489 (0.5891–0.7043)	0.5916 (0.5323–0.6531)
			Spec.	0.1792 (0.1503–0.2073)	0.6867 (0.6514–0.7218)	0.9593 (0.9445–0.9746)	0.9051 (0.8846–0.9259)	0.8690 (0.8422–0.8941)	0.9533 (0.9368–0.9682)
			F1	0.3463 (0.3083–0.3859)	0.6053 (0.5594–0.6455)	0.6072 (0.5436–0.6603)	0.6789 (0.6290–0.7252)	0.6551 (0.6067–0.7002)	0.6920 (0.6380–0.7389)
			Cond. Dice	0.1775 (0.09018–0.2727)	Not supported	Not supported	0.6958 (0.6515–0.7358)	Not supported	0.7600 (0.7281–0.7905)
		liver lesion	Sens.	0.8565 (0.8082–0.9032)	0.3722 (0.3100–0.4398)	0.6861 (0.6232–0.7456)	0.7668 (0.7101–0.8211)	0.7085 (0.6403–0.7663)	0.7668 (0.7091–0.8170)
			Spec.	0.02134 (0.01131–0.03277)	0.8478 (0.8214–0.8750)	0.9516 (0.9343–0.9653)	0.9175 (0.8950–0.9380)	0.9189 (0.8975–0.9379)	0.9161 (0.8956–0.9365)
			F1	0.3466 (0.3114–0.3826)	0.4019 (0.3416–0.4628)	0.7463 (0.6966–0.7937)	0.7566 (0.7113–0.7983)	0.7215 (0.6697–0.7664)	0.7550 (0.7107–0.7965)
			Cond. Dice	0.1392 (0.03089–0.2260)	Not supported	Not supported	0.7294 (0.7031–0.7527)	Not supported	0.6964 (0.6638–0.7267)
		pancreas lesion	Sens.	0.4964 (0.4077–0.5781)	0.4088 (0.3333–0.5000)	0.2920 (0.2170–0.3723)	0.4672 (0.3788–0.5493)	0.2263 (0.1571–0.2986)	0.6642 (0.5789–0.7419)
			Spec.	0.3992 (0.3677–0.4320)	0.7668 (0.7396–0.7959)	1.000 (1.000–1.000)	0.9861 (0.9781–0.9934)	0.9975 (0.9936–1.000)	0.9341 (0.9161–0.9506)
			F1	0.2003 (0.1570–0.2399)	0.2971 (0.2372–0.3609)	0.4520 (0.3567–0.5426)	0.6038 (0.5171–0.6783)	0.3647 (0.2709–0.4531)	0.6500 (0.5789–0.7132)
			Cond. Dice	0.1421 (0.01590–0.2152)	Not supported	Not supported	0.5632 (0.5086–0.6142)	Not supported	0.5514 (0.5014–0.5962)
	LiMT	colon lesion	Sens.	0.8000 (0.6154–0.9524)	0.6800 (0.5000–0.8571)	0.4000 (0.2257–0.5806)	0.7200 (0.5332–0.8929)	0.5200 (0.3181–0.7143)	0.6800 (0.5000–0.8571)
			Spec.	0.04440 (0.03182–0.05778)	0.03885 (0.02655–0.05240)	0.9933 (0.9878–0.9978)	0.9756 (0.9653–0.9855)	0.9889 (0.9811–0.9955)	0.9778 (0.9676–0.9867)
			F1	0.04415 (0.02667–0.06339)	0.03744 (0.02220–0.05507)	0.4878 (0.2939–0.6667)	0.5538 (0.3729–0.6970)	0.5417 (0.3635–0.6977)	0.5484 (0.3793–0.7000)
			Cond. Dice	0.06071 (0.009609–0.1507)	Not supported	Not supported	0.6472 (0.5216–0.7513)	Not supported	0.7291 (0.6624–0.7895)
		liver cyst	Sens.	0.6000 (0.2857–0.9000)	0.0000 (0.0000–0.0000)	0.7000 (0.4000–1.000)	0.7000 (0.4000–1.000)	0.3000 (0.0000–0.6000)	0.6000 (0.2857–0.8892)
			Spec.	0.6000 (0.3810–0.8125)	0.9000 (0.7647–1.000)	0.9000 (0.7616–1.000)	0.8500 (0.6957–1.000)	0.9000 (0.7500–1.000)	0.9000 (0.7500–1.000)
			F1	0.5000 (0.2105–0.7200)	0.0000 (0.0000–0.0000)	0.7368 (0.4444–0.9333)	0.7000 (0.4286–0.9000)	0.4000 (0.0000–0.6667)	0.6667 (0.3636–0.8800)
			Cond. Dice	0.5755 (0.5165–0.6487)	Not supported	Not supported	0.5934 (0.4864–0.6965)	Not supported	0.6870 (0.5900–0.7691)
		liver hemangioma	Sens.	0.6667 (0.2000–1.000)	0.5000 (0.0000–1.000)	0.6667 (0.2000–1.000)	0.6667 (0.2500–1.000)	1.000 (1.000–1.000)	1.000 (1.000–1.000)
			Spec.	0.7083 (0.5200–0.8750)	0.6667 (0.4783–0.8462)	1.000 (1.000–1.000)	1.000 (1.000–1.000)	0.9583 (0.8636–1.000)	0.9583 (0.8571–1.000)
			F1	0.4706 (0.1426–0.7202)	0.3529 (0.0000–0.6316)	0.8000 (0.3333–1.000)	0.8000 (0.4000–1.000)	0.9231 (0.6667–1.000)	0.9231 (0.7143–1.000)
			Cond. Dice	0.5382 (0.4786–0.6393)	Not supported	Not supported	0.8327 (0.7710–0.8788)	Not supported	0.7170 (0.5129–0.8533)
		metastatic liver cancer	Sens.	0.5000 (0.0000–1.000)	0.2500 (0.0000–1.000)	0.5000 (0.0000–1.000)	0.5000 (0.0000–1.000)	0.5000 (0.0000–1.000)	0.5000 (0.0000–1.000)
			Spec.	0.6923 (0.5185–0.8571)	0.6923 (0.5000–0.8750)	1.000 (1.000–1.000)	1.000 (1.000–1.000)	0.9615 (0.8750–1.000)	0.9615 (0.8800–1.000)
			F1	0.2857 (0.0000–0.5885)	0.1538 (0.0000–0.4290)	0.6667 (0.0000–1.000)	0.6667 (0.0000–1.000)	0.5714 (0.0000–1.000)	0.5714 (0.0000–1.000)
			Cond. Dice	0.3953 (0.3170–0.4654)	Not supported	Not supported	0.5243 (0.4417–0.6070)	Not supported	0.6050 (0.5343–0.6758)
		primary liver cancer	Sens.	0.8571 (0.5000–1.000)	0.8571 (0.5000–1.000)	0.8571 (0.5714–1.000)	1.000 (1.000–1.000)	1.000 (1.000–1.000)	1.000 (1.000–1.000)
			Spec.	0.8696 (0.7143–1.000)	0.7826 (0.5998–0.9446)	1.000 (1.000–1.000)	1.000 (1.000–1.000)	0.9565 (0.8571–1.000)	0.9565 (0.8571–1.000)
			F1	0.7500 (0.4000–0.9413)	0.6667 (0.3525–0.8751)	0.9231 (0.7273–1.000)	1.000 (1.000–1.000)	0.9333 (0.7273–1.000)	0.9333 (0.7500–1.000)
			Cond. Dice	0.6659 (0.6137–0.7286)	Not supported	Not supported	0.7219 (0.5662–0.8506)	Not supported	0.6889 (0.4131–0.8564)

Supplementary Table 10 (continued) | Detailed sensitivity, specificity, F1, and conditional Dice scores of HoloCT and competing methods on internal, external-unseen, and private classification-segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. Cond. Dice denotes conditional Dice.

Dataset Type	Dataset Name	Label Name	Metric	nnU-Net	ResNet	HoloCT-4B w/o grounding	HoloCT-4B w/ grounding	HoloCT-8B w/o grounding	HoloCT-8B w/ grounding
External-unseen (zero-shot)	MHB-SEG-multiclass	epidural hemorrhage	Sens.	Not supported	Not supported	0.0000 (0.0000–0.0000)	0.6000 (0.0000–1.000)	0.0000 (0.0000–0.0000)	0.6000 (0.0000–1.000)
			Spec.			1.000 (1.000–1.000)	0.7674 (0.6409–0.8838)	1.000 (1.000–1.000)	0.8140 (0.6889–0.9131)
			F1			0.0000 (0.0000–0.0000)	0.3333 (0.0000–0.6000)	0.0000 (0.0000–0.0000)	0.3750 (0.0000–0.6316)
			Cond. Dice			Not supported	0.07251 (0.0000–0.1890)	Not supported	0.1517 (0.0000–0.4220)
		subdural hemorrhage	Sens.	Not supported	Not supported	0.5000 (0.2778–0.7200)	0.5500 (0.3182–0.7503)	0.6500 (0.4117–0.8423)	0.8000 (0.6111–0.9545)
			Spec.			0.8571 (0.7308–0.9655)	0.6429 (0.4545–0.8078)	0.5714 (0.3846–0.7576)	0.5357 (0.3448–0.7143)
			F1			0.5882 (0.3636–0.7692)	0.5366 (0.3333–0.6957)	0.5778 (0.3636–0.7318)	0.6531 (0.4762–0.7931)
			Cond. Dice			Not supported	0.5082 (0.2943–0.7023)	Not supported	0.4733 (0.3033–0.6348)
		subarachnoid hemorrhage	Sens.	Not supported	Not supported	0.0000 (0.0000–0.0000)	0.1000 (0.0000–0.2384)	0.1500 (0.0000–0.3333)	0.3000 (0.1175–0.5006)
			Spec.			0.9643 (0.8888–1.000)	1.000 (1.000–1.000)	1.000 (1.000–1.000)	0.9643 (0.8889–1.000)
			F1			0.0000 (0.0000–0.0000)	0.1818 (0.0000–0.3850)	0.2609 (0.0000–0.5000)	0.4444 (0.1998–0.6667)
			Cond. Dice			Not supported	0.0003018 (0.0000–0.0006036)	Not supported	0.4099 (0.1943–0.5847)
		intraparenchymal hemorrhage	Sens.	Not supported	Not supported	0.0000 (0.0000–0.0000)	0.05000 (0.0000–0.1739)	0.1000 (0.0000–0.2500)	0.2000 (0.04343–0.3750)
			Spec.			1.000 (1.000–1.000)	0.9643 (0.8788–1.000)	0.9286 (0.8181–1.000)	0.7857 (0.6154–0.9310)
			F1			0.0000 (0.0000–0.0000)	0.09091 (0.0000–0.2857)	0.1667 (0.0000–0.3706)	0.2667 (0.06452–0.4446)
			Cond. Dice			Not supported	0.1675 (0.1675–0.1675)	Not supported	0.09218 (0.0000–0.2251)
		intraventricular hemorrhage	Sens.	Not supported	Not supported	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)
			Spec.			1.000 (1.000–1.000)	1.000 (1.000–1.000)	1.000 (1.000–1.000)	1.000 (1.000–1.000)
			F1			0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)	0.0000 (0.0000–0.0000)
External-unseen (fine-tuning)	MHB-SEG-multiclass	epidural hemorrhage	Sens.	0.6000 (0.0000–1.000)	0.2000 (0.0000–0.6667)	Not supported	0.4000 (0.0000–1.000)	Not supported	0.6000 (0.0000–1.000)
			Spec.	0.6744 (0.5224–0.8050)	0.9070 (0.8049–0.9783)		0.9302 (0.8444–1.000)		0.8837 (0.7727–0.9744)
			F1	0.2727 (0.0000–0.5217)	0.2000 (0.0000–0.5455)		0.4000 (0.0000–0.7500)		0.4615 (0.0000–0.7692)
			Cond. Dice	0.4906 (0.4688–0.5457)	Not supported		0.2990 (0.1455–0.4622)		0.3046 (0.1443–0.4805)
		subdural hemorrhage	Sens.	0.8000 (0.6000–0.9524)	0.6000 (0.3683–0.8182)	Not supported	0.8500 (0.6818–1.000)	Not supported	0.9000 (0.7497–1.000)
			Spec.	0.6786 (0.5000–0.8485)	0.1429 (0.03226–0.2800)		0.8929 (0.7777–1.000)		0.8571 (0.7200–0.9667)
			F1	0.7111 (0.5263–0.8511)	0.4286 (0.2499–0.5862)		0.8500 (0.7179–0.9524)		0.8571 (0.7273–0.9545)
			Cond. Dice	0.6389 (0.5798–0.6881)	Not supported		0.5177 (0.3939–0.6513)		0.5340 (0.4023–0.6668)
		subarachnoid hemorrhage	Sens.	0.8500 (0.6814–1.000)	0.7500 (0.5293–0.9412)	Not supported	0.9000 (0.7500–1.000)	Not supported	0.9500 (0.8333–1.000)
			Spec.	0.7500 (0.5833–0.9091)	0.1429 (0.03125–0.2857)		0.8929 (0.7666–1.000)		0.9286 (0.8148–1.000)
			F1	0.7727 (0.6111–0.8980)	0.5085 (0.3273–0.6571)		0.8780 (0.7442–0.9730)		0.9268 (0.8292–1.000)
			Cond. Dice	0.5750 (0.5144–0.6449)	Not supported		0.3841 (0.2803–0.4974)		0.4089 (0.3092–0.5146)

Supplementary Table 10 (continued) | Detailed sensitivity, specificity, F1, and conditional Dice scores of HoloCT and competing methods on internal, external-unseen, and private classification-segmentation benchmarks.

Each score is reported as mean with 95% confidence interval. Cond. Dice denotes conditional Dice.

Dataset Type	Dataset Name	Label Name	Metric	nnU-Net	ResNet	HoloCT-4B w/o grounding	HoloCT-4B w/ grounding	HoloCT-8B w/o grounding	HoloCT-8B w/ grounding
External-unseen (fine-tuning)	MHB-SEG-multiclass	intraparenchymal hemorrhage	Sens.	0.7000 (0.4706–0.9000)	0.7000 (0.4706–0.9000)	Not supported	0.8500 (0.6818–1.000)	Not supported	0.9000 (0.7500–1.000)
			Spec.	0.5714 (0.3793–0.7407)	0.6429 (0.4544–0.8182)		0.7143 (0.5556–0.8710)		0.6786 (0.5000–0.8485)
			F1	0.6087 (0.4117–0.7600)	0.6364 (0.4499–0.7895)		0.7556 (0.5882–0.8780)		0.7660 (0.6190–0.8889)
			Cond. Dice	0.4188 (0.3323–0.4707)	Not supported		0.1604 (0.09130–0.2585)		0.1802 (0.1199–0.2719)
		intraventricular hemorrhage	Sens.	0.9524 (0.8333–1.000)	0.5714 (0.3529–0.7895)	Not supported	0.8571 (0.7058–1.000)	Not supported	0.9524 (0.8421–1.000)
			Spec.	0.8148 (0.6667–0.9583)	0.1852 (0.04343–0.3478)		0.8519 (0.7096–0.9643)		0.8148 (0.6667–0.9397)
			F1	0.8696 (0.7429–0.9600)	0.4364 (0.2609–0.6000)		0.8372 (0.7111–0.9412)		0.8696 (0.7500–0.9565)
			Cond. Dice	0.4707 (0.4162–0.5493)	Not supported		0.3595 (0.2569–0.4606)		0.3713 (0.2787–0.4859)
Private (fine-tuning)	Private03 (thyroid cancer-multi)	malignant thyroid tumor	Sens.	0.5000 (0.3448–0.6667)	0.5526 (0.3947–0.7105)	Not supported	0.6579 (0.5143–0.8065)	Not supported	0.6842 (0.5333–0.8333)
			Spec.	0.5631 (0.4666–0.6598)	0.5922 (0.4999–0.6882)		0.6699 (0.5818–0.7604)		0.6893 (0.5943–0.7822)
			F1	0.3725 (0.2471–0.4906)	0.4158 (0.2889–0.5377)		0.5155 (0.3953–0.6355)		0.5417 (0.4054–0.6542)
			Cond. Dice	0.2906 (0.1294–0.3550)	Not supported		0.2760 (0.1834–0.3895)		0.3155 (0.2262–0.4279)
		benign thyroid tumor	Sens.	0.3636 (0.2680–0.4600)	0.3939 (0.3000–0.4896)	Not supported	0.5859 (0.4888–0.6863)	Not supported	0.6061 (0.5102–0.7076)
			Spec.	0.5714 (0.4146–0.7250)	0.6190 (0.4706–0.7692)		0.6905 (0.5435–0.8298)		0.7381 (0.5937–0.8684)
			F1	0.4706 (0.3694–0.5679)	0.5065 (0.4082–0.5987)		0.6824 (0.5976–0.7609)		0.7059 (0.6173–0.7865)
			Cond. Dice	0.5451 (0.4786–0.6173)	Not supported		0.6932 (0.6231–0.7659)		0.7309 (0.6735–0.8079)

Supplementary Table 11 | Detailed closed-ended VQA performance of HoloCT and competing methods across internal benchmarks. Each score is reported as mean with 95% confidence interval. Accu. denotes accuracy. For DeepTumorVQA, Recog., Meas., Vis.R., and Med.R. denote recognition, measurement, visual reasoning, and medical reasoning, respectively. For AMOS-MM-VQA, Anat.I., Attr.R., Quant.A., Dis.A., Risk.E., and Ther.S. denote anatomical identification, attribute recognition, quantitative assessment, disease analysis, risk evaluation, and therapeutic suggestion, respectively.

Dataset Type	Dataset Name	Metric		Qwen3-VL-32B	MedGemma1.5-4B	M3D-Phi3-4B	RadFM-14B	HuluMed-32B	HoloCT-4B	HoloCT-8B
Internal	CT-RATE-option	Accuracy		0.6732 (0.6691–0.6867)	0.6117 (0.6029–0.6243)	0.6892 (0.6793–0.6988)	0.3234 (0.3168–0.3308)	0.8473 (0.8393–0.8545)	0.9264 (0.9202–0.9317)	0.9343 (0.9285–0.9395)
	FLARE-2025	Accuracy		0.5988 (0.5887–0.6063)	0.4944 (0.4812–0.5057)	0.5473 (0.5342–0.5592)	0.2767 (0.2675–0.2863)	0.7100 (0.6990–0.7208)	0.7630 (0.7519–0.7725)	0.7997 (0.7900–0.8095)
	3D-RAD Existence	Accuracy		0.6128 (0.6086–0.6299)	0.6874 (0.6781–0.6938)	0.4086 (0.3902–0.4269)	0.3098 (0.2865–0.3184)	0.8226 (0.8117–0.8396)	0.8662 (0.8619–0.8705)	0.8588 (0.8544–0.8631)
	3D-RAD Static.	Accuracy		0.2039 (0.1803–0.2252)	0.3805 (0.3651–0.4026)	0.2551 (0.2426–0.2620)	0.4331 (0.4323–0.4692)	0.6549 (0.6354–0.6737)	0.6947 (0.6780–0.7111)	0.6519 (0.6345–0.6686)
	3D-RAD Longitudinal.	Accuracy		0.4416 (0.4293–0.4662)	0.4287 (0.4060–0.4616)	0.2440 (0.2267–0.2744)	0.4390 (0.4014–0.4617)	0.8193 (0.7931–0.8304)	0.8493 (0.8364–0.8629)	0.8441 (0.8312–0.8583)
	AMOS-MM-VQA	Anat.I.	Accu.	0.4348 (0.3876–0.4740)	0.3832 (0.3205–0.4373)	0.4846 (0.4415–0.5298)	0.3478 (0.3150–0.3786)	0.7269 (0.6879–0.7639)	0.6923 (0.5577–0.8077)	0.7692 (0.6538–0.8846)
		Attr.R.	Accu.	0.4951 (0.4546–0.5332)	0.4458 (0.3940–0.4804)	0.5745 (0.5350–0.6158)	0.4077 (0.3808–0.4329)	0.8600 (0.8312–0.8887)	0.9286 (0.8333–1.000)	1.000 (1.000–1.000)
		Quant.A.	Accu.	0.3777 (0.3189–0.4260)	0.3231 (0.2540–0.3830)	0.4711 (0.4309–0.5113)	0.4479 (0.4206–0.4753)	0.7492 (0.7154–0.7830)	0.7097 (0.5968–0.8226)	0.9032 (0.8226–0.9677)
		Dis.A.	Accu.	0.4062 (0.3560–0.4692)	0.3530 (0.2905–0.4121)	0.4920 (0.4462–0.5424)	0.4053 (0.3755–0.4328)	0.7506 (0.7071–0.7918)	0.8056 (0.6944–0.9167)	0.9167 (0.8056–1.000)
		Risk.E.	Accu.	0.3447 (0.2828–0.3949)	0.3957 (0.3277–0.4542)	0.4537 (0.4073–0.5000)	0.4024 (0.3707–0.4341)	0.7195 (0.6805–0.7610)	0.7561 (0.6341–0.8780)	0.8780 (0.7561–0.9756)
		Ther.S.	Accu.	0.3851 (0.3223–0.4321)	0.4491 (0.3786–0.5062)	0.4124 (0.3503–0.4708)	0.3511 (0.3168–0.3803)	0.8212 (0.7737–0.8650)	0.6333 (0.4667–0.8000)	0.7333 (0.5667–0.8667)
		Recog.	Accu.	0.4557 (0.4365–0.4785)	0.4025 (0.3858–0.4247)	0.4205 (0.3996–0.4404)	0.4309 (0.4098–0.4541)	0.5522 (0.5304–0.5734)	0.7121 (0.6800–0.7409)	0.8128 (0.7852–0.8394)
	DeepTumorVQA	Meas.	Accu.	0.2041 (0.1841–0.2280)	0.1853 (0.1683–0.2038)	0.1706 (0.1547–0.1874)	0.2148 (0.1963–0.2344)	0.2951 (0.2748–0.3159)	0.5686 (0.5421–0.5967)	0.5934 (0.5653–0.6215)
		Vis.R.	Accu.	0.3191 (0.2993–0.3211)	0.2806 (0.2732–0.3028)	0.2907 (0.2768–0.3036)	0.2621 (0.2479–0.2768)	0.4149 (0.3999–0.4302)	0.5812 (0.5610–0.6022)	0.5944 (0.5738–0.6155)
		Med.R.	Accu.	0.2715 (0.2522–0.2980)	0.2448 (0.2266–0.2620)	0.2583 (0.2378–0.2772)	0.2032 (0.1849–0.2222)	0.3318 (0.3108–0.3528)	0.6168 (0.5888–0.6431)	0.6752 (0.6480–0.7015)

Supplementary Table 12 | Detailed open-ended VQA performance of HoloCT and competing methods on internal and external benchmarks. Each score is reported as mean with 95% confidence interval. NLP metrics include BLEU-4, ROUGE-L, and METEOR; the clinical metric is LlamaScore.

Dataset Type	Dataset Name	Metric		Qwen3-VL-32B	MedGemma1.5-4B	M3D-Phi3-4B	RadFM-14B	HuluMed-32B	HoloCT-4B	HoloCT-8B
Internal	CT-RATE-long	NLP	BLEU-4	0.3328 (0.3255–0.3489)	0.2093 (0.1945–0.2181)	0.09760 (0.09634–0.09893)	0.2799 (0.2780–0.2820)	0.3886 (0.3863–0.3908)	0.4928 (0.4904–0.4953)	0.4917 (0.4892–0.4941)
			ROUGE-L	0.3534 (0.3444–0.3692)	0.2257 (0.2117–0.2339)	0.1315 (0.1303–0.1328)	0.2742 (0.2724–0.2761)	0.4122 (0.4102–0.4143)	0.5007 (0.4982–0.5032)	0.4952 (0.4926–0.4978)
			METEOR	0.3682 (0.3566–0.3762)	0.1927 (0.1820–0.2881)	0.08947 (0.08840–0.09045)	0.2803 (0.2782–0.2824)	0.4293 (0.4271–0.4316)	0.5199 (0.5169–0.5227)	0.5208 (0.5181–0.5237)
		Clinical	Llama Score	6.089 (6.054–6.116)	5.127 (5.095–5.152)	4.920 (4.898–4.944)	5.427 (5.401–5.452)	6.873 (6.852–6.895)	7.544 (7.525–7.560)	7.583 (7.565–7.601)
	CT-RATE-short	NLP	BLEU-4	0.1528 (0.1441–0.1691)	0.1272 (0.1121–0.1332)	0.1637 (0.1593–0.1677)	0.09063 (0.08879–0.09251)	0.07747 (0.07611–0.07882)	0.5991 (0.5925–0.6055)	0.5964 (0.5896–0.6029)
			ROUGE-L	0.1949 (0.1885–0.2358)	0.1391 (0.1226–0.1479)	0.2165 (0.2064–0.2263)	0.08163 (0.07888–0.08467)	0.1080 (0.1060–0.1100)	0.5961 (0.5893–0.6030)	0.5940 (0.5866–0.6012)
			METEOR	0.1514 (0.1462–0.1683)	0.1252 (0.1126–0.1375)	0.09350 (0.08951–0.09728)	0.1012 (0.09870–0.1036)	0.1667 (0.1639–0.1698)	0.5108 (0.5043–0.5171)	0.4981 (0.4916–0.5044)
		Clinical	Llama Score	5.248 (5.196–5.339)	4.686 (4.625–4.748)	4.346 (4.249–4.445)	4.518 (4.465–4.570)	5.919 (5.864–5.971)	7.399 (7.348–7.449)	7.442 (7.390–7.492)
	3D-RAD Anomaly Detection	NLP	BLEU-4	0.08660 (0.06120–0.1018)	0.1047 (0.08640–0.1213)	0.1672 (0.1489–0.1717)	0.1239 (0.09490–0.1343)	0.2952 (0.2646–0.3224)	0.5280 (0.5019–0.5549)	0.4967 (0.4639–0.5286)
			ROUGE-L	0.1411 (0.1259–0.1626)	0.1660 (0.1422–0.1861)	0.2330 (0.2109–0.2555)	0.1859 (0.1709–0.2056)	0.4181 (0.3836–0.4343)	0.6214 (0.5977–0.6446)	0.6206 (0.5961–0.6445)
			METEOR	0.1110 (0.09920–0.1386)	0.1309 (0.1170–0.1560)	0.2033 (0.1894–0.2213)	0.1551 (0.1368–0.1629)	0.3595 (0.3285–0.3881)	0.5879 (0.5657–0.6080)	0.5748 (0.5514–0.5955)
		Clinical	Llama Score	4.984 (4.770–5.175)	5.116 (4.914–5.320)	5.513 (5.336–5.689)	5.238 (5.049–5.434)	6.448 (6.258–6.641)	7.746 (7.570–7.918)	8.016 (7.853–8.166)
	3D-RAD Image Observation	NLP	BLEU-4	0.1069 (0.08650–0.1239)	0.1256 (0.1037–0.1472)	0.1667 (0.1493–0.1810)	0.1391 (0.1260–0.1539)	0.2716 (0.2562–0.3041)	0.5209 (0.5011–0.5398)	0.5159 (0.4954–0.5360)
			ROUGE-L	0.1673 (0.1451–0.1812)	0.1859 (0.1606–0.2025)	0.2477 (0.2225–0.2518)	0.2054 (0.1839–0.2106)	0.4026 (0.3864–0.4251)	0.6126 (0.5976–0.6264)	0.6168 (0.6020–0.6305)
			METEOR	0.1341 (0.1140–0.1513)	0.1546 (0.1360–0.1760)	0.2038 (0.1811–0.2209)	0.1608 (0.1415–0.1802)	0.3439 (0.3217–0.3672)	0.5731 (0.5593–0.5867)	0.5734 (0.5589–0.5868)
		Clinical	Llama Score	4.941 (4.804–5.077)	5.064 (4.936–5.201)	5.376 (5.254–5.498)	5.143 (5.017–5.266)	6.200 (6.069–6.327)	7.449 (7.341–7.560)	7.791 (7.681–7.887)
	3D-RAD Medical Computation	NLP	BLEU-4	0.07190 (0.05720–0.09230)	0.1011 (0.08330–0.1325)	0.03370 (0.01180–0.04810)	0.04690 (0.03950–0.05250)	0.3991 (0.3551–0.4335)	0.2171 (0.1933–0.2411)	0.1724 (0.1561–0.1918)
			ROUGE-L	0.1050 (0.08920–0.1346)	0.1575 (0.1323–0.1839)	0.05810 (0.04280–0.07380)	0.06930 (0.06420–0.07460)	0.4099 (0.3772–0.4361)	0.6256 (0.6033–0.6466)	0.6414 (0.6196–0.6643)
			METEOR	0.09060 (0.07320–0.1141)	0.1363 (0.1188–0.1613)	0.05560 (0.04080–0.07190)	0.05400 (0.04150–0.07150)	0.3921 (0.3654–0.4229)	0.5122 (0.4885–0.5341)	0.5695 (0.5474–0.5918)
		Clinical	Llama Score	3.920 (3.682–4.144)	4.128 (3.902–4.332)	3.731 (3.473–3.975)	3.787 (3.535–4.010)	5.414 (5.258–5.576)	7.688 (7.518–7.849)	8.110 (7.949–8.266)
External (zero-shot)	BIMCV-VQA	NLP	BLEU-4	0.1952 (0.1887–0.2763)	0.1441 (0.1397–0.1581)	0.1227 (0.1197–0.1259)	0.2114 (0.2067–0.2163)	0.2817 (0.2763–0.2870)	0.2817 (0.2762–0.2874)	0.2749 (0.2695–0.2802)
			ROUGE-L	0.2198 (0.2148–0.2259)	0.1564 (0.1430–0.1617)	0.1162 (0.1131–0.1194)	0.2332 (0.2289–0.2375)	0.2993 (0.2941–0.3050)	0.2644 (0.2592–0.2694)	0.2628 (0.2577–0.2678)
			METEOR	0.1880 (0.1769–0.1934)	0.1224 (0.1194–0.1352)	0.09925 (0.09661–0.1019)	0.2135 (0.2088–0.2184)	0.2969 (0.2919–0.3019)	0.2639 (0.2586–0.2694)	0.2603 (0.2548–0.2657)
		Clinical	Llama Score	4.551 (4.482–4.622)	4.026 (3.957–4.096)	3.496 (3.429–3.565)	4.197 (4.130–4.264)	4.870 (4.795–4.940)	5.238 (5.163–5.310)	5.357 (5.281–5.426)

Supplementary Table 13 | Detailed narrative report generation performance of HoloCT and competing methods on internal and external zero-shot benchmarks.

Each score is reported as mean with 95% confidence interval. NLP metrics include BLEU-4, ROUGE-L, and METEOR; clinical metrics include RaTEScore, LlamaScore, and GREEN.

Dataset Type	Dataset Name	Metric		Qwen3-VL-32B	MedGemma1.5-4B	M3D-Phi3-4B	RadFM-14B	HuluMed-32B	HoloCT-4B	HoloCT-8B
Internal	CT-RATE-RG	NLP	BLEU-4	0.08220 (0.05110–0.1099)	0.2319 (0.2033–0.2562)	0.02401 (0.02178–0.02628)	0.03029 (0.02762–0.03296)	0.5086 (0.5027–0.5140)	0.5496 (0.5438–0.5556)	0.5481 (0.5424–0.5535)
			ROUGE-L	0.1087 (0.07490–0.1307)	0.1271 (0.09500–0.1513)	0.05858 (0.05717–0.05994)	0.06645 (0.06472–0.06807)	0.3243 (0.3195–0.3295)	0.3848 (0.3785–0.3913)	0.3579 (0.3524–0.3637)
			METEOR	0.09690 (0.06840–0.1237)	0.1207 (0.09770–0.1456)	0.03613 (0.03496–0.03722)	0.04624 (0.04467–0.04777)	0.3731 (0.3683–0.3785)	0.4175 (0.4117–0.4233)	0.4029 (0.3978–0.4076)
		Clinical	RaTE Score	0.4285 (0.4027–0.4442)	0.4571 (0.4390–0.4761)	0.3642 (0.3611–0.3670)	0.3905 (0.3876–0.3935)	0.6384 (0.6353–0.6415)	0.6620 (0.6581–0.6655)	0.6496 (0.6466–0.6527)
			Llama Score	2.434 (2.316–2.549)	4.059 (3.934–4.160)	1.067 (1.025–1.110)	1.794 (1.744–1.843)	6.433 (6.418–6.523)	6.571 (6.506–6.631)	6.430 (6.368–6.494)
			GREEN	0.05540 (0.03650–0.08190)	0.2854 (0.2534–0.3165)	0.005301 (0.004383–0.006346)	0.01819 (0.01614–0.02020)	0.4060 (0.3985–0.4129)	0.4372 (0.4283–0.4462)	0.4197 (0.4115–0.4274)
	AbdomenAtlas3.0	NLP	BLEU-4	0.1524 (0.1106–0.1885)	0.1760 (0.1450–0.2054)	0.07939 (0.07291–0.08591)	0.1253 (0.1174–0.1332)	0.2825 (0.2768–0.2883)	0.5051 (0.4922–0.5185)	0.5704 (0.5593–0.5813)
			ROUGE-L	0.1266 (0.09610–0.1548)	0.1357 (0.1017–0.1613)	0.09772 (0.09481–0.1003)	0.1142 (0.1110–0.1174)	0.1596 (0.1572–0.1621)	0.4305 (0.4219–0.4394)	0.4580 (0.4495–0.4664)
			METEOR	0.1065 (0.07780–0.1322)	0.1161 (0.08480–0.1409)	0.06093 (0.05854–0.06363)	0.08546 (0.08214–0.08878)	0.1698 (0.1661–0.1735)	0.4011 (0.3918–0.4106)	0.4731 (0.4641–0.4825)
		Clinical	RaTE Score	0.3929 (0.3682–0.4242)	0.4066 (0.3790–0.4369)	0.3763 (0.3710–0.3815)	0.3533 (0.3467–0.3600)	0.4602 (0.4528–0.4674)	0.7606 (0.7540–0.7665)	0.7670 (0.7611–0.7729)
			Llama Score	0.6851 (0.5222–0.8501)	0.8290 (0.6778–0.9971)	0.4582 (0.4006–0.5157)	0.2182 (0.1801–0.2610)	1.554 (1.434–1.678)	4.808 (4.711–4.914)	4.663 (4.562–4.774)
			GREEN	0.03510 (0.01880–0.06010)	0.04510 (0.02780–0.07430)	0.008943 (0.006399–0.01175)	0.01228 (0.008561–0.01621)	0.1281 (0.1127–0.1446)	0.3107 (0.2945–0.3254)	0.3087 (0.2953–0.3216)
	INSPECT	NLP	BLEU-4	0.1460 (0.1280–0.1609)	0.1584 (0.1365–0.1742)	0.1126 (0.1083–0.1172)	0.1433 (0.1385–0.1485)	0.1562 (0.1516–0.1609)	0.2609 (0.2532–0.2693)	0.2908 (0.2831–0.2992)
			ROUGE-L	0.1045 (0.08410–0.1252)	0.1111 (0.09580–0.1383)	0.08649 (0.08468–0.08812)	0.1029 (0.1008–0.1047)	0.09737 (0.09589–0.09874)	0.2749 (0.2688–0.2812)	0.2789 (0.2728–0.2857)
			METEOR	0.1213 (0.1087–0.1401)	0.1366 (0.1148–0.1533)	0.08907 (0.08669–0.09144)	0.1135 (0.1113–0.1160)	0.1688 (0.1671–0.1708)	0.2931 (0.2858–0.3000)	0.3151 (0.3074–0.3221)
		Clinical	RaTE Score	0.3516 (0.3358–0.3781)	0.3624 (0.3454–0.3873)	0.2967 (0.2929–0.3007)	0.3430 (0.3394–0.3468)	0.3797 (0.3763–0.3827)	0.5062 (0.4998–0.5126)	0.4942 (0.4882–0.5001)
			Llama Score	2.055 (1.957–2.167)	2.402 (2.290–2.513)	1.072 (1.012–1.130)	1.508 (1.446–1.569)	4.551 (4.473–4.624)	5.208 (5.121–5.297)	5.079 (4.982–5.172)
			GREEN	0.02900 (0.01610–0.04120)	0.03810 (0.01850–0.05900)	0.005848 (0.004210–0.007808)	0.007954 (0.006588–0.009438)	0.03857 (0.03382–0.04414)	0.2247 (0.2173–0.2326)	0.2163 (0.2091–0.2246)
External (zero-shot)	BIMCV-RG	NLP	BLEU-4	0.1560 (0.1201–0.1842)	0.1769 (0.1425–0.2018)	0.1286 (0.1218–0.1356)	0.1348 (0.1275–0.1423)	0.2647 (0.2566–0.2716)	0.1933 (0.1839–0.2026)	0.1982 (0.1887–0.2072)
			ROUGE-L	0.1245 (0.09170–0.1465)	0.1352 (0.1066–0.1577)	0.1069 (0.1041–0.1095)	0.1147 (0.1110–0.1180)	0.1314 (0.1291–0.1337)	0.1706 (0.1661–0.1745)	0.1681 (0.1642–0.1719)
			METEOR	0.1241 (0.09290–0.1563)	0.1451 (0.1123–0.1734)	0.09245 (0.08954–0.09556)	0.1098 (0.1053–0.1144)	0.2042 (0.2011–0.2074)	0.1662 (0.1606–0.1716)	0.1707 (0.1658–0.1753)
		Clinical	RaTE Score	0.3676 (0.3461–0.3966)	0.3877 (0.3622–0.4166)	0.3421 (0.3374–0.3464)	0.3428 (0.3374–0.3482)	0.4454 (0.4414–0.4495)	0.4601 (0.4536–0.4661)	0.4827 (0.4763–0.4889)
			Llama Score	1.853 (1.726–1.996)	2.209 (2.068–2.350)	1.268 (1.191–1.349)	1.189 (1.125–1.248)	4.006 (3.898–4.112)	4.253 (4.136–4.369)	4.069 (3.945–4.184)
			GREEN	0.03550 (0.01450–0.05840)	0.04150 (0.02710–0.05920)	0.01097 (0.008727–0.01360)	0.01595 (0.01312–0.01920)	0.09048 (0.08202–0.09869)	0.1642 (0.1535–0.1758)	0.1824 (0.1705–0.1948)

Supplementary Table 14 | Detailed structured report generation performance of HoloCT and competing methods on internal benchmarks. Each score is reported as mean with 95% confidence interval. NLP metrics include BLEU-4, ROUGE-L, and METEOR; clinical metrics include RaTEScore, LlamaScore, and GREEN.

Dataset Type	Dataset Name	Metric		Qwen3-VL-32B	MedGemma1.5-4B	M3D-Phi3-4B	RadFM-14B	HuluMed-32B	HoloCT-4B	HoloCT-8B
Internal	AMOS-MM-RG	NLP	BLEU-4	0.2180 (0.1696–0.2597)	0.2388 (0.1851–0.2742)	0.02219 (0.01664–0.02840)	0.07494 (0.06268–0.08690)	0.4710 (0.4541–0.4874)	0.5057 (0.4894–0.5226)	0.5489 (0.5319–0.5655)
			ROUGE-L	0.1662 (0.1256–0.2038)	0.1519 (0.1108–0.2006)	0.05091 (0.04573–0.05618)	0.1028 (0.09691–0.1092)	0.2833 (0.2747–0.2912)	0.3323 (0.3232–0.3429)	0.3526 (0.3390–0.3677)
			METEOR	0.1881 (0.1489–0.2211)	0.2006 (0.1666–0.2327)	0.03049 (0.02669–0.03400)	0.07119 (0.06528–0.07731)	0.3162 (0.3061–0.3264)	0.3676 (0.3570–0.3787)	0.3984 (0.3847–0.4117)
		Clinical	RaTE Score	0.4915 (0.4522–0.5358)	0.4579 (0.4146–0.5012)	0.2686 (0.2499–0.2874)	0.3599 (0.3501–0.3701)	0.6314 (0.6212–0.6415)	0.6831 (0.6755–0.6911)	0.6955 (0.6872–0.7046)
			Llama Score	3.524 (3.284–3.764)	3.422 (3.186–3.667)	2.376 (2.062–2.683)	0.4676 (0.3784–0.5568)	4.885 (4.700–5.078)	5.228 (5.065–5.395)	5.160 (5.000–5.345)
			GREEN	0.1409 (0.1081–0.1828)	0.1228 (0.08770–0.1645)	0.01784 (0.01205–0.02436)	0.01490 (0.00891–0.02253)	0.3476 (0.3230–0.3737)	0.3310 (0.3110–0.3501)	0.3458 (0.3237–0.3660)
	RadGenome	NLP	BLEU-4	0.2464 (0.2143–0.2735)	0.2430 (0.2175–0.2780)	0.03642 (0.03192–0.04079)	0.1176 (0.1109–0.1239)	0.5543 (0.5482–0.5604)	0.6096 (0.6017–0.6175)	0.6261 (0.6184–0.6340)
			ROUGE-L	0.1777 (0.1410–0.2019)	0.2092 (0.1785–0.2360)	0.05478 (0.05186–0.05791)	0.1220 (0.1195–0.1242)	0.2832 (0.2800–0.2867)	0.4885 (0.4788–0.4973)	0.4830 (0.4732–0.4918)
			METEOR	0.2077 (0.1782–0.2385)	0.2239 (0.1866–0.2478)	0.03806 (0.03565–0.04061)	0.09699 (0.09411–0.09974)	0.3766 (0.3710–0.3828)	0.5249 (0.5151–0.5340)	0.5440 (0.5349–0.5534)
		Clinical	RaTE Score	0.5598 (0.5325–0.5714)	0.5748 (0.5571–0.5844)	0.2513 (0.2397–0.2627)	0.5037 (0.4990–0.5084)	0.6607 (0.6573–0.6640)	0.7465 (0.7412–0.7514)	0.7419 (0.7370–0.7470)
			Llama Score	4.857 (4.732–4.971)	4.893 (4.741–4.980)	3.972 (3.825–4.137)	2.627 (2.557–2.695)	6.541 (6.493–6.586)	6.820 (6.719–6.911)	7.154 (7.063–7.240)
			GREEN	0.1681 (0.1395–0.1900)	0.1880 (0.1502–0.2050)	0.01946 (0.01603–0.02307)	0.08462 (0.07777–0.09153)	0.3919 (0.3811–0.4028)	0.4716 (0.4580–0.4853)	0.4624 (0.4499–0.4772)
	Merlin	NLP	BLEU-4	0.03657 (0.01616–0.05573)	0.03931 (0.02149–0.05132)	0.02809 (0.02623–0.03018)	0.02491 (0.02305–0.02684)	0.07043 (0.06828–0.07278)	0.3799 (0.3749–0.3849)	0.4393 (0.4345–0.4442)
			ROUGE-L	0.05003 (0.03526–0.07841)	0.05095 (0.03061–0.07650)	0.05689 (0.05557–0.05816)	0.04773 (0.04609–0.04942)	0.09121 (0.08991–0.09252)	0.2953 (0.2924–0.2980)	0.2765 (0.2739–0.2794)
			METEOR	0.03902 (0.01630–0.05542)	0.03273 (0.01305–0.05483)	0.03537 (0.03433–0.03640)	0.03451 (0.03332–0.03580)	0.07548 (0.06420–0.07664)	0.3233 (0.3195–0.3268)	0.3388 (0.3352–0.3421)
		Clinical	RaTE Score	0.3164 (0.2914–0.3478)	0.3108 (0.2825–0.3434)	0.2991 (0.2942–0.3037)	0.2865 (0.2823–0.2906)	0.3257 (0.3227–0.3286)	0.6766 (0.6748–0.6784)	0.6589 (0.6573–0.6606)
			Llama Score	1.627 (1.497–1.759)	1.529 (1.392–1.655)	1.866 (1.814–1.923)	0.8213 (0.7946–0.8487)	1.904 (1.873–1.936)	5.518 (5.475–5.561)	5.114 (5.073–5.155)
			GREEN	0.01973 (0.008451–0.03200)	0.01719 (0.004810–0.03717)	0.01402 (0.01277–0.01543)	0.02245 (0.02021–0.02461)	0.02246 (0.02125–0.02388)	0.3344 (0.3298–0.3393)	0.3178 (0.3133–0.3228)
	FLARE-2025-RG	NLP	BLEU-4	0.2420 (0.2092–0.2816)	0.2188 (0.1700–0.2554)	0.08584 (0.07506–0.09732)	0.1370 (0.1261–0.1485)	0.4895 (0.4791–0.4987)	0.6048 (0.5933–0.6161)	0.6039 (0.5923–0.6170)
			ROUGE-L	0.1820 (0.1419–0.2118)	0.1739 (0.1373–0.2038)	0.1024 (0.09696–0.1072)	0.1438 (0.1381–0.1490)	0.2696 (0.2647–0.2750)	0.4453 (0.4336–0.4574)	0.4551 (0.4430–0.4671)
			METEOR	0.2067 (0.1634–0.2458)	0.1898 (0.1444–0.2260)	0.07372 (0.06861–0.07890)	0.1181 (0.1122–0.1240)	0.3207 (0.3150–0.3264)	0.4979 (0.4868–0.5087)	0.5080 (0.4957–0.5206)
		Clinical	RaTE Score	0.5462 (0.5244–0.5805)	0.5141 (0.4850–0.5507)	0.4091 (0.3953–0.4239)	0.5004 (0.4892–0.5103)	0.6438 (0.6377–0.6499)	0.7570 (0.7512–0.7626)	0.7609 (0.7551–0.7670)
			Llama Score	4.387 (4.062–4.497)	3.958 (3.759–4.101)	2.876 (2.710–3.038)	1.977 (1.874–2.077)	5.916 (5.747–6.068)	5.880 (5.742–6.016)	5.803 (5.662–5.953)
			GREEN	0.1818 (0.1360–0.2105)	0.1621 (0.1284–0.2045)	0.05077 (0.04280–0.05948)	0.08322 (0.07352–0.09426)	0.4217 (0.4028–0.4411)	0.4128 (0.3962–0.4287)	0.4389 (0.4218–0.4541)

Supplementary Table 15 | Close-ended VQA performance of HoloCT with and without grounding on AMOS-MM-VQA and DeepTumorVQA. Each score is reported as mean with 95% confidence interval. Accu. denotes accuracy. For AMOS-MM-VQA, Anat.I., Attr.R., Quant.A., Dis.A., Risk.E., and Ther.S. denote anatomical identification, attribute recognition, quantitative assessment, disease analysis, risk evaluation, and therapeutic suggestion, respectively. For DeepTumorVQA, Recog., Meas., Vis.R., and Med.R. denote recognition, measurement, visual reasoning, and medical reasoning, respectively.

Dataset Type	Dataset Name	Category	Metric	HoloCT-4B w/o grounding	HoloCT-4B w/ grounding	HoloCT-8B w/o grounding	HoloCT-8B w/ grounding
Internal	AMOS-MM-VQA	Anat.I.	Accu.	0.6923 (0.5577–0.8077)	0.8269 (0.7303–0.9231)	0.7692 (0.6538–0.8846)	0.7885 (0.6731–0.8846)
		Attr.R.	Accu.	0.9286 (0.8333–1.000)	0.9286 (0.8333–1.000)	1.000 (1.000–1.000)	1.000 (1.000–1.000)
		Quant.A.	Accu.	0.7097 (0.5968–0.8226)	0.8871 (0.8065–0.9516)	0.9032 (0.8226–0.9677)	0.8710 (0.7903–0.9516)
		Dis.A.	Accu.	0.8056 (0.6944–0.9167)	0.8889 (0.7778–0.9722)	0.9167 (0.8056–1.000)	0.9444 (0.8611–1.000)
		Risk.E.	Accu.	0.7561 (0.6341–0.8780)	0.8293 (0.7073–0.9268)	0.8780 (0.7561–0.9756)	0.9024 (0.8049–0.9756)
		Ther.S.	Accu.	0.6333 (0.4667–0.8000)	0.8333 (0.7000–0.9333)	0.7333 (0.5667–0.8667)	0.8667 (0.7333–0.9667)
		Overall	Accu.	0.7529 (0.6996–0.8024)	0.8669 (0.8251–0.9087)	0.8707 (0.8289–0.9087)	0.8897 (0.8479–0.9278)
	DeepTumorVQA	Recog.	Accu.	0.7121 (0.6800–0.7409)	0.8627 (0.8405–0.8848)	0.8128 (0.7852–0.8394)	0.8571 (0.8339–0.8804)
		Meas.	Accu.	0.5686 (0.5421–0.5967)	0.5818 (0.5529–0.6083)	0.5934 (0.5653–0.6215)	0.5884 (0.5595–0.6149)
		Vis.R.	Accu.	0.5812 (0.5610–0.6022)	0.5770 (0.5551–0.5972)	0.5944 (0.5738–0.6155)	0.5780 (0.5560–0.5981)
		Med.R.	Accu.	0.6168 (0.5888–0.6431)	0.7089 (0.6826–0.7336)	0.6752 (0.6480–0.7015)	0.6982 (0.6719–0.7229)
		Overall	Accu.	0.6077 (0.5948–0.6206)	0.6539 (0.6416–0.6675)	0.6467 (0.6340–0.6593)	0.6525 (0.6405–0.6657)

Supplementary Table 16 | Segmentation performance of HoloCT segmentation and grounded variants on AMOS-MM-VQA and DeepTumorVQA. Each score is reported as mean with 95% confidence interval. Match counts denote matched/reference objects; grounded Dice and HD95 use matched objects. Dice measures volumetric overlap; HD95 denotes the 95th-percentile Hausdorff distance in millimetres.

Dataset Type	Dataset Name	Label Name	Metric	HoloCT-4B Segmentation-only	Match count-4B	HoloCT-4B w/ grounding	HoloCT-8B Segmentation-only	Match count-8B	HoloCT-8B w/ grounding
Internal	AMOS-MM-VQA	spleen	Dice	0.9598 (0.9453–0.9733)	9/12	0.9520 (0.9248–0.9731)	0.9431 (0.9196–0.9631)	10/12	0.9383 (0.9111–0.9625)
			HD95	3.535 (1.379–7.313)		4.394 (1.325–10.13)	4.877 (1.898–8.202)		5.040 (1.842–9.093)
		right kidney	Dice	0.9513 (0.9366–0.9636)	45/45	0.9493 (0.9343–0.9626)	0.9503 (0.9378–0.9614)	42/45	0.9462 (0.9332–0.9582)
			HD95	5.181 (1.681–10.62)		5.410 (1.747–11.21)	3.361 (1.758–5.521)		5.020 (2.240–8.785)
		left kidney	Dice	0.9484 (0.9291–0.9625)	43/44	0.9497 (0.9354–0.9617)	0.9368 (0.9080–0.9595)	42/44	0.9358 (0.9078–0.9580)
			HD95	2.924 (1.672–4.524)		2.930 (1.668–4.566)	3.886 (1.653–7.375)		7.016 (1.559–14.20)
		gallbladder	Dice	0.8687 (0.8289–0.9091)	24/26	0.8728 (0.8300–0.9125)	0.8790 (0.8245–0.9285)	25/26	0.8790 (0.8163–0.9280)
			HD95	7.831 (4.046–12.19)		7.453 (3.611–12.05)	7.239 (3.035–13.00)		7.488 (3.085–14.03)
		esophagus	Dice	0.8823 (0.8823–0.8823)	2/3	0.8869 (0.8869–0.8869)	0.8338 (0.8338–0.8338)	2/3	0.8395 (0.8395–0.8395)
			HD95	2.751 (2.751–2.751)		2.461 (2.461–2.461)	4.922 (4.922–4.922)		4.685 (4.685–4.685)
		liver	Dice	0.9672 (0.9599–0.9725)	84/87	0.9694 (0.9642–0.9735)	0.9684 (0.9651–0.9716)	83/87	0.9694 (0.9662–0.9724)
			HD95	3.555 (2.463–5.295)		3.334 (2.361–4.813)	3.503 (2.880–4.269)		3.115 (2.640–3.646)
		stomach	Dice	0.7155 (0.3183–0.9334)	6/6	0.7184 (0.3191–0.9373)	0.6953 (0.3133–0.9104)	6/6	0.6920 (0.3144–0.9092)
			HD95	24.63 (5.719–59.93)		23.59 (4.641–58.33)	25.91 (8.651–56.16)		26.81 (10.10–54.41)
		aorta	Dice	0.9025 (0.9012–0.9037)	3/3	0.9080 (0.8921–0.9238)	0.9082 (0.9082–0.9082)	2/3	0.9075 (0.9075–0.9075)
			HD95	36.01 (8.463–63.56)		41.30 (19.43–63.17)	2.414 (2.414–2.414)		2.545 (2.545–2.545)
		pancreas	Dice	0.6788 (0.3859–0.8834)	18/20	0.6811 (0.3897–0.8808)	0.7034 (0.4378–0.8558)	19/20	0.7020 (0.4390–0.8553)
			HD95	18.20 (3.661–41.24)		18.18 (3.774–42.11)	15.28 (4.080–39.62)		15.49 (4.092–39.52)
		left adrenal gland	Dice	0.6892 (0.6892–0.6892)	2/2	0.6687 (0.6687–0.6687)	0.8427 (0.8427–0.8427)	2/2	0.8357 (0.8357–0.8357)
			HD95	3.910 (3.910–3.910)		3.910 (3.910–3.910)	1.564 (1.564–1.564)		1.564 (1.564–1.564)
		urinary bladder	Dice	0.8488 (0.7902–0.9073)	3/4	0.8499 (0.8009–0.8989)	0.8435 (0.8114–0.8757)	2/4	0.8454 (0.8299–0.8609)
			HD95	6.087 (4.281–7.894)		5.669 (4.641–6.697)	5.000 (5.000–5.000)		5.000 (5.000–5.000)
	DeepTumorVQA	prostate or uterus	Dice	0.8864 (0.8603–0.9101)	28/30	0.8876 (0.8621–0.9112)	0.8729 (0.8453–0.8978)	29/30	0.8845 (0.8582–0.9089)
			HD95	6.473 (4.138–9.533)		5.730 (3.695–8.594)	56.25 (12.57–107.3)		32.63 (6.226–68.06)
		colon	Dice	0.8973 (0.8776–0.9157)	18/31	0.8823 (0.8589–0.9038)	0.8923 (0.8690–0.9141)	17/31	0.8893 (0.8637–0.9119)
			HD95	11.82 (5.794–18.78)		13.68 (7.184–21.02)	13.99 (7.450–20.91)		15.07 (8.466–22.90)
		intestine	Dice	0.8169 (0.7514–0.8824)	6/10	0.8158 (0.7544–0.8772)	0.8500 (0.7412–0.9044)	4/10	0.8480 (0.7429–0.9008)
			HD95	14.08 (5.651–22.51)		14.48 (6.162–22.79)	12.53 (5.260–23.06)		12.82 (5.000–23.18)
		left kidney	Dice	0.9490 (0.9408–0.9564)	671/749	0.9484 (0.9400–0.9560)	0.9475 (0.9396–0.9546)	710/749	0.9481 (0.9399–0.9554)
			HD95	4.505 (3.353–5.719)		5.129 (3.879–6.437)	4.495 (3.535–5.568)		4.302 (3.299–5.389)
		kidney lesion	Dice	0.6474 (0.6100–0.6865)	620/624	0.6589 (0.6222–0.6975)	0.7025 (0.6706–0.7353)	590/624	0.7061 (0.6724–0.7387)
			HD95	38.35 (30.85–45.69)		42.66 (34.73–50.05)	39.48 (32.20–47.08)		37.48 (31.04–44.71)
		right kidney	Dice	0.9563 (0.9503–0.9615)	660/747	0.9567 (0.9505–0.9624)	0.9567 (0.9498–0.9621)	691/747	0.9562 (0.9492–0.9621)
			HD95	3.588 (2.918–4.390)		3.892 (2.962–5.157)	3.461 (2.742–4.317)		3.250 (2.533–4.085)
		liver	Dice	0.9738 (0.9710–0.9758)	647/721	0.9734 (0.9707–0.9755)	0.9723 (0.9694–0.9744)	645/721	0.9726 (0.9696–0.9748)
			HD95	4.112 (3.490–4.797)		4.252 (3.620–4.954)	4.164 (3.534–4.905)		4.139 (3.492–4.902)

Supplementary Table 16 (continued) | Segmentation performance of HoloCT segmentation and grounded variants on AMOS-MM-VQA and DeepTumorVQA.

Each score is reported as mean with 95% confidence interval. Match counts denote matched/reference objects; grounded Dice and HD95 use matched objects. Dice measures volumetric overlap; HD95 denotes the 95th-percentile Hausdorff distance in millimetres.

Dataset Type	Dataset Name	Label Name	Metric	HoloCT-4B Segmentation-only	Match count-4B	HoloCT-4B w/ grounding	HoloCT-8B Segmentation-only	Match count-8B	HoloCT-8B w/ grounding
Internal	DeepTumorVQA	liver lesion	Dice	0.6981 (0.6684–0.7261)	1047/1063	0.7031 (0.6753–0.7305)	0.6917 (0.6601–0.7194)	1034/1063	0.6873 (0.6554–0.7152)
			HD95	24.92 (20.34–29.58)		25.15 (20.63–29.63)	40.18 (34.29–47.56)		39.42 (33.57–46.52)
		caudate lobe	Dice	0.8221 (0.8058–0.8370)	127/131	0.8153 (0.7982–0.8310)	0.8243 (0.8103–0.8384)	125/131	0.8273 (0.8118–0.8420)
			HD95	6.987 (6.230–7.800)		7.214 (6.421–8.082)	6.820 (6.010–7.629)		6.803 (6.001–7.639)
		left lateral superior segment of liver	Dice	0.9064 (0.8882–0.9210)	111/114	0.9058 (0.8889–0.9198)	0.9014 (0.8798–0.9165)	110/114	0.8979 (0.8753–0.9138)
			HD95	7.427 (6.136–8.953)		7.746 (6.497–9.263)	7.077 (5.739–8.853)		7.918 (6.112–10.15)
		left lateral inferior segment of liver	Dice	0.9090 (0.8976–0.9202)	101/103	0.9091 (0.8976–0.9207)	0.9094 (0.8977–0.9202)	100/103	0.8969 (0.8800–0.9113)
			HD95	5.320 (4.443–6.253)		5.310 (4.423–6.207)	7.250 (5.226–10.29)		10.41 (7.114–14.48)
		left medial segment of liver	Dice	0.8669 (0.8452–0.8833)	154/158	0.8630 (0.8415–0.8800)	0.8585 (0.8341–0.8798)	153/158	0.8648 (0.8422–0.8834)
			HD95	12.01 (9.492–15.42)		14.67 (11.77–18.43)	10.01 (8.014–12.52)		12.80 (9.575–16.77)
		right anterior inferior segment of liver	Dice	0.8724 (0.8408–0.8968)	114/116	0.8754 (0.8442–0.8989)	0.8735 (0.8412–0.8974)	114/116	0.8643 (0.8332–0.8869)
			HD95	19.59 (10.29–33.77)		18.52 (10.27–32.63)	19.17 (10.61–35.37)		22.63 (13.05–38.77)
		right posterior inferior segment of liver	Dice	0.8374 (0.7761–0.8788)	118/119	0.8382 (0.7775–0.8796)	0.8495 (0.7932–0.8899)	118/119	0.8424 (0.7868–0.8829)
			HD95	20.11 (10.42–32.99)		57.19 (38.79–76.77)	15.94 (10.04–22.72)		13.59 (9.850–18.01)
		right posterior superior segment of liver	Dice	0.8789 (0.8521–0.9003)	115/117	0.8843 (0.8603–0.9040)	0.9077 (0.8894–0.9216)	115/117	0.9042 (0.8865–0.9191)
			HD95	17.02 (8.092–31.46)		19.70 (9.431–35.22)	15.36 (6.817–29.56)		16.09 (7.353–30.50)
		right anterior superior segment of liver	Dice	0.8833 (0.8645–0.8967)	149/154	0.8882 (0.8732–0.9006)	0.8929 (0.8801–0.9041)	149/154	0.8923 (0.8805–0.9025)
			HD95	15.54 (8.148–27.19)		15.86 (8.426–27.65)	14.79 (8.687–25.62)		14.93 (8.791–25.82)
		pancreas	Dice	0.8785 (0.8707–0.8854)	1123/1208	0.8781 (0.8703–0.8852)	0.8813 (0.8734–0.8886)	1074/1208	0.8803 (0.8725–0.8878)
			HD95	6.280 (5.173–7.680)		6.217 (5.118–7.553)	6.827 (5.451–8.438)		6.889 (5.527–8.495)
		pancreas lesion	Dice	0.5822 (0.5182–0.6458)	188/267	0.5681 (0.5049–0.6307)	0.6018 (0.5465–0.6523)	224/267	0.5860 (0.5325–0.6380)
			HD95	21.78 (15.80–28.71)		22.69 (16.80–29.70)	27.51 (20.38–35.33)		27.99 (20.83–35.58)
		spleen	Dice	0.9589 (0.9537–0.9634)	1387/1391	0.9583 (0.9531–0.9628)	0.9584 (0.9531–0.9630)	1387/1391	0.9581 (0.9528–0.9627)
			HD95	2.707 (2.152–3.475)		2.733 (2.187–3.502)	2.665 (2.284–3.114)		2.657 (2.277–3.094)
		stomach	Dice	0.9488 (0.9275–0.9632)	7/12	0.9479 (0.9260–0.9631)	0.9560 (0.9429–0.9673)	11/12	0.9519 (0.9389–0.9621)
			HD95	3.047 (2.008–4.208)		3.181 (2.014–4.465)	3.941 (2.848–5.027)		8.419 (3.309–17.17)
		colon lesion	Dice	0.6523 (0.5210–0.7504)	44/44	0.6495 (0.5185–0.7469)	0.6824 (0.5687–0.7675)	40/44	0.6737 (0.5578–0.7659)
			HD95	30.13 (10.46–54.70)		30.82 (10.80–55.84)	28.99 (8.332–59.80)		47.77 (18.86–82.69)

Supplementary Table 17 | Text-quality performance of HoloCT and competing methods for anatomy-grounded and finding-grounded report generation on RadGenome, ReXGroundingCT and private cohorts.

RadGenome and ReXGroundingCT are internal datasets; Private05 and Private06 are private cohorts. ▲ denotes a fine-tuned baseline. Each score is reported as mean with 95% confidence interval. NLP metrics include BLEU-4, ROUGE-L, and METEOR; clinical metrics include RaTEScore, LlamaScore, and GREEN.

Dataset Type	Dataset Name	Metric Type	Metric	HuluMed-32B	HoloCT-4B w/o grounding	HoloCT-4B w/ grounding	HoloCT-8B w/o grounding	HoloCT-8B w/ grounding
Internal	RadGenome	NLP	BLEU-4	0.5543 (0.5482–0.5604)	0.6096 (0.6017–0.6175)	0.6371 (0.6292–0.6445)	0.6261 (0.6184–0.6340)	0.6253 (0.6178–0.6335)
			ROUGE-L	0.2832 (0.2800–0.2867)	0.4885 (0.4788–0.4973)	0.4841 (0.4742–0.4937)	0.4830 (0.4732–0.4918)	0.4859 (0.4763–0.4945)
			METEOR	0.3766 (0.3710–0.3828)	0.5249 (0.5151–0.5340)	0.5460 (0.5367–0.5547)	0.5440 (0.5349–0.5534)	0.5460 (0.5368–0.5552)
		Clinical	RaTEScore	0.6607 (0.6573–0.6640)	0.7465 (0.7412–0.7514)	0.7497 (0.7446–0.7547)	0.7419 (0.7370–0.7470)	0.7482 (0.7434–0.7533)
			Llama Score	6.541 (6.493–6.586)	6.820 (6.719–6.911)	7.230 (7.149–7.310)	7.154 (7.063–7.240)	7.181 (7.093–7.271)
			GREEN	0.3919 (0.3811–0.4028)	0.4716 (0.4580–0.4853)	0.4595 (0.4465–0.4727)	0.4624 (0.4499–0.4772)	0.4740 (0.4606–0.4868)
	ReXGroundingCT	NLP	BLEU-4	0.1456▲ (0.1378–0.1534)	0.2799 (0.2591–0.2998)	0.3288 (0.3046–0.3528)	0.3278 (0.3027–0.3508)	0.3517 (0.3257–0.3803)
			ROUGE-L	0.3678▲ (0.3589–0.3767)	0.3170 (0.3002–0.3355)	0.3313 (0.3090–0.3542)	0.3572 (0.3350–0.3808)	0.3742 (0.3481–0.4012)
			METEOR	0.2567▲ (0.2489–0.2645)	0.2512 (0.2325–0.2709)	0.2776 (0.2543–0.3020)	0.2975 (0.2719–0.3209)	0.3217 (0.2954–0.3505)
		Clinical	RaTEScore	0.5012▲ (0.4898–0.5126)	0.5252 (0.5109–0.5399)	0.5236 (0.5056–0.5414)	0.5509 (0.5345–0.5677)	0.5616 (0.5444–0.5789)
			Llama Score	5.654▲ (5.412–5.896)	5.944 (5.766–6.130)	6.297 (6.121–6.476)	6.179 (6.007–6.362)	6.343 (6.140–6.539)
			GREEN	0.1987▲ (0.1854–0.2120)	0.1788 (0.1553–0.2049)	0.2237 (0.1969–0.2527)	0.2173 (0.1890–0.2465)	0.2424 (0.2128–0.2771)
Private (zero-shot)	Private05(Abdomen Report)	NLP	BLEU-4	0.2759 (0.2660–0.2847)	0.4616 (0.4525–0.4694)	0.4791 (0.4692–0.4851)	0.4596 (0.4497–0.4687)	0.4672 (0.4561–0.4756)
			ROUGE-L	0.1309 (0.1268–0.1350)	0.2807 (0.2750–0.2868)	0.2974 (0.2916–0.3020)	0.3182 (0.3112–0.3251)	0.3277 (0.3211–0.3334)
			METEOR	0.1676 (0.1634–0.1720)	0.4046 (0.3963–0.4129)	0.4274 (0.4195–0.4361)	0.4046 (0.3952–0.4136)	0.4117 (0.4014–0.4210)
		Clinical	RaTEScore	0.4387 (0.4313–0.4456)	0.6435 (0.6396–0.6475)	0.6606 (0.6558–0.6653)	0.6646 (0.6604–0.6691)	0.6711 (0.6671–0.6748)
			Llama Score	3.078 (2.991–3.160)	4.346 (4.259–4.435)	4.436 (4.353–4.523)	4.377 (4.283–4.472)	4.441 (4.347–4.535)
			GREEN	0.09411 (0.08255–0.1071)	0.2735 (0.2630–0.2843)	0.2926 (0.2802–0.3025)	0.3256 (0.3150–0.3357)	0.3346 (0.3228–0.3457)
	Private06(Chest Report)	NLP	BLEU-4	0.3234 (0.3128–0.3344)	0.4178 (0.4084–0.4264)	0.4358 (0.4281–0.4444)	0.4334 (0.4243–0.4427)	0.4412 (0.4317–0.4500)
			ROUGE-L	0.1565 (0.1524–0.1603)	0.2146 (0.2124–0.2171)	0.2327 (0.2322–0.2359)	0.2308 (0.2281–0.2333)	0.2424 (0.2383–0.2462)
			METEOR	0.1965 (0.1915–0.2010)	0.3597 (0.3525–0.3663)	0.3859 (0.3799–0.3920)	0.3881 (0.3808–0.3953)	0.3986 (0.3905–0.4052)
		Clinical	RaTEScore	0.5355 (0.5297–0.5412)	0.6044 (0.6010–0.6081)	0.6241 (0.6201–0.6289)	0.6216 (0.6183–0.6250)	0.6311 (0.6296–0.6336)
			Llama Score	4.019 (3.833–4.199)	4.688 (4.575–4.806)	4.782 (4.669–4.904)	4.556 (4.438–4.676)	4.615 (4.495–4.736)
			GREEN	0.09659 (0.07852–0.1163)	0.08956 (0.08003–0.09852)	0.1039 (0.09320–0.1128)	0.1149 (0.1037–0.1265)	0.1288 (0.1175–0.1416)

Supplementary Table 18 | Segmentation performance on RadGenome for anatomy-grounded report generation.
Each score is reported as mean with 95% confidence interval. Match counts denote matched/reference regions; Dice and HD95 use matched regions. Dice measures volumetric overlap and HD95 denotes the 95th-percentile Hausdorff distance in millimetres.

Dataset Type	Dataset Name	Region Name	Metric	Match count-4B	HoloCT-4B	Match count-8B	HoloCT-8B
Internal	RadGenome	abdomen	Dice	1518/1518	0.9053 (0.9008–0.9094)	1518/1518	0.9117 (0.9075–0.9155)
			HD95		10.21 (9.316–11.41)		10.17 (9.615–10.74)
		bone	Dice	1509/1509	0.9169 (0.9144–0.9192)	1509/1509	0.9192 (0.9168–0.9211)
			HD95		2.091 (1.572–2.900)		1.698 (1.512–1.900)
		breast	Dice	58/58	0.5384 (0.4603–0.6109)	58/58	0.5116 (0.4367–0.5764)
			HD95		64.27 (34.52–98.07)		63.07 (31.95–98.62)
		esophagus	Dice	1328/1328	0.8766 (0.8728–0.8800)	1328/1328	0.8758 (0.8723–0.8789)
			HD95		2.438 (1.914–3.392)		1.962 (1.843–2.103)
		heart	Dice	1433/1433	0.9102 (0.9050–0.9150)	1433/1433	0.9103 (0.9046–0.9157)
			HD95		19.07 (16.65–21.25)		19.55 (17.01–21.97)
		lung	Dice	1514/1514	0.9717 (0.9672–0.9760)	1514/1514	0.9725 (0.9680–0.9766)
			HD95		4.429 (3.411–5.795)		3.606 (3.067–4.233)
		mediastinum	Dice	1523/1523	0.8254 (0.8209–0.8295)	1523/1523	0.8368 (0.8326–0.8404)
			HD95		7.088 (5.523–8.882)		6.082 (4.816–7.520)
		pleura	Dice	1172/1172	0.9711 (0.9655–0.9763)	1172/1172	0.9721 (0.9666–0.9769)
			HD95		5.062 (3.602–6.899)		4.107 (3.230–5.071)
		thyroid	Dice	51/51	0.7054 (0.6395–0.7564)	51/51	0.7182 (0.6578–0.7691)
			HD95		7.121 (4.199–11.72)		4.855 (3.430–6.600)
		trachea and bronchi	Dice	1417/1417	0.9154 (0.9111–0.9191)	1417/1417	0.9175 (0.9134–0.9210)
			HD95		2.954 (1.839–4.399)		3.019 (1.888–4.331)

Supplementary Table 19 | Finding-level referring segmentation performance.
Each score is reported as mean with 95% confidence interval.

Metric	HoloCT-8B	HoloCT-4B	VoxTell	SAT-Pro
Non-focal Dice	0.2517 (0.2237–0.2767)	0.1830 (0.1578–0.2065)	0.1493 (0.1292–0.1831)	0.09780 (0.08140–0.1241)
Focal Dice	0.3010 (0.2864–0.3165)	0.2414 (0.2273–0.2554)	0.2781 (0.2627–0.2979)	0.1849 (0.1781–0.2068)
Average Dice	0.2908 (0.2769–0.3038)	0.2293 (0.2168–0.2420)	0.2591 (0.2404–0.2738)	0.1681 (0.1515–0.1846)
HIT5%	0.7700 (0.7457–0.7943)	0.6936 (0.6684–0.7205)	0.6592 (0.6383–0.6851)	0.5291 (0.5013–0.5630)
HIT10%	0.7118 (0.6866–0.7387)	0.6128 (0.5868–0.6406)	0.5991 (0.5767–0.6251)	0.4863 (0.4619–0.5195)

Supplementary Table 20 | Finding-grounded report generation performance.
Each score is reported as mean with 95% confidence interval.

Metric	HoloCT-8B	HoloCT-4B	HuluMed-32B + VoxTell
Finding F1	0.2776 (0.2475–0.3139)	0.2860 (0.2579–0.3149)	0.2386 (0.2129–0.2777)
Non-focal conditional Dice	0.1789 (0.1233–0.2356)	0.1020 (0.05502–0.1574)	0.1271 (0.08630–0.1758)
Focal conditional Dice	0.2427 (0.2162–0.2702)	0.1503 (0.1284–0.1740)	0.2073 (0.1816–0.2307)
Average conditional Dice	0.2339 (0.2093–0.2579)	0.1432 (0.1218–0.1637)	0.1857 (0.1646–0.2134)
Conditional HIT5%	0.7382 (0.6864–0.7862)	0.5131 (0.4545–0.5715)	0.7386 (0.6900–0.7968)
Conditional HIT10%	0.6691 (0.6127–0.7263)	0.4444 (0.3876–0.5017)	0.5035 (0.4502–0.5633)

Supplementary Table 21 | Zero-shot conditional segmentation performance of HoloCT segmentation and grounded variants across classification targets.

Each score is reported as mean with 95% confidence interval. Cond. Dice denotes conditional Dice. Internal, external-seen and external-unseen objects follow the zero-shot grouping used in the classification–segmentation evaluation.

Dataset Type	Dataset Name	Label Name	HoloCT-4B Segmentation-only	HoloCT-4B w/ grounding	HoloCT-8B Segmentation-only	HoloCT-8B w/ grounding
Internal	Total-SegmentatorV2	left kidney cyst	0.9025 (0.8896–0.9195)	0.9016 (0.8845–0.9240)	0.8882 (0.8882–0.8882)	0.8872 (0.8380–0.9342)
		right kidney cyst	0.8595 (0.7952–0.9316)	0.8536 (0.7821–0.9343)	0.8536 (0.8536–0.8536)	0.8563 (0.7787–0.9328)
	KiTS23	kidney cyst	0.5913 (0.5081–0.6765)	0.5781 (0.4930–0.6636)	0.7056 (0.7056–0.7056)	0.7046 (0.6268–0.7816)
	CT-ICH	intracranial hemorrhage	0.5524 (0.5155–0.5965)	0.5557 (0.5180–0.6021)	0.6682 (0.4940–0.8423)	0.6853 (0.5247–0.8460)
	PanTS	pancreas lesion	0.4899 (0.4223–0.5576)	0.4920 (0.4242–0.5574)	0.5425 (0.5425–0.5425)	0.5412 (0.4820–0.6016)
	AbdomenAtlas 3.0	kidney lesion	0.7349 (0.6924–0.7738)	0.7344 (0.6924–0.7729)	0.8222 (0.8222–0.8222)	0.8221 (0.7941–0.8475)
		liver lesion	0.7240 (0.6950–0.7503)	0.7248 (0.6988–0.7506)	0.7103 (0.7103–0.7103)	0.7102 (0.6796–0.7402)
		pancreas lesion	0.5465 (0.4900–0.6032)	0.5480 (0.4906–0.6055)	0.5722 (0.5722–0.5722)	0.5712 (0.5176–0.6266)
		colon lesion	0.6692 (0.5656–0.7475)	0.6678 (0.5685–0.7530)	0.7250 (0.7250–0.7250)	0.7231 (0.6598–0.7816)
	LiMT	liver cyst	0.6034 (0.4968–0.7306)	0.6003 (0.4689–0.7237)	0.6400 (0.5092–0.7521)	0.6448 (0.5193–0.7476)
		liver hemangioma	0.8304 (0.7592–0.8736)	0.8313 (0.7636–0.8765)	0.7105 (0.5285–0.8361)	0.7106 (0.4914–0.8313)
		metastatic liver cancer	0.6286 (0.6286–0.6286)	0.6313 (0.6313–0.6313)	0.6601 (0.6601–0.6601)	0.6605 (0.6605–0.6605)
		primary liver cancer	0.7394 (0.6040–0.8634)	0.7407 (0.5913–0.8614)	0.6927 (0.4286–0.8604)	0.6904 (0.4280–0.8608)
	LiMT-oneclass	liver lesion	0.4370 (0.2359–0.6160)	0.6100 (0.5105–0.7276)	0.4110 (0.2368–0.5851)	0.5606 (0.4261–0.6810)
External-seen (zero-shot)	QUBIQ2021	pancreas lesion	0.3470 (0.3470–0.3470)	0.3361 (0.3361–0.3361)	0.3855 (0.3363–0.4347)	0.3867 (0.3384–0.4350)
	TotalSeg-subtask-liver-lesion	liver lesion	0.7100 (0.6807–0.7360)	0.7096 (0.6827–0.7361)	0.7441 (0.7163–0.7702)	0.7456 (0.7185–0.7684)
	MHB-SEG-oneclass	intracranial hemorrhage	0.4010 (0.2727–0.5286)	0.4056 (0.2862–0.5260)	0.5052 (0.5052–0.5052)	0.5143 (0.3883–0.6315)
External-unseen (zero-shot)	PANORAMA	pancreatic ductal adenocarcinoma	0.4135 (0.3522–0.4806)	0.4112 (0.3469–0.4737)	0.4515 (0.3745–0.5222)	0.4468 (0.3702–0.5143)
	MHB-SEG-multiclass	subdural hemorrhage	0.6663 (0.4673–0.8423)	0.6676 (0.4534–0.8512)	0.6755 (0.6755–0.6755)	0.6724 (0.4538–0.8441)
		subarachnoid hemorrhage	0.1746 (0.002516–0.4351)	0.1777 (0.0000–0.5956)	0.4967 (0.4967–0.4967)	0.5016 (0.2761–0.7039)

Supplementary Table 22 | Training-strategy ablations S1–S6 across segmentation, grounded lesion screening, grounded visual question answering, and grounded report generation.

For segmentation, Macro Dice denotes the dataset-level macro Dice. For grounded lesion screening, Macro F1 is calculated as the arithmetic mean of label-wise F1 scores within each dataset. For grounded visual question answering, Accu. denotes accuracy. For grounded report generation, BLEU-4, ROUGE-L, METEOR, RaTEScore, LlamaScore, and GREEN are reported. Overall scores retain the supplied task-level summaries and are shown without confidence intervals; the report-generation Overall score is LCS. Dataset scores are reported as mean with 95% confidence interval when available.

Task	Dataset Name	Metric	S1	S2	S3	S4	S5	S6
Segmentation	Total-SegmentatorV2	Macro Dice	Not supported	0.6508 (0.6384–0.6703)	0.6626 (0.6573–0.6733)	0.7467 (0.7351–0.7584)	0.7779 (0.7604–0.7874)	0.7775 (0.7641–0.7892)
	AbdomenAtlas3.0	Macro Dice		0.5514 (0.5255–0.5761)	0.5631 (0.5445–0.5827)	0.6421 (0.6275–0.6673)	0.6949 (0.6700–0.7098)	0.7015 (0.6827–0.7160)
	Overall			0.6011	0.6129	0.6944	0.7364	0.7395
Grounded lesion screening	AbdomenAtlas3.0	Macro F1	0.2955 (0.2557–0.3367)	0.3181 (0.2740–0.3597)	0.3720 (0.3268–0.4169)	0.4241 (0.3753–0.4681)	0.4745 (0.4263–0.5188)	0.4852 (0.4374–0.5296)
	LiMT	Macro F1	0.2172 (0.07629–0.3404)	0.2320 (0.07941–0.3628)	0.3298 (0.1483–0.4713)	0.4287 (0.2525–0.5619)	0.5060 (0.3396–0.6299)	0.5455 (0.3521–0.7089)
	Overall		0.2563	0.2751	0.3509	0.4264	0.4903	0.5153
Grounded visual question answering	AMOS-MM-VQA	Accu.	0.6958 (0.6388–0.7529)	0.7119 (0.6550–0.7643)	0.7338 (0.6806–0.7871)	0.7496 (0.6958–0.7985)	0.7643 (0.7148–0.8137)	0.7795 (0.7262–0.8289)
	DeepTumorVQA	Accu.	0.4553 (0.4417–0.4688)	0.4949 (0.4811–0.5091)	0.5278 (0.5151–0.5414)	0.5549 (0.5422–0.5694)	0.5781 (0.5647–0.5921)	0.5977 (0.5843–0.6112)
	Overall		0.5756	0.6034	0.6308	0.6523	0.6712	0.6886
Grounded report generation	RadGenome	BLEU-4	0.4784 (0.4519–0.4907)	0.4482 (0.4207–0.4650)	0.4908 (0.4702–0.5045)	0.5124 (0.5043–0.5262)	0.5474 (0.5389–0.5534)	0.5637 (0.5534–0.5791)
		ROUGE-L	0.3771 (0.3661–0.3969)	0.3562 (0.3373–0.3715)	0.4119 (0.4015–0.4234)	0.4012 (0.3940–0.4180)	0.4269 (0.4138–0.4341)	0.4303 (0.4244–0.4417)
		METEOR	0.4253 (0.4177–0.4397)	0.4084 (0.3884–0.4208)	0.4439 (0.4332–0.4528)	0.4820 (0.4738–0.4990)	0.4732 (0.4677–0.4813)	0.4800 (0.4792–0.4969)
		RaTEScore	0.5942 (0.5867–0.6170)	0.5636 (0.5404–0.5804)	0.6119 (0.6026–0.6279)	0.6373 (0.6242–0.6425)	0.6584 (0.6489–0.6617)	0.6737 (0.6656–0.6882)
		LlamaScore	5.886 (5.763–6.002)	5.649 (5.511–5.776)	6.104 (5.999–6.217)	6.253 (6.159–6.368)	6.426 (6.326–6.511)	6.553 (6.467–6.644)
		GREEN	0.3456 (0.3226–0.3646)	0.3267 (0.3021–0.3458)	0.3622 (0.3524–0.3821)	0.3748 (0.3632–0.3895)	0.3930 (0.3895–0.4027)	0.4009 (0.3910–0.4149)
	ReXGroundingCT	BLEU-4	0.2422 (0.2058–0.2823)	0.2338 (0.1952–0.2763)	0.2658 (0.2365–0.3002)	0.2706 (0.2441–0.3005)	0.2948 (0.2606–0.3244)	0.3082 (0.2732–0.3329)
		ROUGE-L	0.2743 (0.2470–0.3089)	0.2690 (0.2205–0.3032)	0.2960 (0.2612–0.3281)	0.3071 (0.2780–0.3340)	0.3239 (0.2975–0.3531)	0.3357 (0.3010–0.3523)
		METEOR	0.2473 (0.2144–0.2776)	0.2284 (0.1954–0.2577)	0.2522 (0.2278–0.2882)	0.2771 (0.2489–0.3057)	0.2905 (0.2775–0.3198)	0.2851 (0.2643–0.3002)
		RaTEScore	0.4381 (0.4012–0.4613)	0.4188 (0.3726–0.4439)	0.4546 (0.4236–0.4896)	0.4719 (0.4490–0.5093)	0.4891 (0.4630–0.5049)	0.5070 (0.4885–0.5242)
		LlamaScore	5.312 (5.010–5.598)	5.128 (4.809–5.438)	5.471 (5.190–5.754)	5.657 (5.381–5.925)	5.799 (5.539–6.043)	5.903 (5.672–6.127)
		GREEN	0.1667 (0.1330–0.1925)	0.1538 (0.1221–0.1834)	0.1909 (0.1778–0.2215)	0.1863 (0.1538–0.2158)	0.1913 (0.1740–0.2202)	0.2012 (0.1898–0.2247)
	Overall		0.3924	0.3737	0.4115	0.4260	0.4426	0.4526

Supplementary Table 23 | Masking and grounding-request ablations across segmentation, grounded lesion screening, grounded visual question answering, and grounded report generation.

For segmentation, Macro Dice denotes the dataset-level macro Dice. For grounded lesion screening, Macro F1 is calculated as the arithmetic mean of label-wise F1 scores within each dataset. For grounded visual question answering, Accu. denotes accuracy. For grounded report generation, BLEU-4, ROUGE-L, METEOR, RaTEScore, LlamaScore, and GREEN are reported. Overall scores retain the supplied task-level summaries and are shown without confidence intervals; the report-generation Overall score is LCS. Dataset scores are reported as mean with 95% confidence interval when available.

Task	Dataset Name	Metric	HoloCT-8B Intact w/ grounding	HoloCT-8B Intact w/o grounding	HoloCT-8B Masked w/ grounding	HoloCT-8B Masked w/o grounding
Segmentation	Total-SegmentatorV2	Macro Dice	0.8776 (0.8741–0.8806)	Not supported	0.8300 (0.8265–0.8332)	Not supported
	AbdomenAtlas3.0	Macro Dice	0.8018 (0.7987–0.8046)		0.7912 (0.7880–0.7943)	
	Overall		0.8397		0.8106	
Grounded lesion screening	AbdomenAtlas3.0	Macro F1	0.6383	0.6088	0.05895 (0.03136–0.09357)	0.05536 (0.02878–0.08935)
	LiMT	Macro F1	0.7175 (0.4861–0.9308)	0.7175	0.06250 (0.0000–0.1667)	0.05556 (0.0000–0.1430)
	Overall		0.6779	0.6632	0.06072	0.05546
Grounded visual question answering	AMOS-MM-VQA	Accu.	0.8897 (0.8479–0.9278)	0.8707 (0.8289–0.9087)	0.8707 (0.8289–0.9087)	0.8669 (0.8251–0.9050)
	DeepTumorVQA	Accu.	0.6525 (0.6405–0.6657)	0.6467 (0.6340–0.6593)	0.5475 (0.5344–0.5596)	0.5301 (0.5170–0.5422)
	Overall		0.7711	0.7587	0.7091	0.6985
Grounded report generation	RadGenome	BLEU-4	0.6253 (0.6178–0.6335)	0.6261 (0.6184–0.6340)	0.5969 (0.5888–0.6056)	0.5966 (0.5885–0.6053)
		ROUGE-L	0.4859 (0.4763–0.4945)	0.4830 (0.4732–0.4918)	0.4890 (0.4787–0.4982)	0.4887 (0.4784–0.4979)
		METEOR	0.5460 (0.5368–0.5552)	0.5440 (0.5349–0.5534)	0.5215 (0.5120–0.5315)	0.5212 (0.5117–0.5312)
		RaTEScore	0.7482 (0.7434–0.7533)	0.7419 (0.7370–0.7470)	0.7510 (0.7461–0.7560)	0.7507 (0.7458–0.7557)
		LlamaScore	7.181 (7.093–7.271)	7.154 (7.063–7.240)	6.583 (6.484–6.688)	6.580 (6.481–6.685)
		GREEN	0.4740 (0.4606–0.4868)	0.4624 (0.4499–0.4772)	0.4746 (0.4618–0.4880)	0.4743 (0.4615–0.4877)
	ReXGroundingCT	BLEU-4	0.3517 (0.3257–0.3803)	0.3278 (0.3027–0.3508)	0.2157 (0.1997–0.2315)	0.1943 (0.1777–0.2119)
		ROUGE-L	0.3742 (0.3481–0.4012)	0.3572 (0.3350–0.3808)	0.2413 (0.2276–0.2555)	0.2418 (0.2280–0.2574)
		METEOR	0.3217 (0.2954–0.3505)	0.2975 (0.2719–0.3209)	0.1889 (0.1737–0.2043)	0.1845 (0.1692–0.1994)
		RaTEScore	0.5616 (0.5444–0.5789)	0.5509 (0.5345–0.5677)	0.4530 (0.4413–0.4659)	0.4627 (0.4503–0.4748)
		LlamaScore	6.343 (6.140–6.539)	6.179 (6.007–6.362)	4.853 (4.703–5.007)	4.650 (4.490–4.812)
		GREEN	0.2424 (0.2128–0.2771)	0.2173 (0.1890–0.2465)	0.07198 (0.05739–0.08845)	0.08181 (0.06572–0.09858)
	Overall		0.5070	0.4951	0.4290	0.4266

References

1. Hamamci, I. E. et al. Generalist foundation models from a multimodal dataset for 3D computed tomography. *Nat. Biomed. Eng.* **10**, 1610–1628 (2026).

2. Zhang, X. et al. Development of a large-scale grounded vision language dataset for chest CT analysis. *Sci. Data* **12**, 1636 (2025).

3. Blankemeier, L. et al. Merlin: a computed tomography vision–language foundation model and dataset. *Nature* **652**, 1318–1328 (2026).

4. Chen, Y. et al. Are vision language models ready for clinical diagnosis? A 3D medical benchmark for tumor-centric visual question answering. Preprint at https://doi.org/10.48550/arXiv.2505.18915 (2025).

5. Ji, Y., Ge, C., Li, R. & Luo, P. AMOS-MM: abdominal multimodal analysis challenge: structured description of the challenge design. Zenodo https://doi.org/10.5281/zenodo.10992155 (2024).

6. de la Iglesia Vayá, M. et al. BIMCV COVID-19+: a large annotated dataset of RX and CT images from COVID-19 patients. Preprint at https://doi.org/10.48550/arXiv.2006.01174 (2020).

7. Xin, Y., Ates, G. C., Gong, K. & Shao, W. Med3DVLM: an efficient vision-language model for 3D medical image analysis. *IEEE J. Biomed. Health Inform.* **30**, 2524–2536 (2026).

8. Bai, S. et al. Qwen3-VL technical report. Preprint at https://doi.org/10.48550/arXiv.2511.21631 (2025).

9. Kirillov, A. et al. Segment anything. In *Proc. IEEE/CVF International Conference on Computer Vision*, pp. 4015–4026 (IEEE, 2023).

10. He, Y. et al. VISTA3D: a unified segmentation foundation model for 3D medical imaging. In *Proc. IEEE/CVF Conference on Computer Vision and Pattern Recognition*, pp. 20863–20873 (IEEE, 2025).

11. Wasserthal, J. et al. TotalSegmentator: robust segmentation of 104 anatomic structures in CT images. *Radiol. Artif. Intell.* **5**, e230024 (2023).

12. Bassi, P. R. A. S. et al. RadGPT: constructing 3D image-text tumor datasets. In *Proc. IEEE/CVF International Conference on Computer Vision*, pp. 23720–23730 (IEEE, 2025).

13. Liu, Z. et al. LiMT: a multi-task liver image benchmark dataset. *IEEE J. Biomed. Health Inform.* **30**, 5088–5099 (2026).

14. Baharoon, M. et al. State-of-the-art text-prompted medical segmentation models struggle to ground chest CT findings. In *Proc. Machine Learning for Healthcare Conference*, vol. 298, pp. 1–30 (PMLR, 2025).

15. Isensee, F., Jaeger, P. F., Kohl, S. A. A., Petersen, J. & Maier-Hein, K. H. nnU-Net: a self-configuring method for deep learning-based biomedical image segmentation. *Nat. Methods* **18**, 203–211 (2021).

16. Zhao, Z. et al. Large-vocabulary segmentation for medical images with text prompts. *npj Digit. Med.* **8**, 566 (2025).

17. He, K., Zhang, X., Ren, S. & Sun, J. Deep residual learning for image recognition. In *Proc. IEEE Conference on Computer Vision and Pattern Recognition*, pp. 770–778 (IEEE, 2016).

18. Loshchilov, I. & Hutter, F. Decoupled weight decay regularization. In *Proc. International Conference on Learning Representations* (ICLR, 2019).

19. Bai, F., Du, Y., Huang, T., Meng, M. Q.-H. & Zhao, B. M3D: advancing 3D medical image analysis with multi-modal large language models. Preprint at https://doi.org/10.48550/arXiv.2404.00578 (2024).

20. Jiang, S. et al. Hulu-Med: a transparent generalist model towards holistic medical vision-language understanding. Preprint at https://doi.org/10.48550/arXiv.2510.08668 (2025).

21. Hu, E. J. et al. LoRA: low-rank adaptation of large language models. In *Proc. International Conference on Learning Representations* (ICLR, 2022).

22. Taha, A. A. & Hanbury, A. Metrics for evaluating 3D medical image segmentation: analysis, selection, and tool. *BMC Med. Imaging* **15**, 29 (2015).

23. Rokuss, M. et al. VoxTell: free-text promptable universal 3D medical image segmentation. In *Proc. IEEE/CVF Conference on Computer Vision and Pattern Recognition*, pp. 37538–37557 (IEEE, 2026).

24. Papineni, K., Roukos, S., Ward, T. & Zhu, W.-J. BLEU: a method for automatic evaluation of machine translation. In *Proc. 40th Annual Meeting of the Association for Computational Linguistics*, pp. 311–318 (ACL, 2002).

25. Lin, C.-Y. ROUGE: a package for automatic evaluation of summaries. In *Proc. Text Summarization Branches Out*, pp. 74–81 (ACL, 2004).

26. Grattafiori, A. et al. The Llama 3 herd of models. Preprint at https://doi.org/10.48550/arXiv.2407.21783 (2024).

27. Zhao, W., Wu, C., Zhang, X., Zhang, Y., Wang, Y. & Xie, W. RaTEScore: a metric for radiology report generation. In *Proc. 2024 Conference on Empirical Methods in Natural Language Processing*, pp. 15004–15019 (ACL, 2024).

28. Ma, J. et al. AbdomenCT-1K: is abdominal organ segmentation a solved problem? *IEEE Trans. Pattern Anal. Mach. Intell.* **44**, 6695–6714 (2022).

29. Ji, Y. et al. AMOS: a large-scale abdominal multi-organ benchmark for versatile medical image segmentation. In *Proc. Advances in Neural Information Processing Systems*, vol. 35, pp. 36722–36732 (Curran Associates, 2022).

30. Antonelli, M. et al. The medical segmentation decathlon. *Nat. Commun.* **13**, 4128 (2022).

31. Støverud, K.-H. et al. AeroPath: an airway segmentation benchmark dataset with challenging pathology and baseline method. *PLoS ONE* **19**, e0311416 (2024).

32. Sekuboyina, A. et al. VerSe: a vertebrae labelling and segmentation benchmark for multi-detector CT images. *Med. Image Anal.* **73**, 102166 (2021).

33. Roth, H., Farag, A., Turkbey, E. B., Lu, L., Liu, J. & Summers, R. M. Data from Pancreas-CT (version 2). The Cancer Imaging Archive https://doi.org/10.7937/k9/tcia.2016.tnb1kqbu (2016).

34. Landman, B., Xu, Z., Iglesias, J. E., Styner, M., Langerak, T. R. & Klein, A. (eds). *2015 MICCAI multi-Atlas labeling beyond the cranial vault: workshop and challenge.* Synapse https://doi.org/10.7303/syn3193805 (2015).

35. Simpson, A. L. et al. Preoperative CT and survival data for patients undergoing resection of colorectal liver metastases. *Sci. Data* **11**, 172 (2024).

36. Podobnik, G., Strojan, P., Peterlin, P., Ibragimov, B. & Vrtovec, T. HaN-Seg: the head and neck organ-at-risk CT and MR segmentation dataset. *Med. Phys.* **50**, 1917–1927 (2023).

37. Heller, N., Isensee, F., Tejpaul, R., Wood, A., Papanikolopoulos, N. & Weight, C. 2023 kidney and kidney tumor segmentation challenge. Zenodo https://doi.org/10.5281/zenodo.7840133 (2023).

38. He, Y. et al. Meta grayscale adaptive network for 3D integrated renal structures segmentation. *Med. Image Anal.* **71**, 102055 (2021).

39. Kavur, A. E. et al. CHAOS challenge - combined (CT-MR) healthy abdominal organ segmentation. *Med. Image Anal.* **69**, 101950 (2021).

40. Ma, J. et al. Unleashing the strengths of unlabelled data in deep learning-assisted pan-cancer abdominal organ quantification: the FLARE22 challenge. *Lancet Digit. Health* **6**, e815–e826 (2024).

41. Ma, J. et al. Automatic organ and pan-cancer segmentation in abdomen CT: the FLARE 2023 challenge. Preprint at https://doi.org/10.48550/arXiv.2408.12534 (2024).

42. Rister, B., Yi, D., Shivakumar, K., Nobashi, T. & Rubin, D. L. CT-ORG, a new dataset for multiple organ segmentation in computed tomography. *Sci. Data* **7**, 381 (2020).

43. Armato III, S. G. et al. The lung image database consortium (LIDC) and image database resource initiative (IDRI): a completed reference database of lung nodules on CT scans. *Med. Phys.* **38**, 915–931 (2011).

44. Heimann, T. et al. Comparison and evaluation of methods for liver segmentation from CT datasets. *IEEE Trans. Med. Imaging* **28**, 1251–1265 (2009).

45. Setio, A. A. A. et al. Validation, comparison, and combination of algorithms for automatic detection of pulmonary nodules in computed tomography images: the LUNA16 challenge. *Med. Image Anal.* **42**, 1–13 (2017).

46. Ma, J. et al. Toward data-efficient learning: a benchmark for COVID-19 CT lung and infection segmentation. *Med. Phys.* **48**, 1197–1210 (2021).

47. Liu, P. et al. Deep learning to segment pelvic bones: large-scale CT datasets and baseline models. *Int. J. Comput. Assist. Radiol. Surg.* **16**, 749–756 (2021).

48. Tian, J., Liu, L., Shi, Z. & Xu, F. Automatic Couinaud segmentation from CT volumes on liver using GLC-UNet. In *Proc. International Workshop on Machine Learning in Medical Imaging*, vol. 11861, pp. 274–282 (Springer, 2019).

49. Jaus, A. et al. Towards unifying anatomy segmentation: automated generation of a full-body CT dataset. In *Proc. IEEE International Conference on Image Processing*, pp. 41–47 (IEEE, 2024).

50. Masoudi, M., Pourreza, H.-R., Saadatmand-Tarzjan, M., Eftekhari, N., Zargar, F. S. & Rad, M. P. A new dataset of computed-tomography angiography images for computer-aided detection of pulmonary embolism. *Sci. Data* **5**, 180180 (2018).

51. Pedrosa, J. et al. LNDb challenge on automatic lung cancer patient management. *Med. Image Anal.* **70**, 102027 (2021).

52. Zhuang, X. et al. Evaluation of algorithms for multi-modality whole heart segmentation: an open-access grand challenge. *Med. Image Anal.* **58**, 101537 (2019).

53. Raudaschl, P. F. et al. Evaluation of segmentation methods on head and neck CT: auto-segmentation challenge 2015. *Med. Phys.* **44**, 2020–2036 (2017).

54. Luo, G. et al. Efficient automatic segmentation for multi-level pulmonary arteries: the PARSE challenge. Preprint at https://doi.org/10.48550/arXiv.2304.03708 (2023).

55. Radl, L. et al. AVT: multicenter aortic vessel tree CTA dataset collection with ground truth segmentation masks. *Data Brief* **40**, 107801 (2022).

56. Luo, X. et al. SegRap2023: a benchmark of organs-at-risk and gross tumor volume segmentation for radiotherapy planning of nasopharyngeal carcinoma. *Med. Image Anal.* **101**, 103447 (2025).

57. Lambert, Z., Petitjean, C., Dubray, B. & Kuan, S. SegTHOR: segmentation of thoracic organs at risk in CT images. In *Proc. International Conference on Image Processing Theory, Tools and Applications*, pp. 1–6 (IEEE, 2020).

58. Spahr, A., Ståhle, J., Wang, C. & Kaijser, M. Label-efficient deep semantic segmentation of intracranial hemorrhages in CT-scans. *Front. Neuroimag.* **2**, 1157565 (2023).

59. Aerts, H. J. W. L. et al. Decoding tumour phenotype by noninvasive imaging using a quantitative radiomics approach. *Nat. Commun.* **5**, 4006 (2014).

60. Kiser, K. J. et al. PleThora: pleural effusion and thoracic cavity segmentations in diseased lungs for benchmarking chest CT processing pipelines. *Med. Phys.* **47**, 5941–5952 (2020).

61. Luo, X. et al. WORD: a large scale dataset, benchmark and clinical applicable study for abdominal organ segmentation from CT image. *Med. Image Anal.* **82**, 102642 (2022).

62. Luo, X., Li, Z., Zhang, S., Liao, W. & Wang, G. Rethinking abdominal organ segmentation (RAOS) in the clinical scenario: a robustness evaluation benchmark with challenging cases. In *Proc. International Conference on Medical Image Computing and Computer-Assisted Intervention*, pp. 531–541 (MICCAI, 2024).

63. Hssayeni, M. Computed tomography images for intracranial hemorrhage detection and segmentation (version 1.3.0). PhysioNet https://doi.org/10.13026/w8q8-ky94 (2019).

64. Idris, T. et al. Mediastinal lymph node quantification (LNQ): segmentation of heterogeneous CT data (version 1). The Cancer Imaging Archive https://doi.org/10.7937/qvaz-ja09 (2024).

65. Soler, L. et al. 3D image reconstruction for comparison of algorithm database: a patient-specific anatomical and medical image database. Technical report, IRCAD https://www.ircad.fr/research/data-sets/liver-segmentation-3d-ircadb-01/ (2010).

66. Bakr, S. et al. A radiogenomic dataset of non-small cell lung cancer. *Sci. Data* **5**, 180202 (2018).

67. Li, W. et al. PanTS: the pancreatic tumor segmentation dataset. In *Proc. Advances in Neural Information Processing Systems*, vol. 38, pp. 35806–35828 (Curran Associates, 2025).

68. Lei, W. et al. A synthetic data-driven radiology foundation model for pan-tumor clinical diagnosis. Preprint at https://doi.org/10.48550/arXiv.2502.06171 (2025).

69. de Grauw, M. et al. The ULS23 challenge: a baseline model and benchmark dataset for 3D universal lesion segmentation in computed tomography. *Med. Image Anal.* **102**, 103525 (2025).

70. Roth, H. R. et al. A new 2.5D representation for lymph node detection using random sets of deep convolutional neural network observations. In *Proc. International Conference on Medical Image Computing and Computer-Assisted Intervention*, pp. 520–527 (MICCAI, 2014).

71. Moawad, A. W. et al. Multimodality annotated hepatocellular carcinoma data set including pre- and post-TACE with imaging segmentation. *Sci. Data* **10**, 33 (2023).

72. Li, H. B. et al. QUBIQ: uncertainty quantification for biomedical image segmentation challenge. Preprint at https://doi.org/10.48550/arXiv.2405.18435 (2024).

73. Alves, N. et al. Artificial intelligence and radiologists in pancreatic cancer detection using standard of care CT scans (PANORAMA): an international, paired, non-inferiority, confirmatory, observational study. *Lancet Oncol.* **27**, 116–124 (2026).

74. Wasserthal, J. Training dataset for TotalSegmentator task "liver_lesions". Zenodo https://doi.org/10.5281/zenodo.20272572 (2026).

75. Wang, D. et al. MICCAI2025 MBH-Seg challenge. Zenodo https://doi.org/10.5281/zenodo.15094748 (2025).

76. Jordan, P. et al. Pediatric chest/abdomen/pelvic CT exams with expert organ contours (Pediatric-CT-SEG) (version 2). The Cancer Imaging Archive https://doi.org/10.7937/tcia.x0h0-1706 (2021).

77. Toprak, T. et al. VEELA challenge - vessel extraction and extrication for liver analysis. In *Proc. IEEE International Workshop on Machine Learning for Signal Processing*, pp. 1–6 (IEEE, 2025).

78. Liu, Y. et al. Automatic pelvic fracture segmentation: a deep learning approach and benchmark dataset. *Front. Med. (Lausanne)* **12**, 1511487 (2025).

79. Moawad, A. W. et al. Voxel-level segmentation of pathologically-proven adrenocortical carcinoma with Ki-67 expression (Adrenal-ACC-Ki67-Seg) (version 1). The Cancer Imaging Archive https://doi.org/10.7937/1fpg-vm46 (2023).

80. Gai, X., Liu, J., Li, Y., Meng, Z., Wu, J. & Liu, Z. 3D-RAD: a comprehensive 3D radiology med-VQA dataset with multi-temporal analysis and diverse diagnostic tasks. In *Proc. Advances in Neural Information Processing Systems*, vol. 38, pp. 188177–188218 (Curran Associates, 2025).

81. FLARE-MedFM. FLARE-Task5-MLLM-3D. Hugging Face https://huggingface.co/datasets/FLARE-MedFM/FLARE-Task5-MLLM-3D (2025).

82. Huang, S.-C. et al. INSPECT: a multimodal dataset for patient outcome prediction of pulmonary embolisms. In *Proc. Advances in Neural Information Processing Systems*, vol. 36, pp. 17742–17772 (Curran Associates, 2023).

Supplementary Table 24 | Prompts for report cleaning, question construction and evidence-mask selection. **a,** BIMCV report cleaning and atomic fact extraction. **b,** BIMCV question–answer construction. **c,d,** Supporting-mask assignment for DeepTumorVQA and AMOS-MM-VQA. Bracketed mappings and item fields denote runtime inputs; JSON specifies the requested output format.

Part	Prompt and runtime inputs
a	**BIMCV report cleaning** BIMCV-RG cleaning and fact extraction `You standardize a noisy English radiology report translated from Spanish into a current-CT report-generation target.` `Produce:` `1. a compact list of atomic CT facts;` `2. one fluent cleaned report containing the same information.` `Preserve current CT-visible positive findings, explicit negative findings, diagnostic impressions, uncertainty, anatomy, laterality, distribution, severity, devices, postoperative changes, and reported quantitative values.` `Remove symptoms, medical history, physical examination, vital signs, laboratory or PCR results, treatment information, identifiers, administrative or signature text, acquisition technique, contrast status, reconstruction method, scan-protocol-only text, recommendations, follow-up instructions, and requests for clinical correlation.`  `Remove references to prior examinations and interval change, but retain the current imaging state.`  `Do not introduce a new finding that is unsupported by the report.`

b	**BIMCV question construction**
	BIMCV-VQA generation
	
	`Generate a compact set of natural open-ended question-answer pairs from the provided atomic current-CT facts.`
	
	`Use only the supplied facts.`
	
	`If there are 1-2 facts, generate 1-2 pairs.`
	`If there are 3-5 facts, generate 2-3 pairs.`
	`If there are 6 or more facts, generate 3-5 pairs.`
	
	`Prefer clinically informative and nonredundant questions.`
	
	`Use normal open-ended questions beginning with What, Which, Where, or How.`
	
	`Do not generate yes/no questions.`
	`Do not generate multiple-choice or true/false questions.`
	
	`Preserve laterality, anatomy, uncertainty, negation, numeric values, and units.`
	
	`Do not add information absent from the cited facts.`
	
	`Cite the fact IDs that directly support each answer.`
	`Every answer must have at least one valid evidence ID.`

Continued on next page

Supplementary Table 24 (continued) | Prompts for report cleaning, question construction and evidence-mask selection.

Part	Prompt and runtime inputs

c	**DeepTumorVQA mask matching**
	DeepTumorVQA grounding
	
	`Select CT masks needed to verify each answer.`
	
	`Targets: [ID-to-mask mapping]`
	
	`Each item: q=question, a=answer,`
	`m=available mask IDs.`
	
	`Rules:`
	`- Output only IDs in m.`
	`- Be strict: mentioned anatomy alone is not enough.`
	`- Respect left/right; add both only when both are required.`
	`- Organ measurement -> organ mask.`
	`- Lesion existence/count/size/volume -> lesion mask; never replace missing lesion mask with organ mask.`
	`- Multi-object comparison/sum -> all masks actually needed.`
	`- Prefer specific anatomy over parent anatomy.`
	`- Negative lesion answer does not require parent organ.`
	`- If uncertain, choose fewer.`
	
	`Return JSON only:`
	`{"r":[[ids],...]}`
	
	`One result per input item, same order.`

d	**AMOS-MM-VQA mask matching**
	AMOS-MM-VQA grounding
	
	`Select CT masks needed to visually support each answer.`
	
	`Targets: [ID-to-mask mapping]`
	
	`Semantic synonyms and alternative medical wording count; exact string matching is not required.`
	
	`Rules:`
	`- Return all targets actually needed.`
	`- Respect left/right.`
	`- Add both sides only when bilateral involvement is explicit or clearly implied.`
	`- Do not infer both sides from generic words such as kidney, renal or adrenal.`
	`- Mentioned or nearby anatomy alone is not enough.`
	`- Choose the anatomy providing direct visual evidence, not an organ merely affected as a downstream consequence.`
	`- If none of the targets applies, return [].`
	`- If uncertain, choose fewer.`
	
	`Return JSON only:`
	`{"r":[[ids],...]}`
	
	`One result per input item, same order.`

Supplementary Table 25 | Segmentation, lesion-screening and finding-grounded report generation templates. All 42 entries are shown. Empty braces denote the requested target or finding description. **a,** Target-level segmentation. **b,c,** Presence questions and conditional grounding requests. **d,** Finding-level referring segmentation. **e,** Multiple-target screening. **f,** Finding-grounded and text-only report generation prompts. Numbering is local to each bank.

ID	Template
a | Target-level segmentation	
`segmentation.pure_questions`	
1	`Segment the {} in this CT scan.`
2	`Please segment the {} in this CT volume.`
3	`Generate a 3D mask for the {}.`
4	`Provide the segmentation mask for the {}.`
5	`Delineate the {} in this CT scan.`
6	`Locate and segment the {}.`
b | Single-target presence questions	
`segmentation.classification_questions`	
1	`Is the {} present in this CT scan?`
2	`Can you identify the {} in this CT scan?`
3	`Check whether the {} is present in this CT scan.`
4	`Determine whether the {} is present in this CT volume.`
5	`Does this CT scan contain the {}?`
6	`Can you detect the {} in this CT volume?`
7	`Look for the {} in this CT scan and determine whether it is present.`
8	`Assess whether the {} is visible in this CT scan.`
9	`Is there evidence of the {} in this CT volume?`
10	`Can the {} be found in this scan?`
c | Conditional grounding requests	
`segmentation.classification_grounding_requests`	
1	`If present, please provide its grounded mask.`
2	`If found, also provide the corresponding segmentation mask.`
3	`If it is present, return its 3D mask.`
4	`If detected, provide its voxel-level segmentation.`
5	`If visible, also ground it with a segmentation mask.`
6	`If the target is present, please segment it.`
7	`If identified, provide the corresponding grounded mask.`
8	`If found, also delineate its spatial extent.`
9	`If present, return the corresponding 3D segmentation.`
10	`If detected, also provide its mask.`
d | Finding-level referring segmentation	
`segmentation.referring_segmentation_questions`	
1	`Segment the region described as: "{}"`
2	`Generate a segmentation mask for the following finding: "{}"`
3	`Locate and segment the area corresponding to this description: "{}"`
4	`Provide the segmentation mask for the following text: "{}"`
5	`Identify and segment the target based on this description: "{}"`

Continued on next page

Supplementary Table 25 (continued) | Segmentation, lesion-screening and finding-grounded report generation templates.

ID	Template
6	`Segment the specific pathology or anatomy described here: "{}"`
7	`Ground the following textual description in the 3D volume: "{}"`
8	`Find and segment the region matching this observation: "{}"`
9	`Create a mask for the region described by: "{}"`
10	`Segment the target corresponding to the following report excerpt: "{}"`
e | Multiple-target screening	
`screening_questions`	
`LiMT_Screening`	
1	`Screen the CT and {action} the liver and any potential abnormalities: liver cyst, liver hemangioma, metastatic liver cancer, or primary liver cancer.`
`AbdomenAtlas4Lesion_Screening`	
1	`Screen the CT and {action} any potential abnormalities: kidney lesion, liver lesion, pancreas lesion, or colon lesion.`
`ICH_Screening`	
1	`Screen the CT and {action} any potential intracranial hemorrhage: epidural hemorrhage, subdural hemorrhage, subarachnoid hemorrhage, intraparenchymal hemorrhage, or intraventricular hemorrhage.`
`R3_Screening`	
1	`Screen the CT and {action} any potential thyroid tumors: malignant thyroid tumor or benign thyroid tumor.`
f | Finding-grounded and text-only report generation	
`rex_report_questions`	
Grounded	
1	`Identify and segment all abnormal findings in the lung, pleura, trachea and bronchi.`
Text-only	
1	`Identify all abnormal findings in the lung, pleura, trachea and bronchi.`

Supplementary Table 26 | Question-answering and VQA-grounding templates. All 19 question–answer templates (**a–d**), 20 grounding requests (**e**) and 16 mask-answer templates (**f,g**) are shown. Braced fields specify runtime values. The answer template below each question bank applies to the indicated entries. Grounded VQA combines a base question with a grounding request and the corresponding mask-answer wording.

ID	Template
a | Standard multiple choice	
`vqa.std_mcq`	

1	`{q}`
	`{opts}`
	`Answer with the full option.`

2	`Based on the CT scan, {q}`
	`{opts}`
	`Answer:`

3	`Question: {q}`
	`Options:`
	`{opts}`
	`Select the correct option.`

4	`Looking at this medical image, which of the following is correct?`
	`{q}`
	`{opts}`

Continued on next page

Supplementary Table 26 (continued) | Question-answering and VQA-grounding templates.

ID	Template
5	{q} {opts}

Answer template for all entries: `{ans}`.

b | Multiple choice with reasoning

`vqa.reasoning_mcq`

1	{q} {opts} Select the answer and provide your reasoning based on the scan.
2	{q} {opts} Explain the logic behind your choice.
3	Question: {q} Options: {opts} Answer with explanation.
4	Based on the findings in the imaging, {q} {opts} Give the answer and the reason.
5	Analyze the CT image and answer: {q} {opts}

Answer template for all entries: `{ans}. {reason}`.

c | Binary answers

`vqa.binary`

1	{q} (Yes/No)
2	Based on the medical image, {q}? Yes or No?
3	Please answer with Yes or No: {q}
4	True or False: {q}

Answer templates: entries 1–3, `{ans}`; entry 4, `{ans_tf}`.

d | Open-ended answers

`vqa.open_ended`

1	{q}
2	Question: {q} Answer:
3	Based on the CT volume, {q}
4	Look at the 3D CT scan and answer: {q}
5	According to the 3D scan, {q}

Answer template for all entries: `{ans}`.

e | Optional VQA grounding requests

`vqa_grounding.requests`

1	Ground your answer with the mask.
2	Provide the grounded mask.
3	Include the grounded mask.
4	Give the grounded mask.
5	Also provide the grounded mask.
6	Please provide the grounded mask.
7	Please include the grounded mask.

Continued on next page

Supplementary Table 26 (continued) | Question-answering and VQA-grounding templates.

ID	Template
8	`Please give the grounded mask.`
9	`Ground the answer with the mask.`
10	`Ground your response with the mask.`
11	`Provide the corresponding grounded mask.`
12	`Include the corresponding grounded mask.`
13	`Please respond with the grounded mask.`
14	`Please also provide the grounded mask.`
15	`Please ground your answer with the mask.`
16	`Provide a segmentation mask to ground your answer.`
17	`Include a segmentation mask to ground your answer.`
18	`Please provide a segmentation mask to ground your answer.`
19	`Please accompany your answer with the grounded mask.`
20	`Please accompany your response with the grounded mask.`
f | Single-mask answer wording	
`vqa_grounding.single_mask_answers`	
1	`{grounded} is the grounded mask.`
2	`{grounded} is the corresponding mask.`
3	`{grounded} is the grounding mask.`
4	`{grounded} is the corresponding grounded mask.`
5	`The grounded mask is {grounded}.`
6	`The corresponding mask is {grounded}.`
7	`Grounded mask: {grounded}.`
8	`Grounding: {grounded}.`
g | Multiple-mask answer wording	
`vqa_grounding.multiple_mask_answers`	
1	`{grounded} are the grounded masks.`
2	`{grounded} are the corresponding masks.`
3	`{grounded} are the grounding masks.`
4	`{grounded} are the corresponding grounded masks.`
5	`The grounded masks are {grounded}.`
6	`The corresponding masks are {grounded}.`
7	`Grounded masks: {grounded}.`
8	`Grounding: {grounded}.`

Supplementary Table 27 | Dataset-specific report templates for five source datasets. All 60 templates from the AMOS-MM (**a**), RadGenome (**b**), Merlin (**c**), AbdomenAtlas (**d**) and FLARE-2025 (**e**) banks are shown. Region-list fields specify the annotated anatomical regions requested in the report. INSPECT and BIMCV templates are provided in Supplementary Table 28. Numbering is local to each bank.

ID	Template
a | AMOS-MM	
report_generation.AMOS-MM	
1	Provide a comprehensive radiological narrative for the following regions: {region_list}.
2	Generate a detailed findings report covering the {region_list}, detailing morphology and any lesions found.
3	Describe the radiological appearance across the {region_list} in a continuous, structured format.
4	What are the detailed clinical observations for the organs and tissues in the {region_list}?
5	Deliver a full account of the findings for the following anatomical areas: {region_list}.
6	Provide an in-depth assessment of the {region_list}, focusing on organ size, shape, and parenchymal density.
7	Write a structured findings section for the 3D CT scan, including {region_list}.
8	Outline the visual characteristics and enhancement patterns observed in the {region_list}.
9	Based on this scan, provide a systematic description of findings within the {region_list}.
10	Construct a detailed report summarizing the status of the {region_list} as captured in this volume.
b | RadGenome	
report_generation.Radgenome	
1	Conduct a structural evaluation of the {region_list} based on the CT imaging data.
2	Report the anatomical findings for the following regions: {region_list}.
3	Analyze the {region_list} for any pathological, structural, or sequelae changes.
4	Provide a technical description of the status and integrity of the {region_list}.
5	What are the observations regarding the contours and dimensions of the {region_list}?
6	Evaluate the components of the {region_list} as visualized in this study.
7	Characterize the anatomical appearance of the {region_list} in this scan.
8	Detail the findings for the {region_list}, focusing on their internal structures and boundaries.
9	Provide a systematic assessment of the {region_list} in the specified imaging windows.
10	What is the radiological interpretation for the following anatomical sections: {region_list}?
c | Merlin	
report_generation.Merlin	
1	Assess the radiological status of the following: {region_list}.
2	Provide a concise clinical summary of findings for the {region_list}.
3	What is the condition of the following anatomical structures: {region_list}?
4	Report the imaging observations for the {region_list} in a systematic manner.
5	Summarize the status of the {region_list}, noting any significant abnormalities or normal findings.
6	Document the clinical status for each of the following areas: {region_list}.

Continued on next page

Supplementary Table 27 (continued) | Dataset-specific report templates for five source datasets.

ID	Template
7	Provide a structured overview of the scan findings for the {region_list}.
8	Identify and state any notable findings within the {region_list}.
9	What does the 3D CT study reveal regarding the {region_list}?
10	Give a brief and professional assessment of the following regions: {region_list}.
d | AbdomenAtlas	
report_generation.Abdomenatlas	
1	Provide a quantitative analysis of the abdominal organs, reporting volumes in cc and mean HU values.
2	Generate a morphometric report detailing the size and density of the pancreas, spleen, liver, and kidneys.
3	Extract the volumetric measurements and attenuation characteristics for the major abdominal organs.
4	List the quantitative metrics (volume and mean HU) for the abdominal anatomy and summarize the impression.
5	Report the objective CT measurements for the patient, including organ volumes and any lesion dimensions.
6	Analyze the CT scan to provide a detailed list of organ volumes and densities, noting any enlargements.
7	Create a data-driven report focusing on the Hounsfield Unit statistics and volumetric data of the abdomen.
8	Identify and measure the volume and density of the liver, spleen, pancreas, and kidneys.
9	Provide a technical assessment of the abdominal organs with precise volumetric and densitometric data.
10	Detail the dimensions and mean attenuation values for the abdominal structures and identified lesions.
11	Perform a quantitative volumetric assessment of the abdominal viscera and summarize the findings.
12	Write a report containing only the objective measurements (Volume/HU) and a concluding impression.
13	Calculate and report the size and density parameters for the pancreas, liver, spleen, and kidneys.
14	Assess the abdominal cavity quantitatively, specifying the volume and mean HU for each major organ.
15	Generate a structured report highlighting organ volumes, HU deviations, and lesion measurements.
16	Provide the volumetric segmentation statistics and density analysis for the abdominal CT.
17	Report the calculated volumes and mean attenuation values for the visceral organs.
18	Summarize the quantitative features of the abdomen, including total kidney volume and liver lesion sizes.
19	Present the morphometric data of the patient's abdominal organs, followed by a diagnostic impression.
20	Detail the quantitative findings, specifically focusing on organ volumes (cc) and radiodensity (HU).
e | FLARE-2025	
report_generation.FLARE-2025	
1	Systematically evaluate the following systems and regions for any evidence of abnormality: {region_list}.
2	Provide a structured report assessing the morphology and density of the {region_list}.

Continued on next page

Supplementary Table 27 (continued) | Dataset-specific report templates for five source datasets.

ID	Template
3	`Are there any significant findings in the following areas: {region_list}? Please describe.`
4	`Report on the status of the {region_list}, noting any pathological or normal features identified.`
5	`Describe the appearance of the {region_list} focusing on size, morphology, and regularity.`
6	`Summarize the imaging observations for the following anatomical groups: {region_list}.`
7	`Examine the {region_list} and state the findings for each system.`
8	`Provide the findings for the structures within the {region_list} regarding their parenchymal integrity.`
9	`Check for abnormalities in the {region_list} and provide a standardized findings report.`
10	`Document the radiological status of the {region_list} in this CT volume.`

Supplementary Table 28 | INSPECT, BIMCV and anatomy-grounded report generation templates. a,b, The 20 INSPECT and 5 BIMCV report instructions. **c,** All 40 RadGenome anatomy-grounded report prompts, jointly requesting masks and report text in a specified anatomical order. Region-order fields specify the requested sequence of anatomical regions. Numbering is local to each bank.

ID	Template
a | INSPECT	
`report_generation.INSPECT`	
1	`Generate a numbered radiological impression for this CT scan.`
2	`Write the diagnostic impression for this CTPA as a numbered list.`
3	`Evaluate the scan for pulmonary embolism and list all findings in a numbered impression.`
4	`Provide a structured, numbered clinical impression based on this CT volume.`
5	`Summarize the CTPA findings into a formal, numbered impression.`
6	`Generate the final IMPRESSION section for this scan in a standard numbered format.`
7	`Detail the PE status and any incidental findings in a numbered impression.`
8	`Write a concise, numbered diagnostic summary for this chest CT.`
9	`Formulate the clinical impression for this scan as a numbered list.`
10	`Provide the final radiological impression, noting all findings in numerical order.`
11	`Assess the CT volume and generate a numbered list of diagnostic impressions.`
12	`Summarize all primary and incidental findings into a numbered CTPA impression.`
13	`Draft the IMPRESSION section, listing all radiological findings systematically.`
14	`Generate a comprehensive, numbered impression for this thoracic CT.`
15	`Provide a standard numbered impression evaluating the pulmonary vasculature and secondary findings.`
16	`List the diagnostic conclusions for this CTPA in a numbered impression format.`
17	`Write the final numbered impression, including any clinical recommendations or summaries.`

Continued on next page

Supplementary Table 28 (continued) | INSPECT, BIMCV and anatomy-grounded report generation templates.

ID	Template
18	Evaluate this 3D CT and summarize the impression in a numbered list.
19	Provide a formal, numbered summary of all relevant findings in this scan.
20	Generate the clinical impression, systematically listing all thoracic and abdominal findings.

b | BIMCV

`report_generation.BIMCV`

1	Generate a radiology report for this CT scan.
2	Please write a radiology report for the given CT volume.
3	Describe the radiological findings in this 3D CT scan.
4	Provide a concise radiology report for this CT volume.
5	Generate a clinical imaging report based on the given 3D CT scan.

c | RadGenome anatomy-grounded report generation

`radgenome_grounded_report_questions`

1	Please analyze the CT scan and generate segmentation masks alongside a descriptive report. I need the findings organized in this specific flow: {region_order}.
2	Starting with {region_order}, please segment the anatomical structures in this 3D CT volume and provide the corresponding radiology narratives for each.
3	I'd like a detailed anatomical segmentation and a written summary for this CT study. Please process the regions according to this sequence: {region_order}.
4	For the provided CT data, can you output both the segmentation masks and the clinical report? It's important to follow the order of: {region_order}.
5	Using the 3D CT volume, please perform a region-by-region segmentation and descriptive write-up, adhering to the structure: {region_order}.
6	Could you generate the anatomical masks and report text from this CT? I need the output to follow the progression from {region_order}.
7	Run an anatomical segmentation on this CT dataset and draft a findings report. Please ensure every part follows the requested order: {region_order}.
8	I need the CT findings and masks delivered in a structured format. Please follow this anatomical priority: {region_order}.
9	Map out the segmentation masks for the CT scan and write a narrative report for each segment, following the list: {region_order}.
10	From this 3D CT, I need a combined output of segmented masks and diagnostic text. Please structure it by: {region_order}.
11	Can you segment the following regions in the CT scan—{region_order}—and provide a clinical interpretation for each in that exact order?
12	Generate a segment-by-segment report and mask set from this CT volume, strictly adhering to the specified sequence: {region_order}.
13	I'm looking for a radiology report paired with segmentation masks. Please arrange the findings and masks based on: {region_order}.
14	Process this CT study to yield anatomical segmentations and narrative descriptions, ensuring the workflow follows: {region_order}.
15	Please extract the anatomical masks and corresponding report entries from this CT, keeping the order consistent with: {region_order}.
16	For this CT examination, deliver a set of segmentation masks and the associated report, structured according to the list: {region_order}.
17	I need a comprehensive analysis of the CT scan. Provide masks and report text, following the anatomical layout of {region_order}.
18	Take the 3D CT data and generate a structured report with masks. Please stick to this sequence for the output: {region_order}.

Continued on next page

Supplementary Table 28 (continued) | INSPECT, BIMCV and anatomy-grounded report generation templates.

ID	Template
19	Analyze the CT volume and provide localized segmentation and text descriptions. The order of operations should be: {region_order}.
20	Could you provide a radiology report and mask output for this CT? Please make sure the structures are addressed as: {region_order}.
21	Using the provided CT, create a detailed map of masks and report text. The sequence of interest is {region_order}.
22	I'd appreciate it if you could segment the CT anatomy and write the report findings, following the specific order of {region_order}.
23	Please handle the CT segmentation and report generation. I need it organized structure-by-structure: {region_order}.
24	Your task is to segment the CT and write a descriptive summary. Please follow the anatomical sequence: {region_order}.
25	Can you provide masks and descriptive sentences for the anatomical regions in this CT scan? The desired order is {region_order}.
26	Generate an anatomical segmentation for this 3D CT. The masks and report text should be presented in the order of {region_order}.
27	For the input CT, I need a mask-based analysis and a written report. Please follow this specific path: {region_order}.
28	Perform a region-specific analysis of this CT volume. Produce masks and text findings following: {region_order}.
29	I need a clinical report and anatomical masks derived from this CT. Ensure the segments follow the list: {region_order}.
30	Working with the provided CT, please output masks and findings descriptions in the following anatomical order: {region_order}.
31	Generate both the segmentation masks and the radiology report text. Please align your output with: {region_order}.
32	I need the CT dataset processed into anatomical masks and descriptive text. Follow this precise order: {region_order}.
33	For the given CT, provide segmentation and matching narrative reports, structured as: {region_order}.
34	Please produce anatomical masks and the corresponding imaging report, maintaining the sequence: {region_order}.
35	Extract the segmentation results and write the report for this CT, following the structural sequence: {region_order}.
36	Can you analyze the CT scan and provide masks and clinical text for each region? Please follow the sequence: {region_order}.
37	Deliver a report and segmentation masks from the CT scan. The structures must be analyzed in this order: {region_order}.
38	I'm requesting a mask-based segmentation and a clinical report for this CT scan, ordered by: {region_order}.
39	Please segment the anatomy and draft the findings for this CT study, following the specified list of {region_order}.
40	Process the 3D CT to output masks and a descriptive report, ensuring the sections follow: {region_order}.

Supplementary Table 29 | Clinical report cleaning, translation and anatomical structuring. a, Prompt template for Chinese reports from centre 5. **b,** Thoracic and abdominal region lists. Placeholders denote runtime inputs; the JSON block specifies the requested output format.

Part	Prompt and runtime inputs

a	**Clinical report processing**
	Cleaning, translation and anatomical structuring template
	`You are an expert radiology report editor.`
	`Clean, translate and structure the provided Chinese CT report into professional English.`
	`Instructions:`
	`- Remove patient identifiers, administrative information, non-imaging clinical details, acquisition protocols, recommendations and prior-study comparisons.`
	`- Preserve current CT findings, diagnostic impressions, anatomy, laterality, morphology, severity, uncertainty and all reported measurements.`
	`- Translate using standardized radiological terminology without adding unsupported findings.`
	`- Assign each finding to the most appropriate anatomical region from the provided list.`
	`- Consolidate findings and impressions within each region, avoiding unnecessary repetition.`
	`- Preserve explicit normal findings; do not infer normality from an unmentioned region.`
	`- Return valid JSON containing the region-wise descriptions.`
	`Required regions:`
	`{region_list}`
	`Original Chinese report:`
	`{original_report}`
	`Output:`
	`{`
	`  "process": {`
	`  "<region_name>": "<English description>"`
	`  }`
	`}`

b	**Region-list values**
	Chest and abdomen use their corresponding list
	`Chest (10): mediastinum, abdomen, lung, bone, heart, trachea and bronchi, esophagus, pleura, breast, thyroid.`
	`Abdomen (13): lower thorax, liver and biliary tree, gallbladder, spleen, pancreas, adrenal glands, kidneys and ureters, gastrointestinal tract, peritoneum, pelvic, vasculature, lymph nodes, musculoskeletal.`
	The relevant list populates `{region_list}`; the source report populates `{original_report}`. Region-wise English descriptions are concatenated to form the structured whole-report reference.