

New Orthogonal Multiwavelet Filters Derived By Matrix Spectral Factorization

Vasil Kolev*, Todor Cooklev and Fritz Keinert

Institute of Information and Communication Technologies, Bulgarian Academy of Sciences, Bl. 2, Acad. G. Bonchev St., 1113 Sofia, Bulgaria

Wireless Technology Center, Purdue University, Fort Wayne, IN 46805, USA

Department of Mathematics, Iowa State University, Ames, IA 50011, USA

ARTICLE INFO

Keywords:

Supercompact multiwavelet
spectral factorization
classical and exact Bauer's method
signal denoising
image compression
subband-based edge detection

ABSTRACT

The paper considers the construction of two new orthogonal multiwavelets with supercompact support by using the Fast Bauer's method for matrix spectral factorization on the matrix product filter of the orthogonal CL multiwavelet filter. The new multiwavelets possess orthogonality, symmetry/antisymmetry, and one of them provides better coding and smoothness than other supercompact multiwavelets. The performance of the new multiwavelet filters in subband-based edge detection, grayscale and color image compression and 1D and 2D signal denoising is compared with the GHM, SA4, CL, Integer Haar and Alpert multifilters. The comparative analysis shows that new multiwavelets can provides better human visual measures, SSIM and MS-SSIM in image compression and denoising applications.

1. Introduction

Multiwavelets, as extensions of scalar wavelets, can possess important features such as orthogonality, compact support, symmetry and high-order vanishing moments. Their structure makes them useful in mathematical theory and many applications. Supercompact multiwavelets have support $[0, 1]$ and retain the spatial compactness and orthogonality of Haar wavelets [1]. They are simple to implement in applications on a finite interval and/or for analysis of signals that are piecewise continuous, since no special treatment of external or internal boundaries is necessary. This compactness becomes more important where strong gradients occur, such as near shock waves and in boundary regions. Numerical evidence also illustrates that these supercompact multiwavelets could be utilized without modification even near boundaries [2].

Supercompact multiwavelets are orthogonal vectors of r scaling functions $\phi(t) = [\phi_0, \phi_1, \dots, \phi_{r-1}]^*$ and r wavelet functions $\psi(t) = [\psi_0, \psi_1, \dots, \psi_{r-1}]^*$, which belong to $L_2([0, 1])^r$ [3] and satisfy matrix recursion equations with only two coefficients

$$\begin{aligned}\phi(t) &= \sqrt{2} [H_0 \phi(2t) + H_1 \phi(2t - 1)], \\ \psi(t) &= \sqrt{2} [G_0 \phi(2t) + G_1 \phi(2t - 1)].\end{aligned}$$

The multiscaling symbols $H(z) = H_0 + H_1 z^{-1}$ and $G(z) = G_0 + G_1 z^{-1}$ are linear matrix polynomials.

The orthogonality conditions $\langle \phi_i, \phi_j \rangle = \langle \psi_i, \psi_j \rangle = \delta_{ij}$ and $\langle \phi_i, \psi_j \rangle = 0$ for $i, j = 0 : r - 1$, can be expressed in terms of the coefficients as

$$H_0 H_0^* + H_1 H_1^* = G_0 G_0^* + G_1 G_1^* = I, \quad H_0 G_0^* + H_1 G_1^* = 0.$$

The Hermitian transpose of a matrix polynomial $A(z) = \sum_j A_j z^j$ is given by $A(z)^* = \sum_j A_j^* z^{-j}$. The matrix product filter $P(z)$ is defined as the autocovariance function

$$P(z) = H(z)H(z)^* = P_1 z^{-1} + P_0 + P_{-1} z, \quad (1)$$

✉ vasil.kolev@iict.bas.bg, kolev_acad@abv.bg (V. Kolev); cooklevt@pfw.edu (T. Cooklev); keinert@iastate.edu (F. Keinert)

ORCID(s): 0000-0002-4619-3579 (T. Cooklev); 0000-0003-3116-967X (F. Keinert)




Figure 1: The main steps in the construction of new supercompact multiscaling and multiwavelet functions

which is a positive semidefinite polynomial matrix. The matrix product coefficients are

$$P_0 = H_0 H_0^* + H_1 H_1^*, \quad P_1 = H_0 H_1^*, \quad P_{-1} = P_1^* = H_1 H_0^*. \quad (2)$$

It is Hermitian, and satisfies the half band condition $P(z) + P(-z) = 2I$ [4].

The stability and causality conditions translate to requiring that $P(z)$ has all its poles and zeros inside the unit circle. *Matrix Spectral Factorization* (MSF) refers to finding a matrix spectral factor $H(z)$, given $P(z)$. The existence of such a factor is guaranteed under a condition on $P(z)$ due to Paley and Wiener, requiring that $\log(\det(P(z)))$ have a finite integral over an interval of length 2π . MSF is a key step for the construction of minimum phase spectral factors from matrix product filters. Unlike the spectral factorization of scalar Hermitian polynomial functions, where an explicit formula exists, there is no explicit expression for matrix spectral factorization. However, for low-degree para-Hermitian polynomial matrix polynomials a fast algorithm has been constructed in [5, 6].

Our main goal in this paper is to present a new approach for the construction of new supercompact orthogonal multiwavelets by matrix spectral factorization of a suitable product filter. The main steps are shown in Fig. 1.

The new multiwavelets are obtained from the product filter of the Chui-Lian (CL) multiwavelet filter [7]. This immediately leads to three main problems:

1. Does the Fast Bauer's method for matrix spectral factorization converge in this case?
2. Is possible to obtain an orthogonal multiscaling filter with exact coefficients from existing multifilters?
3. Do the newly derived multiwavelets have better coding gain than other well-known orthogonal or biorthogonal multifilters?

We have the following main contributions:

- Obtain the matrix product filter of the CL orthogonal multiwavelet filter;
- Apply Fast Bauer's method for spectral factorization to find two new supercompact multiscaling functions with exact recursion coefficients;
- Find the complementary new supercompact multiwavelet functions;
- Compare the performance of new multiwavelet filters in subband-based edge detection, 1D signal denoising and grayscale and color image denoising and compression with the Integer Haar, Alpert, GHM, SA4 and CL multiwavelet filter banks.

2. Theory of Matrix Spectral Factorization

Spectral factorization is an important mathematical tool in signal processing, control theory, and other areas. The fundamental theorem of the spectral factorization is the Fejér-Riesz theorem for positive definite functions. The terminology comes from prediction theory, where the nonnegative function plays the role of a spectral density for a multidimensional stationary stochastic process. Fejér [8] first showed the importance of the class of trigonometric polynomials that admit only positive real values; as a theorem it was proved by Riesz [9]. The definition of a trigonometric univariate polynomial is

$$v(z) = \sum_{k=-N}^N v_k z^k.$$

When the function $v(z)$ is real for all $z \in \mathbb{T}$, where $\mathbb{T}$ is the unit circle in the complex plane, then the coefficients satisfy $\bar{v}_k = v_{-k}$ for all k . If $v(z) \geq 0$ for all $z \in \mathbb{T}$, the *spectral factorization* of the function is

$$v(z) = p(z)p(z)^*,$$

where $p(z) = \sum_{k=0}^N p_k z^k$ is called a *scalar spectral factor*.

Matrix spectral factorization (MSF) refers to finding a matrix spectral factor $H(z)$, given $P(z)$, so that

$$P(z) = H(z)H(z)^*.$$

Such a factorization exists if $P(z)$ is positive definite on $\mathbb{T}$, absolutely integrable and satisfies the Paley-Wiener condition [10, 11, 12]

$$\left| \int_{-\pi}^{\pi} \ln \det P(e^{i\omega}) d\omega \right| < \infty.$$

There is a unique minimal phase solution $H_M(z)$. All other solutions are of the form

$$H(z) = H_M(z)U,$$

where U is a unitary matrix multiplier.

Usually, $P(z)$ is a half-band filter with $P_0 = I$, $P_{2k} = 0$ for $k \neq 0$.

2.1. Classical Bauer's method of Matrix Spectral Factorization

Let the multiscaling function $H(z)$ and its complex conjugate $H^*(z)$ be

$$\begin{aligned} H(z) &= H_0 + H_1 z^{-1} + \dots + H_m z^{-m}, \\ H^*(z) &= H_0^* + H_1^* z + \dots + H_m^* z^m, \end{aligned} \quad (3)$$

where H_k are real matrices of size $r \times r$ and $z \in \mathbb{C}$. Let $\mathbb{T} = \{z : |z| = 1\}$ be the unit circle in the complex plane. For $z \in \mathbb{T}$ we have $z^{-1} = \bar{z}$. Then the matrix product

$$P(z) = H(z)H^*(z) = P_m^* z^{-m} + \dots + P_1^* z^{-1} + P_0 + P_1 z + \dots + P_m z^m \quad (4)$$

is called the *matrix product filter*, where each matrix coefficient is determined by

$$P_k = H_{m-k} H_m^* + H_{m-k-1} H_{m-1}^* + \dots + H_0 H_k^*. \quad (5)$$

Supercompact multifilters represent the simplest case $m = 1$, with the coefficients P_0 and P_1 shown in (2).

Bauer's matrix spectral factorization finds the Cholesky factor L_n of the $(n+1) \times (n+1)$ banded block Toeplitz matrix

$$T_n = \begin{pmatrix} P_0 & P_1 & 0 & 0 & 0 \\ P_1^* & P_0 & P_1 & 0 & 0 \\ 0 & \ddots & \ddots & \ddots & 0 \\ 0 & 0 & 0 & P_1^* & P_0 \end{pmatrix} = L_n L_n^*, \quad (6)$$

where

$$L_n = \begin{pmatrix} H_0^{(0)} & 0 & 0 & 0 \\ H_1^{(1)} & H_0^{(1)} & 0 & 0 \\ 0 & \ddots & \ddots & 0 \\ 0 & 0 & H_1^{(n)} & H_0^{(n)} \end{pmatrix}$$

is a lower bidiagonal matrix.

The last row of L_n converges to the true matrix coefficients $H_0^{(n)} \rightarrow H_0$, $H_1^{(n)} \rightarrow H_1$ as $n \rightarrow \infty$.

The classical Bauer's method is very effective for nonsingular matrix product filters $P(z)$. Convergence can be very fast, so n does not need to be large to accurately estimate the spectral factors.

However, since in our applications the matrix product filters are required to possess multiple zeros on the unit circle, the method converges very slowly and requires a very large Toeplitz matrix. In one example, it required a matrix of size around $10^6 \times 10^6$ for achieving a precision of 10^{-13} .

In addition, for larger multiplicity $r \geq 3$ high precision often cannot be achieved. In one example in [6] with $r=5$, $m=1$ and a decuple zero singularity, the maximum achievable accuracy in the classical Bauer's method was around 10^{-3} .

2.2. Fast Bauer's method of Matrix Spectral Factorization

Obviously the Cholesky decomposition (6) can be realized row by row. If we assume that $H_0^{(n)}$, $H_1^{(n)}$ have already been found, with the initial value $H_0^{(0)}$ obtained as the Cholesky factor of P_0 , then it follows that

$$H_1^{(n+1)} = P_1^* \left[H_0^{(n)} \right]^{-*} \quad (7)$$

from $P_1^* = H_1^{(n+1)} \left[H_0^{(n)} \right]^*$, and

$$H_0^{(n+1)} \left[H_0^{(n+1)} \right]^* = P_0 - H_1^{(n+1)} \left[H_1^{(n+1)} \right]^* \quad (8)$$

from $P_0 = H_0^{(n+1)} \left[H_0^{(n+1)} \right]^* + H_1^{(n+1)} \left[H_1^{(n+1)} \right]^*$.

Then (7) substituted into (8) leads to

$$H_0^{(n+1)} \left[H_0^{(n+1)} \right]^* = P_0 - P_1^* \left[H_0^{(n)} \right]^{-*} \left[H_0^{(n)} \right]^{-1} P_1. \quad (9)$$

If we define $X^{(n+1)} = H_0^{(n+1)} \left[H_0^{(n+1)} \right]^*$, then

$$X^{(n+1)} = P_0 - P_1^* \left[X_0^{(n)} \right]^{-1} P_1. \quad (10)$$

The limit $n \rightarrow \infty$ leads to the solution $X^{(n+1)} \rightarrow X$ of the nonlinear matrix equation (NME)

$$X = P_0 - P_1^* X^{-1} P_1. \quad (11)$$

Our focus in this paper is the supercompact case $m=1$, but we note that the general case can be reduced to this by replacing the original coefficients with block matrices. For $m=3$, for example, we set

$$\hat{P}_{-1} = \hat{P}_1^* = \begin{bmatrix} P_3^* & P_2^* & P_1^* \\ 0 & P_3^* & P_2^* \\ 0 & 0 & P_3^* \end{bmatrix}, \quad \hat{P}_0 = \begin{bmatrix} P_0 & P_1 & P_2 \\ P_1^* & P_0 & P_1 \\ P_2^* & P_1^* & P_0 \end{bmatrix}, \quad \hat{P}_1 = \begin{bmatrix} P_3 & 0 & 0 \\ P_2 & P_3 & 0 \\ P_1 & P_2 & P_3 \end{bmatrix}.$$

The resulting coefficients $\hat{H}_0$, $\hat{H}_1$ have the form

$$\hat{H}_1 = \begin{bmatrix} H_3 & H_2 & H_1 \\ 0 & H_3 & H_2 \\ 0 & 0 & H_3 \end{bmatrix}, \quad \hat{H}_0 = \begin{bmatrix} H_0 & 0 & 0 \\ H_1 & H_0 & 0 \\ H_2 & H_1 & H_0 \end{bmatrix}.$$

Once $\hat{H}_0, \hat{H}_1$ have been computed, we can extract the original H_0, H_1, H_2, H_3 as submatrices.

Setting up the symmetrical nonsingular matrix X from the above Fast Bauer's method with symbolic entries x_{ij} converts (11) into a system of nonlinear equations. For small matrix sizes this can be solved with exact values by a computer algebra system (CAS).

For larger r , (11) could be solved numerically by fixed point iteration, which is identical to the original Bauer's method, but Newton's method converges much faster and gives more accurate results. See [6] for more details.

3. Design of the supercompact multiwavelet filter

3.1. The matrix product filter of the CL multiscaling function

The matrix coefficients of the orthogonal CL scaling functions [7]

$$H_{CL}(z) = \frac{1}{4} \left[\begin{pmatrix} 0 & 2 + \sqrt{7} \\ 0 & 2 - \sqrt{7} \end{pmatrix} + \begin{pmatrix} 3 & 1 \\ 1 & 3 \end{pmatrix} z^{-1} + \begin{pmatrix} 2 - \sqrt{7} & 0 \\ 2 + \sqrt{7} & 0 \end{pmatrix} z^{-2} \right]$$

lead to the matrix product filter

$$P_{CL}(z) = \frac{1}{8} \left[\begin{pmatrix} 4 & 1 + \sqrt{7} \\ -(1 + \sqrt{7}) & -\sqrt{7} \end{pmatrix} z^{-1} + \begin{pmatrix} 8 & 0 \\ 0 & 8 \end{pmatrix} + \begin{pmatrix} 4 & -(1 + \sqrt{7}) \\ 1 + \sqrt{7} & -\sqrt{7} \end{pmatrix} z \right]$$

with

$$\det(P_{CL}(z)) = (4 - \sqrt{7}) \frac{(1 + z)^4}{32z^2},$$

which satisfies the regularity criterion that $\det(P(z))$ has a quadruple zero at $z = -1$.

3.2. New supercompact multiscaling functions from the Exact Bauer's method

In order to obtain multiscaling functions with exact coefficients, we solve eq.(11) by an appropriate CAS, for example the Matlab Symbolic Toolbox. The solution is a symmetric matrix

$$X = \frac{1}{8} \begin{pmatrix} 4 & \sqrt{7} + 1 \\ \sqrt{7} + 1 & 4 \end{pmatrix}.$$

The standard Cholesky factor is

$$H_0^{(1)} = \frac{\sqrt{2}}{8} \begin{pmatrix} 4 & 0 \\ \sqrt{7} + 1 & \sqrt{8 - 2\sqrt{7}} \end{pmatrix}$$

The square root has two possible values

$$\sqrt{8 - 2\sqrt{7}} = \pm(\sqrt{7} - 1)$$

which leads to two solutions

$$H_0^{(1)} = \frac{\sqrt{2}}{8} \begin{pmatrix} 4 & 0 \\ \sqrt{7} + 1 & \sqrt{7} - 1 \end{pmatrix}$$

and

$$H_0^{(2)} = \frac{\sqrt{2}}{8} \begin{pmatrix} 4 & 0 \\ \sqrt{7} + 1 & -(\sqrt{7} - 1) \end{pmatrix}$$




Figure 2: (a) CL multiscaling function ϕ_{CL} (red) and multiwavelet function ψ_{CL} (blue); (b) New supercompact multiscaling function $\phi^{(1)}$ (red) and multiwavelet function $\psi^{(1)}$ (blue); (c) New supercompact multiscaling function $\phi^{(2)}$ (red) and multiwavelet function $\psi^{(2)}$ (blue);

The second coefficients $H_1^{(i)}$ is found from (7) as $H_1^{(i)} = P_1^* [H_0^{(i)}]^{-*}$. We call the resulting supercompact multiscaling function *New1* and *New2*.

The resulting coefficients, as well as the coefficients of the corresponding multiwavelet functions and the determinants of the symbols are tabulated in Tab 1. It is important to note that the determinants of both multiscaling functions include a factor of $(1 + z^{-1})/2$, which according to [4] leads to regular scaling functions and a desirable supercompact multiscaling function.

3.3. New supercompact multiwavelet functions

In order to obtain the wavelet matrix coefficients G_0 and G_1 we can employ the QR decomposition as in [5], or in the present case, the orthogonal vectors method. This multiwavelet possesses the all the desirable properties of the Alpert multiwavelet. The recursion coefficients of the new orthogonal multiscaling and multiwavelet functions obtained by QR decomposition are shown in Table 1; the functions are shown in Fig. 2.

4. Performance Analysis

4.1. Numerical errors

The classical Bauer's method is applied to the spectral factorization of the matrix product filter. The numerical errors obtained for the matrix product filter $P(z)$ and the matrix spectral factor $H(z)$ are defined as

$$\begin{aligned} \epsilon_H^{(n)} &= |H - H^{(n)}| \\ \epsilon_P^{(n)} &= |P - P^{(n)}|. \end{aligned} \tag{12}$$

Table 1

The matrix coefficients of the two new multiscaling and multiwavelet functions *New1* and *New2*.

Multiscaling Functions	
$H^{(1)}(z)$	$H^{(2)}(z)$
$H_0^{(1)} = \frac{\sqrt{2}}{8} \begin{pmatrix} 4 & 0 \\ \sqrt{7}+1 & \sqrt{7}-1 \end{pmatrix}$	$H_0^{(2)} = \frac{\sqrt{2}}{8} \begin{pmatrix} 4 & 0 \\ \sqrt{7}+1 & 1-\sqrt{7} \end{pmatrix}$
$H_1^{(1)} = \frac{\sqrt{2}}{8} \begin{pmatrix} 4 & 0 \\ -(\sqrt{7}+1) & \sqrt{7}-1 \end{pmatrix}$	$H_1^{(2)} = \frac{\sqrt{2}}{8} \begin{pmatrix} 4 & 0 \\ -(\sqrt{7}+1) & 1-\sqrt{7} \end{pmatrix}$
$\det(H^{(1)}(z)) = \det(H_0^{(1)} + H_1^{(1)}z^{-1})$ $= \frac{\sqrt{7}-1}{\sqrt{2}}(1+z^{-1})^2$	$\det(H^{(2)}(z)) = \det(H_0^{(2)} + H_1^{(2)}z^{-1})$ $= \frac{1-\sqrt{7}}{\sqrt{2}}(1+z^{-1})^2$

Multiwavelet Functions	
$G^{(1)}(z)$	$G^{(2)}(z)$
$G_0^{(1)} = \frac{\sqrt{2}}{8} \begin{pmatrix} 0 & -4 \\ -(\sqrt{7}-1) & \sqrt{7}+1 \end{pmatrix}$	$G_0^{(2)} = \frac{\sqrt{2}}{8} \begin{pmatrix} 0 & -4 \\ -(\sqrt{7}-1) & -(\sqrt{7}+1) \end{pmatrix}$
$G_1^{(1)} = \frac{\sqrt{2}}{8} \begin{pmatrix} 0 & 4 \\ \sqrt{7}-1 & \sqrt{7}+1 \end{pmatrix}$	$G_1^{(2)} = \frac{\sqrt{2}}{8} \begin{pmatrix} 0 & 4 \\ \sqrt{7}-1 & -(\sqrt{7}+1) \end{pmatrix}$
$\det(G^{(1)}(z)) = \det(G_0^{(1)} + G_1^{(1)}z^{-1})$ $= \frac{\sqrt{7}-1}{\sqrt{2}}(1-z^{-1})^2$	$\det(G^{(2)}(z)) = \det(G_0^{(2)} + G_1^{(2)}z^{-1})$ $= \frac{\sqrt{7}-1}{\sqrt{2}}(1-z^{-1})^2$

The values are shown in Fig. 3, in a log-log plot. The convergence of the two errors ϵ_H and ϵ_P is well modeled by the linear fit $p_{\epsilon_H(x)} = -1.0101n$ and $p_{\epsilon_P(x)} = -2.0454n$ respectively, where the logarithm of the error $\log(\epsilon_H^{(n)})$ has slope (-1) (a red line), while $\log(\epsilon_H^{(n)})$ has a slope of (-2) (cyan line).

In [6], after applying the fast Bauer's method using a general discrete algebraic Riccati equation (GDARE) in Matlab R2018a, the best achieved precision for the matrix spectral factor $\epsilon_H^{(n)} = 6.16 \times 10^{-5}$ and the matrix product filter $\epsilon_P^{(n)} = 6.66 \times 10^{-14}$, while with Matlab R2010a,b the best precisions are $\epsilon_H^{(n)} = 6.7077 \times 10^{-5}$ and $\epsilon_P^{(n)} = 3.3307 \times 10^{-16}$, respectively.

It is important to note that the classical Bauer's method for spectral factorization obtained errors for $n = 10,000$ of $\epsilon_H^{(n)} = 8.396 \times 10^{-5}$ and $\epsilon_P^{(n)} = 2.055 \times 10^{-9}$ shown in Fig. 3. They can easily be modeled by regression functions. The empirical results show minimal possible errors $\epsilon_H^{(n)} = 9.6145 \times 10^{-17}$ and $\epsilon_H^{(n)} = 6.979 \times 10^{-5}$ that can be achieved for $n = 38.639$.

4.2. Statistical Characteristic Analysis of the Test Images

4.2.1. Histograms

The histogram of an image is a one-dimensional statistical characteristic of the given image and represents the frequency of intensity in separate groups of pixels in the image. The groups usually contain the same number of adjacent intensity levels. The most representative (precise) are histograms on all possible single levels.

As an application, the histogram values are in some (engineering) range of intensities of useful (measured) quantities, of course appropriately quantized. Visually, the histogram allows for different interpretations by pixels and groups of pixels, for example how the brightness (luminance) correlates with the number of pixels in it, as well as some easily apparent characteristics such as contrast, dynamic range and saturation effects, which make problems easily visible and facilitate the analysis of the image (and the problem it reflects).

The color of images is represented by color models. The most popular of them is RGB, but we can also mention many others, YUV, HSV, etc., emphasizing different characteristics of color. While a halftone image is a scalar function over the discrete area of pixels, color images are usually vector functions. RGB is a 3-dimensional function, one vector for each of the 3 channels (all halftones).




Figure 3: Influence of the size n of the Toeplitz matrix on the precision of the matrix spectral factor (in log-log format): The numerical error of matrix spectral factor $\epsilon_H^{(n)}$ (red) and of the matrix product filter $\epsilon_P^{(n)}$ (cyan)

For better decomposition at multiple levels of detail and better encoding as well, in the future we could apply a suitable 3D transform based on 3D convolution and extraction of color channels by a *inflated 3D convNet (I3D) model* [26] instead of extraction of RGB channels.

The abscissa Ox shows the pixel intensities in the range from 0 to 255 for each color channel (R, G, B). The value 0 corresponds to the "black" level, and 255 to the "white" level. If the image is halftone, successive levels of "gray" are placed between them. If the channel is color, the "gray" level represents the corresponding base colors (R, G, B) in the given color image. The ordinate Oy shows the frequency of the corresponding pixel intensity levels. Color histograms are triplets of halftone histograms, one for each color (red (R), green (G), and blue (B) channels). The size of the intervals in the figures is not specified and the value of each pixel is displayed separately. Some characteristic tests for seven color and seven grayscale images with different sizes and textures are illustrated in Fig. 4(a)–(f) and Fig. 5(a)–(g).

- **Bart** Run has an extremely low dynamic range with only 14 intensity values;
- **Cameraman** has 2 isolated spikes, as can be seen from the histogram in Fig. 4(b). The big peak on the far left side of the graph means that the image is made up of a lot of shadows or black tones. Therefore the image has low brightness. The peaks in the middle of the histogram indicate lots of midtones which mean muted color grays;
- **Kiel** has distributed intensity values with 2 isolated peaks. The histogram in Fig. 4(c) shows peaks that are cut off on the left or right side, which means an underexposed image;

- House is underexposed (intensity values are clustered) and has high contrast for all three colors. The blue channel, see Fig. 4(d), has many pixels in the high intensity region, while the red and green spectra remain mostly low. This means that the image has shades of blue, but not much red and green.
- Cara has saturated and underexposed pixel values (intensity values are clustered) and has also an isolated peak for all three colors. This is shown in Fig. 4(e), where the histogram peaks in the middle of each color indicate a large presence of dark pixels for each color. Each color has peaks scattered around the middle, indicating a midtone, which means a muted color.
- Voit has properly exposed normal contrast (intensity values are scattered).
- House has low contrast histogram by a slight increase in contrast. The big peak on the right side shown in Fig. 5(a).
- Couple has low contrast histogram gaps caused by a slight increase in contrast shown in Fig. 5(b).
- Brick wall has a normal contrast and many pixels in the low intensity region which means that the image has shades or dark color tones shown in Fig. 5(c).
- Texture 1 has low high and overexposed contrast for R, G and B channels respectively. The blue channel has many pixels with in the lowest intensity region, while the red and green channels remain mostly low shown in Fig. 5(d).
- Subwater has normal contrast for channels, blur the red and blue channels and isolated peak shown in Fig. 5(e).
- Texture 4 Texture 4 has extremely low dynamic range for R, G, B channels with 2 isolated peaks shown in Fig. 5(f).
- Texture 5 histogram gaps caused by a slight increase in contrast shown in Fig. 5(g).

4.2.2. Adjacent Pixel Correlations

Neighboring pixel correlation refers to the relationship between two adjacent pixels in horizontal and vertical coordinates, respectively. The more the pixel values remain concentrated near the linear fit, the stronger the correlation between neighboring pixels in the image; conversely, the more pixels are scattered throughout the plot, the weaker the correlation between neighboring pixels in the image.

The correlation coefficient R^2 is a primary measure of how well a regression model fits the data. This correlation coefficient is simply the square of Pearson's correlation coefficient R . To check the existence of a high correlation among neighboring pixels in the test images, a pixel correlation analysis test has performed. The correlation analysis of adjacent pixels of this encryption scheme is shown in Fig. 6(a)-(f). High correlation among pixels in the original image is represented by the linear fit it converges to. The scatterplot of Fig. 6 leads to the values of R^2 for the regression models shown in Table 2.

The correlation coefficient R^2 , which is the square of the Pearson correlation coefficient, is a basic measure of how well a regression model fits the data; when $R^2 = 1$, the regression line predicts a perfect fit of the pixel values. The values of the regression lines for all test images are shown in Table 2. Fig. 6 shows the scatter plot of all regression models from Table 2.

The images Bart Run and Cara have quite a bit of scatter, but the scatter plot of Fig. 6(a) and (e) (red, green, and blue channels) shows a clear linear trend with correlation 88% – 98%. The images Kiel and Voit have a moderate positive correlation in the interval 62% – 68%. It's not a very strong relationship, as some values show no correlation. Obviously, for these two images there is quite a lot of scatter, but the trend is reasonably linear, despite the fact that the big number of data points makes it difficult to do a full correlation assessment. Fig. 5(b) and (f) (red, green, and blue channels) shows that there is no correlation in some subset pixels and a strong correlation in the whole pixels data set.

The images Bart Run and Cara have quite a large scatter, but the scatter plot in Fig. 6(a) and (e) (RGB channels) shows a clear linear trend with a correlation of 88% – 98%. The two images Kiel and Voit have a moderate positive correlation in the interval 62% – 68%. This is not a very strong relationship, as some values show no correlation. It is clear that there is quite a bit of scatter for these two images, but the trend is relatively linear, despite the large number




Figure 4: Test images and their histograms; (a)Bart Run(128×128); (b)Cameraman(256×256); (c)Kiel(512×512);(d)House(256×256); (e)Cara(256×256);(f)Voit(256×256).

of data points, which makes it difficult to fully assess the correlation. Fig. 6(b) and (f) (RGB channels) show that there is no correlation in some subsets of pixels and a strong correlation in the entire pixel dataset.

The grayscale images House,Cameraman, Texture 5 and color image (for R,G,B channels)Subwater have a strong correlation shown of Fig. 7. The grayscale image House and color images (for R,G,B channels) Texture 1, Texture 4 there is no correlation of pixels and a strong correlation in the entire pixel dataset shown of Fig. 7.

4.3. General multifilter characteristics

We compare the performance of the proposed new multifilters with some well-known orthogonal and biorthogonal filter banks. The results are given in Table 3, with the criteria being coding gain (CG), Sobolev smoothness (S) [13], symmetry/anti-symmetry (S/A) and length.

The CG is the ratio of arithmetic and geometric means of channel variances and is a good indication of the performance in signal processing. To estimate the CG, the variance is computed using a first order Markov model AR(1) with intersample autocorrelation coefficient $\rho = 0.95$ [20]. As shown in Table 3, the supercompact multiwavelets have smaller CGs than SA4 and Geronimo-Hardin-Massopust (GHM), but better CGs, by at least a quarter dB, than even CL and most other orthogonal and biorthogonal multifilters, including designs with three or more matrix coefficients.




Figure 5: Test images and their histograms; (a)House(256×256); (b)Couple(256×256); (c)Brick Wall(1024×1024); (d) Texture 5 (256×256); (e) Texture 1(512×512); (f) Subwater(256×128); (g)Texture 4(1024×1024).

4.4. Applications

4.4.1. Subband-based edge detection

Edge detection is one of the fundamental operations in image processing and computer vision. It is used to identify the edges in an image, for image segmentation and data extraction, and for easy processing as well.

Since the choice of edge detection operator has to be responsive to gradual changes, the Canny method is used, which provides good edge detection. It uses two different threshold types to detect strong and weak edges [21].

The location and orientation of edges in an image can be determined from the encoded subbands of the original image [22]. Note that in many edge detection algorithms, the detail subbands which can be used in multiwavelet subband-based edge detection are using vector quantization (VQ) via k -means-based deep network architecture [23] to encode the multiwavelet coefficients. The main advantage of VQ is the construction of the code book consisting of code vectors. The goal of quantization is to obtain sufficiently small multiwavelet coefficients with a quantized distribution so that they can be encoded at a low bit rate. This motivated us to consider a simple example of a multiwavelet subband-based edge detection.

The Canny method is applied to the present multiwavelet filters for one decomposition level of a simple image, where we obtain sixteen subbands. The results from the subband-based edge detection are shown in Fig. 8. The contour lines of the SA1 filter show that this is an excellent detector, better than the other six transformations.




Figure 6: Pixel correlations and linear fit lines for the gray and color (RGB channels) images; (a) Bart Run (128×128) (b) Cameraman (256×256) (c) Kiel (512×512) (d) House (256×256); (e) Cara (256×256); (f) Voit (256×256).

4.4.2. Image Compression

The measurement for comparative analysis between the original $\mathbf{I}(x, y)$ and the noise $\tilde{\mathbf{I}}(x, y)$ images of size N^2 is the mean square error (MSE),

$$MSE = \frac{1}{N^2} \sum_{(x,y)} (\mathbf{I} - \tilde{\mathbf{I}})^2,$$

which in decibels on a logarithmic scale is the Peak Signal-to-Noise Ratio (PSNR)

$$PSNR = 10 \log_{10} \left(\frac{255^2}{MSE} \right)$$

for $x, y = 1 : N$. PSNR of the compression ratio (CR) is defined as the ratio between the original and compressed image size. Results are shown in Table 5.

The structural information of natural images is based on the human visual system (HVS). Quality assessment of compressed images is obtained through the structural similarity index measures (SSIM) [24] and its multiscale

Table 2

The coefficients of the linear regression lines for the gray and color images

Images	The coefficients of the linear regression equation $y = p_1x + p_2$		Coefficient of determination R^2	
	p_1	p_2		
Bart Run	0.9912	1.3412	0.9824	
Cameraman	0.9887	1.3424	0.9774	
Kiel	0.8271	23.5374	0.6841	
House	0.9586	5.7160	0.9189	
Couple	0.8927	12.9530	0.7970	
Brick wall	0.9053	9.5085	0.8195	
Texture 5	0.9727	2.9083	0.9461	
House	R	0.9326	9.8814	0.8697
	G	0.9408	7.8723	0.8852
	B	0.9668	4.7158	0.9347
Cara	R	0.9763	2.2668	0.9531
	G	0.9797	2.1628	0.9598
	B	0.9861	1.8061	0.9723
Voit	R	0.7972	24.2278	0.6356
	G	0.8058	21.2176	0.6493
	B	0.7870	22.8056	0.6194
Texture 1	R	0.5361	35.3482	0.2874
	G	0.4606	34.0091	0.2121
	B	0.6607	17.9257	0.4365
Subwater	R	0.9761	2.5480	0.9527
	G	0.9762	3.4688	0.9530
	B	0.9689	3.6210	0.9387
Texture 4	R	0.9927	0.7649	0.9854
	G	0.9947	0.5201	0.9894
	B	0.9923	0.7514	0.9846

Table 3Comparison of CGs, S, S/A, for autocorrelation matrix R_x of AR(1) and $\rho = 0.95$

Multifilter	CG	S	S/A	length
Biorthogonal Hermitian cubic spline (bih34n) [14]	1.51	2.5	yes	3/5
Biorthogonal (dual to Hermitian cubic spline) (bih32s) [15]	1.01	2.5	yes	3/5
Integer Haar [16] (<i>supercompact</i>)	1.83	0.5	yes	2
CL [7]	2.06	1.06	yes	3
Alpert [17] (<i>supercompact</i>)	2.13	0.5	yes	2
New2 (<i>supercompact</i>)	2.24	1.28	yes	2
New1 (<i>supercompact</i>)	2.38	1.28	yes	2
SA4 [18]	3.73	0.99	yes	4
GHM [19]	4.41	1.5	yes	4

extension, the multiscale structural similarity index (MS-SSIM) [25], where the objective quality evaluation at each image scale contributes to the overall score.

The SSIM metric combines local image structure, luminance, and contrast into a single local quality score. In this metric, structures are patterns of pixel intensities, especially among neighboring pixels, after normalizing for luminance and contrast. Because the human visual system is good at perceiving structure, the SSIM quality metric agrees more closely with the subjective quality score. The SSIM perceived quality by quantifying the SSIM between an image and

a reference image [24]:

$$SSIM(\mathbf{I}, \tilde{\mathbf{I}}) = m_K(\mathbf{I}, \tilde{\mathbf{I}}) v_K(\mathbf{I}, \tilde{\mathbf{I}}) = \frac{(2\mu_x\mu_y + C_1)(2\sigma_{xy} + C_2)}{(\mu_x^2 + \mu_y^2 + C_1)(\sigma_x^2 + \sigma_y^2 + C_2)},$$

where $m_K(\mathbf{I}, \tilde{\mathbf{I}})$ is the luminance distortion, $v_K(\mathbf{I}, \tilde{\mathbf{I}})$ are the contrast components, and μ_x , μ_y , σ_x , σ_y and σ_{xy} are the local means, standard deviations, and cross-covariance for images $\mathbf{I}(x, y)$ and $\tilde{\mathbf{I}}(x, y)$ with $C_1 = 2.55$, $C_2 = 7.65$.

MS-SSIM [25] expands on the SSIM index by combining luminance information with the variance and cross-correlation components at scale K . The multiple scales account for variability in the perception of image details caused by factors such as viewing distance from the image, distance from the scene to the sensor, and resolution of the image acquisition sensor.

The MS-SSIM index for scale K is given by [25]

$$MS-SSIM = m_K(\mathbf{I}, \tilde{\mathbf{I}})^{\alpha_K} \prod_{k=1}^K v_k(\mathbf{I}, \tilde{\mathbf{I}})^{\beta_k} r_k(\mathbf{I}, \tilde{\mathbf{I}})^{\gamma_k},$$

where

$$r_K(\mathbf{I}, \tilde{\mathbf{I}}) = \frac{\sigma_{xy} + C_3}{\sigma_x\sigma_y + C_3},$$

and $C_3 = C_2/2$ is the structure component at scale K (where $K = 1$ is the full-resolution image). For $K = 5$, the exponents are $\beta_1 = \gamma_1 = 0.0448$, $\beta_2 = \gamma_2 = 0.2856$, $\beta_3 = \gamma_3 = 0.3001$, $\beta_4 = \gamma_4 = 0.2363$, and $\alpha_5 = \beta_5 = \gamma_5 = 0.1333$. The best quality values for structural information are obtained for $SSIM = MS-SSIM = 1$.

Image compression of the gray level images Bart Run, Cameraman and Kie1, with sizes 128^2 , 256^2 , and 512^2 pixels respectively, is applied using balanced GHM, CL, SA4, Alpert, and the two new supercompact multifilters. The balancing for GHM and CL is according to [17]; for SA4, Alpert, and the new supercompact multifilters it is of the Haar type.

Results are shown in Tables 4–9. All minimal MSE and maximal PSNR, SSIM and MS-SSIM values are marked in bold.

Table 4 shows results for the gray level image Bart Run. When $CR = 4$, all multiwavelet filter banks from decomposition levels 2 through 5 achieve nearly constant SSIM and MS-SSIM. The lowest MSE and highest values for PSNR, SSIM and MS-SSIM are obtained by the Integer Haar multiwavelet. The supercompact multiwavelets compared with other multiwavelets (GHM, SA4 and CL) leads to better PSNR, SSIM and MS-SSIM.

When $CR = 8$, nearly constant SSIM and MS-SSIM are achieved for all multiwavelet filter banks from the third to the fifth level decomposition. The lowest MSE and highest values for PSNR are achieved by using the Alpert multiwavelet. We need to note that better MSE and PSNR do not lead to better SSIM and MS-SSIM. The highest SSIM is by using CL multiwavelet while and highest MS-SSIM values are achieved for 3 and more level decompositions from the New1 supercompact multiwavelet.

The next two $CR = 16$ and 32 show that at the first compression level all the measured values for the Alpert, New1 and New2 multiwavelets are equal. This comes from the similarity in structure of the pairs of scaling and wavelet functions. Moreover, Alpert and New1 multiwavelet filters achieve best PSNRs, SSIMs and MS-SSIMs. For $CR = 16$, best MSEs and PSNRs values are for the New1 multiwavelet at 3 and more decomposition level, while for $CR = 32$ it is the Alpert multiwavelet. With respect to SSIMs and MS-SSIMs for $CR = 16$, the Alpert multiwavelet achieved the best values for 3 and more decomposition levels, while for $CR = 32$ the best values are from 1 to 3 decomposition levels with the New1 multiwavelet filter.

Table 5 shows results for the gray level image Cameraman. For $CR = 4$ GHM has better MS-SSIM, but for $CR = 8$, 16 and 32 the New1 supercompact multifilter filter for 3 and more decomposition levels achieved better values than all the rest. It is important to note that all supercompact multiwavelets have nearly constant values for PSNR, SSIM and MS-SSIM for 4 and more decomposition levels.

Table 6 shows results for the gray level image Kie1. For 4 and more decomposition levels and $CR = 4$, 8 and 16 all multiwavelets achieved nearly constant MSEs and PSNRs. Better results are obtained for the two supercompact Alpert and New1 multiwavelet filters. For the three considered CRs, the New1 multiwavelet has better values for MS-SSIM, MSE and PSNR.

Table 7 shows results for the color image House. The PSNR values of all multiwavelet filters for $CR = 4$ and 8 for 4 and more decomposition levels and for $CR = 16$ and 32 for 5 and 6 decomposition levels are nearly constant. For all CR s, the best PSNR, SSIM and MS-SSIM with 3 and more decomposition levels were achieved again by the New1 multiwavelet filter. Moreover, with $CR = 4$, its MS-SSIM values equal those of the SA4 multiwavelet filter, which is longer with more vanishing moments.

Table 8 shows results for the color image Cara. The best measured values for $CR = 4$ of all decomposition levels are achieved by the New1 multiwavelet. The SSIM and MS-SSIM values of GHM, SA4 and Alpert multiwavelets are equal for 2 and more levels. We need to note that for $CR = 4$ and 8 at 3 and more decomposition levels all multifilters have nearly constant SSIM and MS-SSIM values. For $CR = 8$, best measured values are achieved again by the New1 multiwavelet, except for 1 and 2 decomposition levels, and the MS-SSIM values are equal with GHM and Alpert for 3 and more levels. For $CR = 16$, for 1 and 2 decomposition levels the SA4 multifilter has the best SSIM values, for the rest the best measured values are achieved by the New1 multiwavelet. For $CR = 32$, the New1 multiwavelet achieves all best measured values, except MSE and PSNR at 1, 2 and 3 decomposition levels. The MS-SSIM values of the supercompact multiwavelet filters are nearly constant for more than 4 decomposition levels.

Table 9 shows results for the color image Voit. For $CR = 4$ and 8, all multiwavelet filters at 4 and more decomposition levels have nearly constant measured values. With a minimal difference from the SA4 multiwavelet, all best measured values are achieved again by the New1 multiwavelet filter. For $CR = 4$ and 8, the New1 multiwavelet filter has the best MSE, PSNR and SSIM for 3 and more decomposition levels. The best MS-SSIM, is achieved by the SA4 multiwavelet, except at the 2nd level. For $CR = 32$, from 1 to 3 decomposition level the best SSIM is with the SA4 multiwavelet filter, while the best MS-SSIM values are for CL, New1 and SA4 multiwavelets, respectively. For 4 and more decomposition level, the New1 multiwavelet has the best SSIM and MS-SSIM values, while its PSNR values are equal with Alpert multiwavelet.

Table 10 shows results for the color image Texture 1. For $CR = 2$ for decomposition levels bigger than 1, the New1 multifilter has max PSNR, SSIM and MS-SSIM values. For $CR = 4$, the SA4 multifilter has max PSNR and SSIM values for all decomposition levels, which are slightly higher (0.03dB and 0.020 respectively) than New1, while New1 has max MS-SSIM slightly higher (0.20dB, 0.020dB and 0.03dB, respectively) than New2. For $CR = 8$, SA4 has max PSNR and SSIM values for all decomposition level, which are slightly higher (0.03dB, 0.04dB and 0.012dB, respectively) than New1.

Table 11 shows results for the color image Texture 4. Since the image allows superb values at higher compression levels, the beginning value is $CR = 64$.

For $CR=64$, the max PSNR value at the first decomposition level is achieved by the SA4 multifilter, slightly higher (by 0.04dB) than New1. At the second decomposition level, New1 is better. For decomposition levels 3 to 7, Integer Haar has the max PSNR levels, significantly higher (by 0.64dB – 1.36dB) than New1.

For SSIM and MS-SSIM, the maximal values at the first 3 decomposition levels are for New1 and New2, and at levels 4 to 7 the Integer Haar multifilter has negligibly higher levels (by 0.003dB).

For $CR = 128$, max PSNR at the first decomposition level is for SA4, higher by 0.02dB than New1. The New1 multifilter has superior values for PSNR, SSIM and MS-SSIM at levels 2 to 7 (6.5dB better than CL and SA4, 12dB better than GHM and 6.5dB better than New2).

For $CR = 256$, SA4 has the maximal PSNR values at the first and second decomposition levels, better by 0.03dB and 0.05dB respectively) than New1. New1 has max SSIM and MS-SSIM values. From levels 3 to 7 all maximal values are for New1, bigger by around 5.0dB than SA4 and CL, and around 8.0dB better than GHM.

For $CR = 512$, SA4 achieves the maximal PSNR values at the first and second decomposition levels, negligibly higher (by 0.03dB and 0.04dB, respectively) than New1. At the second level, New1 have max SSIM and MS-SSIM values. From levels 3 to 7 New1 has all maximal values, bigger by around 5.0dB than SA4 and CL, and around 6.0dB better than GHM.

The multifilter New1 applied to the image Texture 4, which has a strongly combed histogram, achieves better results for all metrics, with minimal differences for all compression levels.

Table 12 shows results for the grayscale image Brick wall. For $CR=16$, at the first decomposition level SA4 has max PSNR values, while at levels 3 to 7 CL is best, negligibly higher (by 0.01dB) than New1. The max SSIM is for the GHM multifilter, while the SA4 multifilter has max MS-SSIM, negligibly higher (by 0.004dB) than New1. For $CR=32$ and 64, SA4 has the max PSNR, SSIM and MS-SSIM values at decomposition levels 1 and 2, while for levels 3 to 7 the SA4 multifilter New1 has all max values.

Table 13 shows results for the grayscale image Texture 5. For CR = 4, 8 and 16, the New1 multifilter has max PSNR, SSIM and MS-SSIM values at all decomposition levels. For CR=32, the SA4 multifilter has maximal values for the first and second decomposition levels, while New1 has max values for the third to seventh decomposition levels.

Table 14 shows results for the color image Subwater. For CR= 8 all maximal values are obtained for SA4, slightly higher (by 0.17dB and 0.004dB, respectively) than New1; the values of MS-SSIM are equal. For CR=16, max PSNR, SSIM and MS-SSIM values are obtained by New1, except at the second level, where the values are negligible lower (by 0.06dB, 0.0007dB and 0.0001dB, respectively) than SA4. Note that SA4 and New1 multifilters have constant MS-SSIM from the third to the seventh decomposition level. For CR=32, all maximal values are obtained for New1, significantly higher than the others (by around 1.00dB for CL, around 0.70dB) for SA4 and even by 3.00dB for GHM).

4.4.3. Signal denoising

Multiwavelet denoising is a non-linear process, which depends heavily on the choice of a thresholding parameter. For a better comparison of the new supercompact and orthogonal multifilters New1 and New2 with orthogonal GHM, SA4, CL, Alpert and Integer Haar multifilters, noise reduction in 1D and 2D is considered.

1D denoising We apply denoising with seven orthogonal multifilters (GHM, SA4, CL, Alpert, Integer Haar, New1 and New2) to three 1D test signals $s(t)$ (Piece-Polynomial and Cusp with a length of 1024 values and HeaviSine with 4096), with two different additive white Gaussian noise (AWGN) values ($\sigma=10, 15$). The original, noisy signals are shown in Fig. 9(a)-(f). The denoised signals, with a zoom of some parts are shown in Fig. 10(a)-(f). The Piece-Polynomial noisy signal is more complex; it consists of exponential and others polynomial segments with edges and jumps.

The zoom images of the simulation plots give us enough information to recognize the quality of approximation for all multifilters. For all three test signals and both noise values, Fig. 10(c) and (d) show that the New2 multifilter has the best plots for cruder numerical approximations. The scaling part of the HeaviSine signal shows better approximations of sudden changes by both the New2 and New1 multifilters. The zoom plots of Piece-Polynomial and Cusp signals for $\sigma=10, 15$ show better approximation using the New1 and GHM multifilters, as shown in Fig. 10(a),(b),(c) and (d). Using the Alpert multifilter for Piece-Polynomial and Cusp signals leads to spurious peak values.

To complete the analysis, we use the maximal absolute error (MAE):

$$\epsilon = ||s(t) - \hat{s}(t)||_{\infty}. \quad (13)$$

The MAE values are tabulated in Table 15.

Since the Piece-Polynomial noisy signal is more complex, its MAEs are bigger than for other signals. We see that the minimal MAE values for $\sigma=10$ are obtained by both the New1 and New2 multifilters, for $\sigma=15$ only by New1. The noise suppression results obtained for the Cusp signal shows again minimal MAE values for both New1 and New2 multifilters for $\sigma=10, 15$. The results for the HeaviSine signal show minimal MAE value using for $\sigma=10, 15$. A comparison of the results shows that the proposed New1 and New2 multifilters can provide more effective noise suppression and better approximation than the other five multifilters.

2D denoising Image denoising is the process of removing noise from a noisy image while preserving the structure of the original image. The best way to test the effect of noise on a standard grayscale image is to add AWGN. We apply thresholding methods to suppress noise from grayscale images of size 256×256 and 512×512 pixels shown in Fig.4 and Fig.5 and the multiwavelet decomposition for seven orthogonal multiwavelets GHM, SA4, CL, Alpert, Integer Haar, New1 and New2. The AWGN has variances $\sigma=10, 20$. In order to suppress the noise coefficients in all test images, soft or hard vector thresholding is applied at five decomposition levels. Best results are shown in boldface.

Since a single one criterion is not enough to compare denoising quality, four well-defined criteria are introduced: MSE, PSNR, SSIM, and MS-SSIM.

Table 16 gives denoising results for the soft and hard thresholds of a noisy Cameraman image with $\sigma=10$. The SA4 multifilter has max PSNR, SSIM and MS-SSIM values for all measures only for a soft threshold at the first decomposition level. The New1 multifilter achieves highest PSNR, SSIM and MS-SSIM values from the second to the fifth decomposition level for both thresholding methods, while the New2 multifilter achieves only for the hard thresholding method.

Table 17 gives denoising results with the soft and hard thresholds and $\sigma=10$ for the Couple image. The results shows that for all decomposition levels the highest PSNR and SSIM values are achieved by the SA4 multifilter with

soft thresholding, while New1 achieves the highest MS-SSIM values. With hard thresholding, the New1 multifilter achieves the highest PSNR, SSIM and MS-SSIM values for all decomposition levels.

Table 18 gives denoising results with soft and hard thresholding and $\sigma=10$ for the Kiel image. The results show that, except for the first level of decomposition, the multifilter New1 has highest PSNR, SSIM and MS-SSIM values for both thresholding methods.

Table 19 gives denoising results with soft and hard thresholding and $\sigma=20$ for the House image. The New1 multifilter achieves the highest PSNR, SSIM and MS-SSIM values for soft thresholding from the second to fifth decomposition levels, as well as for hard thresholding from the third to the fifth level.

Table 20 gives denoising results with soft and hard thresholding and $\sigma=20$ for the Kiel image. The results show that for soft thresholding the highest PSNR, SSIM, and MS-SSIM values are achieved by New1 from the second to the fifth decomposition levels, and for hard thresholding from the third to the fifth level.

From the results, we can see that in the case of small noise variance $\sigma=10$ the New1 multifilter has the best denoising. Finally, MS-SSIM, which generalizes SSIM over all decomposition levels, gives us information that the New1 multiwavelet filter has better HVPs for image quality and structure.

5. Conclusions

Two new orthogonal bases for supercompact multiscaling functions with exact recursion coefficients are obtained from the CL matrix product filter by the fast Bauer's method of matrix spectral factorization. The complementary supercompact multiwavelet functions are also found by QR decomposition. The two new multifilters, even with very short support, show relatively good CG performance, and better Sobolev smoothness than other well-known orthogonal and biorthogonal multiwavelets.

Considering the highest MSE, PSNR, SSIM, and MS-SSIM (in bold) for CR at all decomposition levels for image compression and denoising, the new supercompact multiwavelet New1 (in same case also New2) achieves better performance than the other multiwavelet filters. The measure MS-SSIM tells us that by applying New1 multifilters, better HVPs for image quality and structure can be achieved in some cases.

From comparison with the well-known orthogonal Alpert, GHM, SA4, CL and Integer Haar multiwavelets, we can conclude that the multifilters New1 and New2 can provide more effective noise suppression of some one-dimensional signals, can be also useful in image compression and subband-based edge detection applications as well.

In the future, these new supercompact bases could be used for solving other different mathematical and engineering applications.

6. Conflict of interest

The author declares that there is no conflict of interest.

7. Data availability statement

Data available on request from the authors.

8. Acknowledgments

The authors would like to thank the editor, two anonymous reviewers for their careful reading, their constructive and valuable comments and very helpful suggestions which helped us to improve the manuscript.

References

- [1] R. M. Beam, R. F. Warming, Multiresolution analysis and supercompact multiwavelets, *SIAM Journal on Scientific Computing* 22 (4) (2000) 1238–1268.
- [2] D. Lee, R. M. Beam, R. F. Warming, Supercompact multiwavelets for flow field simulation, *Computers & Fluids* 30 (7) (2001) 783–805.
- [3] G. Strang, T. Nguyen, *Wavelets and Filter Banks*, Wellesley-Cambridge Press, Wellesley, MA, 1996.
- [4] T. Cooklev, A. Nishihara, M. Kato, M. Sablatash, Two-channel multifilter banks and multiwavelets, in: 1996 IEEE International Conference on Acoustics, Speech, and Signal Processing Conference Proceedings, Vol. 5, 1996, pp. 2769–2772.
- [5] V. Kolev, T. Cooklev, F. Keinert, Matrix spectral factorization for SA4 multiwavelet, *Multidimensional Systems and Signal Processing* 29 (4) (2018) 1613–1641.

- [6] V. Kolev, T. Cooklev, F. Keinert, Bauer's spectral factorization method for low order multiwavelet filter design, *Journal of Computational and Applied Mathematics* 441, article 115713 (2024).
- [7] C. K. Chui, J.-A. Lian, A study of orthonormal multi-wavelets, *Appl. Numer. Math.* 20 (3) (1996) 273–298.
- [8] L. Fejér, Über trigonometrische Polynome, *J. Reine Angew. Math. (Crelles J.)* 146 (1916) 53–82.
- [9] F. Riesz, Über ein Problem des Herrn Carathéodory, *J. Reine Angew. Math. (Crelles J.)* 146 (1916) 83–87.
- [10] R. E. A. C. Paley, N. Wiener, *Fourier Transforms in the Complex Domain*, Colloquium Publications AMS, New York, 1933.
- [11] N. Wiener, On the factorization of matrices, *Commentarii Mathematici Helvetici* 29 (1) (1955) 97–111.
- [12] A. H. Sayed, T. Kailath, A survey of spectral factorization methods, *Numerical Linear Algebra With Applications* 8 (6–7) (2001) 467–496.
- [13] Q. Jiang, On the regularity of matrix refinable functions, *SIAM J. Math. Anal.* 29 (5) (1998) 1157–1176.
- [14] V. Strela, A note on construction of biorthogonal multi-scaling functions, in: *Wavelets, Multiwavelets, and their Applications* (San Diego, CA, 1997), Vol. 216 of *Contemp. Math.*, Amer. Math. Soc., Providence, RI, 1998, pp. 149–157.
- [15] A. Z. Averbuch, V. A. Zheludev, T. Cohen, Multiwavelet frames in signal space originated from Hermite splines, *IEEE Trans. Signal Process.* 55 (3) (2007) 797–808.
- [16] K. Rajakumar, T. Arivoli, Lossy image compression using multiwavelet transform for wireless transmission, *Wireless Personal Communications* 87 (2) (2016) 315–333.
- [17] V. Strela, A. Walden, Signal and image denoising via wavelet thresholding: Orthogonal and biorthogonal, scalar and multiple wavelet transforms, in: W. J. Fitzgerald, R. L. Smith, A. T. Walden, P. C. Young (Eds.), *Nonlinear and Nonstationary Signal Processing*, Cambridge University Press, 2001, pp. 395–441.
- [18] L. Shen, H. H. Tan, J. Y. Tham, Symmetric-antisymmetric orthonormal multiwavelets and related scalar wavelets, *Appl. Comput. Harmon. Anal.* 8 (3) (2000) 258–279.
- [19] J. S. Geronimo, D. P. Hardin, P. R. Massopust, Fractal functions and wavelet expansions based on several scaling functions, *J. Approx. Theory* 78 (3) (1994) 373–401.
- [20] P. P. Vaidyanathan, Theory of optimal orthonormal subband coders, *IEEE Transactions on Signal Processing* 46 (6) (1998) 1528–1543.
- [21] J. Canny, A computational approach to edge detection, *IEEE Transactions on pattern analysis and machine intelligence* (6) (2009) 679–698.
- [22] N. Mohsenian, N. M. Nasrabadi, Edge-based subband vq techniques for images and video, *IEEE transactions on circuits and systems for video technology* 4 (1) (2002) 53–67.
- [23] S.-K. Im, K.-H. Chan, Vector quantization using k-means clustering neural network, *Electronics Letters* 59 (7) (2023) e12758.
- [24] R. D. M., H. S. S., Understanding and simplifying the structural similarity metric, in: *15th IEEE International Conference on Image Processing*, 2008, pp. 1188–1191.
- [25] Z. Wang, E. P. Simoncelli, A. C. Bovik, Multiscale structural similarity for image quality assessment, in: *The Thirty-Seventh Asilomar Conference on Signals, Systems & Computers*, Vol. 2, 2003, pp. 1398–1402.
- [26] Huang X., Chan K.H., Wu W., Sheng H. and Ke W., Fusion of multi-modal features to enhance dense video caption, *Sensors*, vol. 23, no.12, p.5565, 2023.




Figure 7: Pixel correlations and linear fit lines for the gray and color (RGB channels) images; (a)House (256 × 256); (b)Couple(256×256); (c)Brick Wall(1024×1024); (d)Texture 5(256×256); (e)Texture 1(512×512); (f)Subwater(256×128); (g)Texture 4(1024 × 1024).




Figure 8: Subband-based edge detection for a one decomposition level of following orthogonal transforms by using the Canny method: (a) Original image; (b) GHM multiwavelet; (c) SA4 multiwavelet; (d) CL multiwavelet; (e) Alpert multiwavelet; (f) New1 multiwavelet; (g) New2 multiwavelet; (h) Integer Haar multiwavelet.

Table 4

Results for Bart Run. The four lines for each multiwavelet represent MSE, PSNR, SSIM and MS-SSIM, in this order.

Image	CR	Multi-wavelet	Decomposition level				
			1	2	3	4	5
 Bart Run	4:1	GHM	86.167	3.0603	2.4312	2.3896	2.3896
			28.78	43.27	44.27	44.35	44.35
			0.9031	0.9830	0.9857	0.9859	0.9859
			0.9509	0.9984	0.9986	0.9986	0.9986
		SA4	1.0416	1.0153	1.0305	1.0559	1.0548
			47.95	48.07	48.00	47.89	47.9
			0.9935	0.9932	0.993	0.9929	0.9929
			0.9995	0.9994	0.9994	0.9994	0.9994
		CL	0.5866	0.5192	0.4995	0.5065	0.5111
			50.45	50.98	51.15	51.08	51.05
			0.9967	0.9969	0.9969	0.9969	0.9969
			0.9997	0.9997	0.9997	0.9997	0.9997
		Alpert	0.6630	0.5090	0.4955	0.4993	0.5040
			49.92	51.06	51.18	51.15	51.11
			0.9966	0.9970	0.9971	0.9971	0.997
			0.9997	0.9997	0.9997	0.9997	0.9997
		New1	0.6096	0.4987	0.4923	0.4983	0.5021
			50.28	51.15	51.21	51.16	51.12
			0.9967	0.9972	0.9972	0.9971	0.9971
			0.9997	0.9997	0.9997	0.9997	0.9997
		New2	1.1408	0.9912	0.9570	0.9782	0.9921
			47.56	48.17	48.32	48.23	48.17
			0.9948	0.9949	0.9951	0.9950	0.9949
			0.9995	0.9994	0.9994	0.9994	0.9994
		Integer Haar	0.6234	0.4246	0.3891	0.3919	0.3984
			50.18	51.85	52.23	52.20	52.13
			0.9972	0.9977	0.9978	0.9978	0.9978
			0.9997	0.9998	0.9998	0.9998	0.9998
	8:1	GHM	5385.1	15.463	11.505	11.127	11.071
			10.82	36.24	37.52	37.67	37.69
			0.2538	0.9241	0.9404	0.9425	0.9428
			0.6421	0.9912	0.9934	0.9936	0.9936
		SA4	9.9828	6.9655	6.815	6.9071	6.9234
			38.14	39.70	39.80	39.74	39.73
			0.9507	0.9633	0.9634	0.9634	0.9632
			0.9953	0.9957	0.9955	0.9954	0.9954
		CL	9.8477	6.6463	5.931	5.973	6.0143
			38.20	39.91	40.40	40.37	40.34
			0.9531	0.9668	0.9699	0.9696	0.9695
			0.9936	0.9956	0.9957	0.9956	0.9956
		Alpert	10.4090	6.3780	5.7271	5.7040	5.7534
			37.96	40.08	40.55	40.57	40.53
			0.9458	0.9658	0.9683	0.9682	0.9680
			0.9955	0.9962	0.9961	0.9960	0.9960
		New1	9.9606	6.4132	5.9374	5.9642	6.0197
			38.15	40.06	40.39	40.38	40.34
			0.9494	0.9646	0.9684	0.9682	0.9681
			0.9958	0.9960	0.9962	0.9962	0.9960
		New2	14.023	11.193	10.575	10.776	10.767
			36.66	37.64	37.89	37.81	37.81
			0.9244	0.9381	0.9460	0.944	0.9444
			0.9918	0.9919	0.9932	0.9931	0.9931
		Integer Haar	12.76	8.2607	7.4608	7.5038	7.6042
			37.07	38.96	39.40	39.38	39.32
			0.9411	0.9593	0.9628	0.9625	0.9620
			0.9932	0.9950	0.9953	0.9952	0.9952

Table 4(continued)

Results for Bart Run. The four lines for each multiwavelet represent MSE, PSNR, SSIM and MS-SSIM, in this order.

Image	CR	Multi-wavelet	Decomposition level				
			1	2	3	4	5
 Bart Run	16:1	GHM	11837	189.7	27.636	25.712	25.596
			7.399	25.35	33.72	34.03	34.05
			0.0142	0.7820	0.8719	0.8798	0.8802
			0.4809	0.8622	0.9792	0.9806	0.9806
		SA4	189.23	19.975	18.572	18.818	18.889
			25.36	35.13	35.44	35.39	35.37
			0.766	0.9033	0.9099	0.9089	0.9086
			0.9008	0.9850	0.9869	0.9863	0.9862
		CL	240.87	22.234	18.827	18.763	18.888
			24.31	34.66	35.38	35.40	35.37
			0.7795	0.8946	0.9076	0.9076	0.9072
			0.8490	0.9829	0.9852	0.9848	0.9846
		Alpert	64.561	20.539	17.289	17.243	17.264
			30.03	35.00	35.75	35.76	35.76
			0.7730	0.8943	0.9111	0.9113	0.9113
			0.9622	0.9852	0.9881	0.9879	0.9878
		New1	64.561	20.152	17.055	17.054	17.251
			30.03	35.09	35.81	35.81	35.76
			0.7730	0.8969	0.9107	0.9105	0.9096
			0.9622	0.9856	0.9878	0.9874	0.9872
		New2	64.561	29.797	27.838	28.114	28.25
			30.03	33.39	33.68	33.64	33.62
			0.7730	0.8478	0.8597	0.8586	0.8581
			0.9622	0.9743	0.9777	0.9770	0.9768
		Integer Haar	317.99	26.172	23.221	23.302	23.494
			23.11	33.95	34.47	34.46	34.42
			0.7559	0.8760	0.8878	0.8872	0.8862
			0.8150	0.9791	0.9816	0.9811	0.9809
	32:1	GHM	17368	4704.3	73.563	51.504	50.794
			5.733	11.41	29.46	31.01	31.07
			0.00814	0.2671	0.7643	0.7942	0.7976
			0.4401	0.5499	0.9338	0.9531	0.9542
		SA4	5415.3	47.809	37.967	37.859	38.058
			10.79	31.34	32.34	32.35	32.33
			0.3788	0.8026	0.8412	0.8424	0.842
			0.4941	0.9528	0.9681	0.9664	0.9663
		CL	5269.8	56.575	38.712	37.714	37.993
			10.91	30.60	32.25	32.37	32.33
			0.378	0.7439	0.8311	0.8342	0.8338
			0.5236	0.9347	0.9650	0.9654	0.9652
		Alpert	5018	44.849	32.806	32.498	32.648
			11.13	31.61	32.97	33.01	32.99
			0.3735	0.7993	0.8463	0.8463	0.8462
			0.6502	0.9596	0.9711	0.9722	0.9718
		New1	5018	44.08	32.803	32.601	32.927
			11.13	31.69	32.97	33.00	32.96
			0.3735	0.8029	0.8478	0.8469	0.8462
			0.6502	0.9609	0.9728	0.9717	0.9713
		New2	5018	57.571	51.015	51.224	51.497
			11.13	30.53	31.05	31.04	31.01
			0.3735	0.7503	0.7703	0.7692	0.7693
			0.6502	0.9389	0.9510	0.9500	0.9495
		Integer Haar	5524.4	58.614	48.135	47.572	47.93
			10.71	30.45	31.31	31.36	31.32
			0.3684	0.7495	0.801	0.8023	0.8003
			0.4943	0.9393	0.9586	0.9585	0.9579

Table 5

Results for Cameraman. The four lines for each multiwavelet represent MSE, PSNR, SSIM and MS-SSIM, in this order.

Image	CR	Multi-wavelet	Decomposition level					
			1	2	3	4	5	6
 cameraman	4:1	GHM	65.773	9.814	7.9468	7.7663	7.7578	7.7579
			29.95	38.21	39.13	39.23	39.23	39.23
			0.846	0.9596	0.963	0.9633	0.9633	0.9634
			0.9778	0.9945	0.9947	0.9947	0.9947	0.9947
		SA4	6.4772	5.6798	5.7046	5.7165	5.7208	5.7204
			40.02	40.59	40.57	40.56	40.56	40.56
			0.9646	0.9656	0.9659	0.9659	0.9659	0.9659
			0.9946	0.9945	0.9945	0.9945	0.9945	0.9945
		CL	6.3258	5.2186	5.1314	5.129	5.1226	5.1235
			40.12	40.96	41.03	41.03	41.04	41.04
			0.9637	0.9655	0.9663	0.9664	0.9665	0.9665
			0.9941	0.9939	0.9942	0.9941	0.9941	0.9941
		Alpert	6.1808	5.1158	4.9445	4.9299	4.9375	4.9356
			40.22	41.04	41.19	41.20	41.20	41.20
			0.9637	0.9662	0.9672	0.9672	0.9672	0.9672
			0.9952	0.9941	0.9946	0.9945	0.9945	0.9945
		New1	6.1103	5.0703	4.9072	4.8982	4.9033	4.9044
			40.27	41.08	41.22	41.23	41.23	41.22
			0.964	0.9664	0.9675	0.9675	0.9674	0.9674
			0.9953	0.9942	0.9947	0.9946	0.9946	0.9946
		New2	8.0861	7.0528	7.0942	7.1256	7.1481	7.1492
			39.05	39.65	39.62	39.6	39.59	39.59
			0.9584	0.9585	0.9585	0.9584	0.9583	0.9583
			0.9944	0.9920	0.9923	0.9923	0.9923	0.9923
		Integer Haar	7.045	5.9673	5.8743	5.8631	5.8739	5.874
			39.65	40.37	40.44	40.45	40.44	40.44
			0.9616	0.9628	0.9633	0.9634	0.9633	0.9633
			0.9935	0.9933	0.9935	0.9936	0.9935	0.9935
	8:1	GHM	4488	54.003	37.34	35.658	35.56	35.538
			11.61	30.81	32.41	32.61	32.62	32.62
			0.1478	0.8883	0.9086	0.9111	0.9113	0.9113
			0.6785	0.98	0.9835	0.9833	0.9833	0.9833
		SA4	39.525	27.101	26.908	26.987	27.012	27.013
			32.16	33.8	33.83	33.82	33.82	33.82
			0.9050	0.9197	0.9188	0.919	0.919	0.9189
			0.9826	0.9838	0.9820	0.9818	0.9818	0.9818
		CL	41.284	26.052	25.502	25.39	25.433	25.427
			31.97	33.97	34.06	34.08	34.08	34.08
			0.8995	0.9192	0.9203	0.9209	0.9208	0.9208
			0.9793	0.9819	0.9809	0.9813	0.9815	0.9815
		Alpert	40.38	25.849	24.791	24.635	24.693	24.698
			32.07	34.01	34.19	34.22	34.21	34.20
			0.8986	0.9193	0.9215	0.9222	0.9221	0.922
			0.9859	0.9841	0.9831	0.9839	0.9838	0.9838
		New1	39.706	25.503	24.488	24.351	24.413	24.429
			32.14	34.06	34.24	34.27	34.25	34.25
			0.9004	0.9204	0.9226	0.9231	0.9228	0.9228
			0.9863	0.9844	0.9836	0.9841	0.9841	0.9841
		New2	52.833	37.008	36.229	36.612	36.775	36.796
			30.90	32.45	32.54	32.49	32.48	32.47
			0.8819	0.8995	0.8990	0.8976	0.8971	0.8971
			0.9822	0.9793	0.9743	0.9753	0.975	0.9751
		Integer Haar	48.803	32.369	31.122	31.10	31.151	31.15
			31.25	33.03	33.20	33.20	33.20	33.20
			0.8869	0.9078	0.9080	0.9077	0.9075	0.9074
			0.9742	0.9794	0.9783	0.9787	0.9787	0.9787

Table 5(continued)

Results for Cameraman. The four lines for each multiwavelet represent MSE, PSNR, SSIM and MS-SSIM, in this order.

Image	CR	Multi-wavelet	Decomposition level					
			1	2	3	4	5	6
 cameraman	16:1	GHM	9398.3	206.16	97.873	89.405	88.794	88.729
			8.4	24.99	28.22	28.62	28.65	28.65
			0.05697	0.6939	0.8304	0.837	0.8379	0.8379
			0.5289	0.9337	0.9586	0.959	0.959	0.959
		SA4	180.76	75.635	71.015	71.511	71.684	71.656
			25.56	29.34	29.62	29.59	29.58	29.58
			0.6773	0.8468	0.8472	0.8458	0.8455	0.8456
			0.9446	0.9618	0.9562	0.9564	0.956	0.9559
		CL	203.43	75.691	70.133	70.075	70.233	70.217
			25.05	29.34	29.67	29.68	29.67	29.67
			0.6668	0.8445	0.8466	0.8486	0.8481	0.8482
			0.9383	0.9594	0.9550	0.956	0.9555	0.9556
		Alpert	202.99	75.024	67.809	67.454	67.673	67.704
			25.06	29.38	29.82	29.84	29.83	29.82
			0.665	0.8436	0.8472	0.8492	0.8488	0.8488
			0.9601	0.9641	0.9582	0.962	0.962	0.9622
		New1	199.72	73.971	66.886	66.615	66.842	66.858
			25.13	29.44	29.88	29.90	29.88	29.88
			0.6665	0.8448	0.8497	0.8516	0.8515	0.8516
			0.9605	0.9651	0.9594	0.9631	0.9628	0.9630
		New2	234.83	105.86	99.378	100.62	101.19	101.3
			24.42	27.88	28.16	28.1	28.08	28.07
			0.6541	0.8131	0.8135	0.81	0.8088	0.8087
			0.9566	0.954	0.9460	0.9444	0.9441	0.9442
		Integer Haar	229.89	97.676	87.721	87.61	87.707	87.699
			24.52	28.23	28.7	28.71	28.7	28.7
			0.655	0.8190	0.8212	0.8193	0.8191	0.8191
			0.9284	0.9495	0.9467	0.9479	0.9479	0.9478
	32:1	GHM	12589	3469.7	214.42	179.14	175.67	175.25
			7.131	12.73	24.82	25.60	25.68	25.69
			0.02968	0.1307	0.7325	0.7593	0.7615	0.7616
			0.4926	0.5191	0.9122	0.92	0.9203	0.9201
		SA4	4094.8	180.79	145.72	147.17	147.62	147.59
			12.01	25.56	26.50	26.45	26.44	26.44
			0.4219	0.7478	0.7673	0.7637	0.7636	0.7635
			0.6840	0.9209	0.9179	0.9132	0.9131	0.9133
		CL	4190.9	188.21	150.33	149.38	149.54	149.68
			11.91	25.38	26.36	26.39	26.38	26.38
			0.4191	0.7432	0.7676	0.7672	0.767	0.7670
			0.6926	0.9178	0.9184	0.9156	0.9154	0.9152
		Alpert	4250.4	186.87	144.55	143.11	143.5	143.71
			11.85	25.42	26.53	26.57	26.56	26.56
			0.4214	0.7467	0.7739	0.7753	0.7735	0.7734
			0.7224	0.931	0.9274	0.9283	0.9268	0.9269
		New1	4250.4	183.77	142.56	140.93	141.35	141.41
			11.85	25.49	26.59	26.64	26.63	26.63
			0.4214	0.7503	0.7756	0.7768	0.7755	0.7752
			0.7224	0.9319	0.9276	0.9285	0.9264	0.9263
		New2	4250.4	247.83	209	208.59	211.29	211.87
			11.85	24.19	24.93	24.94	24.88	24.87
			0.4214	0.7057	0.7383	0.7307	0.7273	0.727
			0.7224	0.9141	0.9092	0.8941	0.8962	0.8954
		Integer Haar	4295.2	238.78	188.02	185.13	185.47	185.24
			11.80	24.35	25.39	25.46	25.45	25.45
			0.4115	0.7034	0.7408	0.7336	0.7331	0.733
			0.6747	0.8994	0.9052	0.9036	0.9044	0.9046

Table 6

Results for Kiel. The four lines for each multiwavelet represent MSE, PSNR, SSIM and MS-SSIM, in this order.

Image	CR	Multi-wavelet	Decomposition level					
			1	2	3	4	5	6
 Kiel	4:1	GHM	235.84	34.756	28.619	27.844	27.761	27.758
			24.40	32.72	33.56	33.68	33.70	33.70
			0.9432	0.9844	0.9853	0.9855	0.9855	0.9855
			0.9688	0.9911	0.9917	0.9917	0.9916	0.9916
		SA4	24.975	21.198	20.825	20.849	20.854	20.859
			34.16	34.87	34.94	34.94	34.94	34.94
			0.9846	0.9835	0.983	0.983	0.983	0.983
			0.9913	0.9913	0.9909	0.9907	0.9908	0.9908
		CL	25.139	20.194	19.694	19.679	19.70	19.698
			34.13	35.08	35.19	35.19	35.19	35.19
			0.9805	0.9826	0.9824	0.9824	0.9823	0.9823
			0.9903	0.9911	0.9908	0.9907	0.9906	0.9907
		Alpert	25.241	20.372	19.713	19.634	19.631	19.619
			34.11	35.04	35.18	35.20	35.20	35.20
			0.9844	0.9842	0.9841	0.9841	0.9842	0.9842
			0.9926	0.9918	0.9914	0.9914	0.9914	0.9914
		New1	25.028	20.25	19.605	19.547	19.542	19.539
			34.15	35.07	35.21	35.22	35.22	35.22
			0.9847	0.9844	0.9842	0.9842	0.9843	0.9843
			0.9927	0.9919	0.9914	0.9915	0.9915	0.9915
		New2	29.828	26.691	26.337	26.378	26.423	26.422
			33.38	33.87	33.93	33.92	33.91	33.91
			0.9803	0.9788	0.9783	0.9781	0.978	0.978
			0.9911	0.9893	0.9886	0.9884	0.9883	0.9883
		Integer Haar	27.443	23.167	22.629	22.644	22.664	22.668
			33.75	34.48	34.58	34.58	34.58	34.58
			0.9824	0.9818	0.9817	0.9817	0.9817	0.9817
			0.99	0.9902	0.9899	0.9898	0.9899	0.9899
	8:1	GHM	5841.7	156.69	109.85	104.61	104.08	104.06
			10.47	26.18	27.72	27.93	27.96	27.96
			0.2796	0.9397	0.9482	0.9491	0.9492	0.9492
			0.6034	0.9669	0.9722	0.9724	0.9723	0.9723
		SA4	118.98	82.231	78.496	78.299	78.345	78.355
			27.38	28.98	29.18	29.19	29.19	29.19
			0.9411	0.9476	0.9475	0.9469	0.9468	0.9467
			0.9677	0.9732	0.9722	0.971	0.9709	0.9708
		CL	123.00	80.573	76.112	75.721	75.791	75.785
			27.23	29.07	29.32	29.34	29.33	29.33
			0.9271	0.9441	0.9440	0.9435	0.9434	0.9434
			0.9645	0.9719	0.9710	0.9703	0.9701	0.9702
		Alpert	123.4	81.216	76.271	75.632	75.664	75.654
			27.22	29.03	29.31	29.34	29.34	29.34
			0.9408	0.9482	0.9486	0.9484	0.9484	0.9483
			0.9748	0.9747	0.9734	0.9731	0.9732	0.9732
		New1	122.38	80.805	75.999	75.447	75.487	75.451
			27.25	29.06	29.32	29.35	29.35	29.35
			0.9420	0.9490	0.9492	0.9491	0.9489	0.9488
			0.9751	0.9751	0.9738	0.9736	0.9735	0.9734
		New2	137.67	102.07	98.629	98.492	98.688	98.733
			26.74	28.04	28.19	28.20	28.19	28.19
			0.9344	0.9322	0.931	0.9307	0.9305	0.9303
			0.9724	0.9672	0.9652	0.9645	0.9643	0.9643
		Integer Haar	132.02	90.962	86.343	85.976	86.114	86.134
			26.92	28.54	28.77	28.79	28.78	28.78
			0.9351	0.9408	0.9416	0.941	0.9408	0.9408
			0.964	0.9693	0.9687	0.968	0.9679	0.9679

Table 6(continued)

Results for Kiel. The four lines for each multiwavelet represent MSE, PSNR, SSIM and MS-SSIM, in this order.

Image	CR	Multi-wavelet	Decomposition level					
			1	2	3	4	5	6
 Kiel	16:1	GHM	11418	549	259.13	238.48	236.06	235.88
			7.555	20.74	24.00	24.36	24.40	24.40
			0.1151	0.8160	0.8871	0.8902	0.8908	0.8909
			0.4868	0.8957	0.9384	0.9393	0.9392	0.9391
		SA4	536.64	202.31	182.95	181.18	181.18	181.26
			20.83	25.07	25.51	25.55	25.55	25.55
			0.8016	0.8867	0.8895	0.8881	0.8875	0.8876
			0.8826	0.9405	0.9393	0.9365	0.9358	0.9358
		CL	548.81	201.50	180.29	178.04	177.97	177.99
			20.74	25.09	25.57	25.63	25.63	25.63
			0.7703	0.8840	0.8857	0.8855	0.8850	0.8849
			0.8838	0.9395	0.9381	0.9371	0.9365	0.9365
		Alpert	560.92	201.99	180.07	176.93	176.94	176.88
			20.64	25.08	25.58	25.65	25.65	25.65
			0.7958	0.8878	0.8930	0.8932	0.8929	0.8929
			0.9193	0.9451	0.9446	0.9434	0.9431	0.9433
		New1	558.71	201.49	180.21	177.25	177.28	177.19
			20.66	25.09	25.57	25.64	25.64	25.65
			0.7982	0.8875	0.8931	0.8934	0.8927	0.8928
			0.9201	0.9451	0.9445	0.9436	0.9428	0.9429
		New2	571.14	242.55	225.26	223.77	224.12	224.32
			20.56	24.28	24.60	24.63	24.63	24.62
			0.7904	0.8645	0.8655	0.8651	0.8636	0.8634
			0.917	0.9334	0.9298	0.9285	0.927	0.9268
		Integer Haar	578.87	223.72	201.15	198.84	199.08	199.13
			20.5	24.63	25.10	25.15	25.14	25.14
			0.7879	0.8767	0.8810	0.8794	0.8787	0.8785
			0.8696	0.9341	0.9342	0.9322	0.9315	0.9316

Table 7

Results for House. The four lines for each multiwavelet represent MSE, PSNR, SSIM and MS-SSIM, in this order.

Image	CR	Multi-wavelet	Decomposition level					
			1	2	3	4	5	6
 House	4:1	GHM	62.454	6.664	5.9222	5.8497	5.8459	5.8459
			30.43	39.95	40.46	40.51	40.51	40.51
			0.8683	0.9547	0.9593	0.9598	0.9599	0.9599
			0.9746	0.9949	0.9952	0.9952	0.9952	0.9952
		SA4	4.9972	4.3791	4.3717	4.3892	4.3903	4.3899
			41.22	41.79	41.80	41.78	41.78	41.78
			0.9643	0.9688	0.9689	0.9688	0.9688	0.9688
			0.9953	0.9957	0.9956	0.9956	0.9957	0.9957
		CL	5.0617	4.3374	4.2743	4.2803	4.2847	4.2846
			41.16	41.82	41.89	41.88	41.88	41.88
			0.9636	0.9686	0.9691	0.9691	0.9690	0.9690
			0.9946	0.9952	0.9951	0.9951	0.9951	0.9951
		Alpert	5.1499	4.3863	4.291	4.2871	4.2909	4.2912
			41.07	41.76	41.85	41.86	41.85	41.85
			0.9630	0.9682	0.9689	0.9690	0.9689	0.9689
			0.9957	0.9956	0.9956	0.9956	0.9956	0.9956
		New1	5.0737	4.3199	4.229	4.2339	4.2352	4.2352
			41.14	41.83	41.92	41.92	41.92	41.92
			0.9636	0.9687	0.9695	0.9694	0.9694	0.9694
			0.9958	0.9957	0.9957	0.9957	0.9957	0.9957
		New2	6.4048	5.848	5.8072	5.8293	5.8379	5.8389
			40.09	40.48	40.51	40.49	40.48	40.48
			0.9543	0.9580	0.9584	0.9582	0.9582	0.9581
			0.9945	0.9940	0.9939	0.9939	0.9939	0.9939
		Integer Haar	6.0433	5.3726	5.3174	5.3297	5.3319	5.3317
			40.35	40.86	40.90	40.89	40.89	40.89
			0.957	0.9615	0.9619	0.9618	0.9617	0.9617
			0.9939	0.9941	0.9942	0.9942	0.9942	0.9942
	8:1	GHM	5510.5	21.225	16.415	15.993	15.972	15.972
			10.93	34.90	36.01	36.12	36.13	36.13
			0.2712	0.8740	0.8956	0.8981	0.8982	0.8982
			0.657	0.9824	0.9853	0.9856	0.9856	0.9856
		SA4	16.127	12.194	12.057	12.15	12.167	12.169
			36.10	37.33	37.38	37.35	37.34	37.34
			0.8964	0.9200	0.9213	0.9207	0.9206	0.9206
			0.9843	0.9874	0.9872	0.9871	0.9872	0.9872
		CL	16.409	12.007	11.671	11.686	11.709	11.707
			36.02	37.39	37.51	37.51	37.50	37.50
			0.8921	0.9188	0.9213	0.9212	0.9211	0.9211
			0.9826	0.9865	0.9866	0.9865	0.9865	0.9865
		Alpert	16.549	12.091	11.612	11.615	11.63	11.634
			35.97	37.34	37.52	37.52	37.51	37.51
			0.8920	0.9188	0.9223	0.9222	0.9221	0.9220
			0.9866	0.9876	0.9878	0.9878	0.9878	0.9878
		New1	16.37	11.96	11.525	11.534	11.537	11.541
			36.02	37.39	37.56	37.55	37.55	37.55
			0.8931	0.9197	0.9229	0.9228	0.9227	0.9227
			0.9868	0.9879	0.9880	0.9879	0.9879	0.9879
		New2	20.271	16.374	16.168	16.27	16.315	16.323
			35.08	36.00	36.06	36.03	36.02	36.02
			0.8770	0.8955	0.8970	0.8965	0.8963	0.8962
			0.9843	0.9828	0.9826	0.9826	0.9825	0.9825
		Integer Haar	19.548	14.883	14.521	14.609	14.609	14.613
			35.24	36.42	36.53	36.51	36.50	36.50
			0.8790	0.9033	0.9058	0.9051	0.9051	0.9051
			0.9812	0.9837	0.9838	0.9837	0.9837	0.9837

Table 7(continued)

Results for House. The four lines for each multiwavelet represent MSE, PSNR, SSIM and MS-SSIM, in this order.

Image	CR	Multi-wavelet	Decomposition level					
			1	2	3	4	5	6
 House	16:1	GHM	11054	170.5	38.224	35.033	34.845	34.84
			7.738	26.07	32.38	32.74	32.77	32.77
			0.0391	0.7398	0.8271	0.8326	0.833	0.833
			0.5900	0.9162	0.9691	0.9697	0.9696	0.9696
		SA4	198.44	26.90	25.683	26.03	26.116	26.126
			25.37	33.88	34.08	34.02	34.01	34.01
			0.7350	0.8481	0.8541	0.8529	0.8527	0.8525
			0.9111	0.9727	0.9728	0.9721	0.9722	0.9722
		CL	190.88	26.63	24.528	24.533	24.663	24.662
			25.51	33.91	34.27	34.27	34.24	34.24
			0.7426	0.8448	0.8527	0.8529	0.8524	0.8524
			0.9088	0.9712	0.9718	0.9715	0.9715	0.9715
		Alpert	196.03	26.219	23.883	23.806	23.877	23.887
			25.38	33.97	34.37	34.39	34.37	34.37
			0.7221	0.8478	0.8573	0.8575	0.8571	0.8571
			0.9476	0.9735	0.9742	0.9741	0.9741	0.9741
		New1	197.17	25.827	23.695	23.707	23.763	23.775
			25.37	34.03	34.41	34.41	34.40	34.39
			0.7222	0.8502	0.8591	0.8590	0.8588	0.8587
			0.9477	0.9741	0.9747	0.9747	0.9748	0.9748
		New2	201.1	37.967	35.901	36.254	36.519	36.56
			25.29	32.36	32.60	32.56	32.53	32.53
			0.7208	0.8214	0.8253	0.8244	0.8236	0.8234
			0.9482	0.9656	0.9636	0.9634	0.9633	0.9633
		Integer Haar	236.19	33.76	31.283	31.52	31.558	31.56
			24.64	32.87	33.20	33.16	33.16	33.16
			0.7222	0.8282	0.8336	0.8326	0.8323	0.8322
			0.8881	0.9664	0.967	0.9666	0.9666	0.9666
	32:1	GHM	15440	4641.1	102.12	80.223	78.789	78.700
			6.261	11.60	28.26	29.27	29.34	29.35
			0.02841	0.2525	0.7605	0.7766	0.7779	0.7779
			0.5604	0.5453	0.9369	0.9433	0.9433	0.9434
		SA4	5465.5	64.412	53.237	53.817	54.285	54.317
			10.97	30.15	30.96	30.92	30.88	30.88
			0.3528	0.7860	0.7956	0.7940	0.7935	0.7935
			0.6279	0.9507	0.9518	0.9485	0.9481	0.9482
		CL	5586.6	73.51	53.266	52.708	53.512	53.516
			10.88	29.58	30.94	30.99	30.93	30.93
			0.3526	0.7792	0.7958	0.7961	0.7948	0.7949
			0.6347	0.9485	0.9510	0.9509	0.9500	0.9500
		Alpert	5710.4	66.371	48.518	47.634	47.921	48.02
			10.79	30.00	31.33	31.41	31.38	31.37
			0.3419	0.7828	0.8017	0.8025	0.802	0.8017
			0.6331	0.9543	0.9561	0.9551	0.9550	0.9550
		New1	5710.4	64.631	47.651	47.051	47.361	47.434
			10.79	30.12	31.41	31.46	31.44	31.43
			0.3419	0.7853	0.8037	0.8041	0.8033	0.8032
			0.6331	0.9551	0.9564	0.9555	0.9553	9551
		New2	5710.4	103.05	80.823	81.065	82.416	82.71
			10.79	28.12	29.14	29.13	29.06	29.04
			0.3419	0.7521	0.7643	0.7627	0.7605	0.7599
			0.6331	0.9425	0.9397	0.9372	0.9358	0.9355
		Integer Haar	5738.7	90.77	68.696	68.174	68.329	68.398
			10.76	28.65	29.83	29.86	29.85	29.85
			0.3423	0.7594	0.7731	0.7706	0.7700	0.7698
			0.6235	0.9421	0.9453	0.9431	0.9426	0.9427

Table 8

Results for Cara. The four lines for each multiwavelet represent MSE, PSNR, SSIM and MS-SSIM, in this order.

Image	CR	Multi-wavelet	Decomposition level					
			1	2	3	4	5	6
 Cara	4:1	GHM	14.669	1.5443	1.3911	1.3727	1.372	1.3719
			36.47	46.27	46.72	46.78	46.78	46.78
			0.9471	0.9829	0.9841	0.9843	0.9843	0.9843
			0.9927	0.998	0.9981	0.9981	0.9981	0.9981
		SA4	1.2479	1.1258	1.1206	1.1208	1.1209	1.121
			47.19	47.64	47.66	47.66	47.66	47.65
			0.9847	0.9860	0.9861	0.9860	0.9860	0.9860
			0.9979	0.9981	0.9981	0.9981	0.9981	0.9981
		CL	1.2791	1.1236	1.1074	1.1065	1.1052	1.1052
			47.08	47.65	47.71	47.71	47.72	47.72
			0.9843	0.9858	0.9860	0.9860	0.9860	0.9860
			0.9976	0.9978	0.9978	0.9978	0.9978	0.9978
		Alpert	1.2582	1.1111	1.0896	1.0852	1.0847	1.0846
			47.16	47.70	47.78	47.80	47.80	47.80
			0.9847	0.9861	0.9864	0.9864	0.9864	0.9864
			0.9980	0.9980	0.9980	0.9980	0.9980	0.9980
		New1	1.2397	1.0961	1.0759	1.072	1.0717	1.0716
			47.22	47.76	47.84	47.85	47.85	47.85
			0.9849	0.9863	0.9866	0.9866	0.9866	0.9866
			0.9980	0.9981	0.9981	0.9981	0.9981	0.9981
		New2	1.5073	1.3779	1.3716	1.3728	1.3728	1.3733
			46.37	46.76	46.79	46.78	46.78	46.78
			0.982	0.9832	0.9832	0.9832	0.9832	0.9832
			0.9975	0.9973	0.9973	0.9973	0.9973	0.9973
		Integer Haar	1.4994	1.332	1.3149	1.3131	1.3131	1.3133
			46.39	46.91	46.96	46.97	46.97	46.97
			0.9821	0.9836	0.9837	0.9837	0.9837	0.9837
			0.9972	0.9974	0.9974	0.9974	0.9974	0.9974
	8:1	GHM	4659.8	5.9153	4.0861	3.9051	3.8908	3.8907
			11.56	40.44	42.05	42.24	42.26	42.26
			0.05861	0.9611	0.968	0.9688	0.9689	0.9689
			0.5297	0.9945	0.9952	0.9952	0.9952	0.9952
		SA4	4.7746	3.2036	3.1461	3.1508	3.1576	3.1579
			41.37	43.10	43.18	43.17	43.17	43.16
			0.9638	0.9708	0.9713	0.9713	0.9712	0.9712
			0.9946	0.9953	0.9951	0.9951	0.9951	0.9951
		CL	5.0704	3.2022	3.0691	3.0491	3.0534	3.0538
			41.11	43.10	43.29	43.32	43.31	43.31
			0.9619	0.9702	0.9711	0.9712	0.9711	0.9711
			0.9940	0.9949	0.9948	0.9949	0.9949	0.9949
		Alpert	4.8966	3.1377	2.9816	2.9554	2.9505	2.9518
			41.26	43.20	43.42	43.46	43.46	43.46
			0.9628	0.9710	0.9718	0.9719	0.9719	0.9719
			0.9951	0.9950	0.9951	0.9952	0.9952	0.9952
		New1	4.7765	3.0599	2.9094	2.8828	2.8794	2.8819
			41.37	43.30	43.52	43.56	43.57	43.56
			0.9634	0.9715	0.9723	0.9724	0.9725	0.9724
			0.9952	0.9951	0.9952	0.9952	0.9952	0.9952
		New2	6.6138	4.7232	4.5804	4.6141	4.619	4.624
			39.96	41.42	41.56	41.53	41.52	41.52
			0.9538	0.9616	0.9623	0.9619	0.9618	0.9618
			0.9935	0.9934	0.9930	0.9930	0.9930	0.9930
		Integer Haar	6.7467	4.3785	4.1929	4.1946	4.1969	4.1973
			39.87	41.75	41.94	41.94	41.93	41.93
			0.9542	0.9637	0.9647	0.9644	0.9644	0.9644
			0.9925	0.9938	0.9938	0.9940	0.9940	0.9940

Table 8(continued)

Results for Cara. The four lines for each multiwavelet represent MSE, PSNR, SSIM and MS-SSIM, in this order.

Image	CR	Multi-wavelet	Decomposition level					
			1	2	3	4	5	6
 Cara	16:1	GHM	8248.7	38.541	11.345	10.24	10.138	10.135
			9.065	32.29	37.62	38.07	38.11	38.12
			0.03618	0.8883	0.9421	0.9453	0.9457	0.9457
			0.503	0.9728	0.9894	0.9899	0.9899	0.9899
		SA4	44.06	9.0035	8.2593	8.3198	8.3657	8.3696
			31.71	38.62	39.00	38.97	38.95	38.94
			0.8648	0.9475	0.9496	0.9495	0.9492	0.9492
			0.9668	0.9902	0.9895	0.9895	0.9895	0.9895
		CL	47.142	9.0648	8.0847	8.0233	8.0603	8.0547
			31.42	38.59	39.09	39.12	39.10	39.10
			0.8598	0.9462	0.9494	0.9496	0.9495	0.9495
			0.9691	0.9896	0.9894	0.9896	0.9897	0.9897
		Alpert	43.916	8.9286	7.7402	7.6756	7.6529	7.6628
			31.73	38.66	39.28	39.32	39.33	39.33
			0.8621	0.9463	0.9503	0.9504	0.9505	0.9504
			0.9751	0.9905	0.9904	0.9909	0.9909	0.9909
		New1	43.581	8.7188	7.5929	7.5075	7.4977	7.5047
			31.77	38.76	39.36	39.41	39.42	39.41
			0.8625	0.9474	0.9512	0.9514	0.9514	0.9514
			0.9753	0.9907	0.9906	0.9910	0.9910	0.9910
		New2	48.679	14.159	13.083	13.242	13.257	13.293
			31.30	36.67	37.01	36.96	36.96	36.94
			0.8571	0.9265	0.9291	0.9283	0.9282	0.9280
			0.9739	0.9852	0.9849	0.9846	0.9846	0.9846
		Integer Haar	54.35	13.133	11.83	11.86	11.861	11.881
			30.81	36.99	37.44	37.43	37.43	37.43
			0.8485	0.9311	0.9344	0.9338	0.9336	0.9336
			0.9602	0.9862	0.9864	0.9864	0.9865	0.9865
	32:1	GHM	10501	3488	27.742	22.497	22.009	21.981
			8.051	12.82	33.75	34.66	34.76	34.76
			0.01585	0.06779	0.9044	0.9153	0.9165	0.9165
			0.521	0.3933	0.9764	0.9796	0.9796	0.9796
		SA4	4664.8	22.996	18.123	18.228	18.414	18.425
			11.57	34.56	35.60	35.58	35.54	35.53
			0.3833	0.9111	0.9221	0.9216	0.9210	0.9210
			0.5896	0.9780	0.9799	0.9783	0.9783	0.9784
		CL	4721.1	24.749	18.57	18.076	18.272	18.254
			11.52	34.25	35.49	35.61	35.56	35.57
			0.3841	0.9051	0.9191	0.9203	0.9197	0.9198
			0.5860	0.9765	0.9800	0.9798	0.9798	0.9798
		Alpert	4761	23.861	17.324	17.001	16.96	17.015
			11.49	34.41	35.80	35.88	35.89	35.88
			0.3867	0.905	0.9204	0.921	0.9211	0.9208
			0.6056	0.9789	0.9823	0.9822	0.9826	0.9826
		New1	4761.00	23.364	17.139	16.803	16.787	16.845
			11.49	34.50	35.84	35.93	35.93	35.92
			0.3867	0.9070	0.9218	0.9228	0.9228	0.9226
			0.6056	0.9795	0.9826	0.9827	0.9831	0.9831
		New2	4761	35.318	29.309	29.573	29.675	29.859
			11.49	32.73	33.53	33.50	33.48	33.45
			0.3867	0.8801	0.8888	0.8871	0.8865	0.886
			0.6056	0.9710	0.9719	0.9696	0.9694	0.9694
		Integer Haar	4770.8	33.309	26.202	26.049	26.042	26.146
			11.48	32.97	34.01	34.03	34.03	34.02
			0.3792	0.8852	0.8964	0.8953	0.8950	0.8946
			0.5835	0.9703	0.9740	0.9729	0.9730	0.9731

Table 9

Results for Voit. The four lines for each multiwavelet represent MSE, PSNR, SSIM and MS-SSIM, in this order.

Image	CR	Multi-wavelet	Decomposition level					
			1	2	3	4	5	6
 Voit	4:1	GHM	192.89	34.812	31.389	31.084	31.061	31.056
			25.29	32.73	33.18	33.22	33.22	33.22
			0.7299	0.8896	0.898	0.8988	0.8989	0.8989
			0.9547	0.9867	0.9876	0.9877	0.9877	0.9877
		SA4	27.374	24.717	24.37	24.369	24.377	24.378
			33.77	34.22	34.28	34.28	34.27	34.27
			0.9055	0.9136	0.9147	0.9148	0.9148	0.9148
			0.9874	0.9881	0.9882	0.9882	0.9882	0.9882
		CL	27.639	24.311	23.967	23.917	23.932	23.931
			33.73	34.29	34.35	34.36	34.35	34.35
			0.9032	0.9134	0.9147	0.9149	0.9148	0.9148
			0.9854	0.9865	0.9865	0.9866	0.9866	0.9866
		Alpert	27.554	24.036	23.544	23.432	23.424	23.429
			33.74	34.33	34.42	34.44	34.45	34.44
			0.9036	0.9136	0.915	0.9155	0.9155	0.9154
			0.9888	0.9881	0.9879	0.9879	0.9878	0.9879
		New1	27.205	23.891	23.427	23.347	23.338	23.342
			33.80	34.36	34.45	34.46	34.46	34.46
			0.9047	0.9144	0.9159	0.9162	0.9163	0.9163
			0.9891	0.9883	0.9882	0.9881	0.9881	0.9881
		New2	30.78	28.92	29.171	29.252	29.260	29.267
			33.26	33.54	33.50	33.49	33.49	33.49
			0.896	0.9014	0.9011	0.9009	0.9009	0.9009
			0.9874	0.9853	0.9845	0.9845	0.9846	0.9846
		Integer Haar	29.85	26.893	26.702	26.65	26.652	26.655
			33.39	33.85	33.88	33.89	33.89	33.89
			0.8971	0.9053	0.906	0.9061	0.9062	0.9061
			0.9847	0.9856	0.9856	0.9856	0.9855	0.9855
	8:1	GHM	3941.9	105.44	82.277	80.29	80.138	80.112
			12.20	27.93	29.01	29.11	29.12	29.12
			0.2293	0.7473	0.7846	0.789	0.7894	0.7894
			0.6774	0.9559	0.9632	0.9637	0.9638	0.9638
		SA4	82.81	64.558	62.664	62.671	62.667	62.688
			28.97	30.06	30.18	30.18	30.18	30.18
			0.7800	0.8197	0.8242	0.8242	0.8242	0.8241
			0.9581	0.9668	0.9669	0.9669	0.9669	0.9669
		CL	85.127	64.228	62.134	61.891	61.934	61.933
			28.86	30.08	30.22	30.24	30.24	30.24
			0.7739	0.8167	0.8216	0.8223	0.8222	0.8222
			0.9543	0.9630	0.9636	0.9636	0.9636	0.9636
		Alpert	85.654	63.808	61.202	60.714	60.674	60.696
			28.83	30.10	30.29	30.32	30.32	30.32
			0.7712	0.8172	0.8228	0.8239	0.824	0.8239
			0.9665	0.9665	0.9661	0.966	0.966	0.966
		New1	84.412	63.328	60.746	60.379	60.354	60.379
			28.89	30.14	30.32	30.35	30.35	30.35
			0.774	0.8191	0.8247	0.8256	0.8257	0.8256
			0.9672	0.9672	0.9666	0.9667	0.9667	0.9667
		New2	93.636	75.327	74.947	75.224	75.275	75.314
			28.44	29.39	29.41	29.40	29.39	29.39
			0.757	0.7939	0.7959	0.7953	0.7952	0.7951
			0.9626	0.9588	0.957	0.9567	0.9567	0.9567
		Integer Haar	90.456	69.434	67.948	67.661	67.633	67.637
			28.59	29.74	29.83	29.85	29.85	29.85
			0.7632	0.804	0.8067	0.8072	0.8073	0.8073
			0.9514	0.9604	0.9603	0.9605	0.9605	0.9605

Table 9(continued)

Results for Voit. The four lines for each multiwavelet represent MSE, PSNR, SSIM and MS-SSIM, in this order.

Image	CR	Multi-wavelet	Decomposition level					
			1	2	3	4	5	6
 Voit	16:1	GHM	7554.6	375.82	164.29	154.67	153.91	153.84
			9.367	22.39	26.00	26.27	26.29	26.29
			0.0826	0.5001	0.6575	0.6707	0.6718	0.6719
			0.6229	0.8494	0.9211	0.9228	0.9232	0.9232
		SA4	395.03	129.48	121.36	121.23	121.23	121.27
			22.17	27.04	27.32	27.32	27.32	27.32
			0.5099	0.6940	0.7097	0.7098	0.7099	0.7098
			0.8262	0.9278	0.9293	0.9295	0.9295	0.9295
		CL	405.90	130.05	120.54	119.55	119.59	119.62
			22.06	27.02	27.35	27.38	27.38	27.38
			0.5011	0.6914	0.7073	0.7092	0.7095	0.7094
			0.8423	0.9208	0.9236	0.9235	0.9233	0.9233
		Alpert	415.65	128.04	118.22	116.63	116.52	116.58
			21.97	27.08	27.43	27.49	27.49	27.49
			0.4804	0.6921	0.7084	0.7112	0.7115	0.7113
			0.8864	0.9269	0.9293	0.9300	0.9299	0.9299
		New1	414.61	127.11	117.48	116.14	116.05	116.12
			21.98	27.12	27.46	27.51	27.51	27.51
			0.4808	0.6939	0.7108	0.7132	0.7134	0.7133
			0.8861	0.9280	0.9305	0.9308	0.9307	0.9306
		New2	417.16	149.21	143.56	144.00	144.33	144.43
			21.95	26.42	26.59	26.58	26.57	26.57
			0.4808	0.6636	0.6741	0.6736	0.673	0.6729
			0.8871	0.9144	0.9121	0.9112	0.9106	0.9107
		Integer Haar	422.81	139.29	131.03	130.01	129.68	129.72
			21.88	26.72	26.98	27.02	27.03	27.03
			0.4923	0.6754	0.6887	0.6903	0.6907	0.6908
			0.8197	0.9178	0.9189	0.9201	0.9199	0.9199
	32:1	GHM	9930.8	3519.9	283.21	248.98	246.14	245.9
			8.178	12.69	23.64	24.20	24.25	24.25
			0.0484	0.1632	0.5327	0.5633	0.5664	0.5667
			0.5877	0.5275	0.8622	0.8701	0.871	0.8711
		SA4	3805.5	236.61	201.48	199.9	199.67	199.86
			12.35	24.42	25.12	25.15	25.16	25.15
			0.2487	0.5628	0.6022	0.6038	0.6043	0.6042
			0.6103	0.8725	0.8811	0.8805	0.8804	0.8803
		CL	3939.5	244.59	201.47	197.79	197.64	197.75
			12.20	24.27	25.11	25.20	25.20	25.20
			0.24	0.5619	0.6009	0.6052	0.6054	0.6053
			0.6241	0.8595	0.8706	0.8716	0.8713	0.8713
		Alpert	4044.6	236.81	197.71	192.21	191.70	191.90
			12.08	24.41	25.20	25.32	25.33	25.33
			0.2242	0.5604	0.5999	0.6061	0.6069	0.6067
			0.6053	0.8754	0.8799	0.8817	0.8817	0.8816
		New1	4044.6	236.46	197.54	192.33	191.93	192.16
			12.08	24.42	25.20	25.32	25.33	25.32
			0.2242	0.5611	0.6008	0.6067	0.6075	0.6073
			0.6053	0.8760	0.8805	0.8820	0.8819	0.8818
		New2	4044.6	273.36	238.9	238.69	239.17	239.67
			12.08	23.79	24.38	24.39	24.38	24.37
			0.2242	0.5342	0.5636	0.5634	0.563	0.5628
			0.6053	0.8639	0.8544	0.8527	0.8520	0.8522
		Integer Haar	4029.1	264.94	222.82	218.29	216.56	216.82
			12.10	23.93	24.68	24.77	24.8	24.8
			0.2329	0.5406	0.5734	0.5753	0.5771	0.5768
			0.6029	0.8535	0.8621	0.8637	0.8631	0.863

Table 10

Results for Texture 1. The four lines for each multiwavelet represent MSE, PSNR, SSIM and MS-SSIM, in this order.


Image	CR	Multi-wavelet	Decomposition level					
			1	2	3	4	5	6
 Texture 1	2:1	GHM	79.713	53.973	51.129	50.784	50.761	50.754
			29.31	30.99	31.22	31.25	31.25	31.25
			0.9875	0.9898	0.9900	0.9900	0.9900	0.9900
			0.9919	0.9939	0.9941	0.9941	0.9941	0.9941
		SA4	46.476	43.135	42.608	42.554	42.556	42.554
			31.63	31.95	32.00	32.01	32.01	32.01
			0.9854	0.9847	0.9848	0.9848	0.9848	0.9848
			0.9922	0.9924	0.9924	0.9924	0.9924	0.9924
		CL	46.341	43.049	42.432	42.344	42.334	42.33
			31.65	31.97	32.03	32.04	32.04	32.04
			0.9788	0.98	0.98	0.98	0.98	0.98
			0.9905	0.9909	0.9908	0.9908	0.9908	0.9908
		Alpert	213.6	274.26	312.27	361.87	437.23	485.49
			24.97	23.88	23.31	22.64	21.79	21.31
			0.9768	0.8952	0.8301	0.7752	0.7414	0.7329
			0.9829	0.9545	0.9156	0.8599	0.7942	0.7681
		New1	46.024	42.953	42.32	42.242	42.223	42.227
			31.67	31.97	32.03	32.04	32.04	32.04
			0.9853	0.9848	0.9848	0.9847	0.9847	0.9847
			0.9938	0.9928	0.9926	0.9926	0.9926	0.9926
		New2	46.177	44.069	44.158	44.243	44.24	44.238
			31.66	31.85	31.84	31.83	31.83	31.83
			0.9854	0.9843	0.9844	0.9845	0.9845	0.9845
			0.9922	0.9920	0.9920	0.9920	0.9920	0.9920
		Integer Haar	45.865	45.222	45.112	45.123	45.112	45.108
			31.69	31.74	31.75	31.75	31.75	31.75
			0.9844	0.9825	0.9825	0.9825	0.9825	0.9825
			0.9936	0.9918	0.9916	0.9916	0.9916	0.9916
	4:1	GHM	432.58	230.44	212.9	210.92	210.76	210.73
			21.94	24.66	25	25.04	25.05	25.05
			0.9051	0.9466	0.9488	0.9493	0.9493	0.9493
			0.9442	0.9702	0.9717	0.9718	0.9718	0.9718
		SA4	204.65	184.38	181.18	180.82	180.81	180.78
			25.17	25.62	25.7	25.71	25.71	25.71
			0.9301	0.9288	0.9296	0.9296	0.9296	0.9296
			0.9626	0.9648	0.9650	0.9649	0.9649	0.9649
		CL	206.16	185.71	182.11	181.55	181.53	181.52
			25.14	25.59	25.68	25.69	25.69	25.69
			0.9049	0.9114	0.9120	0.9122	0.9122	0.9122
			0.9568	0.9586	0.9586	0.9586	0.9586	0.9586
		Alpert	326.72	366.72	401.02	450.07	525.38	573.66
			23.13	22.62	22.22	21.70	21.00	20.60
			0.9435	0.8529	0.7824	0.725	0.6897	0.6807
			0.9658	0.9328	0.8902	0.833	0.7671	0.7410
		New1	205.68	185.71	182.16	181.61	181.53	181.53
			25.15	25.59	25.67	25.69	25.69	25.69
			0.9299	0.9272	0.9277	0.9276	0.9277	0.9277
			0.9705	0.9657	0.9649	0.9648	0.9648	0.9648
		New2	206.37	190.66	190.61	191.07	191.08	191.07
			25.14	25.47	25.47	25.46	25.46	25.46
			0.9296	0.9243	0.9246	0.9247	0.9248	0.9248
			0.9622	0.9617	0.9612	0.9613	0.9613	0.9613
		Integer Haar	207.13	189.59	187.42	187.32	187.29	187.27
			25.12	25.50	25.55	25.55	25.55	25.55
			0.9245	0.9201	0.9203	0.9205	0.9205	0.9205
			0.9690	0.9625	0.9614	0.9614	0.9614	0.9614

Table 11

Results for Texture 4. The four lines for each multiwavelet represent MSE, PSNR, SSIM and MS-SSIM, in this order.

Image	CR	Multi-wavelet	Decomposition level						
			1	2	3	4	5	6	7
 Texture 4	64:1	GHM	1.10E+04 7.54 0.2919 0.7652	4985.0 11.16 0.3080 0.6724	58.32 30.52 0.9815 0.9716	15.81 36.19 0.9976 0.9921	12.58 37.20 0.9957 0.9926	12.00 37.41 0.9953 0.9928	12.00 37.41 0.9952 0.9928
			5995.0 10.36 0.4384 0.8289	28.03 33.72 0.9844 0.9828	4.139 42.01 0.9983 0.9962	3.3107 42.99 0.9963 0.9962	3.5373 42.70 0.9945 0.9954	3.77 42.42 0.9945 0.9951	3.745 42.45 0.9944 0.9951
			6035 10.33 0.4443 0.8326	37.53 32.48 0.9838 0.9789	1.913 45.40 0.9995 0.9989	1.191 47.50 0.9995 0.9991	1.193 47.49 0.9986 0.9989	1.222 47.39 0.9985 0.9988	1.251 47.29 0.9985 0.9988
			6039 10.32 0.4362 0.8236	19.85 35.16 0.9875 0.9975	37.37 32.41 0.9937 0.9936	86.73 28.75 0.9758 0.9813	119.8 27.35 0.9497 0.9672	141 26.64 0.9135 0.9625	179.2 25.60 0.8859 0.9658
		New1	6039 10.32 0.4362 0.8236	7.116 39.72 0.9882 0.9983	0.5232 50.95 0.9998 0.9996	0.4045 52.07 0.9998 0.9996	0.4921 51.22 0.9998 0.9995	0.4427 51.68 0.9998 0.9996	0.4368 51.74 0.9998 0.9996
			6064 10.31 0.4421 0.8311	40.49 32.14 0.9830 0.9790	1.642 46.09 0.9996 0.9992	0.9366 48.52 0.9997 0.9995	0.9678 48.39 0.9996 0.9995	1.135 47.70 0.9995 0.9994	1.108 47.80 0.9995 0.9994
			6039 10.32 0.4362 0.8236	8.795 38.80 0.9880 0.9981	0.4513 51.59 0.9998 0.9996	0.3109 53.21 0.9999 0.9998	0.2893 53.54 0.9999 0.9998	0.3237 53.05 0.9999 0.9997	0.3197 53.10 0.9999 0.9997
			Integer Haar						
	128:1	GHM	1.30E+04 6.99 0.2381 0.7586	8189 9.00 0.2386 0.6887	1352 16.90 0.5278 0.7073	57.81 30.53 0.9768 0.9683	44.90 31.63 0.9841 0.9734	42.65 31.85 0.9831 0.9742	42.63 31.85 0.9826 0.9741
			9496 8.36 0.3120 0.7853	1607 16.18 0.6811 0.8825	29.78 33.41 0.9748 0.9739	20.07 35.13 0.9777 0.9779	20.75 34.99 0.9757 0.9757	22.44 34.64 0.9749 0.9740	22.18 34.69 0.9741 0.9742
			9540 8.34 0.3168 0.7950	1710 15.88 0.6488 0.8553	28.20 33.66 0.9839 0.9820	16.94 35.87 0.9934 0.9871	15.61 36.23 0.9899 0.9860	15.62 36.22 0.9865 0.9856	15.91 36.14 0.9864 0.9854
			9551 8.34 0.3114 0.7847	1554 16.32 0.6895 0.9100	41.02 32.01 0.9930 0.9925	88.57 28.66 0.9751 0.9805	121.6 27.29 0.9490 0.9664	143.1 26.58 0.9125 0.9614	181.2 25.55 0.8850 0.9648
		New1	9551 8.34 0.3114 0.7847	1553 16.32 0.6895 0.9102	5.55 40.69 0.9987 0.9979	2.956 43.43 0.9987 0.9981	2.869 43.56 0.9985 0.9980	3.475 42.73 0.9979 0.9974	3.353 42.89 0.9979 0.9974
			9575 8.33 0.3130 0.7922	1707 15.91 0.6827 0.8808	37.31 32.45 0.9795 0.9774	20.02 35.18 0.9903 0.9854	20.08 35.17 0.9869 0.9840	23.18 34.54 0.9843 0.9813	22.46 34.68 0.9826 0.9816
			9551 8.34 0.3114 0.7847	1555 16.31 0.6893 0.9096	6.976 39.70 0.9984 0.9975	3.422 42.80 0.9983 0.9979	3.308 42.95 0.9982 0.9978	3.863 42.28 0.9969 0.9971	3.732 42.43 0.9969 0.9971
			Integer Haar						

Table 11(continued)

Results for Texture 4. The four lines for each multiwavelet represent MSE, PSNR, SSIM and MS-SSIM, in this order.

Image	CR	Multi-wavelet	Decomposition level						
			1	2	3	4	5	6	7
 Texture 4	256:1	GHM	1.41E+04 6.66 0.1936 0.6961	1.10E+04 7.71 0.2070 0.7198	4983 11.16 0.2806 0.5966	188.4 25.41 0.8938 0.8823	94.64 28.38 0.9424 0.9365	86.37 28.78 0.9505 0.9428	86.37 28.78 0.9468 0.9425
			1.18E+04 7.42 0.2406 0.7621	6094 10.29 0.4089 0.7943	129.1 27.06 0.9504 0.9373	62.03 30.22 0.9417 0.9405	64.08 30.08 0.9329 0.9323	72.09 29.57 0.9268 0.9253	70.10 29.69 0.9241 0.9266
			1.18E+04 7.41 0.2422 0.7600	6175 10.23 0.4048 0.7877	155.1 26.26 0.9314 0.9183	61.75 30.24 0.9692 0.9614	52.84 30.92 0.9685 0.9617	52.23 30.97 0.9637 0.9616	53.44 30.87 0.9617 0.9606
			1.19E+04 7.39 0.2370 0.7431	6164 10.24 0.4230 0.8101	78.59 29.18 0.9756 0.9858	98.25 28.21 0.9721 0.9776	130.5 26.98 0.9458 0.9632	153.9 26.26 0.9085 0.9568	191.1 25.32 0.8807 0.9604
		New1	1.19E+04 7.39 0.2370 0.7431	6164 10.24 0.4230 0.8101	53.05 30.90 0.9788 0.9883	15.89 36.12 0.9929 0.9926	15.05 36.36 0.9902 0.9909	18.74 35.41 0.9839 0.9864	17.72 35.65 0.9839 0.9870
			1.19E+04 7.39 0.2414 0.7546	6223 10.19 0.4131 0.7978	208.1 24.98 0.9390 0.9253	81.83 29.02 0.9520 0.9537	79.52 29.14 0.9408 0.9466	90.38 28.59 0.9184 0.9329	87.29 28.74 0.9202 0.9362
			1.19E+04 7.39 0.2370 0.7431	6164 10.24 0.4230 0.8101	67.16 29.87 0.9767 0.9863	19.00 35.35 0.9916 0.9919	17.91 35.60 0.9892 0.9905	20.98 34.92 0.9808 0.9855	19.78 35.17 0.9810 0.9862
			Integer Haar						
	512:1	GHM	1.48E+04 6.44 0.1744 0.6838	1.27E+04 7.11 0.1956 0.7316	8213 8.99 0.2181 0.6184	1631 16.06 0.4446 0.6104	199.0 25.15 0.8571 0.8629	159.7 26.11 0.8906 0.8875	158.3 26.15 0.8844 0.8874
			1.36E+04 6.81 0.2092 0.7466	9585 8.32 0.2908 0.7514	1751 15.78 0.6348 0.8420	130.7 26.98 0.9128 0.9097	134.2 26.87 0.8702 0.8847	155.9 26.21 0.8515 0.8666	149.6 26.39 0.8526 0.8703
			1.36E+04 6.79 0.2094 0.7419	9684 8.28 0.2820 0.7520	2083 15.00 0.5581 0.7533	144.5 26.55 0.9317 0.9184	108 27.81 0.9426 0.9365	105.2 27.92 0.9382 0.9373	108.2 27.80 0.9333 0.9352
			1.37E+04 6.78 0.2026 0.7207	9680 8.28 0.2956 0.7580	1747 15.79 0.6476 0.8690	124.6 27.18 0.9558 0.9721	152.5 26.3 0.9376 0.9559	179.8 25.59 0.8982 0.9460	214.2 24.83 0.8714 0.9511
		New1	1.37E+04 6.78 0.2026 0.7207	9680 8.28 0.2956 0.7580	1745 15.80 0.6485 0.8704	49.30 31.21 0.9734 0.9841	44.09 31.69 0.9696 0.9755	56.05 30.65 0.9436 0.9588	51.34 31.03 0.9448 0.9614
			1.37E+04 6.78 0.2056 0.7302	9728 8.26 0.2953 0.7685	1905 15.41 0.6384 0.8447	193.4 25.28 0.9318 0.9242	178.1 25.64 0.8936 0.9051	199.3 25.15 0.8371 0.8780	193.2 25.28 0.8451 0.8819
			1.37E+04 6.78 0.2026 0.7207	9680 8.28 0.2956 0.7580	1750 15.78 0.6470 0.8678	60.02 30.35 0.9713 0.9827	53.95 30.82 0.9662 0.9730	62.12 30.21 0.9386 0.9589	56.73 30.60 0.9414 0.9619
			Integer Haar						

Table 12

Results for Brick wall. The four lines for each multiwavelet represent MSE, PSNR, SSIM and MS-SSIM, in this order.

Image	CR	Multi-wavelet	Decomposition level						
			1	2	3	4	5	6	7
 Brick wall	16:1	GHM	5617	278.9	141.7	135	133.8	133.6	133.6
			10.64	23.68	26.62	26.83	26.87	26.87	26.87
			0.5177	0.9870	0.9860	0.9859	0.9859	0.9859	0.9859
			0.7018	0.9006	0.9383	0.9414	0.9420	0.9421	0.9421
		SA4	288.7	111.8	106.7	105.8	105.5	105.5	105.4
			23.53	27.65	27.85	27.89	27.90	27.90	27.90
			0.9648	0.9854	0.9850	0.9852	0.9851	0.9851	0.9851
			0.8836	0.9463	0.9493	0.9499	0.9501	0.9501	0.9501
		CL	319.4	109.1	103.4	102.5	102.2	102.2	102.2
			23.09	27.75	27.99	28.02	28.04	28.04	28.04
			0.9231	0.9856	0.9843	0.9842	0.9841	0.9841	0.9841
			0.8869	0.9432	0.9463	0.9467	0.9469	0.9469	0.9469
		Alpert	298.8	167.6	252.1	366.5	459.3	495.9	511.3
			23.38	25.89	24.12	22.49	21.51	21.18	21.04
			0.9594	0.9782	0.9285	0.8381	0.8100	0.8058	0.8046
			0.9162	0.9340	0.8850	0.8149	0.7861	0.7817	0.7801
		New1	296.1	109.1	103.5	102.7	102.5	102.5	102.5
			23.42	27.75	27.98	28.01	28.02	28.03	28.03
			0.9593	0.9848	0.9844	0.9846	0.9845	0.9845	0.9845
			0.9165	0.9467	0.9491	0.9496	0.9497	0.9497	0.9497
		New2	332	112.2	107.1	106.8	106.8	106.8	106.8
			22.92	27.63	27.83	27.84	27.85	27.85	27.85
			0.9533	0.9835	0.9829	0.9830	0.9830	0.9830	0.9830
			0.8750	0.9431	0.9463	0.9468	0.9469	0.9469	0.9469
		Integer Haar	307.2	112.4	106.2	105.3	105.2	105.3	105.2
			23.26	27.62	27.87	27.91	27.91	27.91	27.91
			0.9581	0.9831	0.9830	0.9832	0.9832	0.9832	0.9832
			0.9138	0.9422	0.9449	0.9454	0.9456	0.9454	0.9456
	32:1	GHM	7665	2719	232.3	200	195.4	194.5	194.4
			9.29	13.79	24.47	25.12	25.22	25.24	25.24
			0.3534	0.5730	0.9697	0.9713	0.9716	0.9716	0.9717
			0.7400	0.6433	0.8972	0.9075	0.9092	0.9096	0.9097
		SA4	3205	171.8	149.9	146.6	145.8	145.7	145.7
			13.07	25.78	26.37	26.47	26.49	26.50	26.50
			0.5909	0.9726	0.9735	0.9742	0.9742	0.9742	0.9742
			0.6891	0.9117	0.9214	0.9234	0.9238	0.9239	0.9239
		CL	3327	175.2	145.4	142.5	141.6	141.5	141.5
			12.91	25.7	26.5	26.59	26.62	26.62	26.62
			0.6012	0.9751	0.9733	0.9733	0.9734	0.9735	0.9735
			0.6929	0.9109	0.9191	0.9204	0.9209	0.9210	0.9210
		Alpert	3344	217.7	282.6	394.9	487	523.5	538.9
			12.89	24.75	23.62	22.17	21.26	20.94	20.82
			0.5870	0.9664	0.9193	0.8290	0.7997	0.7953	0.7941
			0.6540	0.9086	0.8656	0.7967	0.7673	0.7629	0.7612
		New1	3344	173	145.3	142.5	141.7	141.6	141.5
			12.89	25.75	26.51	26.59	26.62	26.62	26.62
			0.587	0.9722	0.9735	0.9738	0.9738	0.9738	0.9738
			0.654	0.9166	0.9215	0.9228	0.9231	0.9231	0.9232
		New2	3451	191.2	155	152.1	151.7	151.7	151.6
			12.75	25.32	26.23	26.31	26.32	26.32	26.32
			0.5871	0.968	0.9699	0.9703	0.9702	0.9703	0.9703
			0.6808	0.9093	0.9163	0.9175	0.9177	0.9177	0.9177
		Integer Haar	3344	192.1	152.4	148.5	147.9	147.8	147.8
			12.89	25.3	26.3	26.41	26.43	26.43	26.43
			0.5871	0.9686	0.9707	0.9711	0.9709	0.9710	0.9710
			0.6539	0.9115	0.9156	0.9170	0.9171	0.9171	0.9171

Table 13(continued)

Results for Brick wall. The four lines for each multiwavelet represent MSE, PSNR, SSIM and MS-SSIM, in this order.


Image	CR	Multi-wavelet	Decomposition level						
			1	2	3	4	5	6	7
 Brick wall	64:1	GHM	9198	5278	576.6	308.7	288.3	284.4	283.8
			8.494	10.91	20.52	23.24	23.53	23.59	23.60
			0.1961	0.4234	0.8255	0.9446	0.9481	0.9493	0.9495
			0.6883	0.6511	0.7417	0.8630	0.8697	0.8715	0.8718
		SA4	5513.0	624.0	216.5	201.3	198	197.5	197.4
			10.72	20.18	24.78	25.09	25.16	25.18	25.18
			0.4275	0.7833	0.9552	0.9591	0.9595	0.9597	0.9597
			0.6928	0.7199	0.8904	0.8968	0.8981	0.8984	0.8984
		CL	5624	649.3	218.5	203	199.2	198.6	198.5
			10.63	20.01	24.74	25.06	25.14	25.15	25.15
			0.4359	0.7845	0.9614	0.9608	0.9609	0.9611	0.9611
			0.7105	0.7042	0.8912	0.8948	0.8958	0.8960	0.8961
		Alpert	5644	650.1	337.1	439.2	528.9	564.9	580.2
			10.62	20.00	22.85	21.70	20.90	20.61	20.50
			0.4280	0.7709	0.9047	0.8160	0.7858	0.7813	0.7801
			0.6696	0.7310	0.8469	0.7807	0.7510	0.7465	0.7449
		New1	5644	641.2	215.7	201.1	197.6	197.0	196.9
			10.62	20.06	24.79	25.10	25.17	25.19	25.19
			0.4280	0.7674	0.9596	0.9608	0.9608	0.9609	0.9609
			0.6696	0.7299	0.8987	0.9012	0.9017	0.9019	0.9019
		New2	5779	692	279	242.1	235.6	234.9	234.7
			10.51	19.73	23.67	24.29	24.41	24.42	24.43
			0.4263	0.7541	0.9380	0.9473	0.9487	0.9487	0.9488
			0.6910	0.6892	0.8691	0.8841	0.8867	0.8871	0.8871
		Integer Haar	5644	647	244.6	222.2	218.3	217.9	217.8
			10.62	20.02	24.25	24.66	24.74	24.75	24.75
			0.4280	0.7617	0.9530	0.9557	0.9557	0.9558	0.9558
			0.6695	0.7253	0.8910	0.8943	0.8946	0.8946	0.8947

Table 13

Results for Texture 5. The four lines for each multiwavelet represent MSE, PSNR, SSIM and MS-SSIM, in this order.


Image	CR	Multi-wavelet	Decomposition level					
			1	2	3	4	5	6
 Texture 5	4:1	GHM	67.96	16.95	16.25	16.22	16.21	16.21
			29.81	35.84	36.02	36.03	36.03	36.03
			0.9055	0.9605	0.9616	0.9617	0.9617	0.9617
			0.9922	0.9967	0.9968	0.9968	0.9968	0.9968
		SA4	11.63	7.95	7.82	7.82	7.81	7.81
			37.48	39.13	39.20	39.20	39.20	39.20
			0.9693	0.9788	0.9793	0.9793	0.9793	0.9793
			0.9972	0.9981	0.9981	0.9981	0.9981	0.9981
		CL	21.75	13.21	12.89	12.89	12.89	12.89
			34.76	36.92	37.03	37.03	37.03	37.03
			0.9462	0.9661	0.9669	0.9669	0.9669	0.9669
			0.9949	0.9967	0.9968	0.9968	0.9968	0.9968
		Alpert	29.08	63.98	213.98	575.5	1043	1412.5
			33.50	30.07	24.83	20.53	17.95	16.63
			0.9633	0.9298	0.8160	0.7121	0.6869	0.6741
			0.9965	0.9905	0.9495	0.8567	0.7826	0.7496
		New1	5.71	3.73	3.65	3.65	3.65	3.65
			40.57	42.41	42.51	42.51	42.51	42.51
			0.9835	0.9898	0.9900	0.9900	0.9900	0.9900
			0.9985	0.9991	0.9991	0.9991	0.9991	0.9991
		New2	8.28	8.67	8.91	8.92	8.92	8.92
			38.95	38.75	38.63	38.63	38.63	38.63
			0.9766	0.9758	0.9755	0.9755	0.9755	0.9755
			0.9978	0.9977	0.9977	0.9977	0.9977	0.9977
		Integer Haar	7.81	7.9	7.91	7.89	7.89	7.89
			39.20	39.15	39.15	39.16	39.16	39.16
			0.9777	0.9780	0.9780	0.9780	0.9780	0.9780
			0.9976	0.9977	0.9977	0.9977	0.9977	0.9977
	8:1	GHM	1569.8	59.01	51.79	51.48	51.46	51.462
			16.17	30.42	30.99	31.01	31.02	31.02
			0.4348	0.9100	0.9141	0.9146	0.9146	0.9146
			0.8711	0.99100	0.9913	0.9914	0.9914	0.9914
		SA4	54.59	29.02	27.94	27.90	27.87	27.86
			30.76	33.50	33.67	33.67	33.68	33.68
			0.8931	0.9410	0.9435	0.9435	0.9435	0.9435
			0.9860	0.9936	0.9939	0.9939	0.9939	0.9939
		CL	95.17	46.01	43.40	43.32	43.32	43.32
			28.35	31.50	31.76	31.76	31.76	31.76
			0.8366	0.9143	0.9183	0.9182	0.9182	0.9182
			0.9816	0.9900	0.9905	0.9905	0.9905	0.9905
		Alpert	61.65	80.89	229.31	590.77	1058.3	1427.7
			30.23	29.05	24.53	20.42	17.88	16.58
			0.9070	0.9002	0.7934	0.6947	0.6709	0.6588
			0.9882	0.9865	0.9448	0.8515	0.7771	0.7441
		New1	40.33	20.62	19.31	19.27	19.27	19.27
			32.07	34.99	35.27	35.28	35.28	35.28
			0.9136	0.9537	0.9555	0.9556	0.9556	0.9556
			0.9889	0.9950	0.9952	0.9952	0.9952	0.9952
		New2	44.37	44.23	47.84	47.99	47.968	47.968
			31.66	31.67	31.33	31.32	31.32	31.32
			0.9074	0.9107	0.9051	0.9051	0.9052	0.9052
			0.9864	0.9867	0.9857	0.9857	0.9857	0.9857
		Integer Haar	49.65	47.34	46.84	46.88	46.90	46.889
			31.17	31.38	31.42	31.42	31.42	31.42
			0.9002	0.9003	0.9012	0.9011	0.9010	0.9011
			0.9856	0.9862	0.9868	0.9868	0.9868	0.9868

Table 13(continued)

Results for Texture 5. The four lines for each multiwavelet represent MSE, PSNR, SSIM and MS-SSIM, in this order.


Image	CR	Multi-wavelet	Decomposition level					
			1	2	3	4	5	6
 Texture 5	16:1	GHM	5720	170.5	118.2	116.5	116.4	116.4
			10.56	25.81	27.41	27.47	27.47	27.47
			0.1313	0.8197	0.8578	0.8594	0.8596	0.8596
			0.6833	0.9729	0.9810	0.9812	0.9813	0.9813
		SA4	221.13	80.21	71.77	71.32	71.22	71.2
			24.68	29.09	29.57	29.60	29.61	29.61
			0.7388	0.8777	0.8902	0.8908	0.8910	0.8910
			0.9544	0.9819	0.9845	0.9845	0.9845	0.9845
		CL	341.3	116	102.3	101.7	101.6	101.6
			22.80	27.49	28.03	28.06	28.06	28.06
			0.6459	0.8391	0.8558	0.8560	0.8563	0.8564
			0.9508	0.9751	0.9786	0.9787	0.9787	0.9787
		Alpert	208.5	129.4	267.8	629.1	1097	1466
			24.94	27.01	23.85	20.14	17.73	16.47
			0.7655	0.8446	0.7545	0.6628	0.6411	0.6299
			0.9610	0.9752	0.9347	0.8403	0.7653	0.7323
		New1	195.64	72.66	61.17	60.82	60.85	60.84
			25.22	29.52	30.27	30.29	30.29	30.29
			0.7651	0.8808	0.8967	0.8971	0.8972	0.8971
			0.9609	0.9818	0.9849	0.9850	0.9850	0.9850
		New2	215.9	149.6	166.1	168.9	168.7	168.7
			24.79	26.38	25.93	25.85	25.86	25.86
			0.7518	0.7897	0.7724	0.7691	0.7692	0.7691
			0.9521	0.9556	0.9499	0.9493	0.9494	0.9495
		Integer Haar	214.67	149.7	148.24	148.5	148.8	148.6
			24.81	26.38	26.42	26.41	26.41	26.41
			0.7568	0.7772	0.7764	0.7760	0.7756	0.7757
			0.9577	0.9588	0.9612	0.9613	0.9613	0.9613
	32:1	GHM	9408	1443	217.3	206.2	205.4	205.4
			8.4	16.54	24.76	24.99	25.01	25.01
			0.0538	0.4411	0.7761	0.7836	0.7843	0.7843
			0.5223	0.8358	0.9554	0.9579	0.9582	0.9582
		SA4	1699	205.3	154.4	152.7	152.3	152.1
			15.83	25.01	26.24	26.29	26.30	26.31
			0.4132	0.7512	0.8132	0.8157	0.8161	0.8162
			0.8201	0.9469	0.9634	0.9644	0.9645	0.9646
		CL	1932	264.5	207	203.7	203.5	203.4
			15.27	23.91	24.97	25.04	25.05	25.05
			0.3508	0.7098	0.7720	0.7757	0.7761	0.7760
			0.8272	0.9350	0.9545	0.9556	0.9557	0.9557
		Alpert	1732	260.1	341.5	700.4	1168	1538
			15.74	23.98	22.8	19.68	17.46	16.26
			0.4198	0.7234	0.7021	0.6192	0.5989	0.5881
			0.8061	0.9380	0.9161	0.8205	0.7446	0.7112
		New1	1732	215.8	150.5	148.9	148.7	148.7
			15.74	24.79	26.36	26.4	26.41	26.41
			0.4198	0.7344	0.8105	0.8126	0.8127	0.8127
			0.8061	0.9387	0.9616	0.9624	0.9623	0.9623
		New2	1823	371.5	398.3	411.7	412.2	412.1
			15.52	22.43	22.13	21.98	21.98	21.98
			0.4163	0.6328	0.6186	0.6038	0.6031	0.6031
			0.8126	0.8987	0.8844	0.8824	0.8822	0.8823
		Integer Haar	1732	347.1	327.9	329.9	331.3	331
			15.74	22.73	22.97	22.95	22.93	22.93
			0.4196	0.6340	0.6382	0.6331	0.6319	0.6323
			0.8061	0.9101	0.9176	0.9171	0.9169	0.9170

Table 14

Results for Subwater. The four lines for each multiwavelet represent MSE, PSNR, SSIM and MS-SSIM, in this order.

Image	CR	Multi-wavelet	Decomposition level				
			1	2	3	4	5
 Subwater	8:1	GHM	4140	0.9046	0.703	0.6931	0.6917
			6.91	31.22	31.95	32.00	32.00
			0.0569	0.6589	0.6601	0.6601	0.6602
			0.3204	0.6656	0.6657	0.6657	0.6657
		SA4	0.8359	0.4812	0.4567	0.4561	0.4569
			31.45	33.05	33.19	33.20	33.19
			0.6581	0.6617	0.6619	0.6619	0.6619
			0.6651	0.6659	0.6659	0.6659	0.6659
		CL	1.004	0.5448	0.5036	0.5002	0.499
			30.91	32.68	32.91	32.93	32.94
			0.6566	0.6611	0.6614	0.6615	0.6615
			0.6652	0.6657	0.6658	0.6658	0.6658
		Alpert	1.443	2.786	7.515	15.13	39.26
			29.86	27.95	25.11	23.11	20.45
			0.6554	0.6518	0.6399	0.6309	0.6255
			0.6651	0.6641	0.6561	0.6420	0.6150
		New1	0.9442	0.5117	0.4716	0.4686	0.4687
			31.10	32.87	33.11	33.13	33.13
			0.6569	0.6613	0.6616	0.6617	0.6617
			0.6650	0.6659	0.6659	0.6659	0.6659
		New2	1.114	0.8457	0.844	0.8431	0.8438
			30.61	31.41	31.41	31.42	31.41
			0.6557	0.6585	0.6585	0.6585	0.6585
			0.6646	0.6651	0.6651	0.6651	0.6651
		Integer Haar	1.385	0.9685	0.9087	0.9023	0.9031
			29.98	31.01	31.20	31.22	31.22
			0.6532	0.6572	0.6577	0.6577	0.6577
			0.6644	0.6650	0.6651	0.6651	0.6651
	16:1	GHM	7027	29.38	2.214	2.063	2.053
			5.35	21.28	28.63	28.83	28.84
			0.0103	0.6278	0.6512	0.6520	0.6521
			0.3318	0.6234	0.6634	0.6635	0.6635
		SA4	43.59	1.647	1.372	1.355	1.36
			20.07	29.48	30.01	30.04	30.03
			0.6121	0.6532	0.6556	0.6557	0.6557
			0.6073	0.6636	0.6642	0.6642	0.6642
		CL	37.81	1.914	1.523	1.482	1.474
			20.50	29.05	29.70	29.78	29.80
			0.6109	0.6513	0.6543	0.6545	0.6546
			0.6218	0.6631	0.6639	0.6639	0.6639
		Alpert	13.6	3.72	8.119	15.7	39.82
			23.38	27.11	24.88	23.00	20.40
			0.6079	0.6451	0.6355	0.6268	0.6214
			0.6555	0.6625	0.6549	0.6407	0.6137
		New1	13.6	1.681	1.322	1.284	1.282
			23.38	29.42	30.11	30.20	30.20
			0.6079	0.6525	0.6552	0.6555	0.6555
			0.6555	0.6635	0.6643	0.6643	0.6643
		New2	47.96	3.24	3.304	3.273	3.29
			19.80	27.52	27.46	27.48	27.47
			0.6038	0.6406	0.6405	0.6405	0.6403
			0.6047	0.6599	0.6597	0.6598	0.6598
		Integer Haar	13.6	2.953	2.553	2.516	2.519
			23.38	27.78	28.20	28.24	28.24
			0.60790	0.6430	0.6460	0.6463	0.6462
			0.6555	0.6608	0.6622	0.6623	0.6622

Table 14(continued)

Results for Subwater. The four lines for each multiwavelet represent MSE, PSNR, SSIM and MS-SSIM, in this order.

Image	CR	Multi-wavelet	Decomposition level				
			1	2	3	4	5
 Subwater	32:1	GHM	8976	3250	9.484	6.745	6.603
			4.63	7.62	24.46	25.42	25.48
			0.0065	0.0652	0.6322	0.6362	0.6366
			0.3774	0.2303	0.6523	0.6553	0.6555
		SA4	3944	6.575	4.264	4.018	4.036
			7.09	25.48	26.72	26.89	26.88
			0.2598	0.6276	0.6409	0.6423	0.6422
			0.3466	0.6531	0.6578	0.6583	0.6583
		CL	3977.4	7.37	4.771	4.271	4.222
			7.08	25.14	26.39	26.71	26.74
			0.2638	0.6221	0.6382	0.6407	0.6409
			0.3683	0.6523	0.6572	0.6585	0.6586
		Alpert	3955.2	6.846	9.572	16.943	41.073
			7.1	25.35	24.4	22.77	20.31
			0.2626	0.6251	0.6275	0.6203	0.6150
			0.3969	0.6543	0.6523	0.6382	0.6112
		New1	3955	5.416	3.4	3.155	3.137
			7.10	26.02	27.37	27.59	27.60
			0.2626	0.6282	0.6430	0.6444	0.6446
			0.3968	0.6542	0.6603	0.6606	0.6607
		New2	4046	11.35	10.11	9.914	10.04
			7.02	23.88	24.22	24.27	24.23
			0.2552	0.6038	0.6070	0.6066	0.6059
			0.3440	0.6463	0.6452	0.646	0.6458
		Integer Haar	3955.2	7.689	5.9762	5.6887	5.6662
			7.10	25.01	25.74	25.88	25.89
			0.2626	0.6164	0.6275	0.6289	0.6290
			0.3968	0.6507	0.6557	0.6562	0.6563

Table 15

The ϵ values for denoising of 1D Piece-Polynomial, Cusp and HeaviSine signals by using orthogonal multiwavelet filters.

Signal	Multiwavelet	ϵ	
		$\sigma = 10$	$\sigma = 15$
Piece-Polynomial	GHM	28.60	41.67
	SA4	28.36	41.54
	CL	31.47	46.49
	Alpert	69.66	74.91
	Integer Haar	29.52	36.00
	New1	26.32	31.80
	New2	27.34	36.89
Cusp	GHM	0.0900	0.0887
	SA4	0.0974	0.1086
	CL	0.0905	0.0919
	Alpert	0.0495	0.0534
	Integer Haar	0.0562	0.0687
	New1	0.0440	0.0506
	New2	0.0398	0.0464
HeaviSine	GHM	0.7560	0.8731
	SA4	0.6690	0.8544
	CL	0.7089	0.8032
	Alpert	1.1430	1.2355
	Integer Haar	0.5114	0.7404
	New1	0.5496	0.7658
	New2	0.4250	0.6944




Figure 9: Original and Noisy 1D signals with added AWGN with variance $\sigma = 10, 15$; (a) Piece-Polynomial with variance $\sigma = 10$; (b) Piece-Polynomial with variance $\sigma = 15$; (c) Cusp with variance $\sigma = 10$; (d) Cusp with variance $\sigma = 15$; (e) HeaviSine with variance $\sigma = 10$; (f) HeaviSine with variance $\sigma = 15$.




Figure 10: Denoising of 1D signals with added AWGN with variance $\sigma = 10, 15$ by using orthogonal multiwavelet banks and applied soft thresholding; (a) Piece-Polynomial with variance $\sigma = 10$; (b) Piece-Polynomial with variance $\sigma = 15$; (c) Cusp with variance $\sigma = 10$; (d) Cusp with variance $\sigma = 15$; (e) HeaviSine with variance $\sigma = 10$; (f) HeaviSine with variance $\sigma = 15$.

Table 16

Comparative results of MSE, PSNR, SSIM and MS-SSIM of denoised image 'Cameraman' of size 256×256 pixels for seven orthogonal multiwavelets through $J = 5$ decomposition levels with AWGN and variance $\sigma = 10$.

Image	Threshold	Multi-wavelet	Decomposition Levels				
			1	2	3	4	5
 Cameraman	Soft	GHM	71.08	72.02	75.95	78.16	78.99
			29.61	29.56	29.33	29.2	29.16
			0.7803	0.8450	0.8539	0.8542	0.8542
			0.8077	0.8125	0.8101	0.8043	0.8026
		SA4	66.5	70.00	74.23	76.64	77.24
			29.90	29.68	29.42	29.29	29.25
			0.7892	0.8495	0.8566	0.8563	0.8565
			0.8025	0.8165	0.8149	0.7991	0.7960
		CL	69.57	72.49	76.70	78.86	79.48
			29.71	29.53	29.28	29.16	29.13
			0.7851	0.8467	0.8553	0.8554	0.8558
			0.8117	0.8100	0.8104	0.7972	0.7953
		Alpert	129.73	212.88	318.79	431.91	550.35
			27.00	24.85	23.10	21.78	20.72
			0.7508	0.751	0.7010	0.6738	0.6599
			0.7972	0.7809	0.7246	0.6410	0.5467
		New1	67.77	69.45	73.07	75.02	75.66
			29.82	29.71	29.49	29.38	29.34
			0.7855	0.8458	0.8552	0.8558	0.8559
			0.8013	0.8140	0.8277	0.8207	0.8168
		New2	67.77	72.416	76.08	77.99	78.77
			29.82	29.53	29.32	29.21	29.17
			0.7855	0.8397	0.8494	0.8508	0.8513
			0.8013	0.8043	0.8016	0.7870	0.7862
		Integer Haar	72.33	74.82	78.29	80.28	80.97
			29.54	29.39	29.19	29.08	29.05
			0.7784	0.8351	0.8451	0.8459	0.8451
			0.8044	0.8174	0.8190	0.8088	0.8033
	hard	GHM	62.96	56.45	55.52	55.43	55.46
			30.14	30.61	30.69	30.69	30.69
			0.7733	0.8337	0.8430	0.8435	0.8435
			0.8121	0.8270	0.8352	0.8374	0.8370
		SA4	59.7	53.38	52.64	52.65	52.65
			30.37	30.86	30.92	30.92	30.92
			0.7806	0.8393	0.8463	0.8464	0.8464
			0.8033	0.8285	0.8336	0.8326	0.8324
		CL	59.84	53.12	52.14	52.06	52.07
			30.36	30.88	30.96	30.97	30.97
			0.7783	0.8390	0.8480	0.8483	0.8483
			0.8170	0.8228	0.8311	0.8320	0.8318
		Alpert	95.12	146.3	228.29	326.6	435.01
			28.35	26.48	24.55	22.99	21.75
			0.7715	0.8003	0.7649	0.7392	0.7255
			0.8004	0.8006	0.7708	0.7074	0.6176
		New1	59.42	52.45	51.35	51.36	51.36
			30.39	30.93	31.03	31.03	31.02
			0.7797	0.8388	0.8491	0.8494	0.8493
			0.8028	0.8201	0.8383	0.8367	0.8364
		New2	59.42	52.49	51.57	51.49	51.49
			30.39	30.93	31.01	31.01	31.01
			0.7797	0.8346	0.8449	0.8457	0.8457
			0.8028	0.8142	0.8207	0.8244	0.8241
		Integer Haar	61.24	54.26	53.17	53.21	53.2
			30.26	30.79	30.87	30.87	30.87
			0.7727	0.8292	0.8385	0.8389	0.8389
			0.8075	0.8264	0.8342	0.8311	0.8311

Table 17

Comparative results of MSE, PSNR, SSIM and MS-SSIM of denoised image 'Couple' of size 256×256 pixels for seven orthogonal multiwavelets through $J = 5$ decomposition levels with AWGN and variance $\sigma=10$.

Image	Threshold	Multi-wavelet	Decomposition Levels				
			1	2	3	4	5
 Couple	Soft	GHM	92.29	100.35	107.19	110.23	111.19
			28.48	28.12	27.83	27.71	27.67
			0.8162	0.8049	0.7955	0.7936	0.7935
			0.9488	0.9370	0.9238	0.9181	0.9181
		SA4	87.91	98.62	105.69	108.22	109.12
			28.69	28.19	27.89	27.79	27.75
			0.8206	0.8060	0.7967	0.7953	0.7951
			0.9505	0.9398	0.9275	0.9241	0.9236
		CL	91.44	102.99	110.09	112.76	113.7
			28.52	28.00	27.71	27.61	27.57
			0.8169	0.7998	0.7895	0.7885	0.7883
			0.9460	0.9339	0.9181	0.9133	0.9130
		Alpert	155.72	239.46	320.22	394.98	479.76
			26.21	24.34	23.08	22.17	21.32
			0.7496	0.6423	0.5611	0.5394	0.5347
			0.9435	0.8982	0.7995	0.7082	0.6526
		New1	89.19	100.26	106.99	109.52	110.37
			28.63	28.12	27.84	27.74	27.70
			0.8164	0.8002	0.7904	0.7896	0.7895
			0.9509	0.9425	0.9290	0.9245	0.9238
		New2	89.19	106.46	113.41	116.14	117.09
			28.63	27.86	27.58	27.48	27.45
			0.8164	0.7885	0.7820	0.7828	0.7832
			0.9509	0.9339	0.9162	0.9116	0.9113
		Integer Haar	95.04	107.04	114.19	116.87	117.73
			28.35	27.84	27.55	27.45	27.42
			0.8051	0.7864	0.7782	0.7775	0.7777
			0.9493	0.9364	0.9197	0.9135	0.9130
	hard	GHM	77.31	75.38	75.63	75.66	75.66
			29.25	29.36	29.34	29.34	29.34
			0.8324	0.8384	0.8379	0.8378	0.8378
			0.9542	0.9549	0.9539	0.9538	0.9538
		SA4	73.9	72.06	72.1	72.12	72.12
			29.44	29.55	29.55	29.55	29.55
			0.8366	0.8420	0.8421	0.8420	0.8420
			0.9529	0.9547	0.9546	0.9545	0.9545
		CL	74.33	72.47	72.65	72.68	72.68
			29.42	29.53	29.52	29.52	29.52
			0.8357	0.8408	0.8402	0.8402	0.8402
			0.9550	0.9539	0.9522	0.9520	0.9520
		Alpert	109.94	155.68	210.01	269.04	344.7
			27.72	26.21	24.91	23.83	22.76
			0.7959	0.7447	0.6960	0.6792	0.6743
			0.9488	0.9268	0.8716	0.8082	0.7615
		New1	73.91	71.85	71.88	71.893	71.89
			29.44	29.57	29.56	29.56	29.56
			0.8360	0.8423	0.8425	0.8424	0.8424
			0.9533	0.9547	0.9548	0.9548	0.9548
		New2	73.91	73.08	73.29	73.34	73.34
			29.44	29.49	29.48	29.48	29.48
			0.8360	0.8382	0.8384	0.8383	0.8383
			0.9533	0.9521	0.9522	0.9521	0.9521
		Integer Haar	76.616	74.57	74.75	74.78	74.78
			29.29	29.4	29.39	29.39	29.39
			0.8308	0.8361	0.8359	0.8358	0.8358
			0.9541	0.9548	0.9540	0.9540	0.9540

Table 18

Comparative results of MSE, PSNR, SSIM and MS-SSIM of denoised image 'Kiel' of size 512×512 pixels for seven orthogonal multiwavelets through $J = 5$ decomposition levels with AWGN and variance $\sigma=10$.

Image	Threshold	Multi-wavelet	Decomposition Levels				
			1	2	3	4	5
 Kiel	Soft	GHM	105.02	114.96	121.17	123.68	124.52
			27.92	27.53	27.3	27.21	27.18
			0.9139	0.9247	0.9236	0.9221	0.9220
			0.8155	0.8069	0.7988	0.7941	0.7924
		SA4	102.14	114.43	120.9	123.5	124.3
			28.04	27.55	27.31	27.21	27.19
			0.9026	0.9271	0.9268	0.9250	0.9249
			0.8183	0.8085	0.7996	0.7949	0.7922
		CL	104.94	116.99	123.39	125.82	126.59
			27.92	27.45	27.22	27.13	27.11
			0.9178	0.9200	0.9175	0.9164	0.9162
			0.8120	0.8042	0.7944	0.7924	0.7894
		Alpert	248.68	369.09	470.63	555.71	630.52
			24.17	22.46	21.40	20.68	20.13
			0.9000	0.8484	0.7684	0.7140	0.6970
			0.8081	0.7658	0.6911	0.6106	0.5539
		New1	103.02	114.26	120.64	122.91	123.66
			28.00	27.55	27.32	27.23	27.21
			0.9069	0.9263	0.9261	0.9252	0.9252
			0.8184	0.8121	0.8085	0.8095	0.8109
		New2	103.02	118.5	124.74	127.12	127.87
			28.00	27.39	27.17	27.09	27.06
			0.9069	0.9192	0.9195	0.9192	0.9198
			0.8184	0.8063	0.7977	0.7937	0.7878
		Integer Haar	107.76	119.93	126.47	128.7	129.46
			27.81	27.34	27.11	27.03	27.01
			0.9162	0.9227	0.9191	0.9181	0.9181
			0.8164	0.8091	0.8058	0.8047	0.8057
	hard	GHM	75.26	71.62	71.21	71.12	71.11
			29.37	29.58	29.61	29.61	29.61
			0.9209	0.9426	0.9504	0.9517	0.9518
			0.8244	0.8296	0.8345	0.8384	0.8385
		SA4	72.91	69.49	69.17	69.16	69.16
			29.50	29.71	29.73	29.73	29.73
			0.9032	0.9400	0.9482	0.9487	0.9487
			0.8222	0.8291	0.8306	0.8312	0.8316
		CL	73.15	69.15	68.64	68.57	68.59
			29.49	29.73	29.76	29.77	29.77
			0.9322	0.9416	0.9490	0.9502	0.9501
			0.8257	0.8277	0.8308	0.8350	0.8341
		Alpert	167.83	241.1	314.31	384.3	451.25
			25.88	24.31	23.16	22.28	21.59
			0.9063	0.8913	0.8462	0.8069	0.7918
			0.8147	0.7932	0.7522	0.6975	0.6503
		New1	72.757	68.89	68.35	68.26	68.24
			29.51	29.75	29.78	29.79	29.79
			0.9088	0.9408	0.9503	0.9517	0.9518
			0.8221	0.8249	0.8323	0.8374	0.8379
		New2	72.76	69.58	69.00	68.91	68.9
			29.51	29.71	29.74	29.75	29.75
			0.9088	0.9384	0.9481	0.9496	0.9497
			0.8221	0.8223	0.8270	0.8289	0.8283
		Integer Haar	74.41	70.74	70.22	70.13	70.11
			29.41	29.63	29.67	29.67	29.67
			0.9234	0.9445	0.9512	0.9525	0.9526
			0.8227	0.8272	0.8354	0.8384	0.8386

Table 19

Comparative results of MSE, PSNR, SSIM and MS-SSIM of denoised image 'House' of size 256×256 pixels for seven orthogonal multiwavelets through $J = 5$ decomposition levels with AWGN and variance $\sigma=20$.

Image	Threshold	Multi-wavelet	Decomposition Levels				
			1	2	3	4	5
 House	Soft	GHM	142.45	96.18	96.29	101.75	104.51
			26.59	28.30	28.30	28.06	27.94
			0.5531	0.7296	0.7803	0.7856	0.7872
			0.6539	0.6569	0.6459	0.6264	0.6176
		SA4	130.61	87.73	90.28	95.79	98.45
			26.97	28.70	28.57	28.32	28.20
			0.5651	0.7388	0.7842	0.7878	0.7887
			0.6466	0.6643	0.6473	0.6351	0.6320
		CL	138.84	100.21	99.7	105.1	107.66
			26.71	28.12	28.14	27.91	27.81
			0.5577	0.7272	0.7799	0.7862	0.7875
			0.6554	0.6531	0.6477	0.6297	0.6172
		Alpert	141.88	132.45	200.02	283.49	384.37
			26.61	26.91	25.12	23.61	22.28
			0.5528	0.6944	0.6912	0.6868	0.6927
			0.6480	0.6505	0.5989	0.5288	0.4256
		New1	133.63	87.54	85.925	89.234	91.48
			26.87	28.71	28.79	28.63	28.52
			0.5597	0.7335	0.7837	0.7917	0.7932
			0.6476	0.6644	0.6758	0.6801	0.6740
		New2	133.63	100.62	101.43	105.88	108.33
			26.87	28.10	28.07	27.88	27.78
			0.5597	0.7095	0.7612	0.7754	0.7771
			0.6476	0.6520	0.6508	0.6391	0.6247
		Integer Haar	143.67	101.04	100.53	104.21	106.35
			26.56	28.09	28.11	27.95	27.86
			0.5475	0.7072	0.7585	0.7668	0.7706
			0.6517	0.6625	0.6673	0.6627	0.6537
	hard	GHM	166.77	118.57	110.57	109.79	109.77
			25.91	27.39	27.69	27.73	27.73
			0.5216	0.6514	0.6826	0.6850	0.6850
			0.6563	0.6721	0.6790	0.6796	0.6801
		SA4	156.48	109.25	102.48	102.17	102.2
			26.19	27.75	28.02	28.04	28.04
			0.5336	0.6620	0.6876	0.6884	0.6884
			0.6457	0.6739	0.6786	0.6792	0.6795
		CL	161.72	113.67	105.08	104.22	104.28
			26.04	27.57	27.92	27.95	27.95
			0.5262	0.6575	0.6907	0.6930	0.6930
			0.6589	0.6681	0.6772	0.6768	0.6767
		Alpert	137.42	101.72	139.34	200.79	284.32
			26.75	28.06	26.69	25.10	23.59
			0.5559	0.721	0.7417	0.7385	0.7405
			0.6485	0.6607	0.6396	0.5943	0.5093
		New1	157.12	107.16	99.04	98.1	98.07
			26.17	27.83	28.17	28.21	28.22
			0.5307	0.6669	0.6996	0.7034	0.7035
			0.6463	0.6640	0.6785	0.6861	0.6870
		New2	157.12	113.38	106.22	105.33	105.27
			26.17	27.59	27.87	27.91	27.91
			0.5307	0.6502	0.6859	0.6915	0.6918
			0.6463	0.6579	0.6674	0.6665	0.6646
		Integer Haar	163.3	115.52	107.55	106.88	106.86
			26.00	27.50	27.81	27.84	27.84
			0.5218	0.6471	0.6795	0.6830	0.6831
			0.6509	0.6662	0.6805	0.6826	0.6806

Table 20

Comparative results of MSE, PSNR, SSIM and MS-SSIM of denoised image 'Kiel' of size 512×512 pixels for seven orthogonal multiwavelets through $J = 5$ decomposition levels with AWGN and variance $\sigma=20$.

Image	Threshold	Multi-wavelet	Decomposition Levels				
			1	2	3	4	5
 Kiel	Soft	GHM	272.71	268.66	281.78	289.61	292.65
			23.77	23.84	23.63	23.51	23.47
			0.7821	0.8254	0.8385	0.8360	0.8356
			0.7432	0.7304	0.7120	0.6992	0.6950
		SA4	261.74	264.28	278.39	286.46	289.35
			23.95	23.91	23.68	23.56	23.52
			0.7550	0.8282	0.8447	0.8416	0.8413
			0.7447	0.7331	0.7145	0.7010	0.7007
		CL	269.97	271.63	285.83	293.45	296.15
			23.82	23.79	23.57	23.46	23.42
			0.8041	0.8211	0.8283	0.8275	0.8273
			0.7408	0.7282	0.7063	0.6973	0.6955
		Alpert	397.34	507.77	625.36	723.45	805.9
			22.14	21.07	20.17	19.54	19.07
			0.7625	0.7572	0.6895	0.6297	0.6121
			0.7376	0.6973	0.6087	0.5148	0.4615
		New1	266.38	266.5	281.11	288.16	290.8
			23.88	23.87	23.64	23.53	23.49
			0.7635	0.8250	0.8398	0.8394	0.8398
			0.7448	0.7374	0.7212	0.7160	0.7216
		New2	266.38	282.67	296.43	303.67	306.29
			23.88	23.62	23.41	23.31	23.27
			0.7635	0.8061	0.8221	0.8246	0.8269
			0.7448	0.7269	0.7024	0.6943	0.6920
		Integer Haar	279.29	281.94	297.11	304.12	306.81
			23.67	23.63	23.40	23.30	23.26
			0.7870	0.8229	0.8262	0.8249	0.8254
			0.7435	0.7312	0.7130	0.7051	0.7106
	hard	GHM	242.43	215.75	212.99	212.68	212.67
			24.28	24.79	24.85	24.85	24.85
			0.7892	0.8453	0.8722	0.8773	0.8775
			0.7518	0.7577	0.7605	0.7642	0.7644
		SA4	232.79	206.91	204.37	204.5	204.49
			24.46	24.97	25.03	25.02	25.02
			0.7550	0.8418	0.8710	0.8738	0.8740
			0.7479	0.7572	0.7577	0.7588	0.7597
		CL	235.51	207.86	204.95	204.51	204.5
			24.41	24.95	25.01	25.02	25.02
			0.8170	0.8441	0.8682	0.8734	0.8734
			0.7536	0.7549	0.7555	0.7610	0.7611
		Alpert	303.59	344.75	413.51	482.18	548.84
			23.31	22.76	21.97	21.30	20.74
			0.7678	0.8029	0.7818	0.7474	0.7329
			0.7428	0.7245	0.6792	0.6277	0.5912
		New1	232.7	205.71	202.83	202.36	202.26
			24.46	25.00	25.06	25.07	25.07
			0.7650	0.8412	0.8717	0.8780	0.8786
			0.7485	0.7533	0.7575	0.7658	0.7698
		New2	232.7	211.29	207.95	207.64	207.57
			24.46	24.88	24.95	24.96	24.96
			0.7650	0.8313	0.8623	0.8686	0.8694
			0.7485	0.7482	0.7509	0.7556	0.7551
		Integer Haar	239.53	213.74	210.94	210.72	210.63
			24.34	24.83	24.89	24.89	24.90
			0.7947	0.8507	0.8729	0.8778	0.8784
			0.7504	0.7550	0.7594	0.7645	0.7689