

SAMI3D-DW: Interactive Segmentation of Any 3D Medical Images

Ping Gong^{1,*}, Shiyuan Su^{1,*}, Fandong Zhang¹, Xinchun Han¹,
Haowei Sun¹, Yiming Li¹, Yizhou Yu^{1,2}

¹Artificial Intelligence Laboratory, Deepwise Healthcare

²The University of Hong Kong

*Equal contribution.

Abstract

Interactive segmentation of 3D medical images supports quantitative analysis of anatomical structures and disease while allowing users to specify and refine their targets. Despite substantial progress by nnInteractive and VISTA3D, reliable segmentation across diverse clinical targets remains challenging, particularly for complex anatomical structures and the heterogeneous, long-tailed spectrum of pathology. We present SAMI3D-DW V1 (hereafter SAMI3D-DW), an interactive 3D segmentation model trained on Deepwise’s large-scale proprietary medical image datasets. We evaluate the model under simulated user interactions on a CT/MR benchmark comprising 4,326 cases from 219 source datasets, spanning 107 anatomical and pathological categories, organized by a medical taxonomy and evaluated with a category-balanced DSC score. SAMI3D-DW achieves the highest category-macro Dice among evaluated methods in both interaction modes. With one point, it scores 0.5764 versus 0.5315 for nnInteractive, the strongest baseline, rising to 0.7771 versus 0.7494 with five points. With bounding-box initialization, the scores are 0.7130 versus 0.6530. After five corrective clicks, SAMI3D-DW reaches 0.8002 versus 0.7868, making it the only evaluated box-compatible model to exceed 0.80. For radiologists and clinicians, SAMI3D-DW enables segmentation of complex anatomical structures, including intracranial vessel trees on CT and MR angiography, with a few clicks. In a preliminary in-house comparison involving neurofibromatosis type 1 (NF1), SAMI3D-DW-assisted tumor annotation took minutes per case and approximately one-fifteenth of the time required for manual annotation, highlighting its potential to support volumetric treatment-response assessment.


Category-macro Dice ↑ · Higher is better

Axes start at 0.2 · Gains vs nnInteractive


CT/MR coverage: 107 categories, 405 specific targets, 219 datasets · 4,326 cases

Point-only

1st point




5 total points



Box-initialized

BBox initialization



BBox + 5 corrective points




Figure 1. Benchmark-wide performance with point-only and box-initialized interaction. Top: initial prompts. Bottom: five total points or five corrective points after a BBox. Bar endpoints show category-macro Dice on shared axes starting at 0.2; colors identify methods. Gains over nnInteractive use unrounded scores and are in percentage points (pp). Comparisons are descriptive where coverage of valid scores differs (Tables 3 and 4).

Contents

1	Introduction	3
2	Related Work	4
3	Data and Training Overview	6
3.1	Training Dataset Curation	6
3.2	Model Training	7
4	Evaluation Protocol	7
4.1	Test Dataset	7
4.2	Interaction Modes and Recommended Use	8
4.3	Simulated Interaction	11
4.4	Models and Protocol Compatibility	11
4.5	Metrics and Aggregation	11
5	Main Results	12
5.1	Point-Only Segmentation	12
5.2	Box-Initialized Segmentation and Refinement	12
5.3	Category-Level Results	13
5.4	Stratified Results	16
6	Limitations and Responsible Use	16
7	Conclusion	17
	Acknowledgments	17
A	Detailed Medical Target Evaluation	21

1. Introduction

Medical image segmentation delineates anatomical structures and pathological regions for quantitative analysis, treatment planning, and longitudinal assessment. Task-specific methods such as U-Net and nnU-Net have enabled accurate segmentation across many applications (Ronneberger et al., 2015; Isensee et al., 2021), but extending these systems to new targets or imaging settings generally requires additional annotation, training, and validation. Interactive segmentation offers a flexible way to specify the intended target and refine its boundaries at inference time.

SAM and MedSAM established widely used spatial prompting interfaces (Kirillov et al., 2023; Ma et al., 2024). SAM-Med3D and SegVol extended promptable segmentation to volumetric medical images (Wang et al., 2025; Du et al., 2024). VISTA3D integrates automatic segmentation with interactive editing, MedSAM2 adapts memory-based propagation to medical volumes and videos, and nnInteractive supports diverse prompts and iterative correction in 3D (He et al., 2025; Ma et al., 2025; Isensee et al., 2025). A central question for general-purpose use is how accurately these models segment diverse targets, including complex anatomical structures and heterogeneous pathology, from an initial prompt and how they respond to subsequent correction.

We present SAMI3D-DW, a model for interactive segmentation of 3D medical images, trained on Deepwise’s proprietary medical image datasets. The model produces full-volume masks from spatial prompts and supports point-only interaction as well as initialization with a two-dimensional (2D) bounding box followed by corrective points.

Evaluation must reflect both target diversity and aggregation choices. A few familiar datasets may leave relevant targets unrepresented, while dataset-level averages alone do not reveal repeated coverage of the same structures. Pooling all objects gives more weight to categories with more examples. A medical taxonomy supplies common reporting categories across sources, making anatomical and pathological coverage explicit. Full-volume evaluation and explicit interaction budgets are also essential for comparing interactive methods (Ulrich et al., 2025).

We therefore evaluate SAMI3D-DW on a benchmark comprising 4,326 cases from 219 source datasets and 107 medical categories, focusing on computed tomography (CT) and magnetic resonance (MR), the most common modalities for clinical volumetric imaging. Dataset labels are harmonized through a medical taxonomy. Controlled sampling limits contributions from large source datasets, while category-macro Dice averages object scores within each category and weights category means equally. Both interaction modes use simulated prompts, with performance reported from initialization through successive corrections (Section 4).

This report makes two principal contributions:

- An evaluation framework guided by a medical taxonomy, combining controlled sampling across diverse datasets with equal category weighting to complement dataset-level evaluation.
- SAMI3D-DW achieves the highest reported category-macro Dice among the methods evaluated under each interaction protocol. Compared with nnInteractive, the strongest evaluated baseline, SAMI3D-DW achieves 0.5764 versus 0.5315 with one point and 0.7771 versus 0.7494 with five total points. With box initialization, it achieves 0.7130 versus 0.6530. Results at the reported interaction budgets are reported in Section 5.


2. Related Work

From spatial prompts to richer spatial, semantic, and free-text interfaces. Figure 2A summarizes the expansion of target specification. SAM established a reusable point-, box-, and mask-prompted interface (Kirillov et al., 2023). Medical evaluations exposed domain gaps (Mazurowski et al., 2023; Huang et al., 2024), motivating MedSAM’s large-scale, box-conditioned medical adaptation and adapter-based methods (Ma et al., 2024; Cheng et al., 2023; Wu et al., 2025). ScribblePrompt, μ SAM, and CellSAM further developed biomedical interaction and automatic prompt generation (Wong et al., 2024; Archit et al., 2025; Marks et al., 2025). Semantic specification emerged through SAM 3’s concept prompts (Carion et al., 2025), MedCLIP-SAMv2’s text-to-spatial prompting (Koleilat et al., 2025), BioMedParse’s joint segmentation, detection, and recognition across biomedical modalities (Zhao et al., 2025a), and PathSegmentor’s pathology-specific natural-language interface (Chen et al., 2025). SAT and VoxTell extend medical terminology and free-text queries to volumes (Zhao et al., 2025b; Rokuss et al., 2025). These developments broaden how users express *what* to segment, but semantic intent does not necessarily determine an exact instance or boundary. Customer feedback at Deepwise and current technical limitations motivate retaining spatial prompts as a necessary tool for fast, high-accuracy interactive annotation: points, boxes, and other spatial cues provide direct localization and precise error correction, even when language supplies the initial target description.

From 2D images to 3D volumetric segmentation and correction. An equally important trend is the transition from slice-wise masks to volume-level prediction and refinement (Figure 2B). Earlier geodesic and memory-based systems already studied correction and unseen-target transfer (Wang et al., 2019; Luo et al., 2021; Zhou et al., 2023). MA-SAM, 3DSAM-adapter, and MedLSAM extend pretrained features or localization to volumes (Chen et al., 2024; Gong et al., 2024; Lei et al., 2025); SAM-Med3D, SegVol, VISTA3D, and SAT3D develop native volumetric prompting (Wang et al., 2025; Du et al., 2024; He et al., 2025; Peiris et al., 2026). SAM 2 and MedSAM2 reduce repeated initialization through memory-based slice propagation (Ravi et al., 2025; Ma et al., 2025). However, propagation remains slice-based: changes in anatomical structures and accumulated errors can still require inspection and correction across many slices. It does not by itself adequately resolve the multi-slice annotation burden motivating this report. nnInteractive is the state-of-the-art reference most closely aligned with this volume-level correction objective (Isensee et al., 2025). Its native 3D model turns points, boxes, scribbles, and lassos placed in 2D orthogonal views into full-volume masks, using previous-mask feedback and error-driven training to support iterative refinement. SAMI3D-DW continues this second line of development, focusing on *3D volumetric segmentation and correction driven by 2D spatial prompts*. We evaluate point-only and box-initialized corrective-point workflows, measuring full-volume quality over successive interactions rather than isolated prompted masks, consistent with interactive benchmarking practice (Ulrich et al., 2025).


A | From spatial prompts to richer spatial, semantic, and free-text interfaces

Dates = cited versions; branches group capabilities, not a uniform timeline



B | From 2D images to 3D volumetric segmentation and correction

Earlier interaction: DeepGeoS (2019) → MIDeepSeg (2021) → Volumetric Memory Network (2023)



→ interface / conceptual transition

---→ selected SAM adaptation

Not an inheritance or performance ranking

Figure 2. Two major trends in interactive medical segmentation: **A**, from spatial prompts to richer spatial, semantic, and free-text interfaces; **B**, from 2D images to 3D volumetric segmentation and correction, the direction pursued by SAMI3D-DW. Arrows indicate conceptual transitions or selected adaptations, not performance rankings or direct architectural inheritance. Dates follow the cited publication or preprint version; grouped branches are not a uniform timeline.

3. Data and Training Overview

3.1. Training Dataset Curation

The candidate training collection included 849 Deepwise in-house datasets. Three-stage curation yielded 115k 3D volumes (Figure 3).




Figure 3. Training data curation. FN and FP denote false-negative and false-positive annotation errors.

Annotation screening. Three radiologists of attending rank or above assessed false-negative (FN) and false-positive (FP) annotation errors, graded contour quality from 1 to 5 (Table 1), and described each target. Datasets with contour grades below 3 or extensive FN/FP errors were excluded.

Category balancing. Datasets were grouped by target description. Sampling caps limited each dataset’s contribution to balance target categories.

Paired review. SAMI3D researchers paired with a radiologist reviewed sampled annotations for quality and training suitability. Datasets judged too difficult for the training objective (e.g., soft-plaque delineation) or reliant on expert anatomical conventions (e.g., hepatic segments) were excluded.

Table 1. Contour-quality grading. Cases graded 1–2 were excluded.

Grade	Criteria
5: Excellent	Accurate boundaries with fine anatomical detail preserved.
4: Good	Minor focal contour deviations.
3: Acceptable	Overall anatomical structures preserved; localized errors requiring correction.
2: Poor	Major contour errors or substantial anatomical distortion.
1: Unusable	Absent required annotations, invalid masks, or image–mask mismatch.

3.2. Model Training

The architecture and detailed training configuration of SAMI3D-DW are not publicly disclosed at this stage. The model was trained from scratch on the Deepwise-curated datasets described above, without using any pretrained weights.

Training comprised two stages: initial training on the full curated dataset collection, followed by fine-tuning at a reduced learning rate on a subset of challenging datasets, particularly those involving lesions.

4. Evaluation Protocol

We evaluate SAMI3D-DW retrospectively under two simulated interaction modes. The point-only mode uses one to five cumulative foreground/background points and supports the broadest comparison across model interfaces. The BBox-initialized mode begins with one two-dimensional (2D) box and permits up to five corrective point prompts. Its correction sequence follows the structure of the [CVPR 2025 Foundation Models for Interactive 3D Biomedical Image Segmentation Challenge](#); the prompt construction and aggregation rules below define the protocol used in this report, and the two protocols are not identical. Results from the two interaction modes are reported separately because a box and a point convey different amounts of spatial information and need not require equal user effort.

4.1. Test Dataset

Medical taxonomy. Combining medical segmentation datasets requires more than concatenating their label lists. The same structure may have different names across sources, whereas similar names may denote different anatomical structures, pathology, laterality, or annotation policies. We harmonize source annotations with a medical taxonomy independent of dataset-specific label IDs. Each dataset label is mapped to a *Specific Target* (hereafter target), which identifies the medical object together with its modality, laterality, anatomical or pathological subdivision, and verified imaging sequence or phase, where available. Targets are grouped into pre-defined reporting categories:

$$\text{dataset label} \longrightarrow \text{specific target} \longrightarrow \text{medical category.} \quad (1)$$

Thus, 405 specific targets refine the underlying medical objects rather than represent 405 independent object types. These distinctions do not multiply the 107 reporting categories; category-balanced scoring is unchanged.

The taxonomy comprises three domains. *A: Anatomical and physiological structures* covers normal human anatomical structures and physiological constituents, including the liver, kidneys, heart, vascular trees, vertebrae, and muscles. *B: Pathology and abnormalities* covers disease-related lesions, abnormal tissues, and abnormal regions, including tumours such as neurofibromas, haemorrhage, infarction, and infectious lesions. *C: Other segmentation targets* covers targets outside normal human anatomical structures and human pathology, including non-human structures, medical implants, and foreign bodies. Assignment follows the annotated target: a liver mask belongs to A, whereas a liver tumour mask belongs to B, even when both originate from the same image.

Within A, structures are grouped by organ system and tissue family. Within B, lesions are grouped by pathological process, including neoplastic, infectious, inflammatory and immune-mediated, vascular, traumatic, focal non-neoplastic, and indeterminate abnormalities. These domain definitions describe the scope of the taxonomy; benchmark coverage is determined by the observed

annotations and evaluation mappings. Table 2 summarizes the domain definitions, illustrative examples, and counts for the represented families.

Benchmark coverage and sampling. The benchmark is designed to cover diverse anatomical and pathological targets across body regions in CT and MR. Sampling is controlled across source datasets to limit dominance by large, commonly used targets. Medical categories, rather than dataset identities, define the principal reporting units, and category means receive equal weight. This combination broadens coverage beyond a small set of familiar datasets and keeps less frequent target categories visible in the overall evaluation. Balanced weighting does not imply equal sample counts across categories or modalities, or a test distribution matching clinical prevalence.

Because this report studies interactive 3D segmentation, the frozen mixed-dimensional candidate pool is filtered at the source-dataset level using an audited dimensionality registry and, where necessary, lightweight spatial metadata. This removes 22 two-dimensional datasets, 754 cases, and 931 physical objects; no image or label voxels are read to make the dimensionality decision. We further restrict evaluation to CT and MR, removing two additional 3D datasets, 36 cases, and 418 objects from other modalities.

The resulting benchmark comprises 219 source datasets, 4,326 test cases, and 13,826 physical target objects. Curated mapping covers 405 specific targets and 107 categories with 13,969 associated instances: 63 categories of anatomical structures with 7,550 associated instances, 43 pathology categories with 6,394 associated instances, and one Other category with 25 associated instances. Each physical object is counted once per category, so the category-level instance total can exceed the physical-object count when an object maps to multiple medical concepts.

The benchmark contains CT (9,289 objects from 2,373 cases) and MR (4,537 objects from 1,953 cases), covering major body regions from the brain and head and neck through the thorax, abdomen, pelvis, spine, and extremities. Representative coverage is shown in Figure 4.

Training–test separation. None of the 4,326 test cases is used for training. Some source datasets also contribute different cases to model development. The results therefore measure performance on unseen cases from a heterogeneous multi-source benchmark; they do not establish transfer to entirely unseen datasets. We do not describe this evaluation as external, zero-shot, fully independent, or prospective validation. Dataset, institution, and case identifiers are withheld from the public report.

4.2. Interaction Modes and Recommended Use

A point is quick to place, straightforward to interpret as foreground or background, and available in all six displayed entries. Point-only prompting is therefore a practical interface and an important common comparison, not merely a diagnostic. Its first positive point localizes a target but does not necessarily identify it. This limitation is intrinsic to nested and adjacent structures: a point inside a liver tumour, for example, is simultaneously inside the tumour and the liver, so the intended mask cannot be inferred from that coordinate alone.

A BBox supplies approximate extent and generally narrows the set of plausible interpretations before refinement. It does not encode class identity or eliminate every ambiguity, but it provides a more informative spatial initialization across heterogeneous anatomical structures and pathological targets. We therefore recommend BBox initialization followed by corrective points when approximate extent is convenient to provide; point-only interaction remains the simpler option and enables comparison with point-only systems such as VISTA3D. The recommendation concerns the accuracy trajectories measured here and is not a claim about user time.

Table 2. Medical taxonomy and coverage of the scored test set. Domain definitions and illustrative examples describe the classification scope; counts refer only to represented families. Instance counts are deduplicated within each category; a physical object can contribute to more than one category.

Code	Taxonomy family	Categories	Instances
A: Anatomical and physiological structures			
<i>Scope:</i> Normal human anatomical structures and physiological constituents. <i>Examples:</i> Liver, kidneys, heart, vascular trees, vertebrae, and muscles.			
A01	Nervous system	4	224
A02	Cardiovascular system	15	1,107
A03	Respiratory system	2	463
A04	Digestive system	9	1,263
A05	Urinary system	4	481
A06	Reproductive system	2	97
A07	Endocrine system	3	296
A08	Lymphatic, immune, and hematopoietic systems	2	232
A09	Skeletal and articular system	14	2,813
A10	Muscular system	6	462
A12	Sensory system	2	112
<i>A subtotal</i>		63	7,550
B: Pathology and abnormalities			
<i>Scope:</i> Disease-related lesions, abnormal tissues, and abnormal regions. <i>Examples:</i> Tumours (including neurofibromas), haemorrhage, infarction, and infectious lesions.			
B01	Neoplastic lesions	23	3,579
B02	Infectious lesions	1	428
B03	Non-infectious inflammatory and immune-mediated lesions	1	149
B04	Vascular and circulatory disorders	2	634
B05	Trauma and tissue injury	2	238
B07	Non-neoplastic focal lesions	1	84
B10	Indeterminate imaging abnormalities	13	1,282
<i>B subtotal</i>		43	6,394
C: Other segmentation targets			
<i>Scope:</i> Targets outside normal human anatomical structures and human pathology. <i>Examples:</i> Non-human structures, medical implants, and foreign bodies.			
C99	Other uncategorized targets	1	25
Total		107	13,969




Figure 4. Anatomical, pathological, and modality coverage of the CT/MR benchmark. Medical illustrations show representative anatomical structures for all 11 represented anatomical families (A). The pathology panel (B) lists seven families, with bar lengths indicating category counts; the Other domain (C) retains non-human structures for complete accounting. Category-level instance counts are deduplicated within each category; a physical instance can contribute to more than one category. The overall count and modality shares use the 13,826 unique physical instances. Illustrations are cropped and arranged from [Servier Medical Art](#), licensed under [CC BY 4.0](#); they are not to scale and do not imply exhaustive organ coverage.

4.3. Simulated Interaction

The point-only protocol omits the BBox and applies five cumulative EDT-maximum points. The first point is generated from the target mask; later points use the same largest-error-component rule. Results are reported after one, three, and five points. All methods use the same initial point prompts. SAMI3D-DW and nnInteractive share the same cohort, point-selection rule, interaction budget, and metric implementation in both protocols. Subsequent locations can differ because they are generated from each model’s current error mask.

For the BBox-initialized protocol, the simulator places a tight 2D box around the largest connected component on the axial slice with the greatest target area; no in-plane margin is added. It then performs five correction rounds. At each round, the evaluator compares the current prediction with the reference mask, selects the largest 26-connected error component, and places a point at a maximally interior location determined by the Euclidean distance transform. False-negative and false-positive regions yield positive and negative points, respectively. The BBox, preceding points, and previous prediction are retained.

Every prompt is derived from a reference mask. The evaluation therefore models a deterministic, error-aware annotator under controlled conditions; it does not reproduce the imprecision, stopping decisions, timing, or variability of real users. Unless stated otherwise, all interactions in this report are simulated.

4.4. Models and Protocol Compatibility

The complete paired comparison in each interaction mode is between SAMI3D-DW and nnInteractive, for which all 13,826 objects and all 107 categories have valid inference results (Dice scores) at every stage. The point-only table additionally reports MedSAM2, SAM-Med3D, SegVol, and the VISTA3D CVPR 2025 checkpoint. VISTA3D is included here because its published interactive branch accepts positive and negative 3D points, but it is absent from the BBox table because that branch does not define a BBox input compatible with this protocol (He et al., 2025). For overlapping targets such as pancreas and pancreatic tumour, its formulation also assumes that target class x is known and uses that information in an ambiguity embedding. Its point-only values are therefore useful descriptive references but are not treated as a fully input-equivalent inferential comparison.

MedSAM2, SAM-Med3D, and SegVol also provide descriptive reference points under the BBox-plus-correction sequence. They have valid scores for 13,420, 13,823, and 13,540 objects, respectively, at each displayed stage. All three still represent all 107 categories. Missing or invalid scores are never replaced with zero, and method-specific coverage accompanies both comparisons (Tables 3 and 4).

The comparison concerns the evaluated configurations under our simulator. Published workflows differ: MedSAM2 initializes a box on the target’s middle slice and propagates bidirectionally, while SegVol’s main comparison uses box and text prompts together (Ma et al., 2025; Du et al., 2024). Rankings here are specific to the evaluated interaction modes rather than a summary of every prompting option offered by each method.

4.5. Metrics and Aggregation

The principal metric is Dice on the $[0, 1]$ scale. For each method and stage, we first average valid object scores within every category and then weight the available category means equally. For point-only interaction, we report category-macro Dice after one, three, and five points, making both immediate response and fixed-budget refinement visible. For BBox-initialized interaction,

we report initialization and predictions after one, three, and five corrections. We do not combine the two modes into an equal-effort ranking.

The size, spacing, and modality analyses in Section 5.4 are descriptive category-macro summaries. Within each subgroup, the comparison uses the categories with at least one valid score at each of the BBox and +5 stages for all five methods, so every method in a table row has the same category denominator. Size is defined by reference-mask voxel count (small: $< 10,000$; medium: $10,000\text{--}99,999$; large: $\geq 100,000$). Acquisition spacing is the maximum voxel spacing across three image axes (thin: < 1 mm; intermediate: $[1, 5)$ mm; thick: ≥ 5 mm). Modality comes from the audited object metadata and target mappings. These analyses are not adjusted for multiple comparisons.

5. Main Results

We first compare segmentation accuracy under point-only and box-initialized interaction, then examine differences across medical categories and imaging strata. The paired comparison with nnInteractive covers all 13,826 objects and 107 categories. Additional baselines provide descriptive comparisons under the evaluated configurations, with method-specific coverage of valid scores (Section 4.4 and Tables 3 and 4).

5.1. Point-Only Segmentation

SAMI3D-DW achieves the highest observed category-macro Dice among the evaluated point-only configurations (Table 3). It leads nnInteractive, clearly the strongest baseline, from the first point, scoring 0.5764 versus 0.5315, and reaches 0.7771 versus 0.7494 with five total points.

Table 3. Point-only prompt comparison. Scores are category-macro Dice; invalid object scores are excluded rather than replaced with zero. VISTA3D values are descriptive because its published interface can use target-type information that is absent from the spatial-only protocols.

Method	1 point	3 points	5 points	Valid objects	Categories
SAMI3D-DW	0.5764	0.7398	0.7771	13,826	107
nnInteractive	0.5315	0.6917	0.7494	13,826	107
MedSAM2	0.2795	0.3233	0.3656	13,715	107
SAM-Med3D	0.3408	0.3928	0.4055	13,823	107
SegVol	0.2767	0.2965	0.3128	13,785	107
VISTA3D (CVPR’25)	0.3689	0.4186	0.4300	13,747	107

5.2. Box-Initialized Segmentation and Refinement

With box initialization, SAMI3D-DW has the highest observed category-macro Dice among the five methods evaluated under this protocol (Table 4). Its advantage over nnInteractive is largest before correction and narrows as corrective points are added. At initialization, it scores 0.7130 versus 0.6530; after five corrections, the scores are 0.8002 versus 0.7868. These accuracy trajectories describe distinct prompting interfaces; the box and point budgets are not matched for user effort (Section 4.2).

Table 4. BBox-initialized point-refinement comparison. Scores are category-macro Dice computed from each method’s valid object scores. Object and category coverage is identical across the four displayed stages; invalid scores are excluded, not replaced with zero.

Method	BBox	+1	+3	+5	Valid objects	Categories
SAMI3D-DW	0.7130	0.7604	0.7853	0.8002	13,826	107
nnInteractive	0.6530	0.7154	0.7638	0.7868	13,826	107
MedSAM2	0.3766	0.3780	0.4099	0.4352	13,420	107
SAM-Med3D	0.3408	0.3796	0.4013	0.4081	13,823	107
SegVol	0.2408	0.2530	0.2582	0.2617	13,540	107

5.3. Category-Level Results

Detailed category comparisons focus on nnInteractive, for which complete paired results are available. Tables 5 and 6 list all 63 categories of anatomical structures and 43 pathology categories, with their full names, object counts, and both methods’ Dice at BBox initialization and after five corrective points. Detailed results for SAMI3D-DW at BBox, +1, +3, and +5 appear in Section A, using Combined Targets that pool specific-target variants for compact presentation.

Vascular targets and the renal collecting system. The largest final gains among anatomical structures occur in pulmonary vessels of unspecified artery/vein type, head and lower-limb vascular trees, and the renal collecting system (Table 5). For head vascular trees, SAMI3D-DW reaches 0.8228 versus 0.4455 after five corrections; the renal collecting system improves by +0.2299. Neck vascular trees also benefit, reaching 0.9576 versus 0.8805 over 50 objects. Pulmonary vessels, lower-limb vascular trees, and the renal collecting system are represented by only 10, 11, and 20 objects, respectively. The large gains in these categories are based on small samples and need confirmation in larger cohorts.

Gains across neoplastic categories. Pelvic, gastrointestinal, and bone tumors show final BBox-refinement gains of +0.2693, +0.1492, and +0.1527, respectively (Table 6).

NF1-related tumors can occur throughout the body and exhibit complex morphologies, including nodular, diffuse, and mixed forms. In this category (381 objects from 31 cases), SAMI3D-DW achieves 0.6441 Dice at BBox initialization, compared with 0.5512 for nnInteractive (+0.0929). After five corrective clicks, SAMI3D-DW reaches 0.7351 versus 0.7077, retaining an advantage of +0.0275.

Observed Relative Weaknesses. Brain favors SAMI3D-DW at five total points (+0.1187) but nnInteractive after BBox refinement (−0.1282). Jaw bones show the largest final deficit among anatomical structures with a box (−0.6104, 14 objects), despite an approximate tie in point-only mode. CNS demyelinating and immune-mediated lesions favor nnInteractive in both modes. These category comparisons are descriptive; no multiplicity-adjusted inference is made, and the means alone do not explain failure mechanisms or establish vascular branch completeness.

Table 5. Anatomical and Physiological Structures: BBox and BBox + 5 Clicks (Dice)

Code	Category	Samples (instances)	SAMI3D-DW		nnInteractive	
			BBox	+5	BBox	+5
	Mean		0.6864	0.7912	0.6322	0.7784
A01	Brain	72	0.6170	0.6767	0.7260	0.8049
	Ventricular & Cerebrospinal Fluid System	22	0.2870	0.4918	0.1857	0.3331
	Spinal Cord	89	0.6410	0.8487	0.6227	0.8381
	Cranial Nerves	41	0.4998	0.6043	0.3197	0.4840
A02	Heart	119	0.8257	0.8545	0.8011	0.8830
	Other Arterial Vessels	400	0.6981	0.8601	0.6855	0.8806
	Other Venous Vessels	61	0.6526	0.8323	0.6284	0.8505
	Pulmonary Vessels, Artery/Vein Unspecified	10	0.5637	0.8288	0.1181	0.3597
	Hepatic Vessels, Artery/Vein Unspecified	16	0.2849	0.5908	0.2455	0.5367
	Aortic Arterial Tree	29	0.8252	0.9173	0.6289	0.8397
	Head Vascular Tree	36	0.6611	0.8228	0.0747	0.4455
	Neck Vascular Tree	50	0.8657	0.9576	0.7659	0.8805
	Coronary Arterial Tree	50	0.8152	0.9293	0.8555	0.9113
	Lower-limb Vascular Tree	11	0.4800	0.7672	0.1854	0.5057
	Pulmonary Circulation Vessels	29	0.7214	0.8265	0.4141	0.8346
	Hepatic Arteries and Veins	24	0.8088	0.8518	0.7026	0.7984
	Portal Venous System	48	0.6355	0.7598	0.5167	0.7697
	Vena Caval System	156	0.7454	0.8746	0.6124	0.8911
	Brachiocephalic Veins	68	0.8186	0.8987	0.8073	0.9202
A03	Lung	387	0.9061	0.9290	0.8887	0.9465
	Respiratory Tract and Airway	76	0.6650	0.7730	0.5933	0.7629
A04	Oral Cavity & Pharynx	42	0.6106	0.7039	0.5165	0.6936
	Salivary Glands	56	0.7905	0.8497	0.7521	0.8528
	Esophagus	146	0.5984	0.8076	0.5918	0.8213
	Stomach	130	0.8050	0.8800	0.8050	0.9035
	Small Intestine	224	0.5992	0.7458	0.5480	0.7678
	Large Intestine	207	0.5588	0.7431	0.5362	0.7729
	Liver	151	0.9284	0.9335	0.9319	0.9546
	Gallbladder & Biliary System	121	0.7798	0.8453	0.7711	0.8602
A05	Pancreas	186	0.7382	0.8149	0.6492	0.8052
	Kidney	349	0.9279	0.9404	0.8829	0.9151
	Renal Collecting System	20	0.7158	0.7766	0.3857	0.5467
	Ureter	18	0.6432	0.7711	0.3397	0.6369
	Urinary Bladder	94	0.8111	0.8361	0.8062	0.8482
A06	Prostate	82	0.6867	0.7703	0.6607	0.7860
	Prostate or Uterus, Sex-conditional Composite	15	0.8346	0.8822	0.8130	0.8879
A07	Pituitary Gland	14	0.7247	0.7816	0.7075	0.7732
	Thyroid Gland	45	0.7059	0.8337	0.6523	0.8448
	Adrenal Gland	237	0.7391	0.7924	0.7308	0.8160
A08	Lymph Nodes	103	0.7035	0.7660	0.6838	0.7818
	Spleen	129	0.9271	0.9380	0.9239	0.9473
A09	Cartilage & Fibrocartilage	336	0.4218	0.6184	0.4180	0.6611
	Ligament	101	0.3348	0.4950	0.2337	0.4022
	Intervertebral Disc	182	0.8028	0.8190	0.7593	0.8173
	Skull	20	0.5389	0.7919	0.7700	0.8359
	Jaw Bones	14	0.0404	0.2028	0.6107	0.8132
	Cervical Vertebrae	117	0.6829	0.8021	0.8660	0.9114
	Thoracic Vertebrae	372	0.9114	0.9359	0.9315	0.9518
	Lumbar Vertebrae	118	0.9049	0.9273	0.9347	0.9583
	Ribs	779	0.8498	0.9121	0.8924	0.9363
	Sternum	37	0.8408	0.9247	0.6633	0.9351
	Bony Pelvis	118	0.8038	0.8763	0.7771	0.8795
	Shoulder-girdle Bones	139	0.8627	0.9074	0.8861	0.9169
	Limb Bones	335	0.8677	0.8945	0.8036	0.8937
	Spinal Column, Spinal Canal and Level-unspecified Vertebra	145	0.6140	0.7692	0.5225	0.7352
A10	Tendons & Aponeuroses	39	0.4233	0.5355	0.4104	0.5632
	Fascia & Muscular Compartments	13	0.1836	0.2153	0.2003	0.2207
	Head and Neck Muscles	13	0.6187	0.7473	0.4063	0.7293
	Abdominal Muscles	92	0.7339	0.8480	0.7227	0.8703
	Paraspinal Muscles	125	0.7365	0.8726	0.7193	0.8779
	Lower-limb Muscles	180	0.7622	0.8659	0.7892	0.8665
A12	Eye & Ocular Adnexa	84	0.7420	0.8014	0.7249	0.7813
	Ear & Auditory-Vestibular Structures	28	0.7173	0.7732	0.7192	0.7899

CT/MR cohort; taxonomy codes follow Table 2. BBox is the initial 2D box; +5 denotes five corrective points. Category scores average valid instance Dice, counting each physical object once per category. Mean weights categories equally. Missing scores are excluded. Bold marks the higher score at the same stage.

Table 6. Pathological Findings and Abnormalities: BBox and BBox + 5 Clicks (Dice)

Code	Category	Samples (instances)	SAMI3D-DW		nnInteractive	
			BBox	+5	BBox	+5
	Mean		0.7545	0.8185	0.6775	0.7954
B01	Lung Cancer or Pulmonary Mass	238	0.7213	0.7913	0.6599	0.7798
	Adrenal Tumor	51	0.8268	0.9320	0.9000	0.9294
	Kidney Cancer	39	0.8295	0.8747	0.7200	0.8454
	Kidney Tumor	126	0.8931	0.9228	0.8566	0.9162
	Gastric Cancer	206	0.7046	0.7763	0.5376	0.6940
	Gastrointestinal Tumor	151	0.6481	0.7473	0.3679	0.5981
	Gastrointestinal Stromal Tumor	31	0.8255	0.8743	0.7737	0.8535
	Thymic Cancer	70	0.7540	0.8380	0.6504	0.7475
	Other Abdominal Tumor	10	0.7861	0.8488	0.7089	0.8247
	Bone Tumor	41	0.7372	0.8377	0.5183	0.6849
	Whole-body NF1-related Tumor Burden	381	0.6441	0.7351	0.5512	0.7077
	Cervical Cancer	67	0.7505	0.7977	0.6441	0.7694
	Pelvic Tumor	64	0.7687	0.8251	0.4528	0.5558
	Liver Tumor	339	0.7050	0.8012	0.7049	0.7969
	Brain and Intracranial Tumor	318	0.7942	0.8340	0.6950	0.7801
	Parotid Tumor	33	0.8381	0.8686	0.7922	0.8662
	Soft-tissue Tumor	48	0.7896	0.8383	0.7741	0.8469
	Breast Tumor	390	0.8039	0.8493	0.7217	0.8313
	Prostate Tumor	14	0.7311	0.7847	0.6673	0.8057
	Pancreatic Tumor	94	0.8430	0.8832	0.7665	0.8732
	Other Head and Neck Tumor	167	0.7219	0.7986	0.7147	0.8362
	Metastatic Tumor	404	0.7366	0.8178	0.7354	0.8096
	Other or Site-unspecified Tumor	297	0.7316	0.8032	0.6848	0.8231
B02	Pneumonia	428	0.6880	0.7778	0.6567	0.7836
B03	Central Nervous-system Demyelinating and Immune-mediated Lesion	149	0.7420	0.7363	0.7531	0.8566
B04	Intracranial Hemorrhage	461	0.6421	0.7051	0.5823	0.6870
	Cerebral Ischemia, Infarction and Hypoxic-ischemic Injury	173	0.6256	0.7130	0.6082	0.7407
B05	Anterior Cruciate Ligament Injury	54	0.7102	0.7557	0.5101	0.6724
	Fracture	184	0.9279	0.9606	0.8935	0.9491
B07	Renal Cyst	84	0.8961	0.9258	0.8751	0.9230
B10	Indeterminate Breast Lesion	77	0.7067	0.7823	0.5649	0.7427
	Breast Nodule	76	0.8424	0.8696	0.7516	0.8507
	Thyroid Nodule	30	0.7753	0.8326	0.6685	0.8043
	Indeterminate Liver Lesion	121	0.7687	0.8432	0.7488	0.8343
	Pulmonary Nodule	405	0.8029	0.8493	0.7616	0.8387
	Indeterminate Pancreatic Lesion	12	0.7402	0.7850	0.6387	0.8164
	Indeterminate Prostate Lesion	91	0.6363	0.7476	0.6590	0.7724
	Indeterminate Ovarian Lesion	87	0.8264	0.8669	0.7922	0.8864
	Perirectal Lymph-node Target of Indeterminate Status	15	0.8573	0.8806	0.5408	0.7826
	Indeterminate Nervous-system Imaging Abnormality	20	0.7854	0.8550	0.7005	0.7963
	Other Indeterminate Abdominopelvic Lesion	145	0.7810	0.8376	0.7351	0.8303
	Indeterminate Lymph-node Target	188	0.6914	0.7643	0.6817	0.7838
	Pathologic Fluid Collection and Abnormal Gas in Body Cavity	15	0.4129	0.6253	0.4126	0.6734

CT/MR cohort; taxonomy codes follow Table 2. BBox is the initial 2D box; +5 denotes five corrective points. Category scores average valid instance Dice, counting each physical object once per category. Mean weights categories equally. Missing scores are excluded. Bold marks the higher score at the same stage.

5.4. Stratified Results

Imaging modality. SAMI3D-DW has the highest observed BBox-initialized category-macro score in both evaluated modalities. After five corrections, it remains highest on CT (0.8300 versus 0.8195 for nnInteractive) and MR (0.7455 versus 0.7433) (Table 7). CT and MR contribute 85 and 69 represented categories, respectively; these sets overlap and differ in target composition.

Table 7. Category-macro Dice by 3D imaging modality. Each method cell reports BBox / +5 corrections. Categories denotes the common denominator of categories with valid scores for all five methods; bold marks the best value at each stage within a row. The cohort is restricted to CT and MR.

Modality	Categories (used/mapped)	Objects	SAMI3D-DW	nnInteractive	MedSAM2	SAM-Med3D	SegVol
CT	85/85	9,289	0.7447 / 0.8300	0.6872 / 0.8195	0.3768 / 0.4155	0.3584 / 0.4374	0.2291 / 0.2500
MR	69/69	4,537	0.6405 / 0.7455	0.5997 / 0.7433	0.3384 / 0.4080	0.3378 / 0.4162	0.2382 / 0.2741

Object size and voxel spacing. SAMI3D-DW has the highest initial and final category-macro Dice in each reported size and maximum-spacing stratum. The final margins over nnInteractive are small: +0.0034 to +0.0097 across size strata and +0.0059 to +0.0178 across spacing strata. Full five-method results and category denominators appear in Tables 8 and 9. These exploratory comparisons use common category denominators based on valid scores within each stratum, but do not control for differences in target composition or support causal claims about size, spacing, or modality.

Table 8. Category-macro Dice by reference-object size. Each method cell reports BBox / +5 corrections. Categories denotes the common denominator of categories with valid scores for all five methods; bold marks the best value at each stage within a row.

Size by target voxels	Categories (used/mapped)	Objects	SAMI3D-DW	nnInteractive	MedSAM2	SAM-Med3D	SegVol
Small (< 10,000)	97/97	7,855	0.6764 / 0.7654	0.6141 / 0.7558	0.3649 / 0.4735	0.2972 / 0.3256	0.3429 / 0.3585
Medium (10,000–99,999)	90/90	4,009	0.7443 / 0.8268	0.7089 / 0.8230	0.4136 / 0.4282	0.4400 / 0.5255	0.1650 / 0.2132
Large (≥ 100,000)	68/69	1,962	0.7523 / 0.8392	0.7237 / 0.8358	0.4782 / 0.4181	0.4711 / 0.6068	0.0555 / 0.1128

Table 9. Category-macro Dice by maximum voxel spacing. Each method cell reports BBox / +5 corrections. Categories denotes the common denominator of categories with valid scores for all five methods; bold marks the best value at each stage within a row.

Maximum spacing	Categories (used/mapped)	Objects	SAMI3D-DW	nnInteractive	MedSAM2	SAM-Med3D	SegVol
Thin (< 1 mm)	46/47	981	0.7826 / 0.8403	0.7282 / 0.8343	0.3616 / 0.3402	0.3810 / 0.4828	0.1127 / 0.1064
Intermediate (1–< 5 mm)	104/104	9,657	0.7169 / 0.8028	0.6518 / 0.7850	0.3635 / 0.4149	0.3367 / 0.4072	0.2207 / 0.2375
Thick (≥ 5 mm)	70/70	3,188	0.6801 / 0.7762	0.6334 / 0.7685	0.4418 / 0.5389	0.3860 / 0.4367	0.3418 / 0.3702

6. Limitations and Responsible Use

Evaluation scope. Training and test cases are disjoint, but some source datasets contribute to both splits; transfer to unseen datasets and institutions remains untested. The current evaluation covers CT and MR only. Future work will include PET, 3D ultrasound, and independent external validation.

Simulated interaction. Reference masks determine the targets and prompts, so this benchmark does not measure human prompt variability or annotation time. The preliminary NF1 timing comparison is task-specific and requires fuller reporting. Predicted masks and subsequent corrections require qualified human review.

Baseline comparability. Coverage of valid scores differs across baselines. VISTA3D can also use target-type information. Their comparisons are descriptive and specific to the evaluated configurations. Category averages do not establish uniform superiority across targets or clinical effectiveness.

7. Conclusion

We present SAMI3D-DW and evaluate interactive 3D segmentation on a CT/MR benchmark of 219 source datasets, 4,326 cases, and 107 medical categories. With equal category weighting, SAMI3D-DW achieves the highest observed mean Dice among the evaluated configurations: 0.5764 with one point, 0.7771 with five points, 0.7130 with BBox initialization, and 0.8002 after five corrective clicks. These results support point-only interaction as a simple interface and BBox initialization when approximate target extent is available. Independent external validation, real-user studies, and evaluation on additional imaging modalities remain necessary to establish broader applicability and workflow benefits.

Acknowledgments

We would like to thank Professor Zhichao Wang and Dr. Jun Liu, from Shanghai Jiao Tong University School of Medicine and its affiliated Shanghai Ninth People’s Hospital, for their valuable feedback during the early development of SAMI3D-DW.

References

- A. Archi, L. Freckmann, S. Nair, N. Khalid, P. Hilt, V. Rajashekar, M. Freitag, C. Teuber, M. Spitzner, C. Tapia Contreras, G. Buckley, S. von Haaren, S. Gupta, M. Grade, M. Wirth, G. Schneider, A. Dengel, S. Ahmed, and C. Pape. Segment anything for microscopy. *Nature Methods*, 22(3):579–591, 2025. doi: 10.1038/s41592-024-02580-4.
- N. Carion, L. Gustafson, Y.-T. Hu, S. Debnath, R. Hu, D. Suris, C. Ryali, K. V. Alwala, H. Khedr, et al. SAM 3: Segment anything with concepts, 2025. URL <https://arxiv.org/abs/2511.16719>.
- C. Chen, J. Miao, D. Wu, A. Zhong, Z. Yan, S. Kim, J. Hu, Z. Liu, L. Sun, X. Li, T. Liu, P.-A. Heng, and Q. Li. MA-SAM: Modality-agnostic SAM adaptation for 3D medical image segmentation. *Medical Image Analysis*, 98:103310, 2024. doi: 10.1016/j.media.2024.103310.
- Z. Chen, J. Hou, L. Lin, Y. Wang, Y. Bie, X. Wang, Y. Zhou, R. C. K. Chan, and H. Chen. Segment anything in pathology images with natural language. arXiv preprint arXiv:2506.20988v2, 2025. URL <https://arxiv.org/abs/2506.20988v2>.
- J. Cheng, J. Ye, Z. Deng, J. Chen, T. Li, H. Wang, Y. Su, Z. Huang, J. Chen, L. Jiang, J. He, S. Zhang, M. Zhu, and Y. Qiao. SAM-Med2D, 2023. URL <https://arxiv.org/abs/2308.16184>.
- Y. Du, F. Bai, T. Huang, and B. Zhao. SegVol: Universal and interactive volumetric medical image segmentation. In *Advances in Neural Information Processing Systems*, volume 37, pages 110746–110783, 2024.
- S. Gong, Y. Zhong, W. Ma, J. Li, Z. Wang, J. Zhang, P.-A. Heng, and Q. Dou. 3DSAM-adapter: Holistic adaptation of SAM from 2D to 3D for promptable tumor segmentation. *Medical Image Analysis*, 98:103324, 2024. doi: 10.1016/j.media.2024.103324.
- Y. He, P. Guo, Y. Tang, A. Myronenko, V. Nath, Z. Xu, D. Yang, C. Zhao, B. Simon, M. Belue, S. Harmon, B. Turkbey, D. Xu, and W. Li. VISTA3D: A unified segmentation foundation model for 3D medical imaging. In *Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition*, pages 20863–20873, 2025. doi: 10.1109/CVPR52734.2025.01943. URL <https://arxiv.org/abs/2406.05285>.
- Y. Huang, X. Yang, L. Liu, H. Zhou, A. Chang, X. Zhou, R. Chen, J. Yu, J. Chen, C. Chen, et al. Segment anything model for medical images? *Medical Image Analysis*, 92:103061, 2024. doi: 10.1016/j.media.2023.103061.
- F. Isensee, P. F. Jaeger, S. A. A. Kohl, J. Petersen, and K. H. Maier-Hein. nnU-Net: A self-configuring method for deep learning-based biomedical image segmentation. *Nature Methods*, 18(2):203–211, 2021. doi: 10.1038/s41592-020-01008-z.
- F. Isensee, M. Rokuss, L. Krämer, S. Dinkelacker, A. Ravindran, F. Stritzke, B. Hamm, T. Wald, M. Langenberg, C. Ulrich, J. Deissler, R. Floca, and K. Maier-Hein. nnInteractive: Redefining 3D promptable segmentation, 2025. URL <https://arxiv.org/abs/2503.08373>.
- A. Kirillov, E. Mintun, N. Ravi, H. Mao, C. Rolland, L. Gustafson, T. Xiao, S. Whitehead, A. C. Berg, W.-Y. Lo, P. Dollár, and R. Girshick. Segment anything. In *Proceedings of the IEEE/CVF International Conference on Computer Vision*, pages 4015–4026, 2023. doi: 10.1109/ICCV51070.2023.00371.
- T. Koleilat, H. Asgariandehkordi, H. Rivaz, and Y. Xiao. MedCLIP-SAMv2: Towards universal text-driven medical image segmentation. *Medical Image Analysis*, 106:103749, 2025. doi: 10.1016/j.media.2025.103749.

- W. Lei, W. Xu, K. Li, X. Zhang, and S. Zhang. MedLSAM: Localize and segment anything model for 3D CT images. *Medical Image Analysis*, 99:103370, 2025. doi: 10.1016/j.media.2024.103370.
- X. Luo, G. Wang, T. Song, J. Zhang, M. Aertsen, J. Deprest, S. Ourselin, T. Vercauteren, and S. Zhang. MIDeepSeg: Minimally interactive segmentation of unseen objects from medical images using deep learning. *Medical Image Analysis*, 72:102102, 2021. doi: 10.1016/j.media.2021.102102.
- J. Ma, Y. He, F. Li, L. Han, C. You, and B. Wang. Segment anything in medical images. *Nature Communications*, 15(1):654, 2024. doi: 10.1038/s41467-024-44824-z.
- J. Ma, Z. Yang, S. Kim, B. Chen, M. Baharoon, A. Fallahpour, R. Asakereh, H. Lyu, and B. Wang. MedSAM2: Segment anything in 3D medical images and videos, 2025. URL <https://arxiv.org/abs/2504.03600>.
- M. Marks, U. Israel, R. Dilip, Q. Li, C. Yu, E. Laubscher, A. Iqbal, E. Pradhan, A. Ates, M. Abt, C. Brown, E. Pao, S. Li, A. Pearson-Goulart, P. Perona, G. Gkioxari, R. Barnowski, Y. Yue, and D. Van Valen. CellSAM: A foundation model for cell segmentation. *Nature Methods*, 22(12):2585–2593, 2025. doi: 10.1038/s41592-025-02879-w.
- M. A. Mazurowski, H. Dong, H. Gu, J. Yang, N. Konz, and Y. Zhang. Segment anything model for medical image analysis: An experimental study. *Medical Image Analysis*, 89:102918, 2023. doi: 10.1016/j.media.2023.102918.
- H. Peiris, S. Wang, G. Egan, M. Harandi, M. Law, and Z. Chen. Segment any tumour: An uncertainty-aware vision foundation model for whole-body analysis. *Nature Communications*, 17(1):9640, 2026. doi: 10.1038/s41467-026-76531-2.
- N. Ravi, V. Gabeur, Y.-T. Hu, R. Hu, C. Ryali, T. Ma, H. Khedr, R. Rädle, C. Rolland, L. Gustafson, E. Mintun, J. Pan, K. V. Alwala, N. Carion, C.-Y. Wu, R. Girshick, P. Dollár, and C. Feichtenhofer. SAM 2: Segment anything in images and videos. In *International Conference on Learning Representations*, 2025. URL <https://arxiv.org/abs/2408.00714>.
- M. Rokuss, M. Langenberg, Y. Kirchhoff, F. Isensee, B. Hamm, C. Ulrich, S. Regnery, L. Bauer, E. Katsigiannopoulos, T. Norajitra, and K. Maier-Hein. VoxTell: Free-text promptable universal 3D medical image segmentation. arXiv preprint arXiv:2511.11450v1, 2025. URL <https://arxiv.org/abs/2511.11450v1>.
- O. Ronneberger, P. Fischer, and T. Brox. U-Net: Convolutional networks for biomedical image segmentation. In *Medical Image Computing and Computer-Assisted Intervention – MICCAI 2015*, volume 9351 of *Lecture Notes in Computer Science*, pages 234–241. Springer, 2015. doi: 10.1007/978-3-319-24574-4_28.
- C. Ulrich, T. Wald, E. Tempus, M. Rokuss, P. F. Jaeger, and K. Maier-Hein. RadioActive: 3D radiological interactive segmentation benchmark. arXiv preprint arXiv:2411.07885v3, 2025. URL <https://arxiv.org/abs/2411.07885v3>.
- G. Wang, M. A. Zuluaga, W. Li, R. Pratt, P. A. Patel, M. Aertsen, T. Doel, A. L. David, J. Deprest, S. Ourselin, and T. Vercauteren. DeepIGeoS: A deep interactive geodesic framework for medical image segmentation. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 41(7):1559–1572, 2019. doi: 10.1109/TPAMI.2018.2840695.
- H. Wang, S. Guo, J. Ye, Z. Deng, J. Cheng, T. Li, J. Chen, Y. Su, Z. Huang, Y. Shen, B. Fu, S. Zhang, and J. He. SAM-Med3D: A vision foundation model for general-purpose segmentation on volumetric medical images. *IEEE Transactions on Neural Networks and Learning*

- Systems*, 36(10):17599–17612, 2025. doi: 10.1109/TNNLS.2025.3586694. URL <https://arxiv.org/abs/2310.15161>.
- H. E. Wong, M. Rakic, J. Guttag, and A. V. Dalca. ScribblePrompt: Fast and flexible interactive segmentation for any biomedical image. In *Computer Vision – ECCV 2024*, pages 207–229. Springer, 2024. doi: 10.1007/978-3-031-73661-2_12. URL <https://arxiv.org/abs/2312.07381>.
- J. Wu, Z. Wang, M. Hong, W. Ji, H. Fu, Y. Xu, M. Xu, and Y. Jin. Medical SAM adapter: Adapting segment anything model for medical image segmentation. *Medical Image Analysis*, 102:103547, 2025. doi: 10.1016/j.media.2025.103547.
- T. Zhao, Y. Gu, J. Yang, N. Usuyama, H. H. Lee, S. Kiblawi, T. Naumann, J. Gao, A. Crabtree, J. Abel, C. Moungh-Wen, B. Piening, C. Bifulco, M. Wei, H. Poon, and S. Wang. A foundation model for joint segmentation, detection and recognition of biomedical objects across nine modalities. *Nature Methods*, 22(1):166–176, 2025a. doi: 10.1038/s41592-024-02499-w.
- Z. Zhao, Y. Zhang, C. Wu, X. Zhang, X. Zhou, Y. Zhang, Y. Wang, and W. Xie. Large-vocabulary segmentation for medical images with text prompts. *npj Digital Medicine*, 8(1):566, 2025b. doi: 10.1038/s41746-025-01964-w. URL <https://arxiv.org/abs/2312.17183>.
- T. Zhou, L. Li, G. Bredell, J. Li, J. Unkelbach, and E. Konukoglu. Volumetric memory network for interactive medical image segmentation. *Medical Image Analysis*, 83:102599, 2023. doi: 10.1016/j.media.2022.102599.

A. Detailed Medical Target Evaluation

The benchmark distinguishes 405 *Specific Targets* by modality, laterality, anatomical or pathological subdivisions, and verified imaging sequences or phases where available. These refine the underlying medical objects. To avoid listing every variant, this appendix reports *Combined Targets*, pooling modality, laterality, and finer distinctions within each listed medical object. The examples below illustrate the refinement; knee laterality and medial/lateral meniscal position are recorded separately.

Combined Target	Specific Target example	Added distinctions
Rib	CT–left–first rib	Modality, laterality, rib number
Rib	CT–right–twelfth rib	Modality, laterality, rib number
Meniscus	MR–left knee–medial meniscus	Knee laterality and medial/lateral position
Meniscus	MR–right knee–lateral meniscus	Knee laterality and medial/lateral position

Table 10. Anatomical and Physiological Targets: BBox + Point Segmentation (Dice)

Taxonomy	Category	Combined Targets	Samples (instances)	SAMI3D-DW			
				BBox	+1	+3	+5
Mean				0.6933	0.7408	0.7671	0.7888
A01 Nervous System	Brain	Brain	14	0.9121	0.9063	0.9174	0.9279
		Brainstem	14	0.7552	0.8059	0.8275	0.8416
	Ventricular & Cerebrospinal Fluid System	Cerebral Gray Matter	22	0.4740	0.6228	0.4513	0.4913
		Cerebral White Matter	22	0.4841	0.3970	0.5267	0.5973
		Cerebrospinal Fluid	22	0.2870	0.3118	0.4743	0.4918
		Spinal Cord	89	0.6410	0.7856	0.8316	0.8487
	Cranial Nerves	Optic Nerve	28	0.5084	0.5770	0.6204	0.6418
		Optic Chiasm	13	0.4811	0.4886	0.5142	0.5236
A02 Cardiovascular System	Heart	Heart	51	0.8642	0.8830	0.8888	0.8972
		Left Atrium	10	0.9302	0.9364	0.9400	0.9435
		Left Ventricular Cavity	10	0.8859	0.9104	0.9158	0.9195
		Right Ventricular Cavity	10	0.8073	0.8399	0.8586	0.8623
		Myocardium	10	0.4342	0.4637	0.4329	0.5267
		Left Atrial Appendage	28	0.8429	0.8288	0.8234	0.8360
	Other Arterial Vessels	Aorta	140	0.8141	0.8668	0.8887	0.9005
		Common Carotid Artery	94	0.5746	0.7321	0.8041	0.8346
		Subclavian Artery	69	0.6631	0.8024	0.8515	0.8678
		Brachiocephalic Trunk	34	0.8072	0.8332	0.8714	0.8836
		Iliac Artery	63	0.6043	0.6830	0.7329	0.7874
	Other Venous Vessels	Iliac Vein	61	0.6526	0.7595	0.8098	0.8323
	Pulmonary Vessels, Artery/Vein Unspecified	Pulmonary Vessels, Artery/Vein Unspecified	10	0.5637	0.7347	0.8170	0.8288
	Hepatic Vessels, Artery/Vein Unspecified	Hepatic Vessels, Artery/Vein Unspecified	16	0.2849	0.4818	0.5621	0.5908
	Aortic Arterial Tree	Aortic Vessel Tree	29	0.8252	0.8968	0.9189	0.9173
	Head Vascular Tree	Head Vascular Tree, Artery/Vein Unspecified	36	0.6611	0.7902	0.8201	0.8228
		Neck Vascular Tree	50	0.8657	0.9535	0.9600	0.9576
	Coronary Arterial Tree	Coronary Arterial Tree	50	0.8152	0.8500	0.9050	0.9293

Continued on next page

Table 10 (continued)

Taxonomy	Category	Combined Targets	Samples (instances)	SAMI3D-DW			
				BBox	+1	+3	+5
A03 Respiratory System	Lower-limb Vascular Tree	Lower-limb Vascular Tree, Artery/Vein Unspecified	11	0.4800	0.5886	0.6836	0.7672
	Pulmonary Circulation Vessels	Pulmonary Vein	29	0.7214	0.8061	0.8178	0.8265
	Hepatic Arteries and Veins	Hepatic Artery	12	0.8549	0.8739	0.8788	0.8857
		Hepatic Vein	12	0.7627	0.7788	0.7946	0.8180
	Portal Venous System	Portal Vein	12	0.7301	0.7643	0.7342	0.7754
		Portal and Splenic Vein Union	36	0.6040	0.7125	0.7358	0.7547
	Vena Caval System	Superior Vena Cava	32	0.8622	0.9031	0.9167	0.9196
		Inferior Vena Cava	124	0.7153	0.8264	0.8471	0.8630
	Brachiocephalic Veins	Brachiocephalic Vein	68	0.8186	0.8641	0.8880	0.8987
	Lung	Lung	204	0.9079	0.9312	0.9319	0.9344
A04 Digestive System	Respiratory Tract and Airway	Lung Lobe	183	0.9041	0.9140	0.9229	0.9231
		Trachea	36	0.8526	0.9104	0.9206	0.9291
	Oral Cavity & Pharynx	Glottis	13	0.4361	0.4476	0.4783	0.5471
		Supraglottic Larynx	14	0.6307	0.6588	0.7115	0.7136
		Arytenoid Cartilage	13	0.4113	0.5602	0.6070	0.6304
	Salivary Glands	Oral Cavity	14	0.8531	0.8729	0.8824	0.8836
		Buccal Mucosa	14	0.4474	0.5591	0.5914	0.5915
		Lips	14	0.5314	0.5899	0.6288	0.6365
	Esophagus	Parotid Gland	28	0.7945	0.8080	0.8193	0.8339
		Submandibular Gland	28	0.7866	0.8364	0.8557	0.8655
A05 Urinary System	Stomach	Esophagus	146	0.5984	0.7423	0.7922	0.8076
	Small Intestine	Stomach	130	0.8050	0.8608	0.8803	0.8800
	Large Intestine	Duodenum	122	0.6030	0.7138	0.7626	0.7816
		Small Intestine	102	0.5947	0.6611	0.6955	0.7029
	Liver	Appendix	5	0.3531	0.4432	0.4606	0.4813
		Cecum	10	0.6300	0.6736	0.7097	0.7218
		Colon	97	0.5538	0.6681	0.7467	0.7484
		Ascending Colon	10	0.7523	0.8036	0.7881	0.7940
		Transverse Colon	10	0.3130	0.3730	0.4901	0.5753
		Descending Colon	10	0.5422	0.6765	0.7365	0.7614
		Sigmoid Colon	10	0.2989	0.4344	0.6082	0.6523
	Gallbladder & Biliary System	Rectum	55	0.6331	0.7429	0.7782	0.7958
		Liver	151	0.9284	0.9348	0.9416	0.9335
	Pancreas	Gallbladder	113	0.7993	0.8296	0.8478	0.8589
A06 Reproductive System	Kidney	Bile Duct	8	0.5042	0.5942	0.6218	0.6536
		Pancreas	186	0.7382	0.7819	0.8048	0.8149
	Renal Collecting System	Kidney	309	0.9341	0.9427	0.9484	0.9497
		Renal Cortex	20	0.9060	0.9036	0.8249	0.8766
	Ureter	Renal Medulla	20	0.8542	0.8592	0.8607	0.8615
		Renal Pelvis and Calyces	20	0.7158	0.7426	0.7620	0.7766
	Urinary Bladder	Ureter	18	0.6432	0.7004	0.7444	0.7711
		Urinary Bladder	94	0.8111	0.8149	0.8308	0.8361
	Prostate	Prostate	63	0.7040	0.7413	0.7692	0.7892
		Prostate Peripheral Zone	10	0.4955	0.5394	0.5777	0.5984
		Prostate Transition Zone	9	0.7774	0.7954	0.8029	0.8284

Continued on next page

Table 10 (continued)

Taxonomy	Category	Combined Targets	Samples (instances)	SAMI3D-DW			
				BBox	+1	+3	+5
A07 Endocrine System	Prostate or Uterus, Sex-conditional Composite	Prostate or Uterus, Sex-conditional Label	15	0.8346	0.8787	0.8742	0.8822
	Pituitary Gland	Pituitary Gland	14	0.7247	0.7701	0.7813	0.7816
	Thyroid Gland	Thyroid Gland	45	0.7059	0.7992	0.8291	0.8337
	Adrenal Gland	Adrenal Gland	237	0.7391	0.7711	0.7877	0.7924
A08 Lymphatic, Immune & Hematopoietic System	Lymph Nodes	Lymph Node	103	0.7035	0.7369	0.7638	0.7660
A09 Skeletal & Articular System	Spleen	Spleen	129	0.9271	0.9256	0.9326	0.9380
	Cartilage & Fibrocartilage	Costal Cartilage	39	0.4945	0.5977	0.6483	0.6735
		Meniscus	102	0.4237	0.5862	0.6742	0.6862
		Patellar Cartilage	51	0.6030	0.7396	0.7875	0.7885
	Ligament	Femoral Articular Cartilage	63	0.1429	0.2351	0.2685	0.3003
		Tibial Articular Cartilage	81	0.4871	0.5973	0.6218	0.6470
		Anterior Cruciate Ligament	20	0.5456	0.6292	0.6666	0.6881
		Posterior Cruciate Ligament	20	0.3739	0.4596	0.5032	0.5484
		Patellar Ligament	7	0.2775	0.2355	0.3582	0.3847
		Knee Collateral Ligament	26	0.2223	0.2462	0.2586	0.2756
		Patellar Retinaculum	28	0.2751	0.3169	0.4211	0.5502
	Intervertebral Disc	Intervertebral Disc	182	0.8028	0.8109	0.8160	0.8190
	Skull	Skull	20	0.5389	0.5824	0.6369	0.7919
	Jaw Bones	Mandible	14	0.0404	0.1079	0.1252	0.2028
	Cervical Vertebrae	C1 Vertebra	11	0.7951	0.8458	0.8647	0.8838
		C2 Vertebra	12	0.7914	0.7541	0.7659	0.8219
		C3 Vertebra	10	0.5887	0.4703	0.4697	0.7186
		C4 Vertebra	12	0.5029	0.4822	0.5243	0.7901
		C5 Vertebra	15	0.5221	0.5466	0.6156	0.6658
		C6 Vertebra	25	0.6277	0.6740	0.7239	0.7615
		C7 Vertebra	32	0.8189	0.8368	0.8919	0.8929
	Thoracic Vertebrae	T1 Vertebra	32	0.9318	0.9401	0.9438	0.9464
		T2 Vertebra	33	0.8969	0.8807	0.9117	0.9282
		T3 Vertebra	34	0.8644	0.8615	0.8779	0.8991
		T4 Vertebra	33	0.8736	0.9118	0.9016	0.9161
		T5 Vertebra	32	0.8462	0.8462	0.8714	0.8841
		T6 Vertebra	29	0.9073	0.9144	0.9132	0.9209
		T7 Vertebra	27	0.9402	0.9341	0.9507	0.9551
		T8 Vertebra	28	0.9334	0.9533	0.9577	0.9580
		T9 Vertebra	30	0.9414	0.9593	0.9626	0.9635
		T10 Vertebra	31	0.9401	0.9562	0.9582	0.9612
		T11 Vertebra	32	0.9350	0.9498	0.9521	0.9524
		T12 Vertebra	31	0.9419	0.9477	0.9561	0.9568
	Lumbar Vertebrae	L1 Vertebra	30	0.9582	0.9216	0.9483	0.9457
		L2 Vertebra	26	0.8810	0.9235	0.9285	0.9264
		L3 Vertebra	21	0.9059	0.9642	0.9652	0.9676
		L4 Vertebra	20	0.9368	0.8921	0.8766	0.9152
		L5 Vertebra	21	0.8268	0.7900	0.8565	0.8730
	Ribs	Rib	779	0.8498	0.8984	0.9049	0.9121
	Sternum	Sternum	37	0.8408	0.8954	0.9189	0.9247
	Bony Pelvis	Sacrum	33	0.8452	0.8515	0.8711	0.8736
		Hip Bone	65	0.7626	0.8065	0.8460	0.8712
		S1 Vertebra	20	0.8694	0.8847	0.8934	0.8972
	Shoulder-girdle Bones	Scapula	68	0.8145	0.8632	0.8729	0.8782
		Clavicle	71	0.9088	0.9248	0.9300	0.9353

Continued on next page

Table 10 (continued)

Taxonomy	Category	Combined Targets	Samples (instances)	SAMI3D-DW			
				BBox	+1	+3	+5
A10 Muscular System	Limb Bones	Humerus	59	0.8848	0.9012	0.9146	0.9171
		Femur	157	0.8866	0.8988	0.9044	0.9074
		Femoral Head	21	0.8736	0.8688	0.9051	0.9159
		Tibia	34	0.8763	0.8708	0.8676	0.8633
		Fibula	34	0.7726	0.8137	0.8210	0.8404
		Patella	30	0.8289	0.8452	0.8547	0.8646
	Spinal Column, Spinal Canal and Level-unspecified Vertebra	Vertebra, Level Unspecified	122	0.5892	0.7020	0.7386	0.7496
		Vertebral Column	23	0.7456	0.8518	0.8666	0.8730
	Tendons & Aponeuroses	Quadriceps Tendon	7	0.5419	0.5582	0.6181	0.6656
		Biceps Femoris Tendon	13	0.2150	0.2541	0.2785	0.3011
		Popliteus Tendon	19	0.5221	0.5709	0.5928	0.6479
		Iliotibial Band	13	0.1836	0.1945	0.2058	0.2153
	Fascia & Muscular Compartments						
	Head and Neck Muscles	Cricopharyngeus Muscle	13	0.6187	0.6939	0.7283	0.7473
	Abdominal Muscles	Iliopsoas	92	0.7339	0.8188	0.8376	0.8480
	Paraspinal Muscles	Intrinsic Back Muscle	125	0.7365	0.8286	0.8586	0.8726
	Lower-limb Muscles	Gluteus Maximus	59	0.8458	0.8706	0.8817	0.8897
		Gluteus Medius	62	0.6933	0.8489	0.8625	0.8803
		Gluteus Minimus	59	0.7510	0.7859	0.8194	0.8269
A12 Sensory System	Eye & Ocular Adnexa	Anterior Eye Segment	28	0.7485	0.7989	0.8169	0.8243
		Posterior Eye Segment	28	0.8789	0.8739	0.8763	0.8763
		Lacrimal Gland	28	0.5985	0.6945	0.6951	0.7035
	Ear & Auditory-Vestibular Structures	Cochlea	28	0.7173	0.7641	0.7706	0.7732

BBox denotes the initial tight 2D box on the axial slice with the largest target area. +1/3/5 denote 1/3/5 cumulative point refinements after the initial box. The cohort contains CT and MR only. Each row averages valid instance Dice within one Combined Target; a physical object can contribute to multiple Combined Target rows. Mean is the unweighted average of Combined Target means with valid scores.

Table 11. Pathological and Abnormal Targets: BBox + Point Segmentation (Dice)

Taxonomy	Category	Combined Targets	Samples (instances)	SAMI3D-DW				
				BBox	+1	+3	+5	
Mean				0.7483	0.7846	0.8043	0.8132	
B01 Neoplastic Lesion	Lung Cancer or Pulmonary Mass	Lung Cancer Lesion	156	0.7978	0.8296	0.8477	0.8579	
		Primary Non-small-cell Lung Cancer	82	0.5757	0.6196	0.6643	0.6646	
	Adrenal Tumor	Adrenal Tumor	51	0.8268	0.9070	0.9275	0.9320	
	Kidney Cancer	Kidney Cancer Lesion	39	0.8295	0.8463	0.8747	0.8747	
	Kidney Tumor	Kidney Tumor	126	0.8931	0.9094	0.9218	0.9228	
	Gastric Cancer	Gastric Cancer Lesion	206	0.7046	0.7422	0.7636	0.7763	
	Gastrointestinal Tumor	Colorectal Cancer Lesion	151	0.6481	0.7004	0.7338	0.7473	
	Gastrointestinal Stromal Tumor	Gastrointestinal Stromal Tumor	31	0.8255	0.8479	0.8653	0.8743	
	Thymic Cancer	Thymic Tumor	70	0.7540	0.7837	0.8197	0.8380	
	Other Abdominal Tumor	Abdominal Tumor, Site Unspecified	10	0.7861	0.7836	0.8251	0.8488	
	Bone Tumor	Osteosarcoma	7	0.7620	0.8199	0.8330	0.8554	
		Bone Tumor, Type Unspecified	34	0.7321	0.7694	0.7932	0.8340	
	Whole-body NF1-related Tumor Burden	Neurofibroma	381	0.6441	0.6949	0.7213	0.7351	
	Cervical Cancer	Cervical Cancer Lesion	67	0.7505	0.7737	0.7859	0.7977	
	Pelvic Tumor	Pelvic Tumor, Site Unspecified	64	0.7687	0.7834	0.8057	0.8251	
	Liver Tumor	Liver Tumor	61	0.7657	0.8228	0.8341	0.8479	
		Liver Cancer Lesion	183	0.6963	0.7313	0.7545	0.7702	
		Hepatocellular Carcinoma Lesion	95	0.6827	0.7923	0.8263	0.8310	
	Brain and Intracranial Tumor	Meningioma	4	0.8394	0.9101	0.8987	0.9131	
		Glioma	68	0.6727	0.7063	0.7334	0.7524	
		Pituitary Tumor	10	0.8247	0.8522	0.8647	0.8689	
		Brain Tumor, Type Unspecified	207	0.8282	0.8451	0.8542	0.8558	
		Medulloblastoma	24	0.8470	0.8674	0.8653	0.8711	
		Brain Tumor with Edema Composite	5	0.6897	0.7410	0.7535	0.7265	
		Parotid Tumor	Parotid Tumor	33	0.8381	0.8706	0.8670	0.8686
	Soft-tissue Tumor	Soft-tissue Sarcoma	41	0.8042	0.8235	0.8429	0.8485	
		Soft-tissue Tumor, Type Unspecified	7	0.7040	0.7531	0.7817	0.7789	
	Breast Tumor	Breast Tumor	257	0.7935	0.8126	0.8341	0.8442	
		Breast Cancer Lesion	133	0.8239	0.8461	0.8573	0.8592	
	Prostate Tumor	Prostate Cancer Lesion	14	0.7311	0.7755	0.7854	0.7847	
	Pancreatic Tumor	Pancreatic Cancer Lesion	30	0.8270	0.8483	0.8658	0.8708	
		Pancreatic Tumor	64	0.8504	0.8761	0.8844	0.8890	
	Other Head and Neck Tumor	Head and Neck Primary Tumor	150	0.7244	0.7648	0.7859	0.8014	
		Tongue Cancer Lesion	17	0.6995	0.7264	0.7742	0.7742	
	Metastatic Tumor	Metastatic Lymph-node Tumor	26	0.8189	0.8635	0.8805	0.8823	
		Liver Metastasis	378	0.7309	0.7811	0.7976	0.8134	
		Tumor Lesion, Site Unspecified	297	0.7316	0.7618	0.7888	0.8032	
	B02 Infectious Lesion	Pneumonia	COVID-19 Pulmonary Infection Lesion	342	0.6834	0.7440	0.7611	0.7692
			Pneumonia Lesion	86	0.7063	0.7593	0.8043	0.8120

Continued on next page

Table 11 (continued)

Taxonomy	Category	Combined Targets	Samples (instances)	SAMI3D-DW			
				BBox	+1	+3	+5
B03 Non-infectious Inflammatory & Immune- mediated Lesion	Central Nervous-system Demyelinating and Immune-mediated Lesion	Multiple Sclerosis Lesion	149	0.7420	0.7095	0.7210	0.7363
B04 Vascular & Circulatory Disorder	Intracranial Hemorrhage	Intracranial Hemorrhage	51	0.6524	0.6583	0.6730	0.6697
		Intraparenchymal Hemorrhage	73	0.7457	0.7744	0.7796	0.7842
		Intraventricular Hemorrhage	58	0.6453	0.6931	0.7299	0.7439
		Epidural Hemorrhage	8	0.5720	0.6750	0.7085	0.7403
		Subdural Hemorrhage	36	0.4734	0.4976	0.4973	0.5028
		Subarachnoid Hemorrhage	235	0.6351	0.6802	0.7016	0.7084
	Cerebral Ischemia, Infarction and Hypoxic-ischemic Injury	Cerebral Infarction Lesion	173	0.6256	0.6737	0.6985	0.7130
	Anterior Cruciate Ligament Injury	Anterior Cruciate Ligament Injury	54	0.7102	0.7351	0.7521	0.7557
	Fracture	Pelvic Fracture Fragment	184	0.9279	0.9295	0.9556	0.9606
B07 Non-neoplastic Focal Lesion	Renal Cyst	Renal Cyst	84	0.8961	0.9131	0.9223	0.9258
B10 Indeterminate Imaging Abnormality	Indeterminate Breast Lesion	Indeterminate Breast Lesion	77	0.7067	0.7494	0.7748	0.7823
	Breast Nodule	Breast Nodule	76	0.8424	0.8565	0.8666	0.8696
	Thyroid Nodule	Indeterminate Thyroid Lesion	23	0.7831	0.8104	0.8244	0.8296
		Thyroid Nodule	7	0.7500	0.8012	0.8370	0.8428
	Indeterminate Liver Lesion	Indeterminate Liver Lesion	121	0.7687	0.8158	0.8320	0.8432
	Pulmonary Nodule	Pulmonary Ground-glass Nodule	51	0.8133	0.8450	0.8591	0.8626
		Pulmonary Nodule	350	0.8010	0.8277	0.8414	0.8470
		Pulmonary Nodule and Mass Composite	4	0.8343	0.8702	0.8798	0.8816
	Indeterminate Pancreatic Lesion	Indeterminate Pancreatic Lesion	12	0.7402	0.7583	0.7493	0.7850
	Indeterminate Prostate Lesion	Prostate Nodule	91	0.6363	0.7065	0.7370	0.7476
	Indeterminate Ovarian Lesion	Indeterminate Ovarian Lesion	87	0.8264	0.8406	0.8626	0.8669
	Perirectal Lymph-node Target of Indeterminate Status	Perirectal Lymph Node of Indeterminate Status	15	0.8573	0.8680	0.8589	0.8806
	Indeterminate Nervous-system Imaging Abnormality	Indeterminate Brain Lesion	20	0.7854	0.8069	0.8205	0.8550
	Other Indeterminate Abdominopelvic Lesion	Indeterminate Kidney Lesion	78	0.7840	0.8131	0.8261	0.8318
		Indeterminate Uterine Lesion	39	0.7452	0.8097	0.8280	0.8375
		Indeterminate Bladder Lesion	26	0.8247	0.8401	0.8519	0.8549
		Adrenal Nodule	2	0.7937	0.8525	0.8536	0.8433
	Indeterminate Lymph-node Target	Indeterminate Lymph Node	188	0.6914	0.7386	0.7529	0.7643

Continued on next page

Table 11 (continued)

Taxonomy	Category	Combined Targets	Samples (instances)	SAMI3D-DW			
				BBox	+1	+3	+5
	Pathologic Fluid Collection and Abnormal Gas in Body Cavity	Pleural Effusion	15	0.4129	0.5041	0.6242	0.6253

BBox denotes the initial tight 2D box on the axial slice with the largest target area. +1/3/5 denote 1/3/5 cumulative point refinements after the initial box. The cohort contains CT and MR only. Each row averages valid instance Dice within one Combined Target; a physical object can contribute to multiple Combined Target rows. Mean is the unweighted average of Combined Target means with valid scores.