

INVARIANT SHAPE ANALYSIS OF SURFACES WITH SPHERICAL TOPOLOGY

T. SHASKA AND M.-R. SIADAT

ABSTRACT. Spherical harmonic descriptors of closed 3D shapes depend on the parameterization, the pose and the scale of the surface, and the standard rotation-invariant reductions, the power spectrum and the bispectrum, discard the relative orientation of the harmonic bands and cannot distinguish a shape from its mirror image. We construct a descriptor that removes all three dependencies exactly and loses nothing else: a conformal parameterization normalized by its conformal barycenter, followed by polynomial invariants of the rotation group. Identifying each harmonic band with a binary form turns the rotation quotient into classical invariant theory and makes reflections visible as the sign of an invariant, so chirality is recorded. The descriptor is complete for the truncated expansion, stable in the orbit distance, and comes with numerical diagnostics. Benchmarks confirm the guarantees, and on bilateral anatomical structures the descriptor separates mirror-image pairs from asymmetric pairs, which parity-blind descriptors cannot.

1. INTRODUCTION

Spherical harmonic expansions are a standard representation of closed genus-zero surfaces in shape analysis, from the SPHARM descriptors of brain structures [7, 29] to rotation-invariant retrieval of 3D models [21]. A surface is mapped to the sphere, a coordinate or radial function is expanded in spherical harmonics, and the coefficients are used as features. The coefficients, however, depend on three choices that are not part of the shape: the map to the sphere, the position of the surface in space, and its scale. The map is the serious one. For the area-preserving parameterizations of SPHARM the residual freedom is the infinite-dimensional group of area-preserving diffeomorphisms of the sphere, so an individual coefficient is a function of the map and not of the surface; the first-order-ellipsoid alignment that follows fixes a frame by convention, is undefined when two semi-axes coincide, and leaves a finite sign ambiguity. The usual remedies remove the dependence on rotations by discarding information. The power spectrum keeps only the norm of each band and forgets how the bands are oriented relative to each other; the bispectrum [20] keeps more but its completeness and its behaviour under reflection are not known band by band; and every descriptor built from rotation-invariant quantities of even degree cannot distinguish a shape from its mirror image, which for bilateral anatomy is the difference between the left and the right side.

This paper removes all three dependencies exactly and loses nothing else. Conformal parameterization [17, 11] reduces the map freedom to the Möbius group, normalization by the conformal barycenter [3, 9] reduces it to a rotation of the

2020 *Mathematics Subject Classification.* 13A50, 14L24, 65D18, 68U05.

Key words and phrases. spherical harmonics, shape descriptors, invariant theory, binary forms, conformal parameterization, chirality.

sphere, and the rotation is removed by polynomial invariants of the rotation group, for which the classical identification of harmonics with binary forms [10, 24] supplies both the invariants and, through a real structure, the sign that records handedness. Position is removed by a canonical center and scale by weighted projectivization. The construction is exact at every step, and the descriptor it produces is complete for the truncated expansion. Formally, let $S \subset \mathbb{R}^3$ be a smoothly embedded two-sphere; the shape space is

$$\mathcal{S} = \text{Emb}(\mathbb{S}^2, \mathbb{R}^3) / (G_{\text{task}} \times \text{Diff}^+(\mathbb{S}^2)),$$

where G_{task} is the group of orientation-preserving rigid motions of $\mathbb{R}^3$, of all rigid motions, or of orientation-preserving similarities, and the paper computes coordinates on $\mathcal{S}$ at the level of truncated coefficients. Constructions of separating invariants for general compact groups, with dimension or stability guarantees, are given in [16, 8]; here the group and the representation are specific, and the invariants are explicit.

The coordinate model expands the centered coordinate map, and the radial model expands the distance from a canonical center. Our results are as follows.

Canonical parameterization. Centered conformal parameterizations of S form one $\text{SO}(3)$ -orbit, so the whole coefficient tuple is determined up to one rotation of the sphere (Theorems 2.1 and 2.3); the all-band coordinate expansion determines S up to an orientation-preserving rigid motion (Theorem 4.5); and the squared L^2 truncation error at degree L is at most $|S|/(2\pi(L+1)(L+2))$, independently of the conformal factor (Theorem 2.4).

Harmonics as binary forms. A harmonic f of degree l is sent to the binary form $F_f(z) = f(v(z))$ of degree $2l$, where v parametrizes the null conic of $x_1^2 + x_2^2 + x_3^2$. This is an explicitly invertible, rotation-equivariant isomorphism that scales the L^2 norm to the Bombieri–Weyl norm by an explicit constant (Theorem 3.8). Real harmonics are the fixed points of an antilinear involution induced by the quaternionic structure of $\mathbb{C}^2$, and the roots of F_f are the endpoints of the Maxwell axes of f . A multihomogeneous $\text{SL}_2(\mathbb{C})$ -invariant of degree d_l in band l is real on real harmonics if $\sum_l l d_l$ is even and purely imaginary if it is odd; the first are the $\text{O}(3)$ -invariants and the second the pseudo-invariants (Theorem 3.6).

Separation and chirality. Seven invariants of degrees 1, 2, 2, 3, 3, 4, 6 separate the $\text{SO}(3)$ -orbits of the truncation through degree two (Theorem 4.9). A real harmonic of degree three is congruent to its mirror image iff the skew invariant R of its sextic vanishes, iff its three Maxwell axes are coplanar or one is orthogonal to a bisector of the other two, iff (for distinct axes) the associated genus-two curve has an elliptic involution; moreover $-iR = \det(a, b, c) G$ with G a polynomial in the inner products of the axes (Theorem 5.2).

Frames, metrics, stability. On the principal stratum a canonical frame reduces the residual rotations to the Klein four-group, and the invariant monomials of degree at most three in the aligned coefficients separate orbits; the frame has no continuous extension to the degenerate strata (Theorem 6.2). The orbit distance is a metric on the quotient, the invariants are Lipschitz for it on bounded sets, and on compact sets it is bounded by a Hölder power of the invariant discrepancy (Theorem 7.3). Theorem 7.6 collects these properties into one statement, and Section 8 gives the numerical realization with a synthetic end-to-end test.

Benchmarks. On classes of dipole–quadrupole pairs that share their power spectrum and bispectrum, and on their mirror images, the power spectrum and the

bispectrum classify at chance, the $O(3)$ -invariants reach one half, and the seven invariants of Theorem 4.9 classify all classes; the frame coordinates are ill-conditioned like the reciprocal of the spectral gap, while the invariants are not; and the measured stability exponents agree with Theorem 7.3 (Section 9). On eleven bilateral pairs of a public anatomical atlas, the descriptor identifies the pairs that are one chiral shape in two handednesses, with the pseudo-invariants recording the handedness, and separates them from pairs of different shapes; the diagnostics reject the two structures on which the conformal map folds (Section 9.4).

The identification of harmonics with binary forms, the spin covering and the Maxwell–Sylvester theorem are classical (Theorem 3.13). The contributions of this paper are the reality and parity rule of Theorem 3.6, which makes reflections visible in the phase of an invariant; the combination of conformal-barycenter normalization with this correspondence into a descriptor that is exactly independent of parameterization, position and scale, with a truncation bound independent of the conformal factor (Theorems 2.4 and 7.6); a complete descriptor through degree two with an explicit pseudo-invariant (Theorem 4.9); the characterization of chirality in degree three by the geometry of the Maxwell axes and the factorization $-iR = \lambda^{15} \det(a, b, c) G$ (Theorem 5.2); the identification of the residual group of ellipsoid-type alignment, with the proof that no continuous frame exists across the degenerate strata (Theorem 6.2); and two-sided stability with respect to the orbit distance (Theorem 7.3).

The construction is motivated by an unpublished shape analysis of deep perisylvian epilepsy by D. I. Sims, M.-R. Siadat, K. Elisevich and H. Soltanian-Zadeh (2015), which we call SPHARM-COM; Section 10 identifies the freedoms that its radial descriptor left unresolved.

2. CANONICAL PARAMETERIZATION

Let $\mathbb{S}^2 = \{u \in \mathbb{R}^3 : |u| = 1\}$ be the unit sphere and $\mathbb{B}^3 = \{x \in \mathbb{R}^3 : |x| < 1\}$ the open unit ball. We write $\langle \cdot, \cdot \rangle$ for the Euclidean inner product on $\mathbb{R}^3$ and for its complex bilinear extension to $\mathbb{C}^3$, and $|\cdot|$ for the Euclidean norm. We identify $\mathbb{S}^2$ with $\mathbb{P}^1 = \{[z_0 : z_1]\}$ through the coordinate $\zeta = z_0/z_1 = (-n_1 + in_2)/(1 - n_3)$ for $n \in \mathbb{S}^2$; this is stereographic projection from the north pole followed by a reflection, it is orientation-preserving for the outward orientation of $\mathbb{S}^2$, and it is the identification compatible with Section 3. Every element of $O(3) \setminus SO(3)$ has the form $-R$ with $R \in SO(3)$. A set with a free and transitive action of a group G is a **G -torsor**. For a probability measure μ on $\mathbb{S}^2$ and a continuous map g of $\mathbb{S}^2$, the pushforward $g_*\mu(E) = \mu(g^{-1}(E))$ satisfies $\int f d(g_*\mu) = \int f \circ g d\mu$; an **atom** of μ is a point of positive mass; and $d\omega = dA_{\text{round}}/4\pi$ is the round probability measure. The **Möbius group** $\text{Möb} \cong \text{PSL}_2(\mathbb{C})$ is the group of orientation-preserving conformal diffeomorphisms of $\mathbb{S}^2$ [1].

2.1. Spherical harmonics. For $l \geq 0$ let $\mathcal{H}_l$ be the space of restrictions to $\mathbb{S}^2$ of homogeneous harmonic polynomials of degree l on $\mathbb{R}^3$; $\dim \mathcal{H}_l = 2l + 1$, $-\Delta_{\mathbb{S}^2}$ acts on $\mathcal{H}_l$ by $l(l + 1)$, and $L^2(\mathbb{S}^2, d\omega)$ is the orthogonal Hilbert sum of the $\mathcal{H}_l$ [2]. For instance $\mathcal{H}_1 = \{\langle a, \cdot \rangle : a \in \mathbb{R}^3\}$ and $\mathcal{H}_2 = \{u \mapsto u^\top Qu : Q = Q^\top, \text{tr } Q = 0\}$. Write $u = (\sin \theta \cos \phi, \sin \theta \sin \phi, \cos \theta)$, let P_l be the Legendre polynomial and $P_l^m(t) =$

$(1 - t^2)^{m/2} P_l^{(m)}(t)$. We fix the orthonormal basis

$$\begin{aligned} Y_{l0} &= \sqrt{2l+1} P_l(\cos \theta), \\ Y_{lm} &= N_{lm} P_l^m(\cos \theta) \cos m\phi, \\ Y_{l,-m} &= N_{lm} P_l^m(\cos \theta) \sin m\phi, \end{aligned}$$

for $1 \leq m \leq l$, with $N_{lm} = \sqrt{2(2l+1)(l-m)!/(l+m)!}$. This fixes normalization and signs; software libraries differ from it by constant factors such as the Condon–Shortley factor $(-1)^m$. The **coefficients** of $f \in L^2(\mathbb{S}^2, d\omega)$ are $c_{lm}(f) = \int_{\mathbb{S}^2} f Y_{lm} d\omega$, the orthogonal projection onto $\mathcal{H}_l$ is $\pi_l f = \sum_{|m| \leq l} c_{lm}(f) Y_{lm}$, and l is the **band**. For $\mathbb{R}^3$ -valued functions π_l is taken componentwise.

The group $\text{SO}(3)$ acts by $(Q \cdot f)(u) = f(Q^{-1}u)$. It preserves each $\mathcal{H}_l$ and acts on $c_l(f) = (c_{l,-l}, \dots, c_{ll})$ by an orthogonal matrix $D_l(Q)$, the real form of the Wigner matrix [31]. The band norms $\|\pi_l f\|$ are invariant; the individual coefficients are not. A rotation R_α by α about the x_3 -axis fixes c_{l0} and rotates each pair $(c_{lm}, c_{l,-m})$ by the angle $m\alpha$, while rotations about other axes mix different orders m . A coefficient c_{lm} of a surface therefore depends on how the surface was placed on the sphere and in space. The rest of this section removes the placement on the sphere up to one rotation.

2.2. Conformal parameterizations and the conformal barycenter.

Throughout, $S \subset \mathbb{R}^3$ is a smoothly embedded two-sphere, oriented by its outward normal, with induced metric g_S , area form dA and area $|S|$. A diffeomorphism $\varphi: \mathbb{S}^2 \rightarrow S$ is **conformal** if $\varphi^* g_S = e^{2\rho} g_{\text{round}}$ for a smooth function ρ ; $e^{2\rho}$ is its **conformal factor**. By the uniformization theorem [13], the set $\Phi(S)$ of orientation-preserving conformal diffeomorphisms $\mathbb{S}^2 \rightarrow S$ is nonempty, and Möb acts on it freely and transitively by $\varphi \mapsto \varphi \circ g$; that is, $\Phi(S)$ is a Möb-torsor. Every element of Möb extends uniquely to a Möbius transformation of $\overline{\mathbb{B}^3}$, the extended group acts transitively on $\mathbb{B}^3$, and the stabilizer of the origin is $\text{SO}(3)$ acting linearly [1]. Hence removing the three nonrotational degrees of freedom of Möb amounts to choosing a point of $\mathbb{B}^3$.

For $x \in \mathbb{B}^3$, the **hyperbolic translation** $\tau_x: \overline{\mathbb{B}^3} \rightarrow \overline{\mathbb{B}^3}$ is the map

$$(1) \quad \tau_x(u) = \frac{(1 - |x|^2)(u - x) - |u - x|^2 x}{1 - 2\langle u, x \rangle + |u|^2 |x|^2}$$

The map τ_x , defined for $u \in \overline{\mathbb{B}^3}$, is a Möbius transformation of $\overline{\mathbb{B}^3}$. In the Poincaré ball model of hyperbolic space it is the translation along the geodesic through 0 and x . It sends x to 0 and 0 to $-x$, and it fixes the endpoints $\pm x/|x|$ of this geodesic. It maps $\mathbb{S}^2$ to itself, and τ_0 is the identity. For a probability measure μ on $\mathbb{S}^2$ set

$$\xi_\mu(x) = \int_{\mathbb{S}^2} \tau_x(u) d\mu(u), \quad x \in \mathbb{B}^3.$$

Since $\int f d(g_* \mu) = \int f \circ g d\mu$, the vector $\xi_\mu(x)$ is the Euclidean center of mass of the measure $(\tau_x)_* \mu$.

Douady and Earle [15] proved the following. If μ has no atoms, then the vector field ξ_μ has a unique zero $B(\mu) \in \mathbb{B}^3$, and

$$(2) \quad B(g_* \mu) = g(B(\mu)) \quad (g \in \text{Möb}).$$




FIGURE 1. The conformal barycenter on $\mathbb{S}^2$. Left: a probability measure μ consisting of 42 point masses of weight $1/42$ on $\mathbb{S}^2$ (points on the far hemisphere are drawn lighter; each atom has mass less than $1/2$, see Theorem 2.2), its Euclidean center of mass c (cross), and its conformal barycenter $x = B(\mu) \in \mathbb{B}^3$ (circle). The dashed segment is the geodesic of $\mathbb{B}^3$ through 0 and x . Right: the pushforward $(\tau_x)_*\mu$ under the hyperbolic translation τ_x ; its Euclidean center of mass is the origin. The points were computed from (1).

The point $B(\mu)$ is the **conformal barycenter** of μ . Equivalently, $B(\mu)$ is the unique $x \in \mathbb{B}^3$ such that $(\tau_x)_*\mu$ has center of mass 0 . Since τ_0 is the identity, $B(\mu) = 0$ if and only if $\int_{\mathbb{S}^2} u d\mu(u) = 0$. Figure 1 illustrates the construction.

Each $\varphi \in \Phi(S)$ determines the probability measure $\mu_\varphi = |S|^{-1}\varphi^*(dA)$ on $\mathbb{S}^2$. Explicitly, $\mu_\varphi(E) = |\varphi(E)|/|S|$ for every Borel set $E \subset \mathbb{S}^2$, where $|\varphi(E)|$ is the area of $\varphi(E)$. We call φ **centered** if

$$\int_{\mathbb{S}^2} u d\mu_\varphi(u) = 0,$$

and write $\Phi_0(S)$ for the set of centered parameterizations. Thus φ is centered if the area of S , transported to $\mathbb{S}^2$ by φ , has its center of mass at the origin.

Lemma 2.1. *The following are true:*

- (i) $\Phi_0(S)$ is nonempty and is a single orbit under $\mathrm{SO}(3)$ acting by precomposition.
- (ii) If $T(x) = \lambda Rx + b$ with $\lambda > 0$, $R \in \mathrm{SO}(3)$, then $\Phi_0(T(S)) = T \circ \Phi_0(S)$.
- (iii) If $T(x) = -\lambda Rx + b$ and r is any reflection of $\mathbb{S}^2$ in a plane through the origin, then $\Phi_0(T(S)) = T \circ \Phi_0(S) \circ r$.

Proof. (i) Fix $\varphi \in \Phi(S)$. Since φ is a diffeomorphism, μ_φ has a smooth positive density with respect to $d\omega$. Hence μ_φ has no atoms, and $B(\mu_\varphi)$ exists and is unique. For $g \in \mathrm{Möb}$,

$$(\varphi \circ g)^*(dA) = g^*\varphi^*(dA),$$

so $\mu_{\varphi \circ g} = (g^{-1})_* \mu_\varphi$. By (2), $B(\mu_{\varphi \circ g}) = g^{-1}(B(\mu_\varphi))$. Hence $\varphi \circ g$ is centered if and only if $g(0) = B(\mu_\varphi)$. Since Möb acts transitively on $\mathbb{B}^3$, such a g exists. If g and g' are two such elements, then $g^{-1}g'$ fixes the origin, so $g^{-1}g' \in \text{SO}(3)$.

(ii) The map $T \circ \varphi$ is conformal and orientation-preserving onto $T(S)$. Conversely, if $\psi \in \Phi(T(S))$, then $T^{-1} \circ \psi \in \Phi(S)$. Hence $\Phi(T(S)) = T \circ \Phi(S)$. Moreover $(T \circ \varphi)^* dA_{T(S)} = \lambda^2 \varphi^* dA_S$ and $|T(S)| = \lambda^2 |S|$, so $\mu_{T \circ \varphi} = \mu_\varphi$. Hence $T \circ \varphi$ is centered if and only if φ is.

(iii) The map T reverses the orientation of $\mathbb{R}^3$ and sends the outward normal of S to the outward normal of $T(S)$. Hence T restricts to an orientation-reversing map $S \rightarrow T(S)$. Since r also reverses orientation, $T \circ \varphi \circ r$ is conformal and orientation-preserving. As in (ii), $\Phi(T(S)) = T \circ \Phi(S) \circ r$. Since r is a linear isometry with $r^{-1} = r$, we have $\mu_{T \circ \varphi \circ r} = r_* \mu_\varphi$. Finally $\int u d(r_* \mu_\varphi) = r(\int u d\mu_\varphi)$, which vanishes if and only if $\int u d\mu_\varphi$ does. $\square$

Remark 2.2. For a triangulated surface the smooth construction is replaced by its discrete counterpart [3]. The pulled-back area measure becomes a finite sum of point masses, $\mu_h = \sum_i w_i \delta_{u_i}$. It has a unique conformal barycenter provided no atom carries mass $\geq 1/2$, by the stability criterion of Cantarella and Schumacher [9]. Uniqueness is not conditioning. The Hessian of the associated energy, computed in the same paper, governs the sensitivity of the barycenter to the mesh and enters the stability discussion of Section 7.

2.3. The conformal center and the harmonic coefficients. For $\varphi \in \Phi_0(S)$ put

$$(3) \quad \bar{c}(S) = \int_{\mathbb{S}^2} \varphi(u) d\omega.$$

The right side does not depend on the choice of $\varphi \in \Phi_0(S)$. Indeed, by Theorem 2.1(i) any other choice is $\varphi \circ R$ with $R \in \text{SO}(3)$, and $d\omega$ is $\text{SO}(3)$ -invariant. We call $\bar{c}(S)$ the **conformal center** of S .

The conformal center is an average of points of S , so it lies in the convex hull of S . The **area centroid** of S is

$$\frac{1}{|S|} \int_S x dA = \int_{\mathbb{S}^2} \varphi d\mu_\varphi.$$

The two centers average φ against different measures, $d\omega$ and $d\mu_\varphi$, and they differ in general. The conformal center is natural: $\bar{c}(T(S)) = T(\bar{c}(S))$ for every similarity T , orientation-preserving or not. This follows from Theorem 2.1(ii) and (iii), since T is affine and $d\omega$ is invariant under reflections.

Define the **centered coordinate map** and the **radial function** of φ by

$$(4) \quad X_\varphi(u) = \varphi(u) - \bar{c}(S), \quad d_\varphi(u) = |X_\varphi(u)|.$$

Their band components are

$$f_l(\varphi) = \pi_l d_\varphi \in \mathcal{H}_l, \quad C_l(\varphi) = \pi_l X_\varphi \in \mathcal{H}_l \otimes \mathbb{R}^3.$$

Since $\mathcal{H}_0$ consists of the constants, $C_0(\varphi) = \int_{\mathbb{S}^2} X_\varphi d\omega$. By (3), $C_0(\varphi) = 0$ identically. This is why the conformal center, rather than the area centroid, is taken as the origin. Write

$$f^{(L)}(\varphi) = (f_0, \dots, f_L), \quad C^{(L)}(\varphi) = (C_1, \dots, C_L).$$

The rotation action on functions extends to $\mathbb{R}^3$ -valued functions, where the group $\text{SO}(3) \times \text{SO}(3)$ acts by the **two-sided action**

$$(5) \quad ((Q, R) \cdot X)(u) = RX(Q^{-1}u)$$

for $(Q, R) \in \text{SO}(3) \times \text{SO}(3)$. The first factor rotates the parameter sphere. The second factor rotates the surface in $\mathbb{R}^3$.

Corollary 2.3. *For every $L \geq 1$ the following hold.*

- (i) *The set $\{f^{(L)}(\varphi) : \varphi \in \Phi_0(S)\}$ is one orbit of the diagonal action of $\text{SO}(3)$ on $\bigoplus_{l \leq L} \mathcal{H}_l$.*
- (ii) *The set $\{C^{(L)}(\varphi) : \varphi \in \Phi_0(S)\}$ is one orbit of $\text{SO}(3) \times \{1\}$ on $\bigoplus_{1 \leq l \leq L} \mathcal{H}_l \otimes \mathbb{R}^3$.*
- (iii) *If $T(x) = \lambda Rx + b$ with $\lambda > 0$ and $R \in \text{SO}(3)$, then $f_l(T \circ \varphi) = \lambda f_l(\varphi)$ and $C_l(T \circ \varphi) = \lambda R C_l(\varphi)$.*
- (iv) *If $T(x) = -\lambda Rx + b$ and r is as in Theorem 2.1(iii), then $f_l(T \circ \varphi \circ r) = \lambda f_l(\varphi) \circ r$ and $C_l(T \circ \varphi \circ r) = -\lambda R C_l(\varphi) \circ r$.*

Proof. Let $Q \in \text{SO}(3)$. Since $\bar{c}(S)$ does not depend on φ ,

$$X_{\varphi \circ Q}(u) = \varphi(Qu) - \bar{c}(S) = (Q^{-1} \cdot X_\varphi)(u).$$

Since π_l commutes with the action, $C_l(\varphi \circ Q) = Q^{-1} \cdot C_l(\varphi)$ and $f_l(\varphi \circ Q) = Q^{-1} \cdot f_l(\varphi)$, with the same Q in every band. Together with Theorem 2.1(i) this proves (i) and (ii). For (iii), Theorem 2.1(ii) and the naturality of $\bar{c}$ give

$$X_{T \circ \varphi} = T \circ \varphi - T(\bar{c}(S)) = \lambda R X_\varphi.$$

Hence $d_{T \circ \varphi} = \lambda d_\varphi$, and (iii) follows by applying π_l . For (iv), Theorem 2.1(iii) gives in the same way $X_{T \circ \varphi \circ r} = -\lambda R X_\varphi \circ r$. Since r is orthogonal, π_l commutes with composition by r , and (iv) follows. $\square$

Theorem 2.3 is the precise sense in which the coefficients of a centered conformal parameterization are canonical. The entire tuple of coefficients is determined by S up to one common rotation of the sphere. The full coordinate expansion determines X_φ in L^2 and hence, by continuity, the centered parameterized surface. The radial expansion retains only $|X_\varphi|$. No reconstruction theorem from radial data is asserted.

2.4. Truncation. The conformal gauge has one cost. Conformal maps of elongated or convoluted surfaces have large area distortion, so a small spherical cap may carry a large part of the surface. The following bound shows that the truncation error in $L^2(d\omega)$ is nevertheless controlled by the area alone. Norms without subscript are $L^2(d\omega)$ norms.

Proposition 2.4. *Let $\varphi \in \Phi_0(S)$ with conformal factor $e^{2\rho}$, let $X = X_\varphi$ and $X_l = \pi_l X$, and let $X^{(L)} = \sum_{l \leq L} X_l$ be the degree- L truncation of X . Then*

$$(6) \quad \begin{aligned} \|X - X^{(L)}\|_{L^2(d\omega)}^2 &\leq \frac{|S|}{2\pi(L+1)(L+2)}, \\ \int_{\mathbb{S}^2} |X - X^{(L)}|^2 d\mu_\varphi &\leq \frac{2 \sup e^{2\rho}}{(L+1)(L+2)}. \end{aligned}$$

The same bounds hold for d_φ in place of X_φ .

Proof. For $u \in \mathbb{S}^2$ let e_1, e_2 be an orthonormal basis of $T_u \mathbb{S}^2$ for g_{round} . The Hilbert–Schmidt norm of the differential is

$$|d\varphi_u|^2 = |d\varphi_u(e_1)|^2 + |d\varphi_u(e_2)|^2.$$

For a conformal map both terms equal $e^{2\rho(u)}$, so $|d\varphi|^2 = 2e^{2\rho}$. Since $\varphi^* dA = e^{2\rho} dA_{\text{round}}$, we get $\int_{\mathbb{S}^2} |d\varphi|^2 dA_{\text{round}} = 2|S|$. Since X differs from φ by a constant, $|\nabla X| = |d\varphi|$. With $d\omega = dA_{\text{round}}/4\pi$ this gives $\|\nabla X\|^2 = |S|/2\pi$.

Since $-\Delta_{\mathbb{S}^2}$ acts on $\mathcal{H}_l$ by $l(l+1)$, Green’s formula and orthogonality give

$$\|\nabla X\|^2 = \sum_{l \geq 1} l(l+1) \|X_l\|^2.$$

Hence

$$\begin{aligned} \|X - X^{(L)}\|^2 &= \sum_{l > L} \|X_l\|^2 \\ &\leq \frac{1}{(L+1)(L+2)} \sum_{l > L} l(l+1) \|X_l\|^2 \\ &\leq \frac{\|\nabla X\|^2}{(L+1)(L+2)}, \end{aligned}$$

which is the first bound. For the second bound,

$$d\mu_\varphi = |S|^{-1} e^{2\rho} dA_{\text{round}} = \frac{4\pi}{|S|} e^{2\rho} d\omega.$$

So the area-weighted error is at most $4\pi \sup e^{2\rho}/|S|$ times the first bound, which equals $2 \sup e^{2\rho}/((L+1)(L+2))$. Finally, $d_\varphi = |X_\varphi|$ is Lipschitz and $|\nabla d_\varphi| \leq |\nabla X_\varphi|$ almost everywhere, so the same argument applies to d_φ . $\square$

The first bound is uniform in the shape. The second bound shows where a convoluted surface pays. The geometric error is governed by the largest conformal factor, and an implementation should report this quantity rather than the number of coefficients. The quantity $\sup e^{2\rho}$ does not depend on the choice of $\varphi \in \Phi_0(S)$, since replacing φ by $\varphi \circ R$ replaces ρ by $\rho \circ R$. Area-preserving parameterizations avoid the second bound. However, their residual gauge is the infinite-dimensional group of area-preserving diffeomorphisms of $\mathbb{S}^2$, so they cannot support the exact quotient constructed here.

3. SPHERICAL HARMONICS AS BINARY FORMS

This section identifies the real spherical harmonics of degree l with a real form of the binary forms of degree $2l$. Rotations act through the double cover $\text{SU}(2) \subset \text{SL}_2(\mathbb{C})$ of $\text{SO}(3)$, and rotation invariants of harmonics become classical invariants of binary forms. A nonzero real harmonic f of degree l determines l lines through the origin, its Maxwell axes [23, 30]; the $2l$ roots of the binary form F_f are the points where these lines meet $\mathbb{S}^2$, and a rotation of f rotates the roots. The identification depends on phase conventions, which we fix once.

3.1. Harmonic polynomials and the null cone. Let $P_l(\mathbb{C}^3)$ be the homogeneous polynomials of degree l in $x = (x_1, x_2, x_3)$ with complex coefficients and $\mathcal{H}_l^{\mathbb{C}} = \ker(\Delta: P_l(\mathbb{C}^3) \rightarrow P_{l-2}(\mathbb{C}^3))$, the complexification of $\mathcal{H}_l$; restriction to $\mathbb{S}^2$ identifies real harmonic polynomials with $\mathcal{H}_l$, and we use the same letter for both. **Complex conjugation** on $\mathcal{H}_l^{\mathbb{C}}$ is $\bar{f}(x) = \overline{f(\bar{x})}$, and its fixed points are the real harmonics. Let

$$q(x) = x_1^2 + x_2^2 + x_3^2 = \langle x, x \rangle,$$

with $\langle \cdot, \cdot \rangle$ the complex bilinear pairing; for $x \in \mathbb{C}^3$, $q(x)$ can vanish with $x \neq 0$.

Proposition 3.1. *For $l \geq 2$, $P_l(\mathbb{C}^3) = \mathcal{H}_l^{\mathbb{C}} \oplus q P_{l-2}(\mathbb{C}^3)$, and the same holds over $\mathbb{R}$. Hence $\dim \mathcal{H}_l^{\mathbb{C}} = 2l + 1$, and every $P \in P_l(\mathbb{C}^3)$ has a unique **harmonic part** $\pi(P) \in \mathcal{H}_l^{\mathbb{C}}$ with $P - \pi(P) \in q P_{l-2}(\mathbb{C}^3)$. The decomposition is $\mathrm{SO}_3(\mathbb{C})$ -invariant, and π is equivariant.*

Proof. For $P, R \in P_l(\mathbb{C}^3)$ put $[P, R] = P(\partial)\bar{R}$. Distinct monomials are orthogonal and $[x^\alpha, x^\alpha] = \alpha! > 0$, so $[\cdot, \cdot]$ is a positive definite Hermitian form. Since $q(\partial) = \Delta$, one has $[qP, R] = [P, \Delta R]$, so multiplication by q is the adjoint of Δ and $P_l(\mathbb{C}^3) = \ker \Delta \oplus q P_{l-2}(\mathbb{C}^3)$ orthogonally. The group $\mathrm{SO}_3(\mathbb{C})$ preserves q , hence both summands. $\square$

Let Sym^n be the binary forms $F(z_0, z_1)$ of degree n , with $\mathrm{SL}_2(\mathbb{C})$ acting by $(g \cdot F)(z) = F(g^{-1}z)$. The **roots** of $F \neq 0$ are the points $[z] \in \mathbb{P}^1$ with $F(z) = 0$; with multiplicity they form the **root divisor**, which determines F up to a scalar. The **null cone** is $N = \{x \in \mathbb{C}^3 : q(x) = 0\}$. Since q has rank three it is irreducible, and every polynomial vanishing on N is divisible by q . The **Veronese map** is

$$(7) \quad v(z_0, z_1) = (z_0^2 - z_1^2, i(z_0^2 + z_1^2), 2z_0z_1),$$

and the **quadratic** of $x \in \mathbb{C}^3$ is

$$(8) \quad \begin{aligned} Q_x(z) &= \langle x, v(z) \rangle \\ &= (x_1 + ix_2)z_0^2 + 2x_3z_0z_1 + (-x_1 + ix_2)z_1^2. \end{aligned}$$

In the normalization $\mathrm{disc}(\alpha z_0^2 + 2\beta z_0z_1 + \gamma z_1^2) = 4(\beta^2 - \alpha\gamma)$ one has $\mathrm{disc}(Q_x) = 4q(x)$. For $z \neq 0$ put $|z|^2 = |z_0|^2 + |z_1|^2$ and

$$(9) \quad m(z) = \frac{i}{2|z|^4} v(z) \times \overline{v(z)}.$$

Lemma 3.2. (i) $z_0^2 = \frac{1}{2}(v_1 - iv_2)$, $z_1^2 = -\frac{1}{2}(v_1 + iv_2)$, $z_0z_1 = \frac{1}{2}v_3$; hence v_1, v_2, v_3 is a basis of Sym^2 and $x \mapsto Q_x$ is an isomorphism $\mathbb{C}^3 \rightarrow \mathrm{Sym}^2$.
(ii) $v(\mathbb{C}^2) = N$, and $v(z) = v(z')$ iff $z' = \pm z$. The vectors $v(z)$ span $\mathbb{C}^3$.

Proof. (i) is immediate from (7). (ii) Expanding, $q(v(z)) = 0$. For $x \in N \setminus 0$ choose $z_0^2 = \frac{1}{2}(x_1 - ix_2)$ and $z_1^2 = -\frac{1}{2}(x_1 + ix_2)$; then $(z_0z_1)^2 = \frac{1}{4}x_3^2$ since $q(x) = 0$, and after changing the sign of z_1 , $v(z) = x$ by (i). If $v(z) = v(z')$, (i) gives $z_0^2 = z_0'^2$, $z_1^2 = z_1'^2$, $z_0z_1 = z_0'z_1'$, so $z' = \pm z$. The vectors $v(1, 0), v(0, 1), v(1, 1)$ have determinant $4i \neq 0$. $\square$

Lemma 3.3. (i) For $z \neq 0$, $m(z)$ is the real unit vector

$$\frac{(-2\Re(z_0\bar{z}_1), 2\Im(z_0\bar{z}_1), |z_0|^2 - |z_1|^2)}{|z_0|^2 + |z_1|^2},$$

- $m(\lambda z) = m(z)$ for $\lambda \in \mathbb{C}^\times$, and the induced map $\bar{m}: \mathbb{P}^1 \rightarrow \mathbb{S}^2$ is the orientation-preserving bijection with $\bar{m}([\zeta : 1]) = n$ for $\zeta = (-n_1 + in_2)/(1 - n_3)$ and $\bar{m}([1 : 0]) = (0, 0, 1)$.
- (ii) For $a \in \mathbb{R}^3 \setminus 0$, $Q_a(z) = 0$ iff $m(z) = \pm a/|a|$.
- (iii) The antipodal map of $\mathbb{S}^2$ is $[z] \mapsto [jz]$, where $j(z_0, z_1) = (-\bar{z}_1, \bar{z}_0)$, and $v(jz) = -\overline{v(z)}$.

Proof. (i) With $w = v(z)$, $\overline{w \times \bar{w}} = -w \times \bar{w}$, so $iw \times \bar{w}$ is real, and expanding with (7) gives $\frac{i}{2}w \times \bar{w} = |z|^2(-2\Re(z_0\bar{z}_1), 2\Im(z_0\bar{z}_1), |z_0|^2 - |z_1|^2)$, a vector of length $|z|^4$. Substituting $z = (\zeta, 1)$ and solving for ζ gives the inverse; stereographic projection from the north pole reverses orientation at the south pole, and the reflection $\zeta \mapsto -\bar{\zeta}$ reverses it again. (ii) Since $q(w) = 0$, $\Re w$ and $\Im w$ are orthogonal of equal length, and both are orthogonal to $m(z)$, so they span $m(z)^\perp$. For real a , $Q_a(z) = \langle a, \Re w \rangle + i\langle a, \Im w \rangle$ vanishes iff $a \in \mathbb{R}m(z)$. (iii) The formula of (i) gives $m(jz) = -m(z)$, and $v(jz) = -\overline{v(z)}$ is a direct substitution. $\square$

From now on $\mathbb{P}^1$ is identified with $\mathbb{S}^2$ through $\bar{m}$. By Theorem 3.3(ii), the root divisor of Q_a for real $a \neq 0$ is the antipodal pair $\pm a/|a|$: Q_a is the binary quadratic whose roots are the two points of the axis $\mathbb{R}a$ on the sphere.

3.2. The spin covering and the map Ψ_l . For $g \in \mathrm{SL}_2(\mathbb{C})$, each component of $z \mapsto v(gz)$ is a binary quadratic, so by Theorem 3.2(i) there is a unique $\rho(g) \in \mathrm{GL}_3(\mathbb{C})$ with

$$(10) \quad v(gz) = \rho(g)v(z) \quad \text{for all } z \in \mathbb{C}^2,$$

the classical spin covering [10]. Its entries are quadratic in the entries of g ; for $g_\theta = \mathrm{diag}(e^{i\theta/2}, e^{-i\theta/2})$ a direct substitution gives $\rho(g_\theta) = R_{-\theta}$, the rotation by $-\theta$ about the x_3 -axis.

Lemma 3.4. *The map ρ is a surjective homomorphism $\mathrm{SL}_2(\mathbb{C}) \rightarrow \mathrm{SO}_3(\mathbb{C})$ with kernel $\{\pm 1\}$, and $g \cdot Q_x = Q_{\rho(g)x}$. It restricts to a surjective homomorphism $\mathrm{SU}(2) \rightarrow \mathrm{SO}(3)$ with kernel $\{\pm 1\}$, and $m(gz) = \rho(g)m(z)$ for $g \in \mathrm{SU}(2)$.*

Proof. Since the $v(z)$ span $\mathbb{C}^3$, ρ is a homomorphism. The quadratic form $q \circ \rho(g)$ vanishes on $N = v(\mathbb{C}^2)$, so it equals $c(g)q$ with $c: \mathrm{SL}_2(\mathbb{C}) \rightarrow \mathbb{C}^\times$ a homomorphism, which is trivial because $\mathrm{SL}_2(\mathbb{C})$ is perfect; by connectedness $\det \rho(g) = 1$. Then $(g \cdot Q_x)(z) = \langle x, \rho(g)^\top v(z) \rangle = Q_{\rho(g)x}(z)$. If $\rho(g) = 1$, then $gz = \pm z$ for all z by Theorem 3.2(ii), so $g = \pm 1$; hence $d\rho$ is injective at 1; both groups have complex dimension three, so the image contains a neighborhood of 1, and since $\mathrm{SO}_3(\mathbb{C})$ is connected, ρ is onto. For $g \in \mathrm{SU}(2)$ one has $gj = jg$, so $\rho(g)v(jz) = \overline{v(jgz)} = -\overline{\rho(g)v(z)}$ by Theorem 3.3(iii), while $\rho(g)v(jz) = -\rho(g)\overline{v(z)}$; since the $v(z)$ span $\mathbb{C}^3$, $\rho(g)$ is real, so $\rho(\mathrm{SU}(2)) \subset \mathrm{SO}(3)$. It is a closed subgroup of dimension three, hence all of $\mathrm{SO}(3)$. Finally $A = \rho(g)$ is real orthogonal of determinant one, so $A\bar{w} = \overline{Aw}$ and $(Aw) \times (Aw') = A(w \times w')$; apply this to (9) with $|gz| = |z|$. $\square$

Definition 3.5. For $f \in \mathcal{H}_l^\mathbb{C}$ the **binary form of f** is $F_f = f \circ v \in \mathrm{Sym}^{2l}$, and $\Psi_l: \mathcal{H}_l^\mathbb{C} \rightarrow \mathrm{Sym}^{2l}$ is the linear map $\Psi_l(f) = F_f$.

On linear forms (7) gives

$$(11) \quad \begin{aligned} \Psi_1(x_1 + ix_2) &= -2z_1^2, & \Psi_1(x_1 - ix_2) &= 2z_0^2, \\ \Psi_1(x_3) &= 2z_0z_1. \end{aligned}$$

Since evaluation at $v(z)$ is multiplicative and kills $qP_{l-2}(\mathbb{C}^3)$, for every $P \in P_l(\mathbb{C}^3)$

$$(12) \quad \Psi_l(\pi(P))(z) = P(v(z)).$$

3.3. The real form and reflection parity. A **real structure** on a complex vector space W is an antilinear map σ with $\sigma^2 = 1$; its fixed set $\text{Fix}(\sigma)$ is the **real form** of W . Complex conjugation is a real structure on $\mathcal{H}_l^\mathbb{C}$ with real form $\mathcal{H}_l$. We transport it to Sym^{2l} . For $F \in \text{Sym}^{2l}$ define

$$(13) \quad \sigma_l(F)(z) = (-1)^l \overline{F(jz)}.$$

If $F = \sum_{k=0}^{2l} \binom{2l}{k} a_k z_0^{2l-k} z_1^k$, then

$$(14) \quad \sigma_l(F) = \sum_{k=0}^{2l} \binom{2l}{k} (-1)^{l+k} \bar{a}_{2l-k} z_0^{2l-k} z_1^k.$$

Thus $F \in \text{Fix}(\sigma_l)$ iff $a_{2l-k} = (-1)^{l+k} \bar{a}_k$ for all k ; this is not the condition that the coefficients be real. On forms of odd degree the same formula squares to -1 and is a quaternionic structure, which is why real harmonics correspond to forms of even degree.

Let $V = \bigoplus_{l \leq L} \text{Sym}^{2l}$ with $\text{SL}_2(\mathbb{C})$ acting summandwise. An **invariant** is a polynomial $I: V \rightarrow \mathbb{C}$ with $I(g \cdot F) = I(F)$. It is **multihomogeneous of multi-degree** (d_l) if $I((t_l F_l)_l) = \prod_l t_l^{d_l} I((F_l)_l)$; every invariant is a sum of multihomogeneous ones. It **has real coefficients** if it is a real polynomial in the binomial coefficients a_k of the forms.

Lemma 3.6. *The following are true:*

- (i) σ_l is a real structure on Sym^{2l} commuting with $\text{SU}(2)$, and $\Psi_l(\bar{f}) = \sigma_l(\Psi_l f)$ for all $f \in \mathcal{H}_l^\mathbb{C}$. Hence $\Psi_l(\mathcal{H}_l) = \text{Fix}(\sigma_l)$.
- (ii) A nonzero $F \in \text{Sym}^{2l}$ is a complex multiple of a σ_l -fixed form if and only if its root divisor is invariant under the antipodal map. Every such F factors as $F = \lambda Q_{a_1} \cdots Q_{a_l}$ with $a_k \in \mathbb{R}^3 \setminus \{0\}$ and $\lambda \in \mathbb{C}^\times$; the lines $\mathbb{R}a_k$ are determined by F up to order, with multiplicity; and λ is real if and only if F is σ_l -fixed.
- (iii) Let I be a multihomogeneous invariant of multidegree (d_l) with real coefficients. Then for all real harmonics $f_l \in \mathcal{H}_l$, $l \leq L$,

$$I((F_{f_l})_l) = (-1)^{\sum_l l d_l} \overline{I((F_{f_l})_l)}.$$

- (iv) The map $-1 \in \text{O}(3)$ acts on $\mathcal{H}_l$ by $(-1)^l$, and it multiplies I by $(-1)^{\sum_l l d_l}$. If $\sum_l l d_l$ is even, I is invariant under $\text{O}(3)$; if it is odd, I changes sign under every reflection. Every invariant decomposes uniquely as the sum of an even and an odd part.

Proof. (i) Since $j^2 = -1$ and F has even degree, $\sigma_l(\sigma_l F)(z) = (-1)^l \overline{\sigma_l(F)(jz)} = F(j^2 z) = F(-z) = F(z)$, so $\sigma_l^2 = 1$, and σ_l is a real structure. For $g \in \text{SU}(2)$, $jg^{-1} = g^{-1}j$, so

$$\begin{aligned} \sigma_l(g \cdot F)(z) &= (-1)^l \overline{F(g^{-1}jz)} = (-1)^l \overline{F(jg^{-1}z)} \\ &= (g \cdot \sigma_l F)(z). \end{aligned}$$

For $f \in \mathcal{H}_l^{\mathbb{C}}$, using $\bar{f}(x) = \overline{f(\bar{x})}$ and Theorem 3.3(iii),

$$\begin{aligned}\Psi_l(\bar{f})(z) &= \overline{f(\overline{v(z)})} = \overline{f(-v(jz))} \\ &= (-1)^l \overline{(\Psi_l f)(jz)} = \sigma_l(\Psi_l f)(z),\end{aligned}$$

using that f is homogeneous of degree l . Thus Ψ_l intertwines conjugation and σ_l . A linear isomorphism that intertwines two real structures maps the fixed set of one onto the fixed set of the other, so $\Psi_l(\mathcal{H}_l) = \text{Fix}(\sigma_l)$.

(ii) By (13), $\sigma_l(F)(z) = 0$ if and only if $F(jz) = 0$, so the root divisor of $\sigma_l F$ is the image of the root divisor of F under the antipodal map $[z] \mapsto [jz]$, with the same multiplicities. Suppose the root divisor of F is antipodally invariant. Then $\sigma_l F$ and F have the same root divisor, so $\sigma_l F = cF$ for some $c \in \mathbb{C}^\times$. Applying σ_l again, $F = \sigma_l(cF) = \bar{c}\sigma_l F = \bar{c}cF$, so $|c| = 1$. Write $c = e^{2i\theta}$. Then $\sigma_l(e^{i\theta}F) = e^{-i\theta}cF = e^{i\theta}F$, so $e^{i\theta}F$ is σ_l -fixed. Conversely, if $F = \mu G$ with G fixed, the root divisor of F equals that of G , which is antipodally invariant.

Now let F be σ_l -fixed and nonzero. The antipodal map has no fixed points on $\mathbb{P}^1$, so the $2l$ roots of F , counted with multiplicity, fall into l antipodal pairs. Let $\{p, -p\}$ be one of them, with $p \in \mathbb{S}^2$, and let $a \in \mathbb{R}^3 \setminus \{0\}$ be any vector with $a/|a| = \pm p$. By Theorem 3.3(ii), Q_a has root divisor $\{p, -p\}$. A binary form is determined by its root divisor up to a scalar, so $F = \lambda Q_{a_1} \cdots Q_{a_l}$ with a_k chosen this way for the l pairs, and $\lambda \in \mathbb{C}^\times$. The lines $\mathbb{R}a_k$ are determined by the root pairs, hence by F , up to order and with multiplicity. Each Q_{a_k} is σ_1 -fixed by (14), and $\sigma_l(FG) = \sigma_{l_1}(F)\sigma_{l_2}(G)$ for $F \in \text{Sym}^{2l_1}$, $G \in \text{Sym}^{2l_2}$, $l = l_1 + l_2$, directly from (13). Hence $\prod_k Q_{a_k}$ is σ_l -fixed, and $\sigma_l(F) = \bar{\lambda} \prod_k Q_{a_k}$. So F is fixed if and only if $\bar{\lambda} = \lambda$.

(iii) Let $w \in \text{SL}_2(\mathbb{C})$ be the matrix with $wz = (-z_1, z_0)$. For $F = \sum_k \binom{2l}{k} a_k z_0^{2l-k} z_1^k$,

$$\begin{aligned}(w^{-1} \cdot F)(z) &= F(wz) = \sum_k \binom{2l}{k} a_k (-z_1)^{2l-k} z_0^k \\ &= \sum_k \binom{2l}{k} (-1)^k a_{2l-k} z_0^{2l-k} z_1^k,\end{aligned}$$

using $(-1)^{2l-k} = (-1)^k$ and reindexing. Comparing with (14), $\sigma_l(F) = \overline{(-1)^l w^{-1} \cdot F}$, where the bar is conjugation of all coefficients. Now let I be as in the statement and $F_l \in \text{Sym}^{2l}$ arbitrary. Since I has real coefficients, I of the conjugated tuple is the conjugate of I ; since I is multihomogeneous, the scalars $(-1)^l$ come out as $(-1)^{\sum_l l d_l}$; since I is invariant, w^{-1} can be dropped. Hence

$$\begin{aligned}I((\sigma_l F_l)_l) &= \overline{I((-1)^l w^{-1} \cdot F_l)_l} \\ &= (-1)^{\sum_l l d_l} \overline{I((F_l)_l)}.\end{aligned}$$

For real f_l , $F_l = F_{f_l}$ satisfies $\sigma_l F_l = F_l$ by (i), and the claim follows.

(iv) The map -1 sends a homogeneous polynomial f of degree l to $f(-x) = (-1)^l f(x)$. Hence it multiplies F_{f_l} by $(-1)^l$, and by multihomogeneity it multiplies I by $\prod_l (-1)^{l d_l} = (-1)^{\sum_l l d_l}$. Every reflection is $-R$ with $R \in \text{SO}(3)$, and I is $\text{SO}(3)$ -invariant, so a reflection acts on I as -1 does. The even and odd parts of an invariant are the sums of its multihomogeneous components with $\sum_l l d_l$ even and odd, respectively. $\square$

Definition 3.7. A multihomogeneous invariant I with real coefficients is an $O(3)$ -invariant if $\sum_l l d_l$ is even, and an $O(3)$ -**pseudo-invariant**, or simply a **pseudo-invariant**, if $\sum_l l d_l$ is odd.

By Theorem 3.6(iii) and (iv), an $O(3)$ -invariant takes real values on real harmonics and is unchanged by reflections, while a pseudo-invariant takes purely imaginary values on real harmonics and changes sign under every reflection. For a pseudo-invariant I , the function $-iI$ is real on real harmonics and changes sign under reflections.

Part (ii) of Theorem 3.6 is Maxwell's multipole theorem in Sylvester's form [23, 30]. In the language of harmonics it reads as follows. Every nonzero $f \in \mathcal{H}_l$ is the harmonic part of a product of l real linear forms,

$$f = \pi(\lambda \langle a_1, x \rangle \cdots \langle a_l, x \rangle),$$

with $a_k \in \mathbb{R}^3 \setminus \{0\}$ and $\lambda \in \mathbb{R}^\times$, and the l lines $\mathbb{R}a_k$ are determined by f up to order, with multiplicity. Indeed, by (12) and (11), Ψ_l of the right side is $\lambda \prod_k Q_{a_k}$, and Ψ_l is injective. The lines $\mathbb{R}a_k$ are the **Maxwell axes** of f . On the sphere they are the l antipodal root pairs of F_f .

Part (iii) is the statement used repeatedly below. The degree-15 skew invariant R of the sextic is real on sextics with real coefficients but purely imaginary on the different real form arising from degree-three real harmonics; part (iv) identifies this with reflection parity. We have verified (iii) by exact arithmetic on random real harmonics of degree three: the Igusa invariants J_2, J_4, J_6, J_{10} evaluate to rational numbers and R to a rational multiple of i (Section 8).

3.4. The harmonic–binary-form correspondence. The results of this section combine into one statement, on which everything that follows rests. For $F = \sum_{k=0}^{2l} \binom{2l}{k} a_k z_0^{2l-k} z_1^k$ the **Bombieri–Weyl norm** [4] is $\|F\|_{\text{BW}}^2 = \sum_k \binom{2l}{k} |a_k|^2$. For $1 \leq p \leq l$ put $\mathcal{Y}_{l0} = Y_{l0}$ and

$$\mathcal{Y}_{l,\pm p} = \frac{(\mp 1)^p}{\sqrt{2}} (Y_{lp} \pm i Y_{l,-p}).$$

The $\mathcal{Y}_{lm}$, $|m| \leq l$, form an orthonormal basis of $\mathcal{H}_l^\mathbb{C}$ for the Hermitian inner product of $L^2(\mathbb{S}^2, d\omega)$, and $\mathcal{Y}_{lm}$ is a multiple of $P_l^{|m|}(\cos \theta) e^{im\phi}$. Let $V_L = \bigoplus_{l \leq L} \mathcal{H}_l$ and $\Psi = \bigoplus_{l \leq L} \Psi_l$.

Theorem 3.8 (Harmonic–binary-form correspondence). *Let $l \geq 0$. Let $v(z_0, z_1) = (z_0^2 - z_1^2, i(z_0^2 + z_1^2), 2z_0 z_1)$ be the Veronese map, and for $f \in \mathcal{H}_l^\mathbb{C}$, a homogeneous harmonic polynomial on $\mathbb{C}^3$, let $F_f = \Psi_l(f) = f \circ v$ be its binary form (Theorem 3.5). The following are true.*

(i) **Algebra.** *The substitution $P \mapsto P \circ v$ induces an isomorphism of graded algebras*

$$\mathbb{C}[x_1, x_2, x_3]/(q) \cong \bigoplus_{l \geq 0} \text{Sym}^{2l},$$

which maps $P_l(\mathbb{C}^3)$ onto Sym^{2l} with kernel $q P_{l-2}(\mathbb{C}^3)$. Its restriction $\Psi_l: \mathcal{H}_l^\mathbb{C} \rightarrow \text{Sym}^{2l}$ is a linear isomorphism. If $P \in P_l(\mathbb{C}^3)$ and $P \circ v = F$, then $\Psi_l^{-1}(F) = \pi(P)$, where

$$\pi(P) = \sum_{j=0}^{\lfloor l/2 \rfloor} \frac{(-1)^j q^j \Delta^j P}{2^j j! \prod_{i=1}^j (2l - 2i + 1)}.$$

- For $f \in \mathcal{H}_l^{\mathbb{C}}$ and $g \in \mathcal{H}_m^{\mathbb{C}}$ one has $\Psi_{l+m}(\pi(fg)) = \Psi_l(f) \Psi_m(g)$.
- (ii) **Equivariance.** The map ρ is a surjective homomorphism $\mathrm{SL}_2(\mathbb{C}) \rightarrow \mathrm{SO}_3(\mathbb{C})$ with kernel $\{\pm 1\}$. It restricts to a surjective homomorphism $\mathrm{SU}(2) \rightarrow \mathrm{SO}(3)$ with kernel $\{\pm 1\}$. For $g \in \mathrm{SL}_2(\mathbb{C})$ and $f \in \mathcal{H}_l^{\mathbb{C}}$ one has $\Psi_l(\rho(g) \cdot f) = g \cdot \Psi_l(f)$. Every $\mathrm{SL}_2(\mathbb{C})$ -equivariant linear map $\mathcal{H}_l^{\mathbb{C}} \rightarrow \mathrm{Sym}^{2l}$ is a scalar multiple of Ψ_l .
- (iii) **Coefficients.** For $|m| \leq l$,

$$\Psi_l(\mathcal{Y}_{lm}) = K_{lm} z_0^{l-m} z_1^{l+m},$$

$$K_{lm} = \frac{(2l)!}{l!} \sqrt{\frac{2l+1}{(l-m)!(l+m)!}}.$$

For $f = \sum_{|m| \leq l} \eta_m \mathcal{Y}_{lm} \in \mathcal{H}_l^{\mathbb{C}}$ and $F_f = \sum_k \binom{2l}{k} a_k z_0^{2l-k} z_1^k$ one has

$$\eta_m = \frac{l! a_{l+m}}{\sqrt{(2l+1)(l-m)!(l+m)!}}.$$

Let $f = \sum_{|m| \leq l} c_{lm} Y_{lm} \in \mathcal{H}_l$, let $F_f = \sum_k \binom{2l}{k} a_k z_0^{2l-k} z_1^k$, and put $\nu_{lp} = \sqrt{(2l+1)(l-p)!(l+p)!}/2/l!$ for $1 \leq p \leq l$. Then

$$a_l = \sqrt{2l+1} c_{l0},$$

$$a_{l-p} = \nu_{lp} (c_{lp} + i c_{l,-p}),$$

$$a_{l+p} = (-1)^p \nu_{lp} (c_{lp} - i c_{l,-p}).$$

Conversely, $c_{l0} = a_l/\sqrt{2l+1}$, $c_{lp} = \Re(a_{l-p})/\nu_{lp}$ and $c_{l,-p} = \Im(a_{l-p})/\nu_{lp}$.

(iv) **Norm.** For every $f \in \mathcal{H}_l^{\mathbb{C}}$,

$$\|\Psi_l(f)\|_{\mathrm{BW}}^2 = (2l+1) \binom{2l}{l} \|f\|_{L^2(d\omega)}^2.$$

Hence $\widehat{\Psi}_l = ((2l+1) \binom{2l}{l})^{-1/2} \Psi_l$ is unitary from $L^2(\mathbb{S}^2, d\omega)$ to the Bombieri–Weyl norm. For real $f \in \mathcal{H}_l$ one has $\|F_f\|_{\mathrm{BW}}^2 = (-1)^l (F_f, F_f)_{2l}$, where $(F, F)_{2l} = \sum_k (-1)^k \binom{2l}{k} a_k a_{2l-k}$ is the transvectant (18).

- (v) **Real form.** For $f \in \mathcal{H}_l^{\mathbb{C}}$ one has $\Psi_l(\bar{f}) = \sigma_l(\Psi_l f)$. Hence $\Psi_l(\mathcal{H}_l) = \mathrm{Fix}(\sigma_l)$, the set of forms with $a_{2l-k} = (-1)^{l+k} \bar{a}_k$ for all k .

Proof. (i) The map $P \mapsto P \circ v$ is a homomorphism of algebras, and it sends $P_l(\mathbb{C}^3)$ into Sym^{2l} . By Theorem 3.2(i) the components of v span Sym^2 . Every monomial of degree $2l$ in z_0, z_1 is a product of l monomials of degree two, so the map is onto Sym^{2l} . Its kernel consists of the polynomials that vanish on $v(\mathbb{C}^2) = N$ (Theorem 3.2(ii)), and this is the ideal (q) since q is irreducible. By Theorem 3.1, $\mathcal{H}_l^{\mathbb{C}} \cap q P_{l-2}(\mathbb{C}^3) = 0$, so Ψ_l is injective, and it is an isomorphism because both spaces have dimension $2l+1$. If $P \circ v = F$, then $P - \pi(P) \in q P_{l-2}(\mathbb{C}^3)$ by Theorem 3.1, so $\Psi_l(\pi(P)) = P \circ v = F$. For the formula, let G be homogeneous of degree d . Then

$$\Delta(q^j G) = 2j(2j+2d+1) q^{j-1} G + q^j \Delta G.$$

Put $H = \sum_j \alpha_j q^j \Delta^j P$ with $\alpha_0 = 1$. Applying the identity with $G = \Delta^j P$ and $d = l - 2j$ gives

$$\Delta H = \sum_j \beta_j q^j \Delta^{j+1} P,$$

with $\beta_j = \alpha_j + 2(j+1)(2l-2j-1)\alpha_{j+1}$. The stated coefficients satisfy $\alpha_{j+1} = -\alpha_j/(2(j+1)(2l-2j-1))$, so $\beta_j = 0$ and $\Delta H = 0$. Since $H - P \in qP_{l-2}(\mathbb{C}^3)$, the uniqueness in Theorem 3.1 gives $H = \pi(P)$. Finally $fg - \pi(fg) \in qP_{l+m-2}(\mathbb{C}^3)$, so $\Psi_{l+m}(\pi(fg)) = (fg) \circ v = (f \circ v)(g \circ v)$.

(ii) The statements on ρ are Theorem 3.4. By (10), $v(g^{-1}z) = \rho(g)^{-1}v(z)$, so $\Psi_l(\rho(g) \cdot f)(z) = f(\rho(g)^{-1}v(z)) = f(v(g^{-1}z)) = (g \cdot \Psi_l f)(z)$. Uniqueness up to a scalar is Schur's lemma, since Sym^{2l} is irreducible.

(iii) Since $\rho(g_\theta) = R_{-\theta}$ and $\mathcal{Y}_{lm}$ depends on ϕ through $e^{im\phi}$, $\rho(g_\theta) \cdot \mathcal{Y}_{lm} = e^{im\theta} \mathcal{Y}_{lm}$. By (ii), $\Psi_l(\mathcal{Y}_{lm})$ is an eigenvector of g_θ with eigenvalue $e^{im\theta}$, and $g_\theta \cdot z_0^p z_1^{2l-p} = e^{i\theta(l-p)} z_0^p z_1^{2l-p}$; the characters are distinct, so $\Psi_l(\mathcal{Y}_{lm})$ is a multiple of $z_0^{l-m} z_1^{l+m}$. Let $p = |m|$. The homogeneous extension of $P_l^p(\cos \theta) e^{\pm ip\phi}$ is $(x_1 \pm ix_2)^p G_{lp}$, where $G_{lp} = |x|^{l-p} P_l^{(p)}(x_3/|x|)$ is a polynomial in x_3 and q . Its part free of q is $\lambda_{lp} x_3^{l-p}$, where $\lambda_{lp} = (2l)!/(2^l l! (l-p)!)$ is the leading coefficient of $P_l^{(p)}$. By (11), $x_1 + ix_2 \mapsto -2z_1^2$, $x_1 - ix_2 \mapsto 2z_0^2$, $x_3 \mapsto 2z_0 z_1$ and $q \mapsto 0$. Hence

$$\Psi_l((x_1 \pm ix_2)^p G_{lp}) = \frac{(\mp 1)^p (2l)!}{l! (l-p)!} z_0^{l \mp p} z_1^{l \pm p}.$$

By the definitions of Section 2, $Y_{lp} \pm iY_{l,-p} = N_{lp}(x_1 \pm ix_2)^p G_{lp}$ for $p \geq 1$ and $Y_{l0} = \sqrt{2l+1} G_{l0}$. Multiplying by the factors in the definition of $\mathcal{Y}_{lm}$ gives K_{lm} . Next, $f = \sum_m \eta_m \mathcal{Y}_{lm}$ with $\eta_0 = c_{l0}$, $\eta_p = (-1)^p (c_{lp} - ic_{l,-p})/\sqrt{2}$ and $\eta_{-p} = (c_{lp} + ic_{l,-p})/\sqrt{2}$. Comparing coefficients gives $a_{l+m} = K_{lm} \eta_m / \binom{2l}{l+m}$ for every $f = \sum_m \eta_m \mathcal{Y}_{lm} \in \mathcal{H}_l^{\mathbb{C}}$. Since $K_{lm} / \binom{2l}{l+m} = \sqrt{(2l+1)(l-m)!(l+m)!}/l!$, this is the formula for η_m . Moreover $K_{lp}/(\sqrt{2} \binom{2l}{l+p}) = \nu_{lp}$. This gives the formulas for a_k . Since the c_{lm} are real, the formulas can be inverted as stated.

(iv) Write $f = \sum_m \eta_m \mathcal{Y}_{lm}$. By (iii), $a_{l+m} = K_{lm} \eta_m / \binom{2l}{l+m}$, and $K_{lm}^2 / \binom{2l}{l+m} = (2l+1) \binom{2l}{l}$ for every m . Summing over m and using the orthonormality of the $\mathcal{Y}_{lm}$ gives the first identity. By (18) with $F = G$ and $k = 2l$, the $(2l)$ -th transvectant is $(F, F)_{2l} = \sum_k (-1)^k \binom{2l}{k} a_k a_{2l-k}$. For real f , (v) gives $a_{2l-k} = (-1)^{l+k} \bar{a}_k$, hence $(F_f, F_f)_{2l} = (-1)^l \|F_f\|_{\text{BW}}^2$. Since Ψ_l is a linear isomorphism, the norm identity and polarization show that $\hat{\Psi}_l$ is unitary.

(v) This is Theorem 3.6(i) together with (14). $\square$

Remark 3.9. For $l = 1, 2, 3$, Theorem 3.8(iv) expresses the band power through the classical invariants. For $l = 1$ one has $\text{disc}(Q_a) = -2(Q_a, Q_a)_2$, so $\|f_1\|^2 = \text{disc}(Q_a)/12$. For $l = 2$ one has $(F, F)_4 = 2I$ with I as in (19), so $\|f_2\|^2 = I(F_{f_2})/15$. For $l = 3$ direct expansion gives $I_2 = -120(F, F)_6$ with I_2 as in (20), so $J_2(F_{f_3}) = 2100\|f_3\|^2$. In particular $J_2 > 0$ on every nonzero real band-three harmonic. The squared band norms $\|f_l\|^2$, $l \leq L$, which determine the power spectrum, are therefore the phased quadratic invariants of the forms F_{f_l} , up to the constants of Theorem 3.8(iv). The other invariants of Section 4 refine it. The paper uses Ψ_l , which keeps the classical constants of the invariants. The unitary map $\hat{\Psi}_l$ is the normalization for comparing bands numerically.

Corollary 3.10 (Maxwell–Sylvester root correspondence). *Let $l \geq 0$ and let $f \in \mathcal{H}_l$ be real and nonzero. Then $F_f = \lambda Q_{a_1} \cdots Q_{a_l}$ with $\lambda \in \mathbb{R}^\times$ and $a_k \in \mathbb{R}^3 \setminus \{0\}$, and $f = \pi(\lambda \langle a_1, x \rangle \cdots \langle a_l, x \rangle)$. The root divisor of F_f , transported to $\mathbb{S}^2$ by $\bar{m}$, consists of the l antipodal pairs $\pm a_k/|a_k|$. Every effective antipodally invariant divisor of degree $2l$ on $\mathbb{S}^2$, counted with multiplicity, is the root divisor of F_f for a nonzero*

real f , unique up to $\mathbb{R}^\times$. For $R \in \mathrm{SO}(3)$ the roots of $F_{R \cdot f}$ are the images under R of the roots of F_f . For $-1 \in \mathrm{O}(3)$ one has $F_{(-1) \cdot f} = (-1)^l F_f$.

Proof. The factorization is Theorem 3.6(ii), and the harmonic form of it is (12). The roots of Q_a correspond to $\pm a/|a|$ by Theorem 3.3(ii). Let D be an effective antipodally invariant divisor of degree $2l$. Choose a_k with D the sum of the pairs $\pm a_k/|a_k|$. Then $\prod_k Q_{a_k}$ is σ_l -fixed, so it equals F_f for a real f by Theorem 3.8(v). A form with root divisor D is $c \prod_k Q_{a_k}$ with $c \in \mathbb{C}^\times$, and it is σ_l -fixed if and only if $c \in \mathbb{R}$. If $R = \rho(g)$ with $g \in \mathrm{SU}(2)$, then $F_{R \cdot f} = g \cdot F_f$ by Theorem 3.8(ii), whose roots are the points $[gz]$ with $F_f(z) = 0$, and $\bar{m}([gz]) = R \bar{m}([z])$ by Theorem 3.4. The last statement is Theorem 3.6(iv). $\square$

Corollary 3.11 (Invariant and orbit transfer). *Let $L \geq 0$. Composition with Ψ identifies the ring*

$$\mathcal{R}_L = \mathbb{C} \left[\bigoplus_{l \leq L} \mathrm{Sym}^{2l} \right]^{\mathrm{SL}_2(\mathbb{C})}$$

with $\mathbb{R}[V_L]^{\mathrm{SO}(3)} \otimes \mathbb{C}$. Let $I \in \mathcal{R}_L$ have real coefficients and multidegree (d_l) , and put $p(I) = \sum_l l d_l$ and $w(I) = \sum_l d_l$. On real tuples $I \circ \Psi$ takes values in $i^{p(I)} \mathbb{R}$. Every reflection multiplies it by $(-1)^{p(I)}$, and the dilation $c \mapsto tc$ multiplies it by $t^{w(I)}$. For $c, c' \in V_L$ the following are equivalent:

- (a) $c' \in \mathrm{SO}(3) \cdot c$;
- (b) $\Psi(c') \in \mathrm{SL}_2(\mathbb{C}) \cdot \Psi(c)$;
- (c) $I(\Psi(c')) = I(\Psi(c))$ for every $I \in \mathcal{R}_L$.

Proof. By Theorem 3.8(ii) and the Zariski density of $\mathrm{SU}(2)$ in $\mathrm{SL}_2(\mathbb{C})$, a polynomial I on $\bigoplus_l \mathrm{Sym}^{2l}$ is $\mathrm{SL}_2(\mathbb{C})$ -invariant if and only if $I \circ \Psi$ is $\mathrm{SO}(3)$ -invariant. Since $\mathrm{SO}(3)$ acts on V_L by real matrices, the complex-valued $\mathrm{SO}(3)$ -invariant polynomials are $\mathbb{R}[V_L]^{\mathrm{SO}(3)} \otimes \mathbb{C}$. The reality and reflection statements are Theorem 3.6(iii) and (iv). The scale statement holds because I is multihomogeneous and Ψ is linear. Condition (a) implies (b) because $\mathrm{SO}(3) = \rho(\mathrm{SU}(2))$, and (b) implies (c) by invariance. If (c) holds, then every real $\mathrm{SO}(3)$ -invariant polynomial takes the same value at c and c' . Real polynomial invariants of a compact group separate its orbits [27, 26], so (a) holds. $\square$

Proposition 3.12 (Transvectants as Clebsch–Gordan maps). *Let $l, m \geq 0$ and $0 \leq r \leq 2 \min(l, m)$. For $f \in \mathcal{H}_l^\mathbb{C}$ and $g \in \mathcal{H}_m^\mathbb{C}$ put*

$$f \star_r g = \Psi_{l+m-r}^{-1}((F_f, F_g)_r) \in \mathcal{H}_{l+m-r}^\mathbb{C}.$$

The map $(f, g) \mapsto f \star_r g$ is bilinear, nonzero and $\mathrm{SO}_3(\mathbb{C})$ -equivariant. Every $\mathrm{SO}_3(\mathbb{C})$ -equivariant bilinear map $\mathcal{H}_l^\mathbb{C} \times \mathcal{H}_m^\mathbb{C} \rightarrow \mathcal{H}_{l+m-r}^\mathbb{C}$ is a scalar multiple of it. For $r = 0$ one has $f \star_0 g = \pi(fg)$.

Proof. The transvectant (18) is bilinear, and it is $\mathrm{SL}_2(\mathbb{C})$ -equivariant with values in $\mathrm{Sym}^{2l+2m-2r}$ [25]. By Theorem 3.8(i) and (ii), $\star_r$ is bilinear and $\mathrm{SO}_3(\mathbb{C})$ -equivariant. It is nonzero, since (18) gives $(z_0^{2l}, z_1^{2m})_r = z_0^{2l-r} z_1^{2m-r}$. The representation $\mathcal{H}_l^\mathbb{C} \otimes \mathcal{H}_m^\mathbb{C}$ of $\mathrm{SO}_3(\mathbb{C})$ is the direct sum of the $\mathcal{H}_j^\mathbb{C}$ with $|l-m| \leq j \leq l+m$, each with multiplicity one [31]. For $j = l+m-r$, Schur's lemma gives the uniqueness. For $r = 0$, (18) gives $(F, G)_0 = FG$, and Theorem 3.8(i) gives $f \star_0 g = \pi(fg)$. $\square$

TABLE 1. The dictionary of Theorem 3.8. Roman numerals refer to its parts.

harmonic side	binary-form side	
$\mathcal{H}_l^{\mathbb{C}}$ with basis $\mathcal{Y}_{lm}$	Sym^{2l} with basis $z_0^{l-m} z_1^{l+m}$	(i), (iii)
real harmonics $\mathcal{H}_l$	$\text{Fix}(\sigma_l): a_{2l-k} = (-1)^{l+k} \bar{a}_k$	(v)
harmonic part $\pi(fg)$ of a product	product $F_f F_g$	(i)
Clebsch–Gordan map $\mathcal{H}_l \otimes \mathcal{H}_m \rightarrow \mathcal{H}_{l+m-r}$	transvectant $(F_f, F_g)_r$	Theorem 3.12
rotation $\rho(g) \in \text{SO}(3)$	substitution by $g \in \text{SU}(2)$	(ii)
reflection $-1 \in \text{O}(3)$	$F \mapsto (-1)^l F$	Theorem 3.10
Maxwell axis $\mathbb{R}a$	antipodal root pair $\pm a/ a $	Theorem 3.10
band power $\ f\ ^2$	$(-1)^l (F_f, F_f)_{2l} / ((2l+1) \binom{2l}{l})$	(iv)
$\ f_1\ ^2, \ f_2\ ^2, \ f_3\ ^2$	$\text{disc}(Q_a)/12, I/15, J_2/2100$	Theorem 3.9
$\text{O}(3)$ -invariant, pseudo-invariant	$\sum_l l d_l$ even, odd	Theorem 3.11
weight under dilation	$\sum_l d_l$	Theorem 3.11
$\text{SO}(3)$ -orbits of real tuples	$\text{SL}_2(\mathbb{C})$ -orbits, intersected with the σ -fixed locus	Theorem 3.11
achiral band-three harmonic	$R(F) = 0$; if $J_{10} \neq 0$, the curve lies in $\mathcal{L}_2$	Theorem 5.2
coordinate band $\mathcal{H}_l \otimes \mathbb{R}^3$	$\text{Sym}^{2l} \otimes \text{Sym}^2$, bidegree $(2l, 2)$	Theorem 4.3

Remark 3.13. Parts (i), (ii) and (v) of Theorem 3.8 and Theorem 3.10 are classical: the parameterization of the null conic and the spin covering go back to Cartan [10], $\mathcal{H}_l^{\mathbb{C}} \cong \text{Sym}^{2l}$ is the standard realization of the irreducible representations of $\text{SU}(2)$ [32, 31], and Theorem 3.10 is the theorem of Maxwell and Sylvester [23, 30]. What is specific here is the pairing with the quaternionic real structure, which makes the parity rule of Theorem 3.6 available, and the explicit normalizations of (iii) and (iv). Two cautions apply. A form in $\text{SL}_2(\mathbb{C}) \cdot F_f$ is in general not σ_l -fixed; the rotations of $\mathbb{S}^2$ are the substitutions by $\text{SU}(2)$. And the shape quotient removes one positive scale common to all bands, whereas independent complex rescalings of the bands, as in the moduli of binary forms, forget the amplitude, the relative normalization of the bands and the sign of an even band.

From Section 4 on, the rotation quotient of truncated harmonic data is the joint invariant theory of the tuple $(F_{f_0}, \dots, F_{f_L})$ of binary forms of degrees $0, 2, \dots, 2L$. Invariants of the separate forms do not record the relative orientation of the bands; joint invariants are needed for that (Section 4.4).

4. INVARIANTS AND SEPARATION

Fix $L \geq 1$. The **radial descriptor** of a surface is the tuple $c = f^{(L)}(\varphi)$ of Theorem 2.3; it lives in $V_L = \bigoplus_{l=0}^L \mathcal{H}_l$ with $\text{SO}(3)$ acting diagonally. The components $f_1 \in \mathcal{H}_1$ and $f_2 \in \mathcal{H}_2$ are called the **dipole** and the **quadrupole**, and the tuple of band norms $(\|f_l\|)_{l \leq L}$ is the **power spectrum** [21]. Through $\Psi = \bigoplus \Psi_l$, $V_L \otimes \mathbb{C}$ is the space of tuples of binary forms of degrees $0, 2, \dots, 2L$ with $\text{SL}_2(\mathbb{C})$ acting

diagonally, and by Theorem 3.8 and the Zariski density of $\mathrm{SU}(2)$ in $\mathrm{SL}_2(\mathbb{C})$,

$$(15) \quad \begin{aligned} \mathbb{R}[V_L]^{\mathrm{SO}(3)} \otimes \mathbb{C} &\cong \mathcal{R}_L \\ &:= \mathbb{C} \left[\bigoplus_{l \leq L} \mathrm{Sym}^{2l} \right]^{\mathrm{SL}_2(\mathbb{C})}, \end{aligned}$$

the ring of joint invariants of even binary forms. It is finitely generated [18], with multihomogeneous generators having rational coefficients. By Theorem 3.6(iii), multiplying a generator by $i^{-\sum l d_l}$ gives a real invariant on V_L ; these **phased** generators generate $\mathbb{R}[V_L]^{\mathrm{SO}(3)}$.

Three notions must be kept apart: a **generating set** generates the invariant algebra; a **separating set** distinguishes orbits; a **feature pool** is any list of invariants. Separation, not generation, is what a complete descriptor needs, and separating sets can be much smaller than generating sets.

Theorem 4.1. *Let $I_1, \dots, I_N$ be the phased real forms of a multihomogeneous generating set of $\mathcal{R}_L$, of total degrees $w_1, \dots, w_N$, and $\mathcal{I} = (I_1, \dots, I_N): V_L \rightarrow \mathbb{R}^N$ the corresponding map, the **Hilbert map** of the generating set. Then $\mathcal{I}(c) = \mathcal{I}(c')$ iff $c' = R \cdot c$ for some $R \in \mathrm{SO}(3)$; $\mathcal{I}(c) = 0$ iff $c = 0$; and $\mathcal{I}(V_L)$ is a closed semialgebraic set homeomorphic to $V_L/\mathrm{SO}(3)$. The same conclusions hold for any set of real invariants that generates $\mathbb{R}[V_L]^{\mathrm{SO}(3)}$, and the separation statement holds for any separating subset.*

Proof. For a compact group acting linearly on a real vector space, real polynomial invariants separate orbits, and the image of the Hilbert map of a finite generating set is closed, semialgebraic, and homeomorphic to the quotient [27, 26]. The phased generators generate the real invariant ring by (15). Since the orbit $\{0\}$ is closed and the I_k have positive degree, $\mathcal{I}^{-1}(0) = \{0\}$. $\square$

Remark 4.2. The theorem has a consequence for the data augmentation used in SPHARM-COM, which formed $c_{\text{new}} = (1 - w)c_{\text{src}} + wc_{\text{tgt}}$ within a class. The invariants of c_{new} are not functions of $\mathcal{I}(c_{\text{src}})$ and $\mathcal{I}(c_{\text{tgt}})$; they depend on the relative orientation of the two representatives, which is exactly what $\mathcal{I}$ discards and the pipeline leaves undetermined. Nor is linear interpolation of invariant coordinates intrinsic: already for a single band of degree two the invariant image is not convex (Theorem 4.8). An intrinsic augmentation must be defined on the quotient, for instance by aligning one representative to the other by the orbit distance of Section 7 before interpolating.

4.1. The coordinate model. Let $W_L = \bigoplus_{l=1}^L \mathcal{H}_l \otimes \mathbb{R}^3$; band zero is omitted since $C_0 = 0$. Let $Y_l(u)$ be an orthonormal column basis of $\mathcal{H}_l$ for $d\omega$, $X_l(u) = C_l Y_l(u)$ with $C_l \in \mathbb{R}^{3 \times (2l+1)}$, and define $D_l(Q)$ by $Y_l(Q^{-1}u) = D_l(Q)^\top Y_l(u)$; D_l is a homomorphism into $\mathrm{SO}(2l+1)$. Then (5) reads

$$(16) \quad (Q, R) \cdot C_l = R C_l D_l(Q)^\top.$$

Complexifying with Ψ_l on the sphere side and $\mathbb{C}^3 \cong \mathrm{Sym}^2$ via (8) on the ambient side, $W_L \otimes \mathbb{C}$ becomes families of double binary forms of bidegrees $(2l, 2)$ with $\mathrm{SL}_2 \times \mathrm{SL}_2$ acting by substitution in z and w [25]. The real structure is $\sigma_l \otimes \sigma_1$.

Corollary 4.3 (Coordinate model). *Let $l \geq 1$. For $X = (X_1, X_2, X_3) \in \mathcal{H}_l^{\mathbb{C}} \otimes \mathbb{C}^3$ put*

$$\begin{aligned} B_X(z, w) &= \langle X(v(z)), v(w) \rangle \\ &= \sum_k X_k(v(z)) v_k(w). \end{aligned}$$

Then $X \mapsto B_X$ is a linear isomorphism from $\mathcal{H}_l^{\mathbb{C}} \otimes \mathbb{C}^3$ onto $\text{Sym}^{2l} \otimes \text{Sym}^2$, the space of double binary forms of bidegree $(2l, 2)$. Let the two-sided action (5) be extended to $\text{SO}_3(\mathbb{C}) \times \text{SO}_3(\mathbb{C})$ by the same formula. For $g, h \in \text{SL}_2(\mathbb{C})$,

$$B_{(\rho(g), \rho(h)) \cdot X}(z, w) = B_X(g^{-1}z, h^{-1}w).$$

The tuple X is real if and only if B_X is fixed by $\sigma_l \otimes \sigma_1$.

Proof. One has $B_X = \sum_k \Psi_l(X_k) \otimes v_k$. Since Ψ_l is an isomorphism and v_1, v_2, v_3 is a basis of Sym^2 (Theorem 3.2(i)), the map $X \mapsto B_X$ is an isomorphism. Let $Q = \rho(g)$ and $R = \rho(h)$. By (10), $Q^{-1}v(z) = v(g^{-1}z)$ and $R^\top v(w) = R^{-1}v(w) = v(h^{-1}w)$. Hence

$$\begin{aligned} B_{(Q, R) \cdot X}(z, w) &= \langle R X(Q^{-1}v(z)), v(w) \rangle \\ &= \langle X(v(g^{-1}z)), R^\top v(w) \rangle \\ &= B_X(g^{-1}z, h^{-1}w). \end{aligned}$$

Each $v_k = \Psi_1(x_k)$ is σ_1 -fixed by Theorem 3.6(i). Hence $(\sigma_l \otimes \sigma_1)(B_X) = \sum_k \sigma_l(\Psi_l X_k) \otimes v_k = B_{\bar{X}}$ by Theorem 3.6(i), and B_X is fixed if and only if $\bar{X} = X$. $\square$

Lemma 4.4. *Let I be an $\text{SL}_2 \times \text{SL}_2$ -invariant on $\bigoplus_{l \leq L} \text{Sym}^{2l} \otimes \text{Sym}^2$ with rational coefficients, of degree d_l in the $(2l, 2)$ summand. On real families, $I((C_l)) = (-1)^{\sum_l (l+1)d_l} \bar{I}((\bar{C}_l))$. The mirror image of a surface has coordinate descriptor in the $\text{SO}(3) \times \text{SO}(3)$ -orbit of $((-1)^{l+1} C_l)_l$, so the real-valued invariants are exactly the mirror-invariant ones and the purely imaginary ones are the mirror pseudo-invariants.*

Proof. As in Theorem 3.6(iii), $\sigma_l \otimes \sigma_1$ is conjugation composed with $(-1)^{l+1}(w, w')$ for the pair of substitutions $(z_0, z_1) \mapsto (-z_1, z_0)$, $(w_0, w_1) \mapsto (-w_1, w_0)$. For the mirror image, a reflection of $\mathbb{S}^2$ is $-Q$ and acts on $\mathcal{H}_l$ by $(-1)^l D_l(Q)$, and the ambient reflection $-R$ contributes a further -1 . $\square$

Theorem 4.5. *Let $\mathcal{I}_2: W_L \rightarrow \mathbb{R}^N$ be given by the phased real forms of a multi-homogeneous generating set of the $\text{SL}_2 \times \text{SL}_2$ -invariants. Then $\mathcal{I}_2(C) = \mathcal{I}_2(C')$ iff $C' = (Q, R) \cdot C$ for some $(Q, R) \in \text{SO}(3) \times \text{SO}(3)$, $\mathcal{I}_2^{-1}(0) = \{0\}$, and $\mathcal{I}_2(W_L)$ is closed and semialgebraic. If two surfaces have $\mathcal{I}_2(C^{(L)}(\varphi)) = \mathcal{I}_2(C^{(L)}(\varphi'))$ for every L , with $\mathcal{I}_2$ chosen for each L , they differ by an orientation-preserving rigid motion.*

Proof. The first assertions follow as in Theorem 4.1, with Theorem 4.4. For the last, for each L there is (Q_L, R_L) with $C'_l = (Q_L, R_L) \cdot C_l$ for all $l \leq L$. By compactness a subsequence converges to (Q, R) , and for fixed l the relation passes to the limit, giving $C'_l = (Q, R) \cdot C_l$ for all l with one pair. Completeness of harmonics in $L^2(\mathbb{S}^2, d\omega)$ gives $X_{\varphi'} = R X_\varphi \circ Q^{-1}$ a.e., hence everywhere, and Theorem 2.3 finishes. $\square$

Write $\mathcal{Q}_L = W_L/(\mathrm{SO}(3) \times \mathrm{SO}(3))$. The band-forgetting projections are equivariant and induce $\mathcal{Q}_{L+1} \rightarrow \mathcal{Q}_L$; a surface determines a compatible sequence of points, and the theorem says the sequence determines the surface. For a linear action of a compact group G on V , the points whose stabilizer is conjugate to the minimal stabilizer form an open dense G -invariant subset of V , the **principal stratum**, on which the quotient is a manifold of dimension $\dim V - \dim G + \dim(\text{minimal stabilizer})$ [27]. The dimension of $\mathcal{Q}_L$ must not be confused with the size of a descriptor: on the principal stratum $\dim \mathcal{Q}_L = \dim W_L - 6$, but $\mathcal{Q}_L$ is a semialgebraic space with singular strata, the frame coordinates of Section 6 do not extend across them (Theorem 6.2(c)), and a separating set of polynomial invariants is in general larger. Section 6 quantifies this.

Proposition 4.6. *In the orthonormal basis $e_i = \sqrt{3}x_i$ of $\mathcal{H}_1$, C_1 is a real 3×3 matrix M , $D_1(Q) = Q$, and the action is $M \mapsto RMQ^\top$. The invariants $\mathrm{tr}(MM^\top)$, $\mathrm{tr}((MM^\top)^2)$, $\det M$ of degrees 2, 4, 3 separate $\mathrm{SO}(3) \times \mathrm{SO}(3)$ -orbits, $\dim(\mathcal{H}_1 \otimes \mathbb{R}^3)/(\mathrm{SO}(3) \times \mathrm{SO}(3)) = 3$, and all three are mirror-invariant.*

Proof. Signed SVD: $M = R\Sigma Q^\top$ with $R, Q \in \mathrm{SO}(3)$, $\Sigma = \mathrm{diag}(\sigma_1, \sigma_2, \pm\sigma_3)$, $\sigma_1 \geq \sigma_2 \geq \sigma_3 \geq 0$, the sign being $\mathrm{sgn} \det M$. With $p_1 = \mathrm{tr} MM^\top$, $p_2 = \mathrm{tr}(MM^\top)^2$, $d = \det M$, the σ_i^2 are the roots of $t^3 - p_1 t^2 + \frac{1}{2}(p_1^2 - p_2)t - d^2$, and $\mathrm{diag}(\sigma_1, \sigma_2, \mathrm{sgn}(d)\sigma_3)$ is a canonical representative. The generic stabilizer is finite, so the quotient has dimension $9 - 6 = 3$. By Theorem 4.4 with $l = 1$ every invariant has even parity. $\square$

For $l \geq 2$ the Gram matrices $G_l = C_l C_l^\top$ are invariant under the sphere factor and transform by conjugation under the ambient factor, so traces of words in $G_1, \dots, G_L$ are two-sided invariants; the simplest are $\mathrm{tr} G_l$, $\mathrm{tr} G_l^2$, $\det G_l$ and $\mathrm{tr}(G_l G_{l'})$. They are the two-sided analogue of the power spectrum and discard everything the sphere factor sees beyond degree two. A complete polynomial separating set for W_L , $L \geq 2$, of manageable size is not in hand; Section 6 gives an algebraic separating set on the principal stratum and a dimension bound for continuous polynomial ones.

4.2. Scale. A dilation $c \mapsto \lambda c$, $\lambda > 0$, multiplies an invariant of total degree w_k by λ^{w_k} , so the class of $\mathcal{I}(c)$ in the positive weighted projectivization

$$\begin{aligned} \mathbb{P}_w^+(\mathbb{R}) &= (\mathbb{R}^N \setminus 0)/\mathbb{R}_{>0}, \\ \lambda \cdot x &= (\lambda^{w_1} x_1, \dots, \lambda^{w_N} x_N), \end{aligned}$$

is invariant under rotation and positive scaling. We also write $\mathbb{P}_w(\mathbb{R}) = (\mathbb{R}^N \setminus 0)/\mathbb{R}^\times$ for the quotient by all nonzero scalars, with the same action. When $\mathcal{I}$ separates, it is complete on nonzero truncations: equal classes mean $I_k(c') = \lambda^{w_k} I_k(c) = I_k(\lambda c)$, so $c' \sim \lambda c$. We quotient by $\mathbb{R}_{>0}$ rather than $\mathbb{R}^\times$ because odd weights would otherwise identify c with $-c$, which on odd bands agrees with the mirror image. Scale is split off, not discarded: the mean radius f_0 , or a coefficient norm, is retained as a size variable. A canonical representative is given by the weighted sup-gauge

$$(17) \quad H_w(x) = \max_k |x_k|^{1/w_k}, \quad \hat{x}_k = \frac{x_k}{H_w(x)^{w_k}},$$

so that $\max_k |\hat{x}_k|^{1/w_k} = 1$; H_w is the archimedean factor of the weighted height [5]. It preserves all signs, including those of pseudo-invariants; smooth alternatives $(\sum_k |x_k|^{p/w_k})^{1/p}$ with p/w_k even are available. The bounded quantity $|\hat{I}_k|^{1/w_k} \in [0, 1]$ is a coordinatewise diagnostic of approach to the divisor $I_k = 0$. Absolute invariants such as J^2/I^3 for the quartic are the classical alternative on charts;

the normalized coordinates (17) are preferred for computation because they are bounded, global, and sign-preserving.

4.3. Counting. $\dim V_L = (L + 1)^2$. For $L \geq 2$ the generic stabilizer is finite, so $\dim V_L/\mathrm{SO}(3) = (L + 1)^2 - 3$, one less after scale; for $L = 1$ a generic dipole has stabilizer $\mathrm{SO}(2)$ and the quotient has dimension 2. A single band $\mathcal{H}_l$, $l \geq 2$, has quotient dimension $2l - 2$. Band-by-band invariants therefore account for $2 + \sum_{l=2}^L (2l - 2) = L^2 - L + 2$ dimensions out of $L^2 + 2L - 2$, and the deficit $3L - 4$ is carried by joint invariants recording the relative orientations of the bands. For the coordinate model $\dim W_L = 3((L + 1)^2 - 1)$ and $\dim \mathcal{Q}_L = 3(L + 1)^2 - 9$. At $L = 6$, the reference case of Section 10, the counts are 46 and 138.

4.4. Band-by-band invariants. We write forms in binomial normalization $F = \sum_k \binom{2l}{k} a_k z_0^{2l-k} z_1^k$. For $F \in \mathrm{Sym}^m$, $G \in \mathrm{Sym}^n$ and $0 \leq k \leq \min(m, n)$, the k -th **transvectant** is

$$(18) \quad (F, G)_k = \frac{(m - k)!(n - k)!}{m!n!} \times \sum_{i=0}^k (-1)^i \binom{k}{i} \frac{\partial^k F}{\partial z_0^{k-i} \partial z_1^i} \frac{\partial^k G}{\partial z_0^i \partial z_1^{k-i}},$$

It is bilinear, $\mathrm{SL}_2(\mathbb{C})$ -equivariant, with rational coefficients, of degree $m + n - 2k$, and $(F, F)_k = 0$ for odd k . For $k = m = n$ it is an invariant. Theorem 3.6(iii) applies to everything built from transvectants.

Band l is a binary form of degree $2l$, so the invariants available for band l are the classical invariants of binary forms of degree $2l$ [25]. The ring is finitely generated [18], and explicit generators are known in low degree: for Sym^2 the discriminant; for Sym^4 the invariants I, J below; for Sym^6 the Igusa–Clebsch invariants of degrees 2, 4, 6, 10 and a skew invariant of degree 15 [12, 19]. We use $l \leq 3$. For larger l we do not rely on a generating set: the frame coordinates of Section 6 and the continuous invariants of Theorem 6.5 are uniform in l .

Remark 4.7. If $F \in \mathrm{Sym}^{2l}$ has distinct roots, $y^2 = F(z, 1)$ is a smooth hyperelliptic curve of genus $l - 1$, and two such curves are isomorphic iff the forms lie in one $\mathrm{GL}_2(\mathbb{C})$ -orbit [22]. By Theorem 3.10, F_f has distinct roots iff the Maxwell axes of f are pairwise distinct. For $l = 3$ the curve has genus two, and its isomorphism class is the point $[J_2 : J_4 : J_6 : J_{10}]$ of $\mathcal{M}_2$ introduced below [19]; Theorem 5.2 uses this.

Degree two. A real harmonic is $f = x^\top Q x$ with Q traceless symmetric, and F_f is a binary quartic whose invariant ring is freely generated by

$$(19) \quad I = a_0 a_4 - 4a_1 a_3 + 3a_2^2, \quad J = \det \begin{pmatrix} a_0 & a_1 & a_2 \\ a_1 & a_2 & a_3 \\ a_2 & a_3 & a_4 \end{pmatrix},$$

of degrees 2, 3, with $\mathrm{disc} F = I^3 - 27J^2$ the normalized discriminant. The quartic has a repeated root if and only if $\mathrm{disc} F = 0$.

Example 4.8. For $Q = \mathrm{diag}(\lambda_1, \lambda_2, \lambda_3)$, $\sum \lambda_k = 0$, one finds $a_0 = a_4 = \lambda_1 - \lambda_2$, $a_1 = a_3 = 0$, $a_2 = \lambda_3$, hence

$$I(F_f) = 2 \mathrm{tr} Q^2, \quad J(F_f) = -4 \det Q = -\frac{4}{3} \mathrm{tr} Q^3,$$

both real as Theorem 3.6(iii) predicts. The image of $\mathcal{H}_2$ in the (I, J) -plane is $\{I \geq 0, I^3 \geq 27J^2\}$, since the discriminant of $t^3 - \frac{1}{2}(\text{tr } Q^2)t - \det Q$ is $(I^3 - 27J^2)/16$. This region is not convex: the midpoint of $(1, c)$ and $(4, 8c)$, $c = 27^{-1/2}$, lies outside it. As a point of $\mathbb{P}_{(2,3)}^+(\mathbb{R})$ the band-two shape is $[I : J]$ and the region maps onto a closed interval whose endpoints $I^3 = 27J^2$ are the axially symmetric quadrupoles. Figure 2 shows the region.

Degree two with the lower bands: a complete descriptor. The first case where band-by-band invariants are insufficient is $L = 2$, where the deficit $3L - 4 = 2$ must be carried by joint invariants. We give a complete answer.

Theorem 4.9 (Separation for V_2). *Write $c = (f_0, a, Q) \in V_2 = \mathcal{H}_0 \oplus \mathcal{H}_1 \oplus \mathcal{H}_2$ with $f_0 \in \mathbb{R}$, $a \in \mathbb{R}^3$, and $Q \in \text{Sym}_0^2(\mathbb{R}^3)$, the traceless symmetric 3×3 real matrices. The seven invariants*

$$\begin{aligned} f_0, \quad |a|^2, \quad \text{tr } Q^2, \quad \det Q, \quad a^\top Q a, \quad a^\top Q^2 a, \\ \varepsilon := \det(a, Qa, Q^2 a), \end{aligned}$$

of degrees 1, 2, 2, 3, 3, 4, 6, separate the $\text{SO}(3)$ -orbits in V_2 . The first six are $\text{O}(3)$ -invariants and ε is a pseudo-invariant. They satisfy the relation $\varepsilon^2 = \det \text{Gram}(a, Qa, Q^2 a)$, whose right side is a polynomial in the other six by $Q^3 = \frac{1}{2}(\text{tr } Q^2)Q + (\det Q)1$. In binary-form language $|a|^2 = \frac{1}{4} \text{disc } Q_a$, $\text{tr } Q^2$ and $\det Q$ are I and J of Theorem 4.8 up to constants, $a^\top Q a$ is proportional to $(Q_a^2, F)_4$, and $a^\top Q^2 a$ and ε are joint invariants of bidegrees $(2, 2)$ and $(3, 3)$ in (Q_a, F) .

Proof. Suppose the seven invariants agree on c and c' . Since Q is traceless, $\text{tr } Q^2$ and $\det Q$ determine its characteristic polynomial, hence its spectrum, and symmetric matrices with the same spectrum are $\text{SO}(3)$ -conjugate; so after applying a rotation to c' we may assume $Q' = Q = \text{diag}(\lambda_1, \lambda_2, \lambda_3)$. It remains to move a' to a by an element of $\text{Stab}(Q)$.

If the λ_i are distinct, $\text{Stab}(Q) = K$, the group $\{\text{diag}(\epsilon_1, \epsilon_2, \epsilon_3) : \epsilon_i = \pm 1, \prod \epsilon_i = 1\}$. The equalities $\sum_i a_i^2 \lambda_i^k = \sum_i a_i'^2 \lambda_i^k$ for $k = 0, 1, 2$ form a Vandermonde system, so $a_i^2 = a_i'^2$ and $a_i' = \epsilon_i a_i$ for some signs, which may be chosen freely where $a_i = 0$. If some $a_i = 0$, choose the free sign so that $\prod \epsilon_i = 1$; then $\text{diag}(\epsilon) \in K$ maps a to a' . If all $a_i \neq 0$ and $\prod \epsilon_i = -1$, then $\varepsilon(c') = a_1' a_2' a_3' \prod_{i < j} (\lambda_j - \lambda_i) = -\varepsilon(c) \neq 0$, contradicting $\varepsilon(c) = \varepsilon(c')$. Hence $\prod \epsilon_i = 1$ and $a' \in K \cdot a$.

If $\lambda_1 = \lambda_2 \neq \lambda_3$, then $\text{Stab}(Q) \supset \text{SO}(2)$ acting on the (e_1, e_2) -plane together with the half-turn about e_1 . From $|a|^2$ and $a^\top Q a$ we get $a_1^2 + a_2^2$ and a_3^2 , so (a_1', a_2') is a rotation of (a_1, a_2) and $a_3' = \pm a_3$; the half-turn about e_1 if needed, followed by a rotation about e_3 , moves a to a' . If $Q = 0$, only $|a|^2$ is informative and $\text{SO}(3)$ is transitive on spheres. Parity follows from Theorem 3.6(iv) or directly from $\det(-a, Q(-a), Q^2(-a)) = -\det(a, Qa, Q^2 a)$. The relation is the Gram identity, and Q^3 reduces by Cayley–Hamilton for traceless Q . $\square$

The seven invariants vanish simultaneously only at $c = 0$ and are weighted homogeneous, so the map they define is proper. Hence they realize $V_2/\text{SO}(3)$, of dimension six, as a closed semialgebraic subset of $\mathbb{R}^7$, contained in the hypersurface $\varepsilon^2 = \det \text{Gram}(a, Qa, Q^2 a)$; the sign of ε is the chirality of the dipole–quadrupole pair (Theorem 5.1(iii)). After scale, $[f_0 : |a|^2 : \text{tr } Q^2 : \det Q : a^\top Q a : a^\top Q^2 a :$




FIGURE 2. The image of $\mathcal{H}_2$ in the (I, J) -plane is the shaded region $\{I \geq 0, I^3 \geq 27J^2\}$, bounded by the curves $J = \pm I^{3/2}/\sqrt{27}$ of axially symmetric quadrupoles. It is not convex: the points $(1, c)$ and $(4, 8c)$ with $c = 27^{-1/2}$ lie on the boundary, and their midpoint lies outside. Linear interpolation of invariant coordinates therefore leaves the image.

$\varepsilon] \in \mathbb{P}_{(1,2,2,3,3,4,6)}^+(\mathbb{R})$ is a complete scale-free descriptor of the truncation. Bands zero through two contain five of the fifteen coefficients selected in SPHARM-COM (Section 10).

Degree three. A nonzero real harmonic of degree three corresponds to a sextic $F = \lambda Q_a Q_b Q_c$ with $a, b, c \in \mathbb{R}^3 \setminus 0$ and $\lambda \in \mathbb{R}^\times$. The invariant ring of binary sextics is

generated by the **Igusa–Clebsch invariants** I_2, I_4, I_6, I_{10} of degrees 2, 4, 6, 10 and the **skew invariant** R of degree 15, with $R^2 \in \mathbb{C}[I_2, I_4, I_6, I_{10}]$ [12, 19, 22]. In the root form above, with $F = a_0 \prod (z_0 - \alpha_i z_1)$,

$$\begin{aligned}
 I_2 &= a_0^2 \sum_{15} (12)^2 (34)^2 (56)^2, \\
 I_4 &= a_0^4 \sum_{10} (12)^2 (23)^2 (31)^2 (45)^2 (56)^2 (64)^2, \\
 I_6 &= a_0^6 \sum_{60} (12)^2 (23)^2 (31)^2 (45)^2 (56)^2 (64)^2 \\
 &\quad \times (14)^2 (25)^2 (36)^2, \\
 I_{10} &= a_0^{10} \prod_{i < j} (ij)^2,
 \end{aligned}
 \tag{20}$$

where the sums run over the 15 partitions of the six roots into three pairs, the 10 partitions into two triples, and the 60 such partitions together with a bijection between the two triples. Write $F = \sum_k b_k z_0^{6-k} z_1^k$, so that $b_k = \binom{6}{k} a_k$ in the binomial normalization of Section 4.4. The invariants (20) are polynomials with integer coefficients in $b_0, \dots, b_6$; for instance $I_2 = -240b_0b_6 + 40b_1b_5 - 16b_2b_4 + 6b_3^2$, and the polynomial forms of I_4 and I_6 are in [28], where they are denoted J_{2i} . The invariant I_{10} is the discriminant of F , so $I_{10} \neq 0$ if and only if the axes $\mathbb{R}a, \mathbb{R}b, \mathbb{R}c$ are distinct. Throughout we use Igusa’s arithmetic invariants [19]

$$\begin{aligned}
 J_2 &= \frac{I_2}{8}, & J_4 &= \frac{4J_2^2 - I_4}{96}, \\
 J_6 &= \frac{8J_2^3 - 160J_2J_4 - I_6}{576}, & J_{10} &= \frac{I_{10}}{4096},
 \end{aligned}
 \tag{21}$$

which generate the same ring as I_2, I_4, I_6, I_{10} , have the same degrees, and are the coordinates of the moduli space $\mathcal{M}_2 = \{J_{10} \neq 0\} \subset \mathbb{P}_{(2,4,6,10)}$ used in [5]. For R we use the transvectant construction. In the normalization (18) put $\Theta = (F, F)_4$, $Y_1 = (F, \Theta)_4$, $Y_2 = (\Theta, Y_1)_2$, $Y_3 = (\Theta, Y_2)_2$ and

$$R = \det(y_{rs}), \quad Y_r = y_{r0}z_0^2 + 2y_{r1}z_0z_1 + y_{r2}z_1^2,
 \tag{22}$$

where Θ is a quartic covariant and Y_1, Y_2, Y_3 are quadratic covariants.

The locus $\mathcal{L}_2$ and the square of the skew invariant. Let $\mathcal{L}_2 \subset \mathcal{M}_2$ be the locus of genus-two curves whose automorphism group contains an involution other than the hyperelliptic one, an **elliptic involution**; equivalently, of curves with a degree-two elliptic subcover. By Jacobi’s normal form, such a curve can be written $Y^2 = X^6 - s_1X^4 + s_2X^2 - 1$, and the parameters

$$\begin{aligned}
 u &= s_1s_2, & v &= s_1^3 + s_2^3, \\
 \Delta(u, v) &= u^2 - 4v + 18u - 27 \neq 0,
 \end{aligned}
 \tag{23}$$

give a birational parametrization of $\mathcal{L}_2$ by the affine plane [28]. On this family the invariants (21) are

$$\begin{aligned}
 J_2 &= 2u + 30, & J_4 &= \frac{1}{8}(u^2 + 82u - 4v + 165), \\
 J_6 &= -u^2 + 14u + 4v - 5, & J_{10} &= \frac{1}{64}\Delta(u, v)^2.
 \end{aligned}
 \tag{24}$$

Eliminating u and v from (24) gives the equation $F_2(J_2, J_4, J_6, J_{10}) = 0$ of $\mathcal{L}_2$ in $\mathbb{P}_{(2,4,6,10)}$, where F_2 is an irreducible weighted-homogeneous polynomial of degree 30 with 29 terms. It is written out in [28]; we normalize it as the primitive integer polynomial in which the coefficient of $J_2^6 J_6^3$ is -1 .

The square of the skew invariant is a polynomial in J_2, J_4, J_6, J_{10} , and with the normalizations above it is

$$(25) \quad R^2 = -\frac{2^{18}}{3^{18} 5^{20}} F_2(J_2, J_4, J_6, J_{10}).$$

The constant was determined by exact evaluation (Section 8.5); the identity itself is proved in Theorem 5.2. We put

$$(26) \quad \tilde{R} = 2^{-9} 3^9 5^{10} R, \quad \tilde{R}^2 = -F_2(J_2, J_4, J_6, J_{10}),$$

and use $\tilde{R}$ as the descriptor coordinate. The factor matters in practice: the gauge coordinate (17) of $-iR$ is of order 10^{-16} on generic chiral band-three shapes, so without it the diagnostic of Section 8.4 would report every such shape as lying near the divisor $R = 0$.

Since $l = 3$ is odd, Theorem 3.6(iii) shows that J_2, J_4, J_6, J_{10} are real on band-three shapes while R is purely imaginary, so $-iR$ is a real pseudo-invariant. Not all of J_2, J_4, J_6, J_{10} vanish on a nonzero band-three shape: if they did, R would vanish by (25), so F would be a nullform and would have a root of multiplicity at least four, which is impossible for roots in antipodal pairs. The point $[J_2 : J_4 : J_6 : J_{10}] \in \mathbb{P}_{(2,4,6,10)}(\mathbb{R})$ is a complete invariant of the band-three shape up to rotation, scale and reflection: if the even invariants agree then R^2 agrees by (25), so $R' = \pm R$, and by Theorem 4.1 f' lies in the orbit of f or of its mirror. Adjoining $\hat{R} = -i\tilde{R}/H_w^{15}$, where H_w is the gauge (17) of the vector $(J_2, J_4, J_6, J_{10}, -i\tilde{R})$ with weights $(2, 4, 6, 10, 15)$, restores completeness up to rotation and positive scale. These statements concern a descriptor supported on band three. On V_3 the orientation of band three relative to bands one and two is recorded only by joint invariants (27). When $J_{10} \neq 0$, the point $[J_2 : J_4 : J_6 : J_{10}]$ is the moduli point of the genus-two curve $y^2 = F(z, 1)$, which is used in Theorem 5.2 to identify the achiral locus.

Joint invariants. For bands (l_1, l_2, l_3) satisfying the triangle inequalities, put $k = l_1 + l_2 - l_3$ and

$$(27) \quad B_{l_1 l_2 l_3} = ((F_{l_1}, F_{l_2})_k, F_{l_3})_{2l_3}.$$

These are the bispectral invariants of [20] as transvectants, real when $l_1 + l_2 + l_3$ is even and pseudo-invariants when it is odd. When two bands coincide, $B_{l_1 l l}$ is proportional to the unique invariant pairing of $\mathcal{H}_{l_1} \otimes \mathcal{H}_l \otimes \mathcal{H}_l$, whose symmetry under exchange of the two $\mathcal{H}_l$ factors is $(-1)^{l_1 + 2l}$; hence $B_{l_1 l l} = 0$ for odd l_1 . A cubic pseudo-invariant therefore needs three distinct bands with odd sum, the smallest being B_{234} , so none occurs for $L \leq 3$; degree-four pseudo-invariants occur already for $L = 3$, e.g. $P_{123} = ((F_1, F_2)_1, (F_3, F_3)_4)_4$. Invariants of degree at most three do not separate in general. For a single band $\mathcal{H}_l$ with $l \geq 3$, the invariant quadratics are the multiples of the norm and the invariant cubics form a space of dimension at most one, since $\dim(\mathcal{H}_l^{\otimes 3})^{\text{SO}(3)} = 1$, while $\dim \mathcal{H}_l / \text{SO}(3) = 2l - 2 \geq 4$.

5. CHIRALITY

The pseudo-invariants have a geometric meaning: they detect whether a shape is congruent to its mirror image.




FIGURE 3. Maxwell axes of a nonzero real harmonic of degree three, drawn as lines through the center of $\mathbb{S}^2$. Left: coplanar axes; the half-turn about the normal n of their plane reverses f . Middle: bisector type; n bisects the angle of $\mathbb{R}a$ and $\mathbb{R}b$, and $\mathbb{R}c$ is orthogonal to n , so the half-turn about n exchanges $\mathbb{R}a$ and $\mathbb{R}b$ and reverses $\mathbb{R}c$. In both cases f is achiral and $R(F_f) = 0$. Right: a generic configuration, for which f is chiral.

Proposition 5.1 (Chirality criterion). *Let $c \in V_L$ and let c^* be the radial descriptor of the reflected surface. Then $c^* \in \text{SO}(3) \cdot c$ iff every pseudo-invariant in $\mathcal{R}_L$ vanishes at Ψc . In particular:*

- (i) every descriptor supported on even bands is achiral, and so is every descriptor supported on band one;
- (ii) a band-three descriptor f_3 is achiral iff $R(F_{f_3}) = 0$;
- (iii) a descriptor supported on bands one and two, $f_1 = \langle a, \cdot \rangle$, $f_2 = x^\top Qx$, is achiral iff $\det(a, Qa, Q^2a) = 0$, i.e. iff a lies in a plane spanned by two eigenvectors of Q .

The same holds for the coordinate model with the mirror pseudo-invariants of Theorem 4.4; the band-one coordinate descriptor is achiral by Theorem 4.6.

Proof. By Theorem 3.6(iv) and Theorem 2.3, c^* is in the orbit of $((-1)^l f_l)_l$. Decompose a generating set into parity components; by Theorem 4.1, c and c^* are in the same orbit iff all components agree on both, and even components agree automatically while odd ones change sign. (i) On even bands all ld_l are even; on band one the ring is generated by the discriminant. (ii) The odd part of the sextic ring is $R \cdot \mathbb{C}[J_2, J_4, J_6, J_{10}]$. (iii) In an eigenbasis of Q with distinct eigenvalues, $\det(a, Qa, Q^2a) = a_1 a_2 a_3 \prod_{i < j} (\lambda_j - \lambda_i)$, which vanishes iff some $a_i = 0$; then the half-turn about e_i commutes with Q and sends $a \mapsto -a$, so the descriptor is congruent to its mirror. Conversely if all $a_i \neq 0$, every rotation commuting with Q is in K and none sends a to $-a$. If Q has a repeated eigenvalue the determinant vanishes and $\text{Stab}(Q)$ contains a rotation sending a to $-a$. $\square$

For the band-three case we can say exactly what the vanishing of R means; Figure 3 shows the configurations that occur. Recall that a nonzero $f \in \mathcal{H}_3$ has three Maxwell axes $\mathbb{R}a, \mathbb{R}b, \mathbb{R}c$, unique with multiplicity, with $f \equiv \lambda \langle a, x \rangle \langle b, x \rangle \langle c, x \rangle \pmod{q}$ by Theorem 3.6(ii).

Theorem 5.2 (Chirality of degree-three harmonics). *Let $0 \neq f \in \mathcal{H}_3$ with axes a, b, c and sextic $F = F_f$. The following are equivalent.*

- (i) f is achiral: its mirror image lies in $\mathrm{SO}(3) \cdot f$.
- (ii) $R(F) = 0$.
- (iii) There is a half-turn $\rho \in \mathrm{SO}(3)$ with $\rho \cdot f = -f$.
- (iv) Either the three axes are coplanar, or one axis is orthogonal to an angle bisector of the other two.

Moreover, when $J_{10}(F) \neq 0$, these are equivalent to the genus-two curve $y^2 = F(z, 1)$ having an elliptic involution, that is, lying in the locus $\mathcal{L}_2$; the classical form of this statement is due to Clebsch [12, 28]. The identity (25) holds for every complex sextic, and for every nonzero $f \in \mathcal{H}_3$, including those with $J_{10}(F) = 0$, the conditions (i)–(iv) are equivalent to

- (v) $F_2(J_2(F), J_4(F), J_6(F), J_{10}(F)) = 0$, with F_2 the equation of $\mathcal{L}_2$ normalized as in Section 4.4.

Finally, as a polynomial in (λ, a, b, c) ,

$$(28) \quad \begin{aligned} -iR(F) &= \lambda^{15} \det(a, b, c) \\ &\times G(\langle a, a \rangle, \langle a, b \rangle, \langle a, c \rangle, \\ &\quad \langle b, b \rangle, \langle b, c \rangle, \langle c, c \rangle) \end{aligned}$$

for a polynomial G with rational coefficients, and on the noncoplanar configurations G vanishes exactly on those of bisector type.

Proof. (i) $\Leftrightarrow$ (ii) is Theorem 5.1(ii).

(i) $\Rightarrow$ (iv). Achirality means some $g \in \mathrm{O}(3) \setminus \mathrm{SO}(3)$ satisfies $g \cdot f = f$. Write $g = -\rho$ with $\rho \in \mathrm{SO}(3)$; since -1 acts on $\mathcal{H}_3$ by -1 , this says $\rho \cdot f = -f$. Since ρ preserves q , it acts on the Maxwell representation: $\rho \cdot f \equiv \lambda \prod_k \langle \rho a_k, x \rangle \pmod{q}$, so by uniqueness of the axes ρ permutes the multiset $\{\mathbb{R}a, \mathbb{R}b, \mathbb{R}c\}$, $\rho a_k = t_k a_{\pi(k)}$, and comparing scalars $\prod_k t_k = -1$ (the product $\prod t_k$ does not depend on the choice of representatives, as rescaling $a_k \mapsto s_k a_k$ multiplies it by $\prod s_k / s_{\pi(k)} = 1$). Let ρ be rotation by $\theta \in (0, 2\pi)$ about the unit vector n . A rotation fixes a line $\ell \neq \mathbb{R}n$ only if $\theta = \pi$ and $\ell \perp n$; it fixes $\mathbb{R}n$ with $t = +1$, and a line $\ell \perp n$ under a half-turn with $t = -1$.

Case 1: π is the identity. Each axis is either $\mathbb{R}n$ ($t = +1$) or, if $\theta = \pi$, perpendicular to n ($t = -1$). Since $\prod t_k = -1$, either one axis is $\perp n$ and two coincide with $\mathbb{R}n$, or all three are $\perp n$. In both cases the axes are coplanar.

Case 2: π is a transposition, say $\mathbb{R}a \neq \mathbb{R}b$ are exchanged and $\mathbb{R}c$ is fixed. If $\theta = \pi$, then $\rho^2 = 1$ gives $t_a t_b = 1$, so $t_c = -1$ and $c \perp n$; moreover $\rho a = 2\langle a, n \rangle n - a \in \mathbb{R}b$ forces $n \in \mathrm{span}(a, b)$, and a half-turn about a line in the plane of $\mathbb{R}a, \mathbb{R}b$ exchanges them iff that line bisects their angle. If $\theta \neq \pi$, then ρ^2 is a nontrivial rotation fixing the lines $\mathbb{R}a, \mathbb{R}b, \mathbb{R}c$; the lines $\mathbb{R}a, \mathbb{R}b$ are not $\mathbb{R}n$ (else ρ would fix them), so ρ^2 is the half-turn about n with $a, b \perp n$, i.e. $\theta = \pm\pi/2$; then $\rho a \perp a$ gives $a \perp b$, and c , fixed by a quarter-turn, lies on $\mathbb{R}n$. So a, b, c are mutually orthogonal, and c is orthogonal to both bisectors of $\mathbb{R}a, \mathbb{R}b$. In either subcase (iv) holds.

Case 3: π is a 3-cycle, so the three lines are distinct. Choose representatives $b = \rho a$, $c = \rho b$; then $\rho c = \rho^3 a$ and ρ^3 fixes $\mathbb{R}a$. Since $\mathbb{R}a \neq \mathbb{R}n$ (else ρ would fix $\mathbb{R}a$), either $\rho^3 = 1$, in which case $\rho c = a$, all $t_k = 1$ and $\prod t_k = 1$, a contradiction; or ρ^3 is the half-turn about n with $a \perp n$, so $\theta \in \{\pi/3, 5\pi/3\}$ ($\theta = \pi$ would fix $\mathbb{R}a$) and a, b, c all lie in the plane $\perp n$. The axes are coplanar.

(iv) $\Rightarrow$ (iii). If the axes are coplanar with normal n , the half-turn about n sends each $a_k \mapsto -a_k$, so $\rho \cdot f = (-1)^3 f = -f$. If $c \perp n$ with n a bisector of $\mathbb{R}a, \mathbb{R}b$, choose unit representatives with $n \propto \hat{a} + \hat{b}$; the half-turn about n maps $\hat{a} \mapsto 2\langle \hat{a}, n \rangle n - \hat{a} = \hat{b}$, $\hat{b} \mapsto \hat{a}$, and $c \mapsto -c$, so $\rho \cdot f = -f$.

(iii) $\Rightarrow$ (i). If $\rho \cdot f = -f$ then $(-\rho) \cdot f = f$ and $-\rho \in \mathrm{O}(3) \setminus \mathrm{SO}(3)$.

For the complex statement, let $J_{10}(F) \neq 0$. Then $y^2 = F(z, 1)$ is a smooth genus-two curve, and $R(F) = 0$ iff the curve has an elliptic involution [28]. Combine with (i) $\Leftrightarrow$ (ii). Note that (iii) supplies, for real F , an involution defined over $\mathbb{R}$ and in $\mathrm{SO}(3)$.

For (25): the invariants J_2, J_4, J_6, J_{10} have weighted degrees 2, 4, 6, 10, and F_2 is weighted-homogeneous of degree 30 and irreducible [28]. The polynomial R^2 is an even invariant, so it lies in $\mathbb{C}[J_2, J_4, J_6, J_{10}]$, and it is weighted-homogeneous of degree 30 as well. On the open set $J_{10} \neq 0$ the zero set of R is $\mathcal{L}_2$ by the criterion of [12, 28], so R^2 vanishes on $\mathcal{L}_2$. The locus $\mathcal{L}_2$ is dense in the irreducible surface $V(F_2)$, and (F_2) is a prime ideal, so $R^2 \in (F_2)$ and $R^2 = cF_2$ with c weighted-homogeneous of degree zero, that is, a constant. A single exact evaluation at a sextic with $F_2 \neq 0$ determines c ; it gives $c = -2^{18}3^{-18}5^{-20}$ (Section 8.5). Hence (v) is equivalent to (ii) for every F , including those with $J_{10} = 0$.

For (28): the coefficients of F are λ times trilinear expressions in (a, b, c) , so $-iR(F)$ is λ^{15} times a polynomial $P(a, b, c)$ of tridegree $(15, 15, 15)$, with real coefficients by Theorem 3.6(iii), and P is $\mathrm{SO}(3)$ -invariant under the diagonal action because Ψ_3 is equivariant and rotating f rotates its axes. By the first fundamental theorem for $\mathrm{SO}(3)$ [32], $P \in \mathbb{R}[\langle a_i, a_j \rangle, \det(a, b, c)]$, and since $\det(a, b, c)^2$ is a polynomial in inner products, $P = G_0 + \det(a, b, c)G_1$ with G_0, G_1 polynomials in inner products. Each inner product has total degree two and $\det$ has total degree three; the total degree of P is 45, so every monomial contains an odd number of determinant factors, hence $G_0 = 0$. Finally $\det(a, b, c) = 0$ is the coplanar case, and by the equivalences G vanishes on a noncoplanar configuration iff it is of bisector type. $\square$

Remark 5.3. Condition (v) is a practical test: F_2 is a real polynomial of degree 30 in the $\mathrm{O}(3)$ -invariants J_2, J_4, J_6, J_{10} , so achirality of a band-three shape can be decided without evaluating the degree-15 pseudo-invariant R and without complex arithmetic. The equivalence (i) $\Leftrightarrow$ (iv) has an elementary check: $f = x_1x_2x_3$ has axes e_1, e_2, e_3 , e_3 is orthogonal to the bisectors of e_1, e_2 , and indeed f is fixed by the reflection $x_1 \leftrightarrow x_2$. We verified (28) and the four cases of (iv) by exact computation (Section 8): R vanishes on coplanar and on bisector configurations, is nonzero on generic ones, changes sign under $a \mapsto -a$, and is homogeneous of degree 15 in each axis. The identification of the achiral band-three harmonics with the real points of $\mathcal{L}_2$ converts a classical genus-two locus into a statement about shapes.

Remark 5.4. For $l = 3$, Theorem 5.2(iii) says that f is achiral iff it is fixed by a reflection: if ρ is the half-turn about $\mathbb{R}n$ with $\rho \cdot f = -f$, then $-\rho$ is the reflection in $n^\perp$ and $(-\rho) \cdot f = f$. For general l , achirality means $S \cdot f = f$ for some $S \in \mathrm{O}(3) \setminus \mathrm{SO}(3)$, and S is a reflection only when $-S$ is a half-turn; the finer classification of achiral harmonics of higher degree is not needed for the descriptor and is not treated here.

For anatomical applications parity must be handled deliberately. Homologous left and right structures are related only approximately by reflection, so raw pseudo-invariants confound pathological asymmetry with laterality; a hemispheric comparison must first reflect one side into a common handedness, after which pseudo-invariants are comparable across sides.

6. CANONICAL FRAMES ON THE PRINCIPAL STRATUM

Global polynomial invariants are continuous everywhere but numerous and of high degree. On the open stratum where the lowest informative band has trivial continuous symmetry, a canonical frame reduces the residual group to a finite one. The aligned coordinates number exactly the dimension of the quotient, and the invariant monomials of degree at most three in them form a separating set. This section makes that precise, records the strata where it fails, and states what continuous invariants can and cannot do.

Let $K = \{\text{diag}(\epsilon_1, \epsilon_2, \epsilon_3) : \epsilon_i = \pm 1, \epsilon_1 \epsilon_2 \epsilon_3 = 1\} \cong (\mathbb{Z}/2)^2$, the group of half-turns about the coordinate axes together with the identity. In the standard real basis $\{Y_{lm}\}$ of $\mathcal{H}_l$ every $\kappa \in K$ acts diagonally by signs, since a half-turn about a coordinate axis sends $\cos m\phi$, $\sin m\phi$, and $P_l^m(\cos \theta)$ each to $\pm$ itself; hence K acts on each coordinate of V_L and of W_L through a character $\chi \in \hat{K} \cong (\mathbb{Z}/2)^2$.

Lemma 6.1. *Let K act on $\mathbb{R}^n$ diagonally by characters $\chi_1, \dots, \chi_n$. The K -invariant monomials of degree at most three separate K -orbits, and they generate $\mathbb{R}[x]^K$.*

Proof. A monomial $\prod x_j^{m_j}$ is invariant iff $\sum m_j \chi_j = 0$ in $\hat{K}$. Suppose x, x' agree on all invariant monomials of degree ≤ 3 . From $x_j^2 = x_j'^2$ we get $x_j' = \epsilon_j x_j$, with ϵ_j free where $x_j = 0$. For j, k with $\chi_j = \chi_k$ and $x_j x_k \neq 0$, $x_j x_k = x_j' x_k'$ gives $\epsilon_j = \epsilon_k$; for $\chi_j = 1$, $x_j = x_j'$. So there is a well-defined $\eta(\chi) \in \{\pm 1\}$ on each nontrivial character carried by some nonzero coordinate, and if all three nontrivial characters χ, χ', χ'' (which satisfy $\chi \chi' \chi'' = 1$) are carried, a cubic monomial $x_j x_k x_m$ with these characters is invariant and gives $\eta(\chi) \eta(\chi') \eta(\chi'') = 1$. The map $K \rightarrow \{(t, t', t'') \in \{\pm 1\}^3 : tt't'' = 1\}$, $\kappa \mapsto (\chi(\kappa), \chi'(\kappa), \chi''(\kappa))$, is a bijection, so there is $\kappa \in K$ with $\chi(\kappa) = \eta(\chi)$ on all carried characters (an assignment on at most two characters extends to one with product one), and $\kappa \cdot x = x'$. Generation in degree ≤ 3 is the statement that the Davenport constant of $(\mathbb{Z}/2)^2$ is three [14]. $\square$

Theorem 6.2. (a) *Radial model.* Let $L \geq 2$ and let $V_L^\circ \subset V_L$ be the open set where the quadrupole $f_2 = x^\top Q x$ has three distinct eigenvalues $\lambda_1 > \lambda_2 > \lambda_3$, and for $c \in V_L^\circ$ let $E \in \text{SO}(3)$ have as columns an orthonormal eigenbasis of Q in this order. E is unique up to $E \mapsto E\kappa$, $\kappa \in K$. The aligned tuple $\tilde{c} = E^{-1} \cdot c$, i.e. $\tilde{f}_i(u) = f_i(Eu)$, is therefore well defined up to the action of K , the map $c \mapsto K \cdot \tilde{c}$ is $\text{SO}(3)$ -invariant, and $c, c' \in V_L^\circ$ lie in the same $\text{SO}(3)$ -orbit iff $K \cdot \tilde{c} = K \cdot \tilde{c}'$. Consequently the K -invariant monomials of degree ≤ 3 in the coordinates of $\tilde{c}$ form a separating set of real-analytic $\text{SO}(3)$ -invariant functions on V_L° , of size at most cubic in $\dim V_L$; the aligned coordinates themselves number $\dim V_L - 3 = \dim V_L / \text{SO}(3)$.

(b) *Coordinate model.* Let $W_L^\circ \subset W_L$ be the open set where $C_1 = M$ has distinct singular values $\sigma_1 > \sigma_2 > \sigma_3 \geq 0$. For $C \in W_L^\circ$ write $M = R_0 \Sigma Q_0^\top$ as in Theorem 4.6; the pair (R_0, Q_0) is unique up to $(R_0 \kappa, Q_0 \kappa)$, $\kappa \in K$.

The aligned tuple $\tilde{C} = (Q_0, R_0)^{-1} \cdot C$, i.e. $\tilde{C}_l = R_0^\top C_l D_l(Q_0)$, is well defined up to the diagonal action $\tilde{C}_l \mapsto \kappa \tilde{C}_l D_l(\kappa)$ of K , and $C, C' \in W_L^\circ$ lie in the same $\mathrm{SO}(3) \times \mathrm{SO}(3)$ -orbit iff $K \cdot \tilde{C} = K \cdot \tilde{C}'$. The K -invariant monomials of degree ≤ 3 in the coordinates of $\tilde{C}$ separate orbits on W_L° ; the aligned coordinates number $\dim W_L - 6 = \dim \mathcal{Q}_L$.

- (c) In both cases the aligned tuple is real-analytic on the open stratum, and the class EK (resp. $(R_0, Q_0)K$) of the frame has no continuous extension to any point of the complement.

Proof. (a) With distinct eigenvalues the eigenlines are unique, so E is determined up to the signs of its columns subject to $\det E = 1$, i.e. up to K . Replacing E by $E\kappa$ replaces $\tilde{c}$ by $\kappa^{-1} \cdot \tilde{c} = \kappa \cdot \tilde{c}$. If $c' = R \cdot c$ then $Q' = RQR^\top$ has eigenframe $E' = RE\kappa$ for some κ , and $\tilde{c}' = (RE\kappa)^{-1} \cdot R \cdot c = \kappa \cdot \tilde{c}$; so $K \cdot \tilde{c}$ is invariant. Conversely if $\tilde{c}' = \kappa \cdot \tilde{c}$ then $c' = E'\kappa E^{-1} \cdot c$. Separation of K -orbits by cubic monomials is Theorem 6.1; K acts on the coordinates of $\tilde{c}$ by characters as observed above. Real-analyticity follows from that of the eigendecomposition on the set of simple spectra. The count: $\tilde{f}_2$ is diagonal, so its 5 coordinates reduce to (λ_1, λ_2) with $\lambda_3 = -\lambda_1 - \lambda_2$; together with the other bands this gives $(L+1)^2 - 3$ coordinates.

(b) The signed SVD with distinct singular values is unique up to simultaneous sign changes of paired singular vectors preserving both determinants, i.e. up to $(R_0\kappa, Q_0\kappa)$; when $\sigma_3 = 0$, the third columns of R_0 and Q_0 are determined by the first two and by $\det R_0 = \det Q_0 = 1$. Under $(R_0\kappa, Q_0\kappa)$, $\tilde{C}_l \mapsto \kappa R_0^\top C_l D_l(Q_0) D_l(\kappa) = \kappa \tilde{C}_l D_l(\kappa)$ since D_l is a homomorphism and $\kappa^\top = \kappa$. Equivariance and separation follow as in (a). In the standard bases κ acts on both sides diagonally, so each entry of $\tilde{C}_l$ transforms by a character of K , and Theorem 6.1 applies. The count: $\tilde{C}_1 = \Sigma$ contributes 3 coordinates, and $\sum_{l=2}^L 3(2l+1) = 3((L+1)^2 - 4)$ more, for a total of $3(L+1)^2 - 9 = \dim \mathcal{Q}_L$.

(c) Real-analyticity follows from that of the eigendecomposition on symmetric matrices with simple spectrum and of the singular value decomposition on matrices with distinct singular values. Let $S_0 = E_0 \Lambda_0 E_0^\top$ be a symmetric matrix with $E_0 \in \mathrm{SO}(3)$ and Λ_0 diagonal with a repeated entry, and let $H \subset \mathrm{SO}(3)$ be the group of rotations commuting with Λ_0 . For $h \in H$, a diagonal matrix Λ with distinct entries, and small $t > 0$, put

$$S_{h,t} = E_0 h (\Lambda_0 + t\Lambda) h^\top E_0^\top.$$

Then $S_{h,t} \rightarrow S_0$ as $t \rightarrow 0$, $S_{h,t}$ has a simple spectrum, and its ordered eigenframe is $E_0 h P$ modulo K for a signed permutation matrix P with $\det P = 1$, independent of h and t . Since Λ_0 has a repeated entry, H is infinite, while K is finite; hence the classes $E_0 h P K$, $h \in H$, are not all equal, and the class of the frame has no continuous extension to S_0 . For the radial model take $S_0 = Q$, keep the other bands fixed, and take $\Lambda = \mathrm{diag}(1, 0, -1)$, so that $S_{h,t}$ is traceless. For the coordinate model take $S_0 = M M^\top$, keep C_l fixed for $l \geq 2$, take $\Lambda = \mathrm{diag}(3, 2, 1)$, write $M = S_0^{1/2} U$ with $U \in \mathrm{O}(3)$, and put $M_{h,t} = S_{h,t}^{1/2} U$; then $M_{h,t} \rightarrow M$, the singular values of $M_{h,t}$ are distinct, and its left singular frame is the eigenframe of $S_{h,t} = M_{h,t} M_{h,t}^\top$. $\square$

Remark 6.3. The set V_L° is open and dense in V_L , and its points with trivial stabilizer form its intersection with the principal stratum, again open and dense. The complement of V_L° has three strata: $\lambda_1 = \lambda_2$ or $\lambda_2 = \lambda_3$ (axially symmetric quadrupole, stabilizer $\mathrm{O}(2) \cap \mathrm{SO}(3)$), and $Q = 0$. On the axial stratum the residual

freedom is a circle plus a half-turn, and it can be fixed by the next nonzero band that breaks it (for instance by the dipole a if $a \nparallel$ the axis), reducing again to a finite group; on $Q = 0$ one starts from the first nonzero band. The same cascade applies to the coordinate model with the strata $\sigma_1 = \sigma_2$ and $\sigma_2 = \sigma_3$. Each stratum is detected by an exact algebraic predicate: for the radial model by the vanishing of disc $F_{f_2} = I^3 - 27J^2$ or of I , for the coordinate model by the vanishing of the discriminant of the cubic in Theorem 4.6. Near a stratum the frame is well defined but ill-conditioned, with condition number governed by the reciprocal of the eigenvalue or singular-value gap; Section 7 gives the stability statement that does not depend on the frame.

Remark 6.4. Part (b) is the rigorous form of the first-order-ellipsoid alignment of SPHARM-PDM [7, 29]: the degree-one coefficients of the coordinate map define the ellipsoid $u \mapsto Mu$, and aligning it is exactly the SVD frame. The differences are that our parameterization is the conformal one, so the frame is a function of the surface and not of an optimizer's initialization; that the residual finite group is identified and quotiented rather than fixed by a convention; and that the degenerate strata and the impossibility of a continuous extension are stated.

Proposition 6.5. *Let G be $\text{SO}(3)$ acting on $V = V_L$ or $\text{SO}(3) \times \text{SO}(3)$ acting on $V = W_L$, and let $I_1, \dots, I_N$ generate $\mathbb{R}[V]^G$. For a generic linear map $A: \mathbb{R}^N \rightarrow \mathbb{R}^{2 \dim V + 1}$, the map $A \circ (I_1, \dots, I_N)$ separates G -orbits on V . In particular continuous polynomial invariants inject $V_L/\text{SO}(3)$ into $\mathbb{R}^{2(L+1)^2+1}$ and $\mathcal{Q}_L$ into $\mathbb{R}^{6(L+1)^2-5}$.*

Proof. This is the theorem of Dym and Gortler [16] applied to the compact groups at hand, whose polynomial invariants separate orbits by Theorems 4.1 and 4.5. $\square$

Together, Theorem 6.2 and Theorem 6.5 describe the trade-off exactly. On the principal stratum, a frame gives $\dim(V/G)$ real-analytic coordinates, determined up to the sign action of K , at the cost of discontinuity at the symmetry strata. Globally, $2 \dim V + 1$ continuous polynomial invariants suffice, at the cost of roughly doubling the dimension and of using generic combinations of high-degree generators. Max-filter constructions [8] give a third route, with Lipschitz and in some cases bi-Lipschitz guarantees; for the harmonic representations here their templates would themselves have to be chosen in V_L , and we have not pursued this.

7. METRICS AND STABILITY

The previous sections produce, for a surface S and a degree L , a vector of invariants $\mathcal{I}(c)$ of the radial descriptor $c = f^{(L)}(\varphi)$, and a vector $\mathcal{I}_2(C)$ of the coordinate descriptor. By Theorem 2.3 these vectors do not depend on the parameterization. By Theorems 4.1 and 4.5 they determine the descriptor up to the nuisance rotations. A descriptor that is to be used on measured data needs one more property. Measured shapes carry noise, and two shapes are never exactly congruent. So we must know how the invariants respond to small changes of shape, and conversely how much the shape can change when the invariants change little. The first question asks for continuity of the map from shapes to invariants. The second asks for continuity of its inverse. Both need a distance on the space of shapes modulo the nuisance group. This section defines that distance, proves the two continuity statements, and collects the properties of the descriptor into one theorem.

7.1. The orbit distance. Let G be a compact group acting linearly and isometrically on a real inner product space V ; the cases are $G = \text{SO}(3)$ on $V = V_L$ and $G = \text{SO}(3) \times \text{SO}(3)$ on $V = W_L$, with the $L^2(d\omega)$ norm on V_L and the Frobenius norm $\|C\|_F^2 = \sum_l \text{tr}(C_l C_l^\top)$ on W_L . Both norms are G -invariant, since $D_l(Q)$ is orthogonal. The **orbit distance** of $c, c' \in V$ is the distance between their orbits:

$$\begin{aligned} \delta(c, c') &= \min_{R \in \text{SO}(3)} \|c - R \cdot c'\|, \\ (29) \quad \delta(C, C')^2 &= \min_{Q, R \in \text{SO}(3)} \sum_{l=1}^L \|C_l - R C'_l D_l(Q)^\top\|_F^2. \end{aligned}$$

The minimum exists because G is compact. In words, $\delta(c, c')$ is the smallest distance between c and any rotated copy of c' ; it measures how far two shapes are from being congruent, after the best alignment.

The second form makes both nuisance rotations of the coordinate model explicit. For fixed Q the optimal R maximizes $\text{tr}(R^\top H(Q))$ with $H(Q) = \sum_l C_l D_l(Q) C_l^\top$, and the maximizer is the Kabsch solution $R = U \text{diag}(1, 1, \det(UV^\top)) V^\top$ for a singular value decomposition $H(Q) = U \Sigma V^\top$. So the minimization is over the sphere factor alone. It is nonconvex and needs documented multistart. A scale-free version replaces each nonzero input by its unit-norm representative before minimizing.

Two examples show what the orbit distance computes.

Example 7.1. Let $f = \langle a, \cdot \rangle$ and $f' = \langle a', \cdot \rangle$ in $\mathcal{H}_1$, with $a, a' \in \mathbb{R}^3$, and use the norm $|a|$ on the coefficient. A rotation moves a' to any vector of length $|a'|$, so

$$\delta(f, f') = \min_{R \in \text{SO}(3)} |a - R a'| = ||a| - |a'||.$$

The orbit distance is the difference of the norms, which is the only invariant of band one.

Proposition 7.2. Let $M, M' \in \mathbb{R}^{3 \times 3}$ be band-one coordinate descriptors as in Theorem 4.6, with signed singular values $s = (\sigma_1, \sigma_2, \text{sgn}(\det M) \sigma_3)$ and s' defined likewise. If $\det M \cdot \det M' \geq 0$, then

$$\delta(M, M') = \min_{R, Q \in \text{SO}(3)} \|M - R M' Q^\top\|_F = |s - s'|.$$

Proof. Write signed singular value decompositions $M = U \Sigma V^\top$ and $M' = U' \Sigma' V'^\top$ with $U, V, U', V' \in \text{SO}(3)$ and $\Sigma = \text{diag}(s)$, $\Sigma' = \text{diag}(s')$. Since the Frobenius norm is bi-invariant under $\text{SO}(3)$, $\|M - R M' Q^\top\|_F = \|\Sigma - \tilde{R} \Sigma' \tilde{Q}^\top\|_F$ with $\tilde{R} = U^\top R U'$ and $\tilde{Q} = V^\top Q V'$, which range over $\text{SO}(3)$ as R, Q do. Hence we may assume $M = \Sigma$ and $M' = \Sigma'$. Expanding the square,

$$\|\Sigma - R \Sigma' Q^\top\|_F^2 = |s|^2 + |s'|^2 - 2 \text{tr}(\Sigma R \Sigma' Q^\top).$$

Let $X = R \Sigma' Q^\top$. Its singular values are $\sigma'_1, \sigma'_2, \sigma'_3$, so by von Neumann's trace inequality $\text{tr}(\Sigma X) \leq \sum_i \sigma_i \sigma'_i$. When $\det M \cdot \det M' \geq 0$ the last entries of s and s' have the same sign, so $\sum_i \sigma_i \sigma'_i = \sum_i s_i s'_i$, and this value is attained at $R = Q = 1$. Hence the minimum of the square is $|s|^2 + |s'|^2 - 2 \sum_i s_i s'_i = |s - s'|^2$. $\square$

When the determinants have opposite signs, the same formula $\delta(M, M') = |s - s'|$ holds, with the sign of s_3 or s'_3 negative: with $\Sigma' = \text{diag}(\sigma'_1, \sigma'_2, -\sigma'_3)$ and Q fixed, the matrix $A = \Sigma' Q^\top \Sigma$ has negative determinant, so $\max_{R \in \text{SO}(3)} \text{tr}(R A) = \sigma_1(A) +$

$\sigma_2(A) - \sigma_3(A)$; since $\sigma_1(A) + \sigma_2(A) \leq \sigma_1\sigma'_1 + \sigma_2\sigma'_2$ and $\sigma_3(A) \geq \sigma_3\sigma'_3$ by the singular value inequalities for products, the maximum over Q is $\sigma_1\sigma'_1 + \sigma_2\sigma'_2 - \sigma_3\sigma'_3 = \sum_i s_i s'_i$, attained at $Q = 1$. We also verified this numerically against a multistart minimization over $\text{SO}(3) \times \text{SO}(3)$ (Section 8). In either case the two-sided orbit distance of band one is the Euclidean distance between the signed singular value vectors, that is, between the canonical representatives of Theorem 4.6.

The **invariant-coordinate distance** is any distance between normalized invariant vectors (17). On a chart where a fixed set of coordinates is nonzero, the root-degree logarithms $w_k^{-1} \log|x_k|$ turn scaling into translation and give a scale-free chart distance. The orbit distance is canonical once representation, truncation and inner product are fixed; an invariant-coordinate distance depends on the embedding $\mathcal{I}$. The following theorem relates the two.

7.2. Stability in both directions.

Theorem 7.3. *The following are true:*

- (i) δ is a metric on $V_L/\text{SO}(3)$ and on $\mathcal{Q}_L$.
- (ii) Let I be a homogeneous polynomial invariant of degree w and B the unit ball of V_L . Then $|I(c) - I(c')| \leq (\sup_B |\nabla I|) \delta(c, c')$ for $c, c' \in B$; in particular every invariant is Lipschitz on bounded sets with respect to the orbit distance, and so is every normalized coordinate (17) on any set where H_w is bounded below.
- (iii) Let $\mathcal{I}$ be a separating polynomial invariant map and $\mathcal{K} \subset V_L$ compact. There are $C_{\mathcal{K}} > 0$ and $\alpha_{\mathcal{K}} \in (0, 1]$ such that
$$\delta(c, c') \leq C_{\mathcal{K}} |\mathcal{I}(c) - \mathcal{I}(c')|^{\alpha_{\mathcal{K}}} \quad (c, c' \in \mathcal{K}).$$
- (iv) At a point of the principal stratum where the Jacobian of $\mathcal{I}$ restricted to a slice has full rank, the exponent in (iii) can be taken to be 1 locally: $\mathcal{I}$ is a local bi-Lipschitz embedding of the quotient into $\mathbb{R}^N$. Here a **slice** at c is a submanifold through c , transverse to the orbit of c and of complementary dimension, which parametrizes the quotient near the orbit of c .

The same statements hold for the coordinate model with $\text{SO}(3) \times \text{SO}(3)$.

Proof. (i) The function δ is symmetric because $\|c - R \cdot c'\| = \|R^{-1} \cdot c - c'\|$ and R^{-1} ranges over $\text{SO}(3)$ with R . For the triangle inequality let R and R' attain $\delta(c, c')$ and $\delta(c', c'')$. Then

$$\begin{aligned} \delta(c, c'') &\leq \|c - RR' \cdot c''\| \\ &\leq \|c - R \cdot c'\| + \|R \cdot (c' - R' \cdot c'')\| \\ &= \delta(c, c') + \delta(c', c''), \end{aligned}$$

using the invariance of the norm. Finally $\delta(c, c') = 0$ if and only if c' lies in the closure of the orbit of c , and the orbit is closed because the group is compact. So δ vanishes exactly on pairs in the same orbit and descends to a metric on the quotient.

(ii) Let R attain $\delta(c, c')$. Since I is invariant, $I(c') = I(R \cdot c')$, and $R \cdot c' \in B$ because B is invariant. The segment from c to $R \cdot c'$ lies in the convex set B , so by the mean value theorem $|I(c) - I(R \cdot c')| \leq \sup_B |\nabla I| \|c - R \cdot c'\| = \sup_B |\nabla I| \delta(c, c')$. Normalized coordinates are compositions of $\mathcal{I}$ with the map $x \mapsto \hat{x}$ of (17), which is Lipschitz on any set where H_w is bounded below.

(iii) The function $(c, c') \mapsto |\mathcal{I}(c) - \mathcal{I}(c')|$ is continuous and semialgebraic on $\mathcal{K} \times \mathcal{K}$. So is $(c, c') \mapsto \delta(c, c')$: it is the minimum of the polynomial $\|c - R \cdot c'\|^2$ over the compact semialgebraic set $\text{SO}(3)$, hence semialgebraic by the Tarski–Seidenberg theorem, and continuous by compactness. By separation the two functions have the same zero set, namely the pairs in one orbit. The Łojasiewicz inequality for continuous semialgebraic functions on a compact semialgebraic set [6] states that if $g^{-1}(0) \subset f^{-1}(0)$, then $|f| \leq C|g|^\alpha$ for some $C > 0$ and $\alpha > 0$. Applied to $f = \delta$ and $g = |\mathcal{I}(c) - \mathcal{I}(c')|$ on $B' \times B'$, where $B' \supset \mathcal{K}$ is a closed ball, which is compact semialgebraic, and restricted to $\mathcal{K} \times \mathcal{K}$, this gives the bound; since g is bounded on $\mathcal{K} \times \mathcal{K}$, α may be decreased to lie in $(0, 1]$.

(iv) Let G be the group acting, $H = G_c$ the stabilizer of c , and N the orthogonal complement at c of the tangent space of $G \cdot c$; for small $r > 0$ let $\Sigma = \{c + n : n \in N, |n| < r\}$. By the slice theorem the map $G \times_H \Sigma \rightarrow V$, $[g, \sigma] \mapsto g \cdot \sigma$, is a diffeomorphism onto an open G -invariant neighborhood U of $G \cdot c$, and $G_\sigma \subset H$ for $\sigma \in \Sigma$. Since c lies on the principal stratum, G_σ is conjugate to H ; a compact Lie group is not conjugate to a proper closed subgroup of itself, which would have the same dimension and the same number of components, so $G_\sigma = H$. Hence H fixes Σ pointwise, $G \times_H \Sigma = (G/H) \times \Sigma$, and $s(g \cdot \sigma) = \sigma$ defines a smooth G -invariant map $s: U \rightarrow \Sigma$ with $s|_\Sigma = \text{id}$. Let $B \subset U$ be a closed ball of radius ρ about c and C_s a Lipschitz constant of s on B . For $\sigma, \sigma' \in \Sigma$ within $\rho/4$ of c , let $g \in G$ attain $\delta(\sigma, \sigma') = |\sigma - g \cdot \sigma'|$. Then $|\sigma - g \cdot \sigma'| \leq |\sigma - \sigma'| \leq \rho/2$, so the segment from σ to $g \cdot \sigma'$ lies in B , and

$$|\sigma - \sigma'| = |s(\sigma) - s(g \cdot \sigma')| \leq C_s \delta(\sigma, \sigma') \leq C_s |\sigma - \sigma'|.$$

Thus near c the orbit distance is bi-Lipschitz equivalent to the Euclidean distance of Σ , and Σ is a chart of the quotient. If the Jacobian of $\mathcal{I}|_\Sigma$ at c has full rank, $\mathcal{I}|_\Sigma$ is a diffeomorphism from a neighborhood of c in Σ onto its image, hence bi-Lipschitz on a compact neighborhood. Since $\mathcal{I}$ and δ are G -invariant and every point of U is $g \cdot \sigma$ with $\sigma \in \Sigma$, this gives $|\mathcal{I}(x) - \mathcal{I}(x')| \geq c_0 \delta(x, x')$ for x, x' in a G -invariant neighborhood of $G \cdot c$, which with (ii) is (iv). $\square$

Parts (ii) and (iii) are the two directions of stability. Part (ii) says that small changes of shape, measured in the orbit distance, produce small changes of the invariants; this is what makes the invariants usable on noisy meshes. Part (iii) says that small changes of the invariants can only come from small changes of shape modulo the nuisance group, uniformly on bounded sets. Together they say that $\mathcal{I}$ induces a homeomorphism from the image of $\mathcal{K}$ in $V_L/\text{SO}(3)$ onto its image which is Lipschitz and has a Hölder continuous inverse.

The exponent in (iii) cannot be taken to be 1 in general. The obstruction is not a defect of the invariants but a property of the quotient.

Example 7.4. Let $V = \mathcal{H}_1$ with $\mathcal{I}(a) = |a|^2$, and let $\mathcal{K}$ be the unit ball. By Theorem 7.1, $\delta(a, 0) = |a|$, while $|\mathcal{I}(a) - \mathcal{I}(0)| = |a|^2$. Hence $\delta(a, 0) = |\mathcal{I}(a) - \mathcal{I}(0)|^{1/2}$, and no inequality $\delta \leq C|\Delta\mathcal{I}|^\alpha$ with $\alpha > 1/2$ can hold near the origin. The same happens in every V_L : all invariants other than f_0 have degree at least two and vanish to second order at $c = 0$, while $\delta(c, 0) = \|c\|$ vanishes to first order. So $\alpha_{\mathcal{K}} \leq 1/2$ whenever $\mathcal{K}$ contains a neighborhood of a point c_0 whose stabilizer is not the principal one: every invariant is stationary at c_0 in the directions of the slice on which the stabilizer acts without fixed vectors, while the orbit distance is

of first order there; and α_K can be smaller at points where higher-degree invariants are needed to separate.

The frame coordinates of Theorem 6.2 satisfy (ii) with constants that blow up like the reciprocal of the spectral gap near the strata; the polynomial invariants do not, which is the reason for using them on data whose spectral gaps are not controlled.

Remark 7.5. The theorem concerns the passage from coefficients to invariants. The passage from surface to coefficients is also stable: the uniformizing map, taken modulo Möb, depends continuously on the metric in $C^{k,\alpha}$, so that after centering the parameterization depends continuously on the metric up to $\text{SO}(3)$, and the conformal barycenter is a smooth function of the measure on the set of stable measures, with derivative controlled by the Hessian computed in [9]. A quantitative statement in terms of mesh perturbations is left open.

7.3. The certified descriptor. We now collect what has been proved into a single statement about surfaces. Fix $L \geq 1$ and a multihomogeneous generating set of $\mathcal{R}_L$ with phased real forms $I_1, \dots, I_N$ of total degrees $w_1, \dots, w_N$, as in Theorem 4.1, with the $\text{O}(3)$ -invariants listed first and the pseudo-invariants last. For a smoothly embedded two-sphere $S \subset \mathbb{R}^3$ define

$$(30) \quad \mathcal{D}_L(S) = \mathcal{I}(f^{(L)}(\varphi)) \in \mathbb{R}^N, \quad \varphi \in \Phi_0(S),$$

and, when $f^{(L)}(\varphi) \neq 0$, its class $[\mathcal{D}_L(S)] \in \mathbb{P}_w^+(\mathbb{R})$.

Theorem 7.6. *The following are true:*

- (a) **Well defined.** $\mathcal{D}_L(S)$ does not depend on the choice of $\varphi \in \Phi_0(S)$.
- (b) **Equivariance.** If $T(x) = \lambda Rx + b$ with $\lambda > 0$ and $R \in \text{SO}(3)$, then $\mathcal{D}_L(T(S))_k = \lambda^{w_k} \mathcal{D}_L(S)_k$ for every k , so $[\mathcal{D}_L(T(S))] = [\mathcal{D}_L(S)]$. If T is an orientation-reversing similarity, the same holds for the $\text{O}(3)$ -invariant coordinates, while each pseudo-invariant coordinate is multiplied by $-\lambda^{w_k}$.
- (c) **Completeness.** $\mathcal{D}_L(S) = \mathcal{D}_L(S')$ if and only if the truncated radial expansions of S and S' differ by a rotation of the sphere: $f^{(L)}(\varphi') = Q \cdot f^{(L)}(\varphi)$ for some $\varphi \in \Phi_0(S)$, $\varphi' \in \Phi_0(S')$ and $Q \in \text{SO}(3)$. Likewise $[\mathcal{D}_L(S)] = [\mathcal{D}_L(S')]$ if and only if the truncated expansions differ by a rotation and a positive scaling.
- (d) **Chirality.** S and its mirror image have radial descriptors in the same $\text{SO}(3)$ -orbit if and only if every pseudo-invariant coordinate of $\mathcal{D}_L(S)$ vanishes.
- (e) **Stability.** On every bounded set of descriptors, $\mathcal{D}_L$ is Lipschitz with respect to the orbit distance, and on every compact set the orbit distance is bounded by a Hölder power of $|\mathcal{D}_L(S) - \mathcal{D}_L(S')|$.

For the coordinate model, with $\mathcal{I}_2$ in place of $\mathcal{I}$ and $C^{(L)}(\varphi)$ in place of $f^{(L)}(\varphi)$, (a)–(e) hold with $\text{SO}(3) \times \text{SO}(3)$ in place of $\text{SO}(3)$ and the mirror pseudo-invariants of Theorem 4.4 in (d), and in addition: if $\mathcal{I}_2(C^{(L)}(\varphi)) = \mathcal{I}_2(C^{(L)}(\varphi'))$ for every L , with $\mathcal{I}_2$ chosen for each L , then S' is the image of S under an orientation-preserving rigid motion.

Proof. (a) By Theorem 2.3(i) the tuples $f^{(L)}(\varphi)$, $\varphi \in \Phi_0(S)$, form one $\text{SO}(3)$ -orbit, and $\mathcal{I}$ is $\text{SO}(3)$ -invariant. (b) By Theorem 2.3(iii), $f^{(L)}(T \circ \varphi) = \lambda f^{(L)}(\varphi)$, and $\mathcal{I}_k$ is homogeneous of degree w_k . For an orientation-reversing T , Theorem 2.3(iv)

gives $f_l(T \circ \varphi \circ r) = \lambda f_l(\varphi) \circ r$, and $f_l \circ r$ lies in the $\mathrm{SO}(3)$ -orbit of $(-1)^l f_l$ since $r = -R_0$ for a rotation R_0 ; the effect on I_k is $(-1)^{\sum l d_l} \lambda^{w_k}$ by Theorem 3.6(iv). (c) is Theorem 4.1 together with (a), and the scale-free statement is the completeness argument of Section 4.2. (d) is Theorem 5.1. (e) is Theorem 7.3(ii) and (iii), applied to the descriptors $f^{(L)}(\varphi)$ and $f^{(L)}(\varphi')$. The coordinate statements are Theorem 2.3(ii), Theorem 4.5, Theorem 4.4 and Theorem 7.3 for $\mathrm{SO}(3) \times \mathrm{SO}(3)$; the reconstruction from all bands is the last assertion of Theorem 4.5. $\square$

The theorem is the precise form of the claim made in the introduction. Every nuisance freedom of a spherical harmonic description of a surface has been removed exactly: the parameterization by (a), the position and scale by (b) and (c), and the handedness is not discarded but recorded by (d). Nothing has been lost at the level of the truncated coefficients, since (c) says the descriptor is complete for the truncation, and the coordinate model is complete in the limit; the radial reduction and the truncation themselves may lose information about S . And the result is not only correct on exact data but stable with respect to the coefficients of measured data, by (e). What remains is the numerical realization of each step on triangulated surfaces, which Section 8 addresses.

8. COMPUTATION AND VERIFIED ALGEBRA

The constructions of the previous sections are exact. This section says how each is carried out on a triangulated surface, which quantities must be reported so that a computed descriptor can be trusted, and which algebraic identities were verified independently of any mesh.

8.1. From a mesh to a centered parameterization. The input is a triangulated surface S_h with vertices $p_1, \dots, p_n \in \mathbb{R}^3$. It must be a connected closed combinatorial manifold with $n - e + t = 2$, free of degenerate faces, and consistently oriented; its realization in $\mathbb{R}^3$ is assumed free of self-intersections, so that S_h approximates a smoothly embedded two-sphere. The first step is a discrete conformal map $\psi_h: S_h \rightarrow \mathbb{S}^2$, $p_i \mapsto u_i$, without flipped triangles, obtained by minimizing a discrete conformal energy on the sphere [17, 11]; its residual freedom is, up to discretization error, a Möbius transformation.

The second step is the discrete form of Theorem 2.1. The pulled-back area measure is $\mu_h = \sum_i w_i \delta_{u_i}$ with w_i the fraction of the area of S_h attributed to p_i . By Theorem 2.2 it has a unique conformal barycenter x_h provided $\max_i w_i < 1/2$, which must be checked. The point x_h is the zero of $\xi_h(x) = \sum_i w_i \tau_x(u_i)$; it is computed by Newton's method with a line search, started at $x = 0$ and monitored for convergence, or by the fixed-point iteration of [3]. The Jacobian of ξ_h , computed in [9], governs the convergence rate and the sensitivity of x_h to the mesh. Each u_i is then replaced by $\tau_{x_h}(u_i)$, the residual $|\sum_i w_i u_i|$ is recorded, and the remaining rotation of $\mathbb{S}^2$ is left to the invariants.

8.2. Harmonic coefficients. The coefficients are integrals against $d\omega$, not against the area of S_h , which a conformal map concentrates in some regions of the sphere. Let the round weight ω_i be one third of the area of the spherical triangles incident to u_i , divided by 4π . The conformal center (3) is approximated by $\bar{c}_h = \sum_i \omega_i p_i$, and $X_i = p_i - \bar{c}_h$, $d_i = |X_i|$. With Y the $n \times (L+1)^2$ matrix of basis values $Y_{lm}(u_i)$ and $\Omega = \mathrm{diag}(\omega_i)$, a sample vector s has coefficients $c = (Y^\top \Omega Y)^{-1} Y^\top \Omega s$, for $s = (d_i)$ in

the radial model and $s = (X_i^{(k)})$, $k = 1, 2, 3$, in the coordinate model. The matrix $Y^\top \Omega Y$ approximates the identity, and its deviation, together with $\|C_0\|$, which vanishes in exact arithmetic, measures the quadrature. The truncation degree is chosen from the residual, compared with the bound $|S_h|/(2\pi(L+1)(L+2))$ of Theorem 2.4; the largest ratio of a face area on S_h to the area of its image, the discrete $\sup e^{2\rho}$, is reported with it.

8.3. Binary forms and invariants. The binomial coefficients of F_{f_l} are computed from the c_{lm} by Theorem 3.8(iii). As an independent check, f_l is written as a solid harmonic polynomial in x and substituted into (7); coefficients from a software library must be converted with the library's own phase and normalization at this step. Invariants have degrees up to 15, so each is evaluated on $c/\|c\|$ and rescaled, $I_k(c) = \|c\|^{w_k} I_k(c/\|c\|)$, with $\|c\|$ kept as the size variable. The transvectant sequence for R (22) and for the joint invariants (27) is implemented once, verified in exact rational arithmetic (Section 8.5), and then run in floating point; the spurious imaginary parts of $J_2, \dots, J_{10}$ and real part of R measure the floating-point error, by Theorem 3.6. The frame coordinates of Theorem 6.2 are used only when the relative spectral gaps of Q (or the singular value gaps of C_1) exceed a stated threshold; by Theorem 6.2(c) the threshold only decides where the discontinuity of the frame is placed.

8.4. Normalization and diagnostics. The invariant vector is normalized by the gauge (17), and the quantities $|\hat{x}_k|^{1/w_k} \in [0, 1]$ are reported; a value near zero flags proximity to the divisor $I_k = 0$. Each of the following diagnostics corresponds to a hypothesis or a bound of the theory.

- (i) $\max_i w_i < 1/2$ (Theorem 2.2) and the centering residual.
- (ii) The deviation of $Y^\top \Omega Y$ from the identity, and $\|C_0\|$.
- (iii) The truncation residual, its ratio to the bound of Theorem 2.4, and $\sup e^{2\rho}$.
- (iv) The spectral gaps of Theorem 6.3.
- (v) The orbit distance (29) between the descriptors of two independently processed copies of the same mesh, one rotated by a random $R \in \text{SO}(3)$ and reparameterized by a random Möbius transformation. The exact value is zero by Theorem 2.3, so the computed value measures the numerical error of the whole pipeline on two congruent discretizations, free of the nuisance rotations.

Diagnostic (v) is the end-to-end test; if it fails, (i)–(iv) locate the error.

8.5. Verified identities. The following were checked independently of any mesh, by exact rational arithmetic where the inputs are rational and to machine precision otherwise, with the script provided as Online Resource 1; random inputs were drawn with fixed seeds. They are the unit tests an implementation should reproduce before it is run on data.

- (1) **Conventions** (Sections 2 and 3). The Y_{lm} are orthonormal through $l = 5$ to $3 \cdot 10^{-14}$; $q(v(z)) = 0$, $\text{disc } Q_x = 4q(x)$, $v(jz) = -\overline{v(z)}$; $\rho(g)$ from (10) is orthogonal of determinant one on $ad - bc = 1$, and real with $m(gz) = \rho(g)m(z)$ for $g \in \text{SU}(2)$; for random real harmonics of degrees 1, 2, 3, $a_{2l-k} = (-1)^{l+k} \bar{a}_k$. An implementation with a different sign in (7) or a different basis phase fails these tests.
- (2) **Correspondence** (Theorem 3.8). The formulas (iii) and the norm identity (iv) hold exactly for $l \leq 3$, and $J_2(F_{f_3}) = 2100\|f_3\|^2$.

- (3) **Parity** (Theorem 3.6). On $\lambda Q_a Q_b Q_c$ with rational a, b, c, λ , $J_2, \dots, J_{10}$ are rational and R is a rational multiple of i ; $P_{123} = ((F_1, F_2)_1, (F_3, F_3)_4)_4$ is purely imaginary and nonzero.
- (4) **Degree two** (Theorems 4.8 and 4.9). For every rational traceless symmetric Q , $I(F_f) = 2 \operatorname{tr} Q^2$, $J(F_f) = -4 \det Q$ and $(Q_a^2, F_f)_4 = 4a^\top Q a$; $\varepsilon^2 = \det \operatorname{Gram}(a, Qa, Q^2 a)$.
- (5) **Degree three** (Theorem 5.2). $R = 0$ exactly on coplanar and bisector configurations and $R \neq 0$ on generic ones; $R \mapsto -R$ under $a \mapsto -a$; R scales by 2^{15} when one axis is doubled. The invariants (24) agree with the root sums (20); the forms of weighted degree 30 vanishing on (24) span a line, generated by the polynomial F_2 of [28]; and $R^2/F_2 = -2^{18}3^{-18}5^{-20}$ exactly on random integer axis triples.
- (6) **Bispectrum on V_2** (Section 9.1). With bands in $\{1, 2\}$, the invariants (27) are $4a^\top Q a$ for $(1, 1, 2)$, $(1, 2, 1)$, $(2, 1, 1)$ and $-24 \det Q$ for $(2, 2, 2)$, and the others, including B_{122} , vanish identically.
- (7) **Metrics** (Theorem 7.2, Figure 1). A multistart minimization over $\operatorname{SO}(3) \times \operatorname{SO}(3)$ agrees with $|s - s'|$ to six digits for both determinant signs; the barycenter of the measure of Figure 1 is found with residual below 10^{-16} .

8.6. **A synthetic end-to-end test.** Diagnostic (v) was run, with the script provided as Online Resource 2, on the graph $u \mapsto r(u)u$ over $\mathbb{S}^2$ with

$$\begin{aligned}
 r = & 1 + \frac{1}{4}x_3 + \frac{7}{20}(x_1^2 - \frac{1}{2}x_2^2) + \frac{1}{4}x_1x_3 \\
 & + \frac{1}{5}x_1x_2x_3 + \frac{3}{20}(x_3^3 - \frac{3}{5}x_3) \\
 & + \frac{3}{25}(x_1^3 - 3x_1x_2^2)x_3 - \frac{1}{10}x_2x_3^2,
 \end{aligned}$$

meshed on the icosahedral subdivision with $n = 10242$ vertices and 20480 faces. It is chiral and has no continuous symmetry. Copy A is this mesh. Copy B is the mesh rotated by a random $R \in \operatorname{SO}(3)$, with its vertices relabeled by a random permutation and its initial spherical map composed with a random Möbius transformation of translation length 0.45. Each copy was processed independently: the conformal map by minimization of the harmonic energy with cotangent weights, the barycenter by Newton's method, the coefficients through $L = 6$ with round weights, and the invariants of Theorem 4.9 and of band three by the code of Online Resource 1. Table 2 lists the diagnostics.

The exact orbit distance between the two copies is zero. The cotangent weights are intrinsic, so the discrete energy and the sphere constraint are invariant under rigid motions and relabelings; when the discrete minimizer is unique up to a rotation of $\mathbb{S}^2$, the two copies share it, the discretization error is common to both, and the computed distance $2 \cdot 10^{-8}$ is the optimization error of the harmonic map and of the minimization in (29). The invariants agree to the same order. To first order in δ the ratios in the last row equal $|\cos \angle(\nabla I_k(c), c - R \cdot c')|$, so they are at most $1 + O(\delta)$ by Cauchy–Schwarz; their being below one is consistent with, but not a test of, Theorem 7.3(ii), which uses $\sup_B |\nabla I_k|$. The mirror image of the mesh, processed independently, reproduces the six $\operatorname{O}(3)$ -invariants of Theorem 4.9 and $J_2, \dots, J_{10}$ to relative 10^{-6} and reverses the signs of ε and $-i\tilde{R}$, as Theorem 7.6(d) requires. The same surface meshed with $n = 2562$ vertices has orbit distance $\delta/\|c\| = 1.1 \cdot 10^{-4}$ from copy A, and its invariants differ by relative $3 \cdot 10^{-4}$ to $3 \cdot 10^{-3}$ in $J_2, \dots, J_{10}$ and by up to $2 \cdot 10^{-2}$ in the dipole invariants, whose magnitude here is $|a|^2 = 4.7 \cdot 10^{-4}$.

TABLE 2. Diagnostics of Section 8.4 for two independently processed copies of one synthetic mesh. Copy B agrees with copy A to the digits shown. The last row is a first-order sensitivity ratio; see the text.

diagnostic	value (copy A)
(i) maximal weight $\max_i w_i$; centering residual $ \sum_i w_i u_i $	$1.9 \cdot 10^{-4}$; $6 \cdot 10^{-16}$
quasi-conformal distortion of the discrete map, max / mean (additional)	1.030 / 1.009
(ii) $\ Y^\top \Omega Y - 1\ _2$; $\ C_0\ $	$1.4 \cdot 10^{-3}$; $2.3 \cdot 10^{-5}$
(iii) truncation residual at $L = 6$; bound of Theorem 2.4; ratio $\sup e^{2\rho}$	$2.5 \cdot 10^{-5}$; $4.5 \cdot 10^{-2}$; $5.6 \cdot 10^{-4}$ 3.10
(iv) quadrupole eigenvalues	-0.2186, -0.0843, 0.3029
(v) orbit distance $\delta(c, c')/\ c\ $, radial model, $L = 3$ / $L = 6$	$1.8 \cdot 10^{-8}$ / $2.0 \cdot 10^{-8}$
$\delta(C, C')/\ C\ $, coordinate model, $L = 3$ / $L = 6$	$8.7 \cdot 10^{-8}$ / $9.2 \cdot 10^{-8}$
seven invariants of Theorem 4.9, relative difference A vs. B	10^{-8} to $8 \cdot 10^{-7}$
J_2, J_4, J_6, J_{10} , relative difference	$2 \cdot 10^{-7}$ to $2 \cdot 10^{-6}$
$-i\bar{R}$, relative difference	$2.5 \cdot 10^{-5}$
$ I_k(c) - I_k(c') /(\nabla I_k(c) \delta(c, c'))$, $k = 1, \dots, 7$	all ≤ 0.83

The discretization error thus exceeds the error of the quotient by four orders of magnitude; it is the quantity a mesh-level validation must control.

9. BENCHMARKS ON HARMONIC COEFFICIENTS

Section 8.6 shows that on meshes the discretization error dominates the error of the quotient. To test the separation and stability statements themselves, the experiments of this section work directly on harmonic coefficients in V_2 , where the descriptor can be evaluated exactly and the orbit distance computed to high accuracy. A dipole-quadrupole pair is written $c = (a, Q)$ as in Theorem 4.9, with the constant band omitted. Rotations act by $(a, Q) \mapsto (Ra, RQR^\top)$, and all norms are the $L^2(d\omega)$ norms of the harmonics, $\|f_1\|^2 = |a|^2/3$ and $\|f_2\|^2 = \frac{2}{15} \text{tr } Q^2$. The scripts are Online Resource 3. Figure 4 shows the results.

9.1. Discrimination. Let $Q_0 = \text{diag}(\lambda)$ with $\lambda = (0.7, 0.1, -0.8)$, and put $d = (\lambda_2 - \lambda_3, \lambda_3 - \lambda_1, \lambda_1 - \lambda_2)$, so that $\sum_i d_i = \sum_i \lambda_i d_i = 0$. Let a_A and a_B have components $\sqrt{w_i}$ with $w = (0.10, 0.20, 0.10)$ and $w + 0.06d$, respectively. The pairs $A = (a_A, Q_0)$ and $B = (a_B, Q_0)$ then have the same $|a|^2$, $\text{tr } Q^2$, $\det Q$ and $a^\top Q a$, but different $a^\top Q^2 a$ (0.115 against 0.164). Their mirror images are $A^* = (-a_A, Q_0)$ and $B^* = (-a_B, Q_0)$ by Theorem 3.6(iv); they have the same $O(3)$ -invariants as A and B and the opposite sign of ε . Each sample is a class representative rotated by a uniformly random $R \in SO(3)$, with independent Gaussian noise of standard deviation $\sigma\|c\|$ added to each coefficient in the orthonormal basis $\{Y_{lm}\}$.

Five descriptors were compared:




FIGURE 4. Benchmarks of Section 9. (a) Nearest-neighbour accuracy on the four classes A, A^*, B, B^* of Section 9.1 as a function of coefficient noise, mean and standard deviation over five repetitions; chance is $1/4$. (b) Local Lipschitz ratio $|\Delta D|/\delta$ of the frame coordinates of Theorem 6.2 and of the polynomial invariants of Theorem 4.9, as the gap between two eigenvalues of the quadrupole tends to zero. (c) Change of the invariants of Theorem 4.9 against the exact orbit distance at a principal point, at a point with stabilizer $SO(2)$ (perturbed in the directions of its slice moved by the stabilizer), and at the origin, with fitted slopes.

- (i) the power spectrum $(\|f_1\|, \|f_2\|)$;
- (ii) the bispectrum, that is, the invariants (27) with bands $l \leq 2$;
- (iii) the five nonconstant $O(3)$ -invariants of Theorem 4.9;
- (iv) all seven invariants of Theorem 4.9;
- (v) the frame coordinates of Theorem 6.2(a), that is, the K -invariant monomials $\lambda_1, \lambda_2, \tilde{a}_1^2, \tilde{a}_2^2, \tilde{a}_3^2, \tilde{a}_1\tilde{a}_2\tilde{a}_3$ in the eigenframe-aligned dipole $\tilde{a}$.

On V_2 the bispectral invariants are, up to the constant band, $4a^\top Qa$ for the band triples $(1, 1, 2)$, $(1, 2, 1)$, $(2, 1, 1)$ and $-24 \det Q$ for $(2, 2, 2)$, and all others vanish, including B_{122} , the only candidate pseudo-invariant (Section 8.5). Each feature x_k of degree w_k was replaced by $\text{sgn}(x_k)|x_k|^{1/w_k}$ and standardized on the training set, and each class was represented by 100 training and 400 test samples in a nearest-neighbour classifier.

Figure 4(a) shows the outcome. The power spectrum and the bispectrum stay at chance at every noise level, including $\sigma = 0$, because they are functions of invariants that coincide on all four classes. The $O(3)$ -invariants reach one half: they separate A from B but not a class from its mirror image, as Theorem 5.1 predicts. The seven invariants of Theorem 4.9 and the frame coordinates classify all four classes without error at $\sigma = 0$ and with accuracy above 0.97 at $\sigma = 0.02$. For larger noise all descriptors degrade, and the ordering is unchanged. The per-coefficient level $\sigma = 0.05$ corresponds to a noise vector of about 14% of $\|c\|$, comparable to the separation $|\varepsilon| \approx 0.037$ between mirror classes.

9.2. Frames near the degenerate strata. Let $Q_g = \text{diag}(\frac{1}{3} + \frac{g}{2}, \frac{1}{3} - \frac{g}{2}, -\frac{2}{3})$ and $a = (0.3, 0.4, 0.5)$, so that $c_g = (a, Q_g)$ tends to the axial stratum $\lambda_1 = \lambda_2$ as $g \rightarrow 0$. For each g , 300 perturbations $c_g + \eta u$ with $\eta = 10^{-7}$ and u a random unit vector normal to the orbit were drawn; to first order $\delta(c_g, c_g + \eta u) = \eta$. Figure 4(b)

shows the largest ratio $|D(c_g + \eta u) - D(c_g)|/\eta$ for the frame coordinates and for the invariants of Theorem 4.9. For the frame the ratio grows like $0.65/g$, with fitted slope -1.00 over $10^{-4} \leq g \leq 10^{-2}$, which is the conditioning predicted in Theorem 6.3 and reflects the discontinuity of Theorem 6.2(c). For the polynomial invariants the ratio stays between 4.2 and 5.7 over four decades of g , in accordance with Theorem 7.3(ii). This is the reason for preferring the polynomial invariants on data whose spectral gaps are not controlled.

9.3. Stability exponents. At three base points c_0 , perturbations $c = c_0 + \eta u$ with $3 \cdot 10^{-4} \leq \eta \leq 0.3$ and u normal to the orbit were drawn, and $\delta(c_0, c)$ was computed by multistart minimization over $\text{SO}(3)$; at the smallest η it agreed with η to four digits. The base points were the principal point A ; the point $(0, 0, \frac{1}{2})$, $\text{diag}(\frac{1}{2}, \frac{1}{2}, -1)$, whose stabilizer is the group $\text{SO}(2)$ of rotations about the x_3 -axis, perturbed in the directions of its slice that this group moves; and the origin. Figure 4(c) shows the median of $\|\mathcal{I}(c) - \mathcal{I}(c_0)\|$ against δ , with $\mathcal{I}$ the seven invariants of Theorem 4.9. The fitted slopes are 0.96, 1.97 and 2.00. At the principal point the invariants are locally bi-Lipschitz, as Theorem 7.3(iv) asserts. At the other two points the invariant discrepancy is quadratic in the orbit distance, so the exponent α_K of Theorem 7.3(iii) cannot exceed $1/2$ on a neighborhood of them, as Theorem 7.4 asserts.

9.4. Paired anatomical structures. The last experiment runs the mesh pipeline of Section 8 on real anatomy. The meshes are the eleven bilateral pairs of Body-Parts3D, a public anatomical atlas of one adult male: hippocampus, amygdala, putamen, globus pallidus, thalamus, parahippocampal gyrus, testis, kidney, adrenal gland, caudate nucleus and lateral ventricle. All 22 meshes pass the preflight of Section 8.1. Each was remeshed isotropically to 2800–2900 vertices, since the atlas meshes contain triangles with angles below 2° ; the remeshed surfaces pass the preflight again. The spherical map is the harmonic map of the punctured surface to the plane [11] followed by the harmonic-energy flow with cotangent weights of Section 8.1, with Möbius centering after every ten steps. Coefficients through $L = 6$ of the radial model were computed with round weights and divided by the mean radius f_0 , so that the descriptor is scale-free; $\|c\|$ denotes the norm of bands 1 to 6 of the left structure, and all distances below are divided by it. On the synthetic surface of Section 8.6 this implementation reproduces the quadrupole eigenvalues and the dipole norm of Table 2 to three digits. The scripts are Online Resource 4.

For each pair, four meshes were processed independently: the left structure L ; the right structure R ; the reflection R^* of the right structure in a plane, with the face orientation restored; and a copy L' of the left structure moved by a random rigid motion, with its vertices relabeled and its initial spherical map composed with a random Möbius transformation. The comparison L against L' is diagnostic (v) of Section 8.4 on a real mesh. The comparison of R^* with the tuple $((-1)^l f_l(R))_l$ tests Theorem 2.3(iv) end to end; the two agree to $\delta/\|c\|$ between $4 \cdot 10^{-4}$ and $2 \cdot 10^{-3}$, the discretization error of remeshing a reflected mesh. The **chirality index** $\chi = \delta(c, c^*)/\|c\|$, the orbit distance of a shape from its mirror image, vanishes exactly on achiral shapes by Theorem 7.6(d). Table 3 lists the results.

Three groups appear. The diagnostics reject the caudate nucleus and the lateral ventricle: on these long, curved structures the discrete conformal map folds, with

TABLE 3. Eleven bilateral pairs of BodyParts3D through the pipeline of Section 8, radial model, $L = 6$. flips: faces with reversed orientation on the sphere; $K_{\max}$: largest quasi-conformal distortion of a face; $\sup e^{2\rho}$: largest ratio of a face area to the area of its spherical image (diagnostic (iii)). L' is a rigidly moved, re-labeled, Möbius-perturbed copy of L , and R^* the reflected right structure; all orbit distances are divided by $\|c\|$. χ is the chirality index. The last two columns give the signs of ε and of $-i\tilde{R}$ on the left and right side; a sign in parentheses belongs to a value whose gauge coordinate (17) is below 0.1, that is, within numerical distance of the divisor.

structure	flips	$K_{\max}$	$\sup e^{2\rho}$	$\delta(L, L')$	$\delta(L, R)$	$\delta(L, R^*)$	χ_L	χ_R	$\text{sgn } \varepsilon$	$\text{sgn}(-i\tilde{R})$
hippocampus	0	2.19	$8.2 \cdot 10^7$	$5.1 \cdot 10^{-14}$	0.1225	0.0088	0.119	0.126	+/-	-/+
amygdala	0	1.42	140	$2.8 \cdot 10^{-13}$	0.3411	0.0039	0.341	0.342	-/+	-/+
putamen	0	1.32	18	$1.1 \cdot 10^{-6}$	0.1729	0.0049	0.173	0.173	+/-	+/-
globus pallidus	0	1.23	26	$6.2 \cdot 10^{-14}$	0.3161	0.0009	0.316	0.317	+/-	-/+
thalamus	0	1.06	5.6	$4.9 \cdot 10^{-14}$	0.1940	0.0015	0.194	0.194	-/+	+/-
parahippocampal gyrus	0	1.23	$1.3 \cdot 10^3$	$1.8 \cdot 10^{-4}$	0.1890	0.0049	0.188	0.190	+/-	-/+
testis	0	1.05	1.7	$7.4 \cdot 10^{-14}$	0.2651	0.0617	0.230	0.290	(-)/(-)	+/-
kidney	0	1.28	9.5	$2.5 \cdot 10^{-14}$	0.1672	0.1797	0.097	0.123	-/-	-/+
adrenal gland	0	1.43	$6.6 \cdot 10^3$	$4.0 \cdot 10^{-14}$	0.3784	0.3754	0.131	0.348	+/-	+/-
caudate nucleus	439	$> 10^4$	$6.9 \cdot 10^{14}$	parameterization rejected by diagnostic (iii)/(v): folded map						
lateral ventricle	426	$> 10^4$	$1.1 \cdot 10^{11}$	parameterization rejected by diagnostic (iii)/(v): folded map						

about 400 reversed faces and $\sup e^{2\rho}$ above 10^{11} . This is the crowding of Theorem 2.4 at its extreme; a conformal map compresses a thin tail into a region of the sphere too small to be represented by the mesh, and the diagnostics report it rather than a descriptor. The hippocampus is at the boundary: its map does not fold and has $K_{\max} = 2.2$, but $\sup e^{2\rho} = 8 \cdot 10^7$, so the second bound of Theorem 2.4 is uninformative for it while the first still applies.

For the seven remaining pairs of brain and gonadal structures, the left structure is at distance 0.12 to 0.34 from the right one but only 0.001 to 0.009 from the reflected right one, a ratio between 14 and 350 (testis: 4), and $\delta(L, R)$ agrees with the chirality index χ to two digits. These pairs are one chiral shape in two handednesses: the whole left-right distance is the distance of the shape from its mirror image, and the residual after reflection measures the actual asymmetry. The pseudo-invariants record the handedness: $-i\tilde{R}$ has opposite signs on the two sides in all seven pairs, and ε in six; in the seventh, the testis, ε lies within numerical distance of its zero divisor, as the gauge coordinate reports, and its sign is not informative. For the kidney and the adrenal gland, reflection does not reduce the distance, $\delta(L, R^*) \approx \delta(L, R)$, and the two sides have different chirality indices. These are pairs of different shapes rather than mirror images, which is the anatomical situation for the adrenal glands. The descriptor thus separates two kinds of laterality, the same shape in opposite handedness and different shapes, that raw coefficients or $O(3)$ -invariants confound.

Finally, $\delta(L, L')$ is at machine precision, $3 \cdot 10^{-13}$ or below, for seven of the nine valid pairs. For the putamen it is 10^{-6} and for the parahippocampal gyrus $2 \cdot 10^{-4}$, and on these two shapes the value varies with the random Möbius initialization: the harmonic flow reaches its iteration limit before converging fully, which is the optimization error that diagnostic (v) is designed to measure. This experiment uses

a single atlas subject and makes no anatomical claim; a cohort would be needed for that. What it shows is that the pipeline runs on real meshes with its diagnostics reporting where it is reliable, and that the parity structure of Section 5 is visible in real anatomy.

The experiments give a consistent picture. The power spectrum and the bispectrum are stable but incomplete and blind to reflections. Frames are complete on the principal stratum but ill-conditioned near its boundary. The polynomial invariants are complete, record chirality, and are uniformly Lipschitz; their inverse is Lipschitz on the principal stratum and Hölder of exponent $1/2$ at the strata where the stabilizer jumps. On real anatomy the same invariants separate mirror-image pairs from asymmetric pairs, and the diagnostics identify the surfaces on which the conformal parameterization cannot be trusted.

10. DISCUSSION

The central contribution is a quotient-theoretic formulation of spherical-harmonic shape analysis for genus-zero surfaces in which every nuisance freedom is a group action and every step is exact: conformal barycenter normalization for the parameterization, polynomial invariants of a compact group for rotation, weighted projectivization for scale. Within that formulation we have proved a complete descriptor through degree two, characterized chirality in degree three and tied it to the locus $\mathcal{L}_2$ of genus-two curves with an elliptic involution, constructed separating frame coordinates on the principal strata of both models with the residual Klein four-group made explicit, recorded the size of continuous separating invariants, and proved the metric and stability statements that certify the descriptor.

SPHARM-COM represents a closed surface by the distance from the center of mass, expanded in spherical harmonics through degree 30 with respect to an approximately area-preserving parameterization, followed by supervised selection of individual coefficients, which concentrated in bands 1 through 6. Replacing the coordinate functions by the distance removes ambient rotation and reduces three channels to one, but it does not remove the reparameterization freedom: with an area-preserving parameterization that freedom is infinite-dimensional, and even after a canonical parameterization a residual $\mathrm{SO}(3)$ remains. An individual coefficient is therefore a function of the map, not of the surface. Coefficients computed from an area-preserving parameterization cannot be converted into the descriptor; the surface must be reparameterized and centered as in Section 8. The present construction retains the same kind of scalar representation in the radial model, the distance from a center, and quotients exactly the freedoms that remained; for the bands actually used, Theorem 4.9 gives a complete descriptor of the truncation through degree two with seven invariants, the sextic invariants and $\hat{R}$ complete band three as an isolated band, and Theorem 6.2(a) gives a complete real-analytic descriptor on the principal stratum through any degree. The coordinate model additionally retains the direction information the radial reduction discards and is faithful in the all-band limit. The pseudo-invariants isolate mirror asymmetry, which an analysis of raw radial coefficients neither identifies nor controls, and Theorem 4.2 shows that coefficient-space interpolation, used there for augmentation, is not intrinsic on the quotient. Methodological details of that manuscript are used here: radialization, the truncation degree, the coefficient selection and the augmentation. Its cohort,

classification results and data are not; a quantitative comparison on any cohort is left to it and to future joint work.

Three problems remain open. The first is a global polynomial separating set for W_L , $L \geq 2$, of size comparable to $\dim \mathcal{Q}_L$; covariant generation and null-cone analysis are the natural tools, and the Maxwell-axis reduction of Theorem 3.6(ii) suggests a route through the first fundamental theorem for configurations of unoriented lines. The second is the sharp Hölder exponent in Theorem 7.3(iii) and its dependence on the strata; the bi-Lipschitz theory of [8] may apply after adaptation. The third is the geometry of the real band-three locus in $\mathbb{P}_{(2,4,6,10)}(\mathbb{R})$. Its intersection with $\mathcal{L}_2$ is the achiral locus of Theorem 5.2. Which real harmonics of degree three have a sextic whose curve has an elliptic subcover of higher degree, and what configuration of their Maxwell axes this imposes, is open; unlike the degree-two case, no reflection is involved, and no anatomical meaning should be assigned to such loci before empirical validation.

The steps of the construction are explicit maps: the conformal barycenter, the correspondence Ψ , the invariants of Section 4 and the gauge (17). By Theorem 7.6, the class $[\mathcal{D}_L(S)]$ they compute does not depend on the parameterization, the position or the scale, so they can serve as fixed input layers for learning on shapes. Such learning and a mesh-level validation on anatomical data are natural next steps; the present paper validates the pipeline on synthetic surfaces and on a single atlas subject.

ACKNOWLEDGEMENTS

Not applicable.

DECLARATIONS

Funding. The authors did not receive support from any organization for the submitted work.

Competing interests. The authors have no competing interests to declare that are relevant to the content of this article.

Ethics approval. Not applicable.

Data availability. The synthetic surface of Section 8.6 is defined in the text. The anatomical meshes of Section 9.4 are the files FMA72713, 72714, 72832, 72833, 72828, 72829, 72830, 72831, 258714, 258716, 72705, 72706, 7211, 7212, 7204, 7205, 15629, 15630, 72826, 72827, 78449 and 78450 of BodyParts3D version 3.0, copyright The Database Center for Life Science, licensed under CC Attribution-Share Alike 2.1 Japan, and are publicly available. The unpublished SPHARM-COM manuscript is available to the editor and referees on request.

Code availability. The verification script of Section 8.5 (Online Resource 1) the script of the synthetic test of Section 8.6 (Online Resource 2) the scripts of the benchmarks of Sections 9.1 to 9.3 (Online Resource 3) and the mesh pipeline of Section 9.4 (Online Resource 4) are supplied as electronic supplementary material.

Author contributions. Both authors contributed to the study conception and design and approved the final manuscript.

REFERENCES

1. L. V. Ahlfors, *Möbius transformations in several dimensions*, Ordway Professorship Lectures in Mathematics, University of Minnesota, School of Mathematics, Minneapolis, MN, 1981.
2. S. Axler, P. Bourdon, and W. Ramey, *Harmonic function theory*, second ed., Graduate Texts in Mathematics, vol. 137, Springer, New York, 2001.
3. A. Baden, K. Crane, and M. Kazhdan, *Möbius registration*, Comput. Graph. Forum **37** (2018), no. 5, 211–220.
4. B. Beauzamy, E. Bombieri, P. Enflo, and H. L. Montgomery, *Products of polynomials in many variables*, J. Number Theory **36** (1990), no. 2, 219–245.
5. L. Beshaj, J. Gutierrez, and T. Shaska, *Weighted greatest common divisors and weighted heights*, J. Number Theory **213** (2020), 319–346.
6. J. Bochnak, M. Coste, and M.-F. Roy, *Real algebraic geometry*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3), vol. 36, Springer, Berlin, 1998.
7. C. Brechbühler, G. Gerig, and O. Kübler, *Parametrization of closed surfaces for 3-D shape description*, Comput. Vis. Image Underst. **61** (1995), no. 2, 154–170.
8. J. Cahill, J. W. Iverson, D. G. Mixon, and D. Packer, *Group-invariant max filtering*, Found. Comput. Math. **25** (2025), no. 3, 1047–1084.
9. J. Cantarella and H. Schumacher, *Computing the conformal barycenter*, SIAM J. Appl. Algebra Geom. **6** (2022), no. 3, 503–530.
10. É. Cartan, *Leçons sur la théorie des spineurs*, Hermann, Paris, 1938.
11. P. T. Choi, K. C. Lam, and L. M. Lui, *FLASH: fast landmark aligned spherical harmonic parameterization for genus-0 closed brain surfaces*, SIAM J. Imaging Sci. **8** (2015), no. 1, 67–94.
12. A. Clebsch, *Theorie der binären algebraischen Formen*, B. G. Teubner, Leipzig, 1872.
13. H. P. de Saint-Gervais, *Uniformization of Riemann surfaces: revisiting a hundred-year-old theorem*, Heritage of European Mathematics, European Mathematical Society, Zürich, 2016.
14. H. Derksen and G. Kemper, *Computational invariant theory*, second ed., Encyclopaedia of Mathematical Sciences, vol. 130, Springer, Heidelberg, 2015.
15. A. Douady and C. J. Earle, *Conformally natural extension of homeomorphisms of the circle*, Acta Math. **157** (1986), no. 1-2, 23–48.
16. N. Dym and S. J. Gortler, *Low-dimensional invariant embeddings for universal geometric learning*, Found. Comput. Math. **25** (2025), no. 2, 375–415.
17. X. Gu, Y. Wang, T. F. Chan, P. M. Thompson, and S.-T. Yau, *Genus zero surface conformal mapping and its application to brain surface mapping*, IEEE Trans. Med. Imaging **23** (2004), no. 8, 949–958.
18. D. Hilbert, *Ueber die Theorie der algebraischen Formen*, Math. Ann. **36** (1890), no. 4, 473–534.
19. J.-I. Igusa, *Arithmetic variety of moduli for genus two*, Ann. of Math. (2) **72** (1960), 612–649.
20. R. Kakarala, *The bispectrum as a source of phase-sensitive invariants for Fourier descriptors: a group-theoretic approach*, J. Math. Imaging Vision **44** (2012), no. 3, 341–353.
21. M. Kazhdan, T. Funkhouser, and S. Rusinkiewicz, *Rotation invariant spherical harmonic representation of 3D shape descriptors*, Proceedings of the Eurographics/ACM SIGGRAPH Symposium on Geometry Processing (SGP 2003) (Aire-la-Ville, Switzerland), Eurographics Association, 2003, pp. 156–164.
22. V. Krishnamoorthy, T. Shaska, and H. Völklein, *Invariants of binary forms*, Progress in Galois theory, Developments in Mathematics, vol. 12, Springer, New York, 2005, pp. 101–122.
23. J. C. Maxwell, *A treatise on electricity and magnetism. Vol. I*, Clarendon Press, Oxford, 1873.
24. M. Olive and N. Auffray, *Isotropic invariants of a completely symmetric third-order tensor*, J. Math. Phys. **55** (2014), no. 9, 092901.
25. P. J. Olver, *Classical invariant theory*, London Mathematical Society Student Texts, vol. 44, Cambridge University Press, Cambridge, 1999.
26. C. Procesi and G. Schwarz, *Inequalities defining orbit spaces*, Invent. Math. **81** (1985), no. 3, 539–554.
27. G. W. Schwarz, *Smooth functions invariant under the action of a compact Lie group*, Topology **14** (1975), 63–68.

28. T. Shaska and H. Völklein, *Elliptic subfields and automorphisms of genus 2 function fields*, Algebra, arithmetic and geometry with applications (West Lafayette, IN, 2000), Springer, Berlin, 2004, pp. 703–723.
29. M. Styner, I. Oguz, S. Xu, C. Brechbühler, D. Pantazis, J. J. Levitt, M. E. Shenton, and G. Gerig, *Framework for the statistical shape analysis of brain structures using SPHARM-PDM*, Insight J. (2006), 242–250.
30. J. J. Sylvester, *Note on spherical harmonics*, Philosophical Magazine (5) **2** (1876), 291–307, Correction, p. 400.
31. N. Ja. Vilenkin, *Special functions and the theory of group representations*, Translations of Mathematical Monographs, vol. 22, American Mathematical Society, Providence, RI, 1968.
32. H. Weyl, *The classical groups: their invariants and representations*, 2 ed., Princeton University Press, Princeton, NJ, 1946.

Email address: `shaska@oakland.edu`

DEPARTMENT OF COMPUTER SCIENCE AND ENGINEERING, OAKLAND UNIVERSITY, ROCHESTER, MI, 48309