

MOVEBENCH: A Benchmark for Global-Scale Wildlife Movement Forecasting

Justin Kay^{1,*} Shir Bar¹ Ellen O. Aikens^{2,α} Martin Becker^{3,α}
 Francesca Cagnacci^{4,α} Juliet Cohen^{5,α} Scott W. Forrest^{6,α} Jessica Kendall-Bar^{7,α}
 Madeleine Lucas^{8,α} Macon Overcast^{18,α} Meredith S. Palmer^{9,α} Will Rogers^{9,α}
 Nicholas J. Russo^{10,α} Christian Rutz^{11,α} Larissa T. Beumer^{12,β} Michael Brown^{20,β}
 Ying-Chi Chan^{13,β} Sarah C. Davidson^{14,β} Diego Ellis Soto^{15,β} Anne G. Hertel^{16,β}
 Roland Kays^{17,β} Benjamin Koger^{2,β} Guram Mikaberidze^{2,β} Thomas Mueller^{16,β}
 Ruth Oliver^{5,β} Thorsten Papenbrock^{3,β} Robert Patchett^{11,β} Jared A. Stabach^{18,β}
 Dane Taylor^{2,β} Scott W. Yanco^{19,β} Sara Beery¹

¹MIT ²University of Wyoming ³Marburg University, Hessian.AI ⁴Fondazione Edmund Mach, Research and Innovation Centre, Animal Ecology Unit ⁵UCSB ⁶Queensland University of Technology ⁷Scripps Institution of Oceanography, UCSD ⁸Colorado State University ⁹Yale ¹⁰Harvard ¹¹University of St. Andrews ¹²UNIS ¹³NTU Singapore ¹⁴Max Planck Institute of Animal Behavior ¹⁵UC Berkeley ¹⁶Senckenberg Biodiversity and Climate Research Centre ¹⁷NCSU and NC Museum of Natural Sciences ¹⁸Smithsonian NCBI ¹⁹Smithsonian Institution ²⁰Giraffe Conservation Foundation

Abstract

Understanding and predicting wildlife movement is critical for ecology and conservation. While trajectory forecasting has advanced for human and vehicle movement, wildlife trajectories present distinct challenges: they are unconstrained in space, highly stochastic, and influenced by environmental conditions. We introduce MOVEBENCH, the first large-scale benchmark for probabilistic wildlife movement forecasting, containing 2.6M GPS locations from 800+ individuals across 110 species in 127 countries, paired with 1.6B environmental raster tiles capturing 160 covariates known or hypothesized to influence movement. We propose a probabilistic evaluation protocol for movement trajectory forecasts, addressing limitations of point-prediction metrics for inherently stochastic phenomena. Through comprehensive empirical evaluation of four method families across multiple temporal and spatial scales, we reveal that: (1) existing predictive methods generalize better to future timepoints than to unseen individuals, (2) deep learning approaches do not consistently outperform simpler baselines, and (3) environmental covariate selection significantly impacts performance. MOVEBENCH enables standardized evaluation of movement forecasting methods and provides a foundation for methodological advances on this ecologically important task.

1 Introduction

Movement is a fundamental process of life on Earth and is at the basis of key ecological functions, from pollination and nutrient cycling to disease transmission and carbon sequestration [168, 235, 303]. Advancements in animal tracking technology (*e.g.*, GPS-based radio-telemetry [48]) combined with real-time data transmission have allowed scientists to track and record wildlife movement at unprecedented scales and resolutions and for an increasingly diverse array of species [173]. This has

*Correspondence to kayj@mit.edu

^{α,β} Equal contribution within groups, alphabetically ordered




Figure 1: MOVEBENCH is the first global benchmark for animal movement forecasting. (A) The benchmark contains 2.6 million curated GPS locations from over 800 individuals of 110 wildlife species, with observations from 127 countries spanning all continents, and (B) 1.6 billion raster tiles representing 160 associated environmental variables known or hypothesized to influence movement. (C) Five example species are shown that illustrate the diversity of wildlife movement: from northern elephant seals and turkey vultures that migrate thousands of kilometers over land and sea, to plains zebras that migrate over land, to mew gulls and water buffalo that spend most of their lives centered around one location. (D) The predictive challenge is *probabilistic forecasting*: models map from an individual's movement history, environmental conditions, and other covariates to a probability distribution over possible future locations of that individual at a specified time horizon. (E) The data in MOVEBENCH covers 110 species across the tree of life with (F) movement patterns spanning over five orders of magnitude in spatial scale. Bars in (F) extend vertically from the median 1-hour movement distance to the median 7-day movement distance of each species. Trajectories in (A), silhouettes in (C), and bars in (F) correspond by color to species in (E).

catalyzed an explosion in animal movement data collection over the past decade, as evidenced by the over ten billion logged animal locations on the animal tracking data repository Movebank [226]. This breadth of observational data has deepened our understanding of how and why animals move through the world [7, 171, 225], helped prioritize areas for protection [33, 148, 224], and informed strategies for mitigating human-wildlife conflict [32, 33, 128, 248]. Beyond observational insights, the ability to predict how future animal movement might respond to various changes in natural and built environments—for instance, how new roads or energy infrastructure may alter movement corridors, how changing land use may reshape landscape connectivity [67, 78, 150], and how migratory routes may shift under climate change [151]—could revolutionize conservation decision making [77, 88, 117, 126, 307, 336].

Currently, we do not have the ability to robustly forecast animal movement. While many methods have been proposed for analyzing animal movement data [13, 15, 54, 85, 87, 104, 107, 152, 180, 181, 227, 228, 239, 264, 283, 328], their predominant use in the literature is for description, inference, and retrospective analysis rather than prediction [134]. There is great potential for the field of movement ecology to move towards a predictive science, and growing interest in the development of predictive methods [61, 104, 105, 134, 264, 271, 306]. To realize this potential, the field needs clear and shared definitions of success. This requires benchmarking frameworks that evaluate models across the dimensions that matter for real-world use: different species, movement modes (*e.g.*, walking, swimming, flying), spatial and temporal scales, and environmental contexts. No such benchmark exists for animal movement forecasting.

Crucially, it is not possible to simply transfer lessons from other trajectory prediction benchmarks onto animal movement data. The most common trajectory data sources in the machine learning literature—human and vehicle trajectories—are fundamentally different, and in many ways simpler, than wildlife trajectories. For instance, they are almost always map-aligned, *i.e.*, explicitly map-matched to known linear transportation infrastructures supporting human mobility, allowing movement to be modeled as a simplified process evolving over a constrained set of paths [238, 342]. There are no such maps for animal movement—in fact, discovering what routes are taken through a landscape is often an explicit goal of collecting the data in the first place [235, 241, 321]. Further, existing trajectory forecasting benchmarks primarily evaluate only a single best-guess future trajectory with metrics such as average displacement error, rather than probabilistic forecasts representing the range of possible future trajectories. Animal movement is highly stochastic [247], with several plausible futures at any given point in time, so accurate prediction requires both probabilistic forecasts and metrics that can assess the distribution of plausible futures. Finally, animal movement is heavily influenced by local environmental conditions [235, 241, 247], and a key goal of forecasting is to predict movement under changing environments. Incorporating this environmental context into predictive methods requires synthesizing heterogeneous, multi-scale, multi-resolution, and time-varying data sources. This relationship to covariates is a complexity missing from existing human and vehicle trajectory methods, and addressing these challenges will require significant methodological innovation.

We introduce MOVEBENCH, the first large-scale benchmark dataset for animal movement forecasting. The dataset contains 2.6 million curated GPS locations from over 800 individuals of 110 wildlife species, with observations from 127 countries spanning all continents. Alongside the GPS trajectory data, MOVEBENCH incorporates 160 publicly-available and global environmental covariates, which are preprocessed into 1.6 billion environmental raster tiles comprised of over two trillion environmental measurements. We propose a set of metrics for evaluating probabilistic movement forecasting methods and conduct the first large-scale empirical comparison of predictive movement methods for this task. Our experiments demonstrate there is much room for improvement in probabilistic forecasting methods, particularly in out-of-domain generalization, covariate selection, and multiscale spatiotemporal reasoning. Altogether, MOVEBENCH represents both the largest dataset for animal movement forecasting in existence and the most extensive empirical comparison of animal movement prediction methods ever conducted, while providing a novel environmentally-conditioned probabilistic spatiotemporal forecasting challenge to the ML community. The benchmark and codebase will be made publicly available, enabling future research and methods development.²

² Paper currently under review. The benchmark will be officially released following acceptance, at which time this preprint will be updated with links to code and data.

Table 1: Trajectory benchmarks. Prior large-scale trajectory benchmarks focus on predicting vehicle movement on roads. MOVEBENCH expands this to enable probabilistic evaluation on a continuous global spatial domain $\mathcal{X}$ rather than point forecasts on a constrained road map. MOVEBENCH also includes more classes with diverse movement patterns, more covariates that influence movement, and larger spatial and temporal ranges than prior work. *IQR computed for per-trajectory cumulative distances and time ranges.*

Benchmark	# GPS	# Cls	# Cov	Dist IQR (km)	Time IQR (h)	Task	$\mathcal{X}$	Probabilistic?
GeoLife [358]	23.6M	10	0	3 – 22	0.3 – 3	Forecast, Classify	Roads	
Porto [242]	83M	1	4	2 – 7	0.1 – 0.2	Forecast	Roads	
T-Drive [357]	790M	1	0	860 – 2000	145 – 148	Forecast	Roads	
WorldTrace [359]	8.8B	1	0	0.8 – 10	0.03 – 0.2	Forecast	Roads	
MOVEBENCH	2.6M	110	160	283 – 3500	818 – 8000	Forecast	Globe	✓

2 Related work

Compared to existing benchmarks for trajectory prediction (see Tab. 1), MOVEBENCH offers innovations along three axes (for additional related work, see Appendix Sec. A):

Data: The wildlife movement data in MOVEBENCH is more complex, heterogeneous, and covariate-dependent than existing benchmarks focusing on human movement [55, 201, 203, 242, 356, 358, 359]. It is unconstrained in space, varies drastically in pattern and scale across species and geographies, and requires synthesizing large-scale geospatial datasets in order to provide methods with the environmental context needed to predict where animals will move [241, 321]. Existing large datasets with tracked animals do not support probabilistic movement forecasting—BEBE [149] is a benchmark for behavior classification in accelerometry data, and Noonan et al. [240] evaluates methods for space use estimation over a large collection of datasets, anonymizing both GPS locations and timestamps.

Task: MOVEBENCH focuses on probabilistic forecasting to reflect the inherently stochastic nature of animal movement (see Sec. 3). Prior work primarily evaluates forecasting based on either: (i) point-prediction metrics (e.g., L2 error between ground truth and predicted locations), which do not assess the quality of entire predictive distributions [251, 301, 302], (ii) qualitative inspection [306], or (iii) log loss [61, 105], which is probabilistic but cannot be used to compare across arbitrary model classes because it requires an explicit predictive density, which is unavailable for deterministic predictors and may be intractable for some methods. As discussed in Sec. 5, MOVEBENCH introduces an evaluation methodology based on a *proper scoring rule* [124, 337] called the *energy score* that enables robust evaluation of probabilistic forecasts from any trajectory prediction model [299].

Methods: We compare commonly-used and state-of-the-art methods for predicting animal movement, which model the movement process as a function of *habitat selection*, whereby an individual chooses where to go next based on environmental conditions at candidate future locations [13, 61, 105, 107] coupled with their estimated movement capacity based on past movement. MOVEBENCH enables empirical comparison at scale between these methods for the first time.

3 Movement forecasting

The high-level goal is to predict the next step an animal will take at any time point, given its movement history and current environment. The following precisely defines this movement forecasting task (see Fig. 1d), describing observed trajectories, forecast targets, and evaluation settings.

Trajectory observations. We consider a spatiotemporal domain where $\mathcal{X} \subseteq \mathbb{R}^2$ and $\mathcal{T} \subseteq \mathbb{R}$ denote the spatial and temporal extents available to an animal, respectively. For a single tracked animal, let $\tau = ((\ell_n, t_n))_{n=1}^N$ denote an observed trajectory of length N , where $\ell_n \in \mathcal{X}$ is the recorded location at observation time $t_n \in \mathcal{T}$ for $t_1 < t_2 < \dots < t_N$. For $n = 1, \dots, N - 1$, define the elapsed time between consecutive observations by $\Delta t_n = t_{n+1} - t_n$. We do not assume that Δt_n is constant in n , although in many trajectories it is approximately regular.

Covariates. Each spatiotemporal point is associated with a set of covariates, *i.e.* a feature vector. These may include movement-, time-, and environment-derived features. Formally, we represent these features by a vector-valued spatiotemporal field $c : \mathcal{X} \times \mathcal{T} \rightarrow \mathbb{R}^K$, where K is the number of features. For any location–time pair $(\ell, t) \in \mathcal{X} \times \mathcal{T}$, the vector $c(\ell, t)$ represents corresponding features. c can be evaluated for any observed point $c_n = c(\ell_n, t_n)$ with $n = 1, \dots, N$, or at any

Table 2: Covariates in MOVEBENCH. The benchmark includes 160 environmental layers representing context at observed trajectories and at candidate locations for future movement. Covariates are included as rasters, centered at observed GPS points and aligned across spatial and temporal scales. Resolution indicates the spatial extent represented by one pixel. Cadence indicates the temporal sampling rate. Movement- and time-derived covariates are also included for all observed and candidate locations.

Category	# Layers	Resolution	Cadence	Examples
Weather & climate	51	30m – 28km	1hr – 8d	Temperature, precipitation, wind, snow depth, land surface temperature, BioClim [95, 219, 353]
Vegetation & land cover	34	10m – 500m	3d – 1yr	Vegetation cover, vegetation indices, land cover, distance to water [40, 250, 279, 353]
DEM	4	30m	static	Elevation, slope, aspect, rugosity [91]
Marine	6	250m – 28km	static – 30d	Bathymetry, distance to coast, sea surface temp, chlorophyll, surface current [53, 63, 156]
Human impact	1	300m	annual	WCS Human Footprint Index [281]
Embeddings	64	10m	annual	AlphaEarth Foundations [41]
Movement	3	–	–	Step length, turn angle [13]
Time	4	–	–	Time of day, day of year [104]

candidate future location $\tilde{\ell} \in \mathcal{X}$ and candidate future time $\tilde{t} \in \mathcal{T}$. See Tab. 2 for a summary of the covariates available in MOVEBENCH.

Forecasting task. We evaluate probabilistic movement forecasting at forecast horizons $\Delta > 0$. For a forecast origin (ℓ, t) , the prediction target at time horizon Δ is the animal’s location at time $t + \Delta$, denoted $\ell_{t+\Delta}$. For an observed trajectory, define the history up to time t by $\tau_{\leq t} = ((\ell_i, t_i))_{i=1}^{n(t)}$ and any associated covariates by $c_{\leq t} = (c(\ell_i, t_i))_{i=1}^{n(t)}$, with $n(t) = \max\{i : t_i \leq t\}$ representing the number of observed steps up to and including time t .

Then, for a given forecast horizon Δ and origin time t , a movement forecasting method estimates a conditional predictive spatial distribution $P_{t,\Delta}(\cdot \mid \tau_{\leq t}, c_{\leq t})$ over $\mathcal{X}$ representing the probability the animal will move to any spatial location at time $t + \Delta$. In practice, for tractability, this is typically defined over a reasonable spatial range surrounding the animal’s current location (*e.g.*, its 99th percentile next-step movement distance) rather than all of $\mathcal{X}$. Deterministic predictors are a special case where $P_{t,\Delta}$ is a point mass at a single predicted location.

4 MOVEBENCH

MOVEBENCH (see Fig. 1) contains 2.6 million GPS locations from over 800 individuals of 110 wildlife species, accompanied by over 1.6 billion environmental raster tiles. Here, we briefly describe how we select this data and construct the benchmark. More details can be found in Appendix Sec. B.

Trajectory curation. We aggregate trajectory data from: (1) the Movebank Data Repository [2], a subset of data from Movebank [226] that have Creative Commons licensing, DOIs, and have already undergone some quality assurance and anomaly filtering, (2) a literature review of recent movement methods papers with empirical experiments on public data, and (3) a survey of domain experts in the Movement Biodiversity Observation Network [1]. This yields 147 raw trajectory datasets, from which we apply a coarse set of filtering criteria followed by manual curation (see Appendix Sec. B for details and source acknowledgments).

Covariate rasters. MOVEBENCH contains 160 publicly-available environmental covariates that are known or hypothesized to influence movement (see Tab. 2). These include features from public environmental and climatic datasets [95, 219, 279, 353] as well as learned geospatial embeddings [41]. Covariates are downloaded and processed into 101×101 pixel rasters, centered at observed GPS points and scaled to include appropriate spatial context for a given species and forecast horizon, as in previous raster-based work [105] (see Appendix Sec. C). Because we evaluate each method at four different spatiotemporal scales (see Sec. 6), we include the full 160-layer raster stack at four different spatial resolutions for each of the 2.6M points in the dataset. In total, the benchmark therefore contains 10.4M raster stacks containing 1.6B individual raster layers representing over

two trillion environmental measurements. This is at least four orders of magnitude larger than existing raster-based datasets for animal movement [105], enabling evaluation of predictive methods at a drastically larger scale than was previously possible. We also support raster-based versions of common movement- and time-derived covariates such as movement distance and time of movement. See Tab. 2 for a summary of all covariates and Appendix Sec. B for the full set and acknowledgments.

The use of rasters supports arbitrary definitions of candidate future movement locations (see Appendix Sec. E) as well as emerging raster-based methods [105], but introduces significant engineering complexity. The key challenges involve querying, batching, aligning, and efficiently storing data of different spatial and temporal resolutions from large-scale satellite and climate reanalysis datasets, as well as efficiently loading data during training and evaluation. Alongside the processed raster dataset, upon publication we will release the MOVEBENCH raster download and preparation pipeline as a Python package to support future research.

Train and test splits. For each study, we construct one training set and two hold-out test sets representing a *temporal* split and an *individual* split, each capturing a distinct source of ecological variation. The *temporal* split evaluates generalization to future time by testing on later observations from the same individuals in the training set, reflecting the challenge of predicting movement under changing environmental conditions and possibly different animal life stages. The *individual* split evaluates generalization across animals by testing on individuals not seen during training, capturing behavioral and environmental heterogeneity among individuals of the same species but within the same population. We select the held-out time points and individuals in order to mitigate train-test leakage; see Appendix Sec. B.3. In total, MOVEBENCH contains 1,120,127 training points, 965,233 temporal test points, and 537,781 individual test points.

MOVEBENCH-Mini. To enable quicker experimentation, we also make available a “mini” set of ten studies across taxonomic groups and movement types. MOVEBENCH-Mini includes two migratory birds, two non-migratory birds, two migratory mammals, two non-migratory mammals, one marine species, and one terrestrial reptile. We report study specifics in Appendix Sec. B.5.

5 Metrics

Animal movement is inherently *probabilistic*: future motion is often uncertain and multimodal, and thus forecasting methods are designed to return a distribution over possible next steps rather than a single best guess. To evaluate methods, we would ideally compare predicted distributions to the true distribution of the animal’s future movement, of which the observed movement is one realization. This is not possible with commonly used metrics for trajectory forecasting, such as average displacement error [251, 301, 302], which evaluate each ground truth point against a single point prediction. Further, it is well-known that optimizing for point prediction metrics rewards models that predict a central moment of the true distribution rather than its full shape (*e.g.*, L2 error is minimized by deterministically predicting the geometric median [76]). Thus, when several distinct future destinations are plausible, point prediction metrics can reward an averaged prediction that lies between modes, even if that location itself is implausible [124, 326]. See Fig. 2a.

Proper scoring rules offer a robust framework for assessing the quality of probabilistic forecasts [125, 337]. A *scoring rule* is a loss or utility function that can be used to compare predicted distributions with realized observations from a ground truth distribution. A scoring rule is *proper* if it is maximized by predicting the true target distribution rather than hedging toward the median or some otherwise strategically distorted forecast, addressing the problems of point prediction metrics described above (see Fig. 2b and c). Negative log likelihood (NLL) is one example of a proper scoring




Figure 2: Probabilistic forecasting metrics.

(a) Consider an animal with two modes in its next-step movement distribution, *i.e.* it may travel to one of two regions with equal probability. A probabilistic forecast should try to reflect this, but L2-based metrics reward methods that instead deterministically predict the mean or median of the true movement distribution. (b) We see this empirically on MOVEBENCH: according to expected L2 error, the best forecast is to always predict “don’t move”. This is not very useful in practice. (c) Proper scoring rules like the energy score reward accurate and calibrated probabilistic predictions. In (b) and (c), a “win” means that a method has the best score for the given metric on the temporal split for a single study at a particular time horizon.

rule that is already used widely for comparing wildlife movement forecasts [61, 105]. However a key limitation of NLL is that it requires access to predicted probability densities, preventing comparisons with deterministic baselines or more complex models where this is expensive or intractable to obtain.

The energy score [125] is a proper scoring rule that is particularly well suited to evaluating trajectory forecasts for two reasons: (1) It is multivariate, enabling evaluation on $\mathcal{X} \subseteq \mathbb{R}^2$, and (2) it requires only samples from the predictive distribution, not access to the corresponding predictive density. It is a popular choice for evaluating spatial probabilistic forecasts in other applications, such as weather forecasting [172, 194, 208, 286], and has recently been introduced for evaluating probabilistic vehicle trajectory predictions [299]. We adopt it as the main evaluation criterion in MOVEBENCH.

For a given trajectory step (ℓ, t) and forecast distribution $P_{t,\Delta}$ (see Sec. 3), the energy score evaluated against $\ell_{t+\Delta}$, the ground truth location at time $t + \Delta$, is

$$\text{ES}(P_{t,\Delta}, \ell_{t+\Delta}) = \mathbb{E}_{X \sim P_{t,\Delta}} \|X - \ell_{t+\Delta}\|_2 - \frac{1}{2} \mathbb{E}_{X, X' \sim P_{t,\Delta}} \|X - X'\|_2, \quad (1)$$

where X and X' are i.i.d. draws from the predictive distribution. Lower values are better. For a deterministic predictor, $P_{t,\Delta}$ is a point mass, so the ES reduces to the L2 error.

Like many proper scoring rules, the energy score is a combination of two terms. The first can be interpreted as measuring the *accuracy* of a model, and is the average distance between the ground truth location and draws from the predictive distribution (*i.e.*, the expected L2 distance). The second term can be interpreted as measuring the *spread* of a model, computed as half of the average distance between two separate draws from the predictive distribution, and encourages accurate representation of model uncertainty. A model that is performing well will be both accurate and appropriately calibrated, resulting in a low energy score. See Appendix Sec. D for more details on its derivation.

6 Experiments

Baselines. We select a set of representative methods to benchmark on MOVEBENCH, including both methods that are widely used in practice as well as recent methods incorporating deep learning. The methods are: (1) **iSSF Linear**, an integrated step selection function with a standard linear head [13], (2) **iSSF MLP**, an iSSF with a multilayer perceptron head, (3) **deepSSF** [105], a CNN-based method that operates on spatial rasters, and (4) **MoveFormer** [61], a transformer-based method that incorporates longer temporal context and is trained across studies/species. As points of reference, we compare methods with two simple baselines: (1) **don’t move**, which always predicts that the animal does not move (*i.e.*, $\hat{\ell}_{t,\Delta} = \ell_t$); and (2) **correlated random walk (CRW)**, which samples from the same parametric distributions used to select candidates for SSF-style methods. We defer detailed descriptions to Appendix Sec. E.

Main results. We train and evaluate each baseline method at a fixed set of forecast horizons $\Delta \in \{1 \text{ hour}, 12 \text{ hours}, 1 \text{ day}, 7 \text{ days}\}$, compute the energy score (Eq. (1)) for each test step at each forecast horizon, and compute a mean energy score for each time horizon for each study.

We aggregate results across studies by either method (Fig. 3 top) or species/range groups (Fig. 3 bottom). All reported results are macro-averages of per-study energy scores. However, because energy score scales with distance, animals that move longer distances on average would dominate a simple average energy score. Before aggregating across studies we therefore first normalize each study by the average movement distance of the individuals in that study. We display the mean $\pm$ the standard deviation of these normalized energy scores over each grouping. Note that normalizing by average movement distance is equivalent to normalizing by the energy score of the “don’t move” baseline (see Eq. (1)), so “don’t move” always has a normalized ES of 1. See Appendix Sec. F for more details on evaluation methodology and complete hyperparameter settings.

Covariate comparison. We first investigate performance of all methods on MOVEBENCH-Mini using different sets of environmental covariates in Fig. 3 (top). We compare the full set of covariates available in the benchmark (“alphaearth + env all”) with smaller sets to understand which covariates contribute positively or negatively toward forecasting performance for different models. We see that this is not consistent between models, and that more covariates is not always better, even for deep learning methods. For most methods, as well as on average, AlphaEarth embeddings are the best-performing, while the MLP-based iSSF performs best with a combination of AlphaEarth and environmental covariates. While it is not surprising that adding additional highly-correlated covariates




Figure 3: Main results. We report energy scores (lower is better, see Sec. 5), normalized by average movement distance at the study level to account for inter-study variation and aggregated by method (top) or taxonomic group and travel distance (bottom) (see Sec. 6). **Top (Covariate comparison):** For each model, we compare four different sets of input covariates on MOVEBENCH-Mini: “env lite,” a small set of common covariates (NDVI and elevation for terrestrial studies, bathymetry and chlorophyll-a for marine studies); “env all,” the full set of 96 environmental covariates excluding AlphaEarth; AlphaEarth only; and AlphaEarth + “env all”. We see that, on average across all methods, AlphaEarth covariates outperform environmental covariates, and that combining AlphaEarth with environmental covariates does not seem to add benefit for most methods. These trends are not consistent across methods, however, and in particular iSSF MLP performs best when trained with environmental covariates. **Bottom (Method comparison):** We compare all methods across the entirety of MOVEBENCH using AlphaEarth covariates (the best-performing set on average), aggregating studies by taxonomic group and median per-individual maximum movement distance. n indicates the number of individuals in each panel. Most methods outperform simple baselines on the temporal split, but not on the individual split, indicating that it is more difficult to generalize to new individuals who were not present in the training set. deepSSF in particular shows a large generalization gap—it is the best-performing overall on the temporal split but worse than the CRW baseline on the individual split. More work is needed to develop methods that perform well across different generalization challenges.

to the linear model worsens performance, The fact that combining both sources of covariates does not improve performance for other methods—in particular, the deep learning based methods deepSSF and MoveFormer—indicates that more work is needed to enable models to handle large complementary input sets, for instance with additional regularization. Rankings of covariates remain fairly stable across the temporal and individual test splits. Overall, this comparison and the lack of consistency across models point to the importance of covariate selection as a component of methods development.

Method comparison. We then train and evaluate all methods on all 147 datasets in MOVEBENCH using AlphaEarth covariates since they were the highest performing on average in the covariate comparison. In Fig. 3 (bottom) we aggregate energy scores over studies based on broad taxonomic group and movement characteristics. Distance-based groupings are based on the median per-individual maximum displacement of each study.

We see that no method is consistently best across all taxonomic-range groups or timescales. On in-sample individuals (*i.e.*, the temporal split), most methods usually—but not always—outperform




Figure 4: Prediction probability surfaces. Here we show qualitative examples of forecast distributions of all four methods for all four timescales in our main experiments for two sample points of two species (white stork and jaguar). The black dot represents the animals’ starting position and the black star represents their true end position at each time horizon. The map shows Landsat RGB overlaid on Copernicus DEM for the spatial extent available at prediction time—notice that this varies with both study and timescale, and that data from some environmental layers is sometimes missing. Prediction heatmaps represent the log predicted probability that the animal moves to any spatial location at the next timestep. Energy scores are displayed for each prediction, computed on the singular sample (lower is better). Notice that some methods’ predictions are highly influenced by environmental features (*e.g.*, iSSF, deepSSF), while others focus on broad spatial trends (*e.g.*, MoveFormer).

the simple correlated random walk (CRW) baseline, indicating they are leveraging environmental information to improve upon simple movement statistics. deepSSF—the only method that makes use of full-raster spatial context during training—consistently performs the best on future points of training individuals, though MoveFormer is a very close second-best. All deep learning methods are inconsistent in whether or not they outperform a simple linear iSSF, indicating that deep learning on its own is not enough to guarantee better performance.

We see that trained models generalize better to held-out points in time of individuals in the training set (top row of both plots) compared to held out individuals (bottom row of both plots). For held out individuals, most methods do not consistently outperform the correlated random walk baseline. Note that the movement parameters of the correlated random walk are also fit to individuals in the training data, so the CRW baseline faces distribution shift as well with held-out individuals. Therefore, methods that outperform CRW in-sample but underperform CRW out-of-sample appear to be leveraging environmental information in a way that is individual-specific and does not generalize well to new individuals. In particular, deepSSF experiences a large performance drop between the temporal and individual splits, making it the best performing for in-sample individuals but the worst performing for out-of-sample individuals. This indicates that it has learned individual-specific features that are helpful in-distribution and harmful out-of-distribution. MoveFormer—the only method to condition on multiple past timesteps and to train across studies—has the best overall average performance across both hold-out sets.

Qualitative analysis. In Fig. 4 we visualize example outputs from an iSSF model on two start points in the dataset to illustrate common qualities of predictive distributions. For each start point,

we have a ground truth “next step” at each timescale {1 hour, 12 hours, 1 day, 7 days}. The plot shows methods’ predicted log probability of moving to any available location in the spatial extent. Notice that the spatial extent available to predictive methods increases with timescale according to the movement characteristics of the particular species—for white stork, a migratory bird, its 7 day spatial extent is $26\times$ its 1 hour spatial extent, covering a large swath of Northern African coastline. We see that different methods incorporate environmental context as well as movement characteristics differently into their predictions—for instance, MoveFormer shows strong learned directionality in its predictions for both species, and most methods display a preference toward water for the jaguar.

7 Discussion and Conclusion

We introduce MOVEBENCH, the first benchmark for global-scale probabilistic animal movement forecasting. The benchmark enables comparisons across the many axes of variation present in animal movement data, including diverse species, spatiotemporal scales, and movement types, and represents a key step towards advancing state-of-the-art in animal movement forecasting.

Research directions MOVEBENCH highlights as key priorities. First, *cross-individual generalization* is a significant challenge across methods. Opportunities include adapting and developing OOD generalization methods, developing mechanisms to better capture individual variation across populations, and further investigating methods to learn movement patterns not just within but also across populations and species. Additionally, there is potential to further improve prediction by *incorporating additional biologically meaningful information* into movement prediction, such as life history, individual ID, or inferred behavioral state, as suggested by recent ecology literature [271, 272]. We also show that *covariate design and selection* is a key component of performance. Opportunities include methods to automatically identify informative covariates per-species, further develop and improve environmental representations from geospatial foundation models with movement forecasting as a target, and address the tradeoffs between interpretability and predictive capacity when using black-box covariates. Finally, investigating *prediction with long context at any given time horizon* could improve both accuracy and usability of forecasting models, by, *e.g.*, sharing information across spatial and temporal scales via memory (like MoveFormer) or spatial reasoning (like deepSSF).

Opportunities to expand movement ecology benchmarking beyond MOVEBENCH. We see several exciting future opportunities to build on MOVEBENCH, including incorporating larger populations to understand scaling laws in cross-individual generalization, expanding to cover an even broader set of species and regions, expanding to include additional target tasks such as behavior segmentation, corridor mapping, or home range estimation, and expanding to include additional modalities often captured concurrently, such as accelerometry.

Limitations. MOVEBENCH is designed for benchmarking probabilistic movement forecasting methods, and should not be used for drawing general *ecological* conclusions without further analysis. Further, data is sourced from existing published movement studies, inheriting any existing taxonomic and geographical sampling biases, and some nuance is inherently lost when averaging across axes of biological variation in the data. We include further limitations discussion in the Croissant metadata.

Acknowledgments

We thank all data owners (see Appendix Sec. B), without whom none of this work would be possible. We also thank the organizers and members of MoveBON for support and feedback. JK and SaB were supported by NSF award 2330423 and NSERC award 585136. JK was additionally supported by NSF fellowship award 2313998. ShB was supported by a CHE Fellowship for Marine Sciences. JKB was supported by NSF awards 1643532 and NSF 2333609. WR was supported by a Gruber Science Foundation Fellowship. NR was supported by the Harvard University Dean’s Fund for Competitive Scientific Research. RK was supported by National Aeronautics and Space Administration Award 80NSSC21K1182. RYO was supported by the Kuni Endowed Junior Faculty Fellowship, Hellman Family Faculty Fellowship, and the University of California Early Career Faculty Research Excellence Award. DT, GM, BK and EA were supported by the Wyoming Center for Wildlife, Technology and Computing (WyldTech). MO was supported by the Smithsonian Didden Fellowship. CR acknowledges funding from the Gordon and Betty Moore Foundation (GBMF9881; GBMF 11414.01; GBMF12975), and CR and RP from the National Geographic Society (NGS-29784).

References

- [1] Move BON — geobon.org. <https://geobon.org/move-bon/>. [Accessed 18-04-2026].
- [2] Movebank data repository. <https://datarepository.movebank.org/home>. [Accessed 18-04-2026].
- [3] Marta Acácio, Inês Catry, Andrea Soriano-Redondo, João Paulo Silva, Philip W. Atkinson, and Aldina M. A. Franco. Timing is critical: consequences of asynchronous migration for the performance and destination of a long-distance migrant. *Movement Ecology*, 10(1), 2022. ISSN 2051-3933. doi: 10.1186/s40462-022-00328-3. URL <http://dx.doi.org/10.1186/s40462-022-00328-3>.
- [4] Marta Acácio, Inês Catry, Andrea Soriano-Redondo, João Paulo Silva, Philip W. Atkinson, and Aldina M.A. Franco. Data from: Timing is critical: consequences of asynchronous migration for the performance and destination of a long-distance migrant, 2023. URL <https://datarepository.movebank.org/handle/10255/move.258>.
- [5] Ellen Aikens, Jessica Speiser, Karma Choki, Michele Lovara, Anna Weesies, Jeffrey Tillery, Sean Ryder, Erica Lafferty, Amanda Cheeseman, William Severud, and Hall Sawyer. Data from: Challenging conventional views on the elevational limits of pronghorn habitat, 2024. URL <https://datadryad.org/dataset/doi:10.5061/dryad.w9ghx3fz8>.
- [6] Ellen Aikens, Jerod Merkle, Wenjing Xu, and Hall Sawyer. Pronghorn movements and mortality during extreme weather highlight the importance of landscape connectivity, 2025. URL <https://data.mendeley.com/datasets/wgyts8vyrf/1>.
- [7] Ellen O Aikens, Iris D Bontekoe, Lara Blumenstiel, Anna Schlicksupp, and Andrea Flack. Viewing animal migration through a social lens. *Trends in Ecology & Evolution*, 37(11): 985–996, 2022.
- [8] Ellen O. Aikens, Elham Nourani, Wolfgang Fiedler, Martin Wikelski, and Andrea Flack. Learning shapes the development of migratory behavior. *Proceedings of the National Academy of Sciences*, 121(12), March 2024. ISSN 1091-6490. doi: 10.1073/pnas.2306389121. URL <http://dx.doi.org/10.1073/pnas.2306389121>.
- [9] Ellen O. Aikens, Jessica Speiser, Karma Choki, Michele Lovara, Anna Weesies, Jeffrey Tillery, Sean Ryder, Erica Lafferty, Amanda E. Cheeseman, William J. Severud, and Hall Sawyer. Challenging conventional views on the elevational limits of pronghorn habitat. *Ecology*, 105(11), 2024. ISSN 1939-9170. doi: 10.1002/ecy.4422. URL <http://dx.doi.org/10.1002/ecy.4422>.
- [10] Ellen O. Aikens, Jerod A. Merkle, Wenjing Xu, and Hall Sawyer. Pronghorn movements and mortality during extreme weather highlight the critical importance of connectivity. *Current Biology*, 35(8):1927–1934.e2, April 2025. ISSN 0960-9822. doi: 10.1016/j.cub.2025.03.010. URL <http://dx.doi.org/10.1016/j.cub.2025.03.010>.
- [11] Alexandre Alahi, Kratarth Goel, Vignesh Ramanathan, Alexandre Robicquet, Li Fei-Fei, and Silvio Savarese. Social lstm: Human trajectory prediction in crowded spaces. In *Proceedings of the IEEE Conference on Computer Vision and Pattern Recognition (CVPR)*, pages 961–971, 2016.
- [12] Christine M. Anderson, H. Grant Gilchrist, Robert A. Ronconi, Katherine R. Shlepr, Daniel E. Clark, David A. Fifield, Gregory J. Robertson, and Mark L. Mallory. Both short and long distance migrants use energy-minimizing migration strategies in north american herring gulls. *Movement Ecology*, 8(1), 2020. ISSN 2051-3933. doi: 10.1186/s40462-020-00207-9. URL <http://dx.doi.org/10.1186/s40462-020-00207-9>.
- [13] Tal Avgar, Jonathan R Potts, Mark A Lewis, and Mark S Boyce. Integrated step selection analysis: bridging the gap between resource selection and animal movement. *Methods in Ecology and Evolution*, 7(5):619–630, 2016.

- [14] Laurie L Baker, Joanna E Mills Flemming, Ian D Jonsen, Damian C Lidgard, Sara J Iver-son, and W Don Bowen. A novel approach to quantifying the spatiotemporal behavior of instrumented grey seals used to sample the environment. *Movement Ecology*, 3(1), 2015. ISSN 2051-3933. doi: 10.1186/s40462-015-0047-4. URL <http://dx.doi.org/10.1186/s40462-015-0047-4>.
- [15] Nicholas W Bakner, Bret A Collier, and Michael J Chamberlain. Behavioral-dependent recursive movements and implications for resource selection. *Scientific Reports*, 13(1):16632, 2023.
- [16] Hattie L. A. Bartlam-Brooks, Pieter S. A. Beck, Gil Bohrer, and Stephen Harris. In search of greener pastures: Using satellite images to predict the effects of environmental change on zebra migration. *Journal of Geophysical Research: Biogeosciences*, 118(4):1427–1437, October 2013. ISSN 2169-8961. doi: 10.1002/jgrg.20096. URL <http://dx.doi.org/10.1002/jgrg.20096>.
- [17] Hattie L.A. Bartlam-Brooks and Stephen Harris. Data from: In search of greener pastures: using satellite images to predict the effects of environmental change on zebra migration, 2013. URL <https://www.datarepository.movebank.org/handle/10255/move.342>.
- [18] Guillaume Bastille-Rousseau, Jonathan R. Potts, Charles B. Yackulic, Jacqueline L. Frair, E. Hance Ellington, and Stephen Blake. Data from: Flexible characterization of animal movement pattern using net squared displacement and a latent state model, 2016. URL <https://www.datarepository.movebank.org/handle/10255/move.618>.
- [19] Guillaume Bastille-Rousseau, Jonathan R. Potts, Charles B. Yackulic, Jacqueline L. Frair, E. Hance Ellington, and Stephen Blake. Flexible characterization of animal movement pattern using net squared displacement and a latent state model. *Movement Ecology*, 4(1), 2016. ISSN 2051-3933. doi: 10.1186/s40462-016-0080-y. URL <http://dx.doi.org/10.1186/s40462-016-0080-y>.
- [20] Guillaume Bastille-Rousseau, Charles B. Yackulic, Jacqueline L. Frair, Freddy Cabrera, and Stephen Blake. Data from: Allometric and temporal scaling of movement characteristics in galapagos tortoises, 2016. URL <https://www.datarepository.movebank.org/handle/10255/move.549>.
- [21] Guillaume Bastille-Rousseau, Charles B. Yackulic, Jacqueline L. Frair, Freddy Cabrera, and Stephen Blake. Allometric and temporal scaling of movement characteristics in galapagos tortoises. *Journal of Animal Ecology*, 85(5):1171–1181, 2016. ISSN 1365-2656. doi: 10.1111/1365-2656.12561. URL <http://dx.doi.org/10.1111/1365-2656.12561>.
- [22] Guillaume Bastille-Rousseau, Charles B. Yackulic, James Gibbs, Jacqueline L. Frair, Freddy Cabrera, and Stephen Blake. Data from: Migration triggers in a large herbivore: Galápagos giant tortoises navigating resource gradients on volcanoes, 2019. URL <https://www.datarepository.movebank.org/handle/10255/move.834>.
- [23] Guillaume Bastille-Rousseau, Charles B. Yackulic, James P. Gibbs, Jacqueline L. Frair, Freddy Cabrera, and Stephen Blake. Migration triggers in a large herbivore: Galápagos giant tortoises navigating resource gradients on volcanoes. *Ecology*, 100(6), April 2019. ISSN 1939-9170. doi: 10.1002/ecy.2658. URL <http://dx.doi.org/10.1002/ecy.2658>.
- [24] Nyambayar Batbayar, Batbayar Galtbalt, Tseveenmyadag Natsagdorj, Tuvshintugs Sukhbaatar, and Martin Wikelski. Data from: Study "lifetrack mongolia demoiselle cranes", 2024. URL <https://datarepository.movebank.org/handle/10255/move.3263>.
- [25] Nyambayar Batbayar, Batbayar Galtbalt, Tseveenmyadag Natsagdorj, Tuvshintugs Sukhbaatar, and Martin Wikelski. Data from: Study "white-naped crane mongolia wsc", 2024. URL <https://datarepository.movebank.org/handle/10255/move.3264>.
- [26] Jeffrey L. Beck, Jacob D. Hennig, and John D. Scasta. Data from: Study "pronghorn (antilocapra americana), wyoming/colorado, usa", 2022. URL <https://www.datarepository.movebank.org/handle/10255/move.1578>.

- [27] Christopher Beirne, Mark Thomas, Arianna Basto, Eleanor Flatt, Giancarlo Inga Diaz, Diego Rolim Chulla, Flor Perez Mullisaca, Rosio Vega Quispe, Caleb Jonatan Quispe Quispe, Adrian Forsyth, and Andrew Whitworth. Scouts vs. usurpers: alternative foraging strategies facilitate coexistence between neotropical cathartid vultures. *Ibis*, 166(4):1368–1383, April 2024. ISSN 1474-919X. doi: 10.1111/ibi.13327. URL <http://dx.doi.org/10.1111/ibi.13327>.
- [28] Steven E. Bellan and Wayne M. Getz. Data from: Resource-driven encounters among consumers and implications for the spread of infectious disease, 2017. URL <https://www.datarepository.movebank.org/handle/10255/move.706>.
- [29] Lorena Benitez, J. Werner Kilian, George Wittemyer, Lacey F. Hughey, Chris H. Fleming, Peter Leimgruber, Pierre du Preez, and Jared A. Stabach. Precipitation, vegetation productivity, and human impacts control home range size of elephants in dryland systems in northern namibia. *Ecology and Evolution*, 12(9), 2022. ISSN 2045-7758. doi: 10.1002/ece3.9288. URL <http://dx.doi.org/10.1002/ece3.9288>.
- [30] Keith L. Bildstein, David Barber, and Marc J. Bechard. Data from: Environmental drivers of variability in the movement ecology of turkey vultures (*cathartes aura*) in north and south america, 2014. URL <https://www.datarepository.movebank.org/handle/10255/move.362>.
- [31] Keith L. Bildstein, David Barber, Marc J. Bechard, Maricel Graña Grilli, and Jean-François Therrien. Data from: Study "vultures acopian center usa gps" (2003-2021), 2021. URL <https://www.datarepository.movebank.org/handle/10255/move.1306>.
- [32] Barbara A Block, Ian D Jonsen, Salvador J Jorgensen, Arliss J Winship, Scott A Shaffer, Steven J Bograd, Elliott Lee Hazen, David G Foley, GA Breed, A-L Harrison, et al. Tracking apex marine predator movements in a dynamic ocean. *Nature*, 475(7354):86–90, 2011.
- [33] Hannah Blondin, Briana Abrahms, Larry B Crowder, and Elliott L Hazen. Combining high temporal resolution whale distribution and vessel tracking data improves estimates of ship strike risk. *Biological Conservation*, 250:108757, 2020.
- [34] Wayne S J Boardman, David Roshier, Terry Reardon, Kathryn Burbidge, Adam McKeown, David A Westcott, Charles G B Caraguel, and Thomas A A Prowse. Spring foraging movements of an urban population of grey-headed flying foxes (*pteropus poliocephalus*). *Journal of Urban Ecology*, 7(1), 2021. ISSN 2058-5543. doi: 10.1093/jue/juaa034. URL <http://dx.doi.org/10.1093/jue/juaa034>.
- [35] Rebecca K. Borchering, Steve E. Bellan, Jason M. Flynn, Juliet R. C. Pulliam, and Scott A. McKinley. Resource-driven encounters among consumers and implications for the spread of infectious disease. *Journal of The Royal Society Interface*, 14(135):20170555, October 2017. ISSN 1742-5662. doi: 10.1098/rsif.2017.0555. URL <http://dx.doi.org/10.1098/rsif.2017.0555>.
- [36] Rahel M. Borrmann, Richard A. Phillips, Thomas A. Clay, and Stefan Garthe. Data from: High foraging site fidelity and spatial segregation among individual great black-backed gulls, 2019. URL <https://www.datarepository.movebank.org/handle/10255/move.1003>.
- [37] Rahel M. Borrmann, Richard A. Phillips, Thomas A. Clay, and Stefan Garthe. High foraging site fidelity and spatial segregation among individual great black-backed gulls. *Journal of Avian Biology*, 50(12), December 2019. ISSN 1600-048X. doi: 10.1111/jav.02156. URL <http://dx.doi.org/10.1111/jav.02156>.
- [38] Mark S. Boyce and Simone Ciuti. Data from: Human selection of elk behavioural traits in a landscape of fear, 2020. URL <https://www.datarepository.movebank.org/handle/10255/move.1067>.
- [39] Christopher F. Brown, Steven P. Brumby, B. Guzder-Williams, Tyler Birch, Stephen B. Hyde, Joseph Mazzariello, William Czerwinski, Valerie J. Pasquarella, Robert Haertel, Sergey Ilyushchenko, Kurt Schwehr, M. J. Weisse, Fred Stolle, Craig Hanson, Oliver Guinan, and Ryan Moore. Dynamic world, near real-time global 10 m land use land cover mapping.

Scientific Data, 9:251, 2022. doi: 10.1038/s41597-022-01307-4. Accessed: 2026-04-01, This dataset is produced for the Dynamic World Project by Google in partnership with National Geographic Society and the World Resources Institute.

- [40] Christopher F Brown, Steven P Brumby, Brookie Guzder-Williams, Tanya Birch, Samantha Brooks Hyde, Joseph Mazzariello, Wanda Czerwinski, Valerie J Pasquarella, Robert Haertel, Simon Ilyushchenko, et al. Dynamic world, near real-time global 10 m land use land cover mapping. *Scientific data*, 9(1):251, 2022.
- [41] Christopher F Brown, Michal R Kazmierski, Valerie J Pasquarella, William J Rucklidge, Masha Samsikova, Chenhui Zhang, Evan Shelhamer, Estefania Lahera, Olivia Wiles, Simon Ilyushchenko, et al. Alphaearth foundations: An embedding field model for accurate and efficient global mapping from sparse label data. *arXiv preprint arXiv:2507.22291*, 2025.
- [42] Nelleke H. Buitendijk, Monique de Jager, Menno Hornman, Helmut Kruckenberg, Andrea Kölzsch, Sander Moonen, and Bart A. Nolet. More grazing, more damage? assessed yield loss on agricultural grassland relates nonlinearly to goose grazing pressure. *Journal of Applied Ecology*, 59(12):2878–2889, November 2022. ISSN 1365-2664. doi: 10.1111/1365-2664.14306. URL <http://dx.doi.org/10.1111/1365-2664.14306>.
- [43] Nelleke H. Buitendijk, Monique De Jager, Helmut Kruckenberg, Andrea Kölzsch, Sander Moonen, Gerhard J.D.M. Müskens, and Bart A. Nolet. Data from: More grazing, more damage? assessed yield loss on agricultural grassland relates non-linearly to goose grazing pressure, 2023. URL <https://datarepository.movebank.org/handle/10255/move.241>.
- [44] Patrik Byholm, Martin Beal, Natalie Isaksson, Ulrik Lötberg, and Susanne Åkesson. Paternal transmission of migration knowledge in a long-distance bird migrant. *Nature Communications*, 13(1), March 2022. ISSN 2041-1723. doi: 10.1038/s41467-022-29300-w. URL <http://dx.doi.org/10.1038/s41467-022-29300-w>.
- [45] Patrik Byholm, Martin Beal, Ulrik Lötberg, and Susanne Åkesson. Data from: Paternal transmission of migration knowledge in a long-distance bird migrant, 2022. URL <https://www.datarepository.movebank.org/handle/10255/move.1310>.
- [46] Patrik Byholm, Paweł Mirski, and Wouter Vansteelant. Data from: Study "gps tracking of honey buzzards in finland", 2025. URL <https://datarepository.movebank.org/handle/10255/move.3162>.
- [47] David Cabot. Data from: Forecasting spring from afar? timing of migration and predictability of phenology along different migration routes of an avian herbivore [greenland data], 2014. URL <https://www.datarepository.movebank.org/handle/10255/move.380>.
- [48] Francesca Cagnacci, Luigi Boitani, Roger A Powell, and Mark S Boyce. Animal ecology meets gps-based radiotelemetry: a perfect storm of opportunities and challenges. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 365(1550):2157–2162, 2010.
- [49] M. Carroll, C. DiMiceli, M. Wooten, A. Hubbard, R. Sohlberg, and J. Townshend. Mod44w modis/terra land water mask derived from modis and srtm l3 global 250m sin grid v006, 2017. Accessed: 2026-04-01.
- [50] Andrew Carter and Roshier David A. Data from: Toward reliable population density estimates of partially marked populations using spatially explicit mark-resight methods, 2020. URL <https://www.datarepository.movebank.org/handle/10255/move.827>.
- [51] Andrew Carter, Joanne M. Potts, and David A. Roshier. Toward reliable population density estimates of partially marked populations using spatially explicit mark-resight methods. *Ecology and Evolution*, 9(4):2131–2141, January 2019. ISSN 2045-7758. doi: 10.1002/ece3.4907. URL <http://dx.doi.org/10.1002/ece3.4907>.
- [52] Samuel Chambers, Joshua von Nonn, Matthew A. Burgess, Lance R. Brady, Jeffrey Bracewell, Daniel A. Guerra, and Miguel L. Villarreal. The tortoise and the antilocaprid: adapting gps tracking and terrain data to model wildlife walking functions. *Landscape Ecology*,

- 40(5), April 2025. ISSN 1572-9761. doi: 10.1007/s10980-025-02092-2. URL <http://dx.doi.org/10.1007/s10980-025-02092-2>.
- [53] Eric P Chassignet, Harley E Hurlburt, Ole Martin Smedstad, George R Halliwell, Patrick J Hogan, Alan J Wallcraft, Remy Baraille, and Rainer Bleck. The hycom (hybrid coordinate ocean model) data assimilative system. *Journal of Marine Systems*, 65(1-4):60–83, 2007.
 - [54] Nilanjan Chatterjee, David Wolfson, Dongmin Kim, Juliana Velez, Smith Freeman, Nathan M Bacheler, Kyle Shertzer, J Christopher Taylor, and John Fieberg. Modelling individual variability in habitat selection and movement using integrated step-selection analysis. *Methods in Ecology and Evolution*, 15(6):1034–1047, 2024.
 - [55] Eunjoon Cho, Seth A Myers, and Jure Leskovec. Friendship and mobility: user movement in location-based social networks. In *Proceedings of the 17th ACM SIGKDD international conference on Knowledge discovery and data mining*, pages 1082–1090, 2011.
 - [56] Kinley Choden, Sébastien Ravon, Jonathan H. Epstein, Thavry Hoem, Neil Furey, Marie Gely, Audrey Jolivot, Vibol Hul, Chhoeuth Neung, Annelise Tran, and Julien Cappelle. Data from: Pteropus lylei primarily forages in residential areas in kandal, cambodia, 2019. URL <https://www.datarepository.movebank.org/handle/10255/move.869>.
 - [57] Kinley Choden, Sébastien Ravon, Jonathan H. Epstein, Thavry Hoem, Neil Furey, Marie Gely, Audrey Jolivot, Vibol Hul, Chhoeuth Neung, Annelise Tran, and Julien Cappelle. Pteropus lylei primarily forages in residential areas in kandal, cambodia. *Ecology and Evolution*, 9(7):4181–4191, March 2019. ISSN 2045-7758. doi: 10.1002/ece3.5046. URL <http://dx.doi.org/10.1002/ece3.5046>.
 - [58] Beirne Christopher, Mark Thomas, Arianna Basto, Eleanor Flatt, Giancarlo Inga Diaz, Diego Rolim Chulla, Flor Perez Mullisaca, Rosio Vega Quispe, Caleb Jonatan Quispe Quispe, Adrian Forsyth, and Andrew Whitworth. Data from: Scouts vs usurpers: alternative foraging strategies facilitate coexistence between neotropical cathartid vultures, 2024. URL <https://datarepository.movebank.org/handle/10255/move.3080>.
 - [59] Mark W Chynoweth. Data from: Home range use and movement patterns of non-native feral goats in a tropical island montane dry landscape, 2015. URL <https://www.datarepository.movebank.org/handle/10255/move.410>.
 - [60] Mark W. Chynoweth, Christopher A. Lepczyk, Creighton M. Litton, Steven C. Hess, James R. Kellner, and Susan Cordell. Home range use and movement patterns of non-native feral goats in a tropical island montane dry landscape. *PLOS ONE*, 10(3):e0119231, March 2015. ISSN 1932-6203. doi: 10.1371/journal.pone.0119231. URL <http://dx.doi.org/10.1371/journal.pone.0119231>.
 - [61] Ondřej Cífka, Simon Chamaillé-Jammes, and Antoine Liutkus. Moveformer: a transformer-based model for step-selection animal movement modelling. *bioRxiv*, pages 2023–03, 2023.
 - [62] Simone Ciuti, Tyler B. Muhly, Dale G. Paton, Allan D. McDevitt, Marco Musiani, and Mark S. Boyce. Human selection of elk behavioural traits in a landscape of fear. *Proceedings of the Royal Society B: Biological Sciences*, 279(1746):4407–4416, 2012. ISSN 1471-2954. doi: 10.1098/rspb.2012.1483. URL <http://dx.doi.org/10.1098/rspb.2012.1483>.
 - [63] CCS Copernicus. Copernicus climate change service, 2020.
 - [64] Copernicus Climate Change Service. Era5 hourly data on single levels from 1940 to present, 2026. Accessed: 2026-04-01.
 - [65] Daniel Costa, Rachel Holser, Theresa Keates, Taiki Adachi, Roxanne Beltran, Cory Champagne, Crocker Daniel, Arina Favilla, Melinda Fowler, Juan Pablo Gallo-Reynoso, Chandra Goetsch, Jason Hassrick, Luis Hückstädt, Jessica Kendall-Bar, Sarah Kienle, Carey Kuhn, Jennifer Maresh, Sara Maxwell, Birgitte McDonald, Elizabeth McHuron, Patricia Morris, Yasuhiko Naito, Logan Pallin, Sarah Peterson, Patrick Robinson, Samantha Simmons, Akinori Takahashi, Nicole Teuschel, Michael Tift, Yann Tremblay, Stella Villegas-Amtman, and Ken Yoda. Northern elephant seal tracking and diving - raw and curated data, 2025. URL <https://datadryad.org/dataset/doi:10.7291/D10D61>.

- [66] Daniel P. Costa, Rachel R. Holser, Theresa R. Keates, Taiki Adachi, Roxanne S. Beltran, Cory D. Champagne, Daniel E. Crocker, Arina B. Favilla, Melinda A. Fowler, Juan Pablo Gallo-Reynoso, Chandra Goetsch, Jason L. Hassrick, Luis A. Hückstädt, Jessica M. Kendall-Bar, Sarah S. Kienle, Carey E. Kuhn, Jennifer L. Maresh, Sara M. Maxwell, Birgitte I. McDonald, Elizabeth A. McHuron, Patricia A. Morris, Yasuhiko Naito, Logan J. Pallin, Sarah H. Peterson, Patrick W. Robinson, Samantha E. Simmons, Akinori Takahashi, Nicole M. Teuschel, Michael S. Tift, Yann Tremblay, Stella Villegas-Amtmann, and Ken Yoda. Two decades of three-dimensional movement data from adult female northern elephant seals. *Scientific Data*, 11(1), December 2024. ISSN 2052-4463. doi: 10.1038/s41597-024-04084-4. URL <http://dx.doi.org/10.1038/s41597-024-04084-4>.
- [67] Mitchell A Cowan, Scott W Forrest, Samantha A Setterfield, Judy A Dunlop, Lesley A Gibson, and Dale G Nimmo. The impact of mining on animal movement and landscape connectivity revealed through simulations and scenarios. *Ecological Applications*, 35(7):e70134, 2025.
- [68] Gabriele Cozzi, Femke Broekhuis, J. Weldon McNutt, and Bernhard Schmid. Comparison of the effects of artificial and natural barriers on large african carnivores: Implications for interspecific relationships and connectivity. *Journal of Animal Ecology*, 82(3):707–715, February 2013. ISSN 1365-2656. doi: 10.1111/1365-2656.12039. URL <http://dx.doi.org/10.1111/1365-2656.12039>.
- [69] Paul C. Cross, Justin A. Bowers, Craig T. Hay, Julie Wolhuter, Peter Buss, Markus Hofmeyr, Johan T. Du Toit, and Wayne M. Getz. Data from: Nonparametric kernel methods for constructing home ranges and utilization distributions, 2016. URL <https://www.datarepository.movebank.org/handle/10255/move.609>.
- [70] Sebastian Cruz, Carolina B. Proaño, Dave Anderson, Kate Huyvaert, and Martin Wikelski. Data from: The environmental-data automated track annotation (env-data) system: Linking animal tracks with environmental data, 2013. URL <https://www.datarepository.movebank.org/handle/10255/move.331>.
- [71] J. A. Cummings and O. M. Smedstad. Variational data assimilation for the global ocean. In *Data Assimilation for Atmospheric, Oceanic and Hydrologic Applications II*, chapter 13, pages 303–343. 2013.
- [72] KJN d’Entremont, GK Davoren, CJ Walsh, SI Wilhelm, and WA Montevecchi. Intra- and inter-annual shifts in foraging tactics by parental northern gannets *morus bassanus* indicate changing prey fields. *Marine Ecology Progress Series*, 698:155–170, October 2022. ISSN 1616-1599. doi: 10.3354/meps14164. URL <http://dx.doi.org/10.3354/meps14164>.
- [73] Kyle J.N. D’Entremont, Gail K. Davoren, and William A. Montevecchi. Data from: Study "northern gannet breeding season gps data from cape st. mary’s, nl, canada: 2019 to 2022", 2023. URL <https://www.datarepository.movebank.org/handle/10255/move.1592>.
- [74] Somayeh Dodge, Gil Bohrer, Rolf Weinzierl, Sarah C Davidson, Roland Kays, David Douglas, Sebastian Cruz, Jiawei Han, David Brandes, and Martin Wikelski. The environmental-data automated track annotation (env-data) system: linking animal tracks with environmental data. *Movement Ecology*, 1(1), 2013. ISSN 2051-3933. doi: 10.1186/2051-3933-1-3. URL <http://dx.doi.org/10.1186/2051-3933-1-3>.
- [75] Somayeh Dodge, Gil Bohrer, Keith Bildstein, Sarah C. Davidson, Rolf Weinzierl, Marc J. Bechard, David Barber, Roland Kays, David Brandes, Jiawei Han, and Martin Wikelski. Environmental drivers of variability in the movement ecology of turkey vultures (*cathartes aura*) in north and south america. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 369(1643):20130195, May 2014. ISSN 1471-2970. doi: 10.1098/rstb.2013.0195. URL <http://dx.doi.org/10.1098/rstb.2013.0195>.
- [76] Yadolah Dodge and Valentin Rousson. Multivariate ll mean. *Metrika*, 49(2):127–134, 1999.
- [77] Tim S Doherty, Graeme C Hays, and Don A Driscoll. Human disturbance causes widespread disruption of animal movement. *Nature Ecology & Evolution*, 5(4):513–519, 2021.

- [78] Martin Dorber, Manuela Panzacchi, Olav Strand, and Bram van Moorter. New indicator of habitat functionality reveals high risk of underestimating trade-offs among sustainable development goals: The case of wild reindeer and hydropower. *Ambio*, 52(4):757–768, 2023.
- [79] Alyssa Eby, Allison Patterson, Shannon Whelan, Kyle H. Elliott, H. Grant Gilchrist, and Oliver P. Love. Data from: Influence of sea ice concentration, sex, and chick age on foraging flexibility and success in an arctic seabird, 2024. URL <https://datarepository.movebank.org/handle/10255/move.3178>.
- [80] Alyssa Eby, Allison Patterson, Shannon Whelan, Kyle H Elliott, H Grant Gilchrist, and Oliver P Love. Influence of sea ice concentration, sex and chick age on foraging flexibility and success in an arctic seabird. *Conservation Physiology*, 12(1), January 2024. ISSN 2051-1434. doi: 10.1093/conphys/coae057. URL <http://dx.doi.org/10.1093/conphys/coae057>.
- [81] Ron Efrat, Yael Lehnardt, Alexander Bragin, Evgeny Bragin, Tal Avgar, Todd Katzner, and Nir Sapir. Data from: Age-dependent response to anthropogenic habitat during migration of an endangered raptor, 2025. URL <https://datarepository.movebank.org/handle/10255/move.3548>.
- [82] Ron Efrat, Yael Lehnardt, Alexandr Bragin, Evgeny Bragin, Tal Avgar, Todd Katzner, and Nir Sapir. Age-dependent response to anthropogenic habitat during migration of an endangered raptor. *Current Biology*, 35(17):4301–4308.e3, 2025. ISSN 0960-9822. doi: 10.1016/j.cub.2025.07.061. URL <http://dx.doi.org/10.1016/j.cub.2025.07.061>.
- [83] Michael Egan, Nicole Gorman, Storm Crews, Michael W. Eichholz, Daniel Skinner, Peter Schlichting, Nathaniel Rayl, Eric J. Bergman, E. Hance Ellington, and Guillaume Bastille-Rousseau. Estimating encounter-habitat relationships with scale-integrated resource selection functions, 2024. URL https://figshare.com/collections/Estimating_Encounter-Habitat_Relationships_with_Scale-Integrated_Resource_Selection_Functions/7271164.
- [84] Michael E. Egan, Nicole T. Gorman, Storm Crews, Michael W. Eichholz, Dan Skinner, Peter E. Schlichting, Nathaniel D. Rayl, Eric J. Bergman, E. Hance Ellington, and Guillaume Bastille-Rousseau. Estimating encounter-habitat relationships with scale-integrated resource selection functions. *Journal of Animal Ecology*, 93(8):1036–1048, 2024. ISSN 1365-2656. doi: 10.1111/1365-2656.14133. URL <http://dx.doi.org/10.1111/1365-2656.14133>.
- [85] Michael E Egan, Nicole T Gorman, Michael W Eichholz, Dan Skinner, Peter E Schlichting, and Guillaume Bastille-Rousseau. Accounting for spatiotemporal patterns of long-term recursion in estimating local-scale step selection. *Methods in Ecology and Evolution*, 16(3):584–595, 2025.
- [86] Scott L. Eggeman, Mark Hebblewhite, Holger Bohm, Jesse Whittington, and Evelyn H. Merrill. Behavioural flexibility in migratory behaviour in a long-lived large herbivore. *Journal of Animal Ecology*, 85(3):785–797, March 2016. ISSN 1365-2656. doi: 10.1111/1365-2656.12495. URL <http://dx.doi.org/10.1111/1365-2656.12495>.
- [87] Joseph M Eisaguirre, Perry J Williams, and Mevin B Hooten. Rayleigh step-selection functions and connections to continuous-time mechanistic movement models. *Movement Ecology*, 12: 14, 2024.
- [88] Diego Ellis-Soto, Ruth Y Oliver, Vanessa Brum-Bastos, Urška Demšar, Brett Jesmer, Jed A Long, Francesca Cagnacci, Federico Ossi, Nuno Queiroz, Mark Hindell, et al. A vision for incorporating human mobility in the study of human–wildlife interactions. *Nature Ecology & Evolution*, 7(9):1362–1372, 2023.
- [89] Nicole Esteban, Jeanne A. Mortimer, and Graeme C. Hays. How numbers of nesting sea turtles can be overestimated by nearly a factor of two. *Proceedings of the Royal Society B: Biological Sciences*, 284(1849):20162581, February 2017. ISSN 1471-2954. doi: 10.1098/rspb.2016.2581. URL <http://dx.doi.org/10.1098/rspb.2016.2581>.
- [90] European Space Agency. Copernicus digital elevation model (dem) glo-30, 2023. Global 30 m digital surface model derived from WorldDEM; accessed 2026-04-01.

- [91] European Space Agency (ESA). Copernicus dem – global digital elevation model glo-30, 2019. URL <https://doi.org/10.5270/ESA-c5d3d65>.
- [92] European Union-Copernicus Marine Service. Global ocean waves reanalysis waverys, 2019. URL https://resources.marine.copernicus.eu/product-detail/GLOBAL_MULTIYEAR_WAV_001_032/INFORMATION. This study has been conducted using E.U. Copernicus Marine Service Information.
- [93] European Union-Copernicus Marine Service. Global ocean colour (copernicus-globcolour), bio-geo-chemical, l4 (monthly and interpolated) from satellite observations (1997-ongoing), 2022. URL https://resources.marine.copernicus.eu/product-detail/OCEANCOLOUR_GLO_BGC_L4_MY_009_104/INFORMATION.
- [94] Stephen E. Fick and Robert J. Hijmans. Worldclim 2: new 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology*, 37(12):4302–4315, 2017. doi: <https://doi.org/10.1002/joc.5086>. URL <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/joc.5086>.
- [95] Stephen E Fick and Robert J Hijmans. Worldclim 2: new 1-km spatial resolution climate surfaces for global land areas. *International journal of climatology*, 37(12):4302–4315, 2017.
- [96] Wolfgang Fiedler, Andrea Flack, Wolfgang Schäfle, Brigitta Keeves, Michael Quetting, Babette Eid, Heidi Schmid, and Martin Wikelski. Data from: Study "lifetrack white stork sw germany" (2013-2023), 2024. URL <https://datarepository.movebank.org/handle/10255/move.1685>.
- [97] Wolfgang Fiedler, Christiane Hilsendegen, Christian Reis, Jessica Lehmann, Pirmin Hilsendegen, Heidi Schmid, and Martin Wikelski. Data from: Study "lifetrack white stork rheinland-pfalz" (2015-2023), 2024. URL <https://datarepository.movebank.org/handle/10255/move.1687>.
- [98] Wolfgang Fiedler, Elke Leppelsack, Hans Leppelsack, Thomas Stahl, Oda Wieding, and Martin Wikelski. Data from: Study "lifetrack white stork bavaria" (2014-2023), 2024. URL <https://datarepository.movebank.org/handle/10255/move.1686>.
- [99] David A. Fifield, Robert A. Ronconi, and Gregory J. Robertson. Data from: Study "hering gulls (larus argentatus); fifield; witless bay, canada", 2020. URL <https://www.datarepository.movebank.org/handle/10255/move.1089>.
- [100] Manuela Fischer, Julian Di Stefano, Pierre Gras, Stephanie Kramer-Schadt, Duncan R. Sutherland, Graeme Coulson, and Milena Stillfried. Circadian rhythms enable efficient resource selection in a human-modified landscape. *Ecology and Evolution*, 9(13):7509–7527, 2019. ISSN 2045-7758. doi: [10.1002/ece3.5283](https://doi.org/10.1002/ece3.5283). URL <http://dx.doi.org/10.1002/ece3.5283>.
- [101] Manuela Fischer, Julian Di Stefano, Pierre Gras, Stephanie Kramer-Schadt, Duncan R. Sutherland, Graeme Coulson, and Milena Stillfried. Data from: Circadian rhythms enable efficient resource selection in a human-modified landscape, 2020. URL <https://www.datarepository.movebank.org/handle/10255/move.894>.
- [102] E. Fleishman, J. Anderson, B. G. Dickson, D. Krolick, J. A. Estep, R.L. Anderson, C. S. Elphick, D. S. Dobkin, and D. A. Bell. Space use by swainson's hawk (*buteo swainsoni*) in the natomas basin, california. *Collabra*, 2(1), January 2016. ISSN 2376-6832. doi: [10.1525/collabra.35](https://doi.org/10.1525/collabra.35). URL <http://dx.doi.org/10.1525/collabra.35>.
- [103] Erica Fleishman, Jesse Anderson, Brett G. Dickson, David Krolick, Jim A. Estep, Richard L. Anderson, Chris S. Elphick, David S. Dobkin, and Doug A. Bell. Data from: Space use by swainson's hawk (*buteo swainsoni*) in the natomas basin, california, 2017. URL <https://www.datarepository.movebank.org/handle/10255/move.673>.
- [104] Scott W Forrest, Dan Pagendam, Michael Bode, Christopher Drovandi, Jonathan R Potts, Justin Perry, Eric Vanderduys, and Andrew J Hoskins. Predicting fine-scale distributions and emergent spatiotemporal patterns from temporally dynamic step selection simulations. *Ecography*, 2025(2):e07421, 2025.

- [105] Scott W Forrest, Dan Pagendam, Conor Hassan, Jonathan R Potts, Christopher Drovandi, Michael Bode, and Andrew J Hoskins. Predicting animal movement with deepSSF: A deep learning step selection framework. *Methods in Ecology and Evolution*, 17(2):371–391, 2026.
- [106] Scott William Forrest. Predicting fine-scale distributions and emergent spatiotemporal patterns from temporally dynamic step selection simulations, 2024. URL <https://zenodo.org/doi/10.5281/zenodo.13749246>.
- [107] Daniel Fortin, Hawthorne L Beyer, Mark S Boyce, Douglas W Smith, Thierry Duchesne, and Julie S Mao. Wolves influence elk movements: behavior shapes a trophic cascade in yellowstone national park. *Ecology*, 86(5):1320–1330, 2005.
- [108] Guilad Friedemann, Yossi Leshem, and Ido Izhaki. Data from: Multidimensional differentiation in foraging resource use during breeding of two sympatric raptors: space, habitat type, time and food, 2016. URL <https://www.datarepository.movebank.org/handle/10255/move.483>.
- [109] Guilad Friedemann, Yossi Leshem, Lior Kerem, Boaz Shacham, Avi Bar-Massada, Krystaal M. McClain, Gil Bohrer, and Ido Izhaki. Multidimensional differentiation in foraging resource use during breeding of two sympatric top predators. *Scientific Reports*, 6(1), October 2016. ISSN 2045-2322. doi: 10.1038/srep35031. URL <http://dx.doi.org/10.1038/srep35031>.
- [110] Thibault Fronville, Maximilian Pichler, Johannes Signer, Marius Grabow, Stephanie Kramer-Schadt, Viktoriia Radchuk, and Florian Hartig. Analyzing animal movement using deep learning. *arXiv preprint arXiv:2603.24009*, 2026.
- [111] Ryo Fujii, Hideo Saito, and Ryo Hachiuma. Towards predicting any human trajectory in context. *arXiv preprint arXiv:2506.00871*, 2025.
- [112] Stefan Garthe, Philipp Schwemmer, Vitor H. Paiva, Anna-Marie Corman, Heino O. Fock, Christian C. Voigt, and Sven Adler. Data from: Terrestrial and marine foraging strategies of an opportunistic seabird species breeding in the wadden sea, 2016. URL <https://www.datarepository.movebank.org/handle/10255/move.573>.
- [113] Stefan Garthe, Philipp Schwemmer, Vitor H. Paiva, Anna-Marie Corman, Heino O. Fock, Christian C. Voigt, and Sven Adler. Terrestrial and marine foraging strategies of an opportunistic seabird species breeding in the wadden sea. *PLOS ONE*, 11(8):e0159630, August 2016. ISSN 1932-6203. doi: 10.1371/journal.pone.0159630. URL <http://dx.doi.org/10.1371/journal.pone.0159630>.
- [114] Stefan Garthe, Philipp Schwemmer, Ulrike Kubetzki, and Bernd Heinze. Data from: Effects of agricultural practices on foraging habitats of a seabird species in the baltic sea, 2022. URL <https://www.datarepository.movebank.org/handle/10255/move.1552>.
- [115] Stefan Garthe, Philipp Schwemmer, Ulrike Kubetzki, and Bernd Heinze. Effects of agricultural practices on foraging habitats of a seabird species in the baltic sea. *Ecology and Evolution*, 12(11), November 2022. ISSN 2045-7758. doi: 10.1002/ece3.9551. URL <http://dx.doi.org/10.1002/ece3.9551>.
- [116] Marie Claire Gatt, Paulo Lago, Martin Austad, Anne-Sophie Bonnet-Lebrun, and Benjamin J. Metzger. Pre-laying movements of yelkouan shearwaters (*puffinus yelkouan*) in the central mediterranean. *Journal of Ornithology*, 160(3):625–632, March 2019. ISSN 2193-7206. doi: 10.1007/s10336-019-01646-x. URL <http://dx.doi.org/10.1007/s10336-019-01646-x>.
- [117] Kaitlyn M Gaynor, Briana Abrahms, Kezia R Manlove, William K Oestreich, and Justine A Smith. Anthropogenic impacts at the interface of animal spatial and social behaviour. *Philosophical Transactions B*, 379(1912):20220527, 2024.
- [118] Brock Geary, Scott T. Walter, Paul L. Leberg, and Jordan Karubian. Data from: Condition-dependent foraging strategies in a coastal seabird: evidence for the rich get richer hypothesis, 2018. URL <https://www.datarepository.movebank.org/handle/10255/move.808>.

- [119] Brock Geary, Scott T Walter, Paul L Leberg, and Jordan Karubian. Condition-dependent foraging strategies in a coastal seabird: evidence for the rich get richer hypothesis. *Behavioral Ecology*, 30(2):356–363, December 2018. ISSN 1465-7279. doi: 10.1093/beheco/ary173. URL <http://dx.doi.org/10.1093/beheco/ary173>.
- [120] Wayne M. Getz, Scott Fortmann-Roe, Paul C. Cross, Andrew J. Lyons, Sadie J. Ryan, and Christopher C. Wilmers. Locoh: Nonparametric kernel methods for constructing home ranges and utilization distributions. *PLoS ONE*, 2(2):e207, February 2007. ISSN 1932-6203. doi: 10.1371/journal.pone.0000207. URL <http://dx.doi.org/10.1371/journal.pone.0000207>.
- [121] Thomas W. Glass and Martin D. Robards. Data from: Spatiotemporally variable snow properties drive habitat use of an arctic mesopredator, 2023. URL <https://datarepository.movebank.org/handle/10255/move.1651>.
- [122] Thomas W. Glass, Greg A. Breed, Glen E. Liston, Adele K. Reinking, Martin D. Robards, and Knut Kielland. Spatiotemporally variable snow properties drive habitat use of an arctic mesopredator. *Oecologia*, 195(4):887–899, March 2021. ISSN 1432-1939. doi: 10.1007/s00442-021-04890-2. URL <http://dx.doi.org/10.1007/s00442-021-04890-2>.
- [123] Tilmann Gneiting. Making and evaluating point forecasts. *Journal of the American Statistical Association*, 106(494):746–762, 2011.
- [124] Tilmann Gneiting and Adrian E Raftery. Strictly proper scoring rules, prediction, and estimation. *Journal of the American statistical Association*, 102(477):359–378, 2007.
- [125] Tilmann Gneiting, Fadoua Balabdaoui, and Adrian E Raftery. Probabilistic forecasts, calibration and sharpness. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 69(2):243–268, 2007.
- [126] Sara Gomez, Holly M English, Vanesa Bejarano Alegre, Paul G Blackwell, Anna M Bracken, Eloise Bray, Luke C Evans, Jelaine L Gan, W James Grecian, Catherine Gutmann Roberts, et al. Understanding and predicting animal movements and distributions in the anthropocene. *Journal of Animal Ecology*, 94(6):1146–1164, 2025.
- [127] Valentin Graf, Marjorie C. Sorensen, Thomas Mueller, and Elke Lena Neuschulz. Data from: Study "movement patterns of seed dispersing spotted nutcrackers (*nucifraga caryocatactes*)", 2024. URL <https://datarepository.movebank.org/handle/10255/move.3092>.
- [128] Max D Graham, Iain Douglas-Hamilton, William M Adams, and Phyllis C Lee. The movement of african elephants in a human-dominated land-use mosaic. *Animal conservation*, 12(5):445–455, 2009.
- [129] Larry Griffin. Data from: Forecasting spring from afar? timing of migration and predictability of phenology along different migration routes of an avian herbivore [svalbard data], 2014. URL <https://www.datarepository.movebank.org/handle/10255/move.377>.
- [130] Daniel A. Guerra, Todd C. Esque, Drew R. Davis, and Joseph A. Veech. Home range, seasonality, and the importance of canopy cover for texas tortoises (*gopherus berlandieri*). *Herpetologica*, 81(3), August 2025. ISSN 0018-0831. doi: 10.1655/herpetologica-d-24-00045. URL <http://dx.doi.org/10.1655/herpetologica-d-24-00045>.
- [131] Daniel A. Guerra, Joseph A. Veech, Todd C. Esque, and Drew R. Davis. Data from: Study "gps tracking of texas tortoises in south texas usa (guerra et al. 2025)", 2025. URL <https://datarepository.movebank.org/handle/10255/move.3446>.
- [132] Pratik Rajan Gupte, Christine E Beardsworth, Orr Spiegel, Emmanuel Lourie, Sivan Toledo, Ran Nathan, and Allert I Bijleveld. A guide to pre-processing high-throughput animal tracking data. *Journal of Animal Ecology*, 91(2):287–307, 2022.
- [133] S. E. Gutowsky, J. E. Baak, S. R. Craik, M. L. Mallory, N. Knutson, A. A. d’Entremont, and K. A. Allard. Seasonal and circadian patterns of herring gull (*larus smithsonianus*) movements reveal temporal shifts in industry and coastal island interaction. *Ecological Solutions and Evidence*, 4(3), 2023. ISSN 2688-8319. doi: 10.1002/2688-8319.12274. URL <http://dx.doi.org/10.1002/2688-8319.12274>.

- [134] Franziska Hacker, Charlotte Christensen, Grace H Davis, Marlee A Tucker, and Damien R Farine. Making movement ecology into a predictive science. 2026.
- [135] Roi Harel and Ran Nathan. Data from: The characteristic time scale of perceived information for decision-making: departure from thermal columns in soaring birds, 2018. URL <https://www.datarepository.movebank.org/handle/10255/move.751>.
- [136] Roi Harel and Ran Nathan. The characteristic time-scale of perceived information for decision-making: Departure from thermal columns in soaring birds. *Functional Ecology*, 32(8): 2065–2072, 2018. ISSN 1365-2435. doi: 10.1111/1365-2435.13136. URL <http://dx.doi.org/10.1111/1365-2435.13136>.
- [137] Graeme C. Hays, Jeanne A. Mortimer, Alex Rattray, Takahiro Shimada, and Nicole Esteban. Data from: High accuracy tracking reveals how small conservation areas can protect marine megafauna, 2021. URL <https://www.datarepository.movebank.org/handle/10255/move.1347>.
- [138] Graeme C. Hays, Jeanne A. Mortimer, Alex Rattray, Takahiro Shimada, and Nicole Esteban. High accuracy tracking reveals how small conservation areas can protect marine megafauna. *Ecological Applications*, 31(7), August 2021. ISSN 1939-5582. doi: 10.1002/eap.2418. URL <http://dx.doi.org/10.1002/eap.2418>.
- [139] Graeme C. Hays, Nicole Esteban, and Alex Rattray. Data from: Study "green turtles (chelonias mydas); hays; chagos archipelago, western indian ocean", 2024. URL <https://datarepository.movebank.org/handle/10255/move.1688>.
- [140] Mark Hebblewhite. Data from: Study "hebblewhite alberta-bc wolves", 2025. URL <https://datarepository.movebank.org/handle/10255/move.3469>.
- [141] Mark Hebblewhite and Evelyn H. Merrill. Multiscale wolf predation risk for elk: does migration reduce risk? *Oecologia*, 152(2):377–387, February 2007. ISSN 1432-1939. doi: 10.1007/s00442-007-0661-y. URL <http://dx.doi.org/10.1007/s00442-007-0661-y>.
- [142] Mark Hebblewhite, Evelyn Merrill, Hans Martin, Jodi E. Berg, Holger Bohm, and Scott L. Eggeman. Data from: Study "ya ha tinda elk project, banff national park, 2001-2020 (females)", 2020. URL <https://www.datarepository.movebank.org/handle/10255/move.1129>.
- [143] Jacob D. Hennig, J. D. Scasta, Aaron C. Pratt, Caitlyn P. Wanner, and Jeffrey L. Beck. Habitat selection and space use overlap between feral horses, pronghorn, and greater sage-grouse in cold arid steppe. *The Journal of Wildlife Management*, 87(1), November 2022. ISSN 1937-2817. doi: 10.1002/jwmg.22329. URL <http://dx.doi.org/10.1002/jwmg.22329>.
- [144] Jesús Hernández-Pliego, Carlos Rodríguez, and Javier Bustamante. Data from: Why do kestrels soar?, 2015. URL <https://www.datarepository.movebank.org/handle/10255/move.486>.
- [145] Jesús Hernández-Pliego, Carlos Rodríguez, and Javier Bustamante. Why do kestrels soar? *PLOS ONE*, 10(12):e0145402, December 2015. ISSN 1932-6203. doi: 10.1371/journal.pone.0145402. URL <http://dx.doi.org/10.1371/journal.pone.0145402>.
- [146] Hans Hersbach. Decomposition of the continuous ranked probability score for ensemble prediction systems. *Weather and Forecasting*, 15(5):559–570, 2000. doi: 10.1175/1520-0434(2000)015<0559:DOTCRP>2.0.CO;2.
- [147] Hans Hersbach, Bill Bell, Paul Berrisford, G. Biavati, A. Horányi, J. Muñoz Sabater, J. Nicolas, C. Peubey, R. Radu, I. Rozum, D. Schepers, A. Simmons, C. Soci, Dick Dee, and Jean-Noël Thépaut. Era5 hourly data on single levels from 1940 to present, 2018. Accessed: 2026-04-01.
- [148] Mark A Hindell, Ryan R Reisinger, Yan Ropert-Coudert, Luis A Hückstädt, Philip N Trathan, Horst Bornemann, Jean-Benoît Charrassin, Steven L Chown, Daniel P Costa, Bruno Danis, et al. Tracking of marine predators to protect southern ocean ecosystems. *Nature*, 580(7801): 87–92, 2020.

- [149] Benjamin Hoffman, Maddie Cusimano, Vittorio Baglione, Daniela Canestrari, Damien Chevalier, Dominic L. DeSantis, Lorène Jeantet, Monique A. Ladds, Takuya Maekawa, Vicente Mata-Silva, Víctor Moreno-González, Anthony Pagano, Eva Trapote, Outi Vainio, Antti Vehkaoja, Ken Yoda, Katherine Zacarian, and Ari S. Friedlaender. A benchmark for computational analysis of animal behavior, using animal-borne tags. *Movement Ecology*, 12(1), 2024. doi: 10.1186/s40462-024-00511-8.
- [150] David D Hofmann, Gabriele Cozzi, John W McNutt, Arpat Ozgul, and Dominik M Behr. A three-step approach for assessing landscape connectivity via simulated dispersal: African wild dog case study. *Landscape Ecology*, 38(4):981–998, 2023. doi: 10.1007/s10980-023-01602-4.
- [151] David D Hofmann, Dominik M Behr, John W McNutt, Arpat Ozgul, and Gabriele Cozzi. Dispersal and connectivity in increasingly extreme climatic conditions. *Global Change Biology*, 30(5):e17299, 2024.
- [152] David D Hofmann, Gabriele Cozzi, and John Fieberg. Methods for implementing integrated step-selection functions with incomplete data. *Movement Ecology*, 12(1):37, 2024.
- [153] Amanda E. Holland and Michael E. Byrne. Data from: Fine-scale assessment of home ranges and activity patterns for resident black vultures (*coragyps atratus*) and turkey vultures (*cathartes aura*), 2017. URL <https://www.datarepository.movebank.org/handle/10255/move.677>.
- [154] Amanda E. Holland, Michael E. Byrne, A. Lawrence Bryan, Travis L. DeVault, Olin E. Rhodes, and James C. Beasley. Fine-scale assessment of home ranges and activity patterns for resident black vultures (*coragyps atratus*) and turkey vultures (*cathartes aura*). *PLOS ONE*, 12(7):e0179819, 2017. ISSN 1932-6203. doi: 10.1371/journal.pone.0179819. URL <http://dx.doi.org/10.1371/journal.pone.0179819>.
- [155] Mevin B. Hooten, Xinyi Lu, Martha J. Garlick, and James A. Powell. Animal movement models with mechanistic selection functions. *Spatial Statistics*, 37:100406, 2020. ISSN 2211-6753. doi: 10.1016/j.spasta.2019.100406. URL <http://dx.doi.org/10.1016/j.spasta.2019.100406>.
- [156] Boyin Huang, Chunying Liu, Viva F. Banzon, Eric Freeman, Garrett Graham, William Hankins, Thomas M. Smith, and Huai-Min Zhang. Noaa 0.25-degree daily optimum interpolation sea surface temperature (oisst), version 2.1, 2020. URL <https://doi.org/10.25921/RE9P-PT57>.
- [157] Boyin Huang, Chunying Liu, Viva F. Banzon, Eric Freeman, Garrett Graham, William Hankins, Thomas M. Smith, and Huai-Min Zhang. Noaa 0.25-degree daily optimum interpolation sea surface temperature (oisst), version 2.1, 2020. [indicate subset used]; Accessed: 2026-05-04.
- [158] Elena I. Ilyashenko, Ivan Pokrovsky, Andrey E. Gavrilov, Syrymgul Zaripova, Wolfgang Fiedler, and Martin Wikelski. Data from: Study "1000 cranes. southern kazakhstan.", 2024. URL <https://datarepository.movebank.org/handle/10255/move.3255>.
- [159] Elena I. Ilyashenko, Ivan Pokrovsky, Valentin Yu. Ilyashenko, Mikhail Korepov, Wolfgang Fiedler, and Martin Wikelski. Data from: Study "1000 cranes. russia. common crane.", 2024. URL <https://datarepository.movebank.org/handle/10255/move.3257>.
- [160] Ladd Irvine, Daniel M. Palacios, Jorge Urbán, and Bruce Mate. Sperm whale dive behavior characteristics derived from intermediate-duration archival tag data. *Ecology and Evolution*, 7(19):7822–7837, August 2017. ISSN 2045-7758. doi: 10.1002/ece3.3322. URL <http://dx.doi.org/10.1002/ece3.3322>.
- [161] Ladd M. Irvine, Bruce R. Mate, Daniel M. Palacios, and Tomas M. Follett. Data from: Sperm whale dive behavior characteristics derived from intermediate-duration archival tag data, 2019. URL <https://www.datarepository.movebank.org/handle/10255/move.940>.
- [162] Ladd M. Irvine, Tomas M. Follett, Martha H. Winsor, Bruce R. Mate, and Daniel M. Palacios. Data from: Study "sperm whales gulf of mexico 2011-2013 - fastloc gps data", 2020. URL <https://www.datarepository.movebank.org/handle/10255/move.1058>.

- [163] Ladd M. Irvine, Martha H. Winsor, Tomas M. Follett, Bruce R. Mate, and Daniel M. Palacios. An at-sea assessment of argos location accuracy for three species of large whales, and the effect of deep-diving behavior on location error. *Animal Biotelemetry*, 8(1), 2020. ISSN 2050-3385. doi: 10.1186/s40317-020-00207-x. URL <http://dx.doi.org/10.1186/s40317-020-00207-x>.
- [164] LA Isbell and LR Bidner. Data from: Study "leopards, vervets, and baboons in laikipia, kenya" [baboons], 2024. URL <http://dx.doi.org/10.5441/001/1.625>.
- [165] Lynne A. Isbell, Laura R. Bidner, Margaret C. Crofoot, Akiko Matsumoto-Oda, and Damien R. Farine. Gps-identified, low-level nocturnal activity of vervets (*chlorocebus pygerythrus*) and olive baboons (*papio anubis*) in laikipia, kenya. *American Journal of Physical Anthropology*, 164(1):203–211, 2017. ISSN 1096-8644. doi: 10.1002/ajpa.23259. URL <http://dx.doi.org/10.1002/ajpa.23259>.
- [166] Jana W. E. Jeglinski, Jude V. Lane, Steven C. Votier, Robert W. Furness, Keith C. Hamer, Dominic J. McCafferty, Ruedi G. Nager, Maggie Sheddian, Sarah Wanless, and Jason Matthiopoulos. Hpaiv outbreak triggers short-term colony connectivity in a seabird metapopulation. *Scientific Reports*, 14(1), February 2024. ISSN 2045-2322. doi: 10.1038/s41598-024-53550-x. URL <http://dx.doi.org/10.1038/s41598-024-53550-x>.
- [167] JWE Jeglinski, JV Lane, SC Votier, RW Furness, KC Hamer, DJ McCafferty, R Nager, M Sheddian, S Wanless, and J Matthiopoulos. Data from: Hpaiv outbreak triggers short-term colony connectivity in a seabird metapopulation, 2024. URL <http://dx.doi.org/10.5441/001/1.607>.
- [168] Florian Jeltsch, Dries Bonte, Guy Pe'er, Björn Reineking, Peter Leimgruber, Niko Balkenhol, Boris Schröder, Carsten M Buchmann, Thomas Mueller, Niels Blaum, et al. Integrating movement ecology with biodiversity research-exploring new avenues to address spatiotemporal biodiversity dynamics. *Movement Ecology*, 1(1):6, 2013.
- [169] Scott Jennings, David Lumpkin, Nils Warnock, T. Emiko Condeso, and John P. Kelly. Great egret (*ardea alba*) habitat selection and foraging behavior in a temperate estuary: Comparing natural wetlands to areas with shellfish aquaculture. *PLOS ONE*, 16(12):e0261963, December 2021. ISSN 1932-6203. doi: 10.1371/journal.pone.0261963. URL <http://dx.doi.org/10.1371/journal.pone.0261963>.
- [170] Scott Jennings, David Lumpkin, Nils Warnock, and Emiko Condeso. Data from: Study "audubon canyon ranch egret telemetry project", 2025. URL <https://datarepository.movebank.org/handle/10255/move.3395>.
- [171] Brett R Jesmer, Jerod A Merkle, Jacob R Goheen, Ellen O Aikens, Jeffrey L Beck, Alyson B Courtemanch, Mark A Hurley, Douglas E McWhirter, Hollie M Miyasaki, Kevin L Monteith, et al. Is ungulate migration culturally transmitted? evidence of social learning from translocated animals. *Science*, 361(6406):1023–1025, 2018.
- [172] Arpit Kapoor, Anshul Negi, Lucy Marshall, and Rohitash Chandra. Cyclone trajectory and intensity prediction with uncertainty quantification using variational recurrent neural networks. *Environmental Modelling & Software*, 162:105654, 2023.
- [173] Roland Kays, Margaret C Crofoot, Walter Jetz, and Martin Wikelski. Terrestrial animal tracking as an eye on life and planet. *Science*, 348(6240):aaa2478, 2015.
- [174] Roland Kays, Ben Hirsch, Damien Caillaud, Rafael Mares, Shauhin Alavi, Rasmus Worsøe Havmøller, and Margaret Crofoot. Multi-scale movement syndromes for comparative analyses of animal movement patterns. *Movement Ecology*, 11(1), October 2023. ISSN 2051-3933. doi: 10.1186/s40462-022-00365-y. URL <http://dx.doi.org/10.1186/s40462-022-00365-y>.
- [175] Roland Kays, Ben T. Hirsch, Damien Caillaud, Rafael Mares, Shauhin Alavi, Rasmus Worsøe Havmøller, and Margaret C. Crofoot. Data from: Multi-scale movement syndromes for comparative analyses of animal movement patterns, 2023. URL <https://datarepository.movebank.org/handle/10255/move.1664>.

- [176] Kentaro Kazama, Katsuhide Fujita, Yushin Shinoda, and Shinsuke Koike. Sika deer trajectory prediction considering environmental factors by timeseries transformer-based architecture. *Expert systems with applications*, 250:123630, 2024.
- [177] Supawat Khaewphakdee, Laurel E.K. Serieys, and Rattapan Pattanarangsarn. Data from: Study "fishing cat (*prionairulus viverrinus*) study in khao sam roi yot, thailand", 2025. URL <https://datarepository.movebank.org/handle/10255/move.3400>.
- [178] Anjelika Kidd-Weaver, Jeffrey Hepinstall-Cymerman, Catharine C. Welch, Maureen H. Murray, Henry C. Adams, Taylor J. Ellison, Michael J. Yabsley, and Sonia M. Hernandez. Data from: The movements of a recently urbanized wading bird reveal changes in season timing and length related to resource use, 2020. URL <https://www.datarepository.movebank.org/handle/10255/move.1014>.
- [179] Anjelika Kidd-Weaver, Jeffrey Hepinstall-Cymerman, Catharine N. Welch, Maureen H. Murray, Henry C. Adams, Taylor J. Ellison, Michael J. Yabsley, and Sonia M. Hernandez. The movements of a recently urbanized wading bird reveal changes in season timing and length related to resource use. *PLOS ONE*, 15(3):e0230158, March 2020. ISSN 1932-6203. doi: 10.1371/journal.pone.0230158. URL <http://dx.doi.org/10.1371/journal.pone.0230158>.
- [180] Natasha J Klappstein, Théo Michelot, John Fieberg, Eric J Pedersen, and Joanna Mills Fleming. Step selection functions with non-linear and random effects. *Methods in ecology and evolution*, 15(8):1332–1346, 2024.
- [181] Natasha Jean Klappstein, Len Thomas, and Theo Michelot. Flexible hidden markov models for behaviour-dependent habitat selection. *Movement Ecology*, 11(1):30, 2023.
- [182] Erik Kleyheeg, Jacintha G. B. van Dijk, Despina Tsopoglou-Gkina, Tara Y. Woud, Dieuwertje K. Boonstra, Bart A. Nolet, and Merel B. Soons. Movement patterns of a keystone waterbird species are highly predictable from landscape configuration. *Movement Ecology*, 5(1), February 2017. ISSN 2051-3933. doi: 10.1186/s40462-016-0092-7. URL <http://dx.doi.org/10.1186/s40462-016-0092-7>.
- [183] Erik Kleyheeg, Jacintha G.B. Van Dijk, Bart A. Nolet, and Merel B. Soons. Data from: Movement patterns of a keystone waterbird species are highly predictable from landscape configuration, 2017. URL <https://www.datarepository.movebank.org/handle/10255/move.644>.
- [184] A. Kölzsch, G. J. D. M. Müskens, P. Szinai, S. Moonen, P. Glazov, H. Kruckenberg, M. Wikelski, and B. A. Nolet. Flyway connectivity and exchange primarily driven by moult migration in geese. *Movement Ecology*, 7(1), January 2019. ISSN 2051-3933. doi: 10.1186/s40462-019-0148-6. URL <http://dx.doi.org/10.1186/s40462-019-0148-6>.
- [185] A. Kölzsch, A. Flack, G. J. D. M. Müskens, H. Kruckenberg, P. Glazov, and M. Wikelski. Goose parents lead migration v. *Journal of Avian Biology*, 51(3), March 2020. ISSN 1600-048X. doi: 10.1111/jav.02392. URL <http://dx.doi.org/10.1111/jav.02392>.
- [186] Andrea Kölzsch, Silke Bauer, Rob de Boer, Larry Griffin, David Cabot, Klaus-Michael Exo, Henk P. van der Jeugd, and Bart A. Nolet. Forecasting spring from afar? timing of migration and predictability of phenology along different migration routes of an avian herbivore. *Journal of Animal Ecology*, 84(1):272–283, 2014. ISSN 1365-2656. doi: 10.1111/1365-2656.12281. URL <http://dx.doi.org/10.1111/1365-2656.12281>.
- [187] Andrea Kölzsch, Gerhard J.D.M. Müskens, Sander Moonen, Helmut Kruckenberg, Peter Glazov, and Martin Wikelski. Data from: Flyway connectivity and exchange primarily driven by moult migration in geese [north sea population], 2019. URL <https://www.datarepository.movebank.org/handle/10255/move.848>.
- [188] Andrea Kölzsch, Gerhard J.D.M. Müskens, Peter Glazov, Helmut Kruckenberg, and Martin Wikelski. Data from: Goose parents lead migration v, 2020. URL <https://www.datarepository.movebank.org/handle/10255/move.1030>.

- [189] Andrea Kölzsch, Thomas K. Lameris, Gerhard J. D. M. Müskens, Kees H. T. Schreven, Nelleke H. Buitendijk, Helmut Kruckenberg, Sander Moonen, Thomas Heinicke, Lei Cao, Jesper Madsen, Martin Wikelski, and Bart A. Nolet. Wild goose chase: Geese flee high and far, and with aftereffects from new year's fireworks. *Conservation Letters*, 16(1), November 2022. ISSN 1755-263X. doi: 10.1111/conl.12927. URL <http://dx.doi.org/10.1111/conl.12927>.
- [190] Andrea Kölzsch, Thomas K. Lameris, Gerhard J.D.M. Müskens, Kees H.T. Schreven, Nelleke H. Buitendijk, Helmut Kruckenberg, Sander Moonen, Thomas Heinicke, Lei Cao, Jesper Madsen, Martin Wikelski, and Bart A. Nolet. Data from: Wild goose chase: geese flee high and far, and with aftereffects from new year's fireworks, 2022. URL <https://www.datarepository.movebank.org/handle/10255/move.1526>.
- [191] Paulo Lago, Martin Austad, Benjamin Metzger, and Marie Claire Gatt. Data from: Pre-laying movements of yelkouan shearwaters (puffinus yelkouan) in the central mediterranean [2016-2017], 2019. URL <https://www.datarepository.movebank.org/handle/10255/move.866>.
- [192] Juliet S. Lamb, Yvan G. Satgé, and Patrick G. R. Jodice. Influence of density-dependent competition on foraging and migratory behavior of a subtropical colonial seabird. *Ecology and Evolution*, 7(16):6469–6481, 2017. ISSN 2045-7758. doi: 10.1002/ece3.3216. URL <http://dx.doi.org/10.1002/ece3.3216>.
- [193] Juliet S. Lamb, Yvan G. Satgé, and Patrick G.R. Jodice. Data from: Influence of density-dependent competition on foraging and migratory behavior of a subtropical colonial seabird, 2017. URL <https://www.datarepository.movebank.org/handle/10255/move.680>.
- [194] Moritz N Lang, Georg J Mayr, Reto Stauffer, and Achim Zeileis. Bivariate gaussian models for wind vectors in a distributional regression framework. *Advances in Statistical Climatology, Meteorology and Oceanography*, 5(2):115–132, 2019.
- [195] Liam P Langley, Stephen DJ Lang, Luke Ozsanlav-Harris, and Alice M Trevail. Exmove: An open-source toolkit for processing and exploring animal-tracking data in r. *Journal of Animal Ecology*, 93(7):784–795, 2024.
- [196] Roland Langrock, Ruth King, Jason Matthiopoulos, Len Thomas, Daniel Fortin, and Juan M. Morales. Flexible and practical modeling of animal telemetry data: Hidden markov models and extensions. *Ecology*, 93(11):2336–2342, 2012. doi: 10.1890/11-2241.1.
- [197] Scott LaPoint, Paul Gallery, Martin Wikelski, and Roland Kays. Data from: Animal behavior, cost-based corridor models, and real corridors, 2013. URL <https://www.datarepository.movebank.org/handle/10255/move.328>.
- [198] Scott LaPoint, Paul Gallery, Martin Wikelski, and Roland Kays. Animal behavior, cost-based corridor models, and real corridors. *Landscape Ecology*, 28(8):1615–1630, 2013. ISSN 1572-9761. doi: 10.1007/s10980-013-9910-0. URL <http://dx.doi.org/10.1007/s10980-013-9910-0>.
- [199] A. David M. Latham and Stan Boutin. Caribou, primary prey and wolf spatial relationships in northeastern alberta. *Rangifer*, page 155, 2011. ISSN 1890-6729. doi: 10.7557/2.31.2.2006. URL <http://dx.doi.org/10.7557/2.31.2.2006>.
- [200] A. David M. Latham and Stan Boutin. Data from: Wolf ecology and caribou-primary prey-wolf spatial relationships in low productivity peatland complexes in northeastern alberta, 2019. URL <https://www.datarepository.movebank.org/handle/10255/move.948>.
- [201] Songwei Li, Jie Feng, Jiawei Chi, Xinyuan Hu, Xiaomeng Zhao, and Fengli Xu. Limp: Large language model enhanced intent-aware mobility prediction. *arXiv preprint arXiv:2408.12832*, 2024.
- [202] Damian C. Lidgard, W. Don Bowen, and Sara J. Iverson. Data from: A novel approach to quantifying the spatiotemporal behavior of instrumented grey seals used to sample the environment, 2015. URL <https://www.datarepository.movebank.org/handle/10255/move.449>.

- [203] Yu Liu, Qian Ge, Wei Luo, Qiang Huang, Lei Zou, Haixu Wang, Xin Li, and Chang Liu. Graphmm: Graph-based vehicular map matching by leveraging trajectory and road correlations. *IEEE Transactions on Knowledge and Data Engineering*, 36(1):184–198, 2023.
- [204] Julie M. Mallon, Keith L. Bildstein, and William F. Fagan. Inclement weather forces stopovers and prevents migratory progress for obligate soaring migrants. *Movement Ecology*, 9(1), 2021. ISSN 2051-3933. doi: 10.1186/s40462-021-00274-6. URL <http://dx.doi.org/10.1186/s40462-021-00274-6>.
- [205] Mark L. Mallory, Shawn Craik, Karel A. Allard, and Sarah Gutowsky. Data from: Study "american herring gulls - gps - lobster bay, southwest nova scotia, canada", 2023. URL <https://datarepository.movebank.org/handle/10255/move.1654>.
- [206] Doris Matthes, Neus Latorre-Margalef, Andreas Schmidt, Jonas Waldenström, Martin Wikelski, and Mariëlle L. Van Toor. Data from: Flexibility of continental navigation and migration in european mallards, 2013. URL <https://www.datarepository.movebank.org/handle/10255/move.339>.
- [207] Bernie J. McConnell. Data from: State-switching continuous-time correlated random walks, 2019. URL <https://www.datarepository.movebank.org/handle/10255/move.844>.
- [208] J McLean Sloughter, Tilmann Gneiting, and Adrian E Raftery. Probabilistic wind vector forecasting using ensembles and bayesian model averaging. *Monthly Weather Review*, 141(6): 2107–2119, 2013.
- [209] Laura A. McMahon, Janet L. Rachlow, Lisa A. Shipley, and Jennifer S. Forbey. Data from: Evaluation of micro-gps receivers for tracking small-bodied mammals, 2017. URL <https://www.datarepository.movebank.org/handle/10255/move.659>.
- [210] Laura A. McMahon, Janet L. Rachlow, Lisa A. Shipley, Jennifer S. Forbey, Timothy R. Johnson, and Peter J. Olsoy. Evaluation of micro-gps receivers for tracking small-bodied mammals. *PLOS ONE*, 12(3):e0173185, March 2017. ISSN 1932-6203. doi: 10.1371/journal.pone.0173185. URL <http://dx.doi.org/10.1371/journal.pone.0173185>.
- [211] E. P. Medici, S. Mezzini, C. H. Fleming, J. M. Calabrese, and M. J. Noonan. Movement ecology of vulnerable lowland tapirs between areas of varying human disturbance. *Movement Ecology*, 10(1), March 2022. ISSN 2051-3933. doi: 10.1186/s40462-022-00313-w. URL <http://dx.doi.org/10.1186/s40462-022-00313-w>.
- [212] E. Patricia Medici. Data from: Study "lowland tapirs, tapirus terrestris, in southern brazil", 2023. URL <https://www.datarepository.movebank.org/handle/10255/move.1416>.
- [213] Jorge Medina Hernández, Jorge P Rodríguez, Clive R McMahon, Ana MM Sequeira, and Víctor M Eguíluz. Improving prediction region accuracy in marine animal movement with temporal fusion transformer. *Scientific Reports*, 2025.
- [214] Katherine Mertes, Marta A. Jarzyna, and Walter Jetz. Hierarchical multi-grain models improve descriptions of species' environmental associations, distribution, and abundance. *Ecological Applications*, 30(6), May 2020. ISSN 1939-5582. doi: 10.1002/eap.2117. URL <http://dx.doi.org/10.1002/eap.2117>.
- [215] Katherine Mertes, Walter Jetz, and Martin Wikelski. Data from: Hierarchical multi-grain models improve descriptions of species' environmental associations, distribution, and abundance, 2020. URL <https://www.datarepository.movebank.org/handle/10255/move.1052>.
- [216] Julie Mestre, Ana Rita Patrício, Ebaye Sidina, Cheibani Senhoury, Nahi El'bar, Martin Beal, Aissa Regalla, and Paulo Catry. Movement patterns of green turtles at a key foraging site: the banc d'arguin, mauritania. *Marine Biology*, 172(1), November 2024. ISSN 1432-1793. doi: 10.1007/s00227-024-04558-4. URL <http://dx.doi.org/10.1007/s00227-024-04558-4>.

- [217] Théo Michelot and Paul G. Blackwell. State-switching continuous-time correlated random walks. *Methods in Ecology and Evolution*, 10(5):637–649, February 2019. ISSN 2041-210X. doi: 10.1111/2041-210x.13154. URL <http://dx.doi.org/10.1111/2041-210x.13154>.
- [218] Paweł Mirski, Wouter M.G. Vansteelant, and Patrik Byholm. Dispersal, maturation and recruitment in a long-lived, intercontinental migrant bird. *Animal Behaviour*, 230:123366, December 2025. ISSN 0003-3472. doi: 10.1016/j.anbehav.2025.123366. URL <http://dx.doi.org/10.1016/j.anbehav.2025.123366>.
- [219] Franco Molteni, Roberto Buizza, Tim N Palmer, and Thomas Petroligis. The ecmwf ensemble prediction system: Methodology and validation. *Quarterly journal of the royal meteorological society*, 122(529):73–119, 1996.
- [220] Ronaldo G. Morato, Jeffrey J. Thompson, Agustin Paviolo, Jesus A. de La Torre, Fernando Lima, Roy T. McBride, Rogerio C. Paula, Laury Cullen, Leandro Silveira, Daniel L. Z. Kantek, Emiliano E. Ramalho, Louise Maranhão, Mario Haberkfeld, Denis A. Sana, Rodrigo A. Medellin, Eduardo Carrillo, Victor Montalvo, Octavio Monroy-Vilchis, Paula Cruz, Anah T. Jacomo, Natalia M. Torres, Giselle B. Alves, Ivonne Cassaigne, Ron Thompson, Carolina Saens-Bolanos, Juan Carlos Cruz, Luiz D. Alfaro, Isabel Hagnauer, Xavier Marina da Silva, Alexandre Vogliotti, Marcela F. D. Moraes, Selma S. Miyazaki, Thadeu D. C. Pereira, Gedionson R. Araujo, Leanes Cruz da Silva, Lucas Leuzinger, Marina M. Carvalho, Lilian Rampin, Leonardo Sartorello, Howard Quigley, Fernando Tortato, Rafael Hoogesteijn, Peter G. Crawshaw, Allison L. Devlin, Joares A. May, Fernando C. C. de Azevedo, Henrique V. B. Concone, Veronica A. Quiroga, Sebastian A. Costa, Juan P. Arrabal, Ezequiel Vanderhoeven, Yamil E. Di Blanco, Alexandre M. C. Lopes, Cynthia E. Widmer, and Milton Cezar Ribeiro. Jaguar movement database: a gps-based movement dataset of an apex predator in the neotropics. *Ecology*, 99(7):1691–1691, 2018. ISSN 1939-9170. doi: 10.1002/ecy.2379. URL <http://dx.doi.org/10.1002/ecy.2379>.
- [221] Ronaldo Gonçalves Morato, Daniel Luiz Zanella Kantek, Samiko Miyazaki, Thadeu Deluque, and Rogerio Cunha De Paula. Data from: Jaguar movement database-a gps-based movement dataset of an apex predator in the neotropics, 2021. URL <https://www.datarepository.movebank.org/handle/10255/move.1364>.
- [222] Michelangelo Morganti, Enrico Viganò, Alessandro Berlusconi, Sara Cioccarelli, and Mauro Fasola. Data from: Post-fledging habitat selection of a purple heron ardea purpurea revealed by gps/gsm telemetry, 2021. URL <https://www.datarepository.movebank.org/handle/10255/move.1367>.
- [223] Michelangelo Morganti, Enrico Viganò, Alessandro Berlusconi, Sara Cioccarelli, and Mauro Fasola. Post-fledging habitat selection of a purple heron ardea purpurea revealed by gps/gsm telemetry. *Avocetta*, 45(2), 2021. ISSN 0404-4266. doi: 10.30456/avo.2021201. URL <http://dx.doi.org/10.30456/avo.2021201>.
- [224] CD Morrison, MS Boyce, and SE Nielsen. Space-use, movement and dispersal of sub-adult cougars in a geographically isolated population. *peerj* 3: e1118, 2015.
- [225] Henrik Mouritsen. Long-distance navigation and magnetoreception in migratory animals. *Nature*, 558(7708):50–59, 2018.
- [226] MoveBank. Movebank for animal tracking, 2025. URL <https://www.movebank.org/cms/movebank-main>. <https://www.movebank.org/cms/movebank-main>.
- [227] Stefanie Muff, Johannes Signer, and John Fieberg. Accounting for individual-specific variation in habitat-selection studies: Efficient estimation of mixed-effects models using bayesian or frequentist computation. *Journal of Animal Ecology*, 89(1):80–92, 2020.
- [228] Rhys Munden, Luca Börger, Rory P Wilson, James Redcliffe, Rowan Brown, Mathieu Garel, and Jonathan R Potts. Why did the animal turn? time-varying step selection analysis for inference between observed turning-points in high frequency data. *Methods in Ecology and Evolution*, 12(5):921–932, 2021.

- [229] Joaquín Muñoz-Sabater, Emanuel Dutra, Anna Agustí-Panareda, Clément Albergel, Gabriele Arduini, Gianpaolo Balsamo, Souhail Boussetta, Margarita Choulga, Shaun Harrigan, Hans Hersbach, Brecht Martens, Diego G. Miralles, María Piles, Nemesio J. Rodríguez-Fernández, Ervin Zsoter, Carlo Buontempo, and Jean-Noël Thépaut. Era5-land: a state-of-the-art global reanalysis dataset for land applications. *Earth System Science Data*, 13(9):4349–4383, 2021. ISSN 1866-3516. doi: 10.5194/essd-13-4349-2021. URL <http://dx.doi.org/10.5194/essd-13-4349-2021>.
- [230] Gerhard J.D.M. Müskens, Péter Szinai, Tamas Sapi, Andrea Kölzsch, Martin Wikelski, and Bart A. Nolet. Data from: Flyway connectivity and exchange primarily driven by moult migration in geese [pannonic population], 2019. URL <https://www.datarepository.movebank.org/handle/10255/move.851>.
- [231] Rohit Naniwadekar, Akanksha Rathore, Ushma Shukla, Saniya Chaplod, and Aparajita Datta. Data from: How far do asian forest hornbills disperse seeds?, 2019. URL <https://www.datarepository.movebank.org/handle/10255/move.990>.
- [232] Rohit Naniwadekar, Akanksha Rathore, Ushma Shukla, Saniya Chaplod, and Aparajita Datta. How far do asian forest hornbills disperse seeds? *Acta Oecologica*, 101:103482, November 2019. ISSN 1146-609X. doi: 10.1016/j.actao.2019.103482. URL <http://dx.doi.org/10.1016/j.actao.2019.103482>.
- [233] NASA LP DAAC. Mcd12q1 version 6.1: Modis land cover type yearly global 500 m, 2024. Accessed: 2026-04-01.
- [234] NASA LP DAAC. Mod44b version 6.1: Vegetation continuous fields yearly global 250 m, 2024. Accessed: 2026-04-01.
- [235] Ran Nathan, Wayne M Getz, Eloy Revilla, Marcel Holyoak, Ronen Kadmon, David Saltz, and Peter E Smouse. A movement ecology paradigm for unifying organismal movement research. *Proceedings of the National Academy of Sciences*, 105(49):19052–19059, 2008.
- [236] Blaise A. Newman, Jordan R. Dyal, Karl V. Miller, Michael J. Cherry, and Gino J. D’Angelo. Data from: Influence of visual perception on movement decisions by an ungulate prey species, 2023. URL <https://datarepository.movebank.org/handle/10255/move.1657>.
- [237] Blaise A. Newman, Jordan R. Dyal, Karl V. Miller, Michael J. Cherry, and Gino J. D’Angelo. Influence of visual perception on movement decisions by an ungulate prey species. *Biology Open*, 12(10), October 2023. ISSN 2046-6390. doi: 10.1242/bio.059932. URL <http://dx.doi.org/10.1242/bio.059932>.
- [238] Paul Newson and John Krumm. Hidden markov map matching through noise and sparseness. In *Proceedings of the 17th ACM SIGSPATIAL International Conference on Advances in Geographic Information Systems*, 2009. doi: 10.1145/1653771.1653818.
- [239] Aurélien Nicosia, Thierry Duchesne, Louis-Paul Rivest, and Daniel Fortin. A multi-state conditional logistic regression model for the analysis of animal movement. 2017.
- [240] Michael J Noonan, Marlee A Tucker, Christen H Fleming, Thomas S Akre, Susan C Alberts, Abdullahi H Ali, Jeanne Altmann, Pamela Castro Antunes, Jerrold L Belant, Dean Beyer, et al. A comprehensive analysis of autocorrelation and bias in home range estimation. *Ecological Monographs*, 89(2):e01344, 2019.
- [241] Joseph M. Northrup, Eric Vander Wal, Melinda Bonar, John Fieberg, Martin P. Laforge, Mylène Leclerc, Claire M. Prokopenko, and Brian D. Gerber. Conceptual and methodological advances in habitat-selection modeling: Guidelines for ecology and evolution. *Ecological Applications*, 32(1):e02470, 2022. doi: 10.1002/eap.2470.
- [242] Meghan O’Connell, moreiraMatias, and Wendy Kan. Ecml/pkdd 15: Taxi trajectory prediction (i). <https://kaggle.com/competitions/pkdd-15-predict-taxi-service-trajectory-i>, 2015. Kaggle.

- [243] Oliver Padget, Geoff Stanley, Jay K. Willis, Annette L. Fayet, Sara Bond, Louise Maurice, Akiko Shoji, Ben Dean, Holly Kirk, Ignacio Juarez-Martinez, Robin Freeman, Mark Bolton, and Tim Guilford. Data from: Shearwaters know the direction and distance home but fail to encode intervening obstacles after free-ranging foraging trips, 2019. URL <https://www.datarepository.movebank.org/handle/10255/move.983>.
- [244] Oliver Padget, Geoff Stanley, Jay K. Willis, Annette L. Fayet, Sarah Bond, Louise Maurice, Akiko Shoji, Ben Dean, Holly Kirk, Ignacio Juarez-Martinez, Robin Freeman, Mark Bolton, and Tim Guilford. Shearwaters know the direction and distance home but fail to encode intervening obstacles after free-ranging foraging trips. *Proceedings of the National Academy of Sciences*, 116(43):21629–21633, October 2019. ISSN 1091-6490. doi: 10.1073/pnas.1903829116. URL <http://dx.doi.org/10.1073/pnas.1903829116>.
- [245] Ana Rita Patrício, Castro Barbosa, Martin Beal, Julie Mestre, Cheibani Senhoury, and Paulo Catry. Data from: Study "chelonias mydas banc arguin 2020 2021", 2024. URL <https://datarepository.movebank.org/handle/10255/move.3380>.
- [246] Bruce D Patterson, Samuel M Kasiki, Edwin Selempo, and Roland W Kays. Livestock predation by lions (*panthera leo*) and other carnivores on ranches neighboring tsavo national parks, kenya. *Biological Conservation*, 119(4):507–516, October 2004. ISSN 0006-3207. doi: 10.1016/j.biocon.2004.01.013. URL <http://dx.doi.org/10.1016/j.biocon.2004.01.013>.
- [247] Toby A Patterson, Len Thomas, Chris Wilcox, Otso Ovaskainen, and Jason Matthiopoulos. State-space models of individual animal movement. *Trends in ecology & evolution*, 23(2): 87–94, 2008.
- [248] Sasha Pekarsky, Ingo Schiffner, Yuri Markin, and Ran Nathan. Using movement ecology to evaluate the effectiveness of multiple human-wildlife conflict management practices. *Biological Conservation*, 262:109306, 2021.
- [249] Jean-Francois Pekel, Andrew Cottam, Noel Gorelick, and Alan S. Belward. High-resolution mapping of global surface water and its long-term changes. *Nature*, 540:418–422, 2016. doi: 10.1038/nature20584.
- [250] Jean-François Pekel, Andrew Cottam, Noel Gorelick, and Alan S Belward. High-resolution mapping of global surface water and its long-term changes. *Nature*, 540(7633):418–422, 2016.
- [251] Stefano Pellegrini, Andreas Ess, Konrad Schindler, and Luc Van Gool. You’ll never walk alone: Modeling social behavior for multi-target tracking. In *2009 IEEE 12th international conference on computer vision*, pages 261–268. IEEE, 2009.
- [252] Francesco Pezzo, Marco Zenatello, Giulia Cerritelli, Augusto Navone, Dimitri Giunchi, Giovanna Spano, Enrica Pollonara, Alessandro Massolo, Anna Gagliardo, and Nicola Baccetti. Data from: Productivity changes in the mediterranean sea drive foraging movements of yelkouan shearwater *puffinus yelkouan* from the core of its global breeding range, 2021. URL <https://www.datarepository.movebank.org/handle/10255/move.1326>.
- [253] Francesco Pezzo, Marco Zenatello, Giulia Cerritelli, Augusto Navone, Dimitri Giunchi, Giovanna Spano, Enrica Pollonara, Alessandro Massolo, Anna Gagliardo, and Nicola Baccetti. Productivity changes in the mediterranean sea drive foraging movements of yelkouan shearwater *puffinus yelkouan* from the core of its global breeding range. *Marine Ecology*, 42(4), 2021. ISSN 1439-0485. doi: 10.1111/maec.12668. URL <http://dx.doi.org/10.1111/maec.12668>.
- [254] Antti Piironen, Antti Paasivaara, and Toni Laaksonen. Data from: Birds of three worlds: moult migration to high arctic expands a boreal-temperate flyway to a third biome, 2021. URL <https://www.datarepository.movebank.org/handle/10255/move.1266>.
- [255] Antti Piironen, Antti Paasivaara, and Toni Laaksonen. Birds of three worlds: moult migration to high arctic expands a boreal-temperate flyway to a third biome. *Movement Ecology*, 9(1), 2021. ISSN 2051-3933. doi: 10.1186/s40462-021-00284-4. URL <http://dx.doi.org/10.1186/s40462-021-00284-4>.

- [256] Ruthmery Pillco Huarcaya, Andrew Whitworth, Norma Mamani, Mark Thomas, and Elias Condori. Through the eyes of the andean bear: Camera collar insights into the life of a threatened south american ursid. *Ecology and Evolution*, 14(12), December 2024. ISSN 2045-7758. doi: 10.1002/ece3.70304. URL <http://dx.doi.org/10.1002/ece3.70304>.
- [257] Ruthmery Pillco Huarcaya, Andrew Whitworth, and Mark Thomas. Data from: Study "using conservation technologies to understand the life of andean bears in and near manu national park" [part], 2024. URL <https://datarepository.movebank.org/handle/10255/move.3359>.
- [258] Jennifer Pohle, Johannes Signer, Jana A Eccard, Melanie Dammhahn, and Ulrike E Schlägel. How to account for behavioral states in step-selection analysis: a model comparison. *PeerJ*, 12:e16509, 2024.
- [259] Ivan Pokrovsky, Olga Kulikova, and Martin Wikelski. Data from: Longer days enable higher diurnal activity for migratory birds [rough-legged buzzards], 2021. URL <https://www.datarepository.movebank.org/handle/10255/move.1289>.
- [260] Ivan Pokrovsky, Andrea Kölzsch, Sherub Sherub, Wolfgang Fiedler, Peter Glazov, Olga Kulikova, Martin Wikelski, and Andrea Flack. Longer days enable higher diurnal activity for migratory birds. *Journal of Animal Ecology*, 90(9):2161–2171, March 2021. ISSN 1365-2656. doi: 10.1111/1365-2656.13484. URL <http://dx.doi.org/10.1111/1365-2656.13484>.
- [261] Tom A. Porteus, Mike J. Short, Andrew N. Hoodless, and Jonathan C. Reynolds. Movement ecology and minimum density estimates of red foxes in wet grassland habitats used by breeding wading birds. *European Journal of Wildlife Research*, 70(1), December 2023. ISSN 1439-0574. doi: 10.1007/s10344-023-01759-y. URL <http://dx.doi.org/10.1007/s10344-023-01759-y>.
- [262] Tom A. Porteus, Mike J. Short, Andrew N. Hoodless, and Jonathan C. Reynolds. Data from: Study "red fox (vulpes vulpes) in uk wet grasslands", 2024. URL <https://datarepository.movebank.org/handle/10255/move.1677>.
- [263] Jonathan R Potts and Luca Börger. How to scale up from animal movement decisions to spatiotemporal patterns: An approach via step selection. *Journal of Animal Ecology*, 92(1): 16–29, 2023.
- [264] Jonathan R Potts, Luca Börger, Bronson K Strickland, and Garrett M Street. Assessing the predictive power of step selection functions: How social and environmental interactions affect animal space use. *Methods in Ecology and Evolution*, 13(8):1805–1818, 2022.
- [265] R. John Power and Stephen Dell. Data from: A note on the reestablishment of the cheetah population in the pilanesberg national park, south africa, 2020. URL <https://www.datarepository.movebank.org/handle/10255/move.1215>.
- [266] R. John Power and Leanne Venter. Data from: Repatriating leopards into novel landscapes of a south african province, 2021. URL <https://www.datarepository.movebank.org/handle/10255/move.1247>.
- [267] R. John Power, Vincent Van der Merwe, Samantha Page-Nicholson, Mia V. Botha, Stephen Dell, and Pieter Nel. A note on the reestablishment of the cheetah population in the pilanesberg national park, south africa. *African Journal of Wildlife Research*, 49(1), February 2019. ISSN 2410-7220. doi: 10.3957/056.049.0012. URL <http://dx.doi.org/10.3957/056.049.0012>.
- [268] R. John Power, Leanne Venter, Mia-Vasti Botha, and Paul Bartels. Repatriating leopards into novel landscapes of a south african province. *Ecological Solutions and Evidence*, 2(1), January 2021. ISSN 2688-8319. doi: 10.1002/2688-8319.12046. URL <http://dx.doi.org/10.1002/2688-8319.12046>.
- [269] Laura R. Prugh. Data from: Study "gps tracking of bobcats and coyotes in northern washington", 2023. URL <https://datarepository.movebank.org/handle/10255/move.239>.

- [270] Laura R. Prugh, Calum X. Cunningham, Rebecca M. Windell, Brian N. Kertson, Taylor R. Ganz, Savanah L. Walker, and Aaron J. Wirsing. Fear of large carnivores amplifies human-caused mortality for mesopredators. *Science*, 380(6646):754–758, May 2023. ISSN 1095-9203. doi: 10.1126/science.adf2472. URL <http://dx.doi.org/10.1126/science.adf2472>.
- [271] Nathan Ranc, Francesca Cagnacci, and Paul R Moorcroft. Memory drives the formation of animal home ranges: evidence from a reintroduction. *Ecology Letters*, 25(4):716–728, 2022.
- [272] Nathan Ranc, John W Cain, Francesca Cagnacci, and Paul R Moorcroft. The role of memory-based movements in the formation of animal home ranges. *Journal of Mathematical Biology*, 88(5):59, 2024.
- [273] Maria L Rizzo and Gábor J Székely. Energy distance. *wiley interdisciplinary reviews: Computational statistics*, 8(1):27–38, 2016.
- [274] Will Rogers, Scott Yanco, and Walter Jetz. Choices to landscapes: Mechanisms of animal movement scale to landscape patterns of space use. *Ecology Letters*, 28(12):e70279, 2025.
- [275] Robert A. Ronconi and Katherine R. Shlepr. Herring gulls (*larus argentatus*); ronconi; brier island canada, 2020. URL <https://www.datarepository.movebank.org/handle/10255/move.1087>.
- [276] Robert A. Ronconi and Katherine R. Shlepr. Data from: Study "herring gulls (*larus argentatus*); ronconi; brier island, canada", 2020. URL <https://www.datarepository.movebank.org/handle/10255/move.1086>.
- [277] Robert A. Ronconi and Philip D. Taylor. Data from: Study "herring gulls (*larus argentatus*); ronconi; sable island, canada", 2020. URL <https://www.datarepository.movebank.org/handle/10255/move.1080>.
- [278] David Roshier and Wayne S.J. Boardman. Data from: Spring foraging movements of an urban population of grey-headed flying foxes (*pteropus poliocephalus*), 2021. URL <https://www.datarepository.movebank.org/handle/10255/move.1218>.
- [279] Vincent V Salomonson, WL Barnes, Peter W Maymon, Harry E Montgomery, and Harvey Ostrow. Modis: Advanced facility instrument for studies of the earth as a system. *IEEE Transactions on geoscience and remote sensing*, 27(2):145–153, 1989.
- [280] Eric W. Sanderson, Kim Fisher, Adam Duncan, Lucinda Royte, Nathaniel Robinson, and Dustin Sampson. The human footprint: 20-year time series dataset, 2020. Accessed: 2026-04-01, <https://wchumanfootprint.org/data-access>.
- [281] Eric Wayne Sanderson, Kim Fisher, Nathaniel Robinson, Dustin Sampson, Adam Duncan, and Lucinda Royte. The march of the human footprint. 2022.
- [282] John D. Scasta, Jacob D. Hennig, and Jeffrey L. Beck. Data from: Study "feral horse (*equus ferus caballus*), wyoming/colorado, usa", 2022. URL <https://www.datarepository.movebank.org/handle/10255/move.1581>.
- [283] Ulrike E Schlägel, Johannes Signer, Antje Herde, Sophie Eden, Florian Jeltsch, Jana A Eccard, and Melanie Dammhahn. Estimating interactions between individuals from concurrent animal movements. *Methods in Ecology and Evolution*, 10(8):1234–1245, 2019.
- [284] Elodie Schloesing, Alexandre Caron, Rémi Chambon, Nicolas Courbin, Morgane Labadie, Roch Nina, Frida Mouti Mbadinga, Wilfrid Ngoubili, Danficy Sandiala, N’Kaya Tobi, Mathieu Bourgarel, Hélène M. De Nys, and Julien Cappelle. Foraging and mating behaviors of *hypsingnathus monstrosus* at the bat-human interface in a central african rainforest. *Ecology and Evolution*, 13(7), 2023. ISSN 2045-7758. doi: 10.1002/ece3.10240. URL <http://dx.doi.org/10.1002/ece3.10240>.
- [285] Elodie Schloesing, Alexandre Caron, Rémi Chambon, Nicolas Courbin, Morgane Labadie, Roch Nina, Frida Mouti Mdadinga, Wilfrid Ngoubili, Danficy Sandial, N’Kaya-Tobi, Mathieu Bourgarel, Hélène M. De Nys, and Julien Cappelle. Data from: Foraging and mating behaviors of *hypsingnathus monstrosus* at the bat-human interface in central african rainforest, 2024. URL <https://datarepository.movebank.org/handle/10255/move.1634>.

- [286] Nina Schuhen, Thordis L Thorarinsdottir, and Tilmann Gneiting. Ensemble model output statistics for wind vectors. *Monthly weather review*, 140(10):3204–3219, 2012.
- [287] Sara Schweitzer, A. Lawrence Bryan, John Brzorad, and Roland Kays. Data from: Study "nc wood stork tracking", 2023. URL <https://datarepository.movebank.org/handle/10255/move.1676>.
- [288] Philipp Schwemmer and Stefan Garthe. Data from: Migrating curlews on schedule: departure and arrival patterns of a long-distance migrant depend on time and breeding location rather than on wind conditions, 2021. URL <https://www.datarepository.movebank.org/handle/10255/move.1283>.
- [289] Philipp Schwemmer, Moritz Mercker, Klaus Heinrich Vanselow, Pierrick Bocher, and Stefan Garthe. Migrating curlews on schedule: departure and arrival patterns of a long-distance migrant depend on time and breeding location rather than on wind conditions. *Movement Ecology*, 9(1), March 2021. ISSN 2051-3933. doi: 10.1186/s40462-021-00252-y. URL <http://dx.doi.org/10.1186/s40462-021-00252-y>.
- [290] Laurel E.K. Serieys and Jacqueline M. Bishop. Data from: Study "caracal movement ecology study in cape town, south africa", 2025. URL <https://datarepository.movebank.org/handle/10255/move.3458>.
- [291] Laurel E.K. Serieys and Wai-Ming Wong. Data from: Study "bobcat spatial ecology study in the olympic peninsula, washington", 2025. URL <https://datarepository.movebank.org/handle/10255/move.3452>.
- [292] Laurel E.K. Serieys, Matthew S. Rogan, Stephani S. Matsushima, and Christopher C. Wilmers. Road-crossings, vegetative cover, land use and poisons interact to influence corridor effectiveness. *Biological Conservation*, 253:108930, January 2021. ISSN 0006-3207. doi: 10.1016/j.biocon.2020.108930. URL <http://dx.doi.org/10.1016/j.biocon.2020.108930>.
- [293] Laurel E.K. Serieys, Jacqueline M. Bishop, Matthew S. Rogan, Justine A. Smith, Justin P. Suraci, M. Justin O’Riain, and Christopher C. Wilmers. Anthropogenic activities and age class mediate carnivore habitat selection in a human-dominated landscape. *iScience*, 26(7):107050, 2023. ISSN 2589-0042. doi: 10.1016/j.isci.2023.107050. URL <http://dx.doi.org/10.1016/j.isci.2023.107050>.
- [294] Laurel E.K. Serieys, Stephani S. Matsushima, and Christopher C. Wilmers. Data from: Study "bobcat habitat connectivity study in central california", 2024. URL <https://datarepository.movebank.org/handle/10255/move.3087>.
- [295] Laurel E.K. Serieys, Christopher J. Hickling, Daniel Fortin, Matthew S. Rogan, Shannon Kachel, L. Mark Elbroch, Kimberly Sager-Fradkin, Andrew Stratton, Caitlin Kubar, Axel Moehrensclager, Wai-Ming Wong, and Kathleen Carroll. Bobcats select young forests and avoid clear-cut and mature forests in a timber-logged landscape. *Forest Ecology and Management*, 588:122764, 2025. ISSN 0378-1127. doi: 10.1016/j.foreco.2025.122764. URL <http://dx.doi.org/10.1016/j.foreco.2025.122764>.
- [296] Laurel E.K. Serieys, Supawat Khaewphakdee, Wiroon Mongkonsin, Kritsana Kaewplang, Marnoch Yindee, Warong Suksavate, Chaiwat Klakhaeng, Thaksin Wongson, Ronglarp Sukmasuang, Wai-Ming Wong, and Rattapan Pattanarangsarn. Adapting to change: A wetland-specialist carnivore’s habitat selection in a human-dominated wetland. *Biological Conservation*, 310:111382, October 2025. ISSN 0006-3207. doi: 10.1016/j.biocon.2025.111382. URL <http://dx.doi.org/10.1016/j.biocon.2025.111382>.
- [297] Edy Setyawan, Abraham Sianipar, Ronald Mambrasar, and Mark Erdmann. Data from: Staying close to home: horizontal movements of satellite-tracked reef manta rays *mobula alfredi* (krefft, 1868) in the world’s largest manta sanctuary, 2025. URL <https://datarepository.movebank.org/handle/10255/move.3339>.
- [298] Edy Setyawan, Abraham B. Sianipar, Ronald Mambrasar, Muhamad Izuan, Abdy W. Hasan, Fahmi, Mujiyanto, Orgenes Ambafen, Imanuel Mofu, Mochamad Iqbal Herwata Putra, and Mark V. Erdmann. Staying close to home: Horizontal movements of satellite-tracked reef

- manta rays *mobula alfredi* (krefft, 1868) in the world's largest manta sanctuary. *Fishes*, 10(2):66, February 2025. ISSN 2410-3888. doi: 10.3390/fishes10020066. URL <http://dx.doi.org/10.3390/fishes10020066>.
- [299] Novin Shahroudi, Mihkel Lepson, and Meelis Kull. Evaluation of trajectory distribution predictions with energy score. In *Forty-first International Conference on Machine Learning*, 2024.
 - [300] Sherub Sherub and Martin Wikelski. Data from: Study "black-necked crane bhutan (uwice-mpiab)", 2024. URL <https://datarepository.movebank.org/handle/10255/move.3265>.
 - [301] Liushuai Shi, Le Wang, Chengjiang Long, Sanping Zhou, Mo Zhou, Zhenxing Niu, and Gang Hua. Sgcnn: Sparse graph convolution network for pedestrian trajectory prediction. In *Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition (CVPR)*, pages 8994–9003, 2021.
 - [302] Liushuai Shi, Le Wang, Chengjiang Long, Sanping Zhou, Mo Zhou, Zhenxing Niu, and Gang Hua. Trajectory unified transformer for pedestrian trajectory prediction. In *Proceedings of the IEEE/CVF International Conference on Computer Vision (ICCV)*, pages 20660–20669, 2023.
 - [303] David A Siegel, Timothy DeVries, Ivona Cetinić, and Kelsey M Bisson. Quantifying the ocean's biological pump and its carbon cycle impacts on global scales. *Annual review of marine science*, 15(1):329–356, 2023.
 - [304] J Signer, J Fieberg, and T Avgar. Estimating utilization distributions from fitted step-selection functions. *ecosphere*. 2017; 8: e01771, 1771.
 - [305] Johannes Signer, John Fieberg, and Tal Avgar. Animal movement tools (amt): R package for managing tracking data and conducting habitat selection analyses. *Ecology and Evolution*, 9(2):880–890, January 2019. ISSN 2045-7758. doi: 10.1002/ece3.4823. URL <http://dx.doi.org/10.1002/ece3.4823>.
 - [306] Johannes Signer, John Fieberg, Björn Reineking, Ulrike Schlägel, B Smith, Niko Balkenhol, and Tal Avgar. Simulating animal space use from fitted integrated step-selection functions (issf). *Methods in Ecology and Evolution*, 15(1):43–50, 2024.
 - [307] Navinder J Singh and Eleanor J Milner-Gulland. Conserving a moving target: planning protection for a migratory species as its distribution changes. *Journal of Applied Ecology*, 48(1):35–46, 2011.
 - [308] Jan Škrábal, Šimon Krejčí, Rainer Raab, Esther Sebastián-González, and Ivan Literák. Soaring over open waters: horizontal winds provide lift to soaring migrants in weak thermal conditions. *Movement Ecology*, 11(1):76, 2023.
 - [309] Rob Slotow, Maria Thaker, and Abi Tamim Vanak. Data from: Fine-scale tracking of ambient temperature and movement reveals shuttling behavior of elephants to water, 2019. URL <https://www.datarepository.movebank.org/handle/10255/move.980>.
 - [310] Leonard A Smith, Emma B Suckling, Erica L Thompson, Trevor Maynard, and Hailiang Du. Towards improving the framework for probabilistic forecast evaluation. *Climatic Change*, 132(1):31–45, 2015.
 - [311] Marjorie C. Sorensen, Thomas Mueller, Isabel Donoso, Valentin Graf, Dominik Merges, Marco Vanoni, Wolfgang Fiedler, and Eike Lena Neuschulz. Scatter-hoarding birds disperse seeds to sites unfavorable for plant regeneration. *Movement Ecology*, 10(1), 2022. ISSN 2051-3933. doi: 10.1186/s40462-022-00338-1. URL <http://dx.doi.org/10.1186/s40462-022-00338-1>.
 - [312] Orr Spiegel, Wayne M. Getz, and Ran Nathan. Factors influencing foraging search efficiency: Why do scarce lappet-faced vultures outperform ubiquitous white-backed vultures? *The American Naturalist*, 181(5):E102–E115, May 2013. ISSN 1537-5323. doi: 10.1086/670009. URL <http://dx.doi.org/10.1086/670009>.

- [313] Orr Spiegel, Roi Harel, Alejandro Centeno-Cuadros, Ohad Hatzofe, Wayne M. Getz, and Ran Nathan. Moving beyond curve fitting: Using complementary data to assess alternative explanations for long movements of three vulture species. *The American Naturalist*, 185(2): E44–E54, February 2015. ISSN 1537-5323. doi: 10.1086/679314. URL <http://dx.doi.org/10.1086/679314>.
- [314] Orr M. Spiegel, Wayne M. Getz, and Ran Nathan. Data from: Factors influencing foraging search efficiency: Why do scarce lappet-faced vultures outperform ubiquitous white-backed vultures?, 2014. URL <https://www.datarepository.movebank.org/handle/10255/move.392>.
- [315] Orr M. Spiegel, Roi Harel, Alejandro Centeno-Cuadros, Ohad Hatzofe, Wayne M. Getz, and Ran Nathan. Data from: Moving beyond curve-fitting: using complementary data to assess alternative explanations for long movements of three vulture species, 2015. URL <https://www.datarepository.movebank.org/handle/10255/move.389>.
- [316] Jared A. Stabach, Lacey F. Hughey, Robin S. Reid, Jeffrey S. Worden, Peter Leimgruber, and Randall B. Boone. Data from: Comparison of movement strategies of three populations of white-bearded wildebeest, 2020. URL <https://www.datarepository.movebank.org/handle/10255/move.1095>.
- [317] Jared A. Stabach, Lacey F. Hughey, Ramiro D. Crego, Christen H. Fleming, J. Grant C. Hopcraft, Peter Leimgruber, Thomas A. Morrison, Joseph O. Ogutu, Robin S. Reid, Jeffrey S. Worden, and Randall B. Boone. Increasing anthropogenic disturbance restricts wildebeest movement across east african grazing systems. *Frontiers in Ecology and Evolution*, 10, March 2022. ISSN 2296-701X. doi: 10.3389/fevo.2022.846171. URL <http://dx.doi.org/10.3389/fevo.2022.846171>.
- [318] Arnaud Tarroux, Henri Weimerskirch, Sheng-Hung Wang, David H. Bromwich, Yves Cherel, Akiko Kato, Yan Ropert-Coudert, Øystein Varpe, Nigel G. Yoccoz, and Sébastien Descamps. Data from: Flexible flight response to challenging wind conditions in a commuting antarctic seabird: do you catch the drift?, 2016. URL <https://www.datarepository.movebank.org/handle/10255/move.498>.
- [319] Arnaud Tarroux, Henri Weimerskirch, Sheng-Hung Wang, David H. Bromwich, Yves Cherel, Akiko Kato, Yan Ropert-Coudert, Øystein Varpe, Nigel G. Yoccoz, and Sébastien Descamps. Flexible flight response to challenging wind conditions in a commuting antarctic seabird: do you catch the drift? *Animal Behaviour*, 113:99–112, March 2016. ISSN 0003-3472. doi: 10.1016/j.anbehav.2015.12.021. URL <http://dx.doi.org/10.1016/j.anbehav.2015.12.021>.
- [320] Maria Thaker, Pratik R. Gupte, Herbert H. T. Prins, Rob Slotow, and Abi T. Vanak. Fine-scale tracking of ambient temperature and movement reveals shuttling behavior of elephants to water. *Frontiers in Ecology and Evolution*, 7, January 2019. ISSN 2296-701X. doi: 10.3389/fevo.2019.00004. URL <http://dx.doi.org/10.3389/fevo.2019.00004>.
- [321] Henrik Thurfjell, Simone Ciuti, and Mark S. Boyce. Applications of step-selection functions in ecology and conservation. *Movement Ecology*, 2:4, 2014. doi: 10.1186/2051-3933-2-4.
- [322] Christopher M. Todd, David A. Westcott, John M. Martin, Karrie Rose, Adam McKeown, Jane Hall, and Justin A. Welbergen. Data from: Study "movements of the christmas island flying fox, australia", 2022. URL <https://www.datarepository.movebank.org/handle/10255/move.1405>.
- [323] Christopher M. Todd, David A. Westcott, John M. Martin, Karrie Rose, Adam McKeown, Jane Hall, and Justin A. Welbergen. Body-size dependent foraging strategies in the christmas island flying-fox: implications for seed and pollen dispersal within a threatened island ecosystem. *Movement Ecology*, 10(1), April 2022. ISSN 2051-3933. doi: 10.1186/s40462-022-00315-8. URL <http://dx.doi.org/10.1186/s40462-022-00315-8>.
- [324] Pyry Toivonen, Mikko Toivola, and Vesa Selonen. Data from: Ice cover and partner removal increase movements of invasive mesopredator in the coastal island mosaic, 2025. URL <https://datarepository.movebank.org/handle/10255/move.3457>.

- [325] Pyry Toivonen, Mikko Toivola, and Vesa Selonen. Ice cover and partner removal increase movements of invasive mesopredator in the coastal island mosaic. *Ecological Solutions and Evidence*, 6(3), 2025. ISSN 2688-8319. doi: 10.1002/2688-8319.70068. URL <http://dx.doi.org/10.1002/2688-8319.70068>.
- [326] Hristos Tyralis and Georgia Papacharalampous. A review of predictive uncertainty estimation with machine learning. *Artificial Intelligence Review*, 57(4):94, 2024.
- [327] U.S. Geological Survey. Landsat collection 2 level-2 surface reflectance, 2020. 30 m surface reflectance data used to derive 8-day NDVI, EVI, NDSI, and BAI composites; accessed 2023-03-07.
- [328] Denis Valle, Nina Attias, Joshua A Cullen, Mevin B Hooten, Aline Giroux, Luiz Gustavo R Oliveira-Santos, Arnaud LJ Desbiez, and Robert J Fletcher Jr. Bridging the gap between movement data and connectivity analysis using the time-explicit habitat selection (tehs) model. *Movement Ecology*, 12(1):19, 2024.
- [329] Henk Van Der Jeugd, Kees Osterbeek, Bruno J Ens, Judy Shamoun-Baranes, and Klaus-Michael Exo. Data from: Forecasting spring from afar? timing of migration and predictability of phenology along different migration routes of an avian herbivore [barents sea data], 2014. URL <https://www.datarepository.movebank.org/handle/10255/move.383>.
- [330] Mariëlle L. van Toor, Anders Hedenström, Jonas Waldenström, Wolfgang Fiedler, Richard A. Holland, Kasper Thorup, and Martin Wikelski. Flexibility of continental navigation and migration in european mallards. *PLoS ONE*, 8(8):e72629, August 2013. ISSN 1932-6203. doi: 10.1371/journal.pone.0072629. URL <http://dx.doi.org/10.1371/journal.pone.0072629>.
- [331] Mariëlle L. van Toor, Alexis Avril, Guohui Wu, Scott H. Holan, and Jonas Waldenström. As the duck flies-estimating the dispersal of low-pathogenic avian influenza viruses by migrating mallards. *Frontiers in Ecology and Evolution*, 6, December 2018. ISSN 2296-701X. doi: 10.3389/fevo.2018.00208. URL <http://dx.doi.org/10.3389/fevo.2018.00208>.
- [332] Mariëlle L. Van Toor, Ulf Ottosson, Tim Van Der Meer, Sita Van Hoorn, and Jonas Waldenström. Data from: As the duck flies: estimating the dispersal of low-pathogenic avian influenza viruses by migrating mallards, 2018. URL <https://www.datarepository.movebank.org/handle/10255/move.800>.
- [333] Mariëlle L. Van Toor, Sergey Kharitonov, Saulius Švažas, Mindaugas Dagys, Eric Kleyheeg, Gerard Müskens, Ulf Ottosson, Ramunas Žydelis, and Jonas Waldenström. Data from: Study "eurasian wigeon (mareca penelope) netherlands lithuania 2018-2019", 2021. URL <https://www.datarepository.movebank.org/handle/10255/move.1399>.
- [334] Mariëlle L. van Toor, Sergey Kharitonov, Saulius Švažas, Mindaugas Dagys, Erik Kleyheeg, Gerard Müskens, Ulf Ottosson, Ramunas Žydelis, and Jonas Waldenström. Migration distance affects how closely eurasian wigeons follow spring phenology during migration. *Movement Ecology*, 9(1), December 2021. ISSN 2051-3933. doi: 10.1186/s40462-021-00296-0. URL <http://dx.doi.org/10.1186/s40462-021-00296-0>.
- [335] Ashish Vaswani, Noam Shazeer, Niki Parmar, Jakob Uszkoreit, Llion Jones, Aidan N Gomez, Łukasz Kaiser, and Illia Polosukhin. Attention is all you need. *Advances in neural information processing systems*, 30, 2017.
- [336] Tana Verzuh, Martha Torstenson, Yun Tao, John Fryxell, Christian Rutz, and Roxanne S Beltran. Aligning tools and terminology to integrate movement ecology with conservation science. *Conservation Biology*, page e70209, 2026.
- [337] Kartik Waghmare and Johanna Ziegel. Proper scoring rules for estimation and forecast evaluation. *Annual Review of Statistics and Its Application*, 13, 2025.
- [338] Hermann Wagner, Sebastian Cruz, Gustavo Jiménez-Uzcategui, Katherine Albán, Galo Quezada, and Paolo Piedrahita. Data from: Abundance, diet and foraging of galápagos barn owls (tyto furcata punctatissima), 2025. URL <https://datarepository.movebank.org/handle/10255/move.3575>.

- [339] Hermann Wagner, Sebastian Cruz, Gustavo Jiménez-Uzcátegui, Katherine Albán, Galo Quezada, and Paolo Piedrahita. Abundance, diet and foraging of galápagos barn owls (*tyto furcata punctatissima*). *Animals*, 15(15):2283, August 2025. ISSN 2076-2615. doi: 10.3390/ani15152283. URL <http://dx.doi.org/10.3390/ani15152283>.
- [340] Jake Wall, George Wittemyer, Valerie LeMay, Iain Douglas-Hamilton, and Brian Klinkenberg. Data from: Elliptical time-density model to estimate wildlife utilization distributions, 2014. URL <https://www.datarepository.movebank.org/handle/10255/move.372>.
- [341] Jake Wall, George Wittemyer, Valerie LeMay, Iain Douglas-Hamilton, and Brian Klinkenberg. Elliptical time-density model to estimate wildlife utilization distributions. *Methods in Ecology and Evolution*, 5(8):780–790, August 2014. ISSN 2041-210X. doi: 10.1111/2041-210x.12218. URL <http://dx.doi.org/10.1111/2041-210x.12218>.
- [342] Jiankai Wang, Meng Meng, Longkai Guo, Jiayi Xu, Lucio Dai, Yu Jiang, and Ming Yang. Ltp: Lane-based trajectory prediction for autonomous driving. In *Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition (CVPR)*, 2022.
- [343] Chloé Warret Rodrigues and James D. Roth. Data from: Coexistence of two sympatric predators in a transitional ecosystem under constraining environmental conditions: a perspective from space and habitat use, 2023. URL <https://datarepository.movebank.org/handle/10255/move.1652>.
- [344] Chloé Warret Rodrigues and James D. Roth. Coexistence of two sympatric predators in a transitional ecosystem under constraining environmental conditions: a perspective from space and habitat use. *Movement Ecology*, 11(1), October 2023. ISSN 2051-3933. doi: 10.1186/s40462-023-00421-1. URL <http://dx.doi.org/10.1186/s40462-023-00421-1>.
- [345] V. Warwick-Evans, P. W. Atkinson, J. P. Y. Arnould, R. Gauthier, L. Soanes, L. A. Robinson, and J. A. Green. Changes in behaviour drive inter-annual variability in the at-sea distribution of northern gannets. *Marine Biology*, 163(7), 2016. ISSN 1432-1793. doi: 10.1007/s00227-016-2922-y. URL <http://dx.doi.org/10.1007/s00227-016-2922-y>.
- [346] Victoria Warwick-Evans, Louise M. Soanes, Roland D. Gauthier, Philip W. Atkinson, John P.Y. Arnould, and Jon A. Green. Data from: Changes in behaviour drive inter-annual variability in the at-sea distribution of northern gannets, 2017. URL <https://www.datarepository.movebank.org/handle/10255/move.697>.
- [347] Tonglong Wei, Yan Lin, Youfang Lin, Shengnan Guo, Jilin Hu, Haitao Yuan, Gao Cong, and Huaiyu Wan. Plmtrajrec: A scalable and generalizable trajectory recovery method with pre-trained language models. *arXiv preprint arXiv:2410.14281*, 2024.
- [348] Dylan M Westaway, Dale G Nimmo, Scott W Forrest, and Mitchell A Cowan. Navigating fragmentation: movement ecology and habitat selection of a large, long-lived skink in an agricultural landscape. *Landscape ecology*, 40(12):232, 2025. doi: 10.1007/s10980-025-02253-3.
- [349] Sarah Wicks, Christopher Beirne, Cristina Azzopardi Schellmann, Eleanor Flatt, Sandy Quirós Beita, Rigoberto Pereira Rocha, and Andrew Whitworth. Back to the wild: Post-translocation gps monitoring of a rehabilitated ocelot (*leopardus pardalis*) in a forest-agriculture matrix in the osa peninsula, costa rica. *Neotropical Biology and Conservation*, 19(3): 379–392, August 2024. ISSN 2236-3777. doi: 10.3897/neotropical.19.e124324. URL <http://dx.doi.org/10.3897/neotropical.19.e124324>.
- [350] Sarah Wicks, Beirne Christopher, Cristina Azzopardi Schellmann, Eleanor Flatt, Sandy Quirós Beita, Rigoberto Pereira Rocha, and Andrew Whitworth. Data from: Back to the wild: Post-translocation gps monitoring of a rehabilitated ocelot (*leopardus pardalis*) in a high competition, forest-agriculture matrix, 2024. URL <https://datarepository.movebank.org/handle/10255/move.3195>.
- [351] Dhanushi A Wijeyakulasuriya, Elizabeth W Eisenhauer, Benjamin A Shaby, and Ephraim M Hanks. Machine learning for modeling animal movement. *PloS one*, 15(7):e0235750, 2020.

- [352] Wildlife Conservation Society (WCS) and Center for International Earth Science Information Network (CIESIN), Columbia University. Last of the wild project, version 2 (2005): Global human footprint dataset, 2005. Accessed: 2026-04-01.
- [353] Darrel L Williams, Samuel Goward, and Terry Arvidson. Landsat. *Photogrammetric Engineering & Remote Sensing*, 72(10):1171–1178, 2006.
- [354] Kenady Wilson, Ephraim Hanks, and Devin Johnson. Estimating animal utilization densities using continuous-time markov chain models. *Methods in Ecology and Evolution*, 9(5):1232–1240, 2018.
- [355] Scott W. Yanco, Ruth Y. Oliver, Fabiola Iannarilli, Ben S. Carlson, Georg Heine, Uschi Mueller, Nina Richter, Bernd Vorneweg, Yuriy Andryushchenko, Nyambayar Batbayar, Mindaugas Dagys, Mark Desholm, Batbayar Galtbalt, Andrey E. Gavrilov, Oleg A. Goroshko, Elena I. Ilyashenko, Valentin Yu Ilyashenko, Johan Månsson, Elena A. Mudrik, Tseveenmyadag Natsagdorj, Lovisa Nilsson, Sherub Sherub, Henrik Skov, Tuvshintugs Sukhbaatar, Ramunas Zydelis, Martin Wikelski, Walter Jetz, and Ivan Pokrovsky. Migratory birds modulate niche tradeoffs in rhythm with seasons and life history. *Proceedings of the National Academy of Sciences*, 121(41), 2024. ISSN 1091-6490. doi: 10.1073/pnas.2316827121. URL <http://dx.doi.org/10.1073/pnas.2316827121>.
- [356] Dingqi Yang, Daqing Zhang, and Bingqing Qu. Participatory cultural mapping based on collective behavior data in location-based social networks. *ACM Transactions on Intelligent Systems and Technology (TIST)*, 7(3):1–23, 2016.
- [357] Jing Yuan, Yu Zheng, Chengyang Zhang, Wenlei Xie, Xing Xie, Guangzhong Sun, and Yan Huang. T-drive: driving directions based on taxi trajectories. In *Proceedings of the 18th SIGSPATIAL International conference on advances in geographic information systems*, pages 99–108, 2010.
- [358] Yu Zheng, Xing Xie, Wei-Ying Ma, et al. Geolife: A collaborative social networking service among user, location and trajectory. *IEEE Data Eng. Bull.*, 33(2):32–39, 2010.
- [359] Yuanshao Zhu, James Jianqiao Yu, Xiangyu Zhao, Xun Zhou, Liang Han, Xuetao Wei, and Yuxuan Liang. Unitraj: Learning a universal trajectory foundation model from billion-scale worldwide traces. *arXiv preprint arXiv:2411.03859*, 2024.
- [360] Kristín Ágústsdóttir and Skarphéðinn G. Þórisson. Data from: Study "reindeer movement in east iceland", 2022. URL <https://www.datarepository.movebank.org/handle/10255/move.1425>.
- [361] J Škrábal, I Literák, and R Raab. Data from: Study "milvus milvus soaring over adriatic sea", 2023. URL <http://dx.doi.org/10.5441/001/1.300>.
- [362] Ramunas Žydelis, Mark Desholm, Johan Månsson, Lovisa Nilsson, and Henrik Skov. Data from: Study "gps telemetry of common cranes, sweden", 2024. URL <https://datarepository.movebank.org/handle/10255/move.3261>.

Appendix

A Additional related work	39
B Trajectory datasets	40
B.1 Full list of included studies	40
B.2 Trajectory curation and filtering details	44
B.3 Additional details on train/test split	45
B.4 Data modification records	45
B.5 MOVEBENCH-Mini	45
C Environmental covariates	45
C.1 Full list of included covariates	45
C.2 Details on raster preparation	49
D Metrics	52
E Method details	52
E.1 Implementation	53
F Hyperparameters	53
G Additional experiment details	55
G.1 All experimental results	55
G.2 Per-study MoveFormer	71
G.3 Compute and storage requirements	71

A Additional related work

Movement data. We focus on trajectory data, sequences of location-time tuples that record the movement of some object of interest over time. The most common trajectory datasets in machine learning are GPS “mobility traces” of vehicles and pedestrians, typically collected from onboard transmitters [203, 242, 359], or social media check-ins [55, 201, 356, 358]. Locations are often projected onto a known transportation network/map, reducing the dimensionality of possible movement [238, 342].

Animal movement data is difficult and costly to collect, clean, and prepare for analysis. Tag deployment often takes place in remote locations, may require veterinary expertise, and typically requires government permits and approved ethical protocols. Significant data cleaning is often required to deal with problems such as irregular or inaccurate data collection due to limited GPS reception [132, 195]. Further, animal trajectory datasets are often analyzed together with environmental covariates such as land cover, elevation, temperature, precipitation, or oceanographic variables, which must be derived and aligned from GIS layers, remote sensing products, or weather and climate reanalyses [241, 321].

Analyzing and modeling movement. Advancements in machine-learning-based trajectory analysis have paralleled broader developments in sequence modeling. Early methods drew on statistical state-space and hidden Markov models, later adopting recurrent neural networks and LSTMs for direct sequence prediction [11, 196]. More recent approaches use graph neural networks and transformer architectures to improve predictive performance on tasks such as next-step forecasting, multi-step trajectory prediction, and behavior classification [111, 301, 302, 347, 359].

In contrast, recent methods for wildlife movement have modeled the movement process as a function of *habitat selection*, whereby an individual chooses where to go next based in part on the environmental conditions at candidate future locations. Such methods are called *step-selection functions* [13, 107], and are most frequently implemented as a binary classification task where the objective is to differentiate between locations where an animal was actually observed and other nearby locations where it was not, based on environmental conditions. A range of extensions have been proposed for step-selection functions to model additional movement processes, including change-points and behavioral states [15, 181, 228, 239, 258], individual differences or interactions between individuals [54, 180, 227, 264, 283], irregular sampling [87, 152, 228], and variation across time and space [85, 104, 180, 328]. Newer methods have also proposed moving beyond linear models toward deep learning architectures [61, 105, 110]. Recently, step-selection functions have been used to predict future movement and space use [67, 150, 263, 274, 304, 306, 348], however analysis of their performance remains mostly qualitative and limited to a small number of case studies. Beyond step selection, there have been a small number of papers focused on forecasting movement for single studies, including southern elephant seals [213], sika deer [176], and migratory gulls [351], though none have been assessed more broadly and compared across species.

Evaluating predicted trajectories. Trajectory forecasting is commonly evaluated with point-prediction metrics such as the mean L2 distance between predicted points and their associated ground truth observation [251, 301, 302]. However, when future movement is inherently stochastic, point prediction metrics are inappropriate and can reward models that hedge toward the mean or some otherwise strategically distorted forecast [123, 124, 125, 310]. For this reason, probabilistic predictions are commonly evaluated with *proper scoring rules*, which assess the quality of an entire predictive distribution rather than only a single point forecast [124]. For continuous outcomes, standard examples include the continuous ranked probability score (CRPS) [146].

Forecast evaluation in animal movement remains limited, often relying on qualitative inspection [306] or log loss [61, 105]. While log loss is indeed a proper scoring rule, it requires a likelihood and therefore cannot be used across arbitrary model classes, including deterministic baselines. We therefore utilize the energy score [124], a multivariate sample-based generalization of CRPS, as a probabilistic forecasting metric for wildlife movement data in MOVEBENCH.

Movement benchmarks. We provide a summary of existing trajectory benchmarks in Tab. 1. Human and vehicle trajectory datasets including GeoLife [358], Porto [242], T-Drive [357], and WorldTrace [359] have large numbers of GPS points, however apart from T-Drive they are quite short, spanning small distances and temporal extents. All human and vehicle trajectory datasets evaluate deterministic map-aligned predictions of future movement. The two existing benchmarks for animal movement data, BEBE [149] and Noonan et al. [240], also do not support probabilistic forecasting. BEBE [149] focuses on behavior classification using animal-borne accelerometer tags and does not

include GPS data, while Noonan et al. [240] anonymizes both GPS locations and timestamps. Neither includes environmental covariates. To our knowledge, MOVEBENCH is the first benchmark that supports standardized evaluation of probabilistic animal movement forecasting with open-access GPS trajectories and environmental covariates.

B Trajectory datasets

B.1 Full list of included studies

Table 3: MoveBench studies included in the analysis.

Index	Study species	Authors	Data	License	N Indiv	GPS fixes	Median fix interval (min)
1	<i>Mareca penelope</i>	van Toor et al. [334]	Van Toor et al. [333]	CC0	6	42,660	61
2	<i>Anas platyrhynchos</i>	Kleyheeg et al. [182]	Kleyheeg et al. [183]	CC0	6	3,740	60
3	<i>Anas platyrhynchos</i>	van Toor et al. [330]	Matthes et al. [206]	CC0	6	3,987	360
4	<i>Anas platyrhynchos</i>	van Toor et al. [331]	Van Toor et al. [332]	CC0	6	13,785	61
5	<i>Anser albifrons</i>	Buitendijk et al. [42]	Buitendijk et al. [43]	CC0	6	22,337	60
6	<i>Anser albifrons</i>	Kölzsch et al. [184] Kölzsch et al. [185]	Kölzsch et al. [187] Kölzsch et al. [188]	CC0	6	36,919	60
7	<i>Anser albifrons</i>	Kölzsch et al. [184]	Müskens et al. [230]	CC0	6	13,297	60
8	<i>Anser anser</i>	Buitendijk et al. [42]	Buitendijk et al. [43]	CC0	6	53,183	60
9	<i>Anser brachyrhynchus</i>	Kölzsch et al. [189]	Kölzsch et al. [190]	CC0	6	9,149	60
10	<i>Anser fabalis</i>	Piironen et al. [255]	Piironen et al. [254]	CC0	6	35,979	60
11	<i>Anthropoides virgo</i>	Yanco et al. [355]	Batbayar et al. [24]	CC-BY-NC-4.0	2	12,826	67
12	<i>Anthropoides virgo</i>	Yanco et al. [355]	Ilyashenko et al. [158]	CC-BY-4.0	1	374	1439
13	<i>Aquila nipalensis</i>	Efrat et al. [82]	Efrat et al. [81]	CC0	6	60,623	60
14	<i>Ardea alba</i>	Jennings et al. [169]	Jennings et al. [170]	CC0	6	44,661	60
15	<i>Ardea purpurea</i>	Morganti et al. [223]	Morganti et al. [222]	CC0	1	1,969	60
16	<i>Branta leucopsis</i>	Buitendijk et al. [42]	Buitendijk et al. [43]	CC0	6	29,165	60
17	<i>Branta leucopsis</i>	Kölzsch et al. [186]	Cabot [47]	CC0	6	5,829	120
18	<i>Branta leucopsis</i>	Kölzsch et al. [186]	Griffin [129]	CC0	6	7,509	120
19	<i>Branta leucopsis</i>	Kölzsch et al. [189]	Kölzsch et al. [190]	CC0	6	6,299	60
20	<i>Branta leucopsis</i>	Kölzsch et al. [186]	Van Der Jeugd et al. [329]	CC0	6	12,134	180
21	<i>Buceros bicornis</i>	Naniwadekar et al. [232]	Naniwadekar et al. [231]	CC0	5	3,953	60
22	<i>Buteo lagopus</i>	Pokrovsky et al. [260]	Pokrovsky et al. [259]	CC0	5	66,221	60
23	<i>Buteo rufinus</i>	Friedemann et al. [109]	Friedemann et al. [108]	CC0	6	16,906	60
24	<i>Buteo swainsoni</i>	Fleishman et al. [102]	Fleishman et al. [103]	CC0	6	7,305	60
25	<i>Cathartes aura</i>	Mallon et al. [204]	Bildstein et al. [31]	CC-BY-4.0	11	131,764	60
26	<i>Cathartes aura</i>	Dodge et al. [75]	Bildstein et al. [30]	CC0	5	22,506	180
27	<i>Cathartes melambrotus</i>	Beirne et al. [27]	Christopher et al. [58]	CC0	4	23,145	60
28	<i>Ciconia ciconia</i>	Acácio et al. [3]	Acácio et al. [4]	CC0	5	3,136	61
29	<i>Ciconia ciconia</i>	Aikens et al. [8]	Fiedler et al. [98] Fiedler et al. [97] Fiedler et al. [96]	CC0	6	75,185	60

Continued on next page

Table 3: MoveBench studies included in the analysis (continued).

Index	Study species	Authors	Data	License	N Indiv	GPS fixes	Median fix interval (min)
30	<i>Circaetus gallicus</i>	Friedemann et al. [109]	Friedemann et al. [108]	CC0	6	26,302	60
31	<i>Coragyps atratus</i>	Mallon et al. [204]	Bildstein et al. [31]	CC-BY-4.0	6	33,372	69
32	<i>Coragyps atratus</i>	Holland et al. [154]	Holland and Byrne [153]	CC0	2	10,480	60
33	<i>Eudocimus albus</i>	Kidd-Weaver et al. [179]	Kidd-Weaver et al. [178]	CC0	6	28,835	120
34	<i>Falco naumanni</i>	Hernández-Pliego et al. [145]	Hernández-Pliego et al. [144]	CC0	6	1,967	60
35	<i>Grus grus</i>	Yanco et al. [355]	Ilyashenko et al. [159]	CC-BY-4.0	6	3,960	1434
36	<i>Grus grus</i>	Yanco et al. [355]	Žydelis et al. [362]	CC-BY-4.0	6	75,239	63
37	<i>Grus nigricollis</i>	Yanco et al. [355]	Sherub and Wikelski [300]	CC-BY-4.0	4	22,830	60
38	<i>Grus vipio</i>	Yanco et al. [355]	Batbayar et al. [25]	CC-BY-NC-4.0	3	31,731	61
39	<i>Gyps africanus</i>	Spiegel et al. [312]	Spiegel et al. [314]	CC0	6	24,953	60
40	<i>Gyps africanus</i>	Spiegel et al. [313]	Spiegel et al. [315]	CC0	2	9,858	60
41	<i>Gyps fulvus</i>	Harel and Nathan [136]	Harel and Nathan [135]	CC0	4	13,239	60
42	<i>Hydroprogne caspia</i>	Byholm et al. [44]	Byholm et al. [45]	CC0	5	27,655	60
43	<i>Larus argentatus</i>	Anderson et al. [12]	Fifield et al. [99]	CC0	6	30,519	62
44	<i>Larus argentatus</i>	Anderson et al. [12]	Ronconi and Taylor [277]	CC0	6	30,732	120
45	<i>Larus argentatus</i>	Anderson et al. [12]	Ronconi and Shlepr [276]	CC0	6	28,174	63
46	<i>Larus canus</i>	Garthe et al. [115]	Garthe et al. [114]	CC0	6	2,452	60
47	<i>Larus fuscus</i>	Garthe et al. [113]	Garthe et al. [112]	CC0	6	1,713	59
48	<i>Larus marinus</i>	Borrmann et al. [37]	Borrmann et al. [36]	CC0	6	10,613	60
49	<i>Larus smithsonianus</i>	Gutowsky et al. [133]	Mallory et al. [205]	CC0	6	52,359	60
50	<i>Milvus milvus</i>	Škrábal et al. [308]	Škrábal et al. [361]	CC0	6	2,329	120
51	<i>Morus bassanus</i>	Jeglinski et al. [166]	Jeglinski et al. [167]	CC-BY-NC-4.0	6	8,970	60
52	<i>Morus bassanus</i>	Warwick-Evans et al. [345]	Warwick-Evans et al. [346]	CC0	6	2,150	60
53	<i>Morus bassanus</i>	d'Entremont et al. [72]	D'Entremont et al. [73]	CC0	6	3,384	61
54	<i>Mycteria americana</i>		Schweitzer et al. [287]	CC0	1	13,481	60
55	<i>Nucifraga cyathactes</i>	Sorensen et al. [311]	Graf et al. [127]	CC0	6	1,488	60
56	<i>Numenius arquata</i>	Schwemmer et al. [289]	Schwemmer and Garthe [288]	CC0	6	31,290	60
57	<i>Pelecanus occidentalis</i>	Geary et al. [119]	Geary et al. [118]	CC0	6	5,558	60
58	<i>Pelecanus occidentalis</i>	Lamb et al. [192]	Lamb et al. [193]	CC0	6	27,725	90
59	<i>Pernis apivorus</i>	Mirski et al. [218]	Byholm et al. [46]	CC-BY-NC-4.0	5	27,577	120
60	<i>Phoebastria irrorata</i>	Dodge et al. [74]	Cruz et al. [70]	CC0	10	11,564	90
61	<i>Puffinus puffinus</i>	Padget et al. [244]	Padget et al. [243]	CC0	6	3,486	60
62	<i>Puffinus yelkouan</i>	Gatt et al. [116]	Lago et al. [191]	CC0	6	2,572	60
63	<i>Puffinus yelkouan</i>	Pezzo et al. [253]	Pezzo et al. [252]	CC0	2	667	60

Continued on next page

Table 3: MoveBench studies included in the analysis (continued).

Index	Study species	Authors	Data	License	N Indiv	GPS fixes	Median fix interval (min)
64	<i>Rhyticeros undulatus</i>	Naniwadekar et al. [232]	Naniwadekar et al. [231]	CC0	1	1,266	60
65	<i>Sarcoramphus papa</i>	Beirne et al. [27]	Christopher et al. [58]	CC0	3	20,434	60
66	<i>Thalassoica antarctica</i>	Tarroux et al. [319]	Tarroux et al. [318]	CC0	6	2,274	60
67	<i>Tockus deckeni</i>	Mertes et al. [214]	Mertes et al. [215]	CC0	6	68,753	64
68	<i>Torgos tracheliotus</i>	Spiegel et al. [312]	Spiegel et al. [314]	CC0	3	13,691	60
69	<i>Tyto furcata</i>	Wagner et al. [339]	Wagner et al. [338]	CC0	1	231	60
70	<i>Uria lomvia</i>	Eby et al. [80]	Eby et al. [79]	CC0	4	967	60
71	<i>Acinonyx jubatus</i>	Power et al. [267]	Power and Dell [265]	CC0	1	1,182	360
72	<i>Antilocapra americana</i>	Aikens et al. [10]	Aikens et al. [6]	CC-BY-4.0	6	73,328	120
73	<i>Antilocapra americana</i>	Aikens et al. [9]	Aikens et al. [5]	CC0	6	50,729	120
74	<i>Antilocapra americana</i>	Hennig et al. [143]	Beck et al. [26]	CC0	6	25,845	240
75	<i>Ateles geoffroyi</i>	Kays et al. [174]	Kays et al. [175]	CC0	6	8,191	60
76	<i>Brachylagus idahoensis</i>	McMahon et al. [210]	McMahon et al. [209]	CC0	4	964	60
77	<i>Bubalus bubalis</i>	Forrest et al. [104]	Forrest [106]	CC-BY-4.0	6	53,931	60
78	<i>Canis latrans</i>	Prugh et al. [270]	Prugh [269]	CC0	12	42,045	240
79	<i>Canis lupus</i>	Hebblewhite and Merrill [141]	Hebblewhite [140]	CC-BY-4.0	6	28,362	120
80	<i>Canis lupus</i>	Latham and Boutin [199]	Latham and Boutin [200]	CC0	6	8,615	120
81	<i>Canis mesomelas</i>	Borchering et al. [35]	Bellan and Getz [28]	CC0	6	60,475	60
82	<i>Capra hircus</i>	Chynoweth et al. [60]	Chynoweth [59]	CC0	6	16,101	120
83	<i>Capreolus capreolus</i>	–	EURODEER, Bondone	Not specified	6	11,103	240
84	<i>Caracal caracal</i>	Serieys et al. [293]	Serieys and Bishop [290]	CC0	6	11,743	180
85	<i>Cebus capucinus</i>	Kays et al. [174]	Kays et al. [175]	CC0	6	6,091	60
86	<i>Cervus elaphus</i>	Signer et al. [305]	–	Not specified	1	826	360
87	<i>Cervus elaphus</i>	Ciuti et al. [62]	Boyce and Ciuti [38]	CC-BY-4.0	6	66,401	120
88	<i>Cervus elaphus</i>	Eggeman et al. [86]	Hebblewhite et al. [142]	CC0	6	88,391	120
89	<i>Cervus elaphus</i>	–	EURODEER, Alta Val di Non	Not specified	6	51,666	60
90	<i>Cervus elaphus</i>	–	EURODEER, Val Rendena	Not specified	6	17,075	180
91	<i>Connochaetes taurinus</i>	Stabach et al. [317]	Stabach et al. [316]	CC0	15	177,521	60
92	<i>Crocota crocuta</i>	Cozzi et al. [68]	–	CC0	6	19,996	120
93	<i>Equus burchellii</i>	Bartlam-Brooks et al. [16]	Bartlam-Brooks and Harris [17]	CC0	6	36,766	60
94	<i>Equus ferus</i>	Hennig et al. [143]	Scasta et al. [282]	CC0	6	25,628	240
95	<i>Gulo gulo</i>	Glass et al. [122]	Glass and Robards [121]	CC0	3	6,584	80
96	<i>Hypsignathus monstrosus</i>	Schloesing et al. [284]	Schloesing et al. [285]	CC0	6	1,426	60
97	<i>Leopardus pardalis</i>	Wicks et al. [349]	Wicks et al. [350]	CC0	1	695	300

Continued on next page

Table 3: MoveBench studies included in the analysis (continued).

Index	Study species	Authors	Data	License	N Indiv	GPS fixes	Median fix interval (min)
98	<i>Loxodonta africana</i>	Benitez et al. [29]		Not specified	13	222,624	60
99	<i>Loxodonta africana</i>	Thaker et al. [320]	Slotow et al. [309]	CC0	6	65,747	60
100	<i>Loxodonta africana</i>	Wall et al. [341]	Wall et al. [340]	CC0	1	381	60
101	<i>Lynx rufus</i>	Prugh et al. [270]	Prugh [269]	CC0	12	25,643	240
102	<i>Lynx rufus</i>	Serieys et al. [295]	Serieys and Wong [291]	CC0	6	15,706	60
103	<i>Lynx rufus</i>	Serieys et al. [292]	Serieys et al. [294]	CC0	10	48,201	60
104	<i>Martes pennanti</i>	LaPoint et al. [198]	LaPoint et al. [197]	CC0	6	3,320	60
105	<i>Nasua narica</i>	Kays et al. [174]	Kays et al. [175]	CC0	6	7,546	60
106	<i>Nyctereutes procyonoides</i>	Toivonen et al. [325]	Toivonen et al. [324]	CC-BY-4.0	6	4,406	240
107	<i>Odocoileus virginianus</i>	Egan et al. [84]	Egan et al. [83]	Not specified	11	24,475	60
108	<i>Odocoileus virginianus</i>	Newman et al. [237]	Newman et al. [236]	CC0	6	2,889	60
109	<i>Panthera leo</i>	Patterson et al. [246]		CC-BY	3	2,099	420
110	<i>Panthera onca</i>	Morato et al. [220]	Morato et al. [221]	CC0	6	19,363	60
111	<i>Panthera pardus</i>	Power et al. [268]	Power and Venter [266]	CC0	3	1,067	1440
112	<i>Papio anubis</i>	Isbell et al. [165]	Isbell and Bidner [164]	CC-BY-4.0	6	38,758	60
113	<i>Pecari tajacu</i>	Kays et al. [174]	Kays et al. [175]	CC0	1	555	60
114	<i>Potos flavus</i>	Kays et al. [174]	Kays et al. [175]	CC0	6	4,202	60
115	<i>Prionailurus viverrinus</i>	Serieys et al. [296]	Khaewphakdee et al. [177]	CC0	6	8,908	60
116	<i>Pteropus lylei</i>	Choden et al. [57]	Choden et al. [56]	CC0	1	295	60
117	<i>Pteropus melanotus</i>	Todd et al. [323]	Todd et al. [322]	CC0	2	477	60
118	<i>Pteropus poliocephalus</i>	Boardman et al. [34]	Roshier and Boardman [278]	CC0	3	2,456	60
119	<i>Puma concolor</i>	Hooten et al. [155]		GNU General Public License v3.0	1	825	181
120	<i>Rangifer tarandus</i>		Ágústsdóttir and Þórisson [360]	CC0	6	15,723	180
121	<i>Sus scrofa</i>	Potts et al. [264]		CC-BY-4.0	6	17,446	120
122	<i>Syncerus caffer</i>	Getz et al. [120]	Cross et al. [69]	CC0	6	24,153	60
123	<i>Tapirus terrestris</i>	Medici et al. [211]	Medici [212]	CC0	12	125,278	60
124	<i>Tremarctos ornatus</i>	Pillco Huarcaya et al. [256]	Pillco Huarcaya et al. [257]	CC-BY-NC-4.0	1	830	61
125	<i>Urocyon cinereoargenteus</i>	Serieys et al. [292]	Serieys et al. [294]	CC0	1	909	60
126	<i>Vulpes lagopus</i>	Warret Rodrigues and Roth [344]	Warret Rodrigues and Roth [343]	CC0	6	3,769	90
127	<i>Vulpes vulpes</i>	Carter et al. [51]	Carter and David A. [50]	CC0	6	12,457	60
128	<i>Vulpes vulpes</i>	Porteus et al. [261]	Porteus et al. [262]	CC0	12	16,409	60
129	<i>Vulpes vulpes</i>	Warret Rodrigues and Roth [344]	Warret Rodrigues and Roth [343]	CC0	6	6,239	90
130	<i>Wallabia bicolor</i>	Fischer et al. [100]	Fischer et al. [101]	CC0	6	1,277	66
131	<i>Chelonia mydas</i>	Esteban et al. [89]	Hays et al. [139]	CC0	6	12,560	119
132	<i>Chelonia mydas</i>	Mestre et al. [216]	Patrício et al. [245]	CC-BY-NC-4.0	6	2,252	173

Continued on next page

Table 3: MoveBench studies included in the analysis (continued).

Index	Study species	Authors	Data	License	N Indiv	GPS fixes	Median fix interval (min)
133	<i>Eretmochelys imbricata</i>	Hays et al. [138]	Hays et al. [137]	CC0	6	24,017	64
134	<i>Eumetopias jubatus</i>	Wilson et al. [354]		Not specified	3	2,146	103
135	<i>Halichoerus grypus</i>	Baker et al. [14]	Lidgard et al. [202]	CC0	6	20,819	61
136	<i>Halichoerus grypus</i>	Michelot and Blackwell [217]	McConnell [207]	CC0	1	1,949	88
137	<i>Mirounga angustirostris</i>	Costa et al. [66]	Costa et al. [65]	CC0	6	26,595	72
138	<i>Mobula alfredi</i>	Setyawan et al. [298]	Setyawan et al. [297]	CC-BY-NC-4.0	5	1,296	123
139	<i>Physeter macrocephalus</i>	Irvine et al. [160]	Irvine et al. [161]	CC0	4	1,388	62
140	<i>Physeter macrocephalus</i>	Irvine et al. [163]	Irvine et al. [162]	CC0	5	2,767	62
141	<i>Chelonoidis donfaustoi</i>	Bastille-Rousseau et al. [23]	Bastille-Rousseau et al. [22]	CC0	6	78,733	60
142	<i>Chelonoidis hoodensis</i>	Bastille-Rousseau et al. [19]	Bastille-Rousseau et al. [18]	CC0	1	8,791	60
143	<i>Chelonoidis hoodensis</i>	Bastille-Rousseau et al. [23]	Bastille-Rousseau et al. [22]	CC0	6	11,147	720
144	<i>Chelonoidis porteri</i>	Bastille-Rousseau et al. [19]	Bastille-Rousseau et al. [18]	CC0	3	31,017	60
145	<i>Chelonoidis porteri</i>	Bastille-Rousseau et al. [23]	Bastille-Rousseau et al. [22]	CC0	6	60,011	60
146	<i>Gopherus berlandieri</i>	Chambers et al. [52] Guerra et al. [130]	Guerra et al. [131]	CC-BY-NC-4.0	6	50,641	61
147	<i>Chelonoidis</i> spp.	Bastille-Rousseau et al. [21]	Bastille-Rousseau et al. [20]	CC0	6	73,524	60

B.2 Trajectory curation and filtering details

We start by aggregating movement *studies*—single datasets that have been previously released by movement researchers (*i.e.*, a row in Tab. 3). A single study contains GPS trajectory data for one or more individuals of one or more species. Because different species exhibit different movement patterns, they are often tracked using different hardware with different settings. If a single dataset contained multiple species, we split the dataset and consider each species a separate “study” for all data aggregation and analysis, and refer to these as so in the main text.

We begin with the 215 studies containing GPS data on the Movebank Data Repository [2], a subset of data from Movebank [226] that have Creative Commons licensing, DOIs, and have already undergone some quality assurance and anomaly filtering. We then add an additional 14 studies sourced from a literature review of recent movement methods papers that performed some kind of empirical study, as well as 5 additional suggested studies from domain experts in the Movement Biodiversity Observation Network (Move BON)[1] Working Group on AI. In total this results in 234 studies and 89M raw GPS points that we filter down to the final set as follows.

To make this more tractable for benchmarking, we apply a set of coarse filtering criteria followed by manual inspection of all points. We log all dataset operations for reproducibility and report the full set of modifications in the dataset Croissant file (see Sec. B.4).

We first thin the GPS points to a maximum of one point per hour, matching the minimum forecast horizon we evaluate (see Sec. 5), leaving us with 14M points. We then filter out individuals with < 200 GPS points, and keep a maximum of 17, 520 GPS points per individual. We chose the latter number as the number of hours in two years, allowing for a temporal train/test split to contain a full annual cycle of behavioral patterns in both train and test (see “Train and test splits” below). If an

individual has more than the maximum number of points, we keep the most recent 17,520. Many datasets did not meet these criteria; our final set for the benchmark includes 147 studies.

We then inspect the remaining data by hand with the goal of selecting six individuals per species per study. To do this, we use a custom web application we developed, which visualizes trajectories spatially (on a map) and temporally (on a timeline). We inspect for remaining anomalous data, such as trajectories that are stationary for extended periods of time due to falling off the animal, and remove affected individuals. We then keep all individuals whose temporal coverage allows for a temporal train/test split (see below). Finally, if there are more than six individuals remaining, we randomly sample six to keep, discarding the rest.

Note that these decisions were made specifically to keep the dataset of a tractable size. In the future, users may choose to train on larger, less filtered raw datasets.

B.3 Additional details on train/test split

Train and test splits. For each study pair, we aim to create one training set and two hold-out test sets representing a *temporal* split and an *individual* split.

The temporal split defines a hold-out test set that includes test data that comes from future points in time with respect to the training data. We define a single split timestamp for each individual—any observations before that time goes in the training set, and any observations after go in the temporal test set. We aim for this split to be roughly 50/50. When individuals have overlapping spatial and temporal ranges, we avoid train-test leakage by enforcing that every individual in the study population has the same split time, so that models cannot leverage memorized information from nearby individuals during test time. For some studies, observations did not cover a large enough temporal range that would allow movement patterns to appear in both the training and temporal test set—for instance, splitting the data from long-distance migrants with only one recorded migration would mean that training data may come from completely disjoint countries (or even continents) than temporal test data. For these individuals, we only include an individual test set, and do not include a temporal test set.

The individual split defines a hold-out test set containing individuals that do not appear in the training set. We select one-third of the individuals from each study to be held-out, rounding down if there are an odd number of individuals. To avoid train-test leakage, for hold-out individuals we test on data from the same time range that would otherwise be in their temporal hold-out set—*i.e.*, any data for hold-out individuals that occurs before the temporal split point remains unused during both training and testing.

To avoid two training sets per species-study pair, the final training set is the intersection of the training data that would result from either split being created on its own—that is, it includes all data before the temporal split point for all individuals not in the held-out individuals set.

B.4 Data modification records

We built a custom application for inspecting and processing data from its raw state into the version in MOVEBENCH. The application logs atomic records of every data modification operation to maintain full data provenance. We include the complete set of data modification records alongside the release in the Croissant file.

B.5 MOVEBENCH-Mini

The studies in MOVEBENCH-Mini are: Forrest et al. [104], Borchering et al. [35], Garthe et al. [115], Aikens et al. [9], Costa et al. [66], Acácio et al. [3], Bartlam-Brooks et al. [16], Gatt et al. [116], Dodge et al. [75], and Bastille-Rousseau et al. [23].

C Environmental covariates

C.1 Full list of included covariates

Weather & climate

Table 4: MoveBench environmental covariates in the Weather & climate group, with data source, resolution, temporal coverage, and license. ERA5 entries cite Hersbach et al. [147] and were downloaded from Copernicus Climate Change Service [64]. The results contain modified Copernicus Climate Change Service information 2020. Neither the European Commission nor ECMWF is responsible for any use that may be made of the Copernicus information or data it contains.

Source	Covariate	Spatial	Temporal	Coverage	License
ERA5-Land (Muñoz-Sabater et al. [229])	Hourly Temperature 2 m	11km	1 hour	1950-2026	CC-BY-4.0
	Daily Temperature 2 m	11km	1 day	1950-2026	
	Hourly Total Precipitation	11km	1 hour	1950-2026	
	Daily Total Precipitation	11km	1 day	1950-2026	
	Hourly Dewpoint Temperature 2 m	11km	1 hour	1950-2026	
	Daily Dewpoint Temperature 2 m	11km	1 day	1950-2026	
	Hourly Surface Pressure	11km	1 hour	1950-2026	
	Daily Surface Pressure	11km	1 day	1950-2026	
	Hourly U Component of Wind 10 m	11km	1 hour	1950-2026	
	Daily U Component of Wind 10 m	11km	1 day	1950-2026	
	Hourly V Component of Wind 10 m	11km	1 hour	1950-2026	
	Daily V Component of Wind 10 m	11km	1 day	1950-2026	
	Hourly Surface Solar Radiation Downwards	11km	1 hour	1950-2026	
	Daily Surface Solar Radiation Downwards	11km	1 day	1950-2026	
	Hourly Surface Thermal Radiation Downwards	11km	1 hour	1950-2026	
	Daily Surface Thermal Radiation Downwards	11km	1 day	1950-2026	
	Hourly Surface Net Solar Radiation	11km	1 hour	1950-2026	
	Daily Surface Net Solar Radiation	11km	1 day	1950-2026	
	Hourly Surface Net Thermal Radiation	11km	1 hour	1950-2026	
	Daily Surface Net Thermal Radiation	11km	1 day	1950-2026	
	Hourly Surface Latent Heat Flux	11km	1 hour	1950-2026	
	Daily Surface Latent Heat Flux	11km	1 day	1950-2026	
	Hourly Surface Sensible Heat Flux	11km	1 hour	1950-2026	
	Daily Surface Sensible Heat Flux	11km	1 day	1950-2026	
	Hourly Snow Depth	11km	1 hour	1950-2026	
	Daily Snow Depth	11km	1 day	1950-2026	
ERA5 (Hersbach et al. [147]; downloaded from Copernicus Climate Change Service [64])	Boundary Layer Height	28km	1 hour	1950-2026	CC-BY-4.0
	Total Cloud Cover	28km	1 hour	1950-2026	
	Mean Sea Level Pressure	28km	1 hour	1950-2026	
	Thermal uplift	11km	Hourly	1950-2026 (derived)	
Landsat (U.S. Geological Survey [327])	Land surface temperature	30m	8 day	1984-2026	public domain
Bioclim (Fick and Hijmans [94])	BIO1: Annual Mean Temperature	1km	Static	1970-2000 climatology; static layer	CC-BY-4.0
	BIO2: Mean Diurnal Range	1km	Static	1970-2000 climatology; static layer	
	BIO3: Isothermality	1km	Static	1970-2000 climatology; static layer	
	BIO4: Temperature Seasonality	1km	Static	1970-2000 climatology; static layer	
	BIO5: Max Temperature of Warmest Month	1km	Static	1970-2000 climatology; static layer	
	BIO6: Min Temperature of Coldest Month	1km	Static	1970-2000 climatology; static layer	

Continued on next page

Table 4: MoveBench environmental covariates in the Weather & climate group (continued).

Source	Covariate	Spatial	Temporal	Coverage	License
Copernicus DEM GLO-30 (European Space Agency [90]) + ERA5-Land (Muñoz-Sabater et al. [229])	BIO7: Temperature Annual Range	1km	Static	1970-2000 climatology; static layer	Copernicus DEM GLO-30 & Free and open Copernicus WorldDEM-30 licence; attribution required.; CC-BY-4.0
	BIO8: Mean Temperature of Wettest Quarter	1km	Static	1970-2000 climatology; static layer	
	BIO9: Mean Temperature of Driest Quarter	1km	Static	1970-2000 climatology; static layer	
	BIO10: Mean Temperature of Warmest Quarter	1km	Static	1970-2000 climatology; static layer	
	BIO11: Mean Temperature of Coldest Quarter	1km	Static	1970-2000 climatology; static layer	
	BIO12: Annual Precipitation	1km	Static	1970-2000 climatology; static layer	
	BIO13: Precipitation of Wettest Month	1km	Static	1970-2000 climatology; static layer	
	BIO14: Precipitation of Driest Month	1km	Static	1970-2000 climatology; static layer	
	BIO15: Precipitation Seasonality	1km	Static	1970-2000 climatology; static layer	
	BIO16: Precipitation of Wettest Quarter	1km	Static	1970-2000 climatology; static layer	
	BIO17: Precipitation of Driest Quarter	1km	Static	1970-2000 climatology; static layer	
	BIO18: Precipitation of Warmest Quarter	1km	Static	1970-2000 climatology; static layer	
	BIO19: Precipitation of Coldest Quarter	1km	Static	1970-2000 climatology; static layer	
	Orographic uplift	30m	Hourly	1950-2026 (derived from ERA5-Land)	

Vegetation & land cover

Table 5: MoveBench environmental covariates in the Vegetation & land cover group, with data source, resolution, temporal coverage, and license.

Source	Covariate	Spatial	Temporal	Coverage	License
Landsat (U.S. Geological Survey [327])	NDVI	30m	8 day	1984-2026	public domain
	EVI	30m	8 day	1984-2026	
	NDSI (snow)	30m	8 day	1984-2026	
	BAI (burn)	30m	8 day	1984-2026	
MODIS MOD44B.061 (NASA LP DAAC [234])	Percent Tree Cover	250m	1 year	2000-2024	CC0
	Percent Non-tree Vegetation	250m	1 year	2000-2024	
	Percent Non-vegetated	250m	1 year	2000-2024	
MODIS MCD12Q1.061 LC_Type1 (NASA LP DAAC [233])	LC Type1 01: Evergreen Needle-leaf Forests	500m	1 year	2001-2024	CC0
	LC Type1 02: Evergreen Broadleaf Forests	500m	1 year	2001-2024	
	LC Type1 03: Deciduous Needle-leaf Forests	500m	1 year	2001-2024	
	LC Type1 04: Deciduous Broadleaf Forests	500m	1 year	2001-2024	

Continued on next page

Table 5: MoveBench environmental covariates in the Vegetation & land cover group (continued).

Source	Covariate	Spatial	Temporal	Coverage	License
Dynamic World (Brown et al. [39])	LC Type1 05: Mixed Forests	500m	1 year	2001-2024	CC-BY-4.0
	LC Type1 06: Closed Shrublands	500m	1 year	2001-2024	
	LC Type1 07: Open Shrublands	500m	1 year	2001-2024	
	LC Type1 08: Woody Savannas	500m	1 year	2001-2024	
	LC Type1 09: Savannas	500m	1 year	2001-2024	
	LC Type1 10: Grasslands	500m	1 year	2001-2024	
	LC Type1 11: Permanent Wetlands	500m	1 year	2001-2024	
	LC Type1 12: Cropland.	500m	1 year	2001-2024	
	LC Type1 13: Urban and Built-up Lands	500m	1 year	2001-2024	
	LC Type1 14: Cropland/Natural Vegetation Mosaics	500m	1 year	2001-2024	
	LC Type1 15: Permanent Snow and Ice	500m	1 year	2001-2024	
	LC Type1 16: (sand, rock, soil) areas with less than 10% vegetation.	500m	1 year	2001-2024	
	LC Type1 17: Water Bodies	500m	1 year	2001-2024	
	Water	10m	3 to 5 days	2015-2026	
	Trees	10m	3 to 5 days	2015-2026	
	Grass	10m	3 to 5 days	2015-2026	
	Flooded Vegetation	10m	3 to 5 days	2015-2026	
JRC Global Surface Water (Pekel et al. [249])	Crops	10m	3 to 5 days	2015-2026	JRC Global Surface Water & Free and open use under Copernicus / European Commission reuse terms; attribution required.
	Shrub and Scrub	10m	3 to 5 days	2015-2026	
	Built	10m	3 to 5 days	2015-2026	
	Bare	10m	3 to 5 days	2015-2026	
	Snow and Ice	10m	3 to 5 days	2015-2026	
	Distance to water	30m	Mostly static	1984-2021 source observations; under derived layer used as static	

DEM

Table 6: MoveBench environmental covariates in the DEM group, with data source, resolution, temporal coverage, and license.

Source	Covariate	Spatial	Temporal	Coverage	License
Copernicus DEM GLO-30 (European Space Agency [90])	Elevation	30m	Static	2010-2015 acquisition window; static layer	Copernicus DEM GLO-30 & Free and open Copernicus WorldDEM-30 licence; attribution required.
	Slope	30m	Static	2010-2015 acquisition window; static layer	
	Aspect	30m	Static	2010-2015 acquisition window; static layer	
	Rugosity/ruggedness	30m	Static	2010-2015 acquisition window; static layer	

Human impact

Table 7: MoveBench environmental covariates in the Human impact group, with data source, resolution, temporal coverage, and license.

Source	Covariate	Spatial	Temporal	Coverage	License
WCS (Wildlife Conservation Society (WCS) and Center for International Earth Science Information Network (CIESIN), Columbia University [352]; Sanderson et al. [280])	Human impact index	300m	Annual	2001-2020	CC BY-NC-SA 3.0

Marine

Table 8: MoveBench environmental covariates in the Marine group, with data source, resolution, temporal coverage, and license.

Source	Covariate	Spatial	Temporal	Coverage	License
Copernicus Global Ocean Waves (European Union-Copernicus Marine Service [92])	Bathymetry (sea floor depth)	0.9km	Static	2022-2026	Copernicus Marine Service & Free and open use under Copernicus Marine Service licence; attribution required.
MOD44W.006 (Carroll et al. [49])	Distance to coast	250m	Static	2000-2015 source observations; derived layer used as static	CC0
NOAA CDR (Huang et al. [157])	Sea surface temperature	28km	30 day	1981-2026	NOAA OISST v2.1 & Free NOAA/NCEI data; citation requested; no warranty
Copernicus Satellite Ocean Color (European Union-Copernicus Marine Service [93])	Ocean surface chlorophyll	4000m	30 day	1997-2026	Global Ocean Colour & Free Copernicus Marine Service licence; redistribution and derivative products permitted.
HYCOM (Cummings and Smedstad [71])	Surface current eastward	8900m	1 day	1992-2024	Freely available with no restrictions
	Surface current northward	8900m	1 day	1992-2024	

Embeddings

Table 9: MoveBench environmental covariates in the Embeddings group, with data source, resolution, temporal coverage, and license.

Source	Covariate	Spatial	Temporal	Coverage	License
AlphaEarth (Brown et al. [41])	AlphaEarth - 64 bands	10m	Annual	2017-2025	CC-BY 4.0

C.2 Details on raster preparation

Raster preparation begins by defining a *raster recipe* for a given benchmark setting. This recipe defines the spatial characteristics of the rasters that will be downloaded for each study. In particular, it specifies four things: which environmental variables we need; which of them are time-varying versus static; how much spatial context we retain around each movement location; and what spatial resolution is practical for storing and reusing the resulting rasters. We define this recipe before any download occurs so that all later stages, including feature extraction and raster-based models, operate on a consistent set of bands, extents, and spatial grids. We can therefore reuse the same raw raster cache across multiple downstream experiments whenever they share the same spatial recipe, even if they use different models or different subsets of the bands.

We resolve the recipe separately for each benchmark timescale. We determine the scale (meters per pixel) and extent (meters per raster) based on each study’s movement statistics at that timescale. We elaborate on this more below. We do this because the relevant movement radius grows with forecast horizon, and storing rasters at their native resolution for very large extents would not be tractable.

Raster sources and layer families. Most layers are downloaded from Google Earth Engine. The layers that are not available on Google Earth Engine—Human Footprint Index and BioClim—are downloaded in full as local GeoTIFF archives.

Timescale-specific extent and scale. We determine both raster extent and raster scale from movement statistics in the training split at the target forecast horizon. We first identify valid interval steps after partitioning each trajectory into temporally coherent “bursts” (segments of uninterrupted movement). By default, we treat inter-fix gaps above the 0.999 quantile of an individual’s revisit-time distribution as burst boundaries, so very large temporal gaps do not create artificial long-distance steps. We then compute a conservative spatial radius b from the valid training-step lengths L :

$$b = \min(2.0 \times Q_{0.95}(L), Q_{0.99}(L)),$$

where $Q_q(L)$ denotes the q -quantile. This is the spatial range we consider as “available” to the animal during its next step, and is equal to the minimum of double its 95th percentile step length and its 99th percentile step length. This ensures that nearly all observed steps will have raster coverage while reducing the influence of any outliers. We use b as the half-width of the square spatial context around a focal point, so each local raster patch covers a square of side length $2b$.

We choose the stored resolution for each layer by anchoring it to a canonical downstream chip size of 101×101 pixels to match prior work [105]. If a chip has half-width b , then a 101×101 grid spans the full width $2b$ with 101 pixels across each side, so the corresponding pixel size is $2b/101$. For a layer with native resolution s_{native} , we therefore store it at

$$s = \text{clip}\left(s_{\text{native}}, \frac{2b}{101}, \frac{2b}{3}\right).$$

This rule makes the raw cache match the scale of the canonical chip whenever the native raster is finer than necessary, while still enforcing a minimum of about 3×3 pixels of spatial support for very coarse layers. The lower clamp, $2b/101$, prevents fine-resolution products such as AlphaEarth (10m resolution) or DEM (30m resolution) from producing source chips that are much denser than the eventual 101×101 grid at long horizons, which would waste storage. The upper clamp, $2b/3$, prevents coarse products such as ERA5 fields (11–28km resolution) from collapsing to one or two pixels across the chip at short horizons. We apply this rule independently to each layer because their native resolutions differ by orders of magnitude.

We use two storage modes, depending on how a layer varies in time and how much reuse we expect across points: (1) We store *chips* for layers whose value depends on the focal timestamp and therefore must be queried separately for each point-time pair. In our benchmark, this includes layers with high temporal resolution such as ERA5-Land hourly and daily fields, ERA5 hourly fields, Landsat composites and their derived indices, Dynamic World probabilities, and time-varying marine layers. Each chip is a point-centered square crop with half-width b . (2) We store study-level *tiles*, reusable larger rasters for layers that are static or annual and can therefore serve many points within the same study. In our benchmark, this includes DEM, distance-to-water, bathymetry, distance-to-shore, annual MODIS land cover, annual MODIS vegetation fractions, and annual AlphaEarth embeddings.

Earth Engine query rules. We query different layers with source-specific temporal rules. For ERA5-Land hourly layers, we select the nearest image within ± 1 hour of the focal timestamp. If ERA5-Land leaves any pixels masked, we fill those masked pixels from the corresponding bands in the ERA5 hourly collection. For ERA5-Land daily aggregates, we select the nearest image within ± 1 day. If the daily aggregate is masked, we fill the six meteorological bands that also exist in the ERA5 daily collection—2 m air temperature, 2 m dewpoint temperature, total precipitation, surface pressure, and the two 10 m wind components—from ERA5 daily. The radiative flux bands and snow depth remain missing when no ERA5-Land daily aggregate is available.

For layers that are composites over some time range T (e.g. for Landsat, $T = 8$ -days), we query the most recent composite in the interval $[t - T, t]$. For Landsat, if the 8-day composite is missing or masked at some pixels, we fill the missing pixels from the most recent 32-day composite in $[t - 32 \text{ days}, t]$. We then derive NDVI, EVI, NDWI, NDSI, and BAI from the multispectral bands after loading.

For NOAA OISST, we compute a 30-day mean and a 30-day standard deviation of sea-surface temperature over the interval $[t - 30 \text{ days}, t]$. For Copernicus chlorophyll, we compute a 30-day

mean over the same interval. For HYCOM currents, we select the latest image in $[t - 1 \text{ day}, t]$ and derive current speed as $\sqrt{u^2 + v^2}$ from the eastward and northward surface components.

Dynamic World requires special handling because the product begins on 2015-06-23 and because cloud-free daily coverage is irregular. If the focal timestamp is earlier than 2015-06-23, we use the median Dynamic World probability image over the first year of product availability, 2015-06-23 through 2016-06-23. Otherwise, we compute a median over a ± 15 -day window around the focal timestamp, and if that window contains no imagery we widen it to ± 30 days.

For annual shared layers, we resolve the year from the focal timestamp and then snap to the nearest available year in the source archive when the exact year is missing. This rule applies to annual MODIS land cover, annual MODIS vegetation fractions, and annual AlphaEarth embeddings. Within the resolved year, we use a earliest available annual image.

For shared static layers, we query fixed products directly. We obtain elevation by mosaicking the Copernicus GLO-30 DEM. We derive distance to water from the JRC Global Surface Water `max_extent` mask using cumulative geodesic cost and clip the result at 30 km. We derive distance to shore from the latest MOD44W `water_mask`, computing cumulative cost from land and clipping the result at 100 km. We read bathymetric depth from the static Copernicus marine layer.

Planning and batching shared tiles. For *tiles* that are shared between many training/evaluation points, we do not center a shared raster tile on every point. Instead, we first group points into coarse longitude-latitude buckets, separately within each data split. We then expand each bucket by the movement radius b , handle antimeridian crossing explicitly when needed, and convert the resulting buffered rectangles into a regular global tile grid whose cell size is $350 \times s$ square meters. We keep only tiles that intersect at least one buffered rectangle. For annual layers, we reuse the same spatial tile plan for each resolved year.

We batch local dynamic downloads by point and shared static or annual downloads by tile, or by tile-year pair. Each point worker requests all multiband point-centered exports needed for that point. Each tile worker requests one shared tile for one layer, or one shared tile for one layer-year pair. This batching strategy makes the download process parallel, resumable, and easy to deduplicate.

Handling large exports. Before we request a GeoTIFF from Earth Engine, we estimate its size from the requested bounds, meters-per-pixel scale, number of bands, and float32 storage cost. We compare this estimate to Earth Engine’s practical direct-download limit of about 50 MB. If the estimate is near this limit, we subdivide the request into temporary subtiles, download those subtiles separately, and mosaic them locally into one final raster. We use this procedure for both point-centered chips and shared study-level tiles. We write masked pixels to disk as a dedicated nodata sentinel value (-9999) so that missingness survives the GeoTIFF export unchanged.

Reading windows and aligning layers. Raw downloads from different sources do not share a common grid, CRS, or resolution, so we align them only when we build model inputs. For shared tiles, we identify all stored tiles whose bounding boxes intersect the requested lat-lon window and merge them on demand. For raster-based datasets, we reproject all layers for each sample onto a shared grid in a WGS84 azimuthal equidistant (AEQD) projection centered on the step’s starting location. We warp every other layer onto the same extent, affine transform, and output array shape. We use bilinear interpolation for continuous layers and nearest-neighbor interpolation for categorical layers.

After reprojection, we convert the -9999 nodata sentinel back to NaN immediately after reading and construct a boolean validity mask so downstream models can distinguish missing pixels from observed zeros.

Some derived layers depend on spatial gradients and therefore require nontrivial spatial support. We enforce a minimum read window of 2×2 pixels for this reason. We derive slope, aspect, and terrain ruggedness from DEM after loading. We derive orographic uplift from slope, aspect, and ERA5-Land hourly wind. We derive thermal uplift from ERA5-Land hourly surface variables together with ERA5 hourly atmospheric variables. We derive spectral indices from the Landsat bands after loading rather than treating them as independent raw downloads.

Model-specific data preparation and storage. We store raster data in three layers.

First, we store a raw raster cache for each timescale. This cache contains multiband point-centered chips for local dynamic predictors, reusable shared tiles for static and annual predictors, one index per shared layer (and one per year for annual layers), and request metadata that records which downloads completed successfully. Because the cache key depends on the timescale-specific extent and scale, these raw caches are distinct across benchmark horizons even when they reference the same environmental products.

Second, for SSF-style models, we sample each layer at the candidate coordinates and store dense point-feature arrays to reduce computation during training and evaluation. Candidate coordinates are transformed into each raster’s CRS for sampling; terrain-derived layers are first computed on a local metric working grid. For each layer, we store one floating-point array of sampled values and one boolean array of sampled validity flags. We also store one structured shared array containing candidate coordinates, timestamps, step lengths, turn angles, individual identifiers, burst identifiers, and point identifiers. We memory-map these arrays during reading so that later stages can load only the needed rows and bands.

Third, for deepSSF, we store fully aligned raster tensors in streaming tar shards to reduce data loading time during training and evaluation, as loading many small GeoTIFF files for every forward pass is otherwise extremely slow. Each sample contains the spatial tensor, scalar time features, start bearing, target grid, and the start-point identifier.

D Metrics

Here we provide some additional information about the energy score (Sec. 5) and its derivation. Recall that, for a given trajectory step (ℓ, t) and forecast distribution $P_{t,\Delta}$ (see Sec. 3), the energy score evaluated against $\ell_{t+\Delta}$, the ground truth location at time $t + \Delta$, is:

$$\text{ES}(P_{t,\Delta}, \ell_{t+\Delta}^{\text{obs}}) = \underbrace{\mathbb{E}_{X \sim P_{t,\Delta}} \|X - \ell_{t+\Delta}^{\text{obs}}\|_2}_{\text{expected } L_2 \text{ error of forecast}} - \frac{1}{2} \underbrace{\mathbb{E}_{X, X' \sim P_{t,\Delta}} \|X - X'\|_2}_{\text{spread of forecast}}.$$

To see why ES takes the form that it does, consider the *energy distance* [273] between a predicted distribution P and a true distribution Q : $D^2(P, Q) = 2\mathbb{E}\|X - Y\| - \mathbb{E}\|X - X'\| - \mathbb{E}\|Y - Y'\|$, with $X, X' \sim P$ and $Y, Y' \sim Q$. If we had access to the target distribution Q (i.e., the true conditional distribution of an animal’s future location), we could compute and try to minimize the energy distance between Q and our predicted distribution P . However, for any given forecast instance we only observe a single realization y , which we treat as a draw from Q . In expectation, the energy score satisfies $\mathbb{E}_{Y \sim Q} \text{ES}(P, Y) = \frac{1}{2}D^2(P, Q) + \frac{1}{2}\mathbb{E}\|Y - Y'\|$, i.e., in expectation over Q the energy score is the energy distance between P and Q up to a term that does not depend on P . Therefore, minimizing the expected energy score is equivalent to minimizing the energy distance.

E Method details

Integrated step-selection functions (iSSF). iSSFs [13] and their extensions model the probability of an animal’s next step as a function of both movement features (e.g. velocity) and environmental features (see Sec. 4). In practice this is implemented as a conditional logistic regression. At training time, a set of future candidates $C_{t,\Delta} = \{\ell_{t+\Delta}, t + \Delta\}_{i=1}^{n_{\text{cand}}} \in \mathcal{X}$ is sampled from parametric distributions fit to the training data. A model then predicts movement probabilities $P_{t,\Delta}$ over the discrete set of candidates $C_{t,\Delta}$.

In our experiments, we benchmark two iSSF baselines: **(1) iSSF Linear** uses a simple linear layer as the prediction function, matching prior work. Depending on the experiment, inputs may include movement-derived, time-derived, and environmental covariates, as well as linear “interactions” between them. **(2) iSSF MLP** uses a multilayer perceptron (MLP) in place of the linear layer used by standard SSFs. We include the same set of covariates, but no interaction terms, instead allowing the MLP to learn any linear or nonlinear interactions between input variables.

MoveFormer. MoveFormer [61] utilizes the same candidate-scoring framework as iSSFs, but incorporates several attention-based [335] components. First, it adds additional trajectory history before the most recent step by summarizing past steps using a transformer encoder, with one token per

historical step. It then embeds the current set of candidates, performs dot-product attention between the historical embedding and each candidate embedding, and outputs candidate selection probabilities over the same discrete support as iSSFs. MoveFormer also includes token representing the taxonomy of the species of interest, and is the only baseline designed to be trained across studies and across species.

deepSSF. deepSSF [105] replaces the candidate scoring of iSSFs with raster inputs processed by (1) a convolutional habitat-selection head, which captures spatial structure over the range of possible next steps, and (2) a movement head with convolutional, pooling, and fully connected layers that predicts the parameters of a step-length and turn-angle movement kernel. The resulting habitat-selection and movement surfaces are then combined to produce a predicted next-step probability surface.

E.1 Implementation

We implement iSSFs in PyTorch. For each valid step start, we construct one discrete choice set containing n_{cand} sampled candidates plus the observed endpoint, so each training example has $P = n_{\text{cand}} + 1$ alternatives. We fit the candidate proposal distributions on the training split only: step lengths are sampled from a Gamma distribution fit to valid training-step lengths (with location fixed at zero), and turn angles are sampled from a von Mises distribution fit to valid training turn angles, matching prior work [13]. If either parametric fit fails, we fall back to empirical resampling from the corresponding training distribution. We seed candidate generation deterministically from a global random seed and the step identifier, which makes the sampled choice sets exactly reproducible. We append the observed endpoint as the final alternative in each choice set.

For each alternative, we compute three movement features—step length, log step length, and $\cos(\text{turn angle})$ —and we sample all environmental covariates at the candidate endpoint. We standardize environmental covariates with training-split means and standard deviations, and we standardize the three movement features using the corresponding training-split moments. When time covariates are enabled, we encode time of year and time of day as four cyclic features ($\sin$, $\cos$); when individual identity is enabled, we include an indexed individual-specific effect.

For **iSSF Linear**, we use a linear utility function with three groups of terms: (i) one coefficient for each environmental covariate at the candidate endpoint, (ii) one coefficient for each movement feature (step length, log step length, and $\cos(\text{turn angle})$), and (iii) one coefficient for each pairwise product between a movement feature and an environmental covariate. When time covariates are present, we also include time main effects together with time-by-environment and time-by-movement interactions. The resulting choice probability is

$$\Pr(y_i = p \mid C_i) = \frac{\exp(\eta_{ip})}{\sum_{j=1}^P \exp(\eta_{ij})}.$$

We train this model with cross-entropy loss over the P alternatives in each choice set, taking the observed endpoint as the target class. This is algebraically the same likelihood as conditional logistic regression. Standard R implementations typically fit the same model by reshaping the data into one row per alternative, defining one stratum per step, and optimizing the equivalent conditional logit / Cox partial likelihood with `clogit`. Our implementation differs in data layout and optimizer. We use tensorized batches of whole choice sets and optimize the same conditional likelihood with Adam, which lets us share one training and evaluation pipeline across the PyTorch baselines and scale to repeated large-benchmark runs.

For **iSSF MLP**, we keep the same candidate-generation procedure, feature construction, and conditional choice loss, but we replace the linear score η_{ip} with an MLP applied candidate-wise to the concatenated feature vector. In this variant, we do not manually specify interaction terms. Instead, the network receives the base movement, environmental, temporal, and optional individual-level inputs and learns linear and nonlinear interactions directly. We train both iSSF variants with the same PyTorch training scaffold, using minibatches of choice sets, Adam optimization, checkpointing, and early stopping on held-out negative log-likelihood.

F Hyperparameters

Covariates. For all experiments we include some experiment-specific set of the environmental covariates in Tab. 2, as well as movement and time-derived covariates for all methods, which have

been shown to improve performance in prior work [13, 104, 105]. First, for two locations ℓ and ℓ' , we define a *step* beginning at ℓ and ending at ℓ' with displacement vector $\delta\ell' = \ell' - \ell$. Then, the corresponding *step length* is $d' = \|\delta\ell'\|_2$. When $d' > 0$, we define the corresponding *bearing* as the angle $\theta' = \text{atan2}((\delta\ell')_2, (\delta\ell')_1) \in (-\pi, \pi]$. Finally the *turn angle* is the signed angular increment between successive bearings, $\alpha' = \text{wrap}(\theta' - \theta) \in (-\pi, \pi]$.

We then include as covariates for each observed and candidate step: (1) step length, (2) log step length, (3) cosine of turn angle, (4-5) sine and cosine of normalized day-of-year, and (6-7) sine and cosine of normalized time-of-day [13, 104]. For iSSF Linear, we include linear interaction terms between all movement- and time-derived variables and all environmental variables, following prior work. That is, for each movement- or time-derived variable and each environmental variable, we construct an additional feature equal to their product. This results in $(7 + 1) \times 160 = 1280$ input features for iSSF Linear and $7 + 160 = 167$ input features for the other methods. We z-score normalize environmental and movement covariates using training set statistics.

Candidate selection. To be consistent with prior work, we fit a gamma distribution for step-lengths and a Von Mises distribution to turn-angles for all experiments. We use $n_{cand} = 100$ candidates for all steps for all experiments.

Time horizons. We evaluate forecasting methods at a fixed set of forecast horizons $\Delta \in \{1 \text{ hour}, 12 \text{ hours}, 1 \text{ day}, 7 \text{ days}\}$. For each horizon Δ , we evaluate Eq. (1) at every test-set observation (ℓ_n, t_n) for which a valid target location is available within a tolerance of time $t_n + \Delta$. We call this set of observations $\tau_\Delta \subseteq \tau$. We then average these scores over forecast origins for each individual to obtain a per-individual per-timescale mean score $\text{ES}_\Delta = \frac{1}{|\tau_\Delta|} \sum_{i=1}^{|\tau_\Delta|} \text{ES}(P_{t_i, \Delta}, \ell_{t_i + \Delta})$. For our experiments, we set this tolerance to $\pm\Delta/3$.

There are a small number of studies that have no valid test set observations for a given timescale due to either no matching end steps within this tolerance or a native sample rate larger than the specified timescale (*e.g.* studies that were originally sampled with fix rates longer than one hour). During evaluation, we simply skip these study-timescale pairs. Any relevant aggregations (*e.g.* Fig. 3) do not consider skipped study-timescale pairs in their mean and variance calculations.

Evaluation extent. To standardize evaluation, we generate predictions for each method over the entire 101×101 pixel raster of every test point at every time horizon, and compute energy scores on this 101×101 discrete domain. For methods that are trained with candidate points sampled from parametric distributions (*i.e.* iSSF and MoveFormer), we first compute the renormalized probability assigned to each raster cell under the fitted step-length and turn-angle distributions and then combine this with the model output for each cell to obtain the final predictive surface. This corrects for the mismatch between training on non-uniformly sampled candidate locations and evaluating on a fixed uniform grid.

Hyperparameters. We first perform a small hyperparameter sweep of all methods on MOVEBENCH-Mini for fair comparison. For iSSF linear, we swept over two batch sizes and three learning rates; for iSSF MLP, we swept over batch size, learning rate, number of hidden layers, and hidden layer size; for deepSSF, we swept over learning rate, output channels, and hidden dimension; for MoveFormer, we swept over learning rate, number of transformer layers, number of transformer heads, and context length. We used the best setting from each method for all of our main results. We train the iSSF methods on CPU and deepSSF and MoveFormer on a GPU cluster. We use the temporal hold-out sets as validation during training. We evaluate methods during training roughly every 10,000 training samples and perform early stopping if validation loss has not decreased in 10 evaluation runs. We then evaluate the best checkpoint according to validation loss on both test sets.

G Additional experiment details

G.1 All experimental results

Table 10: Raw energy scores for each study and method. Lower is better. Dashes indicate unavailable results. MoveFormer denotes the shared model.

Study	Method	ES, Temporal split ↓				ES, Individual split ↓			
		1h	12h	1d	7d	1h	12h	1d	7d
Signer et al. [305]	don't move	–	686	878	1482	–	–	–	–
	crw	–	538	682	1100	–	–	–	–
	issf linear	–	537	668	1071	–	–	–	–
	issf mlp	–	522	662	1056	–	–	–	–
	deepssf	–	529	664	1153	–	–	–	–
	moveformer	–	496	627	1124	–	–	–	–
Acácio et al. [3]	don't move	14709	102696	185534	1150541	15027	114012	188756	1098839
	crw	11345	79685	130827	776299	11635	88671	134543	735285
	issf linear	10693	72715	98796	431919	11223	82599	111520	502535
	issf mlp	10133	69791	91115	381937	10847	80573	102372	578088
	deepssf	7924	60409	117259	396321	8408	71629	122539	639376
	moveformer	6610	56057	94750	422788	6055	68048	112552	817330
Wilson et al. [354]	don't move	2470	17918	23475	25343	2276	14058	19956	35036
	crw	1913	13968	18544	19991	1823	11476	16371	29322
	issf linear	1842	12980	17145	18195	1696	10464	15184	29658
	issf mlp	1853	12711	17231	17059	1696	10422	15359	30897
	deepssf	1846	13700	18682	20175	1707	11645	17946	29853
	moveformer	1902	13736	18486	19139	1788	11634	16123	29929
Tarroux et al. [319]	don't move	–	–	–	–	11422	102081	163748	651097
	crw	–	–	–	–	8327	83602	136743	511375
	issf linear	–	–	–	–	8269	81923	130108	634303
	issf mlp	–	–	–	–	8167	82158	135286	452419
	deepssf	–	–	–	–	8151	80929	123860	689207
	moveformer	–	–	–	–	5638	66264	127909	429473
Forrest et al. [104]	don't move	182	1194	1028	1747	131	1395	956	1154
	crw	156	878	823	1430	124	1062	785	983
	issf linear	157	895	829	1436	123	1146	845	979
	issf mlp	166	840	869	1409	137	1053	843	986
	deepssf	153	837	814	1403	118	986	1161	1296
	moveformer	144	675	774	1518	120	871	726	1170
Bartlam-Brooks et al. [16]	don't move	451	3455	4911	15903	433	3308	4295	12271
	crw	352	2637	3795	12912	336	2535	3395	10518
	issf linear	348	2922	4145	13661	331	2682	3473	10780
	issf mlp	352	2620	3823	13734	336	2481	3309	11319
	deepssf	344	2608	3907	12587	327	2458	3360	10793
	moveformer	317	2524	3993	13852	300	2399	3480	10877
Bastille-Rousseau et al. [19]	don't move	15.4	33.8	61.7	125	–	–	–	–
	crw	12.3	29.2	52.9	116	–	–	–	–
	issf linear	12.1	28.3	49.4	96	–	–	–	–
	issf mlp	11.8	27.2	47.8	87.5	–	–	–	–
	deepssf	11.9	26.5	43.2	81.6	–	–	–	–
	moveformer	11.6	27.9	50.7	118	–	–	–	–
Bastille-Rousseau et al. [19]	don't move	15	33.3	41.1	80.8	–	–	–	–
	crw	11.2	25.1	32.6	65.7	–	–	–	–
	issf linear	11.1	24.4	32	61.9	–	–	–	–
	issf mlp	11.2	23.7	32	61	–	–	–	–
	deepssf	11.3	23.3	30.3	59.7	–	–	–	–
	moveformer	10.7	22.3	31.6	69.7	–	–	–	–

Continued on next page

Table 10: Energy score results (continued).

Study	Method	ES, Temporal split ↓				ES, Individual split ↓			
		1h	12h	1d	7d	1h	12h	1d	7d
Bastille-Rousseau et al. [23]	don't move	13.5	36.6	64.6	230	16.7	49.5	79.2	302
	crw	10.6	30.6	53.4	195	13	40.3	64.3	253
	issf linear	10.7	30.2	52.8	192	13	40.2	65	260
	issf mlp	10.5	30	52.4	192	13	40	63.4	256
	deepssf	10.5	29.6	53.4	283	13	39.7	69.3	519
	moveformer	9.97	29	50.8	223	12.7	39.4	64.3	367
Bastille-Rousseau et al. [23]	don't move	–	23.2	39.2	105	–	26.8	49.6	128
	crw	–	20	32.8	84.1	–	23	41.2	102
	issf linear	–	20	32.7	80.3	–	24.6	43.7	111
	issf mlp	–	19.7	32.3	77	–	22.9	41.4	101
	deepssf	–	19.3	32.1	78.3	–	22.1	40.5	99.7
	moveformer	–	19	33.2	91.5	–	22	41	109
Bastille-Rousseau et al. [23]	don't move	14.7	42	74.8	383	11.7	30.8	50.6	182
	crw	11.3	34.5	62.5	330	9.13	25.6	42.7	157
	issf linear	11.3	34.3	61.9	330	9.75	26.4	45	177
	issf mlp	11.2	33.3	61	323	9.07	26.1	44.8	180
	deepssf	11.2	33.5	61.5	325	9.11	28.3	45	154
	moveformer	10.6	32.7	59.8	426	8.59	25.4	43.5	182
Bastille-Rousseau et al. [21]	don't move	14.1	45.1	95.7	404	16	51.7	107	502
	crw	11.4	39.8	83	370	13.3	44.9	91.6	447
	issf linear	11.3	39	79	350	13.1	44.1	89.6	436
	issf mlp	11.2	38.2	78	348	13	43.5	91.6	485
	deepssf	11	36.5	75.5	303	12.9	43.8	97.8	522
	moveformer	10.5	36.8	74.6	415	12.3	42.2	85.5	499
Yanco et al. [355]	don't move	723	2611	8714	66096	528	4048	13736	70021
	crw	704	2664	10894	78447	556	3740	14914	76633
	issf linear	718	2631	10056	75157	561	3968	13886	72646
	issf mlp	763	2908	20730	154404	634	3947	27104	183411
	deepssf	665	2311	10383	86162	522	3738	17134	47692
	moveformer	665	3359	12250	77812	489	3814	24228	79664
Yanco et al. [355]	don't move	2878	14508	24895	153212	2126	10729	19061	120209
	crw	2993	14374	26894	160346	2308	10735	20603	130433
	issf linear	2416	13664	21357	130317	3483	12105	21275	118599
	issf mlp	2642	13987	28710	273745	2987	10994	28628	248380
	deepssf	2469	12204	23979	76281	2059	10566	19069	143504
	moveformer	2262	17350	24338	127266	2251	12008	29656	107292
Hennig et al. [143]	don't move	–	1347	1964	4774	–	1098	1545	3276
	crw	–	1010	1468	3584	–	845	1201	2637
	issf linear	–	1010	1467	3592	–	848	1208	2716
	issf mlp	–	1004	1458	3553	–	846	1203	2646
	deepssf	–	988	1460	3549	–	842	1212	2664
	moveformer	–	1001	1466	3679	–	800	1153	2589
Borchering et al. [35]	don't move	458	1691	1726	1867	492	2029	1732	1933
	crw	416	1380	1438	1532	442	1647	1417	1557
	issf linear	416	1370	1420	1510	446	1672	1435	1585
	issf mlp	460	1330	1406	1496	501	1603	1500	1539
	deepssf	402	1282	1378	1481	445	2186	2061	1580
	moveformer	406	1208	1298	1445	438	1306	1232	1407
Mallon et al. [204]	don't move	3643	14843	30107	162001	3223	10709	22685	119029
	crw	3486	13930	27807	155217	3080	10207	21749	122251
	issf linear	3370	13809	26201	140192	3043	10031	21774	114165
	issf mlp	3589	14036	26351	167281	3266	12185	32572	260620
	deepssf	3007	13086	25329	127195	2812	16918	33140	144993
	moveformer	2558	10990	20137	112809	2453	8610	16205	94851

Continued on next page

Table 10: Energy score results (continued).

Study	Method	ES, Temporal split ↓				ES, Individual split ↓			
		1h	12h	1d	7d	1h	12h	1d	7d
Mallon et al. [204]	don't move	1189	3735	6025	14106	6006	6325	8962	12001
	crw	1289	3720	5718	12244	4329	5613	7889	9960
	issf linear	1228	3526	5826	12184	4448	6647	7816	9923
	issf mlp	1521	3921	6179	11599	4820	7789	9349	13917
	deepssf	1056	3230	5178	12137	5153	5577	8989	10925
	moveformer	1067	3305	6154	11472	3907	6006	8071	10989
Dodge et al. [75]	don't move	123	13632	25986	127958	–	11197	18583	88166
	crw	119	12979	24505	129894	–	10849	18677	101220
	issf linear	119	12818	23200	111820	–	11560	19451	104428
	issf mlp	119	13649	24150	135738	–	12810	20398	94295
	deepssf	91.3	11080	28284	120875	–	10543	19727	94188
	moveformer	119	11129	19836	106666	–	9309	14309	80871
Borrmann et al. [37]	don't move	1912	5355	4453	7265	2312	4131	3203	4483
	crw	1824	4680	3955	6209	2011	3314	2642	3545
	issf linear	1823	4500	3861	5785	2020	3394	2774	3586
	issf mlp	2139	4562	4185	5680	2299	4037	3472	4136
	deepssf	1692	4043	3800	5836	2070	3177	2861	3964
	moveformer	2410	3676	3649	5127	2266	2636	2767	3446
Cozzi et al. [68]	don't move	–	3579	5391	8582	–	4145	5907	7785
	crw	–	2902	4230	6642	–	3268	4543	6012
	issf linear	–	2844	4141	6454	–	3267	4558	5845
	issf mlp	–	2845	4119	8031	–	3330	4521	7257
	deepssf	–	2913	4055	6414	–	3728	4948	6137
	moveformer	–	2838	4008	6054	–	3252	4200	5603
Ciuti et al. [62]	don't move	–	951	1350	3361	–	1002	1293	3002
	crw	–	727	1039	2540	–	760	989	2225
	issf linear	–	737	1052	2525	–	763	996	2248
	issf mlp	–	719	1021	2397	–	753	980	2172
	deepssf	–	707	1016	2289	–	750	1020	2736
	moveformer	–	698	1046	2580	–	714	996	2243
Buitendijk et al. [42]	don't move	431	4162	4043	10255	305	2912	2547	5714
	crw	413	3235	3377	8288	302	2518	2390	5274
	issf linear	414	3144	3325	7989	313	2592	2445	5120
	issf mlp	449	3128	3310	8040	332	2479	2633	5768
	deepssf	362	2918	3264	8380	264	2573	2965	6902
	moveformer	361	2720	3346	8096	292	2353	2376	7992
Buitendijk et al. [42]	don't move	163	748	427	773	152	871	510	869
	crw	159	626	391	678	147	701	449	732
	issf linear	152	604	374	649	143	679	444	698
	issf mlp	168	623	400	677	154	687	452	719
	deepssf	148	515	382	667	137	594	450	809
	moveformer	143	385	354	663	133	441	416	675
Buitendijk et al. [42]	don't move	–	–	–	–	377	2981	2426	28188
	crw	–	–	–	–	344	2342	2032	29501
	issf linear	–	–	–	–	344	2280	2094	30219
	issf mlp	–	–	–	–	359	2261	2012	34154
	deepssf	–	–	–	–	327	2244	2328	28645
	moveformer	–	–	–	–	320	1794	2087	43477
Byholm et al. [44]	don't move	5008	35676	40293	205251	3181	28272	26795	216145
	crw	4779	29832	42184	207371	3365	24647	31974	224411
	issf linear	4650	29799	41279	194589	3096	25041	27414	183623
	issf mlp	5647	28434	49286	220742	4674	25272	36811	198411
	deepssf	4573	36180	36643	157740	2996	32143	29063	201837
	moveformer	4060	23480	42025	245204	2843	20129	30754	198982

Continued on next page

Table 10: Energy score results (continued).

Study	Method	ES, Temporal split ↓				ES, Individual split ↓			
		1h	12h	1d	7d	1h	12h	1d	7d
Mirski et al. [218]	don't move	6462	40770	94905	513749	10364	35663	83577	429162
	crw	6150	37229	83172	445432	9665	32406	72183	368096
	issf linear	6099	35397	76265	432178	9600	31570	67520	367457
	issf mlp	7682	38112	75552	409089	10511	34312	65428	336234
	deepssf	6101	35243	79282	472710	9993	30422	71474	356192
	moveformer	3537	29220	62453	400605	7569	24800	52287	296514
Kölzsch et al. [186]	don't move	–	–	–	–	2663	3505	7918	154158
	crw	–	–	–	–	2784	4157	10578	150882
	issf linear	–	–	–	–	2039	4459	9026	104694
	issf mlp	–	–	–	–	2502	5596	18843	142402
	deepssf	–	–	–	–	2297	4125	9582	94533
	moveformer	–	–	–	–	2480	3702	10912	143962
Carter et al. [51]	don't move	631	1920	1980	2335	400	1194	1264	1319
	crw	518	1423	1521	1806	339	932	994	1045
	issf linear	514	1446	1536	1810	338	943	1001	1027
	issf mlp	534	1371	1523	1806	399	945	1020	1046
	deepssf	502	1406	1450	1818	348	995	1178	1045
	moveformer	501	1273	1367	1724	343	808	805	985
Choden et al. [57]	don't move	1686	5752	1592	–	–	–	–	–
	crw	1530	4416	1549	–	–	–	–	–
	issf linear	1463	2358	1524	–	–	–	–	–
	issf mlp	1734	1342	2270	–	–	–	–	–
	deepssf	1460	511	1491	–	–	–	–	–
	moveformer	1446	3009	1534	–	–	–	–	–
Beirne et al. [27]	don't move	1493	4340	8141	10870	1669	4088	8024	11345
	crw	1353	3766	6337	8307	1511	3563	6231	8518
	issf linear	1339	3749	6357	8001	1497	3552	6362	8683
	issf mlp	1476	3973	6169	7896	1611	3879	6407	8604
	deepssf	1244	3525	6218	7810	1404	3332	6356	8607
	moveformer	1251	3494	6105	8377	1369	3352	6146	10098
Beirne et al. [27]	don't move	1459	6764	12986	41183	1320	6917	12662	37614
	crw	1457	6123	10665	29754	1338	6293	10454	27362
	issf linear	1448	6134	10610	27707	1330	6345	10393	24667
	issf mlp	1811	6726	10731	29163	1713	6859	10488	26418
	deepssf	1297	5760	10630	28512	1186	5847	10410	26013
	moveformer	1268	5827	11574	30638	1171	6026	11026	29280
Chynoweth et al. [60]	don't move	–	1006	628	959	–	908	783	1228
	crw	–	781	507	758	–	695	622	968
	issf linear	–	739	495	729	–	687	617	958
	issf mlp	–	780	499	730	–	712	612	939
	deepssf	–	687	517	745	–	744	644	970
	moveformer	–	571	491	761	–	577	602	936
Hooten et al. [155]	don't move	–	1474	2791	12136	–	–	–	–
	crw	–	1219	2313	9215	–	–	–	–
	issf linear	–	1239	2332	10187	–	–	–	–
	issf mlp	–	1218	2304	9115	–	–	–	–
	deepssf	–	1215	2327	8930	–	–	–	–
	moveformer	–	1259	2396	9323	–	–	–	–
Getz et al. [120]	don't move	281	1948	2848	8471	229	1448	2138	5186
	crw	225	1422	2111	6195	188	1084	1588	3751
	issf linear	225	1423	2098	5833	188	1087	1583	3963
	issf mlp	229	1404	2093	5796	201	1082	1624	4714
	deepssf	216	1409	2093	5351	188	1134	1820	10758
	moveformer	201	1430	2247	6246	182	1082	1753	3757

Continued on next page

Table 10: Energy score results (continued).

Study	Method	ES, Temporal split ↓				ES, Individual split ↓			
		1h	12h	1d	7d	1h	12h	1d	7d
Dodge et al. [74]	don't move	5353	34587	62573	280107	6607	43497	65811	286256
	crw	4826	30316	56025	246481	5821	36262	57045	250505
	issf linear	4342	28181	57592	231380	5384	34584	61173	229734
	issf mlp	5260	29365	55301	227068	6005	34634	56091	224691
	deepssf	4100	27690	51053	212281	5304	34457	57276	216689
	moveformer	4532	27214	50692	237828	5758	33513	55062	249763
Aikens et al. [10]	don't move	–	1318	1580	4205	–	1477	1735	3683
	crw	–	1020	1237	3416	–	1118	1321	2917
	issf linear	–	1022	1238	3388	–	1113	1329	2951
	issf mlp	–	1014	1226	3363	–	1115	1322	2943
	deepssf	–	1007	1221	3338	–	1105	1341	2836
	moveformer	–	922	1210	3538	–	1005	1277	2911
Eby et al. [80]	don't move	3066	20725	9704	–	1627	8116	3739	–
	crw	2914	16668	8374	–	1697	6088	3381	–
	issf linear	2987	11879	8380	–	1797	4996	3413	–
	issf mlp	3178	11620	8489	–	2057	4095	4377	–
	deepssf	2705	16013	7516	–	1440	6018	3144	–
	moveformer	2963	12259	8081	–	1581	4475	3183	–
Efrat et al. [82]	don't move	5959	30718	52850	248370	6853	36989	65163	259212
	crw	5510	28702	46733	215023	6287	34047	56827	224338
	issf linear	5389	28169	45107	198791	6314	34800	56663	214723
	issf mlp	5554	29808	45488	195210	6419	35923	56517	233247
	deepssf	4938	26666	44358	204936	5652	33400	56523	225518
	moveformer	4178	26550	39034	196523	4724	32044	50285	203906
Costa et al. [66]	don't move	4841	31827	55339	307797	4599	22654	37745	168978
	crw	3039	20943	35263	205351	3915	16767	28432	135987
	issf linear	3029	20876	35151	199787	3910	16519	28135	135558
	issf mlp	4841	31693	54079	197181	4599	22542	36800	140227
	deepssf	2999	20214	32596	240515	3931	16179	29177	221675
	moveformer	2837	17389	27430	165246	3974	16053	26556	138921
Aikens et al. [8] Aikens et al. [8] Aikens et al. [8]	don't move	808	5506	10795	67780	719	4931	11354	71754
	crw	787	4881	10506	65843	718	4582	11259	67626
	issf linear	819	5024	10645	69071	733	4792	11434	67676
	issf mlp	889	4930	11328	73088	826	4795	12494	70328
	deepssf	753	4774	10203	66530	678	4544	11578	68617
	moveformer	781	4981	10559	83443	697	4828	12313	70387
Anderson et al. [12]	don't move	–	–	–	–	2165	13498	13949	21607
	crw	–	–	–	–	2070	11443	12318	18874
	issf linear	–	–	–	–	2067	12105	13136	19266
	issf mlp	–	–	–	–	2383	11570	12665	18721
	deepssf	–	–	–	–	1937	12436	11989	17901
	moveformer	–	–	–	–	1942	9559	11708	17833
Fischer et al. [100]	don't move	42.7	156	109	134	46.1	193	144	165
	crw	33.8	112	78.4	96	37.4	138	107	123
	issf linear	33.7	97.2	75.8	74.2	38.2	147	104	134
	issf mlp	33.5	78	76.4	36.5	37.4	142	105	166
	deepssf	32.7	106	74.6	29.3	36.9	108	104	256
	moveformer	32.9	66.7	68	97.3	36.5	97.4	101	144
Fleishman et al. [102]	don't move	708	797	978	912	984	1269	1745	2935
	crw	585	653	825	754	856	1076	1517	2655
	issf linear	584	629	798	723	881	1088	1504	2536
	issf mlp	605	652	742	727	886	1133	1453	2684
	deepssf	541	655	699	628	1118	1078	1725	2892
	moveformer	563	651	752	735	885	1115	1458	2673

Continued on next page

Table 10: Energy score results (continued).

Study	Method	ES, Temporal split ↓				ES, Individual split ↓			
		1h	12h	1d	7d	1h	12h	1d	7d
Friedemann et al. [109]	don't move	1028	2025	2830	13611	664	1332	1794	15917
	crw	955	1757	2572	15212	728	1340	1892	17952
	issf linear	947	1933	2717	16778	986	1825	2203	28653
	issf mlp	992	1514	2544	18566	914	2680	2318	27992
	deepssf	943	1524	2995	25396	2307	3149	3591	88155
	moveformer	936	2172	2658	15843	659	1645	1711	19499
Friedemann et al. [109]	don't move	2661	4879	11011	55817	2488	4716	13515	70319
	crw	2472	4519	11116	65167	2405	4632	13696	79131
	issf linear	2270	3722	9585	53301	2160	4276	11124	66584
	issf mlp	2327	3482	9923	130847	4222	8442	14390	166910
	deepssf	2223	3417	9731	34528	5538	12806	29388	163309
	moveformer	2186	4333	8784	57907	3408	8916	10553	74879
Garthe et al. [113]	don't move	3779	25219	24103	18892	3214	18151	18754	5819
	crw	3466	20319	19777	15689	3039	15036	15594	5227
	issf linear	3276	19554	18928	15809	2830	14741	15401	8230
	issf mlp	3644	17444	17842	15618	3226	13349	14328	10611
	deepssf	3089	19231	19076	14826	2660	14623	15570	5919
	moveformer	3324	16591	16532	16355	2832	12957	15410	4935
Garthe et al. [115]	don't move	2279	10575	5399	4822	1710	4426	3668	4468
	crw	2102	8816	4685	4161	1633	3853	3259	3908
	issf linear	1970	8387	4359	3520	1558	4170	3150	2979
	issf mlp	2493	8508	5422	4169	2115	5255	4623	3775
	deepssf	1906	6379	4424	3465	1480	3112	2996	2959
	moveformer	2151	5169	4063	4192	1641	3006	2786	4295
Geary et al. [119]	don't move	5112	18980	24054	25301	4042	19343	29067	35798
	crw	4574	15706	19901	20945	3773	16452	24698	30099
	issf linear	4336	14438	17731	18786	3491	14912	21420	26141
	issf mlp	5027	13782	16803	18506	4396	14899	20579	23426
	deepssf	4328	14484	17691	15609	3415	15010	21933	22956
	moveformer	4374	15215	18772	20458	3543	16228	22638	28152
Glass et al. [122]	don't move	854	5205	7613	10142	716	6028	8949	11309
	crw	718	3860	5580	7309	626	4547	6563	7918
	issf linear	710	3870	5594	7478	623	4512	6446	7983
	issf mlp	831	4012	6597	7379	798	4714	7807	7942
	deepssf	712	3838	5568	7114	615	4512	6582	8042
	moveformer	690	3972	5652	6884	596	4915	6695	8662
Sorensen et al. [311]	don't move	959	1332	1137	1129	1201	1731	1123	1765
	crw	798	1048	939	925	1031	1394	952	1438
	issf linear	733	945	851	820	962	1191	874	1339
	issf mlp	681	977	719	746	911	1194	817	1415
	deepssf	798	1082	944	879	1028	1429	1010	1772
	moveformer	736	1015	923	1171	921	1297	904	1682
Kölzsch et al. [186]	don't move	–	–	–	–	2939	2838	6778	120881
	crw	–	–	–	–	2900	2807	8142	132570
	issf linear	–	–	–	–	2920	2983	8471	104223
	issf mlp	–	–	–	–	2956	3112	17175	147094
	deepssf	–	–	–	–	2802	2671	6919	93783
	moveformer	–	–	–	–	2839	2525	7017	146374
Chambers et al. [52] Guerra et al. [130]	don't move	6.33	11.3	13.9	25.6	7.99	17.5	23.4	40
	crw	4.59	8.81	11	20.2	5.85	13.6	18.4	31
	issf linear	4.62	8.78	11.1	21.7	5.84	14	18.5	31.8
	issf mlp	4.55	8.66	10.8	19.4	5.81	13.5	18.1	30.1
	deepssf	4.55	8.5	10.6	19.5	5.89	13.5	18	30.5
	moveformer	4.24	8.1	10.6	20.6	5.51	12.9	17.2	31.5

Continued on next page

Table 10: Energy score results (continued).

Study	Method	ES, Temporal split ↓				ES, Individual split ↓			
		1h	12h	1d	7d	1h	12h	1d	7d
Harel and Nathan [136]	don't move	6735	7690	11595	11900	5988	8344	15210	15390
	crw	5939	6765	9890	10188	5377	7166	12554	12650
	issf linear	5795	6551	9443	9571	5340	6443	11860	12114
	issf mlp	5976	6262	10187	10922	5654	6495	12458	12547
	deepssf	5733	5232	9465	10059	5076	6089	12028	12269
	moveformer	5368	11658	11329	9612	4921	12281	13286	12648
Hays et al. [138]	don't move	136	376	201	299	118	204	168	204
	crw	109	273	160	236	91.8	170	134	171
	issf linear	107	265	155	224	90.4	181	129	168
	issf mlp	104	244	153	207	91.9	194	135	213
	deepssf	108	234	164	199	113	438	140	429
	moveformer	103	192	151	210	91.7	139	120	153
Esteban et al. [89]	don't move	–	–	–	–	473	2878	3181	11626
	crw	–	–	–	–	404	2454	3643	18485
	issf linear	–	–	–	–	401	2517	3672	18628
	issf mlp	–	–	–	–	397	2459	4041	34229
	deepssf	–	–	–	–	399	2593	3084	12825
	moveformer	–	–	–	–	410	3676	3638	27827
Eggeman et al. [86]	don't move	–	1125	1388	2868	145	1100	1399	3394
	crw	–	829	1040	2137	115	815	1062	2606
	issf linear	–	846	1056	1797	108	850	1108	2489
	issf mlp	–	800	983	1745	116	792	1014	2323
	deepssf	–	726	966	1629	109	743	1009	2439
	moveformer	–	724	982	1725	108	725	1016	2395
Hebblewhite and Merrill [141]	don't move	677	5173	7887	14701	376	4861	8172	14525
	crw	600	4121	6142	11103	369	3904	6451	10979
	issf linear	599	4157	6163	10958	360	3935	6502	10767
	issf mlp	607	4136	6169	14319	410	3891	6444	14237
	deepssf	577	4162	6096	11480	374	3969	6535	11221
	moveformer	559	4097	6139	10565	319	3896	6363	10679
Hernández-Pliego et al. [145]	don't move	1718	2721	1842	2551	1584	2717	1900	2351
	crw	1432	2161	1521	2011	1303	2123	1557	1849
	issf linear	1377	1892	1432	2082	1250	1944	1405	1907
	issf mlp	1362	1561	1476	2028	1257	1727	1419	1822
	deepssf	1369	1627	1431	1731	1231	1743	1346	1616
	moveformer	1426	1973	1562	1995	1304	2071	1584	1733
Holland et al. [154]	don't move	130	1937	4181	3027	170	3243	5175	4741
	crw	122	1659	3592	2324	160	2812	4409	3719
	issf linear	124	1637	3438	2007	162	2999	4355	3809
	issf mlp	125	1562	3314	1812	160	2761	4256	4088
	deepssf	116	1501	3166	2168	155	2802	4304	3734
	moveformer	116	1547	3373	2113	153	2613	4119	3452
Egan et al. [84]	don't move	91.5	355	329	338	79.8	354	326	341
	crw	71.4	253	246	255	63.1	255	239	256
	issf linear	70.9	237	240	244	63.4	250	236	253
	issf mlp	70.8	323	226	234	62.9	318	237	254
	deepssf	69.6	211	229	232	63.4	242	239	251
	moveformer	70.7	203	208	233	63.4	203	208	241
Yanco et al. [355]	don't move	–	–	16198	98420	–	–	–	–
	crw	–	–	12750	85361	–	–	–	–
	issf linear	–	–	12803	81344	–	–	–	–
	issf mlp	–	–	12073	80550	–	–	–	–
	deepssf	–	–	12791	85215	–	–	–	–
	moveformer	–	–	11866	74131	–	–	–	–

Continued on next page

Table 10: Energy score results (continued).

Study	Method	ES, Temporal split ↓				ES, Individual split ↓			
		1h	12h	1d	7d	1h	12h	1d	7d
Yanco et al. [355]	don't move	–	–	11411	186905	–	–	10234	175636
	crw	–	–	9086	172691	–	–	8247	161920
	issf linear	–	–	8362	175155	–	–	7814	162830
	issf mlp	–	–	8256	173834	–	–	7554	157535
	deepssf	–	–	9257	142483	–	–	8388	133910
	moveformer	–	–	8092	146321	–	–	7548	158764
Irvine et al. [160]	don't move	3251	18460	21104	21331	2993	17532	18663	24525
	crw	1791	13141	15788	15672	1660	13084	14376	18446
	issf linear	1793	12706	15168	14855	1669	12839	13881	17042
	issf mlp	1950	12706	15167	14814	1799	12941	14106	16898
	deepssf	1812	13130	15213	14915	1687	13544	16580	17041
	moveformer	1746	12913	17227	15475	1686	12846	16880	21205
Irvine et al. [163]	don't move	2724	20015	27696	47859	2860	25814	42057	–
	crw	1310	14075	20517	37511	1100	12667	23248	–
	issf linear	1308	13986	20392	37041	1100	12755	23064	–
	issf mlp	2724	20015	23533	37481	2860	25814	34810	–
	deepssf	1311	13975	21121	39867	1092	12671	23032	–
	moveformer	1287	14190	21099	40779	1080	13357	21995	–
Isbell et al. [165]	don't move	266	1370	1407	1950	188	977	1020	2094
	crw	231	1080	1199	1685	171	802	905	1652
	issf linear	195	1080	1140	1597	174	945	944	1642
	issf mlp	195	927	1214	1700	191	911	1160	1857
	deepssf	180	861	1088	1484	159	914	1344	2370
	moveformer	179	926	1018	1643	147	716	909	1584
Jeglinski et al. [166]	don't move	7933	74597	123847	152248	5721	50160	86768	93622
	crw	7127	62580	101318	122207	5414	43883	74042	76827
	issf linear	6559	59435	95048	113674	4846	41648	72053	73864
	issf mlp	8522	65422	102894	111216	7831	54370	89146	91012
	deepssf	6219	62196	97990	114924	4458	42584	65663	64557
	moveformer	5668	60833	104078	110257	3955	45333	69722	71461
Jennings et al. [169]	don't move	348	2374	1314	2076	399	3179	2592	9726
	crw	333	1879	1290	2882	383	2477	2316	8462
	issf linear	330	1774	1247	2458	394	2598	2326	9095
	issf mlp	365	1810	1353	3725	443	2486	2355	8932
	deepssf	283	1896	1147	2110	452	2703	2345	8820
	moveformer	324	1085	1245	2706	394	1927	2531	8517
Kays et al. [174]	don't move	145	506	803	1044	153	570	919	1141
	crw	112	379	609	794	117	432	710	879
	issf linear	111	347	538	622	118	431	676	852
	issf mlp	109	326	657	632	115	427	793	752
	deepssf	108	318	511	596	115	466	672	832
	moveformer	108	329	536	708	116	423	650	840
Kays et al. [174]	don't move	163	406	434	423	144	398	442	449
	crw	109	289	311	301	96.7	288	318	321
	issf linear	109	276	317	295	97.4	294	348	321
	issf mlp	154	261	414	419	136	310	423	445
	deepssf	109	259	304	296	96.3	327	304	318
	moveformer	105	285	263	289	93.8	257	256	308
Kays et al. [174]	don't move	126	298	399	410	149	361	420	447
	crw	103	246	322	331	110	273	306	331
	issf linear	102	233	296	318	131	296	321	344
	issf mlp	99.3	220	289	335	108	277	324	366
	deepssf	99	232	280	324	111	262	450	339
	moveformer	97.5	248	269	321	105	272	264	313

Continued on next page

Table 10: Energy score results (continued).

Study	Method	ES, Temporal split ↓				ES, Individual split ↓			
		1h	12h	1d	7d	1h	12h	1d	7d
Kays et al. [174]	don't move	151	331	321	389	–	–	–	–
	crw	118	233	229	284	–	–	–	–
	issf linear	116	185	186	235	–	–	–	–
	issf mlp	110	183	181	208	–	–	–	–
	deepssf	113	191	190	281	–	–	–	–
	moveformer	111	195	199	250	–	–	–	–
Kays et al. [174]	don't move	143	244	232	270	127	254	223	248
	crw	108	176	172	194	97.2	188	172	188
	issf linear	106	164	170	194	99.8	172	174	188
	issf mlp	102	155	161	188	93.4	163	164	180
	deepssf	104	161	167	183	93.1	168	169	183
	moveformer	101	165	156	194	89	163	155	189
Stabach et al. [317]	don't move	242	1392	1401	4578	232	1132	1036	2212
	crw	200	1100	1196	4082	194	906	909	2172
	issf linear	198	1062	1173	3874	197	890	947	2570
	issf mlp	198	1055	1169	3816	198	895	939	2750
	deepssf	190	1062	1164	3893	189	883	962	5041
	moveformer	187	964	1150	4325	188	801	879	2243
Serieys et al. [296]	don't move	108	321	406	766	–	–	–	–
	crw	89.8	249	319	594	–	–	–	–
	issf linear	87.6	234	305	531	–	–	–	–
	issf mlp	87	232	305	529	–	–	–	–
	deepssf	86.1	236	312	571	–	–	–	–
	moveformer	87	241	315	634	–	–	–	–
Kidd-Weaver et al. [179]	don't move	444	2900	1369	3828	489	2059	1061	3081
	crw	415	2374	1272	4314	455	1692	1000	3838
	issf linear	414	2366	1265	4800	464	2271	1065	4589
	issf mlp	424	2218	1411	5659	464	1542	1499	5719
	deepssf	402	1971	1227	3677	460	1611	1209	9861
	moveformer	448	1617	1189	4379	462	1047	905	3571
Kleyheeg et al. [182]	don't move	132	1252	644	2458	84.4	420	195	355
	crw	118	1024	558	2193	78	341	207	681
	issf linear	122	1006	566	2234	78.4	349	218	451
	issf mlp	121	855	549	2267	90	344	204	699
	deepssf	116	925	553	2906	75.7	400	184	330
	moveformer	126	690	539	1362	77.6	316	220	812
Kölzsch et al. [184] Kölzsch et al. [185]	don't move	–	–	–	–	405	4513	13188	180050
	crw	–	–	–	–	417	4239	13395	168265
	issf linear	–	–	–	–	421	4091	13513	180228
	issf mlp	–	–	–	–	482	4469	17229	180101
	deepssf	–	–	–	–	382	4462	13103	163205
	moveformer	–	–	–	–	370	3712	13548	150099
Kölzsch et al. [189]	don't move	424	7415	8461	44711	355	7032	6827	28668
	crw	424	6208	7860	37976	361	5794	5913	24589
	issf linear	421	6190	7798	36730	371	5968	6070	26875
	issf mlp	469	6004	8038	36191	407	5727	6422	24322
	deepssf	375	5173	7790	39985	316	5589	6147	26142
	moveformer	376	4234	7779	48554	317	4562	5676	31497
Kölzsch et al. [189]	don't move	427	3505	3007	7204	401	3802	2621	8439
	crw	386	2763	2456	5557	368	3066	2190	7074
	issf linear	391	2502	2353	5186	370	2991	2197	6671
	issf mlp	416	2468	2368	4713	395	2977	2085	5921
	deepssf	363	1890	2329	4256	327	2704	2019	5861
	moveformer	325	2057	2306	5900	320	2057	2092	7651

Continued on next page

Table 10: Energy score results (continued).

Study	Method	ES, Temporal split ↓				ES, Individual split ↓			
		1h	12h	1d	7d	1h	12h	1d	7d
LaPoint et al. [198]	don't move	218	931	1317	1487	218	726	1003	1120
	crw	196	725	969	1071	185	560	747	815
	issf linear	195	690	896	975	181	528	711	741
	issf mlp	206	744	1317	1002	191	602	1003	752
	deepssf	184	661	873	982	176	626	926	753
	moveformer	181	678	878	967	177	533	685	740
Gatt et al. [116]	don't move	4171	37629	61817	145956	7987	58991	97182	139981
	crw	3545	29231	46839	111110	6112	44923	72523	103931
	issf linear	3473	29098	46829	106858	6098	44372	71895	120201
	issf mlp	3397	28630	45686	142649	6003	44224	71556	137129
	deepssf	3043	28709	47440	101629	5452	43569	71650	100570
	moveformer	3014	33125	53812	137562	5011	40375	77315	138842
Lamb et al. [192]	don't move	3125	7787	11885	43521	3513	9134	22657	36404
	crw	2804	6602	10440	39229	3146	7283	19488	31136
	issf linear	2803	6627	10384	38183	3116	7430	19646	29878
	issf mlp	2915	6908	10871	41397	3305	7082	19677	30945
	deepssf	2754	6967	11349	59069	3186	7373	20300	44089
	moveformer	2727	6520	10849	42998	3133	7552	18993	28370
Latham and Boutin [199]	don't move	597	4336	6731	9806	716	4009	5397	5962
	crw	548	3524	5405	7745	639	3219	4245	4577
	issf linear	537	3550	5497	7867	646	3234	4304	4659
	issf mlp	567	3491	5345	7553	649	3250	4292	4743
	deepssf	533	3520	5333	7411	628	3199	4270	5979
	moveformer	526	3522	5172	7184	604	3203	4007	4437
Baker et al. [14]	don't move	1256	8291	10450	34526	1337	9567	13403	59423
	crw	893	6427	8646	28527	929	7288	10429	48277
	issf linear	878	6405	8643	26974	920	7298	10509	45015
	issf mlp	861	6322	8560	27553	897	7218	10579	49793
	deepssf	858	6501	8723	34711	954	8042	10937	54964
	moveformer	789	6006	9178	30460	819	7054	11031	54303
Gutowsky et al. [133]	don't move	768	3766	2477	3148	800	4744	2510	2861
	crw	561	2913	2069	2524	564	3496	2070	2340
	issf linear	482	2955	2046	2498	479	3509	2034	2279
	issf mlp	562	2695	2129	2484	556	3219	2128	2334
	deepssf	510	2201	1894	2146	520	2122	1875	1909
	moveformer	358	1792	1787	2121	335	1823	1744	1987
van Toor et al. [330]	don't move	–	–	–	–	–	483	447	5327
	crw	–	–	–	–	–	537	536	10458
	issf linear	–	–	–	–	–	525	515	10065
	issf mlp	–	–	–	–	–	584	823	21280
	deepssf	–	–	–	–	–	502	521	8568
	moveformer	–	–	–	–	–	416	505	11321
Michelot and Blackwell [217]	don't move	2898	18299	28639	104206	–	–	–	–
	crw	1887	13797	22018	79735	–	–	–	–
	issf linear	1704	13113	21377	79287	–	–	–	–
	issf mlp	1570	12788	20881	76939	–	–	–	–
	deepssf	1828	12647	21132	90770	–	–	–	–
	moveformer	1726	13155	21654	87859	–	–	–	–
McMahon et al. [210]	don't move	18.6	32.7	25	28.9	29.9	59.9	54.2	87.2
	crw	14.3	24.7	19.1	23.2	22.3	45	42.3	64.5
	issf linear	13.4	24.7	18.6	28.9	21.9	45	41.6	87.1
	issf mlp	13.3	24.6	18.3	18.3	21.8	44.5	41.7	74.2
	deepssf	14.3	24.5	19	23.3	22.4	44	41.1	65.2
	moveformer	13.5	22.9	19.3	22.9	22.5	50	41.1	66.3

Continued on next page

Table 10: Energy score results (continued).

Study	Method	ES, Temporal split ↓				ES, Individual split ↓			
		1h	12h	1d	7d	1h	12h	1d	7d
Medici et al. [211]	don't move	204	937	975	998	210	1056	1097	1059
	crw	173	693	749	764	177	773	842	806
	issf linear	170	677	736	752	174	765	831	795
	issf mlp	171	931	732	773	176	1049	822	813
	deepssf	156	619	718	737	163	749	843	814
	moveformer	153	575	636	693	158	635	681	696
Mertes et al. [214]	don't move	157	381	289	356	117	339	199	219
	crw	129	278	222	271	101	256	162	178
	issf linear	126	267	216	264	99.5	239	155	169
	issf mlp	126	275	214	258	101	239	156	172
	deepssf	121	266	220	264	91.7	225	174	194
	moveformer	119	233	205	267	92.3	194	146	177
Potts et al. [264]	don't move	–	905	993	1219	–	860	922	1245
	crw	–	695	790	935	–	661	738	964
	issf linear	–	642	731	859	–	610	686	894
	issf mlp	–	630	738	847	–	596	690	879
	deepssf	–	631	736	863	–	601	690	909
	moveformer	–	595	708	864	–	558	637	875
Morato et al. [220]	don't move	127	940	1454	3022	79.9	617	1059	3118
	crw	120	766	1147	2247	81.8	538	867	2456
	issf linear	119	762	1145	2293	79.6	541	865	2511
	issf mlp	119	749	1118	3002	83.8	555	863	3103
	deepssf	117	766	1164	2143	75.4	526	855	2509
	moveformer	106	788	1178	2026	69.6	602	890	2364
Morganti et al. [223]	don't move	55.9	144	187	270	–	–	–	–
	crw	42	111	147	205	–	–	–	–
	issf linear	41.6	108	147	204	–	–	–	–
	issf mlp	41.5	106	141	188	–	–	–	–
	deepssf	41.4	111	143	161	–	–	–	–
	moveformer	40.7	107	141	205	–	–	–	–
Aikens et al. [9]	don't move	–	1292	1910	5446	–	1170	1613	4149
	crw	–	985	1474	4307	–	891	1242	3331
	issf linear	–	988	1497	4315	–	893	1256	3393
	issf mlp	–	977	1613	4195	–	881	1347	3364
	deepssf	–	963	1465	4201	–	882	1263	3617
	moveformer	–	963	1460	4438	–	848	1218	3681
Kölzsch et al. [184]	don't move	–	–	–	–	125	21658	48963	417997
	crw	–	–	–	–	209	21224	48021	355670
	issf linear	–	–	–	–	170	21221	47697	318805
	issf mlp	–	–	–	–	325	23339	56157	331584
	deepssf	–	–	–	–	147	21709	46260	323761
	moveformer	–	–	–	–	196	20544	60735	339856
Benitez et al. [29]	don't move	468	3482	5118	13056	419	3299	5061	12616
	crw	347	2580	3911	10375	313	2437	3846	9880
	issf linear	346	2592	3943	10258	309	2455	3882	9887
	issf mlp	342	2543	3859	9999	305	2363	3773	11009
	deepssf	337	2580	3889	9735	305	2439	3898	13372
	moveformer	301	2458	3915	11268	270	2333	3870	11173
Naniwadekar et al. [232]	don't move	251	577	604	1450	245	353	326	380
	crw	199	433	487	1296	187	264	261	555
	issf linear	196	443	483	1250	188	298	338	683
	issf mlp	191	479	478	1208	190	289	451	622
	deepssf	194	424	461	1180	191	339	334	373
	moveformer	196	444	522	1355	214	443	232	493

Continued on next page

Table 10: Energy score results (continued).

Study	Method	ES, Temporal split ↓				ES, Individual split ↓			
		1h	12h	1d	7d	1h	12h	1d	7d
Naniwadekar et al. [232]	don't move	1160	2357	1743	2022	–	–	–	–
	crw	934	1667	1383	1579	–	–	–	–
	issf linear	931	1698	1343	1437	–	–	–	–
	issf mlp	875	1366	1209	1457	–	–	–	–
	deepssf	935	1621	1244	1289	–	–	–	–
	moveformer	884	1346	1377	1713	–	–	–	–
Newman et al. [237]	don't move	96	454	514	801	92.1	460	455	499
	crw	82.4	352	408	608	77.8	339	356	390
	issf linear	81.9	336	400	608	80.7	339	354	396
	issf mlp	82.1	333	399	608	77	331	355	390
	deepssf	81.6	341	397	616	76.7	333	354	405
	moveformer	81	342	409	629	77.4	292	303	394
Padget et al. [244]	don't move	5262	29011	40293	135491	6374	33488	49407	161104
	crw	4176	23694	33348	102634	5105	27116	40478	117479
	issf linear	4183	23299	32110	97593	5104	26621	39491	114606
	issf mlp	4116	23057	32437	96379	5063	26272	39554	110477
	deepssf	4067	25772	32702	84768	4937	28950	39094	100360
	moveformer	3916	24327	33376	121848	4798	27113	40288	121353
Mestre et al. [216]	don't move	480	2667	1469	2360	886	1939	1073	1480
	crw	448	2159	1233	1962	878	1712	940	1367
	issf linear	431	1971	1235	2120	893	2182	977	1251
	issf mlp	518	1876	1199	1932	983	2615	989	1331
	deepssf	433	2152	1173	1984	855	2441	1304	1396
	moveformer	435	1716	1221	1866	860	1654	946	1331
Pezzo et al. [253]	don't move	4376	22543	29582	–	5308	38295	46250	–
	crw	3740	19973	27563	–	4526	30188	37519	–
	issf linear	3746	19376	29185	–	4519	29854	39911	–
	issf mlp	3870	20498	31098	–	4566	30838	42171	–
	deepssf	3669	19315	24201	–	4462	30197	37931	–
	moveformer	3172	26565	42848	–	4011	31262	55931	–
Piironen et al. [255]	don't move	260	3139	4710	104642	245	1945	3585	83899
	crw	233	2810	4972	111662	217	1877	4090	96556
	issf linear	234	2762	5355	106671	217	2059	4485	90096
	issf mlp	238	2760	5743	149005	220	1943	5124	126453
	deepssf	231	2705	4759	146902	213	1748	4066	140524
	moveformer	224	2235	5285	152317	209	1912	5086	127755
Pillco Huarcaya et al. [256]	don't move	92	951	1362	2568	–	–	–	–
	crw	80.6	750	1065	2166	–	–	–	–
	issf linear	81.6	743	1034	2091	–	–	–	–
	issf mlp	79.6	752	1037	1951	–	–	–	–
	deepssf	78.4	758	1072	1457	–	–	–	–
	moveformer	81.7	773	1113	2450	–	–	–	–
Pokrovsky et al. [260]	don't move	450	7585	17409	125350	427	6721	14448	78103
	crw	459	7951	17749	117720	428	7348	15564	80299
	issf linear	470	7667	16459	108337	438	6812	14377	78585
	issf mlp	528	8771	20690	135682	492	7808	18540	100959
	deepssf	415	7092	15990	91810	375	6222	13403	82680
	moveformer	475	8142	15968	98510	449	6553	15183	117513
Porteus et al. [261]	don't move	149	354	314	325	152	414	283	347
	crw	129	269	247	252	129	313	219	268
	issf linear	127	267	240	245	126	317	218	268
	issf mlp	129	267	242	247	127	308	218	262
	deepssf	116	266	245	246	112	344	241	281
	moveformer	119	222	217	230	121	217	200	246

Continued on next page

Table 10: Energy score results (continued).

Study	Method	ES, Temporal split ↓				ES, Individual split ↓			
		1h	12h	1d	7d	1h	12h	1d	7d
Power et al. [268]	don't move	–	–	1474	2397	–	–	2286	5216
	crw	–	–	1213	1886	–	–	1845	4026
	issf linear	–	–	1139	1590	–	–	1943	3790
	issf mlp	–	–	1147	1481	–	–	1817	3826
	deepssf	–	–	1204	1516	–	–	1842	3838
	moveformer	–	–	1179	1617	–	–	1912	4448
Power et al. [267]	don't move	–	721	1024	2156	–	–	–	–
	crw	–	651	959	2442	–	–	–	–
	issf linear	–	622	944	1875	–	–	–	–
	issf mlp	–	659	931	2387	–	–	–	–
	deepssf	–	619	861	1870	–	–	–	–
	moveformer	–	620	973	2943	–	–	–	–
Prugh et al. [270]	don't move	–	1599	1866	2020	–	1529	1751	2012
	crw	–	1203	1417	1519	–	1152	1325	1514
	issf linear	–	1203	1371	1460	–	1188	1324	1504
	issf mlp	–	1186	1696	1447	–	1166	1557	1505
	deepssf	–	1174	1342	1432	–	1170	1341	1567
	moveformer	–	1069	1211	1340	–	1062	1187	1398
Prugh et al. [270]	don't move	–	1544	2346	3554	–	1794	2773	3552
	crw	–	1180	1787	2657	–	1376	2137	2667
	issf linear	–	1164	1747	2591	–	1451	2236	2710
	issf mlp	–	1190	2211	3158	–	1458	2672	3273
	deepssf	–	1140	1668	2522	–	1337	2098	2813
	moveformer	–	1134	1685	2455	–	1319	1959	2475
Anderson et al. [12]	don't move	1496	13181	16022	40419	3381	15738	24221	89569
	crw	1506	11651	15103	38866	3135	14089	22226	82579
	issf linear	1461	10776	13983	41822	2980	13138	20769	84920
	issf mlp	1555	11454	15891	45585	2972	13581	21320	77892
	deepssf	1297	10615	13799	35720	2813	12655	20959	74779
	moveformer	1286	9985	13915	38517	2772	12878	21313	82189
Ronconi and Shlepr [275]	don't move	3476	14286	10302	13435	3147	14146	10426	11578
	crw	3313	11564	9030	11541	3033	11631	9016	9821
	issf linear	3062	10180	8899	11339	2878	9821	8613	9580
	issf mlp	3570	9685	9454	11052	3530	9710	9085	9805
	deepssf	2982	7219	8204	12024	2707	6845	8070	10022
	moveformer	2680	7368	7769	10896	2787	7421	7461	9103
Boardman et al. [34]	don't move	1237	3678	1014	1942	975	3030	1076	1792
	crw	1100	2973	932	1682	908	2471	996	1603
	issf linear	1069	1699	849	1567	927	2065	893	1385
	issf mlp	1140	836	1025	1588	1145	1052	1144	1482
	deepssf	974	1030	876	1678	756	1263	945	1572
	moveformer	961	1624	837	1680	868	2169	928	1701
Hennig et al. [143]	don't move	–	1618	1685	2389	–	1540	1698	2452
	crw	–	1201	1261	1795	–	1149	1281	1888
	issf linear	–	1182	1257	1787	–	1159	1291	1906
	issf mlp	–	1176	1248	1776	–	1150	1282	1893
	deepssf	–	1168	1251	1726	–	1156	1295	2021
	moveformer	–	1012	1165	1671	–	1011	1176	1808
Schloesing et al. [284]	don't move	1002	1950	1105	2784	1542	2763	1660	2518
	crw	930	1736	1031	2395	1402	2240	1445	2120
	issf linear	941	1757	1019	2784	1387	2311	1445	2518
	issf mlp	1011	1551	1131	2768	1416	2422	1575	2504
	deepssf	882	1690	971	2330	1428	2747	1538	2062
	moveformer	935	1620	953	2572	1436	2293	1327	2374

Continued on next page

Table 10: Energy score results (continued).

Study	Method	ES, Temporal split ↓				ES, Individual split ↓			
		1h	12h	1d	7d	1h	12h	1d	7d
Schweitzer et al. [287]	don't move	296	5325	9166	20655	–	–	–	–
	crw	327	5068	8615	18521	–	–	–	–
	issf linear	323	5083	8657	17890	–	–	–	–
	issf mlp	416	5283	9069	18600	–	–	–	–
	deepssf	295	6259	9911	18577	–	–	–	–
	moveformer	329	5151	7774	21592	–	–	–	–
Schwemmer et al. [289]	don't move	514	10626	20471	196612	510	9545	17654	217026
	crw	524	12667	24973	201233	516	11837	22824	221971
	issf linear	552	13345	23575	179645	561	12709	22364	212204
	issf mlp	577	24789	51157	306610	591	23672	50636	359234
	deepssf	481	12145	24804	279114	533	13532	37551	176767
	moveformer	539	23740	31287	234637	577	13933	24884	157809
Serieys et al. [295]	don't move	264	876	993	1196	371	1570	1932	1750
	crw	211	663	758	863	297	1168	1405	1244
	issf linear	209	650	767	840	291	1184	1494	1349
	issf mlp	208	719	993	1186	287	1336	1932	1739
	deepssf	209	654	756	857	284	1120	1328	1232
	moveformer	205	628	712	768	281	1103	1272	1208
Serieys et al. [292]	don't move	180	864	1078	1014	211	960	1121	1131
	crw	147	645	797	746	171	722	847	841
	issf linear	145	631	787	732	170	750	945	852
	issf mlp	146	638	773	738	166	713	853	832
	deepssf	141	689	817	775	169	874	1032	901
	moveformer	138	576	661	664	158	635	685	756
Serieys et al. [292]	don't move	172	499	526	420	–	–	–	–
	crw	139	360	380	301	–	–	–	–
	issf linear	132	306	304	292	–	–	–	–
	issf mlp	132	276	291	419	–	–	–	–
	deepssf	122	284	294	251	–	–	–	–
	moveformer	128	315	324	301	–	–	–	–
Serieys et al. [293]	don't move	157	1101	1789	4090	161	639	721	1020
	crw	138	927	1476	3172	140	555	720	1321
	issf linear	136	907	1404	2856	134	561	753	1831
	issf mlp	133	917	1455	2879	134	582	835	1972
	deepssf	136	953	1462	3054	135	586	726	1528
	moveformer	129	886	1425	2826	130	591	607	999
Setyawan et al. [298]	don't move	771	2902	3212	7373	1978	10325	11631	15687
	crw	625	2571	2873	6451	1458	8100	9094	11366
	issf linear	571	2310	2634	5752	1329	8014	9108	12840
	issf mlp	561	2271	2655	5615	1331	8115	9055	15216
	deepssf	645	2311	2567	5702	1435	9235	9660	15202
	moveformer	590	2263	2695	5907	1566	7974	9221	11503
Yanco et al. [355]	don't move	660	3728	4160	10720	339	2183	2000	5597
	crw	582	3070	3662	10004	314	1783	1721	6135
	issf linear	569	2927	3545	9520	320	1802	1716	5985
	issf mlp	574	2849	3530	13272	318	2253	2014	9266
	deepssf	555	2746	3394	7816	285	2737	3287	16170
	moveformer	573	2769	3411	14107	326	1978	1811	7109
Thaker et al. [320]	don't move	419	2924	3861	9946	319	2281	3185	8799
	crw	298	2090	2817	7373	230	1661	2351	6644
	issf linear	294	2084	2809	7142	229	1656	2354	6934
	issf mlp	294	2097	2808	6776	231	1660	2354	6886
	deepssf	277	2012	2778	6721	228	1667	2371	8356
	moveformer	268	2055	2975	8430	215	1665	2566	7485

Continued on next page

Table 10: Energy score results (continued).

Study	Method	ES, Temporal split ↓				ES, Individual split ↓			
		1h	12h	1d	7d	1h	12h	1d	7d
Spiegel et al. [312]	don't move	5944	23875	39390	74083	8158	28338	39893	45743
	crw	5581	20812	31414	57064	7276	23555	30726	34846
	issf linear	5495	20655	30763	55020	7159	22935	29236	32358
	issf mlp	6528	21657	30805	52314	7738	23220	28017	30671
	deepssf	5305	19448	31601	58567	6884	21318	28914	37740
	moveformer	5018	19711	32400	63549	6381	22163	28378	36685
Spiegel et al. [312]	don't move	7382	19596	28669	37143	4522	15600	23294	28064
	crw	6505	16756	22967	30665	4337	13487	18440	22256
	issf linear	6406	16294	21991	27772	4570	13815	18807	23487
	issf mlp	6955	16592	20980	25097	5683	16958	22023	26421
	deepssf	6007	15844	21578	27596	4038	24269	30279	35623
	moveformer	5809	16171	22970	30213	3915	12393	18743	20181
Spiegel et al. [313]	don't move	7765	24048	37006	57605	6965	21582	36137	55551
	crw	6958	20486	29412	44750	6312	18631	29146	43773
	issf linear	6714	19062	27331	39248	6280	18973	30803	43798
	issf mlp	7548	19114	26032	37478	7169	19764	31633	45615
	deepssf	6823	19238	28817	39075	6233	17943	28478	48622
	moveformer	6174	19381	29366	46919	5973	17814	29438	47774
Todd et al. [323]	don't move	21.4	63.9	40.4	38.3	28.5	164	124	30.7
	crw	17	46.1	31.7	35	23.2	130	96.9	26.7
	issf linear	17.4	42.7	26.6	33.5	24.9	182	122	27.2
	issf mlp	16.6	38.5	31.8	43.1	23.3	229	96.7	287
	deepssf	16.9	47.2	32.8	15.2	22.9	134	95.8	215
	moveformer	17.8	38.5	37.5	39.5	24.1	137	104	43.3
Toivonen et al. [325]	don't move	16.1	650	663	852	–	1071	1275	2368
	crw	51.1	478	515	658	–	830	1019	1968
	issf linear	51.1	474	520	674	–	819	1034	1912
	issf mlp	51.1	473	513	641	–	813	985	1900
	deepssf	20.2	470	518	670	–	832	1002	1920
	moveformer	51.1	440	466	662	–	820	1008	1834
Patterson et al. [246]	don't move	–	3964	6363	14732	–	835	827	–
	crw	–	3197	4972	10918	–	946	1190	–
	issf linear	–	3167	4874	10722	–	938	1158	–
	issf mlp	–	3126	4711	14732	–	1174	1359	–
	deepssf	–	3176	4936	11263	–	842	1148	–
	moveformer	–	3262	5479	9946	–	972	1779	–
Wagner et al. [339]	don't move	125	170	190	–	–	–	–	–
	crw	96.6	119	152	–	–	–	–	–
	issf linear	95.2	78.2	144	–	–	–	–	–
	issf mlp	92	84.6	141	–	–	–	–	–
	deepssf	98.6	82.7	153	–	–	–	–	–
	moveformer	88.2	106	143	–	–	–	–	–
Wall et al. [341]	don't move	288	2201	2208	6412	–	–	–	–
	crw	206	1673	1619	4277	–	–	–	–
	issf linear	218	1566	1587	4024	–	–	–	–
	issf mlp	201	1547	1592	4269	–	–	–	–
	deepssf	197	1607	1647	4188	–	–	–	–
	moveformer	175	1542	1906	7481	–	–	–	–
Warret Rodrigues and Roth [344]	don't move	975	2245	2189	2273	982	2378	2776	2608
	crw	796	1690	1643	1633	774	1741	2029	1871
	issf linear	784	1598	1559	1494	770	1789	2033	1810
	issf mlp	818	2028	1662	2273	811	2205	2197	2608
	deepssf	750	1540	1523	1456	763	1635	1978	1899
	moveformer	733	1378	1446	1571	794	1605	1768	1787

Continued on next page

Table 10: Energy score results (continued).


Study	Method	ES, Temporal split ↓				ES, Individual split ↓			
		1h	12h	1d	7d	1h	12h	1d	7d
Warret Rodrigues and Roth [344]	don't move	676	2540	3729	4377	830	2437	2538	2528
	crw	593	2052	2921	3326	711	1815	1868	1869
	issf linear	588	2159	2994	3492	689	1721	1704	1705
	issf mlp	619	2093	3062	3406	696	1634	1685	1643
	deepssf	590	2006	2786	3048	680	1664	1893	1637
	moveformer	566	1877	2546	3065	645	1517	1617	1784
Warwick-Evans et al. [345]	don't move	7271	52558	90473	73525	9733	64352	90686	122624
	crw	6499	43669	71126	60020	8441	52375	71783	98686
	issf linear	5907	41072	59664	71314	7857	49231	62384	97933
	issf mlp	7002	39171	45982	73525	8458	52348	63781	122623
	deepssf	5582	41393	65891	59772	7614	49717	65980	99022
	moveformer	4955	34198	42965	64683	7035	53870	70076	104950
Wicks et al. [349]	don't move	–	535	835	2422	–	–	–	–
	crw	–	443	695	1951	–	–	–	–
	issf linear	–	444	686	1764	–	–	–	–
	issf mlp	–	442	698	1707	–	–	–	–
	deepssf	–	425	662	1949	–	–	–	–
	moveformer	–	434	780	1923	–	–	–	–
EURODEER: Alta Val di Non (<i>Cervus elaphus</i>)	don't move	98.7	584	462	684	–	–	–	–
	crw	81.4	433	362	535	–	–	–	–
	issf linear	80.9	424	355	520	–	–	–	–
	issf mlp	79.7	392	352	514	–	–	–	–
	deepssf	79.1	377	352	508	–	–	–	–
	moveformer	74.8	288	329	513	–	–	–	–
EURODEER: Bondone (<i>Capreolus capreolus</i>)	don't move	52.3	250	212	367	–	–	–	–
	crw	42.4	181	158	303	–	–	–	–
	issf linear	43.4	179	159	315	–	–	–	–
	issf mlp	42.2	175	158	301	–	–	–	–
	deepssf	43.4	171	161	308	–	–	–	–
	moveformer	41.5	142	148	291	–	–	–	–
EURODEER: Val Rendena (<i>Cervus elaphus</i>)	don't move	–	239	256	357	–	–	–	–
	crw	–	182	200	277	–	–	–	–
	issf linear	–	187	198	267	–	–	–	–
	issf mlp	–	178	195	266	–	–	–	–
	deepssf	–	181	197	270	–	–	–	–
	moveformer	–	158	187	270	–	–	–	–
d'Entremont et al. [72]	don't move	6776	41347	54513	61573	7349	51197	64690	54927
	crw	5784	31337	40833	46754	6126	39427	49138	40988
	issf linear	5841	29193	36462	45079	6207	38599	45114	41468
	issf mlp	5742	27334	33526	52714	6087	36630	42956	46648
	deepssf	5437	27802	32051	46754	5760	35824	42235	41864
	moveformer	5632	30090	38185	47647	5847	38612	46287	39377
Ágústisdóttir and Þórisson [360]	don't move	93.1	1707	2787	8333	162	1553	2474	7035
	crw	78.7	1342	2188	6441	140	1205	1925	5654
	issf linear	79.8	1366	2263	6841	143	1231	2001	5799
	issf mlp	78.9	1337	2179	6455	138	1196	1875	5622
	deepssf	78.2	1343	2153	6495	138	1212	1875	5710
	moveformer	82.7	1287	2146	7202	138	1196	1909	5903
Škrábal et al. [308]	don't move	8772	46423	97432	311740	8522	53622	101891	502389
	crw	7337	38241	74771	232074	6961	43427	73605	366348
	issf linear	6886	35309	63346	137801	6672	42596	64294	184205
	issf mlp	6961	37504	66286	134687	6392	41563	64997	135069
	deepssf	6384	29905	62736	131940	6418	36298	70729	203753
	moveformer	4658	29057	61781	183790	5328	31420	59382	251480

Continued on next page

Table 10: Energy score results (continued).

Study	Method	ES, Temporal split ↓				ES, Individual split ↓			
		1h	12h	1d	7d	1h	12h	1d	7d
van Toor et al. [331]	don't move	–	–	–	–	78.2	929	748	1576
	crw	–	–	–	–	78.6	862	825	1644
	issf linear	–	–	–	–	77.2	765	696	1605
	issf mlp	–	–	–	–	84.7	894	1027	1667
	deepssf	–	–	–	–	77	897	875	1795
	moveformer	–	–	–	–	77.9	579	835	2256
van Toor et al. [334]	don't move	–	–	–	–	111	688	3297	49638
	crw	–	–	–	–	118	723	3859	59314
	issf linear	–	–	–	–	108	636	3711	66730
	issf mlp	–	–	–	–	177	717	5105	71088
	deepssf	–	–	–	–	171	678	3424	220520
	moveformer	–	–	–	–	131	665	4032	82869
Kölzsch et al. [186]	don't move	9499	2030	6081	106578	10065	1327	3777	85952
	crw	8489	1724	6220	110392	9140	1193	4234	94397
	issf linear	6823	1685	6399	103331	7644	1231	4596	88331
	issf mlp	7633	1689	6883	144270	7959	1263	5183	136868
	deepssf	9120	1637	5460	78603	9896	1147	5106	69450
	moveformer	8780	1608	5630	134593	7415	1071	3907	146546
Yanco et al. [355]	don't move	750	3495	3432	36654	708	3755	3879	27280
	crw	705	2831	3199	38501	672	3082	3555	29470
	issf linear	720	2774	3305	39410	691	3036	3809	30053
	issf mlp	735	2721	3337	51747	707	2943	3745	37218
	deepssf	667	2671	3124	37624	637	2868	3496	29063
	moveformer	694	2452	3276	41766	622	2609	3563	29954

G.2 Per-study MoveFormer

**Figure 5:** Training one MoveFormer model per study per time horizon vs. training one MoveFormer model across all studies for each time horizon. The latter is the published default that is used in our main experiments.

MoveFormer is the only method we benchmarked that is designed to train across studies. For comparison with the other methods, which train one model per-study, we also experimented with per-study MoveFormer training. On average, cross-study training was beneficial. See Fig. 5.

G.3 Compute and storage requirements

We trained 3,360 study-specific models across 147 studies, each evaluating one method at one timescale, and four shared MoveFormer training runs that were trained across all studies at each timescale as in the original implementation. We ran experiments on a heterogeneous SLURM cluster.

All jobs were allocated 16 CPU cores and 64 GB RAM, however this is more than required for most experiments. Baseline and iSSF variants ran CPU-only, whereas deepssf and moveformer used a single GPU on either A100, V100, L40S, H100, H200, RTX3090/3080, A6000, or RTX6000 Ada nodes. Unless noted otherwise, the numbers below are median wall-clock times over completed tasks. Note that jobs were assigned to GPU nodes randomly, and run time is affected by the size of the study, so the times below may not be indicative of run time on a particular GPU generally.

- The “don’t move” baseline took 1.1 min, and the correlated random walk baseline took 1.9 min.
- Linear iSSF took 6.9 min, and MLP iSSF took 11.2 min.
- deepSSF took 11.0 min overall. Medians by GPU were 8.4 min on L40S, 9.7 min on H200, 10.7 min on H100, 12.1 min on A100, 13.1 min on A6000, and 17.3 min on V100.
- Training the four shared MoveFormer models took 7.3 h (1 h timescale), 13.4 h (12 h), 5.1 h (1 d), and 6.0 h (7 d), each on one A100, for a total of 31.7 GPU-hours. Subsequent per-study scoring took 4.6 min overall: 4.7 min on A100 and 4.4 min on L40S.
- The 90th-percentile runtimes were 28.5 min for linear iSSF, 36.0 min for MLP iSSF, 107.1 min for deepSSF, and 55.7 min for MoveFormer scoring.

The full dataset, including all covariates for all timescales for all studies, totals 7 terabytes using 32-bit floating point representations. We foresee that most users of the benchmark will not require all of this data. For instance, they may train models with a subset of covariates and/or for a subset of studies. Our Python package enables download of arbitrary subsets of covariates, studies, and timescales directly from a hosted archive. For covariates or timescales not included in our experiments, the package also enables downloading arbitrary layers, timescales, and studies (including custom studies that a user can import themselves), via Google Earth Engine. For reference, all covariates for all timescales of MOVEBENCH-Mini totals approximately 500 gigabytes, and the “lite” set of covariates (Sec. 4) for all timescales of MOVEBENCH-Mini totals approximately 20 gigabytes. Quantized versions of all covariates will also be released for additional space savings.