

Convergence of Practical Muon with Finite Newton–Schulz Iterations and Nesterov Momentum

Hanyang Peng, Hui Wang, Yue Yu
Pengcheng Laboratory, Shenzhen, China
{penghy, wangh06, yuy}@pcl.ac.cn

Abstract

Practical Muon maintains momentum and performs a small, fixed number of Newton–Schulz iterations separately for each parameter matrix, often with a Nesterov correction. We analyze these layer-wise finite-step updates jointly on a coupled nonconvex objective, rather than replacing them by exact polar factors or one global orthogonalization. Under gradient-dependent $(\mathcal{L}_0, \mathcal{L}_1, q)$ -smoothness and conditionally unbiased stochastic gradients with bounded layer-wise variance, we establish an $\mathcal{O}(T^{-1/4})$ bound on the expected average Frobenius gradient norm. The analysis retains the Nesterov recursion and requires neither bounded stochastic gradients, symmetric noise, nor a uniform positive lower bound on the nonzero output singular values. Its constants contain no explicit matrix-dimension or rank factors when the number of blocks and problem constants are fixed. The proof follows a descent inequality and a decomposition of the momentum tracking error into initialization, noise, and drift. For the original five-step quintic, we verify the required scalar-map bounds analytically; the result also allows step-dependent coefficients satisfying the same bounds. A complementary nuclear-norm result quantifies rank dependence under a stronger spectral condition. The vanishing rate uses coupled learning-rate and momentum schedules, including the standard single-coefficient Nesterov rule.

1 Introduction

Muon orthogonalizes a momentum-based update separately for each matrix-valued parameter. The reference implementation maintains a momentum buffer for each parameter matrix, applies Frobenius normalization and a small number of quintic Newton–Schulz iterations to that matrix, and includes an optional Nesterov correction (Jordan, 2024). These three features—layer-wise updates, finite orthogonalization, and Nesterov momentum—all matter for convergence analysis. A single global normalization is not the layer-wise operation, a finite polynomial need not return an exact polar factor, and the Nesterov correction changes the update’s noise and gradient-tracking error.

Our analysis preserves this layer-wise structure. Each matrix block has its own momentum, normalization, and orthogonalization, and the blocks may have different dimensions and noise levels. Their updates nevertheless act on one coupled objective: all stochastic gradient blocks are evaluated at the same iterate, and changing one block may change the gradients in other blocks. We therefore use joint smoothness to control simultaneous updates, while retaining layer-specific variance bounds and tracking errors. The analysis does not assume a separable loss or independent noise across layers. Here a “layer” indexes a parameter matrix; several such matrices may belong to the same architectural layer.

Existing analyses address different aspects of Muon. Pethick et al. (2025) place non-Nesterov Muon in an exact-LMO framework and develop layer-wise norm choices. Riabinin et al. (2025) explicitly analyze layer-wise LMOs under block generalized smoothness. Shen et al. (2025) study

exact-polar Muon and its matrix geometry, whereas Kim and Oh (2025) analyze finite Newton–Schulz iterations through polar-approximation bounds. Choudhury et al. (2026) already combine Nesterov momentum with inexact polar decomposition and allow gradient-dependent heavy-tailed noise. Li and Tsuchiya (2026) exploit finite-iteration smoothing through online-to-nonconvex conversion. Table 1 compares the algorithms, assumptions, and guarantees in these results. Our contribution is not any one of these ingredients in isolation, but a direct analysis of their combination: layer-wise Frobenius-normalized finite polynomial updates with Nesterov momentum, under joint gradient-dependent smoothness and bounded layer-wise conditional variance, yielding an average Frobenius stationarity bound without explicit matrix-size or rank factors. The assumptions are not uniformly weaker than those of all prior work; in particular, our bounded-variance model does not cover the heavier-tailed regimes of Choudhury et al. (2026).

Table 1: Comparison of representative convergence analyses relevant to practical Muon. NS denotes Newton–Schulz iteration; LMO denotes a linear minimization oracle. Entries describe the cited theoretical results, not every implementation studied in each paper.

Work	Block structure	Orthogonalization in theory	Nesterov	Smoothness and noise	Stationarity and dimension factors
Pethick et al. (2025) (Scion/uSCG)	General-norm theorem; layer-wise design	Exact LMO; polar factor for Muon	No ^a	Norm smoothness; bounded Euclidean variance	Dual-norm gradient; geometry-dependent constants
Riabinin et al. (2025) (Gluon)	Explicitly layer-wise	Exact block LMOs	No	Block generalized smoothness; block dual-norm variance	Weighted block dual norms; norm-specific constants
Shen et al. (2025)	Single matrix	Exact SVD-polar factor	No	Frobenius or spectral smoothness; bounded Frobenius variance	Frobenius/nuclear gradient; rank factors in stochastic bounds
Kim and Oh (2025)	Single matrix	Finite Taylor NS; polar-error analysis	No	Spectral smoothness; bounded Frobenius variance	Average nuclear gradient; explicit rank factors
Choudhury et al. (2026)	Single matrix	Inexact polar, including NS; relative alignment bound	Yes	Frobenius L -smoothness; gradient-dependent α -moment noise, $1 < \alpha \leq 2$	Best-iterate Frobenius gradient; d_0 -dependent constants ^b
Li and Tsuchiya (2026)	Single-matrix online learner	Finite Taylor NS; fixed normalization	No	Nonsmooth objectives allowed; moment bounds and almost-sure operator-norm gradient bound	Online-to-nonconvex stationarity; rank-dependent bounds
This work Theorem 3.4	Explicitly layer-wise	Fixed finite NS; includes the practical five-step quintic	Yes	Joint $(\mathcal{L}_0, \mathcal{L}_1, q)$ -smoothness; layer-wise conditional Frobenius variance	Average Frobenius gradient, $\mathcal{O}(T^{-1/4})$; no size/rank factors^c

^aThe Muon specialization in the uSCG theory uses non-Nesterov momentum; Scion also develops layer-wise norm choices.

^b $d_0 = \min\{m, n\}$. ^cWith fixed block count, NS depth and coefficients, and problem constants; the vanishing rate uses coupled momentum and learning-rate schedules. “Single matrix” describes the theorem’s formulation, not an impossibility of extending it. The norms, noise models, and output criteria differ, so the rows are not ordered by a common strength of assumptions. For our implementation scope, see Remark 4.5.

The main obstacle is not only the approximation error relative to a polar factor. For a fixed number J of iterations, let ϕ_J be the composite scalar polynomial acting on normalized singular values. Since $\phi_J(s) \rightarrow 0$ as $s \downarrow 0$, a uniform positive lower bound on all nonzero output singular values is unavailable over arbitrary spectra. However, the ratio $\phi_J(s)/s$ can remain bounded above and away from zero. Together with Frobenius normalization of each block, this yields descent and update-energy bounds that do not introduce a rank factor. We use these bounds without replacing the finite map by an exact polar direction.

The resulting theorem controls the expected average Frobenius gradient norm along the original stochastic iterates. Its proof retains the same three parts of each block’s Nesterov tracking error: the initial momentum bias, the accumulated stochastic noise, and the drift of the true gradient under joint updates. Layer-specific noise levels remain visible through $S_F = \sum_{l=1}^L \sigma_{(l)}$, rather than through a matrix-rank conversion. The $\mathcal{O}(T^{-1/4})$ rate follows by coupling the learning rate and momentum parameters. Taking the same coefficient in the momentum and Nesterov correction is included, not removed for the analysis. For the original quintic, the scalar-map requirements hold for every fixed $J \geq 1$, in particular for $J = 5$; they can also be checked for schedules using different coefficients at different steps.

Section 3 develops these results within one proof framework. It first gives a rank-sensitive nuclear-norm bound under a positive output singular-value floor, then removes that condition by using the finite map and measuring stationarity in Frobenius norm. The latter result uses neither a bounded-gradient nor a noise-symmetry assumption. Dimension independence here concerns the optimization bound with fixed block count and problem constants; it is not a claim about how those constants change across neural-network architectures.

2 Preliminary

We minimize $F(X) = \mathbb{E}_\xi[f(X; \xi)]$ over matrix blocks $X = (X^{(1)}, \dots, X^{(L)})$. Algorithm 1 uses $M_t^{(l)}$ for the first-order momentum and $N_t^{(l)}$ for the direction passed to the orthogonalization routine. The setting $\beta_2 = 1$ gives ordinary momentum, whereas $\beta_2 = \beta_1 = \beta$ gives

$$N_t^{(l)} = \beta M_t^{(l)} + (1 - \beta) G_t^{(l)}, \quad (1)$$

the single-coefficient Nesterov correction used in the reference implementation.¹ The routine may be inexact; its finite Newton–Schulz specialization is specified in Section 3.1.

Layer-wise refers to the separate matrix operations in Algorithm 1: the normalization in each call to orthogonalization uses that block’s $N_t^{(l)}$, not the norm of the full momentum tuple. All $G_t^{(l)}$ are evaluated at X_t before the joint iterate X_{t+1} is formed. This is not cyclic block-coordinate descent, and no cross-block derivatives are set to zero. The common sample ξ_t may also correlate the noise across blocks.

Here “practical” refers to retaining the layer-wise matrix updates, finite Newton–Schulz map, and Nesterov correction, rather than analyzing an exact-polar or globally orthogonalized surrogate. The mathematical iteration is studied in exact arithmetic, without decoupled weight decay or a separate optimizer for other parameter groups. Remark 4.5 treats normalization stabilizers and fixed output scaling separately.

Proposition 2.1 (Spectral Structure of Inexact Orthogonalization). *Let the compact SVD of the extrapolated momentum be $N_t = U\Sigma V^\top \in \mathbb{R}^{m \times n}$ with $\Sigma = \text{diag}(\sigma_1, \dots, \sigma_r)$, where $r = \text{rank}(N_t)$. A*

¹The reference code uses an exponential moving average for the momentum and a second interpolation with the same coefficient: <https://github.com/KellerJordan/Muon/blob/master/muon.py>.

Algorithm 1 Practical layer-wise Muon with Nesterov momentum and inexact orthogonalization

Require: learning rate $\gamma > 0$, momentum $\beta_1 \in [0, 1)$, $\beta_2 \in [0, 1]$, total iterations T , number of layers L .

```
1: Initialize:  $M_0^{(l)} \leftarrow 0$ ,  $X_1^{(l)} \in \mathcal{S}_l$  for each layer  $l \in \{1, \dots, L\}$ .
2: for  $t = 1$  to  $T$  do
3:   for each layer  $l = 1$  to  $L$  do
4:      $G_t^{(l)} \leftarrow \nabla_{(l)} f(X_t; \xi_t)$  ▷ Compute layer-wise batch gradients
5:      $M_t^{(l)} \leftarrow \beta_1 M_{t-1}^{(l)} + (1 - \beta_1) G_t^{(l)}$  ▷ Update first-order momentum
6:      $N_t^{(l)} \leftarrow \beta_2 M_t^{(l)} + (1 - \beta_2) G_t^{(l)}$  ▷ Apply Nesterov correction
7:      $O_t^{(l)} \leftarrow \text{Inexact-Orthogonalization}(N_t^{(l)})$  ▷ Generate approximate polar proxy
8:      $X_{t+1}^{(l)} \leftarrow X_t^{(l)} - \gamma O_t^{(l)}$  ▷ Update parameters
9:   end for
10: end for
```

singular-vector-preserving orthogonalization routine gives $O_t = U\varsigma V^\top$, where $\varsigma = \text{diag}(\varsigma_1, \dots, \varsigma_r)$. For a finite polynomial routine with input normalization $\nu_t > 0$, the modified values are $\varsigma_i = \phi_J(\sigma_i/\nu_t)$, where J denotes the number of polynomial steps and is distinct from the smoothness exponent q . Suppose these values are nonnegative. Then:

1. **Directional Alignment:** $\langle N_t, O_t \rangle \geq \varsigma_{\min} \|N_t\|_*$, where $\varsigma_{\min} = \min_i \varsigma_i$.

2. **Frobenius Norm Bound:** $\|O_t\|_F \leq \varsigma_{\max} \sqrt{r}$, where $\varsigma_{\max} = \max_i \varsigma_i$.

If $N_t = 0$, take $O_t = 0$ and interpret both inequalities directly, without a minimum over an empty spectrum. Uniform positive lower bounds on the nonzero ς_i are an additional condition in Theorem 3.1; they do not follow merely from using finitely many Newton–Schulz steps.

Assumption 2.2 (Objective Lower Boundedness). The objective function F is lower bounded on $\mathcal{S}$; that is, there exists a constant $F^* > -\infty$ such that

$$F(X) \geq F^*, \quad \forall X \in \mathcal{S}. \quad (2)$$

Assumption 2.3 (Global $(\mathcal{L}_0, \mathcal{L}_1, q)$ -Smoothness). Let $\mathcal{S} = \mathbb{R}^{n_1 \times m_1} \times \dots \times \mathbb{R}^{n_L \times m_L}$, and equip the complete parameter tuple with $\|X\|_F^2 = \sum_{l=1}^L \|X^{(l)}\|_F^2$. The differentiable objective $F : \mathcal{S} \rightarrow \mathbb{R}$ satisfies, for some $\mathcal{L}_0 > 0$, $\mathcal{L}_1 \geq 0$, and $q \in [0, 1]$, and all $X, Y \in \mathcal{S}$,

$$\begin{aligned} \|\nabla F(X) - \nabla F(Y)\|_F \\ \leq (\mathcal{L}_0 + \mathcal{L}_1 \|\nabla F(X)\|_F^q) \|X - Y\|_F. \end{aligned} \quad (3)$$

For $q = 0$, the power is interpreted as 1, including at a zero gradient. The condition applies when several blocks change simultaneously, as in Algorithm 1. A bound only for X, Y differing in one block would not justify the joint descent and momentum-drift estimates used below.

Assumption 2.4 (Layer-wise Variance). The stochastic gradient oracle is assumed to have a layer-wise variance bound, where the parameter space $\mathcal{S} = \mathbb{R}^{n_1 \times m_1} \times \dots \times \mathbb{R}^{n_L \times m_L}$ is the Cartesian product of the weight spaces of all L layers. Specifically, for each layer $l \in \{1, \dots, L\}$, there exists a constant $\sigma_{(l)} \geq 0$ such that, for any $X \in \mathcal{S}$, the noisy gradient $\nabla_{(l)} f(X; \xi)$ with respect to the l -th layer satisfies:

$$\begin{aligned} \mathbb{E}_\xi [\nabla_{(l)} f(X; \xi)] &= \nabla_{(l)} F(X), \\ \mathbb{E}_\xi [\|\nabla_{(l)} f(X; \xi) - \nabla_{(l)} F(X)\|_F^2] &\leq \sigma_{(l)}^2, \end{aligned} \quad (4)$$

where ξ denotes the random sampling index and $\|\cdot\|_F$ denotes the Frobenius norm. At iteration t , ξ_t is sampled independently of the past $\mathcal{F}_{t-1}$, which includes X_t . Thus the same statements hold conditionally on $\mathcal{F}_{t-1}$. Independence of the noise across layers is not required.

Scope of the assumptions. Assumption 2.3 includes ordinary global $\mathcal{L}_0$ -smoothness when $\mathcal{L}_1 = 0$ and otherwise allows the smoothness bound to depend on the gradient norm. We use the joint condition because all blocks are updated together. Assumption 2.4 controls conditional variance, not the magnitude of every stochastic gradient; no almost-sure gradient bound, noise symmetry, or sub-Gaussian tail assumption is imposed. The finite-step result below further dispenses with a positive lower bound on nonzero output singular values. These are the specific relaxations used in the analysis; no additional architecture assumption is needed.

3 Main Results

Throughout, $\nabla F_{(l)}(X)$ and $\nabla_{(l)} F(X)$ denote the same gradient block. We develop one descent-tracking argument for Algorithm 1, first under output singular-value bounds and then with the geometry of the finite Newton–Schulz map. The block sums retain layer-specific noise levels, while gradient drift is controlled for the joint move $X_{t+1} - X_t$ on the coupled objective.

3.1 A common descent and tracking argument

The proof has four steps: derive and telescope a descent inequality; expand the Nesterov tracking error; bound its initialization, noise, and gradient-drift terms; and substitute these bounds to absorb the remaining gradient term. The orthogonalization routine enters through alignment and update-energy bounds. We establish these bounds in two forms below.

Control through output singular values. The first form uses Proposition 2.1 and gives a nuclear-norm stationarity bound with explicit rank dependence. The positive output singular-value floor is an additional spectral condition. The finite-map estimates later in this subsection lead to the Frobenius result in Section 3.2 without that condition.

Theorem 3.1 (Convergence of Layer-wise Muon). *Suppose that Assumptions 2.2, 2.3, and 2.4 hold. Let $\{X_t\}_{t=1}^{T+1}$ be generated by Algorithm 1 with $\beta_1 \in [0, 1)$, $\beta_2 \in [0, 1]$ and $\gamma > 0$. For each layer l , let $r^{(l)}$ bound both $\text{rank}(N_t^{(l)})$ and $\text{rank}(N_t^{(l)} - \nabla_{(l)} F(X_t))$ along the iterates (the choice $r^{(l)} = \min\{n_l, m_l\}$ always suffices), and let the inexact orthogonalization satisfy Proposition 2.1 with fixed constants $\varsigma_{\min} > 0$ and $\varsigma_{\max} > 0$ uniformly over all layers and iterates. Define $\hat{\mathcal{L}} = \mathcal{L}_0 + (1-q)\mathcal{L}_1$, $R_0 = \sum_{l=1}^L r^{(l)}$, $S_0 = \sum_{l=1}^L \sqrt{r^{(l)}} \sigma_{(l)}$, $g_l = \sqrt{r^{(l)}} \|\nabla F_{(l)}(X_1)\|_F$, and $G_0 = \sum_{l=1}^L g_l$. Set*

$$\Delta_{\gamma, \beta_1} := \varsigma_{\min} - \frac{\gamma \varsigma_{\max}^2 q \mathcal{L}_1 R_0}{2} - \frac{2\gamma \varsigma_{\max}^2 q \mathcal{L}_1 R_0}{1 - \beta_1}.$$

Then

$$\begin{aligned}
& \Delta_{\gamma, \beta_1} \frac{1}{T} \sum_{t=1}^T \sum_{l=1}^L \mathbb{E} \left[\left\| \nabla F_{(l)}(X_t) \right\|_* \right] \\
& \leq \frac{F(X_1) - F^*}{\gamma T} + \frac{2\beta_2 \varsigma_{\max} G_0}{T(1 - \beta_1)} \\
& \quad + 2 \left(\sqrt{1 - \beta_1} + \sqrt{2\beta_1(1 - \beta_1\beta_2)(1 - \beta_2)} \right) \varsigma_{\max} S_0 \\
& \quad + \frac{2\gamma\beta_2 \varsigma_{\max}^2 \hat{\mathcal{L}}}{1 - \beta_1} R_0 + \frac{\gamma \varsigma_{\max}^2 \hat{\mathcal{L}} R_0}{2}.
\end{aligned} \tag{5}$$

Corollary 3.2 (Rate of Layer-wise Muon with scheduled Nesterov correction). *Suppose that the conditions of Theorem 3.1 hold, with $F(X_1) - F^* > 0$ and $S_0 > 0$ for the optimized parameter formulas below. Denote $R_0 = \sum_{l=1}^L r^{(l)}$, $S_0 = \sum_{l=1}^L \sqrt{r^{(l)}} \sigma_{(l)}$, and $G_0 = \sum_{l=1}^L \sqrt{r^{(l)}} \left\| \nabla F_{(l)}(X_1) \right\|_F$. Let $\beta_2 = 1 - (1 - \beta_1)^a$ for $a > \frac{1}{2}$. Since $\beta_2 \leq 1$, choose $C_1, C_2, \gamma, \beta_1$ by minimizing the leading upper envelope*

$$\begin{aligned}
\Phi(\gamma, \beta_1) := & \frac{F(X_1) - F^*}{\gamma T} + 2\sqrt{1 - \beta_1} \varsigma_{\max} S_0 \\
& + \frac{2\gamma \varsigma_{\max}^2 \hat{\mathcal{L}} R_0}{1 - \beta_1},
\end{aligned}$$

whose Young equality condition is

$$\frac{4(F(X_1) - F^*)}{\gamma T} = 4\sqrt{1 - \beta_1} \varsigma_{\max} S_0 = \frac{8\gamma \varsigma_{\max}^2 \hat{\mathcal{L}} R_0}{1 - \beta_1}.$$

This gives

$$\begin{aligned}
C_1 &= \frac{(F(X_1) - F^*)^{3/4}}{2^{1/4} \varsigma_{\max} S_0^{1/2} \hat{\mathcal{L}}^{1/4} R_0^{1/4}}, \\
C_2 &= \frac{2^{1/2} \hat{\mathcal{L}}^{1/2} (F(X_1) - F^*)^{1/2} R_0^{1/2}}{S_0}, \\
\gamma &= \frac{C_1}{T^{3/4}}, \quad \beta_1 = 1 - \frac{C_2}{T^{1/2}}.
\end{aligned}$$

Assume that T is large enough such that $\beta_1 \in [0, 1)$ and

$$T \geq \left(\left(\frac{C_1 \varsigma_{\max}^2 q \mathcal{L}_1 R_0}{\varsigma_{\min}} \right)^{1/3} + \frac{4C_1 \varsigma_{\max}^2 q \mathcal{L}_1 R_0}{C_2 \varsigma_{\min}} \right)^4. \tag{6}$$

Then the iterates generated by Algorithm 1 satisfy

$$\begin{aligned}
& \frac{1}{T} \sum_{t=1}^T \sum_{l=1}^L \mathbb{E} \left[\left\| \nabla F_{(l)}(X_t) \right\|_* \right] \leq \frac{2 \cdot 512^{1/4} \varsigma_{\max} \hat{\mathcal{L}}^{1/4} S_0^{1/2} R_0^{1/4}}{\varsigma_{\min} T^{1/4}} \\
& \quad \cdot (F(X_1) - F^*)^{1/4} + \frac{4\varsigma_{\max} G_0}{C_2 \varsigma_{\min} T^{1/2}} \\
& \quad + \frac{4\sqrt{2} \varsigma_{\max} S_0}{\varsigma_{\min}} \left(\frac{C_2^{\gamma^{(1+a)/2}}}{T^{(1+a)/4}} + \frac{C_2^a}{T^{a/2}} \right) \\
& \quad + \frac{C_1 \varsigma_{\max}^2 \hat{\mathcal{L}} R_0}{\varsigma_{\min} T^{3/4}}.
\end{aligned} \tag{7}$$

Consequently,

$$\frac{1}{T} \sum_{t=1}^T \sum_{l=1}^L \mathbb{E} \left[\|\nabla F_{(l)}(X_t)\|_* \right] = \mathcal{O} \left(T^{-1/4} \right). \quad (8)$$

Control through the finite Newton–Schulz map. To use the same proof for a fixed number of polynomial steps, we derive alignment and energy bounds from the actual scalar map. Its output can approach zero on small input singular values, so the preceding theorem’s positive output floor is not generally available. The ratio of output to input singular values provides the required control instead. For $N \neq 0$, initialize $Z_0 = N/\|N\|_F$ and take a fixed number J of polynomial steps

$$\begin{aligned} Z_j &= a_j Z_{j-1} + b_j (Z_{j-1} Z_{j-1}^\top) Z_{j-1} \\ &\quad + c_j (Z_{j-1} Z_{j-1}^\top)^2 Z_{j-1}, \quad j = 1, \dots, J, \end{aligned} \quad (9)$$

with $O = Z_J$; set $O = 0$ when $N = 0$. The coefficients may vary with j , but J and their values are fixed independently of matrix size and T . All statements concern exact arithmetic. Write

$$p_j(s) = a_j s + b_j s^3 + c_j s^5, \quad \phi_J = p_J \circ \dots \circ p_1. \quad (10)$$

We require the following property of this scalar map:

$$\begin{aligned} 0 < \ell_J &:= \inf_{0 < s \leq 1} \frac{\phi_J(s)}{s}, \quad u_J := \sup_{0 < s \leq 1} \frac{\phi_J(s)}{s} < \infty, \\ c_J &:= \max_{0 \leq s \leq 1} \phi_J(s). \end{aligned} \quad (11)$$

The ratio has the continuous extension $\phi'_J(0) = \prod_{j=1}^J a_j$ at zero. Thus these are scalar constants, not bounds on a matrix’s smallest nonzero singular value. Positive lower bounds and finite upper bounds can also be used in place of the extrema in (11).

Proposition 3.3 (Finite-map alignment and energy). *Let $N = U\Sigma V^\top \neq 0$, $R = \|N\|_F$, and $s_i = \sigma_i(N)/R$. Then $\sum_i s_i^2 = 1$ and the direction in (9) is $O = U \operatorname{diag}(\phi_J(s_i)) V^\top$. Define*

$$A(N) = \sum_i s_i \phi_J(s_i), \quad B(N) = \sum_i \phi_J(s_i)^2. \quad (12)$$

The exact identities and bounds are

$$\begin{aligned} \langle N, O \rangle &= RA(N), \quad \|O\|_F^2 = B(N), \\ \ell_J &\leq A(N) \leq u_J, \quad B(N) \leq u_J A(N) \leq u_J^2, \\ \|O\|_{\text{op}} &\leq c_J. \end{aligned} \quad (13)$$

Moreover, with

$$\bar{\rho}_J := \frac{u_J + \ell_J}{2\sqrt{u_J \ell_J}}, \quad (14)$$

we have $\sqrt{B(N)} \leq \bar{\rho}_J A(N)$. Consequently, for any matrix H and $E = N - H$,

$$\langle H, O \rangle \geq A(N) (\|N\|_F - \bar{\rho}_J \|E\|_F). \quad (15)$$

No rank factor appears in these statements.

An explicit five-step instance.

For the fixed quintic coefficients $(a, b, c) = (3.4445, -4.775, 2.0315)$, as used in the original Muon implementation (Jordan, 2024), one may use the fully analytic bounds

$$\begin{aligned} \ell_J &\geq \frac{17}{25}, & u_J &= 3.4445^J, \\ c_J &\leq \frac{5}{4}, & J &\geq 1. \end{aligned} \tag{16}$$

The lower bound uses invariant intervals of the whole scalar map; it is stronger than multiplying a separate worst-case lower bound at every step. For $J = 5$, $u_5 = 3.4445^5 \simeq 484.8763$, and the displayed bounds imply $\bar{\rho}_5 < 13.371$. These are certified bounds, not an assertion that a numerical scan has found the sharp lower or angular constant. The details are given in Appendix D.2.

The bounds $\langle N, O \rangle \geq \ell_J \|N\|_F$ and $\|O\|_F \leq u_J$ now replace the nuclear-norm alignment and rank-dependent update bound in the same four-step argument. The Nesterov error expansion is unchanged; joint gradient drift is controlled using $\|X_{t+1} - X_t\|_F \leq \gamma u_J \sqrt{L}$.

3.2 Dimension-independent convergence of practical Muon

We apply the argument of Section 3.1 with its finite-map estimates. The resulting stationarity measure is the sum of block Frobenius norms, which also bounds the Frobenius norm of the complete gradient. No positive output singular-value floor or rank bound for the momentum or tracking error is required.

Theorem 3.4 (Dimension-independent Frobenius stationarity). *Suppose Assumptions 2.2, 2.3, and 2.4 hold. Use (9) in Algorithm 1, with (11), $\beta_1 \in [0, 1]$, $\beta_2 \in [0, 1]$, and $\gamma > 0$. Define*

$$\begin{aligned} \delta &= 1 - \beta_1, & D &= F(X_1) - F^*, \\ S_F &= \sum_{l=1}^L \sigma_{(l)}, \\ G_F &= \sum_{l=1}^L \|\nabla_{(l)} F(X_1)\|_F, \\ v_\beta &= \sqrt{\delta} + \sqrt{2\beta_1(1 - \beta_1\beta_2)(1 - \beta_2)}, \\ \hat{\mathcal{L}} &= \mathcal{L}_0 + (1 - q)\mathcal{L}_1. \end{aligned} \tag{17}$$

Set

$$\Delta_{\gamma, \beta_1}^F := \ell_J - \frac{\gamma L u_J^2 q \mathcal{L}_1}{2} - \frac{\gamma L u_J (\ell_J + u_J) q \mathcal{L}_1}{\delta}. \tag{18}$$

Then

$$\begin{aligned} &\Delta_{\gamma, \beta_1}^F \frac{1}{T} \sum_{t=1}^T \sum_{l=1}^L \mathbb{E} \|\nabla_{(l)} F(X_t)\|_F \\ &\leq \frac{D}{\gamma T} + \frac{\beta_2 (\ell_J + u_J) G_F}{T \delta} + (\ell_J + u_J) v_\beta S_F \\ &\quad + \frac{\gamma \beta_2 L u_J (\ell_J + u_J) \hat{\mathcal{L}}}{\delta} + \frac{\gamma L u_J^2 \hat{\mathcal{L}}}{2}. \end{aligned} \tag{19}$$

In particular, a positive $\Delta_{\gamma, \beta_1}^F$ gives a bound on the original, unstopped sequence of iterates.

Corollary 3.5 (Dimension-independent $T^{-1/4}$ rate). *Under Theorem 3.4, fix $\gamma_0, \delta_0 > 0$ and $a > 1/2$, and set*

$$\gamma = \gamma_0 T^{-3/4}, \quad 1 - \beta_1 = \delta_0 T^{-1/2}, \quad \beta_2 = 1 - (1 - \beta_1)^a. \quad (20)$$

For all sufficiently large T , $\beta_1, \beta_2 \in [0, 1]$ and $\Delta_{\gamma, \beta_1}^F \geq \ell_J/2$. Writing $b_J = \ell_J + u_J$,

$$\begin{aligned} & \frac{1}{T} \sum_{t=1}^T \sum_{l=1}^L \mathbb{E} \|\nabla_{(l)} F(X_t)\|_F \\ & \leq \frac{2}{\ell_J} \left[\left(\frac{D}{\gamma_0} + b_J S_F \sqrt{\delta_0} + \frac{\gamma_0 L u_J b_J \hat{\mathcal{L}}}{\delta_0} \right) T^{-1/4} \right. \\ & \quad + \frac{b_J G_F}{\delta_0 T^{1/2}} \\ & \quad + \sqrt{2} b_J S_F \left(\frac{\delta_0^{(1+a)/2}}{T^{(1+a)/4}} + \frac{\delta_0^a}{T^{a/2}} \right) \\ & \quad \left. + \frac{\gamma_0 L u_J^2 \hat{\mathcal{L}}}{2T^{3/4}} \right]. \end{aligned} \quad (21)$$

The same upper bound holds for $T^{-1} \sum_t \mathbb{E} \|\nabla F(X_t)\|_F$. Its constant contains no $n_l, m_l, r^{(l)}$, or $d = \sum_l n_l m_l$.

The standard Nesterov choice is included. Setting $a = 1$ in (20) gives $\beta_2 = \beta_1$ and hence (1). In this case,

$$v_\beta = \sqrt{\delta} + \delta \sqrt{2\beta_1(1 + \beta_1)} \leq \sqrt{\delta} + 2\delta. \quad (22)$$

Thus the extra Nesterov noise term is $\mathcal{O}(T^{-1/2})$, smaller than the leading $\mathcal{O}(T^{-1/4})$ term. This specialization uses the usual Nesterov algebra, with a horizon-dependent momentum coefficient; it does not assert the same vanishing bound for a fixed numerical momentum coefficient and a fixed batch size.

A sufficient explicit horizon in this corollary is

$$\begin{aligned} T & \geq \max\{1, \delta_0^2, (A_J + B_J^{1/3})^4\}, \\ A_J & = \frac{2\gamma_0 L u_J (\ell_J + u_J) q \mathcal{L}_1}{\delta_0 \ell_J}, \quad B_J = \frac{\gamma_0 L u_J^2 q \mathcal{L}_1}{\ell_J}. \end{aligned} \quad (23)$$

For $q \mathcal{L}_1 = 0$, the absorption condition is automatic. The initial-gradient constant can also be eliminated: joint generalized smoothness and lower boundedness imply

$$G_F \leq \sqrt{L} \left[D q \mathcal{L}_1 + \sqrt{D^2 q^2 \mathcal{L}_1^2 + 2D \hat{\mathcal{L}}} \right]. \quad (24)$$

Thus, for fixed L, J , coefficient schedule, $\gamma_0, \delta_0, a, \mathcal{L}_0, \mathcal{L}_1, q, D$, and S_F , the complete displayed bound is independent of parameter dimension. This is not a proof that these problem quantities stay fixed when a model family changes.

4 Discussion

We discuss the coupled momentum and learning-rate schedules, the meaning of dimension independence, and the scope of the orthogonalization and normalization choices covered by the analysis.

4.1 Momentum schedules

Remark 4.1 (Coupled Muon schedules versus fixed momentum). There is a growing body of evidence that stable and transferable learning rates are not architecture-agnostic. The μ P framework gives width-aware hyperparameter transfer rules for large neural networks (Yang et al., 2022), and recent Muon pretraining studies combine Muon with μ P-style transfer and carefully chosen matrix update scales (Liu et al., 2025; Shah et al., 2025). For SGD/GD-style training, prior analyses have also identified nontrivial depth dependence of maximal or effective learning rates, including maximal initial learning rates in deep ReLU networks (Iyer et al., 2023), depth-dependent μ P learning rates (Jelassi et al., 2023), and architecture-aware learning-rate rules that depend on depth, width, kernel size, and graph topology (Chen et al., 2024). In Transformer training with Adam, Xiong et al. (2020) also showed that depth-dependent gradient behavior is closely related to learning-rate warmup and normalization design.

The rank-sensitive bound makes explicit how matrix structure enters one sufficient Muon stepsize. Corollary 3.2 should be interpreted as a theoretical coupled hyperparameter schedule for Muon. The learning-rate rule

$$\gamma \asymp \frac{1}{S_0^{1/2} R_0^{1/4} T^{3/4}}$$

is obtained together with the momentum schedule

$$1 - \beta_1 \asymp \frac{R_0^{1/2}}{S_0 \sqrt{T}}, \quad 1 - \beta_2 = (1 - \beta_1)^a, \quad a > \frac{1}{2}.$$

Thus, the depth-rank dependence of γ is not a stand-alone law under an arbitrary fixed momentum coefficient. It is the result of balancing the three leading terms in the upper envelope

$$\frac{F(X_1) - F^*}{\gamma T} + 2\sqrt{1 - \beta_1} \varsigma_{\max} S_0 + \frac{2\gamma \varsigma_{\max}^2 \hat{\mathcal{L}} R_0}{1 - \beta_1}.$$

In particular, if one fixes $\delta := 1 - \beta_1$ as in conventional training (e.g., $\delta = 0.1$ or 0.05), then the same envelope becomes

$$\frac{F(X_1) - F^*}{\gamma T} + 2\sqrt{\delta} \varsigma_{\max} S_0 + \frac{2\gamma \varsigma_{\max}^2 \hat{\mathcal{L}} R_0}{\delta}.$$

Optimizing only over γ yields the different scale $\gamma \asymp \sqrt{\delta/(R_0 T)}$, and the stochastic term $2\sqrt{\delta} \varsigma_{\max} S_0$ does not vanish with T . Therefore, the rate and the learning-rate schedule in Corollary 3.2 rely on the scheduled first-momentum gap $1 - \beta_1 = \Theta(R_0^{1/2}/(S_0 T^{1/2}))$. The rank bound R_0 directly enters the learning-rate scale because Muon’s orthogonalized matrix direction satisfies $\|O_t^{(l)}\|_F \lesssim \sqrt{r^{(l)}}$, so the smoothness cost accumulates as $\sum_l r^{(l)}$. This rank-sensitive term is specific to the matrix geometry of Muon and is not present in the same form for coordinate-wise Adam or Euclidean SGD. By contrast, the leading schedule for $1 - \beta_1$ depends on the layer-wise noise aggregate S_0 , the accumulated rank R_0 , and the horizon T ; remaining problem-dependent effects enter through constants such as $\hat{\mathcal{L}}$ and the initial optimality gap.

These rank factors belong to the nuclear-norm bound in Corollary 3.2. The finite-step Frobenius bound in Corollary 3.5 uses different spectral estimates and needs no explicit rank factor in its schedule.

This theoretical schedule should not be confused with empirical neural scaling laws of the Kaplan/Chinchilla type (Kaplan et al., 2020; Hoffmann et al., 2022). Our analysis characterizes how Muon’s hyperparameters should depend on depth, accumulated rank bound, and training horizon to

guarantee stationarity; it does not by itself imply a power-law relation between test loss, dataset size, model size, and compute.

Remark 4.2 (Role of the Nesterov momentum correction). The Nesterov momentum correction plays a different role from the coupled learning-rate and first-momentum schedules above. When $\beta_2 = 1$, $N_t^{(l)}$ reduces to the standard momentum $M_t^{(l)}$. From Theorem 3.1, the Nesterov coefficient enters the initialization term and the following noise–drift terms

$$2\sqrt{2\beta_1(1-\beta_1\beta_2)(1-\beta_2)}\varsigma_{\max}S_0 + \frac{2\gamma\beta_2\varsigma_{\max}^2\hat{\mathcal{L}}R_0}{1-\beta_1}.$$

The bound therefore exhibits a trade-off: a smaller β_2 reduces the initialization and drift coefficients, while changing the stochastic term. This does not by itself prove an empirical early- or late-stage advantage. The extra stochastic contribution and the change in the drift coefficient relative to $\beta_2 = 1$ are lower-order under $\beta_2 = 1 - (1 - \beta_1)^a$ with $a > \frac{1}{2}$; the drift term itself remains leading-order. Thus the Nesterov correction does not change the leading $\mathcal{O}(T^{-1/4})$ convergence rate.

4.2 Dimension independence

The bound in Corollary 3.5 contains no explicit matrix-dimension or rank factor when the block count and problem constants are fixed. The constants ℓ_J and u_J depend on the finite polynomial map; dimension independence does not imply that these constants are small or that the noise and smoothness constants stay fixed as an architecture changes.

Remark 4.3 (What the sharper spectral identities do and do not imply). Equations (12)–(15) retain the full finite map and give a useful descent-or-small-gradient test for an individual block. However, $\|O\|_{\text{op}} \leq c_J$ cannot replace $\|O\|_{\text{F}} \leq u_J$ in a Frobenius-smoothness bound without an additional curvature condition. Similarly, an expected tracking-error bound at each deterministic time does not imply the same bound at the first time that alignment becomes unreliable. We therefore prove the averaged stochastic result directly, using the same three-term tracking decomposition, rather than relying on a stopping-time conversion. Appendix D.5 gives the precise geometric certificate and explains this distinction. The present theorem establishes dimension independence; it does not claim a uniformly small NS constant.

4.3 Implementation scope

Remark 4.4 (Generality beyond exact Newton–Schulz orthogonalization). Our analysis does not require $O_t^{(l)}$ to be an exact orthogonal or polar factor. Exact orthogonalization corresponds to the special case $\varsigma_i = 1$ for all nonzero singular directions. In contrast, the proof of Theorem 3.1 only uses the two inequalities in Proposition 2.1,

$$\langle N_t^{(l)}, O_t^{(l)} \rangle \geq \varsigma_{\min} \|N_t^{(l)}\|_*, \quad \|O_t^{(l)}\|_{\text{F}} \leq \varsigma_{\max} \sqrt{r^{(l)}}.$$

Therefore, exact polar decomposition is not needed. It is sufficient that the orthogonalization routine preserves enough directional alignment with $N_t^{(l)}$ and keeps the update magnitude uniformly bounded.

This viewpoint is related to, but more algorithm-agnostic than, the recent analysis of Muon with Newton–Schulz orthogonalization (Kim and Oh, 2025). That work studies the practical Newton–Schulz iteration directly and shows that, for a fixed number of Newton–Schulz steps, Muon converges at the same rate as the ideal SVD-polar version up to a constant factor, with the factor converging

to one rapidly as the number of Newton–Schulz steps increases. Our argument does not require the modified singular values to be symmetric around 1, nor does it require $\varsigma_{\min}$ and $\varsigma_{\max}$ to approach 1 as the number of Newton–Schulz iterations grows. The constants in our bound depend on $\varsigma_{\min}$ and $\varsigma_{\max}$, and the same rate is preserved as long as $\varsigma_{\min} > 0$ and the distortion ratio $\varsigma_{\max}/\varsigma_{\min}$ remains controlled. Consequently, the proof applies not only to Newton–Schulz approximations, but also to other inexact orthogonalization or spectral-flattening procedures satisfying the same alignment and magnitude conditions. For Frobenius-normalized finite polynomial steps, however, $\phi_J(s) \rightarrow 0$ as $s \rightarrow 0$. Hence a uniform $\varsigma_{\min} > 0$ need not exist over arbitrary spectra. Sections 3.1 and 3.2 avoid this condition by using $\phi_J(s)/s$ and proving Frobenius, rather than nuclear-norm, stationarity.

Remark 4.5 (Normalization and implementation scope). If the initial normalization is $N/(\kappa\|N\|_F)$ for a fixed $\kappa \geq 1$, apply the results to the effective map $\tilde{\phi}_J(s) = \phi_J(s/\kappa)$ on $[0, 1]$. In particular, the slope at zero is $(\prod_j a_j)/\kappa$, not $\prod_j a_j$. For $N/(\|N\|_F + \varepsilon)$ with fixed $\varepsilon > 0$, the same energy bound holds and

$$\langle N, O \rangle \geq \ell_J \frac{\|N\|_F^2}{\|N\|_F + \varepsilon} \geq \ell_J(\|N\|_F - \varepsilon).$$

Accordingly, add $L\ell_J\varepsilon$ to the right-hand side of (19); (21) acquires the residual $2L\varepsilon$.

If the output in block l is multiplied by a fixed $\alpha_l > 0$, the same proof applies with ℓ_J replaced by $\alpha_{\min}\ell_J$ and u_J by $\alpha_{\max}u_J$, where $\alpha_{\min} = \min_l \alpha_l$ and $\alpha_{\max} = \max_l \alpha_l$; the operator-norm bound becomes $\alpha_{\max}c_J$. This follows by applying the blockwise inequalities with these common bounds. Such scaling preserves dimension independence only when $\alpha_{\max}$ and $1/\alpha_{\min}$ remain bounded independently of matrix size. Unbounded size-dependent multipliers cannot be absorbed into a dimension-independent constant. No finite-precision error guarantee is asserted here.

5 Conclusion

We established convergence guarantees for layer-wise Muon with finite Newton–Schulz updates and Nesterov momentum under joint gradient-dependent smoothness and bounded conditional variance. A common descent-and-tracking argument separates the momentum error into initialization, stochastic noise, and gradient drift. Controlling the finite polynomial through $\phi_J(s)/s$ yields an $\mathcal{O}(T^{-1/4})$ bound on the expected average Frobenius gradient norm under coupled learning-rate and momentum schedules, including the standard equal-coefficient Nesterov rule. The bound has no explicit matrix-dimension or rank factors when the block count and problem constants are fixed. We also verified the scalar-map conditions analytically for the original five-step quintic iteration.

The central implication is that finite orthogonalization provides sufficient alignment and update-energy control for convergence without a positive lower bound on its nonzero output singular values. The complementary nuclear-norm result identifies where rank dependence enters under stronger spectral control. Extending the analysis to finite-precision arithmetic and to fixed momentum with increasing batch sizes would further connect these guarantees to training practice.

References

- Chen, W., Wu, J., Wang, Z., and Hanin, B. (2024). Principled architecture-aware scaling of hyperparameters. In *International Conference on Learning Representations*.
- Choudhury, S., Cheng, X., Takáč, M., Na, S., and Kolar, M. (2026). Muon with Nesterov momentum: Heavy-tailed noise and (randomized) inexact polar decomposition. *arXiv preprint arXiv:2605.06884*.

- Hoffmann, J., Borgeaud, S., Mensch, A., Buchatskaya, E., Cai, T., Rutherford, E., Casas, D. d. L., Hendricks, L. A., Welbl, J., Clark, A., Hennigan, T., Noland, E., Millican, K., van den Driessche, G., Damoc, B., Guy, A., Osindero, S., Simonyan, K., Elsen, E., Vinyals, O., Rae, J. W., and Sifre, L. (2022). Training compute-optimal large language models. *arXiv preprint arXiv:2203.15556*.
- Iyer, G., Hanin, B., and Rolnick, D. (2023). Maximal initial learning rates in deep ReLU networks. In *International Conference on Machine Learning*, volume 202 of *Proceedings of Machine Learning Research*, pages 14500–14530. PMLR.
- Jelassi, S., Hanin, B., Ji, Z., Reddi, S. J., Bhojanapalli, S., and Kumar, S. (2023). Depth dependence of μP learning rates in ReLU MLPs. *arXiv preprint arXiv:2305.07810*.
- Jordan, K. (2024). Muon: An optimizer for hidden layers in neural networks. Online technical report. Published December 8, 2024.
- Kaplan, J., McCandlish, S., Henighan, T., Brown, T. B., Chess, B., Child, R., Gray, S., Radford, A., Wu, J., and Amodei, D. (2020). Scaling laws for neural language models. *arXiv preprint arXiv:2001.08361*.
- Kim, G. Y. and Oh, M.-h. (2025). Convergence of Muon with Newton–Schulz. *arXiv preprint arXiv:2510.20377*.
- Li, M. and Tsuchiya, T. (2026). Muon with finite Newton–Schulz: The smoothing benefit in nonsmooth nonconvex optimization. *arXiv preprint arXiv:2608.26288*.
- Liu, J., Su, J., Yao, X., Jiang, Z., Lai, G., Du, Y., Qin, Y., Xu, W., Lu, E., Yan, J., et al. (2025). Muon is scalable for llm training. *arXiv preprint arXiv:2502.16982*.
- Pethick, T., Xie, W., Antonakopoulos, K., Zhu, Z., Silveti-Falls, A., and Cevher, V. (2025). Training deep learning models with norm-constrained LMOs. *arXiv preprint arXiv:2502.07529*.
- Riabinin, A., Shulgin, E., Grutkowska, K., and Richtárik, P. (2025). Gluon: Making Muon & Scion great again! (bridging theory and practice of LMO-based optimizers for LLMs). *arXiv preprint arXiv:2505.13416*.
- Shah, I., Polloreno, A. M., Stratos, K., Monk, P., Chaluvareja, A., Hojel, A., Ma, A., Thomas, A., Tanwer, A., Shah, D. J., et al. (2025). Practical efficiency of muon for pretraining. *arXiv preprint arXiv:2505.02222*.
- Shen, W., Huang, R., Huang, M., Shen, C., and Zhang, J. (2025). On the convergence analysis of Muon. *arXiv preprint arXiv:2505.23737*.
- Xiong, R., Yang, Y., He, D., Zheng, K., Zheng, S., Xing, C., Zhang, H., Lan, Y., Wang, L., and Liu, T. (2020). On layer normalization in the transformer architecture. In *International Conference on Machine Learning*, pages 10524–10533. PMLR.
- Yang, G., Hu, E. J., Babuschkin, I., Sidor, S., Liu, X., Farhi, D., Ryder, N., Pachocki, J., Chen, W., and Gao, J. (2022). Tensor programs V: Tuning large neural networks via zero-shot hyperparameter transfer. *arXiv preprint arXiv:2203.03466*.

A Useful Lemmas

Lemma A.1. *Under Assumption 2.3, for any $X, Y \in \mathcal{S}$,*

$$F(Y) \leq F(X) + \langle \nabla F(X), Y - X \rangle + \frac{\mathcal{L}_0 + \mathcal{L}_1 \|\nabla F(X)\|_{\mathbb{F}}^q}{2} \sum_{l=1}^L \|Y^{(l)} - X^{(l)}\|_{\mathbb{F}}^2. \quad (25)$$

Proof. Writing $V = Y - X$ and applying the fundamental theorem of calculus gives

$$\begin{aligned} F(Y) - F(X) - \langle \nabla F(X), V \rangle &= \int_0^1 \langle \nabla F(X + tV) - \nabla F(X), V \rangle dt \\ &\leq \int_0^1 \|\nabla F(X + tV) - \nabla F(X)\|_{\mathbb{F}} \|V\|_{\mathbb{F}} dt \\ &\leq \frac{\mathcal{L}_0 + \mathcal{L}_1 \|\nabla F(X)\|_{\mathbb{F}}^q}{2} \|V\|_{\mathbb{F}}^2. \end{aligned} \quad (26)$$

The last identity $\|V\|_{\mathbb{F}}^2 = \sum_l \|V^{(l)}\|_{\mathbb{F}}^2$ proves the claim. $\square$

Lemma A.2. *Let $\{N_t^{(l)}\}_{t=1}^T$ be the generalized Nesterov momentum sequence generated by Algorithm 1. Then, for $t \geq 2$, $N_t^{(l)}$ admits the following equivalent recursive form (with $N_1^{(l)} = (1 - \beta_1\beta_2)G_1^{(l)}$):*

$$N_t^{(l)} = \beta_1 N_{t-1}^{(l)} + (1 - \beta_1\beta_2)G_t^{(l)} - \beta_1(1 - \beta_2)G_{t-1}^{(l)} = \beta_1 N_{t-1}^{(l)} + (1 - \beta_1)G_t^{(l)} + \beta_1(1 - \beta_2)(G_t^{(l)} - G_{t-1}^{(l)}). \quad (27)$$

Proof. By the update rule for $N_t^{(l)}$, we obtain

$$\begin{aligned} N_t^{(l)} - \beta_1 N_{t-1}^{(l)} &= \beta_2 M_t^{(l)} + (1 - \beta_2)G_t^{(l)} - \beta_1(\beta_2 M_{t-1}^{(l)} + (1 - \beta_2)G_{t-1}^{(l)}) \\ &= \beta_2(\beta_1 M_{t-1}^{(l)} + (1 - \beta_1)G_t^{(l)}) + (1 - \beta_2)G_t^{(l)} - \beta_1\beta_2 M_{t-1}^{(l)} - \beta_1(1 - \beta_2)G_{t-1}^{(l)} \\ &= (1 - \beta_1\beta_2)G_t^{(l)} - \beta_1(1 - \beta_2)G_{t-1}^{(l)} \\ &= (1 - \beta_1)G_t^{(l)} + \beta_1(1 - \beta_2)(G_t^{(l)} - G_{t-1}^{(l)}). \end{aligned} \quad (28)$$

Rearranging terms gives the desired recursion. $\square$

Lemma A.3. *Let $a, b > 0$ and $0 < \alpha < \beta$. Define*

$$\bar{x} := \left(a + b^{\alpha/\beta}\right)^{1/\alpha}.$$

Then, for every $x \geq \bar{x}$,

$$\frac{a}{x^\alpha} + \frac{b}{x^\beta} \leq 1.$$

Moreover, if $x_\star$ denotes the smallest positive solution of $a/x^\alpha + b/x^\beta \leq 1$, then

$$x_\star \leq \bar{x} \leq 2^{1/\alpha} x_\star.$$

Equivalently, the corresponding threshold in the variable $s = x^\alpha$ is within a factor of 2 of the optimal threshold.

Proof. Let $s = x^\alpha$ and $\gamma = \beta/\alpha > 1$. The desired inequality is equivalent to

$$\frac{a}{s} + \frac{b}{s^\gamma} \leq 1 \quad (29)$$

with $s > 0$. Any feasible s for (29) must satisfy

$$s \geq \max\{a, b^{1/\gamma}\}, \quad (30)$$

since each term on the left-hand side of (29) is nonnegative.

Now set $\bar{s} := a + b^{1/\gamma}$. Since $\bar{s} \geq b^{1/\gamma}$, we have $b/\bar{s}^\gamma \leq b^{1/\gamma}/\bar{s}$. Therefore,

$$\frac{a}{\bar{s}} + \frac{b}{\bar{s}^\gamma} \leq \frac{a}{\bar{s}} + \frac{b^{1/\gamma}}{\bar{s}} = 1. \quad (31)$$

Thus $\bar{s}$ is feasible, and consequently every $x \geq \bar{x} := \bar{s}^{1/\alpha}$ satisfies the claimed inequality.

Let $s_\star$ be the smallest feasible value of s in (29). By (30), $s_\star \geq \max\{a, b^{1/\gamma}\}$. Hence

$$\bar{s} = a + b^{1/\gamma} \leq 2 \max\{a, b^{1/\gamma}\} \leq 2s_\star.$$

Taking the power $1/\alpha$ gives $\bar{x} \leq 2^{1/\alpha} x_\star$, where $x_\star = s_\star^{1/\alpha}$. The inequality $x_\star \leq \bar{x}$ follows from the feasibility of $\bar{x}$. $\square$

B Proof of Theorem 3.1

Proof. **Step 1: Descent and telescoping.** Applying Lemma A.1 with $Y = X_{t+1}$ and $X = X_t$ gives

$$F(X_{t+1}) \leq F(X_t) + \langle \nabla F(X_t), X_{t+1} - X_t \rangle + \sum_{l=1}^L \frac{\mathcal{L}_0 + \mathcal{L}_1 \|\nabla F(X_t)\|_{\mathbb{F}}^q}{2} \|X_{t+1}^{(l)} - X_t^{(l)}\|_{\mathbb{F}}^2. \quad (32)$$

For compactness in this proof, write $K_t = \mathcal{L}_0 + \mathcal{L}_1 \|\nabla F(X_t)\|_{\mathbb{F}}^q$. Substituting the update rule $X_{t+1}^{(l)} = X_t^{(l)} - \gamma O_t^{(l)}$ into the preceding inequality yields

$$\begin{aligned}
F(X_{t+1}) &\leq F(X_t) + \langle \nabla F(X_t), X_{t+1} - X_t \rangle \\
&\quad + \frac{K_t}{2} \sum_{l=1}^L \|X_{t+1}^{(l)} - X_t^{(l)}\|_{\mathbb{F}}^2 \\
&= F(X_t) - \gamma \sum_{l=1}^L \langle \nabla F_{(l)}(X_t), O_t^{(l)} \rangle \\
&\quad + \frac{\gamma^2 K_t}{2} \sum_{l=1}^L \|O_t^{(l)}\|_{\mathbb{F}}^2 \\
&= F(X_t) - \gamma \sum_{l=1}^L \langle N_t^{(l)}, O_t^{(l)} \rangle \\
&\quad + \gamma \sum_{l=1}^L \langle N_t^{(l)} - \nabla F_{(l)}(X_t), O_t^{(l)} \rangle \\
&\quad + \frac{\gamma^2 K_t}{2} \sum_{l=1}^L \|O_t^{(l)}\|_{\mathbb{F}}^2 \\
&\leq F(X_t) - \gamma \sum_{l=1}^L \langle N_t^{(l)}, O_t^{(l)} \rangle \\
&\quad + \gamma \sum_{l=1}^L \|N_t^{(l)} - \nabla F_{(l)}(X_t)\|_{\mathbb{F}} \|O_t^{(l)}\|_{\mathbb{F}} \\
&\quad + \frac{\gamma^2 K_t}{2} \sum_{l=1}^L \|O_t^{(l)}\|_{\mathbb{F}}^2.
\end{aligned} \tag{33}$$

By Proposition 2.1, the inexact orthogonalization direction satisfies $\langle N_t^{(l)}, O_t^{(l)} \rangle \geq \varsigma_{\min} \|N_t^{(l)}\|_*$ and $\|O_t^{(l)}\|_{\mathbb{F}} \leq \varsigma_{\max} \sqrt{r^{(l)}}$. Write $D_t^{(l)} = N_t^{(l)} - \nabla F_{(l)}(X_t)$ and $H_t = \mathcal{L}_0 + (1 - q)\mathcal{L}_1 + q\mathcal{L}_1 \|\nabla F(X_t)\|_{\mathbb{F}}$.

Then

$$\begin{aligned}
F(X_{t+1}) &\leq F(X_t) - \gamma \varsigma_{\min} \sum_{l=1}^L \|N_t^{(l)}\|_* + \gamma \varsigma_{\max} \sum_{l=1}^L \sqrt{r^{(l)}} \|D_t^{(l)}\|_{\mathbb{F}} \\
&\quad + \frac{\gamma^2 \varsigma_{\max}^2 K_t}{2} \sum_{l=1}^L r^{(l)} \\
&\stackrel{(i)}{\leq} F(X_t) - \gamma \varsigma_{\min} \sum_{l=1}^L \|N_t^{(l)}\|_* + \gamma \varsigma_{\max} \sum_{l=1}^L \sqrt{r^{(l)}} \|D_t^{(l)}\|_{\mathbb{F}} \\
&\quad + \frac{\gamma^2 \varsigma_{\max}^2 H_t}{2} \sum_{l=1}^L r^{(l)} \\
&\stackrel{(ii)}{\leq} F(X_t) - \gamma \varsigma_{\min} \sum_{l=1}^L \|\nabla F_{(l)}(X_t)\|_* \\
&\quad + \gamma \varsigma_{\min} \sum_{l=1}^L \|D_t^{(l)}\|_* + \gamma \varsigma_{\max} \sum_{l=1}^L \sqrt{r^{(l)}} \|D_t^{(l)}\|_{\mathbb{F}} \\
&\quad + \frac{\gamma^2 \varsigma_{\max}^2 H_t}{2} \sum_{l=1}^L r^{(l)} \\
&\stackrel{(iii)}{\leq} F(X_t) - \gamma \varsigma_{\min} \sum_{l=1}^L \|\nabla F_{(l)}(X_t)\|_* \\
&\quad + 2\gamma \varsigma_{\max} \sum_{l=1}^L \sqrt{r^{(l)}} \|D_t^{(l)}\|_{\mathbb{F}} + \frac{\gamma^2 \varsigma_{\max}^2 H_t}{2} \sum_{l=1}^L r^{(l)}.
\end{aligned} \tag{34}$$

Here, (i) follows from $a^q \leq (1-q) + qa$ for $a \geq 0$ and $q \in [0, 1]$; (ii) follows from the reverse triangle inequality $\|A\|_* \geq \|B\|_* - \|B - A\|_*$; and (iii) uses $\|A\|_* \leq \sqrt{r} \|A\|_{\mathbb{F}}$ where r bounds the rank of the error matrix $A = \nabla_{(l)} F(X_t) - N_t^{(l)}$, as stipulated in Theorem 3.1, and $\varsigma_{\min} \leq \varsigma_{\max}$.

Taking expectations and summing the resulting descent inequality over $t = 1, \dots, T$, the intermediate terms telescope and give

$$\begin{aligned}
\mathbb{E}[F(X_{T+1})] &\leq F(X_1) - \gamma \varsigma_{\min} \sum_{t=1}^T \sum_{l=1}^L \mathbb{E} \|\nabla F_{(l)}(X_t)\|_* \\
&\quad + 2\gamma \varsigma_{\max} \sum_{t=1}^T \sum_{l=1}^L \sqrt{r^{(l)}} \mathbb{E} \|D_t^{(l)}\|_{\mathbb{F}} \\
&\quad + \frac{\gamma^2 \varsigma_{\max}^2}{2} \sum_{t=1}^T \sum_{l=1}^L r^{(l)} \mathbb{E}[H_t].
\end{aligned} \tag{35}$$

Rearranging the terms and using Assumption 2.2, we obtain

$$\begin{aligned}
&\frac{\varsigma_{\min}}{T} \sum_{t=1}^T \sum_{l=1}^L \mathbb{E} \left[\|\nabla F_{(l)}(X_t)\|_* \right] - \frac{\gamma \varsigma_{\max}^2 q}{2T} \sum_{t=1}^T \sum_{l=1}^L r^{(l)} \mathcal{L}_1 \mathbb{E} [\|\nabla F(X_t)\|_{\mathbb{F}}] \\
&\leq \frac{F(X_1) - F^*}{\gamma T} + \frac{2\varsigma_{\max}}{T} \sum_{t=1}^T \sum_{l=1}^L \sqrt{r^{(l)}} \mathbb{E} \left[\|N_t^{(l)} - \nabla F_{(l)}(X_t)\|_{\mathbb{F}} \right] + \frac{\gamma \varsigma_{\max}^2}{2} \sum_{l=1}^L r^{(l)} \hat{\mathcal{L}},
\end{aligned} \tag{36}$$

where $\hat{\mathcal{L}} := \mathcal{L}_0 + (1 - q)\mathcal{L}_1$.

Step 2: The Nesterov tracking-error expansion. We next control the discrepancy between the extrapolated momentum and the corresponding true gradient block. From the recursion $N_t^{(l)} = \beta_1 N_{t-1}^{(l)} + (1 - \beta_1 \beta_2) G_t^{(l)} - \beta_1 (1 - \beta_2) G_{t-1}^{(l)}$, we have

$$\begin{aligned} N_t^{(l)} - \nabla F_{(l)}(X_t) = & \beta_1 \left(N_{t-1}^{(l)} - \nabla F_{(l)}(X_{t-1}) \right) + (1 - \beta_1 \beta_2) \left(G_t^{(l)} - \nabla F_{(l)}(X_t) \right) \\ & - \beta_1 (1 - \beta_2) \left(G_{t-1}^{(l)} - \nabla F_{(l)}(X_{t-1}) \right) + \beta_1 \beta_2 \left(\nabla F_{(l)}(X_{t-1}) - \nabla F_{(l)}(X_t) \right) \end{aligned} \quad (37)$$

Unrolling this recursion, together with the initialization $M_0^{(l)} = 0$, gives

$$\begin{aligned} N_t^{(l)} - \nabla F_{(l)}(X_t) = & -\beta_1^t \beta_2 \nabla F_{(l)}(X_1) + (1 - \beta_1 \beta_2) (G_t^{(l)} - \nabla F_{(l)}(X_t)) \\ & + \beta_2 (1 - \beta_1) \sum_{k=1}^{t-1} \beta_1^{t-k} (G_k^{(l)} - \nabla F_{(l)}(X_k)) \\ & + \beta_1 \beta_2 \sum_{k=2}^t \beta_1^{t-k} (\nabla F_{(l)}(X_{k-1}) - \nabla F_{(l)}(X_k)), \end{aligned} \quad (38)$$

Taking norms, expectations, and then averaging over $t = 1, \dots, T$, we decompose the error into three terms: Write $\varepsilon_t^{(l)} = G_t^{(l)} - \nabla F_{(l)}(X_t)$.

$$\frac{1}{T} \sum_{t=1}^T \mathbb{E} \|D_t^{(l)}\|_{\mathbb{F}} \leq \mathcal{T}_1 + \mathcal{T}_2 + \mathcal{T}_3, \quad (39)$$

where the initialization, noise, and drift terms are, respectively,

$$\begin{aligned} \mathcal{T}_1 &= \frac{\beta_2}{T} \sum_{t=1}^T \beta_1^t \|\nabla F_{(l)}(X_1)\|_{\mathbb{F}}, \\ \mathcal{T}_2 &= \frac{1}{T} \sum_{t=1}^T \mathbb{E} \left\| (1 - \beta_1 \beta_2) \varepsilon_t^{(l)} \right. \\ &\quad \left. + \beta_2 (1 - \beta_1) \sum_{k=1}^{t-1} \beta_1^{t-k} \varepsilon_k^{(l)} \right\|_{\mathbb{F}}, \\ \mathcal{T}_3 &= \frac{1}{T} \sum_{t=1}^T \mathbb{E} \left\| \beta_1 \beta_2 \sum_{k=2}^t \beta_1^{t-k} \right. \\ &\quad \left. \cdot (\nabla F_{(l)}(X_{k-1}) - \nabla F_{(l)}(X_k)) \right\|_{\mathbb{F}}. \end{aligned}$$

Step 3: Initialization, noise, and gradient drift. The first term is controlled directly by the geometric series:

$$\mathcal{T}_1 = \frac{\beta_2}{T} \sum_{t=1}^T \beta_1^t \|\nabla F_{(l)}(X_1)\|_{\mathbb{F}} \leq \frac{\beta_2 \|\nabla F_{(l)}(X_1)\|_{\mathbb{F}}}{T(1 - \beta_1)}. \quad (40)$$

For the stochastic-gradient noise term $\mathcal{T}_2$, Jensen's inequality and the martingale-difference

property of the stochastic gradients yield

$$\begin{aligned}
\mathcal{T}_2 &= \frac{1}{T} \sum_{t=1}^T \mathbb{E} \left\| (1 - \beta_1 \beta_2) \varepsilon_t^{(l)} + (1 - \beta_1) \beta_2 \sum_{k=1}^{t-1} \beta_1^{t-k} \varepsilon_k^{(l)} \right\|_{\mathbb{F}} \\
&\stackrel{(i)}{\leq} \frac{1}{T} \sum_{t=1}^T \sqrt{\mathbb{E} \left\| (1 - \beta_1 \beta_2) \varepsilon_t^{(l)} + (1 - \beta_1) \beta_2 \sum_{k=1}^{t-1} \beta_1^{t-k} \varepsilon_k^{(l)} \right\|_{\mathbb{F}}^2} \\
&\stackrel{(ii)}{=} \frac{1}{T} \sum_{t=1}^T \left((1 - \beta_1 \beta_2)^2 \mathbb{E} \|\varepsilon_t^{(l)}\|_{\mathbb{F}}^2 \right. \\
&\quad \left. + (1 - \beta_1)^2 \beta_2^2 \sum_{k=1}^{t-1} \beta_1^{2(t-k)} \mathbb{E} \|\varepsilon_k^{(l)}\|_{\mathbb{F}}^2 \right)^{1/2} \\
&\stackrel{(iii)}{\leq} \frac{1}{T} \sum_{t=1}^T \sqrt{(1 - \beta_1 \beta_2)^2 + (1 - \beta_1)^2 \beta_2^2 \sum_{k=1}^{t-1} \beta_1^{2(t-k)} \sigma_{(l)}} \\
&\leq \sqrt{(1 - \beta_1 \beta_2)^2 + \frac{(1 - \beta_1)^2 \beta_1^2 \beta_2^2}{1 - \beta_1^2} \sigma_{(l)}} \\
&= \sqrt{\frac{1 - \beta_1 + 2\beta_1(1 - \beta_1 \beta_2)(1 - \beta_2)}{1 + \beta_1} \sigma_{(l)}} \\
&\stackrel{(iv)}{\leq} \left(\sqrt{1 - \beta_1} + \sqrt{2\beta_1(1 - \beta_1 \beta_2)(1 - \beta_2)} \right) \sigma_{(l)}.
\end{aligned} \tag{41}$$

Here, (i) is Jensen's inequality in the form $\mathbb{E}[Z] \leq \sqrt{\mathbb{E}[Z^2]}$; (ii) uses conditional unbiasedness $\mathbb{E}[G_k^{(l)} - \nabla F_{(l)}(X_k) \mid \mathcal{F}_{k-1}] = 0$ from Assumption 2.4, which eliminates the cross terms; (iii) applies the variance bound in Assumption 2.4; and (iv) follows from $\sqrt{a+b} \leq \sqrt{a} + \sqrt{b}$.

It remains to bound the drift term $\mathcal{T}_3$, which is caused by the change of the true gradient along the trajectory. Write $\mathcal{T}_{3,l}$ for this term in layer l . Cauchy-Schwarz across layers and the triangle inequality give

$$\begin{aligned}
\sum_{l=1}^L \sqrt{r^{(l)}} \mathcal{T}_{3,l} &\leq \frac{\beta_1 \beta_2 \sqrt{R_0}}{T} \sum_{t=1}^T \sum_{k=2}^t \beta_1^{t-k} \mathbb{E} \|\nabla F(X_{k-1}) - \nabla F(X_k)\|_{\mathbb{F}} \\
&\leq \frac{\gamma \beta_1 \beta_2 \varsigma_{\max} R_0}{T} \sum_{t=1}^T \sum_{k=2}^t \beta_1^{t-k} \mathbb{E} [\mathcal{L}_0 + \mathcal{L}_1 \|\nabla F(X_k)\|_{\mathbb{F}}^q] \\
&\leq \frac{\gamma \beta_2 \varsigma_{\max} R_0}{1 - \beta_1} \left(\hat{\mathcal{L}} + \frac{q \mathcal{L}_1}{T} \sum_{k=1}^T \mathbb{E} \|\nabla F(X_k)\|_{\mathbb{F}} \right).
\end{aligned} \tag{42}$$

The second inequality applies Assumption 2.3 with base point X_k and uses $X_k - X_{k-1} = -\gamma O_{k-1}$ and $\|O_{k-1}\|_{\mathbb{F}} \leq \varsigma_{\max} \sqrt{R_0}$. The last inequality uses $z^q \leq (1-q) + qz$, exchanges the finite sums, and bounds the geometric series by $(1 - \beta_1)^{-1}$. In particular, no one-layer bound is applied to a displacement in all layers.

Combining the bounds for $\mathcal{T}_1$, $\mathcal{T}_2$, and $\mathcal{T}_3$ in Eq. (39), we obtain

$$\begin{aligned} \frac{1}{T} \sum_{t=1}^T \sum_{l=1}^L \sqrt{r^{(l)}} \mathbb{E} \|N_t^{(l)} - \nabla_{(l)} F(X_t)\|_F &\leq \frac{\beta_2 G_0}{T(1-\beta_1)} + \left(\sqrt{1-\beta_1} + \sqrt{2\beta_1(1-\beta_1\beta_2)(1-\beta_2)} \right) S_0 \\ &\quad + \frac{\gamma\beta_2\varsigma_{\max}R_0\hat{\mathcal{L}}}{1-\beta_1} + \frac{\gamma\beta_2\varsigma_{\max}R_0q\mathcal{L}_1}{1-\beta_1} \frac{1}{T} \sum_{t=1}^T \mathbb{E} \|\nabla F(X_t)\|_F. \end{aligned} \quad (43)$$

Step 4: Substitution and absorption. Finally, substituting Eq. (43) into Eq. (36) and using $\|\nabla F(X_t)\|_F \leq \sum_l \|\nabla_{(l)} F(X_t)\|_*$ and $\beta_2 \leq 1$ gives

$$\begin{aligned} &\left(\varsigma_{\min} - \frac{\gamma\varsigma_{\max}^2 q\mathcal{L}_1 R_0}{2} - \frac{2\gamma\varsigma_{\max}^2 q\mathcal{L}_1 R_0}{1-\beta_1} \right) \frac{1}{T} \sum_{t=1}^T \sum_{l=1}^L \mathbb{E} \left[\|\nabla_{F(l)}(X_t)\|_* \right] \\ &\leq \frac{F(X_1) - F^*}{\gamma T} + \frac{2\beta_2\varsigma_{\max}}{T(1-\beta_1)} \sum_{l=1}^L \sqrt{r^{(l)}} \|\nabla_{F(l)}(X_1)\|_F \\ &\quad + 2 \left(\sqrt{1-\beta_1} + \sqrt{2\beta_1(1-\beta_1\beta_2)(1-\beta_2)} \right) \varsigma_{\max} \sum_{l=1}^L \sqrt{r^{(l)}} \sigma_{(l)} \\ &\quad + \frac{2\gamma\beta_2\varsigma_{\max}^2 \hat{\mathcal{L}} \sum_{l=1}^L r^{(l)}}{1-\beta_1} + \frac{\gamma\varsigma_{\max}^2 \hat{\mathcal{L}} \sum_{l=1}^L r^{(l)}}{2}, \end{aligned} \quad (44)$$

where $R_0 = \sum_{l=1}^L r^{(l)}$. This is precisely the claimed convergence bound. The use of R_0 in the absorption coefficient accounts for simultaneous, coupled block updates. $\square$

C Proof of Corollary 3.2

Proof. Applying $\beta_2 = 1 - (1 - \beta_1)^a$ for $a > \frac{1}{2}$, we have

$$\sqrt{\beta_1(1-\beta_1\beta_2)(1-\beta_2)} = \sqrt{\beta_1(1-\beta_1 + \beta_1(1-\beta_1)^a)(1-\beta_1)^a} = \sqrt{\beta_1(1-\beta_1)^{1+a} + \beta_1^2(1-\beta_1)^{2a}}. \quad (45)$$

We need the term $\sqrt{\beta_1(1-\beta_1)^{1+a} + \beta_1^2(1-\beta_1)^{2a}}$ to be of no larger order than $\sqrt{1-\beta_1}$ so that the Nesterov momentum projection in Muon does not affect the final convergence rate. This is equivalent to $\beta_1(1-\beta_1)^{1+a} + \beta_1^2(1-\beta_1)^{2a} = O(1-\beta_1)$, which holds for $a > \frac{1}{2}$.

Since $\beta_2 \leq 1$, we minimize the following upper envelope of the three leading terms in Eq. (44). By Young's inequality,

$$\begin{aligned} &\frac{F(X_1) - F^*}{\gamma T} + 2\sqrt{1-\beta_1}\varsigma_{\max}S_0 + \frac{2\gamma\varsigma_{\max}^2\hat{\mathcal{L}}R_0}{1-\beta_1} \\ &\geq \left(\frac{4(F(X_1) - F^*)}{\gamma T} \right)^{1/4} \cdot \left(4\sqrt{1-\beta_1}\varsigma_{\max}S_0 \right)^{1/2} \cdot \left(\frac{8\gamma\varsigma_{\max}^2\hat{\mathcal{L}}R_0}{1-\beta_1} \right)^{1/4} \\ &= \frac{512^{1/4}\varsigma_{\max}\hat{\mathcal{L}}^{1/4}(F(X_1) - F^*)^{1/4}S_0^{1/2}R_0^{1/4}}{T^{1/4}}. \end{aligned} \quad (46)$$

The equality condition is

$$\frac{4(F(X_1) - F^*)}{\gamma T} = 4\sqrt{1-\beta_1}\varsigma_{\max}S_0 = \frac{8\gamma\varsigma_{\max}^2\hat{\mathcal{L}}R_0}{1-\beta_1}. \quad (47)$$

Solving these two balancing equations gives

$$\gamma = \frac{(F(X_1) - F^*)^{3/4}}{2^{1/4} \varsigma_{\max} S_0^{1/2} \hat{\mathcal{L}}^{1/4} R_0^{1/4} T^{3/4}}, \quad \beta_1 = 1 - \frac{(2\hat{\mathcal{L}}(F(X_1) - F^*)R_0)^{1/2}}{S_0 T^{1/2}}. \quad (48)$$

At this equality point, the minimized upper envelope is

$$\frac{512^{1/4} \varsigma_{\max} S_0^{1/2} R_0^{1/4} \hat{\mathcal{L}}^{1/4} (F(X_1) - F^*)^{1/4}}{T^{1/4}}. \quad (49)$$

Define C_1, C_2 as in Corollary 3.2. Choosing the horizon in (6), Lemma A.3 (or a direct check if $q\mathcal{L}_1 = 0$) gives

$$\frac{\gamma \varsigma_{\max}^2 q \mathcal{L}_1 R_0}{2} + \frac{2\gamma \varsigma_{\max}^2 q \mathcal{L}_1 R_0}{1 - \beta_1} \leq \frac{\varsigma_{\min}}{2}. \quad (50)$$

Then, we reformulate Eq. (5) as

$$\begin{aligned} \frac{1}{T} \sum_{t=1}^T \sum_{l=1}^L \mathbb{E}[\|\nabla F_{(l)}(X_t)\|_*] &\leq \frac{2}{\varsigma_{\min}} \left(\frac{512^{1/4} \varsigma_{\max} S_0^{1/2} R_0^{1/4} \hat{\mathcal{L}}^{1/4} (F(X_1) - F^*)^{1/4}}{T^{1/4}} \right. \\ &\quad + \frac{2\varsigma_{\max} G_0}{C_2 T^{1/2}} \\ &\quad + 2\sqrt{2} \varsigma_{\max} S_0 \left(\frac{C_2^{(1+a)/2}}{T^{(1+a)/4}} + \frac{C_2^a}{T^{a/2}} \right) \\ &\quad \left. + \frac{C_1 \varsigma_{\max}^2 \hat{\mathcal{L}} R_0}{2T^{3/4}} \right). \end{aligned} \quad (51)$$

□

D Proofs of the dimension-independent results

D.1 Proof of Proposition 3.3

Proof. Each polynomial step preserves the representation in the original left and right singular-vector bases and applies p_j to its signed diagonal entries. The final entries are positive by (11), giving the displayed SVD formula for O . For $s_i > 0$, put $r_i = \phi_J(s_i)/s_i$ and $w_i = s_i^2$, so $\sum_i w_i = 1$ and $r_i \in [\ell_J, u_J]$. Direct calculation gives

$$A = \sum_i w_i r_i, \quad B = \sum_i w_i r_i^2, \quad \langle N, O \rangle = \|N\|_F A. \quad (52)$$

The energy estimates follow from $\ell_J \leq r_i \leq u_J$ and $r_i^2 \leq u_J r_i$. The operator-norm estimate follows from the largest output singular value.

For the angular estimate, $(r_i - \ell_J)(u_J - r_i) \geq 0$ implies

$$B \leq (u_J + \ell_J)A - u_J \ell_J.$$

Consequently,

$$\frac{B}{A^2} \leq \frac{u_J + \ell_J}{A} - \frac{u_J \ell_J}{A^2} \leq \frac{(u_J + \ell_J)^2}{4u_J \ell_J}. \quad (53)$$

The last expression is the maximum over $A > 0$, attained at $A = 2u_J\ell_J/(u_J + \ell_J)$. This is the Kantorovich angle bound, derived here directly. Finally,

$$\langle H, O \rangle = \langle N, O \rangle - \langle E, O \rangle \geq \|N\|_{\mathbb{F}} A - \|E\|_{\mathbb{F}} \sqrt{B} \geq A(\|N\|_{\mathbb{F}} - \bar{\rho}_J \|E\|_{\mathbb{F}}).$$

The estimates needed by the convergence theorem also hold for $N = 0$, where $O = 0$. $\square$

D.2 Analytic constants for the fixed quintic

Let $p(s) = as + bs^3 + cs^5$, with the exact decimal coefficients in (16), and let $h(z) = a + bz + cz^2$. Completing the square gives

$$h(z) \geq a - \frac{b^2}{4c} = 0.638614570514\dots > 0, \quad z \geq 0. \quad (54)$$

For completeness, the two positive stationary points of p are $\sqrt{z_-}$ and $\sqrt{z_+}$, where

$$z_{\pm} = \frac{-3b \pm \sqrt{9b^2 - 20ac}}{10c}.$$

Evaluating p at these points and at the endpoints proves the interval inclusions

$$p([0, 5/4]) \subseteq [0, 5/4], \quad p([17/25, 5/4]) \subseteq [17/25, 5/4]. \quad (55)$$

For example, the local maximum is between 1.2023 and 1.2024, the local minimum is between 0.6818 and 0.6819, $p(17/25) = 1.1362138043392$, and $p(5/4) = 1.17909912109375$. These loose rational brackets suffice for (55); no optimization of a high-degree composite polynomial is needed. Also, h is decreasing on $[0, (17/25)^2]$ and $h((17/25)^2) > 1$, so $p(s) \geq s$ for $0 \leq s \leq 17/25$.

Consider an initial $s \in (0, 1]$. While an iterate is below $17/25$, it cannot decrease. Once it reaches $[17/25, 5/4]$, it remains there by (55). If $s \geq 17/25$, its later iterates are at least $17/25 \geq (17/25)s$; if $s < 17/25$, they are at least s . Thus $\phi_J(s) \geq (17/25)s$ for every $J \geq 1$. All iterates lie in $[0, 5/4]$, and on this interval $h(s^2) \leq a$, because $(5/4)^2 < -b/c$. It follows that

$$\frac{\phi_J(s)}{s} = \prod_{j=0}^{J-1} h(\phi_j(s)^2) \leq a^J.$$

The limit at $s \downarrow 0$ is a^J , proving $u_J = a^J$. This proves all of (16). The general theorem permits different coefficients at different steps; for such schedules positivity and reachable intervals must be checked for that schedule rather than inferred from this fixed-coefficient example.

D.3 Proof of Theorem 3.4

Proof. We follow the four steps of the proof of Theorem 3.1 in Appendix B. Only the alignment, update-energy, and joint-drift bounds change. Write

$$E_t^{(l)} = N_t^{(l)} - \nabla_{(l)} F(X_t), \quad H_t = \sum_{l=1}^L \|\nabla_{(l)} F(X_t)\|_{\mathbb{F}}, \quad b_J = \ell_J + u_J,$$

and set $\bar{H} = T^{-1} \sum_{t=1}^T \mathbb{E} H_t$ and $\bar{E} = T^{-1} \sum_{t=1}^T \sum_{l=1}^L \mathbb{E} \|E_t^{(l)}\|_{\mathbb{F}}$.

Step 1: Descent and telescoping. Proposition 3.3 gives $\langle N_t^{(l)}, O_t^{(l)} \rangle \geq \ell_J \|N_t^{(l)}\|_F$ and $\|O_t^{(l)}\|_F \leq u_J$. Splitting the true gradient into $N_t^{(l)} - E_t^{(l)}$ and applying the reverse triangle inequality, just as in the nuclear-norm proof, yields

$$\begin{aligned} \langle \nabla_{(l)} F(X_t), O_t^{(l)} \rangle &\geq \ell_J \|N_t^{(l)}\|_F - u_J \|E_t^{(l)}\|_F \\ &\geq \ell_J \|\nabla_{(l)} F(X_t)\|_F - b_J \|E_t^{(l)}\|_F. \end{aligned} \quad (56)$$

These inequalities also hold when $N_t^{(l)} = 0$ and $O_t^{(l)} = 0$. Moreover, $\sum_l \|O_t^{(l)}\|_F^2 \leq Lu_J^2$. Lemma A.1, $z^q \leq (1-q) + qz$, and $\|\nabla F(X_t)\|_F \leq H_t$ therefore imply

$$F(X_{t+1}) \leq F(X_t) - \gamma \ell_J H_t + \gamma b_J \sum_l \|E_t^{(l)}\|_F + \frac{\gamma^2 Lu_J^2}{2} (\hat{\mathcal{L}} + q\mathcal{L}_1 H_t).$$

Taking expectations, summing over t , and using $F(X_{T+1}) \geq F^*$ gives

$$\left(\ell_J - \frac{\gamma Lu_J^2 q \mathcal{L}_1}{2} \right) \bar{H} \leq \frac{D}{\gamma T} + b_J \bar{E} + \frac{\gamma Lu_J^2 \hat{\mathcal{L}}}{2}. \quad (57)$$

This is the counterpart of (36), with Frobenius stationarity and an unweighted sum of block tracking errors.

Step 2: The Nesterov tracking-error expansion. Put $\varepsilon_t^{(l)} = G_t^{(l)} - \nabla_{(l)} F(X_t)$. The momentum recursion and $M_0^{(l)} = 0$ give exactly the expansion (38), which we write as

$$E_t^{(l)} = -\beta_1^t \beta_2 \nabla_{(l)} F(X_1) + Z_t^{(l)} + V_t^{(l)},$$

where

$$\begin{aligned} Z_t^{(l)} &= (1 - \beta_1 \beta_2) \varepsilon_t^{(l)} + \beta_2 \delta \sum_{k=1}^{t-1} \beta_1^{t-k} \varepsilon_k^{(l)}, \\ V_t^{(l)} &= \beta_1 \beta_2 \sum_{k=2}^t \beta_1^{t-k} (\nabla_{(l)} F(X_{k-1}) - \nabla_{(l)} F(X_k)). \end{aligned}$$

Thus the same three terms as in (39) satisfy

$$\frac{1}{T} \sum_{t=1}^T \mathbb{E} \|E_t^{(l)}\|_F \leq \mathcal{T}_{1,l} + \mathcal{T}_{2,l} + \mathcal{T}_{3,l},$$

where $\mathcal{T}_{1,l}$ is the averaged norm of the initialization term, $\mathcal{T}_{2,l} = T^{-1} \sum_t \mathbb{E} \|Z_t^{(l)}\|_F$, and $\mathcal{T}_{3,l} = T^{-1} \sum_t \mathbb{E} \|V_t^{(l)}\|_F$.

Step 3: Initialization, noise, and gradient drift. The initialization bound uses the same geometric series:

$$\sum_l \mathcal{T}_{1,l} = \frac{\beta_2}{T} \sum_{t=1}^T \beta_1^t G_F \leq \frac{\beta_2 G_F}{T \delta}.$$

For the noise term, conditional unbiasedness eliminates temporal cross terms within each block. The variance bound and geometric series give

$$\begin{aligned} \mathbb{E} \|Z_t^{(l)}\|_F^2 &\leq \left((1 - \beta_1 \beta_2)^2 + \beta_2^2 \delta^2 \sum_{k=1}^{t-1} \beta_1^{2(t-k)} \right) \sigma_{(l)}^2 \\ &\leq \frac{\delta + 2\beta_1(1 - \beta_1 \beta_2)(1 - \beta_2)}{1 + \beta_1} \sigma_{(l)}^2. \end{aligned}$$

Jensen's inequality and $\sqrt{x+y} \leq \sqrt{x} + \sqrt{y}$ imply $\sum_l \mathcal{T}_{3,l} \leq v_\beta S_F$. This estimate does not require independence across layers.

For drift, apply Cauchy–Schwarz across blocks before invoking joint smoothness. In place of the rank-weighted bound in (42), use $\sum_l \|W^{(l)}\|_F \leq \sqrt{L}\|W\|_F$ and $\|X_k - X_{k-1}\|_F \leq \gamma u_J \sqrt{L}$:

$$\begin{aligned} \sum_l \mathcal{T}_{3,l} &\leq \frac{\beta_1 \beta_2 \sqrt{L}}{T} \sum_{t=1}^T \sum_{k=2}^t \beta_1^{t-k} \mathbb{E} \|\nabla F(X_{k-1}) - \nabla F(X_k)\|_F \\ &\leq \frac{\gamma \beta_1 \beta_2 L u_J}{T} \sum_{t=1}^T \sum_{k=2}^t \beta_1^{t-k} \mathbb{E} [\hat{\mathcal{L}} + q \mathcal{L}_1 H_k] \\ &\leq \frac{\gamma \beta_2 L u_J}{\delta} (\hat{\mathcal{L}} + q \mathcal{L}_1 \bar{H}). \end{aligned} \tag{58}$$

The second inequality uses Assumption 2.3 with base point X_k and $X_k - X_{k-1} = -\gamma O_{k-1}$. The last exchanges the finite sums and bounds $\beta_1 \sum_{t=k}^T \beta_1^{t-k} \leq \delta^{-1}$. Combining the three terms gives the counterpart of (43):

$$\bar{E} \leq \frac{\beta_2 G_F}{T \delta} + v_\beta S_F + \frac{\gamma \beta_2 L u_J \hat{\mathcal{L}}}{\delta} + \frac{\gamma \beta_2 L u_J q \mathcal{L}_1}{\delta} \bar{H}. \tag{59}$$

Step 4: Substitution and absorption. Substituting (59) into (57) and moving its last term to the left gives

$$\begin{aligned} &\left(\ell_J - \frac{\gamma L u_J^2 q \mathcal{L}_1}{2} - \frac{\gamma \beta_2 L u_J b_J q \mathcal{L}_1}{\delta} \right) \bar{H} \\ &\leq \frac{D}{\gamma T} + \frac{\beta_2 b_J G_F}{T \delta} + b_J v_\beta S_F + \frac{\gamma \beta_2 L u_J b_J \hat{\mathcal{L}}}{\delta} + \frac{\gamma L u_J^2 \hat{\mathcal{L}}}{2}. \end{aligned}$$

Using $\beta_2 \leq 1$ in the coefficient on the left and $\bar{H} \geq 0$ yields (19). This completes the same descent, tracking, and absorption argument as for Theorem 3.1, with the finite-map bounds supplying all geometric constants. $\square$

D.4 Proof of Corollary 3.5

Proof. As in the proof of Corollary 3.2, the schedules balance the three leading terms of the convergence bound. With $b_J = \ell_J + u_J$, the corresponding envelope is

$$\Phi_F(\gamma, \delta) = \frac{D}{\gamma T} + b_J S_F \sqrt{\delta} + \frac{\gamma L u_J b_J \hat{\mathcal{L}}}{\delta}.$$

For $D, S_F > 0$, weighted Young's inequality gives

$$\Phi_F(\gamma, \delta) \geq \left(\frac{64 D (b_J S_F)^2 L u_J b_J \hat{\mathcal{L}}}{T} \right)^{1/4},$$

with equality when

$$\frac{4D}{\gamma T} = 2 b_J S_F \sqrt{\delta} = \frac{4 \gamma L u_J b_J \hat{\mathcal{L}}}{\delta}.$$

These balancing equations give $\gamma \asymp T^{-3/4}$ and $\delta \asymp T^{-1/2}$. The corollary allows any positive γ_0, δ_0 with these exponents; the verification below also covers $D = 0$ or $S_F = 0$ without using the equality conditions.

Under (20), $\delta = \delta_0 T^{-1/2}$ and

$$1 - \beta_1 \beta_2 = \delta + \beta_1 \delta^a, \quad v_\beta \leq \sqrt{\delta} + \sqrt{2}(\delta^{(1+a)/2} + \delta^a).$$

The condition $\Delta_{\gamma, \beta_1}^F \geq \ell_J/2$ follows from $A_J T^{-1/4} + B_J T^{-3/4} \leq 1$. The sufficient threshold (23) follows by the same elementary argument as Lemma A.3; if $A_J = B_J = 0$ there is nothing to check. Substitution into (19), followed by division by $\ell_J/2$, proves (21). For $a > 1/2$, all exponents except the leading $1/4$ are strictly larger than $1/4$. Finally $\|\nabla F(X_t)\|_F \leq H_t$, so the Euclidean/Frobenius stationarity claim follows directly.

To prove (24), put $g = \|\nabla F(X_1)\|_F$ and $L_g = \mathcal{L}_0 + \mathcal{L}_1 g^q > 0$. Applying Lemma A.1 to $Y = X_1 - \nabla F(X_1)/L_g$ and using $F(Y) \geq F^*$ gives $g^2 \leq 2DL_g \leq 2D(\hat{\mathcal{L}} + q\mathcal{L}_1 g)$. Solving this quadratic inequality and using $G_F \leq \sqrt{L_g}$ proves the claim. This argument concerns the stated optimization assumptions, not a network-width limit. $\square$

D.5 Scope of the angular refinement

For one nonzero block momentum N with true gradient H and error $E = N - H$, (15) gives the valid implication

$$\|N\|_F \geq 2\bar{\rho}_J \|E\|_F \implies \langle H, O \rangle \geq \frac{1}{2} \ell_J \|N\|_F.$$

If the displayed sufficient condition fails, the triangle inequality gives

$$\|H\|_F < (2\bar{\rho}_J + 1) \|E\|_F.$$

Failure of this test is not equivalent to actual negative alignment; it only means that this sufficient lower bound is unavailable. Nor does smallness of one block gradient imply stationarity of the complete parameter tuple.

To make precise the stronger small-constant argument, consider the *single-block* case and separately assume spectral-direction smoothness

$$F(X + V) \leq F(X) + \langle \nabla F(X), V \rangle + \frac{\mathcal{L}_{\text{sp}}}{2} \|V\|_{\text{op}}^2. \quad (60)$$

This is an additional hypothesis, not a consequence of Assumption 2.3 with a dimension-independent constant. Define $h_\gamma = \mathcal{L}_{\text{sp}} \gamma c_J^2 / \ell_J$. If $\|N_t\|_F \geq 2\bar{\rho}_J \|E_t\|_F + h_\gamma$, (15) and (60) yield

$$F(X_{t+1}) \leq F(X_t) - \frac{1}{2} \gamma \ell_J \|N_t\|_F.$$

On any sample path for which $\max_{t \leq T} \|E_t\|_F \leq \bar{e}$, either this condition first fails, or it holds at every step. In the first case the gradient at the failed test is at most $(2\bar{\rho}_J + 1)\bar{e} + h_\gamma$; in the second case telescoping gives $\min_t \|\nabla F(X_t)\|_F \leq 2D/(\gamma \ell_J T) + \bar{e}$. Therefore the precise deterministic certificate is

$$\min_{t \leq T} \|\nabla F(X_t)\|_F \leq \frac{2D}{\gamma \ell_J T} + (2\bar{\rho}_J + 1)\bar{e} + \frac{\mathcal{L}_{\text{sp}} \gamma c_J^2}{\ell_J}. \quad (61)$$

An averaged or pointwise expected error bound is not a bound on $\max_t \|E_t\|_F$, and it cannot simply be substituted for $\bar{e}$. Initialization transients must also be included in any such uniform bound. Equation (61) is thus a conditional geometric certificate, not a second stochastic convergence claim under the three stated assumptions.

The size of u_J cannot in general be removed from a uniform Frobenius update bound by sharpening a scalar inequality. Let $A_0 = \phi'_J(0) > 0$ and take rank r with all normalized singular values $s_i = r^{-1/2}$. Then

$$A(N) = \sqrt{r} \phi_J(r^{-1/2}) \longrightarrow A_0, \quad B(N) = r \phi_J(r^{-1/2})^2 \longrightarrow A_0^2.$$

Thus any uniform bound $\|O\|_{\mathbb{F}} \leq U$ needs $U \geq A_0$, and any $B \leq KA$ needs $K \geq A_0$. For the fixed quintic, $A_0 = u_J$. The large universal magnitude constant and the absence of dimension factors are distinct issues.