

A POSITIVE RESOLUTION OF THE GAP-ENTROPY CONJECTURE

P. M. ARONOW, NATHAN KALLUS, AND PATRICK LOPATTO

ABSTRACT. We prove the gap-entropy conjecture for fixed-confidence best-arm identification with independent unit-variance Gaussian arms, means in $[0, 1]$, and a unique optimal arm. For each suboptimal arm i , let $\Delta_i = \mu_* - \mu_i$ be its gap from the optimal mean, and write $H = \sum_{i \neq *} \Delta_i^{-2}$. Let p_r be the fraction of H contributed by arms with $2^{-(r+1)} < \Delta_i \leq 2^{-r}$, and let $\text{Ent}(I) = \sum_{r: p_r > 0} p_r \log(1/p_r)$. Among all algorithms that identify the optimal arm with probability at least $1 - \delta$ on every Gaussian instance, the optimal expected number of samples on a given instance, averaged over all permutations of the arm labels, is within absolute constant factors of $H(\log(1/\delta) + \text{Ent}(I))$. Moreover, there is an algorithm, independent of the instance, whose expected number of samples is bounded by a constant multiple of this quantity plus $g^{-2} \log \log(e^e/g)$, where $g = \min_{i \neq *} \Delta_i$ is the gap to the closest competitor.

1. INTRODUCTION

Experiments often compare several alternatives in order to choose the one with the best mean reward. Adaptive sampling, wherein at each step an algorithm pulls one of $K \geq 2$ arms and immediately receives a sample from its unknown reward distribution to inform future pulls, can significantly improve performance. In regret minimization, avoiding apparently inferior arms in future sampling can help attain higher cumulative rewards over the experiment, and in best-arm identification, concentrating future sampling on apparently close competitors for best can find the best faster. Fixed-confidence best-arm identification formalizes this problem by seeking to identify the best arm with as few samples as possible while controlling the probability of returning the wrong arm for all admissible reward distributions [5, 10].

The difficulty depends on the gaps between the mean rewards $(\mu_1, \dots, \mu_K) \in [0, 1]^K$, where $\mu_* = \max_i \mu_i$ is uniquely attained. In the unit-variance Gaussian model, a suboptimal arm $i \neq *$ with gap $\Delta_i = \mu_* - \mu_i$ from the optimal arm requires $\Omega(\Delta_i^{-2} \log(1/\delta))$ samples to distinguish it from the optimal arm at error level δ . Summing these costs gives the lower bound $\Omega(H \log(1/\delta))$, where $H = \sum_{i \neq *} \Delta_i^{-2}$ is the sum of the inverse squared gaps, capturing the baseline cost of distinguishing the best arm if we know up to orders of magnitude how well separated it is from each other arm [10, 12]. This does not, however, reveal the cost of discovering this scale of the gaps, which matters even when the error level δ is fixed.

Exponential-Gap Elimination [9] and lil'UCB [8] achieved upper bounds with iterated-log dependence on gaps of the form $O(H \log(1/\delta) + \sum_{i \neq *} \Delta_i^{-2} \log \log(e^e/\Delta_i))$. Chen and Li [1] subsequently improved this bound, capping each arm's iterated-logarithmic factor at $\log \log K$, apart from a $g^{-2} \log \log(1/g)$ term where $g = \min_{i \neq *} \Delta_i$. They also proved that for sufficiently small δ and large K , any δ -correct algorithm has a Gaussian instance requiring $\Omega(H \log \log K)$ samples. This raised the question of how this unavoidable adaptation cost depends on the particular configuration of gaps.

Chen and Li [2] proposed that the additional instance-dependent cost is described by *gap entropy*. For an instance I , peel the suboptimal arms into geometrically spaced shells according to their gaps,

Date: September 9, 2026.

This paper was written by the authors with the assistance of large language models, which were used to suggest mathematical arguments, draft and revise exposition, and write code for computational verification. P. L. was partially supported by NSF grant DMS-2450004.

and assign to each shell the fraction of H contributed by its arms. The entropy of the resulting probability vector, $\text{Ent}(I)$, measures how widely the required sampling effort is distributed across scales.

Chen and Li conjectured (their conjecture 3.5) that, among δ -correct algorithms, the smallest expected sample size achievable on a given instance I (averaged over permutations of its arm labels) is of order $H(\log(1/\delta) + \text{Ent}(I))$. (A different algorithm may attain the benchmark for each I , provided it remains δ -correct on all instances.) They further conjectured (their conjecture 3.2) that a single algorithm can attain this instance-wise benchmark simultaneously on all instances, up to an additive term of order $g^{-2} \log \log(1/g)$. Such an adaptation cost is unavoidable already with two arms: adapting to an unknown gap that may be arbitrarily small incurs an iterated-logarithmic cost along some sequences of instances, even though this cost is not required at every fixed instance [2, 6].

The conjectures seek nonasymptotic bounds that hold uniformly in the instance and δ . By contrast, Track-and-Stop [7] already achieves asymptotic optimality as $\delta \downarrow 0$ with the instance fixed. In that limit, the gap entropy of the fixed instance is eventually dominated by $\log(1/\delta)$. An asymptotically optimal leading constant therefore does not determine the additional cost when the gap configuration and the confidence parameter vary together. Simchowitz, Jamieson, and Recht [13] studied the complexity distinctions between fixed δ and $\delta \downarrow 0$ asymptotics, including the cost of learning a sampling allocation in structured models. Cho and Kallus [4] relax exact error control to asymptotic validity in the $\delta \downarrow 0$ regime, enabling the use of strong invariance principles to match the optimal exact-error Gaussian benchmark with known variances even under nonparametric reward distributions.

Chen, Li, and Qiao [3] made progress toward the conjectures, obtaining the upper bound $O(H(\log(1/\delta) + \text{Ent}(I)) + g^{-2} \log \log(1/g) \text{polylog}(K, 1/\delta))$ (their theorem 1.11) and, for Gaussian instances whose gaps are powers of two, a lower bound of $\Omega(H(\log(1/\delta) + \text{Ent}(I)))$ for algorithms whose expected sample cost does not increase when suboptimal arms are deleted (their theorem 1.12 and corollary 1.13).

In this paper, we answer both conjectures in the positive. We remove both the extra polylogarithmic factor from the upper bound (which applies to arbitrary 1-sub-Gaussian reward distributions) and the restrictions on instances and algorithms in the lower bound (for the Gaussian model).

1.1. Main results. For $K \geq 2$, a K -armed bandit instance $I = (\nu_1, \dots, \nu_K)$ has reward distributions ν_i with means $\mu_i \in [0, 1]$. The index of the largest mean, denoted by $*$, is assumed unique. Each arm has an sequence of independent samples with law ν_i , and the sequences are independent across arms. Pulling an arm reveals its next sample.

Let $\mathcal{S}_K$ be the class of such unique-optimum instances with 1-sub-Gaussian rewards, meaning that

$$\mathbb{E}_{X \sim \nu_i} \exp(\lambda(X - \mu_i)) \leq \exp(\lambda^2/2) \quad (\lambda \in \mathbb{R}, i \in [K]).$$

The algorithm knows that the rewards are 1-sub-Gaussian, but not their distributions or means. Let $\mathcal{G}_K \subset \mathcal{S}_K$ be the Gaussian subclass with $\nu_i = N(\mu_i, 1)$.

For either class, write

$$m = \mu_*, \quad \Delta_i = m - \mu_i \ (i \neq *), \quad H = \sum_{i \neq *} \Delta_i^{-2}, \quad g = \min_{i \neq *} \Delta_i, \quad D = g^{-2}.$$

For $r = 0, 1, \dots$, let

$$G_r = \{i \neq * : 2^{-(r+1)} < \Delta_i \leq 2^{-r}\}, \quad H_r = \sum_{i \in G_r} \Delta_i^{-2}, \quad p_r = H_r/H.$$

For a probability vector $q \in [0, 1]^{\mathbb{N}_0}$, write $\text{Ent}(q) = \sum_{r: q_r > 0} q_r \log(1/q_r)$ for its Shannon entropy. The *gap entropy* of instance I is defined as

$$\text{Ent}(I) = \text{Ent}(p). \quad (1.1)$$

It is zero when all gaps lie on the same geometric scale. If d groups contribute equally to H , it equals $\log d$. More generally, writing $\mathcal{U} = \{r : H_r > 0\}$ for the finite set of indices of the nonempty gap groups, we have $0 \leq \text{Ent}(I) \leq \log |\mathcal{U}|$.

All logarithms are natural unless otherwise indicated. Constants denoted by c or C are positive and absolute and may change between occurrences. Constants used as algorithm parameters are fixed within each construction. We write $[K] = \{1, \dots, K\}$.

An algorithm is δ -correct on a model class if, on every instance in that class, it stops almost surely and returns the optimal arm with probability at least $1 - \delta$. Write T_A for its total number of samples. For a permutation $\pi \in S_K$ of the labels, let πI be the relabeled instance. For $I \in \mathcal{G}_K$, following [2, definition 3.1], define

$$\mathcal{L}(I, \delta) = \inf_{A: \delta\text{-correct on } \mathcal{G}_K} \frac{1}{K!} \sum_{\pi \in S_K} \mathbb{E}_{\pi I} T_A. \quad (1.2)$$

The averaging removes any advantage from a favorable labeling. The infimum is taken separately for each Gaussian instance I , but every algorithm in it must be δ -correct on the Gaussian class $\mathcal{G}_K$. [Appendix A](#) compares these conventions with those of [2] and explains why our results imply their original conjectures.

We now present our main results.

Theorem 1.1. *There exist absolute constants $c, C > 0$ such that, for every $I \in \mathcal{G}_K$ and $0 < \delta < 0.1$,*

$$cH(\log(1/\delta) + \text{Ent}(I)) \leq \mathcal{L}(I, \delta) \leq CH(\log(1/\delta) + \text{Ent}(I)).$$

The lower and upper bounds are proved in [Sections 2](#) and [3](#), respectively. The lower bound is proved directly for the original Gaussian instance, after averaging over permutations of the arm labels, without first passing to a subinstance obtained by deleting arms. The matching upper bound uses an algorithm selected for that target instance. More generally, [Proposition 3.1](#) constructs, for each instance in $\mathcal{S}_K$, an algorithm that is δ -correct on $\mathcal{S}_K$ and attains the same bound on every relabeling of the instance. The next theorem gives one algorithm for the whole sub-Gaussian class.

Theorem 1.2. *There exists one algorithm, given only K and δ , that is δ -correct on $\mathcal{S}_K$ and satisfies, for every $I \in \mathcal{S}_K$ and $0 < \delta < 0.1$,*

$$\mathbb{E}_I T_A \leq C(H(\log(1/\delta) + \text{Ent}(I)) + D \log \log(e^e/g)).$$

The constant C is absolute.

The algorithm is defined in [Section 4](#); its correctness and bounds on the expected sample size are proved in [Sections 5](#) and [6](#), respectively. The additive term depends only on the smallest gap, with no further factor involving K or δ . Specializing the uniform algorithm to $\mathcal{G}_K$ and applying [Theorem 1.1](#) gives the near-optimality on every individual instance conjectured in [2, conjecture 3.2].

Corollary 1.3. *For every $I \in \mathcal{G}_K$ and $0 < \delta < 0.1$, the algorithm in [Theorem 1.2](#) satisfies*

$$\mathbb{E}_I T_A \leq C(\mathcal{L}(I, \delta) + D \log \log(e^e/g)).$$

The proof is given at the end of [Section 6](#). [Theorem 1.1](#) proves [2, conjecture 3.5], and together with [Corollary 1.3](#) proves [3, conjecture 1.9]; see [Appendix A](#) for details.

1.2. Ideas of the proof. A short allocation calculation explains why entropy appears. Suppose the group weights H_r were known and group G_r were assigned error budget δ_r . The corresponding sampling cost would have the form $\sum_r H_r \log(1/\delta_r)$. Under the constraint $\sum_r \delta_r \leq \delta$, this expression is minimized by $\delta_r = \delta p_r$, giving

$$\inf_{\substack{\delta_r > 0 \\ \sum_{r \in \mathcal{U}} \delta_r \leq \delta}} \sum_{r \in \mathcal{U}} H_r \log(1/\delta_r) = H(\log(1/\delta) + \text{Ent}(I)). \quad (1.3)$$

In particular, if d groups contribute equally to H , then $\delta_r = \delta/d$ and the additional cost is $H \log d$.

This calculation, used to motivate the conjecture in [2, section 4], does not by itself prove a lower bound for an arbitrary algorithm or give an algorithm when the weights are unknown. We first show that every algorithm correct on the Gaussian class must pay the entropy cost. We then attain this cost using an algorithm selected for a fixed target instance. Finally, we construct an algorithm that does not depend on the instance and bound the additional cost of adaptation.

1.2.1. The lower bound. The term $H \log(1/\delta)$ follows from the standard bandit change-of-measure argument; see [10, lemma 1]. The remaining task is to establish the additional entropy cost for an arbitrary correct algorithm. For each gap group G_r , we choose one of its arms uniformly and examine how many times the algorithm samples that arm. Since arms in G_r have gaps of order 2^{-r} , their sampling complexity is on the scale 4^r , and we associate G_r with an event in which this sample count lies in a suitable range around that scale. Widely separated scales give disjoint ranges, and the main difficulty is to compare the probabilities of these events under one common reference law.

Fix a Gaussian instance and an arbitrary algorithm A that is δ -correct on the Gaussian class. Define a new algorithm by first applying a uniformly random permutation to the arm labels, running A on the relabeled instance, and then applying the inverse permutation to A 's output. The new algorithm is still δ -correct, treats all labelings symmetrically, and has expected sample cost on every labeling equal to the average cost of A over all labelings of the original instance. Let u_i be its expected number of samples from arm i of the target instance, and let

$$\alpha_r = \frac{\sum_{i \in G_r} u_i}{|G_r| 4^r}, \quad r \in \mathcal{U}.$$

An arm in G_r has inverse squared gap comparable to 4^r . Thus α_r is the average sample count in the group divided by this sampling scale.

Fix $a \geq 1$. For each group with $\alpha_r \leq a$, let P_r be the law of the algorithm's run when the optimal arm has label 1, a uniformly chosen arm from G_r has label 2, and the remaining arms are uniformly relabeled. Let N_2 count the samples from label 2, and let $E_r(a)$ be the event that the algorithm returns label 1 with $c_0 4^r \leq N_2 \leq 4a 4^r$, for a sufficiently small absolute constant c_0 . Since $\mathbb{E}_{P_r} N_2 = \alpha_r 4^r \leq a 4^r$, Markov's inequality gives $P_r(N_2 > 4a 4^r) \leq 1/4$.

For the lower tail, raise label 2's mean to m , producing a tied reference law Q_r . Correctness on nearby instances in which label 2 is uniquely optimal forces the probability of returning label 1 under Q_r to be at most δ . Before label 2 has been sampled $c_0 4^r$ times, the KL divergence between the corresponding history distributions is bounded by an absolute constant times c_0 . A finite-horizon change-of-measure argument therefore shows that returning label 1 this early also has small probability under P_r . Combining the two tail bounds with correctness gives $P_r(E_r(a)) \geq 1/2$; the details are in [Section 2](#).

For sufficiently separated indices r, r' , the ranges $[c_0 4^r, 4a 4^r]$ and $[c_0 4^{r'}, 4a 4^{r'}]$ are disjoint. But their event probabilities have been bounded under different laws P_r , and the tied reference laws Q_r also depend on r . To compare different groups, we construct a reference law that can be used for all of them. Fix a nonempty group G_s . For $r > s$, let i denote the arm placed at label 2, choose j uniformly from G_s , independently of the labeling, and change the two means by

$$(\mu_i, \mu_j) \mapsto (m, \mu_i).$$

The second change restores the suboptimal mean removed by the first. Labels 1 and 2 are now tied, and the other labels carry exactly the original suboptimal means with μ_j omitted. After averaging over the random choices, those remaining means are uniformly arranged on labels $3, \dots, K$. The resulting law, denoted by Q_s , therefore depends on s but not on r . For $r = s$, raising the selected arm to m alone gives the same Q_s . Note that each of these resulting instances is inadmissible because labels 1 and 2 are tied for best. While δ -correctness or almost-sure stopping are therefore not required for A on these, the law of the algorithm's run is still well-defined and its finite-time answer probabilities are controlled by its required δ -correct behavior on nearby admissible instances where label 1's mean is slightly reduced, making label 2 uniquely optimal.

A Gaussian change-of-measure calculation bounds the KL divergence from P_r to Q_s , at any finite horizon, by $(\alpha_r + \alpha_s)/2$. If any group satisfies $\alpha_r \leq a$, choose $s = \min\{r \in \mathcal{U} : \alpha_r \leq a\}$. The divergence bound is then at most a for every such group, and $P_r(E_r(a)) \geq 1/2$ implies $Q_s(E_r(a)) \geq e^{-2a}/4$. Under this one law, events from sufficiently separated groups are disjoint. Since each has probability at least $e^{-2a}/4$, any such collection contains at most $4e^{2a}$ groups. Partitioning the indices into $O(a)$ separated collections gives the required counting bound for $a \geq 1$. The standard per-arm lower bound gives $\alpha_r > 1$ for every r , so integrating this counting bound, as carried out in [Section 2](#), yields

$$\sum_r e^{-8\alpha_r} \leq 1.$$

Since the weights $e^{-8\alpha_r}$ have total mass at most one, normalizing them and applying Gibbs's inequality gives

$$H \text{Ent}(I) \leq 8 \sum_r H_r \alpha_r \leq 32 \sum_{i \neq *} u_i.$$

Since $\sum_{i \neq *} u_i$ is the expected number of samples from the suboptimal arms, it is at most the expected total number of samples. Hence, writing T for the total number of samples

$$\mathbb{E}T \geq \frac{1}{32} H \text{Ent}(I).$$

Together with the standard lower bound $\mathbb{E}T \geq cH \log(1/\delta)$, this gives

$$\mathbb{E}T \geq c'H(\log(1/\delta) + \text{Ent}(I))$$

for an absolute constant $c' > 0$, proving the lower half of [Theorem 1.1](#).

1.2.2. The instance-wise upper bound. For each fixed target instance $I \in \mathcal{S}_K$, we may choose an algorithm specifically for I , provided it remains δ -correct on all of $\mathcal{S}_K$. The algorithm's definition includes the indices and sizes of the target's nonempty gap groups and a threshold $\tau \in (m - g/2, m - g/3)$. Thus τ lies strictly between the largest mean m and the second-largest mean $m - g$, at distance comparable to g from both. On the target, the optimal arm is the unique arm with mean above τ . Thus it suffices to eliminate all arms whose means lie below τ . We do this through a sequence of independent trials. Each trial may return an arm or return without an answer.

Choose probability weights q_r proportional to $|G_r|4^r$, so that q_r is comparable to p_r . Given a parameter η that will serve as the trial's error allowance, the trial starts with all arms active and cycles through the active arms, taking one sample from each in turn. For each arm, we combine one exponential test statistic for each supplied gap scale, using mixture weight q_r for scale r . An arm is eliminated when this combined statistic reaches its rejection threshold. For a suboptimal target arm $i \in G_r$, let U_i be the number of samples from that arm before it is eliminated. We show that

$$\mathbb{E}U_i \leq C\Delta_i^{-2}[\log(1/\eta) + \log(1/q_r)].$$

Thus the mixture weight q_r enters the sample bound through the additive term $\log(1/q_r)$. Since q_r is comparable to p_r , summing over the suboptimal arms gives

$$\sum_{i \neq *} \mathbb{E} U_i \leq CH(\log(1/\eta) + \text{Ent}(I)).$$

Because the procedure takes one sample from every active arm in each cycle through the active set, the optimal arm is sampled at most $\max_{i \neq *} U_i$ times before at most one arm remains. Hence the total number of samples used during elimination is at most

$$\sum_{i \neq *} U_i + \max_{i \neq *} U_i \leq 2 \sum_{i \neq *} U_i.$$

Although the algorithm is chosen using information from the target instance, it must remain δ -correct when run on every instance in $\mathcal{S}_K$, including instances whose means and gap groups differ from those of the target. There are two cases. If the input has an arm with mean at least τ , then the optimal arm also has mean at least τ . An incorrect output is therefore possible only if the tests eliminate the optimal arm, which happens with probability at most η . If instead every input mean is below τ , then before returning the sole remaining arm the trial takes fresh samples from it and requires its empirical mean to exceed a level slightly above τ . We choose the number of confirmation samples so that, by the sub-Gaussian tail bound, this event has probability at most η . Thus in this case the trial returns any arm with probability at most η , and in particular returns an incorrect arm with probability at most η .

We impose a deterministic sample limit on each trial and repeat independent trials with geometrically decreasing error allowances whose sum is at most $\delta/2$. On the target instance, each trial returns the optimal arm with probability at least $3/4$, so the probability of reaching later trials decreases geometrically while their sample limits grow only linearly with the trial index. The repeated-trial procedure need not stop on every other instance. To guarantee almost-sure stopping on the full class, we interleave it with an independent $\delta/2$ -correct confidence-interval procedure and return the first answer. This at most doubles the target-instance cost, up to one sample, while the two error guarantees together give δ -correctness.

1.2.3. The uniform upper bound. The preceding instance-wise construction uses the instance's gap distribution to choose its testing scales and error allocation. We now construct one algorithm, given only K and δ , that must discover this information from the data. It runs independent stages with geometrically increasing budgets $B_t = 100^t$. Each stage begins with all K arms and either returns an arm or rejects and passes control to the next stage. A deterministic sample limit of order B_t bounds the cost of the stage on every sample path.

Within a stage, round r works at accuracy $\varepsilon_r = 2^{-r}$. A Median call selects an arm whose mean is within $\varepsilon_r/8$ of the largest active mean, with probability at least 0.99. The algorithm estimates this arm's mean and uses a Fraction call to determine whether sufficiently many active arms lie substantially below it. If so, an Eliminate call removes arms with low estimated means; otherwise the active set is unchanged. The stage returns as soon as only one arm remains. It rejects if either of its two budget tests is triggered, its final permitted round is completed without a singleton, or its sample limit is reached.

For an active set S_r , the sampling-cost scale at accuracy ε_r is

$$w_r = |S_r| \varepsilon_r^{-2} = |S_r| 4^r.$$

This observable quantity replaces the unknown gap-group weights used by the instance-wise algorithm. Set $\eta = c\delta$, where $c > 0$ is a sufficiently small absolute constant. An Eliminate call in round r receives an error allowance proportional to w_r/B_t . The stage controls both the sum of these allowances and the cumulative sum of

$$z_r = w_r \log \frac{B_t}{\eta w_r}.$$

Up to an absolute constant, z_r bounds the expected cost of the Median and Eliminate calls in that round.

The main cost estimate controls how long each suboptimal arm remains active. In the comparison process of [Section 5](#), if Median succeeds in round r and every Eliminate call through that round is valid, the next active set has size at most an absolute constant times

$$1 + |\{i \neq * : \Delta_i \leq 2\varepsilon_r\}|.$$

If Median fails for j additional rounds, the probability of that delay is at most 0.01^j , whereas the sampling scale grows by 4^j . The series

$$\sum_{j \geq 0} (4 \cdot 0.01)^j$$

is finite, so these delays change the expected cost by only an absolute factor. Consequently, arms with gaps of order 2^{-r} contribute mainly near round r . For $B_t \geq H$, sum z_r over executed rounds up to and including the first invalid Eliminate call, or until an earlier stage termination. The active-set and entropy estimates in [Lemma 5.1](#) give

$$\mathbb{E} \sum_r z_r \leq CH [\log(1/\delta) + \text{Ent}(I) + 1 + \log(B_t/H)].$$

The outer Mean and Fraction calls require a separate summation. The separation between the two fraction thresholds is indexed by the number of rounds remaining. Reindexing from the final round turns the resulting polynomial factor into a summable correction to the geometric round costs; see [Lemma 6.1](#). The resulting sample costs also grow logarithmically with the stage index. The calculation in [Lemma 6.2](#) accounts for this dependence through the term $CH(\log(1/\delta) + \text{Ent}(I))$ together with

$$CD \log \log(e^e/g).$$

Correctness across stages requires more than a direct union bound, because each stage receives the same total budget for elimination errors. We again use the comparison process. For a budget B , call the optimal arm and every suboptimal arm satisfying $\Delta_i^{-2} > B$ hard; call the remaining suboptimal arms easy. A wrong output requires the optimal arm and all but possibly one of the hard suboptimal arms to be removed. Every removal of a hard suboptimal arm occurs before its gap scale is reached. The product guarantee for Eliminate controls these joint premature removals. When there are few hard arms, a localized estimate instead uses

$$B^{-1} \sum_{i: i \text{ easy}} \Delta_i^{-2}.$$

When this ratio is large, an exponential-moment argument shows that acceptance itself is unlikely. Together with the bound on disagreement between the actual and comparison processes, these estimates control the total error across stages; see [Section 5](#).

Finally, [Lemma 6.2](#) shows that a stage accepts with probability at least 199/200 once

$$B_t \geq C \{H(\log(1/\delta) + \text{Ent}(I)) + D \log \log(e^e/g)\}.$$

Beyond this point, the probability of reaching the next stage decreases by a factor at most 1/200, while its sample limit increases by a factor of 100. The successive expected costs therefore decrease by a factor at most 1/2. This proves both almost-sure stopping and the claimed bound on the expected sample size. The construction modifies [[3](#), algorithms 4–5] and is specified in [Section 4](#); its correctness and cost are proved in [Sections 5](#) and [6](#).

2. THE INSTANCE-WISE LOWER BOUND

Throughout this section the instance belongs to $\mathcal{G}_K$, and δ -correctness is required on $\mathcal{G}_K$. We prove the lower bound in [Theorem 1.1](#). The term $H \log(1/\delta)$ follows from the usual Gaussian change of measure. For the entropy term, we compare events defined by the number of samples from a selected arm across different gap groups, using a common law with two tied optimal arms.

The conversion of these sample-count bounds into an entropy lower bound parallels the entropy lower-bound argument of Chen, Li and Qiao [\[3, lemma 4.1 and appendix D\]](#). The additional issue here is to construct a common reference law while evaluating all expected sample counts on permutations of the original instance.

The proof of the following proposition appears at the end of this section.

Proposition 2.1. *For every $I \in \mathcal{G}_K$ and $0 < \delta < 0.1$, write $b_\delta = \text{kl}(1 - \delta, \delta)$. Then*

$$\mathcal{L}(I, \delta) \geq \max\{2b_\delta H, H \text{Ent}(I)/32\}, \quad (2.1)$$

where $\text{kl}(p, q) = p \log(p/q) + (1 - p) \log((1 - p)/(1 - q))$.

Since $b_\delta = (1 - 2\delta) \log((1 - \delta)/\delta) \asymp \log(1/\delta)$ uniformly over the stated confidence range, the lower half of [Theorem 1.1](#) follows.

2.1. Symmetry and changes of measure. Fix $I \in \mathcal{G}_K$ and let A be an arbitrary algorithm that is δ -correct on $\mathcal{G}_K$. Randomly permuting labels before running A , and undoing the permutation on its answer, gives a permutation-equivariant algorithm B such that

$$\mathbb{E}_{\pi I} T_B = \frac{1}{K!} \sum_{\rho \in S_K} \mathbb{E}_{\rho I} T_A \quad \text{for every } \pi. \quad (2.2)$$

The algorithm B is still δ -correct. It is useful to distinguish an arm's identity from its label: the identities refer to the fixed instance, whereas the labels specify where the arms are presented to the algorithm. Distinct identities are retained even when their means coincide. Let u_i be the expected number of samples from identity i under B . By permutation equivariance, the expected number of samples allocated to a given arm identity does not depend on the label at which that identity is placed. Thus, for every permutation π ,

$$\mathbb{E}_{\pi I} N_{\pi(i)} = u_i.$$

Consequently, averaging over any of the random labelings used below does not alter these expectations.

If the right side of [\(2.2\)](#) is infinite there is nothing to prove. Henceforth assume $\mathbb{E}_I T_B < \infty$, and suppress the subscript B .

We use the standard finite-horizon likelihood-ratio calculation underlying the bandit change-of-measure inequality; see [\[10, lemma 1 and appendix A.1\]](#). For a deterministic integer M , write $P_\mu^{(M)}$ for the distribution of the recorded history up to $T \wedge M$ under mean vector μ . For two fixed mean vectors μ, λ , the chain rule gives

$$\text{KL}(P_\mu^{(M)} \| P_\lambda^{(M)}) = \frac{1}{2} \sum_i (\mu_i - \lambda_i)^2 \mathbb{E}_\mu N_i(T \wedge M). \quad (2.3)$$

Here $N_i(t)$ counts samples from the arm at label i by time t . Write $N_i = N_i(T)$ for its total number of samples, and let $\hat{i}$ denote the returned label. On $\{T = \infty\}$, set $\hat{i} = 0$ and interpret $N_i(T)$ as $\lim_{t \rightarrow \infty} N_i(t)$. The history at time $T \wedge M$ records whether the algorithm has stopped and, if so, its output, as well as whether truncation occurred. Since the algorithm uses the same sampling, randomization, and stopping rules under both models, these decisions do not contribute additional KL divergence.

We will also use [\(2.3\)](#) after truncation at bounded stopping times and after introducing auxiliary random variables having the same law under both models; these follow from the same chain-rule

calculation and data processing. Whenever we use a tied reference law, we first work at a fixed finite horizon and only then pass to the limit. Thus we never need to assume that the algorithm stops when the two arms are tied.

We first derive the standard lower bound for the expected number of samples from a single suboptimal arm; see [10, lemma 1]. We claim that

$$u_i \geq 2b_\delta \Delta_i^{-2}, \quad \mathbb{E}_I T \geq 2b_\delta H. \quad (2.4)$$

For an alternative instance with a different optimal arm, apply data processing to the event that the algorithm returns the original optimal arm to obtain

$$\frac{1}{2} \sum_i u_i (\mu_i - \lambda_i)^2 \geq b_\delta. \quad (2.5)$$

(To justify (2.5) from (2.3), first use the event $\{T \leq M, \hat{i} = *\}$ and then let $M \rightarrow \infty$.) If $m < 1$, raise suboptimal arm i to $m + \varepsilon$ and send $\varepsilon \downarrow 0$ in (2.5) to get (2.4). If $m = 1$, instead raise arm i to 1 and lower the original optimal arm to $1 - \varepsilon$, with $0 < \varepsilon < g$, and take $\varepsilon \rightarrow 0$ in (2.5), using $u_* < \infty$.

For each nonempty gap group $r \in \mathcal{U}$, put

$$n_r = |G_r|, \quad t_r = \sum_{i \in G_r} u_i, \quad \alpha_r = \frac{t_r}{n_r 4^r}. \quad (2.6)$$

Since $\Delta_i^{-2} \geq 4^r$ in G_r , (2.4) implies

$$\alpha_r \geq 2b_\delta > 1. \quad (2.7)$$

2.2. A common reference law. For $r \in \mathcal{U}$, let P_r be the law of the algorithm's run under the following random relabeling of the original instance I , with all arm distributions left unchanged: the optimal arm has label 1; an identity i chosen uniformly from G_r has label 2; the other identities are placed uniformly on labels $3, \dots, K$. Then

$$P_r(\hat{i} = 1) \geq 1 - \delta, \quad \mathbb{E}_{P_r} N_2 = \alpha_r 4^r. \quad (2.8)$$

The expectation identity follows from permutation equivariance, since the algorithm is not told which instance identities occupy labels 1 and 2.

Under P_r , raising only the mean at label 2 to m would produce a tied law whose remaining suboptimal arms omit the chosen identity from G_r ; this law would therefore still depend on r . To obtain a common reference law, fix $s \in \mathcal{U}$ and define Q_s as follows. Give labels 1 and 2 mean m , choose an identity j uniformly from G_s , and assign all original suboptimal identities other than j uniformly to labels $3, \dots, K$. Thus Q_s has exactly two tied optimal arms and does not depend on r . Although this tied law lies outside the unique-optimum model class, its finite-time answer probabilities are controlled by perturbations with a unique optimum, as the next lemma shows. Let $P_r^{(M)}$ and $Q_s^{(M)}$ denote the corresponding laws of the recorded histories up to $T \wedge M$.

Lemma 2.2. *For every $s \in \mathcal{U}$, $Q_s(T < \infty, \hat{i} = 1) \leq \delta$.*

Proof. Since the optimal arm is unique and all means are nonnegative, $m > 0$. For $0 < \varepsilon < m$, let $Q_{s,\varepsilon}$ be obtained from Q_s by lowering the mean at label 1 from m to $m - \varepsilon$. Every component of $Q_{s,\varepsilon}$ has label 2 as its unique optimal arm. Therefore, for every fixed M ,

$$Q_{s,\varepsilon}(T \leq M, \hat{i} = 1) \leq \delta.$$

The KL divergence between the laws under $Q_{s,\varepsilon}$ and Q_s of the history up to $T \wedge M$ is at most $M\varepsilon^2/2$. Pinsker's inequality consequently gives

$$Q_s(T \leq M, \hat{i} = 1) \leq \delta + \frac{\varepsilon \sqrt{M}}{2}.$$

First let $\varepsilon \downarrow 0$ with M fixed, and then let $M \rightarrow \infty$ to obtain the claim. $\square$

Lemma 2.3. *For $r, s \in \mathcal{U}$ with $s \leq r$ and every finite horizon M ,*

$$\text{KL}(P_r^{(M)} \| Q_s^{(M)}) \leq \frac{\alpha_r + \alpha_s}{2}. \quad (2.9)$$

When $r = s$, the sharper bound $\alpha_s/2$ holds.

Proof. Suppose first that $s < r$. Start from the random labeling defining P_r : choose $i \in G_r$ uniformly, place it at label 2, and arrange the remaining identities uniformly on the other labels. Independently choose $j \in G_s$ uniformly. Modify two means: raise the mean at label 2 from μ_i to m , and change the mean at the label occupied by j from μ_j to μ_i . The first change makes labels 1 and 2 tied for the largest mean, while the second replaces the omitted identity i among labels $3, \dots, K$ by the omitted identity j .

We verify that the modified model has law Q_s . Conditional on i and j , the identities on labels $3, \dots, K$ are initially a uniform arrangement of all suboptimal identities except i . Replacing j by i gives a bijection from these arrangements to the arrangements of all suboptimal identities except j . The latter arrangement is therefore uniform and independent of i and r . Averaging over the uniform choice of $j \in G_s$ gives Q_s . This identity-level argument also applies when distinct arms have equal means.

Adjoin the choices of i, j and the random labeling to the histories under both models. Conditional on these variables, the two models differ only at label 2 and at the label occupied by j . Equation (2.3) therefore bounds their conditional KL divergence by

$$\frac{1}{2} \mathbb{E}_{P_r} \left[\Delta_i^2 N_2(T \wedge M) + (\Delta_j - \Delta_i)^2 N_{\text{label}(j)}(T \wedge M) \right].$$

Forgetting the auxiliary variables can only decrease KL divergence. Since $s < r$, the definitions of G_r and G_s give

$$0 < \Delta_i < \Delta_j, \quad \Delta_i^2 \leq 4^{-r}, \quad (\Delta_j - \Delta_i)^2 \leq 4^{-s}.$$

Moreover, equivariance and uniform choice within the two groups give expected total sample counts t_r/n_r for i and t_s/n_s for j . Hence

$$\text{KL}(P_r^{(M)} \| Q_s^{(M)}) \leq \frac{1}{2} \left(4^{-r} \frac{t_r}{n_r} + 4^{-s} \frac{t_s}{n_s} \right) = \frac{\alpha_r + \alpha_s}{2}.$$

When $r = s$, raise only the mean at label 2 from μ_i to m . The resulting marginal law is Q_s , and

$$\text{KL}(P_s^{(M)} \| Q_s^{(M)}) \leq \frac{1}{2} 4^{-s} \mathbb{E}_{P_s} N_2 = \frac{\alpha_s}{2}.$$

□

2.3. Sample counts and entropy. Recall that under P_r , label 2 contains an arm chosen uniformly from G_r , and

$$\mathbb{E}_{P_r} N_2 = \alpha_r 4^r.$$

Set $c_0 = 1/256$. For $a \geq 1$ and $r \in \mathcal{U}$ with $\alpha_r \leq a$, define

$$E_r(a) = \{T < \infty, \hat{i} = 1, c_0 4^r \leq N_2 \leq 4a 4^r\}. \quad (2.10)$$

Lemma 2.4. *If $\alpha_r \leq a$, then $P_r(E_r(a)) \geq 1/2$.*

Proof. Put $L = c_0 4^r$ and $n = \lceil L \rceil - 1$. Since N_2 is integer,

$$N_2 < L \iff N_2 \leq n.$$

Fix a total horizon M . Starting from the history under P_r , stop recording at the first of the following times: when the algorithm stops, when it has taken M samples in total, or immediately after it selects label 2 for the $(n+1)$ st time and before the corresponding reward is revealed. Record which

of these three stopping conditions occurred. The resulting history contains at most n rewards from label 2 and determines the event

$$A_M = \{T \leq M, \hat{i} = 1, N_2 \leq n\}.$$

Compare this censored history under P_r with the corresponding history under Q_r . Conditional on the selected identity $i \in G_r$ and the random labeling, the two models differ only in the mean at label 2, which changes from μ_i to m . Each observed reward from that label contributes Gaussian KL divergence $\Delta_i^2/2$, and at most n such rewards are observed. Since $\Delta_i^2 \leq 4^{-r}$,

$$\text{KL}(\bar{P}_{r,M,n} \| \bar{Q}_{r,M,n}) \leq \frac{1}{2} n 4^{-r} \leq \frac{c_0}{2},$$

where $\bar{P}_{r,M,n}$ and $\bar{Q}_{r,M,n}$ denote the laws of the censored histories. The same bound holds after averaging over the selected identity and labeling, since forgetting these auxiliary variables can only decrease KL divergence.

By Lemma 2.2,

$$Q_r(A_M) \leq Q_r(T < \infty, \hat{i} = 1) \leq \delta.$$

Pinsker's inequality therefore gives

$$P_r(A_M) \leq Q_r(A_M) + \sqrt{\frac{1}{2} \text{KL}(\bar{P}_{r,M,n} \| \bar{Q}_{r,M,n})} \leq \delta + \frac{\sqrt{c_0}}{2}.$$

Letting $M \rightarrow \infty$ and using $c_0 = 1/256$, we obtain

$$P_r(T < \infty, \hat{i} = 1, N_2 < c_0 4^r) \leq \delta + \frac{1}{32}.$$

For the upper tail, (2.8) and $\alpha_r \leq a$ give

$$\mathbb{E}_{P_r} N_2 = \alpha_r 4^r \leq a 4^r.$$

Hence Markov's inequality yields

$$P_r(N_2 > 4a 4^r) \leq \frac{1}{4}.$$

Finally, $P_r(T < \infty, \hat{i} = 1) \geq 1 - \delta$. Removing from this event the lower- and upper-tail events bounded above gives

$$P_r(E_r(a)) \geq 1 - \delta - \left(\delta + \frac{1}{32} \right) - \frac{1}{4} = 1 - 2\delta - \frac{1}{32} - \frac{1}{4} > \frac{1}{2},$$

because $\delta < 0.1$. □

Lemma 2.5. *For $a \geq 0$, let*

$$N(a) = |\{r \in \mathcal{U} : \alpha_r \leq a\}|.$$

Then

$$N(a) \leq 32ae^{2a} \quad (a \geq 1), \quad \sum_{r \in \mathcal{U}} e^{-8\alpha_r} \leq 1. \quad (2.11)$$

Proof. Fix $a \geq 1$. If $N(a) = 0$, the first bound is immediate. Otherwise let

$$s = \min\{r \in \mathcal{U} : \alpha_r \leq a\}.$$

Every index counted by $N(a)$ satisfies $r \geq s$. Since both α_r and α_s are at most a , Lemma 2.3 gives

$$\text{KL}(P_r^{(M)} \| Q_s^{(M)}) \leq a$$

for every finite horizon M .

We use this inequality to transfer the event $E_r(a)$ from P_r to the common law Q_s . Set

$$B_{r,M} = E_r(a) \cap \{T \leq M\}, \quad p_M = P_r(B_{r,M}), \quad q_M = Q_s(B_{r,M}).$$

The event $B_{r,M}$ is determined by the history up to $T \wedge M$, so data processing and the binary-divergence bound

$$\text{kl}(p, q) \geq p \log(1/q) - \log 2$$

give

$$a \geq \text{kl}(p_M, q_M) \geq p_M \log(1/q_M) - \log 2.$$

As $M \rightarrow \infty$, p_M increases to $P_r(E_r(a))$, which satisfies

$$P_r(E_r(a)) \geq \frac{1}{2}$$

by [Lemma 2.4](#), while q_M increases to $Q_s(E_r(a))$. Therefore

$$\log \frac{1}{Q_s(E_r(a))} \leq 2a + 2 \log 2,$$

and hence

$$Q_s(E_r(a)) \geq \frac{1}{4} e^{-2a}. \quad (2.12)$$

It remains to use the fact that sufficiently separated indices impose disjoint conditions on the same random variable N_2 . Let

$$d(a) = \lceil \log_4(1024a) \rceil + 1.$$

If $r' > r$ and $r' \equiv r \pmod{d(a)}$, then $r' - r \geq d(a)$ and

$$c_0 4^{r'} \geq c_0 4^{r+d(a)} > 4a 4^r,$$

because $c_0 = 1/256$ and $4^{d(a)} > 1024a$. Thus the intervals

$$[c_0 4^r, 4a 4^r]$$

are disjoint within each residue class modulo $d(a)$. The corresponding events $E_r(a)$ are therefore disjoint under the common law Q_s . Since each has probability at least $\frac{1}{4} e^{-2a}$, each residue class contains at most $4e^{2a}$ indices counted by $N(a)$. Consequently,

$$N(a) \leq 4d(a)e^{2a}.$$

Finally,

$$d(a) \leq \log_4 a + 7 \leq 8a \quad (a \geq 1),$$

which proves

$$N(a) \leq 32ae^{2a}.$$

By [\(2.7\)](#), every α_r is greater than 1, so $N(a) = 0$ for $a \leq 1$. Using

$$e^{-8\alpha_r} = \int_{\alpha_r}^{\infty} 8e^{-8a} da$$

and summing over r gives

$$\sum_{r \in \mathcal{U}} e^{-8\alpha_r} = \int_1^{\infty} 8e^{-8a} N(a) da \leq 256 \int_1^{\infty} ae^{-6a} da = \frac{448}{9} e^{-6} < 1.$$

□

Proof of [Proposition 2.1](#). Recall that T denotes the stopping time of the symmetrized algorithm. Set

$$z = \sum_{r \in \mathcal{U}} e^{-8\alpha_r}.$$

By [Lemma 2.5](#), $0 < z \leq 1$, so

$$q_r = \frac{e^{-8\alpha_r}}{z}, \quad r \in \mathcal{U},$$

defines a probability vector. Gibbs's inequality gives

$$0 \leq \sum_{r \in \mathcal{U}} p_r \log \frac{p_r}{q_r} = -\text{Ent}(I) + 8 \sum_{r \in \mathcal{U}} p_r \alpha_r + \log z.$$

Since $\log z \leq 0$, it follows that

$$\text{Ent}(I) \leq 8 \sum_{r \in \mathcal{U}} p_r \alpha_r. \quad (2.13)$$

We now convert the right side into an expected sample count. Using $p_r = H_r/H$ and $\alpha_r = t_r/(n_r 4^r)$ gives

$$H \sum_{r \in \mathcal{U}} p_r \alpha_r = \sum_{r \in \mathcal{U}} \frac{H_r}{n_r 4^r} t_r.$$

For every $i \in G_r$, the inequality $\Delta_i > 2^{-(r+1)}$ implies $\Delta_i^{-2} < 4^{r+1}$. Hence

$$H_r = \sum_{i \in G_r} \Delta_i^{-2} \leq 4 n_r 4^r,$$

and therefore

$$H \sum_{r \in \mathcal{U}} p_r \alpha_r \leq 4 \sum_{r \in \mathcal{U}} t_r = 4 \sum_{i \neq *} u_i \leq 4 \mathbb{E}_I T.$$

Combining this estimate with (2.13) yields

$$\mathbb{E}_I T \geq \frac{H \text{Ent}(I)}{32}.$$

The per-arm change-of-measure bound (2.4) also gives

$$\mathbb{E}_I T \geq 2b_\delta H.$$

Finally, (2.2) identifies $\mathbb{E}_I T$ with the average expected cost of the original algorithm over all relabelings of I . Since that algorithm was arbitrary, taking the infimum in (1.2) proves (2.1). $\square$

3. THE INSTANCE-WISE UPPER BOUND

The following sub-Gaussian proposition implies the upper bound in Theorem 1.1 when restricted to $I \in \mathcal{G}_K$.

Proposition 3.1. *For every $I \in \mathcal{S}_K$ and $0 < \delta < 0.1$, there exists an algorithm A that is δ -correct on $\mathcal{S}_K$ and satisfies*

$$\mathbb{E}_{\pi_I} T_A \leq CH(\log(1/\delta) + \text{Ent}(I)) \quad (\pi \in S_K).$$

The constant C is absolute.

The remainder of this section proves Proposition 3.1. We first describe the target-instance-dependent parameters and a single trial, then prove its correctness and expected-cost bounds, and finally repeat the trials and add an auxiliary procedure to ensure almost-sure stopping on every instance in $\mathcal{S}_K$.

It suffices to prove the upper bounds for $0 < \delta < 0.01$. For $0.01 \leq \delta < 0.1$, run the corresponding algorithm with error parameter 0.005. It remains δ -correct, while

$$\log 200 \leq \frac{\log 200}{\log 10} \log(1/\delta),$$

so the sample bound changes only by an absolute constant factor.

3.1. Instance-specific parameters and a trial. Fix a target instance $I \in \mathcal{S}_K$ and assume $0 < \delta < 0.01$. The algorithm constructed in this section may depend on the target's unlabeled collection of means, but not on the label of its optimal arm. For $r \in \mathcal{U}$, set

$$n_r = |G_r|, \quad w_r = n_r 4^r, \quad W = \sum_{r \in \mathcal{U}} w_r, \quad q_r = \frac{w_r}{W},$$

and let

$$R = \max \mathcal{U}, \quad h = 2^{-R}.$$

For $i \in G_r$,

$$4^r \leq \Delta_i^{-2} < 4^{r+1},$$

so summing over G_r gives $w_r \leq H_r < 4w_r$. Consequently,

$$w_r \leq H_r < 4w_r, \quad W \leq H < 4W, \quad g \leq h < 2g, \quad h^{-2} \leq W. \quad (3.1)$$

Here the bounds involving h follow because an arm with gap g belongs to G_R , while G_R is nonempty and therefore $W \geq w_R \geq 4^R = h^{-2}$.

We will allocate the mixture weights according to $q = (q_r)$. To compare the entropy of these weights with $\text{Ent}(I)$, define, for a finite nonnegative vector v ,

$$\Phi(v) = \left(\sum_r v_r \right) \log \left(\sum_r v_r \right) - \sum_r v_r \log v_r, \quad 0 \log 0 = 0. \quad (3.2)$$

If $V = \sum_r v_r > 0$, then

$$\Phi(v) = V \text{Ent}(v/V).$$

Moreover, Φ is homogeneous of degree one and coordinatewise nondecreasing: at a positive coordinate,

$$\frac{\partial \Phi}{\partial v_r} = \log \frac{\sum_s v_s}{v_r} \geq 0,$$

and the same monotonicity at zero follows by continuity. Applying these properties to the coordinatewise inequalities $w_r \leq H_r \leq 4w_r$ gives

$$W \text{Ent}(q) = \Phi(w) \leq \Phi((H_r)_r) = H \text{Ent}(I) \leq \Phi(4w) = 4W \text{Ent}(q). \quad (3.3)$$

Choose a rational threshold

$$\tau \in (m - g/2, m - g/3) \subset (0, 1). \quad (3.4)$$

On the target instance, the optimal mean m lies above τ , while every suboptimal mean is at most $m - g < \tau$. The algorithm stores the finite set $\mathcal{U}$, the integers $(n_r)_{r \in \mathcal{U}}$, and the rational number τ . These parameters have a finite description and do not identify the optimal arm's label.

Fix an error allowance $0 < \eta < 0.01$. A trial attempts to eliminate arms whose means are below τ while retaining any arm whose mean is at least τ . For each arm i , after observing t samples $X_{i,1}, \dots, X_{i,t}$, define

$$M_i(t) = \sum_{r \in \mathcal{U}} q_r \exp \left(-\lambda_r \sum_{\ell=1}^t (X_{i,\ell} - \tau) - \frac{\lambda_r^2 t}{2} \right), \quad \lambda_r = 2^{-r-2}. \quad (3.5)$$

The component indexed by r is tuned to departures below τ on the scale 2^{-r} . Under the hypothesis $\mu_i \geq \tau$, each component, and hence their mixture, is a nonnegative supermartingale.

Initialize all arms as active. Repeatedly perform a complete sweep, drawing one new sample from each arm active at the start of the sweep. At the end of the sweep, simultaneously discard every arm i whose updated statistic satisfies $M_i(t) \geq 1/\eta$. If no arm remains, end the trial without an answer. If exactly one arm c remains, take

$$n_\eta = \left\lceil 288h^{-2} \log(1/\eta) \right\rceil$$

fresh samples from it. Output c if their average exceeds $\tau + h/12$; otherwise end the trial without an answer. If every input mean lies below τ , this confirmation step bounds the probability that the trial returns an arm by η .

Finally, impose the deterministic sample limit

$$B_\eta = \lceil 16000W[\log(1/\eta) + \text{Ent}(q) + 1] \rceil. \quad (3.6)$$

The limit includes both elimination and confirmation samples. If the next sample would make the total exceed B_η , end the trial without an answer.

3.2. Correctness and cost of a trial. We first show that the trial has error probability at most η on every input instance, even though its parameters were chosen from the fixed target instance. Let an input arm have mean $\mu \geq \tau$. For each $r \in \mathcal{U}$ and a fresh sample X from this arm, the sub-Gaussian assumption gives

$$\mathbb{E}e^{-\lambda_r(X-\tau)-\lambda_r^2/2} = e^{-\lambda_r(\mu-\tau)}\mathbb{E}e^{-\lambda_r(X-\mu)-\lambda_r^2/2} \leq e^{-\lambda_r(\mu-\tau)} \leq 1.$$

The same inequality holds conditionally on all preceding samples. Hence each exponential process in (3.5) is a nonnegative supermartingale. Their weighted sum M_i is therefore also a nonnegative supermartingale, with $M_i(0) = \sum_r q_r = 1$. The maximal inequality for nonnegative supermartingales gives

$$\mathbb{P}\left(\sup_{t \geq 0} M_i(t) \geq \frac{1}{\eta}\right) \leq \eta;$$

see [11, theorem 3.9]. Thus an arm with mean at least τ is eliminated with probability at most η .

Now consider an arbitrary input instance. If its optimal mean is at least τ , the trial can return a suboptimal arm only if it first eliminates the optimal arm. The preceding bound shows that this event has probability at most η . If every arm has mean below τ , condition on the history up to the confirmation step and on the remaining arm c . The confirmation samples are fresh, and $\mu_c < \tau$, so (B.1) gives

$$\mathbb{P}\left(\bar{X}_c > \tau + \frac{h}{12} \mid \text{preceding history}\right) \leq \exp\left(-\frac{n_\eta h^2}{288}\right) \leq \eta.$$

The deterministic sample limit can only end the trial without an answer. Therefore, on every instance in $\mathcal{S}_K$, the probability that the trial returns an incorrect arm is at most η .

We now return to the target instance I and bound the expected cost. Temporarily remove the deterministic sample limit, retaining all other parts of the trial. For each arm, generate an infinite independent sample sequence and define

$$U_i = \inf \left\{ t \geq 1 : M_i(t) \geq \frac{1}{\eta} \right\},$$

with $U_i = \infty$ if the threshold is never reached. This definition allows us to bound the cost even on paths where the optimal arm is erroneously eliminated.

Fix a suboptimal target arm $i \in G_r$ and put

$$d_i = \tau - \mu_i.$$

Since $m - \tau < g/2 \leq \Delta_i/2$,

$$d_i = \Delta_i - (m - \tau) > \frac{\Delta_i}{2}.$$

The definition of G_r and the choice $\lambda_r = 2^{-r-2}$ also give

$$\frac{\Delta_i}{4} \leq \lambda_r < \frac{\Delta_i}{2} < d_i.$$

The logarithm of the unweighted r th exponential component after t samples is

$$a_i t - \lambda_r \sum_{\ell=1}^t (X_{i,\ell} - \mu_i), \quad a_i = \lambda_r d_i - \frac{\lambda_r^2}{2}.$$

The preceding inequalities imply

$$a_i = \lambda_r \left(d_i - \frac{\lambda_r}{2} \right) \geq \frac{\Delta_i^2}{16}, \quad \frac{a_i}{\lambda_r} = d_i - \frac{\lambda_r}{2} \geq \frac{\Delta_i}{4}.$$

Set

$$b_r = \log \frac{1}{\eta q_r}.$$

If $U_i > t$, then $M_i(t) < 1/\eta$, so its r th component satisfies

$$\exp \left(a_i t - \lambda_r \sum_{\ell=1}^t (X_{i,\ell} - \mu_i) \right) < \frac{1}{\eta q_r}.$$

Consequently, for every integer $t \geq 2b_r/a_i$,

$$\begin{aligned} \mathbb{P}(U_i > t) &\leq \mathbb{P} \left(\lambda_r \sum_{\ell=1}^t (X_{i,\ell} - \mu_i) > a_i t - b_r \right) \\ &\leq \exp \left(-\frac{(a_i t - b_r)^2}{2\lambda_r^2 t} \right) \\ &\leq \exp \left(-\frac{a_i^2 t}{8\lambda_r^2} \right) \leq \exp \left(-\frac{\Delta_i^2 t}{128} \right). \end{aligned}$$

Here the penultimate inequality uses $a_i t - b_r \geq a_i t/2$, and the last uses $a_i/\lambda_r \geq \Delta_i/4$.

Let $t_0 = \lceil 2b_r/a_i \rceil$. Summing the tail probabilities gives

$$\mathbb{E}U_i \leq t_0 + \sum_{t=t_0}^{\infty} e^{-\Delta_i^2 t/128} \leq 130\Delta_i^{-2}(b_r + 1).$$

Indeed, $a_i \geq \Delta_i^2/16$ gives $t_0 \leq 32b_r\Delta_i^{-2} + 1$, while $\Delta_i \leq 1$ gives

$$\sum_{t=t_0}^{\infty} e^{-\Delta_i^2 t/128} \leq \frac{1}{1 - e^{-\Delta_i^2/128}} \leq 129\Delta_i^{-2}.$$

Summing over the suboptimal arms and using $b_r = \log(1/\eta) + \log(1/q_r)$ yields

$$\begin{aligned} \sum_{i \neq *} \mathbb{E}U_i &\leq 130 \sum_{r \in \mathcal{U}} H_r [\log(1/\eta) + \log(1/q_r) + 1] \\ &\leq 520 \sum_{r \in \mathcal{U}} w_r [\log(1/\eta) + \log(1/q_r) + 1] \\ &= 520W [\log(1/\eta) + \text{Ent}(q) + 1]. \end{aligned}$$

The second inequality uses $H_r \leq 4w_r$, while the last equality uses $q_r = w_r/W$.

It remains to account for samples from the optimal arm and for the confirmation step. During the elimination phase, suboptimal arm i receives at most U_i samples. If the optimal arm receives t elimination samples, then in its final sweep at least one suboptimal arm is also active and receives its t th sample. Hence $U_i \geq t$ for that arm, so the optimal arm receives at most $\max_{i \neq *} U_i$ samples before the elimination phase ends. This remains true if the optimal arm is itself eliminated. Hence the elimination cost is bounded pathwise by

$$\sum_{i \neq *} U_i + \max_{i \neq *} U_i \leq 2 \sum_{i \neq *} U_i.$$

The confirmation step uses at most n_η further samples. Since $h^{-2} \leq W$, the expected cost of the uncapped trial is therefore at most

$$2000W [\log(1/\eta) + \text{Ent}(q) + 1].$$

Comparing this with (3.6), Markov's inequality shows that the probability of reaching the deterministic sample limit is at most $1/8$.

Finally, on the target instance,

$$m - \tau > \frac{g}{3} > \frac{h}{6}.$$

Thus the confirmation threshold $\tau + h/12$ lies at least $h/12$ below m , and (B.1) gives

$$\mathbb{P}\left(\bar{X}_* \leq \tau + \frac{h}{12}\right) \leq e^{-n_\eta h^2/288} \leq \eta.$$

The bounds on $\mathbb{E}U_i$ imply that $U_i < \infty$ almost surely for every suboptimal arm i . If the optimal arm is never eliminated, the uncapped elimination phase therefore ends with that arm as the sole survivor. Therefore, unless the optimal arm is eliminated, the sample limit is reached, or the confirmation fails, the trial returns the optimal arm. Its success probability is at least

$$1 - \eta - \frac{1}{8} - \eta = 1 - \frac{1}{8} - 2\eta > \frac{3}{4}.$$

All estimates are uniform over relabelings of the target instance.

3.3. Repetition and almost-sure stopping. Run independent trials with error levels

$$\eta_j = \delta 2^{-j-2}, \quad j \geq 0,$$

and stop at the first trial that returns an arm. On every input instance, the probability that some trial returns an incorrect arm is at most

$$\sum_{j \geq 0} \eta_j = \frac{\delta}{2}.$$

For any relabeling of the target instance, each trial returns the optimal arm with probability at least $3/4$. Independence therefore implies that the probability of reaching trial j is at most 4^{-j} . Moreover,

$$B_{\eta_j} \leq CW [\log(1/\delta) + \text{Ent}(q) + j + 1].$$

Consequently, the expected number of samples used by the sequence of trials is at most

$$\sum_{j \geq 0} 4^{-j} B_{\eta_j} \leq CW [\log(1/\delta) + \text{Ent}(q) + 1].$$

The sequence of trials need not stop almost surely when its fixed parameters do not describe the input instance. To ensure almost-sure stopping on every instance, run it in parallel with an independent auxiliary procedure that is $\delta/2$ -correct on $\mathcal{S}_K$. Alternate between the next sample requested by the trial sequence and the next sample requested by the auxiliary procedure, and return the first arm produced by either procedure. Before the trial sequence returns an arm, the combined algorithm uses at most twice as many samples, up to one additional sample. Thus its expected cost on any relabeling of the target instance remains bounded by

$$CW [\log(1/\delta) + \text{Ent}(q) + 1].$$

We construct the auxiliary procedure using simultaneous confidence intervals. Put $\beta = \delta/2$. At stage $k \geq 0$, draw 2^k fresh samples from each arm and let $\bar{X}_{i,k}$ be their average. Form the intervals

$$[\bar{X}_{i,k} - c_k, \bar{X}_{i,k} + c_k], \quad c_k = \sqrt{2^{1-k} \log \frac{8K(k+1)^2}{\beta}}.$$

Stop and return arm i if its lower endpoint exceeds the upper endpoint of every other arm. For each arm and stage, the sub-Gaussian tail bound gives

$$\mathbb{P}(|\bar{X}_{i,k} - \mu_i| > c_k) \leq \frac{\beta}{4K(k+1)^2}.$$

A union bound over all arms and stages shows that the probability of any confidence-interval failure is at most

$$\frac{\beta}{4} \sum_{k \geq 0} \frac{1}{(k+1)^2} = \frac{\beta\pi^2}{24} < \beta.$$

Whenever all intervals contain their respective means, any arm returned by the procedure is optimal. The auxiliary procedure is therefore β -correct.

It remains to verify almost-sure stopping. For every fixed $e > 0$,

$$\mathbb{P}(|\bar{X}_{i,k} - \mu_i| > e) \leq 2 \exp(-2^{k-1}e^2),$$

and the right side is summable in k . Applying Borel–Cantelli for each arm and for $e = 1, \frac{1}{2}, \frac{1}{3}, \dots$ gives

$$\bar{X}_{i,k} \longrightarrow \mu_i \quad \text{almost surely for every arm } i.$$

Since $c_k \rightarrow 0$ and the optimal arm is unique, its lower endpoint eventually exceeds every suboptimal arm's upper endpoint. Hence the auxiliary procedure, and therefore the combined algorithm, stops almost surely on every instance in $\mathcal{S}_K$.

The two component procedures have error probabilities at most $\delta/2$, so the combined algorithm is δ -correct. Finally, (3.1) and (3.3) give

$$W \leq H, \quad W \text{Ent}(q) \leq H \text{Ent}(I).$$

Since $\log(1/\delta) > \log 100$, the additive W term can be absorbed into $H \log(1/\delta)$. Thus, uniformly over all relabelings of the target instance,

$$\mathbb{E}_I T \leq CH [\log(1/\delta) + \text{Ent}(I)],$$

which proves [Proposition 3.1](#).

4. THE UNIFORM ALGORITHM

We construct the algorithm in [Theorem 1.2](#) for $\mathcal{S}_K$ by modifying the Entropy-Elimination and Complexity-Guessing procedures of Chen, Li and Qiao [[3](#), algorithms 4–5]. We retain their geometrically increasing stage budgets, two cumulative budget tests, and elimination thresholds. We adjust the maximum number of rounds and fraction-threshold increments to account for all sampling costs, and impose a deterministic sample limit at each stage. The analysis uses the comparison process in [Section 5](#). Each stage starts with all K arms. One cumulative sum limits the total elimination error budget, and the other limits its expected sampling cost.

4.1. Sampling subroutines. By the reduction in [Section 3](#), it suffices to consider $0 < \delta < 0.01$. Each subroutine call uses samples independent of the history before the call. Consequently, the guarantees below hold conditionally on that history and on the possibly random inputs to the call. They apply to 1-sub-Gaussian reward laws and use only the sample-mean bound ([B.1](#)) and independence between sample blocks.

We refer to Mean and Fraction calls made directly by the main algorithm as *outer tests*, to distinguish them from calls made internally by other subroutines.

Median and Fraction require nonempty finite input sets, whereas Eliminate also accepts the empty set. Throughout, accuracy parameters and differences between mean thresholds lie in $(0, 1]$, error levels lie in $(0, 0.1)$, and differences between fraction thresholds lie in $(0, 0.1]$. These ranges cover every call made by the algorithm. The implementations and proofs are given in [Appendix B](#). Here and below, C denotes an absolute constant.

- (i) *Mean estimation.* The call $\text{Mean}(i, e, \alpha)$ uses $O(e^{-2} \log(1/\alpha))$ samples deterministically and returns an estimate of μ_i whose error is at most e , except with probability α .
- (ii) *Median elimination.* The call $\text{Median}(S, e)$ uses $O(|S|e^{-2})$ samples deterministically and, with conditional probability at least 0.99, returns an arm $i \in S$ satisfying

$$\mu_i \geq \max_{j \in S} \mu_j - e.$$

- (iii) *Fraction testing.* For $l < u$ and $\theta_- < \theta_+$, the call $\text{Fraction}(S, l, u, \theta_-, \theta_+, \alpha)$ has deterministic cost at most

$$C(u-l)^{-2} \log(1/\alpha) (\theta_+ - \theta_-)^{-2} \log \frac{1}{\theta_+ - \theta_-}.$$

Except with probability α , its output obeys the following implications:

$$\text{True} \implies |\{i \in S : \mu_i < u\}| > \theta_- |S|,$$

$$\text{False} \implies |\{i \in S : \mu_i < l\}| < \theta_+ |S|.$$

Fraction is not required to estimate either proportion. It need only return an answer whose corresponding implication is valid; on some inputs, both answers are valid.

- (iv) *Elimination.* The call $\text{Eliminate}(S, l, u, \alpha)$ returns a subset $S' \subseteq S$ and terminates almost surely. If $S = \emptyset$, it returns immediately. Its conditional expected cost is at most

$$C|S|(u-l)^{-2} \log(1/\alpha).$$

If $S \neq \emptyset$, fix, before the call, a designated arm $i^\dagger \in S$ attaining $\max_{i \in S} \mu_i$. Except with probability α , the returned set simultaneously satisfies

$$|\{i \in S' : \mu_i < l\}| \leq 0.1|S'|,$$

and, if $\mu_{i^\dagger} \geq u$, it contains $i^\dagger$. When $S = \emptyset$, the displayed inequality holds deterministically. In addition, for any k arms designated before the call whose means are all at least u , the probability that all k are removed is at most α^k .

We use a product bound to control the joint loss of several good arms across stages. Conditional on the inputs and the history before an Eliminate call, generate independent estimation blocks in advance for every arm and every internal stage. If an arm has mean at least u , then it can be removed only if one of its own estimates is inaccurate. The error budgets for that arm sum to at most $\alpha/10$, so this event has probability at most $\alpha/10$. Because the estimation blocks are independent across arms, the probability that k fixed arms of mean at least u are all removed is at most

$$\left(\frac{\alpha}{10}\right)^k \leq \alpha^k.$$

This conclusion does not require the Fraction calls to be correct: an incorrect Fraction answer may cause an unnecessary deletion step, but no such step removes an arm of mean at least u when that arm's estimate is accurate. This is the argument of [3, lemma B.9]. The full construction, including its behavior on the empty set, is given in [Appendix B.4](#).

4.2. Stages and stopping rules. Let J and M be the absolute constants specified in [Lemmas 6.1](#) and [6.2](#), respectively. After fixing them, choose a sufficiently small absolute constant $c > 0$ and set

$$\eta = c\delta, \quad \rho = \eta^2, \quad a = \log(1/\eta), \quad B_t = 100^t, \quad L_t = a + \log(t+1), \quad t = 1, 2, \dots$$

Each stage starts with all K arms and uses samples and internal randomness independent of every other stage. We say that a stage *accepts* if it returns an arm and *rejects* if it ends without doing so.

For stage t , set

$$R_t = \max\{0, \lfloor \log_4(B_t/L_t) \rfloor - J\}.$$

If $R_t = 0$, reject immediately. Otherwise, write $B = B_t$ and $R = R_t$, and initialize

$$S_1 = [K], \quad U_1 = V_1 = 0.$$

The variables U_r and V_r are running totals used by the two budget checks. At the start of round r ,

$$U_r = 4 \sum_{\substack{k < r \\ \text{Eliminate was called in round } k}} w_k, \quad V_r = \sum_{\substack{k < r \\ \text{round } k \text{ passed both budget checks}}} z_k.$$

They do not record the number of samples used, since that quantity is controlled by the separate deterministic limit involving M .

Define the fraction thresholds by

$$\theta_0 = 0.3, \quad \theta_r = \theta_{r-1} + \frac{1}{10(R+1-r)^2}, \quad 1 \leq r \leq R.$$

They are increasing and satisfy

$$\theta_R = 0.3 + \frac{1}{10} \sum_{k=1}^R \frac{1}{k^2} < 0.3 + \frac{\pi^2}{60} < 0.5.$$

Thus $\theta_r \in [0.3, 0.5)$, and $v < r$ implies $\theta_v \leq \theta_{r-1}$.

At the beginning of round r , return the unique arm in S_r if $|S_r| = 1$. Otherwise, reject if $S_r = \emptyset$ or $r > R$. If the stage continues, set

$$\varepsilon_r = 2^{-r}, \quad w_r = |S_r|4^r.$$

The first budget check rejects if

$$U_r + 4w_r \geq B.$$

If it passes, then $w_r < B/4$, and we may define the positive quantity

$$z_r = w_r \log \frac{B}{\eta w_r}.$$

The second budget check rejects if

$$V_r + z_r \geq 100B.$$

If both checks pass, perform the following operations:

(1) Update the second budget by setting

$$V_{r+1} = V_r + z_r, \quad \alpha_{t,r} = \frac{\eta}{50t^2r^2}.$$

(2) Call

$$\hat{i}_r = \text{Median}(S_r, \varepsilon_r/8), \quad \hat{\mu}_r = \text{Mean}(\hat{i}_r, \varepsilon_r/8, \alpha_{t,r}).$$

(3) Let F_r be the output of

$$\text{Fraction} \left(S_r, \hat{\mu}_r - \frac{7}{4}\varepsilon_r, \hat{\mu}_r - \frac{9}{8}\varepsilon_r, \theta_{r-1}, \theta_r, \alpha_{t,r} \right).$$

(4) If $F_r = \text{True}$, set

$$\beta_r = \frac{4w_r\rho}{B}$$

and call

$$S_{r+1} = \text{Eliminate} \left(S_r, \hat{\mu}_r - \frac{3}{4}\varepsilon_r, \hat{\mu}_r - \frac{5}{8}\varepsilon_r, \beta_r \right).$$

Then set $U_{r+1} = U_r + 4w_r$. If $F_r = \text{False}$, make no elimination call and set

$$S_{r+1} = S_r, \quad U_{r+1} = U_r.$$

Maintain a counter for the samples used within the stage. Before taking any sample, including one requested by a subroutine, reject the stage if that sample would make the counter exceed $\lceil MB \rceil$. After a rejection, begin stage $t + 1$ with fresh samples and randomness. If the stage returns an arm, terminate the algorithm and output that arm.

The singleton check precedes the condition $r > R$. Consequently, if round R produces a singleton S_{R+1} , the algorithm returns that arm at the beginning of round $R + 1$; otherwise, it rejects at that point.

5. PROGRESS AND ERROR CONTROL

Fix an instance $I \in \mathcal{S}_K$, with optimal mean m . We analyze the algorithm through a comparison process in which the outer Mean and Fraction outputs are made valid by construction. Median and Eliminate are not corrected. Successful Median calls will control the sizes of the active sets, while invalid Eliminate calls will be handled separately. In particular, we never condition on all future calls being valid.

5.1. A comparison process. For an outer Mean call on arm i with accuracy e , replace its output by its projection onto

$$[\mu_i - e, \mu_i + e].$$

Thus the corrected estimate always satisfies the advertised accuracy guarantee and agrees with the original output whenever that output is accurate.

For an outer Fraction call with parameters $S, l, u, \theta_-, \theta_+$, call True valid if

$$|\{i \in S : \mu_i < u\}| > \theta_- |S|$$

and False valid if

$$|\{i \in S : \mu_i < l\}| < \theta_+ |S|.$$

At least one answer is valid. Indeed, if True is invalid, then

$$|\{i \in S : \mu_i < l\}| \leq |\{i \in S : \mu_i < u\}| \leq \theta_- |S| < \theta_+ |S|,$$

so False is valid. Replace an invalid Fraction output by the other answer, leaving a valid output unchanged.

These corrections are measurable functions of the current inputs, outputs, and true means, and do not use future samples. They are introduced only for the analysis. They do not alter the samples already drawn or the cost of the corrected call, and the outputs of Median and Eliminate are never corrected.

Couple the actual and comparison processes using the same randomness until their first disagreement. Such a disagreement can occur only when an outer Mean or Fraction output is invalid. There are at most two such calls in each round, each with conditional failure probability $\alpha_{t,r} = \eta/(50t^2r^2)$. Hence

$$\mathbb{P}(\text{the two processes ever disagree}) \leq 2 \sum_{t,r \geq 1} \frac{\eta}{50t^2r^2} \leq C\eta. \quad (5.1)$$

For a fixed stage t , the same argument gives a bound of $C\eta/t^2$. We will not condition on agreement of the two processes. Because every Median call uses fresh samples, its conditional failure probability in the comparison process remains at most

$$\kappa = 0.01.$$

For a call Eliminate(S, l, u, α), designate before the call one arm attaining $\max_{i \in S} \mu_i$. Call the returned set S' *valid* if

$$|\{i \in S' : \mu_i < l\}| \leq 0.1|S'|$$

and, whenever the designated arm has mean at least u , that arm belongs to S' .

We first record why the true optimal arm remains active before the first invalid Eliminate call. The corrected estimate in round r satisfies

$$\hat{\mu}_r \leq \mu_{i_r} + \frac{\varepsilon_r}{8} \leq m + \frac{\varepsilon_r}{8}.$$

Consequently, the upper thresholds used by Fraction and Eliminate satisfy

$$\hat{\mu}_r - \frac{9}{8}\varepsilon_r \leq m - \varepsilon_r, \quad \hat{\mu}_r - \frac{5}{8}\varepsilon_r \leq m - \frac{1}{2}\varepsilon_r.$$

Thus, whenever Eliminate is called while the optimal arm is active, its upper threshold is below m . A valid call therefore retains the optimal arm. Induction over the rounds proves the claim.

Call the Median output in round r *successful* if

$$\mu_{i_r} \geq \max_{i \in S_r} \mu_i - \frac{\varepsilon_r}{8}.$$

Before the first invalid Eliminate call, the optimal arm lies in S_r , so the maximum on the right is m . If Median is successful, then the corrected Mean output satisfies

$$\hat{\mu}_r \geq m - \frac{\varepsilon_r}{4}.$$

It follows that the lower thresholds used by Fraction and Eliminate are bounded below by

$$\hat{\mu}_r - \frac{7}{4}\varepsilon_r \geq m - 2\varepsilon_r, \quad \hat{\mu}_r - \frac{3}{4}\varepsilon_r \geq m - \varepsilon_r.$$

Suppose that Median is successful and that every Eliminate call through the current round is valid. If Fraction returns False, then

$$|\{i \in S_r : \mu_i < \hat{\mu}_r - \frac{7}{4}\varepsilon_r\}| < \theta_r |S_r| < \frac{|S_r|}{2}.$$

More than half of S_r therefore consists of the optimal arm and suboptimal arms with $\Delta_i \leq 2\varepsilon_r$. Since a False answer leaves the active set unchanged,

$$|S_{r+1}| \leq 2(1 + |\{i \neq * : \Delta_i \leq 2\varepsilon_r\}|).$$

If Fraction returns True, validity of Eliminate implies that at least $0.9|S_{r+1}|$ returned arms have mean at least $m - \varepsilon_r$. All such arms are included among the optimal arm and the suboptimal arms with $\Delta_i \leq \varepsilon_r$. Hence the same, slightly weaker bound holds:

$$|S_{r+1}| \leq 2(1 + |\{i \neq * : \Delta_i \leq 2\varepsilon_r\}|). \quad (5.2)$$

There is a sharper conclusion once the round scale is below the smallest gap. Let

$$Q = \lfloor \log_4 D \rfloor, \quad D = g^{-2}.$$

Then

$$2^{-(Q+1)} < g \leq 2^{-Q}.$$

For $r \geq Q + 2$,

$$2\varepsilon_r = 2^{1-r} \leq 2^{-(Q+1)} < g.$$

Thus every suboptimal arm lies below both lower thresholds displayed above. If $|S_r| \geq 2$, at least half of S_r then lies below the Fraction lower threshold, so False is not a valid answer and the comparison process must return True. A valid Eliminate call retains the optimal arm. If n suboptimal arms remain afterward, its validity also gives

$$n \leq 0.1(n + 1),$$

which forces $n = 0$. Therefore every successful Median round $r \geq Q + 2$, before any invalid Eliminate call, leaves S_{r+1} equal to the singleton containing the optimal arm.

5.2. Active-set bounds. We now use (5.2) to control the two cumulative quantities appearing in the budget tests: the linear sum of the round weights and its logarithmically weighted analogue. The next lemma is a stopped-process version of [3, lemmas B.4–B.5].

Recall Φ from (3.2). For a summable nonnegative sequence $v = (v_r)$ with $V = \sum_r v_r$, we use the extension

$$\Phi(v) = \sum_{r:v_r>0} v_r \log \frac{V}{v_r},$$

with $\Phi(0) = 0$.

We first remove the deterministic sample limit from the stage. The resulting bounds continue to hold when the limit is restored, because the limit can only terminate the stage earlier. A round is *entered* when the first budget test is reached and *executed* when both budget tests are passed. We adopt the convention

$$c_r \log \left(e + \frac{B}{\eta c_r} \right) = 0 \quad \text{when } c_r = 0.$$

Lemma 5.1. *Consider one stage of the comparison process without its deterministic sample limit. Stop the analysis at the first invalid Eliminate call, at the first budget rejection, or after the last permitted round, whichever occurs first. The round containing an invalid Eliminate call and a round rejected by a budget test are both included.*

For every entered round with at least two active arms, set

$$c_r = 4^r |S_r|,$$

and set $c_r = 0$ for all subsequent rounds. Thus $c_r = w_r$ on executed rounds; for a round rejected immediately by a budget test, c_r records the corresponding round weight. There is a deterministic nonnegative sequence $b = (b_r)_{r \geq 1}$ such that

$$\mathbb{E} c_r \leq b_r, \quad \sum_r b_r \leq CH, \quad \Phi(b) \leq CH(\text{Ent}(I) + 1).$$

Consequently, whenever $B \geq H$,

$$\mathbb{E} \sum_r c_r \leq CH, \quad \mathbb{E} \sum_r c_r \log \left(e + \frac{B}{\eta c_r} \right) \leq CH[a + \text{Ent}(I) + 1 + \log(B/H)]. \quad (5.3)$$

We also have the following localized version. Fix $2 \leq s \leq K$, an integer $q \geq 0$, and a deterministic cutoff $r_0 \in \mathbb{N} \cup \{\infty\}$. Designate $s - 1$ arms, including the optimal arm, and suppose that each designated arm has mean at least the upper threshold of every Eliminate call made in a round $r < r_0$. Assume that every nondesignated arm belongs to a gap group with index at most q , and set

$$H_{\text{tail}} = \sum_{\substack{i \neq * \\ i \text{ not designated}}} \Delta_i^{-2}.$$

For this localized estimate only, assign all $s - 1$ designated arms to group q , regardless of their actual gaps.

Let $\mathcal{R}$ contain the entered rounds $r < r_0$, up to and including the first round $r \geq q + 2$ in which Median succeeds. If no such round occurs, let $\mathcal{R}$ contain every entered round before the cutoff or an earlier termination. A round rejected immediately by a budget test is included in $\mathcal{R}$. Then

$$\mathbb{E} \sum_{r \in \mathcal{R}} c_r \leq C[H_{\text{tail}} + (s - 1)4^q].$$

Proof. Step 1: A deterministic bound for the round weights. For the global estimate, treat the optimal arm as if it had gap g , and hence as if it belonged to group $Q = \lfloor \log_4 D \rfloor$. Define the

augmented group sizes and weights by

$$n'_j = |G_j| + \mathbf{1}_{\{j=Q\}}, \quad v_j = n'_j 4^j, \quad A_\ell = \sum_{j \geq \ell} n'_j.$$

Thus $\sum_j n'_j = K$.

Suppose that Median succeeds in round k . If $k \leq Q + 1$, then (5.2) implies that any subsequently entered round has at most $2A_{k-1}$ active arms. If $k \geq Q + 2$, the successful round leaves a singleton, so no later round is entered.

To bound $\mathbb{E}c_r$, decompose according to the last successful Median call before round r . The probability that all preceding $r - 1$ Median calls fail is at most κ^{r-1} . If the last success occurs in round $k < r$, the subsequent $r - k - 1$ Median calls must all fail, an event of conditional probability at most κ^{r-k-1} . Therefore

$$\mathbb{E}c_r \leq 4^r \left[K \kappa^{r-1} + 2 \sum_{k=1}^{r-1} \kappa^{r-k-1} A_{k-1} \right].$$

An earlier invalid Eliminate call, budget rejection, or other termination can only set some later c_r 's to zero, so the same bound holds for the stopped process.

For a fixed augmented arm in group j , its contribution to the expression in brackets is at most

$$\kappa^{r-1} + 2 \sum_{k=1}^{\min\{r-1, j+1\}} \kappa^{r-k-1} \leq 4\kappa^{\max\{0, r-j-2\}}.$$

Define the two-sided kernel

$$k_d = \begin{cases} 4^d, & d \leq 2, \\ 4^d \kappa^{d-2}, & d > 2. \end{cases}$$

Then

$$\mathbb{E}c_r \leq b_r, \quad b_r = 4 \sum_{j \geq 0} v_j k_{r-j}, \quad r \geq 1.$$

Because $\kappa = 0.01$,

$$\sum_{d \in \mathbb{Z}} k_d = \sum_{d \leq 2} 4^d + \sum_{d \geq 3} 4^d \kappa^{d-2} = \frac{64}{3} + \frac{64\kappa}{1-4\kappa} = 22.$$

In particular,

$$\sum_{r \geq 1} b_r \leq 88 \sum_j v_j.$$

Step 2: Entropy calculation. The probability vector $(k_d/22)_{d \in \mathbb{Z}}$ has finite entropy because both tails decay geometrically. Put

$$V = \sum_j v_j,$$

let X have law $(v_j/V)_j$, and let Y be independent of X with law $(k_d/22)_d$. Extend the convolution to all $r \in \mathbb{Z}$ by setting

$$\tilde{b}_r = 4 \sum_j v_j k_{r-j}.$$

Its total mass is $88V$, and its normalized distribution is the law of $X + Y$. Since $X + Y$ is a function of (X, Y) ,

$$\text{Ent}(X + Y) \leq \text{Ent}(X, Y) = \text{Ent}(X) + \text{Ent}(Y).$$

Consequently,

$$\Phi(\tilde{b}) = 88V \text{Ent}(X + Y) \leq 88\Phi(v) + 88V \text{Ent}(Y).$$

The sequence b is obtained by restricting $\tilde{b}$ to $r \geq 1$. Coordinatewise monotonicity of Φ , first for finite truncations and then by passage to the limit, gives

$$\Phi(b) \leq 88\Phi(v) + 88V \text{Ent}(Y).$$

The closest competitor belongs to G_Q , so $|G_Q| \geq 1$. Adding the artificial optimal arm therefore at most doubles the weight of that group. Since

$$|G_j|4^j \leq H_j,$$

we have $v_j \leq 2H_j$ for every j . The monotonicity and homogeneity of Φ now give

$$V \leq 2H, \quad \Phi(v) \leq \Phi((2H_j)_j) = 2H \text{Ent}(I).$$

It follows that

$$\sum_r b_r \leq CH, \quad \Phi(b) \leq CH(\text{Ent}(I) + 1),$$

which proves the first set of assertions.

Step 3: Logarithmically weighted cost. Set

$$f(z) = z \log \left(e + \frac{B}{\eta z} \right), \quad f(0) = 0.$$

This function is increasing and concave. Hence Jensen's inequality, followed by $\mathbb{E}c_r \leq b_r$, yields

$$\mathbb{E}f(c_r) \leq f(\mathbb{E}c_r) \leq f(b_r).$$

Let $N = \sum_r b_r$. For every r with $b_r > 0$,

$$e + \frac{B}{\eta b_r} \leq \frac{N}{b_r} \left(e + \frac{B}{\eta N} \right).$$

After taking logarithms, multiplying by b_r , and summing,

$$\sum_r f(b_r) \leq N \log \left(e + \frac{B}{\eta N} \right) + \Phi(b).$$

Since $N \leq CH$, $B \geq H$, and $a = \log(1/\eta)$, monotonicity of f gives

$$N \log \left(e + \frac{B}{\eta N} \right) \leq CH[a + 1 + \log(B/H)].$$

Together with the entropy bound for b , this proves (5.3).

Step 4: Localized estimate. Let $\mathcal{D}$ be the set of $s-1$ designated arms and put

$$W_{\text{loc}} = H_{\text{tail}} + (s-1)4^q.$$

For this step, set $c_r = 0$ outside the rounds in $\mathcal{R}$. Thus the cutoff r_0 , any earlier termination, and the first successful Median call at or after round $q+2$ are all incorporated into the definition of c_r .

Assign the designated arms artificially to group q and define

$$n_j^{\text{loc}} = |\{i \in G_j : i \notin \mathcal{D}\}| + (s-1)\mathbf{1}_{\{j=q\}}, \quad A_\ell^{\text{loc}} = \sum_{j \geq \ell} n_j^{\text{loc}}.$$

By the assumptions on the nondesignated arms,

$$\sum_{j=0}^q n_j^{\text{loc}} = K, \quad \sum_{j=0}^q n_j^{\text{loc}} 4^j \leq H_{\text{tail}} + (s-1)4^q = W_{\text{loc}}.$$

For $k \leq q+1$, every designated arm is counted in A_{k-1}^{loc} . Moreover, every nondesignated arm with $\Delta_i \leq 2\varepsilon_k = 2^{1-k}$ belongs to a group of index at least $k-1$. Therefore, if a successful Median round $k \leq q+1$ is followed by another entered round, (5.2) gives

$$|S_{k+1}| \leq 2A_{k-1}^{\text{loc}}.$$

The same decomposition according to the last successful Median call now yields, for $1 \leq r \leq q+2$,

$$\mathbb{E}c_r \leq 4^r \left[K\kappa^{r-1} + 2 \sum_{k=1}^{r-1} \kappa^{r-k-1} A_{k-1}^{\text{loc}} \right].$$

Applying the kernel estimate from Step 1 and summing over these rounds gives

$$\mathbb{E} \sum_{r=1}^{q+2} c_r \leq 4 \sum_{j=0}^q n_j^{\text{loc}} 4^j \sum_{r=1}^{q+2} k_{r-j} \leq 88W_{\text{loc}}.$$

It remains to control rounds after $q+2$. The active set never grows. For $j \geq 1$, in order for c_{q+2+j} to contribute to the localized sum, the Median calls in rounds $q+2, \dots, q+1+j$ must all fail; otherwise the sum stops after the first successful call. Conditionally on the history upon entering round $q+2$, these j failures have probability at most κ^j . Hence

$$\mathbb{E}c_{q+2+j} \leq (4\kappa)^j \mathbb{E}c_{q+2} \leq CW_{\text{loc}}(4\kappa)^j, \quad j \geq 1.$$

Since $4\kappa < 1$, summing this geometric tail and combining it with the estimate for the earlier rounds gives

$$\mathbb{E} \sum_{r \in \mathcal{R}} c_r \leq CW_{\text{loc}} = C[H_{\text{tail}} + (s-1)4^q].$$

□

5.3. Joint and localized errors. Until the end of this section, all probabilities refer to the comparison process. We first control the joint probability that several arms are eliminated before the rounds at which their gaps become distinguishable.

Fix one stage, and let $\mathcal{E}$ be the set of rounds in which Eliminate is called. Whenever such a call is made in round r , the first budget test has ensured $U_r + 4w_r < B$, and the subsequent update is $U_{r+1} = U_r + 4w_r$. Hence, pathwise,

$$\sum_{r \in \mathcal{E}} \beta_r = \frac{\rho}{B} \sum_{r \in \mathcal{E}} 4w_r = \frac{\rho}{B} \sum_{r \in \mathcal{E}} 4|S_r|4^r < \rho. \quad (5.4)$$

For a suboptimal arm $i \in G_q$, define

$$M_i = \mathbf{1}\{i \text{ is removed by an Eliminate call in some round } r < q\}.$$

Thus $M_i = 1$ means that i is removed before the algorithm reaches the scale associated with its gap. For the optimal arm, define

$$M_* = \mathbf{1}\{* \text{ is removed by an Eliminate call in any round}\}.$$

We refer to either event as a *premature elimination*, corresponding to [3, definition B.8].

The next lemma combines (5.4) with the product guarantee for Eliminate. Its backward-induction argument adapts [3, lemma B.9] to the comparison process.

Lemma 5.2. *For a fixed collection of k arms in one stage,*

$$\mathbb{P}(M_i = 1 \text{ for every arm in the collection}) \leq \rho^k. \quad (5.5)$$

Proof. The corrected mean estimate is at most $m + \varepsilon_r/8$, even if the true optimal arm was removed earlier. Thus the upper elimination threshold in round r is at most $m - \varepsilon_r/2$. For an arm in gap group $q > r$, $\Delta_i \leq 2^{-q} \leq \varepsilon_r/2$, so its mean is at least the upper elimination threshold. The optimal arm also lies above this threshold. Hence every arm whose removal in round r would be premature satisfies the hypothesis of the simultaneous-removal bound. Conditional on the past, a specified set of j eligible arms is therefore eliminated in that call with probability at most β_r^j .

Induct backward over the finitely many remaining rounds. If a specified arm has already been removed at or after its gap-group index, the joint event is impossible. Immediately before a call in round r , let

$$u = \rho - \sum_{\substack{v \in \mathcal{E} \\ v < r}} \beta_v.$$

This quantity is determined by the past. By (5.4), the current and future calls together receive total error allowance at most u on every continuation. Suppose that k specified arms still have to be eliminated prematurely. If the current round has reached the gap-group index of any specified suboptimal arm, the joint event is impossible. Otherwise all k specified arms are eligible for premature elimination in the current call. The induction claim is that their joint probability is at most u^k . Condition on the history immediately before the next Eliminate call, including the preceding Mean and Fraction outputs. Its error budget β is then fixed, and the total budget available to later calls is at most $u - \beta$ on every continuation. The conditional probability of any particular exact subset of j premature removals is at most β^j , since it is bounded by the probability that those j arms are all removed. Conditional on the resulting history, the probability that the remaining $k - j$ arms are all eliminated prematurely is at most $(u - \beta)^{k-j}$. Summing over subsets gives

$$\sum_{j=0}^k \binom{k}{j} \beta^j (u - \beta)^{k-j} = u^k.$$

If no further call is made and $k > 0$ specified arms still require premature elimination, the conditional probability is zero. If $k = 0$, the event is already satisfied and its conditional probability is one. These are the terminal cases of the induction. Initially the available budget is ρ , so the induction proves (5.5). $\square$

When only a few arms are close to optimal, we use a sharper bound involving the sum of the inverse squared gaps of the remaining, easier arms. The next lemma, analogous to [3, lemma B.6], controls the probability of an invalid Eliminate call before the algorithm reaches the scale needed to distinguish the difficult arms.

Order the arms so that

$$\mu_{[1]} = m > \mu_{[2]} \geq \dots \geq \mu_{[K]}, \quad \Delta_{[i]} = m - \mu_{[i]}.$$

Ties among the suboptimal arms may be ordered arbitrarily.

Lemma 5.3. *Fix $2 \leq s \leq K$. If $s = 2$, set $r_* = \infty$ and impose no restriction on B . If $s \geq 3$, assume*

$$B < \Delta_{[s-1]}^{-2}$$

and set

$$r_* = \left\lfloor \log_2 \frac{1}{\Delta_{[s-1]}} \right\rfloor.$$

Then

$$\mathbb{P}(\text{an invalid Eliminate call occurs in some round } r < r_*) \leq C s \rho \frac{\sum_{i=s}^K \Delta_{[i]}^{-2}}{B}. \quad (5.6)$$

Proof. Stop the analysis after the first invalid Eliminate call, including the round in which that call occurs. Set

$$q = \left\lfloor \log_2 \frac{1}{\Delta_{[s]}} \right\rfloor, \quad H_{\text{tail}} = \sum_{i=s}^K \Delta_{[i]}^{-2}.$$

Call the arms of ranks $1, \dots, s-1$ the difficult arms and those of ranks $s, \dots, K$ the tail arms.

Step 1: Separate the difficult and tail arms. For $r < r_*$,

$$\Delta_{[s-1]} \leq \frac{\varepsilon_r}{2}.$$

When $s = 2$, this remains true because $\Delta_{[1]} = 0$. The corrected Mean output satisfies

$$\hat{\mu}_r \leq m + \frac{\varepsilon_r}{8}.$$

Consequently, the upper mean thresholds in Fraction and Eliminate are at most

$$\hat{\mu}_r - \frac{9}{8}\varepsilon_r \leq m - \varepsilon_r, \quad \hat{\mu}_r - \frac{5}{8}\varepsilon_r \leq m - \frac{\varepsilon_r}{2}.$$

Every active difficult arm therefore lies above both upper thresholds.

The definition of q gives

$$\Delta_{[s]} > 2^{-(q+1)}.$$

Hence, if $r \geq q + 2$, then

$$\Delta_{[s]} > 2\varepsilon_r.$$

When Median succeeds in such a round, the corrected Mean output satisfies

$$\hat{\mu}_r \geq m - \frac{\varepsilon_r}{4}.$$

The lower mean thresholds in Fraction and Eliminate are therefore at least

$$m - 2\varepsilon_r \quad \text{and} \quad m - \varepsilon_r,$$

respectively. Every tail arm lies below both lower thresholds.

Step 2: Control the calls after a successful Median round. Suppose that the first successful Median call with index at least $q + 2$ occurs in a round $r < r_*$. Let γ be the fraction of the current active set consisting of tail arms.

If Fraction returns False, its validity gives

$$\gamma < \theta_r,$$

because every tail arm is below the lower Fraction threshold. The active set is unchanged. In any later round $v < r_*$, every arm below the upper Fraction threshold is a tail arm, whereas

$$\theta_{v-1} \geq \theta_r.$$

It follows inductively that True is invalid in every such round. All subsequent Fraction outputs before r_* are therefore False.

If Fraction returns True and the resulting Eliminate call is valid, every surviving tail arm is below the lower elimination threshold. The fraction of tail arms in the returned set is therefore at most 0.1. Since $\theta_{v-1} \geq 0.3$ in every later round v , the same induction shows that all subsequent Fraction outputs before r_* are False.

Thus, after the first successful Median call in a round $r \geq q + 2$, either the Eliminate call in that round is invalid or no further Eliminate call occurs before r_* .

Step 3: Bound the total weight of the relevant rounds. Let $\mathcal{R}$ be the collection of entered rounds before r_* , stopped after and including the first successful Median round with index at least $q + 2$, if such a round occurs. Step 2 shows that every possible first invalid Eliminate call before r_* occurs in a round belonging to $\mathcal{R}$.

Apply the localized part of [Lemma 5.1](#) with cutoff $r_0 = r_*$, designating the top $s - 1$ arms and assigning them artificially to group q . Step 1 verifies the required upper-threshold condition. Every tail arm has gap at least $\Delta_{[s]}$, so its gap-group index is at most q . The lemma therefore gives

$$\mathbb{E} \sum_{r \in \mathcal{R}} |S_r| 4^r \leq C[H_{\text{tail}} + (s - 1)4^q].$$

Since

$$4^q \leq \Delta_{[s]}^{-2} \leq H_{\text{tail}},$$

we obtain

$$\mathbb{E} \sum_{r \in \mathcal{R}} |S_r| 4^r \leq C s H_{\text{tail}}.$$

Step 4: Sum the probabilities of the first invalid call. Before the first invalid Eliminate call, the optimal arm remains active. Conditional on the past, the probability that the call in round r is invalid is therefore at most

$$\beta_r = \frac{4\rho |S_r| 4^r}{B}.$$

Summing over the possible locations of the first invalid call and using the preceding estimate,

$$\mathbb{P}(\text{an invalid Eliminate call before round } r_*) \leq \frac{4\rho}{B} \mathbb{E} \sum_{r \in \mathcal{R}} |S_r| 4^r \leq \frac{C s \rho H_{\text{tail}}}{B}.$$

Substituting the definition of H_{tail} proves (5.6). $\square$

5.4. Summing errors across budget scales. For each suboptimal arm, write

$$w_i = \Delta_i^{-2}.$$

At a stage with budget B , define

$$h = 1 + |\{i \neq * : w_i > B\}|, \quad x = \frac{1}{B} \sum_{\substack{i \neq * \\ w_i \leq B}} w_i.$$

Call the optimal arm and the $h-1$ suboptimal arms with $w_i > B$ *hard*; call the remaining suboptimal arms *easy*. Thus Bx is the sum of the inverse squared gaps of the easy arms.

Step 1: A joint bound for the hard arms. Suppose that a hard suboptimal arm $i \in G_q$ is removed in an executed round r . Its weight satisfies

$$B < w_i < 4^{q+1}.$$

Moreover, the first budget test and $|S_r| \geq 2$ give

$$8 \cdot 4^r \leq 4|S_r| 4^r < B < w_i < 4^{q+1}.$$

It follows that $r < q$, so every removal of a hard suboptimal arm is premature. Every removal of the optimal arm is premature by definition.

If a stage returns an incorrect arm, then the optimal arm and at least $h-2$ of the $h-1$ hard suboptimal arms must have been removed. For $h \geq 2$, take a union bound over the hard suboptimal arm that might remain. Applying (5.5) to each resulting collection of $h-1$ arms gives

$$J_1 = \rho, \quad J_h = (h-1)\rho^{h-1} \quad (h \geq 2) \tag{5.7}$$

as upper bounds for the stage's wrong-output probability.

Step 2: A bound using the easy arms. Suppose $h < K$, so at least one easy arm is present. The hard suboptimal arms have ranks $2, \dots, h$, while the easy arms have ranks $h+1, \dots, K$. Apply (5.6) with $s = h+1$; when $h = 1$, use its $s = 2$ case.

For $h \geq 2$, the cutoff r_* in that lemma is the gap-group index of the hard suboptimal arm with the largest gap. The calculation in Step 1 shows that every executed Eliminate call occurs before this cutoff. Since a wrong output requires the optimal arm to be removed, it requires an invalid Eliminate call before r_* . The same conclusion holds for $h = 1$, for which $r_* = \infty$. Because

$$\sum_{i=h+1}^K \Delta_{[i]}^{-2} = Bx,$$

the local bound gives

$$\mathbb{P}(\text{wrong output in the stage}) \leq C(h+1)\rho x. \quad (5.8)$$

If $h = K$, every suboptimal arm satisfies $\Delta_i < B^{-1/2}$. In any executed round,

$$8 \cdot 4^r < B,$$

and hence $\varepsilon_r = 2^{-r} > B^{-1/2}$. The upper Fraction threshold is at most $m - \varepsilon_r$, which is below every arm mean. Thus True is not valid, every corrected Fraction output is False, and the stage cannot accept. Its wrong-output probability is therefore zero.

Step 3: A necessary condition for acceptance. Assign truncated weights

$$v_* = 1, \quad v_i = \min\{w_i/B, 1\} \quad (i \neq *),$$

and write

$$v_{\text{tot}} = \sum_i v_i = h + x.$$

Every accepted stage satisfies

$$\sum_i v_i(1 - M_i) \leq 2. \quad (5.9)$$

To verify this, consider a suboptimal arm $i \in G_q$ removed in a round $r \geq q$. Then

$$Bv_i \leq w_i < 4^{q+1} \leq 4^{r+1}.$$

The total Bv_i over all such removals in round r is therefore at most

$$|S_r|4^{r+1} = 4|S_r|4^r,$$

the increment of U in that round. The first budget test ensures that the sum of these increments over all Eliminate calls is less than B . Hence the total weight of suboptimal arms removed at or after their gap-group indices is less than one.

The single returned arm contributes at most one more. If the output is wrong, then $M_* = 1$, so the original optimal arm contributes zero to the left side of (5.9). If the output is correct, the optimal arm is the returned arm. This proves (5.9).

Step 4: An exponential-moment bound for acceptance. Set

$$u = \log(1/\rho).$$

By choosing the absolute constant c sufficiently small, we may assume $u \geq 4$. Since $M_i \in \{0, 1\}$,

$$\exp\left(u \sum_i v_i M_i\right) = \prod_i [1 + (e^{uv_i} - 1)M_i].$$

All coefficients in this product expansion are nonnegative. Applying (5.5) to each product of indicators gives

$$\mathbb{E} \exp\left(u \sum_i v_i M_i\right) \leq \prod_i [1 + \rho(e^{uv_i} - 1)].$$

For $0 \leq v \leq 1$, convexity gives

$$e^{uv} - 1 \leq v(e^u - 1).$$

Therefore

$$\begin{aligned} \mathbb{E} \exp\left(u \sum_i v_i M_i\right) &\leq \exp\left(\rho(e^u - 1) \sum_i v_i\right) \\ &\leq e^{v_{\text{tot}}}, \end{aligned}$$

because $\rho(e^u - 1) = 1 - \rho \leq 1$.

On acceptance, (5.9) implies

$$\sum_i v_i M_i \geq v_{\text{tot}} - 2.$$

If $v_{\text{tot}} \geq 4$, Markov's inequality consequently gives

$$\begin{aligned} \mathbb{P}(\text{accept}) &\leq e^{v_{\text{tot}}} \rho^{v_{\text{tot}}-2} \\ &\leq \rho^{v_{\text{tot}}/4}. \end{aligned} \tag{5.10}$$

The final inequality uses $u = \log(1/\rho) \geq 4$. Since every incorrect output is an acceptance, (5.10) also bounds the wrong-output probability.

The preceding exponential-moment argument follows [3, lemmas B.11–B.12 and section B.5], with the truncated weights above.

Step 5: Sum over the geometric budget sequence. For a fixed value $h < K$, the corresponding stages form an interval in the sequence $B_t = 100^t$. Throughout this interval the easy-arm set is fixed, and each successive stage replaces x by $x/100$. We divide the stages into three ranges.

If $x < 1$ and $h = 1$, summing (5.8) over the geometric sequence of x 's gives a contribution of at most $C\rho$. If $x < 1$ and $h \geq 2$, then

$$\min\{J_h, C(h+1)\rho x\} \leq \sqrt{J_h C(h+1)\rho x} \leq Ch\rho^{h/2}\sqrt{x}.$$

For fixed h , the sum of $\sqrt{x}$ over these stages is at most $10/9$. Their total contribution is therefore at most $Ch\rho^{h/2}$, and summing over $h \geq 2$ gives $O(\rho)$.

If $1 \leq x < 16$, there is at most one stage for each h , because successive positive values of x differ by a factor of 100. Hence (5.7) gives a total contribution bounded by

$$J_1 + \sum_{h \geq 2} J_h = O(\rho).$$

Finally, suppose $x \geq 16$. For each fixed h , enumerate these stages backward from the last one in this range. At the j th preceding stage,

$$x \geq 16 \cdot 100^j.$$

Using $v_{\text{tot}} = h + x$ in (5.10), the total contribution of this range is at most

$$\sum_{h \geq 1} \rho^{h/4} \sum_{j \geq 0} \rho^{4 \cdot 100^j} = O(\rho).$$

Stages with $h = K$ contribute zero by Step 2. Generate the independent data for every stage in advance, so that the output of each stage is defined even if that stage is never reached. The event that the comparison process returns an incorrect arm is contained in the union of these stagewise wrong-output events. The preceding bounds and the union bound therefore give probability at most $C\rho$. Combining this estimate with (5.1), the error probability of the actual algorithm is at most

$$C\eta + C\rho = C\eta + C\eta^2.$$

Since $\eta = c\delta$, choosing the absolute constant $c > 0$ sufficiently small makes this quantity at most δ .

6. THE EXPECTED SAMPLE COMPLEXITY

It remains to bound the unconditional expected stopping time $\mathbb{E}T$. We first show that the outer Mean and Fraction calls use only a fixed fraction of each stage budget. We then show that, once the budget is sufficiently large, a stage accepts with probability at least $199/200$. Together with the deterministic sample limit and the geometric growth of the stage budgets, this will control the expected cost of all restarts.

6.1. The cost of the outer tests. The increment $\theta_r - \theta_{r-1}$ is indexed by $j = R_t + 1 - r$, the number of permitted rounds from r through R_t . After factoring out 4^{R_t} , its polynomial growth is dominated by the resulting geometric decay. The series is summable independently of the active sets.

Lemma 6.1. *There is an absolute integer J such that, in every stage t and on every sample path, the total number of samples used by the outer Mean and Fraction calls is at most $B_t/100$.*

Proof. For $1 \leq r \leq R_t$, we have

$$R_t \leq \log_4 B_t = (\log_4 100)t.$$

Since

$$\alpha_{t,r} = \frac{\eta}{50t^2r^2},$$

it follows that

$$\log \frac{1}{\alpha_{t,r}} = a + \log 50 + 2 \log t + 2 \log r \leq C[a + \log(t+1)] = CL_t.$$

Fix a permitted round r and put

$$j = R_t + 1 - r.$$

The Mean call has accuracy $\varepsilon_r/8$, so its cost is at most $C4^r L_t$. The two mean thresholds in the Fraction call differ by $\frac{5}{8}\varepsilon_r$, while its fraction thresholds differ by

$$\theta_r - \theta_{r-1} = \frac{1}{10j^2}.$$

The Fraction cost is therefore at most $C4^r L_t j^4 \log(10j^2)$. Since $j \geq 1$, the combined cost of the two outer calls in round r is at most

$$C4^r L_t j^6.$$

Summing over all permitted rounds, including rounds the stage may not reach, gives

$$\begin{aligned} \text{total outer-test cost} &\leq CL_t \sum_{r=1}^{R_t} 4^r (R_t + 1 - r)^6 \\ &= CL_t 4^{R_t} \sum_{j=0}^{R_t-1} 4^{-j} (j+1)^6 \\ &\leq CL_t 4^{R_t} \sum_{j=0}^{\infty} 4^{-j} (j+1)^6. \end{aligned}$$

The last series is finite and yields an absolute constant.

Whenever $R_t \geq 1$, its definition gives

$$L_t 4^{R_t} \leq 4^{-J} B_t.$$

Choose the absolute integer J sufficiently large that the resulting constant multiple of $4^{-J} B_t$ is at most $B_t/100$. If $R_t = 0$, the stage rejects before making any outer call, so its outer-test cost is zero. $\square$

6.2. Acceptance at a sufficient budget. The next lemma fixes the absolute multiplier M in the deterministic sample limit. It also introduces an absolute constant A used only in the analysis.

Lemma 6.2. *Define*

$$F = H(a + \text{Ent}(I)) + D\ell_g, \quad \ell_g = \log \log(e^e/g).$$

There are absolute constants M and A such that every stage with $B_t \geq AF$ accepts with probability at least $199/200$.

Proof. Write $B = B_t$ and $R = R_t$.

Step 1: Control rejection at the sample limit. Temporarily remove the deterministic sample limit, retaining both budget tests and the restriction to R rounds. On every executed round, $w_r < B/4$. If Eliminate is called, then

$$\beta_r = \frac{4\eta^2 w_r}{B},$$

and

$$\log \frac{1}{\beta_r} = \log \frac{B}{4\eta^2 w_r} \leq 2 \log \frac{B}{\eta w_r}.$$

The conditional expected Eliminate cost in round r is therefore at most Cz_r . The Median cost is at most $Cw_r \leq Cz_r$. Since

$$\sum_{\text{executed } r} z_r < 100B,$$

the tower property bounds the expected total cost of all Median and Eliminate calls by CB . By Lemma 6.1, the outer Mean and Fraction calls use at most $B/100$ additional samples. Thus the expected cost of the uncapped stage is at most C_0B on every instance, including paths on which subroutine errors occur.

Choose the absolute constant M sufficiently large. Markov's inequality then shows that the uncapped stage uses more than MB samples with probability at most $1/2000$. Coupling the capped and uncapped stages until the cap is reached shows that imposing the deterministic limit causes rejection with probability at most $1/2000$. The same estimate holds for the comparison process, since the preceding cost bound applies equally after correcting the outer outputs.

Step 2: Control rejection at the budget tests. In the comparison process, an Eliminate call in round r is invalid with conditional probability at most β_r . Consequently, (5.4) and the tower property give

$$\mathbb{P}(\text{some Eliminate call is invalid}) \leq \rho.$$

Stop the comparison process at its first invalid Eliminate call, but include the round containing that call and any round rejected by a budget test. If the U -test rejects, then

$$\sum_r c_r \geq \frac{B}{4}.$$

If the V -test rejects, then

$$\sum_r c_r \log \left(e + \frac{B}{\eta c_r} \right) \geq 100B.$$

Indeed, the corresponding sum without the e is $V_r + z_r$, which is at least $100B$.

For $B \geq H$, Markov's inequality and (5.3) therefore give

$$\mathbb{P}(\text{a budget test rejects before the first invalid Eliminate call}) \leq \frac{CH}{B} \left[a + \text{Ent}(I) + 1 + \log \frac{B}{H} \right]. \quad (6.1)$$

To bound the right side, put

$$\sigma = a + \text{Ent}(I), \quad y = \frac{B}{H}.$$

Since $F \geq H\sigma$, the condition $B \geq AF$ gives $y \geq A\sigma$. The function

$$y \mapsto \frac{\sigma + 1 + \log y}{y}$$

is decreasing for $y \geq 1$. Hence the right side of (6.1) is at most

$$C \frac{\sigma + 1 + \log(A\sigma)}{A\sigma} \leq C \frac{1 + \log A}{A}.$$

It can therefore be made arbitrarily small by choosing the absolute constant A sufficiently large.

Step 3: Ensure sufficiently many rounds. Set

$$f = \frac{F}{D}.$$

Since $H \geq D$,

$$f = \frac{H}{D}[a + \text{Ent}(I)] + \ell_g \geq a + \ell_g.$$

Let t_0 be the first stage index for which $B_{t_0} \geq AF$. By minimality,

$$B_{t_0} < 100AF = 100ADf,$$

and therefore

$$t_0 + 1 \leq 2 + \frac{\log(100A) + \log D + \log f}{\log 100}.$$

The definition of ℓ_g gives

$$\log D = 2\log(1/g) = 2(e^{\ell_g} - e) \leq 2e^f.$$

Since $f \geq 1$, we also have $\log f \leq e^f$. It follows that

$$t_0 + 1 \leq C[1 + \log A + e^f].$$

Using $a \leq f$, we obtain

$$L_{t_0} = a + \log(t_0 + 1) \leq 2f + C + \log(1 + \log A) \leq C_A f,$$

where

$$C_A = O(1 + \log \log(A + e)).$$

Consequently,

$$\frac{B_{t_0}}{L_{t_0}} \geq \frac{A}{C_A} D.$$

The ratio B_t/L_t is increasing in t , so the same bound holds at every later stage.

Fix an integer $b \geq 1$. Because $A/C_A \rightarrow \infty$ as $A \rightarrow \infty$, we may choose A large enough that every stage with $B_t \geq AF$ satisfies

$$R_t \geq Q + 2 + b.$$

Consider the comparison process without the deterministic sample limit. Suppose that no budget test rejects and no Eliminate call is invalid. If the active set has not become a singleton by the last permitted round, then every Median call in rounds

$$Q + 2, Q + 3, \dots, R_t$$

must fail. Each call uses fresh samples and has conditional failure probability at most κ . Since the list contains at least $b + 1$ calls, successive conditioning gives

$$\mathbb{P} \left(\begin{array}{c} \text{no singleton by the last permitted round,} \\ \text{no budget test rejects, and} \\ \text{no Eliminate call is invalid} \end{array} \right) \leq \kappa^{b+1}.$$

Step 4: Combine the rejection probabilities. First choose b so that κ^{b+1} is sufficiently small. Then choose A large enough both to obtain the required number of rounds and to make (6.1) sufficiently small. These choices can ensure that, in the comparison process without the deterministic sample limit, the probability of rejection before the first invalid Eliminate call, either at a budget test or at the final round, is at most $1/2000$.

The probability of an invalid Eliminate call is at most ρ . By (5.1), the within-stage probability that the actual and comparison processes disagree is at most $C\eta/t^2$. Choose the absolute constant c sufficiently small that

$$\rho \leq \frac{1}{2000}, \quad \frac{C\eta}{t^2} \leq \frac{1}{2000}$$

for every $t \geq 1$. Step 1 bounds the probability of rejection at the sample limit by $1/2000$. Summing these four contributions gives a rejection probability below $1/200$. Thus every stage with $B_t \geq AF$ accepts with probability at least $199/200$. $\square$

6.3. Summing the stage costs. Let t_* be the first index such that

$$B_{t_*} \geq AF.$$

By minimality,

$$B_{t_*} < 100AF.$$

Every stage uses fresh data and begins with all K arms. Hence Lemma 6.2, applied conditionally after each rejection, gives

$$\mathbb{P}(\text{the algorithm reaches stage } t_* + k) \leq 200^{-k}, \quad k \geq 0.$$

Since stage t uses at most $\lceil MB_t \rceil$ samples,

$$\begin{aligned} \mathbb{E}T &\leq \sum_{t < t_*} \lceil MB_t \rceil + \sum_{k \geq 0} 200^{-k} \lceil M100^k B_{t_*} \rceil \\ &\leq CB_{t_*} \leq CF. \end{aligned}$$

The reach probabilities tend to zero, so the algorithm also stops almost surely.

Finally,

$$a = \log(1/\eta) = \log(1/\delta) + \log(1/c).$$

Because c is absolute and $\delta < 0.01$, a is bounded by an absolute constant multiple of $\log(1/\delta)$. Therefore

$$\mathbb{E}T \leq C \{H(\log(1/\delta) + \text{Ent}(I)) + D \log \log(e^e/g)\}.$$

Together with the correctness bound from the preceding section, this proves Theorem 1.2.

Proof of Corollary 1.3. Apply the lower bound in Theorem 1.1 to $H(\log(1/\delta) + \text{Ent}(I))$ in Theorem 1.2. $\square$

APPENDIX A. CONVENTIONS IN THE ORIGINAL CONJECTURES

Our formulation follows that of [2, section 1 and definition 3.1] with minor convention differences. We record these below and explain how Theorems 1.1 and 1.2 still establish [2, conjectures 3.2 and 3.5] as formulated therein.

A.1. Gap-group boundary conventions. Choose a suboptimal arm i at random with probability Δ_i^{-2}/H . Let R be the index of its gap group under our convention and S the corresponding index under the convention $[2^{-s}, 2^{-s+1})$ in [2], including the group containing the endpoint 1. Every cell of either partition meets at most two cells of the other. Hence

$$\text{Ent}(S \mid R) \leq \log 2, \quad \text{Ent}(R \mid S) \leq \log 2.$$

The entropy chain rule gives

$$|\text{Ent}(R) - \text{Ent}(S)| \leq \log 2.$$

Thus the contributions $H \text{Ent}(I)$ differ by an additive $O(H)$, absorbed by $H \log(1/\delta)$. The definition of the gap groups in [3, definition 1.7] already agrees with ours.

A.2. Iterated logarithm. For $x = \log(1/g) \geq 0$,

$$0 \leq \log(e + x) - \log \max\{e, x\} \leq \log 2.$$

Our shifted iterated logarithm and the clipped form used in [2] therefore differ by at most a constant. Since $D \leq H$, the resulting $O(H)$ change is absorbed by $H \log(1/\delta)$. The unregularized logarithm is undefined at $g = 1$ and can be negative for large gaps; clipping makes the sample-complexity convention explicit.

A.3. Stopping conventions. The correctness definition in [2] only requires a correct output with probability at least $1 - \delta$, not that the algorithm also almost surely stops. The algorithms proving the upper bounds stop almost surely on every instance in $\mathcal{S}_K$, and therefore satisfy either correctness definition. The lower-bound proof does not require the algorithm to stop on the Gaussian alternative instances, as it uses only their correct-output probabilities at finite horizons. If the expected count on the target instance is infinite, its lower bound is immediate. Otherwise the algorithm stops almost surely on that instance, permitting the limits in the lower-bound proof. Thus the bounds also cover the original correctness convention.

These comparisons preserve all bounds up to absolute constants. Together with [Corollary 1.3](#), they give the stated formulations of the original conjectures.

APPENDIX B. SAMPLING SUBROUTINES

We prove the subroutine guarantees used in [Section 4](#) for 1-sub-Gaussian reward laws. Input sets may have tied optimal arms. Different calls use fresh samples. Within Fraction, arm identities are sampled independently across draws; conditional on these identities, sample blocks are independent. Fraction samples are also independent of the estimation blocks used to delete arms. All sample sizes are rounded up to integers. Accuracy parameters and differences between mean thresholds are at most one, so rounding is absorbed in the cost bounds.

B.1. Mean estimation. For independent samples $X_1, \dots, X_n$ from a 1-sub-Gaussian arm with mean μ , independence gives

$$\mathbb{E} \exp \left(\lambda \sum_{s=1}^n (X_s - \mu) \right) \leq e^{n\lambda^2/2}.$$

Writing $\bar{X}_n = n^{-1} \sum_{s=1}^n X_s$ and applying the Chernoff bound with $\lambda = x$ and $\lambda = -x$ yields, for $x > 0$,

$$\mathbb{P}(\bar{X}_n - \mu \geq x) \leq e^{-nx^2/2}, \quad \mathbb{P}(\bar{X}_n - \mu \leq -x) \leq e^{-nx^2/2}. \quad (\text{B.1})$$

This is the standard sub-Gaussian sample-mean bound; see [11, corollary 5.5]. For accuracy e and error level α , take

$$n(e, \alpha) = \left\lceil 2e^{-2} \log(2/\alpha) \right\rceil$$

samples and return their average. Equation (B.1) gives

$$\mathbb{P}(|\hat{\mu} - \mu| > e) \leq 2e^{-ne^2/2} \leq \alpha.$$

The count is deterministic and of order $e^{-2} \log(1/\alpha)$ on the stated parameter range. Fresh samples give the same bound conditional on the history before a call.

B.2. Median elimination. We use the median-elimination construction of [5], in the form stated in [3, fact 5.2]. Start with $S_1 = S$. While $|S_j| \geq 2$, put

$$e_j = \frac{e}{4} \left(\frac{3}{4}\right)^{j-1}, \quad a_j = 0.01 \cdot 2^{-j}.$$

Estimate each current mean to accuracy $e_j/2$ with individual failure probability at most $a_j/3$, and retain the $\lceil |S_j|/2 \rceil$ arms with the largest estimates, using fixed tie-breaking. Return the remaining arm.

Conditional on the current set, fix one optimal arm. Its estimate is inaccurate with probability at most $a_j/3$. If it is accurate and no arm whose mean is within e_j of its mean survives, then at least $\lceil |S_j|/2 \rceil$ retained arms have inaccurate estimates. The expected number of inaccurate estimates is at most $|S_j|a_j/3$, so Markov's inequality bounds the probability of this event by $2a_j/3$. Thus the maximum true mean drops by more than e_j with conditional probability at most a_j . As

$$\sum_{j \geq 1} e_j = e, \quad \sum_{j \geq 1} a_j = 0.01,$$

the returned arm has mean within e of the original maximum with probability at least 0.99.

Until termination, $|S_j| \leq 2|S|2^{-(j-1)}$. The deterministic total cost is bounded by

$$C|S|e^{-2} \sum_{j \geq 1} \left(\frac{8}{9}\right)^{j-1} (j+1) = O(|S|e^{-2}).$$

Fresh samples give the same guarantee conditional on an arbitrary history before the call.

B.3. Fraction testing. Write $\varepsilon = u - l$ and $d = \theta_+ - \theta_-$. Draw

$$m_F = \lceil 36d^{-2} \log(2/\alpha) \rceil$$

independent uniform arm identities from S , with replacement. For each draw, obtain a fresh mean estimate of accuracy $\varepsilon/2$ and error level $d/6$. Let Y_j indicate that this estimate is below $(l + u)/2$. Return True precisely when

$$m_F^{-1} \sum_{j=1}^{m_F} Y_j > (\theta_- + \theta_+)/2.$$

The variables Y_j are independent and identically distributed. If the fraction of arms with mean below l is at least θ_+ , then

$$\mathbb{E}Y_j \geq \theta_+ - d/6.$$

If the fraction with mean below u is at most θ_- , then

$$\mathbb{E}Y_j \leq \theta_- + d/6.$$

In the first case, False is an invalid answer; in the second, True is invalid. Either invalid answer requires a deviation of at least $d/3$ from the Bernoulli mean. Hoeffding's inequality bounds its probability by

$$\exp(-2m_F d^2/9) \leq \alpha/2.$$

These implications prove the stated True/False guarantees. The deterministic number of samples is

$$O\left(\varepsilon^{-2} \log(1/\alpha) d^{-2} \log(1/d)\right)$$

for $0 < \varepsilon \leq 1$, $0 < d \leq 0.1$, and $0 < \alpha < 0.1$. This is Algorithm 2 and Fact 5.3 of [3] with explicit rounding.

B.4. Elimination. We use the elimination procedure in [3, algorithm 3 and fact 5.4], returning immediately if the active set is empty and retaining arms whose estimates equal the deletion threshold. Fraction testing and deletion use separate samples. Put

$$\varepsilon = u - l, \quad v = (l + u)/2, \quad q_j = \frac{\alpha}{10 \cdot 2^j} \quad (j \geq 1).$$

Start with $S_1 = S$. At internal stage j :

- (1) If S_j is empty, return it. Otherwise call Fraction with mean thresholds l, v , fraction thresholds $0.05, 0.1$, and error level q_j .
- (2) If False, return S_j .
- (3) If True, estimate each current arm mean independently to accuracy $\varepsilon/4$ with error level q_j , and retain precisely the arms whose estimates are at least $v + \varepsilon/4$. Continue with the retained set.

For the cost calculation only, set the active-set size to zero after the procedure returns. Conditional on a current set, a valid True answer implies that more than $0.05|S_j|$ arms have mean below v . Each such arm survives the independent estimation step with probability at most q_j . A False answer makes the number of active arms in the next internal stage zero. On an invalid Fraction answer, the number of active arms in the next internal stage is still at most $|S_j|$. Averaging over this event, whose probability is at most q_j , gives

$$\mathbb{E}[|S_{j+1}| \mid S_j] \leq [1 - 0.05(1 - q_j)^2]|S_j| \leq 0.96|S_j|.$$

The contraction holds without conditioning on correctness. Iterating it gives

$$\mathbb{E}|S_j| \leq 0.96^{j-1}|S|.$$

The conditional stage cost is at most

$$C|S_j|\varepsilon^{-2}[\log(1/\alpha) + j].$$

Here the Fraction cost is absorbed because $|S_j| \geq 1$ while active. Summing the expected internal-stage costs bounds the unconditional expected total number of samples by

$$C|S|\varepsilon^{-2} \sum_{j \geq 1} 0.96^{j-1}[\log(1/\alpha) + j] = O(|S|\varepsilon^{-2} \log(1/\alpha)).$$

The same estimate gives

$$\mathbb{P}(\text{still active at internal stage } j) \leq \mathbb{E}|S_j| \leq 0.96^{j-1}|S|.$$

This tends to zero as $j \rightarrow \infty$, so the subroutine terminates almost surely.

We next verify the fraction guarantee and bound the probability that specified arms of mean at least u are all removed. The probability that any internal Fraction answer is invalid is at most $\sum_j q_j = \alpha/10$. If every such answer is valid, a nonempty set is returned only after a False answer and therefore has low-mean fraction less than 0.1 . The empty set satisfies the non-strict inequality automatically. For each input arm, generate independent estimation blocks in advance for all internal stages, separately from the Fraction samples. If an arm has mean at least u , every accurate estimate is at least

$$u - \varepsilon/4 = v + \varepsilon/4,$$

so the arm cannot be removed unless one of its own estimates is inaccurate. For each arm, the event that at least one estimate is inaccurate has probability at most $\alpha/10$. These events are independent across arms. Consequently the probability that all k specified arms with mean at least u are removed is at most $(\alpha/10)^k \leq \alpha^k$. This argument based on independent estimation errors follows [3, lemma B.9] and permits adaptive use of the blocks.

For one designated optimal arm of mean at least u , the probability of losing it or violating the bound on the number of returned arms with mean below l is at most $\alpha/5 < \alpha$. Fresh samples give these guarantees conditional on any history before the call.

REFERENCES

- [1] Lijie Chen and Jian Li. On the optimal sample complexity for best arm identification, 2015. arXiv:1511.03774.
- [2] Lijie Chen and Jian Li. Open problem: Best arm identification: Almost instance-wise optimality and the gap entropy conjecture. In Vitaly Feldman, Alexander Rakhlin, and Ohad Shamir, editors, *Proceedings of the 29th Annual Conference on Learning Theory*, volume 49 of *Proceedings of Machine Learning Research*, pages 1643–1646. PMLR, 2016.
- [3] Lijie Chen, Jian Li, and Mingda Qiao. Towards instance optimal bounds for best arm identification. In Satyen Kale and Ohad Shamir, editors, *Proceedings of the 2017 Conference on Learning Theory*, volume 65 of *Proceedings of Machine Learning Research*, pages 535–592. PMLR, 2017.
- [4] Brian M. Cho and Nathan Kallus. Exploration in the limit, 2026. arXiv:2601.00084.
- [5] Eyal Even-Dar, Shie Mannor, and Yishay Mansour. Action elimination and stopping conditions for the multi-armed bandit and reinforcement learning problems. *Journal of Machine Learning Research*, 7(39):1079–1105, 2006.
- [6] R. H. Farrell. Asymptotic behavior of expected sample size in certain one sided tests. *The Annals of Mathematical Statistics*, 35(1):36–72, 1964.
- [7] Aurélien Garivier and Emilie Kaufmann. Optimal best arm identification with fixed confidence. In Vitaly Feldman, Alexander Rakhlin, and Ohad Shamir, editors, *Proceedings of the 29th Annual Conference on Learning Theory*, volume 49 of *Proceedings of Machine Learning Research*, pages 998–1027. PMLR, 2016.
- [8] Kevin Jamieson, Matthew Malloy, Robert Nowak, and Sébastien Bubeck. lil’ UCB: An optimal exploration algorithm for multi-armed bandits. In Maria Florina Balcan, Vitaly Feldman, and Csaba Szepesvári, editors, *Proceedings of the 27th Conference on Learning Theory*, volume 35 of *Proceedings of Machine Learning Research*, pages 423–439. PMLR, 2014.
- [9] Zohar Karnin, Tomer Koren, and Oren Somekh. Almost optimal exploration in multi-armed bandits. In Sanjoy Dasgupta and David McAllester, editors, *Proceedings of the 30th International Conference on Machine Learning*, volume 28(3) of *Proceedings of Machine Learning Research*, pages 1238–1246. PMLR, 2013.
- [10] Emilie Kaufmann, Olivier Cappé, and Aurélien Garivier. On the complexity of best-arm identification in multi-armed bandit models. *Journal of Machine Learning Research*, 17(1):1–42, 2016.
- [11] Tor Lattimore and Csaba Szepesvári. *Bandit Algorithms*. Cambridge University Press, 2020.
- [12] Shie Mannor and John N. Tsitsiklis. The sample complexity of exploration in the multi-armed bandit problem. *Journal of Machine Learning Research*, 5:623–648, 2004.
- [13] Max Simchowitz, Kevin Jamieson, and Benjamin Recht. The simulator: Understanding adaptive sampling in the moderate-confidence regime. In Satyen Kale and Ohad Shamir, editors, *Proceedings of the 2017 Conference on Learning Theory*, volume 65 of *Proceedings of Machine Learning Research*, pages 1794–1834. PMLR, 2017.