

ReLU Neural Network Approximation to Smooth Functional Operator: Dimensional Decay and Error Analysis

Shuhao Jiao¹

Abstract

We study the uniform approximation of smooth scalar-valued functionals on an infinite-dimensional separable Hilbert space by deep ReLU neural networks. Writing the functional input as $X(t) = \sum_{d \geq 1} \xi_d \nu_d(t)$, we quantify the importance of coordinate d through $w_d s_d$, where s_d bounds the magnitude of the corresponding basis score and w_d controls the directional Fréchet sensitivity of the target functional. Our constructive analysis combines coordinate truncation, anisotropic partitioning, local Taylor approximation, and ReLU network realization, while allowing unrestricted interactions among the retained coordinates. We establish a general nonasymptotic upper bound for the uniform approximation error and a complementary pseudo-dimension-based lower bound for the worst-case approximation error. Under generalized exponential coordinate decay $w_d s_d \asymp \exp(-cd^\rho)$, with $\rho > 0$, the upper and lower bounds match at the leading order and thus yield the nearly optimal approximation rate, which is stretched-exponential in the logarithm of the network budget. This is the first work to characterize neural network approximation error for infinite-dimensional functional inputs explicitly through the joint dimensional decay of coordinate magnitudes and directional sensitivities.

1. Introduction

The approximation of nonlinear functionals defined on infinite-dimensional spaces is a fundamental problem in functional data analysis. Neural network architectures for functional inputs have a relatively long history. [Rossi & Conan-Guez \(2005\)](#) extended multilayer perceptrons to functional inputs and established universal approximation

and consistency results, while [Rossi et al. \(2005\)](#) investigated basis-based representations of functional observations for multilayer perceptrons and radial-basis-function networks. More recent studies have developed deep architectures that more explicitly preserve and exploit functional structure. [Yao et al. \(2021\)](#) introduced adaptive basis layers that learn task-dependent finite-dimensional representations, and [Thind et al. \(2023\)](#) incorporated functional input layers into densely connected neural networks. [Rao & Reimherr \(2023\)](#) proposed functional neural networks with continuous hidden layers and basis representations, whereas [Rügamer et al. \(2024\)](#) extended semi-structured neural networks to functional covariates. From an approximation-theoretic perspective, [Song et al. \(2023\)](#) derived quantitative approximation rates for smooth functionals using deep ReLU networks, and [Shi et al. \(2025\)](#) developed a kernel-embedding-based functional deep neural network together with corresponding approximation guarantees. These developments demonstrate the growing potential of deep networks for learning nonlinear relationships from infinite-dimensional functional inputs.

Let $\mathcal{H} = L^2[0, 1]$ denote the space of square-integrable functions on $[0, 1]$, and let $f_0: \mathcal{H} \rightarrow \mathbb{R}$ be an unknown scalar-valued functional. Given a fixed set of orthonormal basis functions $\{\nu_d(t): d \geq 1, t \in [0, 1]\}$ of $\mathcal{H}$, each $X(t) \in \mathcal{H}$ admits the representation $X(t) = \sum_{d \geq 1} \xi_d \nu_d(t)$, where $\xi_d = \langle X, \nu_d \rangle$. Although the input $X(t)$ is intrinsically infinite-dimensional, any implementable neural network can depend on only finitely many coordinates. Therefore, for some $D \in \mathbb{Z}_+$, we approximate $f_0(X)$ by a ReLU neural network $f_\phi(\xi_1, \dots, \xi_D)$ based on the first D basis scores and study the resulting uniform approximation error over a prescribed functional input domain. Under a suitable choice of basis (e.g., functional principle components), the leading scores can provide a parsimonious representation of the dominant variation in the functional input. In contrast, pointwise measurements are typically highly correlated, depend on the observation grid, and do not yield an equally transparent characterization of coordinate importance and truncation error.

A central feature of this problem is that the basis coordinates generally contribute unequally to the target functional.

¹Department of Biostatistics, City University of Hong Kong, Hong Kong SAR, China. Correspondence to: Shuhao Jiao <shuhao.jiao@cityu.edu.hk>.

It is found that that, the effective contribution of the d -th coordinate is governed by directional sensitivity of f_0 along $\nu_d(t)$ and the associated score magnitude. Coordinates with both large score magnitudes and strong directional sensitivities must be approximated accurately, whereas those with small score magnitudes or weak directional sensitivities can be truncated with limited loss. We consider a natural setting in which directional sensitivity decays with the coordinate index. Our objective is to characterize how this coordinate-decay structure, together with the smoothness of f_0 and the network’s width–depth budget, determines the approximation power of deep ReLU neural networks.

There is a substantial literature on neural network approximation theory. For deep ReLU networks, [Yarotsky \(2017; 2018\)](#) derived quantitative upper and lower bounds for smooth and continuous functions, while [Lu et al. \(2021\)](#) established nearly optimal width–depth approximation rates using local Taylor expansions. Approximation guarantees under Sobolev and Besov regularity have been developed by [Gühring et al. \(2020\)](#) and [Siegel \(2023\)](#). Complementary results have characterized the roles of sparse connectivity, parameter encoding, and network depth ([Bölcskei et al., 2019; Elbrächter et al., 2021; Yarotsky & Zhevnerchuk, 2020](#)). For structured target functions, [Petersen & Voigtlaender \(2018\)](#) obtained optimal rates for piecewise smooth functions, [Shen et al. \(2019\)](#) studied approximation through compositional structures, and [Shen et al. \(2020\)](#) characterized approximation rates in terms of network width and depth. Although . We refer to [DeVore et al. \(2021\)](#) for a comprehensive review of neural network approximation theory.

However, existing approximation theory does not explicitly characterize how the decay rates in the input scores and the directional sensitivities of the target functional jointly affect network approximation. While [Song et al. \(2023\)](#) and [Shi et al. \(2025\)](#) study neural-network approximation of functional operators, they do not explicitly characterize how coordinate-wise decay of function scores and anisotropic directional sensitivities jointly affect approximation. Although a functional input is intrinsically infinite-dimensional, its coordinates generally do not contribute equally. High-index coordinates often have smaller score magnitudes, and the target functional may be less sensitive to perturbations along these directions. If the d -th score is bounded by s_d and the corresponding directional sensitivity is measured by w_d , then its effective contribution is governed by $w_d s_d$. Consequently, only finitely many coordinates need to be retained to achieve a prescribed approximation accuracy, while the discarded coordinates contribute a truncation tail. Related coordinate-decay phenomena have been studied extensively in functional linear models; see, for example, [Cai & Hall \(2006\)](#) and [Hall & Horowitz \(2007\)](#). Those studies, however, primarily concern linear-functional estimation and do not address the approximation of nonlin-

ear functionals by ReLU networks.

Learning from functional data poses a distinctive capacity-allocation problem: any implementable neural predictor must compress an intrinsically infinite-dimensional input into a finite representation. Retaining too few coordinates causes information loss, whereas retaining too many requires a fixed network budget to be distributed across an increasingly high-dimensional input. We formalize this representation–capacity trade-off for general scalar-valued functionals on separable Hilbert spaces. Working directly with the basis coefficients of the functional input, we quantify the relevance of each coordinate through its magnitude and the directional Fréchet sensitivity of the target functional. This yields a sensitivity-aware effective dimension determined jointly by the geometry of the functional input domain, the target functional, and the available network budget. We establish non-asymptotic upper bounds and complementary lower bounds for deep ReLU networks, thereby characterizing the best attainable representation error under a finite computational budget. These results provide theoretically justified scaling rules for selecting the input truncation dimension jointly with network width and depth. We further quantify the diminishing importance of high-order coordinates through a generalized exponential decay profile, a flexible and natural family that explicitly links the decay parameter to the effective dimension and approximation rate.

The remainder of the paper is organized as follows. Section 2 introduces the problem formulation, network construction, and assumptions on coordinate-wise score decay and anisotropic directional sensitivities. Section 3 develops the approximation error analysis for the proposed FNN class. Section 4 presents the numerical experiments, and Section 5 concludes the paper. Technical proofs are provided in the appendix.

2. Preliminaries

2.1. Basis Representation and Reparametrization

Let $\mathcal{H}$ be a separable Hilbert space, and assume that $X(t) \in \mathcal{H}$ and $f_0: \mathcal{H} \rightarrow \mathbb{R}$. Let $\{\nu_d(t)\}_{d \geq 1}$ be a fixed orthonormal basis of $\mathcal{H}$, and write $X(t) = \sum_{d \geq 1} \xi_d \nu_d(t)$, where $\mathbf{x} = (\xi_1, \xi_2, \dots)$ denotes the corresponding score sequence. Define the coordinate map $\mathcal{T}: \mathcal{H} \rightarrow \ell^2$, $\mathcal{T}(X) = \mathbf{x}$. Since $\mathcal{T}$ is one-to-one, the regression functional f_0 admits a sequence-space representation: there exists a function $g_0: \ell^2 \rightarrow \mathbb{R}$ such that $f_0 = g_0 \circ \mathcal{T}$. Define $P_D: \mathcal{H} \rightarrow \text{span}\{\nu_1, \dots, \nu_D\} \subset \mathcal{H}$ be the orthogonal projection operator, say $P_D X = \sum_{d=1}^D \xi_d \nu_d$, where $D \in \mathbb{N} \cup \{\infty\}$. To evaluate the directional sensitivity of f_0 along different basis

directions, define

$$w_d := \max_{1 \leq s \leq m} \left\{ \sup_{X \in \mathcal{H}} |\mathcal{D}^s f_0(X)[\nu_d, \dots, \nu_d]| \right\}^{1/s}, \quad (2.1)$$

where $\mathcal{D}^s f_0(X)$ denotes the s -th Fréchet derivative of f_0 at $X(t)$. Clearly, large w_d indicates that f_0 varies more rapidly along the d -th basis direction, while small w_d indicates weaker sensitivity.

Remark 1. The power $1/s$ in the definition (2.1) normalizes derivatives of different orders. To see this, consider a perturbation along the d -th basis direction, say $X + h_d \nu_d$. The Taylor expansion of f_0 gives terms of the form $\mathcal{D}^s f_0(X)[\nu_d, \dots, \nu_d] h_d^s$. By the definition of w_d ,

$$|\mathcal{D}^s f_0(X)[\nu_d, \dots, \nu_d]| |h_d|^s \leq (w_d |h_d|)^s.$$

Thus w_d is the basic sensitivity scale paid each time the d -th direction appears in a Taylor expansion component. If $|h_d| \leq s_d$, the effective contribution of the d -th coordinate is therefore measured by $w_d s_d$. This explains why the joint decay of the directional derivative scale w_d and the score envelope s_d governs the approximation error and the effective dimension.

We now introduce the following assumption.

Assumption 1. Assume that $f_0: \mathcal{H} \rightarrow \mathbb{R}$ is m -times continuously Fréchet differentiable on $\mathcal{H}$.

For $s = 1, \dots, m$ and $X(t) \in \mathcal{H}$, let $\mathcal{D}^s f_0(X)[h_1, \dots, h_s]$ denote the s -th Fréchet derivative of f_0 at $X(t)$ applied to the directions $h_1, \dots, h_s$. Let $\Omega \subset \ell^2$ denote the domain of the score vector $\mathbf{x}$, and let $\mathcal{C}^m(\Omega)$ denote the class of functions on Ω with continuous partial derivatives up to order m . To relate the Fréchet derivatives of f_0 on $\mathcal{H}$ to the coordinate derivatives of its score representation, we introduce the following proposition.

Proposition 1. Let $\mathcal{L}: \mathcal{H} \rightarrow \ell^2$ be a bounded linear operator, and let $g: \ell^2 \rightarrow \mathbb{R}$ be m -times Fréchet differentiable on an open set containing $\mathcal{L}X$ for any $X(t) \in \mathcal{H}$. Define $f = g \circ \mathcal{L}$, then f is m -times Fréchet differentiable. Then for $s = 1, \dots, m$, and $h_1, \dots, h_s \in \mathcal{H}$,

$$\mathcal{D}^s f(X)[h_1, \dots, h_s] = \mathcal{D}^s g(\mathcal{L}X)[Lh_1, \dots, Lh_s].$$

In particular, if $\mathcal{L} = \mathcal{T} \circ P_D$, it follows that $\mathcal{L}X = (\xi_1, \dots, \xi_D, 0, 0, \dots)^\top$, and

$$\begin{aligned} \mathcal{D}^s f(X)[h_1, \dots, h_s] &= \sum_{d_1, \dots, d_s=1}^D \frac{\partial^s g}{\partial \xi_{d_1} \dots \partial \xi_{d_s}}(\mathcal{L}X) \\ &\quad \times \prod_{j=1}^s \langle h_j, \nu_{d_j} \rangle. \end{aligned}$$

Consequently, taking $h_j = \nu_{d'_j}$ for $j = 1, \dots, s$, where $d'_j \in \{1, \dots, D\}$, yields

$$\mathcal{D}^s f(X)[\nu_{d'_1}, \dots, \nu_{d'_s}] = \frac{\partial^s g}{\partial \xi_{d'_1} \dots \partial \xi_{d'_s}}(\mathcal{L}X).$$

Proposition 1 shows that directional Fréchet derivatives of f_0 along the basis $\{\nu_d\}$ coincide with coordinatewise partial derivatives of g_0 in the corresponding score representation. This representation is the key device that allows us to transfer directional sensitivity of regression functional into a form amenable to deep network approximation.

2.2. Network Construction

A major practical challenge is that a functional input $X(t) \in \mathcal{H}$ is infinite-dimensional, whereas any implementable neural network must take a finite-dimensional vector as input. To bridge this gap, we employ truncation. For a truncation level $D \in \mathbb{N}_+$, we approximate $X(t)$ by its projection $P_D X = \sum_{d=1}^D \xi_d \nu_d$, and use the truncated score vector $\mathbf{x}^{(D)} := (\xi_1, \dots, \xi_D)$ as the network input. We consider functions $f_\phi: \mathbb{R}^D \rightarrow \mathbb{R}$, parameterized by ϕ , belonging to a class $\mathcal{F}_{M,W,S}$ of feedforward neural networks with depth M , width W , and size S . We adopt the standard multi-layer perceptron (MLP) architecture, where f_ϕ is expressed as a composition of affine transformations and nonlinear activation functions:

$$f_\phi(\mathbf{x}^{(D)}) = \mathcal{L}_M \circ \rho \circ \mathcal{L}_{M-1} \circ \dots \circ \rho \circ \mathcal{L}_1 \circ \rho \circ \mathcal{L}_0(\mathbf{x}^{(D)}).$$

where $\rho(x) = \max\{0, x\}$ is the ReLU activation function applied componentwise, and each affine map is of the form $\mathcal{L}_i(x) = W_i x + b_i$, $i = 0, 1, \dots, M$. Here $W_i \in \mathbb{R}^{p_{i+1} \times p_i}$, $x \in \mathbb{R}^{p_i}$ and $b_i \in \mathbb{R}^{p_{i+1}}$, with $(p_0, p_1, \dots, p_M, p_{M+1})$ denoting the widths of each layer, where $p_0 = D$ and $p_{M+1} = 1$.

3. Error Analysis

3.1. Coordinatewise decay

To obtain error bounds, we need to quantify how much each basis coordinate contributes to the regression functional. This contribution is determined by two quantities: the size of the corresponding score and the sensitivity of the regression functional along that basis direction. Coordinates with small scores or weak directional sensitivity can be truncated with little loss, whereas coordinates with large values of both quantities must be retained by the network. The next assumption formalizes this joint coordinate-wise control through the score envelopes $\{s_d: d \geq 1\}$ and the directional sensitivity weights $\{w_d: d \geq 1\}$.

Assumption 2. There exists a constant $C_0 \geq 1$ and non-increasing sequence $\{w_d\}, \{s_d\}$ such that, for every finite multi-index $\alpha = (\alpha_1, \alpha_2, \dots)$ with $1 \leq \|\alpha\|_1 \leq m$, $\sup_{\mathbf{x} \in \Omega} |\partial^\alpha g_0(\mathbf{x})| \leq C_0 \prod_{d \geq 1} w_d^{\alpha_d}$. In addition, $|\xi_d| \leq s_d$, $d \geq 1$.

As $w_d \rightarrow 0$, the regression functional becomes progressively less sensitive to higher-index basis directions. The

next proposition shows that the directional-derivative decay condition in Assumption 2 is implied by product-type decay of higher-order interaction coefficients, thereby providing a concrete structural justification for this assumption. Assumption 2 holds for many regression function in statistical models, such as functional single-index model [Chen et al. \(2011\)](#), functional additive model [Müller & Yao \(2008\)](#), and functional quadratic regression model [Yao & Müller \(2010\)](#).

Proposition 2. Let $p \geq 1$ be fixed and consider the p -th order interaction model

$$f_0(X) = G(Q_p(X)), \quad \xi_d = \langle X, \nu_d \rangle, \\ Q_p(X) = \sum_{1 \leq j_1 \leq \dots \leq j_p} \eta_{j_1, \dots, j_p} \prod_{\ell=1}^p \xi_{j_\ell}.$$

Assume that G has bounded derivatives up to order m , that $\sum_{d \geq 1} w_d s_d < \infty$, and that the interaction coefficients satisfy the product-type decay condition

$$|\eta_{j_1, \dots, j_p}| \lesssim \prod_{\ell=1}^p w_{j_\ell}, \quad 1 \leq j_1 \leq \dots \leq j_p. \quad (3.1)$$

Then, for every $1 \leq s \leq m$ and every $d_1, \dots, d_s \geq 1$,

$$|\mathcal{D}^s f_0(X)[\nu_{d_1}, \dots, \nu_{d_s}]| \lesssim \prod_{\ell=1}^s w_{d_\ell}.$$

The product-type condition (3.1) can be viewed as a higher-order analogue of coefficient decay in functional linear regression (see e.g., [Cai & Hall \(2006\)](#)). Indeed, when $p = 1$, the interaction index reduces to the linear form $Q_1(X) = \sum_{j \geq 1} \eta_j \xi_j$, and the condition becomes $|\eta_j| \lesssim w_j$. Thus w_j plays the role of the coordinate-wise coefficient scale. For $p \geq 2$, the same principle requires interactions involving weak directions to be small: if any participating coordinate has a small sensitivity scale, then the corresponding higher-order interaction coefficient is also small. This is reasonable because an interaction term should not exert a pronounced effect if it involves a basis direction whose individual contribution is already weak.

3.2. Approximation framework

The network construction follows the local Taylor approximation framework for ReLU networks (see e.g., [Lu et al. \(2021\)](#)). The basic idea is to partition the input domain into small cells, approximate the local Taylor expansion on each cell, fit the Taylor coefficient maps on the resulting discrete grid, and then approximate the monomial terms appearing in the Taylor expansion by product networks. In the present functional setting, this construction must be adapted to the anisotropic sequence domain Ω .

For $D \geq 1$, define the projected regression functional $f_0^{(D)} := f_0 \circ P_D$, $f_0^{(D)}(X) = f_0(P_D X)$. Since P_D is

a bounded linear operator, Proposition 1 implies that, for $s = 1, \dots, m$,

$$\mathcal{D}^s f_0^{(D)}(X)[h_1, \dots, h_s] \\ = \mathcal{D}^s f_0(P_D X)[P_D h_1, \dots, P_D h_s].$$

Consequently, $\mathcal{D}^s f_0^{(D)}(X)[\nu_{d_1}, \dots, \nu_{d_s}] = 0$ whenever at least one of $d_1, \dots, d_s$ is greater than D . The Taylor expansion of the projected functional $f_0^{(D)}$ at X_0 is

$$f_0^{(D)}(X) = f_0^{(D)}(X_0) + \sum_{s=1}^{m-1} \frac{1}{s!} \sum_{d_1, \dots, d_s=1}^D \mathcal{D}^s f_0(P_D X_0) \\ \times [\nu_{d_1}, \dots, \nu_{d_s}] \prod_{j=1}^s \xi_{h, d_j} + R_{m,D}(X, X_0),$$

where

$$R_{m,D}(X, X_0) = \frac{1}{(m-1)!} \int_0^1 (1-\tau)^{m-1} \\ \times \mathcal{D}^m f_0(P_D X_0 + \tau h_D)[h_D, \dots, h_D] d\tau.$$

To approximate the original functional f_0 , we use the decomposition $f_0(X) = f_0^{(D)}(X) + \Delta_D(X)$. Under Assumption 2, the projection error satisfies

$$|\Delta_D(X)| = \left| \int_0^1 \mathcal{D} f_0(P_D X + \tau(I - P_D)X) \right. \\ \left. [(I - P_D)X] d\tau \right| \\ \leq C_0 \sum_{d>D} w_d |\xi_d| \leq C_0 \sum_{d>D} w_d s_d.$$

Thus, the approximation of f_0 consists of a finite-dimensional network approximation error for $f_0^{(D)}$ and a projection error arising from the discarded coordinates. The goal here is to approximate the above Taylor expansion up to order $m - 1$ through the following steps:

1. For a non-increasing sequence $\{L_d: d \geq 1\}$, define $D_0 := \max\{d: L_d \geq 1\}$, and let $D \geq D_0$ denote the dimension of the input scores. Partition $\Omega_{D_0} := [-s_1, s_1] \times \dots \times [-s_{D_0}, s_{D_0}]$ along its first D_0 coordinates. Let $\delta_d \in (0, 2s_d/L_d)$, put $a_{d,i} := -s_d + 2s_d i/L_d$, and construct the trifling region

$$\mathcal{R}(\Omega_{D_0}, L, \delta) = \bigcup_{d=1}^{D_0} \left\{ \xi \in \Omega_{D_0} : \right. \\ \left. \xi_d \in \bigcup_{i=1}^{L_d-1} (a_{d,i} - \delta_d, a_{d,i}] \right\}.$$

We denote the resulting cells in $\Omega_{D_0} \setminus \mathcal{R}(\Omega_{D_0}, \mathbf{L}, \delta)$ by P_θ , where $\theta_d \in \{0, 1, \dots, L_d - 1\}$. Then construct a neural network ψ that approximately maps each $\mathbf{x}^{(D_0)} \in P_\theta$ to

$$\left(-s_1 + \frac{2s_1\theta_1}{L_1}, \dots, -s_{D_0} + \frac{2s_{D_0}\theta_{D_0}}{L_{D_0}}\right). \quad (3.2)$$

Thus, only the first D_0 coordinates are partitioned, and the number of cells is $\prod_{d=1}^{D_0} L_d$.

The decrease in L_d reflects the heterogeneous importance of the coordinate directions. The quantity L_d is the number of subintervals assigned to the d -th coordinate. Since the range of this coordinate has length $2s_d$, its cell width is of order s_d/L_d . Under the directional derivative bound, a perturbation of this magnitude changes the target functional by at most a constant multiple of $w_d s_d/L_d$. Consequently, coordinates with larger $w_d s_d$ require finer partitions.

2. For each cell $P_\theta^{(D_0)}$, define

$$P_\theta^{(D)} := P_\theta^{(D_0)} \times \prod_{d=D_0+1}^D [-s_d, s_d],$$

$$X_\theta(t) := \sum_{d=1}^{D_0} \left(-s_d + \frac{2s_d\theta_d}{L_d}\right) \nu_d(t).$$

Thus, $\mathbf{x}^{(D)} \in P_\theta^{(D)}$ if and only if its first D_0 coordinates $\mathbf{x}^{(D_0)} = (\xi_1, \dots, \xi_{D_0})$ belong to $P_\theta^{(D_0)}$; the remaining coordinates $(\xi_{D_0+1}, \dots, \xi_D)$ do not determine the cell index θ . For $s = 1, \dots, m-1$ and $\mathbf{d} = (d_1, \dots, d_s) \in \{1, \dots, D\}^s$, define the piecewise-constant Taylor coefficient map by

$$\mathcal{Q}_\mathbf{d}^{(s)}(\mathbf{x}^{(D)}) := \sum_{\theta} \mathcal{D}^s f_0^{(D)}(X_\theta)$$

$$[\nu_{d_1}, \dots, \nu_{d_s}] \mathbf{1}_{P_\theta^{(D_0)}}(\mathbf{x}^{(D_0)}).$$

for $\mathbf{x}^{(D)} \in \bigcup_{\theta} P_\theta^{(D)}$. Since $X_\theta \in \text{span}\{\nu_1, \dots, \nu_{D_0}\}$ and $d_1, \dots, d_s \leq D$, $\mathcal{D}^s f_0^{(D)}(X_\theta)[\nu_{d_1}, \dots, \nu_{d_s}] = \mathcal{D}^s f_0(X_\theta)[\nu_{d_1}, \dots, \nu_{d_s}]$. In particular, if $\mathbf{x}^{(D_0)} \in P_\theta^{(D_0)}$, then

$$\mathcal{Q}_\mathbf{d}^{(s)}(\xi_1, \dots, \xi_{D_0}, \xi_{D_0+1}, \dots, \xi_D)$$

$$= \mathcal{Q}_\mathbf{d}^{(s)}(\xi_1, \dots, \xi_{D_0}, 0, \dots, 0)$$

$$= \mathcal{D}^s f_0(X_\theta)[\nu_{d_1}, \dots, \nu_{d_s}].$$

We then construct a network $\phi_\alpha(\mathbf{x}^{(D)})$ to approximate $\mathcal{Q}_\mathbf{d}^{(s)}(\mathbf{x}^{(D)})$ through point fitting, where α indicates the multi-index and $\|\alpha\|_1 = s$. For each fixed $\mathbf{d}$, the coefficient map has only $\prod_{d=1}^{D_0} L_d$ distinct values, one for each D_0 -dimensional cell (3.2).

3. For $\mathbf{x}^{(D_0)} \in P_\theta$, write

$$\xi_{h_\theta, d} := \begin{cases} \xi_d + s_d - \frac{2s_d\theta_d}{L_d}, & d \leq D_0, \\ \xi_d, & D_0 < d \leq D. \end{cases}$$

For $\mathbf{d} = (d_1, \dots, d_s) \in \{1, \dots, D\}^s$, construct a network $P_\alpha(\cdot)$ that approximates the monomial

$$\prod_{j=1}^s \xi_{h_\theta, d_j} = \prod_{d=1}^{D_0} \left\{ \xi_d - \psi_d(\mathbf{x}^{(D_0)}) \right\}^{\alpha_d} \prod_{d=D_0+1}^D \xi_d^{\alpha_d}.$$

In particular, we first construct a network $\phi_\times$ that approximates the scalar product map $(x, y) \mapsto xy$, and then compose copies of $\phi_\times$ to approximate the required monomials.

Then the neural network construction can be written as

$$\phi(\mathbf{x}^{(D)}) = \sum_{\|\alpha\|_1 \leq m-1} \phi_\times \left(\frac{\phi_\alpha(\mathbf{x}^{(D)})}{\alpha!}, \right.$$

$$\left. P_\alpha \left(\mathbf{x}^{(D_0)} - \psi(\mathbf{x}^{(D_0)}), \xi_{D_0+1}, \dots, \xi_D \right) \right),$$

where $\alpha = (\alpha_1, \dots, \alpha_D)$, $\alpha! := \prod_{d=1}^D \alpha_d!$.

Fix integers D define the finite-dimensional slice $\Omega_D := \prod_{d=1}^D [-s_d, s_d]$. Let $\mathbf{w} = (w_1, \dots, w_D)$, and let $\mathcal{C}^m(\Omega_D, \mathbf{w})$ be the class of functions with continuous partial derivatives up to order m satisfying

$$\|\partial^\alpha g\|_{L^\infty(\Omega_D)} \leq C_0 \prod_{d=1}^D w_d^{\alpha_d}$$

for every multi-index $\alpha \in \mathbb{N}_0^D$ with $1 \leq \|\alpha\|_1 \leq m$. The following proposition gives an upper bound for the approximation error of the resulting network on Ω_D .

In the sequel, let $a_n \asymp b_n$ denote $C^{-1}b_n < a_n \leq Cb_n$ for some $C > 1$, $a_n \lesssim b_n$ denote $a_n \leq Cb_n$ for some $C > 0$, and $a_n \gtrsim b_n$ denote $a_n \geq Cb_n$ for some $C > 0$, where “ C ” denotes a generic positive constant. Let

$$H(D) := \sum_{d>D} w_d s_d, \quad \Psi(D) := \sum_{d=1}^D \log\{(w_d s_d)^{-1}\},$$

then we introduce the following results on approximation error.

Theorem 1. Suppose that Assumption 1 and 2 hold. For $W_\phi, M_\phi \in \mathbb{Z}_+$, let $\mathcal{A} = W_\phi M_\phi$ and

$$D_0 = \max_d \left\{ w_d s_d \exp \left[\frac{\log \mathcal{A}^2 + \Psi(d)}{d} \right] \geq 2 \right\}.$$

Suppose that D satisfies $H(D) \asymp H(D_0)^m$. Then there exists a ReLU network class $\mathcal{F}$ with input dimension D , size S , width W , and depth M satisfying $S \lesssim 3^{D_0} \{D^{m-1} W_\phi^2 M_\phi + 1\}$, $W \lesssim 3^{D_0} \{D^{m-1} W_\phi + 1\}$, and $M \lesssim M_\phi + 2D_0$, such that

$$\sup_{f_0 \in \mathcal{C}^m(\Omega, \mathbf{w})} \inf_{\phi \in \mathcal{F}} \|\phi - f_0\|_{L^\infty(\Omega)} \lesssim D_0^m \exp \left[-\frac{m}{D_0} \{\log \mathcal{A}^2 + \Psi(D_0)\} \right].$$

Theorem 1 gives the approximation bound

$$D_0^m \exp \left[-\frac{m}{D_0} \{\log \mathcal{A}^2 + \Psi(D_0)\} \right].$$

We refer to D_0 as the *effective dimension*, as it quantifies the number of input coordinates that can be retained in the network approximation under a given network budget. Ignoring the polynomial prefactor D_0^m , the leading exponential term in Theorem 1 can be written as $\exp\{-m\varphi(\log \mathcal{A}^2)\}$, where $\varphi(t) = \{t + \Psi(D_0(\mathcal{A}))\}/D_0(\mathcal{A})$. To determine whether the resulting approximation rate can be polynomial in the network budget, it is useful to compare $\varphi(t)$ with $t = \log \mathcal{A}^2$. By definition, $\varphi(t)/t = 1/D_0(\mathcal{A}) + \Psi(D_0(\mathcal{A}))/\{tD_0(\mathcal{A})\}$. Thus, when the effective dimension $D_0(\mathcal{A})$ diverges and the cumulative coordinate-decay term $\Psi(D_0(\mathcal{A}))$ grows more slowly than $tD_0(\mathcal{A})$, the exponent satisfies $\varphi(t) = o(t)$ and then the approximation error decay in a subpolynomial rate.

Intuitively, as more coordinates must be retained, the available network budget is distributed over an expanding input space, preventing the leading exponent from being of order t and the polynomial rate cannot be achieved. The generalized exponential decay regime (3.3) verifies these properties explicitly. In this case, $D_0(\mathcal{A}) \asymp t^{1/(\rho+1)} \rightarrow \infty$ and $\Psi(D_0(\mathcal{A})) \asymp D_0(\mathcal{A})^{\rho+1} \asymp t$. Consequently, $\varphi(t) \asymp t/D_0(\mathcal{A}) \asymp t^{\rho/(\rho+1)} = o(t)$. Therefore, up to polynomial factors in $D_0(\mathcal{A})$, the leading approximation profile is sub-polynomial in $\mathcal{A}^2$. In particular, for every $\alpha > 0$, $\exp\{-c\varphi(\log \mathcal{A}^2)\} \gg (\mathcal{A}^2)^{-\alpha}$ as $\mathcal{A}^2$ is sufficiently large. Together with the preceding lower-bound comparison, this shows that the best attainable approximation rate in the general setting is sub-polynomial in the network budget.

The following proposition gives the lower bound of the approximation error.

Theorem 2. Let $J \geq 1$, and let $\mathcal{F}$ be any approximation class with pseudo-dimension $V = \text{PDim}(\mathcal{F})$. Suppose that $\mathcal{F}$ uniformly approximates $\mathcal{C}^m(\Omega, \mathbf{w})$ with error ε , that is,

$$\sup_{f \in \mathcal{C}^m(\Omega, \mathbf{w})} \inf_{\phi \in \mathcal{F}} \|\phi - f\|_{L^\infty(\Omega)} = \varepsilon.$$

For any J such that $(\frac{C_0}{2\varepsilon})^{1/m} w_J s_J \geq 1$, we have

$$\varepsilon \gtrsim \exp \left[-\frac{m}{J} \{\log V + \Psi(J)\} \right].$$

To compare the lower bound with the constructive upper bound, apply Theorem 2 with $J = D_0$ to the restriction of the constructed network class to the D_0 -dimensional slice obtained by setting $\xi_{D_0+1} = \dots = \xi_D = 0$. Restricting the input domain cannot increase the pseudo-dimension. Therefore, Theorem 2 gives $\varepsilon_{\text{low}} \gtrsim \exp[-m\{\log V + \Psi(D_0)\}/D_0]$, provided that $\{C_0/(2\varepsilon)\}^{1/m} w_{D_0} s_{D_0} \geq 1$. If this admissibility condition fails, then $\varepsilon > (C_0/2)(w_{D_0} s_{D_0})^m$. By the definition of D_0 , we have $\varepsilon \gtrsim (w_{D_0} s_{D_0})^m \gtrsim \exp[-m\{\log \mathcal{A}^2 + \Psi(D_0)\}/D_0]$. Thus the failure of the admissibility condition also implies a lower bound with the same leading exponential rate.

For the constructed network class, $V = \text{PDim}(\mathcal{F}) \lesssim SM \log S$; see Bartlett et al. (2019). Since $S \lesssim 3^{D_0} \{D^{m-1} W_\phi^2 M_\phi + 1\}$, $M \lesssim M_\phi + 2D_0$, and $\mathcal{A} = W_\phi M_\phi$, it follows that $V \lesssim 3^{D_0} D^m \mathcal{A}^2 \log(3^{D_0} D^m \mathcal{A}^2)$. Consequently,

$$\varepsilon_{\text{low}} \gtrsim \exp[-m\{\log \mathcal{A}^2 + \Psi(D_0)\}/D_0] \times 3^{-m} D^{-m^2/D_0} \{\log(3^{D_0} D^m \mathcal{A}^2)\}^{-m/D_0}.$$

On the other hand, Theorem 1 gives $\varepsilon_{\text{up}} \lesssim D_0^m \exp[-m\{\log \mathcal{A}^2 + \Psi(D_0)\}/D_0]$. Thus, the upper and lower bounds share the same leading exponential term, while their discrepancy is confined to the multiplicative factor $3^m D_0^m D^{m^2/D_0} \{\log(3^{D_0} D^m \mathcal{A}^2)\}^{m/D_0}$.

Under generalized exponential coordinate decay

$$w_d s_d \asymp \exp(-cd^\rho), \quad (3.3)$$

for each fixed $\rho > 0$, the tail satisfies $\log H(j) = -cj^\rho + o(j^\rho)$. Therefore, $H(D) \asymp H(D_0)^m$ gives $D = m^{1/\rho} D_0 \{1 + o(1)\}$, and therefore $D \asymp D_0$. Moreover, the definition of D_0 gives $D_0 \asymp (\log \mathcal{A}^2)^{1/(\rho+1)}$. It follows that

$$D^{m^2/D_0} = \exp\{m^2 \log(D)/D_0\} = 1 + o(1)$$

Moreover, since $\log(3^{D_0} D^m \mathcal{A}^2) = O(\log \mathcal{A}^2)$, $\{\log(3^{D_0} D^m \mathcal{A}^2)\}^{m/D_0} = 1 + o(1)$. Consequently, the multiplicative discrepancy is of order at most $3^m D_0^m \{1 + o(1)\} \asymp (\log \mathcal{A}^2)^{m/(\rho+1)}$. Its logarithm is $O(\log \log \mathcal{A})$, which is negligible relative to the common leading exponent $(\log \mathcal{A}^2)^{\rho/(\rho+1)}$. Consequently, the constructive upper bound is nearly optimal on the leading exponential scale.

3.3. Approximation Error Analysis under Generalized Exponential Decay Profile

Following the discussion in the previous section, we consider a generalized exponential coordinate decay regime to quantify the coordinate decay rate. We allow the $w_d s_d$ to

decay as a^{-qd^ρ} , where $q := \tau + \tau_\omega$ and $\rho > 0$. This formulation contains three regimes in a unified way: $0 < \rho < 1$ corresponds to stretched-exponential decay, $\rho = 1$ recovers the ordinary exponential case, and $\rho > 1$ corresponds to super-exponential decay. This general exponential-decay regime is particularly natural for smooth functional inputs possessing rapidly decaying basis coefficients.

Let $q = \tau + \tau_\omega$, $c_a = q \log a$, $\mathcal{A}^2 = W_\phi^2 M_\phi^2$, and $t = \log \mathcal{A}^2$. The following theorem gives a stretched-exponential approximation bound and identifies its one-dimensional limiting profile as ρ diverges.

Theorem 3. Suppose that Assumptions 1 and 2 hold. Fix $a > 1$, $\tau > 0$, and $\tau_\omega > 0$. For $\rho > 0$, assume that the score envelopes and directional sensitivity weights satisfy $s_d \asymp a^{-\tau d^\rho}$ and $w_d \asymp a^{-\tau_\omega d^\rho}$, where the comparison constants are independent of d , ρ , and $\mathcal{A}$. Let $t = \log \mathcal{A}^2$, $c_a = (\tau + \tau_\omega) \log a$, and $c' = m c_a^{1/(\rho+1)} \{(\rho+1)/\rho\}^{\rho/(\rho+1)}$, and suppose that $t \geq \log 2$. Let D_0 be defined as in Theorem 1, and choose $D \geq D_0$ such that $H(D) \asymp H(D_0)^m$. Then the ReLU network class $\mathcal{F}$ constructed in Theorem 1 satisfies

$$\inf_{f \in \mathcal{F}} \|f - f_0\|_{L^\infty} \lesssim \exp\{-c' t^{\frac{\rho}{\rho+1}} (1 + \delta_{\mathcal{A},\rho})\}, \quad (3.4)$$

where

$$1 + \delta_{\mathcal{A},\rho} = \frac{D_0^{-1} \{t + \Psi(D_0)\} - \log D_0 - \Psi(1)}{c_a^{1/(\rho+1)} \{t(\rho+1)/\rho\}^{\rho/(\rho+1)}}. \quad (3.5)$$

and $\delta_{\mathcal{A},\rho} \rightarrow 0$ in either of the following two regimes: $\mathcal{A} \rightarrow \infty$ with $\rho > 0$ fixed, or $\rho \rightarrow \infty$ with $\mathcal{A} \geq 2$ fixed.

The bound has two complementary asymptotic interpretations. For every fixed finite $\rho > 0$, the approximation error decays sub-polynomially in the network budget as $\mathcal{A} \rightarrow \infty$, reflecting the intrinsic difficulty of approximating an unrestricted smooth functional with genuinely infinite-dimensional inputs. By contrast, as $\rho \rightarrow \infty$, the coordinate importance decays increasingly rapidly, and the approximation problem progressively approaches a univariate one. The upper-bound rate consequently converges to $\mathcal{A}^{-2m}$, the approximation rate for a C^m -smooth univariate function.

4. Numerical experiment results

Let $\{\nu_d(t) : d \geq 1\}$ be the Fourier sine basis on $[0, 1]$. We approximate the infinite-dimensional functional input by $X(t) = \sum_{d=1}^{D_{\text{total}}} \xi_d \nu_d(t)$, where $D_{\text{total}} = 100$. Write $\xi_d = s_{d,\rho} u_d$, where the standardized scores u_d are generated independently from $\text{Unif}[-1, 1]$. To impose generalized exponential decay, we define $a_{d,\rho} = \exp(-\lambda d^\rho)$, where $\lambda = 0.5$. We then set $s_{d,\rho} = a_{d,\rho}^{1/2}$ and $w_{d,\rho} = a_{d,\rho}^{1/2}$, so that $w_{d,\rho} s_{d,\rho} = a_{d,\rho}$. Thus, both the score magnitude and the directional sensitivity decay with the coordinate index, and there is no non-decaying low-dimensional component.

We consider $\rho \in \{0.5, 1, 2\}$, corresponding respectively to stretched-exponential, ordinary exponential, and super-exponential decay.

Let $z_{d,\rho} = w_{d,\rho} \xi_d = a_{d,\rho} u_d$, and let $\mathcal{I} = \{(j_r, \ell_r) : r = 1, \dots, R\}$ be a fixed collection of $R = 64$ distinct coordinate pairs sampled uniformly without replacement from all pairs $\{(j, \ell) : 1 \leq j < \ell \leq D_{\text{total}}\}$. We define

$$f_{0,\rho}(X) = \sum_{k=1}^K b_k \sin \left\{ \beta_k + \sum_{d=1}^{D_{\text{total}}} \gamma_{kd} z_{d,\rho} + \frac{\eta}{\sqrt{R}} \sum_{r=1}^R \delta_{kr} \sin(z_{j_r,\rho}) \sin(z_{\ell_r,\rho}) \right\}.$$

Here, $K = 128$ and $\eta = 0.75$. The phase parameters β_k are generated independently from $\text{Unif}[0, 2\pi]$, while γ_{kd} and δ_{kr} are independent Rademacher variables taking values in $\{-1, 1\}$ with equal probability. The coefficients b_k are first generated independently from $N(0, 1)$ and then normalized so that $\sum_{k=1}^K |b_k| = 1$. The same realizations of $\mathcal{I}$, β_k , γ_{kd} , δ_{kr} , and b_k are used for all values of ρ , D , and the network width. The products $\sin(z_{j_r,\rho}) \sin(z_{\ell_r,\rho})$ introduce explicit smooth pairwise interactions, while the outer sine introduces additional nonadditive dependence among the coordinates and keeps the target functional uniformly bounded.

For each $D \in \{1, 5, 10, 30, 50, 70\}$, define the truncated target functional by $f_{0,\rho}^{(D)}(X) = f_{0,\rho}(P_D X)$, where $P_D X = \sum_{d=1}^D \xi_d \nu_d$. A fully connected ReLU network is trained to approximate $f_{0,\rho}^{(D)}$ using the standardized scores $(u_1, \dots, u_D)$ as its input. We use four hidden layers with a common hidden-layer width $W \in \{2, 4, 8, 16, 32, 64\}$. For each value of W , the depth ($M = 4$) and all other architectural specifications are held fixed across D and ρ .

We generate 50,000 training observations, 10,000 validation observations, and 50,000 independent test observations. The networks are trained using a smooth approximation to the empirical L^∞ loss, implemented through the log-sum-exp loss with smoothing parameter 0.05. Optimization is performed using Adam with an initial learning rate of 10^{-3} , a batch size of 1024, and at most 2000 optimization steps. Early stopping is determined by the empirical total L^∞ error on the validation sample.

For each combination of ρ , D , and W , we train the network from ten independent initializations and retain the initialization with the smallest validation total error. For a given width budget W , all fitted networks with hidden-layer width no larger than W are regarded as admissible, and the final model is again selected using only the validation error. We report the empirical total L^∞ error of the selected model on the independent test sample, where the error is evaluated against the full target $f_{0,\rho}(X)$. This cumulative selection produces an empirical approximation envelope




Figure 1. Total approximation error as a function of the common hidden-layer width.

over the width-constrained network classes without using the test data for model selection.

Figure 1 illustrates the joint effects of coordinate decay, truncation dimension, and network capacity. When $D = 1$, the total error is nearly insensitive to the network width, except for a modest initial decrease under the fastest decay setting. This indicates that the omitted coordinates constitute the dominant source of error, which cannot be mitigated by increasing network capacity. Once more coordinates are retained, increasing the width generally reduces the total error, although the improvement eventually levels off as the network approximation error becomes small relative to the remaining truncation and optimization errors. Across all values of D and W , faster coordinate decay consistently produces substantially smaller errors. These results support the theoretical conclusion that faster coordinate decay reduces the effective dimension and the network capacity required for accurate approximation, while slower decay leads to a more pronounced trade-off between coordinate truncation and finite-dimensional approximation.

5. Conclusion

This paper studies the fundamental representation–capacity trade-off that arises when deep neural networks are used to approximate scalar-valued functionals. Working with a fixed basis representation, we quantify the importance

of each coordinate through its score magnitude and the directional Fréchet sensitivity of the target functional. By combining coordinate truncation, anisotropic partitioning, local Taylor approximation, and explicit ReLU network constructions, we derive a non-asymptotic uniform approximation bound. Under mild conditions, a complementary lower bound matches the leading exponential term of the constructive upper bound, showing that the resulting rate is nearly optimal. Under generalized exponential coordinate decay regime, the approximation error decays at the stretched-exponential rate. This characterization provides two useful insights. The effective dimension increases with the network size, and unrestricted interactions among the retained coordinates prevent a polynomial approximation rate over the general functional class considered here.

This sub-polynomial approximation rate also has a direct implication for statistical estimation within the corresponding network framework, leading to a sub-polynomial estimation error after balancing the approximation and stochastic errors. Achieving a polynomial estimation rate therefore requires additional structural restrictions on the target functional. Identifying suitable restrictions and developing the corresponding estimation theory are important directions for future work.

References

- Bartlett, P. L., Harvey, N., Liaw, C., and Mehrabian, A. Nearly-tight vc-dimension and pseudodimension bounds for piecewise linear neural networks. *Journal of Machine Learning Research*, 20(63):1–17, 2019.
- Bölcskei, H., Grohs, P., Kutyniok, G., and Petersen, P. Optimal approximation with sparsely connected deep neural networks. *SIAM Journal on Mathematics of Data Science*, 1(1):8–45, 2019.
- Cai, T. T. and Hall, P. Prediction in functional linear regression. *The Annals of Statistics*, 34(5):2159–2179, 2006.
- Chen, D., Hall, P., and Müller, H.-G. Single and multiple index functional regression models with nonparametric link. *The Annals of Statistics*, 39(3):1720–1747, 2011. doi: 10.1214/11-AOS882.
- DeVore, R., Hanin, B., and Petrova, G. Neural network approximation. *Acta Numerica*, 30:327–444, 2021.
- Elbrächter, D., Perekhrenko, D., Grohs, P., and Bölskei, H. Deep neural network approximation theory. *IEEE Transactions on Information Theory*, 67(5):2581–2623, 2021.
- Gühning, I., Kutyniok, G., and Petersen, P. Error bounds for approximations with deep ReLU neural networks in $W^{s,p}$ norms. *Analysis and Applications*, 18(5):803–859, 2020.
- Hall, P. and Horowitz, J. L. Methodology and convergence rates for functional linear regression. *The Annals of Statistics*, 35(1):70–91, 2007.
- Lu, J., Shen, Z., Yang, H., and Zhang, S. Deep network approximation for smooth functions. *SIAM Journal on Mathematical Analysis*, 53(5):5465–5506, 2021.
- Müller, H.-G. and Yao, F. Functional additive models. *Journal of the American Statistical Association*, 103(484):1534–1544, 2008. doi: 10.1198/016214508000000751.
- Petersen, P. and Voigtlaender, F. Optimal approximation of piecewise smooth functions using deep relu neural networks. *Neural Networks*, 108:296–330, 2018.
- Rao, A. R. and Reimherr, M. Nonlinear functional modeling using neural networks. *Journal of Computational and Graphical Statistics*, 32(4):1248–1257, 2023.
- Rossi, F. and Conan-Guez, B. Functional multi-layer perceptron: A non-linear tool for functional data analysis. *Neural Networks*, 18(1):45–60, 2005.
- Rossi, F., Delannay, N., Conan-Guez, B., and Verleysen, M. Representation of functional data in neural networks. *Neurocomputing*, 64:183–210, 2005.
- Rügamer, D., Liew, B. X. W., Altai, Z., and Stöcker, A. A functional extension of semi-structured networks. In *Advances in Neural Information Processing Systems*, volume 37, 2024.
- Shen, Z., Yang, H., and Zhang, S. Nonlinear approximation via compositions. *Neural Networks*, 119:74–84, 2019. doi: 10.1016/j.neunet.2019.07.011.
- Shen, Z., Yang, H., and Zhang, S. Deep network approximation characterized by number of neurons. *Communications in Computational Physics*, 28(5):1768–1811, 2020. doi: 10.4208/cicp.OA-2020-0149.
- Shi, Z., Fan, J., Song, L., Zhou, D.-X., and Suykens, J. A. Nonlinear functional regression by functional deep neural network with kernel embedding. *Journal of Machine Learning Research*, 26(284):1–49, 2025.
- Siegel, J. W. Optimal approximation rates for deep ReLU neural networks on sobolev and besov spaces. *Journal of Machine Learning Research*, 24(357):1–52, 2023.
- Song, L., Liu, Y., Fan, J., and Zhou, D.-X. Approximation of smooth functionals using deep ReLU networks. *Neural Networks*, 166:424–436, 2023.
- Thind, B., Multani, K., and Cao, J. Deep learning with functional inputs. *Journal of Computational and Graphical Statistics*, 32(1):171–180, 2023.
- Yao, F. and Müller, H.-G. Functional quadratic regression. *Biometrika*, 97(1):49–64, 2010. doi: 10.1093/biomet/asp069.
- Yao, J., Mueller, J., and Wang, J.-L. Deep learning for functional data analysis with adaptive basis layers. In *International conference on machine learning*, pp. 11898–11908. PMLR, 2021.
- Yarotsky, D. Error bounds for approximations with deep relu networks. *Neural Networks*, 94:103–114, 2017. doi: 10.1016/j.neunet.2017.07.002.
- Yarotsky, D. Optimal approximation of continuous functions by very deep relu networks. *Proceedings of Machine Learning Research*, 75:639–649, 2018.
- Yarotsky, D. and Zhevnerchuk, A. The phase diagram of approximation rates for deep neural networks. In *Advances in Neural Information Processing Systems*, volume 33, 2020.

Appendix

Proof of Proposition 1. Note that since L is bounded linear, it is Fréchet differentiable everywhere with $\mathcal{D}L(X) = L$ and $\mathcal{D}^s L(X) = 0$ for all $s \geq 2$. We prove by induction on m . For $m = 1$, the chain rule yields $\mathcal{D}f(X) = \mathcal{D}g(LX) \circ \mathcal{D}L(X) = \mathcal{D}g(LX) \circ L$, hence $\mathcal{D}f(X)[h] = \mathcal{D}g(LX)[Lh]$, proving the result for $m = 1$.

Assume the result holds for $s \leq m-1$. Fix $h_1, \dots, h_{m-1} \in \mathcal{H}$ and define

$$G(u) := \mathcal{D}^{m-1}g(u)[Lh_1, \dots, Lh_{m-1}], \quad u \in \Omega.$$

Since f is m -times Fréchet differentiable, G is Fréchet differentiable and

$$\mathcal{D}G(u)[v] = \mathcal{D}^m g(u)[v, Lh_1, \dots, Lh_{m-1}], \quad v \in \Omega.$$

Using the definition of $\mathcal{D}^m f$ and the induction hypothesis,

$$\begin{aligned} \mathcal{D}^m f(X)[h_1, \dots, h_{m-1}, h_m] &= \lim_{t \rightarrow 0} \frac{1}{t} \left\{ \mathcal{D}^{m-1}f(X + th_m)[h_1, \dots, h_{m-1}] \right. \\ &\quad \left. - \mathcal{D}^{m-1}f(X)[h_1, \dots, h_{m-1}] \right\} \\ &= \lim_{t \rightarrow 0} \frac{1}{t} \left\{ G(L(X + th_m)) - G(LX) \right\}. \end{aligned}$$

Since L is linear, $L(X + th_m) = LX + tLh_m$, and the last limit equals

$$\begin{aligned} \lim_{t \rightarrow 0} \frac{G(LX + tLh_m) - G(LX)}{t} &= \mathcal{D}G(LX)[Lh_m] \\ &= \mathcal{D}^m g(LX) \\ &\quad [Lh_m, Lh_1, \dots, Lh_{m-1}]. \end{aligned}$$

By symmetry of $\mathcal{D}^m g(LX)$, we may permute the arguments, proving the result for m . This completes the induction. $\square$

Proof of Proposition 2. We first prove the derivative bound for Q_p . Since $\xi_j = \langle X, \nu_j \rangle$, we have

$$D\xi_j[\nu_d] = \langle \nu_d, \nu_j \rangle = \mathbf{1}\{j = d\}.$$

For a fixed monomial

$$\eta_{j_1, \dots, j_p} \prod_{\ell=1}^p \xi_{j_\ell},$$

taking q directional derivatives along $\nu_{d_1}, \dots, \nu_{d_q}$ amounts to differentiating q of the p scores. Therefore, each nonzero term after differentiation contains the coefficient $\eta_{j_1, \dots, j_p}$, q Kronecker factors that match q indices with $d_1, \dots, d_q$, and $p - q$ remaining score factors. Hence,

$$D^q Q_p(X)[\nu_{d_1}, \dots, \nu_{d_q}]$$

$$= \sum_{k_1, \dots, k_{p-q} \geq 1} \eta_{d_1, \dots, d_q, k_1, \dots, k_{p-q}}^* \xi_{k_1} \cdots \xi_{k_{p-q}},$$

where η^* denotes the coefficient after arranging the indices in nondecreasing order. By the product-type coefficient decay condition,

$$|\eta_{d_1, \dots, d_q, k_1, \dots, k_{p-q}}^*| \lesssim \left(\prod_{a=1}^q w_{d_a} \right) \left(\prod_{b=1}^{p-q} w_{k_b} \right).$$

Using $|\xi_k| \leq s_k$, we obtain

$$\begin{aligned} |D^q Q_p(X)[\nu_{d_1}, \dots, \nu_{d_q}]| &\lesssim \sum_{k_1, \dots, k_{p-q} \geq 1} \left(\prod_{a=1}^q w_{d_a} \right) \left(\prod_{b=1}^{p-q} w_{k_b} s_{k_b} \right) \\ &= \left(\prod_{a=1}^q w_{d_a} \right) \left(\sum_{k \geq 1} w_k s_k \right)^{p-q}. \end{aligned}$$

Since $\sum_{k \geq 1} w_k s_k < \infty$, this gives

$$|D^q Q_p(X)[\nu_{d_1}, \dots, \nu_{d_q}]| \lesssim \prod_{a=1}^q w_{d_a}.$$

It remains to pass from Q_p to $f_0 = G \circ Q_p$. By the chain rule, for $1 \leq s \leq m$,

$$\begin{aligned} D^s f_0(X)[\nu_{d_1}, \dots, \nu_{d_s}] &= \sum_{\pi \in \mathcal{P}_s} G^{(|\pi|)}(Q_p(X)) \prod_{B \in \pi} D^{|B|} Q_p(X)[\nu_{d_i} : i \in B], \end{aligned}$$

where $\mathcal{P}_s$ denotes the set of all partitions of $\{1, \dots, s\}$. Since Q_p is a polynomial of order p , $D^q Q_p(X) = 0$ for $q > p$. Hence only blocks B with $|B| \leq p$ contribute. For each such block, the first part of the proof gives

$$|D^{|B|} Q_p(X)[\nu_{d_i} : i \in B]| \lesssim \prod_{i \in B} w_{d_i}.$$

Since the derivatives of G up to order m are bounded, each product in the partition expansion is bounded by $\prod_{\ell=1}^s w_{d_\ell}$ up to some constant. The number of partitions depends only on $s \leq m$, so summing over $\pi \in \mathcal{P}_s$ yields

$$|D^s f_0(X)[\nu_{d_1}, \dots, \nu_{d_s}]| \lesssim \prod_{\ell=1}^s w_{d_\ell}.$$

This proves the proposition. $\square$

For $D \geq D_0$, define the finite-dimensional trifling region by $\mathcal{R}_D := \mathcal{R}(\Omega_{D_0}, \mathbf{L}, \boldsymbol{\delta}) \times \prod_{d=D_0+1}^D [-s_d, s_d] \subseteq \Omega_D$, where

$$\mathcal{R}(\Omega_{D_0}, \mathbf{L}, \boldsymbol{\delta}) = \bigcup_{d=1}^{D_0} \left\{ \mathbf{x}^{(D_0)} \in \Omega_{D_0} : \xi_d \in \mathcal{I}_d \right\},$$

$$\mathcal{I}_d = \bigcup_{i=1}^{L_d-1} \left(-s_d + \frac{2s_d i}{L_d} - \delta_d, -s_d + \frac{2s_d i}{L_d} \right].$$

For a continuous function g on Ω_D , define its directional modulus of continuity by $\omega_{g,d}(r) := \sup\{|g(\mathbf{x} + te_d) - g(\mathbf{x})| : \mathbf{x}, \mathbf{x} + te_d \in \Omega_D, |t| \leq r\}$.

Proof of Theorem 1. For $X \in \mathcal{H}$ with score vector $\mathbf{x} = (\xi_1, \xi_2, \dots) \in \Omega$, let $X_0 = \sum_{d=1}^{D_0} \xi_{0,d} \nu_d$. Define

$$h_D := P_D X - X_0 = \sum_{d=1}^D \xi_{h,d} \nu_d,$$

where $\xi_{h,d} = \xi_d - \xi_{0,d}$ for $d \leq D_0$ and $\xi_{h,d} = \xi_d$ for $D_0 < d \leq D$. Since $P_D X_0 = X_0$, the Taylor expansion of $f_0^{(D)}$ at X_0 is

$$\begin{aligned} f_0^{(D)}(X) &= f_0(X_0) + \sum_{s=1}^{m-1} \frac{1}{s!} \sum_{d_1, \dots, d_s=1}^D \mathcal{D}^s f_0(X_0) [\nu_{d_1}, \dots, \nu_{d_s}] \\ &\quad \times \prod_{j=1}^s \xi_{h,d_j} + R_{m,D}(X, X_0), \end{aligned} \quad (\text{A.1})$$

where

$$\begin{aligned} R_{m,D}(X, X_0) &= \frac{1}{(m-1)!} \int_0^1 (1-\tau)^{m-1} \\ &\quad \times \mathcal{D}^m f_0(X_0 + \tau h_D) [h_D, \dots, h_D] d\tau. \end{aligned}$$

Write $\Delta_D(X) := f_0(X) - f_0^{(D)}(X)$, then the mean value theorem gives

$$|\Delta_D(X)| \leq C_0 \sum_{d>D} w_d |\xi_d| \leq C_0 H(D). \quad (\text{A.2})$$

We now construct a network approximation of the Taylor polynomial in (A.1). Given $L > 0$ and a positive sequence $\{\ell_d\}$, define $D_0 := \max\{d : L\ell_d s_d \geq 1\}$, $L_d := \lfloor L\ell_d s_d \rfloor$, $d = 1, \dots, D_0$. For $d \leq D_0$, partition $[-s_d, s_d]$ into L_d intervals and choose $\delta_d \in (0, 2s_d/(3L_d)]$. Denote the resulting cells in $\Omega'_{D_0} := \Omega_{D_0} \setminus \mathcal{R}(\Omega_{D_0}, \mathbf{L}, \boldsymbol{\delta})$ by $P_{\boldsymbol{\theta}}^{(D_0)}$, where $\theta_d \in \{0, \dots, L_d - 1\}$, and define

$$P_{\boldsymbol{\theta}}^{(D)} := P_{\boldsymbol{\theta}}^{(D_0)} \times \prod_{d=D_0+1}^D [-s_d, s_d],$$

$$X_{\boldsymbol{\theta}}(t) := \sum_{d=1}^{D_0} \left(-s_d + \frac{2s_d \theta_d}{L_d} \right) \nu_d(t).$$

For $\mathbf{x}^{(D)} \in P_{\boldsymbol{\theta}}^{(D)}$, define

$$\xi_{h_{\boldsymbol{\theta}},d} := \begin{cases} \xi_d + s_d - \frac{2s_d \theta_d}{L_d}, & 1 \leq d \leq D_0, \\ \xi_d, & D_0 < d \leq D. \end{cases}$$

The remainder of the Taylor expansion satisfies

$$\begin{aligned} |R_{m,D}(X, X_{\boldsymbol{\theta}})| &\leq \frac{C_0}{m!} \sum_{d_1, \dots, d_m=1}^D \prod_{j=1}^m w_{d_j} |\xi_{h_{\boldsymbol{\theta}},d_j}| \\ &= \frac{C_0}{m!} \left(\sum_{d=1}^D w_d |\xi_{h_{\boldsymbol{\theta}},d}| \right)^m \\ &= \frac{C_0}{m!} \left\{ \sum_{d=1}^{D_0} w_d \left| \xi_d + s_d - \frac{2s_d \theta_d}{L_d} \right| \right. \\ &\quad \left. + \sum_{d=D_0+1}^D w_d |\xi_d| \right\}^m. \end{aligned} \quad (\text{A.3})$$

For $\mathbf{x}^{(D_0)} \in P_{\boldsymbol{\theta}}^{(D_0)}$,

$$\left| \xi_d + s_d - \frac{2s_d \theta_d}{L_d} \right| \leq \frac{2s_d}{L_d}, \quad d = 1, \dots, D_0.$$

Since $L_d = \lfloor L\ell_d s_d \rfloor$ and $L\ell_d s_d \geq 1$ for $d \leq D_0$, we have $2s_d/L_d \lesssim L^{-1}\ell_d^{-1}$. Consequently, for $\mathbf{x}^{(D)} \in P_{\boldsymbol{\theta}}^{(D)}$,

$$|\xi_{h_{\boldsymbol{\theta}},d}| \lesssim L^{-1}\ell_d^{-1}, \quad d \leq D_0.$$

In addition, since $|\xi_d| \leq s_d$, it follows that

$$\begin{aligned} |R_{m,D}(X, X_{\boldsymbol{\theta}})| &\lesssim \left\{ L^{-1} \sum_{d=1}^{D_0} w_d \ell_d^{-1} + \sum_{d=D_0+1}^D w_d s_d \right\}^m \\ &= \left\{ L^{-1} \sum_{d=1}^{D_0} w_d \ell_d^{-1} + H(D_0) \right\}^m. \end{aligned} \quad (\text{A.4})$$

For $s = 1, \dots, m-1$ and $\mathbf{d} = (d_1, \dots, d_s) \in \{1, \dots, D\}^s$, define the piecewise-constant Taylor coefficient map

$$\mathcal{Q}_{\mathbf{d}}^{(s)}(\mathbf{x}^{(D)}) := \sum_{\boldsymbol{\theta}} \mathcal{D}^s f_0(X_{\boldsymbol{\theta}}) [\nu_{d_1}, \dots, \nu_{d_s}] \mathbf{1}_{P_{\boldsymbol{\theta}}^{(D_0)}}(\mathbf{x}^{(D_0)}).$$

Also define

$$\mathcal{Q}^{(0)}(\mathbf{x}^{(D)}) := \sum_{\boldsymbol{\theta}} f_0(X_{\boldsymbol{\theta}}) \mathbf{1}_{P_{\boldsymbol{\theta}}^{(D_0)}}(\mathbf{x}^{(D_0)}).$$

For each fixed $\mathbf{d}$, these maps have $\prod_{d=1}^{D_0} L_d$ distinct values. The Taylor polynomial to be approximated is

$$\begin{aligned} A_{\boldsymbol{\theta}}^{(D)}(\mathbf{x}^{(D)}) &:= f_0(X_{\boldsymbol{\theta}}) + \sum_{s=1}^{m-1} \frac{1}{s!} \sum_{d_1, \dots, d_s=1}^D \mathcal{D}^s f_0(X_{\boldsymbol{\theta}}) \\ &\quad [\nu_{d_1}, \dots, \nu_{d_s}] \times \prod_{j=1}^s \xi_{h_{\boldsymbol{\theta}},d_j}. \end{aligned}$$

The network construction steps are as follows:

Step 1 (Discretization). By Lemma 1, for $d = 1, \dots, D_0$, there exists a ReLU network ψ_d with width $6W_{\psi,d}$ and depth $3M_{\psi}$ such that $W_{\psi,d}^2 M_{\psi}^2 = L_d$ and

$$\psi_d(\xi_d) = -s_d + \frac{2s_d i}{L_d}$$

whenever

$$\begin{aligned} \xi_d \in & \left[-s_d + \frac{2s_d i}{L_d}, \right. \\ & \left. -s_d + \frac{2s_d(i+1)}{L_d} - \delta_d \mathbf{1}_{\{i < L_d - 1\}} \right], \\ & i = 0, \dots, L_d - 1. \end{aligned}$$

Define $\tilde{\psi}_0(\mathbf{x}^{(D_0)}) := (\psi_1(\xi_1), \dots, \psi_{D_0}(\xi_{D_0}))$. Then, for $\mathbf{x}^{(D_0)} \in P_{\theta}^{(D_0)}$,

$$\tilde{\psi}_0(\mathbf{x}^{(D_0)}) = \left(-s_d + \frac{2s_d \theta_d}{L_d} \right)_{d=1}^{D_0}.$$

The network $\tilde{\psi}_0$ has width $6 \sum_{d=1}^{D_0} W_{\psi,d}$ and depth $3M_{\psi}$. The index set

$$\{0, \dots, L_1 - 1\} \times \dots \times \{0, \dots, L_{D_0} - 1\}$$

is in one-to-one correspondence with $\{0, \dots, \prod_{d=1}^{D_0} L_d - 1\}$ through

$$i_{\theta} := \sum_{d=1}^{D_0} \theta_d \prod_{k=1}^{d-1} L_k, \quad \theta_d \in \{0, 1, \dots, L_d - 1\}.$$

Accordingly, define

$$\psi_{\text{enc}}(\mathbf{x}^{(D_0)}) := \sum_{d=1}^{D_0} \frac{L_d}{2s_d} \{ \psi_d(\xi_d) + s_d \} \prod_{k=1}^{d-1} L_k.$$

For every $\mathbf{x}^{(D_0)} \in P_{\theta}^{(D_0)}$, we have $\psi_{\text{enc}}(\mathbf{x}^{(D_0)}) = i_{\theta}$.

Step 2 (Approximation of the Taylor coefficients). Let

$$\begin{aligned} c_{s,d} &:= C_0 \prod_{j=1}^s w_{d_j}, \\ e_{s,d} &:= c_{s,d} \prod_{j=1}^s s_{d_j} = C_0 \prod_{j=1}^s w_{d_j} s_{d_j}. \end{aligned}$$

Assumption 2 gives $|\mathcal{D}^s f_0(X_{\theta})[\nu_{d_1}, \dots, \nu_{d_s}]| \leq c_{s,d}$. The number of cells satisfies

$$\prod_{d=1}^{D_0} L_d = L^{D_0} \prod_{d=1}^{D_0} \ell_d s_d.$$

Define

$$\Psi(D_0) := - \sum_{d=1}^{D_0} \log(\ell_d s_d),$$

so that we may equivalently write $\prod_{d=1}^{D_0} L_d = L^{D_0} e^{-\Psi(D_0)}$. For each $\mathbf{d} = (d_1, \dots, d_s) \in \{1, \dots, D\}^s$, Lemma 2, composed with the radix encoder ψ_{enc} , gives a ReLU network $\Phi_{\mathbf{d},s}$ satisfying

$$\left| \Phi_{\mathbf{d},s}(\mathbf{x}^{(D)}) - \mathcal{Q}_{\mathbf{d}}^{(s)}(\mathbf{x}^{(D)}) \right| \lesssim c_{s,\mathbf{d}} \left\{ L^{D_0} e^{-\Psi(D_0)} \right\}^{-m}. \quad (\text{A.5})$$

Similarly, there exists a network Φ_0 satisfying

$$\left| \Phi_0(\mathbf{x}^{(D)}) - \mathcal{Q}^{(0)}(\mathbf{x}^{(D)}) \right| \lesssim \left\{ L^{D_0} e^{-\Psi(D_0)} \right\}^{-m}. \quad (\text{A.6})$$

Each $\Phi_{\mathbf{d},s}$ has width $16rW_{\phi} \lceil \log_2(8W_{\phi}) \rceil$ and depth $5(M_{\phi} + 2) \lceil \log_2(4M_{\phi}) \rceil + 3M_{\psi}$, where $W_{\phi}^2 M_{\phi}^2 = L^{D_0} e^{-\Psi(D_0)}$. Since $\mathbf{d} \in \{1, \dots, D\}^s$, the parallel network for all order- s coefficient maps has width $16rD^s W_{\phi} \lceil \log_2(8W_{\phi}) \rceil$. Notice that D^s counts the number of coefficient maps, whereas each coefficient map is fitted on only $\prod_{d=1}^{D_0} L_d$ locations.

Step 3 (Monomial approximation). For $\mathbf{d} = (d_1, \dots, d_s) \in \{1, \dots, D\}^s$, Lemma 4, applied on the corresponding coordinate ranges, gives a ReLU network $J_{\mathbf{d},s}$ with width $9W_J + s - 1$ and depth $7s^2 M_J$ such that

$$\left| J_{\mathbf{d},s}(\mathbf{x}^{(D)}) - \prod_{j=1}^s \xi_{h_{\theta}, d_j} \right| \leq 2^s \left(\prod_{j=1}^s s_{d_j} \right) 9sW_J^{-7sM_J}. \quad (\text{A.7})$$

The coordinate-dependent factor follows by applying the multiplication network on the corresponding bounded intervals; it does not require a rescaling of the input domain. By Lemma 3, the multiplication network $\Phi_{\times}$ can be chosen so that, on the relevant bounded rectangles,

$$|\Phi_{\times}(u, v) - uv| \lesssim e_{s,\mathbf{d}} W_{\times}^{-M_{\times}}. \quad (\text{A.8})$$

Combining the preceding networks, define

$$\begin{aligned} \hat{A}_{\theta}^{(D)}(\mathbf{x}^{(D)}) &= \Phi_0(\mathbf{x}^{(D)}) \\ &+ \sum_{s=1}^{m-1} \sum_{d_1, \dots, d_s=1}^D \Phi_{\times} \left(\frac{\Phi_{\mathbf{d},s}(\mathbf{x}^{(D)})}{s!}, J_{\mathbf{d},s}(\mathbf{x}^{(D)}) \right). \end{aligned}$$

Using (A.5)–(A.8) and the triangle inequality gives

$$\begin{aligned} \left| \hat{A}_{\theta}^{(D)}(\mathbf{x}^{(D)}) - A_{\theta}^{(D)}(\mathbf{x}^{(D)}) \right| &\lesssim \sum_{s=0}^{m-1} \frac{1}{s!} \sum_{d_1, \dots, d_s=1}^D e_{s,\mathbf{d}} \{ (W_{\phi}^2 M_{\phi}^2)^{-m} \\ &\quad + 9sW_J^{-7sM_J} + 6W_{\times}^{-M_{\times}} \}. \end{aligned}$$

Indeed,

$$\sum_{d_1, \dots, d_s=1}^D \prod_{j=1}^s w_{d_j} s_{d_j} = \left(\sum_{d=1}^D w_d s_d \right)^s$$

$$\leq \left(\sum_{d=1}^{\infty} w_d s_d \right)^s < \infty.$$

Since $\sum_{d=1}^{\infty} w_d s_d < \infty$, we obtain

$$\begin{aligned} & \left| \hat{A}_{\theta}^{(D)}(\mathbf{x}^{(D)}) - A_{\theta}^{(D)}(\mathbf{x}^{(D)}) \right| \\ & \lesssim (W_{\phi}^2 M_{\phi}^2)^{-m} + \max_{1 \leq s \leq m-1} 9s W_J^{-7s M_J} + 6W_{\times}^{-M_{\times}}. \end{aligned} \quad (\text{A.9})$$

To attain an approximation error ε , the coefficient-fitting term requires $W_{\phi} M_{\phi} \gtrsim \varepsilon^{-1/(2m)}$, whereas the monomial and multiplication networks require only $M_J \log W_J \gtrsim \log(1/\varepsilon)$, $M_{\times} \log W_{\times} \gtrsim \log(1/\varepsilon)$. Therefore, the coefficient-fitting networks determine the leading complexity. Since the number of coefficient maps is bounded by $\sum_{s=0}^{m-1} D^s = \mathcal{O}(D^{m-1})$, the combined network can be chosen, up to the logarithmic factors appearing above, with

$$\begin{aligned} W &= \mathcal{O}(D^m W_{\phi}), & M &= \mathcal{O}(M_{\phi}), \\ SM &= \mathcal{O}(D^m W_{\phi}^2 M_{\phi}^2). \end{aligned}$$

In particular, with $\mathcal{A}^2 := W_{\phi}^2 M_{\phi}^2$, the total network complexity is $SM = \mathcal{O}(D^m \mathcal{A}^2)$. Thus

$$\left| \hat{A}_{\theta}^{(D)}(\mathbf{x}^{(D)}) - A_{\theta}^{(D)}(\mathbf{x}^{(D)}) \right| \lesssim (W_{\phi}^2 M_{\phi}^2)^{-m}.$$

Consequently, combining (A.4) and (A.9) gives, uniformly over $\mathbf{x}^{(D)} \in \Omega_D \setminus \mathcal{R}_D$,

$$\begin{aligned} & \left| \hat{A}_{\theta}^{(D)}(\mathbf{x}^{(D)}) - f_0^{(D)}(X) \right| \\ & \lesssim \left\{ L^{-1} \sum_{d=1}^{D_0} w_d \ell_d^{-1} + H(D_0) \right\}^m \end{aligned} \quad (\text{A.10})$$

After balancing $L^{-1} \sum_{d=1}^{D_0} w_d \ell_d^{-1} = H(D_0)$ the Taylor remainder is bounded by a constant multiple of $H(D_0)^m$, so the network approximation error is of smaller order. Therefore, the leading scale in (A.10) is $H(D_0)^m$, determined by the Taylor remainder.

We now optimize the grid allocation. We balance the retained local discretization error with $H(D_0)$ by choosing

$$L = \frac{\sum_{d=1}^{D_0} w_d \ell_d^{-1}}{H(D_0)}. \quad (\text{A.11})$$

The coefficient-fitting complexity for each coefficient map satisfies $\mathcal{A}^2 = W_{\phi}^2 M_{\phi}^2 = L^{D_0} \prod_{d=1}^{D_0} \ell_d s_d$. Substituting (A.11) gives

$$\mathcal{A}^2 = H(D_0)^{-D_0} \left(\sum_{d=1}^{D_0} w_d \ell_d^{-1} \right)^{D_0} \times \prod_{d=1}^{D_0} \ell_d s_d. \quad (\text{A.12})$$

For fixed D_0 , minimizing the required point-fitting budget (A.12) is equivalent to minimizing

$$\left(\sum_{d=1}^{D_0} w_d \ell_d^{-1} \right)^{D_0} \prod_{d=1}^{D_0} \ell_d$$

over $\{\ell_d : d \geq 1\}$. Let $a_d := w_d \ell_d^{-1}$. Then

$$\left(\sum_{d=1}^{D_0} w_d \ell_d^{-1} \right)^{D_0} \prod_{d=1}^{D_0} \ell_d = \frac{\left(\sum_{d=1}^{D_0} a_d \right)^{D_0}}{\prod_{d=1}^{D_0} a_d} \prod_{d=1}^{D_0} w_d.$$

By the arithmetic–geometric mean inequality,

$$\frac{\left(\sum_{d=1}^{D_0} a_d \right)^{D_0}}{\prod_{d=1}^{D_0} a_d} \geq D_0^{D_0},$$

with equality if and only if $a_1 = \dots = a_{D_0}$. Therefore, up to a common scaling absorbed into L , the optimal allocation is $\ell_d = w_d$. Consequently,

$$\mathcal{A}^2 = L^{D_0} \prod_{d=1}^{D_0} w_d s_d = L^{D_0} e^{-\Psi(D_0)},$$

The balancing relation (A.11) becomes $L^{-1} D_0 = H(D_0)$. since $\mathcal{A}^2 = \prod_{d=1}^{D_0} L_d \geq L_1 = L$ and the balancing relation gives $H(D_0) = D_0/L$, we have $\mathcal{A}^{-2m} \leq L^{-m} = D_0^{-m} H(D_0)^m \leq H(D_0)^m$. Therefore, the network approximation error is no larger than the Taylor remainder.

It follows that

$$\mathcal{A}^2 = \left(\frac{D_0}{H(D_0)} \right)^{D_0} e^{-\Psi(D_0)},$$

and hence D_0 is determined by

$$\log \mathcal{A}^2 = D_0 \log \left(\frac{D_0}{H(D_0)} \right) - \Psi(D_0).$$

Thus,

$$L = \exp \left[\frac{1}{D_0} \{ \log \mathcal{A}^2 + \Psi(D_0) \} \right].$$

Substituting these bounds into (A.10) yields

$$\begin{aligned} & \|\tilde{f} - f_0^{(D)}\|_{L^\infty(\Omega_D \setminus \mathcal{R}_D)} \\ & \lesssim H(D_0)^m + \mathcal{A}^{-2m} \lesssim H(D_0)^m \\ & = D_0^m \exp \left[-\frac{m}{D_0} \{ \log \mathcal{A}^2 + \Psi(D_0) \} \right]. \end{aligned}$$

For each fixed s and $\mathbf{d} = (d_1, \dots, d_s)$, the coefficient-fitting network $\Phi_{\mathbf{d},s}$ can be chosen, up to logarithmic factors, with size $\mathcal{O}(W_{\phi}^2 M_{\phi})$, width $\mathcal{O}(W_{\phi})$, and depth $\mathcal{O}(M_{\phi})$. The

number of coefficient maps appearing in the Taylor polynomial is

$$\sum_{s=0}^{m-1} D^s \leq mD^{m-1} = \mathcal{O}(D^{m-1}).$$

Since these coefficient-fitting subnetworks are constructed in parallel, their sizes and widths are additive, whereas their depths are given by the maximum depth. Therefore, the combined coefficient-fitting network satisfies

$$\begin{aligned} S_{\text{coef}} &\lesssim D^{m-1} W_\phi^2 M_\phi, \\ W_{\text{coef}} &\lesssim D^{m-1} W_\phi, \\ M_{\text{coef}} &\lesssim M_\phi. \end{aligned}$$

The discretization network is shared by all coefficient maps, while the monomial and multiplication subnetworks require only logarithmic complexity in the target accuracy. Compared with the coefficient-fitting subnetworks, the complexities of the shared discretization network and the monomial and multiplication subnetworks are negligible and therefore do not affect the leading orders of the network size, width, and depth. Before removing the trifling region, the resulting network therefore satisfies

$$\tilde{S} \lesssim D^{m-1} W_\phi^2 M_\phi, \quad \tilde{W} \lesssim D^{m-1} W_\phi, \quad \tilde{M} \lesssim M_\phi.$$

It remains to extend the preceding approximation from $\Omega_D \setminus \mathcal{R}_D$ to all of Ω_D . Because the construction is valid for every positive $\delta_d \leq 2s_d/(3L_d)$, choose

$$\delta_d \leq \min \left\{ \frac{2s_d}{3L_d}, \frac{H(D_0)^m}{C_0 D_0 w_d} \right\}, \quad d = 1, \dots, D_0.$$

Then $C_0 \sum_{d=1}^{D_0} w_d \delta_d \leq H(D_0)^m$. Applying Lemma 5 with $g = f_0^{(D)}$ gives a ReLU network $f^\#$ satisfying

$$\begin{aligned} \|f^\# - f_0^{(D)}\|_{L^\infty(\Omega_D)} &\lesssim H(D_0)^m + C_0 \sum_{d=1}^{D_0} w_d \delta_d \\ &\lesssim H(D_0)^m. \end{aligned}$$

The projection error satisfies $H(D) \asymp H(D_0)^m$. The same lemma gives $S \lesssim 3^{D_0} \{D^{m-1} W_\phi^2 M_\phi + 1\}$, $W \lesssim 3^{D_0} \{D^{m-1} W_\phi + 1\}$, and $M \lesssim M_\phi + 2D_0$. This proves the asserted global uniform approximation bound. $\square$

Proof of Theorem 2. For each $D \geq 1$, consider the finite-dimensional slice $\Omega_D = \prod_{d=1}^D [-s_d, s_d]$, embedded into Ω by setting all coordinates $d > D$ equal to zero. Any family constructed on this slice can be extended to the full sequence domain by ignoring the tail coordinates. Hence the finite-dimensional packing argument can be carried out on Ω_D . We partition the d -th coordinate interval into cells of length of order $L^{-1} \ell_d^{-1}$, where $L > 0$ is a global resolution

parameter and ℓ_d is a coordinate-specific grid allocation factor, with $\ell_1 = 1$. Thus the number of cells in the d -th coordinate is $N_d = \lfloor L s_d \ell_d \rfloor$, provided $L s_d \ell_d \geq 2$. Let x_β denote the center of a grid cell, where $\beta = (\beta_1, \dots, \beta_D)$ and $\beta_d \in \{0, \dots, N_d - 1\}$. Choose a fixed bump function $\varphi \in C^m(\mathbb{R}^D)$ such that $\varphi(0) = 1$, φ is supported in a sufficiently small ball, and

$$\max_{\|\alpha\|_1 \leq m} \|\partial^\alpha \varphi\|_\infty \leq 1.$$

Define the anisotropic bump

$$\varphi_\beta(x) = B_L \varphi(L \operatorname{diag}(\ell_1, \dots, \ell_D)(x - x_\beta)),$$

where $\psi(0) = 1$. For any multi-index α , the chain rule gives

$$|\partial^\alpha \varphi_\beta(x)| \leq B_L L^{|\alpha|_1} \prod_{d=1}^D \ell_d^{\alpha_d}.$$

To ensure $\varphi_\beta \in \mathcal{C}^m(\Omega_D, r)$, it is enough that

$$B_L L^{|\alpha|_1} \prod_{d=1}^D \ell_d^{\alpha_d} \leq C_0 \prod_{d=1}^D w_d^{\alpha_d}.$$

Equivalently,

$$B_L \leq C_0 L^{-|\alpha|_1} \prod_{d=1}^D (w_d \ell_d^{-1})^{\alpha_d}.$$

We now determine the optimal choice of the grid allocation ℓ_d . The derivative constraint requires

$$B_L \leq C_0 \min_{1 \leq \|\alpha\|_1 \leq m} L^{-|\alpha|_1} \prod_{d=1}^D (w_d \ell_d^{-1})^{\alpha_d}.$$

At the same time, the grid cardinality satisfies

$$\prod_{d=1}^D N_d = L^D \prod_{d=1}^D s_d \ell_d,$$

so larger values of ℓ_d lead to a larger shattering set. Hence the optimal construction takes ℓ_d as large as permitted by the derivative constraint.

We consider two regimes.

Case 1: $\ell_d/w_d \rightarrow 0$. In this case, $w_d \ell_d^{-1}$ is increasing across d . The minimum in the derivative constraint is attained by placing all m derivatives on the first coordinate. Hence $B_L = C_0 L^{-m}$. Taking the critical scaling $B_L = 2\varepsilon$ gives $L = C_0^{1/m} (2\varepsilon)^{-1/m}$. The grid cardinality is

$$\prod_{d=1}^D N_d = L^D \prod_{d=1}^D s_d \ell_d.$$

The largest possible entropy is achieved at the boundary value $\ell_d = w_d$.

Case 2: $\ell_d/w_d \rightarrow \infty$. In this case, $w_d \ell_d^{-1}$ is decreasing across d . The derivative constraint is now most restrictive at the largest coordinate D . Hence $B_L = C_0 L^{-m} (w_d \ell_D^{-1})^m$. Taking $B_L = 2\varepsilon$ gives $L = C_0^{1/m} (2\varepsilon)^{-1/m} w_d \ell_D^{-1}$. Substituting this into the grid cardinality,

$$\begin{aligned} \prod_{d=1}^D N_d &= L^D \prod_{d=1}^D s_d \ell_d \\ &= C_0^{D/m} (2\varepsilon)^{-D/m} w_d^D \ell_D^{-D} \times \prod_{d=1}^D s_d \ell_d. \end{aligned}$$

Now fix all coordinates except ℓ_D . Then the above expression becomes

$$C_0^{D/m} (2\varepsilon)^{-D/m} w_d^D \left(\prod_{d=1}^{D-1} s_d \ell_d \right) s_D \ell_D^{-(D-1)}.$$

Since $D \geq 2$, the factor $\ell_D^{-(D-1)}$ is decreasing in ℓ_D . Therefore the grid cardinality is maximized by taking the smallest admissible value of ℓ_D . On the other hand, the present regime assumes $w_d \ell_D^{-1} \leq 1$, whose boundary is $\ell_D = w_d$. Hence the smallest admissible choice is exactly the boundary value $\ell_D = w_d$. Repeating the same argument coordinatewise yields $\ell_d = w_d$, $d = 1, \dots, D$. Combining the two cases, the optimal grid allocation is $\ell_d = w_d$, and the derivative constraint reduces to $B_L \leq C_0 L^{-m}$. Under the envelope representation $w_d s_d = \exp\{-\psi(d)\}$, the grid cardinality becomes

$$\begin{aligned} \prod_{d=1}^D N_d &= L^D \prod_{d=1}^D w_d s_d = L^D \exp\{-\Psi(D)\}, \\ \Psi(D) &= \sum_{d=1}^D \psi(d). \end{aligned}$$

Taking $B_L = 2\varepsilon$ and maximizing grid cardinality give $L = C_0^{1/m} (2\varepsilon)^{-1/m}$.

Define

$$\mathcal{B} := \left\{ \chi : \prod_{d=1}^D \{0, 1, \dots, L_d - 1\} \rightarrow \{-1, 1\} \right\}.$$

For each sign pattern $\chi \in \mathcal{B}$, define

$$f_\chi(\mathbf{x}) := \sum_{\beta} \chi(\beta) \psi_\beta(\mathbf{x}), \quad \mathbf{x} \in \Omega_D.$$

Since the supports of the bumps are disjoint, each f_χ satisfies the anisotropic derivative bounds on Ω_D . By the uniform approximation property on the infinite-dimensional

class, for every $\chi \in \mathcal{B}$, there exists $\phi_\chi \in \mathcal{F}$ such that $\|\phi_\chi - F_\chi\|_{L^\infty(\Omega)} \leq \varepsilon$. At the embedded center $\iota_D(\mathbf{x}_\beta)$, we have $|F_\chi(\iota_D(\mathbf{x}_\beta))| = |f_\chi(\mathbf{x}_\beta)| = B_L$. If $B_L = 2\varepsilon$, then

$$|\phi_\chi(\iota_D(\mathbf{x}_\beta)) - F_\chi(\iota_D(\mathbf{x}_\beta))| \leq \varepsilon < |F_\chi(\iota_D(\mathbf{x}_\beta))|.$$

Therefore, $\text{sign}\{\phi_\chi(\iota_D(\mathbf{x}_\beta))\} = \text{sign}\{F_\chi(\iota_D(\mathbf{x}_\beta))\} = \chi(\beta)$. Thus $\{\phi_\chi : \chi \in \mathcal{B}\} \subseteq \mathcal{F}$ shatters the embedded grid centers

$$\left\{ \mathbf{x}_\beta : \beta \in \prod_{d=1}^D \{0, 1, \dots, N_d - 1\} \right\}.$$

Consequently,

$$\begin{aligned} V = \text{PDim}(\mathcal{F}) &\geq \prod_{d=1}^D N_d \\ &= C_0^{D/m} (2\varepsilon)^{-D/m} \exp\{-\Psi(D)\}. \end{aligned}$$

Thus, for every admissible D satisfying $(\frac{C_0}{2\varepsilon})^{1/m} w_D s_D \geq 1$,

$$\varepsilon \geq \frac{C_0}{2} \exp\left[-\frac{m}{D} \{\log V + \Psi(D)\}\right].$$

Combining the fact that

$$\begin{aligned} \sup_{f \in \mathcal{C}^m(\Omega, \mathbf{w})} \inf_{\phi \in \mathcal{F}} \|\phi - f\|_{L^\infty(\Omega)} \\ \geq \sup_{f \in \mathcal{C}^m(\Omega_D, \mathbf{w})} \inf_{\phi \in \mathcal{F}} \|\phi - f\|_{L^\infty(\Omega_D)} \end{aligned}$$

completes the proof. $\square$

Proof of Theorem 3. Put $a_d = w_d s_d$ and $B(d) = d \log(2/a_d) - \Psi(d)$, then $D_0 = \max\{d : B(d) \leq t\}$. By the assumed uniform decay equivalence, there is a constant C_e , independent of d , ρ , and t , such that

$$|\log(a_d^{-1}) - c_a d^\rho| \leq C_e. \quad (\text{A.13})$$

For every $d \geq 1$ and every $\rho > 0$, monotonicity of x^ρ and integral comparison give

$$\frac{d^{\rho+1}}{\rho+1} \leq \sum_{k=1}^d k^\rho \leq \frac{d^{\rho+1}}{\rho+1} + d^\rho.$$

Consequently,

$$\Psi(d) = \frac{c_a}{\rho+1} d^{\rho+1} + R_\Psi(d), \quad |R_\Psi(d)| \leq C(c_a d^\rho + d), \quad (\text{A.14})$$

where C is independent of d , t , and $\mathcal{A}$. Substituting (A.13) and (A.14) into the definition of $B(d)$ gives

$$B(d) = \frac{\rho c_a}{\rho+1} d^{\rho+1} + R_B(d),$$

$$|R_B(d)| \leq C(c_a d^\rho + d). \quad (\text{A.15})$$

By the definition of D_0 in Theorem 1,

$$B(D_0) \leq t < B(D_0 + 1). \quad (\text{A.16})$$

Combining (A.15) and (A.16) yields the uniform budget-dimension bracket

$$\begin{aligned} t &\geq \frac{\rho c_a}{\rho + 1} D_0^{\rho+1} - C(c_a D_0^\rho + D_0) \\ t &< \frac{\rho c_a}{\rho + 1} (D_0 + 1)^{\rho+1} + C\{c_a (D_0 + 1)^\rho + D_0 + 1\}. \end{aligned} \quad (\text{A.17})$$

Set $G_{t,\rho} = D_0^{-1}\{t + \Psi(D_0)\} - \log D_0$. By Theorem 1,

$$\begin{aligned} \inf_{f \in \mathcal{F}} \|f - f_0\|_{L^\infty} &\lesssim \exp(-m G_{t,\rho}) \\ &\lesssim \exp[-m\{G_{t,\rho} - \Psi(1)\}]. \end{aligned} \quad (\text{A.18})$$

where the last implicit constant is independent of t and ρ . By the definition of $G_{t,\rho}$, the numerator in (3.5) is $G_{t,\rho} - \Psi(1)$. Hence

$$G_{t,\rho} - \Psi(1) = c_a^{1/(\rho+1)} \left\{ \frac{t(\rho+1)}{\rho} \right\}^{\rho/(\rho+1)} (1 + \delta_{\mathcal{A},\rho}).$$

Multiplying both sides by m and recalling that

$$c' = m c_a^{1/(\rho+1)} \left(\frac{\rho+1}{\rho} \right)^{\rho/(\rho+1)},$$

we obtain $m\{G_{t,\rho} - \Psi(1)\} = c' t^{\rho/(\rho+1)} (1 + \delta_{\mathcal{A},\rho})$. Therefore,

$$\exp[-m\{G_{t,\rho} - \Psi(1)\}] = \exp\{-c' t^{\rho/(\rho+1)} (1 + \delta_{\mathcal{A},\rho})\}.$$

Substituting this identity into (A.18) yields (3.4).

It remains only to verify the two asserted limits of $\delta_{\mathcal{A}}$. First fix $\rho > 0$ and let $t \rightarrow \infty$. Let $x_{\rho,t} = \{(\rho+1)t/(\rho c_a)\}^{1/(\rho+1)}$ and $F_\rho(x) = \rho c_a x^{\rho+1}/(\rho+1)$, so that $F_\rho(x_{\rho,t}) = t$. Since ρ is fixed and $x_{\rho,t} \rightarrow \infty$, (A.17) and the mean-value theorem imply

$$D_0^\rho = x_{\rho,t}^\rho (1 + \eta_{\rho,t}), \quad |\eta_{\rho,t}| \lesssim \frac{\rho+1}{x_{\rho,t}} + \frac{1}{c_a x_{\rho,t}^\rho}. \quad (\text{A.19})$$

Taking (A.14), (A.17), and (A.19) together gives

$$\begin{aligned} G_{t,\rho} - \Psi(1) &= c_a x_{\rho,t}^\rho \left\{ 1 + O\left(\frac{\rho+1}{x_{\rho,t}} \right. \right. \\ &\quad \left. \left. + \frac{1 + |\log x_{\rho,t}|}{c_a x_{\rho,t}^\rho} \right) \right\}. \end{aligned} \quad (\text{A.20})$$

Since $m c_a x_{\rho,t}^\rho = m c_a^{1/(\rho+1)} \{(\rho+1)/\rho\}^{\rho/(\rho+1)} t^{\rho/(\rho+1)}$, the definition in (3.5) and (A.20) give

$$\delta_{\mathcal{A},\rho} = O\left\{ \frac{\rho+1}{x_{\rho,t}} + \frac{1 + |\log x_{\rho,t}|}{c_a x_{\rho,t}^\rho} \right\} = o(1)$$

as $t \rightarrow \infty$ for every fixed $\rho > 0$.

We next fix $t \geq \log 2$ and let $\rho \rightarrow \infty$. Write $e_d = \log(a_d^{-1}) - c_a d^\rho$. By (A.13), $|e_d| \leq C_e$, and hence

$$\begin{aligned} B(d) &= d \log 2 + c_a \left\{ d^{\rho+1} - \sum_{k=1}^d k^\rho \right\} \\ &\quad + d e_d - \sum_{k=1}^d e_k. \end{aligned} \quad (\text{A.21})$$

For $d \geq 2$ and $\rho \geq 1$, the mean-value theorem gives

$$\begin{aligned} d^{\rho+1} - \sum_{k=1}^d k^\rho &= \sum_{k=1}^{d-1} (d^\rho - k^\rho) \\ &\geq (d-1)\{d^\rho - (d-1)^\rho\} \\ &\geq \rho(d-1)^\rho \geq \frac{\rho d}{2}. \end{aligned}$$

Since $|d e_d - \sum_{k=1}^d e_k| \leq 2C_e d$, (A.21) implies

$$B(d) \geq d \left(\log 2 + \frac{c_a \rho}{2} - 2C_e \right), \quad d \geq 2. \quad (\text{A.22})$$

For fixed t , the right-hand side exceeds t for all $d \geq 2$ once ρ is sufficiently large. On the other hand, $B(1) = \log 2 \leq t$. The definition of D_0 therefore gives $D_0 = 1$. Consequently, $G_{t,\rho} - \Psi(1) = t$. At the same time,

$$c_a^{1/(\rho+1)} \left(\frac{\rho+1}{\rho} \right)^{\rho/(\rho+1)} t^{\rho/(\rho+1)} \rightarrow t. \quad (\text{A.23})$$

Therefore, (3.5) gives

$$\begin{aligned} 1 + \delta_{\mathcal{A},\rho} &= \frac{t}{c_a^{1/(\rho+1)} \{(\rho+1)/\rho\}^{\rho/(\rho+1)} t^{\rho/(\rho+1)}} \\ &\rightarrow 1. \end{aligned}$$

Thus $\delta_{\mathcal{A},\rho} \rightarrow 0$ as $\rho \rightarrow \infty$ with t fixed. This completes the proof. $\square$

Lemma 1 (Proposition 4.3 in Lu et al. (2021)). For any $W, M \in \mathbb{N}_+$ and $\delta_d \in (0, 3K^{-1}]$ with $K = W^2 M^2$, there exists a one-dimensional function ϕ implemented by a ReLU fully connected neural network with width $6W$ and depth $3M$ such that

$$\phi(x) = j, \quad \text{if } x \in \left[\frac{j}{K}, \frac{j+1}{K} - \delta \cdot \mathbb{1}_{\{j < K-1\}} \right],$$

for $j = 0, 1, \dots, K-1$.

Lemma 2 (Proposition 4.4 in Lu et al. (2021)). Given any $W, M, s \in \mathbb{N}_+$ and $\xi_i \in [0, 1]$ for $i = 0, 1, \dots, W^2 M^2 - 1$, there exists a function ϕ implemented by a ReLU fully connected neural network with width $16s(W + 1)\lceil \log_2(8W) \rceil$ and depth $5(M + 2)\lceil \log_2(4M) \rceil$ such that

$$|\phi(i) - \xi_i| \leq W^{-2s} M^{-2s}, \quad i = 0, 1, \dots, W^2 M^2 - 1,$$

and $0 \leq \phi(x) \leq 1, \forall x \in \mathbb{R}$.

Lemma 3 (Lemma 4.2 in Lu et al. (2021)). For any $W, M \in \mathbb{N}_+$ and $a, b \in \mathbb{R}$ with $a < b$, there exists a function ϕ implemented by a ReLU fully connected neural network with width $9W + 1$ and depth M such that $|\phi(x, y) - xy| \leq 6(b - a)^2 W^{-M}, \forall x, y \in [a, b]$.

Lemma 4 (Proposition 4.1 in Lu et al. (2021)). Assume $P_\alpha(x) = x^\alpha = x_1^{\alpha_1} x_2^{\alpha_2} \dots x_d^{\alpha_d}$ for $\alpha \in \mathbb{N}_0^d$ with $\|\alpha\|_1 \leq s$, where $s \in \mathbb{N}_+$. For any $W, M \in \mathbb{N}_+$, there exists a function ϕ implemented by a ReLU fully connected neural network with width $9W + s - 1$ and depth $7s^2 M$ such that

$$|\phi(x) - P(x)| \leq 9sW^{-7sM}, \quad \forall x \in [0, 1]^d.$$

Lemma 5 (Anisotropic removal of the trifling region). Let $g \in C(\Omega_{D_0})$. Suppose that $0 < \delta_d \leq 2s_d/(3L_d)$ for $d = 1, \dots, D_0$, and let $\tilde{f}$ be a ReLU network of size $\tilde{S}$, width $\tilde{W}$, and depth $\tilde{M}$, defined on Ω_{D_0} . If $|\tilde{f}(\mathbf{x}^{(D_0)}) - g(\mathbf{x}^{(D_0)})| \leq \varepsilon$ for every $\mathbf{x}^{(D_0)} \in \Omega_D \setminus \mathcal{R}_{D_0}$, then there exists a ReLU network $f^\sharp$ on Ω_{D_0} such that

$$\|f^\sharp - g\|_{L^\infty(\Omega_{D_0})} \leq \varepsilon + \sum_{d=1}^{D_0} \omega_{g,d}(\delta_d).$$

Moreover, $f^\sharp$ may be chosen with width at most $3^{D_0}(\tilde{W} + 4)$, depth at most $\tilde{M} + 2D_0$, and size $S \lesssim 3^{D_0}\{\tilde{S} + 1\}$. If $g = f_0^{(D_0)}$, then Assumption 2 gives $\omega_{f_0^{(D_0)},d}(\delta_d) \leq C_0 w_d \delta_d$, and therefore

$$\|f^\sharp - f_0^{(D_0)}\|_{L^\infty(\Omega_{D_0})} \leq \varepsilon + C_0 \sum_{d=1}^{D_0} w_d \delta_d.$$

Proof. The proof is the coordinatewise median-extension argument of Theorem 2.1 in Lu et al. (2021). At step d , use the three versions of the current network obtained by leaving the d -th coordinate unchanged and shifting it by $\pm \delta_d$, with exact ReLU clipping at the endpoints $\pm s_d$, and take their median. Because $\delta_d \leq 2s_d/(3L_d)$, at least two of these three arguments avoid the d -th collection of boundary strips. The median step therefore extends the approximation across those strips and increases the error by at most $\omega_{g,d}(\delta_d)$. Repeating this operation for $d = 1, \dots, D_0$ removes all coordinate strips and gives the displayed error bound. The median of three real numbers and the clipping maps are represented

exactly by fixed-size ReLU networks. Sequential application of the construction triples the width and size at each coordinate and adds two layers, yielding the stated complexity bounds. Finally, Assumption 2 and the one-dimensional mean value theorem give $\omega_{f_0^{(D_0)},d}(\delta_d) \leq C_0 w_d \delta_d$. $\square$