

Convex Optimization with Nested Evolving Feasible Sets (CONES) under Time-Varying Loss Functions

Rahul Vaze

September 11, 2026

Abstract

Convex Optimization with Nested Evolving Feasible Sets (CONES) was introduced in [10] where the objective function f remains fixed but the feasible region evolves over time as a nested sequence $S_1 \supseteq S_2 \supseteq \dots \supseteq S_T$. The goal of an online algorithm is to simultaneously minimize the regret with respect to hindsight static optimal benchmark and the total movement cost $M_{\mathcal{A}}(T)$ while ensuring feasibility at all times. CONES is an optimization-oriented generalization of the well-known *nested convex body chasing* (NCBC). In this paper, we extend CONES to allow for loss functions f'_t s to also change over time. When all loss functions are convex, we show that the projected proximal algorithm achieves $O(T^{1-\beta}), O(T^\beta)$ simultaneous regret and movement cost, respectively, for any $\beta \in [0, 1)$, over a time horizon of T . We also show that any *weakly adaptive* online algorithm with $O(T^\beta)$ regret has a movement cost of $\Omega\left(T^{\frac{1-\beta}{2}}\right)$ for any $\beta \in [0, 1)$. When all loss functions are strongly convex, we show that the projected proximal algorithm simultaneously achieves $O(1)$ regret and a movement cost of $O(\log T)$. To complement this, we show that any online algorithm with sublinear *anytime* regret has a movement cost of $\Omega(\log T)$.

1 Introduction

In this paper, we consider the following problem known as CONES [10]. At each round t , a convex (loss) function $f_t : \mathcal{X} \rightarrow \mathbb{R}$ and a convex set $S_t \subseteq \mathcal{X}$ is revealed such that $S_t \subseteq S_{t-1}$, and $S_0 = \mathcal{X} \subset \mathbb{R}^d$ is a convex, compact, and bounded set. Once S_t is revealed, the objective for an online algorithm $\mathcal{A}$ is to choose action $x_t \in S_t$ (feasible action) so as to **simultaneously** minimize the loss function cost

$$C_{\mathcal{A}}(T) = \sum_{t=1}^T f_t(x_t), \quad (1)$$

and total movement cost

$$M_{\mathcal{A}}(T) = \sum_{t=1}^T \|x_t - x_{t-1}\|, \quad (2)$$

where $x_0 \in \mathcal{X}$ is some fixed action. Throughout, we use $\|\cdot\|$ to denote the 2-norm or the Euclidean distance.

The performance of $\mathcal{A}$ is compared against a *static* optimal benchmark OPT that chooses its action $x^{\text{OPT}} \in S_T$ (assumed non-empty) that minimizes the loss function cost, i.e.,

$$x^{\text{OPT}} \in \arg \min_{x \in S_T} \sum_t f_t(x)$$

and

$$C_{\text{OPT}}(T) = \sum_{t=1}^T f_t(x^{\text{OPT}}).$$

Since S_t 's are nested, x^{OPT} is feasible with respect to all S'_t s. The movement cost of OPT is simply

$$M_{\text{OPT}}(T) = \|x^{\text{OPT}} - x_0\|.$$

The **regret** of $\mathcal{A}$ is then defined as

$$\text{Regret}_{\mathcal{A}}(T) = \sup_{f_t, S_t} \{C_{\mathcal{A}}(T) - C_{\text{OPT}}(T)\}. \quad (3)$$

For $\mathcal{A}$ its two objectives are: minimize $\text{Regret}_{\mathcal{A}}(T)$ and movement cost $M_{\mathcal{A}}(T)$ simultaneously.

1.1 Prior Work with $f_t = f \forall t$

CONES was introduced in [10] for the case when $f_t = f \forall t$. [10] also discussed in detail the motivation to study it, its applications, and its connections to related prior work, e.g. in nested convex body chasing (NCBC) [1–3, 6, 7, 15], convex online constrained optimization (COCO) [8, 11, 12, 17, 18, 20–22], sensitivity/perturbation analysis of optimization problems [5], trajectory tracking [16], etc. With $f_t = f \forall t$ the following results were derived in [10].

1. When f is **strongly convex**, it showed that the greedy algorithm $\mathcal{G}$ that chooses $x_t = x_t^* = \arg \min_{x \in S_t} f(x)$ for all t has minimum regret among all online algorithms but incurs a movement cost of $\Theta(\sqrt{T})$. Next, a ‘lazy’ algorithm called FRUGAL was analyzed, that plays action $x_t = x_t^*$ only when it is forced to do so in order to keep its regret non-positive and otherwise plays the action obtained by projecting its current action x_{t-1} on to the most recently revealed set S_t . The regret of FRUGAL is non-positive by definition and its movement cost is shown to be $O(\log T)$, a fundamental improvement over $\mathcal{G}$ when f is also smooth. This upper bound was complemented by a lower bound, that showed that any online algorithm with sublinear *anytime* regret has movement cost $\Omega\left(\sqrt{\frac{\log T}{\log \log T}}\right)$. Recently, an algorithm that combines FRUGAL and Level Set Projection (LSP) algorithm of [10] has been shown to achieve a non-positive regret and movement cost of $O(\sqrt{\log T})$ in [13].
2. When f is **convex**, [10] showed a negative result that the greedy algorithm $\mathcal{G}$ has a movement cost of $\Omega(T)$. Next, it showed that the LSP algorithm achieves a (regret, movement cost) tuple of $(T^{1-\beta}, T^\beta)$ for any $\beta \in [0, 1]$.

In this paper, with arbitrary time-varying f_t ’s, we derive the following results.

1.2 Contributions

1. When f_t ’s are **strongly convex**, we show that the projected proximal algorithm (we refer to it as Prox) that chooses

$$x_t \in \arg \min_{x \in S_t} \left\{ f_t(x) + \frac{1}{2\eta_t} \|x - x_{t-1}\|^2 \right\}, \quad (4)$$

has regret of $O(1)$ and its movement cost is $O(\log T)$, when $\eta_t = \frac{1}{\mu t}$. Prox is conceptually a different algorithm than FRUGAL [10] and automatically adapts to changing constraints rather than being forced depending on current regret.

We also show that any online algorithm with sublinear *anytime* regret has movement cost $\Omega(\log T)$. Thus obtaining a tight result and showing that Prox is an optimal (order-wise) algorithm.

It is worth noting that compared to the case when $f_t = f$ for all t , the optimal movement cost with sublinear *anytime* regret is order-wise different.

2. When f is **convex**, we show that Prox (4) achieves a (regret, movement cost) tuple of $(T^{1-\beta}, T^\beta)$ for any $\beta \in [0, 1]$ by choosing $\eta = \frac{D^2}{\epsilon T}$ and $\epsilon = T^{-\beta}$.

Moreover, we also show that any *weakly adaptive* online algorithm with $O(T^\beta)$ regret has a movement cost of $\Omega\left(T^{\frac{1-\beta}{2}}\right)$ for $\beta \in [0, 1)$ **even when $f_t = f$ for all t** , the special case studied in [10]. No such lower bound was provided in [10].

2 f_t ’s are strongly convex

In this section, we consider the twin objectives of simultaneously minimizing regret (3) and movement cost (2), when f_t ’s are μ -strongly convex and G -Lipschitz. Throughout, we assume that the diameter of $\mathcal{X}$ is at most D .

2.1 Upper Bound

Theorem 1. When f_t 's are μ -strongly convex, G -Lipschitz, for algorithm Prox (4) with $\eta_t = \frac{1}{\mu t}$,

$$\text{Regret}_{\mathcal{A}}(T) \leq \frac{\mu D^2}{2} = O(1), \quad (5)$$

and

$$M_{\mathcal{A}}(T) \leq C_d \left(D + \frac{G}{\mu} (1 + \log T) \right) = O(\log T),$$

where C_d is a constant depending only on $\|\cdot\|$ and the dimension d . The exact bound on C_d can be extracted from [19].

The main idea of the proof is to exploit the strong convexity of f_t 's to upper bound the regret by an $O(1)$ quantity, while using the fact that (proved in Lemma 2) x_t defined in (4) is equivalent to

$$x_t = \Pi_{S_t}(x_{t-1} - \eta_t g_t). \quad (6)$$

where $g_t \in \partial f_t(x_t)$, and $\partial f_t(x_t)$ denotes the subgradient set of f_t at x_t . This association (6) allows us to utilize a recent result in [14] that upper bounds $\sum_t \|x_t - x_{t-1}\|$ for iterates x_t satisfying (6) when sets S_t 's are nested. Remarkably, it is worth noting that η_t is in the numerator of (6) unlike being in the denominator of (4). Thus, with $\eta_t = \frac{1}{\mu t}$ allows us to bound the movement cost as $O(\log T)$.

Remark 2.1:

The regret bound of Theorem 1

$$\text{Regret}_{\mathcal{A}}(T) \leq \frac{\mu D^2}{2}$$

has a qualitatively different dependence on the strong-convexity parameter μ from the usual $O(1/\mu)$ regret bound for online convex optimization [9] or COCO [14]. This is because algorithm Prox uses the entire loss function f_t in its implicit proximal update and chooses

$$\eta_t = \frac{1}{\mu t}.$$

As μ decreases, the proximal regularization becomes weaker and the update approaches the greedy minimizer of the current loss over S_t . In the present CONES setting, where f_t and S_t are revealed before x_t is chosen, the greedy action satisfies

$$f_t(x_t) \leq f_t(x^{\text{OPT}})$$

at every round, since $x^{\text{OPT}} \in S_t$. Thus the regret can in fact approach zero as $\mu \rightarrow 0$. The dependence

$$\text{Regret}_{\mathcal{A}}(T) = O(\mu D^2)$$

is therefore consistent with the algorithm becoming increasingly greedy as the strong convexity vanishes.

To complement this upper bound result, we derive a matching lower bound as follows.

2.2 Lower Bound

Theorem 2. Suppose that an online algorithm $\mathcal{A}$ satisfies, for every t ,

$$\text{Regret}_{\mathcal{A}}(t) := \sup_{\{f_r, S_r\}_{r=1}^t} \sum_{r=1}^t f_r(x_r) - \min_{x \in S_t} \sum_{r=1}^t f_r(x) \leq R(t). \quad (7)$$

An online algorithm $\mathcal{A}$ is defined to have anytime sublinear regret if for all t ,

$$R(t) = o(t),$$

For any $\mathcal{A}$ with anytime sublinear regret, against an adaptive adversary, there exists a sequence of 1-strongly convex and 1-Lipschitz loss functions f_t for which

$$M_{\mathcal{A}}(T) = \sum_{t=1}^T \|x_t - x_{t-1}\| = \Omega(\log T).$$

Discussion: Theorem 1 and 2 together fully characterize the optimal movement cost for any algorithm that can achieve anytime $o(T)$ regret. Compared to the case when $f_t = f$ for all t , where the movement cost is $O(\sqrt{\log T})$ and $\Omega\left(\sqrt{\frac{\log T}{\log \log T}}\right)$ [10, 13] for any algorithm that can achieve anytime $o(T)$ regret, with changing f_t 's it is $\Theta(\log T)$. Thus, there is a fundamental degradation which is expected in this more general setting.

3 f_t 's are convex

In this section, we consider the twin objectives of minimizing regret (3) and movement cost (2), when f_t 's are just convex and G -Lipschitz.

3.1 Upper Bound

Theorem 3. When f_t 's are convex and G -Lipschitz, for any $\epsilon > 0$, algorithm Prox (4), with

$$\eta = \frac{D^2}{\epsilon T}.$$

satisfies

$$\text{Regret}_{\mathcal{A}}(T) \leq \frac{D^2}{2\eta} = \frac{\epsilon T}{2},$$

and

$$M_{\mathcal{A}}(T) \leq C_d \left(D + \frac{GD^2}{\epsilon} \right),$$

where C_d is the same constant that appears in Theorem 1. In particular, for any $0 \leq \beta \leq 1$, choosing

$$\epsilon = T^{-\beta}$$

yields

$$\text{Regret}_{\mathcal{A}}(T) = O(T^{1-\beta}), \quad M_{\mathcal{A}}(T) = O_d(T^{\beta}).$$

We next derive lower bound on achievable simultaneous regret and movement cost when f_t 's are convex.

3.2 Lower Bounds

3.2.1 Weakly Adaptive Algorithms

Definition 3.1: Weakly adaptive algorithm

An online algorithm $\mathcal{A}$ for CONES with changing losses $f_1, \dots, f_T$ is called *R-weakly adaptive* if, for every CONES instance, every interval

$$I = \{r, r+1, \dots, s\} \subseteq [T],$$

with benchmark

$$x_s^{\text{OPT}} \in \arg \min_{x \in S_s} \sum_{t=1}^s f_t(x)$$

$\mathcal{A}$ satisfies

$$\sum_{t=r}^s (f_t(x_t) - f_t(x_s^{\text{OPT}})) \leq R.$$

Since the feasible sets are nested,

$$S_s \subseteq S_t \quad \text{for all } t \leq s,$$

and hence

$$x_s^{\text{OPT}} \in S_t \quad \text{for all } t \in [a, b].$$

Thus, x_s^{OPT} is feasible throughout the interval I . Taking $r = 1$ and $s = T$ gives

$$\sum_{t=1}^T (f_t(x_t) - f_t(x_T^{\text{OPT}})) \leq R,$$

which is the usual terminal static-regret guarantee.

The notion of weakly adaptive algorithms is quite popular in related literature [4, 14]. We next derive a lower bound on the simultaneous regret and movement cost for any weakly adaptive algorithm **when** $f_t = f$ **for all** t , which directly applies for general f_t 's, and thereafter give examples of weakly adaptive algorithms.

Theorem 4. Fix $T \geq 1$ and let

$$R := \text{Regret}_{\mathcal{A}}(T) \geq 0, \quad \text{satisfy} \quad T \geq 16(R + 1).$$

Consider any online algorithm $\mathcal{A}$ satisfying the weakly adaptive regret guarantee

$$\sum_{t=r}^s (f_t(x_t) - f_t(x_s^{\text{OPT}})) \leq R \quad \text{for every } [r, s] \subseteq [T], \quad \text{where} \quad x_s^{\text{OPT}} \in \arg \min_{x \in S_s} \sum_{t=1}^s f_t(x).$$

Then there exists a two-dimensional CONES instance with

$$\mathcal{X} = [-1, 1] \times [0, 1], \quad \text{diam}(\mathcal{X}) = \sqrt{5}, \quad S_0 = \mathcal{X}, \quad S_t \subseteq S_{t-1},$$

and fixed loss $f : \mathcal{X} \rightarrow \mathbb{R}$ that is convex and 1-Lipschitz, such that

$$M_{\mathcal{A}}(T) \geq \frac{1}{8} \sqrt{\frac{T}{R+1}} - 1.$$

In particular, if

$$\text{Regret}_{\mathcal{A}}(T) = O(T^\beta), \quad 0 \leq \beta < 1,$$

then

$$M_{\mathcal{A}}(T) = \Omega\left(T^{(1-\beta)/2}\right).$$

3.3 Examples of weakly adaptive algorithms for CONES

1. Prox algorithm (4) is R -weakly adaptive, with

$$R = \frac{D^2}{2\eta},$$

when $f_t : \mathcal{X} \rightarrow \mathbb{R}$ are convex and Prox uses a constant stepsize $\eta_t = \eta$ for all t .

We show it as follows. Fix any interval $I = [a, b] \subseteq [T]$, and let

$$x_b^{\text{OPT}} \in \arg \min_{x \in S_b} \sum_{t=1}^b f_t(x).$$

Since the feasible sets are nested,

$$S_b \subseteq S_t \quad \text{for all } t \leq b,$$

and hence

$$x_b^{\text{OPT}} \in S_t \quad \text{for all } t \in [a, b].$$

Let $g_t \in \partial f_t(x_t)$. Since f_t is convex,

$$f_t(x_t) - f_t(x_b^{\text{OPT}}) \leq \langle g_t, x_t - x_b^{\text{OPT}} \rangle.$$

The first-order optimality condition for the proximal update (4) gives

$$\left\langle g_t + \frac{x_t - x_{t-1}}{\eta}, x_b^{\text{OPT}} - x_t \right\rangle \geq 0.$$

Therefore,

$$\langle g_t, x_t - x_b^{\text{OPT}} \rangle \leq \frac{1}{\eta} \langle x_t - x_{t-1}, x_b^{\text{OPT}} - x_t \rangle.$$

Using

$$2\langle x_t - x_{t-1}, x_t - x_b^{\text{OPT}} \rangle = \|x_t - x_{t-1}\|^2 + \|x_t - x_b^{\text{OPT}}\|^2 - \|x_{t-1} - x_b^{\text{OPT}}\|^2,$$

we obtain

$$f_t(x_t) - f_t(x_b^{\text{OPT}}) \leq \frac{\|x_{t-1} - x_b^{\text{OPT}}\|^2 - \|x_t - x_b^{\text{OPT}}\|^2 - \|x_t - x_{t-1}\|^2}{2\eta}.$$

Summing over $t = a, \dots, b$ and dropping the nonpositive movement term gives

$$\sum_{t=a}^b (f_t(x_t) - f_t(x_b^{\text{OPT}})) \leq \frac{\|x_{a-1} - x_b^{\text{OPT}}\|^2 - \|x_b - x_b^{\text{OPT}}\|^2}{2\eta} \leq \frac{D^2}{2\eta}.$$

Since this holds for every interval $[a, b] \subseteq [T]$, the Prox algorithm is R -weakly adaptive with

$$R = \frac{D^2}{2\eta}.$$

2. With changing losses f_t , the GREEDY algorithm plays

$$x_t \in \arg \min_{x \in S_t} f_t(x)$$

at every round.

Consider any interval $I = [a, b] \subseteq [T]$. By definition,

$$x_b^{\text{OPT}} \in \arg \min_{x \in S_b} \sum_{t=1}^b f_t(x).$$

Since the feasible sets are nested,

$$S_b \subseteq S_t \quad \forall t \leq b,$$

and hence

$$x_b^{\text{OPT}} \in S_t \quad \forall t \in [a, b].$$

By the definition of GREEDY,

$$f_t(x_t) = \min_{x \in S_t} f_t(x) \leq f_t(x_b^{\text{OPT}}) \quad \forall t \in [a, b].$$

Therefore,

$$\sum_{t=a}^b (f_t(x_t) - f_t(x_b^{\text{OPT}})) \leq 0.$$

Hence GREEDY is 0-weakly adaptive.

3. Level Set Projection (LSP) algorithm [10]: The LSP algorithm was defined in [10] for the case $f_t = f$ for all t . Its natural extension to changing losses is as follows. Define

$$L_t := \min_{x \in S_t} f_t(x), \quad \Phi_t := \{x \in S_t : f_t(x) \leq L_t + \varepsilon\},$$

and choose some

$$x_t \in \Phi_t.$$

For example, one may choose

$$x_t = \Pi_{\Phi_t}(x_{t-1}).$$

By construction,

$$f_t(x_t) \leq \min_{x \in S_t} f_t(x) + \varepsilon.$$

Lemma 1. *The changing-loss version of LSP is $T\varepsilon$ -weakly adaptive.*

Proof. Consider any interval

$$I = [a, b] \subseteq [T].$$

By definition,

$$x_b^{\text{OPT}} \in \arg \min_{x \in S_b} \sum_{t=1}^b f_t(x).$$

Since the feasible sets are nested, $S_b \subseteq S_t \forall t \leq b$, and hence $x_b^{\text{OPT}} \in S_t \forall t \in [a, b]$.

By construction of x_t ,

$$f_t(x_t) \leq \min_{x \in S_t} f_t(x) + \varepsilon \leq f_t(x_b^{\text{OPT}}) + \varepsilon.$$

Thus,

$$f_t(x_t) - f_t(x_b^{\text{OPT}}) \leq \varepsilon \quad \forall t \in [a, b].$$

Summing over the interval gives

$$\sum_{t=a}^b (f_t(x_t) - f_t(x_b^{\text{OPT}})) \leq |I|\varepsilon \leq T\varepsilon.$$

Hence the changing-loss version of LSP is $T\varepsilon$ -weakly adaptive. □

4 Conclusion

In this paper, we generalized the important optimization paradigm of Convex Optimization with Nested Evolving Feasible Sets (CONES), introduced in [10], from a fixed objective function to a setting in which both the loss functions and the feasible sets may evolve over time, while the benchmark remains a static hindsight optimum. This extension allows us to separate the effects of evolving optimization geometry from those arising from temporal variation in the objective itself.

For the strongly convex case, changing loss functions increases the movement required under $o(T)$ anytime regret from the previously known $O(\sqrt{\log T})$ upper bound in the fixed-objective setting to the tight $\Theta(\log T)$ rate established here. For the convex case, we establish the same upper bound in the evolving objective setting as already known for the fixed-objective setting, while developing new lower bounds that apply even for the fixed-objective setting. With these improvements we have widened the scope of CONES and developed a better theoretical understanding.

References

- [1] C. J. Argue, Sébastien Bubeck, Michael B. Cohen, Anupam Gupta, and Yin Tat Lee. A nearly-linear bound for chasing nested convex bodies, 2018.
- [2] C. J. Argue, Anupam Gupta, Ziye Tang, and Guru Guruganesh. Chasing convex bodies with linear competitive ratio. *J. ACM*, 68(5), August 2021.
- [3] Nikhil Bansal, Martin Böhm, Marek Eliáš, Grigorios Koumoutsos, and Seeun William Umboh. Nested convex bodies are chaseable, 2017.
- [4] Martino Bernasconi, Matteo Castiglioni, and Andrea Celli. No-regret is not enough! bandits with general constraints through adaptive regret minimization. *arXiv preprint arXiv:2405.06575*, 2024.
- [5] J. Frédéric Bonnans and Alexander Shapiro. *Perturbation Analysis of Optimization Problems*. Springer Series in Operations Research. Springer Science & Business Media, New York, 2000.
- [6] Sébastien Bubeck, Bo’az Klartag, Yin Tat Lee, Yuezhi Li, and Mark Sellke. Chasing nested convex bodies nearly optimally, 2021.
- [7] Joel Friedman and Nathan Linial. On convex body chasing. *Discrete & Computational Geometry*, 9(3):293–321, 1993.
- [8] Hengquan Guo, Xin Liu, Honghao Wei, and Lei Ying. Online convex optimization with hard constraints: Towards the best of two worlds and beyond. *Advances in Neural Information Processing Systems*, 35:36426–36439, 2022.
- [9] Elad Hazan. Introduction to online convex optimization. *CoRR*, abs/1909.05207, 2019.
- [10] Karthick M. Krishna, Haricharan Balasundaram, and Rahul Vaze. Convex optimization with nested evolving feasible sets. *arXiv preprint arXiv:2605.07386*, 2026.
- [11] Nikolaos Liakopoulos, Apostolos Destounis, Georgios Paschos, Thrasyvoulos Spyropoulos, and Panayotis Mertikopoulos. Cautious regret minimization: Online optimization with long-term budget constraints. In *International Conference on Machine Learning*, pages 3944–3952. PMLR, 2019.
- [12] Michael J Neely and Hao Yu. Online convex optimization with time-varying constraints. *arXiv preprint arXiv:1702.04783*, 2017.
- [13] Dhruv Sarkar and Aprameyo Chakrabartty. Nested convex-body chasing for online optimization with evolving feasible sets, 2026.
- [14] Dhruv Sarkar and Abhishek Sinha. Improved guarantees for constrained online convex optimization via self-contraction. *arXiv preprint arXiv:2605.21107*, 2026.
- [15] Mark Sellke. *Chasing Convex Bodies Optimally*, pages 313–335. Springer International Publishing, Cham, 2023.
- [16] Andrea Simonetto, Emiliano Dall’Anese, Santiago Paternain, Geert Leus, and Georgios B Giannakis. Time-varying convex optimization: Time-structured algorithms and applications. *Proceedings of the IEEE*, 108(11):2032–2048, 2020.
- [17] Abhishek Sinha and Rahul Vaze. Optimal algorithms for online convex optimization with adversarial constraints. In *The Thirty-eighth Annual Conference on Neural Information Processing Systems*, 2024.
- [18] Abhishek Sinha and Rahul Vaze. Beyond $\tilde{O}(\sqrt{T})$ constraint violation for online convex optimization with adversarial constraints. In *The Thirty-ninth Annual Conference on Neural Information Processing Systems, NeurIPS 2025*, 2025.
- [19] Eugene Stepanov and Yana Teplitskaya. Self-contracted curves have finite length. *Journal of the London Mathematical Society*, 96(2):455–481, 2017.

- [20] Wen Sun, Debadeepta Dey, and Ashish Kapoor. Safety-aware algorithms for adversarial contextual bandit. In *International Conference on Machine Learning*, pages 3280–3288. PMLR, 2017.
- [21] Xinlei Yi, Xiuxian Li, Tao Yang, Lihua Xie, Yiguang Hong, Tianyou Chai, and Karl H Johansson. Distributed online convex optimization with adversarial constraints: Reduced cumulative constraint violation bounds under Slater’s condition. *arXiv preprint arXiv:2306.00149*, 2023.
- [22] Hao Yu, Michael Neely, and Xiaohan Wei. Online convex optimization with stochastic constraints. *Advances in Neural Information Processing Systems*, 30, 2017.

5 Proof of Theorem 1

Proof. Let

$$g_t \in \partial f_t(x_t),$$

where $\partial f_t(x_t)$ denotes the sub-gradient of f_t at x_t . Since f_t is μ -strongly convex,

$$f_t(x_t) - f_t(x^{\text{OPT}}) \leq \langle g_t, x_t - x^{\text{OPT}} \rangle - \frac{\mu}{2} \|x_t - x^{\text{OPT}}\|^2. \quad (8)$$

The first-order optimality condition for algorithm Prox (4) implies

$$\left\langle g_t + \frac{x_t - x_{t-1}}{\eta_t}, x^{\text{OPT}} - x_t \right\rangle \geq 0,$$

since $x^{\text{OPT}} \in S_t$ for all t .

Therefore,

$$\langle g_t, x_t - x^{\text{OPT}} \rangle \leq \frac{1}{\eta_t} \langle x_t - x_{t-1}, x^{\text{OPT}} - x_t \rangle.$$

Using

$$\langle x_t - x_{t-1}, x^{\text{OPT}} - x_t \rangle = -\langle x_t - x_{t-1}, x_t - x^{\text{OPT}} \rangle$$

and the identity

$$2\langle x_t - x_{t-1}, x_t - x^{\text{OPT}} \rangle = \|x_t - x^{\text{OPT}}\|^2 + \|x_t - x_{t-1}\|^2 - \|x_{t-1} - x^{\text{OPT}}\|^2,$$

we obtain

$$\langle g_t, x_t - x^{\text{OPT}} \rangle \leq \frac{1}{2\eta_t} (\|x_{t-1} - x^{\text{OPT}}\|^2 - \|x_t - x^{\text{OPT}}\|^2 - \|x_t - x_{t-1}\|^2). \quad (9)$$

Substituting this into (8) and using $\eta_t = 1/(\mu t)$,

$$f_t(x_t) - f_t(x^{\text{OPT}}) \leq \frac{\mu}{2} (t\|x_{t-1} - x^{\text{OPT}}\|^2 - (t+1)\|x_t - x^{\text{OPT}}\|^2 - t\|x_t - x_{t-1}\|^2).$$

Summing over t yields

$$\text{Regret}_{\mathcal{A}}(T) \leq \frac{\mu}{2} \left(\|x_0 - x^{\text{OPT}}\|^2 - (T+1)\|x_T - x^{\text{OPT}}\|^2 - \sum_{t=1}^T t\|x_t - x_{t-1}\|^2 \right),$$

and therefore

$$\text{Regret}_{\mathcal{A}}(T) \leq \frac{\mu D^2}{2}.$$

For bounding the movement cost $M_{\mathcal{A}}(T)$, we rewrite Prox (4) as follows.

Lemma 2. x_t defined in (4) is equivalent to

$$x_t = \Pi_{S_t}(x_{t-1} - \eta_t g_t). \quad (10)$$

where $g_t \in \partial f_t(x_t)$.

Note that (10) is a fixed point equation where x_t appears on both sides of the equality since $g_t \in \partial f_t(x_t)$. This fixed point property is only a structural result and not used by the Prox algorithm. Moreover, importantly η_t appears in the numerator (10) compared to the denominator (4), which allows a *win-win* on two fronts given that $\eta_t = \frac{1}{\mu t}$: T independent regret bound and $O(\log T)$ movement cost.

Proof of Lemma 2 is elementary and provided later for completeness.

Thus, the update (4) of algorithm Prox is of the form

$$x_t = \Pi_{S_t}(x_{t-1} - e_t), \quad \text{where } e_t := \eta_t g_t, \quad (11)$$

and

$$\|e_t\| \leq \frac{G}{\mu t},$$

since $\eta_t = 1/(\mu t)$ and $\|g_t\| \leq G$.

For a nested projected-gradient trajectory (11) we invoke the following result.

Lemma 3 (Nested-projection movement bound [14, Lemma 4.1]). *Let*

$$K_0 \supseteq K_1 \supseteq \dots \supseteq K_T$$

be a nested sequence of nonempty closed convex subsets of $\mathbb{R}^d$ such that

$$\text{diam}(K_0) \leq D.$$

Let $z_0 \in K_0$, and suppose that

$$z_t = \Pi_{K_t}(z_{t-1} - e_t), \quad t = 1, \dots, T,$$

where $e_t \in \mathbb{R}^d$ is arbitrary. Then

$$\boxed{\sum_{t=1}^T \|z_t - z_{t-1}\| \leq C_d \left(D + \sum_{t=1}^T \|e_t\| \right)},$$

where $C_d > 0$ is a constant depending only on $\|\cdot\|$ and the dimension d .

In our setting $S_t = K_t$ and $S_t \subseteq S_{t-1}$, thus, using Lemma 3, we get from (11), that for algorithm Prox,

$$\sum_{t=1}^T \|x_t - x_{t-1}\| \leq C_d \left(D + \sum_{t=1}^T \|e_t\| \right),$$

with $\|e_t\| \leq \frac{G}{\mu t}$. Hence

$$M_{\mathcal{A}}(T) \leq C_d \left(D + \frac{G}{\mu} \sum_{t=1}^T \frac{1}{t} \right) \leq C_d \left(D + \frac{G}{\mu} (1 + \log T) \right).$$

□

Proof of Lemma 2. Let $y_t := x_{t-1} - \eta_t g_t$. From (4), x_t minimizes the function

$$\phi_t(x) := f_t(x) + \frac{1}{2\eta_t} \|x - x_{t-1}\|^2$$

over the closed convex set S_t . Thus, the first-order optimality condition implies

$$0 \in \partial \phi_t(x_t) + N_{S_t}(x_t),$$

where $N_S(x)$ is the normal cone to S at point $x \in S$. Because

$$\partial\phi_t(x_t) = g_t + \frac{1}{\eta_t}(x_t - x_{t-1}),$$

we obtain

$$0 \in g_t + \frac{1}{\eta_t}(x_t - x_{t-1}) + N_{S_t}(x_t).$$

Thus there exists some

$$n_t \in N_{S_t}(x_t)$$

such that

$$g_t + \frac{1}{\eta_t}(x_t - x_{t-1}) + n_t = 0.$$

Multiplying by η_t gives

$$\eta_t g_t + x_t - x_{t-1} + \eta_t n_t = 0,$$

and hence

$$x_{t-1} - \eta_t g_t - x_t = \eta_t n_t.$$

By the definition of y_t ,

$$y_t - x_t = \eta_t n_t.$$

Since the normal cone is a cone, $n_t \in N_{S_t}(x_t)$ implies

$$\eta_t n_t \in N_{S_t}(x_t).$$

Therefore

$$y_t - x_t \in N_{S_t}(x_t). \tag{12}$$

We next recall the well-known projection characterization, whose proof is included for completeness. Let S be a nonempty closed convex set. We claim that, for $x \in S$,

$$x = \Pi_S(y) \iff y - x \in N_S(x). \tag{13}$$

First suppose that $x = \Pi_S(y)$. By definition,

$$\|y - x\|^2 \leq \|y - z\|^2 \quad \forall z \in S.$$

For any $z \in S$ and any $\lambda \in (0, 1]$, convexity gives

$$x + \lambda(z - x) \in S.$$

Hence

$$\|y - x\|^2 \leq \|y - x - \lambda(z - x)\|^2.$$

Expanding,

$$0 \leq -2\lambda\langle y - x, z - x \rangle + \lambda^2\|z - x\|^2.$$

Dividing by λ and letting $\lambda \downarrow 0$ gives

$$\langle y - x, z - x \rangle \leq 0 \quad \forall z \in S.$$

By the definition of the normal cone,

$$y - x \in N_S(x).$$

Conversely, suppose that $x \in S$ and

$$y - x \in N_S(x).$$

Then, for every $z \in S$,

$$\langle y - x, z - x \rangle \leq 0.$$

Therefore

$$\begin{aligned} \|y - z\|^2 &= \|y - x - (z - x)\|^2 \\ &= \|y - x\|^2 + \|z - x\|^2 - 2\langle y - x, z - x \rangle \\ &\geq \|y - x\|^2. \end{aligned}$$

Thus x minimizes $\|y - z\|$ over $z \in S$, so

$$x = \Pi_S(y).$$

This proves (13).

Finally, $x_t \in S_t$ by construction, and (12) gives

$$y_t - x_t \in N_{S_t}(x_t).$$

Applying (13) with $S = S_t$, $y = y_t$, and $x = x_t$ yields

$$x_t = \Pi_{S_t}(y_t) = \Pi_{S_t}(x_{t-1} - \eta_t g_t).$$

□

6 Proof of Theorem 2

Proof. Let the mother set be

$$\mathcal{X} = [-1/2, 1/2],$$

and let

$$S_t = \mathcal{X} \quad \forall t, i.e.,$$

all sets S_t 's are identical and hence nested.

Let

$$N_k = 33^k, \quad k \geq 0,$$

so that $N_0 = 1$. Define the first phase by

$$\mathcal{I}_1 = \{1, \dots, N_1\},$$

and, for every $k \geq 2$, define

$$\mathcal{I}_k = \{N_{k-1} + 1, \dots, N_k\}.$$

Thus, for every $k \geq 2$, if

$$N = N_{k-1},$$

then phase k consists of exactly

$$N_k - N_{k-1} = 32N$$

rounds.

Fix

$$s_1 = \frac{1}{2}$$

in advance, and for every $t \in \mathcal{I}_1$ define

$$f_t(x) = \frac{1}{2}(x - s_1)^2.$$

At the beginning of phase $k \geq 2$, the phase starts at round $N_{k-1} + 1$. Recall that, in CONES, the action x_t is chosen after f_t and S_t are revealed. Hence, at the beginning of phase k , the adversary can observe $x_{N_{k-1}}$ before choosing s_k .

After observing action $x_{N_{k-1}}$ of $\mathcal{A}$, the adversary chooses

$$s_k \in \{-1/2, 1/2\}$$

such that

$$\|x_{N_{k-1}} - s_k\| \geq \frac{1}{2},$$

and then for every $t \in \mathcal{I}_k$, uses

$$f_t(x) = \frac{1}{2}(x - s_k)^2, \quad t = N_{k-1} + 1, \dots, N_k.$$

Since $x, s_k \in [-1/2, 1/2]$,

$$\|f'_t(x)\| = \|x - s_k\| \leq 1,$$

so every f_t is 1-Lipschitz. Moreover, every f_t is 1-strongly convex.

Next, we show that for any $\mathcal{A}$ with $o(t)$ anytime regret, for all sufficiently large N , any phase of length $32N$ must incur at least $1/4$ movement. Fix a phase $k \geq 2$ and write

$$N = N_{k-1}.$$

Suppose, for contradiction, that the total movement in phase k

$$\sum_{t=N+1}^{N+32N} \|x_t - x_{t-1}\| < \frac{1}{4}.$$

Then, for every

$$t \in \{N+1, \dots, N+32N\},$$

$$\|x_t - x_N\| \leq \sum_{r=N+1}^t \|x_r - x_{r-1}\| < \frac{1}{4}.$$

By the choice of s_k ,

$$\|x_t - s_k\| \geq \|x_N - s_k\| - \|x_t - x_N\| > \frac{1}{2} - \frac{1}{4} = \frac{1}{4}.$$

Hence

$$f_t(x_t) = \frac{1}{2}(x_t - s_k)^2 > \frac{1}{32} \tag{14}$$

for every

$$t = N+1, \dots, N+32N.$$

Consider the prefix $[1, \tau]$ where

$$\tau = N + 32N = 33N.$$

Let

$$x_\tau^{\text{OPT}} \in \arg \min_{x \in \mathcal{X}} \sum_{r=1}^{\tau} f_r(x)$$

be the static benchmark for this prefix, since $S_t = \mathcal{X}$ for all t .

Every loss has the form

$$f_r(x) = \frac{1}{2}(x - s_r)^2, \quad s_r \in \{-1/2, 1/2\}.$$

Hence

$$\sum_{r=1}^{\tau} f_r(x) = \frac{1}{2} \sum_{r=1}^{\tau} (x - s_r)^2,$$

whose minimizer is

$$x_\tau^{\text{OPT}} = \frac{1}{\tau} \sum_{r=1}^{\tau} s_r.$$

Over the first N rounds, the offsets s_r are those prescribed by phases $1, \dots, k-1$, while over the remaining $32N$ rounds, corresponding to the current phase k , the offset is s_k . Therefore,

$$\begin{aligned} x_\tau^{\text{OPT}} - s_k &= \frac{1}{33N} \left(\sum_{r=1}^N s_r + 32N s_k \right) - s_k \\ &= \frac{1}{33N} \left(\sum_{r=1}^N s_r + 32N s_k - 33N s_k \right) \\ &= \frac{1}{33N} \sum_{r=1}^N (s_r - s_k). \end{aligned}$$

Since

$$\|s_r - s_k\| \leq 1,$$

we obtain

$$\|x_\tau^{\text{OPT}} - s_k\| \leq \frac{N}{33N} = \frac{1}{33} < \frac{1}{8}.$$

Consequently, for every

$$\begin{aligned} t &= N+1, \dots, N+32N, \\ f_t(x_\tau^{\text{OPT}}) &= \frac{1}{2} (x_\tau^{\text{OPT}} - s_k)^2 < \frac{1}{128}. \end{aligned}$$

Combining this with (14),

$$f_t(x_t) - f_t(x_\tau^{\text{OPT}}) > \frac{1}{32} - \frac{1}{128} = \frac{3}{128}. \quad (15)$$

For the preceding N rounds,

$$r = 1, \dots, N,$$

each loss satisfies

$$f_r(x_r) \geq 0$$

and, since $x_\tau^{\text{OPT}}, s_r \in [-1/2, 1/2]$,

$$f_r(x_\tau^{\text{OPT}}) = \frac{1}{2} (x_\tau^{\text{OPT}} - s_r)^2 \leq \frac{1}{2}.$$

Thus

$$f_r(x_r) - f_r(x_\tau^{\text{OPT}}) \geq -\frac{1}{2}, \quad r = 1, \dots, N. \quad (16)$$

Splitting the prefix regret into these two parts and using (16) and (15),

$$\begin{aligned} R_{\mathcal{A}}(\tau) &= \sum_{r=1}^N (f_r(x_r) - f_r(x_\tau^{\text{OPT}})) + \sum_{r=N+1}^{N+32N} (f_r(x_r) - f_r(x_\tau^{\text{OPT}})) \\ &> -\frac{N}{2} + 32N \frac{3}{128} = \frac{N}{4}. \end{aligned} \quad (17)$$

By the assumed prefix-regret guarantee,

$$R_{\mathcal{A}}(\tau) \leq R(\tau) = R(33N).$$

Since

$$R(t) = o(t),$$

we have

$$\frac{R(33N)}{N} = 33 \frac{R(33N)}{33N} \longrightarrow 0 \quad \text{as } N \rightarrow \infty.$$

Hence, there exists N_0 such that, for every $N \geq N_0$,

$$R(33N) < \frac{N}{4}.$$

But from (17), we have that if the total movement during the phase is less than $1/4$, then

$$R_{\mathcal{A}}(33N) > \frac{N}{4}.$$

Therefore, for every phase with

$$N = N_{k-1} \geq N_0,$$

we must have

$$\sum_{t=N+1}^{N+32N} \|x_t - x_{t-1}\| \geq \frac{1}{4}.$$

Since

$$N_{k-1} = 33^{k-1},$$

only finitely many phases satisfy

$$N_{k-1} < N_0.$$

In particular, the number of such phases is at most

$$1 + \lceil \log_{33} N_0 \rceil,$$

which is independent of T . Hence all but finitely many complete phases contribute at least $1/4$ movement, where the number of exceptional phases is independent of T .

Finally, the number of complete phases contained in $[T]$ is

$$K = \lfloor \log_{33} T \rfloor.$$

Therefore, denoting $O(1)$ as a constant for the finitely many phase where we do not have a lower bound on the movement cot,

$$M_{\mathcal{A}}(T) \geq \frac{1}{4}K - O(1) = \Omega(\log T).$$

□

7 Proof of Theorem 3

Proof. Let $g_t \in \partial f_t(x_t)$. Since $x^{\text{OPT}} \in S_T \subseteq S_t$ and f_t is convex,

$$f_t(x_t) - f_t(x^{\text{OPT}}) \leq \langle g_t, x_t - x^{\text{OPT}} \rangle. \quad (18)$$

The first-order optimality condition for algorithm Prox (4) implies

$$\left\langle g_t + \frac{x_t - x_{t-1}}{\eta}, x^{\text{OPT}} - x_t \right\rangle \geq 0,$$

since $x^{\text{OPT}} \in S_t$ for all t . Similar to (9), we have

$$\langle g_t, x_t - x^{\text{OPT}} \rangle \leq \frac{\|x_{t-1} - x^{\text{OPT}}\|^2 - \|x_t - x^{\text{OPT}}\|^2 - \|x_t - x_{t-1}\|^2}{2\eta}.$$

Therefore, from (18), we get

$$f_t(x_t) - f_t(x^{\text{OPT}}) \leq \frac{\|x_{t-1} - x^{\text{OPT}}\|^2 - \|x_t - x^{\text{OPT}}\|^2 - \|x_t - x_{t-1}\|^2}{2\eta}.$$

Dropping the nonpositive last term and summing over t gives

$$\text{Regret}_{\mathcal{A}}(T) \leq \frac{\|x_0 - x^{\text{OPT}}\|^2 - \|x_T - x^{\text{OPT}}\|^2}{2\eta} \leq \frac{D^2}{2\eta} = \frac{\epsilon T}{2}.$$

Using Lemma 2, we know that the algorithm Prox's update (4) is of the form

$$x_t = \Pi_{S_t}(x_{t-1} - e_t), \quad e_t := \eta g_t.$$

Since f_t is G -Lipschitz,

$$\|e_t\| \leq \eta G.$$

Since S_t 's are nested, applying Lemma 3, we get

$$M_{\mathcal{A}}(T) \leq C_d \left(D + \sum_{t=1}^T \|e_t\| \right),$$

and therefore

$$M_{\mathcal{A}}(T) \leq C_d(D + \eta GT) = C_d \left(D + \frac{GD^2}{\epsilon} \right).$$

□

8 Proof of Theorem 4

Proof. Let

$$\mathcal{X} = [-1, 1] \times [0, 1],$$

and the fixed loss function be

$$f(x, y) = -y.$$

Thus, $d = 2$, f is linear and hence convex, and

$$\|\nabla f(x, y)\| = 1,$$

so f is 1-Lipschitz on $\mathcal{X}$.

Let

$$\text{conv}(A)$$

denote the convex hull of a set A , i.e., the smallest convex set containing A .

Let

$$R := \text{Regret}_{\mathcal{A}}(T).$$

Define

$$\delta := 2\sqrt{\frac{R+1}{T}}, \quad K := \left\lfloor \frac{1}{2\delta} \right\rfloor.$$

The assumption

$$T \geq 16(R+1)$$

gives

$$\delta \leq \frac{1}{2}.$$

Hence

$$\frac{1}{2\delta} \geq 1,$$

and therefore

$$K = \left\lfloor \frac{1}{2\delta} \right\rfloor \geq \frac{1}{4\delta}.$$

For $k = 1, \dots, K$, define

$$q_k := 1 - (k-1)\delta, \quad s_k := (-1)^k, \quad z_k := (s_k, q_k),$$

and let

$$c := (0, 1 - K\delta).$$

Since $K\delta \leq 1/2$, we have

$$q_k \geq q_K = 1 - (K-1)\delta > \frac{1}{2},$$

and

$$\frac{1}{2} \leq 1 - K\delta \leq 1.$$

Thus,

$$z_1, \dots, z_K, c \in \mathcal{X}.$$

For $k = 1, \dots, K$, define

$$S_k := \text{conv}\{c, z_k, z_{k+1}, \dots, z_K\}.$$

Then

$$S_1 \supseteq S_2 \supseteq \dots \supseteq S_K.$$

Moreover,

$$q_k > q_{k+1} > \dots > q_K > 1 - K\delta.$$

Hence, the unique minimizer of $f(x, y) = -y$ over S_k is z_k . Thus

$$v_k := \min_{x \in S_k} f(x) = f(z_k) = -q_k.$$

For $k = 1, \dots, K$, define the *trigger* set

$$A_k := \left\{ (x, y) \in S_k : s_k x \geq \frac{1}{2} \right\}.$$

See Fig. 1 for an illustration of this input.

Lemma 4. *For every*

$$p = (x, y) \in S_k \setminus A_k,$$

we have

$$f(p) - f(z_k) > \frac{\delta}{4}.$$

Proof of Lemma 4 is provided at the end. The purpose of Lemma 4 is to show that every action outside the trigger set A_k incurs a definite loss relative to the optimal action z_k in S_k . Thus, an algorithm cannot remain outside A_k for too many consecutive rounds: weak adaptivity forces it to enter A_k after a bounded number of rounds. We will use this to define the




Figure 1: Illustration of the nested feasible sets $S_k = \text{conv}\{c, z_k, \dots, z_K\}$. The points alternate between $x = -1$ and $x = 1$ with $q_1 > q_2 > \dots > q_K$. In the illustrated $K = 5$ case, z_3 and z_5 lie on the left boundary of S_1 , while z_2 and z_4 lie on its right boundary. The dashed lines $x = \pm \frac{1}{2}$ indicate the regions used by the trigger sets $A_k = \{(x, y) \in S_k : s_k x \geq \frac{1}{2}\}$.

input in phases as follows.

Phase k uses the feasible set S_k . The first phase starts at round 1. More generally, if phase k ends at round τ_k , then phase $k + 1$ begins at round $\tau_k + 1$ with feasible set S_{k+1} . Thus, if round t belongs to phase k , we set

$$S_t := S_k.$$

Phase k continues until the first round t for which

$$x_t \in A_k.$$

That round is the final round of phase k .

Let

$$L := \left\lceil \frac{4(R+1)}{\delta} \right\rceil. \quad (19)$$

Suppose, for contradiction, that phase k contains L consecutive rounds without the algorithm $\mathcal{A}$ choosing an action in A_k . Let I_k denote these L rounds, and **let b be the last round of I_k** . Since the feasible set is S_k throughout this phase,

$$x_b^{\text{OPT}} \in \arg \min_{x \in S_b} \sum_{t=1}^b f(x) = \arg \min_{x \in S_k} b f(x) = \{z_k\},$$

and hence

$$x_b^{\text{OPT}} = z_k.$$

Therefore, the weakly adaptive regret guarantee applied to I_k gives

$$\sum_{t \in I_k} (f(x_t) - f(z_k)) \leq R.$$

On the other hand, $x_t \notin A_k$ for every $t \in I_k$, so by Lemma 4,

$$\sum_{t \in I_k} (f(x_t) - f(z_k)) > \frac{\delta |I_k|}{4} = \frac{\delta L}{4} \geq R + 1,$$

where the last inequality follows from the definition of L . This is a contradiction. Hence every phase has length at most L .

It remains to verify that all K phases fit within the horizon. Since each phase contains at most L rounds,

$$KL \leq \frac{1}{2\delta} \left(\frac{4(R+1)}{\delta} + 1 \right).$$

Using

$$\delta^2 = \frac{4(R+1)}{T},$$

we have

$$\frac{2(R+1)}{\delta^2} = \frac{T}{2}.$$

Moreover,

$$\frac{1}{2\delta} \leq \frac{\sqrt{T}}{4} \leq \frac{T}{4},$$

where the first inequality follows from

$$\delta \geq \frac{2}{\sqrt{T}}$$

and the second from $T \geq 1$. Therefore,

$$KL \leq \frac{T}{2} + \frac{T}{4} = \frac{3T}{4} < T.$$

Hence all K phases fit within the horizon; the remaining rounds, if any, are filled by continuing with the feasible set S_K .

Finally, since

$$s_{k+1} = -s_k,$$

we have

$$A_k \subseteq \left\{ (x, y) : s_k x \geq \frac{1}{2} \right\},$$

while

$$A_{k+1} \subseteq \left\{ (x, y) : s_{k+1} x \geq \frac{1}{2} \right\} = \left\{ (x, y) : s_k x \leq -\frac{1}{2} \right\}.$$

Thus,

$$\text{dist}(A_k, A_{k+1}) = \min_{p_1 \in A_k, p_2 \in A_{k+1}} \|p_1 - p_2\| \geq 1.$$

Every phase k ends with an action $x_{\tau_k} \in A_k$. Since the next phase ends with an action $x_{\tau_{k+1}} \in A_{k+1}$ and

$$\text{dist}(A_k, A_{k+1}) \geq 1,$$

we have

$$1 \leq \|x_{\tau_k} - x_{\tau_{k+1}}\| \leq \sum_{t=\tau_k+1}^{\tau_{k+1}} \|x_t - x_{t-1}\|.$$

Thus each pair of consecutive phases contributes at least one unit of movement, and hence

$$M_{\mathcal{A}}(T) \geq K - 1.$$

Using

$$K \geq \frac{1}{4\delta},$$

we obtain

$$M_{\mathcal{A}}(T) \geq \frac{1}{4\delta} - 1 = \frac{1}{8} \sqrt{\frac{T}{R+1}} - 1.$$

In particular, if

$$R = o(T),$$

then

$$\sqrt{\frac{T}{R+1}} \rightarrow \infty.$$

Hence, for all sufficiently large T ,

$$\frac{1}{8} \sqrt{\frac{T}{R+1}} - 1 \geq \frac{1}{16} \sqrt{\frac{T}{R+1}},$$

and therefore

$$M_{\mathcal{A}}(T) = \Omega\left(\sqrt{\frac{T}{R+1}}\right).$$

□

Proof of Lemma 4. Since $p \in S_k$, there exist coefficients

$$\lambda_0, \lambda_k, \dots, \lambda_K \geq 0, \quad \lambda_0 + \sum_{j=k}^K \lambda_j = 1,$$

such that

$$p = \lambda_0 c + \sum_{j=k}^K \lambda_j z_j.$$

Thus, p is a convex combination of c and the points $z_k, \dots, z_K$.

Let

$$w := \lambda_0 + \sum_{\substack{j > k \\ s_j = -s_k}} \lambda_j.$$

This is the total weight assigned to c and to those points z_j with $j > k$ whose first coordinate has the opposite sign from s_k . These are precisely the terms that reduce the quantity $s_k x$.

Since

$$p = \lambda_0 c + \sum_{j=k}^K \lambda_j z_j$$

and

$$c = (0, 1 - K\delta), \quad z_j = (s_j, q_j),$$

the first coordinate of $p = (x, y)$ is

$$x = \lambda_0 \cdot 0 + \sum_{j=k}^K \lambda_j s_j = \sum_{j=k}^K \lambda_j s_j.$$

Multiplying both sides by $s_k \in \{-1, 1\}$ gives

$$s_k x = \sum_{j=k}^K \lambda_j s_k s_j.$$

For $j = k$, we have

$$s_k s_k = 1,$$

so the corresponding term is λ_k . For $j > k$, either $s_j = s_k$, in which case

$$s_k s_j = 1,$$

or $s_j = -s_k$, in which case

$$s_k s_j = -1.$$

Therefore,

$$\begin{aligned} s_k x &= \lambda_k + \sum_{\substack{j>k \\ s_j=s_k}} \lambda_j - \sum_{\substack{j>k \\ s_j=-s_k}} \lambda_j, \\ &= 1 - \lambda_0 - 2 \sum_{\substack{j>k \\ s_j=-s_k}} \lambda_j. \end{aligned}$$

Since $p \notin A_k$,

$$s_k x < \frac{1}{2}.$$

Therefore,

$$\lambda_0 + 2 \sum_{\substack{j>k \\ s_j=-s_k}} \lambda_j > \frac{1}{2}.$$

Since

$$2w = 2\lambda_0 + 2 \sum_{\substack{j>k \\ s_j=-s_k}} \lambda_j \geq \lambda_0 + 2 \sum_{\substack{j>k \\ s_j=-s_k}} \lambda_j,$$

we obtain

$$w > \frac{1}{4}.$$

Every point contributing to w has second coordinate at most $q_k - \delta$. Indeed, for every $j > k$,

$$q_j \leq q_k - \delta,$$

while

$$1 - K\delta \leq q_k - \delta$$

because $k \leq K$. All remaining points in the convex combination have second coordinate at most q_k . Hence

$$\begin{aligned} y &\leq w(q_k - \delta) + (1 - w)q_k \\ &= q_k - w\delta \\ &< q_k - \frac{\delta}{4}. \end{aligned}$$

Since

$$f(p) - f(z_k) = -y + q_k,$$

we conclude that

$$f(p) - f(z_k) > \frac{\delta}{4}.$$

□