

Optimal Alternating Regret for Online Learning and Games*

Yixin Tao[†]

Weiqiang Zheng[‡]

Abstract

We settle the minimax-optimal alternating regret, a regret notion motivated by alternating learning dynamics in games, for both online linear optimization (OLO) and online convex optimization (OCO).

For OLO over the probability simplex Δ_d , we give an algorithm with $O(\log d)$ alternating regret that remains a constant for any time horizon T , and a matching lower bound. Our constant regret bound significantly improves previous results with $O(\log^{2/3} d \cdot T^{1/3})$ regret [Cevher, Cutkosky, Kavis, Piliouras, Skoulakis, Viano, NeurIPS 2023, Hait, Li, Luo, Zhang, COLT 2025]. As a result, we obtain alternating learning dynamics with $O(\log d/T)$ convergence to Nash equilibria in two-player zero-sum games and $O(\log d/T)$ convergence to coarse correlated equilibria in two-player general-sum games. This is the first uncoupled learning dynamics with $O(1/T)$ convergence to CCE in two-player general-sum games, while all prior works suffer additional $\log T$ factors.

For general OCO over a d -dimensional compact convex set, we give an algorithm with $O(d \log(1 + T/d))$ alternating regret, improving the previous best of $\tilde{O}(d^{2/3} T^{1/3})$. We also prove a matching lower bound of $\Omega(d \log(1 + T/d))$, showing that the $\Omega(\log T)$ factor is unavoidable.

Contents

1	Introduction	2
1.1	Our results	3
1.1.1	Optimal alternating regret	4
1.1.2	Fast alternating learning dynamics in games	4
1.2	Further related works	6
2	Constant alternating regret for OLO over the simplex	6
2.1	The alternation-aware Hedge algorithm	6
2.2	A potential proof of $O(\log d)$ alternating regret	7
2.3	An $\Omega(\log d)$ lower bound	8
3	Alternating regret for online convex optimization	9
3.1	The continuous alternation-aware Hedge algorithm	9
3.2	Alternating regret bounds	10

*Authors are alphabetically ordered.

[†]ITCS, Key Laboratory of Interdisciplinary Research of Computation and Economics, Shanghai University of Finance and Economics. Email: taoyixin@mail.shufe.edu.cn

[‡]Yale University. Email: weiqiang.zheng@yale.edu

4	Matching lower bounds for OCO	11
4.1	An $\Omega(\log T)$ lower bound for OCO over 2-dimensional ball	11
4.2	An $\Omega(d \log(1 + T/d))$ lower bound for OCO	14
5	Conclusion	16
A	Proofs for Online Convex Optimization	19
A.1	Existence and feasibility of the update	19
A.2	A potential proof of logarithmic alternating regret	19
A.3	Proof of Theorem 7	20

1 Introduction

We consider the standard online convex optimization (OCO) problem [Zinkevich, 2003, Hazan, 2016, Orabona, 2019]. In each iteration $t \in [T]$, the learner first chooses an action $\mathbf{x}_t \in \mathcal{X}$ from a d -dimensional convex set $\mathcal{X}$, then the adversary chooses a bounded convex loss function $f_t : \mathcal{X} \rightarrow [-1, 1]$ and the learner suffers loss $f_t(\mathbf{x}_t)$.

Regret and learning in games. The classic goal is to minimize the (external) regret defined as

$$\text{Reg}^T := \max_{\mathbf{x} \in \mathcal{X}} \left\{ \text{Reg}^T(\mathbf{x}) = \sum_{t=1}^T f_t(\mathbf{x}_t) - \sum_{t=1}^T f_t(\mathbf{x}) \right\}. \quad (\text{standard regret})$$

If the loss functions $\{f_t(\mathbf{x}) = \mathbf{c}_t^\top \mathbf{x}\}$ are linear, the problem is known as the online linear optimization (OLO) problem. If additionally the action set $\mathcal{X} = \Delta_d$ is the probability simplex, the problem reduces to the classic adversarial d -expert problem. The general OCO algorithm has regret $O(\sqrt{T})$, which is tight by the lower bound of $\Omega(\sqrt{T})$ even for the 2-expert problem.

Algorithms for OCO/OLO have applications for learning in large-scale games and computing approximate equilibria. It is well-known that if players *simultaneously* employ OLO algorithms to choose their strategies, then their time-averaged strategy converges to approximate Nash equilibria (NE) in two-player zero-sum games; their empirical distribution of play converges to approximate coarse correlated equilibria (CCE) in general-sum games [Foster and Vohra, 1997, Freund and Schapire, 1999, Hart and Mas-Colell, 2000, Cesa-Bianchi and Lugosi, 2006]. The approximation depends on the average regret of the algorithm. Thus an algorithm with $\sqrt{T}$ regret in the adversarial setting implies $1/\sqrt{T}$ convergence rate in games. However, better convergence rate is possible in the game setting where every player employs the same algorithm. For two-player zero-sum games, the pioneering work of [Daskalakis, Deckelbaum, and Kim, 2011] show $O(\log T/T)$ convergence rate to NE, while optimistic online learning algorithms achieve the optimal $O(1/T)$ convergence rate [Rakhlin and Sridharan, 2013], which can even be extended to last-iterate convergence [Cai and Zheng, 2023]. For general-sum games, Syrgkanis, Agarwal, Luo, and Schapire [2015] show the first fast convergence rate of $O(1/T^{1/4})$, and a long line of work improve the convergence rate to $O(\log T/T)$ [Chen and Peng, 2020, Daskalakis, Deckelbaum, and Kim, 2011, Anagnostides, Daskalakis, Farina, Fishelson, Golowich, and Sandholm, 2022a, Anagnostides, Farina, Kroer, Lee, Luo, and Sandholm, 2022b, Farina, Anagnostides, Luo, Lee, Kroer, and Sandholm, 2022, Cai, Luo, Wei, and Zheng, 2024, Soleymani, Piliouras, and Farina, 2025b,a], some of which even hold for the stronger notion of correlated equilibrium and general convex games. A very recent work reports a $O(\sqrt{\log T}/T)$ convergence rate [Tsuchiya, 2026]. However, a major problem for learning in games

remains open: can we achieve $O(1/T)$ convergence in general-sum games? This problem was open even for two-player games.

Alternating regret and learning dynamics. In this paper, we focus on *alternation*, a simple trick for *two-player* games where the players take turns to update their strategies (see [Section 1.1.2](#) for a formal definition). Compared to simultaneous learning, alternating learning dynamics have strong empirical performance and are used for training superhuman AI models for poker [[Tammelin, 2014](#), [Bowling, Burch, Johanson, and Tammelin, 2015](#), [Brown and Sandholm, 2018, 2019](#)]. A theoretical explanation of the power of alternation, however, remained elusive until recent work. [Wibisono, Tao, and Piliouras \[2022\]](#) show that alternation with the classic Hedge algorithm [[Freund and Schapire, 1997](#)] gives a fast $O(1/T^{2/3})$ convergence rate in two-player zero-sum games. In their analysis, they propose a new regret notion capturing the nature of alternation, later termed as *alternating regret* by [Cevher, Cutkosky, Kavis, Piliouras, Skoulakis, and Viano \[2023\]](#). In the most general setting of OCO, the alternating regret is defined as follows:

$$\text{Reg}_{\text{Alt}}^T := \max_{\mathbf{x} \in \mathcal{X}} \left\{ \text{Reg}_{\text{Alt}}^T(\mathbf{x}) = \sum_{t=1}^T (f_{t-1}(\mathbf{x}_t) + f_t(\mathbf{x}_t)) - \sum_{t=1}^T (f_{t-1}(\mathbf{x}) + f_t(\mathbf{x})) \right\}. \quad (\text{alternating regret})$$

So the alternating regret can be seen as the standard regret with respect to the loss function $f_{t-1} + f_t$ in each iteration t (define $f_0 = 0$). In alternating regret, the learner knows f_{t-1} when they choose the strategy $\mathbf{x}_t$. This indicates that we may get $o(\sqrt{T})$ alternating regret even in the *adversarial* setting where standard regret suffers $\Omega(\sqrt{T})$ bound.

The work by [Cevher, Cutkosky, Kavis, Piliouras, Skoulakis, and Viano \[2023\]](#) show the first $o(\sqrt{T})$ alternating regret in the adversarial setting. They provide algorithms with $O(\log^{4/3}(dT) \cdot T^{1/3})$ regret for OLO over the probability simplex $\mathcal{X} = \Delta_d$ (i.e., the d -expert problem), and even $O(\log T)$ when $\mathcal{X}$ is the ℓ_2 unit ball. Their results were improved by [Hait, Li, Luo, and Zhang \[2025\]](#), who show that the classic Hedge algorithm already achieves an improved $O(\log^{2/3} d \cdot T^{1/3})$ alternating regret for the d -expert problem. Moreover, [Hait, Li, Luo, and Zhang \[2025\]](#) show that the continuous Hedge algorithm [[Narayanan and Rakhlin, 2010](#)] achieves $\tilde{O}(d^{2/3} T^{1/3})$ alternating regret for the general OCO problem over any convex set $\mathcal{X}$.

However, it is unknown whether $T^{1/3}$ is a fundamental limit for alternating regret. In this paper, we attack the main open question on alternating regret:

what is the minimax-optimal alternating regret for OLO/OCO?

As remarked by [Hait, Li, Luo, and Zhang \[2025\]](#), “*This appears to be highly non-trivial even for very special cases—for example, even for 1-dimensional OLO over $[-1, 1]$ where $O(\log T)$ regret is achieved by [Cevher et al. \[2023\]](#), figuring out whether the optimal bound is $\Theta(\log T)$ or $\Theta(1)$ appears to require new techniques.*”¹

1.1 Our results

In this paper, we settle the minimax-optimal alternating regret for OLO and OCO. We provide algorithms with optimal alternating regret with matching lower bounds, and fast alternating learning dynamics in two-player games.

¹One can easily show that 1-dimensional OLO over $[-1, 1]$ reduces to OLO over Δ_2 (the 2-expert problem).

1.1.1 Optimal alternating regret

Online linear optimization over the probability simplex. We propose an online algorithm with $O(\log d)$ alternating regret for OLO over the probability simplex Δ_d . The algorithm is anytime and does not need to know the time horizon T . It is remarkable that the alternating regret does not grow with T .

Theorem 1 (Constant alternating regret). *For any $T \geq 1$ and adaptive chosen linear loss sequence $\mathbf{c}_1, \dots, \mathbf{c}_T \in [-1, 1]^d$, the alternation-aware Hedge (AA-Hedge) algorithm (Algorithm 1) over Δ_d guarantees that $\text{Reg}_{\text{Alt}}^T \leq 8 \log d$.*

We also show that for any $T \geq d - 1$, an adaptive adversary could force the alternating regret to be at least $\text{Reg}_{\text{Alt}}^T \geq 2 \log d - O(1)$ (Theorem 6). Thus the minimax-optimal alternating regret for the d -expert problem is $\Theta(\log d)$.

The AA-Hedge algorithm (Algorithm 1) is a variant of the Hedge algorithm with non-trivial modifications. It has been shown that the vanilla Hedge algorithm can not achieve $o(T^{1/3})$ alternating regret [Hait, Li, Luo, and Zhang, 2025]. In round t , Algorithm 1 fully exploits the fact that the known loss $\mathbf{c}_{t-1}$ is part of that round's loss. The played strategy $\mathbf{x}_t$ guarantees for any unknown $\mathbf{c}_t$, the instant loss $(\mathbf{c}_{t-1} + \mathbf{c}_t)^\top \mathbf{x}_t$ can be charged on a potential function without any additional terms. A potential analysis similar to that of the Hedge algorithm then concludes the constant alternating regret bound in general.

Online convex optimization. Motivated by the constant regret results for OLO over the simplex, we then study the general setting of online convex optimization over d -dimensional compact convex sets. We recall that the previous best is the $\tilde{O}(d^{2/3}T^{1/3})$ regret by Hait, Li, Luo, and Zhang [2025], achieved by the continuous Hedge algorithm. In Section 3, we extend Algorithm 1 and our analysis to the general online convex optimization setting with convex action set $\mathcal{X}$, using ideas similar to the continuous Hedge algorithm. The resulting continuous AA-Hedge (Algorithm 2) enjoys $O(d \log(1 + T/d))$ alternating regret.

Theorem 2 (Informal, alternating regret upper bound for OCO). *For online convex optimization over any compact convex set $\mathcal{X} \subseteq \mathbb{R}^d$ with any bounded convex losses, the continuous AA-Hedge (Algorithm 2) has $O(d \log(1 + T/d))$ alternating regret.*

For the special case of OLO over a convex polytope with N vertices, the continuous AA-Hedge algorithm enjoys a better alternating regret of $8 \log N$, which also recovers Theorem 1.

Somewhat surprisingly, we show that the $\Omega(\log T)$ dependence is necessary even when the set $\mathcal{X}$ is the 2-dimensional unit ball (Theorem 9). Using this lower bound, we further establish a matching lower bound for OCO over d -dimensional convex sets (Theorem 10).

Theorem 3 (Informal, alternating regret lower bound for OCO). *There exists a compact convex set $\mathcal{X}$ that lies in the d -dimensional unit ℓ_2 ball such that for any OCO algorithm, there exists an oblivious adversary such that the algorithm suffers $\Omega(d \log(1 + T/d))$ alternating regret*

Our upper and lower bounds together establish the minimax-optimal $\Theta(d \log(1 + T/d))$ alternating regret for OCO (Corollary 3).

1.1.2 Fast alternating learning dynamics in games

As an important application, our result implies in alternating learning dynamics with fast convergence in two-player games. Consider a two-player *convex game* with loss functions $u_1, u_2 : \mathcal{X} \times \mathcal{Y} \rightarrow [-1, 1]$

for the x -player and the y -player respectively. The loss $u_1(\mathbf{x}, \mathbf{y})$ is convex in $\mathbf{x}$ for any $\mathbf{y} \in \mathcal{Y}$ and the loss $u_2(\mathbf{x}, \mathbf{y})$ is convex in $\mathbf{y}$ for any $\mathbf{x} \in \mathcal{X}$. The game is zero-sum if $u_1 = -u_2$ and otherwise is general-sum. An important subclass is *normal-form games*.

Example 1 (Normal-form games). *In a two-player normal-form game, $\mathcal{X}$ and $\mathcal{Y}$ are probability simplices and the loss functions are linear: $u_1(\mathbf{x}, \mathbf{y}) = \mathbf{x}^\top A \mathbf{y}$, $u_2(\mathbf{x}, \mathbf{y}) = \mathbf{x}^\top B \mathbf{y}$. The game is zero-sum if $A = -B$, otherwise it is general-sum.*

We recall the definition of approximate Nash equilibrium and coarse correlated equilibrium.

Definition 1 (Nash equilibrium and coarse correlated equilibrium). *A joint strategy profile $(\mathbf{x}, \mathbf{y})$ is an ϵ -Nash equilibrium (ϵ -NE) if $u_1(\mathbf{x}, \mathbf{y}) \leq \min_{\mathbf{x}' \in \mathcal{X}} u_1(\mathbf{x}', \mathbf{y}) + \epsilon$ and $u_2(\mathbf{x}, \mathbf{y}) \leq \min_{\mathbf{y}' \in \mathcal{Y}} u_2(\mathbf{x}, \mathbf{y}') + \epsilon$. A distribution σ over joint strategy profiles $\mathcal{X} \times \mathcal{Y}$ is an ϵ -coarse correlated equilibrium (ϵ -CCE) if $\mathbb{E}_{(\mathbf{x}, \mathbf{y}) \sim \sigma} [u_1(\mathbf{x}, \mathbf{y})] \leq \min_{\mathbf{x}' \in \mathcal{X}} \mathbb{E}_{(\mathbf{x}, \mathbf{y}) \sim \sigma} [u_1(\mathbf{x}', \mathbf{y})] + \epsilon$ and $\mathbb{E}_{(\mathbf{x}, \mathbf{y}) \sim \sigma} [u_2(\mathbf{x}, \mathbf{y})] \leq \min_{\mathbf{y}' \in \mathcal{Y}} \mathbb{E}_{(\mathbf{x}, \mathbf{y}) \sim \sigma} [u_2(\mathbf{x}, \mathbf{y}')] + \epsilon$.*

In a simultaneous learning dynamics, for every iteration $t \geq 1$, both players use an OLO/OCO algorithm to choose their actions $\mathbf{x}_t \in \mathcal{X}$ and $\mathbf{y}_t \in \mathcal{Y}$ simultaneously and then they observe their loss functions $u_1(\cdot, \mathbf{y}_t)$ and $u_2(\mathbf{x}_t, \cdot)$. The *alternating learning dynamics* is slightly different: in iteration t , the x -player chooses $\mathbf{x}_t$ first; then after seeing $\mathbf{x}_t$, the y -player chooses $\mathbf{y}_t$. More specifically, in each round $t \geq 1$,

1. the x -player chooses $\mathbf{x}_t$ based on $u_1(\cdot, \mathbf{y}_1), \dots, u_1(\cdot, \mathbf{y}_{t-1})$;
2. the y -player chooses $\mathbf{y}_t$ based on $u_2(\mathbf{x}_1, \cdot), \dots, u_2(\mathbf{x}_{t-1}, \cdot), u_2(\mathbf{x}_t, \cdot)$;

Algorithms for OLO/OCO with $o(T)$ alternating regret imply convergence to approximate equilibria in two-player games, as observed by [Wibisono, Tao, and Piliouras, 2022, Cevher, Cutkosky, Kavis, Piliouras, Skoulakis, and Viano, 2023] for NE in zero-sum games and by [Hait, Li, Luo, and Zhang, 2025] for CCE in general-sum games. Formally, we have

Theorem 4 (Theorem 1 and 2 in [Hait, Li, Luo, and Zhang, 2025]). *Suppose that in the above alternating learning dynamics, the x -player and the y -player use OCO algorithms with alternating regret bound $\text{Reg}_{\text{Alt}}^T(x)$ and $\text{Reg}_{\text{Alt}}^T(y)$, respectively. Then*

1. *for a zero-sum game, the averaged strategy $(1/T \sum_{t=1}^T \mathbf{x}_t, 1/T \sum_{t=1}^T \mathbf{y}_t)$ is an ϵ -NE with $\epsilon = O((\text{Reg}_{\text{Alt}}^T(x) + \text{Reg}_{\text{Alt}}^T(y))/T)$;*
2. *for a general-sum game, the uniform distribution over $\{(\mathbf{x}_t, \mathbf{y}_t), (\mathbf{x}_{t+1}, \mathbf{y}_{t+1})\}_{t \in [T]}$ is an ϵ -CCE with $\epsilon = O(\max\{\text{Reg}_{\text{Alt}}^T(x), \text{Reg}_{\text{Alt}}^T(y)\}/T)$*

As a corollary, our results imply alternating learning dynamics with fast convergence in games. Notably, we give the first learning dynamics with $O(1/T)$ convergence to CCE in two-player general-sum normal-form games, while existing results suffer additional $\log T$ factors.

Corollary 1 (Fast convergence in two-player games). *For two-player normal-form games where each player has at most d actions, there is an alternating learning dynamics with $O(\log d/T)$ convergence to NE in zero-sum games and $O(\log d/T)$ convergence to CCE in general-sum games.*

1.2 Further related works

Alternation has been studied in two-player zero-sum games and the related min-max optimization problem, mostly in the unconstrained setting where the strategy set is the whole Euclidean space. There is a line of work on the convergence of alternating gradient-descent-ascent in the unconstrained setting [Bailey, Gidel, and Piliouras, 2020, Zhang, Wang, Lessard, and Grosse, 2022, Lee, Cho, and Yun, 2024, Feng, Fujii, Skoulakis, Wang, and Cevher, 2025, Shugart and Altschuler, 2025], the simplex-constrained setting [Nan, Das Gupta, Iyengar, and Kroer, 2026], and manifold setting [Xu, Jiang, Liu, and So, 2026]. The difference is that our work considers alternating regret in the more challenging setting of adversarial online learning, which implies convergence in the game setting.

2 Constant alternating regret for OLO over the simplex

In this section, we focus on the important case of online linear optimization over the simplex, i.e., the adversarial expert problem and present an algorithm with constant alternating regret.

We first introduce some notations. The action set is the d -dimensional probability simplex $\Delta_d = \{\mathbf{x} \in \mathbb{R}^d : x_i \geq 0 \text{ and } \sum_{i=1}^d x_i = 1\}$. We define $\mathbf{c}_0 = \mathbf{0}$ to be the all-0 vector. In each iteration $t \geq 1$, the learner chooses $\mathbf{x}_t \in \Delta_d$ based on the history $\{\mathbf{c}_1, \dots, \mathbf{c}_{t-1}\}$. After observing $\mathbf{x}_t$, the adversary *adaptively* chooses $\mathbf{c}_t \in [-1, 1]^d$. The learner then observes $\mathbf{c}_t$ and suffers loss $(\mathbf{c}_{t-1} + \mathbf{c}_t)^\top \mathbf{x}_t$. In this case, the alternating regret can be written as

$$\text{Reg}_{\text{Alt}}^T = \sum_{t=1}^T (\mathbf{c}_{t-1} + \mathbf{c}_t)^\top \mathbf{x}_t - \min_{\mathbf{x} \in \Delta_d} \sum_{t=1}^T (\mathbf{c}_{t-1} + \mathbf{c}_t)^\top \mathbf{x}. \quad (1)$$

Because the objective is linear, the comparator in (1) is one of the d pure actions.

2.1 The alternation-aware Hedge algorithm

We propose the alternation-aware Hedge algorithm, which introduce a bias toward the most recent known loss vector $\mathbf{c}_{t-1}$ when choosing $\mathbf{x}_t$. The algorithm is in Algorithm 1, where we define $\mathbf{L}_0 = \mathbf{0}$ and $\mathbf{L}_t = \sum_{s=1}^t (\mathbf{c}_{s-1} + \mathbf{c}_s)$ as the effective cumulative loss for $t \geq 1$. We first show that each step in Algorithm 1 is well-defined and feasible: (1) θ_t is unique; (2) the played strategy $\mathbf{x}_t$ lies in the simplex Δ_d .

Algorithm 1 alternation-aware Hedge

- 1: Set $\mathbf{L}_0 = \mathbf{0}$, $\eta = 1/8$, and $\beta = 1/32$.
 - 2: **for** $t = 1, 2, \dots$ **do**
 - 3: Compute $\theta_t = \arg \max_{\theta \in \mathbb{R}} \log \sum_{i=1}^d \exp[-\eta L_{t-1,i} - \beta(c_{t-1,i} - \theta)^2]$
 - 4: Set $\mathbf{p}_t$ such that $p_{t,i} \propto \exp[-\eta L_{t-1,i} - \beta(c_{t-1,i} - \theta_t)^2]$
 - 5: Play $\mathbf{x}_t$ such that $x_{t,i} = p_{t,i} \left(1 - \frac{c_{t-1,i} - \theta_t}{2}\right)$.
 - 6: Observe $\mathbf{c}_t$ and update $\mathbf{L}_t \leftarrow \mathbf{L}_{t-1} + \mathbf{c}_{t-1} + \mathbf{c}_t$.
-

Lemma 1. Assume $\beta < 1/2$. In Algorithm 1, θ_t is unique and $\theta_t = \sum_{i=1}^d p_{t,i} c_{t-1,i}$. Moreover, $\mathbf{x}_t \in \Delta_d$.

Proof. Define $F(\theta) = \log \sum_{i=1}^d \exp[-\eta L_{t-1,i} - \beta(c_{t-1,i} - \theta)^2]$ and $\mathbf{p}(\theta) \in \Delta_d$ such that $p_i(\theta) \propto \exp[-\eta L_{t-1,i} - \beta(c_{t-1,i} - \theta)^2]$. We can calculate both the first-order and the second-order derivative

of $F(\theta)$:

$$\begin{aligned} F'(\theta) &= 2\beta(\mathbb{E}_{\mathbf{p}(\theta)}(\mathbf{c}_{t-1}) - \theta) \\ F''(\theta) &= 4\beta^2 \cdot \text{Var}_{\mathbf{p}(\theta)}(\mathbf{c}_{t-1}) - 2\beta \end{aligned}$$

where we define the expectation $\mathbb{E}_{\mathbf{p}(\theta)}(\mathbf{c}_{t-1}) = \sum_i p_i(\theta)c_{t-1,i}$ and $\text{Var}_{\mathbf{p}(\theta)}(\mathbf{c}_{t-1}) = \sum_i p_i(\theta)(c_{t-1,i} - \mathbb{E}_{\mathbf{p}(\theta)}(\mathbf{c}_{t-1}))^2$. Since $c_{t-1,i} \in [-1, 1]$, the variance term $\text{Var}_{\mathbf{p}(\theta)}(\mathbf{c}_{t-1}) \leq 1$. Since $\beta < \frac{1}{2}$, we have $F''(\theta) < 0$ and F is thus strictly concave. Moreover, $F(\theta) \rightarrow -\infty$ as $|\theta| \rightarrow +\infty$. Thus F has a unique maximizer θ_t and $\mathbf{p}_t = \mathbf{p}(\theta_t)$. The first-order optimality condition $F'(\theta_t) = 0$ implies $\theta_t = \sum_{i=1}^d p_{t,i}c_{t-1,i} \in [\min_i c_{t-1,i}, \max_i c_{t-1,i}]$.

Both $c_{t-1,i}$ and θ_t lie in $[-1, 1]$, so $x_{t,i} = p_{t,i}(1 - (c_{t-1,i} - \theta_t)/2) \geq 0$ for all $i \in [d]$. Moreover, $\theta_t = \sum_{i=1}^d p_{t,i}c_{t-1,i}$ implies $\sum_{i=1}^d x_{t,i} = 1$. Hence $\mathbf{x}_t \in \Delta_d$. $\square$

2.2 A potential proof of $O(\log d)$ alternating regret

We use a potential function argument to bound the alternating regret. Define the alternating regret vector $\mathbf{R}_t$ such that $R_{t,i} = \sum_{s=1}^t (\mathbf{c}_{s-1} + \mathbf{c}_s)^\top \mathbf{x}_s - \sum_{s=1}^t (c_{s-1,i} + c_{s,i})$ for all $i \in [d]$.

Potential function. The potential function is

$$\Phi_t := \max_{\theta \in \mathbb{R}} \log \left(\sum_{i=1}^d \exp[\eta R_{t-1,i} - \beta(c_{t-1,i} - \theta)^2] \right) = \log \left(\sum_{i=1}^d \exp[\eta R_{t-1,i} - \beta(c_{t-1,i} - \theta_t)^2] \right). \quad (2)$$

The equality holds by definition of θ_t in [Algorithm 1](#) and the fact that constant terms do not affect the maximizer. Initially, $\Phi_1 = \log d$.

Theorem 5 (Monotonicity of potential). *Assume $\eta \leq \frac{1}{8}$ and $\beta = \eta/4$. For every round $t \geq 1$, we have $\Phi_{t+1} \leq \Phi_t \leq \log d$.*

Proof. Let us define $\boldsymbol{\xi}_t = \mathbf{c}_{t-1} - \theta_t \mathbf{1}$ and $\boldsymbol{\xi}_{t+1} = \mathbf{c}_t - \theta_{t+1} \mathbf{1}$. Applying [Lemma 1](#) at rounds t and $t+1$ gives $\boldsymbol{\xi}_t, \boldsymbol{\xi}_{t+1} \in [-2, 2]^d$, $\mathbf{p}_t^\top \boldsymbol{\xi}_t = 0$, and the update gives $x_{t,i} = p_{t,i}(1 - \xi_{t,i}/2)$. We have the following identity:

$$R_{t,i} - R_{t-1,i} = (\mathbf{c}_{t-1} + \mathbf{c}_t)^\top \mathbf{x}_t - c_{t-1,i} - c_{t,i} = (\boldsymbol{\xi}_t + \boldsymbol{\xi}_{t+1})^\top \mathbf{x}_t - \xi_{t,i} - \xi_{t+1,i}.$$

By definition of Φ_t and $p_{t,i} \propto \exp(\eta R_{t-1,i} - \beta(c_{t-1,i} - \theta_t)^2)$, we have

$$\begin{aligned} \Phi_{t+1} - \Phi_t &= \log \left(\frac{\sum_{i=1}^d \exp(\eta R_{t,i} - \beta(c_{t,i} - \theta_{t+1})^2)}{\sum_{i=1}^d \exp(\eta R_{t-1,i} - \beta(c_{t-1,i} - \theta_t)^2)} \right) \\ &= \log \left(\sum_{i=1}^d p_{t,i} \exp(\eta(R_{t,i} - R_{t-1,i}) + \beta(\xi_{t,i}^2 - \xi_{t+1,i}^2)) \right) \\ &= \log \left(\sum_{i=1}^d p_{t,i} \exp(\eta((\boldsymbol{\xi}_t + \boldsymbol{\xi}_{t+1})^\top \mathbf{x}_t - \xi_{t,i} - \xi_{t+1,i}) + \beta(\xi_{t,i}^2 - \xi_{t+1,i}^2)) \right). \end{aligned}$$

We then use the following lemma to show that $\Phi_{t+1} \leq \Phi_t$.

Lemma 2. *Assume $\eta \leq \frac{1}{8}$ and $\beta = \eta/4$. Then for any $t \geq 1$, we have*

$$\log \left(\sum_{i=1}^d p_{t,i} \exp(\eta((\boldsymbol{\xi}_t + \boldsymbol{\xi}_{t+1})^\top \mathbf{x}_t - \xi_{t,i} - \xi_{t+1,i}) + \beta(\xi_{t,i}^2 - \xi_{t+1,i}^2)) \right) \leq 0.$$

Proof. Let us define the function $H_t : [-2, 2]^d \rightarrow \mathbb{R}$ as follows:

$$H_t(\mathbf{z}) = \log \sum_{i=1}^d p_{t,i} \exp\left(\eta((\boldsymbol{\xi}_t + \mathbf{z})^\top \mathbf{x}_t - \xi_{t,i} - z_i) + \beta(\xi_{t,i}^2 - z_i^2)\right).$$

It suffices to show $H_t(\boldsymbol{\xi}_{t+1}) \leq 0$.

Define $\mathbf{q}(\mathbf{z})$ be the distribution obtained by normalizing the summands:

$$q_j(\mathbf{z}) = \frac{p_{t,j} \exp\left(\eta((\boldsymbol{\xi}_t + \mathbf{z})^\top \mathbf{x}_t - \xi_{t,j} - z_j) + \beta(\xi_{t,j}^2 - z_j^2)\right)}{\sum_{i=1}^d p_{t,i} \exp\left(\eta((\boldsymbol{\xi}_t + \mathbf{z})^\top \mathbf{x}_t - \xi_{t,i} - z_i) + \beta(\xi_{t,i}^2 - z_i^2)\right)}.$$

We can calculate the gradient $\nabla H_t(\mathbf{z}) = \eta \mathbf{x}_t + \mathbf{q}(\mathbf{z}) \odot (-\eta \mathbf{1} - 2\beta \mathbf{z})$, where we use the point-wise product notation $(\mathbf{a} \odot \mathbf{b})_i = a_i b_i$. We then calculate the Hessian

$$\begin{aligned} \nabla^2 H_t(\mathbf{z}) &= \text{Diag}[q_i(\mathbf{z}) ((-\eta - 2\beta z_i)^2 - 2\beta)] \\ &\quad - (\mathbf{q}(\mathbf{z}) \odot (-\eta \mathbf{1} - 2\beta \mathbf{z})) (\mathbf{q}(\mathbf{z}) \odot (-\eta \mathbf{1} - 2\beta \mathbf{z}))^\top, \end{aligned}$$

where we use notation $\text{Diag}[a_i]$ to denote the square $d \times d$ diagonal matrix whose i -th diagonal is a_i for $i \in [d]$. For any $\mathbf{z} \in [-2, 2]^d$, since $2\beta \geq \max\{(\eta - 4\beta)^2, (\eta + 4\beta)^2\}$, we have $\nabla^2 H_t(\mathbf{z})$ is negative semidefinite. Therefore H_t is concave over $[-2, 2]^d$.

Let us analyze $\mathbf{z} = -\boldsymbol{\xi}_t$. By definition, we have $H_t(-\boldsymbol{\xi}_t) = 0$ and $\mathbf{q}(-\boldsymbol{\xi}_t) = \mathbf{p}_t$. Then the gradient $\nabla H_t(-\boldsymbol{\xi}_t)$ satisfies

$$\begin{aligned} \nabla H_t(-\boldsymbol{\xi}_t) &= \eta \mathbf{x}_t + \mathbf{p}_t \odot (-\eta + 2\beta \boldsymbol{\xi}_t) \\ &= \eta \mathbf{p}_t \odot (1 - \boldsymbol{\xi}_t/2) + \mathbf{p}_t \odot (-\eta + 2\beta \boldsymbol{\xi}_t) \\ &= (2\beta - \eta/2) \mathbf{p}_t \odot \boldsymbol{\xi}_t \\ &= \mathbf{0}. \end{aligned}$$

Thus $-\boldsymbol{\xi}_t$ is the maximizer of H_t over $[-2, 2]^d$ since H_t is concave. Then we have $H_t(\boldsymbol{\xi}_{t+1}) \leq H_t(-\boldsymbol{\xi}_t) = 0$. This completes the proof. $\square$

By [Lemma 2](#), we conclude that $\Phi_{t+1} \leq \Phi_t$ for every $t \geq 1$. $\square$

Proof of [Theorem 1](#). The choices $\eta = 1/8$ and $\beta = 1/32$ in [Algorithm 1](#) satisfy the conditions in [Lemma 1](#) and [Theorem 5](#). Fix any $T \geq 1$. By [Theorem 5](#), we have $\Phi_{T+1} \leq \Phi_1 = \log d$. Fix any pure action $i \in [d]$. By setting $\theta = c_{T,i}$ in [\(2\)](#), we have $\Phi_{T+1} \geq \eta R_{T,i}$. Therefore $R_{T,i} \leq \log d / \eta = 8 \log d$ for every i . Thus the alternating regret is at most $8 \log d$ for any $T \geq 1$. $\square$

2.3 An $\Omega(\log d)$ lower bound

We construct an adaptive adversary such that any algorithm suffers $\Omega(\log d)$ alternating regret. Together with [Theorem 1](#), we show that the minimax-optimal alternating regret for OLO over the simplex is $\Theta(\log d)$. In this lower bound construction, the adversary adaptively chooses the loss function so that the alternating regret in each round $t \in [d-1]$ is at least $2/(d+1-t)$. This then implies the $\Omega(\log d)$ bound.

Theorem 6 (Logarithmic dependence is necessary). *For every $d \geq 2$, every horizon $T \geq d-1$, and every online algorithm, an adaptive adversary can choose linear costs in $[-1, 1]^d$ such that*

$$\text{Reg}_{\text{Alt}}^T \geq 2 \left(\sum_{m=1}^d \frac{1}{m} - 1 \right) = 2 \log d - O(1).$$

Proof. Define $S_0 = [d]$. In round $1 \leq t \leq d-1$, after observing $\mathbf{x}_t$, the adversary computes $j_t \in \arg \max_{j \in S_{t-1}} x_{t,j}$, update $S_t := S_{t-1} \setminus \{j_t\}$, and sets the loss $\mathbf{c}_t$

$$c_{t,i} = \begin{cases} -1, & i \in S_t, \\ +1, & i \notin S_t. \end{cases}$$

The set S_{d-1} contains one action, denoted by i^* . For $t \geq d$, the adversary always chooses $\mathbf{c}_t = \mathbf{c}_{d-1}$.

At $t = 1$, we have $\mathbf{c}_1^\top \mathbf{x}_1 - c_{1,i^*} = 2x_{1,j_1} \geq 2/d$. For $2 \leq t \leq d-1$, the effective cost $\mathbf{g}_t := \mathbf{c}_{t-1} + \mathbf{c}_t$ is

$$g_{t,i} = \begin{cases} -2, & i \in S_t, \\ 0, & i = j_t, \\ +2, & i \notin S_{t-1}. \end{cases}$$

Thus the learner's one-round alternating regret relative to i^* is

$$\mathbf{g}_t^\top \mathbf{x}_t - g_{t,i^*} = 2x_{t,j_t} + 4 \sum_{i \notin S_{t-1}} x_{t,i} \geq 2(1 - \sum_{i \notin S_{t-1}} x_{t,i})/(d+1-t) + 4 \sum_{i \notin S_{t-1}} x_{t,i} \geq \frac{2}{d+1-t},$$

where the first inequality is because $x_{t,j_t} = \max_{j \in S_{t-1}} x_{t,j} \geq (1 - \sum_{i \notin S_{t-1}} x_{t,i})/|S_{t-1}| = (1 - \sum_{i \notin S_{t-1}} x_{t,i})/(d+1-t)$. Therefore, we have

$$\text{Reg}_{\text{Alt}}^{d-1} \geq \sum_{t=1}^{d-1} (\mathbf{g}_t^\top \mathbf{x}_t - g_{t,i^*}) \geq \sum_{t=1}^{d-1} \frac{2}{d+1-t} = 2 \left(\sum_{m=1}^d \frac{1}{m} - 1 \right) = 2 \log d - O(1).$$

Moreover, since $\mathbf{c}_t = \mathbf{c}_{d-1}$ for all $t \geq d$, we have $\text{Reg}_{\text{Alt}}^t \geq \text{Reg}_{\text{Alt}}^{d-1} \geq 2 \log d - O(1)$. $\square$

3 Alternating regret for online convex optimization

In this section, we extend our result to the more general setting of online convex optimization.

Online convex optimization. We first introduce some notations. Let $\mathcal{X} \subseteq \mathbb{R}^d$ be a nonempty compact convex set. We denote by $\Delta_{\mathcal{X}}$ the set of Borel probability measures over $\mathcal{X}$. For $\rho, \mu \in \Delta_{\mathcal{X}}$, we say that μ *dominates* ρ if ρ is absolutely continuous with respect to μ , i.e., $\rho(A) = 0$ for every measurable set A such that $\mu(A) = 0$. We write $\rho \ll \mu$ if μ dominates ρ .

At round $t \geq 1$, the learner chooses $\mathbf{x}_t \in \mathcal{X}$ using $f_1, \dots, f_{t-1}$, and then the adversary adaptively chooses a continuous convex loss function $f_t : \mathcal{X} \rightarrow [-1, 1]$. The alternating regret to $\mathbf{u} \in \mathcal{X}$ is

$$\text{Reg}_{\text{Alt}}^T(\mathbf{u}) := \sum_{t=1}^T (f_{t-1}(\mathbf{x}_t) + f_t(\mathbf{x}_t)) - \sum_{t=1}^T (f_{t-1}(\mathbf{u}) + f_t(\mathbf{u})) = \sum_{t=1}^T (g_t(\mathbf{x}_t) - g_t(\mathbf{u})), \quad (3)$$

where we define $f_0 \equiv 0$ and the effective loss function $g_t := f_{t-1} + f_t$ for $t \geq 1$. For a distribution $\rho \in \Delta_{\mathcal{X}}$, we write $\text{Reg}_{\text{Alt}}^T(\rho) := \mathbb{E}_{\mathbf{u} \sim \rho} [\text{Reg}_{\text{Alt}}^T(\mathbf{u})]$. We also use the shorthand $\mathbb{E}_{\mu}[f] := \mathbb{E}_{\mathbf{x} \sim \mu}[f(\mathbf{x})]$.

3.1 The continuous alternation-aware Hedge algorithm

We propose the continuous alternation-aware Hedge algorithm ([Algorithm 2](#)), a continuous generalization of [Algorithm 1](#). Fix a reference distribution $\mu \in \Delta_{\mathcal{X}}$, and define $L_0 \equiv 0$ and $L_t(\mathbf{u}) := \sum_{s=1}^t g_s(\mathbf{u})$. The algorithm replaces the finite weight vector $\mathbf{p}_t$ by a probability measure P_t , corrects it to a

probability measure Q_t , and plays the barycenter of Q_t . When $\mathcal{X} = \Delta_d$, μ is uniform over the pure actions, and the losses are linear, the algorithm reduces to [Algorithm 1](#).

As in [Algorithm 1](#), we first show that the algorithm is well defined and feasible. The proof is deferred to [Section A](#).

Lemma 3. *Assume $\beta < 1/2$. In [Algorithm 2](#), θ_t is unique and satisfies $\theta_t = \mathbb{E}_{P_t}[f_{t-1}]$. Moreover, Q_t is a probability measure and $\mathbf{x}_t \in \mathcal{X}$.*

Algorithm 2 Continuous alternation-aware Hedge (continuous AA-Hedge)

- 1: Given action set $\mathcal{X}$ and the reference distribution $\mu \in \Delta_{\mathcal{X}}$
- 2: Set $f_0 \equiv 0$, $L_0 \equiv 0$, $\eta = 1/8$, and $\beta = 1/32$.
- 3: **for** $t = 1, 2, \dots$ **do**
- 4: Compute θ_t as the unique maximizer of

$$\log \int_{\mathcal{X}} \exp[-\eta L_{t-1}(\mathbf{u}) - \beta(f_{t-1}(\mathbf{u}) - \theta)^2] \mu(d\mathbf{u}).$$

- 5: Set $\xi_t(\mathbf{u}) = f_{t-1}(\mathbf{u}) - \theta_t$ and form the probability measure

$$P_t(d\mathbf{u}) \propto \exp[-\eta L_{t-1}(\mathbf{u}) - \beta \xi_t(\mathbf{u})^2] \mu(d\mathbf{u}).$$

- 6: Set $Q_t(d\mathbf{u}) = (1 - \xi_t(\mathbf{u})/2) P_t(d\mathbf{u})$.
 - 7: Play the barycenter $\mathbf{x}_t = \int_{\mathcal{X}} \mathbf{u} Q_t(d\mathbf{u})$.
 - 8: Observe f_t and update $L_t \leftarrow L_{t-1} + f_{t-1} + f_t$.
-

3.2 Alternating regret bounds

We first prove an upper bound on the distributional alternating regret $\text{Reg}_{\text{Alt}}^T(\rho)$ for any $\rho \ll \mu$. The alternating regret is at most $O(\text{KL}(\rho \parallel \mu))$ where the KL divergence over measures is defined as $\text{KL}(\rho \parallel \mu) := \int_{\mathcal{X}} \log \frac{d\rho}{d\mu} d\rho$. This is the continuous analogue of comparing with a pure action in [Theorem 1](#). The proof is similar to [Theorem 1](#) and is deferred to [Section A](#).

Theorem 7 (Distributional alternating-regret bound). *For every horizon $T \geq 1$ and every $\rho \ll \mu$, [Algorithm 2](#) guarantees*

$$\text{Reg}_{\text{Alt}}^T(\rho) \leq 8 \text{KL}(\rho \parallel \mu) + \frac{\text{Var}_{\rho}(f_T)}{4} \leq 8 \text{KL}(\rho \parallel \mu) + \frac{1}{4}. \quad (4)$$

Then we instantiate the results with μ being the uniform probability measure over $\mathcal{X}^2$, to get alternating regret bound $\max_{\mathbf{u} \in \mathcal{X}} \text{Reg}_{\text{Alt}}^T(\mathbf{u}) = O(d \log T)$.

Theorem 8 ($O(d \log(1 + T/d))$ alternating regret for OCO). *For every horizon $T \geq 1$, [Algorithm 2](#) with the uniform probability measure $\mu \in \Delta_{\mathcal{X}}$ guarantees*

$$\max_{\mathbf{u} \in \mathcal{X}} \text{Reg}_{\text{Alt}}^T(\mathbf{u}) \leq 8d \left(1 + \log_+ \frac{T}{2d} \right) + \frac{1}{4}, \quad (5)$$

where $\log_+ z := \max\{0, \log z\}$. Since $\max_{\mathbf{u} \in \mathcal{X}} \text{Reg}_{\text{Alt}}^T(\mathbf{u}) \leq 4T$ trivially, we have

$$\max_{\mathbf{u} \in \mathcal{X}} \text{Reg}_{\text{Alt}}^T(\mathbf{u}) = O \left(\min \left\{ T, d \left(1 + \log_+ \frac{T}{2d} \right) \right\} \right) = O(d \log(1 + T/d))$$

²This requires $\mathcal{X} \subseteq \mathbb{R}^d$ to be full-dimensional. If $\mathcal{X}$ is not full-dimensional, let μ be the uniform probability measure over its affine hull and define d as the dimension of its affine.

Proof. Fix $\mathbf{u} \in \mathcal{X}$ and $\delta \in (0, 1)$. Let $\rho_{\mathbf{u}, \delta}$ be the uniform probability measure on the set $(1-\delta)\mathbf{u} + \delta\mathcal{X}$. The affine map $\mathbf{z} \mapsto (1-\delta)\mathbf{u} + \delta\mathbf{z}$ scales d -dimensional volume by δ^d , so $\text{KL}(\rho_{\mathbf{u}, \delta} \parallel \mu) = d \log(1/\delta)$. For any $\mathbf{v} = (1-\delta)\mathbf{u} + \delta\mathbf{z}$ with $\mathbf{z} \in \mathcal{X}$, the convexity of $f_t \in [-1, 1]$ gives

$$f_t(\mathbf{v}) \leq (1-\delta)f_t(\mathbf{u}) + \delta f_t(\mathbf{z}) \leq f_t(\mathbf{u}) + 2\delta.$$

Thus, we have $\mathbb{E}_{\rho_{\mathbf{u}, \delta}}[f_t] - f_t(\mathbf{u}) \leq 2\delta$ for any $t \geq 1$.

Combining the above with [Theorem 7](#), we have

$$\begin{aligned} \text{Reg}_{\text{Alt}}^T(\mathbf{u}) &= \text{Reg}_{\text{Alt}}^T(\rho_{\mathbf{u}, \delta}) + 2 \sum_{t=1}^{T-1} (\mathbb{E}_{\rho_{\mathbf{u}, \delta}}[f_t] - f_t(\mathbf{u})) + (\mathbb{E}_{\rho_{\mathbf{u}, \delta}}[f_T] - f_T(\mathbf{u})). \\ &\leq 8\text{KL}(\rho_{\mathbf{u}, \delta} \parallel \mu) + \frac{1}{4} + 4T\delta \\ &\leq 8d \log(1/\delta) + \frac{1}{4} + 4T\delta. \end{aligned}$$

Choosing $\delta = \min\{1, 2d/T\}$ proves the claim. $\square$

Moreover, for OLO over a polytope $\mathcal{X}$ with N vertices, [Algorithm 2](#) with μ being the uniform distribution over these vertices has alternating regret $O(\log N)$.

Corollary 2 (Constant alternating regret for polyhedral OLO). *Let $\mathcal{X} = \text{conv}\{\mathbf{v}_1, \dots, \mathbf{v}_N\}$ be a polytope. Consider linear losses $f_t(\mathbf{x}) = \langle \mathbf{c}_t, \mathbf{x} \rangle$, with $f_t(\mathbf{x}) \in [-1, 1]$ for every $\mathbf{x} \in \mathcal{X}$. Running [Algorithm 2](#) with μ being the uniform distribution on $\{\mathbf{v}_1, \dots, \mathbf{v}_N\}$ gives*

$$\max_{\mathbf{u} \in \mathcal{X}} \text{Reg}_{\text{Alt}}^T(\mathbf{u}) \leq 8 \log N.$$

In particular, for $\mathcal{X} = \Delta_d$ and $\mathbf{c}_t \in [-1, 1]^d$, [Algorithm 2](#) coincides with [Algorithm 1](#) and recovers the bound $8 \log d$.

Proof. The cumulative comparator loss is linear, so it attains its minimum over the polytope at one of its vertices. Applying [Theorem 7](#) with $\rho = \delta_{\mathbf{v}_j}$ and note that $\text{KL}(\delta_{\mathbf{v}_j} \parallel \mu) = \log N$ and $\text{Var}_{\delta_{\mathbf{v}_j}}(f_T) = 0$. Thus we have $\max_{\mathbf{u} \in \mathcal{X}} \text{Reg}_{\text{Alt}}^T(\mathbf{u}) = \max_{j \in [N]} \text{Reg}_{\text{Alt}}^T(\mathbf{v}_j) \leq 8 \log N$. $\square$

4 Matching lower bounds for OCO

In this section, we prove a alternating regret lower bound of $\Omega(d \log(1 + T/d))$ for OCO with bounded convex losses over a d -dimensional convex set $\mathcal{X} \subseteq B_d(1) := \{\mathbf{x} \in \mathbb{R}^d : \|\mathbf{x}\|_2 \leq 1\}$. In [Section 4.1](#), we first prove a $\Omega(\log T)$ lower bound when the action set is the 2-dimensional unit ball $B_2(1)$ ([Theorem 9](#)). Then, in [Section 4.2](#), we extend the 2-dimensional construction to d -dimensional convex sets and prove the $\Omega(d \log(1 + T/d))$ lower bound.

4.1 An $\Omega(\log T)$ lower bound for OCO over 2-dimensional ball

In this section, we show that the $\log T$ dependence in [Theorem 8](#) is unavoidable. The lower bound already holds on the two-dimensional Euclidean unit ball $B_2(1) := \{\mathbf{x} \in \mathbb{R}^2 : \|\mathbf{x}\|_2 \leq 1\}$. Throughout this section, let $\text{Reg}_{\text{Alt}}^T := \max_{\mathbf{u} \in B_2(1)} \text{Reg}_{\text{Alt}}^T(\mathbf{u})$.

Theorem 9 (Logarithmic lower bound for OCO). *For every integer $T \geq 1$ and every possibly randomized online algorithm $\mathcal{A}$, there is an sequence of continuous convex losses $f_1, \dots, f_T : B_2(1) \rightarrow [0, 1]$ such that*

$$\mathbb{E}_{\mathcal{A}}[\text{Reg}_{\text{Alt}}^T] \geq \frac{1}{125} \lfloor \log_{16} T \rfloor.$$

The expectation is only over the randomness of $\mathcal{A}$.

Proof overview. We in fact prove a stronger lower bound:

$$\mathbb{E}_{\mathcal{A}}[\text{Reg}_{\text{Alt}}^T] \geq \mathbb{E}_{\mathcal{A}} \left[\sum_{t=1}^T f_t(\mathbf{x}_t) - 2 \min_{\mathbf{u} \in B_2(1)} \sum_{t=1}^T f_t(\mathbf{u}) \right] \geq \Omega(\log T).$$

The first inequality holds since the losses are non-negative. The factor 2 in the cumulative loss of the best fixed action is thus not essential. Our lower bound can be easily extended to the regret notion where we compete with $c \cdot \min_{\mathbf{u} \in B_2(1)} \sum_{t=1}^T f_t(\mathbf{u})$ for any constant $c > 0$.

The construction of the loss sequence proceeds in $K = O(\log T)$ blocks. Each block $k \in [K]$ has a center $\mathbf{u}_k$ on the boundary of $B_2(1)$ and an angular scale α_k . Within block k , each round randomizes between a *common loss*, which is minimized at $\mathbf{u}_k$, and a *rare loss*, which is largest at $\mathbf{u}_k$ but vanishes after a rotation by α_k . Fixing a small constant ν , we choose the rare-loss probability $O(\alpha_k^2)$ and the block length proportional to $O(\nu/\alpha_k^2)$. This choice ensures that any algorithm incurs expected loss at least ν in every block (Lemma 4) and thus cumulative loss of at least $\Omega(\nu K)$ (Lemma 5).

After a block, we rotate the center by α_k if the block contains a rare loss, and otherwise leave it unchanged. We maintain geometrically decreasing $\alpha_k = \sqrt{\nu}/4^{k-1}$. We use the endpoint $\mathbf{u}_{K+1}$ after all K rotations in these blocks as the hindsight comparator. We prove that $\mathbf{u}_{K+1}$ has 0 loss on every realized rare, while incurring only $O(\nu^2)$ expected common loss per block (Lemma 6). Thus the (expected) alternating regret of any algorithm is at least $\Omega(\nu K - \nu^2 K) = \Omega(K) = \Omega(\log T)$.

One-round loss pair. Fix a unit vector $\mathbf{u} \in \partial B_2(1)$ and an angle $0 < \alpha \leq 1$. Define

$$\begin{aligned} \ell_{\mathbf{u}}(\mathbf{x}) &:= \frac{1 - \langle \mathbf{u}, \mathbf{x} \rangle}{2}, \\ h_{\mathbf{u}, \alpha}(\mathbf{x}) &:= \max \left\{ 0, \frac{\langle \mathbf{u}, \mathbf{x} \rangle - \cos \alpha}{1 - \cos \alpha} \right\}, \\ p_{\alpha} &:= \frac{1 - \cos \alpha}{3 - \cos \alpha}. \end{aligned}$$

We refer to $\ell_{\mathbf{u}}$ as the common loss and $h_{\mathbf{u}, \alpha}$ as the rare loss. We will randomly choose the common loss $\ell_{\mathbf{u}}$ with probability $1 - p_{\alpha}$ and $h_{\mathbf{u}, \alpha}$ with probability p_{α} . We first present some useful properties of these loss functions.

Lemma 4 (Properties of one-round loss pair). *The common loss $\ell_{\mathbf{u}}$ and the rare loss $h_{\mathbf{u}, \alpha}$ are continuous and convex and take values in $[0, 1]$. Moreover, $h_{\mathbf{u}, \alpha}(\mathbf{u}) = 1$, and $h_{\mathbf{u}, \alpha}(\mathbf{v}) = 0$ whenever $\langle \mathbf{u}, \mathbf{v} \rangle \leq \cos \alpha$. For every $\mathbf{x} \in B_2(1)$, we have*

$$(1 - p_{\alpha})\ell_{\mathbf{u}}(\mathbf{x}) + p_{\alpha}h_{\mathbf{u}, \alpha}(\mathbf{x}) \geq p_{\alpha}.$$

Finally, $\alpha^2/9 \leq p_{\alpha} \leq \alpha^2/4$

Proof. The common loss $\ell_{\mathbf{u}}$ is affine, and the rare loss $h_{\mathbf{u},\alpha}$ is the maximum of two affine functions. Both are therefore continuous and convex. Since $\langle \mathbf{u}, \mathbf{x} \rangle \in [-1, 1]$, both take values in $[0, 1]$. By definition, we can verify $h_{\mathbf{u},\alpha}(\mathbf{u}) = 1$, and $h_{\mathbf{u},\alpha}(\mathbf{v}) = 0$ whenever $\langle \mathbf{u}, \mathbf{v} \rangle \leq \cos \alpha$.

It remains to prove the two inequalities. The definition of p_α gives

$$\frac{1 - p_\alpha}{2} = \frac{p_\alpha}{1 - \cos \alpha} = \frac{1}{3 - \cos \alpha}.$$

If $\langle \mathbf{u}, \mathbf{x} \rangle \leq \cos \alpha$, then the rare loss is zero and the common-loss term is at least p_α . Otherwise, the two terms sum to

$$\frac{1 - \langle \mathbf{u}, \mathbf{x} \rangle}{3 - \cos \alpha} + \frac{\langle \mathbf{u}, \mathbf{x} \rangle - \cos \alpha}{3 - \cos \alpha} = p_\alpha.$$

Finally, for $0 < \alpha \leq 1$, we have $\alpha^2/3 \leq 1 - \cos \alpha \leq \alpha^2/2$ and $2 \leq 3 - \cos \alpha \leq 3$, which proves the bounds on p_α . $\square$

Block construction. Fix a positive integer K and set $\nu := 10^{-2}$. Start from $\mathbf{u}_1 = (1, 0)$ and, for $k = 1, \dots, K$, set

$$\alpha_k := \frac{\sqrt{\nu}}{4^{k-1}}, \quad p_k := p_{\alpha_k}, \quad n_k := \left\lceil \frac{\nu}{p_k} \right\rceil.$$

We construct the loss sequence as follows: for each block k with n_k rounds

1. We independently choose the rare loss $h_{\mathbf{u}_k, \alpha_k}$ with probability p_k on each of the n_k rounds, and choose the common loss $\ell_{\mathbf{u}_k}$ otherwise.
2. If no rare loss was sampled in block k , set $\mathbf{u}_{k+1} \leftarrow \mathbf{u}_k$. Otherwise at least one rare loss was sampled in block k , we update $\mathbf{u}_{k+1}$ by rotating $\mathbf{u}_k$ counterclockwise through α_k . That is, if $\mathbf{u}_k = (\cos(\phi), \sin(\phi))$, then update $\mathbf{u}_{k+1} \leftarrow (\cos(\phi + \alpha_k), \sin(\phi + \alpha_k))$.

Let $N_K = \sum_{k=1}^K n_k$. We note that this is an oblivious randomized adversary. We use $\mathbb{E}_{\text{loss}}$ for expectation over the construction and $\mathbb{E}_{\text{loss}, \mathcal{A}}$ for expectation over both the construction and the randomness of $\mathcal{A}$.

Lemma 5 (Cost of the algorithm). *For every online algorithm $\mathcal{A}$,*

$$\mathbb{E}_{\text{loss}, \mathcal{A}} \left[\sum_{t=1}^{N_K} f_t(\mathbf{x}_t) \right] \geq K\nu.$$

Proof. Consider a round in block k and condition on the past and on the private randomness used to choose $\mathbf{x}_t$. This fixes $\mathbf{x}_t$ and $\mathbf{u}_k$, while the current common-or-rare choice remains independent. By Lemma 4, the conditional expected loss is at least p_k . Summing over the blocks gives

$$\mathbb{E}_{\text{loss}, \mathcal{A}} \left[\sum_{t=1}^{N_K} f_t(\mathbf{x}_t) \right] \geq \sum_{k=1}^K n_k p_k \geq K\nu. \quad \square$$

Lemma 6 (Cost of the endpoint). *The endpoint of the block construction $\mathbf{u}_\star := \mathbf{u}_{K+1}$ satisfies*

$$\mathbb{E}_{\text{loss}} \left[\sum_{t=1}^{N_K} f_t(\mathbf{u}_\star) \right] \leq 10K\nu^2.$$

Proof. Let I_j indicate whether block j contains a rare loss. By the union bound and the definition of n_j ,

$$\Pr(I_j = 1) = 1 - (1 - p_j)^{n_j} \leq n_j p_j \leq \nu + p_j < 2\nu,$$

where the last inequality uses $p_j \leq \alpha_j^2/4 \leq \nu/4$ by [Lemma 4](#) and the definition $\alpha_j = \sqrt{\nu}/4^{j-1}$.

Define counterclockwise angle from $\mathbf{u}_k$ to $\mathbf{u}_\star$ as $\Delta_k := \sum_{j=k}^K I_j \alpha_j$. By definition of α_k , we have

$$0 \leq \Delta_k \leq \sum_{j=k}^{\infty} \alpha_j = \frac{4\alpha_k}{3} < \pi.$$

If block k contains a rare loss, then $\Delta_k \geq \alpha_k$. Thus $\langle \mathbf{u}_k, \mathbf{u}_\star \rangle = \cos \Delta_k \leq \cos \alpha_k$ and the rare loss $h_{\mathbf{u}_k, \alpha_k}(\mathbf{u}_\star) = 0$ is zero ([Lemma 4](#)). Each common loss in the block is at most $\ell_{\mathbf{u}_k}(\mathbf{u}_\star) = \frac{1 - \cos \Delta_k}{2} \leq \frac{\Delta_k^2}{4}$. Since $\Delta_k \leq 4\alpha_k/3$, we have

$$\mathbb{E}_{\text{loss}}[\Delta_k^2] \leq \frac{4\alpha_k}{3} \mathbb{E}_{\text{loss}}[\Delta_k] = \frac{4\alpha_k}{3} \sum_{j=k}^K \alpha_j \Pr(I_j = 1) < \frac{4\alpha_k}{3} \cdot 2\nu \cdot \frac{4\alpha_k}{3} < 4\nu\alpha_k^2.$$

Then by $p_k \geq \alpha_k^2/9$ ([Lemma 4](#)) and $\alpha_k^2 \leq \nu$, the expected loss of $\mathbf{u}_\star$ in block k is at most

$$n_k \mathbb{E}_{\text{loss}}[\ell_{\mathbf{u}_k}(\mathbf{u}_\star)] \leq \frac{n_k}{4} \mathbb{E}_{\text{loss}}[\Delta_k^2] \leq n_k \nu \alpha_k^2 \leq \nu \left(\frac{\nu}{p_k} + 1 \right) \alpha_k^2 \leq \nu(9\nu + \alpha_k^2) \leq 10\nu^2.$$

Summing over the blocks proves the claim. $\square$

Proof of Theorem 9. We assume $K := \lfloor \log_{16} T \rfloor \geq 1$ since otherwise the claim is trivial. We use the loss construction described above. Since $\alpha_k^2 = \nu/16^{k-1}$, [Lemma 4](#) gives $n_k \leq \nu/p_k + 1 \leq 9\nu/\alpha_k^2 + 1 \leq 9 \cdot 16^{k-1} + 1$. Consequently,

$$N_K = \sum_{k=1}^K n_k \leq \frac{9(16^K - 1)}{15} + K \leq 16^K \leq T.$$

We pad zero losses for rounds $N_K < t \leq T$.

For every realization of the loss construction and of the algorithm's randomness, we evaluate alternating regret at $\mathbf{u}_\star$. Since all losses are nonnegative and each comparator loss appears at most twice, we have

$$\text{Reg}_{\text{Alt}}^T \geq \sum_{t=1}^{N_K} f_t(\mathbf{x}_t) - 2 \sum_{t=1}^{N_K} f_t(\mathbf{u}_\star).$$

Applying [Lemmas 5](#) and [6](#) and using $\nu = 10^{-2}$ gives

$$\mathbb{E}_{\text{loss}} [\mathbb{E}_{\mathcal{A}} [\text{Reg}_{\text{Alt}}^T \mid f_1, \dots, f_T]] = \mathbb{E}_{\text{loss}, \mathcal{A}} [\text{Reg}_{\text{Alt}}^T] \geq K\nu - 20K\nu^2 = \frac{K}{125}.$$

Thus some fixed realization of the loss sequence satisfies the desired bound. $\square$

4.2 An $\Omega(d \log(1 + T/d))$ lower bound for OCO

[Theorem 9](#) shows that the $\log T$ factor in [Theorem 8](#) is unavoidable even for 2-dimensional unit ball. We now show that the linear dependence on d is also necessary in the worst case over a convex set $\mathcal{X} \subseteq B_d(1)$ and prove the optimal lower bound of $\Omega(d \log(1 + T/d))$.

High-level idea: Roughly speaking, we construct the losses in phases, where in each phase we simulate the lower bound construction in [Theorem 9](#) on 2 coordinates. There are $d/2$ phases and each phase contains $m = \Theta(T/d)$ rounds. By [Theorem 9](#), the alternating regret in each phase is at least $\Omega(\log(1 + T/d))$. Thus the total alternating regret is at least $\Omega(d \log(1 + T/d))$.

Let $r := 1/\sqrt{d}$. Define

$$\mathcal{X} := \underbrace{rB_2(1) \times \cdots \times rB_2(1)}_{\lfloor d/2 \rfloor \text{ times}} \times [-r, r]^{d-2\lfloor d/2 \rfloor} \subseteq \mathbb{R}^d,$$

where $d - 2\lfloor d/2 \rfloor \in \{0, 1\}$. The set $\mathcal{X}$ is compact, convex, and full-dimensional. Moreover, every $\mathbf{x} \in \mathcal{X}$ satisfies

$$\|\mathbf{x}\|_2^2 \leq \lfloor d/2 \rfloor r^2 + (d - 2\lfloor d/2 \rfloor) r^2 = (d - \lfloor d/2 \rfloor) r^2 \leq 1,$$

so $\mathcal{X} \subseteq B_d(1)$. For $j \in [\lfloor d/2 \rfloor]$, let $\pi_j : \mathcal{X} \rightarrow B_2(1)$ be the projection onto the j -th two-dimensional block, rescaled by $1/r$. Thus π_j is linear and $\pi_j(\mathcal{X}) = B_2(1)$. Throughout this subsection, let $\text{Reg}_{\text{Alt}}^T := \max_{\mathbf{u} \in \mathcal{X}} \text{Reg}_{\text{Alt}}^T(\mathbf{u})$.

Theorem 10 (Linear dimension dependence is necessary). *For every $d \geq 2$, every horizon $T \geq 16$, and every possibly randomized online algorithm $\mathcal{A}$, there is a sequence of continuous convex losses $f_1, \dots, f_T : \mathcal{X} \rightarrow [0, 1]$ such that*

$$\mathbb{E}_{\mathcal{A}} [\text{Reg}_{\text{Alt}}^T] \geq \frac{mK}{125} \quad \text{where} \quad m := \min \left\{ \left\lfloor \frac{d}{2} \right\rfloor, \left\lfloor \frac{T}{16} \right\rfloor \right\}, \quad K := \left\lfloor \log_{16} \left\lfloor \frac{T}{m} \right\rfloor \right\}.$$

In particular $\mathbb{E}_{\mathcal{A}} [\text{Reg}_{\text{Alt}}^T] = \Omega(d \log(1 + T/d))$.

Proof. Set $T' := \lfloor T/m \rfloor$. Since $m \leq \lfloor T/16 \rfloor$, we have $T' \geq 16$ and $K = \lfloor \log_{16} T' \rfloor \geq 1$. Since $m \leq \lfloor d/2 \rfloor$, we use the first mT' rounds as m consecutive phases of T' rounds, one for each of the first m two-dimensional blocks.

We construct the losses in these phases successively. For $j \in [m]$, suppose the losses in the first $j-1$ phases have been fixed. This fixed prefix induces a distribution over $\mathcal{A}$'s internal state. Starting from this state, simulate $\mathcal{A}$ for T' more rounds: return its decisions projected by π_j , and feed each two-dimensional loss back to $\mathcal{A}$ after lifting it through π_j . This defines a possibly randomized online algorithm on $B_2(1)$. Inspecting the proof of [Theorem 9](#), a fixed sequence of continuous convex losses $\tilde{f}_1^{(j)}, \dots, \tilde{f}_{T'}^{(j)} : B_2(1) \rightarrow [0, 1]$ can be chosen for this projected algorithm. Let $\mathbf{u}_\star^{(j)} \in B_2(1)$ minimize $\sum_{i=1}^{T'} \tilde{f}_i^{(j)}(\mathbf{u})$. Then

$$\mathbb{E}_{\mathcal{A}} \left[\sum_{i=1}^{T'} \tilde{f}_i^{(j)}(\pi_j(\mathbf{x}_{(j-1)T'+i})) - 2 \sum_{i=1}^{T'} \tilde{f}_i^{(j)}(\mathbf{u}_\star^{(j)}) \right] \geq \frac{K}{125}.$$

For $t = (j-1)T' + i$, set $f_t := \tilde{f}_i^{(j)} \circ \pi_j$. After all m phases have been fixed, set $f_t \equiv 0$ for $mT' < t \leq T$. Each phase is fixed for the randomized algorithm induced by the preceding phases, not for a realization of $\mathcal{A}$'s random choices. Thus the resulting loss sequence is deterministic and oblivious. Since π_j is linear, every f_t is continuous, convex, and takes values in $[0, 1]$.

Let $\mathbf{u}_\star \in \mathcal{X}$ have j -th two-dimensional block $r\mathbf{u}_\star^{(j)}$ for $j \in [m]$ and all remaining coordinates equal to zero. Then $\pi_j(\mathbf{u}_\star) = \mathbf{u}_\star^{(j)}$ for every $j \in [m]$. Since the losses are nonnegative, we may discard the terms $f_{t-1}(\mathbf{x}_t)$ from the learner's loss, while every comparator loss appears at most twice. Therefore,

$$\text{Reg}_{\text{Alt}}^T \geq \sum_{t=1}^T f_t(\mathbf{x}_t) - 2 \sum_{t=1}^T f_t(\mathbf{u}_\star) = \sum_{j=1}^m \left[\sum_{i=1}^{T'} \tilde{f}_i^{(j)}(\pi_j(\mathbf{x}_{(j-1)T'+i})) - 2 \sum_{i=1}^{T'} \tilde{f}_i^{(j)}(\mathbf{u}_\star^{(j)}) \right].$$

Taking expectation and summing the phase guarantees gives

$$\mathbb{E}_{\mathcal{A}}[\text{Reg}_{\text{Alt}}^T] \geq \frac{mK}{125}.$$

It remains to verify the asymptotic bound. If $T \geq 8d$, then $m = \lfloor d/2 \rfloor \geq d/3$ and $T' = \lfloor T/m \rfloor \geq T/(2m) \geq T/d$. Since $T' \geq 16$, we have $K = \Omega(\log(1 + T')) = \Omega(\log(1 + T/d))$. Hence $mK = \Omega(d \log(1 + T/d))$. If $T < 8d$, then $m = \lfloor T/16 \rfloor \geq T/32$ and $K \geq 1$. Since $\log(1 + x) \leq x$, we have $mK \geq T/32 \geq d \log(1 + T/d) / 32$. Thus we conclude $\mathbb{E}_{\mathcal{A}}[\text{Reg}_{\text{Alt}}^T] = \Omega(d \log(1 + T/d))$. $\square$

Combining [Theorem 10](#) with [Theorem 8](#) pins down the worst-case alternating regret over d -dimensional bodies in every regime of T and d .

Corollary 3 (Minimax-optimal alternating regret for OCO). *Let $\mathcal{R}^*(d, T) := \sup_{\mathcal{X}} \inf_{\mathcal{A}} \sup_{f_1, \dots, f_T} \text{Reg}_{\text{Alt}}^T$, where the outer supremum is over compact convex $\mathcal{X} \subseteq \mathbb{R}^d$ and the losses are convex with values in $[-1, 1]$. Then, for all $d \geq 2$ and $T \geq 16$,*

$$\mathcal{R}^*(d, T) = \Theta\left(d \log\left(1 + \frac{T}{d}\right)\right).$$

5 Conclusion

In this paper, we settle the minimax-optimal alternating regret for both OLO and OCO. For OLO over the simplex Δ_d , we give the alternation-aware Hedge (AA-Hedge) algorithm with optimal $O(\log d)$ regret. For OCO, we give the continuous AA-Hedge algorithm with $O(d \log(1 + T/d))$ regret and proves a $\Omega(d \log(1 + T/d))$ lower bound. Our results imply the first alternating learning dynamics with $O(1/T)$ convergence to CCE in two-player general-sum games.

Acknowledgement

The proofs were first obtained with the assistance of ChatGPT 5.6 Sol and Claude Opus 5. The authors subsequently verified and significantly rewrote the proofs for better exposition.

References

- I. Anagnostides, C. Daskalakis, G. Farina, M. Fishelson, N. Golowich, and T. Sandholm. Near-optimal no-regret learning for correlated equilibria in multi-player general-sum games. In *Proceedings of the 54th Annual ACM SIGACT Symposium on Theory of Computing (STOC)*, 2022a.
- I. Anagnostides, G. Farina, C. Kroer, C.-W. Lee, H. Luo, and T. Sandholm. Uncoupled learning dynamics with $o(\log t)$ swap regret in multiplayer games. In *Proceedings of the Annual Conference on Neural Information Processing Systems (NeurIPS)*, 2022b.
- J. P. Bailey, G. Gidel, and G. Piliouras. Finite regret and cycles with fixed step-size via alternating gradient descent-ascent. In *Conference on Learning Theory*, pages 391–407, 2020.
- M. Bowling, N. Burch, M. Johanson, and O. Tammelin. Heads-up limit hold'em poker is solved. *Science*, 347(6218):145–149, 2015.
- N. Brown and T. Sandholm. Superhuman AI for heads-up no-limit poker: Libratus beats top professionals. *Science*, 359(6374):418–424, 2018.

- N. Brown and T. Sandholm. Superhuman AI for multiplayer poker. *Science*, 365(6456):885–890, 2019.
- Y. Cai and W. Zheng. Doubly optimal no-regret learning in monotone games. In *International Conference on Machine Learning (ICML)*, pages 3507–3524, 2023.
- Y. Cai, H. Luo, C.-Y. Wei, and W. Zheng. Near-optimal policy optimization for correlated equilibrium in general-sum markov games. In *International Conference on Artificial Intelligence and Statistics*, pages 3889–3897, 2024.
- N. Cesa-Bianchi and G. Lugosi. *Prediction, Learning, and Games*. Cambridge university press, 2006.
- V. Cevher, A. Cutkosky, A. Kavis, G. Piliouras, S. Skoulakis, and L. Viano. Alternation makes the adversary weaker in two-player games. In *Advances in Neural Information Processing Systems*, volume 36, 2023. doi: 10.52202/075280-0804. URL https://proceedings.neurips.cc/paper_files/paper/2023/hash/3acb49252187efa352a1ae0e4b066ced-Abstract-Conference.html.
- X. Chen and B. Peng. Hedging in games: Faster convergence of external and swap regrets. In *Proceedings of the Annual Conference on Neural Information Processing Systems (NeurIPS)*, 2020.
- C. Daskalakis, A. Deckelbaum, and A. Kim. Near-optimal no-regret algorithms for zero-sum games. In *Proceedings of the twenty-second annual ACM-SIAM symposium on Discrete Algorithms*, pages 235–254. SIAM, 2011.
- G. Farina, I. Anagnostides, H. Luo, C.-W. Lee, C. Kroer, and T. Sandholm. Near-optimal no-regret learning dynamics for general convex games. In *Proceedings of the Annual Conference on Neural Information Processing Systems (NeurIPS)*, 2022.
- Y. Feng, K. Fujii, S. Skoulakis, X. Wang, and V. Cevher. Continuous-time analysis of heavy ball momentum in min-max games. In *Forty-second International Conference on Machine Learning*, 2025. URL <https://openreview.net/forum?id=7vdg7VICsX>.
- D. P. Foster and R. V. Vohra. Calibrated learning and correlated equilibrium. *Games and Economic Behavior*, 21(589):40–55, 1997.
- Y. Freund and R. E. Schapire. A decision-theoretic generalization of on-line learning and an application to boosting. *Journal of computer and system sciences*, 55(1):119–139, 1997.
- Y. Freund and R. E. Schapire. Adaptive game playing using multiplicative weights. *Games and Economic Behavior*, 29(1-2):79–103, 1999.
- S. Hait, P. Li, H. Luo, and M. Zhang. Alternating regret for online convex optimization. In *Proceedings of the Thirty-Eighth Conference on Learning Theory*, volume 291 of *Proceedings of Machine Learning Research*, pages 2632–2633. PMLR, 2025. URL <https://proceedings.mlr.press/v291/hait25a.html>.
- S. Hart and A. Mas-Colell. A simple adaptive procedure leading to correlated equilibrium. *Econometrica*, 68(5):1127–1150, 2000.
- E. Hazan. Introduction to online convex optimization. *Foundations and Trends® in Optimization*, 2(3-4):157–325, 2016.

- J. Lee, H. Cho, and C. Yun. Fundamental benefit of alternating updates in minimax optimization. In *Forty-first International Conference on Machine Learning*, 2024. URL <https://openreview.net/forum?id=s6ZAT8MLKU>.
- T. Nan, S. Das Gupta, G. Iyengar, and C. Kroer. On the $o(1/t)$ convergence of alternating gradient descent–ascent in bilinear games. In *International Conference on Learning Representations*, volume 2026, pages 79281–79320, 2026.
- H. Narayanan and A. Rakhlin. Random walk approach to regret minimization. *Advances in Neural Information Processing Systems*, 23, 2010.
- F. Orabona. A modern introduction to online learning. *arXiv preprint arXiv:1912.13213*, 2019.
- A. Rakhlin and K. Sridharan. Online learning with predictable sequences. In *Proceedings of the 26th Annual Conference on Learning Theory*, volume 30 of *Proceedings of Machine Learning Research*, pages 993–1019. PMLR, 2013. URL <https://proceedings.mlr.press/v30/Rakhlin13.html>.
- H. Shugart and J. M. Altschuler. Negative stepsizes make gradient-descent-ascent converge. *arXiv preprint arXiv:2505.01423*, 2025.
- A. Soleymani, G. Piliouras, and G. Farina. Cautious optimism: A meta-algorithm for near-constant regret in general games. In *Proceedings of the 26th ACM Conference on Economics and Computation*, pages 870–870, 2025a.
- A. Soleymani, G. Piliouras, and G. Farina. Faster rates for no-regret learning in general games via cautious optimism. In *Proceedings of the 57th Annual ACM Symposium on Theory of Computing*, pages 518–529, 2025b.
- V. Syrgkanis, A. Agarwal, H. Luo, and R. E. Schapire. Fast convergence of regularized learning in games. In *Proceedings of the Annual Conference on Neural Information Processing Systems (NeurIPS)*, 2015.
- O. Tammelin. Solving large imperfect information games using cfr+. *arXiv preprint arXiv:1407.5042*, 2014.
- T. Tsuchiya. Sublogarithmic swap regret in multiplayer general-sum games via hybrid regularization. *arXiv preprint arXiv:2608.04149*, 2026.
- A. Wibisono, M. Tao, and G. Piliouras. Alternating mirror descent for constrained min–max games. In *Advances in Neural Information Processing Systems*, volume 35, 2022. doi: 10.52202/068431-2551. URL https://proceedings.neurips.cc/paper_files/paper/2022/hash/e496e0ce207ba9cdcc7d79bd499db67e-Abstract-Conference.html.
- M. Xu, B. Jiang, Y.-F. Liu, and A. M.-C. So. A riemannian alternating descent ascent algorithmic framework for nonconvex-linear minimax problems on riemannian manifolds. *Mathematics of Operations Research*, 2026.
- G. Zhang, Y. Wang, L. Lessard, and R. B. Grosse. Near-optimal local convergence of alternating gradient descent-ascent for minimax optimization. In *International Conference on Artificial Intelligence and Statistics*, pages 7659–7679. PMLR, 2022.
- M. Zinkevich. Online convex programming and generalized infinitesimal gradient ascent. In *Proceedings of the 20th international conference on machine learning (ICML)*, 2003.

A Proofs for Online Convex Optimization

The proof follows the same strategy as in [Section 2](#): we first show that the update is well defined and feasible, and then use a one-step inequality to prove that a potential function is nonincreasing.

A.1 Existence and feasibility of the update

Proof of [Lemma 3](#). Define

$$F_t(\theta) := \log \int_{\mathcal{X}} \exp[-\eta L_{t-1}(\mathbf{u}) - \beta(f_{t-1}(\mathbf{u}) - \theta)^2] \mu(d\mathbf{u}),$$

and let $P_{t,\theta}$ be the probability measure obtained by normalizing the integrand. F_t is differentiable since it is logarithm of a bounded integrand over a compact set. We calculate its first-order and second-order derivative:

$$\begin{aligned} F'_t(\theta) &= 2\beta (\mathbb{E}_{P_{t,\theta}}[f_{t-1}] - \theta), \\ F''_t(\theta) &= 4\beta^2 \text{Var}_{P_{t,\theta}}(f_{t-1}) - 2\beta. \end{aligned}$$

Since $f_{t-1} \in [-1, 1]$, the variance is at most 1. Since $\beta < 1/2$, we have $F''_t(\theta) \leq 4\beta^2 - 2\beta < 0$, so F_t is strictly concave. Moreover, $F_t(\theta) \rightarrow -\infty$ as $|\theta| \rightarrow \infty$, so the maximizer exists and is unique. The first-order condition gives

$$\theta_t = \mathbb{E}_{P_t}[f_{t-1}] \in \left[\min_{\mathbf{u} \in \text{supp}(\mu)} f_{t-1}(\mathbf{u}), \max_{\mathbf{u} \in \text{supp}(\mu)} f_{t-1}(\mathbf{u}) \right].$$

Thus $\mathbb{E}_{P_t}[\xi_t] = 0$ and $\xi_t(\mathbf{u}) \in [-2, 2]$ on $\text{supp}(P_t)$. It follows that $1 - \xi_t(\mathbf{u})/2 \geq 0$ and

$$Q_t(\mathcal{X}) = 1 - \frac{1}{2} \mathbb{E}_{P_t}[\xi_t] = 1.$$

Hence Q_t is a probability measure. Since $\mathcal{X}$ is compact and convex, the barycenter of Q_t , $\mathbf{x}_t$, lies in $\mathcal{X}$. $\square$

A.2 A potential proof of logarithmic alternating regret

Define the learner's cumulative effective loss through round $t-1$ by $A_{t-1} := \sum_{s=1}^{t-1} g_s(\mathbf{x}_s)$, and define its pointwise regret by $R_{t-1}(\mathbf{u}) := A_{t-1} - L_{t-1}(\mathbf{u})$. Consider the potential

$$\Phi_t := \max_{\theta \in \mathbb{R}} \log \int_{\mathcal{X}} \exp(\eta R_{t-1}(\mathbf{u}) - \beta(f_{t-1}(\mathbf{u}) - \theta)^2) \mu(d\mathbf{u}). \quad (6)$$

The factor $e^{\eta A_{t-1}}$ is common to all comparator points, so the maximizing centering parameter and normalized distribution in (6) are exactly θ_t and P_t in [Algorithm 2](#). Initially, $\Phi_1 = 0$.

Theorem 11 (Monotonicity of potential). *Assume $\eta \leq 1/8$ and $\beta = \eta/4$. For every round $t \geq 1$, we have $\Phi_{t+1} \leq \Phi_t \leq 0$.*

Proof. Define $\xi_t(\mathbf{u}) := f_{t-1}(\mathbf{u}) - \theta_t$ and $\xi_{t+1}(\mathbf{u}) := f_t(\mathbf{u}) - \theta_{t+1}$. By [Lemma 3](#), both functions take values in $[-2, 2]$, $\mathbb{E}_{P_t}[\xi_t] = 0$, $Q_t = (1 - \xi_t/2)P_t$, and $\mathbf{x}_t = \mathbb{E}_{Q_t}[\mathbf{u}]$.

Since $g_t = f_{t-1} + f_t$ is convex, Jensen's inequality gives $g_t(\mathbf{x}_t) \leq \mathbb{E}_{Q_t}[g_t]$. The two scalar centers cancel, so for every $\mathbf{u} \in \mathcal{X}$,

$$R_t(\mathbf{u}) - R_{t-1}(\mathbf{u}) \leq \mathbb{E}_{Q_t}[\xi_t + \xi_{t+1}] - \xi_t(\mathbf{u}) - \xi_{t+1}(\mathbf{u}).$$

By the definition of the potential and the fact that P_t is its normalized exponential-weight distribution, we have

$$\begin{aligned}\Phi_{t+1} - \Phi_t &= \log \mathbb{E}_{P_t} \exp\{\eta(R_t - R_{t-1}) + \beta(\xi_t^2 - \xi_{t+1}^2)\} \\ &\leq \log \mathbb{E}_{P_t} \exp\{\eta(\mathbb{E}_{Q_t}[\xi_t + \xi_{t+1}] - \xi_t - \xi_{t+1}) + \beta(\xi_t^2 - \xi_{t+1}^2)\}.\end{aligned}$$

We use the following continuous counterpart of [Lemma 2](#).

Lemma 7 (Functional one-step inequality). *Let P be a probability measure on an arbitrary measurable space. Let ξ, ζ be measurable functions with values in $[-2, 2]$ such that $\mathbb{E}_P[\xi] = 0$, and define $Q(\mathrm{d}u) := (1 - \xi(u)/2)P(\mathrm{d}u)$. If $\eta \leq 1/8$ and $\beta = \eta/4$, then*

$$\log \mathbb{E}_P \exp\{\eta(\mathbb{E}_Q[\xi + \zeta] - \xi - \zeta) + \beta(\xi^2 - \zeta^2)\} \leq 0.$$

Proof. For any measurable function z with values in $[-2, 2]$, define

$$H(z) := \log \mathbb{E}_P \exp\{\eta(\mathbb{E}_Q[\xi + z] - \xi - z) + \beta(\xi^2 - z^2)\}.$$

It suffices to show that $H(\zeta) \leq 0$. Let Π_z be the probability measure obtained by normalizing the exponential in the definition of $H(z)$. To show that H is concave, fix functions z, w such that $z_s := z + sw$ remains in $[-2, 2]$, and write $v_s := -\eta - 2\beta z_s$. $H(z_s)$ is differentiable since it is logarithm of a bounded integrand over a compact set. We note that $\mathbb{E}_Q[\xi + z_s]$ is affine and contributes nothing to the second-order derivative.

$$\frac{\mathrm{d}^2}{\mathrm{d}s^2} H(z_s) = \mathbb{E}_{\Pi_{z_s}}[(v_s^2 - 2\beta)w^2] - (\mathbb{E}_{\Pi_{z_s}}[v_s w])^2.$$

Since $z_s \in [-2, 2]$, $\beta = \eta/4$, and $\eta \leq 1/8$, we have $v_s \in [-2\eta, 0]$ and $v_s^2 \leq 4\eta^2 \leq 2\beta$. Thus H is concave.

At $z = -\xi$, every exponent vanishes, so $H(-\xi) = 0$ and $\Pi_{-\xi} = P$. For every bounded direction w , the directional derivative is

$$\begin{aligned}DH(-\xi)[w] &= \eta \mathbb{E}_Q[w] + \mathbb{E}_P[(-\eta + 2\beta\xi)w] \\ &= (2\beta - \eta/2)\mathbb{E}_P[\xi w] = 0,\end{aligned}$$

where the second equality uses $Q = (1 - \xi/2)P$ and the last uses $\beta = \eta/4$. Taking $w = \zeta + \xi$, the path $-\xi + sw$ remains in $[-2, 2]$ for $s \in [0, 1]$. Concavity therefore gives $H(\zeta) \leq H(-\xi) = 0$. $\square$

Applying [Lemma 7](#) with $P = P_t$, $\xi = \xi_t$, and $\zeta = \xi_{t+1}$ gives $\Phi_{t+1} \leq \Phi_t$. Since $\Phi_1 = 0$, the claim follows. $\square$

A.3 Proof of [Theorem 7](#)

Proof of [Theorem 7](#). The choices $\eta = 1/8$ and $\beta = 1/32$ in [Algorithm 2](#) satisfy the conditions in [Theorem 11](#). Thus $\Phi_{T+1} \leq \Phi_1 = 0$. By the definition of the potential, for every $\theta \in \mathbb{R}$,

$$\log \int_{\mathcal{X}} \exp(\eta R_T(\mathbf{u}) - \beta(f_T(\mathbf{u}) - \theta)^2) \mu(\mathrm{d}\mathbf{u}) \leq 0.$$

By [\(3\)](#), $R_T(\mathbf{u}) = \text{Reg}_{\text{Alt}}^T(\mathbf{u})$.

If $\text{KL}(\rho\|\mu) = \infty$, the claim is immediate. Otherwise, write $r := \text{d}\rho/\text{d}\mu$. Since $r > 0$ ρ -almost surely, for every $\theta \in \mathbb{R}$,

$$\begin{aligned} & \int_{\mathcal{X}} \exp(\eta R_T - \beta(f_T - \theta)^2) \, \text{d}\mu \\ & \geq \int_{\{x \in \mathcal{X} : r > 0\}} \exp(\eta R_T - \beta(f_T - \theta)^2) \, \text{d}\mu \\ & = \int_{\mathcal{X}} \exp(\eta R_T - \beta(f_T - \theta)^2 - \log r) \, \text{d}\rho. \end{aligned}$$

Here the inequality discards the possible μ -mass on $\{x \in \mathcal{X} : r = 0\}$, and the equality uses $\text{d}\rho = r \, \text{d}\mu$. Taking logarithms and applying Jensen's inequality to the probability measure ρ gives

$$\begin{aligned} 0 & \geq \log \int_{\mathcal{X}} \exp(\eta R_T - \beta(f_T - \theta)^2) \, \text{d}\mu \\ & \geq \log \int_{\mathcal{X}} \exp(\eta R_T - \beta(f_T - \theta)^2 - \log r) \, \text{d}\rho \\ & \geq \mathbb{E}_{\rho}[\eta R_T - \beta(f_T - \theta)^2 - \log r] \\ & = \eta \text{Reg}_{\text{Alt}}^T(\rho) - \beta \mathbb{E}_{\rho}[(f_T - \theta)^2] - \text{KL}(\rho\|\mu). \end{aligned}$$

Choose $\theta = \mathbb{E}_{\rho}[f_T]$. Since $1/\eta = 8$ and $\beta/\eta = 1/4$, rearranging proves the first inequality in (4). The second follows from $\text{Var}_{\rho}(f_T) \leq 1$. $\square$