
TC-Next: Zero-Shot Multimodal Cyclone Forecasting

Zhe Wang*

Carnegie Mellon University
Pittsburgh, PA 15213
zhew2@andrew.cmu.edu

Sijie Chen*

Carnegie Mellon University
Pittsburgh, PA 15213
sijieche@andrew.cmu.edu

Yiming Luo

Carnegie Mellon University
Pittsburgh, PA 15213
yimingl1@andrew.cmu.edu

Daehyun Kim

Seoul National University
Seoul, South Korea
daehyun@snu.ac.kr

Chien-Yi Chang

Durham University
Durham, United Kingdom
chien-yi.chang@durham.ac.uk

Abstract

We present TropicalCycloneNext (TC-Next), a multimodal deep learning model that forecasts tropical cyclone track and intensity at 6–24 h leads by leveraging a foundation model’s forecast fields of atmospheric kinematic and thermodynamic fields and GridSat infrared satellite imagery. Trained only on GraphCast forecasts over the Western Pacific (WP), yet reliant only on generic atmospheric variables, TC-Next on GraphCast lowers track error by 15–44% and intensity error by a factor of 3–6 relative to a conventional, rule-based tracker, TempestExtremes; applied without retraining to the forecast fields of Pangu-Weather and IFS HRES, it stays ahead of TempestExtremes on both. Applied zero-shot to the generic weather fields of WeatherNext Cyclones on the 2025 WP season, TC-Next attains lower intensity error at every lead time, and lower or comparable track error, compared to that model’s specialized direct tracker in a deterministic comparison. Our ablation studies show that our multimodal model is able to utilize the additional modality to improve performance in tracking errors at every lead time and in intensity prediction at longer lead times.

1 Introduction

Tropical cyclones (TCs) are among the deadliest climate hazards, a threat accelerating under a warming climate [1, 2]. Mass evacuations depend on accurate track and intensity forecasts, such as those that evacuated over two million people ahead of the 145 kt Super Typhoon Ragasa in 2025 [3]. Improving them is therefore a concrete, high-leverage form of climate adaptation. Recent advances in deep learning have produced weather foundation models, including deterministic ones such as GraphCast [4], Pangu-Weather [5], and Aurora [6], as well as probabilistic ones such as GenCast [7] and FGN [8], that forecast the global atmospheric states, matching or outperforming operational numerical weather prediction (NWP) while running orders of magnitude faster. While they reproduce TC tracks well [4, 7], they struggle with intensity. When intensity is read off their fields using TC trackers like TempestExtremes [9], the results are no better than persistence at short leads [10]. Furthermore, the ERA5 reanalysis [11] they are trained on carries a 12–15 m/s intensity error against best-track data [12] due to resolution constraints and assimilation processes [13].

Prior work addresses this issue in three main ways. First, WeatherNext Cyclones (WN-C) [14] extends the underlying weather model with TC-specific feature channels trained using best-track labels and extracts track and intensity through a specialized heuristic direct tracker. This attains state-of-the-art intensity, but at the cost of leaving the tracker and the TC head locked to the specialized model. Second, Fuxi-TC downscales the atmospheric variables resolution to resolve the storm’s tight inner core [13] by using a diffusion model to map coarse 0.25° FuXi forecasts to 0.1° WRF [15]-simulated

*Equal contribution.

fields. However, its intensity skill is below its teacher model, WRF-0.1. Third, a number of end-to-end models that do not leverage foundation models have been proposed in recent years [e.g., 16–18]. Although these models are relatively successful at intensity prediction, they cannot match the track skill achieved by foundation models. Furthermore, no existing approach achieves zero-shot transfer across diverse forecast models while also leveraging satellite observations of the storm core.

We introduce **TropicalCycloneNext** (TC-Next), a multimodal deep learning model designed to predict TC track and intensity around Western Pacific (WP) basin. TC-Next uses a foundation model’s forecast fields, which supply the large-scale steering flow and thermodynamic environmental conditions that are known affect TC intensity, together with high-resolution (0.07°) GridSat satellite imagery [19] of the storm’s recent history, which provides convective structures associated with the storm. Our contributions are three-fold: **1. State-of-the-Art Zero-Shot Transfer:**² We conduct comprehensive zero-shot out-of-distribution (OOD) evaluations across both AI-based and numerical forecasting models. Despite being trained solely on GraphCast, TC-Next achieved state-of-the-art transfer performance: it outperformed the specialized direct tracker of the current SOTA (WN-C) in a deterministic setup (Table 1), and consistently surpassed the TempestExtremes baseline in both track and intensity errors across multiple AI foundation models as well as the HRES numerical model at all lead times, lowering track error by 15–49% and intensity error by a factor of 2–6 (Table 2). **2. Multimodal Efficacy:** Ablation studies on the satellite infrared (IR) branch demonstrate that multimodal inputs lower track error at every lead time and improve intensity error from +18 h onward. The benefit increases with lead time, peaking at +24 h (Table 3). **3. Cross-Model Portability:** TC-Next requires from the forecast source only generic atmospheric variables (Table 5; SST is persisted from the analysis). Hence, the same trained model can be applied across forecast sources, providing a common learned tracker for evaluating TC forecast skill.

2 Methodology

Problem formulation. We forecast TC track and intensity at lead times $\mathcal{L} = \{6, 12, 18, 24\}$ h. Given observations through time t , a network f_θ emits, for each lead ℓ , raw center and extent updates $\mathbf{a}_\ell, \mathbf{s}_\ell \in \mathbb{R}^2$ and an intensity increment $u_\ell \in \mathbb{R}$:

$$\{(\mathbf{a}_\ell, \mathbf{s}_\ell, u_\ell)\}_{\ell \in \mathcal{L}} = f_\theta(E_{t-24:t}, S_{t-24:t}, \hat{E}_{t+6:t+24}^F, \mathbf{c}_{t-24:t}, \mathbf{v}_{t-24:t}). \quad (1)$$

The center and intensity updates are applied autoregressively from time t :

$$\hat{\mathbf{c}}_{t+\ell} = \hat{\mathbf{c}}_{t+\ell-6} + \delta \tanh(\mathbf{a}_\ell), \quad \hat{v}_{t+\ell} = \hat{v}_{t+\ell-6} + u_\ell, \quad (2)$$

where $\delta = 4^\circ$ bounds the motion per step along each axis; $\hat{\mathbf{c}}_{t+\ell}$ and $\hat{v}_{t+\ell}$ are the predicted center and intensity at $t + \ell$. Here, $E_{t-24:t} \in \mathbb{R}^{5 \times H \times W \times C}$ holds five historical atmospheric frames with $H \times W = 280 \times 420$ grid cells (a $70^\circ \times 105^\circ$ domain at 0.25°) and $C = 12$ channels (Table 5); $S_{t-24:t} \in \mathbb{R}^{5 \times 256 \times 256 \times 2}$ holds five IR frames, each a 16° window of brightness temperature and its validity mask; $\mathbf{c}_{t-24:t} \in \mathbb{R}^{5 \times 2}$ denotes historical centers; and $\mathbf{v}_{t-24:t} \in \mathbb{R}^5$ is the historical intensity. The four future atmospheric frames $\hat{E}_{t+6:t+24}^F \in \mathbb{R}^{4 \times H \times W \times C}$ are forecasts from a weather foundation model F initialized at t . The extent $\mathbf{s}_\ell$ updates an auxiliary box around the storm (from USA_R34 wind radius) that lets the loss judge a position miss by both distance and storm size; see Eq. (3) for its update rule. The history window has is fixed at five steps, but past fixes are optional: only the anchor fix ($\mathbf{c}_t, v_t$) must be observed, so a forecast can issue from a storm’s first fix.³

Data. To train TC-Next, we built a dataset combining atmospheric fields—ERA5 reanalysis for the storm’s observed history and GraphCast forecasts for the lead times ahead—with IR brightness temperatures near $11 \mu\text{m}$ from GridSat-B1 and storm state from IBTrACS. From each atmospheric source we crop a fixed, WP-basin-centered $70^\circ \times 105^\circ$ domain; 0.07° IR is resampled to 0.0625° and cropped to a 16° window centered on the storm’s observed position at each history step. Storm centers follow IBTrACS’s best track, and intensity is the 1-minute maximum sustained wind (USA_WIND).

A training sample is anchored to a single IBTrACS fix at time t . Over the five 6-hourly steps ending at t , it gathers ERA5, IR, and TC positions and intensities, masking any steps that precede the storm’s

²We define *zero-shot transfer* as replacing the training-time forecast source $F = \text{GraphCast}$ (the publicly released 0.25° , 37-level checkpoint trained on ERA5 1979–2017) with an unseen source F' at inference without retraining or modifying any other input. The only exception is the comparison against WN-C in Table 1: there, IFS HRES analysis frames replace the ERA5 history frames to match the analysis distribution WN-C is fine-tuned on, and the TC Vitals block further replaces the IBTrACS anchors with real-time TC Vitals estimates.

³History steps preceding the storm’s first record are zero-filled and masked, in training and at inference alike.

Table 1: WP, 2025; TC-Next against the direct tracker of WN-C. Errors as defined in Section 2; bold marks the lower error in each column within an anchor block. Scored samples: Table 6.

Anchor record	TC model	Track ↓ (km)				Intensity ↓ ($V_{\max}$, kt)			
		+6 h	+12 h	+18 h	+24 h	+6 h	+12 h	+18 h	+24 h
IBTrACS	TC-Next (WN-C)^a	33.55	43.46	51.12	58.70	4.20	5.65	6.69	7.80
IBTrACS	WN-C direct tracker ^b	43.10	46.71	52.36	60.84	7.88	8.19	8.62	9.35
TC Vitals	TC-Next (WN-C)^a	35.11	42.05	49.95	58.00	3.94	5.34	6.44	7.29
TC Vitals	WN-C direct tracker ^b	37.33	42.01	49.40	60.36	7.95	8.23	8.71	9.49

^aZero-shot: applied to WN-C without retraining, reading only WN-C’s generic weather channels.

^bThe direct tracker derives its position and intensity from WN-C’s dedicated TC feature channels.

first record. It also includes the four GraphCast forecasts initialized at t . As targets, we use the storm’s positions and intensities from $t+6$ to $t+24$ h, masking any leads beyond its final record. Every in-domain fix that is followed by another in-domain fix 6 h later is a sample; storms that never reach tropical-storm strength (peak USA_SSHS < 0) are dropped. Details are in Appendix A. We partition data chronologically to prevent temporal leakage: training (1990–2017), validation (2018), and testing (2019–2020), comprising 28,962, 1,171, and 1,815 samples from 801, 34, and 57 storms, respectively. To avoid temporal overlap with WN-C’s training data (through 2024), we evaluate against it on a separate 2025 test set of 867 samples from 29 storms. At evaluation, we score track and intensity separately at each lead time: track error is the mean great-circle distance (km), and intensity error is the mean absolute error (kt). Beyond GraphCast, we draw Pangu-Weather [5] and IFS HRES [20] forecasts from the WeatherBench2 archive [21], and real-time storm estimates from TC Vitals [22].

Model. TC-Next is a multimodal encoder-decoder architecture with 3.6 M parameters, trainable on a TPU v4-8 VM in less than 70 minutes. Its two encoders process atmospheric fields and satellite imagery at their respective input resolutions. A *macro* CNN maps each atmospheric frame to a 2° feature grid. A finer-stride *micro* CNN processes a 16° GridSat window centered on the observed storm position at each historical time step, together with its per-pixel validity mask, producing features on a 0.25° grid. The macro feature map goes into a region-proposal network [23] (RPN) to predict a storm-centered proposal box and the extent, supervised directly against the center and wind-radii box at every history step, to give the macro CNN a gradient path that bypasses the recurrent encoder–decoder. RoIAlign [24] then extracts geographically aligned 10° windows around the proposed center from both feature maps, and low-rank multimodal fusion [25] combines the resulting representations; see Eq. (5).

An encoder LSTM processes the five fused historical representations and initializes a decoder LSTM. At each forecast lead ℓ , the decoder extracts a 10° window from the macro-encoded forecast frame at $t + \ell$, centered on its previous predicted storm position. Separate prediction heads produce the center and intensity updates of Eq. (2) together with an extent update s_ℓ that scales the predicted box extent $\hat{\mathbf{r}}_{t+\ell} \in \mathbb{R}^2$ (width and height):

$$\hat{\mathbf{r}}_{t+\ell} = \text{clip}_{[2^\circ, 10^\circ]} \left(\hat{\mathbf{r}}_{t+\ell-6} \odot \exp(\tanh(s_\ell)/2) \right). \quad (3)$$

The decoder initializes $\hat{\mathbf{c}}_t$ and $\hat{\mathbf{r}}_t$ from the last RPN proposal at t , and sets $\hat{v}_t = v_t$. Each forecast therefore updates an existing estimate of storm position, extent, and intensity—the sense in which the forecast is anchored—the decoder then rolls forward autoregressively over the available forecast frames. Architectural and training details, including the loss in Eq. (6), appear in Appendix B.

3 Experiments and Results

We evaluate TC-Next in three settings: (1) against the direct tracker of WN-C on the WP 2025 season (Table 1); (2) against TempestExtremes [9], configured as in TC Bench [10], on the pooled WP 2019–2020 seasons across three forecast sources (Table 2); and (3) with its satellite infrared branch ablated (Table 3). TC-Next is trained solely on GraphCast fields; Pangu-Weather, IFS HRES, and WN-C are evaluated without retraining. Baseline configuration, sample matching, pooling, and scoring conventions are given in Appendix C.

TC-Next transfers zero-shot to WN-C and outperforms its direct tracker. Reading only generic weather channels, our frozen model outperforms the WN-C direct tracker’s single-member predictions

Table 2: WP, 2019–2020; TC-Next vs. TempestExtremes. Scored samples: Table 7.

Forecast source	TC model	Track ↓ (km)				Intensity ↓ ($V_{\max}$, kt)			
		+6 h	+12 h	+18 h	+24 h	+6 h	+12 h	+18 h	+24 h
GraphCast	TempestExtremes	51.06	46.54	52.44	57.03	22.44	25.30	26.63	28.90
GraphCast	TC-Next^a	28.47	35.59	43.80	48.68	3.68	5.94	7.46	8.93
Pangu [†]	TempestExtremes	53.58	53.07	61.89	62.07	22.71	24.70	26.19	26.88
Pangu [†]	TC-Next (Pangu)^b	29.48	39.63	47.76	51.11	3.75	6.23	8.26	10.15
IFS HRES	TempestExtremes	59.41	63.18	72.50	79.75	15.08	15.69	16.17	16.50
IFS HRES	TC-Next (HRES)^b	30.14	40.88	51.77	62.25	3.56	5.87	7.35	8.77

^aSource: the model was trained on this forecast source. ^bZero-shot: applied without retraining.[†]Pangu-Weather used 2019 for validation, so that year is not strictly held out.

Table 3: WP, 2019–2020; IR ablation study. Scored samples: Table 8.

IR branch	Track ↓ (km)				Intensity ↓ ($V_{\max}$, kt)			
	+6 h	+12 h	+18 h	+24 h	+6 h	+12 h	+18 h	+24 h
Fed	34.86	49.42	60.26	69.22	3.39	5.17	6.37	7.45
IR-ablated	35.60	50.82	61.74	71.37	3.38	5.11	6.43	7.60
Δ (fed – IR-ablated)	–0.74	–1.40	–1.48	–2.15	+0.01	+0.06	–0.06	–0.15

on track and intensity. WN-C’s ensemble members are scored individually against best track and averaged, so Table 1 reflects expected single-member error, not ensemble-mean error. The IBTrACS block uses the same HRES atmospheric source and IBTrACS anchor convention used for WN-C fine-tuning: our track error is lower at every lead, and early-lead intensity error is roughly halved. The TC Vitals block [22] swaps the IBTrACS anchors for the operational, real-time ones. Here, track performance remains on par—leading at +6 h and +24 h, trailing by under 0.6 km in between—while our intensity advantage widens, holding between 3.94 and 7.29 kt against the direct tracker’s 7.95 to 9.49 kt across all leads. These results suggest that the gains survive the shift to operational anchors; fine-tuning directly on real-time inputs (TC Vitals and GOES-R imagery [26]) is a natural next step.

TC-Next outperforms TempestExtremes across all forecast sources and lead times. On GraphCast, the source the model is trained on, track error is 15–44% lower than TempestExtremes’ (Table 2), and intensity error is 3.2–6.1 times lower. The margin survives zero-shot transfer: applied without retraining, TC-Next leads TempestExtremes on track by 18–45% on Pangu-Weather and by 22–49% on IFS HRES, and its intensity error stays below 11 kt on every source at every lead, whereas TempestExtremes’ never falls below 15 kt. These results highlight that our learned model provides an alternative to rule-based detection across forecast sources.

The satellite infrared branch lowers track error at every lead and intensity error at longer leads, with a benefit that grows with lead time. Table 3 ablates the satellite infrared branch on the full test set: the *IR-ablated* variant is a separately trained model with the IR input zeroed in training and evaluation, all else fixed. The fed variant lowers track error at every lead, by 0.7 km at +6 h rising to 2.2 km at +24 h, and intensity error from +18 h onward. These results suggest that the benefit of multimodal input increases with lead time and is largest at +24 h.

4 Conclusion

In this study, we introduce TC-Next, a multimodal deep learning model that predicts TC track and intensity using a foundation weather model’s forecast fields and high-resolution satellite imagery of the storm’s history. Because the model’s forecast input relies exclusively on generic atmospheric variables, it can be applied across forecast sources without retraining. Evaluated on our WP dataset, TC-Next demonstrates robust zero-shot transfer capabilities across diverse foundation models, numerical prediction systems, and real-time input configurations, consistently outperforming traditional rule-based detection methods. Ongoing work focuses on expanding this dataset and extending the zero-shot benchmark to additional TC basins toward a global benchmark.

Acknowledgments and Disclosure of Funding

We gratefully acknowledge the support of Google for providing computing resources via the TPU Research Cloud (TRC) program, which made the extensive training of our models possible. This work contains modified Copernicus Climate Change Service information (ERA5); neither the European Commission nor ECMWF is responsible for any use of the Copernicus information it contains. This work originated from a CMU course project. We thank our original team member, Ziping He, for her initial contributions, and course staff Ralf Brown and Yonatan Bisk for their guidance. DK was supported by Creative-Pioneering Researchers Program through Seoul National University and the National Research Foundation of Korea(NRF) grant funded by the Korea government(MSIT) (RS-2025-02310080). CC was supported by the JustNOW Strategic Research Fund through Durham University and the project ELEVATE-ProClima through British Academy (EPG\100778).

References

- [1] Kerry Emanuel. Increasing destructiveness of tropical cyclones over the past 30 years. *Nature*, 436(7051):686–688, 2005. doi: 10.1038/nature03906.
- [2] Thomas Knutson, Suzana J. Camargo, Johnny C. L. Chan, Kerry Emanuel, Chang-Hoi Ho, James Kossin, Mrutyunjay Mohapatra, Masaki Satoh, Masato Sugi, Kevin Walsh, and Liguang Wu. Tropical cyclones and climate change assessment: Part I: Detection and attribution. *Bulletin of the American Meteorological Society*, 100(10):1987–2007, 2019. doi: 10.1175/BAMS-D-18-0189.1.
- [3] Guy Carpenter. Post-event report: 2025 western north pacific typhoon ragasa. Technical report, Guy Carpenter & Company, LLC, October 2025. <https://www.guycarp.com/insights/2025/10/post-event-report-2025-western-north-pacific-typhoon-rRagasa.html>.
- [4] Remi Lam, Alvaro Sanchez-Gonzalez, Matthew Willson, Peter Wernsberger, Meire Fortunato, Ferran Alet, Suman Ravuri, Timo Ewalds, Zach Eaton-Rosen, Weihua Hu, Alexander Merose, Stephan Hoyer, George Holland, Oriol Vinyals, Jacklynn Stott, Alexander Pritzel, Shakir Mohamed, and Peter Battaglia. Learning skillful medium-range global weather forecasting. *Science*, 382(6677):1416–1421, 2023.
- [5] Kaifeng Bi, Lingxi Xie, Hengheng Zhang, Xin Chen, Xiaotao Gu, and Qi Tian. Accurate medium-range global weather forecasting with 3D neural networks. *Nature*, 619:533–538, 2023.
- [6] Cristian Bodnar, Wessel P. Bruinsma, Ana Lucic, Megan Stanley, Anna Vaughan, Johannes Brandstetter, Patrick Garvan, Maik Riechert, Jonathan A. Weyn, Haiyu Dong, Jayesh K. Gupta, Kit Thambiratnam, Alexander T. Archibald, Chun-Chieh Wu, Elizabeth Heider, Max Welling, Richard E. Turner, and Paris Perdikaris. A foundation model for the Earth system. *Nature*, 641, 2025. Aurora; arXiv:2405.13063.
- [7] Ilan Price, Alvaro Sanchez-Gonzalez, Ferran Alet, Tom R. Andersson, Andrew El-Kadi, Dominic Masters, Timo Ewalds, Jacklynn Stott, Shakir Mohamed, Peter Battaglia, Remi Lam, and Matthew Willson. Probabilistic weather forecasting with machine learning. *Nature*, 2025. GenCast; arXiv:2312.15796.
- [8] Ferran Alet, Ilan Price, Andrew El-Kadi, Dominic Masters, Stratis Markou, Tom R. Andersson, Jacklynn Stott, Remi Lam, Matthew Willson, Alvaro Sanchez-Gonzalez, and Peter Battaglia. Skillful joint probabilistic weather forecasting from marginals. 2025. FGN; arXiv:2506.10772.
- [9] Paul A. Ullrich, Colin M. Zarzycki, Elizabeth E. McClenny, Marielle C. Pinheiro, Alyssa M. Stansfield, and Kevin A. Reed. TempestExtremes v2.1: a community framework for feature detection, tracking, and analysis in large datasets. *Geoscientific Model Development*, 14(8): 5023–5048, 2021. doi: 10.5194/gmd-14-5023-2021.
- [10] Milton Gomez, Saranya Ganesh S., Marie McGraw, Frederick Iat-Hin Tam, Ilia Azizi, Samuel Darmon, Monika Feldmann, Stella Bourdin, Louis Poulain-Auzéau, Suzana J. Camargo, Jonathan Lin, Dan Chavas, Chia-Ying Lee, Ritwik Gupta, Andrea Jenney, and Tom Beucler. TCBench: A benchmark for tropical cyclone track and intensity forecasting at the global scale. 2026. arXiv:2601.23268.
- [11] Hans Hersbach, Bill Bell, Paul Berrisford, Shoji Hirahara, András Horányi, Joaquín Muñoz-Sabater, Julien Nicolas, Carole Peubey, Raluca Radu, Dinand Schepers, et al. The ERA5 global reanalysis. *Quarterly Journal of the Royal Meteorological Society*, 146(730):1999–2049, 2020. doi: 10.1002/qj.3803.
- [12] Kenneth R. Knapp, Michael C. Kruk, David H. Levinson, Howard J. Diamond, and Charles J. Neumann. The international best track archive for climate stewardship (IBTrACS): Unifying tropical cyclone data. *Bulletin of the American Meteorological Society*, 91(3):363–376, 2010. doi: 10.1175/2009BAMS2755.1.

- [13] Shan Guo, Lei Chen, Yangyang Zhao, Yuetan Lin, Zeyi Niu, Xinyan Zhang, Ziyao Sun, Xiaohui Zhong, and Hao Li. FuXi-TC: A generative framework integrating deep learning and physics-based models for improved tropical cyclone forecasts. 2025. arXiv:2508.16168.
- [14] Ferran Alet, Tom R. Andersson, Ilan Price, Stratis Markou, Andrew El-Kadi, Dominic Masters, Amy Li, Samier Merchant, Natalie Williams, Gregory Thornton, Ken MacKay, Olivia Graham, Akib Uddin, Ben Gaiarin, Devaja Shah, Elinor Kruse, Wallace Hogsett, David Zelinsky, John Cangialosi, Jonathan Martinez, James Franklin, Mark DeMaria, Kate Musgrave, Caroline L. Bain, Helen Titley, Jacklynn Stott, Remi Lam, Aaron Bell, Paul Komarek, Matthew Willson, Alvaro Sanchez-Gonzalez, and Peter Battaglia. Operational tropical cyclone forecasting with AI. *Nature*, 2026. doi: 10.1038/s41586-026-10953-2. WeatherNext Cyclones.
- [15] William C. Skamarock, Joseph B. Klemp, Jimy Dudhia, David O. Gill, Zhiqian Liu, Judith Berner, Wei Wang, Jordan G. Powers, Michael G. Duda, Dale M. Barker, and Xiang-Yu Huang. A description of the advanced research WRF model version 4.3. Technical Report NCAR/TN-556+STR, National Center for Atmospheric Research, Boulder, CO, 2021.
- [16] Léonard Boussioux, Cynthia Zeng, Théo Guénais, and Dimitris Bertsimas. Hurricane forecasting: A novel multimodal machine learning framework. *Weather and Forecasting*, 37(6): 817–831, 2022.
- [17] Renlong Hang, Zihao Xu, Jiuwei Zhao, Runling Yu, Leye Cheng, and Qingshan Liu. Cyclone-MAE: A scalable multi-task learning model for global tropical cyclone probabilistic forecasting. 2026. arXiv:2604.12180.
- [18] Zihao Lin, Jung-Eun Chu, and Yoo-Geun Ham. Enhancing tropical cyclone track and intensity predictions with the OWZP-Transformer model. *npj Artificial Intelligence*, 2025. doi: 10.1038/s44387-025-00037-3.
- [19] Kenneth R. Knapp, Steve Ansari, Caroline L. Bain, Mark A. Bourassa, Michael J. Dickinson, Chris Funk, Chip N. Helms, Christopher C. Hennon, Christopher D. Holmes, George J. Huffman, James P. Kossin, Hai-Tien Lee, Alexander Loew, and Gudrun Magnusdottir. Globally gridded satellite observations for climate studies. *Bulletin of the American Meteorological Society*, 92(7):893–907, 2011. doi: 10.1175/2011BAMS3039.1.
- [20] ECMWF. IFS documentation CY49R1 – part IV: Physical processes. Technical report, European Centre for Medium-Range Weather Forecasts, Reading, UK, 2024. <https://www.ecmwf.int/en/elibrary/81626-ifs-documentation-cy49r1-part-iv-physical-processes>.
- [21] Stephan Rasp, Stephan Hoyer, Alexander Merose, Ian Langmore, Peter Battaglia, Tyler Russell, Alvaro Sanchez-Gonzalez, Vivian Yang, Rob Carver, Shreya Agrawal, Matthew Chantry, Zied Ben Bouallegue, Peter Dueben, Carla Bromberg, Jared Sisk, Luke Barrington, Aaron Bell, and Fei Sha. WeatherBench 2: A benchmark for the next generation of data-driven global weather models. *Journal of Advances in Modeling Earth Systems*, 16(6):e2023MS004019, 2024. doi: 10.1029/2023MS004019.
- [22] Sam Trahan and Lynn Sparling. An analysis of NCEP tropical cyclone vitals and potential effects on forecasting models. *Weather and Forecasting*, 27(3):744–756, 2012. doi: 10.1175/WAF-D-11-00063.1.
- [23] Shaoqing Ren, Kaiming He, Ross Girshick, and Jian Sun. Faster R-CNN: Towards real-time object detection with region proposal networks. In *Advances in Neural Information Processing Systems*, 2015.
- [24] Kaiming He, Georgia Gkioxari, Piotr Dollár, and Ross Girshick. Mask R-CNN. In *IEEE International Conference on Computer Vision (ICCV)*, 2017.
- [25] Zhun Liu, Ying Shen, Varun Bharadhwaj Lakshminarasimhan, Paul Pu Liang, Amir Zadeh, and Louis-Philippe Morency. Efficient low-rank multimodal fusion with modality-specific factors. In *Proceedings of the 56th Annual Meeting of the Association for Computational Linguistics (ACL)*, 2018.
- [26] Timothy J. Schmit, Paul Griffith, Mathew M. Gunshor, Jaime M. Daniels, Steven J. Goodman, and William J. Leblair. A closer look at the ABI on the GOES-R series. *Bulletin of the American Meteorological Society*, 98(4):681–698, 2017. doi: 10.1175/BAMS-D-15-00230.1.

A Data Details

Table 4 gives the sample and storm counts per split and the rule behind them; Table 5 lists the channels extracted per frame.

Data sources and preprocessing. GraphCast’s forecasts use the same 0.25° grid and the same variables as the ERA5 history frames; both are read into one array with no regridding. GridSat-B1 is the NOAA climate data record of intercalibrated geostationary imagery, native to a 0.07° grid at 3-h cadence; we take its single infrared-window channel at the 00, 06, 12 and 18 UTC frames that coincide with the reanalysis cadence and resample it by bilinear interpolation. Pixels without a valid source observation, and whole frames absent from the record (43 in 1990–2018, 25 in 2020), are zero-filled and carried as a per-pixel validity mask that the model receives as a second input channel. Each of the 12 atmospheric channels is normalized by its mean and standard deviation, computed in one pass over all five frames of every one of the 28,962 training samples, the statistics are then fixed for validation, testing, and every forecast source. Infrared brightness temperature is scaled by fixed constants, $(T_b - 250 \text{ K})/40 \text{ K}$, and multiplied by the validity mask.

The best-track records are post-season reanalyses for 1990–2020, whereas the 2025 records remain provisional. Storms that never reach tropical-storm strength are excluded, and fixes outside the domain are discarded. Storm positions are kept in one coordinate system throughout: each best-track position (longitude wrapped to $0\text{--}360^\circ$, so the domain stays continuous across the antimeridian) is mapped linearly onto the 0.25° latitude–longitude grid and expressed as a fractional (row, column) location in $[0, 1]^2$ of the domain zero-padded to 288×448 cells ($72^\circ \times 112^\circ$ extent). The model reads, proposes, and predicts positions in these units; multiplying by that extent converts them to the degrees used by the box loss, and forecasts are mapped back to latitude and longitude only for scoring. A fix with no reported wind is masked out of the intensity term only. The extent target is a square box centered on the best-track position whose side is twice the mean of the four quadrant 34 kt wind radii reported by JTWC (IBTrACS USA_R34), converted from nautical miles to degrees and clipped so that the side lies in $[2^\circ, 10^\circ]$. Where no radii are reported, the side is fixed at 4° . When a storm’s record spans less than 24 h at the anchor time, the unavailable history steps are masked rather than dropping the sample.

Array dimensions and storage. Atmospheric fields are cached as one file per initialization time t : five frames $\times 280 \times 420 \times 12$ channels in half precision (about 14.1 MB), holding the ERA5 analysis at t and the four GraphCast forecasts initialized at t (leads 6–24 h). A sample anchored at t reads the analysis frame from the five files at $t-24, \dots, t$ and the four forecast frames from the file at t . Each analysis frame is therefore stored once and reused by every sample whose history window covers it—about 1.5 samples per file, with history windows straddling split boundaries—so we report the dataset as total files rather than per-split counts. GridSat infrared frames are stored separately, one per 6-hourly step (1120×1680 at resampled 0.0625° , compressed, about 1.7 MB each).

Dataset statistics. The WP atmospheric cache (ERA5 and GraphCast) comprises 19,360 files for 1990–2018, 770 for 2019 and 491 for 2020, or 20,621 files and 271 GiB in total; the 2025 WN-C comparison adds a further 601 files (7.9 GiB). The infrared imagery totals 34.7 GiB for 1990–2020 and 1.00 GiB for 2025.

Table 4: Dataset summary. The view is a fixed $70^\circ \times 105^\circ$ domain at 0.25° resolution spanning $5^\circ\text{S--}65^\circ\text{N}$ and $95^\circ\text{E--}160^\circ\text{W}$. A sample is a 6-hourly fix inside the view followed by another fix 6 h later, also inside it; storm counts are of the storms those samples come from. Because the view is a fixed geographic box rather than a basin mask, it also holds storms that IBTrACS codes to adjacent basins: 1,108 of the 28,962 training samples, most of them East Pacific. The test set pools two held-out seasons, 2019 (1,069 samples, 32 storms) and 2020 (746 samples, 25 storms). The 2025 counts come from the per-basin file IBTrACS.WP.v04r01.nc (snapshot of 2026-08-18) and are snapshot-dependent while the 2025 records remain provisional.

Basin	Train (1990–2017)		Val (2018)		Test (2019–2020)		Test (2025)	
	Samples	Storms	Samples	Storms	Samples	Storms	Samples	Storms
WP	28,962	801	1,171	34	1,815	57	867	29

Table 5: Atmospheric channels extracted per frame from the analysis (history) and GraphCast (forecast). Only three pressure levels are used, chosen for the environmental controls on TC intensity: 850 hPa (low-level circulation and moisture), 500 hPa (mid-tropospheric environmental flow and core ascent), and 200 hPa (outflow; paired with 850 hPa it yields the deep-layer vertical wind shear). Sea-surface temperature is not forecast by GraphCast and is taken from the ERA5 analysis at initialization time t in every forecast frame.

Level	Variables	Channels
Surface	Mean sea-level pressure; 10 m wind (u, v)	3
850 hPa	Wind (u, v); specific humidity	3
500 hPa	Geopotential; vertical velocity (ω)	2
200 hPa	Temperature; wind (u, v)	3
Surface (analysis at t)	Sea-surface temperature	1

B Model and Training Details

Architecture. A *macro* CNN encodes each atmospheric frame with three stride-2 blocks of 32, 64, and 96 channels, reducing the 0.25° input to a 2° feature map. Its input is the 12 atmospheric channels of Table 5 plus four static planes: latitude, the sine and cosine of longitude, and the ERA5 land-sea mask. A finer-stride *micro* CNN processes the 16° GridSat window centered on the observed storm position at each history step, together with its per-pixel validity mask, mapping the resampled 0.0625° IR input to a 0.25° feature grid; the two maps then sit on scales RoIAlign can align, but the micro features are drawn from imagery four times finer than the environment fields.

A region-proposal network [23] reads the macro map alongside a Gaussian prior rendered at the storm’s known fix for that step and emits a box: a center offset from the prior, clamped to 2.5° , plus a side length in $[2^\circ, 10^\circ]$. The proposal decides where the model looks: RoIAlign [24] samples a fixed 10° window from both feature maps at the proposed center (8×8 points on the macro map, 16×16 on the micro map), and separate convolutional heads flatten each crop to 2048 dimensions.

Write $z_\tau^m \in \mathbb{R}^{2048}$ for the crop vector of stream $m \in \{\text{env}, \text{ir}\}$ at history step τ . Each is projected to 128 dimensions, layer-normalized with its own learned scale and offset, and extended by a constant:

$$\tilde{z}_\tau^m = [\text{LN}_m(P_m z_\tau^m); 1] \in \mathbb{R}^{129}. \quad (4)$$

The two streams are combined by low-rank multimodal fusion [25],

$$V_\tau = \sum_{r=1}^R w_r (W_{\text{env}}^{(r)} \tilde{z}_\tau^{\text{env}}) \odot (W_{\text{ir}}^{(r)} \tilde{z}_\tau^{\text{ir}}) + b, \quad (5)$$

with $R = 8$ and $\odot$ the element-wise product. The appended 1 retains the unimodal terms, so Eq. (5) is strictly more expressive than passing the concatenation of the two streams through a linear layer. The per-stream normalization is load-bearing: in a product each stream’s gradient is scaled by the other, so without it the infrared branch is starved by the much larger environment signal.

The storm’s own scalars—the prior position, current intensity, availability flags, and a step index—are concatenated to V_τ , then passes through a two-layer MLP of width 256 with ReLU, and an encoder LSTM of width 256 reads the five resulting vectors in order; its final state initializes a decoder LSTM of the same width. At each lead ℓ , the decoder input is the 2048-dimensional macro crop head applied to the RoIAlign window of the forecast frame at $t + \ell$, taken at the previous predicted center, concatenated with that center, the previous extent (divided by 10°), the previous intensity, and a one-hot lead index; a linear layer with ReLU maps this to 256 dimensions before the LSTM. Two linear heads read the LSTM output: a four-dimensional box head giving $\mathbf{a}_\ell$ and $\mathbf{s}_\ell$ of Eqs. (2) and (3), and a scalar head giving u_ℓ . Both heads are initialized near zero, so the untrained model forecasts persistence. The IR data enters the forecast only through the encoder state: the decoder reads the forecast environment fields alone, since no IR imagery exists for future times.

Training. The total loss has three terms of two kinds: a box loss, which scores position and extent, applied to both the forecast boxes and the history proposals; and an intensity loss, which scores maximum sustained wind. Writing $b_\ell = (\mathbf{c}_\ell, \mathbf{r}_\ell)$ for the storm’s box in degrees, the extent target $\mathbf{r}_\ell$

being the best-track gale-force (34 kt) wind radius (R34, the mean of the four reported quadrants; a fixed 4° where unreported), training minimizes

$$\mathcal{L}_{\text{box}}(\mathcal{T}_F) + \beta \mathcal{L}_{\text{int}} + \gamma \mathcal{L}_{\text{box}}(\mathcal{T}_H),$$

$$\mathcal{L}_{\text{box}}(\mathcal{T}) = \frac{1}{|\mathcal{T}|} \sum_{\ell \in \mathcal{T}} \left[\lambda \|\hat{b}_\ell - b_\ell\|_1 + 1 - \text{GIoU}(\hat{b}_\ell, b_\ell) \right], \quad \mathcal{L}_{\text{int}} = \frac{1}{|\mathcal{T}_v|} \sum_{\ell \in \mathcal{T}_v} (\hat{v}_\ell - v_\ell)^2, \quad (6)$$

with $\lambda = 0.2$ and $\gamma = 0.3$. Here $\mathcal{T}_F \subseteq \{t+6, \dots, t+24\}$ and $\mathcal{T}_H \subseteq \{t-24, \dots, t\}$ are the forecast and history steps at which the storm has a best-track fix, and $\mathcal{T}_v \subseteq \mathcal{T}_F$ those at which wind is also reported, so a storm with a partial record contributes only its observed fixes. For $\ell \in \mathcal{T}_H$, $\hat{b}_\ell$ is the region-proposal box at that step. The intensity weight β opens at 20% of its final value of 2.5 and reaches it after one epoch, which lets the network localize before intensity competes for the shared trunk. We optimize with AdamW at a peak learning rate of 3×10^{-4} , warmed up over 300 steps and cosine-decayed thereafter, with weight decay 10^{-4} , gradients clipped to global norm 1, and a batch of 32 sharded data-parallel over a TPU v4-8. The model trains on the 1990–2017 split for 15 epochs in about 65 minutes; we keep the epoch with the best validation score on 2018, track error in km over 100 plus intensity MAE in knots over 10, which selects epoch 6. No held-out season enters selection at any point. Under zero-shot transfer the input normalization statistics remain those of the training source; none are re-estimated.

C Experimental Setup Details

TempestExtremes. For every forecast source we read the storm’s position and intensity directly off its forecast fields with TempestExtremes [9], using the detection parameters of TC-Bench [10]: a sea-level-pressure minimum is accepted only if the pressure rises by 200 Pa within 6.5° great-circle distance (a closed low) and the 300–500 hPa thickness falls by $58.8 \text{ m}^2 \text{ s}^{-2}$ within 5.5° (a warm core), with candidate nodes closer than 6.0° merged. Intensity is measured at each node as the minimum sea-level pressure at the node itself and the maximum 10 m wind speed within 2.0° great-circle distance. The baseline receives the same anchor as TC-Next. Where no node falls inside the gate the tracker returns no forecast: it produces a forecast for 54–70% of samples depending on source, whereas TC-Next covers all samples by construction. All comparisons are therefore restricted to the samples both methods produce. Two caveats apply. First, the 2.0° wind radius is the default of the standard TempestExtremes recipe; all baseline intensities are conditional on this choice. Second, a gridded wind maximum over a finite radius is not the same quantity as the 1-minute sustained wind of the best-track record, so part of the baseline’s intensity error is definitional rather than forecast error.

Forecast sources. TC-Next is trained on GraphCast forecast fields only, together with ERA5 history frames, GridSat IR frames, and best-track anchors from IBTrACS. Every other input appears at evaluation time alone: the model is applied without retraining to the forecast fields of Pangu-Weather, IFS HRES, and WN-C, with the Pangu-Weather and IFS HRES fields taken from the WeatherBench2 archives [21]. Pangu-Weather emits no vertical velocity, so the ω_{500} input channel is zero-filled for its rows; TempestExtremes does not use this field and is unaffected. Table 1 likewise substitutes inputs at evaluation time only. Throughout the table, IFS HRES analysis frames replace the ERA5 history, matching the analyses WN-C is fine-tuned on. In the TC Vitals block, TC Vitals [22]—the real-time storm estimates issued by the operational forecast centers—additionally replace the best-track anchors; that configuration therefore measures zero-shot robustness to the inputs available in real time, not the behavior of a retrained variant.

Scoring. Table 2 pools the 2019 and 2020 seasons, each cell weighted by its own per-season matched count, and uses the 00:00 and 12:00 UTC initialization times, the only times at which Pangu-Weather and IFS HRES are archived. Within each forecast source, all rows are scored on the samples that source’s tracker matched, so each comparison is homogeneous. Table 7 counts, under the Track columns, the forecast samples for which TempestExtremes found a storm near the anchored position at that lead time (TC-Next forecasts every sample with a best-track target); under the Intensity columns, the subset of those matched samples whose best-track record includes the 1-minute wind. In Tables 1 and 2, track and intensity are therefore scored on different samples: samples without a best-track 1-minute wind (USA_WIND) are dropped from intensity scoring only. In Table 1, WN-C is fine-tuned on IFS HRES analyses, so TC-Next is given HRES analysis frames in

place of the ERA5 history to keep it on the same analysis distribution. The WN-C tracker can predict that a storm has dissipated while the IBTrACS still reports it; those rows are dropped for both models, so the IBTrACS block scores fewer than the 867 samples Table 4 counts for 2025. Table 6 lists the scored samples per lead. Each of WN-C’s four ensemble members is scored separately and its errors averaged, rather than the ensemble being collapsed first: this measures the two models rather than the error cancellation an ensemble mean buys. TC-Next is trained against a deterministic backbone, and adapting it to an ensemble would require a different loss, which we leave to future work. Table 3 is scored on every held-out sample at all synoptic times; Table 8 lists those counts.

Table 6: Scored samples n at each lead for Table 1 (WP, 2025; TC-Next against the direct tracker of WN-C). Each row counts the samples on which both models of that anchor block are scored, namely those the WN-C tracker produced: where the WN-C tracker predicts that the storm has dissipated, the sample is dropped for both models. Intensity counts are lower because they also need a reported USA_WIND.

Anchor record	Track				Intensity			
	+6 h	+12 h	+18 h	+24 h	+6 h	+12 h	+18 h	+24 h
IBTrACS	810	774	743	710	602	592	579	564
TC Vitals	639	637	633	623	586	572	559	543

Table 7: Scored samples n at each lead for Table 2 (WP, pooled 2019–2020; TC-Next against TempestExtremes). Each row counts the samples on which both models of that forecast source are scored: the 00:00 and 12:00 UTC samples (the only initialization times at which WeatherBench2 archives Pangu-Weather and IFS HRES forecasts) that have a best-track target at that lead and for which TempestExtremes found the storm; TC-Next forecasts every sample with a target, so the counts are limited only by TempestExtremes. Intensity counts are lower because they also need a reported USA_WIND.

Forecast source	Track				Intensity			
	+6 h	+12 h	+18 h	+24 h	+6 h	+12 h	+18 h	+24 h
GraphCast	562	514	484	458	524	484	456	432
Pangu	562	510	481	450	527	487	459	431
IFS HRES	640	575	538	506	591	542	511	479

Table 8: Scored samples n at each lead for Table 3 (WP, 2019–2020; ablation of the IR branch). Both variants are scored on every held-out sample at all synoptic times, so the counts exceed those of Table 7 and the two tables are not directly comparable. Intensity counts are lower because they also need a reported USA_WIND.

IR branch	Track				Intensity			
	+6 h	+12 h	+18 h	+24 h	+6 h	+12 h	+18 h	+24 h
Fed and IR-ablated	1815	1758	1702	1646	1510	1468	1424	1378