

NEURAL OPERATORS APPROXIMATE STRONGLY CONTINUOUS CONVEX MONOTONE SEMIGROUPS

JONAS BLESSING, PHILIPP SCHMOCKER, AND ALESSANDRO SGARABOTTOLO

ABSTRACT. We approximate strongly continuous convex monotone semigroups by learning their Chernoff-type one-step operators with neural operators. First, we introduce the general class of so-called Chernoff-neural operators and show in a universal approximation theorem that they can approximate the Chernoff one-step operators arbitrarily well. By using stability estimates between weighted Hölder spaces, the one-step approximation error can be propagated through the iterations which yields universal approximation of the corresponding semigroup. Second, we introduce the more specialized class of envelope-neural operators for envelope semigroups which allows us to derive quantitative approximation rates. Finally, we illustrate the effectiveness of these neural operators in several numerical examples arising from non-linear partial differential equations, stochastic optimal control and stochastic processes under model uncertainty.

CONTENTS

1. Introduction	2
2. Setup and notation	3
2.1. Strongly continuous convex monotone semigroups	5
3. Chernoff-neural operators	7
3.1. Universal approximation	7
3.2. Qualitative approximation of the semigroup	10
4. Envelope-neural operators	12
4.1. Universal approximation	13
4.2. Quantitative approximation of the semigroup	17
5. Numerical experiments	18
5.1. Splitting schemes for semilinear PDEs	18
5.2. Stochastic optimal control	20
5.3. Stochastic processes under model uncertainty	22
Appendix A. Weighted Hölder spaces	25
Appendix B. Weighted Stone-Weierstrass theorems for Banach scales	29
Appendix C. Proof of Lemmas 5.1–5.4	33
C.1. Proof of Lemma 5.1	33
C.2. Proof of Lemma 5.2	34
C.3. Proof of Lemma 5.3	34
C.4. Proof of Lemma 5.4	35
References	36

Date: September 3, 2026.

Key words: convex monotone semigroup, Chernoff approximation, optimal control, Hamilton–Jacobi–Bellman equation, neural operator, universal approximation, convergence rates.

MSC2020 Subject Classification: primary 47H20, 65M15, 68T07; secondary 41A81, 49M25, 65J08.

1. INTRODUCTION

In this article, we show that strongly continuous convex monotone semigroups on spaces of continuous functions can be approximated by iterating a sequence of neural operators. Strongly continuous convex monotone semigroups have been studied in a series of articles [8–12, 21, 27, 54]. Starting from a sequence $(I_n)_{n \in \mathbb{N}}$ of Chernoff-type one-step operators $I_n: C_\kappa \rightarrow C_\kappa$ and a sequence $(h_n)_{n \in \mathbb{N}} \subset (0, \infty)$ with $h_n \rightarrow 0$, one can construct a semigroup

$$S(t)f := \lim_{n \rightarrow \infty} I_n^{k_n^t} f = \lim_{n \rightarrow \infty} \underbrace{(I_n \circ \dots \circ I_n)}_{k_n^t \text{ times}} f, \quad (1.1)$$

with $k_n^t := \max\{k \in \mathbb{N}_0 : kh_n \leq t\}$, whose infinitesimal generator is given by

$$Af = I'(0)f := \lim_{n \rightarrow \infty} \frac{I_n f - f}{h_n} \quad \text{for all } f \in C_b^\infty. \quad (1.2)$$

The comparison principle in [8] guarantees that the semigroup $(S(t))_{t \geq 0}$ is uniquely determined by the infinitesimal behavior of the one-step operators in equation (1.2).

The framework of strongly continuous convex monotone semigroups covers various applications such as stochastic optimal control, Hamilton–Jacobi–Bellman (HJB) equations, large deviations and stochastic processes under model uncertainty, see [5, 11–13, 21, 25, 27, 41, 42, 54]. It is also worth mentioning that convex monotone semigroups do not rely on the theory of viscosity solutions [17, 18, 47] but are consistent with it, i.e., $u(t) := S(t)f$ is a viscosity solution of the HJB equation

$$\begin{cases} \partial_t u(t) = Au(t), & t \in (0, \infty), \\ u(0) = f, \end{cases} \quad (1.3)$$

see [27, Theorem 6.3]. In particular, if the viscosity solution is unique, the Chernoff-type approximation (1.1) can be seen as a monotone approximation scheme [1–3, 14, 22, 34, 39] for the solution operator $f \mapsto u(t) := S(t)f$ of the HJB equation (1.3).

In this paper, we approximate the strongly continuous convex monotone semigroup $(S(t))_{t \geq 0}$ by learning, for sufficiently large $n \in \mathbb{N}$, the Chernoff-type one-step operator I_n with a neural operator Φ_n on a sufficiently rich subset of C_κ^α so that equation (1.1) yields

$$S(t)f \approx I_n^{k_n^t} f \approx \Phi_n^{k_n^t} f \quad \text{for all } f \in C_\kappa^\alpha, \quad (1.4)$$

where C_κ^α is a weighted Hölder space. First, we introduce the general class of so-called *Chernoff-neural operators*, which are based on the infinite-dimensional generalization of neural networks on weighted spaces in [19]. These neural operators map the infinite-dimensional input $f \in C_\kappa^\alpha$ via suitable hidden layer maps to the real hidden layer space, on which a scalar non-linear activation function is applied, followed by the multiplication with linear readout functions to obtain an output in C_κ^α . By generalizing the weighted Stone–Weierstrass theorem of [19] to the scale of weighted Hölder spaces, we establish a universal approximation theorem for the one-step operators $(I_n)_{n \in \mathbb{N}}$, see Theorem 3.5. The approximation error can then be propagated through the iterations to derive the approximation of the semigroup in equation (1.4), see Theorem 3.6.

Second, we introduce the more specialized but structure-preserving class of so-called *envelope-neural operators* if the sequence of one-step operators $(I_n)_{n \in \mathbb{N}}$ is of the form

$$(I_n f)(x) = \sup_{\lambda \in \Lambda} ((I_{n,\lambda} f)(x) - \eta_n(\lambda)h_n), \quad (1.5)$$

where Λ is a parameter set, $I_{n,\lambda}$ are linear operators and $\eta_n: \Lambda \rightarrow [0, \infty]$ are penalization functions. This framework covers, for instance, stochastic optimal control problems, HJB equations and transition semigroups of stochastic processes under model uncertainty. To make the supremum on the right-hand side of equation (1.5) tractable, we replace the supremum over Λ by a maximum over

finitely many trainable parameters $\lambda_1, \dots, \lambda_M \in \Lambda$. Since finite maxima can be represented by ReLU networks, we define envelope-neural operators as compositions of ReLU networks with finite subcollections of the operators $(I_{n,\lambda})_{\lambda \in \Lambda}$. Under suitable regularity assumptions on the parameter dependence, we show that envelope-neural operators can approximate the one-step operators $(I_n)_{n \in \mathbb{N}}$ arbitrarily well and subsequently propagate the resulting one-step approximation error through the iterations, see Theorem 4.6 and Theorem 4.7. Moreover, we derive explicit approximation rates in terms of the size of the neural operator and combine them with the convergence rates of the one-step operator in [12] to obtain quantitative error bounds for the approximation of the strongly continuous convex monotone semigroup, see Theorem 4.9 and Theorem 4.12.

Compared to the literature, our approach differs both from deep learning methods that approximate the solution of a partial differential equation (PDE) for fixed initial data and from operator-learning methods that approximate the full solution operator of the underlying PDE. For fixed initial data, our approach is closely related to deep splitting schemes [6, 58] and, more broadly, to neural PDE solvers such as the deep BSDE method [31], the deep Galerkin method [65], the deep Ritz method [23], physics-informed neural networks [62], deep backward dynamic programming [33] and random-feature models [28, 52, 56], see also [15, 29, 30, 51, 60, 69] in the context of HJB equations. For varying initial data, our approach is related to operator-learning methods for PDEs such as Fourier neural operators [46], neural integral operators [35], DeepONets [43, 48], generative equilibrium operators [38] and kernel methods [32], see also [44] for HJB equations. These results rely on universal approximation theorems for neural operators between infinite-dimensional function spaces starting with [16, 50] and followed by results, e.g., on non-Euclidean domains [26, 36, 37], Fréchet spaces [7], and weighted infinite-dimensional spaces [19, 63].

In contrast to these approaches, we do not approximate the solution operator $f \mapsto S(t)f$ directly but instead learn the Chernoff-type one-step operator I_n with a neural operator Φ_n and recover $S(t)$ by iteration. Hence, the training of Φ_n only requires evaluations of I_n , often available in closed form, instead of supervised PDE solution data. We illustrate our approach in three numerical examples from non-linear partial differential equations, stochastic optimal control, and stochastic processes under model uncertainty. While the Chernoff-neural operator provides a more general architecture and is straightforward to evaluate, the envelope-neural operator is particularly efficient in training, achieving accurate approximations with small architecture.

The rest of the article is organized as follows. In Section 2, we introduce strongly continuous convex monotone semigroups. In Section 3, we establish a universal approximation theorem for Chernoff-neural operators followed by the one for envelope-neural operators and their approximation rates in Section 4. Finally, we present several numerical experiments in Section 5 while some proofs are given in Appendices A–C.

2. SETUP AND NOTATION

Let $\kappa: \mathbb{R}^d \rightarrow (0, \infty)$ be a bounded continuous function satisfying

$$\sup_{x \in \mathbb{R}^d} \sup_{|y| \leq 1} \frac{\kappa(x)}{\kappa(x-y)} < \infty. \quad (2.1)$$

Typical examples that we have in mind are polynomial weights $\kappa(x) := (1 + |x|^q)^{-1}$ with $q \geq 0$. We define C_κ as the space of all continuous functions $f: \mathbb{R}^d \rightarrow \mathbb{R}$ satisfying

$$\|f\|_\kappa := \sup_{x \in \mathbb{R}^d} |f(x)|\kappa(x) < \infty.$$

An operator $S: C_\kappa \rightarrow C_\kappa$ is called convex if $S(\lambda f + (1-\lambda)g) \leq \lambda Sf + (1-\lambda)Sg$ for all $\lambda \in [0, 1]$ and $f, g \in C_\kappa$ and monotone if $Sf \leq Sg$ for all $f, g \in C_\kappa$ with $f \leq g$, where $f \leq g$ means that $f(x) \leq g(x)$ for all $x \in \mathbb{R}^d$. Furthermore, we write $f_n \downarrow f$ if a sequence $(f_n)_{n \in \mathbb{N}} \subset C_\kappa$ decreases

pointwise to another function $f \in C_\kappa$ and define $B_{C_\kappa}(r) := \{f \in C_\kappa : \|f\|_\kappa \leq r\}$. Let Lip_b be the space of all bounded Lipschitz functions $f: \mathbb{R}^d \rightarrow \mathbb{R}$ and denote by $\text{Lip}_b(r)$ the set of all r -Lipschitz functions $f: \mathbb{R}^d \rightarrow \mathbb{R}$ with $\|f\|_\infty \leq r$. Moreover, the space C_b^1 (resp., C_b^∞ ; resp. C_c^∞) consists of all bounded differentiable (resp., infinitely differentiable; resp. compactly supported infinitely differentiable) functions $f: \mathbb{R}^d \rightarrow \mathbb{R}$ such that all existing derivatives are bounded.

Throughout this article, we endow C_κ with the mixed topology between $\|\cdot\|_\kappa$ and the topology of uniform convergence on compact sets, i.e., the strongest locally convex topology on C_κ that coincides on $\|\cdot\|_\kappa$ -bounded sets with the topology of uniform convergence on compact subsets, see [68] for a detailed introduction. Choosing the mixed topology rather than the norm topology is crucial for the analysis of strongly continuous convex monotone semigroups, see [8, 11, 12]. Although the mixed topology is not metrizable, for convex monotone operators $S: C_\kappa \rightarrow C_\kappa$, continuity w.r.t. the mixed topology is equivalent to sequential continuity and to continuity from above, see [20, 53]. Moreover, a sequence $(f_n)_{n \in \mathbb{N}} \subset C_\kappa$ converges to $f \in C_\kappa$ if and only if

$$\sup_{n \in \mathbb{N}} \|f_n\|_\kappa < \infty \quad \text{and} \quad \lim_{n \rightarrow \infty} \|f - f_n\|_{\infty, K} = 0$$

for all compact subsets $K \Subset \mathbb{R}^d$, where $\|f\|_{\infty, K} := \sup_{x \in K} |f(x)|$, see [27, Proposition A.4]. If not stated otherwise, all limits in C_κ are taken w.r.t. the mixed topology and compact subsets are denoted by $K \Subset \mathbb{R}^d$. The mixed topology can also be seen as a strict topology, see [24, 40, 64].

In order to derive a universal approximation result, we further introduce weighted Hölder spaces as follows. For every $\alpha \in [0, 1]$, we denote by C_κ^α the space of all functions $f: \mathbb{R}^d \rightarrow \mathbb{R}$ satisfying

$$\|f\|_{\alpha, \kappa} := \max \left\{ \sup_{x \in \mathbb{R}^d} |f(x)| \kappa(x), \sup_{x \neq y} \frac{|f(x) - f(y)|}{|x - y|^\alpha} \bar{\kappa}(x, y) \right\} < \infty,$$

where $\bar{\kappa}(x, y) := \frac{2\kappa(x)\kappa(y)}{\kappa(x) + \kappa(y)} = \frac{2}{\kappa(x)^{-1} + \kappa(y)^{-1}}$ satisfies $\min(\kappa(x), \kappa(y)) \leq \bar{\kappa}(x, y) \leq 2 \min(\kappa(x), \kappa(y))$. By Theorem A.1, the space $(C_\kappa^\alpha, \|\cdot\|_{\alpha, \kappa})$ is a Banach space satisfying $C_\kappa^1 = \text{Lip}_b$ if $\kappa \equiv 1$. Moreover, since the norms $\|\cdot\|_{0, \kappa}$ and $\|\cdot\|_\kappa$ are equivalent, we have $C_\kappa^0 = C_\kappa$ as sets, but we keep the notation C_κ^0 to distinguish the norm topology from the mixed topology on C_κ . In addition, Proposition A.2 guarantees that the embedding $C_\kappa^\alpha \hookrightarrow C_{\kappa'}^{\alpha'}$ is continuous if $0 \leq \alpha' \leq \alpha \leq 1$ and $\kappa' \lesssim \kappa$, where $\kappa' \lesssim \kappa$ means that $\kappa' \leq c\kappa$ for some constant $c \geq 0$. By Theorem A.3, the embedding $C_\kappa^\alpha \hookrightarrow C_{\kappa'}^{\alpha'}$ is even compact if $0 \leq \alpha' < \alpha \leq 1$ and $\kappa' \ll \kappa$, where $\kappa' \ll \kappa$ means that $\lim_{|x| \rightarrow \infty} \frac{\kappa'(x)}{\kappa(x)} = 0$. Moreover, we denote by c_κ^α the little weighted Hölder space consisting of all functions $f \in C_\kappa^\alpha$ with

$$\lim_{|x| \rightarrow \infty} |f(x)| \kappa(x) = 0 \quad \text{and} \quad \lim_{\delta \rightarrow 0} \sup_{\substack{x \neq y, \\ |x - y| < \delta}} \frac{|f(x) - f(y)|}{|x - y|^\alpha} \bar{\kappa}(x, y) = 0.$$

Note that $c_\kappa^\alpha \subset C_\kappa^\alpha$ is a closed linear subspace and therefore itself a Banach space. In particular, the embedding $C_\kappa^\alpha \hookrightarrow c_{\kappa'}^{\alpha'}$ is continuous and dense if $0 \leq \alpha' < \alpha \leq 1$ and $\kappa' \ll \kappa$, see Theorem A.2 and Theorem A.4. We define $B_{C_\kappa^\alpha}(r) := \{f \in C_\kappa^\alpha : \|f\|_{\alpha, \kappa} \leq r\}$ for all $r \geq 0$.

Throughout this article, we also use the following notations. Let $\mathbb{R}_+ := \{x \in \mathbb{R} : x \geq 0\}$. We define $B_{\mathbb{R}^d}(r) := \{x \in \mathbb{R}^d : |x| \leq r\}$ for all $r \geq 0$, where $|\cdot|$ is the Euclidean norm. The transpose of a vector $x \in \mathbb{R}^d$ is denoted by $x^\top$, compact subsets are denoted by $K \Subset \mathbb{R}^d$ and the translation of a function $f: \mathbb{R}^d \rightarrow \mathbb{R}$ by a vector $x \in \mathbb{R}^d$ is denoted by $(\tau_x f)(y) := f(x + y)$ for all $y \in \mathbb{R}^d$. The set $\mathcal{M}_\kappa$ consists of all signed Borel measures μ with $\int_{\mathbb{R}^d} \frac{1}{\kappa(x)} |\mu|(dx) < \infty$. Moreover, for topological spaces X, Y , the space $C_b(X; Y)$ consists of all uniformly bounded continuous functions $f: X \rightarrow Y$, while $C(X)$ consists of all continuous functions $f: X \rightarrow \mathbb{R}$. For a Banach space X , we denote by $C^1(X)$ the space of all continuously differentiable functions $f: X \rightarrow \mathbb{R}$. If (X, d) is a metric space, we define $\mathring{B}_d(x, r) := \{y \in X : d(x, y) < r\}$ for all $r > 0$ and $x \in X$. Finally, the set $\mathbb{S}_+^d \subset \mathbb{R}^{d \times d}$ consists of all symmetric positive semi-definite matrices.

2.1. Strongly continuous convex monotone semigroups. In this section, we recall the results from [8, Section 5] and [12, Section 2] about Chernoff-type approximations of strongly continuous convex monotone semigroups.

Definition 2.1. A family $(S(t))_{t \geq 0}$ of operators $S(t): C_\kappa \rightarrow C_\kappa$ is called *strongly continuous convex monotone semigroup* if the following conditions are satisfied:

- (i) $S(t)$ is convex and monotone with $S(t)f_n \downarrow 0$ for all $t \geq 0$ and $f_n \downarrow 0$.
- (ii) $S(0)f = f$ and $S(s+t)f = S(s)S(t)f$ for all $s, t \geq 0$ and $f \in C_\kappa$.
- (iii) $\sup_{t \in [0, T]} \|S(t)\frac{f}{\kappa}\|_\kappa < \infty$ for all $r, T \geq 0$.
- (iv) $f = \lim_{t \downarrow 0} S(t)f$ for all $f \in C_\kappa$.

Furthermore, the generator of the semigroup is defined by

$$A: D(A) \rightarrow C_\kappa, f \mapsto \lim_{h \downarrow 0} \frac{S(h)f - f}{h},$$

where the domain consists of all $f \in C_\kappa$ such that the previous limit exists.

Let $(I_n)_{n \in \mathbb{N}}$ be a sequence of operators $I_n: C_\kappa \rightarrow C_\kappa$ and $(h_n)_{n \in \mathbb{N}} \subset (0, \infty)$ be a sequence with $h_n \rightarrow 0$. For every $t \geq 0$, $n \in \mathbb{N}$ and $f \in C_\kappa$, we define

$$I(\pi_n^t)f := I_n^{k_n^t}f = \underbrace{(I_n \circ \dots \circ I_n)}_{k_n^t \text{ times}}f,$$

where $k_n^t := \max\{k \in \mathbb{N}_0: kh_n \leq t\}$ and $\pi_n^t := \{0, h_n, \dots, k_n^t h_n\}$. In addition, for every $f \in C_\kappa$ such that the following limit exists, we define

$$I'(0)f := \lim_{n \rightarrow \infty} \frac{I_n f - f}{h_n} \in C_\kappa.$$

The following conditions guarantee that the sequence of iterated operators converges to a strongly continuous convex monotone semigroup which is uniquely determined by the infinitesimal behavior of the one-step operators.

Assumption 2.2. Suppose that the following conditions are satisfied:

- (i) I_n is convex and monotone with $I_n 0 = 0$ for all $n \in \mathbb{N}$.
- (ii) There exists $\omega \geq 0$ with $\|I_n f - I_n g\|_\kappa \leq e^{\omega h_n} \|f - g\|_\kappa$ for all $f, g \in C_\kappa$ and $n \in \mathbb{N}$.
- (iii) For every $\varepsilon > 0$, $r, T \geq 0$, and $K \Subset \mathbb{R}^d$ there exist $c \geq 0$ and $K' \Subset \mathbb{R}^d$ with

$$\|I(\pi_n^t)f - I(\pi_n^t)g\|_{\infty, K} \leq c\|f - g\|_{\infty, K'} + \varepsilon$$

for all $t \in [0, T]$, $n \in \mathbb{N}$ and $f, g \in B_{C_\kappa}(r)$.

- (iv) For every $\varepsilon > 0$, there exists $\delta > 0$ and $n_0 \in \mathbb{N}$ with

$$I_n(\tau_x f) \leq \tau_x I_n f + \frac{r\varepsilon h_n}{\kappa}$$

for all $r \geq 0$, $f \in \text{Lip}_b(r)$, $x \in B_{\mathbb{R}^d}(\delta)$ and $n \geq n_0$.

- (v) It holds $I_n: \text{Lip}_b(r) \rightarrow \text{Lip}_b(e^{\omega h_n} r)$ for all $r \geq 0$ and $n \in \mathbb{N}$.
- (vi) The limit $I'(0)f \in C_\kappa$ exists for all $f \in C_b^\infty$.

It follows from [8, Lemma C.2] that condition (iii) is equivalent to the following: for every $T \geq 0$, $K \Subset \mathbb{R}^d$ and $(f_k)_{k \in \mathbb{N}} \subset C_\kappa$ with $f_k \downarrow 0$, it holds

$$\sup_{(t, x) \in [0, T] \times K} \sup_{n \in \mathbb{N}} (I(\pi_n^t)f_k)(x) \downarrow 0.$$

Moreover, we refer to [12, Section 2.4] for sufficient conditions on the one-step operators I_n which guarantee that condition (vi) is satisfied and which can easily be verified in applications. The next theorem follows immediately from [12, Theorem 2.11 and Corollary 2.15].

Theorem 2.3. *Suppose that Assumption 2.2 is satisfied. Then, there exists a strongly continuous convex monotone semigroup $(S(t))_{t \geq 0}$ on C_κ with generator $A: D(A) \rightarrow C_\kappa$ given by*

$$S(t)f := \lim_{n \rightarrow \infty} I(\pi_n^t)f \quad \text{for all } t \geq 0 \text{ and } f \in C_\kappa.$$

In addition, the following statements are true:

- (i) *It holds $f \in D(A)$ and $Af = I'(0)f$ for all $f \in C_\kappa$ such that $I'(0)f \in C_\kappa$ exists. In particular, this is valid for all $f \in C_b^\infty$.*
- (ii) *It holds $\|S(t)f - S(t)g\|_\kappa \leq e^{\omega t}\|f - g\|_\kappa$ for all $t \geq 0$ and $f, g \in C_\kappa$.*
- (iii) *For every $\varepsilon > 0$, $r, T \geq 0$ and $K \in \mathbb{R}^d$, there exists $c \geq 0$ and $K' \in \mathbb{R}^d$ with*

$$\|S(t)f - S(t)g\|_{\infty, K} \leq c\|f - g\|_{\infty, K'} + \varepsilon$$

for all $t \in [0, T]$ and $f, g \in B_{C_\kappa}(r)$.

- (iv) *For every $\varepsilon > 0$, there exists $\delta > 0$ with*

$$S(t)(\tau_x f) \leq \tau_x S(t)f + \frac{e^{\omega t} r \varepsilon t}{\kappa}$$

for all $r, t \geq 0$, $f \in \text{Lip}_b(r)$ and $x \in B_{\mathbb{R}^d}(\delta)$.

- (v) *It holds $S(t): \text{Lip}_b(r) \rightarrow \text{Lip}_b(e^{\omega t} r)$ for all $r, t \geq 0$.*

Furthermore, let $(T(t))_{t \geq 0}$ be another strongly continuous convex monotone semigroup on C_κ with generator $B: D(B) \rightarrow C_\kappa$ satisfying the conditions (iv) and (v), $C_b^\infty \subset D(B)$ and

$$Af = Bf \quad \text{for all } f \in C_b^\infty.$$

Then, it holds $S(t)f = T(t)f$ for all $t \geq 0$ and $f \in C_\kappa$.

Corollary 2.4. *Let Assumption 2.2 be satisfied and denote by $(S(t))_{t \geq 0}$ the strongly continuous convex monotone semigroup from Theorem 2.3. Assume that there exists $\alpha \in (0, 1]$ and $\tilde{\omega} \geq 0$ with*

$$\|I_n f\|_{\alpha, \kappa} \leq e^{\tilde{\omega} h_n} \|f\|_{\alpha, \kappa} \quad \text{for all } n \in \mathbb{N} \text{ and } f \in C_\kappa^\alpha.$$

Then, it holds $\|S(t)f\|_{\alpha, \kappa} \leq e^{\tilde{\omega} t} \|f\|_{\alpha, \kappa}$ for all $t \geq 0$ and $f \in C_\kappa^\alpha$. Furthermore,

$$\lim_{n \rightarrow \infty} \sup_{f \in B_{C_\kappa^\alpha}(r)} \sup_{t \in [0, T]} \|S(t)f - I(\pi_n^t)f\|_{\alpha', \kappa'} = 0$$

for all $r, T \geq 0$, $\alpha' \in [0, \alpha)$ and $\kappa' \ll \kappa$.

Proof. For every $t \geq 0$ and $f \in C_\kappa^\alpha$, it follows by induction that

$$\|I(\pi_n^t)f\|_{\alpha, \kappa} \leq e^{\tilde{\omega} t} \|f\|_{\alpha, \kappa} \quad \text{for all } n \in \mathbb{N}, \quad (2.2)$$

which, by taking the limit, implies

$$\|S(t)f\|_{\alpha, \kappa} \leq e^{\tilde{\omega} t} \|f\|_{\alpha, \kappa}. \quad (2.3)$$

Now, let $r, T \geq 0$, $\alpha' \in [0, \alpha)$, $\kappa' \ll \kappa$ and $\varepsilon' > 0$. Since $\kappa' \ll \kappa$, Assumption 2.2(iii) and Theorem 2.3(iii) imply that there exists $\delta > 0$ with

$$\|S(t)f - S(t)g\|_{\kappa'} < \frac{\varepsilon'}{5} \quad \text{and} \quad \|I(\pi_n^t)f - I(\pi_n^t)g\|_{\kappa'} < \frac{\varepsilon'}{5} \quad (2.4)$$

for all $t \in [0, T]$, $n \in \mathbb{N}$ and $f, g \in B_{C_\kappa}(r)$ with $\|f - g\|_{\kappa'} < \delta$. By Theorem A.3, there exist $f_1, \dots, f_N \in B_{C_\kappa^\alpha}(r)$ such that, for every $f \in B_{C_\kappa^\alpha}(r)$, there exists $j \in \{1, \dots, N\}$ with

$$\|f - f_j\|_{\kappa'} < \delta. \quad (2.5)$$

In addition, by using $\kappa' \ll \kappa$ and the strong continuity, there exist $t_1, \dots, t_M \in [0, T]$ such that, for every $t \in [0, T]$, there exists $i \in \{1, \dots, M\}$ with

$$\|S(t)f_j - S(t_i)f_j\|_{\kappa'} < \frac{\varepsilon'}{5} \quad \text{and} \quad \|I(\pi_n^t)f_j - I(\pi_n^{t_i})f_j\|_{\kappa'} < \frac{\varepsilon'}{5} \quad \text{for all } j \in \{1, \dots, N\}. \quad (2.6)$$

Theorem A.3 yields the existence of a subsequence $(n_l)_{l \in \mathbb{N}} \subset \mathbb{N}$ and functions $g_{ij} \in C_\kappa^\alpha$ with

$$\lim_{l \rightarrow \infty} \|I(\pi_{n_l}^{t_i})f_j - g_{ij}\|_{\kappa'} = 0 \quad \text{for all } i \in \{1, \dots, M\} \text{ and } j \in \{1, \dots, N\}.$$

Since $I(\pi_{n_l}^{t_i})f_j \rightarrow S(t_i)f_j$ in the mixed topology, we obtain $g_{ij} = S(t_i)f_j$. In particular, the limit does not depend on the choice of the convergent subsequence, whence there exists $n_0 \in \mathbb{N}$ with

$$\|I(\pi_n^{t_i})f_j - S(t_i)f_j\|_{\kappa'} < \frac{\varepsilon'}{5} \quad \text{for all } i \in \{1, \dots, M\}, j \in \{1, \dots, N\} \text{ and } n \geq n_0. \quad (2.7)$$

For every $t \in [0, T]$, $f \in B_{C_\kappa^\alpha}(r)$ and $n \geq n_0$, the inequalities (2.4)-(2.7) imply

$$\begin{aligned} \|I(\pi_n^t)f - S(t)f\|_{\kappa'} &\leq \|I(\pi_n^t)f - I(\pi_n^t)f_j\|_{\kappa'} + \|I(\pi_n^t)f_j - I(\pi_n^{t_i})f_j\|_{\kappa'} \\ &\quad + \|I(\pi_n^{t_i})f_j - S(t_i)f_j\|_{\kappa'} + \|S(t_i)f_j - S(t)f_j\|_{\kappa'} \\ &\quad + \|S(t)f_j - S(t)f\|_{\kappa'} \\ &< \varepsilon'. \end{aligned}$$

In order to derive the convergence w.r.t. $\|\cdot\|_{\alpha', \kappa'}$, we observe that

$$\mathcal{F} := \overline{\mathcal{E}}^{\|\cdot\|_{\alpha', \kappa'}} \cup \{0\} \quad \text{with} \quad \mathcal{E} := \{I(\pi_n^t)f - S(t)f : t \in [0, T], n \in \mathbb{N}, f \in B_{C_\kappa^\alpha}(r)\}$$

is bounded in C_κ^α due to the inequalities (2.2) and (2.3) and therefore compact in $C_{\kappa'}^{\alpha'}$ by Theorem A.3. Since $(\mathcal{F}, \|\cdot\|_{\alpha', \kappa'}) \hookrightarrow (\mathcal{F}, \|\cdot\|_{\kappa'})$ is a continuous bijection from a compact space into a Hausdorff space, it is a homeomorphism. Hence, for every $\varepsilon > 0$, there exists $\varepsilon' > 0$ such that, for every $g \in \mathcal{F}$, we have

$$\|g\|_{\kappa'} < \varepsilon' \quad \Rightarrow \quad \|g\|_{\alpha', \kappa'} < \varepsilon.$$

Combining this estimate with the first part of the proof yields the claim. $\square$

3. CHERNOFF-NEURAL OPERATORS

In this section, we introduce a very general neural operator architecture to approximate the one-step operators $(I_n)_{n \in \mathbb{N}}$ from the previous section. In particular, by learning the one-step operators uniformly over different functions, we can then iterate the neural operators to approximate the corresponding strongly continuous convex monotone semigroup $(S(t))_{t \geq 0}$.

3.1. Universal approximation. Let $\alpha \in (0, 1]$, $p \in (1, \infty)$ and $\kappa, \kappa_0 : \mathbb{R}^d \rightarrow (0, \infty)$ be bounded continuous functions satisfying $\kappa_0 \lesssim \kappa$. Furthermore, we fix a sequence $(I_n)_{n \in \mathbb{N}}$ of operators $I_n : C_\kappa \rightarrow C_\kappa$ and a sequence $(h_n)_{n \in \mathbb{N}} \subset (0, \infty)$ with $h_n \rightarrow 0$ such that Assumption 2.2 is valid.

Assumption 3.1. The operators $(I_n)_{n \in \mathbb{N}}$ can be extended to operators $I_n : C_{\kappa_0} \rightarrow C_{\kappa_0}$. Furthermore, there exists a set $\mathcal{K}$ of bounded continuous functions $\kappa' : \mathbb{R}^d \rightarrow (0, \infty)$ with $\kappa_0 \lesssim \kappa' \lesssim \kappa$ such that $\kappa \in \mathcal{K}$ and the following conditions are satisfied:

- (i) For every $\kappa_1, \kappa_2 \in \mathcal{K}$ with $\kappa_1 \lesssim \kappa_2$, there exists $\kappa_3 \in \mathcal{K}$ with $\kappa_1 \lesssim \kappa_3 \lesssim \kappa_2$.
- (ii) It holds $\|I_n f\|_{\alpha', \kappa'} \leq e^{\omega h_n} \|f\|_{\alpha', \kappa'}$ for all $\alpha' \in (0, \alpha]$, $\kappa' \in \mathcal{K}$, $n \in \mathbb{N}$ and $f \in C_{\kappa'}^{\alpha'}$.
- (iii) For every $\alpha' \in [0, \alpha]$, $\mathcal{K} \ni \kappa' \lesssim \kappa$ and $n \in \mathbb{N}$, there exists $c \geq 0$ with

$$\|I_n f - I_n g\|_{\alpha', \kappa'} \leq c \|f - g\|_{\alpha', \kappa'} \quad \text{for all } f, g \in C_{\kappa'}^{\alpha'}.$$

In order to approximate the operators $(I_n)_{n \in \mathbb{N}}$ on C_κ^α , we follow the infinite-dimensional generalization of neural networks over weighted spaces in [19]. For every $\alpha' \in [0, \alpha]$ and $\mathcal{K} \ni \kappa' \lesssim \kappa$, we consider the normed vector space $C_{\kappa, \kappa'}^{\alpha, \alpha'} := (C_\kappa^\alpha, \|\cdot\|_{\alpha', \kappa'})$ and define the weight function

$$\psi_{\alpha', \kappa', p} : C_{\kappa, \kappa'}^{\alpha, \alpha'} \rightarrow (0, \infty), \quad f \mapsto (1 + \|f\|_{\alpha, \kappa})^p.$$

By Theorem A.3, the set $K_R := \psi_{\alpha', \kappa', p}^{-1}((0, R]) \subset C_{\kappa, \kappa'}^{\alpha, \alpha'}$ is compact and thus $\psi_{\alpha', \kappa', p}$ is an admissible weight in the sense of [19, Definition 2.1]. For a normed space Y , we denote by $\mathcal{B}_p(C_{\kappa, \kappa'}^{\alpha, \alpha'}; Y)$ the completion of the space of all uniformly bounded continuous operators $T: C_{\kappa, \kappa'}^{\alpha, \alpha'} \rightarrow Y$ w.r.t.

$$\|T\|_{\mathcal{B}_p(C_{\kappa, \kappa'}^{\alpha, \alpha'}; Y)} := \sup_{f \in C_{\kappa}^{\alpha}} \frac{\|Tf\|_Y}{(1 + \|f\|_{\alpha, \kappa})^p}.$$

Here, an operator $T: C_{\kappa, \kappa'}^{\alpha, \alpha'} \rightarrow Y$ is called uniformly bounded if $\sup\{\|Tf\|_Y : f \in C_{\kappa, \kappa'}^{\alpha, \alpha'}\} < \infty$. In the case of $Y := \mathbb{R}$, it follows from [19, Lemma 2.7] that $T \in \mathcal{B}_p(C_{\kappa, \kappa'}^{\alpha, \alpha'}; \mathbb{R})$ if and only if $T|_{K_R}$ is continuous for all $R > 0$ and $\lim_{R \rightarrow \infty} \sup_{f \in C_{\kappa}^{\alpha} \setminus K_R} \frac{|Tf|}{\psi_{\alpha', \kappa', p}(f)} = 0$.

Now, we introduce Chernoff-neural operators mapping the input $f \in C_{\kappa}^{\alpha}$ via some mappings $h \in \mathcal{H}$ to the real hidden layer space, on which a non-linear activation function $\rho \in C(\mathbb{R})$ is applied, and then return the output via some linear readouts $\mathcal{L} \subset C_{\kappa}^{\alpha}$ into C_{κ}^{α} , see Figure 1.

Definition 3.2. For a set $\mathcal{H}$ consisting of mappings $h: C_{\kappa_0} \rightarrow \mathbb{R}$, $\rho \in C(\mathbb{R})$ and a subset $\mathcal{L} \subset C_{\kappa}^{\alpha}$, a *Chernoff-neural operator* is a mapping of the form

$$\Phi: C_{\kappa_0} \rightarrow C_{\kappa}^{\alpha}, f \mapsto \sum_{m=1}^M y_m \rho(h_m(f)),$$

where $M \in \mathbb{N}$ is the number of neurons, $h_1, \dots, h_M \in \mathcal{H}$ are hidden layer maps and $y_1, \dots, y_M \in \mathcal{L}$ are linear readouts. We collect all operators of this form in the set $\mathcal{CN}_{C_{\kappa_0}, C_{\kappa}^{\alpha}}^{\mathcal{H}, \rho, \mathcal{L}}$.




FIGURE 1. A Chernoff-neural operator $\Phi: C_{\kappa_0} \rightarrow C_{\kappa}^{\alpha}$ with additive family $\mathcal{H}$, activation function $\rho \in C(\mathbb{R})$, linear readout $\mathcal{L} \subset C_{\kappa}^{\alpha}$ and $M = 3$ neurons.

Assumption 3.3. Suppose that the following conditions are satisfied:

- (i) The set $\mathcal{H}$ is a vector space, contains all constant mappings and is point separating on C_{κ_0} , i.e., for every $f, g \in C_{\kappa_0}$ with $f \neq g$ there exists $h \in \mathcal{H}$ with $h(f) \neq h(g)$. Moreover, for every $0 \leq \alpha' < \alpha'' \leq \alpha$, $\kappa' \lesssim \kappa'' \in \mathcal{K}$ and $h \in \mathcal{H}$, the mapping $h: C_{\kappa'', \kappa'}^{\alpha'', \alpha'} \rightarrow \mathbb{R}$ is continuous and satisfies

$$c_{\alpha'', \kappa'', h} := \sup_{f \in C_{\kappa''}^{\alpha''}} \frac{1 + |h(f)|}{1 + \|f\|_{\alpha'', \kappa''}} < \infty.$$

- (ii) The function $\rho \in C(\mathbb{R})$ is non-polynomial with $\lim_{|x| \rightarrow \infty} \frac{|\rho(x)|}{(1+|x|)^p} = 0$.
- (iii) The set $\mathcal{L} \subset C_{\kappa}^{\alpha}$ is a dense linear subspace of $C_{\kappa'}^{\alpha'}$ for all $0 \leq \alpha' < \alpha$ and $\mathcal{K} \ni \kappa' \lesssim \kappa$.

Example 3.4. As an example for Assumption 3.3(i), we consider the set

$$\mathcal{H} := \left\{ C_{\kappa_0} \rightarrow \mathbb{R}, f \mapsto \int_{\mathbb{R}^d} f(x) \nu(dx) + b : \nu \in \mathcal{M}_{\kappa_0}, b \in \mathbb{R} \right\}.$$

Since $\mathcal{H}$ contains the Dirac measures $\{\delta_x : x \in \mathbb{R}^d\}$, it is point separating. Moreover, for every $h \in \mathcal{H}$ of the form $h(f) := \int_{\mathbb{R}^d} f(x) \nu(dx) + b$, $\alpha'' \in [0, \alpha]$ and $\kappa'' \in \mathcal{K}$, it holds

$$\sup_{f \in C_{\kappa''}^{\alpha''}} \frac{1 + |h(f)|}{1 + \|f\|_{\alpha'', \kappa''}} \leq \sup_{f \in C_{\kappa''}^{\alpha''}} \frac{1 + \|f\|_{\kappa_0} c_\nu + |b|}{1 + \|f\|_{\alpha'', \kappa''}} \leq \sup_{f \in C_{\kappa''}^{\alpha''}} \frac{1 + c_{\alpha'', \kappa''} \|f\|_{\alpha'', \kappa''} c_\nu + |b|}{1 + \|f\|_{\alpha'', \kappa''}} < \infty,$$

where $c_\nu := \int_{\mathbb{R}^d} \frac{1}{\kappa_0(x)} |\nu|(dx)$ and $c_{\alpha'', \kappa''}$ is the operator norm of the embedding $C_{\kappa''}^{\alpha''} \hookrightarrow C_{\kappa_0}$.

We further observe that, for a bounded and non-polynomial $\tilde{\rho} \in C(\mathbb{R})$ and a fixed probability measure $\tilde{\nu} \in \mathcal{M}_{\kappa_0}$ having full support, the subset

$$\tilde{\mathcal{H}} := \left\{ C_{\kappa_0} \rightarrow \mathbb{R}, f \mapsto \int_{\mathbb{R}^d} f(x) \varphi(x) \tilde{\nu}(dx) + b : \varphi \in \mathcal{N}_{\mathbb{R}^d, \mathbb{R}}^{\tilde{\rho}} \right\}$$

also satisfies Assumption 3.3(i). Note that $\tilde{\mathcal{H}}$ can be implemented on a computer by approximating the integral with samples from $\tilde{\nu}$. Here, the space $\mathcal{N}_{\mathbb{R}^d, \mathbb{R}}^{\tilde{\rho}}$ consists of all neural networks

$$\varphi : \mathbb{R}^d \rightarrow \mathbb{R}, x \mapsto \sum_{k=1}^K y_k \tilde{\rho}(a_k^\top x + b_k),$$

where $K \in \mathbb{N}$ is the number of neurons, $a_1, \dots, a_K \in \mathbb{R}^d$ are weights, $b_1, \dots, b_K \in \mathbb{R}$ are biases, $y_1, \dots, y_K \in \mathbb{R}$ are linear readouts and $\tilde{\rho}$ is a (bounded) activation function. Indeed, for every $f, g \in C_{\kappa_0}$ with $f \neq g$, we define $u := f - g \neq 0$, $c := \int_{\mathbb{R}^d} |u| d\tilde{\nu} > 0$ and $d\mu := \frac{|u|}{c} d\tilde{\nu}$. Since $\mathcal{N}_{\mathbb{R}^d, \mathbb{R}}^{\tilde{\rho}}$ is dense in $L^1(\mu)$ by [45, Proposition 1], there is $\varphi \in \mathcal{N}_{\mathbb{R}^d, \mathbb{R}}^{\tilde{\rho}}$ with $\|\varphi - \text{sgn}(u(\cdot))\|_{L^1(\mu)} < 1$. Hence,

$$\left| \frac{1}{c} \int_{\mathbb{R}^d} u(x) \varphi(x) \tilde{\nu}(dx) - 1 \right| = \left| \int_{\mathbb{R}^d} \text{sgn}(u(x)) (\varphi(x) - \text{sgn}(u(x))) \mu(dx) \right| \leq \|\varphi - \text{sgn}(u(\cdot))\|_{L^1(\mu)} < 1$$

and therefore $\int_{\mathbb{R}^d} f(x) \varphi(x) \tilde{\nu}(dx) \neq \int_{\mathbb{R}^d} g(x) \varphi(x) \tilde{\nu}(dx)$.

As an example for Assumption 3.3(iii), we can choose $\mathcal{L} := \mathcal{N}_{\mathbb{R}^d, \mathbb{R}}^{\tilde{\rho}} \subset C_{\kappa}^{\alpha}$. For instance, if $\alpha = 1$, $\kappa(x) = (1 + |x|^q)^{-1}$ with $q > 0$ and $\tilde{\rho} \in C^1(\mathbb{R})$ is non-polynomial with $\lim_{|x| \rightarrow \infty} \frac{|\tilde{\rho}(x)|}{(1+|x|)^q} = 0$, the set $\mathcal{N}_{\mathbb{R}^d, \mathbb{R}}^{\tilde{\rho}}$ is dense in C_{κ}^1 by [57, Theorem 2.7] and therefore also dense in $C_{\kappa'}^{\alpha'}$ for all $0 \leq \alpha' < 1$ and $\mathcal{K} \ni \kappa' \ll \kappa$, see Theorem A.4.

In order to obtain the following universal approximation result, we adapt the arguments from [19, Theorem 4.13] to establish the universality of Chernoff-neural operators.

Theorem 3.5. *Suppose that the Assumptions 3.1 and 3.3 are satisfied. Then, for every $k, n \in \mathbb{N}$, $0 \leq \alpha'_l < \alpha_l \leq \alpha$ and $\kappa'_l \ll \kappa_l \in \mathcal{K}$ for $l = 1, \dots, k$ and $\varepsilon > 0$, there exists $\Phi_n \in \mathcal{CN}_{C_{\kappa_0}, C_{\kappa}}^{\mathcal{H}, \rho, \mathcal{L}}$ with*

$$\max_{l=1, \dots, k} \sup_{f \in C_{\kappa'_l}^{\alpha'_l}} \frac{\|I_n f - \Phi_n f\|_{\alpha'_l, \kappa'_l}}{(1 + \|f\|_{\alpha_l, \kappa_l})^p} < \varepsilon.$$

Proof. First, we show that $I_n \in \mathcal{B}_p(C_{\kappa_l, \kappa'_l}^{\alpha_l, \alpha'_l}; C_{\kappa'_l}^{\alpha'_l})$ for all $n \in \mathbb{N}$ and $l = 1, \dots, k$. To do so, for every $R \in \mathbb{N}$, we define $I_n^{(R)} := T^{(R)} \circ I_n : C_{\kappa_l, \kappa'_l}^{\alpha_l, \alpha'_l} \rightarrow C_{\kappa'_l}^{\alpha'_l}$, where

$$T^{(R)} : C_{\kappa'_l}^{\alpha'_l} \rightarrow C_{\kappa'_l}^{\alpha'_l}, g \mapsto \begin{cases} g & \text{if } \|g\|_{\alpha'_l, \kappa'_l} \leq R, \\ \frac{R}{\|g\|_{\alpha'_l, \kappa'_l}} g & \text{if } \|g\|_{\alpha'_l, \kappa'_l} > R. \end{cases}$$

Assumption 3.1(iii) and Theorem A.2 imply that $I_n^{(R)} := T^{(R)} \circ I_n : C_{\kappa_l, \kappa'_l}^{\alpha_l, \alpha'_l} \rightarrow C_{\kappa'_l}^{\alpha'_l}$ is well-defined and continuous, while the definition of $T^{(R)}$ ensures that $I_n^{(R)}$ is uniformly bounded on $C_{\kappa_l, \kappa'_l}^{\alpha_l, \alpha'_l}$

which implies that $I_n^{(R)} \in C_b(C_{\kappa_l, \kappa'_l}^{\alpha_l, \alpha'_l}; c_{\kappa'_l}^{\alpha'_l})$. Moreover, we use Assumption 3.1(ii), Theorem A.2 with $c \geq 0$ denoting the operator norm of the embedding $C_{\kappa_l}^{\alpha_l} \hookrightarrow C_{\kappa'_l}^{\alpha'_l}$ and $p > 1$ to obtain

$$\begin{aligned} \lim_{R \rightarrow \infty} \|I_n - I_n^{(R)}\|_{\mathcal{B}_p(C_{\kappa_l, \kappa'_l}^{\alpha_l, \alpha'_l}; c_{\kappa'_l}^{\alpha'_l})} &= \lim_{R \rightarrow \infty} \sup_{f \in C_{\kappa_l}^{\alpha_l}} \frac{\|I_n f - I_n^{(R)} f\|_{\alpha'_l, \kappa'_l}}{(1 + \|f\|_{\alpha_l, \kappa_l})^p} \\ &\leq \lim_{R \rightarrow \infty} \sup_{\|I_n f\|_{\alpha'_l, \kappa'_l} \geq R} \frac{\|I_n f\|_{\alpha'_l, \kappa'_l} + \|I_n^{(R)} f\|_{\alpha'_l, \kappa'_l}}{(1 + \|f\|_{\alpha_l, \kappa_l})^p} \\ &\leq \lim_{R \rightarrow \infty} \sup_{e^{\omega h_n} \|f\|_{\alpha'_l, \kappa'_l} \geq R} \frac{e^{\omega h_n} \|f\|_{\alpha'_l, \kappa'_l} + R}{(1 + \|f\|_{\alpha_l, \kappa_l})^p} \\ &\leq 2ce^{\omega h_n} \lim_{R \rightarrow \infty} \sup_{ce^{\omega h_n} \|f\|_{\alpha_l, \kappa_l} \geq R} \frac{\|f\|_{\alpha_l, \kappa_l}}{(1 + \|f\|_{\alpha_l, \kappa_l})^p} = 0. \end{aligned}$$

This shows that $I_n \in \mathcal{B}_p(C_{\kappa_l, \kappa'_l}^{\alpha_l, \alpha'_l}; c_{\kappa'_l}^{\alpha'_l})$. Following the arguments in the proof of [19, Theorem 4.13], one can further show that $y\rho(h(\cdot)) \in \mathcal{B}_p(C_{\kappa_l, \kappa'_l}^{\alpha_l, \alpha'_l}; c_{\kappa'_l}^{\alpha'_l})$ to obtain $\mathcal{CN}_{C_{\kappa_0}, C_{\kappa}^{\alpha}}^{\mathcal{H}, \rho, \mathcal{L}} \subset \mathcal{B}_p(C_{\kappa_l, \kappa'_l}^{\alpha_l, \alpha'_l}; c_{\kappa'_l}^{\alpha'_l})$.

Second, we define $\mathcal{A} := \text{span}(\{\cos(h(\cdot)): h \in \mathcal{H}\} \cup \{\sin(h(\cdot)): h \in \mathcal{H}\})$ and show that the vector space $\mathcal{W} := \text{span}(\{ya(\cdot): a \in \mathcal{A}, y \in \mathcal{L}\})$ is contained in the closure of $\mathcal{CN}_{C_{\kappa_0}, C_{\kappa}^{\alpha}}^{\mathcal{H}, \rho, \mathcal{L}}$ w.r.t.

$$\max_{l=1, \dots, k} \|T\|_{\mathcal{B}_p(C_{\kappa_l, \kappa'_l}^{\alpha_l, \alpha'_l}; c_{\kappa'_l}^{\alpha'_l})} := \max_{l=1, \dots, k} \sup_{f \in C_{\kappa_l}^{\alpha_l}} \frac{\|Tf\|_{\alpha'_l, \kappa'_l}}{(1 + \|f\|_{\alpha_l, \kappa_l})^p}. \quad (3.1)$$

For $y \in \mathcal{L}$, $h \in \mathcal{H}$, and $\varepsilon > 0$, we define $c_h := \max_{l=1, \dots, k} c_{\alpha_l, \kappa_l, h}$ and $c_y := \max_{l=1, \dots, k} \|y\|_{\alpha'_l, \kappa'_l}$. Then, by [19, Proposition 4.4(A3)] or [57, Theorem 2.7], there exists $\varphi_1 \in \mathcal{NN}_{\mathbb{R}, \mathbb{R}}^{\rho}$ with

$$\sup_{t \in \mathbb{R}} \frac{|\cos(t) - \varphi_1(t)|}{(1 + |t|)^p} < \frac{\varepsilon}{\max(1, c_h^p c_y)}.$$

Hence, the Chernoff-neural operator $\varphi := y\varphi_1(h(\cdot)) \in \mathcal{CN}_{C_{\kappa_0}, C_{\kappa}^{\alpha}}^{\mathcal{H}, \rho, \mathcal{L}}$ satisfies

$$\begin{aligned} \max_{l=1, \dots, k} \|y \cos(h(\cdot)) - \varphi(\cdot)\|_{\mathcal{B}_p(C_{\kappa_l, \kappa'_l}^{\alpha_l, \alpha'_l}; c_{\kappa'_l}^{\alpha'_l})} &= \max_{l=1, \dots, k} \sup_{f \in C_{\kappa_l}^{\alpha_l}} \frac{\|y \cos(h(f)) - y\varphi_1(h(f))\|_{\alpha'_l, \kappa'_l}}{(1 + \|f\|_{\alpha_l, \kappa_l})^p} \\ &\leq \left(\max_{l=1, \dots, k} \|y\|_{\alpha'_l, \kappa'_l} \right) \left(\max_{l=1, \dots, k} \sup_{f \in C_{\kappa_l}^{\alpha_l}} \frac{(1 + |h(f)|)^p}{(1 + \|f\|_{\alpha_l, \kappa_l})^p} \right) \sup_{t \in \mathbb{R}} \frac{|\cos(t) - \varphi_1(t)|}{(1 + |t|)^p} < \frac{c_y c_h^p \varepsilon}{\max(1, c_h^p c_y)} \leq \varepsilon. \end{aligned}$$

Since the same estimate holds true for $y \sin(h(\cdot))$, the set $\mathcal{W}$ is contained in the closure of $\mathcal{CN}_{C_{\kappa_0}, C_{\kappa}^{\alpha}}^{\mathcal{H}, \rho, \mathcal{L}}$ w.r.t. the norm defined in equation (3.1).

Third, due to some fundamental trigonometric identities, see [19, Equation 4.3], the set $\mathcal{A}$ is an algebra. By definition, the set $\mathcal{W}$ is an $\mathcal{A}$ -submodule. Since $\mathcal{H}$ is point separating and contains all constant mappings, the set $\mathcal{A}$ is point separating and nowhere vanishing. Finally, by Assumption 3.3(iii), the set $\mathcal{W}(f) := \{w(f): w \in \mathcal{W}\} = \mathcal{L}$ is dense in $c_{\kappa'}^{\alpha'}$ for all $0 \leq \alpha' < \alpha'' \leq \alpha$, $\kappa' \preccurlyeq \kappa'' \in \mathcal{K}$ and $f \in C_{\kappa''}^{\alpha''}$. Hence, the claim follows from Theorem B.3. $\square$

3.2. Qualitative approximation of the semigroup. We now present the main result of this article which shows that iterations of Chernoff-neural operators $\Phi_n \in \mathcal{CN}_{C_{\kappa_0}, C_{\kappa}^{\alpha}}^{\mathcal{H}, \rho, \mathcal{L}}$ are able to approximate the strongly continuous convex monotone semigroup $(S(t))_{t \geq 0}$.

Theorem 3.6. *Suppose that the Assumptions 3.1 and 3.3 are satisfied. Then, for every $\alpha' \in [0, \alpha)$, $\mathcal{K} \ni \kappa' \lesssim \kappa$, $T \geq 0$, $n \in \mathbb{N}$ and $\varepsilon > 0$, there exists $\Phi_n \in \mathcal{CN}_{C_{\kappa_0}, C_{\kappa}}^{\mathcal{H}, \rho, \mathcal{L}}$ with*

$$\|I(\pi_n^t)f - \Phi(\pi_n^t)f\|_{\alpha', \kappa'} \leq (1 + \|f\|_{\alpha, \kappa})^{p^{k_n^t}} \varepsilon \quad \text{for all } t \in [0, T] \text{ and } f \in C_{\kappa}^{\alpha}.$$

Moreover, let Assumption 2.2 be valid and let $(S(t))_{t \geq 0}$ be the semigroup from Theorem 2.3. Then, for every $r, T \geq 0$, $\alpha' \in [0, \alpha)$, $\mathcal{K} \ni \kappa' \lesssim \kappa$ and $\varepsilon > 0$, there exist $n \in \mathbb{N}$ and $\Phi_n \in \mathcal{CN}_{C_{\kappa_0}, C_{\kappa}}^{\mathcal{H}, \rho, \mathcal{L}}$ with

$$\sup_{f \in B_{C_{\kappa}^{\alpha}}(r)} \sup_{t \in [0, T]} \|S(t)f - \Phi(\pi_n^t)f\|_{\alpha', \kappa'} < \varepsilon.$$

Proof. Let $\alpha' \in [0, \alpha)$, $\kappa' \lesssim \kappa \in \mathcal{K}$, $p \in (1, \infty)$, $T \geq 0$, $n \in \mathbb{N}$ and $\varepsilon > 0$. By Assumption 3.1(i), there exist $\alpha' = \alpha_{k_n^T} < \dots < \alpha_0 = \alpha$ and $\kappa' = \kappa_{k_n^T} \lesssim \dots \lesssim \kappa_0 = \kappa$. Since Assumption 3.1(ii) guarantees that $I_n^k f \in C_{\kappa}^{\alpha}$ for all $k \in \mathbb{N}$ and $f \in C_{\kappa}^{\alpha}$, we can further apply Assumption 3.1(ii)–(iii) and Theorem A.2 to obtain a constant $c \geq 1$ with

$$\|I_n f\|_{\alpha_l, \kappa_l} \leq c \|f\|_{\alpha_{l-1}, \kappa_{l-1}}, \quad (3.2)$$

$$\|I_n^k f - I_n^k g\|_{\alpha_l, \kappa_l} \leq c \|f - g\|_{\alpha_l, \kappa_l} \quad (3.3)$$

for all $f, g \in C_{\kappa}^{\alpha}$ and $k, l \in \{1, \dots, k_n^T\}$. By Theorem 3.5, there exists $\Phi_n \in \mathcal{CN}_{C_{\kappa_0}, C_{\kappa}}^{\mathcal{H}, \rho, \mathcal{L}}$ with

$$\max_{1 \leq l \leq k_n^T} \sup_{f \in C_{\kappa_{l-1}}^{\alpha_{l-1}}} \frac{\|I_n f - \Phi_n f\|_{\alpha_l, \kappa_l}}{(1 + \|f\|_{\alpha_{l-1}, \kappa_{l-1}})^p} < \min \left\{ 1, \frac{\varepsilon}{C} \right\}, \quad (3.4)$$

where $C := c \sum_{l=1}^{k_n^T} (2c)^{\frac{p(l-1)}{p-1}}$. We use $\Phi_n^{l-1} f \in C_{\kappa}^{\alpha}$, inequality (3.2) and inequality (3.4) to obtain

$$\begin{aligned} 1 + \|\Phi_n^l f\|_{\alpha_l, \kappa_l} &= 1 + \|\Phi_n \Phi_n^{l-1} f\|_{\alpha_l, \kappa_l} \\ &\leq 1 + \|I_n \Phi_n^{l-1} f\|_{\alpha_l, \kappa_l} + \|I_n \Phi_n^{l-1} f - \Phi_n \Phi_n^{l-1} f\|_{\alpha_l, \kappa_l} \\ &\leq 1 + c \|\Phi_n^{l-1} f\|_{\alpha_{l-1}, \kappa_{l-1}} + (1 + \|\Phi_n^{l-1} f\|_{\alpha_{l-1}, \kappa_{l-1}})^p \\ &\leq 2c(1 + \|\Phi_n^{l-1} f\|_{\alpha_{l-1}, \kappa_{l-1}})^p \end{aligned}$$

for all $l \in \{1, \dots, k_n^T\}$ and $f \in C_{\kappa}^{\alpha}$. Hence, by induction on $l \in \{1, \dots, k_n^T\}$, it follows that

$$1 + \|\Phi_n^l f\|_{\alpha_l, \kappa_l} \leq (2c)^{\frac{p(l-1)}{p-1}} (1 + \|f\|_{\alpha, \kappa})^{p^l} \quad \text{for all } f \in C_{\kappa}^{\alpha}. \quad (3.5)$$

For every $t \in [0, T]$ and $f \in C_{\kappa}^{\alpha}$, we use $\Phi_n^{l-1} f \in C_{\kappa}^{\alpha}$ and the inequalities (3.3)–(3.5) to obtain

$$\begin{aligned} \|I(\pi_n^t)f - \Phi(\pi_n^t)f\|_{\alpha', \kappa'} &= \|I_n^{k_n^t} f - \Phi_n^{k_n^t} f\|_{\alpha', \kappa'} \leq \sum_{l=1}^{k_n^t} \|I_n^{k_n^t-l} I_n \Phi_n^{l-1} f - I_n^{k_n^t-l} \Phi_n \Phi_n^{l-1} f\|_{\alpha', \kappa'} \\ &\leq c \sum_{l=1}^{k_n^t} \|I_n \Phi_n^{l-1} f - \Phi_n \Phi_n^{l-1} f\|_{\alpha_l, \kappa_l} \leq \frac{c\varepsilon}{C} \sum_{l=1}^{k_n^t} (1 + \|\Phi_n^{l-1} f\|_{\alpha_{l-1}, \kappa_{l-1}})^p \\ &\leq \frac{c\varepsilon}{C} \sum_{l=1}^{k_n^t} (2c)^{\frac{p(l-1)}{p-1}} (1 + \|f\|_{\alpha, \kappa})^{p^l} \leq (1 + \|f\|_{\alpha, \kappa})^{p^{k_n^t}} \varepsilon. \end{aligned}$$

The second part follows from the first part and Corollary 2.4. $\square$

4. ENVELOPE-NEURAL OPERATORS

We now introduce a more specific neural operator architecture for the case that the one-step operators are given as the supremum over a family of linear operators. This covers, for instance, the important class of Nisio semigroups which have originally been introduced in [59] and adapted to transition semigroups of stochastic processes under model uncertainty in [21, 54]. In the sequel, we fix $\alpha \in (0, 1]$, a bounded continuous function $\kappa: \mathbb{R}^d \rightarrow (0, \infty)$ satisfying inequality (2.1), an index set Λ , a family $\{I_{n,\lambda}: n \in \mathbb{N}, \lambda \in \Lambda\}$ of operators $I_{n,\lambda}: C_\kappa \rightarrow C_\kappa$, penalization functions $\eta, \eta_n: \Lambda \rightarrow [0, \infty]$ and a sequence $(h_n)_{n \in \mathbb{N}} \subset (0, \infty)$ with $h_n \rightarrow 0$. We define

$$(I_n f)(x) := \sup_{\lambda \in \Lambda} ((I_{n,\lambda} f)(x) - \eta_n(\lambda) h_n) \quad (4.1)$$

for all $n \in \mathbb{N}$, $f \in C_\kappa$ and $x \in \mathbb{R}^d$. Moreover, for every $\lambda \in \Lambda$, we define

$$A_\lambda: D(A_\lambda) \rightarrow C_\kappa, \quad f \mapsto \lim_{n \rightarrow \infty} \frac{I_{n,\lambda} f - f}{h_n},$$

where the domain $D(A_\lambda)$ consists of all $f \in C_\kappa$ such that the previous limit exists. To guarantee that $I_n: C_\kappa \rightarrow C_\kappa$ and that Assumption 2.2 is satisfied, we impose the following conditions.

Assumption 4.1. Suppose that the conditions (i)–(vii) or that the conditions (i)–(vi) and (vii') of the following list are satisfied:

- (i) For every $n \in \mathbb{N}$, there exists $\lambda_{0,n} \in \Lambda$ with $\eta_n(\lambda_{0,n}) = 0$.
- (ii) The operators $I_{n,\lambda}$ are linear and monotone for all $n \in \mathbb{N}$ and $\lambda \in \Lambda$.
- (iii) There exists $\omega \geq 0$ with $\|I_{n,\lambda} f\|_\kappa \leq e^{\omega h_n} \|f\|_\kappa$ for all $n \in \mathbb{N}$, $\lambda \in \Lambda$ and $f \in C_\kappa$.
- (iv) For every $\varepsilon > 0$, there exists $\delta > 0$ and $n_0 \in \mathbb{N}$ with

$$I_{n,\lambda}(\tau_x f) \leq \tau_x(I_{n,\lambda} f) + \frac{r \varepsilon h_n}{\kappa}$$

for all $n \geq n_0$, $r \geq 0$, $f \in \text{Lip}_b(r)$, $x \in B_{\mathbb{R}^d}(\delta)$ and $\lambda \in \Lambda$.

- (v) For every $\varepsilon > 0$, $r, T \geq 0$ and $K \Subset \mathbb{R}^d$, there exist $c \geq 0$ and $K' \Subset \mathbb{R}^d$ with

$$\|I(\pi_n^t) f - I(\pi_n^t) g\|_{\infty, K} \leq c \|f - g\|_{\infty, K'} + \varepsilon$$

for all $t \in [0, T]$, $n \in \mathbb{N}$ and $f, g \in B_{C_\kappa}(r)$.

- (vi) It holds $I_{n,\lambda}: \text{Lip}_b(r) \rightarrow \text{Lip}_b(e^{\omega h_n} r)$ for all $n \in \mathbb{N}$, $\lambda \in \Lambda$ and $r \geq 0$.
- (vii) The limit $I'(0)f \in C_\kappa$ exists for all $f \in C_b^\infty$.
- (vii') It holds $C_b^\infty \subset D(A_\lambda)$ for all $\lambda \in \Lambda$. In addition, for every $f \in C_b^\infty$, there exist $c \geq 0$, $n_0 \in \mathbb{N}$ and $\Lambda_0 \subset \Lambda$ such that the following statements are true:
 - (a) $\sup_{\lambda \in \Lambda} (I_{n,\lambda} f - \eta_n(\lambda) h_n) = \sup_{\lambda \in \Lambda_0} (I_{n,\lambda} f - \eta_n(\lambda) h_n)$ for all $n \geq n_0$,
 - (b) $\sup_{\lambda \in \Lambda} (A_\lambda f - \eta(\lambda)) = \sup_{\lambda \in \Lambda_0} (A_\lambda f - \eta(\lambda))$,
 - (c) $\sup_{n \in \mathbb{N}} \sup_{\lambda \in \Lambda_0} \eta_n(\lambda) < \infty$, $\sup_{\lambda \in \Lambda_0} \|I_{n,\lambda} f - f\|_\kappa \leq c h_n$ for all $n \geq n_0$ and

$$\lim_{n \rightarrow \infty} \sup_{\lambda \in \Lambda_0} \left\| \frac{I_{n,\lambda} f - f}{h_n} - A_\lambda f \right\|_{\infty, K} = 0 \quad \text{for all } K \Subset \mathbb{R}^d.$$

Furthermore, it holds $\lim_{n \rightarrow \infty} \sup_{\lambda \in \Lambda_0} |\eta_n(\lambda) - \eta(\lambda)| = 0$.

Note that condition (vii') immediately implies condition (vii).

Theorem 4.2. Suppose that the Assumptions 4.1(i)–(vii) are valid. Then, there exists a strongly continuous convex monotone semigroup $(S(t))_{t \geq 0}$ on C_κ given by

$$S(t)f := \lim_{n \rightarrow \infty} I(\pi_n^t) f \quad \text{for all } t \geq 0 \text{ and } f \in C_\kappa$$

whose generator satisfies $Af = I'(0)f$ for all $f \in C_b^\infty$. If the Assumption 4.1(i)–(vi) and (vii') are valid, we further obtain

$$Af = \sup_{\lambda \in \Lambda} (A_\lambda f - \eta(\lambda)) \quad \text{for all } f \in C_b^\infty.$$

Proof. This is a particular case of Theorem 2.3. $\square$

Corollary 4.3. Let Assumption 4.1 be satisfied and denote by $(S(t))_{t \geq 0}$ the strongly continuous convex monotone semigroup from Theorem 4.2. Assume that there exists $\tilde{\omega} \geq 0$ with

$$\|I_{n,\lambda}f\|_{\alpha,\kappa} \leq e^{\tilde{\omega}h_n}\|f\|_{\alpha,\kappa} \quad \text{for all } n \in \mathbb{N}, \lambda \in \Lambda \text{ and } f \in C_\kappa^\alpha.$$

Then, it holds $\|S(t)f\|_{\alpha,\kappa} \leq e^{\tilde{\omega}t}\|f\|_{\alpha,\kappa}$ for all $t \geq 0$ and $f \in C_\kappa^\alpha$. Furthermore,

$$\lim_{n \rightarrow \infty} \sup_{f \in B_{C_\kappa^\alpha}(r)} \sup_{t \in [0,T]} \|S(t)f - I(\pi_n^t)f\|_{\alpha',\kappa'} = 0$$

for all $r, T \geq 0$, $\alpha' \in [0, \alpha)$ and $\kappa' \ll \kappa$.

Proof. This is a particular case of Corollary 2.4. $\square$

4.1. Universal approximation. In order to approximate the operators $(I_n)_{n \in \mathbb{N}}$ defined by equation (4.1), we replace the supremum over $\lambda \in \Lambda$ by the maximum over finitely many trainable parameters $\lambda_0, \dots, \lambda_M \in \Lambda$ which can be implemented using a ReLU-neural network, see Figure 2.

Definition 4.4. For every $n \in \mathbb{N}$, an envelope-neural operator is a mapping of the form

$$\Phi_n: C_\kappa \rightarrow C_\kappa, f \mapsto \varphi((I_{n,\lambda_0}f)(\cdot), (I_{n,\lambda_1}f)(\cdot), \dots, (I_{n,\lambda_M}f)(\cdot)),$$

where $\lambda_0 := \lambda_{0,n} \in \Lambda$ is the fixed parameter from Assumption 4.1(i), $\lambda_1, \dots, \lambda_M \in \Lambda$ are the (hidden-layer) parameters and $\varphi \in \mathcal{NN}_{\mathbb{R}^{M+1}, \mathbb{R}}^{\text{ReLU}}$ is a ReLU-neural network. We collect all operators of this form in the set $\mathcal{EN}_{C_\kappa, C_\kappa}^{\text{ReLU}}$.




FIGURE 2. An envelope-neural operator $\Phi_n: C_\kappa \rightarrow C_\kappa$ with hidden-layer maps from the family $(I_{n,\lambda})_{\lambda \in \Lambda}$, ReLU-neural network $\varphi \in \mathcal{NN}_{\mathbb{R}^3, \mathbb{R}}^{\text{ReLU}}$ as linear readout and $M + 1 = 3$ neurons.

Assumption 4.5. Suppose that the following conditions are satisfied:

- (i) It holds $\|I_{n,\lambda}f\|_{\alpha,\kappa} \leq e^{\omega h_n}\|f\|_{\alpha,\kappa}$ for all $n \in \mathbb{N}$, $\lambda \in \Lambda$ and $f \in C_\kappa^\alpha$.
- (ii) For every $r \geq 0$, there exists a totally bounded metric space (Λ_r, d_r) with $\Lambda_r \subset \Lambda$ such that the set $\Lambda_{r,n} := \Lambda_r \cap \{\eta_n < \infty\}$ satisfies

$$\sup_{\lambda \in \Lambda} (I_{n,\lambda}f - \eta_n(\lambda)h_n) = \sup_{\lambda \in \Lambda_{r,n}} (I_{n,\lambda}f - \eta_n(\lambda)h_n) \quad \text{for all } n \in \mathbb{N} \text{ and } f \in B_{C_\kappa^\alpha}(r).$$

In addition, for every $r \geq 0$ and $\varepsilon > 0$, there exists $\delta > 0$ such that

$$\|(I_{n,\lambda_1}f - \eta_n(\lambda_1)h_n) - (I_{n,\lambda_2}f - \eta_n(\lambda_2)h_n)\|_\kappa < \varepsilon$$

for all $n \in \mathbb{N}$, $f \in B_{C_\kappa^\alpha}(r)$ and $\lambda_1, \lambda_2 \in \Lambda_{r,n}$ with $d_r(\lambda_1, \lambda_2) < \delta$. Finally, the element $\lambda_{0,n}$ from Assumption 4.1(i) satisfies $\lambda_{0,n} \in \Lambda_r$ for all $n \in \mathbb{N}$.

Theorem 4.6. *Suppose that the Assumptions 4.1 and 4.5 are satisfied. Then, for every $n \in \mathbb{N}$, $r \geq 0$, $\alpha' \in [0, \alpha)$ and $\varepsilon > 0$, there exists $\Phi_n \in \mathcal{EN}_{C_\kappa, C_\kappa}^{\text{ReLU}}$ with*

$$\sup_{f \in B_{C_\kappa^\alpha}(r)} \|I_n f - \Phi_n f\|_{\alpha', \kappa} < \varepsilon.$$

Proof. Let $\varepsilon > 0$ and $\varepsilon' \in (0, \varepsilon)$ with $(2e^{\omega h_n r})^{\alpha'/\alpha} (2\varepsilon')^{1-\alpha'/\alpha} < \varepsilon/2$. Moreover, let $\Lambda_{r,n}$ and $\delta > 0$ satisfy Assumption 4.5(ii) with ε' . Since $\Lambda_{r,n}$ is also totally bounded, there exist $\lambda_1, \dots, \lambda_M \in \Lambda_{r,n}$ with $\Lambda_{r,n} \subset \bigcup_{m=1}^M \bar{B}_{d_r}(\lambda_m, \delta)$. In particular, choosing $m(\lambda) \in \arg \min_{m=1, \dots, M} d_r(\lambda, \lambda_m)$ yields

$$\|(I_{n,\lambda} f - \eta_n(\lambda) h_n) - (I_{n,\lambda_{m(\lambda)}} f - \eta_n(\lambda_{m(\lambda)}) h_n)\|_\kappa < \varepsilon' < \varepsilon \quad \text{for all } f \in B_{C_\kappa^\alpha}(r).$$

We implement the function $\mathbb{R}^{M+1} \rightarrow \mathbb{R}$, $y \mapsto \max_{m=0, \dots, M} (y_m - \eta_n(\lambda_m) h_n)$ with $\lambda_0 := \lambda_{0,n}$ as ReLU-neural network $\varphi \in \mathcal{NN}_{\mathbb{R}^{M+1}, \mathbb{R}}^{\text{ReLU}}$ and define $\Phi_n \in \mathcal{EN}_{C_\kappa, C_\kappa}^{\text{ReLU}}$ by $\Phi_n(f) := \varphi \circ I_{n,\lambda_{0:M}} f$, where $I_{n,\lambda_{0:M}} f := ((I_{n,\lambda_0} f)(\cdot), \dots, (I_{n,\lambda_M} f)(\cdot))$. Then, for every $f \in B_{C_\kappa^\alpha}(r)$, it holds

$$\|I_n f - \Phi_n f\|_\kappa = \left\| \sup_{\lambda \in \Lambda_{r,n}} (I_{n,\lambda} f - \eta_n(\lambda) h_n) - \max_{m=0, \dots, M} (I_{n,\lambda_m} f - \eta_n(\lambda_m) h_n) \right\|_\kappa < \varepsilon' < \varepsilon. \quad (4.2)$$

Let $f \in B_{C_\kappa^\alpha}(r)$ and define $g_n := I_n f - \varphi \circ I_{n,\lambda_{0:M}} f$. Assumption 4.5(i) implies

$$\sup_{x \neq y} \frac{|g_n(x) - g_n(y)|}{|x - y|^\alpha} \bar{\kappa}(x, y) \leq \sup_{\lambda \in \Lambda} \|I_{n,\lambda} f\|_{\alpha, \kappa} + \max_{m=0, \dots, M} \|I_{n,\lambda_m} f\|_{\alpha, \kappa} \leq 2e^{\omega h_n} \|f\|_{\alpha, \kappa} \leq 2e^{\omega h_n} r.$$

Hence, it follows from inequality (4.2) and $\bar{\kappa}(x, y) \leq 2 \min(\kappa(x), \kappa(y))$ that

$$\begin{aligned} \sup_{x \neq y} \frac{|g_n(x) - g_n(y)|}{|x - y|^\alpha} \bar{\kappa}(x, y) &\leq \sup_{x \neq y} \left(\frac{|g_n(x) - g_n(y)|}{|x - y|^\alpha} \bar{\kappa}(x, y) \right)^{\frac{\alpha'}{\alpha}} (|g_n(x) - g_n(y)| \bar{\kappa}(x, y))^{1-\frac{\alpha'}{\alpha}} \\ &\leq (2e^{\omega h_n} r)^{\frac{\alpha'}{\alpha}} (2\varepsilon')^{1-\frac{\alpha'}{\alpha}} < \varepsilon. \end{aligned} \quad \square$$

The following theorem states that iterations of an envelope-neural operator Φ_n are able to approximate the strongly continuous convex monotone semigroup $(S(t))_{t \geq 0}$.

Theorem 4.7. *Suppose that the Assumptions 4.1 and 4.5 are satisfied. Then, for every $r, T \geq 0$, $n \in \mathbb{N}$, $\alpha' \in [0, \alpha)$ and $\varepsilon > 0$, there exists $\Phi_n \in \mathcal{EN}_{C_\kappa, C_\kappa}^{\text{ReLU}}$ with*

$$\sup_{f \in B_{C_\kappa^\alpha}(r)} \sup_{t \in [0, T]} \|I(\pi_n^t) f - \Phi(\pi_n^t) f\|_{\alpha', \kappa} < \varepsilon.$$

Denote by $(S(t))_{t \geq 0}$ the semigroup from Theorem 4.2. Then, for every $r, T \geq 0$, $\alpha' \in [0, \alpha)$, $\kappa' \lesssim \kappa$ and $\varepsilon > 0$, there exist $n \in \mathbb{N}$ and $\Phi_n \in \mathcal{EN}_{C_\kappa, C_\kappa}^{\text{ReLU}}$ with

$$\sup_{f \in B_{C_\kappa^\alpha}(r)} \sup_{t \in [0, T]} \|S(t) f - \Phi(\pi_n^t) f\|_{\alpha', \kappa'} < \varepsilon.$$

Proof. Let $r, T \geq 0$, $n \in \mathbb{N}$, $\alpha' \in [0, \alpha)$, $\varepsilon > 0$ and $\varepsilon' \in (0, \varepsilon)$ with $(2e^{\omega T r})^{\alpha'/\alpha} (2\varepsilon')^{1-\alpha'/\alpha} < \varepsilon$. Assumption 4.1(iii) guarantees that

$$\|I_{n,\lambda} f\|_\kappa \leq e^{\omega h_n} \|f\|_\kappa \quad \text{and} \quad \|I_n^k f - I_n^k g\|_\kappa \leq e^{\omega k h_n} \|f - g\|_\kappa$$

for all $f, g \in C_\kappa^\alpha$, $k \in \{1, \dots, k_n^T\}$ and $\lambda \in \Lambda$. By Theorem 4.6, there exists $\Phi_n \in \mathcal{EN}_{C_\kappa, C_\kappa}^{\text{ReLU}}$ with

$$\|I_n f - \Phi_n f\|_\kappa < \frac{\varepsilon'}{e^{\omega T k_n^T}} \quad \text{for all } f \in B_{C_\kappa^\alpha}(e^{\omega T r}),$$

where $\Phi_n f := \max_{m=0,\dots,M} (I_{n,\lambda_m} f - \eta_n(\lambda_m) h_n)$ with $\lambda_0 := \lambda_{0,n}$ and some $\lambda_1, \dots, \lambda_M \in \Lambda$. For every $t \in [0, T]$ and $f \in B_{C_\kappa^\alpha}(r)$, we use $\|\Phi_n^{l-1} f\|_\kappa \leq \|\Phi_n^{l-1} f\|_{\alpha,\kappa} \leq e^{\omega(l-1)h_n} r \leq e^{\omega T} r$ to obtain

$$\begin{aligned} \|I(\pi_n^t) f - \Phi(\pi_n^t) f\|_\kappa &= \|I_n^{k_n^t} f - \Phi_n^{k_n^t} f\|_\kappa \leq \sum_{l=1}^{k_n^t} \|I_n^{k_n^t-l} I_n \Phi_n^{l-1} f - I_n^{k_n^t-l} \Phi_n \Phi_n^{l-1} f\|_\kappa \\ &\leq e^{\omega T} \sum_{l=1}^{k_n^t} \|I_n \Phi_n^{l-1} f - \Phi_n \Phi_n^{l-1} f\|_\kappa < \frac{e^{\omega T} k_n^t \varepsilon'}{e^{\omega T} k_n^T} \leq \varepsilon'. \end{aligned} \quad (4.3)$$

Let $f \in B_{C_\kappa^\alpha}(r)$ and define $g_n^t := I(\pi_n^t) f - \Phi(\pi_n^t) f$. It holds

$$\sup_{x \neq y} \frac{|g_n^t(x) - g_n^t(y)|}{|x - y|^\alpha} \bar{\kappa}(x, y) \leq \|I(\pi_n^t) f\|_{\alpha,\kappa} + \|\Phi(\pi_n^t) f\|_{\alpha,\kappa} \leq 2e^{\omega k_n^t h_n} \|f\|_{\alpha,\kappa} \leq 2e^{\omega T} r.$$

Hence, it follows from inequality (4.3) and $\bar{\kappa}(x, y) \leq 2 \min(\kappa(x), \kappa(y))$ that

$$\begin{aligned} \sup_{x \neq y} \frac{|g_n^t(x) - g_n^t(y)|}{|x - y|^\alpha} \bar{\kappa}(x, y) &\leq \sup_{x \neq y} \left(\frac{|g_n^t(x) - g_n^t(y)|}{|x - y|^\alpha} \bar{\kappa}(x, y) \right)^{\frac{\alpha'}{\alpha}} (|g_n^t(x) - g_n^t(y)| \bar{\kappa}(x, y))^{1 - \frac{\alpha'}{\alpha}} \\ &\leq (2e^{\omega T} r)^{\frac{\alpha'}{\alpha}} (2\varepsilon')^{1 - \frac{\alpha'}{\alpha}} < \varepsilon. \end{aligned}$$

This shows the first part of the claim and the second part follows from Corollary 4.3. $\square$

It remains to quantify the approximation error in terms of the number of neurons $M \in \mathbb{N}$.

Assumption 4.8. For every $r \geq 0$, there exists a totally bounded metric space (Λ_r, d_r) with $\Lambda_r \subset \Lambda$ such that the set $\Lambda_{r,n} := \Lambda_r \cap \{\eta_n < \infty\}$ satisfies

$$\sup_{\lambda \in \Lambda} (I_{n,\lambda} f - \eta_n(\lambda) h_n) = \sup_{\lambda \in \Lambda_{r,n}} (I_{n,\lambda} f - \eta_n(\lambda) h_n) \quad \text{for all } n \in \mathbb{N} \text{ and } f \in B_{C_\kappa^\alpha}(r).$$

In addition, for every $r \geq 0$, there exist $L_r, \beta_r \geq 0$ with

$$\|(I_{n,\lambda_1} f - \eta_n(\lambda_1) h_n) - (I_{n,\lambda_2} f - \eta_n(\lambda_2) h_n)\|_\kappa \leq L_r d_r(\lambda_1, \lambda_2)^{\beta_r}$$

for all $n \in \mathbb{N}$, $f \in B_{C_\kappa^\alpha}(r)$ and $\lambda_1, \lambda_2 \in \Lambda_{r,n}$. Finally, the element $\lambda_{0,n}$ from Assumption 4.1(i) satisfies $\lambda_{0,n} \in \Lambda_r$ for all $n \in \mathbb{N}$.

For every $M \in \mathbb{N}$ and $r \geq 0$, the *fill-in distance* of (Λ_r, d_r) is defined by

$$\delta_r(M) := \inf_{\lambda_1, \dots, \lambda_M \in \Lambda_r} \sup_{\lambda \in \Lambda_r} \min_{m=1, \dots, M} d_r(\lambda, \lambda_m).$$

For example, if $\Lambda_r \subset \mathbb{R}^m$ and d_r is equivalent to the Euclidean metric, then $\delta_r(M) \asymp M^{-1/m}$.

Theorem 4.9. Suppose that the Assumptions 4.1, 4.5(i) and 4.8 are satisfied. Then, for every $r \geq 0$, $\alpha' \in [0, \alpha)$ and $M, n \in \mathbb{N}$, there exists $\Phi_{n,M} \in \mathcal{EN}_{C_\kappa, C_\kappa}^{\text{ReLU}}$ with $M+1$ neurons such that

$$\|I_n f - \Phi_{n,M} f\|_{\alpha', \kappa} \leq \max \left\{ L_r (4\delta_r(M))^{\beta_r}, (C_{r,\alpha',\kappa} (4\delta_r(M))^{\beta_r})^{1 - \frac{\alpha'}{\alpha}} \right\} \quad \text{for all } f \in B_{C_\kappa^\alpha}(r),$$

where $C_{r,\alpha',\kappa} > 0$ is a constant depending only on r , α' and κ . Furthermore, the output ReLU-neural network of $\Phi_{n,M}$ has depth $\mathcal{O}(\log_2(M))$ and width $\mathcal{O}(M)$.

Proof. Since (Λ_r, d_r) is totally bounded, by definition of the infimum in $\delta_r(M)$, there exist some $\tilde{\lambda}_1, \dots, \tilde{\lambda}_M \in \Lambda_r$ with $\sup_{\lambda \in \Lambda_r} \min_{m=1, \dots, M} d_r(\lambda, \tilde{\lambda}_m) \leq 2\delta_r(M)$. For every $m \in \{1, \dots, M\}$ with $\Lambda_{r,n} \cap B_{d_r}(\tilde{\lambda}_m, 2\delta_r(M)) \neq \emptyset$, choose some $\lambda_m \in \Lambda_{r,n} \cap B_{d_r}(\tilde{\lambda}_m, 2\delta_r(M))$. By repeating selected points if necessary, we obtain $\lambda_1, \dots, \lambda_M \in \Lambda_{r,n}$, for which the triangle inequality ensures

$$\sup_{\lambda \in \Lambda_{r,n}} \min_{m=1, \dots, M} d_r(\lambda, \lambda_m) \leq 4\delta_r(M).$$

By [61, Lemma 5.11], we can implement $\mathbb{R}^{M+1} \rightarrow \mathbb{R}$, $y \mapsto \max_{m=0,\dots,M} (y_m - \eta_n(\lambda_m)h_n)$ with $\lambda_0 := \lambda_{0,n}$ as ReLU network $\varphi \in \mathcal{NN}_{\mathbb{R}^{M+1}, \mathbb{R}}^{\text{ReLU}}$ of depth $\mathcal{O}(\log_2(M))$ and width $\mathcal{O}(M)$. We define $\Phi_{n,M} \in \mathcal{EN}_{C_\kappa, C_\kappa}^{\text{ReLU}}$ by $\Phi_{n,M}f := \varphi((I_{n,\lambda_0}f)(\cdot), \dots, (I_{n,\lambda_M}f)(\cdot))$ and use Assumption 4.8 to obtain

$$\begin{aligned} \|I_n f - \Phi_{n,M} f\|_\kappa &= \left\| \sup_{\lambda \in \Lambda_{r,n}} (I_{n,\lambda} f - \eta_n(\lambda)h_n) - \max_{m=0,\dots,M} (I_{n,\lambda_m} f - \eta_n(\lambda_m)h_n) \right\|_\kappa \\ &\leq \sup_{\lambda \in \Lambda_{r,n}} \| (I_{n,\lambda} f - \eta_n(\lambda)h_n) - (I_{n,\lambda_{m(\lambda)}} f - \eta_n(\lambda_{m(\lambda)})h_n) \|_\kappa \\ &\leq L_r \sup_{\lambda \in \Lambda_{r,n}} d_r(\lambda, \lambda_{m(\lambda)})^{\beta_r} \leq L_r (4\delta_r(M))^{\beta_r} \end{aligned}$$

for all $f \in B_{C_\kappa}(r)$, where $m(\lambda) \in \arg \min_{m=1,\dots,M} d_r(\lambda, \lambda_m)$. Estimating the Hölder seminorm similarly to the proof of Theorem 4.6 yields the claim. $\square$

Finally, by scaling the number of neurons $(M_n)_{n \in \mathbb{N}}$ depending on the step-size $(h_n)_{n \in \mathbb{N}}$, we obtain a sequence $(\Phi_{n,M_n})_{n \in \mathbb{N}}$ of neural operators which generate the same strongly continuous convex monotone semigroup $(S(t))_{t \geq 0}$ as the Chernoff one-step operators $(I_n)_{n \in \mathbb{N}}$. To this end, we recall that the covering number of Λ_r is defined by

$$\mathfrak{N}_r(\varepsilon) := \min \left\{ M \in \mathbb{N} : \exists \lambda_1, \dots, \lambda_M \in \Lambda_r \text{ s.t. } \bigcup_{m=1}^M \mathring{B}_{d_r}(\lambda_m, \varepsilon) = \Lambda_r \right\} \quad \text{for all } \varepsilon > 0.$$

It holds $\mathfrak{N}_r(\varepsilon) = \min\{M \in \mathbb{N} : \delta_r(M) < \varepsilon\}$ and $\delta_r(M) = \inf\{\varepsilon > 0 : \mathfrak{N}_r(\varepsilon) \leq M\}$.

Proposition 4.10. *Suppose that the Assumptions 4.1 and 4.8 are satisfied and denote by $(S(t))_{t \geq 0}$ the semigroup from Theorem 4.2. Let $(r_n)_{n \in \mathbb{N}} \subset (0, \infty)$ be a sequence with $r_n \rightarrow \infty$. Choose $\varepsilon_n > 0$ with $L_{r_n} \varepsilon_n^{\beta_{r_n}} = o(h_n)$ and $M_n \geq \mathfrak{N}_{r_n}(\varepsilon_n/4)$. Then, there exists a sequence $(\Phi_{n,M_n})_{n \in \mathbb{N}}$ of envelope-neural operators $\Phi_{n,M_n} \in \mathcal{EN}_{C_\kappa, C_\kappa}^{\text{ReLU}}$ with $M_n + 1$ neurons such that*

$$S(t)f = \lim_{n \rightarrow \infty} \Phi_{n,M_n}^{k_n^t} f \quad \text{for all } t \geq 0 \text{ and } f \in C_\kappa.$$

Proof. For every $f \in C_b^\infty$ and $K \Subset \mathbb{R}^d$, Assumption 4.1(vii) and Theorem 4.9 imply

$$\begin{aligned} \lim_{n \rightarrow \infty} \left\| \frac{\Phi_{n,M_n} f - f}{h_n} - Af \right\|_{\infty, K} &\leq c \lim_{n \rightarrow \infty} \frac{\|\Phi_{n,M_n} f - I_n f\|_\kappa}{h_n} + \lim_{n \rightarrow \infty} \left\| \frac{I_n f - f}{h_n} - Af \right\|_{\infty, K} \\ &\leq c \lim_{n \rightarrow \infty} \frac{L_{r_n} (4\delta_{r_n}(M_n))^{\beta_{r_n}}}{h_n} \leq c \lim_{n \rightarrow \infty} \frac{L_{r_n} \varepsilon_n^{\beta_{r_n}}}{h_n} = 0, \end{aligned}$$

where $c := \sup_{x \in K} \frac{1}{\kappa(x)}$. Furthermore, Assumption 4.1 guarantees that $(\Phi_{n,M_n})_{n \in \mathbb{N}}$ is a family of one-step operators $\Phi_{n,M_n} : C_\kappa \rightarrow C_\kappa$ satisfying Assumption 2.2. By Theorem 2.3, there exists another strongly continuous convex monotone semigroup $(T(t))_{t \geq 0}$ on C_κ given by

$$T(t)f := \lim_{n \rightarrow \infty} \Phi_{n,M_n}^{k_n^t} f \quad \text{for all } t \geq 0 \text{ and } f \in C_\kappa$$

with generator $B : D(B) \rightarrow C_\kappa$ satisfying (iv) and (v) of Theorem 2.3, $C_b^\infty \subset D(B)$ and

$$Af = Bf \quad \text{for all } f \in C_b^\infty.$$

Consequently, we obtain $S(t)f = T(t)f$ for all $t \geq 0$ and $f \in C_\kappa$. $\square$

4.2. Quantitative approximation of the semigroup. Finally, by combining Theorem 4.9 with the convergence rates in [9], we obtain an explicit error bound for the approximation of the strongly continuous convex monotone semigroup $(S(t))_{t \geq 0}$ by the envelope-neural operators. The next assumption is a particular case of [9, Assumption 2.5 and 2.7]. Let $\zeta: \mathbb{R}_+ \times \mathbb{R}^d \rightarrow \mathbb{R}$ be an infinitely differentiable function satisfying

$$\text{supp}(\zeta) \subset [0, 1] \times B_{\mathbb{R}^d}(1) \quad \text{and} \quad \int_{\mathbb{R}_+ \times \mathbb{R}^d} \zeta(t, x) dt dx = 1.$$

For every locally bounded measurable function $u: \mathbb{R}_+ \times \mathbb{R}^d \rightarrow \mathbb{R}$, $\varepsilon = (\varepsilon_1, \varepsilon_2) \in (0, \infty)^2$, $t \geq 0$ and $x \in \mathbb{R}^d$, we define $\zeta^\varepsilon(t, x) := \varepsilon_1^{-1} \varepsilon_2^{-d} \zeta(\varepsilon_1^{-1} t, \varepsilon_2^{-1} x)$ and

$$u^\varepsilon(t, x) := (u * \zeta^\varepsilon)(t, x) := \int_{\mathbb{R}_+ \times \mathbb{R}^d} u(s + t, x + y) \zeta^\varepsilon(s, y) ds dy.$$

Assumption 4.11. Suppose that the following conditions are satisfied:

- (i) There exist $c \geq 0$ and $\varepsilon_0 \in (0, 1]$ with

$$\|I_n(\tau_x f) - \tau_x I_n f\|_\kappa \leq cr h_n |x|$$

for all $x \in B_{\mathbb{R}^d}(\varepsilon_0)$, $r \geq 0$, $f \in \text{Lip}_b(r)$ and $n \in \mathbb{N}$.

- (ii) There exist a function $a_1: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ and constants $a_2, p \geq 0$ with

$$\|I_n f - f\|_\kappa \leq \left(a_1(d^{-\frac{1}{2}} \|\nabla f\|_\infty) + a_2(d^{-1} \|\nabla^2 f\|_\infty)^p \right) h_n \quad \text{for all } n \in \mathbb{N} \text{ and } f \in C_b^\infty.$$

- (iii) There exist a function $\theta: \mathbb{R}_+^2 \rightarrow \mathbb{R}_+$ and constants $\gamma_1^\pm, \dots, \gamma_N^\pm \geq 0$ with

$$\begin{aligned} \partial_t u^\varepsilon(t) - A u^\varepsilon(t) - \frac{u^\varepsilon(t) - I_n u^\varepsilon(t - h_n)}{h_n} &\leq \frac{\theta(r, t)}{\kappa} \max_{i=1, \dots, N} h_n^{\gamma_i^+} \varepsilon_2^{-\gamma_i^-}, \\ \partial_t u_n^\varepsilon(t) - A u_n^\varepsilon(t) - \frac{u_n^\varepsilon(t) - I_n u_n^\varepsilon(t - h_n)}{h_n} &\geq -\frac{\theta(r, t)}{\kappa} \max_{i=1, \dots, N} h_n^{\gamma_i^+} \varepsilon_2^{-\gamma_i^-} \end{aligned}$$

for all $n \in \mathbb{N}$, $r \geq 0$, $f \in \text{Lip}_b(r)$, $t \geq h_n$ and $\varepsilon = (\varepsilon_1, \varepsilon_2) \in (0, \varepsilon_0]^2$ with $\varepsilon_1 = \varepsilon_2^{1+p} \geq h_n$, where $u(t) := S(t)f$ and $u_n(t) := I(\pi_n^t)f$.

Theorem 4.12. Let the Assumptions 4.1, 4.5(i), 4.8 and 4.11 be satisfied and denote by $(S(t))_{t \geq 0}$ the semigroup from Theorem 4.2. Then, for every $r, T \geq 0$, there exists $c_{r,T} \geq 0$ such that, for every $M_n \geq \mathfrak{N}_{e^{\omega T r}}(\varepsilon_n/4)$ with $\varepsilon_n^{\beta e^{\omega T r}} = \mathcal{O}(h_n^\gamma/k_n^T)$, there exists $\Phi_{n, M_n} \in \mathcal{EN}_{C_\kappa, C_\kappa}^{\text{ReLU}}$ with $M_n + 1$ neurons such that

$$\|S(t)f - \Phi_{n, M_n}^{k_n^t} f\|_\kappa \leq c_{r,T} h_n^\gamma \quad \text{for all } t \in [0, T] \text{ and } f \in \text{Lip}_b(r),$$

where $\gamma := \min \left\{ \frac{1}{1+p}, \frac{\gamma_1^+}{1+\gamma_1^-}, \dots, \frac{\gamma_N^+}{1+\gamma_N^-} \right\}$.

Proof. We use [9, Theorem 2.9], argue similarly as in the proof of Theorem 4.7 and apply Theorem 4.9 to obtain constants $\tilde{c}_{r,T}, c_{r,T} \geq 0$ with

$$\begin{aligned}
\|S(t)f - \Phi_{n,M_n}^{k_n^t} f\|_\kappa &\leq \|S(t)f - I(\pi_n^t)f\|_\kappa + \|I_n^{k_n^t} f - \Phi_{n,M_n}^{k_n^t} f\|_\kappa \\
&\leq \tilde{c}_{r,T} h_n^\gamma + \sum_{l=1}^{k_n^t} \|I_n^{k_n^t-l} I_n \Phi_{n,M_n}^{l-1} f - I_n^{k_n^t-l} \Phi_{n,M_n} \Phi_{n,M_n}^{l-1} f\|_\kappa \\
&\leq \tilde{c}_{r,T} h_n^\gamma + e^{\omega T} \sum_{l=1}^{k_n^t} \|I_n \Phi_{n,M_n}^{l-1} f - \Phi_{n,M_n} \Phi_{n,M_n}^{l-1} f\|_\kappa \\
&\leq \tilde{c}_{r,T} h_n^\gamma + e^{\omega T} k_n^t L_{e^{\omega T} r} (4\delta_{e^{\omega T} r}(M_n))^{\beta_{e^{\omega T} r}} \\
&\leq \tilde{c}_{r,T} h_n^\gamma + e^{\omega T} k_n^T L_{e^{\omega T} r} \varepsilon_n^{\beta_{e^{\omega T} r}} \leq c_{r,T} h_n^\gamma. \quad \square
\end{aligned}$$

The previous theorem might seem very abstract at first, but in many applications verifying the assumptions and deriving explicit convergence rates is rather straightforward, see [9, Section 4]. For instance, in case of the optimal control problem studied in Subsection 5.2 below, it follows from [9, Theorem 4.3] that $\gamma = \frac{1}{4}$. Furthermore, it holds $\beta_r = \alpha$ and $\Lambda \subset \mathbb{R}^d \times \mathbb{S}_+^d \cong \mathbb{R}^{\frac{d(d+3)}{2}}$. Consequently, one should choose

$$M_n + 1 \geq \mathfrak{N}_{e^{\omega T} r}(\varepsilon_n) + 1 \asymp \varepsilon_n^{-\frac{d(d+3)}{2}} \asymp \left(\frac{h_n^\gamma}{k_n^T} \right)^{-\frac{d(d+3)}{2\alpha}} \asymp h_n^{-\frac{d(d+3)(\gamma+1)}{2\alpha}} = h_n^{-\frac{5d(d+3)}{8\alpha}}$$

neurons in the envelope-neural operator.

5. NUMERICAL EXPERIMENTS

In this section, we illustrate with three numerical examples¹ how Chernoff-neural operators and envelope-neural operators can be effectively used to learn the Chernoff one-step operators $(I_n)_{n \in \mathbb{N}}$ and therefore the corresponding strongly continuous convex monotone semigroup $(S(t))_{t \geq 0}$. The examples follow an increasing level of abstraction and are carefully chosen to highlight the key aspects of our approach.

5.1. Splitting schemes for semilinear PDEs. In the first example, we approximate the solution operator $f \mapsto S(t)f := u(t, \cdot)$ of the semilinear partial differential equation (PDE)

$$\begin{cases} \partial_t u(t, x) = \frac{1}{2} \Delta u(t, x) + H(\nabla u(t, x)), & (t, x) \in (0, \infty) \times \mathbb{R}^d, \\ u(0, x) = f(x), & x \in \mathbb{R}^d, \end{cases} \quad (5.1)$$

where Δ denotes the Laplacian, ∇ represents the gradient and $H: \mathbb{R}^d \rightarrow \mathbb{R}$ is a convex Hamiltonian whose convex conjugate is given by

$$H^*(y) := \sup_{x \in \mathbb{R}^d} (y^\top x - H(x)) \quad \text{for all } y \in \mathbb{R}^d.$$

We split the differential operator in equation (5.1) into the linear diffusion part $A^{\text{diff}} f := \frac{1}{2} \Delta f$ and the non-linear first-order part $A^{\text{ham}} f := H(\nabla f)$. For a fixed sequence $(h_n)_{n \in \mathbb{N}} \subset (0, \infty)$ with $h_n \rightarrow 0$, the corresponding Chernoff one-step operators are given by

$$(J_n^{\text{diff}} f)(x) := \mathbb{E}[f(x + W_{h_n})] \quad \text{and} \quad (J_n^{\text{ham}} f)(x) := \sup_{\lambda \in \Lambda} (f(x + \lambda h_n) - H^*(\lambda) h_n)$$

¹The numerical experiments have been implemented in `Python` and were executed on a high-performance computing (HPC) cluster of ETH Zurich. The code can be found at the URL https://github.com/sgarale/chernoff_neural

for all $n \in \mathbb{N}$, $f \in C_b$ and $x \in \mathbb{R}^d$, where $(W_t)_{t \geq 0}$ is a d -dimensional Brownian motion satisfying $\text{cov}(W_1) = I_d$ and $\Lambda := \{H^* < \infty\}$. For every $f \in C_b^\infty$, Itô's and Taylor's formula imply

$$\lim_{n \rightarrow \infty} \frac{J_n^{\text{diff}} f - f}{h_n} = A^{\text{diff}} f := \frac{1}{2} \Delta f \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{J_n^{\text{ham}} f - f}{h_n} = A^{\text{ham}} f := H(\nabla f).$$

For every $n \in \mathbb{N}$, $f \in C_b$ and $x \in \mathbb{R}^d$, we define

$$(I_n f)(x) := (J_n^{\text{ham}} J_n^{\text{diff}} f)(x) = \sup_{\lambda \in \Lambda} (\mathbb{E}[f(x + \lambda h_n + W_{h_n})] - H^*(\lambda) h_n). \quad (5.2)$$

It follows from [12, Theorem 3.6] that there exists a strongly continuous convex monotone semigroup $(S(t))_{t \geq 0}$ on C_b given by

$$S(t)f := \lim_{n \rightarrow \infty} I(\pi_n^t) f \quad \text{for all } t \geq 0 \text{ and } f \in C_b$$

whose generator is given by

$$Af = \frac{1}{2} \Delta f + H(\nabla f) \quad \text{for all } f \in C_b^\infty.$$

Furthermore, the unique (viscosity) solution of equation (5.1) is given by $u(t, \cdot) := S(t)f$ for all $t \geq 0$. In the following, we implement the splitting scheme $u(t, \cdot) \approx I_n^{k_n} f$, where each diffusion step is followed by a non-linear transport step. We will approximate the one-step operator I_n both with a Chernoff-neural operator $\Phi_n \in \mathcal{CN}_{C_{\kappa_0}, C_{\kappa}}^{\mathcal{H}, \rho, \mathcal{L}}$ and an envelope-neural operator $\Phi_n \in \mathcal{EN}_{C_{\kappa}, C_{\kappa}}^{\text{ReLU}}$.

Lemma 5.1. *Let Λ be bounded and $H^*|_{\Lambda}$ Lipschitz continuous. Furthermore, let $\alpha := 1$, $\kappa \equiv 1$, $\kappa_0(x) := (1 + |x|^{q_0})^{-1}$ and $\mathcal{K} := \{\kappa\} \cup \{(1 + |\cdot|^{q'})^{-1} : q' \in (2, q_0]\}$ for some $q_0 \in (2, \infty)$. Then, the operators $(I_n)_{n \in \mathbb{N}}$ defined by equation (5.2) satisfy the Assumptions 2.2, 3.1, 4.1, 4.5 and 4.8.*

Proof. See Appendix C.1. □

For the numerical experiment, we choose $d = 1$, a bounded time horizon $T = 1$, $n = 30$, $h_n = \frac{1}{30}$ and $H(p) = |p|$ which implies $\Lambda = [-1, 1]$ and $H^* = +\infty \mathbf{1}_{[-1, 1]^c}$. Furthermore, we independently sample the parameters

$$a_i \sim \mathcal{U}(0.5, 1.5), \quad \sigma_i \sim \mathcal{U}(0.5, 1.5), \quad b_i \sim \mathcal{U}(-2, 2) \quad \text{and} \quad c_i \sim \mathcal{U}(-1, 1)$$

for $i = 1, \dots, I$ with $I := 2000$ according to uniform distributions. Using these parameters, we define the functions

$$f_{i,1}(x) := c_i + a_i \left(\Gamma \left(\frac{x - b_i}{\sigma_i} \right) - \frac{1}{2} \right) \quad \text{and} \quad f_{i,2}(x) := c_i - a_i \left(\Gamma \left(\frac{x - b_i}{\sigma_i} \right) - \frac{1}{2} \right) \quad (5.3)$$

which are split up into 90%/10% for training and testing, where Γ denotes the cumulative distribution function of the standard normal distribution $\mathcal{N}(0, 1)$. Since

$$\mathbb{E} \left[\Gamma \left(\frac{x + h_n \lambda + W_{h_n} - b_i}{\sigma_i} \right) \right] = \Gamma \left(\frac{x + h_n \lambda - b_i}{\sqrt{\sigma_i^2 + h_n}} \right)$$

and $f_{i,1}$ and $f_{i,2}$ are increasing and decreasing, respectively, the supremum over $\lambda \in [-1, 1]$ is attained at $\lambda_{i,1}^* = 1$ and $\lambda_{i,2}^* = -1$ such that

$$\begin{aligned} (I_n f_{i,1})(x) &= c_i + a_i \left(\Gamma \left(\frac{x + h_n - b_i}{\sqrt{\sigma_i^2 + h_n}} \right) - \frac{1}{2} \right), \\ (I_n f_{i,2})(x) &= c_i - a_i \left(\Gamma \left(\frac{x - h_n - b_i}{\sqrt{\sigma_i^2 + h_n}} \right) - \frac{1}{2} \right). \end{aligned} \quad (5.4)$$

In particular, the class of functions defined by equation (5.3) is preserved under iteration of I_n . Then, we learn the neural operators Φ_n by minimizing the mean squared error (MSE)

$$\frac{1}{2|\mathcal{I}|L} \sum_{i \in \mathcal{I}} \sum_{j=1}^2 \sum_{l=1}^L |(I_n f_{i,j})(x_l) - (\Phi_n f_{i,j})(x_l)|^2 \quad (5.5)$$

over the training set $\mathcal{I}$, where $(x_l)_{l=1,\dots,L} \sim \mathcal{N}(0, \sigma^2)$ are $L := 200$ independent and identically distributed (i.i.d.) evaluation points with $\sigma := 3$. In order to assess the out-of-sample performance of the trained operators, we evaluate them on the hyperbolic tangent function

$$f^{\text{th}}(x) := \frac{1}{2} \tanh(x)$$

which does not belong to the class of training functions defined by equation (5.3). As reference solutions, we use an explicit representation for $S(t)f_{i,j}$ which is similar to equation (5.4). Moreover, since f^{th} is increasing, it holds $(S(t)f^{\text{th}})(x) = \frac{1}{2}\mathbb{E}[\tanh(x + t + W_t)]$ which we approximate with Gauss–Hermite quadrature. For the Chernoff-neural operator $\Phi_n \in \mathcal{CN}_{C_{\kappa_0}, C_{\kappa}^{\alpha}}^{\mathcal{H}, \rho, \mathcal{L}}$, we use $M = 100$ neurons and $\tanh$ as activation function, where $\mathcal{H}$ and $\mathcal{L}$ consist of neural networks with 20 neurons and $\tanh$ as activation function, and apply the Adam algorithm over 10^4 epochs with learning rate 10^{-5} and batchsize 200 per type $j = 1, 2$. For the envelope-neural operator $\Phi_n \in \mathcal{EN}_{C_{\kappa}, C_{\kappa}}^{\text{ReLU}}$, we choose $M = 64$, a ReLU-neural network $\varphi: \mathbb{R}^M \rightarrow \mathbb{R}$ with 2 hidden layers of size 64 and 32 and apply the Adam algorithm over 10^4 epochs with learning rate 10^{-5} and batchsize 500. The results are reported in Figure 3.

5.2. Stochastic optimal control. In the second example, we approximate the value function

$$(S(t)f)(x) := \sup_{\substack{(\mu_s)_{s \in [0, T]} \subset \Xi, \\ (\sigma_s)_{s \in [0, T]} \subset \Sigma}} \mathbb{E} \left[f \left(x + \int_0^t \mu_s ds + \int_0^t \sigma_s dW_s \right) \right] \quad (5.6)$$

of a stochastic optimal control problem, where $(\mu_s)_{s \in [0, T]}$ and $(\sigma_s)_{s \in [0, T]}$ are predictable processes taking values in bounded subsets $\Xi \subset \mathbb{R}^d$ and $\Sigma \subset \mathbb{S}_+^d$, respectively. By [8, Theorem 6.2], the family $(S(t))_{t \geq 0}$ is a strongly continuous convex monotone semigroup on C_{κ} with $\kappa := (1 + |\cdot|^q)^{-1}$ for any $q \geq 2$ whose generator is given by

$$Af = \sup_{(\mu, \sigma) \in \Xi \times \Sigma} \left(\frac{1}{2} \text{tr}(\sigma^2 \nabla^2 f) + \mu^\top \nabla f \right) \quad \text{for all } f \in C_b^\infty.$$

Furthermore, it holds $S(t)f = \lim_{n \rightarrow \infty} I(\pi_n^t)f$ for all $t \geq 0$ and $f \in C_{\kappa}$, where


$$(I_n f)(x) := \sup_{(\mu, \sigma) \in \Xi \times \Sigma} \mathbb{E}[f(x + \mu h_n + \sigma W_{h_n})] \quad (5.7)$$

and $(h_n)_{n \in \mathbb{N}} \subset (0, \infty)$ is a fixed sequence with $h_n \rightarrow 0$. We learn this Chernoff one-step operator by a Chernoff-neural operator $\Phi_n \in \mathcal{CN}_{C_{\kappa_0}, C_{\kappa}^{\alpha}}^{\mathcal{H}, \rho, \mathcal{L}}$ and an envelope-neural operator $\Phi_n \in \mathcal{EN}_{C_{\kappa}, C_{\kappa}}^{\text{ReLU}}$.


Lemma 5.2. *Let $\alpha \in (0, 1]$, $2 \leq q < q_0 < \infty$, $\kappa(x) := (1 + |x|^q)^{-1}$, $\kappa_0(x) := (1 + |x|^{q_0})^{-1}$ and $\mathcal{K} := \{(1 + |\cdot|^{q'})^{-1} : q' \in [q, q_0]\}$. Then, the family $(I_n)_{n \in \mathbb{N}}$ defined by equation (5.7) satisfies the Assumptions 2.2, 3.1, 4.1, 4.5 and 4.8.*

Proof. See Appendix C.2. □


For the numerical experiment, we choose $d = 1$, a finite time horizon $T = 1$, $n = 30$, $h_n = \frac{1}{30}$, $\Xi := [\underline{\mu}, \bar{\mu}] := [-0.25, 0.25]$, $\Sigma := [\underline{\sigma}, \bar{\sigma}] := [1.5, 2.0]$ and generate $I := 1000$ i.i.d. strike prices




(C1) MSE (5.5) between $\Phi_n^{k_n^t} f_{i,j}$ and $S(t)f_{i,j}$ along training epochs and time.



(E1) MSE (5.5) between $\Phi_n^{k_n^t} f_{i,j}$ and $S(t)f_{i,j}$ along training epochs and time.



(C2) Approximation of $S(t)f^{\text{th}}$ by $\Phi_n^{k_n^t} f^{\text{th}}$.



(E2) Approximation of $S(t)f^{\text{th}}$ by $\Phi_n^{k_n^t} f^{\text{th}}$.

FIGURE 3. Approximation of the solution operator $f \mapsto S(t)f := u(t, \cdot)$ of the PDE (5.1) by using the Chernoff one-step operator I_n defined in equation (5.2) which is learned with a Chernoff-neural operator $\Phi_n \in \mathcal{CN}_{C_{\kappa_0}, C_{\kappa}^{\alpha}}^{\mathcal{H}, \rho, \mathcal{L}}$ in (C1)–(C2) and an envelope-neural operator $\Phi_n \in \mathcal{EN}_{C_{\kappa}, C_{\kappa}}^{\text{ReLU}}$ in (E1)–(E2). In (C1)+(E1), the MSE (5.5) on the train/test set is evaluated. In (C2)+(E2), the approximation of $S(t)f^{\text{th}}$ by $\Phi_n^{k_n^t} f^{\text{th}}$ is shown.

$(K_i)_{i=1, \dots, I} \sim \mathcal{N}(0, \sigma^2)$ with $\sigma = 2$. We use these parameters to define the functions

$$\begin{aligned} f_{i,1}(x) &= (x - K_i)_+, & f_{i,2}(x) &= (K_i - x)_+, \\ f_{i,3}(x) &= -(x - K_i)_+, & f_{i,4}(x) &= -(K_i - x)_+, \end{aligned} \tag{5.8}$$

where $s_+ := \max(s, 0)$, which are split up into 90%/10% for training and testing equally among the different types. Note that the operators $I_{n,(\mu,\sigma)}$ applied to call/put functions are given by

$$\begin{aligned} (I_{n,(\mu,\sigma)}(\cdot - K)_+)(x) &= \mathbb{E}[(x + \mu h_n + \sigma W_{h_n} - K)_+] \\ &= (x + \mu h_n - K) \Gamma\left(\frac{x + \mu h_n - K}{\sigma \sqrt{h_n}}\right) + \sigma \sqrt{h_n} \gamma\left(\frac{x + \mu h_n - K}{\sigma \sqrt{h_n}}\right), \end{aligned} \quad (5.9)$$

$$\begin{aligned} (I_{n,(\mu,\sigma)}(K - \cdot)_+)(x) &= \mathbb{E}[(K - x - \mu h_n - \sigma W_{h_n})_+] \\ &= (K - x - \mu h_n) \Gamma\left(\frac{K - x - \mu h_n}{\sigma \sqrt{h_n}}\right) + \sigma \sqrt{h_n} \gamma\left(\frac{K - x - \mu h_n}{\sigma \sqrt{h_n}}\right), \end{aligned} \quad (5.10)$$

where Γ (resp., γ) denotes the cumulative distribution (resp., probability density) function of the standard normal distribution $\mathcal{N}(0, 1)$. Consequently, the functions $I_n f_{i,j}$ are explicitly given by the equations (5.9) and (5.10), where the supremum over (μ, σ) is attained at the following values: $(\mu, \sigma) = (\bar{\mu}, \bar{\sigma})$ if $j = 1$, $(\mu, \sigma) = (\underline{\mu}, \bar{\sigma})$ if $j = 2$, $(\mu, \sigma) := (\underline{\mu}, \underline{\sigma})$ if $j = 3$ and $(\mu, \sigma) = (\bar{\mu}, \underline{\sigma})$ if $j = 4$. In order to assess the out-of-sample performance of the trained operators, we evaluate them on the absolute value function

$$f^{\text{ab}}(x) := |x|$$

which does not belong to the training functions given by equation (5.8). As reference solution, we compute $S(t)f_{i,j}$ and $S(t)f^{\text{ab}}$ using a finite difference scheme for the corresponding fully nonlinear HJB equation. We learn the neural operators by minimizing the mean squared error (MSE) defined in equation (5.5) over the training set, where $(x_l)_{l=1,\dots,L} \sim \mathcal{N}(0, \sigma^2)$ are $L = 200$ i.i.d. evaluation points with $\sigma = 3$. The other parameters are chosen as in Section 5.1, except that the Chernoff-neural operator has ReLU activation functions and learning rate $2 \cdot 10^{-5}$ and the envelope-neural operator has $M = 8$ neurons and learning rate 10^{-5} with the ReLU-neural network $\varphi: \mathbb{R}^M \rightarrow \mathbb{R}$ having two hidden layers of 8 and 4 neurons. Both operators are trained with batchsize 100 per type $j = 1, \dots, 4$. The results are reported in Figure 4.

5.3. Stochastic processes under model uncertainty. In the third example, we follow [5, 25] to study strongly continuous convex monotone semigroups arising from Wasserstein perturbations of the transition probabilities of Lévy processes. To that end, let $p > 1$ and $(X_t)_{t \geq 0}$ be a Lévy process with transition probabilities

$$p_t(x, B) := \varpi_t(\{y \in \mathbb{R}^d : x + y \in B\}) \quad \text{for all } t \geq 0, x \in \mathbb{R}^d \text{ and } B \in \mathcal{B}(\mathbb{R}^d),$$


where $(\varpi_t)_{t \geq 0}$ is a family of probability measures with finite p -th moment such that

$$\lim_{t \rightarrow 0} \int_{\mathbb{R}^d} |y|^p \varpi_t(dy) = 0. \quad (5.11)$$


In addition, let $\eta: \mathbb{R}_+ \rightarrow [0, \infty]$ be a convex non-decreasing function with $\eta(0) = 0$ and $\eta \not\equiv 0$ which is locally Lipschitz continuous on $\{\eta < \infty\}$. Moreover, the function $\mathbb{R}_+ \rightarrow [0, \infty]$, $v \mapsto \eta(v^{1/p})$ is supposed to be convex which implies $\liminf_{v \rightarrow \infty} \frac{\eta(v)}{v^p} > 0$. For every $n \in \mathbb{N}$, $f \in C_b$ and $x \in \mathbb{R}^d$, we define

$$(I_n f)(x) := \sup_{\nu \in \mathcal{P}_p(\mathbb{R}^d)} \left(\int_{\mathbb{R}^d} f(x + z) \nu(dz) - \eta\left(\frac{\mathcal{W}_p(\varpi_{h_n}, \nu)}{h_n}\right) h_n \right), \quad (5.12)$$


where $\mathcal{P}_p(\mathbb{R}^d)$ consists of all probability measures on $\mathcal{B}(\mathbb{R}^d)$ with finite p -th moment, $\mathcal{W}_p$ denotes the p -Wasserstein distance on $\mathcal{P}_p(\mathbb{R}^d)$ and $(h_n)_{n \in \mathbb{N}} \subset (0, \infty)$ is a fixed sequence with $h_n \rightarrow 0$. This Chernoff one-step operator incorporates nonparametric model uncertainty by taking the fixed transition probabilities of the Lévy process $(X_t)_{t \geq 0}$ as reference model and weighting all other




(C1) MSE (5.5) between $\Phi_n^{k_n^t} f_{i,j}$ and $S(t)f_{i,j}$ along training epochs and time.



(E1) MSE (5.5) between $\Phi_n^{k_n^t} f_{i,j}$ and $S(t)f_{i,j}$ along training epochs and time.



(C2) Approximation of $S(t)f^{ab}$ by $\Phi_n^{k_n^t} f^{ab}$.



(E2) Approximation of $S(t)f^{ab}$ by $\Phi_n^{k_n^t} f^{ab}$.

FIGURE 4. Approximating the stochastic control problem $f \mapsto S(t)f$ in (5.6) by the Chernoff one-step operator $I_n f$ in (5.7) which is learned with a Chernoff-neural operator $\Phi_n \in \mathcal{CN}_{C_{\kappa_0}, C_{\kappa}^{\alpha}}^{\mathcal{H}, \rho, \mathcal{L}}$ in (C1)–(C2) and an envelope-neural operator $\Phi_n \in \mathcal{EN}_{C_{\kappa}, C_{\kappa}}^{\text{ReLU}}$ in (E1)–(E2). In (C1)+(E1), the MSE (5.5) on the train/test set is evaluated. In (C2)+(E2), the approximation of $S(t)f^{ab}$ by $\Phi_n^{k_n^t} f^{ab}$ is shown.

probability measures according to their distance to the reference model. For instance, in case that $\eta := +\infty \mathbb{1}_{\{0\}^c}$, the Chernoff one-step operators coincide with the linear transition semigroup

$$(T(t)f)(x) := \int_{\mathbb{R}^d} f(x+y) \varpi_t(dy)$$

of the Lévy process. Following [25], it is possible to generalize the framework to reference dynamics of the form $X_t^x = \psi_t(x) + Y_t$ with a Lévy process $(Y_t)_{t \geq 0}$ and a deterministic drift $(\psi_t)_{t \geq 0}$ which covers, for instance, Ornstein–Uhlenbeck processes. It follows from [25, Theorem 3.14] that there exists a strongly continuous convex monotone semigroup $(S(t))_{t \geq 0}$ on C_b given by

$$S(t)f := \lim_{n \rightarrow \infty} I(\pi_n^t)f \quad \text{for all } t \geq 0 \text{ and } f \in C_b.$$

Furthermore, the generator of $(S(t))_{t \geq 0}$ is given by

$$Af = Bf + \eta^*(|\nabla f|) \quad \text{for all } f \in D(B) \cap C_b^1, \quad (5.13)$$

where $\eta^*(w) := \sup_{v \geq 0} (vw - \eta(v))$ for all $w \geq 0$ and B denotes the generator of $(T(t))_{t \geq 0}$.

Lemma 5.3. *Let $\alpha := 1$, $\kappa \equiv 1$ and assume $C_b^\infty \subset D(B)$. Then, the operators $(I_n)_{n \in \mathbb{N}}$ defined by equation (5.12) satisfy the Assumptions 4.1(i)–(vii), 4.5 and 4.8.*

Proof. See Appendix C.3. $\square$

Looking at the generator in equation (5.13), it becomes apparent that the same semigroup can be constructed by only taking parametric drift uncertainty into consideration. Indeed, let

$$(J_n f)(x) := \sup_{\lambda \in \mathbb{R}^d} \left(\int_{\mathbb{R}^d} f(x + \lambda h_n + y) \varpi_{h_n}(dy) - \eta(|\lambda|) h_n \right) \quad (5.14)$$

for all $n \in \mathbb{N}$, $f \in C_b$ and $x \in \mathbb{R}^d$. By [25, Section 4.5], there exists another strongly continuous convex monotone semigroup $(\tilde{S}(t))_{t \geq 0}$ on C_b given by

$$\tilde{S}(t)f := \lim_{n \rightarrow \infty} J(\pi_n^t)f \quad \text{for all } t \geq 0 \text{ and } f \in C_b$$

whose generator is given by

$$\tilde{A}f = \sup_{\lambda \in \mathbb{R}^d} (Bf + \lambda^T \nabla f - \eta(|\lambda|)) = Bf + \eta^*(|\nabla f|) \quad \text{for all } f \in D(B) \cap C_b^1.$$

Lemma 5.4. *Let $\alpha := 1$, $\kappa \equiv 1$ and assume $C_b^\infty \subset D(B)$. Then, the operators $(J_n)_{n \in \mathbb{N}}$ defined by equation (5.14) satisfy the Assumptions 4.1, 4.5 and 4.8. In particular,*

$$S(t)f = \tilde{S}(t)f \quad \text{for all } t \geq 0 \text{ and } f \in C_b.$$

Proof. See Appendix C.4. $\square$

By choosing $\nu = \varpi_{h_n} * \delta_{\lambda h_n}$ for all $\lambda \in \mathbb{R}^d$, we see that $J_n f \leq I_n f$ for all $n \in \mathbb{N}$ and $f \in C_b$. It has further been shown in [25, Lemma 3.10 and Section 4.5] that the iterated operators satisfy

$$J_n^{k_n^t} f \leq S(t)f \leq I_n^{k_n^t} f \quad \text{for all } t \geq 0, n \in \mathbb{N} \text{ and } f \in C_b.$$

In addition, if $(h_n)_{n \in \mathbb{N}}$ defines a sequence of refining partitions, e.g., for $h_n := 2^{-n}$, it holds

$$I(\pi_n^t)f \downarrow S(t)f \quad \text{and} \quad J(\pi_n^t)f \uparrow S(t)f \quad \text{for all } t \geq 0 \text{ and } f \in C_b$$

meaning that the first sequence is decreasing and the second one is increasing.

For our numerical example, we choose $d = 1$, a bounded time horizon $T = 1$, $n = 30$, $h_n = \frac{1}{30}$ and consider $(X_t)_{t \geq 0} = (W_t)_{t \geq 0}$. Furthermore, let $p = 2$ and $\eta(v) = \frac{1}{6}v^6$ with convex conjugate $\eta^*(w) = \frac{5}{6}w^{6/5}$. Note that the mapping $v \mapsto \eta(v^{1/2}) = \frac{v^3}{6}$ is convex and $\lim_{v \rightarrow \infty} \frac{\eta(v)}{v^2} = \infty$. Since the rescaled penalty heavily penalizes all measures $\nu \in \mathcal{P}_p(\mathbb{R}^d)$ with $\mathcal{W}_p(\varpi_{h_n}, \nu) > h_n$, the supremum in equation (5.12) can be seen as a smoothed version of the supremum over a Wasserstein ball. We therefore proceed as in Section 5.2. To that end, we generate $I := 1000$ i.i.d. midpoints $(c_i)_{i=1, \dots, I} \sim \mathcal{N}(0, 1)$ and half-widths $(w_i)_{i=1, \dots, I} \sim \mathcal{U}(\frac{1}{4}, \frac{3}{2})$, set $K_{i,1} := c_i - w_i$ and $K_{i,2} := c_i + w_i$ and define the bull and bear spread functions

$$\begin{aligned} f_{i,1}(x) &= (x - K_{i,1})_+ - (x - K_{i,2})_+, & f_{i,2}(x) &= (K_{i,2} - x)_+ - (K_{i,1} - x)_+, \\ f_{i,3}(x) &= -f_{i,1}(x), & f_{i,4}(x) &= -f_{i,2}(x), \end{aligned} \quad (5.15)$$

which are split up into 90%/10% for training and testing equally among the different types.

We implement I_n as an envelope-neural operator using the parametric representation of [55], where the supremum in equation (5.12) is restricted to measures of the form $\varpi_{h_n} \circ (\text{id} + h_n a)^{-1}$ with vector fields $a \in L^p(\varpi_{h_n}; \mathbb{R})$. Restricting the supremum yields the envelope-neural operator

$$(\Phi_n^I f)(x) := \max_{m=1, \dots, M} \left(\int_{\mathbb{R}} f(x + y + a_m(y)) \varpi_{h_n}(\mathrm{d}y) - h_n \eta \left(\frac{\|a_m\|_{L^p(\varpi_{h_n}; \mathbb{R})}}{h_n} \right) \right) \quad (5.16)$$

with finitely many trainable neurons $(a_m)_{m=1, \dots, M} \subset L^p(\varpi_{h_n}; \mathbb{R})$. For the implementation, we choose affine functions $a_{(\varrho, \beta)}(y) := \varrho + \beta y$ with $\varrho, \beta \in \mathbb{R}$ for which the transported measure is given by $\varpi_{h_n} \circ (\text{id} + a_{(\varrho, \beta)})^{-1} = \mathcal{N}(\varrho, (1 + \beta)^2 h_n)$. Hence, for the piecewise linear payoffs defined by equation (5.15) the inner integral in equation (5.16) is a Bachelier price and therefore available in closed form. Moreover, the penalization term is explicitly given by $\|a_{(\varrho, \beta)}\|_{L^2(\varpi_{h_n})}^2 = |\varrho|^2 + \beta^2 h_n$. For the operator J_n defined by equation (5.14), the envelope-neural operator is given by

$$(\Phi_n^J f)(x) := \max_{m=1, \dots, M} \left(\int_{\mathbb{R}} f(x + \lambda_m h_n + y) \varpi_{h_n}(\mathrm{d}y) - \eta(|\lambda_m|) h_n \right) \quad (5.17)$$

with finitely many trainable neurons $(\lambda_m)_{m=1, \dots, M} \subset \mathbb{R}$.

The neural operators are trained by minimizing the MSE (5.5) over $L = 100$ i.i.d. evaluation points $(x_l)_{l=1, \dots, L} \sim \mathcal{N}(0, \sigma^2)$ with $\sigma = 3$. Both envelope-neural operators use $M = 32$ neurons and the maximum readouts of the equations (5.16) and (5.17), respectively, so that Φ_n^I has 64 and Φ_n^J has 32 trainable parameters. Both are trained by applying the Adam algorithm over 10^4 epochs with learning rate 10^{-5} and batchsize 100 per type $j = 1, \dots, 4$. The training labels $I_n f_{i,j}$ of the envelope-neural operator Φ_n^I are computed from the dual representation

$$(I_n f)(x) = \min_{\lambda \geq 0} \left(\int_{\mathbb{R}} \sup_{w \in \mathbb{R}} (f(x + y + w) - \lambda |w|^p) \varpi_{h_n}(\mathrm{d}y) + G^*(\lambda) \right), \quad (5.18)$$

see [4, Theorem 2.4], where G^* denotes the convex conjugate of $G(\delta) := h_n \eta(\delta^{1/p}/h_n)$. In our case, we have $G(\delta) = \frac{\delta^3}{6h_n^5}$ and $G^*(\lambda) = \frac{2}{3} \sqrt{2h_n^5} \lambda^{3/2}$. Note that, for a one-dimensional piecewise linear payoff, the computation of the inner supremum in equation (5.18) reduces to maximizing over a finite set of points and the integral can be computed with a one-dimensional quadrature. The training labels $J_n f_{i,j}$ of the envelope-neural operator Φ_n^J are obtained by evaluating the supremum in equation (5.14) over a discretized grid of drifts $\lambda \in \mathbb{R}$.

In order to assess the out-of-sample performance of the trained operators, we evaluate them on the butterfly option

$$f^{\text{bf}}(x) := (x - K_1)_+ - 2(x - K_2)_+ + (x - K_3)_+ \quad \text{with} \quad K_1 := -1, K_2 := 0 \text{ and } K_3 := 1 \quad (5.19)$$

which is not monotone and therefore does not belong to any of the four classes of training functions defined in (5.15). As reference solution, we compute $S(t)f^{\text{bf}}$ using a finite difference scheme for the corresponding HJB equation.

The results of the training are reported in Figure 5. In Figure 6, we repeat the training of both neural operators for different numbers of steps n with step size $h_n = T/n$ and display the two terminal values $(\Phi_n^I)^{k_n^T} f^{\text{bf}}$ and $(\Phi_n^J)^{k_n^T} f^{\text{bf}}$ together with the finite difference solution $S(T)f^{\text{bf}}$ at the final time T . The gap between the iterated envelope-neural operators is decreasing in n which illustrates how the nonparametric Wasserstein uncertainty and the parametric drift uncertainty lead to the same continuous time limit in accordance with Lemma 5.4.

APPENDIX A. WEIGHTED HÖLDER SPACES

We prove some basic properties of the weighted Hölder spaces from Section 2. Recall that, for bounded continuous functions $\kappa', \kappa: \mathbb{R}^d \rightarrow (0, \infty)$, we write $\kappa' \lesssim \kappa$ if and only if $\kappa' \leq c\kappa$ for some




FIGURE 5. Approximating the Wasserstein-perturbation semigroup $f \mapsto S(t)f$ with two envelope-neural operators, evaluated on the out-of-sample butterfly f^{bf} in (5.19). The operator Φ_n^I in (5.16) in (I1)-(I2), and the operator Φ_n^J in (5.17) in (J1)-(J2). In (I1)+(J1), the MSE between the iterations $(\Phi_n^I)^{k_n^t} f^{\text{bf}}$, respectively $(\Phi_n^J)^{k_n^t} f^{\text{bf}}$, and $S(t)f^{\text{bf}}$ is displayed along the training epochs and the time $t = kh_n$, $k = 1, \dots, n$. In (I2)+(J2), the approximation of $S(t)f^{\text{bf}}$ by the two iterated neural operators is compared with the finite difference solution of the corresponding HJB equation over $[0, T]$.

constant $c \geq 0$, and $\kappa' \lesssim \kappa$ if and only if $\lim_{|x| \rightarrow \infty} \frac{\kappa'(x)}{\kappa(x)} = 0$. Furthermore, we define

$$\bar{\kappa}(x, y) := \frac{2\kappa(x)\kappa(y)}{\kappa(x) + \kappa(y)} = \frac{2}{\kappa(x)^{-1} + \kappa(y)^{-1}} \quad \text{for all } x, y \in \mathbb{R}^d.$$

Proposition A.1. *Let $\alpha \in [0, 1]$ and $\kappa: \mathbb{R}^d \rightarrow (0, \infty)$ be a bounded continuous function. Then, the space $(C_{\kappa}^{\alpha}, \|\cdot\|_{\alpha, \kappa})$ is complete.*




FIGURE 6. Terminal values $(\Phi_n^I)^{k_n^T} f^{\text{bf}}$ and $(\Phi_n^J)^{k_n^T} f^{\text{bf}}$ of the two envelope-neural operators of Section 5.3, evaluated on the out-of-sample butterfly f^{bf} in (5.19) and compared with the finite difference solution $S(T)f^{\text{bf}}$ at the terminal time T . Both operators are retrained for each number of steps $n \in \{10, 20, 30, 40\}$, with step size $h_n = T/n$. The shades run from light to dark as n grows.

Proof. Let $(f_n)_{n \in \mathbb{N}} \subset C_\kappa^\alpha$ be a Cauchy sequence. Since $(f_n)_{n \in \mathbb{N}}$ is also a Cauchy sequence in the Banach space C_κ , there exists $f \in C_\kappa$ with $\|f - f_n\|_\kappa \rightarrow 0$. By definition of the norm, we obtain

$$\|f\|_{\alpha, \kappa} \leq \sup_{n \in \mathbb{N}} \|f_n\|_{\alpha, \kappa} < \infty.$$

Moreover, for every $\varepsilon > 0$, there exist $n_0 \in \mathbb{N}$ with $\|f_m - f_n\|_{\alpha, \kappa} < \varepsilon$ for all $m, n \geq n_0$. Hence, for every $x, y \in \mathbb{R}^d$ with $x \neq y$ and $n \geq n_0$, we obtain

$$\begin{aligned} \frac{|(f - f_n)(x) - (f - f_n)(y)|}{|x - y|^\alpha} \bar{\kappa}(x, y) &= \lim_{m \rightarrow \infty} \frac{|(f_m - f_n)(x) - (f_m - f_n)(y)|}{|x - y|^\alpha} \bar{\kappa}(x, y) \\ &\leq \lim_{m \rightarrow \infty} \|f_m - f_n\|_{\alpha, \kappa} \leq \varepsilon. \end{aligned} \quad \square$$

Theorem A.2. Let $0 \leq \alpha' \leq \alpha \leq 1$ and $\kappa' \lesssim \kappa: \mathbb{R}^d \rightarrow (0, \infty)$ be bounded continuous functions. Then, the embedding $C_\kappa^\alpha \hookrightarrow C_{\kappa'}^{\alpha'}$ is continuous. Moreover, if $\alpha' < \alpha$ and $\kappa' \lesssim \kappa$, then $C_\kappa^\alpha \hookrightarrow C_{\kappa'}^{\alpha'}$.

Proof. Let $f \in C_\kappa^\alpha$. For every $x, y \in \mathbb{R}^d$ with $|x - y| \leq 1$, it follows from $\alpha' \leq \alpha$ and $\kappa' \leq c\kappa$ that $\bar{\kappa}'(x, y) = \frac{2}{\kappa'(x)^{-1} + \kappa'(y)^{-1}} \leq c \frac{2}{\kappa(x)^{-1} + \kappa(y)^{-1}} \leq c \bar{\kappa}(x, y)$ and therefore

$$\frac{|f(x) - f(y)|}{|x - y|^{\alpha'}} \bar{\kappa}'(x, y) \leq c \frac{|f(x) - f(y)|}{|x - y|^\alpha} \bar{\kappa}(x, y).$$

Moreover, for every $x, y \in \mathbb{R}^d$ with $|x - y| > 1$, the inequality $\bar{\kappa}'(x, y) \leq 2 \min(\kappa'(x), \kappa'(y))$ implies

$$\frac{|f(x) - f(y)|}{|x - y|^{\alpha'}} \bar{\kappa}'(x, y) \leq 2|f(x) - f(y)| \min(\kappa'(x), \kappa'(y)) \leq 2(|f(x)|\kappa'(x) + |f(y)|\kappa'(y)).$$

Hence, it holds $\|f\|_{\alpha', \kappa'} \leq 4c\|f\|_{\alpha, \kappa}$ for all $f \in C_\kappa^\alpha$.

Now, let $\alpha' < \alpha$ and $\kappa' \ll \kappa$. For every $f \in C_\kappa^\alpha$, it holds

$$\lim_{|x| \rightarrow \infty} |f(x)|\kappa'(x) \leq \|f\|_{\alpha, \kappa} \lim_{|x| \rightarrow \infty} \frac{\kappa'(x)}{\kappa(x)} = 0.$$

In addition, by using that $\kappa' \leq c\kappa$ implies $\bar{\kappa}'(x, y) \leq c\bar{\kappa}(x, y)$, we obtain

$$\begin{aligned} \lim_{\delta \rightarrow 0} \sup_{\substack{x \neq y, \\ |x-y| < \delta}} \frac{|f(x) - f(y)|}{|x-y|^{\alpha'}} \bar{\kappa}'(x, y) &\leq c \lim_{\delta \rightarrow 0} \left(\delta^{\alpha-\alpha'} \sup_{\substack{x \neq y, \\ |x-y| < \delta}} \frac{|f(x) - f(y)|}{|x-y|^\alpha} \bar{\kappa}(x, y) \right) \\ &\leq c \|f\|_{\alpha, \kappa} \lim_{\delta \rightarrow 0} \delta^{\alpha-\alpha'} = 0. \end{aligned} \quad \square$$

Theorem A.3. *Let $0 \leq \alpha' < \alpha \leq 1$ and $\kappa' \ll \kappa: \mathbb{R}^d \rightarrow (0, \infty)$ be bounded continuous functions. Then, the embedding $C_\kappa^\alpha \hookrightarrow C_{\kappa'}^{\alpha'}$ is compact.*

Proof. Let $(f_n)_{n \in \mathbb{N}} \subset C_\kappa^\alpha$ be bounded and $C := \sup_{n \in \mathbb{N}} \|f_n\|_{\alpha, \kappa}$. By Arzelà-Ascoli's theorem, there exists a diagonal sequence, which is again denoted by $(f_n)_{n \in \mathbb{N}}$, and $f \in C(\mathbb{R}^d)$ such that

$$\lim_{n \rightarrow \infty} \|f - f_n\|_{\infty, K} = 0 \quad \text{for all } K \Subset \mathbb{R}^d.$$

For every $x, y \in \mathbb{R}^d$, we use $\|f_n\|_{\alpha, \kappa} \leq C$ to obtain

$$\begin{aligned} |f(x)|\kappa(x) &= \lim_{n \rightarrow \infty} |f_n(x)|\kappa(x) \leq C, \\ |f(x) - f(y)|\bar{\kappa}(x, y) &= \lim_{n \rightarrow \infty} |f_n(x) - f_n(y)|\bar{\kappa}(x, y) \leq C|x-y|^\alpha \end{aligned}$$

which guarantees that $f \in C_\kappa^\alpha$ and $\|f - f_n\|_{\alpha, \kappa} \leq 2C$. It remains to show that $\|f - f_n\|_{\alpha', \kappa'} \rightarrow 0$. Let $\varepsilon > 0$. We choose $r > 0$ with $\sup_{|x| > r} \frac{\kappa'(x)}{\kappa(x)} < \frac{\varepsilon}{4C}$ and $n_0 \in \mathbb{N}$ with $\|f - f_n\|_{\infty, B_{\mathbb{R}^d}(r)} < \frac{\varepsilon}{2\|\kappa'\|_\infty}$. Then, for every $n \geq n_0$, we obtain

$$\|f - f_n\|_{\kappa'} \leq \|\kappa'\|_\infty \sup_{x \in B_{\mathbb{R}^d}(r)} |(f - f_n)(x)| + 2C \sup_{|x| > r} \frac{\kappa'(x)}{\kappa(x)} < \|\kappa'\|_\infty \frac{\varepsilon}{2\|\kappa'\|_\infty} + 2C \frac{\varepsilon}{4C} = \varepsilon. \quad (\text{A.1})$$

Furthermore, the estimates $\kappa' \leq c\kappa$ and $\bar{\kappa}'(x, y) \leq 2 \min(\kappa'(x), \kappa'(y))$ imply

$$\begin{aligned} &\sup_{x \neq y} \frac{|(f - f_n)(x) - (f - f_n)(y)|}{|x-y|^{\alpha'}} \bar{\kappa}'(x, y) \\ &\leq \left(\sup_{x \neq y} \frac{|(f - f_n)(x) - (f - f_n)(y)|}{|x-y|^\alpha} \bar{\kappa}(x, y) \right)^{\frac{\alpha'}{\alpha}} \left(\sup_{x \neq y} |(f - f_n)(x) - (f - f_n)(y)| \bar{\kappa}'(x, y) \right)^{1 - \frac{\alpha'}{\alpha}} \\ &\leq \left(c \sup_{x \neq y} \frac{|(f - f_n)(x) - (f - f_n)(y)|}{|x-y|^\alpha} \bar{\kappa}(x, y) \right)^{\frac{\alpha'}{\alpha}} \left(2 \sup_{x \in \mathbb{R}^d} |(f - f_n)(x)| \kappa'(x) \right)^{1 - \frac{\alpha'}{\alpha}} \\ &\leq (2Cc)^{\frac{\alpha'}{\alpha}} (2\|f - f_n\|_{\kappa'})^{1 - \frac{\alpha'}{\alpha}}. \end{aligned}$$

Combining the previous estimate with inequality (A.1) yields $\|f - f_n\|_{\alpha', \kappa'} \rightarrow 0$. $\square$

Theorem A.4. *Let $\alpha \in [0, 1)$ and $\kappa: \mathbb{R}^d \rightarrow (0, \infty)$ be a bounded continuous function. Then, the set $C_c^\alpha \subset C_\kappa^\alpha$ is dense.*

Proof. First, we show that every $f \in C_\kappa^\alpha$ can be approximated by a sequence of compactly supported functions. To do so, let $\chi \in C_b^\infty$ with $0 \leq \chi \leq 1$, $\chi \equiv 1$ on $B_{\mathbb{R}^d}(1)$ and $\chi \equiv 0$ on $B_{\mathbb{R}^d}(2)^c$. We define $\chi_r(x) := \chi(x/r)$, $f_r(x) := \chi_r(x)f(x)$ and $h_r(x) := f(x) - f_r(x) = (1 - \chi_r(x))f(x)$ for

all $r > 0$ and $x \in \mathbb{R}^d$. It holds $f_r, h_r \in c_\kappa^\alpha$ for all $r > 0$. Since $f \in c_\kappa^\alpha$, we have

$$m_f(r) := \sup_{|x| > r} |f(x)|\kappa(x) \rightarrow 0 \quad \text{as } r \rightarrow \infty, \quad (\text{A.2})$$

$$\omega_f(\delta) := \sup_{\substack{x \neq y, \\ |x-y| < \delta}} \frac{|f(x) - f(y)|}{|x-y|^\alpha} \bar{\kappa}(x, y) \rightarrow 0 \quad \text{as } \delta \rightarrow 0. \quad (\text{A.3})$$

It follows from equation (A.2) that $\|h_r\|_\kappa = \|f - f_r\|_\kappa \leq m_f(r) \rightarrow 0$ as $r \rightarrow \infty$. Moreover, there exists $L \geq 0$ with $\|\nabla \chi_r\|_\infty \leq L/r$ for all $r > 0$. For every $x, y \in \mathbb{R}^d$ with $0 < |x - y| < \delta$, we use equation (A.3), $\chi_r(x) = \chi_r(y) = 1$ in case that $|y| \leq r - \delta$ and $\bar{\kappa}(x, y) \leq 2\kappa(y)$ to estimate

$$\begin{aligned} |h_r(x) - h_r(y)|\bar{\kappa}(x, y) &\leq (1 - \chi_r(x))|f(x) - f(y)|\bar{\kappa}(x, y) + 2|\chi_r(x) - \chi_r(y)| \cdot |f(y)|\kappa(y) \\ &\leq \omega_f(\delta)|x - y|^\alpha + \frac{2L}{r}|x - y|m_f(r - \delta) \end{aligned}$$

which implies that

$$\sup_{\substack{x \neq y, \\ |x-y| < \delta}} \frac{|h_r(x) - h_r(y)|}{|x-y|^\alpha} \bar{\kappa}(x, y) \leq \omega_f(\delta) + \frac{2L}{r}\delta^{1-\alpha}m_f(r - \delta).$$

On the other hand, for every $x, y \in \mathbb{R}^d$ with $|x - y| \geq \delta$, it holds

$$\sup_{\substack{x \neq y, \\ |x-y| \geq \delta}} \frac{|h_r(x) - h_r(y)|}{|x-y|^\alpha} \bar{\kappa}(x, y) \leq \frac{2 \sup_{x \in \mathbb{R}^d} |h_r(x)|\kappa(x)}{\delta^\alpha} \leq \frac{2m_f(r)}{\delta^\alpha}.$$

Letting first $r \rightarrow \infty$ and then $\delta \rightarrow 0$ yields $\|f - f_r\|_{\alpha, \kappa} = \|h_r\|_{\alpha, \kappa} \rightarrow 0$.

Second, we show that every $f \in c_\kappa^\alpha$ with compact support can be approximated by a sequence of functions in C_c^∞ . Since κ is bounded away from zero on compact subsets, the weighted Hölder norm is equivalent to the classical Hölder norm on $\text{supp}(f)$, whence a standard mollification argument yields a sequence $(f_n)_{n \in \mathbb{N}} \subset C_c^\infty$ with $\|f - f_n\|_{\alpha, \kappa} \rightarrow 0$, see, e.g., [49, Proposition 0.2.1]. $\square$

APPENDIX B. WEIGHTED STONE-WEIERSTRASS THEOREMS FOR BANACH SCALES

Let $(\Theta, \preceq)$ be a pre-ordered set equipped with an additional relation $\prec$ satisfying $\theta_1 \prec \theta_2 \Rightarrow \theta_1 \preceq \theta_2$ such that, for every $k \in \mathbb{N}$ and $\theta_1, \dots, \theta_k \in \Theta$, there exist $l_*, l^* \in \{1, \dots, k\}$ with $\theta_{l_*} \preceq \theta_l \preceq \theta_{l^*}$ for all $l \in \{1, \dots, k\}$. Let X be a Banach space and $(X_\theta, \|\cdot\|_\theta)_{\theta \in \Theta}$ be a family of Banach spaces such that, for every $\theta' \preceq \theta$, it holds that $X_\theta \subset X_{\theta'} \subset X$ and the inclusion $\iota_{X_\theta, X_{\theta'}}: X_\theta \hookrightarrow X_{\theta'}$ is continuous, and for every $\theta' \prec \theta$, the set $B_{X_\theta}(r) := \{u \in X_\theta: \|u\|_\theta \leq r\}$ is compact in $X_{\theta'}$. Moreover, for a strictly increasing continuous function $\psi: [0, \infty) \rightarrow (0, \infty)$ with $\lim_{R \rightarrow \infty} \psi(R) = \infty$, we define the weight functions

$$\psi_\theta: X_\theta \rightarrow (0, \infty), \quad u \mapsto \psi(\|u\|_\theta) \quad \text{for all } \theta \in \Theta.$$

Then, for every $\theta' \prec \theta$ and $R > 0$, the pre-image $\psi_\theta^{-1}((0, R]) = B_{X_\theta}(\psi^{-1}(R)) \subset X_{\theta, \theta'}$ is compact and therefore ψ_θ is an admissible weight on $X_{\theta, \theta'} := (X_\theta, \|\cdot\|_{\theta'})$ in the sense of [19, Definition 2.1]. Furthermore, let Y be a vector space and $(Y_\theta, \|\cdot\|_\theta)_{\theta \in \Theta}$ be a family of Banach spaces such that, for every $\theta' \preceq \theta$, it holds $Y_\theta \subset Y_{\theta'} \subset Y$ and the inclusion $\iota_{Y_\theta, Y_{\theta'}}: Y_\theta \hookrightarrow Y_{\theta'}$ is continuous.

Remark B.1. In the proof of Theorem 3.5, we consider $\Theta := (\beta_q, \eta_q)_{q \in Q} \subset [0, \alpha] \times \mathcal{K}$, with totally ordered $(Q, \leq)$ satisfying $\beta_{q'} < \beta_q$ and $\eta_{q'} \lesssim \eta_q$ for all $q' < q$. We equip Θ with

$$(\beta_{q'}, \eta_{q'}) \preceq (\beta_q, \eta_q) :\Leftrightarrow q' \leq q \quad \text{and} \quad (\beta_{q'}, \eta_{q'}) \prec (\beta_q, \eta_q) :\Leftrightarrow q' < q.$$

Moreover, we set $X_{\theta, \theta'} := C_{\eta_q, \eta_{q'}}^{\beta_q, \beta_{q'}} = (C_{\eta_q}^{\beta_q}, \|\cdot\|_{\beta_{q'}, \eta_{q'}})$ and $Y_{\theta'} := C_{\eta_{q'}}^{\beta_{q'}}$ for $\theta' := (\beta_{q'}, \eta_{q'}) \prec \theta := (\beta_q, \eta_q)$.

We further introduce the intersection mapping spaces

$$\begin{aligned}\bigcap_{\theta' \prec \theta} \mathcal{B}_\psi(X_{\theta, \theta'}; Y_{\theta'}) &:= \{f: X \rightarrow Y: f|_{X_\theta} \in \mathcal{B}_\psi(X_{\theta, \theta'}; Y_{\theta'}) \text{ for all } \theta' \prec \theta \in \Theta\}, \\ \bigcap_{\theta' \prec \theta} \mathcal{C}_b(X_{\theta, \theta'}; Y_{\theta'}) &:= \{f: X \rightarrow Y: f|_{X_\theta} \in \mathcal{C}_b(X_{\theta, \theta'}; Y_{\theta'}) \text{ for all } \theta' \prec \theta \in \Theta\}\end{aligned}$$

which satisfy $\bigcap_{\theta' \prec \theta} \mathcal{C}_b(X_{\theta, \theta'}; Y_{\theta'}) \subset \bigcap_{\theta' \prec \theta} \mathcal{B}_\psi(X_{\theta, \theta'}; Y_{\theta'})$ since $\mathcal{C}_b(X_{\theta, \theta'}; Y_{\theta'}) \subset \mathcal{B}_\psi(X_{\theta, \theta'}; Y_{\theta'})$. We equip $\bigcap_{\theta' \prec \theta} \mathcal{B}_\psi(X_{\theta, \theta'}; Y_{\theta'})$ with the locally convex topology generated by the seminorms

$$\|f\|_{\mathcal{B}_\psi(X_{\theta, \theta'}; Y_{\theta'})} := \sup_{u \in X_\theta} \frac{\|f(u)\|_{\theta'}}{\psi(\|u\|_\theta)} \quad \text{for all } \theta' \prec \theta \in \Theta.$$

A zero-neighborhood basis of $\bigcap_{\theta' \prec \theta} \mathcal{B}_\psi(X_{\theta, \theta'}; Y_{\theta'})$ is given by sets of the form

$$\left\{ f \in \bigcap_{\theta' \prec \theta} \mathcal{B}_\psi(X_{\theta, \theta'}; Y_{\theta'}) : \max_{l=1, \dots, k} \|f\|_{\mathcal{B}_{\psi_{\theta_l}}(X_{\theta_l, \theta'_l}; Y_{\theta'_l})} < \varepsilon \right\},$$

where $\varepsilon > 0$, $k \in \mathbb{N}$ and $\theta'_l \prec \theta_l \in \Theta$, $l = 1, \dots, k$. Recall that a vector space $\mathcal{A}$ of mappings $a: X \rightarrow \mathbb{R}$ is a *subalgebra* if $a_1 \cdot a_2 \in \mathcal{A}$ for all $a_1, a_2 \in \mathcal{A}$, where $(a_1 \cdot a_2)(u) := a_1(u)a_2(u)$. The set $\mathcal{A}$ is *point separating* if, for every $u_1, u_2 \in X$ with $u_1 \neq u_2$, there is $a \in \mathcal{A}$ with $a(u_1) \neq a(u_2)$. Finally, $\mathcal{A}$ is called *nowhere vanishing* if, for every $u \in X$, there is $a \in \mathcal{A}$ with $a(u) \neq 0$.

Theorem B.2 ($\mathbb{R}$ -valued weighted Stone-Weierstrass). *Let $\mathcal{A} \subset \bigcap_{\theta' \prec \theta} \mathcal{C}_b(X_{\theta, \theta'}; \mathbb{R})$ be a point separating nowhere vanishing subalgebra. Then, the set $\mathcal{A}$ is dense in $\bigcap_{\theta' \prec \theta} \mathcal{B}_\psi(X_{\theta, \theta'}; \mathbb{R})$.*

Proof. We follow the proof of [19, Theorem 3.9]. Let $f \in \bigcap_{\theta' \prec \theta} \mathcal{B}_\psi(X_{\theta, \theta'}; \mathbb{R})$, $\varepsilon > 0$, $k \in \mathbb{N}$ and $\theta'_l \prec \theta_l \in \Theta$ for $l = 1, \dots, k$. By [19, Lemma 2.7(i)], there exists $R_1 > 0$ with

$$\max_{l=1, \dots, k} \sup_{u \in X_{\theta_l} \setminus K_{l, R_1}} \frac{1 + |f(u)|}{\psi(\|u\|_{\theta_l})} < \frac{\varepsilon}{5}, \quad (\text{B.1})$$

where $K_{l, R_1} := \psi_{\theta_l}^{-1}((0, R_1])$ denotes the compact pre-image of $\psi_{\theta_l}(\cdot) := \psi(\|\cdot\|_{\theta_l})$. Moreover, we define $M := (\min_{l=1, \dots, k} \inf_{u \in X_{\theta_l}} \psi(\|u\|_{\theta_l}))^{-1}$, let $b := 1 + \max_{l=1, \dots, k} \sup_{u \in K_{l, R_1}} |f(u)|$ and choose $R_2 \geq \max\{R_1, \frac{5b}{\varepsilon}\}$. Furthermore, there exists $\theta' \in \Theta$ with $\theta' \preceq \theta'_l$ for all $l \in \{1, \dots, k\}$. Since the set $K_2 := \bigcup_{l=1}^k K_{l, R_2} \subset X_{\theta'}$ is compact, we can apply the classical Stone-Weierstrass theorem [66] to $\mathcal{A}|_{K_2} \subset C(K_2, \|\cdot\|_{\theta'})$ to obtain $a \in \mathcal{A}$ with

$$\sup_{u \in K_2} |f(u) - a(u)| < \min\left\{1, \frac{\varepsilon}{5M}\right\}. \quad (\text{B.2})$$

We use that $g(s) := \max\{\min(s, b), -b\}$ satisfies $g(a(u)) = a(u)$ for all $u \in K_1 := \bigcup_{l=1}^k K_{l,R_1}$, the inequality $|a(u)| \leq 1 + |f(u)|$ for all $u \in K_2$ and the inequalities (B.1)–(B.2) to estimate

$$\begin{aligned}
& \max_{l=1,\dots,k} \|f - g \circ a\|_{\mathcal{B}_{\psi_{\theta_l}}(X_{\theta_l, \theta'_l}; \mathbb{R})} = \max_{l=1,\dots,k} \sup_{u \in X_{\theta_l}} \frac{|f(u) - g(a(u))|}{\psi(\|u\|_{\theta_l})} \\
& \leq \max_{l=1,\dots,k} \sup_{u \in X_{\theta_l} \setminus K_{l,R_1}} \frac{|f(u)|}{\psi(\|u\|_{\theta_l})} + \max_{l=1,\dots,k} \sup_{u \in K_{l,R_1}} \frac{|f(u) - g(a(u))|}{\psi(\|u\|_{\theta_l})} \\
& \quad + \max_{l=1,\dots,k} \sup_{u \in K_{l,R_2} \setminus K_{l,R_1}} \frac{|g(a(u))|}{\psi(\|u\|_{\theta_l})} + \max_{l=1,\dots,k} \sup_{u \in X_{\theta_l} \setminus K_{l,R_2}} \frac{|g(a(u))|}{\psi(\|u\|_{\theta_l})} \\
& \leq \max_{l=1,\dots,k} \sup_{u \in X_{\theta_l} \setminus K_{l,R_1}} \frac{1 + |f(u)|}{\psi(\|u\|_{\theta_l})} + M \sup_{u \in K_{l,R_1}} |f(u) - a(u)| \\
& \quad + \max_{l=1,\dots,k} \sup_{u \in K_{l,R_2} \setminus K_{l,R_1}} \frac{1 + |f(u)|}{\psi(\|u\|_{\theta_l})} + \max_{l=1,\dots,k} \sup_{u \in X_{\theta_l} \setminus K_{l,R_2}} \frac{b}{\psi(\|u\|_{\theta_l})} \\
& < \frac{\varepsilon}{5} + M \frac{\varepsilon}{5M} + \frac{\varepsilon}{5} + \frac{b}{R_2} \leq \frac{4\varepsilon}{5}. \tag{B.3}
\end{aligned}$$

Moreover, by the Weierstrass theorem [67], there exists a polynomial p on $\mathbb{R}$ with

$$|g(s) - p(s)| < \frac{\varepsilon}{5M} \quad \text{for all } |s| \leq c := \max_{l=1,\dots,k} \sup_{u \in X_{\theta_l}} |a(u)|.$$

Without loss of generality, we assume $p(0) = 0$, implying

$$\begin{aligned}
& \max_{l=1,\dots,k} \|g \circ a - p \circ a\|_{\mathcal{B}_{\psi_{\theta_l}}(X_{\theta_l, \theta'_l}; \mathbb{R})} = \max_{l=1,\dots,k} \sup_{u \in X_{\theta_l}} \frac{|g(a(u)) - p(a(u))|}{\psi(\|u\|_{\theta_l})} \\
& \leq M \max_{l=1,\dots,k} \sup_{u \in X_{\theta_l}} |g(a(u)) - p(a(u))| \leq M \sup_{|s| \leq c} |g(s) - p(s)| < \frac{M\varepsilon}{5M} = \frac{\varepsilon}{5}. \tag{B.4}
\end{aligned}$$

Combining the inequalities (B.3) and (B.4) yields $\max_{l=1,\dots,k} \|f - p \circ a\|_{\mathcal{B}_{\psi}(X_{\theta_l, \theta'_l}; \mathbb{R})} < \varepsilon$. Since $\mathcal{A}$ is an algebra, it holds that $p \circ a \in \mathcal{A}$ and the claim follows. $\square$

Following [19, Section 3.2], Theorem B.2 can be generalized to $\mathcal{A} \subset \bigcap_{\theta' \prec \theta} \mathcal{B}_{\psi}(X_{\theta, \theta'}; \mathbb{R})$ consisting of unbounded mappings. Given a vector space $\mathcal{A}$ of mappings $a: X \rightarrow \mathbb{R}$, a vector space $\mathcal{W}$ of mappings $w: X \rightarrow Y$ is an $\mathcal{A}$ -submodule if $a \cdot w \in \mathcal{W}$ for all $a \in \mathcal{A}$ and $w \in \mathcal{W}$, where $(a \cdot w)(u) := a(u)w(u)$.

Theorem B.3 (Vector-valued weighted Stone–Weierstrass). *Let $\mathcal{A} \subset \bigcap_{\theta' \prec \theta} C_b(X_{\theta, \theta'}; \mathbb{R})$ be a point separating nowhere vanishing subalgebra and let $\mathcal{W} \subset \bigcap_{\theta' \prec \theta} C_b(X_{\theta, \theta'}; Y_{\theta'})$ be an $\mathcal{A}$ -submodule such that $\mathcal{W}(u) := \{w(u) : w \in \mathcal{W}\}$ is a dense subset of $Y_{\theta'}$ for all $u \in X_{\theta}$ and $\theta', \theta \in \Theta$ with $\theta' \prec \theta$. Then, the set $\mathcal{W}$ is dense in $\bigcap_{\theta' \prec \theta} \mathcal{B}_{\psi}(X_{\theta, \theta'}; Y_{\theta'})$.*

Proof. Let $f \in \bigcap_{\theta' \prec \theta} \mathcal{B}_{\psi}(X_{\theta, \theta'}; Y_{\theta'})$, $\varepsilon > 0$, $k \in \mathbb{N}$ and $\theta'_l \prec \theta_l \in \Theta$ for $l = 1, \dots, k$. It follows from [19, Lemma 2.7(i)] that there exists $R_1 > 0$ with

$$\max_{l=1,\dots,k} \sup_{u \in X_{\theta_l} \setminus K_{l,R_1}} \frac{1 + \|f(u)\|_{\theta'_l}}{\psi(\|u\|_{\theta_l})} < \frac{\varepsilon}{4}, \tag{B.5}$$

where $K_{l,R_1} := \psi_{\theta_l}^{-1}((0, R_1])$. Let $M := (\min_{l=1,\dots,k} \inf_{u \in X_{\theta_l}} \psi(\|u\|_{\theta_l}))^{-1}$. We choose $\theta' \in \Theta$ with $\theta' \preceq \theta'_l$ for all $l \in \{1, \dots, k\}$ and observe that $K_1 := \bigcup_{l=1}^k K_{l,R_1}$ is compact in $X_{\theta'}$. Moreover, for

every $u \in K_1$, we define $L_u := \{l \in \{1, \dots, k\} : u \in X_{\theta_l}\}$ and choose $l_u \in L_u$ with $\theta'_l \preceq \theta'_{l_u}$ for all $l \in L_u$. Since $\mathcal{W}(u)$ is a dense subset of $Y_{\theta'_{l_u}}$, there exists $w_u \in \mathcal{W}$ with

$$\|f(u) - w_u(u)\|_{\theta'_{l_u}} < \frac{\varepsilon}{4c_u M},$$

where $c_u := 1 + \max_{l \in L_u} \|\iota_{\theta'_{l_u}, \theta'_l}\|_{L(Y_{\theta'_{l_u}}, Y_{\theta'_l})}$. Consequently, for every $l \in L_u$, it holds that

$$\|f(u) - w_u(u)\|_{\theta'_l} \leq \|\iota_{\theta'_{l_u}, \theta'_l}\|_{L(Y_{\theta'_{l_u}}, Y_{\theta'_l})} \|f(u) - w_u(u)\|_{\theta'_{l_u}} < c_u \frac{\varepsilon}{4c_u M} = \frac{\varepsilon}{4M}$$

and therefore

$$\frac{\|f(u) - w_u(u)\|_{\theta'_l}}{\psi(\|u\|_{\theta_l})} < M \frac{\varepsilon}{4M} = \frac{\varepsilon}{4}.$$

In addition, for every $u \in K_1$ and $l \in \{1, \dots, k\}$, the set

$$F_{u,l} := \left\{ v \in X_{\theta_l} : \frac{\|f(v) - w_u(v)\|_{\theta'_l}}{\psi(\|v\|_{\theta_l})} \geq \frac{\varepsilon}{4} \right\}$$

is compact in $X_{\theta'}$. Indeed, by applying [19, Lemma 2.7(i)] to f and using that w_u is uniformly bounded, we obtain

$$\lim_{R \rightarrow \infty} \sup_{v \in X_{\theta_l} \setminus K_{l,R}} \frac{\|f(v) - w_u(v)\|_{\theta'_l}}{\psi(\|v\|_{\theta_l})} = 0$$

which implies that $F_{u,l} \subset K_{l,R}$ for sufficiently large $R > 0$. Since $K_{l,R}$ is compact and $F_{u,l}$ is closed, the set $F_{u,l}$ is compact. In particular, it follows from $u \notin F_{u,l}$ that $U_u := X_{\theta'} \setminus \bigcup_{l=1}^k F_{u,l}$ is an open neighborhood of u in $X_{\theta'}$ satisfying

$$\frac{\|f(v) - w_u(v)\|_{\theta'_l}}{\psi(\|v\|_{\theta_l})} < \frac{\varepsilon}{4} \quad \text{for all } v \in U_u \cap X_{\theta_l}. \quad (\text{B.6})$$

Since $(U_u)_{u \in K_1}$ is an open cover of the compact set K_1 , there exist $u_1, \dots, u_N \in K_1$ satisfying $K_1 \subset \bigcup_{n=1}^N U_{u_n}$ and a partition of unity $(g_n)_{n=1, \dots, N} \subset C(K_1)$ subordinate to $(U_{u_n})_{n=1, \dots, N}$ which means that $0 \leq g_n \leq 1$, $\sum_{n=1}^N g_n = 1$ and $\text{supp}(g_n) \subset U_{u_n}$. By using again that K_1 is compact and $\text{supp}(g_n) \subset U_{u_n}$, we can extend each $g_n \in C(K_1)$ to some $\tilde{g}_n \in C(X_{\theta'})$ with $0 \leq \tilde{g}_n \leq 1$, $\tilde{g}_n|_{K_1} = g_n$ and $\text{supp}(\tilde{g}_n) \subset U_{u_n}$. By further choosing $\tilde{g}_0 \in C(X_{\theta'})$ with $0 \leq \tilde{g}_0 \leq 1$, $\tilde{g}_0|_{K_1} = 0$ and $\tilde{g}_0|_{X_{\theta'} \setminus K_1} > 0$, the normalized functions $\bar{g}_n := \frac{\tilde{g}_n}{G}$ with $G := \sum_{n=0}^N \tilde{g}_n$ for $n = 0, \dots, N$ form a partition of unity on $X_{\theta'}$ satisfying $\bar{g}_0|_{K_1} = 0$, $\bar{g}_n|_{K_1} = g_n$ and $\text{supp}(\bar{g}_n) \subset U_{u_n}$ for $n = 1, \dots, N$. Let $w_n := w_{u_n}$ for $n = 1, \dots, N$. Since $\bar{g}_0(u) > 0$ implies $u \notin K_1$ and therefore $u \in X_{\theta_l} \setminus K_{l,R_1}$ and $\bar{g}_n(u) > 0$ implies $u \in U_{u_n}$, it follows from inequality (B.5) and inequality (B.6) that

$$\begin{aligned} & \max_{l=1, \dots, k} \sup_{u \in X_{\theta_l}} \frac{\|f(u) - \sum_{n=1}^N \bar{g}_n(u) w_n(u)\|_{\theta'_l}}{\psi(\|u\|_{\theta_l})} \\ & \leq \max_{l=1, \dots, k} \sup_{u \in X_{\theta_l}} \bar{g}_0(u) \frac{\|f(u)\|_{\theta'_l}}{\psi(\|u\|_{\theta_l})} + \max_{l=1, \dots, k} \sup_{u \in X_{\theta_l}} \sum_{n=1}^N \bar{g}_n(u) \frac{\|f(u) - w_n(u)\|_{\theta'_l}}{\psi(\|u\|_{\theta_l})} \\ & < \frac{\varepsilon}{4} + \frac{\varepsilon}{4} = \frac{\varepsilon}{2} \end{aligned} \quad (\text{B.7})$$

Moreover, for every $n \in \{1, \dots, N\}$, we apply Theorem B.2 to obtain $a_n \in \mathcal{A}$ with

$$\max_{l=1, \dots, k} \|\bar{g}_n - a_n\|_{\mathcal{B}_\psi(X_{\theta_l, \theta'_l}; \mathbb{R})} < \frac{\varepsilon}{2c_{w_n} N}, \quad (\text{B.8})$$

where $c_{w_n} := 1 + \max_{l=1, \dots, k} \sup_{u \in X_{\theta_l}} \|w_n(u)\|_{Y_{\theta'_l}}$. Finally, the inequalities (B.7) and (B.8) imply

$$\begin{aligned} & \max_{l=1, \dots, k} \left\| f - \sum_{n=1}^N a_n w_n \right\|_{\mathcal{B}_\psi(X_{\theta_l, \theta'_l}; Y_{\theta'_l})} \\ & \leq \max_{l=1, \dots, k} \left\| f - \sum_{n=1}^N \bar{g}_n w_n \right\|_{\mathcal{B}_\psi(X_{\theta_l, \theta'_l}; Y_{\theta'_l})} + \sum_{n=1}^N \max_{l=1, \dots, k} \|\bar{g}_n - a_n\|_{\mathcal{B}_\psi(X_{\theta_l, \theta'_l}; \mathbb{R})} \sup_{u \in X_{\theta_l}} \|w_n(u)\|_{Y_{\theta'_l}} \\ & < \frac{\varepsilon}{2} + \sum_{n=1}^N \frac{\varepsilon}{2c_{w_n} N} c_{w_n} = \varepsilon. \end{aligned}$$

Since $\mathcal{W}$ is an $\mathcal{A}$ -submodule, it holds that $\sum_{n=1}^N a_n \cdot w_n \in \mathcal{W}$ and the claim follows. $\square$

Following [19, Section 3.3], Theorem B.3 can be generalized to $\mathcal{W} \subset \bigcap_{\theta' \prec \theta} \mathcal{B}_\psi(X_{\theta, \theta'}; Y_{\theta'})$ and $\mathcal{A} \subset \bigcap_{\theta' \prec \theta} \mathcal{B}_\psi(X_{\theta, \theta'}; \mathbb{R})$ consisting of unbounded mappings.

APPENDIX C. PROOF OF LEMMAS 5.1–5.4

Lemma C.1. *Let $q \geq 2$, $\mu \in \mathbb{R}^d$, $\sigma \in \mathbb{R}^{d \times m}$, $\Sigma := \sigma \sigma^\top \in \mathbb{S}_+^d$ and $(W_t)_{t \geq 0}$ be a m -dimensional Brownian motion. Define $X_t := \mu t + \sigma W_t$ for all $t \geq 0$ and $\omega_q := q|\mu| + \frac{q}{2} \text{tr}(\Sigma) + \frac{q(q-2)}{2} |\Sigma|$. Then,*

$$\mathbb{E}[1 + |x + X_t|^q] \leq e^{\omega_q t} (1 + |x|^q) \quad \text{for all } t \geq 0 \text{ and } x \in \mathbb{R}^d.$$

Proof. Define $V_q(x) := 1 + |x|^q$. The generator of the Itô process $(X_t)_{t \geq 0}$ is given by

$$(\mathcal{L}f)(x) := \mu^\top \nabla f(x) + \frac{1}{2} \text{tr}(\Sigma \nabla^2 f(x)).$$

Since $\nabla V_q(x) = q|x|^{q-2}x$ and $\nabla^2 V_q(x) = q|x|^{q-2}I_d + q(q-2)|x|^{q-4}xx^\top$ for all $x \neq 0$, which can be continuously extended to zero, we can use $|x|^{q-1} \leq 1 + |x|^q$ and $|x|^{q-2} \leq 1 + |x|^q$ to obtain

$$\begin{aligned} (\mathcal{L}V_q)(x) &= q|x|^{q-2}\mu^\top x + \frac{q}{2}|x|^{q-2}\text{tr}(\Sigma) + \frac{q(q-2)}{2}|x|^{q-4}x^\top \Sigma x \\ &\leq q|\mu| \cdot |x|^{q-1} + \frac{q}{2}(\text{tr}(\Sigma) + (q-2)|\Sigma|)|x|^{q-2} \leq \omega_q V_q(x). \end{aligned}$$

For every $t \geq 0$ and $x \in \mathbb{R}^d$, it follows from Itô's formula that

$$\mathbb{E}[V_q(x + X_t)] = V_q(x) + \int_0^t \mathbb{E}[\mathcal{L}V_q(x + X_s)] ds \leq V_q(x) + \omega_q \int_0^t \mathbb{E}[V_q(x + X_s)] ds.$$

Finally, Gronwall's lemma implies $\mathbb{E}[V_q(x + X_t)] \leq e^{\omega_q t} V_q(x)$ for all $t \geq 0$ and $x \in \mathbb{R}^d$. $\square$

C.1. Proof of Lemma 5.1. Regarding the verification of the Assumptions 2.2 and 4.1 as well as the extension $I_n : C_{\kappa_0} \rightarrow C_{\kappa_0}$, we refer to [12, Theorem 3.6], whereas Assumption 3.1(i) is clearly satisfied. Furthermore, by Lemma C.1, there exists a constant $\omega \geq 0$ with

$$\begin{aligned} \|I_{n, \lambda} f\|_{\alpha', \kappa'} &\leq \max \left\{ \sup_{x \in \mathbb{R}^d} \mathbb{E}[|f(x + Z_{n, \lambda})|] \kappa'(x), \sup_{x \neq y} \frac{\mathbb{E}[|f(x + Z_{n, \lambda}) - f(y + Z_{n, \lambda})|]}{|x - y|^{\alpha'}} \bar{\kappa}'(x, y) \right\} \\ &\leq \|f\|_{\alpha', \kappa'} \max \left\{ \sup_{x \in \mathbb{R}^d} \mathbb{E} \left[\frac{\kappa'(x)}{\kappa'(x + Z_{n, \lambda})} \right], \sup_{x \neq y} \mathbb{E} \left[\frac{\bar{\kappa}'(x, y)}{\bar{\kappa}'(x + Z_{n, \lambda}, y + Z_{n, \lambda})} \right] \right\} \\ &\leq e^{\omega h_n} \|f\|_{\alpha', \kappa'} \end{aligned}$$

for all $n \in \mathbb{N}$, $\lambda \in \Lambda$, $\alpha' \in [0, \alpha]$, $\kappa' \in \mathcal{K}$ and $f \in C_{\kappa'}^{\alpha'}$, where $(I_{n, \lambda} f)(x) := \mathbb{E}[f(x + Z_{n, \lambda})]$ and $Z_{n, \lambda} := \lambda h_n + W_{h_n}$. Indeed, for $\kappa'(x) = (1 + |x|^q)^{-1}$, it holds $\kappa'(x)^{-1} = 1 + |x|^q$ and

$\bar{\kappa}'(x, y)^{-1} = \frac{\kappa'(x)^{-1} + \kappa'(y)^{-1}}{2} = \frac{2 + |x|^{q'} + |y|^{q'}}{2}$. Using the linearity of $I_{n, \lambda}$ and taking the supremum over $\lambda \in \Lambda$, this shows that the Assumptions 3.1(ii)–(iii) and 4.5(i) are valid. Denoting by $c \geq 0$ the Lipschitz constant of $H^*|_{\Lambda}$, for every $n \in \mathbb{N}$, $\lambda_1, \lambda_2 \in \Lambda$ and $f \in C_{\kappa}^{\alpha} = \text{Lip}_b$, it holds

$$\begin{aligned} & \| (I_{n, \lambda_1} f - \eta(\lambda_1)) - ((I_{n, \lambda_2} f) - \eta(\lambda_2) h_n) \|_{\kappa} \\ & \leq \sup_{x \in \mathbb{R}^d} \mathbb{E}[|f(x + Z_{n, \lambda_1}) - f(x + Z_{n, \lambda_2})|] + |H^*(\lambda_1) - H^*(\lambda_2)| h_n \\ & \leq (\|f\|_{\alpha, \kappa} + c) |\lambda_1 - \lambda_2| h_n. \end{aligned}$$

Hence, Assumption 4.8 is valid with $L_r := (c + r) \sup_{n \in \mathbb{N}} h_n$, $d_r := |\cdot|$ and $\beta_r := 1$ for all $r \geq 0$.

C.2. Proof of Lemma 5.2. Regarding the verification of the Assumptions 2.2 and 4.1 as well as the extension $I_n : C_{\kappa_0} \rightarrow C_{\kappa_0}$, we refer to [8, Theorem 6.2], whereas Assumption 3.1(i) is clearly satisfied. Furthermore, by Lemma C.1, there exists a constant $\omega \geq 0$ with

$$\begin{aligned} & \|I_{n, (\mu, \sigma)} f\|_{\alpha', \kappa'} \\ & \leq \max \left\{ \sup_{x \in \mathbb{R}^d} \mathbb{E}[|f(x + Z_{n, (\mu, \sigma)})|] \kappa'(x), \sup_{x \neq y} \frac{\mathbb{E}[|f(x + Z_{n, (\mu, \sigma)}) - f(y + Z_{n, (\mu, \sigma)})|] \bar{\kappa}'(x, y)}{|x - y|^{\alpha'}} \right\} \\ & \leq \|f\|_{\alpha', \kappa'} \max \left\{ \sup_{x \in \mathbb{R}^d} \mathbb{E} \left[\frac{\kappa'(x)}{\kappa'(x + Z_{n, (\mu, \sigma)})} \right], \sup_{x \neq y} \mathbb{E} \left[\frac{\bar{\kappa}'(x, y)}{\bar{\kappa}'(x + Z_{n, (\mu, \sigma)}, y + Z_{n, (\mu, \sigma)})} \right] \right\} \\ & \leq e^{\omega h_n} \|f\|_{\alpha', \kappa'} \end{aligned}$$

for all $n \in \mathbb{N}$, $(\mu, \sigma) \in \Xi \times \Sigma$, $\alpha' \in [0, \alpha]$, $\kappa' \in \mathcal{K}$ and $f \in C_{\kappa'}^{\alpha'}$, where $Z_{n, (\mu, \sigma)} := \mu h_n + \sigma W_{h_n}$ and $(I_{n, (\mu, \sigma)} f)(x) := \mathbb{E}[f(x + Z_{n, (\mu, \sigma)})]$. Using the linearity of $I_{n, (\mu, \sigma)}$ and taking the supremum over $(\mu, \sigma) \in \Xi \times \Sigma$ yields that the Assumptions 3.1(ii)–(iii) and 4.5(i) are satisfied. Finally, for every $n \in \mathbb{N}$, $(\mu_1, \sigma_1), (\mu_2, \sigma_2) \in \Xi \times \Sigma$ and $f \in C_{\kappa}^{\alpha}$, it follows from $\frac{1 + |x + z|^q}{1 + |x|^q} \leq 2^{q-1}(1 + |z|^q)$, Lemma C.1, Hölder's inequality and Jensen's inequality that

$$\begin{aligned} & \|I_{n, (\mu_1, \sigma_1)} f - I_{n, (\mu_2, \sigma_2)} f\|_{\kappa} \leq \sup_{x \in \mathbb{R}^d} \mathbb{E}[|f(x + Z_{n, (\mu_1, \sigma_1)}) - f(x + Z_{n, (\mu_2, \sigma_2)})|] \kappa(x) \\ & \leq \sup_{x \in \mathbb{R}^d} \|f\|_{\alpha, \kappa} \mathbb{E} \left[\frac{|Z_{n, (\mu_1, \sigma_1)} - Z_{n, (\mu_2, \sigma_2)}|^{\alpha}}{\bar{\kappa}(x + Z_{n, (\mu_1, \sigma_1)}, x + Z_{n, (\mu_2, \sigma_2)})} \right] \kappa(x) \\ & \leq 2^{q-2} \|f\|_{\alpha, \kappa} \mathbb{E}[|Z_{n, (\mu_1, \sigma_1)} - Z_{n, (\mu_2, \sigma_2)}|^{\alpha} (2 + |Z_{n, (\mu_1, \sigma_1)}|^q + |Z_{n, (\mu_2, \sigma_2)}|^q)] \\ & \leq 2^{q-2} \|f\|_{\alpha, \kappa} \mathbb{E}[|Z_{n, (\mu_1, \sigma_1)} - Z_{n, (\mu_2, \sigma_2)}|^{2\alpha}]^{\frac{1}{2}} \mathbb{E}[(2 + |Z_{n, (\mu_1, \sigma_1)}|^q + |Z_{n, (\mu_2, \sigma_2)}|^q)^2]^{\frac{1}{2}} \\ & \leq 2^{q-2} \|f\|_{\alpha, \kappa} \mathbb{E}[|Z_{n, (\mu_1, \sigma_1)} - Z_{n, (\mu_2, \sigma_2)}|^2]^{\frac{\alpha}{2}} \mathbb{E}[(2 + |Z_{n, (\mu_1, \sigma_1)}|^q + |Z_{n, (\mu_2, \sigma_2)}|^q)^2]^{\frac{1}{2}} \\ & \leq C 2^{q-2} \|f\|_{\alpha, \kappa} d((\mu_1, \sigma_1), (\mu_2, \sigma_2))^{\alpha}, \end{aligned}$$

where $d((\mu_1, \sigma_1), (\mu_2, \sigma_2)) := \sqrt{|\mu_1 - \mu_2|^2 + |\sigma_1 - \sigma_2|^2}$ and

$$C := \sup_{n \in \mathbb{N}} \sup_{(\mu_1, \sigma_1), (\mu_2, \sigma_2) \in \Xi \times \Sigma} \mathbb{E}[(2 + |Z_{n, (\mu_1, \sigma_1)}|^q + |Z_{n, (\mu_2, \sigma_2)}|^q)^2]^{\frac{1}{2}} (h_n^2 + h_n)^{\frac{\alpha}{2}}.$$

Hence, Assumption 4.8 is valid with $L_r := 2^{q-2} C r$, $d_r := d$ and $\beta_r := \alpha$ for all $r \geq 0$.

C.3. Proof of Lemma 5.3. Regarding the verification of the Assumptions 4.1(i)–(vii), we refer to [25]. Since the function $\eta : \mathbb{R}_+ \rightarrow [0, \infty]$ is non-decreasing, it holds

$$I_n f = \sup_{(\nu, a) \in \Lambda} (I_{n, (\nu, a)} f - \eta_n(\nu, a) h_n)$$

for all $n \in \mathbb{N}$ and $f \in C_b$, where $\Lambda := \mathcal{P}_p(\mathbb{R}^d) \times \mathbb{R}_+$ and

$$\eta_n(\nu, a) := \begin{cases} \eta(a), & \text{if } \mathcal{W}_p(\varpi_{h_n}, \nu) \leq ah_n, \\ \infty, & \text{otherwise.} \end{cases}$$

Using $\alpha = 1$ and $\kappa \equiv 1$, it holds for every $n \in \mathbb{N}$, $(\nu, a) \in \Lambda$ and $f \in \text{Lip}_b$ that

$$\begin{aligned} \|I_{n,(\nu,a)}f\|_{\alpha,\kappa} &\leq \max \left\{ \sup_{x \in \mathbb{R}^d} \int_{\mathbb{R}^d} |f(x+z)| \nu(dz), \sup_{x \neq y} \frac{\int_{\mathbb{R}^d} |f(x+z) - f(y+z)| \nu(dz)}{|x-y|} \right\} \\ &\leq \max \left\{ \|f\|_\infty, \|f\|_{\alpha,\kappa} \sup_{x \neq y} \frac{|(x+z) - (y+z)|}{|x-y|} \right\} = \|f\|_{\alpha,\kappa}. \end{aligned}$$

This shows that Assumption 4.5(i) is valid. Moreover, for every $r \geq 0$, there exists $R_r \geq 0$ with

$$\begin{aligned} (I_n f)(x) &= \sup_{\{\nu: \mathcal{W}_p(\varpi_{h_n}, \nu) \leq R_r h_n\}} \left(\int_{\mathbb{R}^d} f(x+z) \nu(dz) - \eta \left(\frac{\mathcal{W}_p(\varpi_{h_n}, \nu)}{h_n} \right) h_n \right) \\ &= \sup_{(\nu,a) \in \Lambda_r} ((I_{n,(\nu,a)}f)(x) - \eta_n(\nu, a)h_n) \end{aligned}$$

for all $n \in \mathbb{N}$, $f \in \text{Lip}_b(r)$ and $x \in \mathbb{R}^d$, see [25, Lemma 3.3], where

$$\Lambda_r := \{\nu \in \mathcal{P}_p(\mathbb{R}^d): \mathcal{W}_p(\nu, \delta_0) \leq C_r\} \times [0, R_r] \quad \text{with} \quad C_r := \sup_{n \in \mathbb{N}} (R_r h_n + \mathcal{W}_p(\varpi_{h_n}, \delta_0)).$$

Equation (5.11) guarantees that $C_r < \infty$ and $p > 1$ implies that Λ_r is compact w.r.t. the metric

$$d((\nu_1, a_1), (\nu_2, a_2)) := \mathcal{W}_1(\nu_1, \nu_2) + |a_1 - a_2|.$$

We use the Kantorovich–Rubinstein inequality, that $\eta: [0, R_r] \cap \{\eta < \infty\} \rightarrow \mathbb{R}_+$ is Lipschitz continuous with some constant $c_r \geq 0$ and $\bar{h} := \sup_{n \in \mathbb{N}} h_n < \infty$ to estimate

$$\begin{aligned} &\| (I_{n,(\nu_1,a_1)}f - \eta_n(\nu_1, a_1)h_n) - (I_{n,(\nu_2,a_2)}f - \eta_n(\nu_2, a_2)h_n) \|_\infty \\ &\leq \sup_{x \in \mathbb{R}^d} \left| \int_{\mathbb{R}^d} f(x+z) \nu_1(dz) - \int_{\mathbb{R}^d} f(x+z) \nu_2(dz) \right| + |\eta(a_1) - \eta(a_2)|h_n \\ &\leq r\mathcal{W}_1(\nu_1, \nu_2) + \bar{h}c_r|a_1 - a_2| \\ &\leq \max\{r, \bar{h}c_r\}d((\nu_1, a_1), (\nu_2, a_2)) \end{aligned}$$

for all $n \in \mathbb{N}$, $r \geq 0$, $f \in \text{Lip}_b(r)$ and $(\nu_1, a_1), (\nu_2, a_2) \in \Lambda_{r,n} := \Lambda_r \cap \{\eta_n < \infty\}$. This shows that Assumption 4.8 is valid with $L_r := \max\{r, \bar{h}c_r\}$, $d_r := d$ and $\beta_r := 1$ for all $r \geq 0$.

C.4. Proof of Lemma 5.4. Regarding the verification of Assumption 4.1, we refer to [25], and the verification of Assumption 4.5(i) is the same as in the proof of Lemma 5.3. Furthermore, we use $\eta(0) = 0$ to obtain

$$\begin{aligned} &((J_{n,\lambda}f)(x) - \eta(|\lambda|)h_n) - ((J_{n,0}f)(x) - \eta(0)h_n) \\ &= \int_{\mathbb{R}^d} f(x + \lambda h_n + y) - f(x + y) \varpi_{h_n}(dy) - \eta(|\lambda|)h_n \leq (r|\lambda| - \eta(|\lambda|))h_n \end{aligned}$$

for all $n \in \mathbb{N}$, $r \geq 0$, $f \in \text{Lip}_b(r)$ and $\lambda, x \in \mathbb{R}^d$, where

$$(J_{n,\lambda}f)(x) := \int_{\mathbb{R}^d} f(x + \lambda h_n + y) \varpi_{h_n}(dy).$$

Since $\liminf_{v \rightarrow \infty} \frac{\eta(v)}{v^p} > 0$ with $p > 1$, there exists a totally bounded set $\Lambda_r \subset \mathbb{R}^d$ with

$$J_n f = \sup_{\lambda \in \Lambda_r} (J_{n,\lambda}f - \eta(|\lambda|)h_n)$$

for all $n \in \mathbb{N}$, $r \geq 0$ and $f \in \text{Lip}_b(r)$. We use that $\eta(|\cdot|): \Lambda_r \rightarrow \mathbb{R}_+$ is Lipschitz continuous with some constant $c_r \geq 0$ and $\bar{h} := \sup_{n \in \mathbb{N}} h_n < \infty$ to estimate

$$\begin{aligned} & \left\| \left((J_{n,\lambda_1} f) - \eta(|\lambda_1|)h_n \right) - \left(J_{n,\lambda_2} f - \eta(|\lambda_2|)h_n \right) \right\|_\infty \\ & \leq \sup_{x \in \mathbb{R}^d} \left| \int_{\mathbb{R}^d} f(x + \lambda_1 h_n + y) \varpi_{h_n}(dy) - \int_{\mathbb{R}^d} f(x + \lambda_2 h_n + y) \varpi_{h_n}(dy) \right| + |\eta(|\lambda_1|) - \eta(|\lambda_2|)|h_n \\ & \leq \bar{h}(r + c_r)|\lambda_1 - \lambda_2| \end{aligned}$$

for all $n \in \mathbb{N}$, $r \geq 0$, $f \in \text{Lip}_b(r)$ and $\lambda_1, \lambda_2 \in \Lambda_r$. This shows that Assumption 4.8 is valid with $L_r := \bar{h}(r + c_r)$, $d_r := |\cdot|$ and $\beta_r := 1$ for all $r \geq 0$.

Acknowledgments. P. Schmocker is partly supported by the FinsureTech Hub of ETH Zurich.

REFERENCES

- [1] G. Barles and E. R. Jakobsen. Error bounds for monotone approximation schemes for Hamilton-Jacobi-Bellman equations. *SIAM Journal on Numerical Analysis*, 43(2):540–558, 2005.
- [2] G. Barles and E. R. Jakobsen. Error bounds for monotone approximation schemes for parabolic Hamilton-Jacobi-Bellman equations. *Mathematics of Computation*, 76(260):1861–1893, 2007.
- [3] G. Barles and P. E. Souganidis. Convergence of approximation schemes for fully nonlinear second order equations. *Asymptotic Analysis*, 4:271–283, 1991.
- [4] D. Bartl, S. Drapeau, and L. Tangpi. Computational aspects of robust optimized certainty equivalents and option pricing. *Mathematical Finance*, 30(1):287–309, 2020.
- [5] D. Bartl, S. Eckstein, and M. Kupper. Limits of random walks with distributionally robust transition probabilities. *Electronic Communications in Probability*, 26:13, 2021. Id/No 28.
- [6] C. Beck, S. Becker, P. Cheridito, A. Jentzen, and A. Neufeld. Deep splitting method for parabolic PDEs. *SIAM Journal on Scientific Computing*, 43(5):A3135–A3154, 2021.
- [7] F. E. Benth, N. Detering, and L. Galimberti. Neural networks in Fréchet spaces. *Annals of Mathematics and Artificial Intelligence*, 91:75–103, 2023.
- [8] J. Blessing, R. Denk, M. Kupper, and M. Nendel. Convex monotone semigroups and their generators with respect to Γ -convergence. *Journal of Functional Analysis*, 288(8):110841, 2025.
- [9] J. Blessing, L. Jiang, M. Kupper, and G. Liang. Convergence rates for Chernoff-type approximations of convex monotone semigroups. *Stochastic Processes and their Applications*, 189:104700, 2025.
- [10] J. Blessing and M. Kupper. Nonlinear semigroups built on generating families and their Lipschitz sets. *Potential Analysis*, 59:857–895, 2023.
- [11] J. Blessing and M. Kupper. Nonlinear semigroups and limit theorems for convex expectations. *The Annals of Applied Probability*, 35(2):779–821, 2025.
- [12] J. Blessing, M. Kupper, and M. Nendel. Convergence of infinitesimal generators and stability of convex monotone semigroups. *forthcoming in SIAM Journal on Mathematical Analysis*, 2026.
- [13] J. Blessing, M. Kupper, and A. Sgarabottolo. Discrete approximation of risk-based prices under volatility uncertainty. *Preprint arXiv:2411.00713*, 2024.
- [14] L. A. Caffarelli and P. E. Souganidis. A rate of convergence for monotone finite difference approximations to fully nonlinear, uniformly elliptic PDEs. *Communications on Pure and Applied Mathematics*, 61(1):1–17, 2008.
- [15] W. Cai, S. Fang, and T. Zhou. SOC-MartNet: A martingale neural network for the Hamilton–Jacobi–Bellman equation without explicit $\inf_{u \in U} H$ in stochastic optimal controls. *SIAM Journal on Scientific Computing*, 47(4):C795–C819, 2025.
- [16] T. Chen and H. Chen. Approximation capability to functions of several variables, nonlinear functionals, and operators by radial basis function neural networks. *IEEE Transactions on Neural Networks*, 6(4):904–910, 1995.
- [17] M. G. Crandall, H. Ishii, and P.-L. Lions. User’s guide to viscosity solutions of second order partial differential equations. *American Mathematical Society. Bulletin. New Series*, 27(1):1–67, 1992.
- [18] M. G. Crandall and P.-L. Lions. Viscosity solutions of Hamilton-Jacobi equations. *Transactions of the American Mathematical Society*, 277(1):1–42, 1983.
- [19] C. Cuchiero, P. Schmocker, and J. Teichmann. Global universal approximation of functional input maps on weighted spaces. *Constructive Approximation*, 63:537–612, 2026.
- [20] F. Delbaen. Convex increasing functionals on $C_b(X)$ spaces. *Studia Mathematica*, 271(1):107–120, 2023.

- [21] R. Denk, M. Kupper, and M. Nendel. A semigroup approach to nonlinear Lévy processes. *Stochastic Processes and their Applications*, 130:1616–1642, 2020.
- [22] H. Dong and N. V. Krylov. Rate of convergence of finite-difference approximations for degenerate linear parabolic equations with C^1 and C^2 coefficients. *Electronic Journal of Differential Equations*, pages No. 102, 25, 2005.
- [23] W. E and B. Yu. The deep Ritz method: A deep learning-based numerical algorithm for solving variational problems. *Communications in Mathematics and Statistics*, 6:1–12, 2018.
- [24] D. H. Fremlin, D. J. H. Garling, and R. G. Haydon. Bounded measures on topological spaces. *Proceedings of the London Mathematical Society*, s3-25(1):115–136, 1972.
- [25] S. Fuhrmann, M. Kupper, and M. Nendel. Wasserstein perturbations of Markovian transition semigroups. *Annales de l’Institut Henri Poincaré. Probabilités et Statistiques*, 59(2):904–932, 2023.
- [26] L. Galimberti, A. Kratsios, and G. Livieri. Designing universal causal deep learning models: The case of infinite-dimensional dynamical systems from stochastic analysis. *forthcoming in Constructive Approximation*, 2026.
- [27] B. Goldys, M. Nendel, and M. Röckner. Operator semigroups in the mixed topology and the infinitesimal description of Markov processes. *Journal of Differential Equations*, 412:23–86, 2024.
- [28] L. Gonon, L. Grigoryeva, and J.-P. Ortega. Approximation bounds for random neural networks and reservoir systems. *The Annals of Applied Probability*, 33(1):28–69, 2023.
- [29] P. Grohs and L. Herrmann. Deep neural network approximation for high-dimensional parabolic Hamilton–Jacobi–Bellman equations. *Preprint arXiv:2103.05744*, 2021.
- [30] J. Han and W. E. Deep learning approximation for stochastic control problems. *Preprint arXiv:1611.07422*, 2016.
- [31] J. Han, A. Jentzen, and W. E. Solving high-dimensional partial differential equations using deep learning. *Proceedings of the National Academy of Sciences*, 115(34):8505–8510, 2018.
- [32] J. Hu and J.-P. Ortega. Kernel learning of PDE solution operators. *Preprint arXiv:2605.09643*, 2026.
- [33] C. Huré, H. Pham, and X. Warin. Deep backward schemes for high-dimensional nonlinear PDEs. *Mathematics of Computation*, 89(324):1547–1579, 2020.
- [34] L. Jiang. Discrete-time approximation for stochastic optimal control problems under the G -expectation framework. *Optimal Control Applications & Methods*, 43(2):418–434, 2022.
- [35] N. Kovachki, Z. Li, B. Liu, K. Azizzadenesheli, K. Bhattacharya, A. Stuart, and A. Anandkumar. Neural operator: Learning maps between function spaces with applications to PDEs. *Journal of Machine Learning Research*, 24(89):1–97, 2023.
- [36] A. Kratsios and I. Bilokopytov. Non-Euclidean universal approximation. *Advances in Neural Information Processing Systems*, 33:10635–10646, 2020.
- [37] A. Kratsios, C. Liu, M. Lassas, M. V. de Hoop, and I. Dokmanić. An approximation theory for metric space-valued functions with a view towards deep learning. *Preprint arXiv:2304.12231*, 2023.
- [38] A. Kratsios, A. Neufeld, and P. Schmock. Generative neural operators of log-complexity can simultaneously solve infinitely many convex programs. *Preprint arXiv:2508.14995*, 2025.
- [39] N. V. Krylov. On the rate of convergence of finite-difference approximations for Bellman’s equations with variable coefficients. *Probability Theory and Related Fields*, 117(1):1–16, 2000.
- [40] M. Kunze. Continuity and equicontinuity of semigroups on norming dual pairs. *Semigroup Forum*, 79(3):540–560, 2009.
- [41] M. Kupper, M. Nendel, and A. Sgarabottolo. Risk measures based on weak optimal transport. *Quantitative Finance*, 25(2):163–180, 2025.
- [42] M. Kupper, M. Nendel, and A. Sgarabottolo. Hopf-Lax approximation for value functions of Lévy optimal control problems. *Proceedings of the American Mathematical Society*, 154(1):243–254, 2026.
- [43] S. Lanthaler, S. Mishra, and G. E. Karniadakis. Error estimates for DeepONets: a deep learning framework in infinite dimensions. *Transactions of Mathematics and Its Applications*, 6(1), 2022.
- [44] S. Lanthaler and A. M. Stuart. The parametric complexity of operator learning. *IMA Journal of Numerical Analysis*, 46(2):647–712, 2026.
- [45] M. Leshno, Vladimir Ya. Lin, A. Pinkus, and S. Schocken. Multilayer feedforward networks with a nonpolynomial activation function can approximate any function. *Neural Networks*, 6(6):861–867, 1993.
- [46] Z. Li, N. Kovachki, K. Azizzadenesheli, B. Liu, K. Bhattacharya, A. Stuart, and A. Anandkumar. Fourier neural operator for parametric partial differential equations. *Preprint arXiv:2010.08895*, 2020.
- [47] P.-L. Lions. *Generalized solutions of Hamilton–Jacobi equations*, volume 69 of *Research Notes in Mathematics*. Pitman (Advanced Publishing Program), Boston, Mass.-London, 1982.

- [48] L. Lu, P. Jin, G. Pang, Z. Zhang, and G. E. Karniadakis. Learning nonlinear operators via DeepONet based on the universal approximation theorem of operators. *Nature Machine Intelligence*, 3:218–229, 2021.
- [49] A. Lunardi. *Analytic semigroups and optimal regularity in parabolic problems*. Progress in nonlinear differential equations and their applications, vol. 16. Birkhäuser, Boston, 1995.
- [50] H. N. Mhaskar and N. Hahm. Neural networks for functional approximation and system identification. *Neural Computation*, 9:143–159, 1997.
- [51] T. Nakamura-Zimmerer, Q. Gong, and W. Kang. Adaptive deep learning for high-dimensional Hamilton–Jacobi–Bellman equations. *SIAM Journal on Scientific Computing*, 43(2):A1221–A1247, 2021.
- [52] N. H. Nelsen and A. M. Stuart. The random feature model for input-output maps between Banach spaces. *SIAM Journal on Scientific Computing*, 43(5):A3212–A3243, 2021.
- [53] M. Nendel. Lower semicontinuity of monotone functionals in the mixed topology on C_b . *Finance and Stochastics*, 29:261–287, 2025.
- [54] M. Nendel and M. Röckner. Upper envelopes of families of Feller semigroups and viscosity solutions to a class of nonlinear Cauchy problems. *SIAM Journal on Control and Optimization*, 59(6):4400–4428, 2021.
- [55] M. Nendel and A. Sgarabottolo. A parametric approach to the estimation of convex risk functionals based on Wasserstein distance. *Applied Mathematics and Optimization*, 93(8):44, 2026.
- [56] A. Neufeld and P. Schmock. Universal approximation property of Banach space-valued random feature models including random neural networks. *forthcoming in The Annals of Applied Probability*, 2026.
- [57] A. Neufeld and P. Schmock. Universal approximation results for neural networks with non-polynomial activation function over non-compact domains. *Analysis and Applications*, 24(05):1123–1173, 2026.
- [58] A. Neufeld, P. Schmock, and S. Wu. Full error analysis of the random deep splitting method for nonlinear parabolic PDEs and PIDEs. *Communications in Nonlinear Science and Numerical Simulation*, 143:108556, 2025.
- [59] M. Nisio. On a non-linear semi-group attached to stochastic optimal control. *Kyoto University. Research Institute for Mathematical Sciences. Publications*, 12(2):513–537, 1976/77.
- [60] N. Nüsken and L. Richter. Solving high-dimensional Hamilton–Jacobi–Bellman PDEs using neural networks: perspectives from the theory of controlled diffusions and measures on path space. *Partial Differential Equations and Applications*, 2:48, 2021.
- [61] P. Petersen and J. Zech. Mathematical theory of deep learning. *Preprint arXiv:2407.18384*, 2024.
- [62] M. Raissi, P. Perdikaris, and G. Karniadakis. Physics-informed neural networks: A deep learning framework for solving forward and inverse problems involving nonlinear partial differential equations. *Journal of Computational Physics*, 378:686–707, 2019.
- [63] P. Schmock and J. Teichmann. Weighted universal approximation of differentiable maps on infinite-dimensional manifolds. *Preprint arXiv:2606.09820*, 2026.
- [64] F. D. Sentilles. Bounded continuous functions on a completely regular space. *Transactions of the American Mathematical Society*, 168:311–336, 1972.
- [65] J. Sirignano and K. Spiliopoulos. DGM: A deep learning algorithm for solving partial differential equations. *Journal of Computational Physics*, 375:1339–1364, 2018.
- [66] M. H. Stone. The generalized Weierstrass approximation theorem. *Mathematics Magazine*, 21(4):167–184, 1948.
- [67] K. Weierstrass. Über die analytische Darstellbarkeit sogenannter willkürlicher Functionen einer reellen Veränderlichen. *Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin*, II, 1885.
- [68] A. Wiweger. Linear spaces with mixed topology. *Polska Akademia Nauk. Instytut Matematyczny. Studia Mathematica*, 20:47–68, 1961.
- [69] M. Zhou, J. Han, and J. Lu. Actor-critic method for high dimensional static Hamilton–Jacobi–Bellman partial differential equations based on neural networks. *SIAM Journal on Scientific Computing*, 43(6):A4043–A4066, 2021.

DEPARTMENT OF MATHEMATICS, ETH ZURICH, SWITZERLAND
 Email address: jonas.blessing@math.ethz.ch

DEPARTMENT OF MATHEMATICS, ETH ZURICH, SWITZERLAND
 Email address: philipp.schmock@math.ethz.ch

DEPARTMENT OF MATHEMATICS, LMU MUNICH, GERMANY
 Email address: sgarabottolo@math.lmu.de