

GRADED REPRESENTATION THEORY OF EQUIVARIANT NEURAL NETWORKS

MANI SHAYESTEHFAR

ABSTRACT. Nonlinear activations can create equivariant interactions between irreducible representations that linear maps cannot. We use the Gaussian degree decomposition to extend ordinary polynomial degree to such nonlinear maps, and prove that for a fixed coordinatewise equivariant layer each degree factors into a polynomial determined by the linear maps and a scalar determined by the activation. This separates three distinct obstructions, coming from symmetry, coordinates, and activation.

CONTENTS

Introduction	1
1. Preliminaries	5
2. Degree-resolved equivariant maps	7
3. Coordinatewise nonlinearities	11
4. Activation spectra	17
5. Architectures determined by the degree calculation	20
6. Subset representations of the symmetric group	23
7. Scope and future directions	28
Appendix A. Supplementary proofs	29
Appendix B. Hidden spaces with several coordinate orbits	30
Appendix C. Representation formulae for subset modules	31
References	33

INTRODUCTION

Neural networks alternate linear maps with nonlinear activation functions. When the data carry symmetries, one often asks the network to respect them: rotating, translating or relabelling an input should have a prescribed effect on the output. Mathematically, the feature spaces are then representations of a group G , and the linear parts of the network are G -equivariant maps. This is precisely the viewpoint promoted by geometric deep learning [BBCV21].

Representation theory is very effective for understanding the linear part. A finite-dimensional representation can be decomposed into irreducible pieces, and an equivariant linear map cannot mix two non-isomorphic irreducibles. In a suitable basis, this makes the linear maps much simpler.

The activation function changes the picture. It is nonlinear and is usually applied coordinatewise, so it can create interactions between irreducible pieces which are completely invisible to linear representation theory. The simplest example already occurs for the symmetric group S_3 .

2020 *Mathematics Subject Classification.* Primary: 20C15, 68T07; Secondary: 05E10, 13A50.

Key words and phrases. Equivariant neural networks, representation theory, polynomial chaos, symmetric powers, permutation representations, piecewise-linear maps.

Let S_3 permute the coordinates of

$$\mathbb{R}^3 = L_0 \oplus L_1, \quad L_0 = \text{span}\{(1, 1, 1)\}, \quad L_1 = \{x : x_1 + x_2 + x_3 = 0\}.$$

The first summand is the trivial representation and the second is the two-dimensional standard representation. There is no nonzero equivariant linear map from L_1 to L_0 . Coordinatewise squaring, however, gives

$$p_0(x_1^2, x_2^2, x_3^2) = \frac{\|x\|^2}{3}(1, 1, 1), \quad x \in L_1, \quad (1)$$

where p_0 is orthogonal projection onto L_0 . Thus a coordinatewise nonlinearity creates a quadratic interaction between two representation types which no linear equivariant map can connect.

This is the starting point of the paper. We ask not merely whether a nonlinear equivariant interaction exists, but at which degrees it occurs and whether a given neural network layer actually produces it.

From existence to degree. Gibson, Tubbenhauer and Williamson [GTW25] (GTW) study the first, degree-free version of this question. For irreducible real representations L and K of a finite group, they show that a nonzero equivariant piecewise-linear map $L \rightarrow K$ exists precisely when

$$\ker L \subseteq \ker K. \quad (2)$$

Here piecewise linear means continuous and affine on each of finitely many polyhedral pieces. This is already very different from the linear theory: nonlinear equivariant maps can connect many different irreducible representations.

Condition (2), however, does not say how the interaction occurs. For example, the interaction in (1) first appears in degree two. More generally, one would like to distinguish such interactions by the degree in which they occur. One would also like to distinguish two further questions: symmetry may permit a degree without the chosen coordinate system producing it, and the coordinate system may produce it while the chosen activation suppresses it.

For polynomial maps there is an obvious notion of degree: the degree- k equivariant maps $L \rightarrow K$ are

$$\text{Hom}_G(\text{Sym}^k L, K).$$

Activations such as ReLU are not polynomial, so we need a way to extend this notion of degree. We put the standard Gaussian measure on L . Just as monomials organise ordinary polynomials by degree, Hermite polynomials form an orthogonal basis for functions with respect to the Gaussian measure. Grouping the multivariate Hermite polynomials by their total degree gives an orthogonal decomposition, called the *polynomial chaos* decomposition.

Thus polynomial chaos extends the usual polynomial grading to non-polynomial activations such as ReLU. We write Π_k for the orthogonal projection onto the degree- k part.

This grading is classical. Its usefulness here is that it gives a common language for polynomial and non-polynomial activations. A ReLU layer, for example, has infinitely many Hermite components, and we can ask exactly which representation-theoretic interactions occur in each one.

There is already a useful answer before considering a particular architecture. If $L \neq 0$, then an allowed degree can never occur in isolation:

$$\text{Hom}_G(\text{Sym}^k L, K) \neq 0 \implies \text{Hom}_G(\text{Sym}^{k+2m} L, K) \neq 0 \quad (m \geq 0).$$

Thus the possible degrees consist of at most two tails, one even and one odd. For finite groups we also refine (2): writing $N = \ker L$, some polynomial degree occurs if and only if $K^N \neq 0$ (where K^N denotes the N -fixed subspace of K), and the least such degree

is at most $|G/N| - 1$. For irreducible K , this recovers the GTW kernel condition and additionally locates the interaction in degree.

What does one layer produce? Knowing that a degree- k equivariant map $L \rightarrow K$ exists does not mean that a given neural network layer produces one. We now make this distinction precise.

Let $H = \mathbb{R}^{\mathcal{B}}$ be a representation with a basis $\mathcal{B}$ which is permuted by G . A typical equivariant neural network layer has the form

$$L \xrightarrow{A} H \xrightarrow{\sigma_{\mathcal{B}}} H \xrightarrow{R} K.$$

The first map A expresses the input in the hidden coordinates $\mathcal{B}$, and the last map R sends the hidden features to the output space K . Between them we apply the same scalar activation $\sigma : \mathbb{R} \rightarrow \mathbb{R}$ separately to each hidden coordinate: if

$$h = \sum_{b \in \mathcal{B}} h_b b, \quad \text{then} \quad \sigma_{\mathcal{B}}(h) = \sum_{b \in \mathcal{B}} \sigma(h_b) b.$$

Since G only permutes the basis vectors in $\mathcal{B}$, applying the same function to every coordinate commutes with the G -action. Hence $\sigma_{\mathcal{B}}$ is equivariant, and so is the whole layer whenever A and R are equivariant linear maps.

There are two separate reasons why a degree permitted by representation theory may fail to appear in this layer:

- (a) the linear maps A and R may not produce the relevant polynomial interaction, or
- (b) the activation σ may have no component in that degree.

The main calculation separates these two effects. Let s^2 be the common variance of the coordinates of AX when X is a standard Gaussian vector in L . Let $a_k(\sigma; s)$ be the coefficient of the k th Hermite polynomial in the expansion of the one-variable function $z \mapsto \sigma(sz)$. Finally, for an ordinary input $x \in L$, set

$$Q_k(x) = R((Ax)^{\odot k}),$$

where $(Ax)^{\odot k}$ is the coordinatewise k th power. Then

$$\Pi_k[R\sigma_{\mathcal{B}}(Ax)] = \frac{a_k(\sigma; s)}{s^k} \Pi_k Q_k. \quad (3)$$

This gives three separate questions:

- (a) Does symmetry allow a degree- k equivariant map $L \rightarrow K$?
- (b) Do the chosen linear maps A and R actually produce one?
- (c) Does the activation σ retain degree k ?

The first asks whether $\text{Hom}_G(\text{Sym}^k L, K) \neq 0$. The second asks whether Q_k is nonzero, and the third whether $a_k(\sigma; s)$ is nonzero.

The second question also has an explicit answer. Put $C = AA^*$, written in the coordinate basis of $\mathbb{R}^{\mathcal{B}}$ (with G acting transitively). Then

$$\|\Pi_k Q_k\|_{L^2}^2 = k! \text{tr}(RC^{\odot k} R^*), \quad (4)$$

where $C^{\odot k}$ is obtained by taking the k th power of each entry of C . Hence

$$Q_k \neq 0 \iff \text{tr}(RC^{\odot k} R^*) > 0.$$

For S_4 we give an example where a quadratic equivariant map exists, but the chosen coordinatewise layer produces no quadratic term at all, and changing the activation cannot fix this. For C_8 the opposite phenomenon occurs: the layer does produce the first permitted polynomial interaction, but a positively homogeneous activation such as ReLU removes it. Adding a bias restores it.

Thus a nonlinear interaction can fail for three different reasons: it may be forbidden by symmetry, missed by the chosen linear maps, or removed by the activation. Distinguishing these three mechanisms is the main point of the paper.

Consequences for architectures. The same calculation answers two practical questions. For any fixed collection of layers, an explicit Gram matrix determines exactly which degree- k equivariant maps they can produce. Moreover, shifted ReLUs can realise arbitrary Hermite coefficients up to any prescribed finite degree. Using the regular representation then gives all equivariant polynomial directions, although not efficiently.

For ordinary ReLU there is a simple restriction: all odd Hermite degrees above one vanish. A shared bias removes this restriction generically. These degree-resolved interactions can also be used to design representation-aware architectures. In a companion paper by the author, the resulting source-to-target decomposition is used to build a higher-order graph network on unordered triples, and experiments show an improvement over both a standard GNN baseline and a parameter-matched control with the full linear S_n -equivariant mixing space [Sha26]. This addresses a question left open in [GTW25]: whether organising nonlinear equivariant networks by irreducible representation theory is useful in practice.

Subset features. The final part of the paper treats the action of S_n on r -element subsets. A degree- d monomial in these variables corresponds to an r -uniform multihypergraph with d edges. If the output is indexed by s -subsets, one simply marks an s -subset of vertices. Equivariant polynomial maps are therefore indexed by these marked multihypergraphs, up to relabelling.

This gives an explicit basis whose types stabilise at the sharp bound

$$n = rd + s.$$

For quadratic edge-to-vertex maps there are seven stable types, and all seven can be evaluated in $O(n^2)$ operations, or $O(n + m)$ for m nonzero edges.

Relation to existing work and scope. Our starting point is the piecewise-linear representation theory of [GTW25, Theorem 2H.3]. For irreducible real representations of a finite group, they characterise the existence of a nonzero equivariant piecewise-linear map by inclusion of the representation kernels. We refine this existence question by recording the polynomial degrees in which an interaction can occur. We then ask which of these interactions are produced by a specified coordinatewise layer and activation.

The compact-group structure theorem of [Aie26, Theorem 2.5 and Remark 2.6] gives a broader context for this finite-group analysis. Let Γ be a compact group with identity component Γ° , and equip its finite-dimensional real representations L and K with invariant inner products. Every equivariant piecewise-linear map $F : L \rightarrow K$ has the form

$$F(v + w) = \psi(v) + Tw, \quad v \in L^{\Gamma^\circ}, \quad w \in (L^{\Gamma^\circ})^\perp,$$

where $\psi : L^{\Gamma^\circ} \rightarrow K^{\Gamma^\circ}$ is equivariant and piecewise linear, and $T : (L^{\Gamma^\circ})^\perp \rightarrow (K^{\Gamma^\circ})^\perp$ is an equivariant linear map. The joint action on the two fixed subspaces has finite image. Consequently, our finite-group degree criteria apply to this remaining nonlinear problem, and our layer criteria can test chosen coordinatewise constructions for ψ . For connected groups, nonlinear dependence is confined to the trivial subrepresentations. This restriction also explains why polynomial equivariants alone cannot characterise piecewise-linear existence for general compact groups.

At the level of layer construction, [PDL24] classify the combinations of representations, coordinates and pointwise activations that give an equivariant activation map. We work with hidden permutation representations, where shared coordinatewise activations are equivariant, and analyse the interactions selected by the linear lift, activation and readout.

The analytic tools are classical. Gaussian chaos, Wick ordering and their interpretation through symmetric tensors are developed in [Jan97, Chapters 2–4]. Hermite expansions

and the associated covariance identities also underlie the dual-activation framework of [DFS16]. We use these identities to separate the polynomial selected by an equivariant layer from the scalar coefficient supplied by its activation. This gives explicit Gram criteria for the vanishing of a degree component and for the degree subspace spanned by a specified family of layers.

Our spanning construction also connects with equivariant approximation theory. [Rav20] establishes universality using regular hidden actions, while [Yar22] obtains invariant and equivariant approximation through group averaging. Here regular representations provide an exact spanning construction in each polynomial degree. Matching finitely many Hermite components does not, by itself, control the remaining approximation error.

For subset representations, the marked-hypergraph basis follows the orbit-sum principle used by [PLK⁺23] to construct multigraph-indexed bases of equivariant graph polynomials. We formulate this construction for r -subset inputs and s -subset outputs, and determine its sharp stable range. The distinction between parameter sharing across dimensions and compatibility with embeddings already appears in [LD24]; our marked-orbit criterion makes that compatibility explicit for these polynomial maps.

The Gaussian measure here is only a reference measure for the grading, not an assumption about trained features, and the resulting one-layer obstructions need not persist under arbitrary composition.

Organisation. Sections 1 and 2 introduce polynomial and Gaussian degree and prove the finite-group degree-detection results. Section 3 studies coordinatewise layers and proves (3) and (4). Section 4 determines the relevant activation spectra, and Section 5 gives the resulting architecture-level spanning statements. Section 6 treats polynomial layers on subset features. Supplementary proofs and representation formulae are collected in the appendices.

Acknowledgement. This research was completed during a master’s project supervised by Dani Tubbenhauer, whom I thank for many discussions and for guidance throughout.

Use of AI. The initial backbone of the work was developed independently. Discussions with Claude Fable pointed towards sources such as [Jan97], where the connection between $L^2(\gamma_L)$ and $\text{Sym}^k L$ was found. ChatGPT Sol assisted with polishing arguments and proofs, and with the literature review. After using each tool, the author reviewed and edited the content as needed, and takes full responsibility for the content of the work.

1. PRELIMINARIES

All representations are finite-dimensional and real unless otherwise stated, and are equipped with fixed invariant inner products. The acting group G is finite unless compact groups are explicitly allowed. For a representation L , we write $\ker L$ for the kernel of the action; it is a normal subgroup of G .

1.1. Polynomial maps and symmetric powers. Polynomial degree is the basic grading used throughout the paper. We begin by recalling the standard identification between homogeneous polynomial maps and maps out of symmetric powers.

For a real representation L and $k \geq 1$, define

$$\text{Sym}^k L = L^{\otimes k} / \text{span}\{t - \pi t : t \in L^{\otimes k}, \pi \in S_k\}, \quad \text{Sym}^0 L = \mathbb{R},$$

where S_k acts by permuting tensor factors. The diagonal action of G on $L^{\otimes k}$ commutes with this action, and therefore descends to $\text{Sym}^k L$.

For $x \in L$, we write x^k for the class of $x^{\otimes k}$ in $\text{Sym}^k L$. Pure powers span $\text{Sym}^k L$, and the induced inner product is characterised by

$$\langle x^k, y^k \rangle = \langle x, y \rangle^k.$$

To avoid confusion, coordinatewise powers in a based permutation representation will instead be denoted by $x^{\odot k}$.

For real representations L and K , let $\text{Poly}_k(L, K)$ denote the space of homogeneous polynomial maps $P : L \rightarrow K$ of degree k . The G -action is $(g \cdot P)(x) = gP(g^{-1}x)$, so that

$$\text{Poly}_k(L, K)^G = \{P \in \text{Poly}_k(L, K) : P(gx) = gP(x) \text{ for all } g \in G, x \in L\}.$$

Similarly, G acts on $\text{Hom}(\text{Sym}^k L, K)$ by $(g \cdot T)(y) = gT(g^{-1}y)$.

For a finite index set I , multi-indices are written $\alpha \in \mathbb{N}_0^I$, with $|\alpha| = \sum_{i \in I} \alpha_i$.

Proposition 1.1. *For every $k \geq 0$,*

$$\text{Poly}_k(L, K)^G \cong \text{Hom}_G(\text{Sym}^k L, K).$$

See Section A.1 for a proof. The isomorphism sends T to $x \mapsto T(x^k)$ and follows from polarisation. We use this identification throughout to regard homogeneous degree- k equivariant polynomial maps as elements of $\text{Hom}_G(\text{Sym}^k L, K)$.

1.2. Piecewise-linear maps. Our nonlinear layers are piecewise linear, while the degree decomposition below lives in a Gaussian L^2 space. We first fix the class of maps to which both viewpoints apply. A map $F : L \rightarrow K$ is *piecewise linear* if it is continuous and there is a finite covering of L by polyhedra on each of which F is affine. We write $\text{Hom}_G^{\text{PL}}(L, K)$ for the equivariant ones. We follow the affine-on-pieces convention of [GTW25], which includes constant maps. In particular, applying a continuous piecewise-linear scalar function with finitely many knots coordinatewise again gives a piecewise-linear map.

1.3. Gaussian spaces and polynomial chaos. Let L be an orthogonal representation of dimension m and let γ_L denote the standard Gaussian measure on L . This measure is invariant under $O(L)$, hence under G , and has moments of every order. Its Hermite expansion provides the orthogonal grading used below.

For a representation K , let $L^2(\gamma_L; K)$ be the Hilbert space of square-integrable maps $L \rightarrow K$ modulo null sets, with G acting unitarily by $(g \cdot F)(x) = gF(g^{-1}x)$.

We use the probabilists' Hermite polynomials He_k , defined by the generating function $\exp(tx - t^2/2) = \sum_{k \geq 0} \text{He}_k(x)t^k/k!$ and normalised so that $\mathbb{E}[\text{He}_j(Z)\text{He}_k(Z)] = k! \delta_{jk}$ for $Z \sim N(0, 1)$.

In m variables, the products $\prod_i \text{He}_{k_i}(x_i)$ with $\sum_i k_i = k$ span the k th *polynomial chaos* $\mathcal{H}_k(L)$, and

$$L^2(\gamma_L) = \widehat{\bigoplus_{k \geq 0} \mathcal{H}_k(L)}. \quad (5)$$

Thus the chaos index is a polynomial degree after orthogonal removal of lower-degree terms. For example, $\text{He}_2(x) = x^2 - 1$ lies entirely in the second chaos. We write Π_k for the orthogonal projection onto $\mathcal{H}_k(L)$ and use the same symbol for the induced projection on K -valued maps:

$$\Pi_k : L^2(\gamma_L; K) \longrightarrow \mathcal{H}_k(L) \otimes K.$$

The chaos projection Π_k restricts to an $O(L)$ -equivariant isomorphism from homogeneous degree- k polynomials onto $\mathcal{H}_k(L)$, and hence realises the identification

$$\mathcal{H}_k(L) \cong \text{Sym}^k(L^*) \cong \text{Sym}^k L,$$

using the fixed invariant inner product (see [Jan97]). For a linear form $\ell(x) = \langle x, a \rangle$ with $\|a\| = s > 0$, $\Pi_k(\ell^k) = s^k \text{He}_k(\ell/s)$.

If P is homogeneous of degree k , then $P - \Pi_k P$ has degree strictly less than k . The space $\mathcal{H}_k(L)$ therefore consists of degree-at-most- k polynomials orthogonal to all lower-degree polynomials, rather than homogeneous polynomials themselves.

The following identity will be used in the factorisation and norm calculations.

Lemma 1.2 (Mehler). *Let (X, Y) be jointly standard normal with correlation ρ . Then*

$$\mathbb{E}[\text{He}_a(X)\text{He}_b(Y)] = a! \rho^a \delta_{ab}.$$

Proof. This is a standard identity. See [Jan97, Theorem 3.9]. $\square$

To use this Hilbert-space decomposition for the piecewise-linear maps above, we only need to check square-integrability.

Proposition 1.3. *Let L and K be finite-dimensional. If $F : L \rightarrow K$ is piecewise linear with finitely many polyhedral pieces, then $F \in L^2(\gamma_L; K)$.*

Proof. Since F is affine on finitely many polyhedral pieces, there exist constants $C, D \geq 0$ such that $\|F(x)\| \leq C\|x\| + D$ for all $x \in L$. Hence $\|F(x)\|^2 \leq 2C^2\|x\|^2 + 2D^2$. If $X \sim \gamma_L$, then $\mathbb{E}\|X\|^2 < \infty$, so

$$\int_L \|F(x)\|^2 d\gamma_L(x) \leq 2C^2 \mathbb{E}\|X\|^2 + 2D^2 < \infty.$$

Thus $F \in L^2(\gamma_L; K)$. $\square$

Two continuous maps agreeing γ_L -almost everywhere agree everywhere, because γ_L has full support and the set where they differ is open. Consequently, the chaos components determine a continuous map uniquely.

2. DEGREE-RESOLVED EQUIVARIANT MAPS

We first identify the equivariant part of each polynomial chaos. We then extend the grading to several source summands and relate occurrence at some degree to the finite-group existence criterion.

2.1. The equivariant decomposition. The Gaussian grading is useful here only if it respects equivariance. The next theorem says that it does so, degree by degree.

Theorem 2.1. *Let G be a finite or compact group acting orthogonally on finite-dimensional real spaces L and K . There is a canonical isomorphism of Hilbert spaces*

$$L^2(\gamma_L; K)^G \cong \widehat{\bigoplus_{k \geq 0} \text{Hom}_G(\text{Sym}^k L, K)}.$$

Proof. Tensoring (5) with the finite-dimensional space K gives $L^2(\gamma_L; K) = \widehat{\bigoplus_k (\mathcal{H}_k(L) \otimes K)}$. Each chaos projection is $O(L)$ -equivariant. Since $G \subseteq O(L)$, it is also G -equivariant and hence commutes with the projection onto G -fixed vectors. Therefore taking fixed points commutes with the orthogonal sum. Finally

$$(\mathcal{H}_k(L) \otimes K)^G \cong (\text{Sym}^k(L^*) \otimes K)^G \cong \text{Hom}_G(\text{Sym}^k L, K)$$

using the $O(L)$ -equivariant identification $\mathcal{H}_k(L) \cong \text{Sym}^k L$ above and the tensor–Hom identification. $\square$

The key point is that each chaos projection is equivariant under the full orthogonal group. It therefore preserves the G -fixed subspace, allowing the Gaussian grading to restrict to equivariant maps.

Corollary 2.2. *Every equivariant piecewise-linear $F : L \rightarrow K$ has a unique sequence $(\Pi_k F)_{k \geq 0}$ with $\Pi_k F \in \text{Hom}_G(\text{Sym}^k L, K)$, and $F \mapsto (\Pi_k F)_k$ is injective on continuous square-integrable equivariant maps.*

Proof. Square-integrability follows from Proposition 1.3. Orthogonality gives uniqueness in L^2 , and full support of the Gaussian measure gives pointwise uniqueness for continuous maps. $\square$

Corollary 2.2 establishes injectivity but does not characterise the image of the transform. In particular, not every square-summable sequence of equivariant polynomial components arises from a piecewise-linear map. The finite-group existence questions considered here do not require such a characterisation. The distinction becomes essential for compact connected groups, as discussed in Section 7.

2.2. Multi-sources and multidegrees. Fix a finite orthogonal G -stable decomposition $V = \bigoplus_{i \in I} W_i$ into nonzero subrepresentations. The summands need not be irreducible or pairwise non-isomorphic. The multidegree refinement recorded below separates the contributions of the summands W_i within a single total degree. This allows the coordinatewise results of Section 3 to be applied to a direct-sum input without treating its summands separately.

Multi-source inputs are not used in our constructions; the following decompositions indicate how the theory extends to them.

Symmetric powers of a direct sum split by multidegree. Iterating $\text{Sym}(U \oplus W) \cong \text{Sym} U \otimes \text{Sym} W$ gives the G -equivariant isomorphism

$$\text{Sym}^k V \cong \bigoplus_{|\alpha|=k} \bigotimes_{i \in I} \text{Sym}^{\alpha_i} W_i.$$

Hence for any K ,

$$\text{Hom}_G(\text{Sym}^k V, K) \cong \bigoplus_{|\alpha|=k} \text{Hom}_G\left(\bigotimes_i \text{Sym}^{\alpha_i} W_i, K\right).$$

Example 2.3. For $V = W_1 \oplus W_2$,

$$\text{Sym}^2 V \cong \text{Sym}^2 W_1 \oplus (W_1 \otimes W_2) \oplus \text{Sym}^2 W_2,$$

$$\text{Sym}^3 V \cong \text{Sym}^3 W_1 \oplus (\text{Sym}^2 W_1 \otimes W_2) \oplus (W_1 \otimes \text{Sym}^2 W_2) \oplus \text{Sym}^3 W_2.$$

The mixed summands have no counterpart in a GTW interaction graph whose edges have a single source. Instead, they record simultaneous nonlinear dependence on two summands.

On the Gaussian side, orthogonality of the W_i makes the components of $x \sim \gamma_V$ independent, so $\gamma_V = \bigotimes_i \gamma_{W_i}$, and the Hilbert tensor product gives the *multichaos* decomposition

$$L^2(\gamma_V) \cong \widehat{\bigoplus_{\alpha \in \mathbb{N}_0^I} \bigotimes_i \mathcal{H}_{\alpha_i}(W_i)},$$

whose total-degree- k part is the sum over $|\alpha| = k$. Write Π_α for the projection onto the α th multichaos. Tensoring with K and taking fixed points degreewise, as in Theorem 2.1, gives

$$L^2(\gamma_V; K)^G \cong \widehat{\bigoplus_{\alpha \in \mathbb{N}_0^I} \text{Hom}_G\left(\bigotimes_i \text{Sym}^{\alpha_i} W_i, K\right)}.$$

2.3. Degree sets and detection. We now forget multiplicities and record only the degrees in which the target can occur. This is the information needed for the existence and parity questions below.

Definition 2.4. For representations L, K define the *degree set*

$$D(L, K) := \{k \geq 0 : \text{Hom}_G(\text{Sym}^k L, K) \neq 0\},$$

and $d(L, K) := \min D(L, K) \in \mathbb{N}_0 \cup \{\infty\}$. We use $\min \emptyset = \infty$.

If $K^G \neq 0$, then degree zero records constant equivariant maps. When the question is transfer of input information, the first positive degree may therefore be the more relevant quantity.

Proposition 2.5. *Let $L \neq 0$, and let G be finite or compact. If $k \in D(L, K)$, then $k + 2m \in D(L, K)$ for every $m \in \mathbb{N}_0$. Consequently, for each $\varepsilon \in \{0, 1\}$ define*

$$d_\varepsilon(L, K) := \min\{k \in D(L, K) : k \equiv \varepsilon \pmod{2}\}.$$

Then

$$D(L, K) = (d_0(L, K) + 2\mathbb{N}_0) \cup (d_1(L, K) + 2\mathbb{N}_0),$$

where a term is omitted if $d_\varepsilon(L, K)$ is infinite.

Proof. Since G is finite or compact, L admits a G -invariant inner product, so $q(x) = \|x\|^2$ is a nonzero invariant quadratic polynomial. Hence multiplication by q sends

$$\text{Poly}_k(L, K)^G \longrightarrow \text{Poly}_{k+2}(L, K)^G.$$

This map is injective because the polynomial ring on L is an integral domain. Hence $k \in D(L, K) \implies k + 2 \in D(L, K)$. Thus, within each parity, occurrence persists from the first occurring degree onward, giving the stated $D(L, K)$, with the corresponding term omitted when $d_\varepsilon(L, K) = \infty$. $\square$

Thus, once the first even and odd occurrences are known, the entire degree set is known. The remaining task is to determine those two thresholds.

Theorem 2.6 (Finite-group degree detection). *Let $L \neq 0$ and K be (not necessarily irreducible) real G -representations, and put $N = \ker L$ and $Q = G/N$. Then*

$$D(L, K) \neq \emptyset \iff K^N \neq 0.$$

In this case $d(L, K) \leq |Q| - 1$. If K is irreducible, the condition is equivalent to $\ker L \subseteq \ker K$.

Proof. Suppose $k \in D(L, K)$ and choose $0 \neq T \in \text{Hom}_G(\text{Sym}^k L, K)$. Since $N = \ker L$ acts trivially on $\text{Sym}^k L$, we have $nT(y) = T(ny) = T(y)$ for all $n \in N$ and $y \in \text{Sym}^k L$. Thus $\text{im } T \subseteq K^N$, and since $T \neq 0$ we obtain $K^N \neq 0$. If K is irreducible, then K^N is G -stable because $N \triangleleft G$. Hence $K^N \neq 0$ implies $K^N = K$. Equivalently, $\ker L \subseteq \ker K$.

For the converse, assume $K^N \neq 0$. Both L and K^N descend to $Q = G/N$, and L is faithful as a Q -representation. We argue in three steps.

A free orbit exists. For $1 \neq q \in Q$ the fixed space L^q is a proper subspace, since L is faithful. The finite union of these proper subspaces cannot cover L , so some $v \in L$ has trivial stabiliser, and the orbit Qv consists of $|Q|$ distinct points.

The orbit can be separated by a linear form. A functional $\ell \in L^*$ fails to separate two distinct orbit points qv and $q'v$ precisely when $\ell(qv - q'v) = 0$. Since $qv - q'v \neq 0$, the set of such functionals is a proper hyperplane of L^* . There are only finitely many pairs of orbit points, so a finite union of these hyperplanes cannot cover L^* . Hence some $\ell \in L^*$ separates all points of Qv .

Interpolate. With such an ℓ , the Lagrange polynomial

$$p(x) = \prod_{1 \neq q \in Q} \frac{\ell(x) - \ell(qv)}{\ell(v) - \ell(qv)}$$

has degree at most $|Q| - 1$, with $p(v) = 1$ and $p(qv) = 0$. Fix $0 \neq w \in K^N$ and average:

$$F(x) = \sum_{q \in Q} p(q^{-1}x) qw.$$

Reindexing $q = aq'$ shows $F(ax) = aF(x)$, so F is equivariant, and $F(v) = w \neq 0$, so $F \neq 0$. The action is linear, so each homogeneous component F_j of F is separately equivariant, and some $F_j \neq 0$ with $j \leq |Q| - 1$. So $\text{Poly}_j(L, K^N)^Q \neq 0$. Hence $\text{Hom}_Q(\text{Sym}^j L, K^N) \neq 0$. Inflating along $G \twoheadrightarrow Q$ and composing with the inclusion $K^N \hookrightarrow K$ gives $0 \neq \text{Hom}_G(\text{Sym}^j L, K^N) \subseteq \text{Hom}_G(\text{Sym}^j L, K)$, so $j \in D(L, K)$. $\square$

The key point here is that every irreducible representation of a finite group occurs somewhere in the symmetric algebra of a faithful representation. The interpolation proof gives the explicit bound $d(L, K) \leq |G/\ker L| - 1$. No such argument is available for a connected group, which is consistent with the failure discussed in Section 7.

For irreducible L, K , combining Theorem 2.6 with (2) identifies the GTW existence criterion with $D(L, K) \neq \emptyset$. The following refinement also decides whether each parity occurs.

Corollary 2.7. *Let $L \neq 0$ and let K be irreducible with $\ker L \subseteq \ker K$. Put $Q = G/\ker L$, so that Q acts faithfully on L and naturally on K .*

(a) *If no element of Q acts as $-I_L$, then both even and odd degrees occur, with*

$$d_0(L, K), d_1(L, K) \leq 2|Q| - 1.$$

(b) *If some $z \in Q$ acts as $-I_L$, then z is a central involution and acts on K as either I_K or $-I_K$. In the first case only even degrees occur, and in the second only odd degrees occur. In either case, the occurring parity has threshold at most $|Q| - 1$.*

Proof. Fix $\varepsilon \in \{0, 1\}$ and let $\tilde{Q} = Q \times \{\pm 1\}$ act on L and K by

$$(q, t) \cdot x = t q x, \quad (q, t) \cdot y = t^\varepsilon q y.$$

A homogeneous Q -equivariant polynomial $P : L \rightarrow K$ is $\tilde{Q}$ -equivariant exactly when its degree is congruent to ε modulo 2.

Suppose first that no element of Q acts as $-I_L$. Then the $\tilde{Q}$ -action on L is faithful. By Theorem 2.6, there exists a nonzero $\tilde{Q}$ -equivariant homogeneous polynomial of degree at most $|\tilde{Q}| - 1 = 2|Q| - 1$. Applying this for $\varepsilon = 0$ and $\varepsilon = 1$ gives both parities.

Now suppose $z \in Q$ acts as $-I_L$. Since Q acts faithfully on L , we have $z^2 = 1$, and for every $q \in Q$ the elements qzq^{-1} and z act identically on L . Hence z is central. The kernel of the $\tilde{Q}$ -action on L is therefore $\{(1, 1), (z, -1)\}$.

By Theorem 2.6, a degree of parity ε occurs exactly when this kernel acts trivially on K , that is, when

$$(-1)^\varepsilon zy = y \quad \text{for all } y \in K.$$

Since z is central, its action on K lies in $\text{End}_G(K)$. Moreover $z^2 = 1$, so irreducibility implies that z acts as either I_K or $-I_K$. Hence exactly one parity occurs: even if z acts as I_K , and odd if it acts as $-I_K$. The faithful quotient has order $|Q|$, so the corresponding threshold is at most $|Q| - 1$. $\square$

2.4. Computing degree sets. The detection theorem decides whether some degree occurs, but not where the first occurrences lie. Molien’s formula packages the dimensions in all degrees into one generating series.

Proposition 2.8 (Molien series). *Let G be finite and $c_k = \dim \operatorname{Hom}_G(\operatorname{Sym}^k L, K)$. Then*

$$\sum_{k \geq 0} c_k t^k = \frac{1}{|G|} \sum_{g \in G} \frac{\chi_K(g)}{\det(I - t \rho_L(g))} = \frac{1}{|G|} \sum_{g \in G} \chi_K(g) \exp\left(\sum_{m \geq 1} \frac{t^m}{m} \chi_L(g^m)\right).$$

Here $\rho_L : G \rightarrow \operatorname{GL}(L)$ is the homomorphism defining the action of G on L , and χ_L and χ_K are the characters of L and K .

This is the usual character average; see Section A.2. The exponential form is useful computationally because it uses only the power-map characters $\chi_L(g^m)$, which are fixed-point counts for permutation representations. We use this form for subset representations in Theorem C.2.

When K is irreducible, the coefficient c_k need not equal the multiplicity of K in $\operatorname{Sym}^k L$ over $\mathbb{R}$, but rather

$$c_k = [\operatorname{Sym}^k L : K] \dim_{\mathbb{R}} \operatorname{End}_G(K).$$

In small representations one can often read the degree set directly from the symmetric powers. The next two examples illustrate Theorem 2.6 and Proposition 2.5.

Example 2.9 (One-dimensional sources). Suppose $\dim L = 1$. Then $\operatorname{Sym}^k L = L^{\otimes k}$. If χ is the real character of L , its order is $q \in \{1, 2\}$. For the one-dimensional target χ^a , $a \in \{0, 1\}$,

$$D(L, \chi^a) = \{k \geq 0 : k \equiv a \pmod{q}\}.$$

Thus a trivial source reaches the trivial target in every degree, while a source of order two reaches the trivial target in even degrees and itself in odd degrees.

Example 2.10 (Rotation representations of C_n). Let $C_n = \langle a \rangle$, and let ρ_s and ρ_t be irreducible two-dimensional real rotation representations, with $2s, 2t \not\equiv 0 \pmod{n}$. Writing $(\rho_s)_{\mathbb{C}} \cong W_s \oplus W_{-s}$, define

$$R_k = \{r \in \mathbb{Z} : |r| \leq k, r \equiv k \pmod{2}\}.$$

Then $\operatorname{Sym}^k((\rho_s)_{\mathbb{C}}) \cong \bigoplus_{r \in R_k} W_{rs}$, and therefore

$$D(\rho_s, \rho_t) = \{k \geq 0 : \exists r \in R_k \text{ with } rs \equiv \pm t \pmod{n}\}.$$

This set is nonempty exactly when $\gcd(s, n) \mid t$, or equivalently when $\ker \rho_s \subseteq \ker \rho_t$. Since $R_k \subseteq R_{k+2}$, occurrence persists within each parity, as predicted by Proposition 2.5.

3. COORDINATEWISE NONLINEARITIES

The preceding sections determine which equivariant polynomial maps exist. We now ask which of these maps a coordinatewise layer produces.

The input representation L is arbitrary throughout and need not be irreducible or multiplicity-free. In particular, the results include interactions between components of a direct-sum input, without requiring a separate treatment of those components.

3.1. Coordinatewise layers and their degree components. There are two pieces of data to track: the polynomial produced by the hidden coordinates, and the scalar Hermite coefficient supplied by the activation. We introduce them separately.

Let L and K be finite-dimensional real orthogonal G -representations, regarded as the input and output spaces. Coordinatewise application is equivariant only after choosing coordinates that the group permutes. A *hidden representation* is an intermediate representation $H = \mathbb{R}^{\mathcal{B}}$ with an orthonormal basis $\mathcal{B}$ permuted by G . Throughout this section we assume that the action on $\mathcal{B}$ is transitive.

No inclusion relations between L , H , and K are assumed. They are connected by equivariant linear maps

$$A : L \longrightarrow H, \quad R : H \longrightarrow K.$$

Thus a coordinatewise layer has the form

$$L \xrightarrow{A} H \xrightarrow{\sigma_{\mathcal{B}}} H \xrightarrow{R} K,$$

where the nonlinearity is applied in the distinguished coordinate basis of H .

All adjoints are taken with respect to the fixed inner products. Hidden representations with several coordinate orbits are treated in Section B.

Definition 3.1. For an activation $\sigma : \mathbb{R} \rightarrow \mathbb{R}$, its *coordinatewise extension* to H is

$$\sigma_{\mathcal{B}} \left(\sum_{b \in \mathcal{B}} z_b b \right) = \sum_{b \in \mathcal{B}} \sigma(z_b) b.$$

The associated *coordinatewise layer* is

$$F : L \longrightarrow K, \quad F(x) = R\sigma_{\mathcal{B}}(Ax).$$

Since $\sigma_{\mathcal{B}}$ commutes with permutations of $\mathcal{B}$, the layer F is equivariant. We assume $A \neq 0$.

Definition 3.2. For $b \in \mathcal{B}$, define the coordinate vector $a_b = A^*b \in L$. Thus the b th hidden coordinate is $(Ax)_b = \langle x, a_b \rangle$. The *coordinate Gram matrix* is

$$C = (C_{bb'})_{b, b' \in \mathcal{B}}, \quad C_{bb'} = \langle a_b, a_{b'} \rangle.$$

Thus C is the matrix of AA^* in the basis $\mathcal{B}$.

Lemma 3.3. *There is an $s > 0$ such that $\|a_b\| = s$ for every $b \in \mathcal{B}$. Moreover,*

$$s^2 = C_{bb} = \frac{\text{tr } C}{\dim H}.$$

For $X \sim \gamma_L$, every coordinate $(AX)_b$ therefore has distribution $N(0, s^2)$.

Proof. Equivariance and orthogonality give $a_{gb} = ga_b$. Transitivity implies that all coordinate vectors have the same norm, and $A \neq 0$ implies that this norm is positive. Summing the diagonal entries of C gives the trace identity. Finally, $\langle X, a_b \rangle$ is centred Gaussian with variance $\|a_b\|^2$. $\square$

The activation enters the k th chaos through a single scalar coefficient.

Definition 3.4. Suppose $\sigma(sZ)$ is square-integrable for $Z \sim N(0, 1)$. Its *Hermite coefficients at scale s* are

$$a_k(\sigma; s) = \frac{1}{k!} \mathbb{E}[\sigma(sZ) \text{He}_k(Z)], \quad a_k(\sigma) := a_k(\sigma; 1).$$

We impose this integrability assumption throughout. It holds for every continuous piecewise-linear activation with finitely many knots, since such a function has at most linear growth.

The remaining degree- k information comes from the fixed lift and readout. It is encoded by the coordinatewise k th power.

Definition 3.5. For $z = \sum_b z_b b \in H$, define

$$z^{\odot k} := \sum_b z_b^k b \quad (k \geq 0).$$

The *degree- k coordinate polynomial* associated with A and R is

$$Q_k : L \longrightarrow K, \quad Q_k(x) = R((Ax)^{\odot k}) = \sum_{b \in \mathcal{B}} \langle x, a_b \rangle^k Rb.$$

For matrices, $\circ$ denotes entrywise multiplication, and $C^{\circ k}$ denotes the entrywise power, with $C^{\circ 0} = J$, the all-ones matrix.

Coordinatewise powers commute with permutations, so $Q_k \in \text{Poly}_k(L, K)^G$. On homogeneous degree- k polynomials, Π_k is an isomorphism onto $\mathcal{H}_k(L)$. In particular,

$$\Pi_k Q_k = 0 \iff Q_k = 0.$$

All L^2 norms below use the standard Gaussian on the source and the fixed inner product on the target.

With these definitions, the activation and the coordinate geometry separate as follows.

Theorem 3.6 (Coordinatewise factorisation and Gram formula). *Under the preceding assumptions, for every $k \geq 0$,*

$$\Pi_k F = \frac{a_k(\sigma; s)}{s^k} \Pi_k Q_k, \tag{6}$$

$$\|\Pi_k F\|_{L^2}^2 = \frac{k! |a_k(\sigma; s)|^2}{s^{2k}} \text{tr}(RC^{\circ k} R^*).$$

Consequently,

$$\Pi_k F \neq 0 \iff a_k(\sigma; s) \neq 0 \text{ and } Q_k \neq 0,$$

$$Q_k \neq 0 \iff \text{tr}(RC^{\circ k} R^*) > 0.$$

No irreducibility or multiplicity-free assumption is required.

The transitivity assumption is used only to give all hidden coordinates the same variance. The corresponding formula for several coordinate orbits is recorded in Section B.

Proof. For each b , the random variable $\langle X, a_b \rangle / s$ is standard normal. Hence

$$\sigma(\langle x, a_b \rangle) = \sum_{k \geq 0} a_k(\sigma; s) \text{He}_k(\langle x, a_b \rangle / s)$$

in $L^2(\gamma_L)$. The k th summand belongs to $\mathcal{H}_k(L)$, and

$$\Pi_k(\langle x, a_b \rangle^k) = s^k \text{He}_k(\langle x, a_b \rangle / s).$$

Multiplying by Rb and summing over b proves (6).

Applying Lemma 1.2 to the hidden coordinates gives

$$\left\langle \Pi_k(\langle x, a_b \rangle^k), \Pi_k(\langle x, a_{b'} \rangle^k) \right\rangle_{L^2} = k! C_{bb'}^k.$$

Therefore

$$\|\Pi_k Q_k\|_{L^2}^2 = k! \sum_{b, b'} C_{bb'}^k \langle Rb, Rb' \rangle = k! \text{tr}(RC^{\circ k} R^*).$$

Combining this identity with (6) proves the norm formula. The nonvanishing statements follow from injectivity of Π_k on homogeneous degree- k polynomials. $\square$

The theorem separates the three questions (a)–(c) of the introduction. For fixed A and R , changing the activation only changes the scalar multiplying $\Pi_k Q_k$; it cannot select a different polynomial direction at degree k .

Example 3.7. Let $H = \mathbb{R}^3$ be the S_3 permutation representation, and recall the trivial and standard subrepresentations, L_0 and L_1 .

Let $A : L_1 \hookrightarrow H$ be inclusion, and $R = p_{L_0}$ be orthogonal projection. Then $C = p_{L_1}$ and $s^2 = \dim L_1/3 = 2/3$. As in (1),

$$Q_2(x) = \frac{1}{3} \|x\|^2 \mathbf{1}, \quad \Pi_2 Q_2(x) = \frac{1}{3} (\|x\|^2 - 2) \mathbf{1}.$$

Here $\mathbb{E}[\|X\|^2] = 2$ for $X \sim \gamma_{L_1}$. Thus

$$\Pi_2 F(x) = \frac{a_2(\sigma; \sqrt{2/3})}{2} (\|x\|^2 - 2) \mathbf{1}.$$

The coordinate polynomial is nonzero, so this second-chaos interaction is present precisely when $a_2(\sigma; \sqrt{2/3}) \neq 0$.

3.2. The submodule generated by the coordinates. We next explain why the polynomial Q_k can vanish even when $\text{Hom}_G(\text{Sym}^k L, K)$ is nonzero.

Definition 3.8. For $k \geq 0$, define

$$Y_k = \text{span}\{a_b^k : b \in \mathcal{B}\} \subseteq \text{Sym}^k L.$$

We call Y_k the *coordinate-generated submodule* of degree k . Define also

$$\phi_k : H \longrightarrow \text{Sym}^k L, \quad \phi_k(b) = a_b^k.$$

The relation $a_{gb} = ga_b$ implies that ϕ_k is equivariant. Thus $Y_k = \text{im } \phi_k$ is indeed a submodule.

Proposition 3.9. For every $k \geq 0$, $\phi_k^* \phi_k = C^{\circ k}$, and so

$$Y_k \cong (\ker C^{\circ k})^\perp$$

as G -representations. Proposition 1.1 gives $Q_k(x) = R\phi_k^*(x^k)$. Thus

$$Q_k \neq 0 \iff R\phi_k^* \neq 0.$$

Proof. The tensor inner product gives

$$\langle a_b^k, a_{b'}^k \rangle = \langle a_b, a_{b'} \rangle^k = C_{bb'}^k.$$

Hence $C^{\circ k}$ is the Gram matrix of the vectors a_b^k , proving $\phi_k^* \phi_k = C^{\circ k}$. In particular, $\ker \phi_k = \ker C^{\circ k}$. Restricting ϕ_k to the orthogonal complement of its kernel gives the stated isomorphism onto Y_k .

For $z \in \text{Sym}^k L$,

$$R\phi_k^* z = \sum_b \langle z, a_b^k \rangle Rb.$$

Taking $z = x^k$ gives $Q_k(x)$. Since pure powers span $\text{Sym}^k L$, the polynomial vanishes identically if and only if $R\phi_k^*$ is zero. $\square$

An arbitrary degree- k equivariant polynomial may use all of $\text{Sym}^k L$, whereas the coordinate construction sees only Y_k . A target can therefore occur in $\text{Sym}^k L$ and still be invisible to the chosen coordinates. Even when it occurs in Y_k , the readout R may kill the resulting direction.

3.3. Projection onto irreducible summands. When the lift is an inclusion and the readout is an orthogonal projection, the Gram criterion becomes especially transparent. Let $W, K \subseteq H$ be nonzero subrepresentations and take

$$A : W \hookrightarrow H, \quad R = p_K : H \longrightarrow K.$$

Writing E_W and E_K for the orthogonal projector matrices in $\mathcal{B}$, we have

$$C = E_W, \quad s^2 = \frac{\dim W}{\dim H}, \quad Q_k(x) = p_K(x^{\odot k}) \quad (x \in W).$$

The norm formula becomes

$$\|\Pi_k Q_k\|_{L^2(\gamma_W; K)}^2 = k! \operatorname{tr}(E_K E_W^{\odot k}),$$

which holds without a multiplicity-free assumption.

Suppose now that H is multiplicity-free, $H = \bigoplus_{j=0}^m L_j$ with the L_j pairwise non-isomorphic and irreducible. Let L_0 be the trivial summand, let E_j be orthogonal projection onto L_j , and let $d_j = \dim L_j$. Transitivity gives

$$E_0 = \frac{J}{\dim H}, \quad (E_j)_{bb} = \frac{d_j}{\dim H}.$$

When H is multiplicity-free, the matrix $E_W^{\odot k}$ is determined by one scalar on each irreducible summand. We record these scalars next.

Definition 3.10. For the fixed source subrepresentation W (not necessarily irreducible), define

$$\kappa_j(k) := \frac{1}{d_j} \operatorname{tr}(E_j E_W^{\odot k}), \quad k \geq 0.$$

We call $\kappa_j(k)$ the *projector coefficient* for the source W , target L_j and degree k .

Corollary 3.11. For each $k \geq 0$:

(a) The Hadamard power of the projector decomposes as

$$E_W^{\odot k} = \sum_{j=0}^m \kappa_j(k) E_j, \quad \kappa_j(k) \geq 0.$$

(b) For $Q_k(x) = p_j(x^{\odot k})$ with $x \in W$,

$$Q_k \neq 0 \iff \kappa_j(k) > 0, \quad \|\Pi_k Q_k\|_{L^2}^2 = k! d_j \kappa_j(k).$$

(c) The coordinate-generated submodule for the inclusion $W \hookrightarrow H$ is

$$Y_k \cong \bigoplus_{j: \kappa_j(k) > 0} L_j.$$

Proof. Hadamard products of G -equivariant matrices are again equivariant. Thus $E_W^{\odot k}$ is a self-adjoint G -endomorphism of $H = \bigoplus_{j=0}^m L_j$. By multiplicity-freeness and Schur's lemma, together with self-adjointness, it acts on each L_j by a real scalar. Hence $E_W^{\odot k} = \sum_{j=0}^m \lambda_j(k) E_j$. Multiplying by E_j and taking traces proves $E_W^{\odot k} = \sum_{j=0}^m \kappa_j(k) E_j$.

Since E_W is positive semidefinite, the Schur product theorem implies that $E_W^{\odot k}$ is positive semidefinite. Its scalar on L_j is therefore nonnegative, so $\kappa_j(k) \geq 0$.

For the inclusion $W \hookrightarrow H$ and projection $p_j : H \rightarrow L_j$, the norm formula gives

$$\|\Pi_k Q_k\|_{L^2}^2 = k! \operatorname{tr}(E_j E_W^{\odot k}) = k! d_j \kappa_j(k).$$

Since Π_k is injective on homogeneous degree- k polynomials,

$$Q_k \neq 0 \iff \kappa_j(k) > 0.$$

Finally, Proposition 3.9 gives $Y_k \cong (\ker E_W^{\circ k})^\perp$. Since $E_W^{\circ k}$ acts on L_j by $\kappa_j(k)$, its kernel is the sum of those L_j with $\kappa_j(k) = 0$. Hence

$$Y_k \cong \bigoplus_{j: \kappa_j(k) > 0} L_j.$$

□

When W is an irreducible summand and $k = 2$, these Hadamard coefficients are the usual Krein parameters, the structure constants for entrywise products of primitive idempotents, up to the chosen normalisation. Higher k gives iterated Hadamard products. See also Section C for the Johnson-scheme case.

Lemma 3.12. *For each $k \geq 0$,*

$$\kappa_j(k) > 0 \implies \kappa_j(k+2) > 0.$$

Equivalently, if $p_j(x^{\odot k})|_W$ is nonzero, then $p_j(x^{\odot(k+2)})|_W$ is nonzero. Hence, for each j , the supported degrees form at most one even tail and one odd tail.

Proof. For the trivial summand L_0 ,

$$\kappa_0(2) = \text{tr}(E_0 E_W^{\circ 2}) = \frac{\dim W}{\dim H}.$$

Using the decomposition from Corollary 3.11,

$$E_W^{\circ 2} = \frac{\dim W}{\dim H} E_0 + B = \frac{\dim W}{(\dim H)^2} J + B,$$

where B is positive semidefinite. Therefore

$$E_W^{\circ(k+2)} = E_W^{\circ k} \circ E_W^{\circ 2} = \frac{\dim W}{(\dim H)^2} E_W^{\circ k} + E_W^{\circ k} \circ B.$$

The second term is positive semidefinite by the Schur product theorem. Its projector coefficients are therefore nonnegative, so

$$\kappa_j(k+2) \geq \frac{\dim W}{(\dim H)^2} \kappa_j(k).$$

Since $W \neq 0$, positivity of $\kappa_j(k)$ implies positivity of $\kappa_j(k+2)$. The equivalence with the coordinate polynomial follows from Corollary 3.11. □

This is the coordinate-level analogue of Proposition 2.5: within each parity, coordinate support also persists once it appears.

3.4. Examples of realisation and obstruction. The first example shows that a permitted quadratic map can be missed by the coordinate construction.

Example 3.13. Let S_4 act on the six unordered pairs of $[4] := \{1, 2, 3, 4\}$, and let $H = \mathbb{R}^{\binom{[4]}{2}}$ be the corresponding permutation representation. Its standard subrepresentation is

$$W = \left\{ x_{ab} = u_a + u_b : u \in \mathbb{R}^4, \sum_a u_a = 0 \right\}.$$

The map $u \mapsto x$ is equivariant and injective, since $u_a = \frac{1}{2} \sum_{b \neq a} x_{ab}$ on its image. Thus W is a copy of the Specht module $S^{(3,1)}$. Let J_c be the complementation operator on edge coordinates, defined by

$$J_c e_A = e_{[4] \setminus A}.$$

For $x \in W$, the zero-sum condition gives $x_{A^c} = -x_A$. Thus J_c acts as -1 on W , whereas

$$(x^{\odot 2})_{A^c} = (x^{\odot 2})_A.$$

Since J_c is a self-adjoint involution, its two eigenspaces are orthogonal. Consequently,

$$p_W(x^{\odot 2}) = 0 \quad (x \in W).$$

Nevertheless, the map

$$u \mapsto \left(u_a^2 - \frac{1}{4} \sum_{b=1} u_b^2 \right)^4$$

is a nonzero equivariant quadratic map from the standard vertex representation to itself. Transporting it through the identification above gives $\text{Hom}_{S_4}(\text{Sym}^2 W, W) \neq 0$.

Thus symmetry permits a quadratic interaction that the fixed layer $F(x) = p_W \sigma_B(x)$ cannot produce in its second chaos. This holds for every activation satisfying the integrability assumption, including biased ReLU, because its coordinate polynomial Q_2 is zero. The same argument applies in every even degree.

Equivalently, the Gram criterion in Theorem 3.6 gives $\text{tr}(E_W E_W^{\circ 2}) = 0$ in this example.

For regular representations of finite abelian groups, every permitted degree has a nonzero coordinate polynomial.

Proposition 3.14. *Let G be finite abelian, and let $H = \mathbb{R}G = \bigoplus_j L_j$ be its real regular representation, equipped with the regular coordinate basis. Let $W \subseteq H$ be any nonzero subrepresentation, and let p_j be orthogonal projection onto L_j . Then, for every $k \geq 0$,*

$$p_j(x^{\odot k})|_W \neq 0 \iff \text{Hom}_G(\text{Sym}^k W, L_j) \neq 0.$$

Proof. After complexification, identify $H_{\mathbb{C}}$ with functions $G \rightarrow \mathbb{C}$. Since G is abelian, its regular representation is a direct sum of one-dimensional character spaces. The subrepresentation $W_{\mathbb{C}}$ is therefore spanned by a subset of the character functions.

Consider the equivariant multiplication map

$$\mu_k : \text{Sym}^k W_{\mathbb{C}} \longrightarrow H_{\mathbb{C}}, \quad f_1 \cdots f_k \mapsto f_1 \odot \cdots \odot f_k.$$

A monomial in character functions is sent to their pointwise product, which is a nonzero character function of the same representation type. Thus every character occurring in $\text{Sym}^k W_{\mathbb{C}}$ occurs in $\text{im } \mu_k$, and conversely.

The polynomial $p_j(x^{\odot k})$ corresponds to $p_j \mu_k$. Since pure powers span the symmetric power, this polynomial is nonzero exactly when a character of $(L_j)_{\mathbb{C}}$ occurs in $\text{Sym}^k W_{\mathbb{C}}$. That is, $\text{Hom}_G(\text{Sym}^k W, L_j) \neq 0$. $\square$

The proposition detects nonzero degrees only; it does not say that the selected polynomial spans the full equivariant polynomial space. For these regular abelian representations, the presence of a permitted degree in the layer is therefore decided by the activation coefficient in Theorem 3.6.

4. ACTIVATION SPECTRA

The coordinatewise factorisation separates the polynomial selected by a layer from the coefficient supplied by its activation. We now determine when $a_k(\sigma; s)$ vanishes.

Definition 4.1. Let $s > 0$, and suppose $\sigma(sZ)$ is square-integrable for $Z \sim N(0, 1)$. The *Hermite spectrum of σ at scale s* is the sequence $(a_k(\sigma; s))_{k \geq 0}$. Its *support* is

$$\text{supp } a_{\bullet}(\sigma; s) = \{k \in \mathbb{N}_0 : a_k(\sigma; s) \neq 0\}.$$

At scale one, write $\text{supp } a_{\bullet}(\sigma) = \text{supp } a_{\bullet}(\sigma; 1)$.

For a layer satisfying the assumptions of Theorem 3.6,

$$\{k : \Pi_k F \neq 0\} = \{k : Q_k \neq 0\} \cap \text{supp } a_\bullet(\sigma; s).$$

Thus the activation support determines which of the nonzero coordinate polynomials contribute to the layer.

4.1. Positively homogeneous activations. We start with positively homogeneous activations, which include the identity, absolute value, and ReLU.

Definition 4.2. A function $\sigma : \mathbb{R} \rightarrow \mathbb{R}$ is *positively homogeneous of degree one* if

$$\sigma(tx) = t\sigma(x) \quad (t \geq 0, x \in \mathbb{R}).$$

Writing $m_+ = \sigma(1)$ and $m_- = -\sigma(-1)$, such a function is necessarily of the form

$$\sigma(x) = \begin{cases} m_+x, & x \geq 0, \\ m_-x, & x \leq 0, \end{cases} = \frac{m_+ + m_-}{2}x + \frac{m_+ - m_-}{2}|x|.$$

Every such function is continuous and piecewise linear.

From here on, *homogeneous* means positively homogeneous of degree one.

Theorem 4.3. Let $\sigma \neq 0$ be homogeneous, with slopes m_+ and m_- as above. For every $s > 0$,

$$a_k(\sigma; s) = s a_k(\sigma),$$

and its support is given by

$$\begin{aligned} m_+ = m_- &\implies \text{supp } a_\bullet(\sigma) = \{1\}, \\ m_+ = -m_- &\implies \text{supp } a_\bullet(\sigma) = 2\mathbb{N}_0, \\ m_+ \neq \pm m_- &\implies \text{supp } a_\bullet(\sigma) = \{1\} \cup 2\mathbb{N}_0. \end{aligned}$$

Proof. Positive homogeneity gives $\sigma(sZ) = s\sigma(Z)$, which proves the scaling identity.

The odd part of σ is a multiple of $x = \text{He}_1(x)$, which contributes only in degree one. The function $|x|$ is even, so its odd Hermite coefficients vanish. Its degree-zero coefficient is $\mathbb{E}[|Z|] = 2\varphi(0) > 0$, where $\varphi(t) = (2\pi)^{-1/2}e^{-t^2/2}$ is the standard one-dimensional Gaussian density. For $r \geq 1$, two integrations by parts give

$$a_{2r}(|\cdot|) = \frac{2}{(2r)!} \text{He}_{2r-2}(0)\varphi(0).$$

These coefficients are nonzero because

$$\text{He}_{2m}(0) = (-1)^m \frac{(2m)!}{2^m m!}.$$

Thus $|x|$ has support exactly $2\mathbb{N}_0$. The two parts of σ have disjoint supports, so their nonzero coefficients cannot cancel. $\square$

Example 4.4. The identity activation has support $\{1\}$, absolute value has support $2\mathbb{N}_0$, and $\text{ReLU}(x) := \max(x, 0) = \frac{x+|x|}{2}$ has support $\{1\} \cup 2\mathbb{N}_0$.

The scaling identity shows that changing the input scale cannot create a degree that is absent at scale one. In particular, no odd degree $k > 1$ occurs, for any σ in the class and any input scale.

Corollary 4.5. Let $F(x) = R\sigma_{\mathcal{B}}(Ax)$ satisfy the assumptions of Theorem 3.6, with σ homogeneous. Then

$$\Pi_k F = 0 \quad \text{for every odd } k \geq 3.$$

The same conclusion holds for finite sums of such layers.

Proof. Write $\sigma(z) = \lambda z + \mu |z|$ with $\lambda = \frac{m_+ + m_-}{2}$ and $\mu = \frac{m_+ - m_-}{2}$. The second summand is even, so the odd part of the layer is

$$\frac{F(x) - F(-x)}{2} = \lambda R Ax,$$

which is linear and therefore lies in the first chaos, while an even map has no odd chaos components at all. Hence $\Pi_k F = 0$ for every odd $k \geq 3$, and the same argument applies to a finite sum of such layers. $\square$

This statement applies to a single layer, or a sum of such layers, but it does not imply the same restriction for their compositions.

4.2. Bias and input scale. We now show how a shared additive bias changes this conclusion.

Definition 4.6. For a threshold $c \in \mathbb{R}$, define

$$\sigma_c(x) := \text{ReLU}(x - c).$$

For fixed equivariant maps $A : L \rightarrow H$ and $R : H \rightarrow K$, write $F_c(x) = R(\sigma_c)_B(Ax)$.

The same threshold is applied to every hidden coordinate, so F_c remains equivariant. The identity $\sigma_c(sz) = s \sigma_{c/s}(z)$ gives $a_k(\sigma_c; s) = s a_k(\sigma_{c/s})$. Thus the support depends on the ratio c/s . The following formula determines it explicitly.

Corollary 4.7. For every $s > 0$ and $k \geq 2$,

$$a_k(\sigma_c; s) = \frac{s}{k!} \text{He}_{k-2}(c/s) \varphi(c/s).$$

Moreover, $a_0(\sigma_c; s)$ and $a_1(\sigma_c; s)$ are strictly positive.

Proof. Put $u = c/s$. For $k \geq 2$,

$$a_k(\sigma_c; s) = \frac{s}{k!} \int_u^\infty (z - u) \text{He}_k(z) \varphi(z) dz = \frac{s}{k!} \text{He}_{k-2}(u) \varphi(u),$$

where the second equality follows from two integrations by parts using $(\text{He}_j \varphi)' = -\text{He}_{j+1} \varphi$. The cases $k = 0, 1$ follow directly from the same integral and are strictly positive. $\square$

Corollary 4.8. Fix $A, R \neq 0$ and a degree bound $D \in \mathbb{N}_0$. Let s be the common coordinate scale. For all but finitely many thresholds $c \in \mathbb{R}$,

$$\Pi_k F_c \neq 0 \iff Q_k \neq 0 \quad (0 \leq k \leq D).$$

Proof. The coefficients in degrees zero and one never vanish. For $2 \leq k \leq D$, vanishing occurs precisely when c/s is a root of He_{k-2} . Each of these finitely many polynomials has finitely many roots. Outside the resulting finite set of thresholds, every activation coefficient through degree D is nonzero. Apply Theorem 3.6. $\square$

Thus a suitable shared bias retains every nonzero coordinate polynomial through a chosen degree. It cannot recover a degree for which $Q_k = 0$, or select additional polynomial directions with A and R fixed.

Unlike the positively homogeneous case, changing the input scale at a fixed nonzero threshold can change activation support, because it changes c/s . This concerns only the scalar activation coefficients, so any change to the lift must also be accounted for in the coordinate polynomials.

4.3. An activation obstruction for C_8 . The ungraded ReLU obstruction in this example is already part of the cyclic-group classification in [GTW25, Theorem 3B.4]. Here we identify its first permitted degree and explain the effect of a bias.

The real regular representation of C_8 decomposes as

$$\mathbb{R}[C_8] \cong \mathbf{1} \oplus \varepsilon \oplus \rho_1 \oplus \rho_2 \oplus \rho_3,$$

where ε is the sign character and ρ_j is the two-dimensional representation in which a generator acts by rotation through $2\pi j/8$.

Consider the two projected layers from ρ_1 to ρ_3 and from ρ_3 to ρ_1 .

The element a^4 acts as $-I$ on ρ_1 and on ρ_3 , so Corollary 2.7 already shows that only odd degrees occur in either direction. Example 2.10 locates the first of them:

$$D(\rho_1, \rho_3) = D(\rho_3, \rho_1) = \{3, 5, 7, \dots\}.$$

Indeed, for the first direction the character condition is $r \equiv \pm 3 \pmod{8}$, while for the reverse direction it is $3r \equiv \pm 1 \pmod{8}$, which is equivalent to the same condition. Together with $|r| \leq k$ and $r \equiv k \pmod{2}$, this permits exactly the odd degrees $k \geq 3$.

By Proposition 3.14, the coordinate polynomials are nonzero in every permitted degree. For a positively homogeneous activation, however, Theorem 4.3 gives

$$a_k(\sigma; s) = 0 \quad (k = 3, 5, 7, \dots).$$

Hence every chaos component of either projected layer vanishes. Since these layers are continuous, Gaussian almost-everywhere vanishing implies that they vanish identically.

A shifted ReLU with $c \neq 0$ changes this conclusion. Its third coefficient is nonzero by Corollary 4.7, so both projected layers acquire a nonzero third-chaos component.

This distinguishes the example from Example 3.13. There, the quadratic coordinate polynomial vanishes and changing the activation cannot repair it. Here, the cubic coordinate polynomial is already nonzero and the bias supplies its missing scalar coefficient.

5. ARCHITECTURES DETERMINED BY THE DEGREE CALCULATION

The preceding results concern one component of one layer. We now describe what a collection of such components can span, how to control its activation coefficients, and how these statements lead to explicit polynomial layers.

5.1. The degree subspace of a fixed collection of components. At a fixed degree, each branch contributes at most one polynomial direction. We first compute the span of those directions.

Fix equivariant $A_h : L \rightarrow H_h$ and $R_h : H_h \rightarrow K$ for $1 \leq h \leq M$, where each $H_h = \mathbb{R}^{\mathcal{B}_h}$ has a transitive coordinate basis $\mathcal{B}_h$ and $A_h \neq 0$. Write

$$Q_{h,k}(x) = R_h((A_h x)^{\odot k}), \quad \mathcal{S}_k := \text{span}\{Q_{h,k} : 1 \leq h \leq M\} \subseteq \text{Poly}_k(L, K)^G.$$

Consider the layer $F_\theta(x) = \sum_h \theta_h R_h(\sigma_h)_{\mathcal{B}_h}(A_h x)$, with independent real output coefficients θ_h .

Proposition 5.1. *If $a_k(\sigma_h; s_h) \neq 0$ for all h , where $s_h^2 = \text{tr}(A_h A_h^*) / \dim H_h$, then*

$$\{\Pi_k F_\theta : \theta \in \mathbb{R}^M\} = \Pi_k \mathcal{S}_k.$$

Its dimension is the rank of the positive semidefinite Gram matrix

$$\mathbf{G}_{h\ell}^{(k)} = k! \text{tr} \left(R_h (A_h A_\ell^*)^{\odot k} R_\ell^* \right).$$

Here $A_h A_\ell^ : H_\ell \rightarrow H_h$ is represented in the two coordinate bases; its entrywise power is therefore well-defined even when H_h and H_ℓ have different dimensions. If an activation coefficient vanishes, omit that component before forming the span and Gram matrix. In*

particular, the whole degree- k equivariant space is spanned exactly when this rank equals $\dim \operatorname{Hom}_G(\operatorname{Sym}^k L, K)$.

Proof. Fix the degree k . By Theorem 3.6, the h th branch contributes

$$\Pi_k(R_h(\sigma_h)_{\mathcal{B}_h}(A_h x)) = \frac{a_k(\sigma_h; s_h)}{s_h^k} \Pi_k Q_{h,k}.$$

Thus, when $a_k(\sigma_h; s_h) \neq 0$, the activation only rescales the degree- k direction $\Pi_k Q_{h,k}$ selected by that branch. Since the coefficients θ_h are independent, varying θ therefore gives exactly

$$\{\Pi_k F_\theta : \theta \in \mathbb{R}^M\} = \operatorname{span}\{\Pi_k Q_{h,k} : 1 \leq h \leq M\} = \Pi_k \mathcal{S}_k.$$

It remains to determine the dimension of this span. The matrix $\mathbf{G}^{(k)}$ is precisely the Gram matrix of the vectors $\Pi_k Q_{h,k}$. The same calculation as in Theorem 3.6 gives

$$\mathbf{G}_{h\ell}^{(k)} = k! \operatorname{tr}\left(R_h(A_h A_\ell^*)^{ok} R_\ell^*\right), \quad \text{and} \quad \operatorname{rank} \mathbf{G}^{(k)} = \dim \Pi_k \mathcal{S}_k.$$

Finally, Π_k is an isomorphism on homogeneous degree- k polynomials, so this accessible space equals the full degree- k equivariant polynomial space exactly when

$$\operatorname{rank} \mathbf{G}^{(k)} = \dim \operatorname{Hom}_G(\operatorname{Sym}^k L, K).$$

If $a_k(\sigma_h; s_h) = 0$, the corresponding branch contributes nothing in degree k and may simply be omitted. $\square$

This is the degree- k space accessible for the chosen lifts A_h and readouts R_h . Varying those maps may enlarge it. Moreover, although each degree can be analysed separately, the same coefficients θ_h act across all degrees, so the degree components cannot in general be chosen independently.

This also yields an approximation obstruction. For $T \in L^2(\gamma_L; K)^G$ and any degree bound $D \geq 0$,

$$\inf_{\theta} \|T - F_\theta\|_{L^2}^2 \geq \sum_{k=0}^D \operatorname{dist}_{L^2}(\Pi_k T, \Pi_k \mathcal{S}_k)^2,$$

since the degree- k component of F_θ lies in $\Pi_k \mathcal{S}_k$. Thus any component of $\Pi_k T$ outside this space gives an unavoidable error for the fixed architecture under the Gaussian reference measure.

The preceding proposition treats one degree at a time, but the same branch coefficient acts across all degrees. To control several degrees independently, we need activations with prescribed initial Hermite coefficients.

Proposition 5.2. *Fix $D \geq 0$. There exist real $t_0, \dots, t_D$ such that, for every $(b_0, \dots, b_D) \in \mathbb{R}^{D+1}$, there are $\theta_0, \dots, \theta_D \in \mathbb{R}$ for which the continuous piecewise-linear function*

$$\sigma(z) = \sum_{r=0}^D \theta_r \operatorname{ReLU}(z - t_r)$$

satisfies $a_k(\sigma; 1) = b_k$ for $0 \leq k \leq D$. Thus the first $D+1$ Hermite coefficients can be chosen independently.

Proof. For $0 \leq k \leq D$, set $f_k(t) := a_k(\operatorname{ReLU}(\cdot - t))$. By linearity of the Hermite coefficients, if $\sigma(z) = \sum_{r=0}^D \theta_r \operatorname{ReLU}(z - t_r)$, then

$$a_k(\sigma; 1) = \sum_{r=0}^D \theta_r f_k(t_r).$$

It is therefore enough to show that $(f_0(t), \dots, f_D(t))$, as t varies, spans $\mathbb{R}^{D+1}$. If not, there exist coefficients $c_0, \dots, c_D$, not all zero, such that $\sum_{k=0}^D c_k f_k(t) = 0$ for every t . Differentiating $f_k(t)$ twice gives $f_k''(t) = \frac{1}{k!} \text{He}_k(t) \varphi(t)$. Hence

$$0 = \sum_{k=0}^D c_k f_k''(t) = \varphi(t) \sum_{k=0}^D \frac{c_k}{k!} \text{He}_k(t).$$

Since $\varphi(t) > 0$ and the Hermite polynomials are linearly independent, $c_k = 0$ for every k .

Therefore the vectors $(f_0(t), \dots, f_D(t))$ span $\mathbb{R}^{D+1}$. Choose $t_0, \dots, t_D$ giving a basis. Then, for any prescribed $(b_0, \dots, b_D)$, suitable coefficients θ_r give $a_k(\sigma; 1) = b_k$ for all $0 \leq k \leq D$. $\square$

For a lift with coordinate variance s^2 , the preceding proposition may be applied after rescaling the hidden coordinates by s . Choosing $b_k = \delta_{kj}$ produces a piecewise-linear activation whose chaos components vanish in every degree $0, \dots, D$ except degree j . Components above degree D may still be present.

There are two simpler ways to obtain an exact degree- j layer. Using the polynomial activation He_j coordinatewise gives the pure j th-chaos component

$$R(\text{He}_j)_B(Ax/s) = s^{-j} \Pi_j[R((Ax)^{\odot j})].$$

Alternatively, $R((Ax)^{\odot j})$ is itself an ordinary homogeneous polynomial of degree j .

Thus the piecewise-linear construction controls only finitely many chaos coefficients, whereas the Hermite activation isolates a single chaos degree exactly. In particular, Proposition 5.2 does not control the higher-degree tail.

5.2. Regular lifts span every degree. Regular representations of finite groups give a simple construction showing that coordinatewise powers can realise every equivariant polynomial map of a fixed degree. The obstruction of Example 3.13 is therefore a property of the chosen coordinates, not an intrinsic one.

Proposition 5.3. *Let G be finite and let L, K be real orthogonal G -representations. For $u \in L$ define the linear map*

$$A_u : L \longrightarrow \mathbb{R}G, \quad (A_u x)_g = \langle x, gu \rangle,$$

where $\mathbb{R}G$ carries the left regular action, and for $v \in K$ define $A_v : K \rightarrow \mathbb{R}G$ analogously and put $R_v := |G|^{-1} A_v^*$, that is, $R_v z = |G|^{-1} \sum_{g \in G} z_g gv$. Then A_u and R_v are equivariant, their coordinate polynomial is

$$Q_{u,v,k}(x) = R_v((A_u x)^{\odot k}) = \frac{1}{|G|} \sum_{g \in G} \langle x, gu \rangle^k gv,$$

and for every $k \geq 0$ these polynomials span $\text{Poly}_k(L, K)^G$ as u and v vary.

Proof. Equivariance of A_u is the identity $(A_u(hx))_g = \langle x, h^{-1}gu \rangle = (A_u x)_{h^{-1}g}$, and R_v is a multiple of the adjoint of an equivariant map, hence equivariant. Since $A_v^* e_g = gv$, the displayed formula for $Q_{u,v,k}$ follows.

Pure powers of linear forms span the homogeneous scalar polynomials of degree k , so the maps $P_{u,v}(x) = \langle x, u \rangle^k v$ span $\text{Poly}_k(L, K)$. Averaging, that is,

$$\mathcal{R}(P)(x) = |G|^{-1} \sum_g gP(g^{-1}x),$$

projects $\text{Poly}_k(L, K)$ onto $\text{Poly}_k(L, K)^G$, and orthogonality of the action gives $\mathcal{R}(P_{u,v}) = Q_{u,v,k}$. A projection maps a spanning set of the ambient space to a spanning set of its image. $\square$

The proposition gives a universal, but generally inefficient, way to match any prescribed equivariant polynomial components through a finite degree bound D . For each required degree k , choose finitely many $Q_{u,v,k}$ spanning the desired component, with u normalised so that the coordinates of $A_u X$ have variance one.

By Proposition 5.2, one may choose a piecewise-linear activation whose Hermite coefficients vanish in all degrees $0, \dots, D$ except degree k . Then (6) shows that the corresponding layer contributes, up to degree D , only the chosen degree- k coordinate polynomial. Summing over the required degrees gives the prescribed chaos components.

Thus copies of the regular representation are sufficient to recover every equivariant polynomial direction. The cost is that each regular block has $|G|$ hidden coordinates, and spanning several directions may require several such blocks.

6. SUBSET REPRESENTATIONS OF THE SYMMETRIC GROUP

We now make the ambient polynomial spaces from Section 2 explicit for an important family of permutation representations. Subset representations model vertex, edge, and higher-order subset features in equivariant graph networks. This section describes the full polynomial spaces against which the coordinate-selected spaces of Sections 3 and 5 can be compared.

Definition 6.1. Let $[n] = \{1, \dots, n\}$, with $[0] = \emptyset$, and let S_n be its permutation group. For $r \geq 0$, define

$$X_{n,r} := \binom{[n]}{r} = \{A \subseteq [n] : |A| = r\}.$$

The r -subset representation is the real vector space

$$V_{n,r} := \mathbb{R}[X_{n,r}] = \bigoplus_{A \in X_{n,r}} \mathbb{R}e_A,$$

equipped with the inner product for which the basis $\{e_A : A \in X_{n,r}\}$ is orthonormal. The action of S_n is defined by $g \cdot e_A = e_{gA}$, where $gA = \{g(a) : a \in A\}$.

The space $V_{n,0} \cong \mathbb{R}$ is the trivial representation. Intuitively, $V_{n,1}$ consists of vertex features, $V_{n,2}$ of features on unordered pairs of distinct vertices, and $V_{n,3}$ of features on unordered triples. Pair features may be viewed as real edge weights. Triple features are assigned to all triples, whether or not those triples form triangles in a particular graph. Throughout, these spaces contain arbitrary real-valued features on every subset of the indicated size.

Fix an input subset size $r \geq 1$, an output subset size $s \geq 0$, and a polynomial degree $d \geq 0$. Our objective is to construct an explicit basis of

$$\text{Poly}_d(V_{n,r}, V_{n,s})^{S_n} \cong \text{Hom}_{S_n}(\text{Sym}^d V_{n,r}, V_{n,s}).$$

An element of this space has the form

$$P(x) = \sum_{Q \in X_{n,s}} P_Q(x) e_Q,$$

where each P_Q is a homogeneous polynomial of degree d in the input coordinates x_A . Equivariance means $P(g \cdot x) = g \cdot P(x)$, or equivalently, $P_{gQ}(g \cdot x) = P_Q(x)$ for $g \in S_n$, $Q \in X_{n,s}$.

A degree- d input monomial $x_{A_1} \cdots x_{A_d}$ records d input subsets, with repetitions allowed. An output coordinate also specifies an s -subset Q . The basis constructed below is indexed by these configurations up to simultaneous relabelling of the vertices. Each basis map sums the monomials belonging to one such configuration type.

6.1. Symmetric powers are multihypergraph modules. To describe symmetric powers combinatorially, for a finite G -set X and $d \geq 0$, let $\text{MSet}_d(X)$ denote the size- d multisets on X with the induced action.

Proposition 6.2. *For a finite group G and a finite G -set X ,*

$$\text{Sym}^d(\mathbb{R}[X]) \cong \mathbb{R}[\text{MSet}_d(X)]$$

as G -representations. If m_ω represents an orbit $\omega \in \text{MSet}_d(X)/G$ and $H_\omega = \text{Stab}_G(m_\omega)$, then

$$\text{Sym}^d(\mathbb{R}[X]) \cong \bigoplus_{\omega \in \text{MSet}_d(X)/G} \text{Ind}_{H_\omega}^G \mathbf{1}.$$

Proof. The monomials $e^m = \prod_{x \in X} e_x^{m(x)}$ for $m \in \text{MSet}_d(X)$ form a basis of $\text{Sym}^d(\mathbb{R}[X])$, and $ge^m = e^{gm}$. Each orbit therefore spans a permutation module isomorphic to $\mathbb{R}[G/H_\omega] = \text{Ind}_{H_\omega}^G \mathbf{1}$. $\square$

Definition 6.3. An r -uniform multihypergraph on $[n]$ with d edges is a multiset

$$\mathcal{H} = \{E_1, \dots, E_d\}, \quad E_i \in X_{n,r},$$

where repeated edges are allowed.

- A vertex is *active* if it lies in at least one edge of $\mathcal{H}$. The subhypergraph obtained by deleting all isolated vertices is called the *active core*.
- Two multihypergraphs are *isomorphic* if there is a bijection between their vertex sets that preserves edge multiplicities. We write $\text{Aut}(\mathcal{H})$ for the automorphism group of $\mathcal{H}$.
- Let $\mathfrak{H}_{r,d}$ denote the set of isomorphism classes of r -uniform multihypergraphs with d edges and no isolated vertices. For $\mathcal{H} \in \mathfrak{H}_{r,d}$, write $v(\mathcal{H})$ for its number of vertices.

Take $X = X_{n,r}$. A monomial $e_{A_1} \cdots e_{A_d}$ corresponds to the edge multiset $\{A_1, \dots, A_d\}$, and two monomials lie in the same S_n -orbit exactly when their active cores are isomorphic. Every core satisfies $v(\mathcal{H}) \leq rd$, with equality precisely when its edges are pairwise disjoint.

For $v(\mathcal{H}) \leq n$, let $P_{\mathcal{H}}(n)$ denote the span of all monomials whose active core is isomorphic to $\mathcal{H}$.

The active core is exactly the orbit data, so it gives the desired decomposition of the symmetric power.

Theorem 6.4. *For $n \geq 0$, $r \geq 1$ and $d \geq 0$,*

$$\text{Sym}^d V_{n,r} = \bigoplus_{\substack{[\mathcal{H}] \in \mathfrak{H}_{r,d} \\ v(\mathcal{H}) \leq n}} P_{\mathcal{H}}(n), \quad \text{where} \quad P_{\mathcal{H}}(n) \cong \text{Ind}_{\text{Aut}(\mathcal{H}) \times S_{n-v(\mathcal{H})}}^{S_n} \mathbf{1}.$$

The subgroup acts by automorphisms on a labelled copy of the core and by arbitrary permutations on its complement. For $n \geq rd$, every core type occurs.

For each partition $\lambda \vdash n$, the multiplicity of the irreducible Specht module S^λ is

$$[P_{\mathcal{H}}(n) : S^\lambda] = \dim(S^\lambda)^{\text{Aut}(\mathcal{H}) \times S_{n-v(\mathcal{H})}}.$$

Proof. See Section A.3. $\square$

This is a decomposition into orbit spans, not generally into irreducible representations. Each summand records one overlap pattern among the input subsets. Importantly, the indexing set stabilises for $n \geq rd$, although the modules $P_{\mathcal{H}}(n)$ themselves still depend on n .

Example 6.5. For $r = 2$, $d = 2$ and $n \geq 4$, there are three core types: a repeated edge, two adjacent edges and two disjoint edges. Writing $e_{ab} = e_{\{a,b\}}$, the representatives are e_{12}^2 , $e_{12}e_{23}$ and $e_{12}e_{34}$, respectively. Hence

$$\text{Sym}^2 V_{n,2} = P_{\text{rep}}(n) \oplus P_{\text{adj}}(n) \oplus P_{\text{disj}}(n).$$

For general r , the orbit of $e_A e_B$ is determined by $t = |A \cap B|$, where $\max(0, 2r - n) \leq t \leq r$. If $t < r$, its stabiliser is $S_t \times (S_{r-t} \wr S_2) \times S_{n-2r+t}$. The factors permute the intersection, the two exclusive parts (allowing their exchange), and the unused vertices. If $t = r$, the edge is repeated and the stabiliser is $S_r \times S_{n-r}$.

Symmetric powers of permutation modules have also been studied in the modular setting [Jia21]. Here the orbit description is useful because it keeps the overlap data visible without first decomposing into irreducibles. For example, at $n = rd$ the disjoint-edge summand is the Foulkes module $\text{Ind}_{S_r \wr S_d}^{S_{rd}} \mathbf{1}$, whose general irreducible decomposition is an open problem [PW16].

6.2. Polynomial equivariants: the marked-orbit basis. The marked-orbit basis describes all homogeneous polynomial maps from r -subset to s -subset features. We begin with the classical linear case.

Proposition 6.6. *For finite G -sets X, Y we have $\text{Hom}_G(\mathbb{R}[X], \mathbb{R}[Y]) \cong \mathbb{R}[X \times Y]^G$, with basis the maps*

$$T_{\mathcal{O}}(e_x) = \sum_{y:(x,y) \in \mathcal{O}} e_y$$

indexed by the G -orbits $\mathcal{O} \subseteq X \times Y$.

Proof. A linear map T with matrix $k_T(x, y)$ is equivariant if and only if $k_T(gx, gy) = k_T(x, y)$, that is, if and only if k_T is constant on diagonal orbits. See also [GTW25, Lemma 2E.3] and [MBHSL19]. $\square$

By Proposition 6.2 we may apply this with $X = \text{MSet}_d(X_{n,r})$, which leads to the following description of the orbits.

Definition 6.7. A *marked degree- d (r, s) -interaction* is a pair $(\mathcal{A}, Q)$ with $\mathcal{A} \in \text{MSet}_d(X_{n,r})$ and $Q \in X_{n,s}$, that is, an r -uniform d -edge multihypergraph on $[n]$ together with a distinguished s -subset of vertices. Its *type* is its orbit under simultaneous relabelling. Marked vertices need not be active.

Theorem 6.8. $\text{Poly}_d(V_{n,r}, V_{n,s})^{S_n} \cong \text{Hom}_{S_n}(\text{Sym}^d V_{n,r}, V_{n,s})$, with a basis indexed by the diagonal S_n -orbits $\mathcal{O} \subseteq \text{MSet}_d(X_{n,r}) \times X_{n,s}$. For such an orbit, define

$$\Psi_{\mathcal{O}}(x) := \sum_{Q \in X_{n,s}} \left(\sum_{\mathcal{A}: (\mathcal{A}, Q) \in \mathcal{O}} x^{\mathcal{A}} \right) e_Q, \quad x^{\mathcal{A}} := \prod_{A \in X_{n,r}} x_A^{m_{\mathcal{A}}(A)}.$$

Then the maps $\Psi_{\mathcal{O}}$ form a basis of $\text{Poly}_d(V_{n,r}, V_{n,s})^{S_n}$.

Proof. By Proposition 6.2, a degree- d monomial is indexed by a multiset $\mathcal{A} \in \text{MSet}_d(X_{n,r})$. Hence a polynomial map $V_{n,r} \rightarrow V_{n,s}$ is determined by coefficients $c_{\mathcal{A}, Q}$ indexed by $(\mathcal{A}, Q) \in \text{MSet}_d(X_{n,r}) \times X_{n,s}$. By Proposition 6.6, such a map is S_n -equivariant exactly when these coefficients are constant on diagonal S_n -orbits. The corresponding orbit indicators are precisely the maps $\Psi_{\mathcal{O}}$. $\square$

Under the polarisation identification of Proposition 1.1, the usual multinomial factors only rescale the multiset basis vectors. These factors are constant on S_n -orbits, so they do not change the orbit indexing or the basis statement.

Orbit-indexed graph-polynomial bases also appear in [PLK⁺23]. The orbit basis is explicit, but we also want its dimension and stable range. Fixing the active core reduces the count to the possible positions of the output mark.

Proposition 6.9. *Fix an active core $\mathcal{H}$ with vertex set $U = V(\mathcal{H})$, and write $v = v(\mathcal{H})$. A marked type with core $\mathcal{H}$ is determined by $q := |Q \cap U|$ together with the $\text{Aut}(\mathcal{H})$ -orbit of the active part $Q \cap U \in \binom{U}{q}$. Consequently,*

$$\dim \text{Hom}_{S_n}(\text{Sym}^d V_{n,r}, V_{n,s}) = \sum_{\substack{[\mathcal{H}] \in \mathfrak{H}_{r,d} \\ v(\mathcal{H}) \leq n}} \sum_{q=\max(0, s-(n-v(\mathcal{H})))}^{\min(s, v(\mathcal{H}))} \left| \binom{V(\mathcal{H})}{q} / \text{Aut}(\mathcal{H}) \right|. \quad (7)$$

Proof. By Theorem 6.4 and Frobenius reciprocity, the contribution of a fixed core $\mathcal{H}$ is $\dim(V_{n,s})^{\text{Aut}(\mathcal{H}) \times S_{n-v}}$. Since $V_{n,s} = \mathbb{R}[X_{n,s}]$ is a permutation representation, this dimension is the number of $(\text{Aut}(\mathcal{H}) \times S_{n-v})$ -orbits on s -subsets $Q \subseteq [n]$.

Write $q = |Q \cap U|$. $\text{Aut}(\mathcal{H})$ acts on the active part $Q \cap U$, while S_{n-v} is transitive on all $(s-q)$ -subsets of the inactive vertices. Thus, for fixed q , the orbit of Q is determined exactly by the A -orbit of $Q \cap U \in \binom{U}{q}$.

Such a marking exists precisely when $0 \leq q \leq v$ and $0 \leq s-q \leq n-v$, giving the stated limits. Summing over the core types proves (7). $\square$

By Burnside's lemma, evaluating (7) requires only the cycle structures of the automorphism groups. Theorem C.2 computes the same number by summing over conjugacy classes of S_n instead.

The counting formula also shows exactly when increasing n can no longer create new marked types.

Theorem 6.10. *Let $r, d \geq 1$ and $s \geq 0$, and write $D(n) := \dim \text{Hom}_{S_n}(\text{Sym}^d V_{n,r}, V_{n,s})$. Then:*

- (a) $D(n)$ is non-decreasing in n .
- (b) For $n \geq rd + s$,

$$D(n) = \sum_{[\mathcal{H}] \in \mathfrak{H}_{r,d}} \sum_{q=0}^{\min(s, v(\mathcal{H}))} \left| \binom{V(\mathcal{H})}{q} / \text{Aut}(\mathcal{H}) \right|,$$

so $D(n)$ is independent of n in this range.

- (c) The threshold is sharp: $D(rd + s - 1) < D(rd + s)$.

In the stable range, the basis $\{\Psi_{\mathcal{O}}\}$ is indexed by a set independent of n , so one coefficient vector defines one equivariant polynomial layer simultaneously for every $n \geq rd + s$.

Proof. In (7) the set of admissible types grows with n and each lower summation limit is non-increasing in n , so D is non-decreasing. For $n \geq rd + s$ we have $n - v \geq s$ for every core, so every lower limit is 0 and no term depends on n , which gives (b).

For sharpness, let $\mathcal{H}_0$ be the disjoint type, with $v(\mathcal{H}_0) = rd$. At $n = rd + s - 1$ the value $q = 0$ would require $s - q \leq n - v = s - 1$, which fails, so $q = 0$ is inadmissible for $\mathcal{H}_0$. Also when $s = 0$ the core $\mathcal{H}_0$ does not embed at all. At $n = rd + s$ it becomes admissible and contributes exactly one further orbit, while no other term decreases. Hence the strict inequality. $\square$

6.3. Worked example: seven quadratic edge-to-vertex maps. Take $(r, d, s) = (2, 2, 1)$ and write $x_{uv} = x_{\{u,v\}}$. By Theorem 6.8 we must list the marked types: a repeated edge with the mark on it or off it; an adjacent pair with the mark at the centre, at an endpoint, or outside; a disjoint pair with the mark on an active vertex or outside. These give seven types (figure 1), hence $\dim \text{Hom}_{S_n}(\text{Sym}^2 V_{n,2}, V_{n,1}) = 7$ for $n \geq 5$. At $n = 3$ only four types fit, while at $n = 4$ only the disjoint pair with an outside mark is absent. Thus the dimensions are 4, 6, 7, 7, ... for $n = 3, 4, 5, 6, \dots$




FIGURE 1. The seven marked interaction types for quadratic maps $V_{n,2} \rightarrow V_{n,1}$ in the stable range. Filled vertices carry the output mark v ; double lines denote a repeated input edge.

Writing the v th output coordinate gives the following basis, using the same labels as figure 1.

$$\begin{aligned}
 (\Phi_1 x)_v &= \sum_{u \neq v} x_{uv}^2, & \text{repeated edge, mark on,} \\
 (\Phi_2 x)_v &= \sum_{\{i,j\} \not\ni v} x_{ij}^2, & \text{repeated edge, mark off,} \\
 (\Phi_3 x)_v &= \sum_{\{a,c\} \subseteq [n] \setminus \{v\}} x_{av} x_{vc}, & \text{adjacent edges, centre,} \\
 (\Phi_4 x)_v &= \sum_{u \neq v} \sum_{w \notin \{u,v\}} x_{vu} x_{uw}, & \text{adjacent edges, endpoint,} \\
 (\Phi_5 x)_v &= \sum_{b \neq v} \sum_{\{a,c\} \subseteq [n] \setminus \{v,b\}} x_{ab} x_{bc}, & \text{adjacent edges, outside,} \\
 (\Phi_6 x)_v &= \sum_{u \neq v} \sum_{\{a,b\} \subseteq [n] \setminus \{v,u\}} x_{vu} x_{ab}, & \text{disjoint edges, active,} \\
 (\Phi_7 x)_v &= \sum_{\substack{\{E_1, E_2\} \\ E_1 \cap E_2 = \emptyset \\ v \notin E_1 \cup E_2}} x_{E_1} x_{E_2}, & \text{disjoint edges, outside.}
 \end{aligned}$$

In the last sum, E_1 and E_2 are two-element subsets of $[n]$ and their pair is unordered. Each map is an orbit sum of one marked type. Their monomial supports are disjoint, so the maps are independent and the dimension count shows that they form a basis.

The basis also shows the geometry directly. Φ_1 is a local square statistic at v , while Φ_3 and Φ_4 are quadratic contractions along length-two paths through or ending at v . The other four use configurations that are not local path contractions at v .

6.4. Evaluation of the seven quadratic operations. For implementation, the seven orbit sums can be evaluated without enumerating triples, quadruples or five-tuples. Write $X_{uv} = x_{uv}$ for $u \neq v$ and $X_{vv} = 0$, and define

$$h = X\mathbf{1}, \quad q_v = \sum_{u \neq v} x_{uv}^2, \quad S = \sum_{u < v} x_{uv}, \quad T = \sum_{u < v} x_{uv}^2.$$

Let

$$W = \frac{1}{2} \sum_v (h_v^2 - q_v), \quad D_{\text{disj}} = \frac{1}{2} (S^2 - T) - W.$$

These are the total adjacent-pair and disjoint-pair products, respectively. All quantities refer to scalar, real-valued edge features.

Proposition 6.11. *The seven basis maps of Section 6.3 satisfy*

$$\begin{aligned} (\Phi_1 x)_v &= q_v, & (\Phi_2 x)_v &= T - q_v, \\ (\Phi_3 x)_v &= \frac{1}{2}(h_v^2 - q_v), & (\Phi_4 x)_v &= (Xh)_v - q_v, \\ (\Phi_5 x)_v &= W - (\Phi_3 x)_v - (\Phi_4 x)_v, \\ (\Phi_6 x)_v &= Sh_v - h_v^2 - (Xh)_v + q_v, & (\Phi_7 x)_v &= D_{\text{disj}} - (\Phi_6 x)_v. \end{aligned}$$

For $n \geq 5$, every homogeneous quadratic equivariant map $V_{n,2} \rightarrow V_{n,1}$ is therefore a unique linear combination of these seven maps and can be evaluated in $O(n^2)$ arithmetic operations. With m nonzero edge features supplied sparsely, the cost is $O(n + m)$.

Proof. Expanding h_v^2 separates repeated edges from unordered pairs of distinct edges incident to v , giving Φ_3 . Expanding $(Xh)_v$ separates the returning path v, u, v , which contributes q_v , from paths with three distinct vertices, giving Φ_4 . Every adjacent edge pair places v at its centre, at an endpoint, or outside. Subtracting the first two contributions from W gives Φ_5 .

For a fixed edge $\{v, u\}$, the sum of features on edges disjoint from it is $S - h_v - h_u + x_{vu}$. Multiplying by x_{vu} and summing over u gives Φ_6 . The total product over unordered distinct edge pairs is $(S^2 - T)/2$. Subtract adjacent pairs to obtain D_{disj} , then subtract those containing v to obtain Φ_7 . The first two identities are immediate from the definitions. Only edge sums, the two matrix–vector products $X\mathbf{1}$ and Xh , and vertexwise arithmetic are required. Completeness follows from Theorem 6.8. $\square$

For dense inputs, the $O(n^2)$ cost is of the same order as reading the $\binom{n}{2}$ edge coordinates; for sparse inputs, the $O(n + m)$ bound is linear in the stored input size up to the vertex term.

An explicit quadratic layer is

$$F_\theta(x) = \sum_{a=1}^7 \theta_a \Phi_a(x).$$

Together with the equivariant linear terms and constant output, this spans all equivariant polynomial maps of degree at most two. The seven-dimensional statement concerns the quadratic polynomial space on the full real feature space. The quadratic products in Φ_a are explicit operations of the architecture, so this is not a claim that finite ReLU networks can exactly realise quadratic functions on $\mathbb{R}^{\binom{n}{2}}$. If inputs are restricted to Boolean adjacency matrices, additional identities such as $x_{uv}^2 = x_{uv}$ may introduce dependencies between polynomial expressions.

7. SCOPE AND FUTURE DIRECTIONS

Limitations. It remains open to characterise which sequences of chaos components come from equivariant piecewise-linear maps; see Corollary 2.2. The transform determines each such map uniquely, but equivariance and square-summability of the components do not ensure that the resulting map is piecewise linear. A complete structural picture is therefore still missing. Matching finitely many components with shifted ReLUs also leaves the higher-degree tail uncontrolled, so it gives no approximation bound on its own. For compact connected groups, polynomial existence does not suffice to detect piecewise-linear existence; the additional restrictions are discussed in the introduction.

At the architectural level, the Gram criterion describes the space available in each degree for fixed lifts and readouts. A characterisation of networks with prescribed width and depth, allowing these maps to be trained, remains open here. It must account for shared

parameters across degrees and for interactions created by composition and joint processing of source components.

The general spanning construction can also be costly: each regular hidden block has $|G|$ coordinates. For subset features, the multihypergraph orbit modules can still be reducible, so a full irreducible decomposition requires further work. Restricting the polynomial bases to Boolean inputs can introduce additional relations. Finally, the expressivity results do not establish guarantees about training or generalisation.

Constructing architectures. The main practical task is to turn these constructions into trainable networks with prescribed degree spaces. Choose source and target modules L, K , a degree budget D , and equivariant lifts $A_h : L \rightarrow H_h$ and readouts $R_h : H_h \rightarrow K$, where each H_h is a transitive permutation module. Use the Gram test in Proposition 5.1 to select coordinate polynomials $R_h((A_h x)^{\odot k})$ spanning the intended spaces, and choose activations with nonzero coefficients at the required degrees and scales. Independent branch weights then parametrise the computed span in each degree. For separate degree control, Section 5 provides coordinatewise powers for homogeneous polynomial layers, normalised Hermite activations for pure chaos layers, and shifted-ReLU combinations for matching coefficients through D . The polynomial constructions need not be piecewise linear, while finite ReLU coefficient matching leaves an uncontrolled higher-degree tail.

Graph networks. The companion paper [Sha26] applies this theory to predicting graph diameter and algebraic connectivity. The reported gains over a parameter-matched control with full linear S_n -equivariant mixing concern one synthetic benchmark with a frozen GIN backbone. The subset representations studied here place this construction within a broader family of vertex, edge and higher-order features. In each degree, the marked-orbit basis gives all polynomial maps between these spaces, guiding layer construction and tests of which directions a proposed layer spans (Theorem 6.8). Its stable indexing allows coefficients to be shared across graph orders (Theorem 6.10).

A further direction is to adapt the GNN architecture to classify d -manifolds from their triangulations. Equivariance under vertex relabelling alone is insufficient: subdivisions and retriangulations, such as Pachner moves, can change both the vertex set and the incidence structure. Extending the theory to this setting requires specifying how features are transferred between triangulations and how network layers respect these transfers, with the aim of making the final prediction invariant under the chosen changes of triangulation.

APPENDIX A. SUPPLEMENTARY PROOFS

This appendix supplies the polarisation and character-theoretic arguments used in the main text.

A.1. Proof of Proposition 1.1.

Proof. Polarisation identifies a homogeneous polynomial map P of degree k with a symmetric k -linear map $\tilde{P}$ satisfying $P(x) = \tilde{P}(x, \dots, x)$, and the universal property of $\text{Sym}^k L$ converts $\tilde{P}$ into a unique linear $T_P : \text{Sym}^k L \rightarrow K$ with $T_P(x^k) = P(x)$. The two constructions are mutually inverse and natural in L and K , so $P(gx) = gP(x)$ for all g and x if and only if T_P commutes with the G -action. $\square$

A.2. Proof of Proposition 2.8.

Proof. Diagonalising $\rho_L(g)$ over $\mathbb{C}$ gives $\sum_k \chi_{\text{Sym}^k L}(g)t^k = \det(I - t\rho_L(g))^{-1}$, and taking logarithms,

$$\log \det(I - t\rho_L(g))^{-1} = \sum_{m \geq 1} \frac{t^m}{m} \text{tr } \rho_L(g^m) = \sum_{m \geq 1} \frac{t^m}{m} \chi_L(g^m).$$

Real characters are self-conjugate, so $c_k = \langle \chi_{\text{Sym}^k L}, \chi_K \rangle$, and averaging over G gives both displayed forms. $\square$

A.3. Proof of Theorem 6.4.

Proof. A basis of $\text{Sym}^d V_{n,r}$ is given by the monomials

$$e^m = \prod_{A \in X_{n,r}} e_A^{m(A)}, \quad m : X_{n,r} \rightarrow \mathbb{N}_0, \quad \sum_A m(A) = d.$$

The action of S_n is

$$g \cdot e^m = e^{gm}, \quad (gm)(A) = m(g^{-1}A).$$

Thus each monomial may be identified with an r -uniform multihypergraph on $[n]$ having $m(A)$ copies of the edge A .

A permutation g maps the active core of m isomorphically onto that of gm . Conversely, any isomorphism of active cores extends arbitrarily to a permutation of their complements in $[n]$. Hence two monomials lie in the same S_n -orbit if and only if their active cores are isomorphic, and

$$\text{Sym}^d V_{n,r} = \bigoplus_{\substack{[\mathcal{H}] \in \mathfrak{H}_{r,d} \\ v(\mathcal{H}) \leq n}} P_{\mathcal{H}}(n).$$

Fix a monomial with active core $\mathcal{H}$, active vertex set U , and $v = |U|$. Its stabiliser consists precisely of automorphisms of the core together with arbitrary permutations of the inactive vertices. Therefore $\text{Stab}_{S_n}(m) \cong \text{Aut}(\mathcal{H}) \times S_{n-v}$, and the corresponding orbit module is the associated permutation representation, $P_{\mathcal{H}}(n) \cong \text{Ind}_{\text{Aut}(\mathcal{H}) \times S_{n-v}}^{S_n} \mathbf{1}$.

Since $\mathcal{H}$ has d edges, each of size r , $v(\mathcal{H}) \leq rd$. Thus every core type occurs once $n \geq rd$, so the indexing set is independent of n in this range.

Finally, for $\lambda \vdash n$, absolute irreducibility of the real Specht modules and Frobenius reciprocity give

$$\begin{aligned} [P_{\mathcal{H}}(n) : S^\lambda] &= \dim \text{Hom}_{S_n}(P_{\mathcal{H}}(n), S^\lambda) \\ &= \dim \text{Hom}_{\text{Aut}(\mathcal{H}) \times S_{n-v}}(\mathbf{1}, \text{Res } S^\lambda) \\ &= \dim (S^\lambda)^{\text{Aut}(\mathcal{H}) \times S_{n-v}}. \end{aligned}$$

This proves the result. $\square$

APPENDIX B. HIDDEN SPACES WITH SEVERAL COORDINATE ORBITS

The common scalar in Theorem 3.6 requires equal coordinate variances. With several coordinate orbits there is still an exact formula, but the scalar becomes a diagonal matrix.

Let $A : L \rightarrow \mathbb{R}^{\mathcal{B}}$ and $R : \mathbb{R}^{\mathcal{B}} \rightarrow K$ be equivariant linear maps for an arbitrary permutation basis $\mathcal{B}$. Put $C = AA^*$ and $s_b = \sqrt{C_{bb}}$. Assume $\sigma(s_b \cdot) \in L^2(\gamma_{\mathbb{R}})$ whenever $s_b > 0$. Let D_k be diagonal with

$$(D_k)_{bb} = \begin{cases} a_k(\sigma; s_b)/s_b^k, & s_b > 0, \\ \sigma(0), & s_b = 0, \ k = 0, \\ 0, & s_b = 0, \ k > 0. \end{cases}$$

Each s_b , and hence each diagonal entry, is constant on a coordinate orbit. Therefore D_k is equivariant.

Proposition B.1. For $F(x) = R\sigma_{\mathcal{B}}(Ax)$,

$$\Pi_k F = \Pi_k [RD_k((Ax)^{\odot k})], \quad \|\Pi_k F\|_{L^2}^2 = k! \text{tr}(RD_k C^{\odot k} D_k R^*).$$

Proof. Expand each coordinate using its own variance. If $s_b = 0$, then $A^*b = 0$ and that coordinate is the constant $\sigma(0)$, contributing only to degree zero. For positive variances, the identity from Section 1.3 gives $\Pi_k \langle x, A^*b \rangle^k = s_b^k \text{He}_k(\langle x, A^*b \rangle / s_b)$. By Lemma 1.2, the corresponding cross inner product is $k!C_{bb'}^k$, including degree zero with $C^{00} = J$. Summing gives both assertions. $\square$

Different orbit contributions can cancel after the readout. Thus separate nonzero tests for the orbits do not replace this combined formula. In particular, several feature channels in a hidden space do not require a new theory, but do require retaining their possibly different variances and cross terms.

APPENDIX C. REPRESENTATION FORMULAE FOR SUBSET MODULES

The Specht modules S^λ , $\lambda \vdash n$, are the irreducible real representations of S_n . They are absolutely irreducible, so character inner products below are ordinary multiplicities. For the two-row formula assume $0 \leq r \leq n/2$ and recall that complementation gives $V_{n,r} \cong V_{n,n-r}$ for the remaining nonzero cases.

C.1. The subset module and its Johnson scheme. We finish with formulae specific to subset modules. They support the multiplicity-free and computational statements used in the main text.

Proposition C.1. $V_{n,r} \cong \text{Ind}_{S_r \times S_{n-r}}^{S_n} \mathbf{1}$, and for $r \leq n/2$

$$V_{n,r} \cong \bigoplus_{j=0}^r S^{(n-j,j)}.$$

In particular $V_{n,r}$ is multiplicity-free, and $\chi^{(n-j,j)}(g) = |X_{n,j}^g| - |X_{n,j-1}^g|$ for $0 \leq j \leq n/2$, with $X_{n,-1} = \emptyset$.

Proof. The action of S_n on r -subsets is transitive, with stabiliser $S_r \times S_{n-r}$, so $V_{n,r} \cong \text{Ind}_{S_r \times S_{n-r}}^{S_n} \mathbf{1}$. The decomposition of this Young permutation module is multiplicity-free and standard. See [Jam78, Sag01].

Finally, since $\chi_{V_{n,j}}(g) = |X_{n,j}^g|$ and $V_{n,j} \cong \bigoplus_{i=0}^j S^{(n-i,i)}$, subtracting the decompositions for j and $j-1$ gives

$$\chi^{(n-j,j)}(g) = |X_{n,j}^g| - |X_{n,j-1}^g|.$$

$\square$

Thus $V_{n,r}$ is transitive and multiplicity-free, and Section 3.3 applies without modification. Here the orbit algebra is the Bose–Mesner algebra of the Johnson scheme $J(n, r)$. Its orbit indicator matrices are indexed by the intersection number $|A \cap A'|$, and the primitive idempotents E_i have entries depending only on that number, given by the dual eigenvalues (Q -numbers) of the scheme in the standard normalisation [BCN89, Section 9.1]. The Hadamard coefficients of Corollary 3.11 are therefore obtained from the Johnson Krein parameters in that normalisation.

C.2. A character formula. Theorem 6.4 describes the orbit geometry. For dimension and multiplicity calculations, Pólya theory gives a complementary formula that avoids enumerating hypergraphs up to isomorphism.

Theorem C.2 (Molien–Pólya formula). *For $g \in S_n$ let*

$$f_r(g) := |X_{n,r}^g| = [z^r] \prod_{c \in \text{Cyc}(g)} (1 + z^{|c|}),$$

the product being over cycles of g on $[n]$, and define $h_d(g)$ by

$$\sum_{d \geq 0} h_d(g) t^d = \exp\left(\sum_{m \geq 1} \frac{t^m}{m} f_r(g^m)\right), \quad \text{equivalently} \quad d h_d(g) = \sum_{m=1}^d f_r(g^m) h_{d-m}(g).$$

Then for every $\lambda \vdash n$ and $0 \leq s \leq n$,

$$[\mathrm{Sym}^d V_{n,r} : S^\lambda] = \frac{1}{n!} \sum_{g \in S_n} h_d(g) \chi^\lambda(g),$$

$$\dim \mathrm{Hom}_{S_n}(\mathrm{Sym}^d V_{n,r}, V_{n,s}) = \frac{1}{n!} \sum_{g \in S_n} h_d(g) f_s(g).$$

Both sums depend on g only through its cycle type.

Proof. A subset is fixed by g if and only if it is a union of cycles of g , which gives the generating-function expression for f_r .

By Proposition 6.2, $\chi_{\mathrm{Sym}^d V_{n,r}}(g)$ is the number of g -invariant multisets of size d on $X_{n,r}$, that is, of multiplicity functions constant on the cycles of g acting on $X_{n,r}$. Hence

$$\sum_{d \geq 0} \chi_{\mathrm{Sym}^d V_{n,r}}(g) t^d = \prod_{c \in \mathrm{Cyc}(g \curvearrowright X_{n,r})} (1 - t^{|c|})^{-1},$$

and taking logarithms gives $\sum_{m \geq 1} \frac{t^m}{m} |X_{n,r}^{g^m}|$, since a point of $X_{n,r}$ is fixed by g^m exactly when its cycle length divides m . As $|X_{n,r}^{g^m}| = f_r(g^m)$, this shows $\chi_{\mathrm{Sym}^d V_{n,r}}(g) = h_d(g)$, which is Proposition 2.8 for the permutation representation $V_{n,r}$. The second formula is the first with $\chi_{V_{n,s}} = f_s$. $\square$

The recursion for h_d gives a direct computational procedure.

C.3. Support and stability. This supplementary subsection records how the irreducible support behaves as n grows; it is not needed for the coordinatewise layer calculations above.

Theorem C.3. (a) *If S^λ occurs in $P_{\mathcal{H}}(n)$ then $\lambda_1 \geq n - v(\mathcal{H})$. Hence every constituent of $\mathrm{Sym}^d V_{n,r}$ satisfies $n - \lambda_1 \leq rd$.*

(b) *Fix r, d and a partition $\nu \vdash s$, and put $\lambda[n] = (n - s, \nu)$. For n such that $\lambda[n]$ is a partition, $m_\nu(n) := [\mathrm{Sym}^d V_{n,r} : S^{\lambda[n]}]$ vanishes for all n if $s > rd$, and is independent of n for $n \geq s + rd$ if $s \leq rd$.*

Proof. (a) Since $\mathrm{Aut}(\mathcal{H}) \times S_{n-v} \supseteq S_{n-v}$ we get $[P_{\mathcal{H}}(n) : S^\lambda] \leq \dim(S^\lambda)^{S_{n-v}}$, and iterated branching gives $(S^\lambda)^{S_m} \neq 0$ if and only if $\lambda_1 \geq m$.

(b) The first claim is (a). For the second, take $n \geq rd$ so that the type set is fixed. Write $\mathrm{Ind}_{\mathrm{Aut}(\mathcal{H})}^{S_v} \mathbf{1} = \bigoplus_\mu c_\mu S^\mu$ and induce in stages; the contribution of $\mathcal{H}$ to the multiplicity of $S^{\lambda[n]}$ is $\sum_\mu c_\mu c_{\mu, (n-v)}^{\lambda[n]}$, and by Pieri's rule the Littlewood–Richardson coefficient $c_{\mu, (n-v)}^{\lambda[n]}$ is 1 exactly when $\lambda[n]/\mu$ is a horizontal strip, that is when $\lambda[n]_i \geq \mu_i \geq \lambda[n]_{i+1}$ for all i , and 0 otherwise.

Only one of those inequalities involves n , namely $n - s \geq \mu_1$. Since $\mu_1 \leq v \leq rd$, it holds for every μ once $n \geq s + rd$, so the multiplicity is constant from there on. $\square$

In the language of representation stability, the sequence $n \mapsto \mathrm{Sym}^d V_{n,r}$ is a finitely generated FI-module, which also implies eventual constancy [CEF15]. The argument above gives an explicit sufficient range for each padded Specht multiplicity. The sharpness statement in Theorem 6.10 concerns the full space of maps into s -subset features; it does not assert that every individual Specht multiplicity first stabilises at this bound.

REFERENCES

- [Aie26] Valeriano Aiello. Piecewise linear equivariant maps for compact groups. *arXiv preprint arXiv:2608.21645*, 2026.
- [BBCV21] Michael M. Bronstein, Joan Bruna, Taco Cohen, and Petar Veličković. Geometric Deep Learning: Grids, Groups, Graphs, Geodesics, and Gauges. *arXiv preprint arXiv:2104.13478*, 2021.
- [BCN89] Andries E. Brouwer, Arjeh M. Cohen, and Arnold Neumaier. *Distance-Regular Graphs*, volume 18 of *Ergebnisse der Mathematik und ihrer Grenzgebiete*. Springer, 1989.
- [CEF15] Thomas Church, Jordan S. Ellenberg, and Benson Farb. FI-modules and stability for representations of symmetric groups. *Duke Mathematical Journal*, 164(9):1833–1910, 2015.
- [DFS16] Amit Daniely, Roy Frostig, and Yoram Singer. Toward deeper understanding of neural networks: The power of initialization and a dual view on expressivity. In *Advances in Neural Information Processing Systems*, volume 29, pages 2253–2261, 2016.
- [GTW25] Joel Gibson, Daniel Tubbenhauer, and Geordie Williamson. Equivariant neural networks and piecewise linear representation theory. In *Modern Algebra, Volume 1: Representation Theory*, volume 829 of *Contemporary Mathematics*, pages 157–192. American Mathematical Society, 2025.
- [Jam78] Gordon D. James. *The Representation Theory of the Symmetric Groups*, volume 682 of *Lecture Notes in Mathematics*. Springer, 1978.
- [Jan97] Svante Janson. *Gaussian Hilbert Spaces*, volume 129 of *Cambridge Tracts in Mathematics*. Cambridge University Press, 1997.
- [Jia21] Yu Jiang. On the symmetric and exterior powers of young permutation modules. *Journal of Algebra*, 568:660–704, 2021.
- [LD24] Eitan Levin and Mateo Díaz. Any-dimensional equivariant neural networks. In *Proceedings of the 27th International Conference on Artificial Intelligence and Statistics*, volume 238 of *PMLR*, pages 2773–2781, 2024.
- [MBHSL19] Haggai Maron, Heli Ben-Hamu, Nadav Shamir, and Yaron Lipman. Invariant and equivariant graph networks. In *International Conference on Learning Representations*, 2019.
- [PDLS24] Marco Pacini, Xiaowen Dong, Bruno Lepri, and Gabriele Santin. A characterization theorem for equivariant networks with point-wise activations. *arXiv preprint arXiv:2401.09235*, 2024.
- [PLK⁺23] Omri Puny, Derek Lim, Bobak T. Kiani, Haggai Maron, and Yaron Lipman. Equivariant polynomials for graph neural networks. In *Proceedings of the 40th International Conference on Machine Learning*, volume 202 of *PMLR*, pages 28191–28222, 2023.
- [PW16] Rowena Paget and Mark Wildon. Minimal and maximal constituents of twisted foulkes characters. *Journal of the London Mathematical Society*, 93(2):301–318, 2016.
- [Rav20] Siamak Ravanbakhsh. Universal equivariant multilayer perceptrons. In *Proceedings of the 37th International Conference on Machine Learning*, volume 119 of *Proceedings of Machine Learning Research*, pages 7996–8006, 2020.
- [Sag01] Bruce E. Sagan. *The Symmetric Group: Representations, Combinatorial Algorithms, and Symmetric Functions*, volume 203 of *Graduate Texts in Mathematics*. Springer, 2 edition, 2001.
- [Sha26] Mani Shayestehfar. Nonlinear maps between irreducible representations in graph neural networks. In preparation, 2026.
- [Yar22] Dmitry Yarotsky. Universal approximations of invariant maps by neural networks. *Constructive Approximation*, 55:407–474, 2022.

SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF SYDNEY, NSW 2006, AUSTRALIA
 Email address: mani.shayestehfar@sydney.edu.au
 Personal Webpage: <https://maani.info>