BODYinTRANSIT-Lab: A Multimodal Dataset for Investigating Sensory-Driven Transformations in Body Perception and Affective Experiences

Mohammad Mahdi Dehshibi *Senior Member, IEEE*, Tomás Martínez Cortés, Amar D'Adamo, Daniel de la Prida, Luis-Antonio Azpicueta-Ruíz, Laia Turmo Vidal, Marte Roel Lesur, Joaquín R. Díaz-Durán, Alexander Väljamäe, Angel Sánchez, Temitayo Olugbade, Nadia Bianchi-Berthouze, Fernando Díaz-de-María, and Ana Tajadura-Jiménez

arXiv:2611.14093v1 [cs.HC] 10 Nov 2026

Abstract—Multimodal datasets that combine synchronised behavioural and physiological signals with subjective affective and body perception measures are still rare. This limits both the understanding of embodied experience and the development of computational models that infer subjective states from sensor data. We introduce BODYinTRANSIT-Lab, a multimodal dataset that addresses this gap by capturing affective, perceptual, and behavioural responses to the *Footsteps Illusion*. In this closed-loop paradigm, participants perform walking tasks while receiving real-time manipulated auditory feedback of their own footsteps intended to alter perceived body weight. The dataset comprises synchronised recordings from 101 participants recruited through systematic stratification by eating-disorder symptomatology and physical activity level. BODYinTRANSIT-Lab combines full-body kinematics with electromyography, electrocardiography, respiration, and electrodermal activity. Its annotations capture both immediate and post-trial experience, including within-trial perceived-weight ratings, post-trial assessments of body feelings and affective experience, 3D avatar body visualisations, and quantitative body maps. Psychological, behavioural, demographic, and social measures provide further information about factors that may influence individual responses. We provide a deep learning benchmark that uses walking kinematics to classify relative affective valence and perceived weight within subjects. On within-session, segment-level splits, the baseline model achieved mean balanced accuracies of 45.9% for affective valence and 44.5% for perceived weight. The BODYinTRANSIT-Lab dataset provides an experimentally controlled resource for cognitive neuroscience and affective computing research, with potential applications in affect-aware interactive systems, health and well-being technologies, and creative embodied experiences.

Index Terms—Multimodal Dataset, Body Perception, Affective Computing, Human-Computer Interaction, Wearable Sensors, Kinematics, Physiology, Deep Learning, Multisensory Perception, Sound, Body Illusion, Body Transformation Experiences.

1 Introduction

Our perception of our own body shapes how we move, feel, and interact with the world [22]. Because this perception is continuously updated by sensory integration, it can change even when the physical body does not [4, 16]. By deliberately manipulating multisensory feedback, interactive technologies can reshape mental body representations [43, 44, 45, 32, 30] and induce Body Transformation Experiences (BTEs). BTEs involve temporary changes in how individuals perceive their bodies, including appearance, size, weight, ownership, and physical capabilities [64]. These transformations can alter affective experience, cognition, and behaviour, with potential implications for health and well-being [18, 45, 32]. Understanding how these experiences relate to measurable bodily responses, and why those responses differ across individuals, is therefore an important problem for affective computing.

The Footsteps Illusion offers a compelling setting for investigating this relationship during everyday movement. By modifying the frequency spectrum of self-generated footstep sounds in real time, auditory feedback can alter how heavy or light people perceive their bodies to be, alongside changes in emotional state and gait [55]. Walking has a dual role in this paradigm: it generates the sensory feedback that shapes bodily experience, while also providing a behavioural signal through which that experience can be investigated. This action-dependent feedback loop brings body perception, affect and movement into the same experimental setting, making the paradigm particularly suitable

Manuscript received Month Day, 2026; revised Month Day, Year.
This work involved human subjects. Approval of all ethical and experimental procedures and protocols was granted by the Committee of Ethics in Research of the Universidad Carlos III de Madrid.
M.M. Dehshibi is with the School of Computer Science and Creative Technologies, University of the West of England, Bristol, U.K.
T. Martínez, A. D'Adamo, M. Roel Lesur, J. Díaz and A. Tajadura-Jiménez are with the Department of Informática, Universidad Carlos III de Madrid, Leganés, Spain.
A. Sánchez is with the Department of Mathematics, Universidad Carlos III de Madrid, Leganés, Spain.
F. Díaz and L.A. Azpicueta-Ruíz are with the Department of Signal Theory and Communications, Universidad Carlos III de Madrid, Leganés, Spain.
D. de la Prida is with the Architectural Acoustics Research Group, Universidad Politécnica de Madrid, Madrid, Spain.
L. Turmo is with the Department of Media Technology and Interaction Design, KTH Royal Institute of Technology, Stockholm, Sweden.
A. Väljamäe is with the HCI group, School of Digital Technologies, Tallinn University, Estonia.
T. Olugbade is with the School of Engineering & Informatics, University of Sussex, Brighton, UK.
N. Berthouze and A. Tajadura-Jiménez are with the UCL Interaction Centre, University College London, London, UK.

for examining how subjective bodily experience relates to measurable behavioural responses [60, 12].

However, the experience elicited by altered bodily feedback cannot be inferred from the manipulation alone. Responses to the Footsteps Illusion vary with body-related beliefs and aspirations, eating-disorder symptomatology, physical activity, and social context [57, 12, 13]. The same feedback may therefore have different perceptual and affective significance for different people. For affective computing, this raises two connected questions: how do self-reported changes in body perception and affect relate to recorded kinematic and physiological signals, and which individual characteristics are associated with variation in these responses? Addressing these questions requires an experimental record that connects the applied feedback, reported experience, movement, physiology, and participant characteristics.

Affective and gait datasets provide important foundations for modelling emotion from movement, pain-related expression, and psychological characteristics associated with walking [3, 9, 68]. However, computationally investigating BTEs imposes additional requirements on the relationship between experimental design and annotation. Researchers must distinguish the sensory manipulation from the experience reported in response to it, examine within-person responses across repeated exposures, and relate these responses to characteristics that may explain individual variability [25, 41, 17]. Repeated ratings can capture how perceived weight varies within a trial, whereas post-trial reports capture complementary aspects of the experience. Yet existing resources were designed for different scientific objectives and do not integrate these elements within a single experimental framework.

We introduce **BODYinTRANSIT-Lab**, a multimodal dataset designed to support these investigations. It contains recordings from 101 adults, recruited with systematic stratification by eating-disorder symptomatology and physical activity level. Participants completed six counterbalanced walking trials under three auditory-feedback conditions, while full-body kinematics were synchronised with electrocardiography (ECG), electrodermal activity (EDA), electromyography (EMG), acceleration (ACC), and respiration. Subjective annotations include repeated within-trial perceived-weight ratings, post-trial body feelings and affective valence, arousal, and dominance, quantitative body maps, and avatar-based body-size estimates. Psychological, behavioural, demographic, anthropometric, and social measures provide context for investigating individual variability in these responses. BODYinTRANSIT-Lab therefore links controlled sensory manipulation with synchronised movement, physiological recordings, and complementary descriptions of bodily and affective experience. It provides data for studying BTE-based technologies intended to influence body perception, affect, and behaviour. We further describe the data-curation protocol as a basis for developing comparable multimodal datasets for the computational study of BTEs.

To demonstrate the dataset's computational usability, we introduce a sensor-aggregation deep learning architecture that combines pretrained movement representations with attention pooling, and report initial within-session, segment-level reference performance for classifying auditory condition, relative perceived body weight, and relative affective valence from walking kinematics. The benchmark demonstrates one way of studying the relationship between sensed movement and reported experience, while the multimodal recordings and participant-level information support substantially broader analyses beyond the present tasks. The contributions of this work are threefold:

1) A multimodal dataset for investigating sensory-driven body transformation experiences (BTEs) that links predefined, digitally shaped auditory bodily feedback with time-aligned bodily signals, complementary subjective annotations and systematic participant characterisation.
2) A documented experimental and data-curation protocol covering acquisition, synchronisation, preprocessing, and organisation, which provides a basis for developing comparable multimodal datasets for the computational study of BTEs.
3) An initial computational benchmark reporting reference performance for classifying auditory condition and relative perceptual and affective labels from walking kinematics under within-session, segment-level splits.

BODYinTRANSIT-Lab provides a resource for studying how affective and body perception experiences relate to measurable bodily responses during controlled sensory interactions, and this purpose in turn motivates the theoretical and empirical requirements that guide dataset design. The following section elaborates on these requirements and situates them within existing research on sensory-induced BTEs, affective movement datasets, and computational modelling.

2 Background and Related Work

Computational investigation of BTEs requires linking how a BTE is elicited, reported, and expressed in bodily signals. Figure 2 provides a conceptual framework linking sensory inputs, individual factors, body representation, and affective and behavioural responses. These complementary observations provide the basis for computationally characterising and modelling BTEs. We review the theoretical and empirical work that motivates these requirements, followed by the datasets and modelling approaches that provide foundations for BODYinTRANSIT-Lab.

2.1 Multisensory Body Representations, BTEs and Affective Experience

Mental body representations integrate sensory information about the body with knowledge and beliefs about its properties [4, 22]. One influential account distinguishes body image, which concerns how people perceive and evaluate their bodies, from body schema, which concerns how the body is represented for posture and action [22, 16]. This distinction does not fully capture how perceptual, affective, and motor aspects of bodily experience interact [46]. A report of perceived size, a judgement of body satisfaction, and a change in movement may therefore offer complementary perspectives on a bodily transformation without measuring the same aspect of it [48]. One influential account



Fig. 1. Conceptual framework for Body Transformation Experiences (BTEs), linking multisensory input, social and contextual factors, individual characteristics, body representation, and behavioural and affective responses. Arrows depict conceptual relationships and feedback, rather than causal pathways established by the present dataset.

distinguishes body image, which concerns how people perceive and evaluate their bodies, from body schema, which concerns how the body is represented for posture and action [22, 16]. This distinction does not fully capture how perceptual, affective, and motor aspects of bodily experience interact [46]. A report of perceived size, a judgement of body satisfaction, and a change in movement may therefore offer complementary perspectives on a bodily transformation without measuring the same aspect of it [48].

Experimental body illusions show that multisensory stimulation can alter these representations, leading to BTEs. Botvinick and Cohen [6] demonstrated that synchronised visual and tactile stimulation can produce a sense of ownership over a rubber hand. Petkova and Ehrsson [43] extended this work to whole-body ownership, showing that a first-person view of another body, paired with correlated bodily stimulation, can induce an experience of that body as one's own. They also measured skin conductance responses to threats directed at the artificial body, providing physiological evidence alongside subjective reports. Related experiments induced ownership of differently sized bodies and showed corresponding changes in perceived object size and distance [65]. These studies established experimental methods for altering bodily self-perception by integrating sensory signals.

Further work has examined the affective significance of such changes. Preston and Ehrsson [44] induced ownership of slimmer and wider mannequin bodies and measured both perceived size and body satisfaction. Ownership of a slimmer body increased body satisfaction, whereas responses to a wider body were related to eating-disorder psychopathology. Crucianelli et al. [10] investigated a complementary direction and found that slow, pleasant touch enhanced subjective embodiment during the rubber hand illusion without a corresponding increase in proprioceptive drift, i.e., the shift in perceived hand location toward the artificial hand. These studies illustrate a bidirectional relationship: BTEs can influence affect, while the affective qualities of stimulation can influence embodiment. They also show that different measures of embodiment need not change in parallel.

BTEs can therefore involve several interrelated perceptual, affective, behavioural, and social outcomes. Beyond affective responses, illusory body ownership has been shown to influence behaviour, social cognition, and self-identity. Embodying a child-sized body can alter perceived object size and walking patterns [65], while embodying a body of a different gender or ethnicity can shift implicit social biases and aspects of gender identity [32, 54]. Because perceptual transformation alone does not determine affective meaning or behavioural expression, the embodiment literature [49] points to the need for multimodal BTE measurement that jointly captures body-perception reports, affective assessments, movement, and physiological data.

2.2 Auditory Feedback and BTEs During Action

Auditory approaches indicate that BTEs can arise from the sensory consequences of action. Tajadura-Jiménez et al. [62] demonstrated that altered action sounds can recalibrate perceived tactile distance, consistent with the experience of having a longer arm [61]. In a related investigation of whole-body dimensions, Tajadura-Jiménez et al. [58] manipulated the auditory and tactile feedback associated with a dropped object's impact. A longer apparent falling time led participants to report feeling taller and to behave as if their legs were longer. These studies address different bodily properties, yet both show that action-related feedback can influence estimates of the body beyond the manipulated sensory modality. In a related auditory paradigm, playback of participants' own voice and movement sounds from an allocentric perspective elicited a feeling of a self-identified presence outside the body [47].

The Footsteps Illusion applies this principle to walking. Tajadura-Jiménez et al. [55] altered the relative energy in different frequency bands of participants' footstep sounds and assessed perceived body weight, emotional state, and gait. Enhancing high frequencies made participants feel lighter, whereas emphasising low frequencies made them feel heavier, alongside changes in affect and gait. This established a paradigm in which an everyday full-body action generates both the manipulated sensory input and a measurable behavioural response. Clausen et al. [8] further investigated the self-related and social significance of altered footstep sounds, examining changes in perceived gender identity and associations with gender groups. Their work broadens the outcomes relevant to auditory BTEs beyond perceived weight alone.

This paradigm distinguishes the delivered feedback from the resulting experience. The auditory condition identifies the manipulation, while subjective reports describe its perceptual and affective consequences. Preserving both enables separate analysis of information associated with the experimental manipulation and information associated with participants' reported experience.

2.3 Individual Differences in Responses to BTEs

The consequences of a BTE depend partly on the person experiencing it. In the Footsteps Illusion, Tajadura-Jiménez et al. [60] investigated the use of altered walking sounds to support physical activity and found that responses interacted with participants' body weight and masculinity–femininity aspirations. This links sensory feedback effects to the relationship between one's current and desired body. It

also indicates why a transformation towards feeling lighter or heavier need not carry the same affective significance for every participant.

Eating-disorder symptomatology provides another source of variation. Tajadura-Jiménez et al. [57] examined the Footsteps Illusion in participants with differing symptom levels and in women with anorexia nervosa. Participants with high symptomatology or anorexia nervosa showed gait and body-visualisation patterns in the light-footstep condition that differed from those of participants with low symptomatology. The self-report measures did not show the same group differences. This result matters for both participant characterisation and annotation design, because variation in one response measure does not necessarily imply equivalent variation in the others.

D'Adamo et al. [12] investigated several individual factors simultaneously using SoniWeight Shoes. Their study examined eating-disorder symptomatology, physical activity, body concerns, and imagery vividness in relation to perceptual, affective, physiological, and behavioural responses. Among their findings, less physically active participants reported more pronounced changes in body feelings and happiness across sound conditions, while responses in felt strength and surprise varied with body concerns. In a complementary study, D'Adamo et al. [13] examined social support networks and found stronger illusion effects among participants with smaller networks. This adds social context to the psychological and behavioural characteristics relevant to understanding response variability.

These findings motivate participant characterisation beyond demographics. Relevant dimensions include physical activity, body-related attitudes, imagery vividness, and social relationships. BODYinTRANSIT-Lab combines systematic recruitment stratification with participant-level psychological, behavioural, and social measures, allowing examination of individual variability alongside repeated within-person responses to altered feedback.

2.4 Complementary Measures of Bodily and Affective Experience

Subjective annotation determines which aspects of a BTE can be investigated computationally. Dimensional affect measures and body-specific reports address complementary questions. Bradley and Lang's Self-Assessment Manikin (SAM) [7] provides assessments of valence, arousal, and dominance. Footsteps Illusion studies have combined these affective measures with questionnaires on bodily feelings and avatar-adjustment tasks [55, 57]. The latter allow participants to visually represent perceived body size. Keeping these measures distinct lets us examine whether changes in felt weight accompany changes in visualised size or affective evaluation.

Body maps have been used in qualitative data analysis to quantify and compare changes in body perception across experimental conditions and participant groups [1]. They also indicate where an experience is felt. Nummenmaa et al. [38] asked participants to mark increases and decreases in bodily sensations on silhouettes and identified distinct spatial patterns associated with different emotions. Their work demonstrated how topographical self-reports can support quantitative analysis of bodily experience. In auditory BTE research, D'Adamo et al. [12, 14] used body maps to investigate the distribution of perceived changes across bodily regions. Such maps complement scalar ratings by preserving spatial information that a single whole-body score cannot convey.

Annotation timing introduces an additional distinction. Sharma et al. [52] developed CASE to collect simultaneous, continuous valence and arousal ratings with a joystick while participants watched affective videos, along with synchronised physiological recordings. This enables the investigation of temporal variation that post-stimulus ratings alone cannot describe. In a walking paradigm, annotation must also account for the activity producing the signals of interest. BODYinTRANSIT-Lab therefore combines repeated verbal perceived-weight ratings within each trial with post-trial questionnaires, body maps, and avatar-based estimates. This design provides complementary temporal, spatial, and perceptual descriptions of experience without assuming that a single annotation method captures the entire BTE.

These approaches demonstrate the value of collecting multiple forms of subjective evidence [49]. For computational use, the annotations must also remain explicitly linked to the experimental conditions and to synchronised sensor recordings. BODYinTRANSIT-Lab links these measures within its acquisition and data-curation protocol so that researchers can examine both agreement and divergence between reported experience and bodily responses, while providing a basis for developing comparable multimodal BTE datasets.

2.5 Affective Movement Datasets and Computational Modelling

Research on affective body expression examines how posture and movement convey affective information [28]. Multimodal affect datasets have established methods for relating bodily signals to subjective experience. DEAP [29] combines physiological recordings with music-video elicitation and affective ratings. AMIGOS [36] adds personality, mood, and individual and group viewing contexts. Together with CASE's temporal annotation [52], these resources demonstrate the value of multimodality, participant characterisation, and context. Movement datasets extend these questions to bodily action, but differ in what varies within a participant, the bodily data they provide, and what their annotations describe. Table 1 summarises selected resources along these dimensions.

Acted and affect-elicitation datasets provide complementary foundations. Fourati and Pelachaud's Emilya [21] supports analysis of emotional expression across daily actions, whereas Li et al. [31] examine walking following video-based affect elicitation. Pain-related datasets link movement to bodily experience and its affective context. EmoPain [3] combines rehabilitation activities with pain- and anxiety-related reports, participant-level distress and catastrophising measures, and expert annotations of pain-related expression and behaviour, supporting models such as [41, 25, 17] for pain and protective-behaviour detection. EmoPain@Home [39] extends this line of research to everyday home activities, with repeated reports of pain, worry, and movement confidence. These resources show

TABLE 1
Selected affective-movement resources compared by experimental variation, recorded or released bodily data, and relevant annotations. Participant-level measures do not imply continuous annotation.

Resource	Experimental variation	Bodily data	Annotations & participant measures
Emilya [21]	Acted emotional states across seven daily actions	Full-body inertial motion capture; video; audio	Intended emotion and third-party ratings of expressed emotion.
Li et al. [31]	Walking before and after video-based induction of anger and happiness	Kinect-based 3D joint trajectories	Self-reported anger and happiness around elicitation and walking.
EmoPain [3]	Rehabilitation activities under normal and increased physical demand	Full-body 3D motion capture; EMG; video; audio	Pain and anxiety reports; distress and catastrophising measures; expert annotations of pain-related expression and behaviour.
EmoPain@Home [39]	Naturally occurring everyday activities at home	Wearable inertial motion capture	Activity labels and repeated reports of pain, worry, and movement confidence.
PsyMo [9]	Seven walking variations, including speed, carrying, and dual-task conditions	Video-derived silhouettes; 2D/3D skeletons; 3D body meshes	Personality, self-esteem, fatigue, aggression, and mental-health measures.
Zhao et al. [68]	Between-person variation in depression severity and personality	Video and audio; ankle-worn IMU gait data	Depression severity and personality.
BODYinTRANSIT-Lab	Footsteps Illusion investigating BTEs through repeated walking with controlled changes to self-generated auditory feedback	Full-body inertial kinematics; ECG; EDA; EMG; respiration; backup acceleration	Within-trial perceived weight; post-trial body feelings and affect; body maps; avatar estimates; physical, psychological, behavioural, and social participant characteristics (Table 5).

that intended expression, elicitation condition, bodily experience, and behavioural annotation capture distinct aspects of affective movement and should not be treated as interchangeable.

Psychologically annotated gait resources address another dimension. Cosma and Radoi's PsyMo [9] combines multiple walking conditions with extensive questionnaire-derived psychological attributes, while Zhao et al. [68] examine depression severity and personality within a multimodal gait-assessment framework. These resources demonstrate that neither multimodality nor rich participant characterisation alone distinguishes BODYinTRANSIT-Lab. Within the comparison in Table 1, its distinction for BTE research lies in combining controlled changes to self-generated sensory feedback with synchronised movement and physiological recordings, repeated within-person reports of body perception and affective experience, and participant-level characteristics that can be examined as sources of between-person variability.

Advances in movement representation learning offer methods for investigating these relationships when task-specific labelled data are scarce. Yuan et al. [67] pre-trained activity-recognition models on 700,000 person-days of unlabelled wearable data and demonstrated transfer across eight activity benchmarks. Their work provides reusable representations for sensor-based movement analysis. These representations serve as a useful computational starting point for examining whether movement contains information associated with experimentally manipulated feedback and with reported bodily or affective experience.

The reviewed literature therefore motivates an integrated resource that connects four elements, including (1) controlled action-dependent feedback, (2) synchronised bodily signals, (3) complementary reports of perceptual and affective experience, and (4) characteristics associated with individual variability. BODYinTRANSIT-Lab brings these elements together and documents their acquisition, synchronisation, curation, and organisation so that the resulting relationships can be investigated computationally and the protocol can inform the development of comparable datasets. The following section describes how these requirements informed participant recruitment, the experimental protocol, and the measures collected.

3 BODYinTRANSIT-Lab: Study Design and Experimental Protocol

BODYinTRANSIT-Lab was designed to examine BTE-related responses to altered own auditory feedback during walking, while retaining information about the individuals experiencing it. The experiment combines repeated exposure to three auditory conditions with recordings of movement and physiology, within-trial perceived-weight ratings, and complementary post-trial reports, summarised in Table 5. This structure connects the feedback delivered during a trial with the participant's reported experience and bodily signals, while participant-level measures provide context for investigating variation in these responses.

The design operates at two levels. Within participants, the auditory condition changes across repeated walking trials. Between participants, recruitment stratification and psychological, demographic, anthropometric, and social measures characterise differences relevant to BTE research. We first describe the auditory comparison, then the participant sample and experimental setup, and finally the trial sequence and subjective measures.

3.1 Auditory Conditions and Repeated-Trial Design

The experiment implements the Footsteps Illusion using SoniWeight Shoes [12, 15]. The system records partici-



Fig. 2. **Left:** real-time footstep feedback under control (C), high-frequency (HF), and low-frequency (LF) conditions; curves indicate frequency shaping. **Centre:** intended perceptual changes relative to the physical body, assessed subjectively. **Right:** kinematic and physiological acquisition, aligned by accelerometer cross-correlation, with start/stop coordinated by Node-RED [40]. Sensor locations are schematic.

pants' footstep sounds, modifies their frequency spectra, and plays them back in real time. This preserves the action-dependent relationship between walking and the auditory feedback under investigation [35]. Following the three-condition paradigm used in [57], the spectral manipulation comprised the following conditions (see Fig. 2):

1) **Control (C):** Footstep sounds were played back without spectral alteration.
2) **High frequency (HF):** Frequencies above 1 kHz were amplified by $+12$ dB, while frequencies at 83–250 Hz were attenuated by -12 dB.
3) **Low frequency (LF):** Frequencies at 83–250 Hz were amplified by $+12$ dB, while frequencies above 1 kHz were attenuated by -12 dB.

Consistent with prior Footsteps Illusion studies [55, 60, 57], we selected the HF and LF conditions to elicit sensations of a lighter and heavier body, respectively. These intended effects motivated the manipulation, and the subjective measures captured participants' responses. The control condition preserves playback of participants' own footsteps and serves as a baseline for comparison without spectral modification. Each participant's session comprised six main walking trials, with two repetitions of each condition presented in a counterbalanced order. This repeated-trial structure provides observations for each auditory condition within the same participant. To examine how these responses vary between individuals, the design also incorporates systematic participant characterisation.

3.2 Participant Recruitment and Characterisation

BODYinTRANSIT-Lab contains recordings from 101 adults recruited through the Universidad Carlos III de Madrid participant pool and public advertisements. Eligibility required participants to be aged 18–70 years, proficient in Spanish, self-report normal hearing, be able to perform the walking tasks, and understand the study information and provide informed consent. Exclusion criteria included psychiatric or neurological diagnoses, regular use of medical or recreational psychoactive substances, disabling orthopaedic or neuromuscular conditions requiring assistance during walking, factors interfering with sensor placement, deafness or acute hearing impairment, and inability to provide informed consent.

We stratified recruitment using eating-disorder symptomatology and physical activity level, two dimensions associated with individual differences in responses to the Footsteps Illusion [57]. Potential participants completed Spanish versions of the Eating Disorder Examination Questionnaire (EDE-Q) [42] and the International Physical Activity Questionnaire (IPAQ) [50] before the experimental session. EDE-Q scores were classified as low or high using global-score cut-offs of 1.98 for female participants and 1.09 for male participants, while IPAQ scores were separated at 3000 MET-min/week. We selected these cut-offs according to [12]. Crossing these classifications resulted in four recruitment groups: low EDE-Q/low IPAQ ($n = 20$), high EDE-Q/low IPAQ ($n = 21$), low EDE-Q/high IPAQ ($n = 41$), and high EDE-Q/high IPAQ ($n = 19$).

Further measures describe aspects of participants' bodies, self-perceptions, and everyday context that are not captured by the two stratification variables. Demographic information included age, self-reported sex assigned at birth and gender identity, and highest completed education level. Sex response options were female, male, intersex, and prefer not to say, and gender identity options were man, woman, non-binary, and prefer not to say. Anthropometric and body-related information included height, actual and



Fig. 3. Experimental setting and tasks. Left: participant equipment. Centre: walking between A and B with spoken perceived-weight ratings. Right: post-trial questionnaires, avatar adjustment, and body maps. The sensor locations are shown in Fig.2

ideal weight, waist and hip circumferences, and shoe EU size. Body measurements also informed motion-capture suit sizing. Separate actual and ideal gender-identity ratings recorded participants' self-perceived and preferred positions on a seven-point masculinity–femininity scale.

Participants also completed Spanish versions of three questionnaires. The Multidimensional Body Self Relations Questionnaire (MBSRQ) [51] assessed attitudes towards the physical body through four scales. Betts' Questionnaire upon Mental Imagery (Betts' QMI) [5] assessed imagery vividness across sensory modalities, including visual, auditory, tactile, gustatory, olfactory, and organic imagery. The Social Support Questionnaire (SSQ6) [33] characterised the social environment using measures such as network size, sex ratio, number of social groups, and the average age of network members. These measures support the examination of associations between participant characteristics and responses to the auditory conditions. Full sample characteristics overall and by recruitment group are reported in the Supplementary Table S1.

The study received approval from the Committee of Ethics in Research of the Universidad Carlos III de Madrid under Application No. 'CEI21_04_TAJADURA_Ana' and was conducted in accordance with the Declaration of Helsinki. All participants provided written informed consent and received 20 euros for their time.

3.3 Experimental Setting and Feedback Apparatus

The experimental setting supported both generating footstep sounds and recording bodily responses during walking. Sessions took place in a laboratory measuring 10.0 m long, 4.7 m wide, and 2.7 m high. Participants walked on a wooden path 7 m long and 0.8 m wide, with wood selected for its acoustic properties (see Fig. 3). We minimised external noise during the sessions by controlling the recording environment. We positioned the microphones very close to the footwear and kept them at a consistent distance and location for all participants, ensuring a very high SNR, maintaining standardised recording conditions, and enabling the capture of the footstep sounds used for playback.

SoniWeight Shoes delivered the auditory conditions via a Bela processing board and Sennheiser HDA300 headphones. Participants wore a motion-capture suit and physiological sensors throughout the entire recording session. Figure 2 distinguishes the auditory-feedback loop from the recording systems. The acquisition specifications and sensor montage are detailed in Section 4.1. The hardware and software specifications are provided in Table 2.

TABLE 2
Hardware and software specifications used in the experimental setup, including data acquisition and sound playback systems.

Component	Details
Motion capture suit	Rokoko Smartsuit Pro II with 19 IMU sensors, sampled at 100 Hz.
Physiological sensing	BITalino 8-channel biosignalsplux kit; sampling rate 1000 Hz.
Auditory feedback	SoniWeight Shoes (custom) [15].
Microphones	Core Sound binaural microphone.
DSP board	Bela.io Starter Kit; average latency 1.6 ms.
Headphones	Sennheiser HDA300; closed-back, circumaural.
Control software	Bespoke JavaScript GUI implemented in Node-RED [40].
Operating system	Windows 11.
Analysis software	Python 3.10 and MATLAB R2024a.

3.4 Trial Sequence and Within-Trial Reports

Each trial comprised a brief acclimatisation period followed by repeated walking along the path. Participants first walked in place at Point A for approximately 10 s while exposed to the assigned auditory condition, then provided an initial perceived-weight rating before beginning the walking task. They subsequently walked back and forth three times between Points A and B, 7 m apart, yielding six traversals per trial. Upon reaching each endpoint, participants verbally rated their perceived body weight on a seven-point scale from 1 (light) to 7 (heavy), providing additional ratings, for a total of seven ratings per trial. Figure 3 illustrates the walking directions and spoken ratings at the endpoints.

These repeated reports captured perceived weight at successive time points within each walking trial. Immediately following each trial, participants completed the post-trial assessments described below. Across a complete six-trial session, the protocol thus produced 42 within-trial perceived-weight ratings and six sets of post-trial assessments, together with continuous kinematic and physiological recordings. These two annotation timescales were complementary: repeated ratings captured variation in felt weight during task performance, whereas post-trial measures characterised additional aspects of the experience.

3.5 Post-Trial Measures of Bodily and Affective Experience

Post-trial annotation combined scalar ratings, avatar adjustments, and body maps to capture aspects of the experience that the within-trial weight rating alone cannot describe. The Body Feelings Questionnaire, adapted from [55, 60], collected seven-point ratings of bodily sensations, including perceived weight. The Self-Assessment Manikin (SAM) [7] provided nine-point ratings of valence, arousal, and dominance. These measures retain the distinction between how the body feels and the affective experience accompanying that feeling. The post-trial perceived-weight rating was separate from the seven verbal weight ratings collected during walking. After each trial, participants rated how light or heavy they felt on the same seven-point (1–7) scale.

To obtain a visual estimate of felt body size, participants adjusted the weight of a three-dimensional avatar using the Body Visualizer tool [34] until it matched their felt body size. The resulting avatar weight, expressed in kilograms, provides an alternative measure of perceived body size, consistent with its use in previous studies [55, 60, 56, 57]. In addition to this scalar measure, participants freely marked locations where they experienced changes in bodily sensation or perception after each trial (see Fig. 3, right panel). These drawings were subsequently coded by region. This spatial reporting approach complements the scalar and avatar-based measures. Its conversion into quantitative data is described in Section 4.3.

The protocol therefore produces several descriptions of the same trial: the auditory condition, bodily recordings, repeated perceived-weight reports, and post-trial assessments of bodily and affective experience. The following section describes how the recordings and annotations were acquired and prepared as components of the BODYinTRANSIT-Lab resource.

4 Multimodal Data Acquisition, Curation, and Dataset Structure

The experimental protocol defines the conditions and observations needed to investigate BTEs. Their computational use also depends on how signals from separate acquisition systems are aligned, how subjective reports are represented, and how the resulting data are organised. This section follows these stages from sensor acquisition through signal and annotation preparation to the structure of the BODYinTRANSIT-Lab resource. The documented workflow provides a basis for developing comparable multimodal datasets.

4.1 Kinematic and Physiological Acquisition

Kinematic signals were acquired with the Rokoko Smartsuit Pro II [2] at 100 Hz, using the configuration specified in the Rokoko software [2]. The system provides kinematics for 17 anatomical segments, including three-dimensional position, velocity, and acceleration, as well as joint angles and angular derivatives. The resulting representation contains 306 channels, comprising 153 segment-level linear kinematic channels and 153 joint-angular channels. Segment positions and joint angles are reported in the suit's body-fixed anatomical coordinate system, with segment axes aligned with the corresponding anatomical directions and joint angles defined relative to adjacent segments, following Rokoko's standard conventions. Table 3 summarises how these channels are organised by anatomical location and measurement type.

We acquired physiological signals using the BITalino system at 1000 Hz. EMG was recorded from the cervical trapezius and left vastus medialis oblique, ECG from the chest, EDA from the non-dominant hand, and respiration using a chest belt. The system also included a 3-axis ACC. After recording, we identified intermittent sensor failures affecting individual axes, so each recording contains one, two, or three acceleration axes, depending on the trial. The released dataset retains this signal as backup rather than as a complete 3-axis modality.

These two systems record movement and physiological data during the walking trials, with their acquisition specifications summarised in Table 2.

4.2 Synchronisation and Signal Preparation

Recording across the two systems required both acquisition coordination and subsequent temporal alignment. A bespoke JavaScript interface, implemented in Node-RED [40], sent simultaneous start and stop commands. We aligned streams using cross-correlation of accelerometer signals recorded during the initial walking-in-place period. Thus, the acclimatisation activity described in Section 3.4 also supplied a shared movement signal for alignment. The synchronisation code is available through [11].

Signal preparation was modality-specific, reflecting the spectral characteristics and artefacts of each physiological recording, as summarised in Table 4. The preprocessing pipeline included band-pass or low-pass filtering, as appropriate, and Hampel filtering to remove transient outliers. EMG was additionally filtered with a 49–51 Hz Butterworth notch filter and full-wave rectified. BITalino recordings were

TABLE 3
Kinematic channels exported by the Rokoko Smartsuit Pro II. Segment linear kinematics comprise position, velocity, and acceleration along the X, Y, and Z axes. Each joint-angular group comprises angle, angular velocity, and angular acceleration. ‡ 17 body segments include the head, chest, pelvis, shoulders, upper arms, forearms, hands, upper legs, lower legs, and feet.

Kinematic quantity	Anatomical locations	Groups	Channels
Segment linear kinematics			
Position, velocity, acceleration (X, Y, Z)	17 body segments‡	17 each	**153**
Joint angular kinematics			
Flexion	Hips, knees, shoulders, elbows, wrists, neck	11	33
Abduction	Wrists, elbows, shoulders, hips	6	18
External rotation	Shoulders, hips, knees	6	18
Pronation	Wrists, elbows	4	12
Adduction	Knees, hips	4	12
Dorsiflexion, inversion, internal rotation	Ankles	2 each	18
Extension, lateral flexion/rotation, axial rotation	Thorax, pelvis	2 each	18
Protraction, medial rotation, posterior tilt	Scapulae	2 each	18
Leftward tilt, Rightward rotation	Neck	1 each	6
		Angular subtotal	**153**
		Total kinematic channels	**306**

processed at the original 1000 Hz sampling rate using the modality-specific procedures, then downsampled to 100 Hz. We applied no additional filtering to the Rokoko recordings. These steps describe preparation of the acquired streams. The original acquisition rates differ from the sampling rates of the processed signals, and we provide both raw and preprocessed sensor recordings.

4.3 Subjective Annotation Conversion

Subjective measures retain distinct temporal and representational forms. Verbal perceived-weight ratings provide repeated observations within each trial, whereas body-feeling and SAM scores describe the post-trial assessment. The avatar-adjustment task contributes the selected avatar weight in kilograms. Preserving these distinctions ensures that the reports are interpreted according to the manner and timing of their elicitation, rather than as interchangeable descriptions of experience.

The free-form paper body maps required a subsequent coding step. We visually inspected the completed drawings and manually coded them using a 136-region representation covering the anterior and posterior body surfaces, following the method described in [12, 14]. We assigned each region a value from 0 to 1 based on the proportion of its area the participant marked, with 0 indicating no marking in that region. This representation retains information on the spatial distribution of reported bodily changes, along with scalar ratings and avatar estimates.

4.4 Dataset Organisation and Data Inventory

BODYinTRANSIT-Lab contains recordings from 101 participants, totalling approximately 16 hours of data and about 15 GB. The database is organised by participant. Each participant directory contains separate subfolders for BITalino, Rokoko, and Label. Trial-specific CSV filenames encode the participant identifier, auditory condition, and repetition, linking the physiological, kinematic, and trial-level annotation files across the database. The manually coded body-map annotations are stored separately in CSV files and retain the corresponding participant and trial identifiers.

The data inventory in Table 5 separates participant-level characteristics, experimental identifiers, time-series recordings, and subjective reports. Participant identifiers, trial indices, and auditory-condition codes provide context for interpreting the observations. Distinguishing participant-level information from repeated trial-level measurements supports analysis of between- and within-individual variability.

This organisation brings controlled feedback, participant characteristics, bodily signals, and reported experiences into a common resource. The next section defines a computational benchmark using selected kinematic recordings and subjective measures, and describes its task-specific transformations as part of that analysis.

5 Computational Benchmark

The computational benchmark examines whether walking kinematics contain information associated with the auditory manipulation and the bodily and affective experiences reported in response to it. These relationships provide an initial reference performance for evaluating the computational use of BODYinTRANSIT-Lab. We selected auditory condition, relative perceived weight, and relative affective valence as classification targets to examine the delivered manipulation, the body-perceptual change targeted by the Footsteps Illusion, and the reported affective experience. Auditory-condition labels identify the delivered feedback (C, HF, or LF), whereas the subjective labels derive from the post-trial seven-point perceived-weight rating and nine-point SAM valence rating. The physiological recordings and participant-level measures remain available for broader analyses beyond this kinematic benchmark.

For the subjective tasks, the comparison of interest is each participant's reported experience relative to their control condition. We therefore constructed labels separately for each participant and task, using the mean of the two control-trial ratings as the participant-specific reference. Both control trials were assigned label 0 by definition. For the four manipulated-feedback trials (two HF and two LF), ratings

TABLE 4
Preprocessing of BITalino signals. Filters were applied using zero-phase forward–backward filtering. Hampel filtering removed transient outliers using fixed or signal-dependent window lengths.

Signal	Filtering	Filter parameters	Subsequent processing
Respiration	Butterworth IIR band-pass	Order 2; 0.1–0.35 Hz	Adaptive Hampel filtering with window length derived from the detected breathing rate; 3-SD threshold.
EMG	FIR band-pass; Butterworth IIR notch	Band-pass order 20, 20–450 Hz; notch order 2, 49–51 Hz	Hampel window 10 samples; 3-SD threshold; full-wave rectification.
ECG	FIR band-pass	Order $\lfloor 0.3 f_s \rfloor$; 3–45 Hz	Hampel window 50 samples; 3-SD threshold.
EDA	Butterworth IIR low-pass	Order 4; 5 Hz	Hampel window 150 samples; 3-SD threshold.
Backup acceleration‡	Butterworth IIR band-pass	Order 2; 0.5–10 Hz	Hann window followed by adaptive Hampel filtering with a window length derived from the dominant frequency.

‡The dominant frequency was estimated from the Hann-windowed signal and used to determine the Hampel-window length; Hampel filtering was then applied to the same windowed signal.

TABLE 5
BODYinTRANSIT-Lab data inventory, grouped by participant characteristics, experimental identifiers and auditory manipulation, sensor recordings, and subjective responses. The table distinguishes each entry by its level and collection timing. Detailed acquisition specifications are given in Table 2, and kinematic channels are listed in Table 3.

Component	Measures or identifiers	Level and timing	Representation
Participant characteristics and identifier			
Participant identifier	Unique participant ID	Participant	Integer identifier
Demographic measures	Age, sex, gender, education	Participant; before session	Numeric or categorical values
Anthropometric measures	Height, waist, hip circumferences (cm); weight (kg); shoe size (EU)	Participant; before session	Numeric values in the stated units
Body-related self-perceptions	Actual and ideal gender-identity ratings; ideal weight (kg)	Participant; before session	Ordinal masculinity–femininity ratings, 1–7; numeric weight
Psychological and behavioural measures	EDE-Q, IPAQ, MBSRQ, Betts' QMI	Participant; before session	Questionnaire scores
Social measures	SSQ6	Participant; before session	Numeric network descriptors
Experimental identifiers and auditory manipulation			
Auditory condition and trial index	Control (C), high frequency (HF), low frequency (LF); trial identifier	Trial	Condition code; index 1–6
Sensor recordings			
Kinematic recordings	Segment position, velocity, acceleration; joint angles and angular derivatives	Time series; during trial	306 numeric channels
Physiological recordings	ECG, EDA, EMG, respiration	Time series; during trial	Numeric signal channels
Backup acceleration	Available accelerometer axes	Time series; during trial	One, two, or three numeric channels, depending on the trial
Subjective responses			
Within-trial perceived weight	Seven verbal reports per trial	Trial; during trial	Ordinal ratings, 1–7
Post-trial body feelings	Body Feelings Questionnaire, including perceived weight	Trial; immediately after walking	Ordinal ratings, 1–7
Post-trial affect	SAM valence, arousal, dominance	Trial; immediately after walking	Ordinal ratings, 1–9
Avatar-based estimate	Adjusted avatar weight representing felt body size	Trial; immediately after walking	Numeric value (kg)
Body maps	Regional extent of reported changes in bodily sensation or perception	Trial; immediately after walking	136 regional occupancy values, 0–1; 0 denotes no marking

above this reference received label $+1$, ratings below it received label -1, and ratings equal to it received label 0. Since preliminary inspection indicated that many participants used a narrow portion of the rating scale, we did not apply a tolerance margin. Under this definition, class 0 includes the control trials regardless of their individual ratings and does not denote an independently measured absence of a BTE. Each walking segment, corresponding to one traversal, inherits the auditory condition and the relevant subjective label of its parent trial. The subjective targets therefore describe post-trial reports rather than momentary measurements of experience at each input window. The seven within-trial perceived-weight ratings, avatar estimates, and body maps described in Sections 3.4 and 3.5 provide complementary annotations but were not used to construct these targets.

Individual differences in movement and subjective scale usage motivate a within-participant evaluation, with a separate model trained for each participant and task. The benchmark evaluates prediction on held-out walking segments from the same recorded session. Because segments from a single trial may appear in different subsets, the benchmark is limited to prediction within recorded sessions rather than generalisation to unseen trials, sessions, or participants. Of the 101 participants, 99 completed all six trials, and 98 also provided the questionnaire data required for the subjective tasks. Excluding participants whose labels comprised only one class left 92 participants for relative perceived weight and 83 for relative affective valence. These eligibility criteria apply to the benchmark and do not alter the dataset's composition of 101 participants.

5.1 Model Architecture



Fig. 4. The network architecture used for the computational benchmark.

The proposed architecture, depicted in Figure 4, combines the frozen movement feature extractor of the multi-task self-supervised model proposed by Yuan et al. [67] with newly trained attention pooling and a classification head. The architecture addresses two design considerations: learning from limited participant-specific data and combining information from multiple body locations. We select the five-second checkpoint pretrained on wrist accelerometry from 100,000 UK Biobank participants [19] and replace its original classifier with the pooling and classification components described below. Only these new components are trained for each participant and task.

The shared frozen feature extractor processes each of the 17 triaxial acceleration streams independently, producing a 512-dimensional representation for each body location. A shared linear projection scores each feature vector, and a softmax across the 17 body locations converts these scores into input-dependent attention weights. An attention-pooling layer uses these weights to combine the 17 feature vectors into a single 512-dimensional vector. This mechanism is intended to accommodate differences in signal informativeness across body locations, including those associated with the artefact masking described in Section 5.2.

The classification head comprises a linear projection to 256 dimensions, a ReLU activation, and a final linear layer producing three class logits. The frozen extractor provides a common movement representation, while the trainable pooling and classification layers determine how information from the different locations contributes to the prediction.

5.2 Benchmark Inputs and Evaluation

The benchmark uses the three acceleration components from each of the 17 body segments, giving 51 input channels from the 306 kinematic channels described in Section 4.1. Each observation contains 150 samples per channel, matching the pretrained model's input size. We downsampled the 100 Hz recordings by a factor of three using a normalised triangular kernel $[0.25,\ 0.5,\ 0.25]$. This downsampling was applied independently to each channel using grouped one-dimensional convolution with stride 3 and one-sample padding. At the resulting rate of approximately 33.3 Hz, 150 samples span 4.5 s. The pretrained model used 5 s windows sampled at 30 Hz. These transformations prepare inputs for the benchmark and are separate from the dataset-curation procedures in Section 4.

We applied artefact masking separately to each acceleration axis to limit the influence of unusually low or high signal variation within a window. Masking uses each axis's dynamic range within a window, defined as the difference between its max and min values, relative to an axis-specific reference interval. The interval spans the 5th to 95th percentiles of dynamic ranges from five participants' recordings selected by visual inspection for signal quality. These participants were also included in the benchmark, so the reference recordings were not independent of the evaluation sample. During training and evaluation, an axis with a dynamic range outside its reference interval was set to zero. Min–max normalisation to $[-1, 1]$ then places each axis within each window on a common scale while retaining the temporal pattern of non-constant signals.

A complete session provides 36 walking segments after acclimatisation, corresponding to six traversals in each of

six trials. For each participant and random seed, we partitioned these segments, rather than individual windows, so that every window from a given segment belonged to the same subset. We used Monte Carlo cross-validation with 10 random seeds and a two-stage split into training, validation, and test subsets at target proportions of 60%, 20%, and 20%. For stratification, we considered auditory condition and task label jointly. However, when the available observations did not permit stratification, the split was random. The subjective labels were defined before partitioning, using the mean of both control-trial ratings as the participant-specific reference. Within each subset, we randomly sampled walking segments and window starting positions for training, validation, and testing. To compute the reported metrics, we used individual window predictions without aggregation into segment- or trial-level decisions.

For each participant, task, and random seed, the trainable components of the benchmark model were optimised using a weighted cross-entropy loss. Class weights were computed as the inverse of the class frequencies in the training subset for each run, to increase the contribution of less frequent labels to the loss. Models were optimised with Adam using a learning rate of 1×10^{-4}, a batch size of 64, and a maximum of 100 epochs. Training stopped when validation loss did not improve for 10 consecutive epochs or when validation balanced accuracy reached 1.0. The checkpoint with the lowest validation loss was retained for evaluation on the test set.

Performance was evaluated using balanced accuracy (BA), macro F1, and overall accuracy (Acc). BA averages class-specific recall, whereas macro F1 averages the harmonic mean of precision and recall within each class. These metrics are calculated using Eq. (1), where K denotes the number of evaluated classes and TP_c, FP_c, and FN_c denote true positives, false positives, and false negatives for class c, respectively.

$$\mathrm{BA} = \frac{1}{K}\sum_{c=1}^{K}\frac{\mathrm{TP}_c}{\mathrm{TP}_c + \mathrm{FN}_c},$$
$$\mathrm{F1} = \frac{1}{K}\sum_{c=1}^{K}\frac{2\mathrm{TP}_c}{2\mathrm{TP}_c + \mathrm{FP}_c + \mathrm{FN}_c}. \quad (1)$$

Overall accuracy is the proportion of correctly classified test windows. To summarise BA, macro F1, and overall accuracy across participants, we averaged each participant's scores over the 10 random seeds and then calculated the mean and standard deviation of these participant-level averages. This gives each participant equal weight in the reported metrics. For descriptive comparison, we also report the proportion of participants whose mean BA exceeded the nominal three-class reference of 33.3%. Figure 5 provides the confusion matrices for the subjective tasks, showing predictions pooled across participants and seeds.

5.3 Benchmark Results

Table 6 summarises classification performance and task-specific participant counts. Auditory-condition classification achieved a mean balanced accuracy of 36.6%, compared with the nominal three-class reference of 33.3%. Mean balanced accuracy was higher for relative perceived weight ($44.5 \pm 11.5\%$) and relative affective valence ($45.9 \pm 12.7\%$). The reported $\pm$ values are the sample standard deviations across participants after averaging each participant's balanced accuracy over the 10 random seeds. For the auditory condition, 61.6% of participants had balanced accuracy greater than 33.3%, 81.5% for relative perceived weight, and 84.3% for relative affective valence.



Fig. 5. Confusion matrices for relative (a) affective valence and (b) perceived weight. Window-level predictions and targets were pooled across participants and 10 random seeds. Ref. (0) includes control trials and manipulated-feedback trials rated equal to the participant-specific control mean. Negative/positive labels indicate lower/higher valence or lighter/heavier perceived weight relative to that mean.

Alongside classification performance, the subjective-task attention profiles describe how the benchmark model allocates weights across body locations. For each participant, we averaged attention weights across test windows within each seed and then across the 10 seeds. Figure 6 presents the attention profiles for the two subjective tasks. The models assigned the largest average weights to the foot sensors, followed by contributions from the hands and upper legs. Trunk and proximal locations remained closer to the uniform-weight reference. The two profiles were broadly similar, with differences mainly in the relative weighting of individual locations.

Error bars overlapped for many locations with intermediate weights. The profiles therefore provide a broad description of weight allocation rather than a precise ranking of body regions.

6 Discussion

BODYinTRANSIT-Lab connects controlled auditory feedback with multimodal recordings, complementary subjective annotations, and participant characteristics. The benchmark examines one computational use of this resource: classifying auditory conditions and relative post-trial reports from walking kinematics. We discuss what the dataset and these results support, how the observations may inform further work, and the methodological and ethical considerations that accompany the computational study of BTEs.

6.1 Complementary Annotations and Further Computational Uses

The dataset supports investigation of relationships among sensory manipulation, reported experience, and bodily sig-

TABLE 6
Within-participant classification performance across eligible participants and 10 random seeds. N/pool denotes the number of eligible participants relative to the task-specific pool. Class distributions are reported as average participant-level percentages, with subjective-task class distributions ordered as $[-1, 0, +1]$. >ref. denotes the percentage of participants with balanced accuracy exceeding the nominal three-class reference value of 33.3%. All performance values are percentages.

Task	N/pool	Distribution	Accuracy	BA	F1	>ref.
Auditory condition	99/99	34.4/33.8/31.8	34.8	36.6	28.5	61.6
Relative perceived weight	92/98	23.3/41.4/35.3	43.9	44.5	37.6	81.5
Relative valence	83/98	26.5/47.0/26.5	47.9	45.9	39.4	84.3



(a) (b)

Fig. 6. Attention profiles for (a) relative affective valence and (b) relative perceived weight. Weights were averaged within participants across test windows and then across 10 seeds. Bars show participant-level means averaged across participants, and error bars show population standard deviations across participants.

nals without requiring these measures to be interchangeable. Seven perceived-weight reports per trial provide repeated observations of perceived weight, while post-trial questionnaires, avatar estimates, and body maps capture affective, scalar, visual, and spatial aspects of the experience. These complementary annotations support investigation of how different aspects of a BTE relate to one another and, through the repeated within-trial reports, enable fine-grained analysis of how the BTE emerges and unfolds over time [63]. Participant-level measures add context for examining variation between individuals. These observations also allow researchers to study both convergence and divergence among responses to the same feedback.

The present benchmark uses only acceleration signals and two post-trial rating measures. Further analyses could examine relationships between reported body size and felt weight, and associations between participant characteristics and response patterns. Combining kinematics with ECG, EDA, EMG, and respiration may provide complementary information for these analyses. The dataset's availability, along with its acquisition and curation protocol, supports secondary analyses and the development of comparable resources without repeating the entire experimental procedure [49, 23, 24].

6.2 Interpretation of the Kinematic Benchmark

The subjective tasks achieved higher mean balanced accuracy than auditory-condition classification under the reported evaluation protocol. This pattern aligns with the distinction established in the study design: the delivered auditory feedback and the experience reported under it are related but target different constructs. These results provide initial reference values for predicting control-referenced perceived weight and affective valence labels from walking kinematics within recorded sessions. Because training and test subsets may contain segments from the same trial, the evaluation does not distinguish information associated with reported experience from other trial-specific movement characteristics. They do not establish that subjective experience mediates an effect of auditory feedback on movement, or that condition-related information has been eliminated from the subjective tasks.

Interpretation also depends on how the labels are constructed. Both control trials were assigned class 0, while ratings under HF and LF feedback were compared with the participant's control mean. The subjective targets therefore combine the experimental reference with the relative rating. Moreover, the three tasks differ in class distributions and eligible participant pools. Their performance contrast should therefore be treated as a descriptive comparison under the specified benchmark, rather than a direct comparison of the strengths of perceptual, affective, and motor effects.

The benchmark model provides a baseline for combining pretrained movement representations across body locations. The attention profiles place the largest average weights on the feet for both subjective tasks, motivating evaluation of whether a smaller sensor configuration could retain useful predictive information. This possibility requires direct comparisons of sensor subsets, supported by systematic sensor ablation [26]. Similarly, the contribution of attention pooling to artefact robustness requires comparison with alternative aggregation methods; the present results alone do not isolate that contribution.

6.3 Ethical Considerations for BTE Computing

Although automatic detection of BTEs has the potential to support interventions for health and well-being, it also raises important ethical challenges. Body-perception technologies occupy a particularly sensitive position because they both influence bodily experience and infer internal psychological states. As recently argued in HCI research [37], such systems require careful consideration of agency, informed consent, transparency, and trust, particularly when technologies adapt their behaviour in real time or influence users outside their conscious awareness. Moreover, physiological and behavioural signals should not be treated as objective proxies for subjective experience. Body perception and emotion are inherently contextual, culturally situated, and individually variable, making algorithmic overconfidence and inappropriate personalisation significant risks.

In this regard, relying on biometrics, movement sensing, and computational models risks reducing rich, lived bodily experiences to quantitative metrics or categorical labels. Such reductions become particularly problematic when adaptive systems aim to guide users toward a "preferred" bodily or emotional state, raising questions about who defines these states and according to which values. Because BTEs inevitably intersect with cultural norms surrounding appearance, ability, gender, ageing, and emotional expression, computational models and feedback mechanisms may inadvertently reinforce normative assumptions about what constitutes a "healthy," "proper," or "ideal" body [59].

Addressing these challenges requires human-centred approaches that make underlying assumptions explicit, recognise that sensing and adaptation technologies are not value-neutral, and preserve users' autonomy throughout the interaction [59]. Future systems should enable users to understand and challenge system inferences, minimise the unnecessary collection of sensitive physiological data, evaluate fairness across diverse populations [37], as well as support opportunities for users to define meaningful BTEs on their own terms rather than imposing normative models of embodiment [59]. We therefore argue that ethical reflection should be treated as a core design principle of computational models of BTEs from the outset, particularly as these technologies move from laboratory settings into everyday and clinical environments.

6.4 Limitations and Future Work

The benchmark evaluates held-out walking segments from participants whose other segments are available for training. Because partitioning occurs at segment level, observations from the same trial may appear in different subsets. Performance therefore characterises within-session prediction and does not establish generalisation to unseen trials, sessions, or participants. Evaluation that holds out complete trials, followed by separate-session and participant-independent studies, would address progressively broader prediction settings.

The subjective tasks also depend on a reference estimated from two control ratings. This reference is sensitive to variation in those ratings, and the resulting classes retain direction rather than the magnitude of the difference. Future benchmarks could compare alternative reference and target constructions, including regression, and evaluate their stability.

Annotation timing places an additional constraint on the conclusions. The seven within-trial perceived-weight ratings are repeated discrete observations, while post-trial ratings provide one assessment per trial. Assigning a post-trial label to walking windows does not establish the participant's state at every moment within those windows. More frequent or continuous annotation could support investigation of temporal changes [20], but its use during walking must account for the movement and cognitive demands introduced by reporting. The distinction between repeated and post-trial annotation therefore remains relevant even when the underlying sensor signals are continuous.

The sample consisted predominantly of young adults recruited in Madrid, Spain. Replication across age groups, cultural contexts, and health conditions would help assess the transferability of observations and models [27, 53]. The protocol also concerns short-term exposure to auditory feedback. Longitudinal studies would be needed to examine the persistence of perceptual changes and their relationship to longer-term changes in the physical body [66]. Adaptive, everyday, or clinical uses would additionally require prospective evaluation of prediction reliability and the effects of acting on model outputs.

7 Conclusion

BODYinTRANSIT-Lab provides a multimodal dataset for investigating Body Transformation Experiences (BTEs) through the Footsteps Illusion, in which participants hear their own footstep sounds modified in real time as they walk. Recordings from 101 adults, recruited through stratification by eating-disorder symptomatology and physical activity, link repeated auditory conditions to full-body kinematics, physiology, and subjective assessments. This design allows responses to the same feedback to be examined within individuals and in relation to differences in their characteristics.

Seven within-trial perceived-weight ratings capture variation during walking, whereas post-trial bodily and affective assessments, avatar estimates, and body maps describe additional perceptual and spatial aspects of the experience. Linking these observations to bodily recordings supports investigation of how movement and physiology relate to different aspects of reported experience.

As an initial computational benchmark, we proposed a deep learning architecture that combines pretrained movement representations with attention-based pooling across body locations. Using walking acceleration, it achieved mean balanced accuracies of 36.6% for the auditory condition, 44.5% for relative perceived weight, and 45.9% for relative affective valence when evaluated on held-out walking segments within participants' recorded sessions.

Beyond this kinematic benchmark, the physiological recordings support an investigation into whether combining movement and physiology improves predictions of reported bodily and affective experiences. The accompanying acquisition and curation protocol documents how we collected, synchronised, prepared, and organised the observations to

provide a basis for other researchers to construct comparable multimodal datasets for studying BTEs. Together, the dataset and documented protocol provide a reusable basis for investigating variation in bodily and subjective responses within and across individuals and for collecting comparable multimodal data on BTEs.

8 Data Availability

The BODYinTRANSIT-Lab database is made available to the research community to promote reproducible research. The proposed benchmark implementation is available at https://github.com/tmcortes/BODYinTRANSIT. Due to the sensitive nature of the data, access is provided under specific conditions. Interested researchers are required to complete and sign an End-User Licence Agreement. Upon verification, researchers will be provided with credentials to access the dataset (∼15 GB) from a secure repository at the Universidad Carlos III de Madrid.

Acknowledgement

This work was supported by the European Union's Horizon 2020 program under grant No. 101002711 for the BODYinTRANSIT project.

Appendix I
Participant Demographics

Table S1 shows Demographics, Anthropometric, Psychometric, and Social Characteristics for 101 participants.

References

[1] Karen Anne Cochrane, Kristina Mah, Anna Ståhl, Claudia Núñez-Pacheco, Madeline Balaam, Naseem Ahmadpour, and Lian Loke. Body maps: A generative tool for soma-based design. In *Sixteenth International Conference on Tangible, Embedded, and Embodied Interaction*, New York, NY, USA, 2022. ACM.

[2] Rokoko Electronics ApS. Smartsuit Pro II: Full body motion capture. https://www.rokoko.com/. Accessed: November 11, 2026.

[3] Min S. H. Aung, Sebastian Kaltwang, Bernardino Romera-Paredes, Brais Martinez, Aneesha Singh, Matteo Cella, Michel Valstar, Hongying Meng, Andrew Kemp, Moslem Shafizadeh, et al. The Automatic Detection of Chronic Pain-Related Expression: Requirements, Challenges and the Multimodal EmoPain Dataset. *IEEE Transactions on Affective Computing*, 7(4):435–451, 2016.

[4] Elena Azañón, Luigi Tamè, Angelo Maravita, Sally A. Linkenauger, Elisa R. Ferrè, Ana Tajadura-Jiménez, and Matthew R. Longo. Multimodal Contributions to Body Representation. *Multisensory Research*, 29(6-7):635–661, 2016.

[5] John S. Baranchok. *The linguistic and statistical equivalence of Spanish and English versions of Betts questionnaire upon mental imagery*. PhD thesis, Texas Tech University, 1994. Available on: http://hdl.handle.net/2346/61170.

[6] Matthew Botvinick and Jonathan Cohen. Rubber hands 'feel' touch that eyes see. *Nature*, 391(6669):756, 1998.

[7] Margaret M. Bradley and Peter J. Lang. Measuring emotion: The Self-Assessment Manikin and the Semantic Differential. *Journal of Behavior Therapy and Experimental Psychiatry*, 25(1):49–59, 1994.

[8] Sunje Clausen, Ana Tajadura-Jiménez, Christian P. Janssen, and Nadia Bianchi-Berthouze. Action sounds informing own body perception influence gender identity and social cognition. *Frontiers in Human Neuroscience*, 15:688170, 2021.

[9] Adrian Cosma and Emilian Radoi. Psymo: A dataset for estimating self-reported psychological traits from gait. In *Proceedings of the IEEE/CVF Winter Conference on Applications of Computer Vision*, pages 4603–4613, Waikoloa, HI, USA, 2024. IEEE.

[10] Laura Crucianelli, Nicola Metcalf, Aikaterini Fotopoulou, and Paul M. Jenkinson. Bodily pleasure matters: velocity of touch modulates body ownership during the rubber hand illusion. *Frontiers in Psychology*, 4:703, 2013.

[11] Amar D'Adamo. SoniWeightShoes App: Synchronization Script for OpenSignals and Rokoko Studio APIs, 2025. Available on: https://github.com/hokumi9/RaramuriSync-SoniWeight-Shoes.

[12] A. D'Adamo, M. Roel Lesur, L. Turmo Vidal, M. M. Dehshibi, J. R. Díaz Durán, D. De La Prida, L. A. Azpicueta-Ruiz, A. Väljamäe, and A. Tajadura-Jiménez. SoniWeight Shoes: Investigating Effects and Personalisation of a Wearable Sound Device for Altering Body Perception and Behavior. In *Proceedings of the 2024 CHI Conference on Human Factors in Computing Systems*, pages 1–20, New York, NY, USA, 2024. ACM.

[13] Amar D'Adamo, Angel Sánchez, Lize De Coster, and Ana Tajadura-Jiménez. Sound effects on body perception vary with the social support network of individuals. *iScience*, 28(8):113091, 2025.

[14] Amar D'Adamo, Laia Turmo Vidal, Karunya Srinivasan, Mohammad Mahdi Dehshibi, Daniel De La Prida, and Ana Tajadura-Jiménez. Mapping the Body: Developing Body Maps as Research Tool to Derive Quantifiable and Context-Sensitive Design Insights. In *Proceedings of the Twentieth International Conference on Tangible, Embedded, and Embodied Interaction (TEI '26)*, pages 1–16, Chicago, IL, USA, 2026. ACM.

[15] Daniel de la Prida, Joaquín R. Díaz Durán, Luis Antonio Azpicueta-Ruiz, and Ana Tajadura-Jiménez. As light as your footsteps: Design and evaluation of a portable device for changing body perception through a sound illusion. In *53º Congreso Español de Acústica – XII Congreso Ibérico de Acústica (TECNIACUSTICA)*, Elche, Spain, 2022. Sociedad Española de Acústica.

[16] Frédérique De Vignemont. *Mind the Body: An Exploration of Bodily Self-Awareness*. Oxford University Press, Oxford, UK, 2018.

[17] Mohammad Mahdi Dehshibi, Temitayo Olugbade, Fernando Díaz-de María, Nadia Bianchi-Berthouze, and Ana Tajadura-Jiménez. Pain Level and Pain-Related Behaviour Classification Using GRU-Based Sparsely-Connected RNNs. *IEEE Journal of Selected Topics in Signal Processing*, 17(3):677–688, 2023.

[18] Chris Dijkerman and Bigna Lenggenhager. The body

TABLE S1
Participant Demographics, Anthropometric, Psychometric, and Social Characteristics (N=101). **Sex (F/M)**: Sex assigned at birth (Woman / Man); **Gender (F/M/NB/O)**: Self-reported gender (Feminine / Masculine / Non-binary / Other or Prefer not to say); **Education Level (1-5)**: Ordinal education level (1=Secondary, 2=Vocational, 3=Bachelor's, 4=Master's, 5=PhD); **Gender Identity (1-7)**: Self-perceived gender identity (1=very masculine, 7=very feminine); **Ideal Gender Identity (1-7)**: Desired gender identity (1=very masculine, 7=very feminine); **EDE-Q**: Eating Disorder Examination Questionnaire (Global Score); range 0–6, where 0 indicates no/lowest frequency of eating-disorder-related symptoms and 6 indicates the highest frequency/severity of symptoms; **IPAQ**: International Physical Activity Questionnaire (total score in MET-min/week); quantifies physical activity based on reported activity during the previous 7 days, expressed in Metabolic Equivalent of Task (MET) minutes per week; **MBSRQ**: Multidimensional Body-Self Relations Questionnaire (MBSRQ) subscale scores: Importance of Corporality (ISC), Behaviours aimed at maintaining physical shape (COMF), Self-evaluated physical attractiveness (AFA), and Physical appearance care (CAF); range 1–5, where higher scores indicate greater endorsement of the corresponding dimension; **Betts' QMI**: Overall Imagery Vividness score from the Betts' Questionnaire upon Mental Imagery; range 1–7, where a higher numerical score actually indicates lower imagery vividness; **SSQ6**: Statistics describing the participant's self-reported social network, as measured by the Social Support Questionnaire (Sex Ratio, Network Size, Group Count, Average Age of network, Age Difference with participant, and Mean & Standard deviation (SD) of people per questionnaire item); assesses perceived social support, including the number of available supportive people and satisfaction with that support, where higher scores indicate greater perceived availability of social support. Group 1: Low EDE-Q / Low IPAQ, Group 2: High EDE-Q / Low IPAQ, Group 3: Low EDE-Q / High IPAQ, Group 4: High EDE-Q / High IPAQ.

Characteristic	Group 1 (n=20)	Group 2 (n=21)	Group 3 (n=41)	Group 4 (n=19)	Overall (N=101)
Age (years)	24.25 (8.72)	25.81 (7.38)	25.10 (6.50)	25.11 (7.85)	25.08 (7.32)
Sex (F/M)	16 / 4	11 / 10	19 / 22	8 / 11	54 / 47
Gender (F/M/NB/O)	16 / 3 / 0 / 1	12 / 9 / 0 / 0	18 / 22 / 1 / 0	8 / 10 / 1 / 0	54 / 44 / 2 / 1
Height (cm)	169.55 (9.03)	168.52 (8.30)	171.32 (8.49)	169.58 (7.55)	170.06 (8.35)
Weight (kg)	65.35 (12.21)	65.76 (12.47)	67.85 (10.61)	62.37 (5.86)	65.89 (10.71)
Ideal Weight (kg)	62.06 (11.82)	62.67 (9.17)	65.95 (9.93)	62.76 (6.43)	63.96 (9.65)
Waist (cm)	75.35 (9.17)	78.33 (9.37)	78.05 (11.26)	73.47 (6.11)	76.71 (9.72)
Hip (cm)	91.70 (8.52)	92.19 (14.73)	93.51 (9.73)	90.79 (8.45)	92.37 (10.44)
Shoe Size (EU)	39.30 (2.39)	40.43 (2.98)	40.41 (2.52)	39.47 (3.91)	40.02 (2.90)
Education Level (1-5)	1.95 (1.36)	2.33 (1.49)	2.10 (1.28)	2.21 (1.18)	2.14 (1.31)
Gender Identity (1-7)	5.16 (1.89)	4.48 (1.83)	3.95 (1.80)	4.68 (1.92)	4.43 (1.87)
Ideal Gender Identity (1-7)	5.05 (1.75)	4.19 (1.83)	3.95 (1.80)	4.58 (2.04)	4.33 (1.86)
EDE-Q (0-6)	0.55 (0.44)	2.00 (0.82)	0.60 (0.41)	2.01 (0.66)	1.15 (0.90)
IPAQ (MET-min/week)	1802 (773)	2380 (1479)	7249 (5872)	6881 (5953)	5089 (5191)
MBSRQ ISC (1-5)	1.53 (0.37)	1.42 (0.37)	1.57 (0.26)	1.71 (0.33)	1.56 (0.33)
MBSRQ COMF (1-5)	1.26 (0.35)	1.59 (0.30)	1.56 (0.28)	1.63 (0.26)	1.52 (0.32)
MBSRQ AFA (1-5)	4.33 (0.75)	4.21 (1.02)	4.35 (0.85)	4.30 (0.91)	4.31 (0.87)
MBSRQ CAF (1-5)	4.68 (0.63)	4.28 (0.71)	4.60 (0.76)	4.77 (0.44)	4.58 (0.69)
Betts' QMI (1-7)	2.68 (1.25)	2.82 (0.70)	2.40 (0.63)	2.61 (0.74)	2.58 (0.83)
SSQ6: Sex Ratio	0.60 (0.26)	0.54 (0.19)	0.49 (0.24)	0.48 (0.21)	0.52 (0.23)
SSQ6: Network Size	4.80 (2.24)	4.86 (1.88)	4.66 (2.46)	4.32 (1.80)	4.66 (2.16)
SSQ6: Groups Count	1.25 (0.55)	1.81 (0.87)	2.02 (1.21)	1.95 (0.97)	1.81 (1.03)
SSQ6: Average Age	30.96 (12.35)	36.29 (8.37)	34.50 (7.98)	32.48 (10.01)	33.86 (9.37)
SSQ6: Age Difference	6.23 (7.13)	10.07 (8.95)	9.52 (8.95)	7.36 (7.52)	8.63 (8.37)
SSQ6: People/Item (Mean)	3.03 (1.25)	3.02 (0.96)	3.00 (1.38)	2.85 (0.76)	2.98 (1.16)
SSQ6: People/Item (SD)	1.16 (0.70)	1.20 (0.62)	1.04 (0.62)	1.11 (0.57)	1.11 (0.62)

and cognition: The relation between body representations and higher level cognitive and social processes. *Cortex*, 104:133–139, 2018.

[19] A. Doherty, D. Jackson, N. Hammerla, T. Plötz, P. Olivier, M. H. Granat, T. White, V. T. van Hees, M. I. Trenell, C. G. Owen, S. J. Preece, R. Gillions, S. Sheard, T. Peakman, S. Brage, and N. J. Wareham. Large Scale Population Assessment of Physical Activity Using Wrist Worn Accelerometers: The UK Biobank Study. *PLoS ONE*, 12(2):e0169649, 2017.

[20] Gianluca Finotti, Sara Garofalo, Marcello Costantini, and Dennis R. Proffitt. Temporal dynamics of the Rubber Hand Illusion. *Scientific Reports*, 13(1):6816, 2023.

[21] Nesrine Fourati and Catherine Pelachaud. Emilya: Emotional body expression in daily actions database. In Nicoletta Calzolari, Khalid Choukri, Thierry Declerck, Hrafn Loftsson, Bente Maegaard, Joseph Mariani, Asuncion Moreno, Jan Odijk, and Stelios Piperidis, editors, *Proceedings of the Ninth International Conference on Language Resources and Evaluation (LREC'14)*, pages 3486–3493, Reykjavik, Iceland, 2014. European Language Resources Association (ELRA).

[22] Shaun Gallagher. *How the Body Shapes the Mind*. Oxford University Press, Oxford, 2005.

[23] Rick O. Gilmore, Michele T. Diaz, Brad A. Wyble, and Tal Yarkoni. Progress toward openness, transparency, and reproducibility in cognitive neuroscience. *Annals of the New York Academy of Sciences*, 1396(1):5–18, 2017.

[24] Andrew H. Hales, Eric D. Wesselmann, and Joseph Hilgard. Improving Psychological Science through Transparency and Openness: An Overview. *Perspectives on Behavior Science*, 42(1):13–31, 2019.

[25] Freek Hens, Mohammad Mahdi Dehshibi, Leila Bagheriye, Ana Tajadura-Jiménez, and Mahyar Shahsavari. LAST-PAIN: Learning Adaptive Spike Thresholds for Low Back Pain Biosignals Classification. *IEEE Transactions on Neural Systems and Rehabilitation Engineering*, 33:1038–1047, 2025.

[26] Freek Hens, Amirhossein Sadough, Aleksa Bokšan, Mahyar Shahsavari, and Mohammad Mahdi Dehshibi. When one sensor fails: tolerating dysfunction in multisensor prototypes. *International Journal of Parallel, Emergent and Distributed Systems*, 0(0):1–14, 2026.

[27] Geert Hofstede, Gert Jan Hofstede, and Michael Minkov. *Cultures and organizations: Software of the mind*. McGraw-Hill, New York, 3rd edition, 2011.
[28] Andrea Kleinsmith and Nadia Bianchi-Berthouze. Affective Body Expression Perception and Recognition: A Survey. *IEEE Transactions on Affective Computing*, 4(1):15–33, 2013.
[29] Sander Koelstra, Christian Mühl, Mohammad Soleymani, Jong-Seok Lee, Ashkan Yazdani, Touradj Ebrahimi, Thierry Pun, Anton Nijholt, and Ioannis Patras. DEAP: A Database for Emotion Analysis Using Physiological Signals. *IEEE Transactions on Affective Computing*, 3(1):18–31, 2012.
[30] Bigna Lenggenhager, Tej Tadi, Thomas Metzinger, and Olaf Blanke. Video Ergo Sum: Manipulating Bodily Self-Consciousness. *Science*, 317(5841):1096–1099, 2007.
[31] Shun Li, Liqing Cui, Changye Zhu, Baobin Li, Nan Zhao, and Tingshao Zhu. Emotion recognition using Kinect motion capture data of human gaits. *PeerJ*, 4:e2364, 2016.
[32] Lara Maister, Mel Slater, Maria V. Sanchez-Vives, and Manos Tsakiris. Changing bodies changes minds: owning another body affects social cognition. *Trends in Cognitive Sciences*, 19(1):6–12, 2015.
[33] Zeltia Martínez-López, María Fernanda Páramo Fernández, María Adelina Guisande Couñago, Carolina Tinajero Vacas, Leandro da Silva Almeida, and María Soledad Rodríguez González. Apoyo social en universitarios españoles de primer año: propiedades psicométricas del Social Support Questionnaire-Short Form y el Social Provisions Scale. *Revista Latinoamericana de Psicología*, 46(2):102–110, 2014.
[34] Max Planck Institute for Intelligent Systems. Body Visualizer. https://bodyvisualizer.is.tue.mpg.de/, 2024. Accessed: November 11, 2026.
[35] Fritz Menzer, Anna Brooks, Pär Halje, Christof Faller, Martin Vetterli, and Olaf Blanke. Feeling in control of your footsteps: Conscious gait monitoring and the auditory consequences of footsteps. *Cognitive Neuroscience*, 1(3):184–192, 2010.
[36] Juan Abdon Miranda-Correa, Mojtaba Khomami Abadi, Nicu Sebe, and Ioannis Patras. AMIGOS: A Dataset for Affect, Personality and Mood Research on Individuals and Groups. *IEEE Transactions on Affective Computing*, 12(2):479–493, 2021.
[37] Florian 'Floyd' Mueller, Nadia Berthouze, Misha Sra, Mar Gonzalez-Franco, Henning Pohl, Susanne Boll, Richard Byrne, Arthur Caetano, Masahiko Inami, Jarrod Knibbe, Per Ola Kristensson, Xiang Li, Zhuying Li, Joe Marshall, Louise Petersen Matjeka, Minna Nygren, Rakesh Patibanda, Sara Price, Harald Reiterer, Aryan Saini, Oliver Schneider, Ambika Shahu, Phoebe O Toups Dugas, and Don Samitha Elvitigala. Grand Challenges around Designing Computers' Control Over Our Bodies. In *Proceedings of the 2026 CHI Conference on Human Factors in Computing Systems*, pages 1–22, New York, NY, USA, 2026. Association for Computing Machinery.
[38] Lauri Nummenmaa, Enrico Glerean, Riitta Hari, and Jari K. Hietanen. Bodily maps of emotions. *Proceedings of the National Academy of Sciences*, 111(2):646–651, 2014.
[39] Temitayo Olugbade, Raffaele Andrea Buono, Kyrill Potapov, Alex Bujorianu, Amanda C de C Williams, Santiago de Ossorno Garcia, Nicolas Gold, Catherine Holloway, and Nadia Bianchi-Berthouze. The EmoPain@Home Dataset: Capturing Pain Level and Activity Recognition for People With Chronic Pain in Their Homes. *IEEE Transactions on Affective Computing*, 16(2):471–484, 2025.
[40] OpenJS Foundation and Node-RED contributors. Node-RED: Low-code programming for event-driven applications. https://nodered.org/. Accessed: November 11, 2026.
[41] Jorge Ortigoso-Narro, Fernando Díaz-de María, Mohammad Mahdi Dehshibi, and Ana Tajadura-Jiménez. L-SFAN: Lightweight Spatially Focused Attention Network for Pain Behavior Detection. *IEEE Sensors Journal*, 25(10):18409–18418, 2025.
[42] María Angeles Peláez-Fernández, Francisco Javier Labrador, and Rosa Marí a Raich. Validation of Eating Disorder Examination Questionnaire (EDE-Q) – Spanish Version– for Screening Eating Disorders. *The Spanish Journal of Psychology*, 15(2):817–824, 2012.
[43] Valeria I. Petkova and H. Henrik Ehrsson. If I Were You: Perceptual Illusion of Body Swapping. *PLoS ONE*, 3(12):e3832, 2008.
[44] Catherine Preston and H. Henrik Ehrsson. Illusory Changes in Body Size Modulate Body Satisfaction in a Way That Is Related to Non-Clinical Eating Disorder Psychopathology. *PLOS ONE*, 9(1):e85773, 2014.
[45] Catherine Preston and H. Henrik Ehrsson. Illusory Obesity Triggers Body Dissatisfaction Responses in the Insula and Anterior Cingulate Cortex. *Cerebral Cortex*, 26(12):4450–4460, 2016.
[46] Lisa Raoul and Marie-Hélène Grosbras. Relating different dimensions of bodily experiences: Review and proposition of an integrative model relying on phenomenology, predictive brain and neuroscience of the self. *Neuroscience & Biobehavioral Reviews*, 148:105141, 2023.
[47] Marte Roel Lesur, Elena Bolt, Gianluca Saetta, and Bigna Lenggenhager. The monologue of the double: Allocentric reduplication of the own voice alters bodily self-perception. *Consciousness and Cognition*, 95:103223, 2021.
[48] Marte Roel Lesur, Laia Turmo Vidal, Matthew R. Longo, Jenny Slatman, Aleksander Väljamäe, and Ana Tajadura-Jiménez. Researching body perception: Toward an integration of quantitative and qualitative interdisciplinary approaches to address the multiplicity of bodily experiences. *Psychophysiology*, 63(1):e70234, 2026.
[49] Marte Roel Lesur, Laia Turmo Vidal, Matthew R. Longo, Jenny Slatman, Aleksander Väljamäe, and Ana Tajadura-Jiménez. Researching Body Perception: Toward an Integration of Quantitative and Qualitative Interdisciplinary Approaches to Address the Multiplicity of Bodily Experiences. *Psychophysiology*, 63(1):e70234, 2026.
[50] Blanca Román Viñas, Lourdes Ribas Barba, Joy Ngo, and Lluís Serra Majem. Validación en población catalana del cuestionario internacional de actividad física. *Gaceta Sanitaria*, 27(3):254–257, 2013.
[51] María Roncero, Conxa Perpiñá, Jose H. Marco, and Ser-

gio Sánchez-Reales. Confirmatory factor analysis and psychometric properties of the Spanish version of the Multidimensional Body-Self Relations Questionnaire-Appearance Scales. *Body Image*, 14:47–53, 2015.

[52] Karan Sharma, Claudio Castellini, Egon L. van den Broek, Alin Albu-Schäffer, and Friedhelm Schwenker. A dataset of continuous affect annotations and physiological signals for emotion analysis. *Scientific Data*, 6(1):196, 2019.

[53] V. Swami and D. Barron. Translation and validation of body image instruments: Challenges, good practice guidelines, and reporting recommendations for test adaptation. *Body Image*, 36:204–220, 2021.

[54] Pawel Tacikowski, Marieke L. Weijs, and H. Henrik Ehrsson. Perception of Our Own Body Influences Self-Concept and Self-Incoherence Impairs Episodic Memory. *iScience*, 23(9):101429, 2020.

[55] Ana Tajadura-Jiménez, Maria Basia, Ophelia Deroy, Merle Fairhurst, Nicolai Marquardt, and Nadia Bianchi-Berthouze. As Light As Your Footsteps: Altering Walking Sounds to Change Perceived Body Weight, Emotional State and Gait. In *Proceedings of the 33rd Annual ACM Conference on Human Factors in Computing Systems*, CHI '15, pages 2943–2952, New York, NY, USA, 2015. ACM.

[56] Ana Tajadura-Jiménez, Helen Cohen, and Nadia Bianchi-Berthouze. Bodily Sensory Inputs and Anomalous Bodily Experiences in Complex Regional Pain Syndrome: Evaluation of the Potential Effects of Sound Feedback. *Frontiers in Human Neuroscience*, 11:379, 2017.

[57] Ana Tajadura-Jiménez, Laura Crucianelli, Rebecca Zheng, Chloe Cheng, Judith Ley-Flores, Mercedes Borda-Más, Nadia Bianchi-Berthouze, and Aikaterini Fotopoulou. Body weight distortions in an auditory-driven body illusion in subclinical and clinical eating disorders. *Scientific Reports*, 12(1):20031, 2022.

[58] Ana Tajadura-Jiménez, Ophelia Deroy, Torsten Marquardt, Nadia Bianchi-Berthouze, Tomohisa Asai, Toshitaka Kimura, and Norimichi Kitagawa. Audio-tactile cues from an object's fall change estimates of one's body height. *PLoS ONE*, 13(6):e0199354, 2018.

[59] Ana Tajadura-Jiménez, Elena Márquez Segura, Laia Turmo Vidal, Kristina Höök, Kristi Kuusk, Pedro Lopes, and Mel Slater. Sustainable Body Transformation Experiences. In *Proceedings of the Extended Abstracts of the 2026 CHI Conference on Human Factors in Computing Systems*, pages 1–9, New York, NY, USA, 2026. Association for Computing Machinery.

[60] Ana Tajadura-Jiménez, Joseph Newbold, Linge Zhang, Patricia Rick, and Nadia Bianchi-Berthouze. As Light as You Aspire to Be: Changing Body Perception with Sound to Support Physical Activity. In *Proceedings of the 2019 CHI Conference on Human Factors in Computing Systems*, pages 1–14, New York, NY, USA, 2019. Association for Computing Machinery.

[61] Ana Tajadura-Jiménez, Manos Tsakiris, Torsten Marquardt, and Nadia Bianchi-Berthouze. Action sounds update the mental representation of arm dimension: contributions of kinaesthesia and agency. *Frontiers in Psychology*, 6:689, 2015.

[62] Ana Tajadura-Jiménez, Aleksander Väljamäe, Isao Toshima, Tadashi Kimura, Manos Tsakiris, and Norimichi Kitagawa. Action sounds recalibrate perceived tactile distance. *Current Biology*, 22(13):R516–R517, 2012.

[63] Laia Turmo Vidal, Ana Tajadura-Jiménez, and Judith Ley-Flores. Temporal Trajectories: Characterizing Somatic Experiences that Unfold Over Time. In *Proceedings of the 2025 ACM Designing Interactive Systems Conference*, pages 2931–2949, New York, NY, USA, 2025. Association for Computing Machinery.

[64] Laia Turmo Vidal, Ana Tajadura-Jiménez, José Manuel Vega-Cebrián, Judith Ley-Flores, Joaquin R. Díaz-Durán, and Elena Márquez Segura. Body Transformation: An Experiential Quality of Sensory Feedback Wearables for Altering Body Perception. In *Proceedings of the Eighteenth International Conference on Tangible, Embedded, and Embodied Interaction*, pages 1–19, New York, NY, USA, 2024. ACM.

[65] B. Van der Hoort, A. Guterstam, and H. H. Ehrsson. Being Barbie: the size of one's own body determines the perceived size of the world. *PLoS ONE*, 6(5):e20195, 2011.

[66] Marieke L. Weijs, Marte Roel Lesur, Moritz M. Daum, and Bigna Lenggenhager. Keeping up with ourselves: Multimodal processes underlying body ownership across the lifespan. *Cortex*, 177:209–223, 2024.

[67] Hang Yuan, Shing Chan, Andrew P. Creagh, Catherine Tong, Aidan Acquah, David A. Clifton, and Aiden Doherty. Self-supervised learning for human activity recognition using 700,000 person-days of wearable data. *npj Digital Medicine*, 7(1):91, 2024.

[68] Yuliang Zhao, Huawei Zhang, Jian Li, Siyang Song, Chao Lian, Yinghao Liu, Yulin Wang, and Changzeng Fu. Multimodal Depression Assessment Framework Integrating Personality and Gait for Older Adults With Medical Conditions. *IEEE Transactions on Affective Computing*, 16(3):2048–2061, 2025.