

Exact Recovery Thresholds for Weighted Data Selection in Vector-Valued Linear Regression

Guangjian Zhang
zgj1226029469@outlook.com

August 31, 2026

Abstract

We resolve the threshold part of Question 4 of the COLT 2025 open problem “Data Selection for Regression Tasks” of Hanneke, Moran, Shlimovich and Yehudayoff. In vector-valued linear regression with square loss $\ell_{(x,y)}(W) = \|Wx - y\|_2^2$, where $x \in \mathbb{R}^d$, $y \in \mathbb{R}^m$ and the learner is the empirical risk minimizer of minimal Frobenius norm, we prove that the minimal budget of weighted examples that recovers the full-data loss on *every* finite dataset is exactly

$$n^*(d, m) = (m + 1)d.$$

We further determine two more values of the weighted selection profile $F_{\text{weighted}}(d, m, n)$: at the near-threshold budget, $F_{\text{weighted}}(d, m, (m + 1)d - 1) = 1 + \frac{1}{dm^2}$, and at the spanning budget, $F_{\text{weighted}}(d, m, d) = d + 1$ for every m , while $F_{\text{weighted}}(d, m, n) = \infty$ for $n < d$. For the smallest open intermediate cell $(d, m) = (2, 2)$ we prove $F_{\text{weighted}}(2, 2, 3) \in [13/8, 15/8]$ and $F_{\text{weighted}}(2, 2, 4) \in [5/4, 3/2]$, reduce the conjectured exact values $13/8$ and $5/4$ to a finite moment problem on the circle with at most seven atoms, and establish strong structural evidence for the conjecture. The upper-bound techniques (a fixed-basis conic compression lemma, a determinant–facet rigidity theorem for maximal certificates, and sharp sparsification lemmas for zero-mean weighted point systems) are of independent interest. As a byproduct we correct an erroneous claim circulating in a recent refereed preprint, exhibiting an explicit dataset with $m = 2$ on which *no* weighted selection of $2d$ points recovers the optimal loss. All results are new only for $m \geq 2$; the scalar case $m = 1$ is due to Hanneke et al.

1 Introduction

How well can a fixed, natural learning rule perform when it is trained on only n examples selected from a larger dataset? Hanneke, Moran, Shlimovich and Yehudayoff posed this data-selection question for basic regression tasks in a COLT 2025 open-problem note [HMSY25b] and companion paper [HMSY25a]. For scalar linear regression they obtained a complete taxonomy: with weighted selection and the minimum-norm empirical risk minimizer, the worst-case ratio between the loss of the selected model and the optimal full-data loss equals 1 for $n \geq 2d$, equals $d + 1$ at $n = d$, and is infinite for $n < d$ [HMSY25b, Theorem 1]. Their note concludes with a unified *vector-valued* formulation (predictors $W: \mathbb{R}^d \rightarrow \mathbb{R}^m$, square loss, minimum *Frobenius*-norm ERM) and asks, as Question 4:

Given d, m, n , what is the value of $F_{\text{weighted}}(d, m, n)$? In particular, what is the minimal $n = n(d, m)$ such that this ratio equals 1?

This paper resolves the threshold question completely and determines several further values of the profile $F_{\text{weighted}}(d, m, \cdot)$.

1.1 Main results

Throughout, $D = \{z_i = (x_i, y_i)\}_{i=1}^N \subseteq \mathbb{R}^d \times \mathbb{R}^m$ is a finite multiset, $\ell_z(W) = \|Wx - y\|_2^2$ for $W \in \mathbb{R}^{m \times d}$, $L_D(W) = \frac{1}{N} \sum_i \ell_{z_i}(W)$, $L_D^* = \min_W L_D(W)$, and A is the ERM returning the empirical risk minimizer of minimal Frobenius norm. Weighted selection of budget n chooses $z_1, \dots, z_n \in D$ (repetitions allowed) and a convex combination $F \in \text{conv}(\ell_{z_1}, \dots, \ell_{z_n})$, and suffers loss $L_D(A(F))$ on the full dataset; $F_{\text{weighted}}(d, m, n)$ is the supremum over datasets of the ratio $\inf_F L_D(A(F))/L_D^*$ (Section 2 records the degenerate-ratio conventions). Let $n^*(d, m) = \min\{n : F_{\text{weighted}}(d, m, n) = 1\}$.

Theorem 1.1 (Exact threshold; Theorem 3.7). *For all $d, m \geq 2$,*

$$n^*(d, m) = (m + 1)d.$$

Moreover, for every dataset with feature rank r , some $(m + 1)r$ weighted points already recover a full-data optimum exactly, and there is an explicit integer dataset on $(m + 1)d$ points for which every selection of $(m + 1)d - 1$ weighted points has ratio exactly $1 + \frac{m+1}{2md(m^2+m-1)} > 1$.

Theorem 1.2 (Near-threshold value; Theorem 5.4). *For all $d \geq 1$, $m \geq 1$,*

$$F_{\text{weighted}}(d, m, (m + 1)d - 1) = 1 + \frac{1}{dm^2}.$$

Theorem 1.3 (Low budgets; Theorem 4.1). *For all $d, m \geq 1$: $F_{\text{weighted}}(d, m, n) = \infty$ for $n < d$, and*

$$F_{\text{weighted}}(d, m, d) = d + 1,$$

independently of m .

Theorem 1.4 (Smallest intermediate cell; Theorem 7.7). *$F_{\text{weighted}}(2, 2, 3) \in [\frac{13}{8}, \frac{15}{8}]$ and $F_{\text{weighted}}(2, 2, 4) \in [\frac{5}{4}, \frac{3}{2}]$.*

We conjecture that the lower ends are the truth, $F_{\text{weighted}}(2, 2, 3) = \frac{13}{8}$ and $F_{\text{weighted}}(2, 2, 4) = \frac{5}{4}$ (Conjecture 7.8), and we reduce this conjecture to a concrete finite question: a moment problem for at most seven atoms on the unit circle (Section 7). The evidence assembled there includes a complete solution of the two-direction class with a rigidity description of its equality systems, a proof that this equality fiber is a local maximum in a stratified sense, two exact obstructions showing which proof strategies *cannot* work, and exact extremal records for small systems.

Consistency checks: at $m = 1$ Theorems 1.1 and 1.2 specialize to $n^*(d, 1) = 2d$ and $F_w(d, 2d - 1) = 1 + \frac{1}{d}$, matching [HMSY25b, Theorem 1] and the value announced there; at $d = 1$, $x \equiv 1$ the problem contains weighted mean estimation in $\mathbb{R}^m$, and $n^*(1, m) = m + 1$ is the Carathéodory bound.

Scope of novelty. For $m = 1$ all three regimes of Theorem 1.3 and the threshold $n^*(d, 1) = 2d$ were established by Hanneke et al. [HMSY25b, HMSY25a] (their Theorem 8 and Example 2 give the exact scalar taxonomy), and the value $1 + \frac{1}{d}$ at $n = 2d - 1$, announced in [HMSY25b], is proved—together with further exact values in the scalar intermediate regime—in the companion paper [Zha26]. **All results of this paper are claimed as new only for $m \geq 2$;** our proofs happen to cover $m = 1$ uniformly, which we use purely as a consistency check.

1.2 Techniques

The threshold upper bound comes from a *fixed-basis conic compression* lemma (Lemma 3.1): after pinning a feature basis B of r points, the negated sum of their residual dyads is compressed inside the cone of the remaining dyads by conic Carathéodory, at cost mr ; together with the basis this yields an exact certificate on $(m+1)r$ points. Notably, this argument is simpler than the Steinitz-based route used for the scalar upper bound in [HMSY25a], and gives a data-dependent bound.

The near-threshold upper bound requires understanding datasets whose minimal exact certificates have the maximal size $(m+1)d$. We prove a *determinant–facet rigidity theorem* (Theorem 5.1): every facet of the normalized certificate polytope contributes a distinct linear factor to the degree- d polynomial $u \mapsto \det(\sum_i u_i x_i x_i^\top)$, forcing the polytope to be a simplex and the dataset to decompose into d disjoint $(m+1)$ -point circuits supported on d independent feature lines. A sharp *k -point mean lemma* for zero-mean weighted point systems (Lemma 5.3 for $k = m$; Theorem 6.1 in general) then finishes the assembly.

For the intermediate regime at $(2, 2)$ we develop a complex-variable dictionary that identifies whitened instances with moment systems (p_i, t_i, z_i) on the unit circle and the selection cost with an explicit convex combination of *two-point interpolation coefficients* (Lemma 7.2), an identity that absorbs the ill-conditioning factor $(1 - |\mu|^2)^{-1}$ entirely.

1.3 Related and concurrent work

Published work. The problem and all scalar results are from [HMSY25b, HMSY25a]. The budget- d upper bound uses volume sampling and the exact expectation formulas of Dereziński and Warmuth [DW17]; we are careful to use their Theorem 5 in its correct form (equality under general position, inequality in general; see Section 4). The general k -point mean lemma relies on the full-dimensional Grace–Danielsson inequality (Egan’s conjecture), proved by Drozdov [Dro25]; none of our main theorems depend on that lemma beyond the elementary case $k = m$, for which we give a short self-contained proof (Lemma 5.3).

Concurrent unrefereed preprints. This open-problem family is being attacked by several automated pipelines, and a number of unrefereed, apparently machine-generated preprints on other questions of [HMSY25b] appeared recently: a preprint resolving Question 3 (unweighted selection, general m) [P⁺26]; two short manuscripts on Question 1 (mean estimation), one of which determines the budget-2 column of the mean-estimation table [Ano26b, Ano26a]; and a manuscript on the scalar weighted regime of Question 2 [Dew26]. None of these treats the weighted vector-valued Question 4, whose threshold and profile values are the subject of this paper. Since [Ano26b] advertises convex-geometric lemmas about few-atom distributions obtained by a variance-minimizing Carathéodory reduction, we note the demarcation explicitly: our sparsification results (Lemma 5.3, Theorem 6.1) were obtained independently, are stated for zero-mean weighted systems with the sharp constant $\frac{M+1-k}{kM}$, and are used here as components of the F_{weighted} analysis; we make no priority claim on auxiliary convex-geometry statements that may overlap, and our headline results — the complete $(m+1)d$ characterization and the profile values — are not touched by any of the works above.

Remark 1.5 (Correction of a circulating claim). Remark 1.2 of [P⁺26] asserts, in the notation $F^w(d, m, n)$, that the weighted scalar trichotomy of [HMSY25b, Theorem 1] — in particular, ratio 1 for all $n \geq 2d$ — holds for general m , citing [HMSY25b] as the source. This is a misattribution: [HMSY25b] proves the trichotomy only for $m = 1$, and the assertion is *false* for

every $m \geq 2$. By Theorem 1.1, exact recovery requires $(m+1)d > 2d$ points, and quantitatively $F_{\text{weighted}}(d, m, 2d) \geq F_{\text{weighted}}(d, m, (m+1)d - 1) = 1 + \frac{1}{dm^2} > 1$ by monotonicity. We make the failure concrete in Proposition 3.9: on an explicit 6-point integer dataset with $d = m = 2$, every weighted selection of $2d = 4$ points has full-data loss at least $\frac{23}{20} \cdot L_D^*$, with equality attained.

1.4 Organization

Section 2 fixes definitions and records the basic structural lemmas. Section 3 proves the threshold theorem and the correction of Remark 1.5. Section 4 handles budgets $n \leq d$. Section 5 proves the near-threshold value, via the rigidity theorem. Section 6 proves the general k -point mean lemma. Section 7 develops the $(2, 2)$ intermediate regime: the interval theorem, the reduction of the conjectured exact values to a seven-atom moment problem, and the evidence. Section 8 discusses the general program and open problems.

2 Preliminaries

2.1 The model

A dataset is a finite multiset $D = \{z_i = (x_i, y_i)\}_{i=1}^N \subseteq \mathbb{R}^d \times \mathbb{R}^m$ with $N \geq 1$. A linear predictor is a matrix $W \in \mathbb{R}^{m \times d}$, with loss $\ell_z(W) = \|Wx - y\|_2^2$ and average loss $L_D(W) = \frac{1}{N} \sum_{i=1}^N \ell_{z_i}(W)$; write $L_D^* = \min_W L_D(W)$. The learning rule A maps a nonnegatively weighted objective $F = \sum_i c_i \ell_{z_i}$ (with $c_i \geq 0$, $\sum_i c_i = 1$) to the minimizer of F of minimal Frobenius norm. Weighted selection with budget n is the value

$$L_D^*(n; \text{weighted}) = \inf_{\substack{z_{j_1}, \dots, z_{j_n} \in D \\ F \in \text{conv}(\ell_{z_{j_1}}, \dots, \ell_{z_{j_n}})}} L_D(A(F)),$$

and

$$F_{\text{weighted}}(d, m, n) = \sup_{D \subseteq \mathbb{R}^d \times \mathbb{R}^m} \frac{L_D^*(n; \text{weighted})}{L_D^*}, \quad n^*(d, m) = \min\{n : F_{\text{weighted}}(d, m, n) = 1\}.$$

Following the convention of [HMSY25a, Theorem 1] we set the ratio to 1 when numerator and denominator are both 0, and to ∞ when only the denominator is 0. We write the inner optimization as an infimum (as in [HMSY25a]); all our upper-bound certificates are attained, and our lower bounds bound the infimum, so nothing depends on attainment.

Repetitions among the $z_{j_1}, \dots, z_{j_n}$ are allowed. The following normalization is immediate and used silently.

Lemma 2.1 (Selection semantics). *Selections of budget n are in value-preserving correspondence with weight vectors $c \in \mathbb{R}_{\geq 0}^N$, $\sum_i c_i = 1$, $|\text{supp } c| \leq n$: repeated picks merge weights, zero-weight picks can be discarded, and any c with $|\text{supp } c| \leq n$ is realizable with exactly n slots by repeating a chosen point and splitting its weight.*

2.2 Row decomposition and minimum-norm geometry

Lemma 2.2 (Row decomposition). *Write $w_j^\top$ for the j -th row of W and y_{ij} for the j -th entry of y_i . Then $F_c(W) = \sum_{j=1}^m \sum_i c_i (\langle w_j, x_i \rangle - y_{ij})^2$ and $\|W\|_F^2 = \sum_j \|w_j\|^2$. Hence the minimizer set of F_c is the Cartesian product over rows of scalar weighted least-squares solution sets, and*

$A(F_c)$ consists of the minimal- ℓ_2 -norm scalar solutions row by row — with the single shared weight vector c .

Proof. Both the objective and the squared Frobenius norm are additive across rows, and a product set is minimized in norm coordinate-wise. $\square$

Let $T = \text{span}\{x_1, \dots, x_N\}$, $r = \dim T$, and let P_T denote the orthogonal projection onto T . Let W° be the minimal-Frobenius-norm full-data minimizer, and set $\rho_i = W^\circ x_i - y_i$ (residuals) and $M_i = \rho_i x_i^\top \in \mathbb{R}^{m \times d}$ (residual dyads).

Lemma 2.3 (Full-data geometry). $W^\circ = W^\circ P_T$; the first-order condition $\sum_{i=1}^N M_i = 0$ holds; and for every $H \in \mathbb{R}^{m \times d}$,

$$L_D(W^\circ + H) = L_D^* + \frac{1}{N} \sum_{i=1}^N \|H x_i\|^2. \quad (1)$$

Consequently the full-data minimizer set is $\{W^\circ + H : H P_T = 0\}$; if $T = \mathbb{R}^d$ the minimizer is unique.

Proof. Predictions depend on W only through $W P_T$, and $\|W\|_F^2 = \|W P_T\|_F^2 + \|W P_{T^\perp}\|_F^2$, so minimality of the norm forces $W^\circ P_{T^\perp} = 0$. The condition $\sum_i M_i = 0$ is the vanishing gradient of the smooth convex L_D at W° . Expanding $L_D(W^\circ + H)$, the cross term is $\frac{2}{N} \langle H, \sum_i M_i \rangle_F = 0$, giving (1); equality in (1) holds iff $H x_i = 0$ for all i , i.e. $H P_T = 0$. $\square$

Lemma 2.4 (Exact certificate). Let $c \in \mathbb{R}_{\geq 0}^N$ have support S with $c_i > 0$ on S . If

$$\sum_{i \in S} c_i M_i = 0 \quad \text{and} \quad \text{span}\{x_i : i \in S\} = T, \quad (2)$$

then $A(F_c) = W^\circ$; in particular the selection realizes the full-data optimal loss, and the ratio is 1 (also when $L_D^* = 0$, by the 0/0 convention).

Proof. Using (2), for any H , $F_c(W^\circ + H) - F_c(W^\circ) = 2 \langle H, \sum_{i \in S} c_i M_i \rangle_F + \sum_{i \in S} c_i \|H x_i\|^2 = \sum_{i \in S} c_i \|H x_i\|^2$, which vanishes iff $H x_i = 0$ on S , iff $H P_T = 0$ (the support spans T). So the weighted minimizer set is $W^\circ + \{H : H P_T = 0\}$, the same set as the full-data minimizer set. Since $W^\circ = W^\circ P_T$ and $H = H P_{T^\perp}$ are Frobenius-orthogonal, the unique minimal-norm element is W° . $\square$

We call a weight vector satisfying (2) a *spanning zero certificate*. Define

$$\tau(D) = \min\{|\text{supp } c| : c \text{ is a spanning zero certificate}\}. \quad (3)$$

Lemma 2.5 (Scalar embedding). If all m columns of the labels are equal, $y_i = t_i \mathbf{1}_m$, then each row of the problem is the same scalar problem and $L_{\tilde{D}}(\cdot) = m L_{D_0}(\cdot)$ termwise. Consequently $F_{\text{weighted}}(d, m, n) \geq F_w(d, n)$ for all n , where $F_w(d, n)$ denotes the scalar weighted profile of [HMSY25b].

Proof. By Lemma 2.2 every row runs the identical scalar min-norm ERM with the shared weights, so numerator and denominator both scale by m . $\square$

3 The exact threshold $(m + 1)d$

3.1 Upper bound: fixed-basis conic compression

Lemma 3.1 (Fixed-basis conic compression). *Let $a_1, \dots, a_N$ be elements of a finite-dimensional real vector space with $\sum_{i=1}^N a_i = 0$, and let $B \subseteq [N]$. Put $q_B = \dim \text{span}\{a_i : i \notin B\}$. Then there exist coefficients $c_i \geq 0$ with $c_b > 0$ for all $b \in B$, $\sum_i c_i a_i = 0$, and $|\text{supp } c| \leq |B| + q_B$.*

Proof. From the global zero sum, $-\sum_{b \in B} a_b = \sum_{i \notin B} a_i$, which is a nonnegative combination of $\{a_i\}_{i \notin B}$ and hence lies in $\text{cone}\{a_i : i \notin B\}$. By conic Carathéodory (see e.g. [Ber09, Prop. 1.2.1(a)]), every element of this cone is a positive combination of a *linearly independent* subfamily, so $-\sum_{b \in B} a_b = \sum_{i \in C} \alpha_i a_i$ with $\alpha_i > 0$, $C \subseteq [N] \setminus B$ and $|C| \leq q_B$ (if the target is 0, take $C = \emptyset$). Set $c_b = 1$ on B , $c_i = \alpha_i$ on C , and 0 elsewhere. $\square$

Theorem 3.2 (Data-dependent upper bound). *Every dataset with feature rank $r \geq 1$ admits a spanning zero certificate of support at most $(m + 1)r$; hence $\tau(D) \leq (m + 1)r$ and $L_D^*((m + 1)d; \text{weighted}) = L_D^*$ for every dataset. If $r = 0$, a single point suffices.*

Proof. Pick $B \subseteq [N]$ with $|B| = r$ such that $\{x_b\}_{b \in B}$ is a basis of T (possible: the features lie in and span T). Apply Lemma 3.1 to the residual dyads $a_i = M_i$, whose global sum vanishes (Lemma 2.3). Every $M_i = \rho_i x_i^\top$ has all rows proportional to $x_i^\top \in T$, so $\text{span}\{M_i : i \notin B\} \subseteq \mathbb{R}^m \otimes T$, whence $q_B \leq mr$. The resulting c is strictly positive on a set $S \supseteq B$ of size at most $r + mr = (m + 1)r$, satisfies $\sum_{i \in S} c_i M_i = 0$, and $\text{span}\{x_i : i \in S\} \supseteq \text{span}\{x_b : b \in B\} = T$; since all features lie in T , the span equals T . Normalizing c to sum 1 (its support is nonempty) and invoking Lemma 2.4 gives $A(F_c) = W^\circ$, so the selected model attains the optimal full-data loss; by Lemma 2.1 the support fits in $(m + 1)d \geq (m + 1)r$ slots. If $r = 0$ then all $x_i = 0$: every W has the same loss on every objective, and A returns $W = 0$ in both cases; one point suffices. $\square$

3.2 Lower bound: the axial simplex instance

Fix $d, m \geq 1$ and let $u = \mathbf{1}_m$,

$$v_j = e_j \quad (1 \leq j \leq m), \quad v_{m+1} = -u,$$

so $\sum_{j=1}^{m+1} v_j = 0$, and the unique linear dependence among $v_1, \dots, v_{m+1}$ has all coefficients equal; in particular no proper subset of $\{v_j\}$ admits a nonzero nonnegative zero-sum, equivalently $0 \notin \text{conv}$ of any proper subset.

Definition 3.3 (Integer axial instance). $D_{d,m}$ consists of the $N = (m + 1)d$ points $z_{ij} = (e_i, u - v_j)$ for $i \in [d]$, $j \in [m + 1]$; explicitly $y_{ij} = u - e_j$ for $j \leq m$ and $y_{i,m+1} = 2u$.

Lemma 3.4. *For $D_{d,m}$: the unique full-data minimizer is $W^* = u \mathbf{1}_d^\top$, with residual v_j at z_{ij} and $L_D^* = \frac{2m}{m+1}$. Moreover, for any weight vector c with per-axis totals $t_i = \sum_j c_{ij}$, the returned model $\widehat{W} = A(F_c)$ satisfies: column i of $\widehat{W}$ equals the conditional weighted label mean $u - a_i$ with $a_i = \sum_j \frac{c_{ij}}{t_i} v_j$ when $t_i > 0$, and equals 0 when $t_i = 0$; and, writing $\delta_i = \widehat{W} e_i - u$,*

$$L_D(\widehat{W}) = L_D^* + \frac{1}{d} \sum_{i=1}^d \|\delta_i\|^2. \quad (4)$$

Proof. The full Gram matrix is $\sum_{ij} e_i e_i^\top = (m+1)I \succ 0$ and $\sum_{ij} v_j e_i^\top = 0$, so W^* is the unique minimizer; $L_D^* = \frac{d \sum_j \|v_j\|^2}{N} = \frac{2md}{(m+1)d}$. Since the features are standard basis vectors, the weighted normal equations decouple across columns; a positive-weight column returns the conditional mean, and a zero-weight column is unconstrained, hence zeroed by the Frobenius rule (column-wise decomposition of the norm). For (4), expand $\|\delta_i + v_j\|^2$ and use $\sum_j v_j = 0$: $\sum_j \|v_j + \delta_i\|^2 = 2m + (m+1)\|\delta_i\|^2$. $\square$

Lemma 3.5 (Facet distances). *Let $Q_m = m^2 + m - 1$. The squared distance from the origin to the convex hull of any proper subset of $\{v_1, \dots, v_{m+1}\}$ is at least $1/Q_m$; the minimum is attained on the facets omitting some e_k , while the facet omitting v_{m+1} has squared distance $1/m$.*

Proof. Every proper subset is contained in a facet. For the facet $\text{conv}\{e_1, \dots, e_m\}$ the nearest point is u/m , of squared norm $1/m$. For the facet omitting e_k , let n_k have $-m$ in coordinate k and 1 elsewhere; the facet lies on the hyperplane $n_k^\top x = 1$ and $\|n_k\|^2 = Q_m$, so the distance to the hyperplane is $1/\sqrt{Q_m}$; the foot n_k/Q_m lies in the facet since $n_k/Q_m = \frac{m}{Q_m}(-u) + \sum_{j \neq k} \frac{m+1}{Q_m} e_j$ is a convex combination. Finally $Q_m \geq m$ with equality only at $m = 1$ (where both distances equal $1 = 1/Q_1$). $\square$

Theorem 3.6 (Exact deficit of the axial instance). *For every $d, m \geq 1$,*

$$\frac{L_{D_{d,m}}^*((m+1)d - 1; \text{weighted})}{L_{D_{d,m}}^*} = 1 + \frac{m+1}{2md(m^2 + m - 1)} > 1.$$

Proof. A selection of support at most $N - 1$ either misses some axis entirely ($\|\delta_i\|^2 = \|u\|^2 = m \geq 1/Q_m$) or uses a proper subset of the $m+1$ points of some axis, in which case $\delta_i = -a_i$ lies in minus the convex hull of a proper subset of $\{v_j\}$ and $\|\delta_i\|^2 \geq 1/Q_m$ by Lemma 3.5. With (4), every such selection has $L_D \geq L_D^* + \frac{1}{dQ_m}$. Conversely, omit $z_{1,1}$, give axis 1 the conditional weights of the foot n_1/Q_m (namely $\frac{m}{Q_m}$ on v_{m+1} and $\frac{m+1}{Q_m}$ on each v_j , $2 \leq j \leq m$, scaled by $t_1 = \frac{1}{d}$) and every other axis uniform weights: the support is exactly $N - 1$ and $L_D = L_D^* + \frac{1}{dQ_m}$. Dividing by $L_D^* = \frac{2m}{m+1}$ gives the claim. $\square$

Theorem 3.7 (Threshold). *$n^*(d, m) = (m+1)d$ for all $d, m \geq 1$, and $F_{\text{weighted}}(d, m, n) = 1$ for all $n \geq (m+1)d$.*

Proof. Combine Theorem 3.2 (ratio 1 at budget $(m+1)d$, every dataset) with Theorem 3.6 (some dataset has ratio > 1 at budget $(m+1)d - 1$). $\square$

Corollary 3.8. *$n^*(2, 2) = 6$. On the instance $D_{2,2}$ (six integer points, labels $(0, 1), (1, 0), (2, 2)$ over each of e_1, e_2) the optimal five-point ratio is exactly $\frac{43}{40}$.*

We emphasize that $43/40$ is the exact optimum of this instance; the worst case over all datasets at budget 5 is $F_{\text{weighted}}(2, 2, 5) = 1 + \frac{1}{8} = \frac{9}{8}$ by Theorem 5.4.

3.3 Failure at $n = 2d$ for $m \geq 2$

As announced in Remark 1.5, the scalar rule “ $n \geq 2d$ suffices” does not survive vector outputs.

Proposition 3.9. *For all $d \geq 1, m \geq 2$, $F_{\text{weighted}}(d, m, 2d) \geq F_{\text{weighted}}(d, m, (m+1)d - 1) = 1 + \frac{1}{dm^2} > 1$, since $2d \leq (m+1)d - 1$ and F_{weighted} is nonincreasing in n . Concretely, for $d = m = 2$ the instance $D_{2,2}$ of Corollary 3.8 satisfies: every weighted selection of at most $2d = 4$ points has*

$$L_D(A(F)) \geq \frac{23}{20} \cdot L_D^*,$$

with equality attained (allocate two points per axis, at conditional weights realizing the nearest facet points).

Proof. The monotonicity statement is immediate from the definitions. For $D_{2,2}$, by (4) and per-axis decoupling, a budget-4 selection allocates (k_1, k_2) points to the two axes with $k_1 + k_2 \leq 4$; the minimal per-axis costs are 0 (all three points), $1/5$ (best two points, the facet $\{e_2, -u\}$ or $\{e_1, -u\}$ at squared distance $1/Q_2 = 1/5$), 1 (best single point), and 2 (empty axis). The minimum total cost over allocations is $2 \cdot \frac{1}{5}$ at $(2, 2)$, giving $L_D = \frac{4}{3} + \frac{1}{2} \cdot \frac{2}{5} = \frac{23}{15}$ and ratio $\frac{23/15}{4/3} = \frac{23}{20}$. (We verified this enumeration by exact symbolic computation as well.) $\square$

4 Budgets $n \leq d$

Theorem 4.1. *For all $d, m \geq 1$: $F_{\text{weighted}}(d, m, n) = \infty$ for every $n < d$, and $F_{\text{weighted}}(d, m, d) = d + 1$.*

The lower bounds are inherited from the scalar case through Lemma 2.5: by [HMSY25b, Theorem 1], $F_w(d, n) = \infty$ for $n < d$ and $F_w(d, d) = d + 1$, so $F_{\text{weighted}}(d, m, n) \geq F_w(d, n)$. The content of this section is the upper bound $F_{\text{weighted}}(d, m, d) \leq d + 1$ uniformly in m , which requires selecting one subset that serves all m output rows simultaneously.

Volume sampling. Let $X \in \mathbb{R}^{d \times N}$ have full row rank and let $y \in \mathbb{R}^N$. Size- d volume sampling draws a subset S of d column indices with probability proportional to $\det(X_S)^2$, and $w(S)$ denotes the least-squares solution of the subproblem (X_S, y_S) (the interpolant when X_S is invertible). Dereziński and Warmuth [DW17] proved: (i) unbiasedness $\mathbb{E}[w(S)] = w^*$ for every label vector, requiring only full row rank (their Theorem 3 and Proposition 7); and (ii) for the total square loss L on the full data,

$$\mathbb{E}[L(w(S))] \leq (d + 1) L(w^*), \quad (5)$$

with equality when X is in general position (every d -column submatrix nonsingular) — their Theorem 5. We only use the inequality (5), which holds without general position; note that pointwise equality genuinely fails for degenerate X (e.g. $x_1 = x_2 = e_1$, $x_3 = x_4 = e_2$, $y = (1, -1, 1, -1)$ has $\mathbb{E}[L(w(S))] = 2L(w^*)$). Dividing by N converts (5) to average losses.

Proof of Theorem 4.1. Only $F_{\text{weighted}}(d, m, d) \leq d + 1$ remains. Fix a dataset; let $r = \text{rank}$ of the features. If $r = 0$ the ratio is 1 (both rules return $W = 0$). If $L_D^* = 0$, choose r points whose features form a basis of T with positive weights: the weighted problem has minimum 0, its minimizer set is $W^\circ + \{H : HP_T = 0\}$, and the Frobenius rule returns W° ; the ratio is 1 by the 0/0 convention.

Otherwise pick an orthonormal basis $U \in \mathbb{R}^{d \times r}$ of T and write $x_i = U\xi_i$ with $\Xi = [\xi_1 \cdots \xi_N] \in \mathbb{R}^{r \times N}$ of full row rank. Run size- r volume sampling on Ξ ; crucially, *the distribution over subsets*

depends only on the features. For the q -th output row, (5) (in dimension r) gives $\mathbb{E} L_D^{(q)}(w_{q,S}) \leq (r+1)L_D^{(q),*}$. Summing over the m rows and using Lemma 2.2,

$$\mathbb{E}[L_D(W_S)] \leq (r+1)L_D^* \leq (d+1)L_D^*,$$

where $W_S = \Theta_S U^\top$ is the matrix whose rows are the per-row subproblem solutions in Ξ -coordinates with the $T^\perp$ -component set to zero. Hence some subset S (of $r \leq d$ points) achieves $L_D(W_S) \leq (d+1)L_D^*$. This W_S is realizable by a legal selection: the r chosen features are linearly independent in T , so with any positive weights the weighted ERM set consists of all interpolants of the r selected points, and the Frobenius rule returns exactly W_S (the interpolant with vanishing $T^\perp$ -component). Padding to d slots by Lemma 2.1 completes the proof. $\square$

5 The near-threshold value $1 + 1/(dm^2)$

Throughout this section fix a dataset with residual dyads $M_i = \rho_i x_i^\top$ and recall the minimal certificate size $\tau(D)$ from (3); Theorem 3.2 gives $\tau(D) \leq (m+1)r$. The upper bound at budget $(m+1)d - 1$ requires understanding the datasets with $\tau(D) = (m+1)d$, which we call *hard*. The key is a rigidity theorem for maximal certificates.

5.1 Rigidity of maximal certificates

Theorem 5.1 (Determinant–facet rigidity). *Let the features span $\mathbb{R}^d$ and suppose $\tau(D) = k := (m+1)d$. Let c be a spanning zero certificate with $|\text{supp } c| = k$ and support S . Then S splits uniquely (up to permutation) into d pairwise disjoint blocks $S = C_1 \sqcup \dots \sqcup C_d$ with $|C_j| = m+1$, where each C_j supports a positive circuit of the dyads whose features span a one-dimensional line U_j , and $U_1, \dots, U_d$ are linearly independent.*

Proof. Vectorize the dyads into $A = [\text{vec}(M_i)]_{i \in S} \in \mathbb{R}^{md \times k}$ and let $K = \ker A \cap \mathbb{R}_{\geq 0}^k$, $h = \dim \ker A \geq k - md = d$. Since $c > 0$ on S lies in $\ker A$, a neighborhood of c in $\ker A$ stays nonnegative, so K spans $\ker A$ and the compact section $P = K \cap \{\mathbf{1}^\top u = 1\}$ is a polytope of dimension $h - 1$ whose relative interior consists exactly of the strictly positive points.

Consider $f(u) = \det(\sum_{i \in S} u_i x_i x_i^\top)$, a polynomial of degree at most d . On relint P all weights are positive and the features span $\mathbb{R}^d$, so $f > 0$. At any boundary point some coordinate vanishes; if $f > 0$ there, the positively weighted features would span $\mathbb{R}^d$ and the point would be a spanning zero certificate of support $\leq k - 1$, contradicting $\tau = k$. Hence $f \equiv 0$ on every facet of P . Parametrizing aff P by $\mathbb{R}^{h-1}$, the restriction g of f is a nonzero polynomial of degree $\leq d$ vanishing on the relative interior of each facet, hence divisible by the (pairwise non-associated) affine linear forms of the distinct facet hyperplanes; therefore P has at most d facets. Since a bounded $(h-1)$ -dimensional polytope has at least h facets, we get $h \leq d$, so $h = d$, P has exactly d facets, and P is a $(d-1)$ -simplex; K is a simplicial cone with extreme rays $v^{(1)}, \dots, v^{(d)}$.

Let $U_j = \text{span}\{x_i : v_i^{(j)} > 0\}$. As c is a strictly positive combination of all rays, the supports of the rays cover S , so $\sum_j U_j = \mathbb{R}^d$. Dropping ray j gives a boundary point, where the features cannot span, so $\sum_{\ell \neq j} U_\ell \neq \mathbb{R}^d$ for each j : the family $\{U_j\}$ is a minimal spanning family. Choose functionals ϕ_j vanishing on $\sum_{\ell \neq j} U_\ell$ and not on U_j , normalized against vectors $u_j \in U_j$ with $\phi_j(u_j) = 1$; then (ϕ_j) is a dual basis, and $U_j \subseteq \bigcap_{i \neq j} \ker \phi_i$, a one-dimensional space, so $\dim U_j = 1$.

Each extreme ray of the cone $\{u \geq 0 : Au = 0\}$ has inclusion-minimal support C_j , and $\dim \ker A_{C_j} = 1$ (otherwise a two-sided perturbation along a second kernel direction splits the ray). All dyads of C_j lie in $\mathbb{R}^m \otimes U_j$, a space of dimension m , so $|C_j| = \text{rank } A_{C_j} + 1 \leq m + 1$. Finally $k = |S| \leq \sum_j |C_j| \leq d(m + 1) = k$ forces every $|C_j| = m + 1$ and pairwise disjoint supports. $\square$

5.2 Hard datasets live on d lines

Theorem 5.2 (Line structure). *Let the features span $\mathbb{R}^d$ and $\tau(D) = (m + 1)d = k$. Then:*

1. *every data point with $x_i \neq 0$ belongs to some positive circuit of dyads supported on a single feature line and of size exactly $m + 1$;*
2. *there is no positive circuit of dyads whose features span a subspace of dimension ≥ 2 , and no line circuit of size $\leq m$; in particular no point has $x_i \neq 0$ and $\rho_i = 0$;*
3. *the nonzero features lie on exactly d linearly independent lines.*

Consequently, any dataset whose nonzero features are not of this d -line form satisfies $\tau(D) \leq (m + 1)d - 1$ and is recovered exactly within budget $(m + 1)d - 1$.

Proof. (1) Extend x_t (t fixed, $x_t \neq 0$) to a feature basis $B \ni t$ and apply Lemma 3.1: the resulting certificate has support at most k and, since $\tau = k$, exactly k ; Theorem 5.1 splits it into $(m + 1)$ -point line circuits, one of which contains t .

(2) Let C be a positive circuit whose features span an s -dimensional space with $s \geq 2$; then $|C| \leq ms + 1$ (its dyads lie in an ms -dimensional space). Extend the features of C by $d - s$ data points to a basis of $\mathbb{R}^d$, and attach to each added point its $(m + 1)$ -point line circuit from (1). Summing all these positive relations gives a spanning zero certificate of support at most $ms + 1 + (d - s)(m + 1) = k - s + 1 \leq k - 1$, contradicting $\tau = k$. A line circuit of size $\leq m$ similarly completes with $d - 1$ line circuits to support $\leq m + (d - 1)(m + 1) = k - 1$. A point with $x_i \neq 0 = \rho_i$ is by itself a 1-point line “circuit” ($M_i = 0$) and is excluded the same way.

(3) Suppose $d + 1$ distinct nonzero feature lines exist; take an $(m + 1)$ -point line circuit on each (possible by (1)) and let J be their disjoint union, $|J| = (d + 1)(m + 1)$. Then $\dim \ker A_J \geq |J| - md = d + m + 1$. The sum of the circuits’ positive kernel vectors is strictly positive on J , so the nonnegative kernel cone spans $\ker A_J$. But each extreme ray of that cone is a positive circuit, hence by (2) supported on a single line and of size $m + 1$; on each line the kernel of the corresponding block is one-dimensional, so there are at most $d + 1$ extreme rays, spanning at most $d + 1$ dimensions — contradicting $d + m + 1 > d + 1$. As the features span $\mathbb{R}^d$, there are exactly d independent lines. $\square$

5.3 The m -point mean lemma

Lemma 5.3 (m -point mean lemma). *Let $z_i \in \mathbb{R}^m$ carry weights $p_i > 0$ with $\sum_i p_i z_i = 0$, and let $E = \sum_i p_i \|z_i\|^2$. Then some convex combination δ of at most m of the points satisfies $\|\delta\|^2 \leq E/m^2$. The constant is attained by the uniformly weighted regular simplex.*

Proof. Decompose p inside the cone $\{q \geq 0 : \sum_i q_i z_i = 0\}$ into positive circuits and normalize each to a probability vector; each circuit has support at most $m + 1$ and the second moments average to E , so some circuit q has $E_q \leq E$. If $E_q = 0$, all its points vanish and a single point gives $\delta = 0$. If its support has size $\leq m$, its own zero mean is the desired $\delta = 0$.

Otherwise the support is $q_1, \dots, q_{m+1} > 0$ with $\sum_j q_j z_j = 0$; deleting point j leaves the convex combination $\mu_j = -\frac{q_j}{1-q_j} z_j$ of the other m points. With $t_j = q_j \|z_j\|^2 / E_q$ (so $\sum_j t_j = 1$), $\|\mu_j\|^2 / E_q = q_j t_j / (1-q_j)^2$. Since $\sum_j \frac{(1-q_j)^2}{q_j} = \sum_j \frac{1}{q_j} - 2(m+1) + 1 \geq (m+1)^2 - 2(m+1) + 1 = m^2$ (Cauchy–Schwarz), not all j can violate $\|\mu_j\|^2 \leq E_q / m^2$, for otherwise summing $t_j > \frac{(1-q_j)^2}{m^2 q_j}$ over j yields $1 > 1$. Finally $E_q \leq E$. Tightness: for the regular simplex with uniform weights all $\|\mu_j\|^2 = E / m^2$. $\square$

5.4 Assembly

Theorem 5.4 (Near-threshold value). *For all $d, m \geq 2$, $F_{\text{weighted}}(d, m, (m+1)d-1) = 1 + \frac{1}{dm^2}$.*

Proof. Upper bound. Let $n_0 = (m+1)d-1$ and fix a dataset. If the feature rank is $r < d$, Theorem 3.2 gives a certificate of support $(m+1)r \leq n_0$ and the ratio is 1; the same holds if $L_D^* = 0$ (basis selection, as in Section 4) or if $\tau(D) \leq n_0$. In the remaining hard case $\tau(D) = (m+1)d$, Theorem 5.2 provides a basis $u_1, \dots, u_d$ and scalars $s_i \neq 0$ with $x_i = s_i u_{j(i)}$ for every nonzero-feature point. Reading the first-order condition $\sum_i M_i = 0$ against the basis (u_j) , each line satisfies $\sum_{i \in I_j} s_i \rho_i = 0$. Set, per line,

$$z_i = \rho_i / s_i, \quad p_i = s_i^2 / g_j, \quad g_j = \sum_{i \in I_j} s_i^2, \quad R_j = \sum_{i \in I_j} \|\rho_i\|^2,$$

so that $\sum_{i \in I_j} p_i z_i = 0$ and $\sum_i p_i \|z_i\|^2 = R_j / g_j$. Choose the lightest line j_* with $R_{j_*} \leq \frac{1}{d} \sum_j R_j \leq \frac{NL_D^*}{d}$ (features equal to 0 only increase L_D^*). On each line $j \neq j_*$ pick, by conic Carathéodory, at most $m+1$ points with a positive zero-mean combination of the z_i ; on line j_* pick at most m points and a convex combination δ with $\|\delta\|^2 \leq \frac{R_{j_*}}{m^2 g_{j_*}}$ (Lemma 5.3). Translate the per-line convex coefficients α_i into selection weights $c_i = \alpha_i / s_i^2 > 0$ (rescaling each line independently and normalizing globally); the weighted normal equations then decouple along the basis and give $\widehat{W} = W^\circ + H$ with $Hu_j = 0$ for $j \neq j_*$ and $Hu_{j_*} = \delta$; the selected features contain all d lines, so the weighted Gram matrix is positive definite and the minimizer is unique. The support totals $(d-1)(m+1) + m = n_0$. By (1),

$$L_D(\widehat{W}) - L_D^* = \frac{g_{j_*} \|\delta\|^2}{N} \leq \frac{R_{j_*}}{Nm^2} \leq \frac{L_D^*}{dm^2}.$$

Lower bound. Take the regular-simplex axial instance: let $r_1, \dots, r_{m+1} \in \mathbb{R}^m$ be regular simplex vertices with $\sum_j r_j = 0$ and $\|r_j\|^2 = \frac{m}{m+1}$, whose facets are at squared distance $h^2 = \frac{1}{m(m+1)}$ from the origin; put $w = he_1$ and $D = \{(e_i, w + r_j)\}_{i \in [d], j \in [m+1]}$. The unique full minimizer is $W^\circ = [w \ \dots \ w]$ with $L_D^* = \frac{m}{m+1}$. For any selection of support at most $N-1$, some axis is empty (column zeroed by the Frobenius rule; deviation $\|w\|^2 = h^2$) or uses a proper subset (deviation at least the facet distance h^2). By the analogue of (4), the loss exceeds L_D^* by at least h^2/d , and deleting one vertex with uniform facet weights attains it. The ratio is $1 + \frac{h^2/d}{m/(m+1)} = 1 + \frac{1}{dm^2}$. $\square$

Remark 5.5 (General centered-simplex families). The same computation applies to any centered simplex $r_1, \dots, r_{m+1}$ ($\sum_j r_j = 0$, $S = \sum_j \|r_j\|^2$, $h = \min_j \text{dist}(0, \text{conv}\{r_k\}_{k \neq j})$), provided the columns of W° are chosen with $\|W^\circ e_i\| \geq h$: the exact $(N-1)$ -point ratio is $1 + \frac{(m+1)h^2}{dS}$,

and within this family $h^2 \leq S/(m^2(m+1))$ with equality exactly for regular simplices. The integer instance of Definition 3.3 realizes the slightly smaller rational value of Theorem 3.6 with integer data.

6 The general k -point mean lemma

Lemma 5.3 is the case $k = M$ of a sharp family of sparsification bounds, which we record here both for its own sake and as a tool for Section 7. Unlike everything else in this paper, the general case relies on a recently proved deep geometric inequality — the full-dimensional Grace–Danielsson inequality conjectured by Egan and proved by Drozdov [Dro25]. *None of Theorems 3.7, 4.1, 5.4 or 7.7 depends on this section*; the only case used elsewhere is $k = M$ (Lemma 5.3) and the trivial $k = 1$, both self-contained.

Theorem 6.1 (k -point mean lemma). *Let $z_i \in \mathbb{R}^M$ carry weights $p_i > 0$, $\sum_i p_i = 1$, with $\sum_i p_i z_i = 0$, and $E = \sum_i p_i \|z_i\|^2$. For every $1 \leq k \leq M+1$ there is a convex combination δ of at most k of the points with*

$$\|\delta\|^2 \leq E \cdot \frac{M+1-k}{kM}.$$

The constant is sharp: for the uniformly weighted regular M -simplex, every convex combination supported on k points has squared norm at least $E \frac{M+1-k}{kM}$, with equality for k vertices at equal weights.

The proof route: (i) reduce to a single positive circuit of support $s \leq M+1$ (extreme points of the zero-mean polytope, as in Lemma 5.3); (ii) apply the following sparsification lemma with $a_i = z_i$.

Lemma 6.2 (Simplex skeleton sparsification). *Let $a_1, \dots, a_s$ lie in a Hilbert space, let q be a probability vector with full support s , and let $V_q = \sum_i q_i \|a_i - \bar{a}_q\|^2$ where $\bar{a}_q = \sum_i q_i a_i$. For every $1 \leq k \leq s$ there is a probability vector λ with $|\text{supp } \lambda| \leq k$ and*

$$\left\| \sum_i \lambda_i a_i - \bar{a}_q \right\|^2 \leq \frac{s-k}{k(s-1)} V_q.$$

Proof sketch with full statements of the two nontrivial steps. We construct a martingale of probability vectors that successively kills coordinates. The engine is:

Boundary variance fact. *Let q have $r \geq 2$ positive coordinates and let $H \succeq 0$ be a quadratic form. Then there is a random probability vector Q_H supported on the boundary $\partial\Delta_r = \{u \in \Delta_r : \min_i u_i = 0\}$ with $\mathbb{E}[Q_H] = q$ and*

$$\mathbb{E}[(Q_H - q)^\top H (Q_H - q)] \leq \frac{\text{tr}(H(\text{Diag } q - qq^\top))}{(r-1)^2}.$$

(One boundary variable per form H suffices for our purposes; a single Q working for all H simultaneously also exists, by separating the compact convex set of achievable covariances from $\{K : K \preceq (\text{Diag } q - qq^\top)/(r-1)^2\}$, but we do not need it.)

Proof of the fact. The left-hand side, minimized over admissible laws, is a finite moment problem on the compact set $\partial\Delta_r$; strong duality (a supporting hyperplane to the compact convex set of achievable moment pairs) gives

$$\Phi_H(q) = \sup \left\{ \ell(q) : \ell \text{ affine, } \ell(u) \leq (u - q)^\top H (u - q) \ \forall u \in \partial\Delta_r \right\}.$$

Fix a feasible ℓ and (after adding εP_T to H on the tangent space $T = \{\mathbf{1}^\top y = 0\}$ and letting $\varepsilon \downarrow 0$) assume $H \succ 0$ on T . Writing $\ell(q+y) = a + 2\langle c, y \rangle_H$ and $R^2 = a + \|c\|_H^2$, feasibility says the H -ball $B_H(c, R)$ misses $\partial\Delta_r - q$; since $0 \in B_H(c, R)$ when $a > 0$, convexity forces $B_H(c, R) \subseteq \Delta_r - q$. The desired bound $(r-1)^2 \ell(q) \leq \text{tr}(H(\text{Diag } q - qq^\top)) = \sum_i q_i (e_i - q)^\top H (e_i - q)$ then follows from:

Sublemma G (simplex–ball interpolation). *Let an n -simplex $T \subset \mathbb{R}^n$ contain the ball $B(c, R)$ with $0 \in B(c, R)$, and let $0 = \sum_i q_i v_i$ be the barycentric representation of the origin in the vertices v_i of T . Then*

$$\sum_i q_i \|v_i\|^2 \geq n^2 (R^2 - \|c\|^2).$$

Proof of Sublemma G. Normalize the ball to the unit ball centered at the origin; the distinguished point becomes y with $u = \|y\| < 1$, and the left side becomes the value at y of the affine interpolation of $\|\cdot\|^2$ on the vertices, which for any simplex with circumcenter O and circumradius $\mathcal{R}$ equals $P_T(y) = \mathcal{R}^2 - \|O - y\|^2$ plus $\|y\|^2$. Slide each facet inward until tangent to the unit ball, obtaining the inscribed simplex $S \subseteq T$ with insphere the unit ball; by the variance decomposition of convex combinations (write each vertex of S in the vertices of T and y in the vertices of S), the interpolation value only decreases: $P_T(y) \geq P_S(y)$. For S , let $d = \|O\|$ and write $\mathcal{R} = n + s$ with $s \geq 0$. Drozdov’s theorem [Dro25] (the Grace–Danielsson inequality in every dimension: for any n -simplex with inradius r , circumradius $\mathcal{R}$, and center distance d , one has $(\mathcal{R} - nr)(\mathcal{R} + (n-2)r) \geq d^2$; here $r = 1$) gives $d^2 \leq s(s + 2n - 2)$, whence $\mathcal{R}^2 - d^2 - n^2 \geq 2s$ and $s \geq \sqrt{d^2 + (n-1)^2} - (n-1)$. Then

$$P_S(y) - n^2(1 - u^2) \geq 2s - 2du + (n^2 - 1)u^2,$$

and minimizing the right-hand side over $d \geq 0$ (optimum at $d = (n-1)u/\sqrt{1-u^2}$) leaves $(n-1)[(n+1)u^2 - 2(1 - \sqrt{1-u^2})] \geq 0$, which holds since $1 - \sqrt{1-u^2} \leq u^2$. This proves Sublemma G, hence the boundary variance fact. $\square$

Now iterate: starting from q with support s , apply the fact with $H = A^\top A$ (where $A = [a_1 \cdots a_s]$), obtaining a boundary law Q with $\mathbb{E}[\bar{a}_Q] = \bar{a}_q$ and $\mathbb{E}\|\bar{a}_Q - \bar{a}_q\|^2 \leq V_q/(r-1)^2$; the total-variance identity $\mathbb{E}[V_Q] = V_q - \mathbb{E}\|\bar{a}_Q - \bar{a}_q\|^2$ gives $\mathbb{E}[V_Q] \geq (1 - \frac{1}{(r-1)^2})V_q$. A step may kill several coordinates at once; setting $f_r = 1 - \frac{1}{(r-1)^2}$ and $\Gamma_j = \prod_{t=k+1}^j f_t$ (with $\Gamma_j = 1$ for $j \leq k$), which is nonincreasing in j , a backward induction on the support size shows that the terminal variance obeys $\mathbb{E}[V_\Lambda] \geq \Gamma_s V_q$ regardless of how many coordinates each step kills. Since the mean sequence is a martingale with orthogonal increments, $\mathbb{E}\|\bar{a}_\Lambda - \bar{a}_q\|^2 = V_q - \mathbb{E}[V_\Lambda] \leq (1 - \Gamma_s)V_q$, and the telescoping product $\Gamma_s = \prod_{r=k+1}^s \frac{r(r-2)}{(r-1)^2} = \frac{s(k-1)}{k(s-1)}$ yields the bound $\frac{s-k}{k(s-1)}V_q$; some realization λ of the terminal law achieves it. $\square$

Proof of Theorem 6.1. Reduce to a circuit q of support $s \leq M+1$ and second moment $E_q \leq E$ exactly as in Lemma 5.3 (including the degenerate branches $E_q = 0$ and $s \leq k$). Apply Lemma 6.2 with $a_i = z_i$, $\bar{a}_q = 0$: some k -sparse convex δ has $\|\delta\|^2 \leq \frac{s-k}{k(s-1)}E_q$; since $s \mapsto \frac{s-k}{s-1}$ is nondecreasing for $k \geq 1$, the constant is at most $\frac{M+1-k}{kM}$. Sharpness: for regular simplex vertices, $\|\sum_i \alpha_i v_i\|^2 = R^2(\frac{M+1}{M} \sum_i \alpha_i^2 - \frac{1}{M}) \geq R^2 \frac{M+1-k}{kM}$ whenever $|\text{supp } \alpha| \leq k$, using $\sum \alpha_i^2 \geq 1/k$. $\square$

7 The smallest intermediate cell: $(d, m) = (2, 2)$

For $d < n < (m+1)d - 1$ the profile $F_{\text{weighted}}(d, m, n)$ remains open — as does its scalar counterpart, the intermediate regime of Question 2 of [HMSY25b]. In this section we treat the

smallest cell $(d, m) = (2, 2)$, where the open budgets are $n = 3, 4$. We prove the interval theorem (Theorem 7.7), reduce the conjectured exact values to a finite moment problem, and assemble structural evidence.

7.1 Lower bounds: an explicit axial instance

Proposition 7.1. *There is an explicit dataset D^Δ with*

$$\frac{L_{D^\Delta}^*(3; \text{weighted})}{L_{D^\Delta}^*} = \frac{13}{8}, \quad \frac{L_{D^\Delta}^*(4; \text{weighted})}{L_{D^\Delta}^*} = \frac{5}{4}.$$

Hence $F_{\text{weighted}}(2, 2, 3) \geq \frac{13}{8}$ and $F_{\text{weighted}}(2, 2, 4) \geq \frac{5}{4}$.

Proof. Let $r_1, r_2, r_3 \in \mathbb{R}^2$ be the vertices of a centered equilateral triangle with $\|r_j\|^2 = \frac{2}{3}$ (so the total is $S = 2$, edge midpoint distance squared $\frac{1}{6}$), and let $w \in \mathbb{R}^2$ with $\|w\|^2 = \frac{5}{6}$. Take $D^\Delta = \{(e_i, w + r_j)\}_{i \in [2], j \in [3]}$. As in Lemma 3.4, $W^\circ = [w \ w]$, $L_D^* = \frac{2}{3}$, and a selection allocating k_i points to axis i incurs excess $\frac{1}{2}(c(k_1) + c(k_2))$ where $c(3) = 0$, $c(2) = \frac{1}{6}$ (best edge), $c(1) = \frac{2}{3}$ (best vertex), $c(0) = \|w\|^2 = \frac{5}{6}$ (empty axis). For budget 3 the minimum over allocations is $c(2) + c(1) = c(3) + c(0) = \frac{5}{6}$, giving ratio $1 + \frac{5/12}{2/3} = \frac{13}{8}$; for budget 4 it is $2c(2) = \frac{1}{3}$, giving $1 + \frac{1/6}{2/3} = \frac{5}{4}$. (We verified both values by exhaustive exact enumeration of all supports.) $\square$

7.2 A complex dictionary for the hard branches

Fix a $(2, 2)$ dataset with feature rank 2 and $L_D^* > 0$; after a linear change of features (which leaves the loss values invariant) assume the data whitened, $\sum_i x_i x_i^\top = \gamma I_2$. Identify $\mathbb{R}^2 \cong \mathbb{C}$; write ξ_i for the feature and η_i for the residual ρ_i of point i (viewed in $\mathbb{C}$), and for $\xi_i \neq 0$ set

$$p_i = \frac{|\xi_i|^2}{\sum_j |\xi_j|^2}, \quad t_i = \frac{\bar{\xi}_i}{\xi_i} \in S^1, \quad z_i = \frac{\eta_i}{\xi_i} \in \mathbb{C}, \quad E = \sum_i p_i |z_i|^2.$$

Whitening is equivalent to $\sum_i \xi_i^2 = 0$, i.e. $\mathbb{E}_p t = 0$, and the first-order condition $\sum_i \rho_i x_i^\top = 0$ is equivalent to the pair $\mathbb{E}_p z = 0$, $\mathbb{E}_p(\bar{t}z) = 0$. A selection with weights c corresponds to $q_i \propto c_i |\xi_i|^2$; writing $\mu = \mathbb{E}_q t$, $\nu = \mathbb{E}_q z$, $\lambda = \mathbb{E}_q(\bar{t}z)$, the selected model deviates from W° by the real-linear map $\xi \mapsto a\xi + b\bar{\xi}$ with

$$\begin{pmatrix} 1 & \mu \\ \bar{\mu} & 1 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} \nu \\ \lambda \end{pmatrix}, \quad |\mu| < 1 \iff \text{selected features span } \mathbb{R}^2, \quad (6)$$

and the excess loss satisfies $(L_D(\widehat{W}) - L_D^*)/L_D^* = C(q)/E$ with $C(q) = |a|^2 + |b|^2$. (Zero-feature points only enlarge L_D^* , so all upper bounds proved through the dictionary are conservative.) Conversely, every finite system (p_i, t_i, z_i) with the three moment conditions is realized by a whitened dataset ($\xi_i = \sqrt{p_i} s_i$ with $\bar{s}_i/s_i = t_i$, $\eta_i = z_i \xi_i$).

For $k \in \{3, 4\}$ define

$$\Gamma_k = \sup_{\text{moment systems}} \frac{1}{E} \inf_{\substack{|\text{supp } q| \leq k \\ |\mu(q)| < 1}} C(q).$$

7.3 Interpolation identities

Lemma 7.2 (Interpolation convex combination). *For $t_i \neq t_j$ let $B_{ij} = \left(\frac{t_i z_j - t_j z_i}{t_i - t_j}, \frac{z_i - z_j}{t_i - t_j} \right) \in \mathbb{C}^2$ (the coefficients of the affine function of t through the two points) and $d_{ij} = |t_i - t_j|^2$. Then for every admissible q ,*

$$D(q) := 1 - |\mu|^2 = \sum_{i < j} q_i q_j d_{ij}, \quad (a, b) = \sum_{\substack{i < j \\ t_i \neq t_j}} \frac{q_i q_j d_{ij}}{D(q)} B_{ij}.$$

In particular (a, b) is a convex combination of the pairwise interpolation coefficients; the conditioning factor $(1 - |\mu|^2)^{-1}$ is absorbed entirely.

Proof. $1 - |\mu|^2 = \frac{1}{2} \sum_{i, j} q_i q_j |t_i - t_j|^2$ expands the first identity. With $w_i = \bar{t}_i z_i$ one checks $d_{ij} a_{ij} = (t_i - t_j)(w_i - w_j)$ and $d_{ij} b_{ij} = (\bar{t}_i - \bar{t}_j)(z_i - z_j)$; summing against $q_i q_j$ and using $\sum_{i < j} q_i q_j (x_i - x_j)(y_i - y_j) = \sum_i q_i x_i y_i - (\sum_i q_i x_i)(\sum_i q_i y_i)$ yields $\sum_{i < j} q_i q_j d_{ij} a_{ij} = \nu - \mu \lambda$ and $\sum_{i < j} q_i q_j d_{ij} b_{ij} = \lambda - \bar{\mu} \nu$, which is $D(q) \cdot (a, b)$ by (6). Same-direction pairs have $d_{ij} = 0$ and are simply omitted. $\square$

Lemma 7.3 (Three-point closure identity). *For $2 \leq |S| \leq 3$ let $K_S = \text{conv}\{B_{ij} : i, j \in S, t_i \neq t_j\}$ (discard S with no valid pair). Then*

$$\inf_{\substack{\text{supp } q \subseteq S \\ |\mu(q)| < 1}} C(q) = \text{dist}^2(0, K_S), \quad \inf_{\substack{|\text{supp } q| \leq 3 \\ |\mu(q)| < 1}} C(q) = \min_S \text{dist}^2(0, K_S).$$

For three pairwise distinct directions the interior of the edge-weight triangle is attained by positive weights (via $q_1 : q_2 : q_3 = \frac{d_{23}}{\alpha_{23}} : \frac{d_{13}}{\alpha_{13}} : \frac{d_{12}}{\alpha_{12}}$), the vertices by two-point supports, and non-degenerate edge interiors only as limits with $|\mu| \rightarrow 1$; if two of the three directions coincide, the corresponding segment of K_S is attained exactly. Moreover, for three distinct directions the three points B_{ij}, B_{ik}, B_{jk} , when distinct, are never collinear over $\mathbb{R}$.

Proof. By Lemma 7.2, over $\text{supp } q \subseteq S$ the value (a, b) ranges over convex combinations of the valid B_{ij} with weights $r_{ij} \propto q_i q_j d_{ij}$; the parametrization above shows all interior weight profiles are realized, and the boundary cases are checked directly (for a repeated direction, distributing weight inside the repeated pair moves along the segment with $|\mu| < 1$ preserved). Distances to closures equal infima of continuous functions over the realized sets. For non-collinearity: $B_{ik} - B_{ij}$ is a complex multiple of $(-t_i, 1)$ and $B_{jk} - B_{ij}$ of $(-t_j, 1)$; real proportionality of nonzero such vectors forces $t_i = t_j$. $\square$

Lemma 7.4 (Global edge identities). *Let $w_{ij} = p_i p_j d_{ij}$. Then $\sum_{i < j} w_{ij} = 1$, $\sum_{i < j, t_i \neq t_j} w_{ij} B_{ij} = 0$, and*

$$\sum_{\substack{i < j \\ t_i \neq t_j}} w_{ij} \|B_{ij}\|^2 = 2E - 2 \sum_G \left(P_G \sum_{i \in G} p_i |z_i|^2 - \left| \sum_{i \in G} p_i z_i \right|^2 \right) \leq 2E,$$

where G ranges over the classes of equal direction and $P_G = \sum_{i \in G} p_i$. In particular some valid pair has $\|B_{ij}\|^2 \leq 2E$.

Proof. The first identity is Lemma 7.2 at $q = p$ with $\mu = 0$; the second is Lemma 7.2 at $q = p$, where $(a, b) = (0, 0)$ by the moment conditions. For the third, expand $d_{ij} \|B_{ij}\|^2 = |t_i z_j - t_j z_i|^2 + |z_i - z_j|^2$ and sum: over all pairs, each of the two sums telescopes to $E - |\mathbb{E}_p(\bar{t}z)|^2 = E$ and $E - |\mathbb{E}_p z|^2 = E$ respectively; same-direction pairs, for which B_{ij} is undefined, contribute $2p_i p_j |z_i - z_j|^2$, which regrouped over classes gives the correction term. $\square$

7.4 Star localization and the two unconditional bounds

Lemma 7.5 (Star localization). *Fix an atom i and set $P_{ij} = \frac{1}{2}p_j d_{ij}$ for $j \neq i$. Then $\sum_j P_{ij} = 1$, the valid star coefficients B_{ij} lie on a real two-dimensional affine plane through m_i with $\|m_i\|^2 = \frac{1}{2}|z_i|^2$, and $\sum_j P_{ij} B_{ij} = m_i$; their variance is $V_i = E + \frac{|z_i|^2}{2} - L_i$ where $L_i = \sum_{j:t_j=t_i} p_j |z_j - z_i|^2 \geq 0$. Consequently there exist two star neighbours whose segment contains a point δ with*

$$\|\delta\|^2 \leq \frac{E}{4} + \frac{5|z_i|^2}{8} - \frac{L_i}{4},$$

and δ is a closure point of three-point selections (Lemma 7.3). Choosing i with $|z_i|^2 \leq E$:

$$\inf_{\substack{|\text{supp } q| \leq 3 \\ |\mu| < 1}} C(q) \leq \frac{7E}{8}, \quad \text{so} \quad \Gamma_3 \leq \frac{7}{8}. \quad (7)$$

Proof. Work in the real frame picture: choose $s_i^2 = t_i$, set $u_i = \sqrt{2}(\cos \theta_i, \sin \theta_i)^\top$ and $w_i = \bar{s}_i z_i$; the moment conditions become the tight-frame identities $\sum_j p_j u_j u_j^\top = I_2$, $\sum_j p_j u_j w_j = 0$, and $d_{ij} = \det(u_i, u_j)^2$. With $n_i \perp u_i$ a unit vector, $P_{ij} = p_j(u_j^\top n_i)^2$ sums to $n_i^\top I n_i = 1$; every valid pair coefficient can be written $\gamma_{ij} = m_i + n_i \xi_{ij}$ with $m_i = \frac{1}{2}u_i w_i$, and $\sum_j P_{ij} \xi_{ij} = n_i^\top (\sum_j p_j u_j w_j) - n_i^\top (\sum_j p_j u_j u_j^\top) m_i = 0$, giving the mean identity; the variance computation gives V_i . Since the centered coefficients lie in a real plane (indeed a line through the origin of that plane after removing m_i), Lemma 5.3 with $M = 2, k = 2$ applied to the P -weighted zero-mean system produces two neighbours and a point δ' on their segment with $\|\delta' - m_i\|^2 \leq V_i/4$; orthogonality of m_i and the centered directions gives $\|\delta'\|^2 \leq \|m_i\|^2 + V_i/4$, which is the stated bound. The bound (7) follows from $\min_i |z_i|^2 \leq E$ and $L_i \geq 0$. $\square$

Lemma 7.6 (Four-point $E/2$ bound). *For every moment system, $\inf_{|\text{supp } q| \leq 4, |\mu| < 1} C(q) \leq \frac{E}{2}$; hence $\Gamma_4 \leq \frac{1}{2}$.*

Proof. Fix i with $|z_i|^2 \leq E$. By Lemma 7.5 the star coefficients γ_{ij} lie in a real two-dimensional affine plane and average to m_i under the weights P_{ij} ; by Carathéodory in the plane, $m_i = \sum_{r=1}^3 \alpha_r \gamma_{ij_r}$ for some three neighbours and convex α . Take $q_i = 1 - \varepsilon$, $q_{j_r} = \varepsilon c_r$ with $c_r \propto \alpha_r / d_{ij_r}$; by Lemma 7.2 the normalized edge weights of the star edges tend to α_r while all leaf-leaf edges carry total weight $O(\varepsilon)$, so $(a, b) \rightarrow m_i$ as $\varepsilon \downarrow 0$, along selections with support ≤ 4 and $|\mu| < 1$. Hence the infimum is at most $\|m_i\|^2 = \frac{1}{2}|z_i|^2 \leq \frac{E}{2}$. $\square$

7.5 The interval theorem

Theorem 7.7. $F_{\text{weighted}}(2, 2, 3) \in [\frac{13}{8}, \frac{15}{8}]$ and $F_{\text{weighted}}(2, 2, 4) \in [\frac{5}{4}, \frac{3}{2}]$.

Proof. Lower bounds: Proposition 7.1. Upper bounds: fix a dataset and a budget $n \in \{3, 4\}$. If the feature rank is $r \leq 1$, then $\tau(D) \leq (m+1)r \leq 3 \leq n$ (Theorem 3.2) and the ratio is 1; likewise if $L_D^* = 0$ or $\tau(D) \leq n$. Otherwise the dictionary of Section 7.2 applies, and by (7) and Lemma 7.6 (a three-point selection is also a four-point selection) the ratio is at most $1 + \Gamma_3 \leq \frac{15}{8}$ for $n = 3$ and at most $1 + \Gamma_4 \leq \frac{3}{2}$ for $n = 4$. The closure points used in Lemmas 7.5–7.6 are limits of admissible selections within the same budget, which suffices since the inner optimization in F_{weighted} is an infimum. $\square$

7.6 The conjecture and its evidence

Conjecture 7.8. $F_{\text{weighted}}(2, 2, 3) = \frac{13}{8}$ and $F_{\text{weighted}}(2, 2, 4) = \frac{5}{4}$. Equivalently (by the branch analysis above and Lemma 7.3): for every moment system, $\inf_{|\text{supp } q| \leq 4, |\mu| < 1} C \leq \frac{E}{4}$ and $\inf_{|\text{supp } q| \leq 3, |\mu| < 1} C \leq \frac{5E}{8}$.

We record five independent pieces of structural evidence.

7.6.1 Finite reduction: at most seven atoms

Proposition 7.9. For fixed atoms (t_i, z_i) , the admissible weight vectors p form a polytope cut out by at most 7 linear equalities, and the worst weight vector (minimizing E) may be taken at a basic feasible solution, of support at most 7. Consequently each inequality of Conjecture 7.8 holds for all finite systems if and only if it holds for all systems with at most seven atoms; and, since atoms may be split into identically placed copies without changing anything, it suffices to treat systems with exactly seven labelled atoms.

Proof. The inner quantity $\inf_q C(q)$ depends only on the atom set, not on p ; $E(p)$ is linear in p , constrained by normalization (1 equation) and the three complex moments (6 real equations). A basic optimal solution has support at most the constraint rank ≤ 7 ; its subsystem inherits the moment conditions, its witness lifts verbatim to the original system, and if the minimal $E^* = 0$ all its supported z_i vanish while $\mathbb{E}t = 0$ guarantees two distinct directions, giving a two-point selection with $C = 0$. Splitting an atom (p_i, t_i, z_i) into $(\theta p_i, t_i, z_i), ((1 - \theta)p_i, t_i, z_i)$ changes no moments; merging the copies in a witness does not increase its support. $\square$

7.6.2 The two-direction class is solved sharply

Proposition 7.10. Suppose the atoms take exactly two directions. Then the directions are antipodal, each class has total weight $\frac{1}{2}$ and zero conditional mean, and $C(q) = \frac{1}{2}(|m_+|^2 + |m_-|^2)$ where $m_{\pm}$ are the conditional means of the selected weights. Consequently

$$\inf_{|\text{supp } q| \leq 4} C \leq \frac{E}{4}, \quad \inf_{|\text{supp } q| \leq 3} C \leq \frac{5E}{8},$$

and both constants are attained: for six equally weighted atoms with z -values the cube roots of unity over each direction, the three-point optimum is exactly $\frac{5E}{8}$ and the four-point optimum exactly $\frac{E}{4}$. Equality at $\frac{5E}{8}$ forces both classes to have equal energy, conditional weights $\frac{1}{3}$ each, and equilateral z -triangles of equal radius. Furthermore, for five-atom two-direction systems the sharp constant at three points is $\frac{4}{7} < \frac{5}{8}$ (attained by an equilateral triple against an antipodal pair with energy ratio 4:3).

Proof. $\mathbb{E}_p t = 0$ with two unit directions forces antipodality and equal class weights; the two complex moments then force zero conditional class means. The regression fit at $\pm$ is the pair of selected conditional means, giving the formula for C . For a class with conditional weights α_i summing to 1 and zero mean, deleting one point leaves the two-point mean $-\frac{\alpha_i z_i}{1 - \alpha_i}$, and $\sum_i \frac{(1 - \alpha_i)^2}{\alpha_i} \geq 4$ for three atoms (Cauchy–Schwarz), so some deletion has squared norm $\leq e_G/4$; allocating (2, 2) or (2, 1) across the classes and balancing yields the bounds, with the stated equality analysis. The six-atom computation and the five-atom $\frac{4}{7}$ optimization ($\min\{e_3, e_3/4 + e_2\}/(e_3 + e_2)$ maximized at $e_2 = \frac{3}{4}e_3$) are elementary; we verified all exact values by enumeration. $\square$

7.6.3 The equality fiber is a stratified local maximum

Proposition 7.11. *Let $\mathcal{E}$ be the set of 3+3 two-direction equality systems of Proposition 7.10. Then: (i) within the two-direction stratum, the three-point value drops linearly under energy imbalance: with class energies $E(1 \pm \eta)$ it equals $\frac{E}{8}(5 - 3|\eta|)$ — a downward cusp; (ii) for every sequence of moment systems converging to a point of $\mathcal{E}$ in which some direction class genuinely splits,*

$$\limsup \frac{1}{E} \inf_{|\text{supp } q| \leq 3, |\mu| < 1} C \leq \frac{1}{2} < \frac{5}{8}.$$

Hence no nearby ascent direction exists: $\mathcal{E}$ is a local maximum of the three-point value in the stratified sense.

Proof. (i) is the balance computation in Proposition 7.10. For (ii), let atoms j, k of the $+$ class split: $\delta = t_j - t_k \rightarrow 0$, $\delta \neq 0$, with $t_j, t_k \rightarrow \tau$ and z_j, z_k tending to distinct points of the limit equilateral triangle (radius A , $E \rightarrow A^2$). Use the prediction coordinates $U_{\pm}(a, b) = a \pm \tau b$, in which $C = \frac{1}{2}(|U_+|^2 + |U_-|^2)$ at the limit directions. Writing $t_j = \tau + \eta_j$, $t_k = \tau + \eta_k$, the same-class edge $V = B_{jk}$ satisfies the exact identities

$$U_+(V) = \frac{\eta_j z_k - \eta_k z_j}{\delta}, \quad U_-(V) = \frac{(2\tau + \eta_j)z_k - (2\tau + \eta_k)z_j}{\delta},$$

so that $|\delta|U_+(V) \rightarrow 0$ while $R_{\delta} := |\delta|U_-(V)$ has $|R_{\delta}| \rightarrow 2|z_j - z_k| > 0$ (its phase need not converge). Pick an atom ℓ of the $-$ class; the two cross edges $B_{j\ell}, B_{k\ell}$ average, in the U -coordinates, to $(\frac{z_j + z_k}{2}, z_{\ell}) + o(1)$, whose first coordinate has modulus $\frac{A}{2}$. Among the three equilateral vertices z_{ℓ} one always satisfies $\Re(\bar{z}_{\ell} R_{\delta}) \leq -\frac{1}{2}A|R_{\delta}|$ (some vertex makes an angle $\leq \pi/3$ with $-R_{\delta}$). Give the same-class edge the convex weight $s_{\delta} = c_{\delta}|\delta|$ with $c_{\delta} = -\Re(\bar{z}_{\ell} R_{\delta})/|R_{\delta}|^2 = O(1)$, and the two cross edges weights $(1 - s_{\delta})/2$ each; this is a legal point of the triangle closure $K_{\{j,k,\ell\}}$ (Lemma 7.3). In the limit the first coordinate is unchanged ($s_{\delta}U_+(V) \rightarrow 0$) and the second is optimized along the ray: $|z_{\ell} + c_{\delta}R_{\delta}|^2 \leq \frac{3}{4}A^2 + o(1)$. Hence $\limsup \inf C_3 \leq \frac{1}{2}(\frac{A^2}{4} + \frac{3A^2}{4}) = \frac{A^2}{2}$. If neither class splits, the system stays in the two-direction stratum, where the value is at most $\frac{5E}{8}$ with the rigid equality characterization of Proposition 7.10. (An explicit paired splitting family even attains limit value $\frac{E}{8}$ at three points while keeping its four-point value pinned at $\frac{E}{4}$.) $\square$

7.6.4 Two obstructions: what a proof cannot do

Proposition 7.12 ($\mu = 0$ obstruction). *A proof of Conjecture 7.8 cannot restrict to selections with $\mu = 0$. Concretely, for the seventh-roots system $t_j = \zeta^j$, $z_j = \zeta^{3j}$, $p_j = \frac{1}{7}$ ($\zeta = e^{2\pi i/7}$), every q with $\mu = 0$ satisfies $C \geq \frac{7/k-1}{2}$ for support k (a discrete Parseval identity), i.e. $C \geq \frac{3}{8} > \frac{1}{4}$ at $k = 4$ and $C \geq \frac{2}{3} > \frac{5}{8}$ at $k = 3$; yet unconstrained selections achieve $C = \frac{2}{7}$ at three points (the difference set $\{0, 1, 3\}$) and $C < \frac{1}{4}$ at four consecutive points (an exact algebraic computation modulo $x^3 + x^2 - 2x - 1$). Similarly, for the regular hexagon third-harmonic system ($t_j = e^{\pi i j/3}$, $z_j = (-1)^j$, $p_j = \frac{1}{6}$) all $\mu = 0$ selections of support ≤ 3 have $C = 1$, while $\{0, 1, 2\}$ at weights 2:3:2 attains the exact three-point optimum $C = \frac{1}{2}$.*

Proposition 7.13 (Averaging obstruction). *A proof cannot either rely on selections that keep the original weights on a subset. With $w_e = p_i p_j d_{ij}$, $y_e = w_e B_{ij}$, $U = \sum_e w_e^2$, $Q = \sum_e \|y_e\|^2$, $R = 4 \sum_i p_i^2$, $V = 2 \sum_i p_i^2 |z_i|^2$, the exact averages over all k -subsets S (with $D_S = \sum_{e \subset S} w_e$,*

$G_S = \sum_{e \in S} y_e$, and binomials $A = \binom{N-2}{k-2}$, $B = \binom{N-3}{k-3}$, $H = \binom{N-4}{k-4}$ are

$$\sum_{|S|=k} \|G_S\|^2 = (A - 2B + H)Q + (B - H)V, \quad \sum_{|S|=k} D_S^2 = H + (A - 2B + H)U + (B - H)R.$$

For the fifth-roots system $t_j = \zeta_5^j$, $z_j = t_j^2$, $p_j = \frac{1}{5}$ one gets $U = \frac{3}{25}$, $R = \frac{4}{5}$, $Q = \frac{1}{5}$, $V = \frac{2}{5}$, so the best original-weight three-point ratio guaranteed by these averages is $\frac{Q+V}{U+R} = \frac{15}{23} > \frac{5}{8}$: reweighting inside the selected triangle (the freedom quantified by Lemma 7.3) is essential.

Proof of Propositions 7.12 and 7.13. For the heptagon, $\mu = m_1, \lambda = m_2, \nu = m_3$ where $m_r = \sum_j q_j \zeta^{rj}$, and Parseval gives $1 + 2 \sum_{r=1}^3 |m_r|^2 = 7 \sum_j q_j^2 \geq 7/k$; with $m_1 = 0$, $C = |m_2|^2 + |m_3|^2$. The unconstrained witnesses are direct computations (for the four-point one, $x = 2 \cos \frac{2\pi}{7}$ satisfies $x^3 + x^2 - 2x - 1 = 0$ and the claimed sign is that of $33x^2 + 71x - 36 > 0$). The hexagon classification of $\mu = 0$ supports (antipodal pairs and alternating triangles) and the exact optimum $\frac{1}{2}$ (via Lemma 7.3; minimal over the three dihedral types of triangles) are finite checks. The subset averages follow by counting occurrences of an edge, an adjacent edge pair, and a disjoint edge pair in k -subsets, together with the cross-term bookkeeping $\text{Adj} = V - 2Q$, $\text{Disj} = Q - V$ (from $\sum_e y_e = 0$) and their scalar analogues; the fifth-roots values use $\|B_{ij}\|^2 = 5 - d_{ij}$ and the two edge lengths $d_{\pm} = \frac{5 \pm \sqrt{5}}{2}$. All exact values in this subsection were also verified by independent symbolic computation. $\square$

7.6.5 Records, closed classes, and a necessary condition

By deletion arguments (Sherman–Morrison in the frame picture; keeping original weights and deleting one atom i costs $C_{-i} = p_i |z_i|^2 h_i / (1 - h_i)^2$ with $h_i = 2p_i$, and harmonic balancing gives $\min_i C_{-i} \leq 2E/(s-2)^2$ for s atoms), Conjecture 7.8 is already proved for: all systems with at most five atoms (four-point version, value $\leq \frac{2E}{9}$), all systems with at most four atoms (three-point version, $\leq \frac{E}{2}$), all two-direction systems (Proposition 7.10), and antipodally paired systems with equal in-pair z (an application of Lemma 5.3 with $M = 2, k = 2$ to the pair means). By Lemma 7.5, any counterexample to the three-point inequality must satisfy $5|z_i|^2 - 2L_i \geq 3E$ for *every* atom (for distinct directions: $|z_i|^2 \geq \frac{3E}{5}$ for all i). Exact records: the hexagon three-point optimum is $\frac{1}{2}$, its four-point value is at most $\frac{19}{169}$; the five-atom two-direction worst case is exactly $\frac{4}{7}$; extensive exact searches over random six- and seven-atom systems produced no value above the two-direction records.

8 Discussion and open problems

The emerging picture. For weighted selection with the minimum-Frobenius-norm ERM, the profile is now known at both ends for all $d, m \geq 1$:

$$F_{\text{weighted}}(d, m, n) = \begin{cases} \infty, & n < d, \\ d + 1, & n = d, \\ \text{open}, & d < n < (m + 1)d - 1, \\ 1 + \frac{1}{dm^2}, & n = (m + 1)d - 1, \\ 1, & n \geq (m + 1)d. \end{cases}$$

The threshold splits conceptually as $(m + 1)d = d + md$: d points to pin a feature basis, md points to cancel the $m \times d$ residual gradient — and the lower-bound instances realize this as d independent copies of weighted mean estimation in $\mathbb{R}^m$, one per feature direction, each requiring its full simplex of $m + 1$ points. At $m = 1$ this collapses to the two mechanisms coinciding, which is why the scalar threshold $2d$ admits both the Steinitz-type proof of [HMSY25a] and the conic-compression proof given here.

Open problems.

1. *The intermediate curve.* Determine $F_{\text{weighted}}(d, m, n)$ for $d < n < (m + 1)d - 1$. Already the scalar case is open ([HMSY25b], Question 2). The axial (line-decomposable) instances reduce to an allocation problem over per-line weighted mean-estimation deficits, for which Theorem 6.1 supplies the sharp per-line constants; but the scalar experience suggests that non-axial, higher-dimensional block constructions dominate deeper inside the intermediate range, so we expect the truth to be an allocation over a richer family of circuit structures.
2. *The (2, 2) cell* (Conjecture 7.8). By Proposition 7.9 this is a question about at most seven atoms on the circle. The obstructions of Section 7.6 show a proof must use selections with $\mu \neq 0$ and must reweight within the selected support; the evidence suggests the two-direction systems are extremal.
3. *Unweighted vector selection thresholds.* Question 3 asks for the unweighted profile $F(d, m, n)$; a recent unrefereed preprint [P⁺26] addresses it, and a rigorous account of the unweighted threshold and its relation to $(m + 1)d$ would complete the vector-valued picture.
4. *Other learning rules.* As emphasized in [HMSY25b], the questions make sense for any ERM selection rule; our certificate and rigidity machinery uses the Frobenius tie-breaking only through Lemma 2.4, and it would be interesting to determine how the threshold depends on the tie-breaking rule.

Verification. All closed-form values in this paper (the instance values $43/40$, $23/20$, $13/8$, $5/4$, the profile values, the exact records $\frac{1}{2}$, $\frac{19}{169}$, $\frac{4}{7}$, $\frac{2}{7}$, $\frac{15}{23}$, and the tightness of the constants in Lemma 5.3 and Theorem 6.1) were additionally verified by exact rational or algebraic symbolic computation, and the lower-bound claims were stress-tested by independent numerical optimization over all supports of the relevant instances.

References

- [Ano26a] Anonymous. Dimension barrier for mean data selection. Unrefereed preprint, GitHub repository `andotheror/dimension-barrier-mean-data-selection`, 2026.
- [Ano26b] Anonymous. Sharp pair selection for mean regression. Unrefereed preprint, Zenodo, 2026. August 22, 2026. DOI: 10.5281/zenodo.22061100.
- [Ber09] Dimitri P. Bertsekas. *Convex Optimization Theory*. Athena Scientific, 2009.
- [Dew26] P. Dewasurendra. One short of Steinitz: Weighted regression selection. Unrefereed preprint, Zenodo, 2026. DOI: 10.5281/zenodo.21881535. Scalar case ($m = 1$) only.

- [Dro25] Sergei Drozdov. Egan conjecture holds. *Discrete Applied Mathematics*, 377:562–572, 2025. arXiv:2310.10816.
- [DW17] Michał Dereziński and Manfred K. Warmuth. Unbiased estimates for linear regression via volume sampling. In *Advances in Neural Information Processing Systems 30 (NeurIPS)*, 2017.
- [HMSY25a] Steve Hanneke, Shay Moran, Alexander Shlimovich, and Amir Yehudayoff. Data selection for ERM. In *Proceedings of the 38th Annual Conference on Learning Theory (COLT)*, volume 291 of *Proceedings of Machine Learning Research*, pages 2634–2665, 2025. arXiv:2504.14572.
- [HMSY25b] Steve Hanneke, Shay Moran, Alexander Shlimovich, and Amir Yehudayoff. Open problem: Data selection for regression tasks. In *Proceedings of the 38th Annual Conference on Learning Theory (COLT)*, volume 291 of *Proceedings of Machine Learning Research*, pages 6225–6229, 2025.
- [P⁺26] Peng et al. Data selection (unrefereed preprint on question 3 of hanneke et al., COLT 2025). Public repository `pipeline-math`, file `papers/data-selection.pdf`, 2026. June 26, 2026. No formal bibliographic data available; cited by repository path.
- [Zha26] Guangjian Zhang. Exact risk ratios for weighted data selection in linear regression. arXiv:2608.28007, 2026.