

Toward Optimal Second-Order Path-Length Guarantee for Adversarial Multi-Armed Bandits

Mengxiao Zhang
University of Iowa
mengxiao-zhang@uiowa.edu

Abstract

We study second-order path-length regret in adversarial K -armed bandits against oblivious loss sequences. Bubeck et al. [4] designed an algorithm that achieves $\tilde{\mathcal{O}}(K + \sqrt{KQ_{\infty,1}})$ regret, where $Q_{\infty,1}$ is the first-order path length, and left open whether $\tilde{\mathcal{O}}(\text{poly}(K)\sqrt{1 + Q_{\infty,2}})$ regret is achievable under bandit feedback, where $Q_{\infty,2}$ is the second-order path length. Somewhat surprisingly, we resolve this question positively by showing that with a more involved analysis, the exact same algorithm of Bubeck et al. [4] achieves $\mathcal{O}\left(K \log(KT) + \sqrt{K \log(KT)(1 + Q_{\infty,2})}\right)$ expected regret when $Q_{\infty,2}$ is known, where T is the horizon. This matches the $\Omega(\sqrt{KQ_{\infty,2}})$ lower bound up to logarithmic factors and additive terms. We further remove the knowledge of $Q_{\infty,2}$ using an adaptive restart scheme whose path-length estimator has uniformly bounded increments.

1 Introduction

The adversarial multi-armed bandit problem is a canonical model of sequential decision making under partial feedback. At the beginning of the game, the environment secretly decides a loss sequence $\ell_1, \ell_2, \dots, \ell_T \in [0, 1]^K$ for T rounds where K is the number of actions.¹ In each of T rounds, a learner chooses one of K actions, or arms. Then, she incurs and observes only the selected coordinate ℓ_{t,I_t} ; the other coordinates remain hidden. Performance is measured against the best single arm in hindsight. This elementary protocol captures the central tension between exploration and exploitation and has consequently served as a basic model for online recommendation, routing, allocation, and repeated decision making. Without further structure on the loss sequence, the minimax regret is $\Theta(\sqrt{KT})$ [1, 26].

The minimax guarantee is derived for an environment that may change arbitrarily at every round, and can therefore be pessimistic on more regular data. A substantial literature develops adaptive guarantees that replace the horizon by an observable or instance-dependent measure of difficulty. Examples include first-order bounds governed by the loss of a good comparator [19, 24], best-of-both-worlds guarantees that are minimax in adversarial environments and logarithmic in stochastic ones [2, 26], and bounds depending on empirical variance or related second-order quantities [3, 7, 9]. These results express the same broad principle: the regret should depend on the complexity actually present in the loss sequence rather than automatically scaling as the worst-case rate $\sqrt{KT}$.

This paper concerns adaptivity to temporal variation, also called the path length of the loss sequence. For loss vectors $\ell_1, \dots, \ell_T \in [0, 1]^K$, define the first- and second-order path lengths by

$$Q_{\infty,1} \triangleq \sum_{t=2}^T \|\ell_t - \ell_{t-1}\|_{\infty}, \quad Q_{\infty,2} \triangleq \sum_{t=2}^T \|\ell_t - \ell_{t-1}\|_{\infty}^2.$$

The first quantity measures movement linearly, whereas the second measures it quadratically. This distinction is substantial when the environment moves through many small increments: if each increment has infinity norm $T^{-1/2}$, then $Q_{\infty,2} = \mathcal{O}(1)$ although the corresponding first-order movement can be $\mathcal{O}(\sqrt{T})$.

¹This environment is also commonly referred to as an oblivious adversary.

For adversarial MAB, Wei and Luo [24] proved the guarantee $\tilde{O}(K + \sqrt{KQ_{1,1}})$, where $Q_{1,1} = \sum_{t=2}^T \|\ell_t - \ell_{t-1}\|_1$.² Bubeck et al. [4] subsequently introduced a recent-arm-biased optimistic mirror-descent algorithm and proved the sharper guarantee of $\tilde{O}(K + \sqrt{KQ_{\infty,1}})$, where $Q_{\infty,1} = \sum_{t=2}^T \|\ell_t - \ell_{t-1}\|_{\infty} \leq Q_{1,1}$.³ Both guarantees nevertheless depend linearly on the magnitude of each temporal increment. This is different from the full-information experts setting, in which the complete vector ℓ_t is observed after every round. In that setting, Steinhardt and Liang [21] showed that second-order path-length dependence is achievable. More recently, Chen et al. [5] resolved the associated “impossible tuning” problem: a single expert algorithm can adapt simultaneously to comparator-specific second-order prediction errors without knowing their scales in advance. Thus full information permits such second-order path-length guarantees, while the corresponding possibility under bandit feedback remained open. This leads to the following question left by Bubeck et al. [4], Wei and Luo [24]:

Is $\tilde{O}(\text{poly}(K)(1 + \sqrt{Q_{\infty,2}}))$ regret achievable for adversarial multi-armed bandits?

Contributions. We answer the question affirmatively for oblivious loss sequences. Somewhat surprisingly, when $Q_{\infty,2}$ is known, we prove that *the exact same algorithm* proposed by Bubeck et al. [4], without any change to its sampler, estimator, or mirror-descent update, achieves the regret of

$$\mathcal{O}\left(K \log(KT) + \sqrt{K \log(KT)(1 + Q_{\infty,2})}\right),$$

under oracle tuning. The contribution in Section 3 is therefore a new analysis, not a new algorithm. Standard OMD reduces the proof to a stability divergence plus the bias caused by recent-arm sampling. Instead of taking an absolute value and bounding it by a first-order movement term, as in Bubeck et al. [4], we combine the bias exactly with the prediction-error square and retain a signed difference term. This refined analysis leads to the desired second-order path-length guarantee. We then remove knowledge of $Q_{\infty,2}$ in Section 4 using a different, adaptively tuned algorithm with a modified sampler and a bounded-increment path-length estimator. The resulting parameter-free regret is $\mathcal{O}(K \log(KT) + \log T + \sqrt{KQ_{\infty,2} \log(KT) \log T})$, which is a factor of $\sqrt{\log T}$ worse than the one with oracle tuning.

Related work. Under full-information feedback, path-length guarantees were developed through gradual-variation and predictable-sequence analyses where optimistic online learning can compare the current loss vector with a predictable hint and obtain regret whose data-dependent term is governed by the prediction error [6, 20]. Using the previous loss vector as the hint turns the squared prediction error into squared temporal variation. Steinhardt and Liang [21] developed an adaptive optimistic exponentiated-gradient algorithm and obtained variance and squared-path regret guarantees for the best expert. Chen et al. [5] later solved the impossible-tuning problem for experts by combining mirror descent, a correction term, and a weighted entropy regularizer. Their algorithm adapts simultaneously to the second-order prediction error of every expert. Both results use the full loss vector. They therefore do not directly tune the hidden squared path length in ordinary MAB, where only one coordinate is observed per round and the squared increments are themselves hidden.

The bandit path-length literature replaces the missing vector observation by importance weighting. Wei and Luo [24] developed the BROAD-OMD framework and obtained bounds involving the first-order movement of the best arm or of all arms. The latter comes with a negative stability term that can cancel the opponents’ movement in repeated games. This gives faster convergence of the average play to equilibrium in two-player zero-sum games with bandit feedback, continuing the full-information connection between no-regret learning and game dynamics [22]. Bubeck et al. [4] improved the MAB dependence from the ℓ_1 path measure to the smaller ℓ_{∞} path measure by using a common optimistic prediction and biasing play toward the most recently selected arm. Our oracle-tuned result uses precisely that algorithm. What changes is the analysis: their proof bounds the bias by a first-order selected-coordinate movement, whereas our signed identity and contraction reduce the same algorithm to $Q_{\infty,2}$.

²We use $\tilde{O}(\cdot)$ to suppress poly-logarithmic terms in K and T .

³Bubeck et al. [4] in fact prove this guarantee against a stronger adaptive adversary. They also propose another algorithm achieving $\tilde{O}(K^{1/3}Q_{1,1}^{1/3}T^{1/6} + K^2)$ for an oblivious adversary.

Beyond temporal variation, adaptive adversarial-bandit guarantees have been developed for small comparator loss [19, 24], loss variance [3, 7, 9], sparse loss vectors [3, 13], and best-of-both-worlds adaptation [2, 12, 16, 26]. These guarantees are generally incomparable because they exploit different structure, including cumulative magnitude, dispersion, support size, or stochastic separation. None of these is characterized by temporal smoothness alone. Related instance-adaptive guarantees also extend beyond ordinary MABs, including bandits with feedback graphs [15, 17], adversarial MDPs [10, 11, 14], linear bandits [4, 8], combinatorial semi-bandits [18, 24], and bandit convex optimization [23, 25].

2 Notations and Problem Setting

Notation. For $K \geq 2$, write $[K] = \{1, \dots, K\}$. Let e_i be the i th standard basis vector and for a vector $v \in \mathbb{R}^K$, denote its i th coordinate as v_i . Let $\Delta_K = \{p \in \mathbb{R}^K : \sum_{i=1}^K p_i = 1, p_i \geq 0 \forall i \in [K]\}$ be the probability simplex. Inner products and Euclidean norms are denoted by $\langle \cdot, \cdot \rangle$ and $\|\cdot\|_2$. For a differentiable convex function Ψ , its Bregman divergence is $D_\Psi(x, y) \triangleq \Psi(x) - \Psi(y) - \langle \nabla \Psi(y), x - y \rangle$. For $z \in \mathbb{R}^K$, the notation $z^{\odot 2}$ means coordinatewise squaring: $(z^{\odot 2})_i = z_i^2$ for all $i \in [K]$. By convention, we use $0/0 = 0$.

Problem setting. Before interaction, an oblivious adversary fixes loss vectors $\ell_1, \dots, \ell_T \in [0, 1]^K$. At round t , the learner selects $I_t \in [K]$ using only its internal randomness and the history $\mathcal{F}_{t-1} = \sigma(I_1, c_1, \dots, I_{t-1}, c_{t-1})$, incurs loss $c_t = \ell_{t, I_t}$, and observes only c_t . The performance criterion is expected static (pseudo) regret

$$\text{Reg}_T \triangleq \max_{i \in [K]} \text{Reg}_T(i) \triangleq \max_{i \in [K]} \mathbb{E} \left[\sum_{t=1}^T \ell_{t, I_t} - \sum_{t=1}^T \ell_{t, i} \right]. \quad (2.1)$$

All expectations in the upper bounds are over the learner's internal randomization (and over the adversary's initial randomization if it draws the oblivious loss table from a distribution). The goal is to obtain an adaptive regret bound depending on the second-order path length $Q_{\infty, 2} \triangleq \sum_{t=2}^T \|\ell_t - \ell_{t-1}\|_\infty^2$. For conciseness, we also write $\mathbb{E}_t[\cdot] = \mathbb{E}[\cdot \mid \mathcal{F}_{t-1}]$ when conditioning on all the randomness before I_t is drawn.

Lower bound. Adapting the binary-loss construction used in Bubeck et al. [4], a lower bound of $\Omega(\sqrt{KQ_{\infty, 2}})$ can be directly derived by the standard lower bound instance in adversarial MAB. We include this lower bound in the following for completeness.

Proposition 2.1. *For every $K \geq 2$, $T \geq 2$, and $2K \leq q \leq T$, every algorithm has to suffer $\Omega(\sqrt{Kq})$ regret under certain oblivious loss sequence $\ell_1, \ell_2, \dots, \ell_T \in [0, 1]^K$ where $Q_{\infty, 2} \leq q$.*

Proof. Let $n = \lfloor q \rfloor$. By the standard adversarial MAB lower bound [1, Theorem 5.1], for any algorithm, one can find an oblivious binary loss sequence of length n on which the algorithm suffers $\Omega(\sqrt{Kn})$ expected regret. Following Bubeck et al. [4, Section 2], use this sequence during the first n rounds and set all losses to zero afterward. Since the losses are binary, every $\|\ell_t - \ell_{t-1}\|_\infty^2$ is at most one. There are at most n transitions before the loss sequence becomes identically zero, and hence $Q_{\infty, 2} \leq n \leq q$. Finally, $q \geq 2K \geq 2$ implies $n = \lfloor q \rfloor \geq q/2$, so the regret is $\Omega(\sqrt{Kn}) = \Omega(\sqrt{Kq})$. $\square$

3 Known $Q_{\infty, 2}$: a Refined Analysis of Bubeck et al. [4]

In this section, we first consider the case in which the learner knows the value of $Q_{\infty, 2}$. In Section 4, we will remove this restriction using a more adaptive algorithm. Somewhat surprisingly, no new algorithm is needed when $Q_{\infty, 2}$ is known, and we show that Algorithm 1 of Bubeck et al. [4] in fact already achieves the desired guarantee when its learning rate is chosen using $Q_{\infty, 2}$. Our contribution is then a more refined analysis of that algorithm.

For completeness, here we briefly introduce Algorithm 1 of Bubeck et al. [4]. The algorithm maintains two distributions p_t and x_t at each round t . Specifically, the algorithm first selects an action I_t from distribution p_t . After observing the loss $c_t \triangleq \ell_{t, I_t}$, the algorithm constructs an unbiased loss estimator $\hat{\ell}_{t, i}$ using both c_{t-1} , the loss suffered in the previous round, and c_t . A direct calculation shows that $\mathbb{E}_t[\hat{\ell}_{t, i}] = \ell_{t, i}$ for every

$i \in [K]$ and more importantly, the variance of this estimator can be much smaller when the path length of the loss vectors is small. Next, the algorithm computes x_{t+1} by applying a step of online mirror descent with log-barrier regularizer using $\widehat{\ell}_t$. Finally, the next-round strategy p_{t+1} is obtained by biasing x_{t+1} toward the arm I_t just selected, with weight determined by $\lambda_{t+1} = \alpha(1 - c_t)$. Algorithm 1 shows the full pseudocode for completeness.

Algorithm 1: Oracle-tuned Algorithm 1 of Bubeck et al. [4]

Input: $K, T, \eta > 0$
1 Set $x_1 = p_1 = \frac{1}{K} \mathbf{1}$, $\alpha = 8\eta$, and $c_0 = 0$
2 Set $\Psi(x) \leftarrow \eta^{-1} \sum_{i=1}^K \log(1/x_i)$
3 **for** $t = 1, \dots, T$ **do**
4 Draw $I_t \sim p_t$ and observe $c_t \leftarrow \ell_{t, I_t}$
5 Set $\widehat{\ell}_{t,i} \leftarrow c_{t-1} + \mathbf{1}\{I_t = i\}(c_t - c_{t-1})/p_{t,i}$ for every $i \in [K]$
6 Set $x_{t+1} \leftarrow \arg \min_{x \in \Delta_K} \left(\langle x, \widehat{\ell}_t \rangle + D_\Psi(x, x_t) \right)$
7 Set $\lambda_{t+1} \leftarrow \alpha(1 - c_t)$ and $p_{t+1} \leftarrow (x_{t+1} + \lambda_{t+1} e_{I_t}) / (1 + \lambda_{t+1})$

Bubeck et al. [4] shows that Algorithm 1 achieves $\widetilde{\mathcal{O}}(K + \sqrt{KQ_{\infty,1}})$ regret where $Q_{\infty,1} = \sum_{t=2}^T \|\ell_t - \ell_{t-1}\|_\infty$ is the first-order path-length that can be much larger than $Q_{\infty,2}$. However, we show in the following theorem that when η and α are chosen properly based on the knowledge of $Q_{\infty,2}$, Algorithm 1 in fact achieves the desired $\widetilde{\mathcal{O}}(K + \sqrt{KQ_{\infty,2}})$ regret bound.

Theorem 3.1. *For every $K \geq 2$, $T \geq 2$, and oblivious loss sequence $\ell_1, \dots, \ell_T \in [0, 1]^K$, Algorithm 1 with $\eta = \min \left\{ \frac{1}{162}, \sqrt{\frac{K \log(KT)}{1 + Q_{\infty,2}}} \right\}$ guarantees that*

$$\text{Reg}_T \leq \mathcal{O} \left(K \log(KT) + \sqrt{K \log(KT)(1 + Q_{\infty,2})} \right). \quad (3.1)$$

To our knowledge, Theorem 3.1 gives the first $\widetilde{\mathcal{O}}(\text{poly}(K)(1 + \sqrt{Q_{\infty,2}}))$ guarantee for adversarial MAB, resolving the open problems proposed in Bubeck et al. [4], Wei and Luo [24]. In the following, we will show the proof of Theorem 3.1, which is our key novelty.

3.1 Analysis of Algorithm 1

We begin with the standard OMD analysis and highlight how our analysis is different from Bubeck et al. [4]. Define $D_t \triangleq D_\Psi(x_t, x_{t+1})$ as the Bregman divergence between consecutive OMD updates x_t and x_{t+1} , and $B_t \triangleq \langle p_t - x_t, \ell_t \rangle$ as the loss difference between x_t and p_t at round t . For each arm $i \in [K]$, set $\gamma = 1/(KT)$ and define the interior comparator $u^{(i)} = (1 - (K-1)\gamma)e_i + \gamma \sum_{j \neq i} e_j$. The OMD first-order condition and the three-point identity (e.g. Wei and Luo [24, Lemma 6]) imply that

$$\langle x_t - u^{(i)}, \widehat{\ell}_t \rangle \leq D_\Psi(u^{(i)}, x_t) - D_\Psi(u^{(i)}, x_{t+1}) + D_t. \quad (3.2)$$

Summing (3.2), taking expectations, using $\mathbb{E}_t[\widehat{\ell}_t] = \ell_t$, and adding the sampling bias gives, for every $i \in [K]$,

$$\begin{aligned} \text{Reg}_T(i) &\leq 1 + \mathbb{E} \left[\sum_{t=1}^T \langle p_t - u^{(i)}, \ell_t \rangle \right] \\ &= 1 + \mathbb{E} \left[\sum_{t=1}^T B_t \right] + \mathbb{E} \left[\sum_{t=1}^T \langle x_t - u^{(i)}, \widehat{\ell}_t \rangle \right] \\ &\leq 1 + D_\Psi(u^{(i)}, x_1) + \mathbb{E} \left[\sum_{t=1}^T (D_t + B_t) \right]. \end{aligned} \quad (3.3)$$

A direct calculation from $x_1 = \frac{1}{K} \mathbf{1}$ gives $D_\Psi(u^{(i)}, x_1) \leq \mathcal{O}(K \log(KT)/\eta)$, uniformly over $i \in [K]$. Therefore, it remains to prove that

$$\mathbb{E} \left[\sum_{t=1}^T (D_t + B_t) \right] \leq \mathcal{O}(1 + \eta Q_{\infty,2}). \quad (3.4)$$

This is the exact point at which our analysis departs from Bubeck et al. [4, proof of Theorem 2]. We briefly discuss how they handle this term. Specifically, their OMD calculation first upper-bounds $\mathbb{E}[\sum_{t=1}^T D_t]$ by $4\eta \mathbb{E}[\sum_{t=1}^T (c_t - c_{t-1})^2]$. Then, their subsequent control of the sampling bias $\mathbb{E}[\sum_{t=1}^T B_t]$ contains the negative of the same term shown above and an additional first-order quantity

$$\alpha \mathbb{E} \left[\sum_{t=2}^T |\ell_{t,I_{t-1}} - \ell_{t-1,I_{t-1}}| \right]. \quad (3.5)$$

The two residual-square terms cancel, leaving (3.5). This is precisely why their argument produces a first-order path length.

However, we do not make that relaxation in bounding $\mathbb{E}[\sum_{t=1}^T (D_t + B_t)]$. Instead, we keep the exact divergence D_t , whose coefficient is strictly smaller than the available negative residual square as we will see later, and rewrite B_t by an equality which replaces the first-order term in (3.5) by a squared movement plus a signed difference term.

The next four lemmas implement this argument in three steps: an exact expansion of B_t , the stability control of the OMD movement x_t , and control of the recent-arm correction. For the remainder of the section, we define $r_t \triangleq c_t - c_{t-1}$, $v_t \triangleq \ell_{t,I_{t-1}} - \ell_{t-1,I_{t-1}} = \ell_{t,I_{t-1}} - c_{t-1}$, and $R \triangleq \mathbb{E}[\sum_{t=2}^T r_t^2]$. Also, for notational convenience, we define $\varphi(z) \triangleq z - z^2/2$ and set $\phi_t \triangleq (\varphi(\ell_{t,1}), \dots, \varphi(\ell_{t,K}))$.

Step 1: expand the sampling bias exactly. The first lemma replaces the relaxation leading to (3.5) by an identity. It is a scalar calculation based only on the explicit difference between p_t and x_t .

Lemma 3.2. *Algorithm 1 satisfies that*

$$\mathbb{E} \left[\sum_{t=2}^T B_t \right] = 4\eta \mathbb{E} \left[\sum_{t=2}^T v_t^2 \right] - 4\eta \mathbb{E} \left[\sum_{t=2}^T r_t^2 \right] + \alpha \mathbb{E} \left[\sum_{t=2}^T \langle \phi_t, p_{t-1} - p_t \rangle \right]. \quad (3.6)$$

Proof. Fix any $t \geq 2$ and condition on $\mathcal{F}_{t-1}$. Since $I_t \sim p_t$, we know that $\mathbb{E}_t[r_t^2] = \sum_{i=1}^K p_{t,i} (\ell_{t,i} - c_{t-1})^2$, while $(\ell_{t,I_{t-1}} - c_{t-1})^2 = v_t^2$. Moreover, the definition of p_t gives $p_t = (x_t + \alpha(1 - c_{t-1})e_{I_{t-1}})/(1 + \alpha(1 - c_{t-1}))$. Consequently, $p_t - x_t = \alpha(1 - c_{t-1})(e_{I_{t-1}} - p_t)$, and hence

$$B_t = \alpha(1 - c_{t-1}) \sum_{i=1}^K p_{t,i} (\ell_{t,I_{t-1}} - \ell_{t,i}). \quad (3.7)$$

We next separate the residual-square difference $4\eta(v_t^2 - \mathbb{E}_t[r_t^2])$ from (3.7). This is the quantity whose negative part will later be combined with the OMD stability term D_t . For each $i \in [K]$, add and subtract the corresponding residual-square difference:

$$\begin{aligned} & \alpha(1 - c_{t-1})(\ell_{t,I_{t-1}} - \ell_{t,i}) \\ &= 4\eta(v_t^2 - (\ell_{t,i} - c_{t-1})^2) + [4\eta((\ell_{t,i} - c_{t-1})^2 - v_t^2) + \alpha(1 - c_{t-1})(\ell_{t,I_{t-1}} - \ell_{t,i})] \\ &= 4\eta(v_t^2 - (\ell_{t,i} - c_{t-1})^2) + 4\eta(\ell_{t,I_{t-1}} - \ell_{t,i})(2 - \ell_{t,I_{t-1}} - \ell_{t,i}) \\ &= 4\eta(v_t^2 - (\ell_{t,i} - c_{t-1})^2) + \alpha(\varphi(\ell_{t,I_{t-1}}) - \varphi(\ell_{t,i})). \end{aligned} \quad (3.8)$$

Multiplying (3.8) by $p_{t,i}$ and summing over $i \in [K]$ now yields

$$B_t = 4\eta v_t^2 - 4\eta \mathbb{E}_t[r_t^2] + \alpha(\varphi(\ell_{t,I_{t-1}}) - \mathbb{E}_t[\varphi(\ell_{t,I_t})]). \quad (3.9)$$

It remains to identify the expectation of the final term. As the loss vectors $\ell_1, \dots, \ell_T$ are oblivious, the tower property gives

$$\mathbb{E}[\varphi(\ell_{t,I_{t-1}}) - \mathbb{E}_t[\varphi(\ell_{t,I_t})]] = \mathbb{E}[\langle \phi_t, p_{t-1} \rangle - \langle \phi_t, p_t \rangle] = \mathbb{E}[\langle \phi_t, p_{t-1} - p_t \rangle]. \quad (3.10)$$

Taking expectations in (3.9), combining (3.10), and summing over $t = 2, \dots, T$ proves (3.6). $\square$

Step 2: control the stability of OMD updates x_t . We next control the two terms involving the OMD updates. The first bound in the following lemma compares the stability term D_t with the negative residual square in Lemma 3.2, and the second controls the movement $x_{t+1} - x_t$ in a weighted Euclidean norm. Both are standard in the analysis of OMD; see, for example, Wei and Luo [24, Definition 11 and Lemmas 12–14]. We include their specialization to our update because the orientation of D_t and the strict constant in (3.11) are useful later. The third bound is not inherited from standard OMD analysis and controls the inner product between ϕ_t and the OMD update $x_t - x_{t+1}$. This turns out to be important to bound the part of the final term in Lemma 3.2 arising from the movement of x_t .

Lemma 3.3 (Log-barrier stability). *Suppose $\eta \leq 1/162$. For every t , Algorithm 1 satisfies*

$$D_t \leq 0.6\eta r_t^2, \quad (3.11)$$

$$|\langle z, x_{t+1} - x_t \rangle| \leq 1.5\sqrt{\eta D_t} \sqrt{\langle x_t, z^{\odot 2} \rangle} \quad \text{for every } z \in \mathbb{R}^K, \quad (3.12)$$

$$\mathbb{E}_t [\langle \phi_t, x_t - x_{t+1} \rangle] \leq 8\mathbb{E}_t [D_t]. \quad (3.13)$$

In addition, $x_{t+1,i}/x_{t,i} \in [0.99, 1.01]$ for every $i \in [K]$, and $D_\Psi(x_{t+1}, x_t) \leq 2D_t$.

Proof. Fix the history before round t . For every fixed $i \in [K]$ and $s \in [0, 1]$, let $x^{(i)}(s)$ be the OMD update obtained when the estimator is $c_{t-1}\mathbf{1} + s(\ell_{t,i} - c_{t-1})e_i/p_{t,i}$. Thus $x^{(i)}(0) = x_t$, $x_{t+1} = x^{(I_t)}(1)$, and $r_t = \sum_{i=1}^K \mathbf{1}\{I_t = i\}(\ell_{t,i} - c_{t-1})$. We first prove that $x_j^{(i)}(1)/x_{t,j} \in [0.99, 1.01]$ for every fixed $i, j \in [K]$. The common vector $c_{t-1}\mathbf{1}$ has zero inner product with every simplex-tangent direction. Thus $x^{(i)}(1)$ can be equivalently written as $\arg \min_{x \in \Delta_K} (\langle x, (\ell_{t,i} - c_{t-1})e_i/p_{t,i} \rangle + D_\Psi(x, x_t))$. The dual local norm of $(\ell_{t,i} - c_{t-1})e_i/p_{t,i}$ at x_t is bounded as

$$\eta x_{t,i}^2 \frac{(\ell_{t,i} - c_{t-1})^2}{p_{t,i}^2} \leq \eta(1 + 8\eta)^2 (\ell_{t,i} - c_{t-1})^2 \leq \eta(1 + 8\eta)^2 < \frac{1}{9}, \quad (3.14)$$

where we used $p_{t,i} \geq x_{t,i}/(1 + 8\eta)$ and $|\ell_{t,i} - c_{t-1}| \leq 1$. We next modify the Dikin calculation in Lemmas 12–14 of Wei and Luo [24] only by retaining the exact curvature of the log-barrier. Define the local displacement associated with the fixed index i by $\Delta_{t,i}^{\text{loc}} = \left(\sum_{j=1}^K (x_j^{(i)}(1) - x_{t,j})^2 / x_{t,j}^2 \right)^{1/2}$. Instead of replacing the Hessian along the update segment by the relaxed factor $1/3$ in Wei and Luo [24, Eq. (15)], direct integration gives

$$\begin{aligned} & \left\langle \nabla(\eta\Psi)(x^{(i)}(1)) - \nabla(\eta\Psi)(x_t), x^{(i)}(1) - x_t \right\rangle \\ &= \int_0^1 \left\langle \nabla^2(\eta\Psi)(x_t + s(x^{(i)}(1) - x_t))(x^{(i)}(1) - x_t), x^{(i)}(1) - x_t \right\rangle ds \\ &= \int_0^1 \sum_{j=1}^K \frac{(x_j^{(i)}(1) - x_{t,j})^2}{(x_{t,j} + s(x_j^{(i)}(1) - x_{t,j}))^2} ds \geq \frac{(\Delta_{t,i}^{\text{loc}})^2}{1 + \Delta_{t,i}^{\text{loc}}}. \end{aligned} \quad (3.15)$$

The first equality is the fundamental theorem of calculus applied to $\nabla(\eta\Psi)$ along the segment from x_t to $x^{(i)}(1)$, and the second uses $\nabla^2(\eta\Psi)(x) = \text{diag}(1/x^2)$. To prove the last inequality, the definition of $\Delta_{t,i}^{\text{loc}}$ gives $|x_j^{(i)}(1) - x_{t,j}|/x_{t,j} \leq \Delta_{t,i}^{\text{loc}}$ for every j . Thus the denominator in the integrand is at most $x_{t,j}^2(1 + s\Delta_{t,i}^{\text{loc}})^2$, and hence the integral is at least $\sum_{j=1}^K \frac{(x_j^{(i)}(1) - x_{t,j})^2}{x_{t,j}^2} \int_0^1 \frac{1}{(1 + s\Delta_{t,i}^{\text{loc}})^2} ds = \frac{(\Delta_{t,i}^{\text{loc}})^2}{1 + \Delta_{t,i}^{\text{loc}}}$.

On the other hand, the KKT equation defining $x^{(i)}(1)$ is $\nabla\Psi(x^{(i)}(1)) - \nabla\Psi(x_t) + ((\ell_{t,i} - c_{t-1})/p_{t,i})e_i + \lambda_{t,i}\mathbf{1} = 0$ for a scalar $\lambda_{t,i}$. Taking its inner product with $x^{(i)}(1) - x_t$ eliminates the multiplier because $\langle \mathbf{1}, x^{(i)}(1) - x_t \rangle = 0$. Therefore,

$$\begin{aligned} & \left\langle \nabla(\eta\Psi)(x^{(i)}(1)) - \nabla(\eta\Psi)(x_t), x^{(i)}(1) - x_t \right\rangle \\ &= -\eta \frac{\ell_{t,i} - c_{t-1}}{p_{t,i}} (x_i^{(i)}(1) - x_{t,i}) \leq \eta \frac{|\ell_{t,i} - c_{t-1}|}{p_{t,i}} |x_i^{(i)}(1) - x_{t,i}| \leq \eta x_{t,i} \frac{|\ell_{t,i} - c_{t-1}|}{p_{t,i}} \Delta_{t,i}^{\text{loc}}. \end{aligned} \quad (3.16)$$

Combining (3.15) and (3.16) gives

$$\max_{j \in [K]} \left| \frac{x_j^{(i)}(1) - x_{t,j}}{x_{t,j}} \right| \leq \Delta_{t,i}^{\text{loc}} \leq \frac{\eta x_{t,i} |\ell_{t,i} - c_{t-1}| / p_{t,i}}{1 - \eta x_{t,i} |\ell_{t,i} - c_{t-1}| / p_{t,i}} \leq \frac{\eta(1+8\eta)}{1 - \eta(1+8\eta)} < 0.01. \quad (3.17)$$

The first inequality follows from the definition of $\Delta_{t,i}^{\text{loc}}$. For the second, if $\Delta_{t,i}^{\text{loc}} = 0$ there is nothing to prove; otherwise, divide (3.15)–(3.16) by $\Delta_{t,i}^{\text{loc}}$ and solve the inequality. The third inequality uses $p_{t,i} \geq x_{t,i}/(1+8\eta)$ and $|\ell_{t,i} - c_{t-1}| \leq 1$. The fourth uses $\eta \leq 1/162$. Therefore, $x_j^{(i)}(1)/x_{t,j} \in [0.99, 1.01]$ for every fixed $i, j \in [K]$. The representation of the realized update above therefore implies $x_{t+1,j}/x_{t,j} \in [0.99, 1.01]$ for every $j \in [K]$.

Next, we prove (3.11). We retain the orientation of the Bregman divergence to obtain the strict constant needed below. The self-concordant conjugate inequality used in the proof of Wei and Luo [24, Lemma 14], applied with local decrement $\eta x_{t,i} |\ell_{t,i} - c_{t-1}| / p_{t,i}$, bounds $\eta D_\Psi(x_t, x^{(i)}(1))$ by $-z - \log(1-z)$ at $z = \eta x_{t,i} |\ell_{t,i} - c_{t-1}| / p_{t,i}$. Since $-z - \log(1-z) \leq z^2/(2(1-z))$ for $0 \leq z < 1$, and $p_{t,i} \geq x_{t,i}/(1+8\eta)$, we obtain that

$$D_\Psi(x_t, x^{(i)}(1)) \leq \frac{1}{2\eta(1-\eta(1+8\eta))} \left(\eta x_{t,i} \frac{|\ell_{t,i} - c_{t-1}|}{p_{t,i}} \right)^2 \leq \frac{(1+8\eta)^2}{2(1-\eta(1+8\eta))} \eta (\ell_{t,i} - c_{t-1})^2. \quad (3.18)$$

For $\eta \leq 1/162$, the final coefficient is no more than 0.6, proving (3.11) after substituting the realized index I_t .

We next derive the two consequences of coordinatewise stability used below. By (3.17), $x_{t+1,i}/x_{t,i} \in [0.99, 1.01]$ for every $i \in [K]$. For every $z \in [0.99, 1.01]$, direct differentiation gives

$$\begin{aligned} z - 1 - \ln z &\leq 2 \left(\frac{1}{z} - 1 + \ln z \right), \\ \frac{1}{z} - 1 + \ln z &\geq \frac{(z-1)^2}{2.1}. \end{aligned}$$

Applying the first inequality coordinatewise gives

$$\begin{aligned} D_\Psi(x_{t+1}, x_t) &= \frac{1}{\eta} \sum_{i=1}^K \left(\frac{x_{t+1,i}}{x_{t,i}} - 1 - \ln \frac{x_{t+1,i}}{x_{t,i}} \right) \\ &\leq \frac{2}{\eta} \sum_{i=1}^K \left(\frac{x_{t,i}}{x_{t+1,i}} - 1 + \ln \frac{x_{t+1,i}}{x_{t,i}} \right) \\ &= 2D_\Psi(x_t, x_{t+1}) = 2D_t, \end{aligned}$$

proving the last argument in the statement. Similarly, the second inequality implies

$$D_t = \frac{1}{\eta} \sum_{i=1}^K \left(\frac{x_{t,i}}{x_{t+1,i}} - 1 + \ln \frac{x_{t+1,i}}{x_{t,i}} \right) \geq \frac{1}{2.1\eta} \sum_{i=1}^K \left(\frac{x_{t+1,i} - x_{t,i}}{x_{t,i}} \right)^2.$$

Rearranging the terms gives the following inequality

$$\sum_{i=1}^K \frac{(x_{t+1,i} - x_{t,i})^2}{x_{t,i}^2} \leq 2.1\eta D_t. \quad (3.19)$$

Therefore, for every $z \in \mathbb{R}^K$, applying Cauchy–Schwarz in the log-barrier local norm gives

$$|\langle z, x_{t+1} - x_t \rangle| \leq \sqrt{\sum_{j=1}^K x_{t,j}^2 z_j^2} \sqrt{\sum_{j=1}^K \frac{(x_{t+1,j} - x_{t,j})^2}{x_{t,j}^2}} \leq 1.5\sqrt{\eta D_t} \sqrt{\langle x_t^{\odot 2}, z^{\odot 2} \rangle} \leq 1.5\sqrt{\eta D_t} \sqrt{\langle x_t, z^{\odot 2} \rangle},$$

where the last inequality uses $x_{t,j}^2 \leq x_{t,j}$. Therefore, we prove (3.12).

It remains to prove (3.13), which does not follow from the generic local-norm argument. We prove it by a direct calculation from the KKT equation. Recall that $x^{(i)}(s)$ is defined for each fixed $i \in [K]$ and $s \in [0, 1]$, and write $x_j^{(i)}(s)$ for its j th coordinate.

The KKT equation of $x^{(i)}(s)$ is $\nabla \Psi(x^{(i)}(s)) + s(\ell_{t,i} - c_{t-1})e_i/p_{t,i} + \lambda^{(i)}(s)\mathbf{1} = \nabla \Psi(x_t)$ for a scalar $\lambda^{(i)}(s)$. Since $\nabla^2 \Psi(x) = \eta^{-1} \text{diag}(1/x_1^2, \dots, 1/x_K^2)$, differentiating the j th coordinate gives $\frac{d}{ds}x_j^{(i)}(s) = -\eta(x_j^{(i)}(s))^2((\ell_{t,i} - c_{t-1})\mathbf{1}(j=i)/p_{t,i} + (\lambda^{(i)})'(s))$. Summing this identity over j and using $\sum_{j=1}^K \frac{d}{ds}x_j^{(i)}(s) = 0$ determines $(\lambda^{(i)})'(s)$. Substitution gives

$$-\frac{d}{ds}x_j^{(i)}(s) = \eta \frac{\ell_{t,i} - c_{t-1}}{p_{t,i}} \left((x_i^{(i)}(s))^2 \mathbf{1}(j=i) - \frac{(x_i^{(i)}(s))^2 (x_j^{(i)}(s))^2}{\sum_{k=1}^K (x_k^{(i)}(s))^2} \right). \quad (3.20)$$

Multiplying (3.20) by $\varphi(\ell_{t,j})$, summing over $j \in [K]$, and integrating over $s \in [0, 1]$ gives

$$\begin{aligned} \langle \phi_t, x_t - x^{(i)}(1) \rangle &= - \int_0^1 \left\langle \phi_t, \frac{d}{ds}x^{(i)}(s) \right\rangle ds \\ &= \eta \frac{\ell_{t,i} - c_{t-1}}{p_{t,i}} \int_0^1 \sum_{j=1}^K \varphi(\ell_{t,j}) \left((x_i^{(i)}(s))^2 \mathbf{1}(j=i) - \frac{(x_i^{(i)}(s))^2 (x_j^{(i)}(s))^2}{\sum_{k=1}^K (x_k^{(i)}(s))^2} \right) ds \\ &= \eta \frac{\ell_{t,i} - c_{t-1}}{p_{t,i}} \int_0^1 \left(\varphi(\ell_{t,i}) (x_i^{(i)}(s))^2 - \frac{(x_i^{(i)}(s))^2}{\sum_{k=1}^K (x_k^{(i)}(s))^2} \sum_{j=1}^K (x_j^{(i)}(s))^2 \varphi(\ell_{t,j}) \right) ds \\ &= \eta \frac{\ell_{t,i} - c_{t-1}}{p_{t,i}} \int_0^1 \frac{(x_i^{(i)}(s))^2}{\sum_{k=1}^K (x_k^{(i)}(s))^2} \sum_{j=1}^K (x_j^{(i)}(s))^2 (\varphi(\ell_{t,i}) - \varphi(\ell_{t,j})) ds \\ &= \eta \frac{\ell_{t,i} - c_{t-1}}{p_{t,i}} \int_0^1 \sum_{j \neq i} \frac{(x_i^{(i)}(s))^2 (x_j^{(i)}(s))^2}{\sum_{k=1}^K (x_k^{(i)}(s))^2} (\varphi(\ell_{t,i}) - \varphi(\ell_{t,j})) ds. \end{aligned} \quad (3.21)$$

We first establish coordinatewise stability along the entire scaled update path. Fix $i \in [K]$ and $s \in [0, 1]$. Since $x^{(i)}(s)$ is the OMD update driven by $s(\ell_{t,i} - c_{t-1})e_i/p_{t,i}$, repeating the calculation leading to (3.17) gives

$$\max_{j \in [K]} \left| \frac{x_j^{(i)}(s) - x_{t,j}}{x_{t,j}} \right| \leq \frac{\eta x_{t,i} s |\ell_{t,i} - c_{t-1}| / p_{t,i}}{1 - \eta x_{t,i} s |\ell_{t,i} - c_{t-1}| / p_{t,i}} \leq \frac{\eta(1+8\eta)}{1 - \eta(1+8\eta)} < 0.01. \quad (3.22)$$

Therefore, $0.99x_{t,j} \leq x_j^{(i)}(s) \leq 1.01x_{t,j}$ for every $j \in [K]$ and $s \in [0, 1]$. Consequently,

$$0.94 \frac{x_{t,i}^2 x_{t,j}^2}{\sum_{k=1}^K x_{t,k}^2} \leq \frac{0.99^4}{1.01^2} \frac{x_{t,i}^2 x_{t,j}^2}{\sum_{k=1}^K x_{t,k}^2} \leq \frac{(x_i^{(i)}(s))^2 (x_j^{(i)}(s))^2}{\sum_{k=1}^K (x_k^{(i)}(s))^2} \leq \frac{1.01^4}{0.99^2} \frac{x_{t,i}^2 x_{t,j}^2}{\sum_{k=1}^K x_{t,k}^2} \leq 1.07 \frac{x_{t,i}^2 x_{t,j}^2}{\sum_{k=1}^K x_{t,k}^2}, \quad (3.23)$$

which implies that

$$\left| \frac{(x_i^{(i)}(s))^2 (x_j^{(i)}(s))^2}{\sum_{k=1}^K (x_k^{(i)}(s))^2} - \frac{x_{t,i}^2 x_{t,j}^2}{\sum_{k=1}^K x_{t,k}^2} \right| \leq 0.07 \frac{x_{t,i}^2 x_{t,j}^2}{\sum_{k=1}^K x_{t,k}^2}. \quad (3.24)$$

Since $x_{t+1} = x^{(I_t)}(1)$ and the conditional distribution of I_t is p_t , the definition of conditional expectation gives $\mathbb{E}_t[\langle \phi_t, x_t - x_{t+1} \rangle] = \sum_{i=1}^K p_{t,i} \langle \phi_t, x_t - x^{(i)}(1) \rangle$. Substituting (3.21) into this finite sum cancels each factor $p_{t,i}$ and gives

$$\begin{aligned} &\mathbb{E}_t[\langle \phi_t, x_t - x_{t+1} \rangle] \\ &= \eta \sum_{1 \leq i < j \leq K} (\varphi(\ell_{t,i}) - \varphi(\ell_{t,j})) \int_0^1 \left((\ell_{t,i} - c_{t-1}) \frac{(x_i^{(i)}(s))^2 (x_j^{(i)}(s))^2}{\sum_{k=1}^K (x_k^{(i)}(s))^2} - (\ell_{t,j} - c_{t-1}) \frac{(x_i^{(j)}(s))^2 (x_j^{(j)}(s))^2}{\sum_{k=1}^K (x_k^{(j)}(s))^2} \right) ds. \end{aligned} \quad (3.25)$$

Adding and subtracting $x_{t,i}^2 x_{t,j}^2 / \sum_{k=1}^K x_{t,k}^2$ inside the integral and applying (3.24) yield

$$\begin{aligned} \mathbb{E}_t [\langle \phi_t, x_t - x_{t+1} \rangle] &\leq \eta \sum_{1 \leq i < j \leq K} \frac{x_{t,i}^2 x_{t,j}^2}{\sum_{k=1}^K x_{t,k}^2} ((\varphi(\ell_{t,i}) - \varphi(\ell_{t,j})) (\ell_{t,i} - \ell_{t,j}) \\ &\quad + 0.07 |\varphi(\ell_{t,i}) - \varphi(\ell_{t,j})| (|\ell_{t,i} - c_{t-1}| + |\ell_{t,j} - c_{t-1}|)). \end{aligned} \quad (3.26)$$

Since $\varphi(x)$ is 1-Lipschitz on $[0, 1]$ and direct calculation shows that

$$\begin{aligned} (\ell_{t,i} - \ell_{t,j})^2 &\leq 2 ((\ell_{t,i} - c_{t-1})^2 + (\ell_{t,j} - c_{t-1})^2), \\ |\ell_{t,i} - \ell_{t,j}| (|\ell_{t,i} - c_{t-1}| + |\ell_{t,j} - c_{t-1}|) &\leq 2 ((\ell_{t,i} - c_{t-1})^2 + (\ell_{t,j} - c_{t-1})^2), \end{aligned}$$

substituting these inequalities into (3.26) gives

$$\begin{aligned} \mathbb{E}_t [\langle \phi_t, x_t - x_{t+1} \rangle] &\leq 2.14\eta \sum_{1 \leq i < j \leq K} \frac{x_{t,i}^2 x_{t,j}^2}{\sum_{k=1}^K x_{t,k}^2} ((\ell_{t,i} - c_{t-1})^2 + (\ell_{t,j} - c_{t-1})^2) \\ &= 2.14\eta \sum_{i=1}^K \left(x_{t,i}^2 - \frac{x_{t,i}^4}{\sum_{j=1}^K x_{t,j}^2} \right) (\ell_{t,i} - c_{t-1})^2 \\ &\leq 2.2\eta \sum_{i=1}^K \left(x_{t,i}^2 - \frac{x_{t,i}^4}{\sum_{j=1}^K x_{t,j}^2} \right) (\ell_{t,i} - c_{t-1})^2. \end{aligned} \quad (3.27)$$

It remains to compare the right-hand side of (3.27) with D_t . For each fixed $i \in [K]$, the KKT condition defining $x^{(i)}(1)$ gives

$$\nabla \Psi(x_t) - \nabla \Psi(x^{(i)}(1)) = c_{t-1} \mathbf{1} + \frac{\ell_{t,i} - c_{t-1}}{p_{t,i}} e_i + \lambda_{t,i} \mathbf{1}$$

for some scalar $\lambda_{t,i}$. By the definition of the Bregman divergence, we know that

$$D_\Psi(x_t, x^{(i)}(1)) + D_\Psi(x^{(i)}(1), x_t) = \left\langle \nabla \Psi(x_t) - \nabla \Psi(x^{(i)}(1)), x_t - x^{(i)}(1) \right\rangle = \frac{\ell_{t,i} - c_{t-1}}{p_{t,i}} (x_{t,i} - x_i^{(i)}(1)). \quad (3.28)$$

Next, setting $j = i$ in (3.20) and integrating over $s \in [0, 1]$ gives

$$x_{t,i} - x_i^{(i)}(1) = \eta \frac{\ell_{t,i} - c_{t-1}}{p_{t,i}} \int_0^1 \left((x_i^{(i)}(s))^2 - \frac{(x_i^{(i)}(s))^4}{\sum_{k=1}^K (x_k^{(i)}(s))^2} \right) ds. \quad (3.29)$$

Substituting (3.29) into (3.28) yields

$$\begin{aligned} D_\Psi(x_t, x^{(i)}(1)) + D_\Psi(x^{(i)}(1), x_t) &= \eta \frac{(\ell_{t,i} - c_{t-1})^2}{p_{t,i}^2} \int_0^1 \left((x_i^{(i)}(s))^2 - \frac{(x_i^{(i)}(s))^4}{\sum_{k=1}^K (x_k^{(i)}(s))^2} \right) ds \\ &= \eta \frac{(\ell_{t,i} - c_{t-1})^2}{p_{t,i}^2} \int_0^1 \left(\sum_{j \neq i} \frac{(x_i^{(i)}(s))^2 (x_j^{(i)}(s))^2}{\sum_{k=1}^K (x_k^{(i)}(s))^2} \right) ds \\ &\geq \eta \cdot \frac{0.94(\ell_{t,i} - c_{t-1})^2}{p_{t,i}^2} \left(x_{t,i}^2 - \frac{x_{t,i}^4}{\sum_{k=1}^K x_{t,k}^2} \right) \end{aligned} \quad (3.30)$$

where the last inequality uses (3.23).

Finally, as $D_t = D_\Psi(x_t, x^{(I_t)}(1))$, applying the definition of conditional expectation to both divergences and then using (3.30) gives

$$\begin{aligned}\mathbb{E}_t [D_t + D_\Psi(x_{t+1}, x_t)] &= \eta \sum_{i=1}^K \frac{(\ell_{t,i} - c_{t-1})^2}{p_{t,i}} \int_0^1 \left((x_i^{(i)}(s))^2 - \frac{(x_i^{(i)}(s))^4}{\sum_{k=1}^K (x_k^{(i)}(s))^2} \right) ds \\ &\geq 0.94\eta \sum_{i=1}^K \frac{(\ell_{t,i} - c_{t-1})^2}{p_{t,i}} \left(x_{t,i}^2 - \frac{x_{t,i}^4}{\sum_{k=1}^K x_{t,k}^2} \right) \\ &\geq 0.94\eta \sum_{i=1}^K \left(x_{t,i}^2 - \frac{x_{t,i}^4}{\sum_{k=1}^K x_{t,k}^2} \right) (\ell_{t,i} - c_{t-1})^2.\end{aligned}\tag{3.31}$$

The last inequality uses $p_{t,i} \leq 1$ and its non-negativity: $x_{t,i}^2 - \frac{x_{t,i}^4}{\sum_{k=1}^K x_{t,k}^2} = \sum_{j \neq i} \frac{x_{t,i}^2 x_{t,j}^2}{\sum_{k=1}^K x_{t,k}^2} \geq 0$.

Combining (3.27) and (3.31), and then using $D_\Psi(x_{t+1}, x_t) \leq 2D_t$, gives

$$\mathbb{E}_t [\langle \phi_t, x_t - x_{t+1} \rangle] \leq \frac{2.2}{0.94} \mathbb{E}_t [D_t + D_\Psi(x_{t+1}, x_t)] \leq \frac{6.6}{0.94} \mathbb{E}_t [D_t] \leq 8 \mathbb{E}_t [D_t].$$

□

We now apply the stability lemma to the part of the final term in Lemma 3.2 that contains the OMD iterates x_t .

Lemma 3.4. *Under the conditions of Lemma 3.3, Algorithm 1 guarantees that*

$$\mathbb{E} \left[\sum_{t=2}^T \langle \phi_t, x_{t-1} - x_t \rangle \right] \leq 8\eta \left(1 + R + \sqrt{RQ_{\infty,2}} \right).\tag{3.32}$$

Proof. For every $t \geq 2$, write $\phi_t = \phi_{t-1} + (\phi_t - \phi_{t-1})$. The left-hand side of (3.32) is therefore equal to $S_1 + S_2$, where

$$S_1 = \mathbb{E} \left[\sum_{t=2}^T \langle \phi_{t-1}, x_{t-1} - x_t \rangle \right],\tag{3.33}$$

$$S_2 = \mathbb{E} \left[\sum_{t=2}^T \langle \phi_t - \phi_{t-1}, x_{t-1} - x_t \rangle \right].\tag{3.34}$$

The term S_1 pairs ϕ_{t-1} with the OMD update performed on round $t-1$. Applying (3.13) on round $t-1$ and summing over $t = 2, \dots, T$ gives

$$S_1 \leq 8\mathbb{E} \left[\sum_{t=2}^T D_{t-1} \right] = 8\mathbb{E} \left[\sum_{s=1}^{T-1} D_s \right] \leq 4.8\eta \mathbb{E} \left[\sum_{s=1}^{T-1} r_s^2 \right] \leq 4.8\eta(1 + R),$$

where the second inequality is due to (3.11) and the last inequality follows from $r_1^2 \leq 1$ and $R = \mathbb{E}[\sum_{s=2}^T r_s^2]$.

We next control S_2 . We treat its first summand separately and write

$$S_2 = \mathbb{E} [\langle \phi_2 - \phi_1, x_1 - x_2 \rangle] + \mathbb{E} \left[\sum_{t=3}^T \langle \phi_t - \phi_{t-1}, x_{t-1} - x_t \rangle \right].$$

Applying (3.12) on round 1 with $z = \phi_2 - \phi_1$ gives

$$|\mathbb{E} [\langle \phi_2 - \phi_1, x_1 - x_2 \rangle]| \leq 1.5\mathbb{E} \left[\sqrt{\eta D_1} \sqrt{\langle x_1, (\phi_2 - \phi_1)^{\odot 2} \rangle} \right] \leq 1.5\sqrt{0.6}\eta \leq 1.2\eta.$$

Here (3.11) and $r_1^2 \leq 1$ imply $D_1 \leq 0.6\eta$, while the one-Lipschitz property of φ and $\ell_1, \ell_2 \in [0, 1]^K$ imply $\langle x_1, (\phi_2 - \phi_1)^{\odot 2} \rangle \leq 1$.

For the remaining summands, applying (3.12) on round $t - 1$ and then Cauchy–Schwarz over time and expectation gives

$$\left| \mathbb{E} \left[\sum_{t=3}^T \langle \phi_t - \phi_{t-1}, x_{t-1} - x_t \rangle \right] \right| \leq 1.5 \sqrt{\eta \mathbb{E} \left[\sum_{t=3}^T D_{t-1} \right]} \sqrt{\mathbb{E} \left[\sum_{t=3}^T \langle x_{t-1}, (\phi_t - \phi_{t-1})^{\odot 2} \rangle \right]}. \quad (3.35)$$

By (3.11) and the definition of R ,

$$\mathbb{E} \left[\sum_{t=3}^T D_{t-1} \right] = \mathbb{E} \left[\sum_{s=2}^{T-1} D_s \right] \leq 0.6\eta \mathbb{E} \left[\sum_{s=2}^{T-1} r_s^2 \right] \leq 0.6\eta R.$$

Moreover, since $\varphi'(z) = 1 - z \in [0, 1]$ for $z \in [0, 1]$, φ is 1-Lipschitz. Hence

$$\langle x_{t-1}, (\phi_t - \phi_{t-1})^{\odot 2} \rangle = \sum_{i=1}^K x_{t-1,i} (\varphi(\ell_{t,i}) - \varphi(\ell_{t-1,i}))^2 \leq \|\ell_t - \ell_{t-1}\|_{\infty}^2,$$

and therefore

$$\mathbb{E} \left[\sum_{t=3}^T \langle x_{t-1}, (\phi_t - \phi_{t-1})^{\odot 2} \rangle \right] \leq Q_{\infty,2}.$$

Substituting these two bounds into (3.35) gives

$$|S_2| \leq 1.2\eta + 1.5\sqrt{0.6\eta} \sqrt{RQ_{\infty,2}} \leq 1.2\eta \left(1 + \sqrt{RQ_{\infty,2}} \right).$$

Combining the bounds on S_1 and S_2 yields

$$S_1 + S_2 \leq 4.8\eta(1 + R) + 1.2\eta \left(1 + \sqrt{RQ_{\infty,2}} \right) \leq 8\eta \left(1 + R + \sqrt{RQ_{\infty,2}} \right),$$

which proves (3.32). $\square$

Step 3: control the difference between p_t and x_t . Define $b_t \triangleq p_t - x_t$. Then $p_{t-1} - p_t = (x_{t-1} - x_t) + (b_{t-1} - b_t)$. The first difference is handled by Lemma 3.4. It remains to control the part containing b_t , which arises because p_t places additional mass on the arm selected on the preceding round.

Lemma 3.5. *Let $b_t = p_t - x_t$. Algorithm 1 with $\eta \leq 1/162$ guarantees that*

$$\mathbb{E} \left[\sum_{t=2}^T \langle \phi_t, b_{t-1} - b_t \rangle \right] \leq 40\eta \left(1 + Q_{\infty,2} + \sqrt{RQ_{\infty,2}} \right). \quad (3.36)$$

Proof. We first explain why it suffices to control the squared norm of $\mathbb{E}[b_t]$. Since the loss table is oblivious, ϕ_t is deterministic, and rearranging the summation terms gives

$$\mathbb{E} \left[\sum_{t=2}^T \langle \phi_t, b_{t-1} - b_t \rangle \right] = \langle \phi_2, \mathbb{E}[b_1] \rangle - \langle \phi_T, \mathbb{E}[b_T] \rangle + \sum_{t=2}^{T-1} \langle \phi_{t+1} - \phi_t, \mathbb{E}[b_t] \rangle. \quad (3.37)$$

Moreover, Cauchy–Schwarz inequality and the 1-Lipschitz property of $\varphi(\cdot)$ imply that

$$\sum_{t=2}^{T-1} \langle \phi_{t+1} - \phi_t, \mathbb{E}[b_t] \rangle \leq \left(\sum_{t=2}^{T-1} \|\phi_{t+1} - \phi_t\|_{\infty}^2 \right)^{1/2} \left(\sum_{t=2}^{T-1} \|\mathbb{E}[b_t]\|_1^2 \right)^{1/2} \leq \sqrt{Q_{\infty,2}} \left(\sum_{t=2}^{T-1} \|\mathbb{E}[b_t]\|_1^2 \right)^{1/2}. \quad (3.38)$$

Therefore, it remains only to prove the following

$$\sum_{t=2}^T \|\mathbb{E}[b_t]\|_1^2 \leq 16\alpha^2(1 + R + Q_{\infty,2}). \quad (3.39)$$

For $i \in [K]$, define $\beta_{t,i} \triangleq \alpha(1 - \ell_{t-1,i}) / (1 + \alpha(1 - \ell_{t-1,i}))$. The sampling rule gives $b_t = \beta_{t,I_{t-1}}(e_{I_{t-1}} - x_t)$. Since the loss table is oblivious, $\beta_{t,i}$ is also deterministic. Therefore, direct calculation shows that for each $i \in [K]$,

$$\begin{aligned}
\mathbb{E}[b_{t,i}] &= \mathbb{E}[p_{t,i} - x_{t,i}] && \text{(by definition)} \\
&= \mathbb{E}[\beta_{t,I_{t-1}}(\mathbf{1}\{I_{t-1} = i\} - x_{t,i})] && \text{(by definition of } p_t) \\
&= \beta_{t,i}\mathbb{E}[p_{t-1,i}] - \mathbb{E}[\beta_{t,I_{t-1}}x_{t,i}] && \text{(since } \beta_{t,i} \text{ is deterministic)} \\
&= \beta_{t,i}(\mathbb{E}[p_{t-1,i}] - \mathbb{E}[x_{t,i}]) + \mathbb{E}[(\beta_{t,i} - \beta_{t,I_{t-1}})x_{t,i}] \\
&= \beta_{t,i}(\mathbb{E}[b_{t-1,i}] - (\mathbb{E}[x_{t,i}] - \mathbb{E}[x_{t-1,i}])) + \mathbb{E}[(\beta_{t,i} - \beta_{t,I_{t-1}})x_{t,i}], \tag{3.40}
\end{aligned}$$

where the last equality uses the fact that $p_{t-1} = x_{t-1} + b_{t-1}$. Since $0 \leq \beta_{t,i} \leq \alpha$ for every $i \in [K]$, taking the ℓ_1 norm in (3.40) gives

$$\|\mathbb{E}[b_t]\|_1 \leq \alpha\|\mathbb{E}[b_{t-1}]\|_1 + \alpha\|\mathbb{E}[x_t] - \mathbb{E}[x_{t-1}]\|_1 + \left\| \left(\mathbb{E}[(\beta_{t,i} - \beta_{t,I_{t-1}})x_{t,i}] \right)_{i=1}^K \right\|_1.$$

We next bound the final term. For notational convenience, we draw an auxiliary arm J from x_t conditionally on $\mathcal{F}_{t-1}$ according to x_t . Since the mapping $z \mapsto \alpha(1 - z) / (1 + \alpha(1 - z))$ is α -Lipschitz, we have $|\beta_{t,i} - \beta_{t,j}| \leq \alpha|\ell_{t-1,i} - \ell_{t-1,j}|$. The triangle inequality followed by the conditional law of J and Jensen's inequality therefore gives

$$\begin{aligned}
\left\| \left(\mathbb{E}[(\beta_{t,i} - \beta_{t,I_{t-1}})x_{t,i}] \right)_{i=1}^K \right\|_1 &\leq \mathbb{E} \left[\sum_{i=1}^K x_{t,i} |\beta_{t,i} - \beta_{t,I_{t-1}}| \right] \\
&= \mathbb{E}[|\beta_{t,J} - \beta_{t,I_{t-1}}|] \\
&\leq \alpha\mathbb{E}[|\ell_{t-1,J} - \ell_{t-1,I_{t-1}}|] \\
&\leq \alpha\sqrt{\mathbb{E}[(\ell_{t-1,J} - \ell_{t-1,I_{t-1}})^2]}.
\end{aligned}$$

To control the last expectation, observe directly from the definition of $\beta_{t,I_{t-1}}$ that $p_t = (1 - \beta_{t,I_{t-1}})x_t + \beta_{t,I_{t-1}}e_{I_{t-1}}$. Consequently, conditionally on $\mathcal{F}_{t-1}$, the draw $I_t \sim p_t$ can be generated using a Bernoulli coin that is independent of J conditionally on $\mathcal{F}_{t-1}$: set $I_t = I_{t-1}$ with probability $\beta_{t,I_{t-1}}$, and set $I_t = J$ otherwise. Under this equivalent coupling,

$$\begin{aligned}
\mathbb{E}_t[r_t^2] &= \beta_{t,I_{t-1}}(\ell_{t,I_{t-1}} - \ell_{t-1,I_{t-1}})^2 + (1 - \beta_{t,I_{t-1}})\mathbb{E}[(\ell_{t,J} - \ell_{t-1,I_{t-1}})^2 \mid \mathcal{F}_{t-1}] \\
&\geq (1 - \alpha)\mathbb{E}[(\ell_{t,J} - \ell_{t-1,I_{t-1}})^2 \mid \mathcal{F}_{t-1}].
\end{aligned}$$

Taking the full expectation and using $(a + b)^2 \leq 2a^2 + 2b^2$ now gives

$$\begin{aligned}
\left\| \left(\mathbb{E}[(\beta_{t,i} - \beta_{t,I_{t-1}})x_{t,i}] \right)_{i=1}^K \right\|_1^2 &\leq \alpha^2\mathbb{E}[(\ell_{t-1,J} - \ell_{t-1,I_{t-1}})^2] \\
&\leq \frac{2\alpha^2}{1 - \alpha}\mathbb{E}[r_t^2] + 2\alpha^2\|\ell_t - \ell_{t-1}\|_\infty^2 \\
&\leq 3\alpha^2(\mathbb{E}[r_t^2] + \|\ell_t - \ell_{t-1}\|_\infty^2),
\end{aligned}$$

where the last inequality uses $\alpha \leq 8/162$. Summing over $t = 2, \dots, T$ yields

$$\sum_{t=2}^T \left\| \left(\mathbb{E}[(\beta_{t,i} - \beta_{t,I_{t-1}})x_{t,i}] \right)_{i=1}^K \right\|_1^2 \leq 3\alpha^2(R + Q_{\infty,2}).$$

It remains to bound the movement of x_t in (3.40). For every $t \geq 2$, Jensen's and Cauchy-Schwarz

inequality give

$$\begin{aligned}
\|\mathbb{E}[x_t] - \mathbb{E}[x_{t-1}]\|_1^2 &\leq \mathbb{E}[\|x_t - x_{t-1}\|_1^2] \\
&\leq \mathbb{E}\left[\sum_{i=1}^K \frac{(x_{t,i} - x_{t-1,i})^2}{x_{t-1,i}}\right] \\
&\leq \mathbb{E}\left[\sum_{i=1}^K \frac{(x_{t,i} - x_{t-1,i})^2}{x_{t-1,i}^2}\right] \leq 2.1\eta\mathbb{E}[D_{t-1}],
\end{aligned}$$

where the third inequality uses $x_{t-1,i} \leq 1$, and the last inequality is (3.19) applied on round $t-1$. Summing this display and then applying (3.11) gives

$$\sum_{t=2}^T \|\mathbb{E}[x_t] - \mathbb{E}[x_{t-1}]\|_1^2 \leq 2.1\eta\mathbb{E}\left[\sum_{t=1}^{T-1} D_t\right] \leq 1.26\eta^2\mathbb{E}\left[\sum_{t=1}^{T-1} r_t^2\right] \leq 2\eta^2(1+R).$$

Finally, we square the preceding bound on $\|\mathbb{E}[b_t]\|_1$, use $(a+b+c)^2 \leq 3a^2 + 3b^2 + 3c^2$, and sum over $t = 2, \dots, T$. The two bounds just proved yield

$$\sum_{t=2}^T \|\mathbb{E}[b_t]\|_1^2 \leq 3\alpha^2 \sum_{t=2}^T \|\mathbb{E}[b_{t-1}]\|_1^2 + 6\alpha^2\eta^2(1+R) + 9\alpha^2(R+Q_{\infty,2}).$$

Rearranging the terms and using the fact that $6\eta^2 < 1$, we know that

$$(1 - 3\alpha^2) \sum_{t=2}^T \|\mathbb{E}[b_t]\|_1^2 \leq 10\alpha^2(1+R+Q_{\infty,2}),$$

and $1 - 3\alpha^2 > 99/100$ proves (3.39).

Finally, the endpoint terms in (3.37) are at most α because $b_1 = 0$, $\|\mathbb{E}[b_T]\|_1 \leq 2\alpha$, and $0 \leq \varphi \leq 1/2$. Combining (3.38) and (3.39) gives

$$\mathbb{E}\left[\sum_{t=2}^T \langle \phi_t, b_{t-1} - b_t \rangle\right] \leq \alpha + 4\alpha\sqrt{Q_{\infty,2}(1+R+Q_{\infty,2})} \leq 5\alpha\left(1+Q_{\infty,2} + \sqrt{RQ_{\infty,2}}\right).$$

Since $\alpha = 8\eta$, this proves (3.36). $\square$

3.2 Proof of Theorem 3.1

Proof. Fix an arbitrary arm $i \in [K]$ and the corresponding comparator $u^{(i)}$ defined above. The OMD decomposition (3.3) reduces the proof to bounding $D_t + B_t$. We now combine the three steps above in the same order in which they were proved.

Lemma 3.2 and the identity $p_t = x_t + b_t$ give, up to the $\mathcal{O}(1)$ first-round boundary,

$$\begin{aligned}
\mathbb{E}\left[\sum_{t=1}^T (D_t + B_t)\right] &= \mathbb{E}\left[\sum_{t=2}^T D_t\right] + 4\eta\mathbb{E}\left[\sum_{t=2}^T v_t^2\right] - 4\eta R \\
&\quad + \alpha\mathbb{E}\left[\sum_{t=2}^T \langle \phi_t, x_{t-1} - x_t \rangle\right] + \alpha\mathbb{E}\left[\sum_{t=2}^T \langle \phi_t, b_{t-1} - b_t \rangle\right] + \mathcal{O}(1).
\end{aligned} \tag{3.41}$$

We bound the four terms on the right-hand side separately. First, Lemma 3.3 gives $\mathbb{E}[\sum_{t=2}^T D_t] \leq 0.6\eta R$. Second, $v_t^2 \leq \|\ell_t - \ell_{t-1}\|_\infty^2$ pathwise, and hence $\mathbb{E}[\sum_{t=2}^T v_t^2] \leq Q_{\infty,2}$. Third, Lemma 3.4, together with $\alpha = 8\eta$, bounds the term containing $x_{t-1} - x_t$ by $64\eta^2(1+R+\sqrt{RQ_{\infty,2}})$. Finally, Lemma 3.5 bounds the term containing $b_{t-1} - b_t$ by $320\eta^2(1+Q_{\infty,2}+\sqrt{RQ_{\infty,2}})$.

It remains to absorb the terms depending positively on R . Since $\eta \leq 1/162$, we have $64\eta^2 R \leq 0.4\eta R$ and $384\eta^2 \sqrt{RQ_{\infty,2}} \leq 0.8\eta R + 1.8\eta Q_{\infty,2}$. Therefore, the entire positive R -dependent contribution is at most $1.2\eta R$. Substituting the four bounds into (3.41) therefore gives

$$\mathbb{E} \left[\sum_{t=1}^T (D_t + B_t) \right] \leq (0.6 - 4 + 1.2)\eta R + \mathcal{O}(\eta Q_{\infty,2} + 1) \leq \mathcal{O}(\eta Q_{\infty,2} + 1). \quad (3.42)$$

This proves (3.4). Finally, $D_\Psi(u^{(i)}, x_1) \leq \mathcal{O}(K \log(KT)/\eta)$. Substituting (3.42) into (3.3) and maximizing over $i \in [K]$ give the fixed-rate inequality

$$\text{Reg}_T \leq \mathcal{O} \left(\frac{K \log(KT)}{\eta} + \eta Q_{\infty,2} + 1 \right). \quad (3.43)$$

Tuning η optimally leads to (3.1). $\square$

4 Unknown $Q_{\infty,2}$: Adaptive Tuning Via Bounded-Scale Estimator

Section 3 tunes Algorithm 1 using the value of $Q_{\infty,2}$. To remove this knowledge, a standard approach is the doubling trick: start from an initial guess, run the algorithm with the learning rate corresponding to that guess, and restart with a larger guess once the cumulative variation exceeds the current threshold. Such restart constructions are common in adaptive bandit algorithms (e.g., Lee et al. [15], Wei and Luo [24]). The difficulty here is that the second-order path-length $Q_{\infty,2}$ is not observed under bandit feedback.

Recall that $r_t \triangleq c_t - c_{t-1}$ and $v_t \triangleq \ell_{t,I_{t-1}} - \ell_{t-1,I_{t-1}}$. The quantity v_t^2 is observed whenever $I_t = I_{t-1}$ because $r_t = v_t$ on this event. This suggests the following single-round path-length estimator

$$\hat{v}_t^2 \triangleq \frac{\mathbf{1}(I_t = I_{t-1}) r_t^2}{p_{t,I_t}}. \quad (4.1)$$

A direct calculation gives $\mathbb{E}_t[\hat{v}_t^2] = v_t^2$. It is therefore natural to use the cumulative sum of $\hat{v}_t^2$ to estimate the total second-order path length. Under Algorithm 1, however, the denominator in (4.1) can be arbitrarily small. Indeed, if $c_{t-1} = 1$, then the additional mass on I_{t-1} vanishes and, on the event $I_t = I_{t-1}$, $p_{t,I_t} = x_{t,I_{t-1}}$. Although the log-barrier guarantees that this probability is positive, it does not provide a uniform lower bound. Consequently, $\hat{v}_t^2$ can be arbitrarily large, and the cumulative estimate can overshoot a doubling threshold by an uncontrolled amount.

We therefore make one modification to Algorithm 1: we replace $\lambda_t = \alpha(1 - c_{t-1})$ by $\lambda_t = \alpha(2 - c_{t-1})$. Equivalently, we add an α floor to the additional mass assigned to I_{t-1} , thereby keeping the denominator bounded away from zero. The loss estimator $\hat{\ell}_t$ and the OMD update remain unchanged. Since $c_{t-1} \in [0, 1]$, this modification gives $p_{t,I_{t-1}} \geq \alpha/(1 + \alpha)$ and makes (4.1) uniformly bounded. We later show that this uniform bound controls the overshoot of the cumulative estimator at every phase transition. The following lemma records the two properties of the estimator that are needed by the doubling argument.

Lemma 4.1. *Suppose $p_t = (x_t + \lambda_t e_{I_{t-1}})/(1 + \lambda_t)$ with $\lambda_t = \alpha(2 - c_{t-1})$. Then the estimator in (4.1) satisfies*

$$\mathbb{E}_t[\hat{v}_t^2] = v_t^2 \quad \text{and} \quad 0 \leq \hat{v}_t^2 \leq \frac{1 + \alpha}{\alpha}. \quad (4.2)$$

Proof. The definition of p_t gives $p_{t,I_{t-1}} \geq \lambda_t/(1 + \lambda_t) \geq \alpha/(1 + \alpha)$. Moreover, $r_t = v_t$ on the event $I_t = I_{t-1}$. Therefore,

$$\mathbb{E}_t[\hat{v}_t^2] = \sum_{i=1}^K p_{t,i} \frac{\mathbf{1}(i = I_{t-1})(\ell_{t,i} - c_{t-1})^2}{p_{t,i}} = v_t^2.$$

For the upper bound, the numerator is nonzero only when $I_t = I_{t-1}$, in which case $p_{t,I_t} = p_{t,I_{t-1}} \geq \alpha/(1 + \alpha)$. The result now follows from $|r_t| \leq 1$. $\square$

We are now ready to present our adaptive algorithm. Define $A_K = 64K \log(KT)$ and $\eta_0 = 10^{-5}$. The algorithm starts with the threshold $H_0 = 4A_K/\eta_0^2$. During phase j , it uses $\eta_j \triangleq 2^{-j}\eta_0$, $\alpha_j \triangleq 8\eta_j$, and $H_j \triangleq 4A_K/\eta_j^2$. It samples from p_t and updates x_{t+1} exactly as in Algorithm 1, except for the α_j floor in Line 16. After every noninitial round of the phase, it adds $\hat{v}_t^2$ to its estimate $\hat{H}_j$. If $\hat{H}_j$ exceeds H_j , phase j ends and the outer loop begins phase $j+1$ on the next round with a uniform initial distribution. The scalar $\hat{H}_j$ is initialized to zero when phase j begins. The following theorem shows that Algorithm 2 achieves $\tilde{\mathcal{O}}(K + \sqrt{KQ_{\infty,2}})$ without knowing $Q_{\infty,2}$. The overhead is a $\sqrt{\log T}$ factor compared to the guarantee in Theorem 3.1.

Algorithm 2: Adaptive tuning without knowledge of $Q_{\infty,2}$

Input: $K > 0, T > 0$

- 1 Set $A_K = 64K \log(KT)$, $\eta_0 = 10^{-5}$, $j = 0$, $s_0 = t = 1$, and $c_0 = 0$
- 2 **while** $t \leq T$ **do**
- 3 Set $\eta_j = 2^{-j}\eta_0$, $\alpha_j = 8\eta_j$, $H_j = 4A_K/\eta_j^2$, and $\hat{H}_j = 0$
- 4 Phase j starts at round $s_j = t$
- 5 Set $x_t = p_t = \frac{1}{K}\mathbf{1}$, and $\hat{v}_{s_j}^2 = 0$
- 6 **while** $t \leq T$ **do**
- 7 Draw $I_t \sim p_t$ and observe $c_t = \ell_{t,I_t}$
- 8 Set $\hat{\ell}_{t,i} = c_{t-1} + \mathbf{1}(I_t = i)(c_t - c_{t-1})/p_{t,i}$ for every $i \in [K]$
- 9 Set $x_{t+1} = \arg \min_{x \in \Delta_K} \left(\langle x, \hat{\ell}_t \rangle + D_{\Psi_j}(x, x_t) \right)$, where $\Psi_j(x) = \eta_j^{-1} \sum_{i=1}^K \log(1/x_i)$
- 10 **if** $t > s_j$ **then**
- 11 Set $r_t = c_t - c_{t-1}$, $\hat{v}_t^2 = \mathbf{1}(I_t = I_{t-1})r_t^2/p_{t,I_t}$, and $\hat{H}_j \leftarrow \hat{H}_j + \hat{v}_t^2$
- 12 **if** $\hat{H}_j > H_j$ **and** $t < T$ **then**
- 13 Set $t \leftarrow t + 1$, $j \leftarrow j + 1$
- 14 **break**
- 15 **else if** $t < T$ **then**
- 16 Set $\lambda_{t+1} = \alpha_j(2 - c_t)$ and $p_{t+1} = (x_{t+1} + \lambda_{t+1}e_{I_t})/(1 + \lambda_{t+1})$
- 17 Set $t \leftarrow t + 1$

Theorem 4.2. *For every $K \geq 2$, $T \geq 2$, and oblivious loss sequence $\ell_1, \dots, \ell_T \in [0, 1]^K$, Algorithm 2 requires no knowledge of $Q_{\infty,2}$ and guarantees that*

$$\text{Reg}_T \leq \mathcal{O} \left(K \log(KT) + \sqrt{KQ_{\infty,2} \log(KT) \log T} \right). \quad (4.3)$$

4.1 Analysis of Algorithm 2

We first relate the cumulative estimate $\hat{H}_j$ to the two squared quantities in the analysis of Section 3. Let J_t denote the value of the algorithm's phase counter immediately before I_t is drawn. Thus J_t is $\mathcal{F}_{t-1}$ -measurable. In the analysis below, j is always a deterministic phase label; the randomness of the active phase is carried by J_t . Consistently with Algorithm 2, define the first round of phase j by $s_j \triangleq \inf\{t \in [T] : J_t = j\}$, with the convention $\inf \emptyset = \infty$, and define $\chi_{j,t} \triangleq \mathbf{1}(J_t = j, t > s_j)$. Hence $\chi_{j,t}$ is the indicator that round t is a noninitial round of phase j , and it is $\mathcal{F}_{t-1}$ -measurable. Let $\rho_j \triangleq \Pr(s_j \leq T)$ be the probability that phase j is reached. For notational convenience, we define the following two quantities:

$$R_j \triangleq \mathbb{E} \left[\sum_{t=2}^T \chi_{j,t} r_t^2 \right], \quad (4.4)$$

$$P_j \triangleq \mathbb{E} \left[\sum_{t=2}^T \chi_{j,t} v_t^2 \right]. \quad (4.5)$$

Specifically, the quantity R_j is the expected cumulative residual square over the noninitial rounds of phase j and the quantity P_j is the expected cumulative squared loss movement along the previously selected arm. By the unbiasedness of $\widehat{v}_t^2$, it is also the expected value of the doubling statistic accumulated during phase j . The next lemma shows that the doubling statistic has expectation P_j , has controlled overshoot, and that the sum of these phase-wise expectations is at most the true squared path-length.

Lemma 4.3. *For every deterministic phase label $j \in \{0, \dots, T-1\}$, Algorithm 2 guarantees that $H_j \rho_{j+1} \leq P_j \leq 2H_j \rho_j$ and $\sum_{j=0}^{T-1} P_j \leq Q_{\infty,2}$.*

Proof. For a deterministic j , let $\widehat{H}_j^{\text{fin}} \triangleq \sum_{t=2}^T \chi_{j,t} \widehat{v}_t^2$ denote the final value accumulated by phase j ; it is zero on the event $s_j = \infty$. Since $\chi_{j,t}$ is $\mathcal{F}_{t-1}$ -measurable, Lemma 4.1 and the tower property give $\mathbb{E}[\chi_{j,t} \widehat{v}_t^2] = \mathbb{E}[\chi_{j,t} v_t^2]$. Summing over the deterministic set $t = 2, \dots, T$ therefore yields $\mathbb{E}[\widehat{H}_j^{\text{fin}}] = P_j$.

Pathwise, reaching phase $j+1$ requires phase j to cross its threshold, so $H_j \mathbf{1}(s_{j+1} \leq T) \leq \widehat{H}_j^{\text{fin}}$. Before the triggering round the accumulated value is at most H_j , while Lemma 4.1 bounds the triggering increment by $(1 + \alpha_j)/\alpha_j$. Since $\alpha_j = 8\eta_j$ and $\eta_j \leq 10^{-5}$, this increment is at most $1/(4\eta_j)$ as

$$\frac{1 + \alpha_j}{\alpha_j} = \frac{1}{8\eta_j} + 1 \leq \frac{1}{4\eta_j}.$$

Moreover, $H_j = 4A_K/\eta_j^2 \geq 1/(4\eta_j)$. Consequently, $\widehat{H}_j^{\text{fin}} \leq 2H_j \mathbf{1}(s_j \leq T)$ pathwise. Taking expectations in these two pathwise inequalities proves $H_j \rho_{j+1} \leq P_j \leq 2H_j \rho_j$.

Finally, for every t , at most one indicator $\chi_{j,t}$ equals one. Since the loss table is fixed and $v_t^2 \leq \|\ell_t - \ell_{t-1}\|_\infty^2$, we have

$$\sum_{j=0}^{T-1} P_j = \mathbb{E} \left[\sum_{t=2}^T \sum_{j=0}^{T-1} \chi_{j,t} v_t^2 \right] \leq \sum_{t=2}^T \|\ell_t - \ell_{t-1}\|_\infty^2 = Q_{\infty,2}.$$

This proves the second claim. $\square$

We next show in the following lemma how to control the contribution of the noninitial rounds of each phase. Specifically, as shown in (3.3), in order to bound the regret, the relevant quantity after the standard OMD decomposition is the sum of $D_t + B_t$, where we recall that $B_t \triangleq \langle p_t - x_t, \ell_t \rangle$ and with a slight abuse of notation, we define $D_t \triangleq D_{\Psi_{j_t}}(x_t, x_{t+1})$. Here and throughout the phasewise analysis, if round t ends a phase, x_{t+1} denotes the OMD iterate computed in Line 9 before the subsequent reset. The initial round of each phase is excluded from $\chi_{j,t}$ because its sampling distribution is reset to the uniform distribution and we do not count the path-length quantity across phase boundaries. We will handle its contribution separately in the proof of Theorem 4.2.

Lemma 4.4. *For every deterministic phase label $j \in \{0, \dots, T-1\}$,*

$$\mathbb{E} \left[\sum_{t=2}^T \chi_{j,t} (D_t + B_t) \right] \leq \mathcal{O}(\eta_j P_j + \rho_j). \quad (4.6)$$

Proof. Fix a phase label $j \in \{0, 1, \dots, T-1\}$. The additional α_j in the sampling weight $\lambda_t = \alpha_j(2 - c_{t-1})$ changes the scalar function used in Lemma 3.2. Therefore, we define $\varphi_2(z) \triangleq 2z - z^2/2$, so that $\varphi_2(z) = \varphi(z) + z$, and set

$$\phi_{2,t} \triangleq (\varphi_2(\ell_{t,1}), \dots, \varphi_2(\ell_{t,K})).$$

We decompose B_t at every noninitial round of phase j by following the same add-and-subtract calculation as in Lemma 3.2. Specifically, the definition of the sampling distribution gives

$$B_t = \langle p_t - x_t, \ell_t \rangle = \alpha_j(2 - c_{t-1}) \sum_{i=1}^K p_{t,i} (\ell_{t,I_{t-1}} - \ell_{t,i}). \quad (4.7)$$

Direct calculation shows that

$$\begin{aligned} & 4\eta_j ((\ell_{t,i} - c_{t-1})^2 - (\ell_{t,I_{t-1}} - c_{t-1})^2) + \alpha_j (2 - c_{t-1})(\ell_{t,I_{t-1}} - \ell_{t,i}) \\ &= 8\eta_j (\ell_{t,I_{t-1}} - \ell_{t,i}) \left(2 - \frac{\ell_{t,I_{t-1}} + \ell_{t,i}}{2} \right) = \alpha_j (\varphi_2(\ell_{t,I_{t-1}}) - \varphi_2(\ell_{t,i})). \end{aligned} \quad (4.8)$$

Multiplying (4.8) by $p_{t,i}$, summing over $i \in [K]$, and using the fact that $\mathbb{E}_t[r_t^2] = \sum_{i=1}^K p_{t,i}(\ell_{t,i} - c_{t-1})^2$ and $v_t^2 = (\ell_{t,I_{t-1}} - c_{t-1})^2$ yield

$$B_t = 4\eta_j v_t^2 - 4\eta_j \mathbb{E}_t[r_t^2] + \alpha_j (\varphi_2(\ell_{t,I_{t-1}}) - \mathbb{E}_t[\varphi_2(\ell_{t,I_t})]). \quad (4.9)$$

Since $\chi_{j,t}$ is $\mathcal{F}_{t-1}$ -measurable, the tower property gives $\mathbb{E}[\chi_{j,t} \mathbb{E}_t[\varphi_2(\ell_{t,I_t})]] = \mathbb{E}[\chi_{j,t} \varphi_2(\ell_{t,I_t})]$. It remains to bound

$$\alpha_j \mathbb{E} \left[\sum_{t=2}^T \chi_{j,t} (\varphi_2(\ell_{t,I_{t-1}}) - \varphi_2(\ell_{t,I_t})) \right].$$

Two deterministic-horizon steps from Section 3 cannot be obtained by simply multiplying each summand by $\chi_{j,t}$. First, $\chi_{j,t}$ is $\mathcal{F}_{t-1}$ -measurable but need not be $\mathcal{F}_{t-2}$ -measurable. Although

$$\mathbb{E}[\mathbf{1}\{I_{t-1} = i\} \mid \mathcal{F}_{t-2}] = p_{t-1,i},$$

the indicator $\chi_{j,t}$ cannot in general be taken outside this conditional expectation. Consequently, in general, $\mathbb{E}[\chi_{j,t} \mathbf{1}\{I_{t-1} = i\}] \neq \mathbb{E}[\chi_{j,t} p_{t-1,i}]$.

The second issue concerns the phasewise tuning of η_j using the path-length estimator. With $d_{t,i} = \ell_{t,i} - \ell_{t-1,i}$, define

$$Q_j^{\max} \triangleq \mathbb{E} \left[\sum_{t=2}^T \chi_{j,t} \max_{i \in [K]} d_{t,i}^2 \right]. \quad (4.10)$$

The phase indicators are disjoint, and hence $\sum_j Q_j^{\max} \leq Q_{\infty,2}$. Even after handling the first issue through stopping-boundary corrections, a direct phasewise application of the maximum-norm argument from Section 3 would give a bound of the form

$$\mathbb{E} \left[\sum_{t=2}^T \chi_{j,t} (D_t + B_t) \right] \leq \mathcal{O}(\eta_j P_j + \eta_j^2 Q_j^{\max} + \rho_j). \quad (4.11)$$

Summing the additional term over phases yields only $\sum_j \eta_j^2 Q_j^{\max} \leq \eta_0^2 \sum_j Q_j^{\max} \leq \eta_0^2 Q_{\infty,2}$, which depends linearly on $Q_{\infty,2}$ and is therefore insufficient for the desired bound. The phasewise estimator, however, controls

$$P_j \triangleq \sum_{t=2}^T \sum_{i=1}^K \mathbb{E}[\chi_{j,t} \mathbf{1}\{I_{t-1} = i\}] d_{t,i}^2. \quad (4.12)$$

Therefore, to exploit the phasewise choice of η_j , the analysis must replace $Q_j^{\max}$ by P_j .

The first issue can be handled by a one-round correction. If $A_{j,t} = \mathbf{1}\{J_t = j\}$, then

$$A_{j,t-1} = \chi_{j,t} + \delta_{j,t}, \quad \mathbb{E} \left[\sum_{t=2}^T \delta_{j,t} \right] \leq \rho_j, \quad (4.13)$$

where $A_{j,t-1}$ is measurable before I_{t-1} is drawn. At the boundary we use the OMD update computed before the subsequent reset. This extension is sufficient for the OMD part of Lemma 3.4: if $g_t = \phi_{2,t} - \phi_{2,t-1}$, then

$$\mathbb{E} \left[\sum_{t=2}^T A_{j,t-1} \langle x_{t-1}, g_t^{\odot 2} \rangle \right] \leq 5(P_j + \rho_j). \quad (4.14)$$

Here we use $|g_{t,i}| \leq 2|d_{t,i}|$ by the 2-Lipschitzness of $\varphi_2(\cdot)$ and $p_{t-1,i} \geq x_{t-1,i}/(1 + 2\alpha_j)$.

The second issue requires a self-normalized replacement for the unavailable maximum-coordinate bound. Following the notation in Section 3, we set $b_t \triangleq p_t - x_t$. With $q_{j,t,i} \triangleq \mathbb{E}[\chi_{j,t}(\mathbf{1}\{I_t = i\} + x_{t,i})]$, Cauchy-Schwarz inequality gives that

$$\left| \sum_{t=2}^{T-1} \sum_{i=1}^K g_{t+1,i} \mathbb{E}[\chi_{j,t} b_{t,i}] \right| \leq \left(\sum_{t,i} \frac{(\mathbb{E}[\chi_{j,t} b_{t,i}])^2}{q_{j,t,i}} \right)^{1/2} \left(\sum_{t,i} q_{j,t,i} g_{t+1,i}^2 \right)^{1/2}. \quad (4.15)$$

With a more careful calculation, we bound the right-hand side terms as follows

$$\begin{aligned} \sum_{t,i} \frac{(\mathbb{E}[\chi_{j,t} b_{t,i}])^2}{q_{j,t,i}} &\leq \mathcal{O}(\eta_j^2 R_j + \alpha_j^2 P_j + \alpha_j \rho_j), \\ \sum_{t,i} q_{j,t,i} g_{t+1,i}^2 &\leq \mathcal{O}(\eta_j^2 R_j + P_j + \rho_j). \end{aligned} \quad (4.16)$$

The rest of the proof uses $p_t = x_t + b_t$ in the same order as Lemmas 3.4 and 3.5. It gives, for every j ,

$$\alpha_j \mathbb{E} \left[\sum_{t=2}^T \chi_{j,t} (\varphi_2(\ell_{t,I_{t-1}}) - \varphi_2(\ell_{t,I_t})) \right] \leq 2000 \eta_j^2 R_j + \mathcal{O}(\eta_j^2 P_j + \rho_j). \quad (4.17)$$

The full proof of (4.17) is deferred to Appendix A. The $\mathcal{O}(\rho_j)$ additional term bounds the initialization term and the single stopping-boundary term of phase j .

It remains to control D_t . Since $\lambda_t = \alpha_j(2 - c_{t-1}) \leq 2\alpha_j$, the modified sampling rule satisfies $p_{t,i} \geq x_{t,i}/(1 + 2\alpha_j)$. Applying the direct calculation in (3.18) with this lower bound gives

$$D_t \leq \frac{(1 + 2\alpha_j)^2}{2(1 - \eta_j(1 + 2\alpha_j))} \eta_j r_t^2 \leq 0.6 \eta_j r_t^2, \quad (4.18)$$

where the last inequality follows from $\alpha_j = 8\eta_j$ and $\eta_j \leq 10^{-5}$.

Multiplying (4.9) by $\chi_{j,t}$, taking expectations, and summing over t now give

$$\mathbb{E} \left[\sum_{t=2}^T \chi_{j,t} (D_t + B_t) \right] \leq (-3.4\eta_j + 2000\eta_j^2) R_j + \mathcal{O}(\eta_j P_j + \rho_j).$$

Since $2000\eta_j \leq 0.02$, the coefficient of R_j is negative. Dropping this term proves (4.6), which is exactly the phasewise counterpart of (3.42). $\square$

We are now ready to prove Theorem 4.2.

Proof of Theorem 4.2. Fix an arbitrary arm $i \in [K]$. On every realization for which phase j is reached, its rounds form a contiguous interval. Sum (3.2) over this interval and compare with the smoothed version of e_i used in Section 3. The OMD divergences telescope pathwise. Since the phase starts from the uniform distribution, the comparator term is at most A_K/η_j on the event that the phase is reached. Moreover, $p_{s_j} = x_{s_j}$, so the initial round has no sampling bias, and its remaining OMD and smoothing terms are at most 3. Lemma 4.4 therefore gives

$$\mathbb{E} \left[\sum_{t=1}^T \mathbf{1}(J_t = j) (\ell_{t,I_t} - \ell_{t,i}) \right] \leq \underbrace{\frac{A_K \rho_j}{\eta_j}}_{\text{comparator term}} + \underbrace{\mathbb{E} \left[\sum_{t=2}^T \chi_{j,t} (D_t + B_t) \right]}_{\text{noninitial rounds}} + 3\rho_j \leq \frac{A_K \rho_j}{\eta_j} + \mathcal{O}(\eta_j P_j + \rho_j). \quad (4.19)$$

It remains to sum this bound over the phases. From Lemma 4.3 and $H_j = 4A_K/\eta_j^2$, we know that

$$\eta_j P_j \leq \sqrt{8A_K \rho_j P_j}. \quad (4.20)$$

The lower bound in the same lemma also gives $4A_K\rho_{j+1}/\eta_j^2 \leq P_j$. Since $\eta_{j+1} = \eta_j/2$, we obtain $A_K\rho_{j+1}/\eta_{j+1} \leq \eta_j P_j/2$. Thus the movement term of phase j also controls the comparator term of phase $j+1$.

Next, we show that the number of phases is logarithmic. Indeed, Lemma 4.1 gives $\hat{v}_t^2 \leq 1/(4\eta_j)$ during phase j . A completed phase must cross $H_j = 4A_K/\eta_j^2$ and hence contains at least $16A_K/\eta_j$ noninitial rounds. Let M denote the random number of phases reached. Pathwise, summing this lower bound over the first $M-1$ completed phases gives $T \geq 16A_K(2^{M-1}-1)/\eta_0$, and therefore $M \leq 2\log_2(2T)$. Since $M = \sum_{j=0}^{T-1} \mathbf{1}(s_j \leq T)$, taking expectations yields

$$\sum_{j=0}^{T-1} \rho_j = \mathbb{E}[M] \leq 2\log_2(2T). \quad (4.21)$$

Finally, the phase indicators partition the horizon. Summing (4.19), bounding every comparator term except the first one by the preceding-phase movement term as above, and using (4.20) yield

$$\begin{aligned} \text{Reg}_T(i) &\leq \frac{A_K}{\eta_0} + \mathcal{O}\left(\sum_{j=0}^{T-1} \sqrt{A_K\rho_j P_j} + \sum_{j=0}^{T-1} \rho_j\right) \\ &\leq \frac{A_K}{\eta_0} + \mathcal{O}\left(\sqrt{A_K \log(T) \sum_{j=0}^{T-1} P_j} + \log T\right) \\ &\leq \mathcal{O}\left(K \log(KT) + \sqrt{KQ_{\infty,2} \log(KT) \log T}\right), \end{aligned}$$

where the second inequality uses Cauchy–Schwarz and the last one uses Lemma 4.3, $A_K = 64K \log(KT)$, and $\eta_0 = 10^{-5}$. The bound holds for every $i \in [K]$. Taking the maximum over i proves (4.3). $\square$

5 Conclusion

We study second-order path-length guarantees for adversarial multi-armed bandits under ordinary bandit feedback. Our main result shows that, when $Q_{\infty,2}$ is known, the recent-arm optimistic log-barrier algorithm of Bubeck et al. [4] already achieves $O\left(K \log(KT) + \sqrt{K \log(KT)(1 + Q_{\infty,2})}\right)$ expected regret against oblivious loss sequences, but requires a more refined analysis. We further remove the knowledge of $Q_{\infty,2}$ through an adaptive restart scheme. Together with the $\Omega(\sqrt{KQ_{\infty,2}})$ lower bound, these results establish the optimal dependence on the second-order path length up to logarithmic factors.

Several questions remain open. First, the additional $\sqrt{\log T}$ factor in the adaptive guarantee arises from the phase-based tuning argument, and it would be interesting to determine whether the oracle rate can be achieved without knowing $Q_{\infty,2}$. Second, our analysis relies essentially on the loss sequence being oblivious. Extending the second-order guarantee to adaptive adversaries would require a different treatment of the signed difference terms used in the proof. More broadly, it would be interesting to understand whether similar second-order path-length guarantees can be obtained in richer partial-information models, such as linear bandits or MDPs.

Acknowledgments. The authors used GPT-5.6 for assistance with writing and to explore proof strategies. The authors have checked the arguments and take full responsibility for the content of the paper.

References

- [1] Peter Auer, Nicolò Cesa-Bianchi, Yoav Freund, and Robert E. Schapire. The nonstochastic multiarmed bandit problem. *SIAM Journal on Computing*, 32(1), 2002.
- [2] Sébastien Bubeck and Aleksandrs Slivkins. The best of both worlds: Stochastic and adversarial bandits. In *Conference on Learning Theory*, 2012.

- [3] Sébastien Bubeck, Michael B. Cohen, and Yuanzhi Li. Sparsity, variance and curvature in multi-armed bandits. In *Algorithmic Learning Theory*, 2018.
- [4] Sébastien Bubeck, Yuanzhi Li, Haipeng Luo, and Chen-Yu Wei. Improved path-length regret bounds for bandits. In *Conference on Learning Theory*, 2019.
- [5] Liyu Chen, Haipeng Luo, and Chen-Yu Wei. Impossible tuning made possible: A new expert algorithm and its applications. In *Conference on Learning Theory*, 2021.
- [6] Chao-Kai Chiang, Tianbao Yang, Chia-Jung Lee, Mehrdad Mahdavi, Chi-Jen Lu, Rong Jin, and Shenghuo Zhu. Online optimization with gradual variations. In *Conference on Learning Theory*, 2012.
- [7] Elad Hazan and Satyen Kale. Better algorithms for benign bandits. *Journal of Machine Learning Research*, 12, 2011.
- [8] Shinji Ito and Kei Takemura. Best-of-three-worlds linear bandit algorithm with variance-adaptive regret bounds. In *Conference on Learning Theory*, 2023.
- [9] Shinji Ito, Taira Tsuchiya, and Junya Honda. Adversarially robust multi-armed bandit algorithm with variance-dependent regret bounds. In *Conference on Learning Theory*, 2022.
- [10] Tiancheng Jin and Haipeng Luo. Simultaneously learning stochastic and adversarial episodic MDPs with known transition. In *Advances in Neural Information Processing Systems*, volume 33, 2020.
- [11] Tiancheng Jin, Longbo Huang, and Haipeng Luo. The best of both worlds: stochastic and adversarial episodic mdps with unknown transition. *Advances in Neural Information Processing Systems*, 34, 2021.
- [12] Tiancheng Jin, Junyan Liu, and Haipeng Luo. Improved best-of-both-worlds guarantees for multi-armed bandits: Ftrl with general regularizers and multiple optimal arms. *Advances in Neural Information Processing Systems*, 36, 2023.
- [13] Joon Kwon and Vianney Perchet. Gains and losses are fundamentally different in regret minimization: The sparse case. *Journal of Machine Learning Research*, 17, 2016.
- [14] Chung-Wei Lee, Haipeng Luo, Chen-Yu Wei, and Mengxiao Zhang. Bias no more: High-probability data-dependent regret bounds for adversarial bandits and MDPs. In *Advances in Neural Information Processing Systems*, volume 33, 2020.
- [15] Chung-Wei Lee, Haipeng Luo, and Mengxiao Zhang. A closer look at small-loss bounds for bandits with graph feedback. In *Conference on Learning Theory*, 2020.
- [16] Jongyeong Lee, Junya Honda, Shinji Ito, and Min-hwan Oh. Follow-the-perturbed-leader with fréchet-type tail distributions: Optimality in adversarial bandits and best-of-both-worlds. In *Conference on Learning Theory*, 2024.
- [17] Thodoris Lykouris, Karthik Sridharan, and Éva Tardos. Small-loss bounds for online learning with partial information. In *Conference on Learning Theory*, 2018.
- [18] Gergely Neu. First-order regret bounds for combinatorial semi-bandits. In *Conference on Learning Theory*, 2015.
- [19] Roman Pogodin and Tor Lattimore. On first-order bounds, variance and gap-dependent bounds for adversarial bandits. In *Uncertainty in Artificial Intelligence Conference*, volume 115 of *Proceedings of Machine Learning Research*. PMLR, 2020.
- [20] Alexander Rakhlin and Karthik Sridharan. Online learning with predictable sequences. In *Conference on Learning Theory*, 2013.
- [21] Jacob Steinhardt and Percy Liang. Adaptivity and optimism: An improved exponentiated gradient algorithm. In *International Conference on Machine Learning*, 2014.

- [22] Vasilis Syrgkanis, Alekh Agarwal, Haipeng Luo, and Robert E. Schapire. Fast convergence of regularized learning in games. In *Advances in Neural Information Processing Systems*, volume 28, 2015.
- [23] Dirk van der Hoeven, Ashok Cutkosky, and Haipeng Luo. Comparator-adaptive convex bandits. In *Advances in Neural Information Processing Systems*, volume 33, 2020.
- [24] Chen-Yu Wei and Haipeng Luo. More adaptive algorithms for adversarial bandits. In *Conference on Learning Theory*, 2018.
- [25] Hang Yu, Yu-Hu Yan, and Peng Zhao. Improved dimension dependence for bandit convex optimization with gradient variation. In *International Conference on Machine Learning*, 2026.
- [26] Julian Zimmert and Yevgeny Seldin. Tsallis-INF: An optimal algorithm for stochastic and adversarial bandits. *Journal of Machine Learning Research*, 22(28), 2021.

A Omitted Details in Section 4

In this section, we show the omitted details in Section 4. Specifically, we provide the proof of (4.17), which is the key to control the regret in each phase j . For notational convenience, since we will consider each phase index j separately, in the proofs in this section, we will fix a deterministic phase label $j \in \{0, \dots, T-1\}$ and abbreviate

$$\begin{aligned} \chi_t &= \chi_{j,t}, & \rho &= \rho_j, & \eta &= \eta_j, & \alpha &= \alpha_j, & R &= R_j, & P &= P_j, \\ s &= s_j, & \tau &= \max\{u \in [T] : J_u = j\}, & \bar{\chi}_t &= \mathbf{1}\{J_{t-1} = j\}, & \delta_t &= \bar{\chi}_t - \chi_t. \end{aligned}$$

Here $\tau = 0$ if phase j is not reached. We also recall that $\phi_{2,t,i} = \varphi_2(\ell_{t,i})$, $b_t = p_t - x_t$, $\alpha = 8\eta$, and $\eta \leq 10^{-5}$. For the theorem statement, we still keep the index j . The following lemma shows the bound on (4.17).

Lemma A.1. *For every deterministic phase label $j \in \{0, \dots, T-1\}$, Algorithm 2 satisfies*

$$\alpha_j \mathbb{E} \left[\sum_{t=2}^T \chi_{j,t} (\varphi_2(\ell_{t,I_{t-1}}) - \varphi_2(\ell_{t,I_t})) \right] \leq 2000\eta_j^2 R_j + \mathcal{O}(\eta_j^2 P_j + \rho_j). \quad (\text{A.1})$$

To prove this, following the analysis in Section 3, we need to first bound the summation of $\langle \phi_{2,t}, b_{t-1} - b_t \rangle$ within each phase. The following lemma shows that this term is bounded by

Lemma A.2. *For every deterministic phase label $j \in \{0, \dots, T-1\}$, Algorithm 2 satisfies*

$$\alpha_j \mathbb{E} \left[\sum_{t=2}^T \chi_{j,t} \langle \phi_{2,t}, b_{t-1} - b_t \rangle \right] \leq 1700\eta_j^2 R_j + 2500\eta_j^2 P_j + 20\rho_j. \quad (\text{A.2})$$

Proof. If phase j is not reached, the left-hand side of (A.2) is zero. On the event that it is reached, s and τ are its first and last rounds. Since $b_s = 0$, the left-hand side can be written as

$$\sum_{t=2}^T \chi_t \langle \phi_{2,t}, b_{t-1} - b_t \rangle = -\langle \phi_{2,\tau}, b_\tau \rangle + \sum_{t=2}^{T-1} \chi_{t+1} \langle \phi_{2,t+1} - \phi_{2,t}, b_t \rangle.$$

Since $0 \leq \varphi_2(z) \leq 3/2$ on $[0, 1]$, we have $\|\phi_{2,\tau}\|_\infty \leq 3/2$. Moreover, the sampling rule gives $\|b_\tau\|_1 \leq 4\alpha$. Since phase j is reached with probability ρ , we have

$$\begin{aligned} & \alpha \mathbb{E} \left[\sum_{t=2}^T \chi_t \langle \phi_{2,t}, b_{t-1} - b_t \rangle \right] \\ & \leq 6\alpha^2 \rho + \alpha \mathbb{E} \left[\sum_{t=2}^{T-1} (\chi_{t+1} - \chi_t) \langle \phi_{2,t+1} - \phi_{2,t}, b_t \rangle \right] + \alpha \left| \sum_{t=2}^{T-1} \sum_{i=1}^K (\phi_{2,t+1,i} - \phi_{2,t,i}) \mathbb{E}[\chi_t b_{t,i}] \right| \\ & \leq 12\alpha^2 \rho + \alpha \left| \sum_{t=2}^{T-1} \sum_{i=1}^K (\phi_{2,t+1,i} - \phi_{2,t,i}) \mathbb{E}[\chi_t b_{t,i}] \right|, \end{aligned}$$

where the last inequality follows from the fact that $|\langle \phi_{2,t+1} - \phi_{2,t}, b_t \rangle| \leq \|\phi_{2,t+1} - \phi_{2,t}\|_\infty \|b_t\|_1 \leq \frac{3}{2} \cdot 4\alpha = 6\alpha$ and

$$\begin{aligned} \left| \mathbb{E} \left[\sum_{t=2}^{T-1} (\chi_{t+1} - \chi_t) \langle \phi_{2,t+1} - \phi_{2,t}, b_t \rangle \right] \right| &= \left| \mathbb{E} \left[\sum_{t=2}^{T-1} \chi_t (1 - \chi_{t+1}) \langle \phi_{2,t+1} - \phi_{2,t}, b_t \rangle \right] \right| \\ &\leq 6\alpha \mathbb{E} \left[\sum_{t=2}^{T-1} \chi_t (1 - \chi_{t+1}) \right] \leq 6\alpha \rho, \end{aligned}$$

where the equality uses $b_s = 0$, and the last inequality follows from $\sum_{t=2}^{T-1} \chi_t(1 - \chi_{t+1}) \leq \mathbf{1}\{s \leq T\}$. Further applying Cauchy–Schwarz inequality gives

$$\begin{aligned} & \alpha \left| \sum_{t=2}^{T-1} \sum_{i=1}^K (\phi_{2,t+1,i} - \phi_{2,t,i}) \mathbb{E}[\chi_t b_{t,i}] \right| \\ & \leq \alpha \left(\sum_{t=2}^{T-1} \sum_{i=1}^K \frac{(\mathbb{E}[\chi_t b_{t,i}])^2}{\mathbb{E}[\chi_t(\mathbf{1}\{I_t = i\} + x_{t,i})]} \right)^{1/2} \left(\sum_{t=2}^{T-1} \sum_{i=1}^K \mathbb{E}[\chi_t(\mathbf{1}\{I_t = i\} + x_{t,i})] (\phi_{2,t+1,i} - \phi_{2,t,i})^2 \right)^{1/2} \\ & \leq 2 \sum_{t=2}^{T-1} \sum_{i=1}^K \frac{(\mathbb{E}[\chi_t b_{t,i}])^2}{\mathbb{E}[\chi_t(\mathbf{1}\{I_t = i\} + x_{t,i})]} + \frac{\alpha^2}{8} \sum_{t=2}^{T-1} \sum_{i=1}^K \mathbb{E}[\chi_t(\mathbf{1}\{I_t = i\} + x_{t,i})] (\phi_{2,t+1,i} - \phi_{2,t,i})^2. \end{aligned}$$

Because χ_t is measurable before I_t is drawn, we have $\mathbb{E}[\chi_t b_{t,i}] = \mathbb{E}[\chi_t(\mathbf{1}\{I_t = i\} - x_{t,i})]$. Therefore, using Lemma A.6, the first term is bounded as

$$2 \sum_{t=2}^{T-1} \sum_{i=1}^K \frac{(\mathbb{E}[\chi_t b_{t,i}])^2}{\mathbb{E}[\chi_t(\mathbf{1}\{I_t = i\} + x_{t,i})]} = 2 \sum_{t=2}^{T-1} \sum_{i=1}^K \frac{(\mathbb{E}[\chi_t(\mathbf{1}\{I_t = i\} - x_{t,i})])^2}{\mathbb{E}[\chi_t(\mathbf{1}\{I_t = i\} + x_{t,i})]} \leq 1030\eta^2 R + 18\alpha^2 P + 680\alpha\rho.$$

Lemma A.7 further bounds the second term as follows:

$$\frac{\alpha^2}{8} \sum_{t=2}^{T-1} \sum_{i=1}^K \mathbb{E}[\chi_t(\mathbf{1}\{I_t = i\} + x_{t,i})] (\phi_{2,t+1,i} - \phi_{2,t,i})^2 \leq \frac{\alpha^2}{8} (12360\eta^2 R + 13P + 200\rho).$$

Combining these two bounds, we have

$$\begin{aligned} \alpha \mathbb{E} \left[\sum_{t=2}^T \chi_t \langle \phi_{2,t}, b_{t-1} - b_t \rangle \right] & \leq 12\alpha^2 \rho + 2(515\eta^2 R + 9\alpha^2 P + 340\alpha\rho) + \frac{\alpha^2}{8} (12360\eta^2 R + 13P + 200\rho) \\ & \leq 1700\eta^2 R + 2500\eta^2 P + 20\rho, \end{aligned}$$

where the last inequality uses $\alpha = 8\eta$ and $\eta \leq 10^{-5}$. $\square$

Next, we control the summation of $\langle \phi_{2,t}, x_{t-1} - x_t \rangle$ within the phase.

Lemma A.3. *For every deterministic phase label $j \in \{0, \dots, T-1\}$, Algorithm 2 satisfies*

$$\alpha_j \mathbb{E} \left[\sum_{t=2}^T \chi_{j,t} \langle \phi_{2,t}, x_{t-1} - x_t \rangle \right] \leq 100\eta_j^2 R_j + 20\eta_j^2 P_j + 4\rho_j. \quad (\text{A.3})$$

Proof. If phase j is not reached, the left-hand side of (A.3) is zero. On $\{\bar{\chi}_t = 1\}$, let $\bar{x}_t$ be the OMD update computed at the end of round $t-1$, before a possible phase reset; when $\bar{\chi}_t = 0$, set $\bar{x}_t = x_{t-1}$. Thus $\bar{x}_t = x_t$ whenever $\chi_t = 1$. Define

$$S_x \triangleq \mathbb{E} \left[\sum_{t=2}^T \chi_t \langle \phi_{2,t}, x_{t-1} - x_t \rangle \right]$$

and

$$\bar{S}_x \triangleq \mathbb{E} \left[\sum_{t=2}^T \bar{\chi}_t \langle \phi_{2,t}, x_{t-1} - \bar{x}_t \rangle \right].$$

Since $\bar{\chi}_t = \chi_t + \delta_t$ and $\bar{x}_t = x_t$ on $\{\chi_t = 1\}$, their exact difference is

$$S_x - \bar{S}_x = -\mathbb{E} \left[\sum_{t=2}^T \delta_t \langle \phi_{2,t}, x_{t-1} - \bar{x}_t \rangle \right].$$

Using $\|\phi_{2,t}\|_\infty \leq 3/2$, $\mathbb{E}[\sum_{t=2}^T \delta_t] \leq \rho$, we have

$$|S_x - \bar{S}_x| \leq \mathbb{E} \left[\sum_{t=2}^T \delta_t \|\phi_{2,t}\|_\infty \|x_{t-1} - \bar{x}_t\|_1 \right] \leq 3 \mathbb{E} \left[\sum_{t=2}^T \delta_t \right] \leq 3\rho.$$

Following the decomposition (3.33)–(3.34), write $g_t = \phi_{2,t} - \phi_{2,t-1}$ and define

$$\bar{S}_{x,1} \triangleq \mathbb{E} \left[\sum_{t=2}^T \bar{\chi}_t \langle \phi_{2,t-1}, x_{t-1} - \bar{x}_t \rangle \right]$$

and

$$\bar{S}_{x,2} \triangleq \mathbb{E} \left[\sum_{t=2}^T \bar{\chi}_t \langle g_t, x_{t-1} - \bar{x}_t \rangle \right].$$

We have $\bar{S}_x = \bar{S}_{x,1} + \bar{S}_{x,2}$.

We first control $\bar{S}_{x,1}$ by making the comparison with Section 3 explicit. Equation (3.20) is unchanged. In the passage from (3.26) to (3.27), replacing the 1-Lipschitz function φ by the 2-Lipschitz function φ_2 doubles the coefficient 2.2 to 4.4. Equation (3.31) is unchanged. The phase-sampler bound $p_{t-1,i} \geq x_{t-1,i}/(1+2\alpha)$ and $\eta \leq 10^{-5}$ also preserve the 0.01 coordinatewise stability used in (3.23). Therefore the final comparison used to prove (3.13) becomes

$$\mathbb{E}_{t-1}[\langle \phi_{2,t-1}, x_{t-1} - \bar{x}_t \rangle] \leq \frac{4.4}{0.94} \mathbb{E}_{t-1}[D_{t-1} + D_\Psi(\bar{x}_t, x_{t-1})] \leq 16 \mathbb{E}_{t-1}[D_{t-1}].$$

Here the second inequality uses $D_\Psi(\bar{x}_t, x_{t-1}) \leq 2D_{t-1}$, as shown in the last step of the proof of (3.13). Further, since $\bar{\chi}_t = \mathbf{1}\{J_{t-1} = j\}$ is $\mathcal{F}_{t-2}$ -measurable, the tower property and (4.18) give

$$\begin{aligned} \bar{S}_{x,1} &= \mathbb{E} \left[\sum_{t=2}^T \bar{\chi}_t \mathbb{E}_{t-1} [\langle \phi_{2,t-1}, x_{t-1} - \bar{x}_t \rangle] \right] && (\bar{\chi}_t \text{ is } \mathcal{F}_{t-2}\text{-measurable}) \\ &\leq 16 \mathbb{E} \left[\sum_{t=2}^T \bar{\chi}_t \mathbb{E}_{t-1}[D_{t-1}] \right] \\ &\leq 9.6\eta \mathbb{E} \left[\sum_{t=2}^T \bar{\chi}_t r_{t-1}^2 \right] && (\text{using (4.18)}) \\ &\leq 9.6\eta \mathbb{E} \left[\mathbf{1}\{s \leq T\} + \sum_{t=2}^T \chi_t r_t^2 \right] \\ &= 9.6\eta(R + \rho). \end{aligned}$$

We next control $\bar{S}_{x,2}$. Set $q_t \triangleq \langle x_{t-1}, g_t^{\odot 2} \rangle$. Applying (3.12) on round $t-1$ with $z = g_t$, and then applying (4.18), gives

$$|\langle g_t, x_{t-1} - \bar{x}_t \rangle| \leq 1.5\sqrt{\eta D_{t-1}}\sqrt{q_t} \leq 1.5\sqrt{0.6\eta}|r_{t-1}|\sqrt{q_t} \leq 1.2\eta|r_{t-1}|\sqrt{q_t}.$$

Therefore, we have

$$\begin{aligned} |\bar{S}_{x,2}| &\leq 1.2\eta \mathbb{E} \left[\sum_{t=2}^T \bar{\chi}_t |r_{t-1}|\sqrt{q_t} \right] \\ &\leq 1.2\eta \sqrt{\mathbb{E} \left[\sum_{t=2}^T \bar{\chi}_t r_{t-1}^2 \right] \mathbb{E} \left[\sum_{t=2}^T \bar{\chi}_t q_t \right]} && (\text{Cauchy-Schwarz}) \\ &\leq 1.2\eta \sqrt{(R + \rho) \mathbb{E} \left[\sum_{t=2}^T \bar{\chi}_t q_t \right]}. && (\text{A.4}) \end{aligned}$$

The 2-Lipschitz property of φ_2 gives $|g_{t,i}| = |\varphi_2(\ell_{t,i}) - \varphi_2(\ell_{t-1,i})| \leq 2|\ell_{t,i} - \ell_{t-1,i}|$ and $g_{t,i}^2 \leq 4(\ell_{t,i} - \ell_{t-1,i})^2$. Together with $x_{t-1,i} \leq (1 + 2\alpha)p_{t-1,i}$, this yields

$$q_t = \langle x_{t-1}, g_t^{\odot 2} \rangle = \sum_{i=1}^K x_{t-1,i} g_{t,i}^2 \leq 4 \sum_{i=1}^K x_{t-1,i} (\ell_{t,i} - \ell_{t-1,i})^2 \leq 4(1 + 2\alpha) \sum_{i=1}^K p_{t-1,i} (\ell_{t,i} - \ell_{t-1,i})^2.$$

Since $\bar{\chi}_t$ is $\mathcal{F}_{t-2}$ -measurable, we have

$$\begin{aligned} \mathbb{E} \left[\sum_{t=2}^T \bar{\chi}_t q_t \right] &\leq 4(1 + 2\alpha) \mathbb{E} \left[\sum_{t=2}^T \bar{\chi}_t \sum_{i=1}^K p_{t-1,i} (\ell_{t,i} - \ell_{t-1,i})^2 \right] \\ &= 4(1 + 2\alpha) \mathbb{E} \left[\sum_{t=2}^T \bar{\chi}_t \mathbb{E}_{t-1} [(\ell_{t,I_{t-1}} - \ell_{t-1,I_{t-1}})^2] \right] \\ &= 4(1 + 2\alpha) \mathbb{E} \left[\sum_{t=2}^T \bar{\chi}_t (\ell_{t,I_{t-1}} - \ell_{t-1,I_{t-1}})^2 \right] \quad (\bar{\chi}_t \text{ is } \mathcal{F}_{t-2}\text{-measurable}) \\ &= 4(1 + 2\alpha) \left(P + \mathbb{E} \left[\sum_{t=2}^T \delta_t (\ell_{t,I_{t-1}} - \ell_{t-1,I_{t-1}})^2 \right] \right) \\ &\leq 4(1 + 2\alpha)(P + \rho) \\ &\leq 5(P + \rho). \end{aligned} \tag{A.5}$$

Combining the above inequalities together, we obtain that

$$S_x \leq \bar{S}_x + 3\rho = \bar{S}_{x,1} + \bar{S}_{x,2} + 3\rho \leq 9.6\eta(R + \rho) + 1.2\eta\sqrt{5(R + \rho)(P + \rho)} + 3\rho. \tag{A.6}$$

Since $\alpha = 8\eta$, the AM-GM inequality gives

$$9.6\sqrt{5}\eta^2\sqrt{(R + \rho)(P + \rho)} \leq 4.8\sqrt{5}\eta^2(R + P + 2\rho) \leq 10.8\eta^2(R + P + 2\rho).$$

Further using the fact that $\eta \leq 10^{-5}$ gives

$$\alpha S_x \leq 76.8\eta^2(R + \rho) + 10.8\eta^2(R + P + 2\rho) + 24\eta\rho \leq 100\eta^2R + 20\eta^2P + 4\rho,$$

□

Now we are ready to prove Lemma A.1.

Proof of Lemma A.1. By the definitions of $\bar{\chi}_t$ and δ_t , $\bar{\chi}_t = \chi_t + \delta_t$. The indicator δ_t marks the unique round immediately after phase j ends. Therefore we have $\mathbb{E} \left[\sum_{t=2}^T \delta_t \right] \leq \rho$.

Next, as $\bar{\chi}_t$ is measurable before I_{t-1} is drawn, but χ_t is not, the two conditional-expectation identities are $\mathbb{E}[\bar{\chi}_t \varphi_2(\ell_{t,I_{t-1}})] = \mathbb{E}[\bar{\chi}_t \langle \phi_{2,t}, p_{t-1} \rangle]$ and $\mathbb{E}[\chi_t \varphi_2(\ell_{t,I_t})] = \mathbb{E}[\chi_t \langle \phi_{2,t}, p_t \rangle]$. Further using $\bar{\chi}_t = \chi_t + \delta_t$ gives

$$\alpha \mathbb{E} \left[\sum_{t=2}^T \chi_t (\varphi_2(\ell_{t,I_{t-1}}) - \varphi_2(\ell_{t,I_t})) \right] = \alpha \mathbb{E} \left[\sum_{t=2}^T \chi_t \langle \phi_{2,t}, p_{t-1} - p_t \rangle \right] + \alpha \mathbb{E} \left[\sum_{t=2}^T \delta_t (\langle \phi_{2,t}, p_{t-1} \rangle - \varphi_2(\ell_{t,I_{t-1}})) \right]. \tag{A.7}$$

Thus (A.7) is the stopped counterpart of (3.10): the second term is precisely the boundary residual created when the predictable indicator $\bar{\chi}_t$ is replaced by χ_t . Since $0 \leq \varphi_2(z) \leq 3/2$ on $[0, 1]$, the absolute value of its summand is at most $3\delta_t/2$, and $\mathbb{E}[\sum_{t=2}^T \delta_t] \leq \rho$ bounds the entire residual by $3\alpha\rho/2$. Substituting $p_t = x_t + b_t$ therefore gives

$$\alpha \mathbb{E} \left[\sum_{t=2}^T \chi_t (\varphi_2(\ell_{t,I_{t-1}}) - \varphi_2(\ell_{t,I_t})) \right] \leq \alpha \mathbb{E} \left[\sum_{t=2}^T \chi_t \langle \phi_{2,t}, x_{t-1} - x_t \rangle \right] + \alpha \mathbb{E} \left[\sum_{t=2}^T \chi_t \langle \phi_{2,t}, b_{t-1} - b_t \rangle \right] + \frac{3}{2}\alpha\rho. \tag{A.8}$$

Lemma A.3 bounds the first term on the right-hand side of (A.8), and Lemma A.2 bounds the second. Their sum is at most $1800\eta^2R + 2520\eta^2P + (24 + 3\alpha/2)\rho$, which is bounded by the right-hand side of (A.1). This proves the lemma. □

A.1 Auxiliary Lemmas

The following auxiliary lemmas are used in the proof of Lemma A.1. Throughout this subsection, $\mathbb{E}$ denotes expectation with respect to all randomness generated by Algorithm 2, with the oblivious loss sequence fixed.

The first lemma records the properties of the ratio functional used in the proof of Lemma A.2.

Lemma A.4. *For any jointly distributed random arm $I \in [K]$, random probability vector $x \in \Delta_K$, and random weight $w \in [0, 1]$, define*

$$\mathcal{N}(w; I, x) \triangleq \sum_{i=1}^K \frac{(\mathbb{E}[w(\mathbf{1}\{I = i\} - x_i)])^2}{\mathbb{E}[w(\mathbf{1}\{I = i\} + x_i)]}. \quad (\text{A.9})$$

If $0 \leq w' \leq w \leq 1$, then we have $\mathcal{N}(w'; I, x) \leq \mathcal{N}(w; I, x) + 6\mathbb{E}[w - w']$ and $\mathcal{N}(w; I, x) \leq 2\mathbb{E}[w]$. Moreover, whenever $w_1 + w_2 \leq 1$, we have $\mathcal{N}(w_1 + w_2; I, x) \leq \mathcal{N}(w_1; I, x) + \mathcal{N}(w_2; I, x)$.

Proof. For every nonnegative random variable u , define $\mu_i(u) \triangleq \mathbb{E}[u(\mathbf{1}\{I = i\} - x_i)]$ and $\nu_i(u) \triangleq \mathbb{E}[u(\mathbf{1}\{I = i\} + x_i)]$. The pointwise inequality $|\mathbf{1}\{I = i\} - x_i| \leq \mathbf{1}\{I = i\} + x_i$ gives

$$|\mu_i(u)| \leq \mathbb{E}[u|\mathbf{1}\{I = i\} - x_i|] \leq \nu_i(u).$$

Direct calculation shows that $\mu_i(w) = \mu_i(w') + \mu_i(w - w')$ and $\nu_i(w) = \nu_i(w') + \nu_i(w - w')$. If $\nu_i(w') = 0$, then $\mu_i(w') = 0$ and for any $w \geq w'$

$$\frac{\mu_i(w')^2}{\nu_i(w')} - \frac{\mu_i(w)^2}{\nu_i(w)} = -\frac{\mu_i(w)^2}{\nu_i(w)} \leq 0 \leq 3\nu_i(w - w').$$

If $\nu_i(w') > 0$, then we have

$$\begin{aligned} \frac{\mu_i(w')^2}{\nu_i(w')} - \frac{\mu_i(w)^2}{\nu_i(w)} &= \frac{\nu_i(w - w')\mu_i(w')^2 - 2\nu_i(w')\mu_i(w')\mu_i(w - w') - \nu_i(w')\mu_i(w - w')^2}{\nu_i(w')\nu_i(w)} \\ &\leq \frac{\nu_i(w - w')|\mu_i(w')|^2 + 2\nu_i(w')|\mu_i(w')||\mu_i(w - w')|}{\nu_i(w')\nu_i(w)} \\ &\leq \frac{3\nu_i(w')^2\nu_i(w - w')}{\nu_i(w')\nu_i(w)} \leq \frac{3\nu_i(w')^2\nu_i(w - w')}{\nu_i(w')^2} = 3\nu_i(w - w'). \end{aligned}$$

Summing over i gives $\mathcal{N}(w'; I, x) - \mathcal{N}(w; I, x) \leq 3 \sum_{i=1}^K \nu_i(w - w') = 6\mathbb{E}[w - w']$. The second inequality follows from

$$\mathcal{N}(w; I, x) = \sum_{i=1}^K \frac{\mu_i(w)^2}{\nu_i(w)} \leq \sum_{i=1}^K \nu_i(w) = 2\mathbb{E}[w],$$

where we use the fact that $x \in \Delta_K$. Finally, Cauchy-Schwarz inequality leads to the following

$$\mathcal{N}(w_1 + w_2; I, x) = \sum_{i=1}^K \frac{(\mu_i(w_1) + \mu_i(w_2))^2}{\nu_i(w_1) + \nu_i(w_2)} \leq \sum_{i=1}^K \left(\frac{\mu_i(w_1)^2}{\nu_i(w_1)} + \frac{\mu_i(w_2)^2}{\nu_i(w_2)} \right) = \mathcal{N}(w_1; I, x) + \mathcal{N}(w_2; I, x).$$

□

Define $\mathcal{N}_t^{\text{cur}} \triangleq \mathcal{N}(\chi_t; I_t, x_t)$ and $\mathcal{N}_t^{\text{prev}} \triangleq \mathcal{N}(\chi_t; I_{t-1}, x_t)$. Since χ_t is $\mathcal{F}_{t-1}$ -measurable, we have

$$\mathbb{E}[\chi_t(\mathbf{1}\{I_t = i\} - x_{t,i})] = \mathbb{E}[\chi_t \mathbb{E}_t[\mathbf{1}\{I_t = i\} - x_{t,i}]] = \mathbb{E}[\chi_t(p_{t,i} - x_{t,i})] = \mathbb{E}[\chi_t b_{t,i}].$$

The following lemma bounds the current squared mean norm in terms of its previous-arm counterpart.

Lemma A.5. *For every fixed round $t \in \{2, \dots, T\}$,*

$$\mathcal{N}_t^{\text{cur}} \leq 13\alpha \mathcal{N}_t^{\text{prev}} + 8\alpha^2 \mathbb{E}[\chi_t(r_t^2 + v_t^2)]. \quad (\text{A.10})$$

Proof. On the event $\{\chi_t = 1\}$, the sampling rule in Algorithm 2 gives

$$\begin{aligned}\lambda_t &= \alpha(2 - c_{t-1}) = \alpha(2 - \ell_{t-1, I_{t-1}}), \\ p_{t,i} &= \frac{x_{t,i} + \lambda_t \mathbf{1}\{I_{t-1} = i\}}{1 + \lambda_t} = x_{t,i} + \frac{\alpha(2 - \ell_{t-1, I_{t-1}})}{1 + \alpha(2 - \ell_{t-1, I_{t-1}})} (\mathbf{1}\{I_{t-1} = i\} - x_{t,i}).\end{aligned}$$

Define $\gamma_{t,i} \triangleq \alpha(2 - \ell_{t-1,i})/(1 + \alpha(2 - \ell_{t-1,i}))$. Thus,

$$\begin{aligned}p_{t,i} - x_{t,i} &= \gamma_{t, I_{t-1}} (\mathbf{1}\{I_{t-1} = i\} - x_{t,i}), \\ p_{t,i} + x_{t,i} &= (2 - \gamma_{t, I_{t-1}})x_{t,i} + \gamma_{t, I_{t-1}} \mathbf{1}\{I_{t-1} = i\}.\end{aligned}$$

Since χ_t is measurable before I_t is drawn, we have

$$\begin{aligned}\mathbb{E}[\chi_t (\mathbf{1}\{I_t = i\} - x_{t,i})] &= \mathbb{E}[\chi_t (p_{t,i} - x_{t,i})] = \mathbb{E}[\chi_t \gamma_{t, I_{t-1}} (\mathbf{1}\{I_{t-1} = i\} - x_{t,i})], \\ \mathbb{E}[\chi_t (\mathbf{1}\{I_t = i\} + x_{t,i})] &= \mathbb{E}[\chi_t (p_{t,i} + x_{t,i})] = \mathbb{E}[\chi_t ((2 - \gamma_{t, I_{t-1}})x_{t,i} + \gamma_{t, I_{t-1}} \mathbf{1}\{I_{t-1} = i\})].\end{aligned}$$

Therefore, by the definition of $\mathcal{N}_t^{\text{cur}}$,

$$\mathcal{N}_t^{\text{cur}} = \sum_{i=1}^K \frac{(\mathbb{E}[\chi_t \gamma_{t, I_{t-1}} (\mathbf{1}\{I_{t-1} = i\} - x_{t,i})])^2}{\mathbb{E}[\chi_t ((2 - \gamma_{t, I_{t-1}})x_{t,i} + \gamma_{t, I_{t-1}} \mathbf{1}\{I_{t-1} = i\})]}.$$

We then decompose $\mathcal{N}_t^{\text{cur}}$ using $(u + v)^2 \leq 2u^2 + 2v^2$:

$$\mathcal{N}_t^{\text{cur}} \leq \underbrace{2 \sum_{i=1}^K \frac{\gamma_{t,i}^2 (\mathbb{E}[\chi_t (\mathbf{1}\{I_{t-1} = i\} - x_{t,i})])^2}{\mathbb{E}[\chi_t ((2 - \gamma_{t, I_{t-1}})x_{t,i} + \gamma_{t, I_{t-1}} \mathbf{1}\{I_{t-1} = i\})]}}_{\text{previous-arm contribution}} + \underbrace{2 \sum_{i=1}^K \frac{(\mathbb{E}[\chi_t x_{t,i} (\gamma_{t,i} - \gamma_{t, I_{t-1}})])^2}{\mathbb{E}[\chi_t ((2 - \gamma_{t, I_{t-1}})x_{t,i} + \gamma_{t, I_{t-1}} \mathbf{1}\{I_{t-1} = i\})]}}_{\text{remainder}}.$$

We first bound the previous-arm contribution. Since $\mathbb{E}[\chi_t ((2 - \gamma_{t, I_{t-1}})x_{t,i} + \gamma_{t, I_{t-1}} \mathbf{1}\{I_{t-1} = i\})] \geq \frac{\alpha}{1+\alpha} \mathbb{E}[\chi_t (\mathbf{1}\{I_{t-1} = i\} + x_{t,i})]$, and $\gamma_{t,i} \leq 2\alpha/(1 + 2\alpha)$, we have

$$2 \sum_{i=1}^K \frac{\gamma_{t,i}^2 (\mathbb{E}[\chi_t (\mathbf{1}\{I_{t-1} = i\} - x_{t,i})])^2}{\mathbb{E}[\chi_t ((2 - \gamma_{t, I_{t-1}})x_{t,i} + \gamma_{t, I_{t-1}} \mathbf{1}\{I_{t-1} = i\})]} \leq 2 \frac{(2\alpha/(1 + 2\alpha))^2}{\alpha/(1 + \alpha)} \mathcal{N}_t^{\text{prev}} \leq 12\alpha \mathcal{N}_t^{\text{prev}}.$$

It remains to control the remainder. Since $\mathbb{E}[\chi_t (\mathbf{1}\{I_t = i\} + x_{t,i})] \geq \mathbb{E}[\chi_t x_{t,i}]$, Cauchy-Schwarz inequality gives

$$\begin{aligned}2 \sum_{i=1}^K \frac{(\mathbb{E}[\chi_t x_{t,i} (\gamma_{t,i} - \gamma_{t, I_{t-1}})])^2}{\mathbb{E}[\chi_t (\mathbf{1}\{I_t = i\} + x_{t,i})]} &\leq 2 \sum_{i=1}^K \frac{(\mathbb{E}[\chi_t x_{t,i} (\gamma_{t,i} - \gamma_{t, I_{t-1}})])^2}{\mathbb{E}[\chi_t x_{t,i}]} \\ &\leq 2 \mathbb{E} \left[\chi_t \sum_{i=1}^K x_{t,i} (\gamma_{t,i} - \gamma_{t, I_{t-1}})^2 \right] \\ &\leq 2\alpha^2 \mathbb{E} \left[\chi_t \sum_{i=1}^K x_{t,i} (\ell_{t-1,i} - \ell_{t-1, I_{t-1}})^2 \right].\end{aligned}$$

Combining the two parts yields

$$\mathcal{N}_t^{\text{cur}} \leq 12\alpha \mathcal{N}_t^{\text{prev}} + 2\alpha^2 \mathbb{E} \left[\chi_t \sum_{i=1}^K x_{t,i} (\ell_{t-1,i} - \ell_{t-1, I_{t-1}})^2 \right]. \quad (\text{A.11})$$

To bound the remaining loss discrepancy, conditionally on $\mathcal{F}_{t-1}$ draw $I'_t \sim x_t$ and realize the sampling rule by setting $I_t = I_{t-1}$ with probability $\gamma_{t, I_{t-1}}$ and $I_t = I'_t$ otherwise. Since $\gamma_{t, I_{t-1}} \leq 2\alpha$, we have

$$\mathbb{E}[\chi_t r_t^2] = \mathbb{E}[\chi_t \gamma_{t, I_{t-1}} v_t^2 + \chi_t (1 - \gamma_{t, I_{t-1}}) (\ell_{t, I'_t} - \ell_{t-1, I_{t-1}})^2] \geq (1 - 2\alpha) \mathbb{E}[\chi_t (\ell_{t, I'_t} - \ell_{t-1, I_{t-1}})^2].$$

Applying Proposition A.8 with $A = \mathbb{E}[\chi_t x_{t,i}]$ and $B = \mathbb{E}[\chi_t \mathbf{1}\{I_{t-1} = i\}]$, multiplying the resulting inequality by $(\ell_{t,i} - \ell_{t-1,i})^2$, and summing over i give

$$\begin{aligned} \mathbb{E}[\chi_t (\ell_{t,I'_t} - \ell_{t-1,I'_t})^2] &= \sum_{i=1}^K \mathbb{E}[\chi_t x_{t,i} (\ell_{t,i} - \ell_{t-1,i})^2] \\ &\leq \sum_{i=1}^K \left(2\mathbb{E}[\chi_t \mathbf{1}\{I_{t-1} = i\}] + 6 \frac{(\mathbb{E}[\chi_t (x_{t,i} - \mathbf{1}\{I_{t-1} = i\}])^2]}{\mathbb{E}[\chi_t (x_{t,i} + \mathbf{1}\{I_{t-1} = i\}])} \right) (\ell_{t,i} - \ell_{t-1,i})^2 \\ &\leq 2 \sum_{i=1}^K \mathbb{E}[\chi_t \mathbf{1}\{I_{t-1} = i\}] (\ell_{t,i} - \ell_{t-1,i})^2 + 6 \sum_{i=1}^K \frac{(\mathbb{E}[\chi_t (\mathbf{1}\{I_{t-1} = i\} - x_{t,i})])^2}{\mathbb{E}[\chi_t (\mathbf{1}\{I_{t-1} = i\} + x_{t,i})]} \\ &= 2\mathbb{E}[\chi_t v_t^2] + 6\mathcal{N}_t^{\text{prev}}, \end{aligned}$$

where the second inequality uses $(\ell_{t,i} - \ell_{t-1,i})^2 \leq 1$. Consequently, we can obtain that

$$\begin{aligned} \mathbb{E} \left[\chi_t \sum_{i=1}^K x_{t,i} (\ell_{t-1,i} - \ell_{t-1,I_{t-1}})^2 \right] &= \mathbb{E}[\chi_t (\ell_{t-1,I'_t} - \ell_{t-1,I_{t-1}})^2] \\ &\leq 2\mathbb{E}[\chi_t (\ell_{t,I'_t} - \ell_{t-1,I_{t-1}})^2] + 2\mathbb{E}[\chi_t (\ell_{t,I'_t} - \ell_{t-1,I'_t})^2] \\ &\leq 3\mathbb{E}[\chi_t r_t^2] + 4\mathbb{E}[\chi_t v_t^2] + 12\mathcal{N}_t^{\text{prev}}. \end{aligned} \quad (\text{A.12})$$

Here the last inequality uses $2/(1-2\alpha) \leq 3$. Substituting (A.12) into (A.11) gives

$$\mathcal{N}_t^{\text{cur}} \leq (12\alpha + 24\alpha^2)\mathcal{N}_t^{\text{prev}} + 6\alpha^2\mathbb{E}[\chi_t r_t^2] + 8\alpha^2\mathbb{E}[\chi_t v_t^2] \leq 13\alpha\mathcal{N}_t^{\text{prev}} + 8\alpha^2\mathbb{E}[\chi_t (r_t^2 + v_t^2)],$$

where the last inequality uses $24\alpha \leq 1$. $\square$

The next lemma propagates the squared mean norm by one round and sums the result over the phase.

Lemma A.6. *Algorithm 2 satisfies*

$$\sum_{t=2}^T \mathcal{N}_t^{\text{prev}} \leq 3 \sum_{t=2}^T \mathcal{N}_t^{\text{cur}} + 3\eta^2 R + 25\rho, \quad (\text{A.13})$$

$$\sum_{t=2}^T \mathcal{N}_t^{\text{cur}} \leq 515\eta^2 R + 9\alpha^2 P + 340\alpha\rho. \quad (\text{A.14})$$

Proof. By definition, $\chi_t = \mathbf{1}\{s < t \leq \tau\}$ if the phase is entered and otherwise, $\chi_t = 0$ for every t . Therefore, pathwise, we have $\sum_{t=1}^{T-1} (1 - \chi_t)\chi_{t+1} \leq \mathbf{1}\{s \leq T\}$ and $\sum_{t=1}^{T-1} \chi_t(1 - \chi_{t+1}) \leq \mathbf{1}\{s \leq T\}$.

Using $\chi_{t+1} = \chi_t \chi_{t+1} + (1 - \chi_t)\chi_{t+1}$ and $\chi_t \chi_{t+1} = \chi_t - \chi_t(1 - \chi_{t+1})$, Lemma A.4 gives that

$$\begin{aligned} \mathcal{N}(\chi_{t+1}; I_t, x_t) &\leq \mathcal{N}(\chi_t \chi_{t+1}; I_t, x_t) + \mathcal{N}((1 - \chi_t)\chi_{t+1}; I_t, x_t) \\ &\leq \mathcal{N}_t^{\text{cur}} + 6\mathbb{E}[\chi_t(1 - \chi_{t+1})] + 2\mathbb{E}[(1 - \chi_t)\chi_{t+1}]. \end{aligned} \quad (\text{A.15})$$

We next compare $\mathcal{N}_{t+1}^{\text{prev}} = \mathcal{N}(\chi_{t+1}; I_t, x_{t+1})$ with $\mathcal{N}(\chi_{t+1}; I_t, x_t)$. For every i ,

$$\mathbb{E}[\chi_{t+1}(\mathbf{1}\{I_t = i\} - x_{t+1,i})] = \mathbb{E}[\chi_{t+1}(\mathbf{1}\{I_t = i\} - x_{t,i})] - \mathbb{E}[\chi_{t+1}(x_{t+1,i} - x_{t,i})].$$

On $\{\chi_{t+1} = 1\}$, applying (3.17) on round t gives $x_{t+1,i} \geq 0.99x_{t,i}$ and hence

$$\mathbb{E}[\chi_{t+1}(\mathbf{1}\{I_t = i\} + x_{t+1,i})] \geq 0.99\mathbb{E}[\chi_{t+1}(\mathbf{1}\{I_t = i\} + x_{t,i})].$$

Therefore, applying AM-GM inequality and the above inequality give the following

$$\mathcal{N}_{t+1}^{\text{prev}} \leq \frac{2}{0.99}\mathcal{N}(\chi_{t+1}; I_t, x_t) + \frac{2}{0.99} \sum_{i=1}^K \frac{(\mathbb{E}[\chi_{t+1}(x_{t+1,i} - x_{t,i})])^2}{\mathbb{E}[\chi_{t+1}(\mathbf{1}\{I_t = i\} + x_{t,i})]}.$$

To further bound the second term, (3.19) applied on round t and Cauchy-Schwarz inequality yield

$$\begin{aligned}
\sum_{i=1}^K \frac{(\mathbb{E}[\chi_{t+1}(x_{t+1,i} - x_{t,i})])^2}{\mathbb{E}[\chi_{t+1}(\mathbf{1}\{I_t = i\} + x_{t,i})]} &\leq \sum_{i=1}^K \frac{(\mathbb{E}[\chi_{t+1}|x_{t+1,i} - x_{t,i}|])^2}{\mathbb{E}[\chi_{t+1}x_{t,i}]} \\
&\leq \sum_{i=1}^K \mathbb{E}\left[\chi_{t+1} \frac{(x_{t+1,i} - x_{t,i})^2}{x_{t,i}}\right] \quad (\text{using Cauchy-Schwarz inequality}) \\
&\leq \mathbb{E}\left[\chi_{t+1} \sum_{i=1}^K \frac{(x_{t+1,i} - x_{t,i})^2}{x_{t,i}^2}\right] \\
&\leq 2.1\eta\mathbb{E}[\chi_{t+1}D_t].
\end{aligned}$$

Combining the above two inequalities gives

$$\mathcal{N}_{t+1}^{\text{prev}} \leq 2.03\mathcal{N}(\chi_{t+1}; I_t, x_t) + 4.3\eta\mathbb{E}[\chi_{t+1}D_t]. \quad (\text{A.16})$$

Moreover, since $\sum_{t=1}^{T-1} \chi_{t+1}r_t^2 \leq \mathbf{1}\{s \leq T\} + \sum_{t=2}^T \chi_t r_t^2$, (4.18) applied on round t gives

$$\mathbb{E}\left[\sum_{t=1}^{T-1} \chi_{t+1}D_t\right] \leq 0.6\eta\mathbb{E}\left[\sum_{t=1}^{T-1} \chi_{t+1}r_t^2\right] \leq 0.6\eta(R + \rho).$$

Summing (A.15) and (A.16) over $t = 1, \dots, T-1$, using $\chi_1 = 0$ and the preceding bounds, yields

$$\begin{aligned}
\sum_{t=2}^T \mathcal{N}_t^{\text{prev}} &\leq 2.03 \sum_{t=1}^{T-1} \mathcal{N}(\chi_{t+1}; I_t, x_t) + 4.3\eta\mathbb{E}\left[\sum_{t=1}^{T-1} \chi_{t+1}D_t\right] \\
&\leq 2.03 \sum_{t=2}^T \mathcal{N}_t^{\text{cur}} + 12.18\mathbb{E}\left[\sum_{t=1}^{T-1} \chi_t(1 - \chi_{t+1})\right] + 4.06\mathbb{E}\left[\sum_{t=1}^{T-1} (1 - \chi_t)\chi_{t+1}\right] + 4.3\eta\mathbb{E}\left[\sum_{t=1}^{T-1} \chi_{t+1}D_t\right] \\
&\leq 2.03 \sum_{t=2}^T \mathcal{N}_t^{\text{cur}} + 16.24\rho + 2.58\eta^2(R + \rho) \\
&\leq 3 \sum_{t=2}^T \mathcal{N}_t^{\text{cur}} + 3\eta^2R + 25\rho,
\end{aligned}$$

which proves (A.13).

Finally, summing (A.10) over t and applying (A.13) give

$$\sum_{t=2}^T \mathcal{N}_t^{\text{cur}} \leq 13\alpha \sum_{t=2}^T \mathcal{N}_t^{\text{prev}} + 8\alpha^2(R + P) \leq 39\alpha \sum_{t=2}^T \mathcal{N}_t^{\text{cur}} + (39\alpha\eta^2 + 8\alpha^2)R + 8\alpha^2P + 325\alpha\rho.$$

Rearranging and using $\alpha = 8\eta$ and $\eta \leq 10^{-5}$ yield

$$\sum_{t=2}^T \mathcal{N}_t^{\text{cur}} \leq \frac{39\alpha\eta^2 + 8\alpha^2}{1 - 39\alpha}R + \frac{8\alpha^2}{1 - 39\alpha}P + \frac{325\alpha}{1 - 39\alpha}\rho \leq 515\eta^2R + 9\alpha^2P + 340\alpha\rho,$$

which proves (A.14). $\square$

The following lemma bounds the weighted sum of squared differences used after summation by parts.

Lemma A.7. *Let $g_{t+1,i} \triangleq \phi_{2,t+1,i} - \phi_{2,t,i}$ and $q_{t,i} \triangleq \mathbb{E}[\chi_t(\mathbf{1}\{I_t = i\} + x_{t,i})]$. Then*

$$\sum_{t=2}^{T-1} \sum_{i=1}^K q_{t,i} g_{t+1,i}^2 \leq 12360\eta^2R + 13P + 200\rho. \quad (\text{A.17})$$

Proof. The inequalities $\chi_t \leq \chi_{t+1} + \chi_t(1 - \chi_{t+1})$, $|g_{t+1,i}| \leq 3/2$, and $\sum_{i=1}^K (\mathbf{1}\{I_t = i\} + x_{t,i}) = 2$ give

$$\mathbb{E} \left[\sum_{t=2}^{T-1} \chi_t(1 - \chi_{t+1}) \sum_{i=1}^K (\mathbf{1}\{I_t = i\} + x_{t,i}) g_{t+1,i}^2 \right] \leq \frac{9}{2} \mathbb{E} \left[\sum_{t=2}^{T-1} \chi_t(1 - \chi_{t+1}) \right] \leq 7\rho.$$

By the definition of $q_{t,i}$ and $\chi_t \leq \chi_{t+1} + \chi_t(1 - \chi_{t+1})$,

$$\begin{aligned} \sum_{t=2}^{T-1} \sum_{i=1}^K q_{t,i} g_{t+1,i}^2 &= \sum_{t=2}^{T-1} \sum_{i=1}^K \mathbb{E}[\chi_t(\mathbf{1}\{I_t = i\} + x_{t,i})] g_{t+1,i}^2 \\ &\leq \sum_{t=2}^{T-1} \sum_{i=1}^K \mathbb{E}[\chi_{t+1}(\mathbf{1}\{I_t = i\} + x_{t,i})] g_{t+1,i}^2 + \mathbb{E} \left[\sum_{t=2}^{T-1} \chi_t(1 - \chi_{t+1}) \sum_{i=1}^K (\mathbf{1}\{I_t = i\} + x_{t,i}) g_{t+1,i}^2 \right] \\ &\leq 7\rho + \sum_{t=2}^{T-1} \sum_{i=1}^K \mathbb{E}[\chi_{t+1} \mathbf{1}\{I_t = i\}] g_{t+1,i}^2 + \sum_{t=2}^{T-1} \sum_{i=1}^K \mathbb{E}[\chi_{t+1} x_{t,i}] g_{t+1,i}^2, \end{aligned}$$

Since φ_2 is 2-Lipschitz, $g_{t+1,i}^2 \leq 4(\ell_{t+1,i} - \ell_{t,i})^2$. Hence, we have

$$\sum_{t=2}^{T-1} \sum_{i=1}^K \mathbb{E}[\chi_{t+1} \mathbf{1}\{I_t = i\}] g_{t+1,i}^2 \leq 4\mathbb{E} \left[\sum_{t=2}^{T-1} \chi_{t+1} (\ell_{t+1,I_t} - \ell_{t,I_t})^2 \right] = 4\mathbb{E} \left[\sum_{u=3}^T \chi_u (\ell_{u,I_{u-1}} - \ell_{u-1,I_{u-1}})^2 \right] \leq 4P.$$

Applying Proposition A.8 with $A = \mathbb{E}[\chi_{t+1} x_{t,i}]$ and $B = \mathbb{E}[\chi_{t+1} \mathbf{1}\{I_t = i\}]$, and using $g_{t+1,i}^2 \leq 4(\ell_{t+1,i} - \ell_{t,i})^2 \leq 4$, give

$$\mathbb{E}[\chi_{t+1} x_{t,i}] g_{t+1,i}^2 \leq 8\mathbb{E}[\chi_{t+1} \mathbf{1}\{I_t = i\}] (\ell_{t+1,i} - \ell_{t,i})^2 + 24 \frac{(\mathbb{E}[\chi_{t+1}(\mathbf{1}\{I_t = i\} - x_{t,i})])^2}{\mathbb{E}[\chi_{t+1}(\mathbf{1}\{I_t = i\} + x_{t,i})]}.$$

Summing over $t \in \{2, \dots, T-1\}$ and $i \in [K]$ yields

$$\begin{aligned} \sum_{t=2}^{T-1} \sum_{i=1}^K \mathbb{E}[\chi_{t+1} x_{t,i}] g_{t+1,i}^2 &\leq 8\mathbb{E} \left[\sum_{t=3}^T \chi_t (\ell_{t,I_{t-1}} - \ell_{t-1,I_{t-1}})^2 \right] + 24 \sum_{t=2}^{T-1} \mathcal{N}(\chi_{t+1}; I_t, x_t) \\ &\leq 8P + 24 \sum_{t=2}^{T-1} \mathcal{N}(\chi_{t+1}; I_t, x_t). \end{aligned}$$

Summing (A.15) over $t = 2, \dots, T-1$ gives

$$\sum_{t=2}^{T-1} \mathcal{N}(\chi_{t+1}; I_t, x_t) \leq \sum_{t=2}^{T-1} \mathcal{N}_t^{\text{cur}} + 6\mathbb{E} \left[\sum_{t=2}^{T-1} \chi_t(1 - \chi_{t+1}) \right] + 2\mathbb{E} \left[\sum_{t=2}^{T-1} (1 - \chi_t) \chi_{t+1} \right] \leq \sum_{t=2}^{T-1} \mathcal{N}_t^{\text{cur}} + 8\rho.$$

Combining the preceding bounds and applying (A.14) yield

$$\begin{aligned} \sum_{t=2}^{T-1} \sum_{i=1}^K q_{t,i} g_{t+1,i}^2 &\leq 7\rho + 4P + 8P + 24 \sum_{t=2}^{T-1} \mathcal{N}(\chi_{t+1}; I_t, x_t) \\ &\leq 12P + 24 \sum_{t=2}^{T-1} \mathcal{N}_t^{\text{cur}} + 199\rho \\ &\leq 12360\eta^2 R + (12 + 216\alpha^2)P + (199 + 8160\alpha)\rho \\ &\leq 12360\eta^2 R + 13P + 200\rho, \end{aligned}$$

where $\alpha = 8\eta \leq 8 \cdot 10^{-5}$ implies $216\alpha^2 \leq 1$ and $8160\alpha \leq 1$. □

Proposition A.8. For any $A, B \geq 0$, we have $A \leq 2B + 6 \frac{(A-B)^2}{A+B}$.

Proof. The inequality is immediate when $A \leq 2B$. When $A > 2B$, we have $6 \frac{(A-B)^2}{A+B} \geq 6 \frac{(A/2)^2}{3A/2} = A$. □