

Policy Iteration Is Not Strongly Polynomial for Deterministic Markov Decision Processes: The Price of Algorithmic Anarchy

Han Zhong*

Yinyu Ye[†]

Abstract

We establish an exponential iteration lower bound in the number of states for Howard’s policy iteration on deterministic discounted Markov decision processes, with at most two actions per state. This rules out strong polynomiality of Howard’s policy iteration when the discount factor is part of the input and yields an exponential separation from the simplex method with Dantzig’s pivoting rule, which is proved to be strongly polynomial on this class. Even when each reward is restricted to logarithmic bit length, we obtain a stretched-exponential iteration lower bound. The gap between Howard’s decentralized and simultaneous selfish improvements and Dantzig’s coordinated selection of a single action with the largest gain across all states reveals a “price” of algorithmic anarchy.

Keywords. deterministic Markov decision processes; policy iteration; strongly polynomial algorithms.

1 Introduction

Markov decision processes (MDPs) provide a standard framework for sequential decision-making and reinforcement learning (Puterman, 1994; Sutton and Barto, 2018). A deterministic discounted MDP has a finite state space $\mathcal{S}$ and a nonempty finite action set $\mathcal{A}_s$ at each state s . Taking action $a \in \mathcal{A}_s$ yields a reward $r(s, a) \in \mathbb{Q}$ and moves to the successor state $f(s, a) \in \mathcal{S}$. Future rewards are discounted by a common factor $0 < \gamma < 1$. Write $N = |\mathcal{S}|$. A policy π selects an action $\pi(s) \in \mathcal{A}_s$ at each state. Its value $V^\pi(s)$ is the total discounted reward obtained from s by following π . The objective is to find a policy maximizing these values for all initial states. The action value $Q^\pi(s, a)$ is the discounted return from using a once and then following π . We have

$$V^\pi(s) = r(s, \pi(s)) + \gamma V^\pi(f(s, \pi(s))), \quad Q^\pi(s, a) = r(s, a) + \gamma V^\pi(f(s, a)). \quad (1.1)$$

Policy iteration, also known as Howard’s policy iteration (Howard, 1960), is a foundational algorithm for solving MDPs. Let π_0 be the initial policy and π_t the policy after t iterations. Given π_t , the algorithm computes V^{π_t} and uses the corresponding action values Q^{π_t} from (1.1) to select π_{t+1} :

$$\pi_{t+1}(s) \in \arg \max_{a \in \mathcal{A}_s} Q^{\pi_t}(s, a), \quad \forall s \in \mathcal{S}. \quad (1.2)$$

If $\pi_t(s)$ attains the maximum in (1.2), we set $\pi_{t+1}(s) = \pi_t(s)$. Otherwise, we choose the first maximizer in a fixed ordering of $\mathcal{A}_s$. The algorithm stops when $\pi_{t+1} = \pi_t$, which holds if and only if π_t is optimal.

For a fixed discount factor, policy iteration is strongly polynomial (Ye, 2011; Scherrer, 2016), a guarantee previously established for an interior-point algorithm (Ye, 2005). These bounds for policy iteration depend on the discount and therefore do not establish strong polynomiality when the discount is part of the input. Whereas Howard’s policy iteration makes simultaneous local improvements, the simplex method with Dantzig’s pivoting rule updates only one state per iteration, selecting an action with the largest positive gain $Q^\pi(s, a) - V^\pi(s)$ across all states (Ye, 2011). For deterministic MDPs, Post and Ye (2015) prove that this method is strongly polynomial independently of the discount. Whether Howard’s policy iteration is strongly polynomial on deterministic MDPs has remained open (Goenka et al., 2026).

*Shanghai Jiao Tong University. Email: han.zhong@sjtu.edu.cn.

[†]Shanghai Jiao Tong University, SIMIS, and Stanford University. Email: yinyu-ye@stanford.edu.

Exponential lower bounds for policy iteration are known for general MDPs (Fearnley, 2010; Hollanders et al., 2012). These constructions use stochastic transitions and a number of actions per state that grows with the instance size. Hansen and Zwick (2010) and Hansen (2012) establish quadratic iteration lower bounds for deterministic MDPs. For MDPs with a constant number of actions per state, no superpolynomial lower bound for Howard’s policy iteration was previously known, even with stochastic transitions (Mukherjee and Kalyanakrishnan, 2025).

We establish an exponential iteration lower bound for deterministic discounted MDPs, even when each state has at most two actions.

Theorem 1.1. *There is a family of deterministic discounted MDPs with N states and at most two actions per state, on which Howard’s policy iteration performs at least $\exp(\Omega(N))$ iterations from a specified initial policy. Each instance has encoding length $O(N^2)$.*

In the construction for Theorem 1.1, each reward uses $O(N)$ bits. We next consider integer rewards encoded with $O(\log N)$ bits. Under this restriction, Asadi et al. (2025) give a quadratic lower bound for deterministic average-reward MDPs. For deterministic discounted MDPs with at most two actions per state and nonnegative integer rewards encoded with $O(\log N)$ bits, Mukherjee and Kalyanakrishnan (2025) give an iteration upper bound of $\exp(O(\sqrt{N}(\log N)^{3/2})) = \exp(o(N))$, independently of the discount. This rules out the exponential behavior in Theorem 1.1 but leaves open whether a polynomial iteration bound holds. The next theorem gives a stretched-exponential lower bound even with nonnegative integer rewards smaller than $4N$, and hence with $O(\log N)$ bits per reward.

Theorem 1.2. *There is a family of deterministic discounted MDPs with N states, at most two actions per state, and nonnegative integer rewards smaller than $4N$, on which Howard’s policy iteration performs at least $\exp(\Omega(N^{1/4}))$ iterations from a specified initial policy. Each instance has encoding length $O(N \log N)$.*

In summary, Theorem 1.1 rules out strong polynomiality of Howard’s policy iteration on deterministic MDPs even with at most two actions per state, and Theorem 1.2 shows that superpolynomial iteration complexity persists with $O(\log N)$ bits per reward. The exponential separation from Dantzig’s rule reveals a “price” of algorithmic anarchy: Howard’s rule makes simultaneous local greedy updates using the same policy values, whereas Dantzig’s rule coordinates updates across states and re-evaluates after each switch (Remark 3.2).

2 Construction of the MDPs

Section 2.1 gives the formal construction of the MDPs, and Section 2.2 describes a family of policies that encode binary numbers and compares their values. We first introduce the parameters and polynomials used in the construction.

Fix an integer $d \geq 3$, and let $F_i, G_i \in \mathbb{Z}[x]$, $0 \leq i < d$. Assume that $F_0(x) > 0$ for x sufficiently close to 1 from below. Define

$$f_i(\gamma) := \frac{F_i(\gamma)}{F_0(\gamma)}, \quad f_i := \lim_{\gamma \rightarrow 1^-} f_i(\gamma), \quad g_i(\gamma) := \frac{G_i(\gamma)}{F_0(\gamma)}, \quad g_i := \lim_{\gamma \rightarrow 1^-} g_i(\gamma), \quad (2.1)$$

where the limits are assumed to exist and satisfy

$$f_0 = 1, \quad f_i \geq 2f_{i-1}, \quad g_0 \geq 4f_{d-1}, \quad g_i \geq 2g_{i-1} \quad \forall 1 \leq i < d. \quad (2.2)$$

Set $L = 2 + \max_{0 \leq i < d} \{\deg F_i, \deg G_i\} \geq 2$, where $\deg$ denotes polynomial degree. Write

$$F_i(x) = \sum_{j=1}^{L-1} F_{i,j} x^{j-1}, \quad G_i(x) = \sum_{j=1}^{L-1} G_{i,j} x^{j-1},$$

and set $F_{i,j} = G_{i,j} = 0$ for $j \notin \{1, \dots, L-1\}$. Throughout this section, $0 \leq i < d$ unless otherwise specified.

2.1 Formal construction

Given the parameters and polynomials above, we define the MDP as follows.

States and actions. The state space $\mathcal{S}$ is partitioned into four sets:

1. $\mathcal{S}_1 = \{s_i, s_{i,1}, \dots, s_{i,5}, \bar{s}_i, \bar{s}_{i,1}, \dots, \bar{s}_{i,4}\}_{i=0}^{d-1} \cup \{s_d, \bar{s}_d\}$, with $|\mathcal{S}_1| = 11d + 2$.
2. $\mathcal{S}_2 = \{s_i^c, s_i^z\}_{i=0}^{d-1} \cup \{s_{-1}^c, s_{-1}^z\}$, with $|\mathcal{S}_2| = 2d + 2$.
3. $\mathcal{S}_3 = \{s_{i,0}^r, s_{i,1}^r, s_{i,2}^r\}_{i=0}^{d-1}$, with $|\mathcal{S}_3| = 3d$.
4. $\mathcal{S}_4$ consists of the $(67d + 5)L - 32d - 5$ intermediate states introduced in the transition construction below.

The action sets are

$$\mathcal{A}_s = \begin{cases} \{a_0\}, & s \in \mathcal{S}_4 \cup \{\bar{s}_d, s_{-1}^c, s_{-1}^z\}, \\ \{a_0, a_1\}, & \text{otherwise.} \end{cases}$$

For the total number of states, we obtain

$$N = \sum_{j=1}^4 |\mathcal{S}_j| = (67d + 5)L - 16d - 1 = \Theta(dL). \quad (2.3)$$

Transitions. For each $s \in \mathcal{S}_1 \cup \mathcal{S}_2 \cup \mathcal{S}_3$ and $a \in \mathcal{A}_s$, Table 1 specifies an endpoint $s' \in \mathcal{S}_1 \cup \mathcal{S}_2 \cup \mathcal{S}_3$ and a path length m , a positive multiple of L . Introduce $m - 1$ intermediate states $\{s_{a,j}^\circ\}_{j=1}^{m-1}$ and set

$$f(s, a) = s_{a,1}^\circ, \quad f(s_{a,j}^\circ, a_0) = s_{a,j+1}^\circ \quad \forall 1 \leq j < m - 1, \quad f(s_{a,m-1}^\circ, a_0) = s'.$$

Thus, starting from s , we reach $s_{a,j}^\circ$ after j transitions. The intermediate states are distinct for different pairs (s, a) and together constitute $\mathcal{S}_4$.

1. *Paths starting in $\mathcal{S}_1$.* Figure 1 shows the $6L$ -transition paths from s_i to s_{i+1} and from $\bar{s}_i$ to $\bar{s}_{i+1}$, together with the branches at $s_{i,4}, s_{i,5}, \bar{s}_{i,4}$. The paths from s_d have length L and end at $\bar{s}_0$ under a_0 and at s_0 under a_1 . The sole action at $\bar{s}_d$ gives an L -transition path to $\bar{s}_0$.
2. *Paths starting in $\mathcal{S}_2$.* For $0 \leq i < d$, the a_0 paths from s_i^c and s_i^z both end at $s_{i,0}^r$. The a_1 paths end at $s_{i,1}^r$ and $s_{i,2}^r$, respectively. The sole actions at s_{-1}^c, s_{-1}^z give paths to $\bar{s}_0$. Every path from $\mathcal{S}_2$ has length L .
3. *Paths starting in $\mathcal{S}_3$.* Under a_0 , the path from $s_{i,1}^r$ ends at s_{i-1}^c , and those from $s_{i,0}^r$ and $s_{i,2}^r$ end at s_{i-1}^z . Under a_1 , the path from $s_{i,0}^r$ ends at s_{i+1} , and those from $s_{i,1}^r$ and $s_{i,2}^r$ end at $s_{i,1}$. All paths have length L except the a_1 path from $s_{i,0}^r$, which has length $6L$.

The $32d + 5$ paths in Table 1 have total length $(67d + 5)L$. Each path of length m adds $m - 1$ states, so $|\mathcal{S}_4| = (67d + 5)L - 32d - 5$.

Rewards. We first define integer rewards, allowing negative values. Before the uniform shift below, nonzero rewards occur only at the intermediate states in $\mathcal{S}_4$. For $s \in \mathcal{S}_1 \cup \mathcal{S}_2 \cup \mathcal{S}_3$, each action $a \in \mathcal{A}_s$ has immediate reward $r(s, a) = 0$ and selects a path through $\mathcal{S}_4$. The following rules specify the intermediate rewards according to the set containing the starting state of each path. For a path of length m in Table 1, the formulas apply to $1 \leq j < m$. All unspecified rewards are zero.

1. *Paths starting in $\mathcal{S}_1$.* For $s \in \mathcal{S}_1$, set

$$r(s_{a,j}^\circ, a_0) = \begin{cases} (96d + 16)G_{i,j}, & s = s_{i,4}, a = a_0, & (2.4a) \\ 4F_{i,j} - 16F_{d-1,j} - 2F_{0,j}, & s = s_{i,4}, a = a_1, & (2.4b) \\ -(96d + 16)G_{i,j}, & s = s_{i,5}, a = a_0, & (2.4c) \\ -(96d + 16)G_{i,j} - 4F_{i,j} - F_{0,j}, & s = s_{i,5}, a = a_1, & (2.4d) \\ (96d + 16)(G_{i,j} - G_{i,j-L}), & s = \bar{s}_{i,4}, a = a_0, & (2.4e) \\ -32F_{d-1,j}, & s = \bar{s}_{i,4}, a = a_1, & (2.4f) \\ 16F_{d-1,j}, & s = s_d, a = a_0, & (2.4g) \\ 0, & \text{otherwise.} \end{cases}$$




Figure 1: Top: $6L$ -transition paths from s_i to s_{i+1} and from $\bar{s}_i$ to $\bar{s}_{i+1}$, with labels before and after each slash, respectively. Arrows without a length label represent paths of length L . The upper $2L$ arrow represents either the path through $s_{i,5}$ with a_0 at both $s_{i,4}$ and $s_{i,5}$, or the a_0 path from $\bar{s}_{i,4}$ to $\bar{s}_{i+1}$. Bottom: action choices and path lengths at $s_{i,4}$, $s_{i,5}$, $\bar{s}_{i,4}$.

2. *Paths starting in $\mathcal{S}_2$.* For $s \in \mathcal{S}_2$, set

$$r(s_{a,j}^\circ, a_0) = \begin{cases} 16F_{d-1,j}, & s = s_{-1}^c, a = a_0, \\ 0, & \text{otherwise.} \end{cases} \quad (2.5)$$

3. *Paths starting in $\mathcal{S}_3$.* Set

$$r(s_{a,j}^\circ, a_0) = \begin{cases} 4F_{i,j}, & s = s_{i,0}^r, a = a_0, \\ -32F_{d-1,j} - 4F_{i,j} + 4F_{0,j} - 4F_{0,j-L}, & s = s_{i,0}^r, a = a_1, \\ -32F_{d-1,j} - 4F_{i,j}, & s \in \{s_{i,1}^r, s_{i,2}^r\}, a = a_1, \\ 0, & \text{otherwise.} \end{cases} \quad \begin{matrix} (2.6a) \\ (2.6b) \\ (2.6c) \end{matrix}$$

Let $r_{\max} = \max_{s \in \mathcal{S}, a \in \mathcal{A}_s} |r(s, a)|$ be the maximum absolute unshifted immediate reward. Add $r_{\max}$ to every reward, so the final rewards lie in $[0, 2r_{\max}]$. This adds $r_{\max}/(1 - \gamma)$ to every $V^\pi(s)$ and $Q^\pi(s, a)$, leaving all comparisons and optimality gaps unchanged. The proof uses the unshifted rewards.

Discount and initial policy. We choose the discount factor γ and the initial policy π_0 as follows:

$$\gamma = 1 - \frac{1}{4^N r_{\max}}, \quad \pi_0(s) = \begin{cases} a_1, & s \in \{s_d\} \cup \{s_{i,j} : 0 \leq i < d, 1 \leq j \leq 3\}, \\ a_0, & \text{otherwise.} \end{cases} \quad (2.7)$$

2.2 Binary encoding and policy values

A policy π encodes the same vector $\mathbf{b} \in \{0, 1\}^d$ in the first group $\{s_i\}_{i=0}^{d-1}$ and the second group $\{\bar{s}_i\}_{i=0}^{d-1}$ if

$$\pi(s_i) = \pi(\bar{s}_i) = a_{[\mathbf{b}]_i}, \quad \forall 0 \leq i < d.$$

The states s_i and $\bar{s}_i$ represent the i -th bit, with $i = 0$ denoting the least significant bit. Write $\mathbf{0} = (0, \dots, 0)$ and $\mathbf{1} = (1, \dots, 1)$. For $\mathbf{b} \neq \mathbf{1}$, let $\mathbf{b}^+$ denote the result of adding one in binary. For example, when $d = 5$,

Set	Starting state	Action a_0		Action a_1	
		Endpoint	Length	Endpoint	Length
$\mathcal{S}_1$	$s_i/\bar{s}_i$	$s_{i+1}/\bar{s}_{i+1}$	$6L$	$s_{i,1}/\bar{s}_{i,1}$	L
	$s_{i,j}/\bar{s}_{i,j}$	$s_{i+1}/\bar{s}_{i+1}$	$(6-j)L$	$s_{i,j+1}/\bar{s}_{i,j+1}$	L
	$s_{i,4}$	$s_{i,5}$	L	s_i^c	$2L$
	$s_{i,5}$	s_{i+1}	L	s_i^z	L
	$\bar{s}_{i,4}$	$\bar{s}_{i+1}$	$2L$	s_i	L
	s_d	$\bar{s}_0$	L	s_0	L
	$\bar{s}_d$	$\bar{s}_0$	L	—	—
$\mathcal{S}_2$	s_i^c	$s_{i,0}^r$	L	$s_{i,1}^r$	L
	s_i^z	$s_{i,0}^r$	L	$s_{i,2}^r$	L
	s_{-1}^c, s_{-1}^z	$\bar{s}_0$	L	—	—
$\mathcal{S}_3$	$s_{i,0}^r$	s_{i-1}^z	L	s_{i+1}	$6L$
	$s_{i,1}^r$	s_{i-1}^c	L	$s_{i,1}$	L
	$s_{i,2}^r$	s_{i-1}^z	L	$s_{i,1}$	L

Table 1: Paths from $\mathcal{S}_1$, $\mathcal{S}_2$, and $\mathcal{S}_3$. Here $0 \leq i < d$ except in the row explicitly indexed by -1 , and $1 \leq j \leq 3$ in the second row of the $\mathcal{S}_1$ block. Slashes separate corresponding states, as in $s_i/\bar{s}_i$. A dash indicates an unavailable action.

$\mathbf{b} = 01011$ represents 11, and $\mathbf{b}^+ = 01100$ represents 12. For each $\mathbf{b} \in \{0, 1\}^d$, define the policy $\pi[\mathbf{b}]$ by

$$\pi[\mathbf{b}](s) = \begin{cases} a_{[\mathbf{b}]_i}, & s \in \{s_i, \bar{s}_i, s_i^c, s_i^z, \bar{s}_{i,1}, \bar{s}_{i,2}, \bar{s}_{i,3}\}, \quad 0 \leq i < d, \\ \pi_0(s), & \text{otherwise,} \end{cases} \quad (2.8)$$

where π_0 is defined in (2.7). In particular, $\pi[\mathbf{0}] = \pi_0$. For $0 \leq i < d$, $\pi[\mathbf{b}]$ selects the following paths from s_i to s_{i+1} and from $\bar{s}_i$ to $\bar{s}_{i+1}$:

- If $[\mathbf{b}]_i = 0$, the policy selects a_0 at s_i and $\bar{s}_i$. Each path reaches its endpoint in $6L$ transitions.
- If $[\mathbf{b}]_i = 1$, the policy follows the horizontal paths in the top panel of Figure 1. From s_i , it passes through $s_{i,1}, \dots, s_{i,5}$ to s_{i+1} : four a_1 paths followed by two a_0 paths, all of length L . From $\bar{s}_i$, it passes through $\bar{s}_{i,1}, \dots, \bar{s}_{i,4}$ to $\bar{s}_{i+1}$: four a_1 paths of length L , followed by one a_0 path of length $2L$. Both paths therefore have length $6L$.

The policy selects a_1 at s_d . The selected paths form a cycle from s_0 through $s_1, \dots, s_d$ and back to s_0 , and another from $\bar{s}_0$ through $\bar{s}_1, \dots, \bar{s}_d$ and back to $\bar{s}_0$. Each cycle consists of d paths of length $6L$ followed by an L -transition path back to its starting state, giving total length $(6d + 1)L$.

Lemma 2.1 (Value ordering). *There exists $\gamma_0 \in (0, 1)$ such that, for all $\gamma \in (\gamma_0, 1)$ and $\mathbf{b} \in \{0, 1\}^d \setminus \{\mathbf{1}\}$,*

$$V^{\pi[\mathbf{b}^+]}(s_0) > V^{\pi[\mathbf{b}]}(s_0), \quad V^{\pi[\mathbf{b}^+]}(\bar{s}_0) > V^{\pi[\mathbf{b}]}(\bar{s}_0).$$

Proof. We use the unshifted rewards, which give the same value differences. Fix any $\mathbf{b} \in \{0, 1\}^d$. For $0 \leq i < d$, consider the paths from s_i to s_{i+1} and from $\bar{s}_i$ to $\bar{s}_{i+1}$ selected by $\pi[\mathbf{b}]$. If $[\mathbf{b}]_i = 0$, the policy selects a_0 at s_i and $\bar{s}_i$, and every reward along both paths is zero by the last case of (2.4). If $[\mathbf{b}]_i = 1$, both paths have zero rewards for the first $4L$ transitions. By (2.4a), (2.4c), and (2.4e), the remaining $2L$ transitions on each path carry the following two L -term reward sequences, in order:

$$(0, (96d + 16)G_{i,1}, \dots, (96d + 16)G_{i,L-1}), \quad (0, -(96d + 16)G_{i,1}, \dots, -(96d + 16)G_{i,L-1}).$$

On each $6L$ -transition path, the second sequence starts L transitions after the first and contributes $-\gamma^L$ times

the discounted reward of the first. For the total discounted reward along each path, we therefore obtain

$$(96d + 16)\gamma^{4L}(1 - \gamma^L) \sum_{j=1}^{L-1} \gamma^j G_{i,j} = (96d + 16)\gamma^{4L+1}(1 - \gamma^L)G_i(\gamma). \quad (2.9)$$

Each bit i with $[\mathbf{b}]_i = 1$ contributes γ^{6iL} times (2.9) to the discounted reward over one cycle starting from s_0 or $\bar{s}_0$. The paths selected at s_d and $\bar{s}_d$ have zero rewards. Summing over i with $[\mathbf{b}]_i = 1$ and repeating the cycle every $(6d + 1)L$ transitions, we obtain

$$V^{\pi[\mathbf{b}]}(s_0) = V^{\pi[\mathbf{b}]}(\bar{s}_0) = \frac{(96d + 16)\gamma^{4L+1}(1 - \gamma^L)}{1 - \gamma^{(6d+1)L}} \sum_{i=0}^{d-1} \gamma^{6iL} G_i(\gamma) [\mathbf{b}]_i. \quad (2.10)$$

For $\mathbf{b} \neq \mathbf{1}$, we compare the cycle values in (2.10) for $\mathbf{b}$ and $\mathbf{b}^+$. Let $k = \min\{i : [\mathbf{b}]_i = 0\}$ be the index of the least significant zero bit of $\mathbf{b}$. Adding one changes bit k from zero to one and every lower bit from one to zero, leaving higher bits unchanged. For $s \in \{s_0, \bar{s}_0\}$, we combine the polynomial limits in (2.1) with $(1 - \gamma^L)/(1 - \gamma^{(6d+1)L}) \rightarrow 1/(6d + 1)$ and the condition $\mathbf{g}_i \geq 2\mathbf{g}_{i-1}$ in (2.2) to obtain

$$\begin{aligned} \lim_{\gamma \rightarrow 1^-} \frac{V^{\pi[\mathbf{b}^+]}(s) - V^{\pi[\mathbf{b}]}(s)}{4\gamma F_0(\gamma)} &= 4 \sum_{i=0}^{d-1} \mathbf{g}_i ([\mathbf{b}^+]_i - [\mathbf{b}]_i) \\ &= 4 \left(\mathbf{g}_k - \sum_{i=0}^{k-1} \mathbf{g}_i \right) \geq 4 \left(\mathbf{g}_k - \sum_{i=0}^{k-1} (\mathbf{g}_{i+1} - \mathbf{g}_i) \right) = 4\mathbf{g}_0 > 0. \end{aligned} \quad (2.11)$$

We conclude that the two value inequalities in Lemma 2.1 hold for $\gamma < 1$ sufficiently close to one. Since there are finitely many encodings, a common threshold $\gamma_0 \in (0, 1)$ suffices for all $\mathbf{b} \neq \mathbf{1}$. $\square$

By Lemma 2.1, for $\gamma < 1$ sufficiently close to one, the 2^d policies $\pi[\mathbf{b}]$, listed in binary order, have strictly increasing values at s_0 and $\bar{s}_0$. In Section 3, we show that policy iteration, starting from $\pi[\mathbf{0}]$, performs the first $2^d - 2$ increments in this order, using five iterations per increment. This yields $\Omega(2^d)$ iterations. Section 4 derives the corresponding lower bounds in the number of states.

3 Binary counting under policy iteration

Lemma 3.1 (Binary counting). *For the construction in Section 2, with γ, π_0 in (2.7), Howard's policy iteration in (1.2) produces $\pi[\mathbf{b}]$ after $5 \sum_{i=0}^{d-1} 2^i [\mathbf{b}]_i$ iterations for every $\mathbf{b} \in \{0, 1\}^d \setminus \{\mathbf{1}\}$.*

Proof overview. Starting from $\pi[\mathbf{b}]$, where $\mathbf{b}$ represents an integer in $0, \dots, 2^d - 3$, five iterations produce $\pi[\mathbf{b}^+]$. Figure 2 tracks the selected paths and bit choices: the first group encodes $\mathbf{b}^+$ after Iteration 2 but forms its own cycle after Iteration 3. Each update is simultaneous and uses action values under the preceding policy.

Iteration 1. Every s_i selects a_1 , while the second group retains $\mathbf{b}$ and its cycle. The comparisons under $\pi[\mathbf{b}]$ at $s_{i,4}, s_{i,5}$ select branches through $\mathcal{S}_2$ at exactly the zero positions of $\mathbf{b}^+$. These branches determine which s_i will select a_0 in Iteration 2. As Figure 2(b) shows, the path from s_0 now enters the old cycle through $\mathcal{S}_2$, and s_d selects a_0 toward $\bar{s}_0$. The first group therefore has no cycle of its own.

Iteration 2. The states s_i select a_0 at the positions chosen in Iteration 1, bypassing the branches through $\mathcal{S}_2$, and retain a_1 elsewhere. The first group therefore encodes $\mathbf{b}^+$. The selected path from s_0 reaches s_d and then enters the old cycle through $\bar{s}_0$, traversing the new encoding only once (Figure 2(c)).

Iteration 3. The state s_d selects a_1 , returning to s_0 . The first-group path now repeats, forming a cycle for $\mathbf{b}^+$, while the second group retains its cycle for $\mathbf{b}$ (Figure 2(d)). The new cycle's value advantage over the old cycle outweighs the reward losses along the connecting paths, making them preferable in Iteration 4.

Iteration 4. The states $\bar{s}_{i,4}$ and those in $\mathcal{S}_3$ select a_1 , establishing the connections into the first group. All $\bar{s}_i$ also select a_1 , but their paths enter the first-group cycle through $\bar{s}_{i,4}$, as in Figure 2(e).

Iteration 5. Through these connections, the action comparisons at $\bar{s}_i, s_i^c, s_i^z$ reduce to the comparison at s_i , with a small bias toward a_0 . When $[\mathbf{b}^+]_i = 0$, the actions at s_i tie, so these states select a_0 . When $[\mathbf{b}^+]_i = 1$, the extra reward from a_1 outweighs the bias, so they select a_1 . The states $\bar{s}_{i,4}$ and those in $\mathcal{S}_3$ return to a_0 . Together with the updates at $\{s_{i,j}\}_{j=1}^3$ and $\{\bar{s}_{i,j}\}_{j=1}^3$, these changes give the full policy $\pi[\mathbf{b}^+]$. Both groups again have their own cycles, now encoding $\mathbf{b}^+$ (Figure 2(f)).

The states $\{s_{i,j}\}_{j=1}^3$ and $\{\bar{s}_{i,j}\}_{j=1}^3$ delay changes at s_i and $\bar{s}_i$, respectively. Within each set of three states, switches from a_0 to a_1 proceed in the order $j = 3, 2, 1$, one per iteration, while ties under the preceding policy keep the corresponding bit at zero. This preserves the old second-group zeros through Iteration 3 and the new first-group zeros through Iteration 5. Table 2 records all action choices, which are verified in the proof below.




Figure 2: Selected paths and bit choices during the five iterations from $\pi[\mathbf{b}]$ to $\pi[\mathbf{b}^+]$. Circles denote states, and arrows represent paths selected by the current policy. Within each panel, the upper and lower rows show the first and second groups, respectively.

Proof of Lemma 3.1. Suppose $\pi_t = \pi[\mathbf{b}]$, where $\mathbf{b}$ encodes an integer in $0, \dots, 2^d - 3$. Let

$$k = \min\{i : [\mathbf{b}]_i = 0\}. \quad (3.1)$$

We show that the next five iterations produce $\pi_{t+5} = \pi[\mathbf{b}^+]$. In Sections 3.1–3.5, we determine the actions selected at each iteration by comparing action values as $\gamma \rightarrow 1^-$. Section 3.6 verifies these comparisons for the discount in (2.7). Table 2 records the choices at all states with two actions throughout the increment.

We divide the unshifted rewards by $4\gamma F_0(\gamma)$, which is positive for $\gamma < 1$ sufficiently close to one. For a path starting with action a at s and then following π , suppose its first visit to s' at a positive time occurs after m transitions. When $s' = s$, this is the first return. Let $r_0, \dots, r_{m-1}$ be the unshifted rewards before this visit and define

$$\tilde{R}^\pi((s, a) \rightsquigarrow s') := \frac{\sum_{\ell=0}^{m-1} \gamma^\ell r_\ell}{4\gamma F_0(\gamma)}, \quad \tilde{R}^\pi(s \rightsquigarrow s') = \tilde{R}^\pi((s, \pi(s)) \rightsquigarrow s'). \quad (3.2)$$

For $0 \leq i < d$, the selected $6L$ -transition paths under $\pi[\mathbf{b}]$ have scaled rewards given by (2.4) and (2.9):

$$\tilde{R}^{\pi[\mathbf{b}]}(s_i \rightsquigarrow s_{i+1}) = \tilde{R}^{\pi[\mathbf{b}]}(\bar{s}_i \rightsquigarrow \bar{s}_{i+1}) = [\mathbf{b}]_i \gamma^{4L} R_i(\gamma), \quad R_i(\gamma) := (24d + 4)g_i(\gamma)(1 - \gamma^L). \quad (3.3)$$

State	π_t	π_{t+1}	π_{t+2}	π_{t+3}	π_{t+4}	π_{t+5}
s_i	$[\mathbf{b}]_i$	1	$[\mathbf{b}^+]_i$	$[\mathbf{b}^+]_i$	$[\mathbf{b}^+]_i$	$[\mathbf{b}^+]_i$
$s_{i,1}$	1	1	$[\mathbf{b}^+]_i$	$[\mathbf{b}^+]_i$	$[\mathbf{b}^+]_i$	1
$s_{i,2}$	1	1	$[\mathbf{b}^+]_i$	$[\mathbf{b}^+]_i$	1	1
$s_{i,3}$	1	1	$[\mathbf{b}^+]_i$	1	1	1
$s_{i,4}$	0	$\mathbb{1}\{i < k\}$	0	0	0	0
$s_{i,5}$	0	$\mathbb{1}\{i > k, [\mathbf{b}]_i = 0\}$	0	0	0	0
s_d	1	0	0	1	1	1
$\bar{s}_i$	$[\mathbf{b}]_i$	$[\mathbf{b}]_i$	$[\mathbf{b}]_i$	$[\mathbf{b}]_i$	1	$[\mathbf{b}^+]_i$
$\bar{s}_{i,1}$	$[\mathbf{b}]_i$	$[\mathbf{b}]_i$	$[\mathbf{b}]_i$	1	1	$[\mathbf{b}^+]_i$
$\bar{s}_{i,2}$	$[\mathbf{b}]_i$	$[\mathbf{b}]_i$	1	1	1	$[\mathbf{b}^+]_i$
$\bar{s}_{i,3}$	$[\mathbf{b}]_i$	1	1	1	1	$[\mathbf{b}^+]_i$
$\bar{s}_{i,4}$	0	0	0	0	1	0
s_i^c	$[\mathbf{b}]_i$	$\mathbb{1}\{i \leq k\}$	$\mathbb{1}\{i \leq k+1\}$	$\mathbb{1}\{i \leq k+2\}$	$\mathbb{1}\{i \leq k+3\}$	$[\mathbf{b}^+]_i$
s_i^z	$[\mathbf{b}]_i$	0	0	0	0	$[\mathbf{b}^+]_i$
$s_{i,j}^r$	0	0	0	0	1	0

Table 2: Policy choices at all states with two actions during one binary increment. Entries 0, 1 denote a_0, a_1 . Here $0 \leq i < d$, $0 \leq j \leq 2$, $k = \min\{i : [\mathbf{b}]_i = 0\}$, and $\mathbb{1}\{\cdot\}$ denotes the indicator. All remaining states have only action a_0 .

Here $R_i(\gamma)$ is the scaled reward along the $2L$ -transition a_0 path from $\bar{s}_{i,4}$ to $\bar{s}_{i+1}$. For any policy π , define the scaled values

$$\tilde{V}^\pi(s) := \frac{V^\pi(s)}{4\gamma F_0(\gamma)}, \quad \tilde{Q}^\pi(s, a) := \frac{Q^\pi(s, a)}{4\gamma F_0(\gamma)}, \quad \tilde{Q}^\pi(s, a) = \tilde{R}^\pi((s, a) \rightsquigarrow s') + \gamma^m \tilde{V}^\pi(s'), \quad (3.4)$$

where the decomposition in (3.4) follows by applying (1.1) along the path in (3.2). Using (3.3) and (3.4), we obtain

$$\tilde{V}^{\pi[\mathbf{b}]}(s_i) = \gamma^{6L} \tilde{V}^{\pi[\mathbf{b}]}(s_{i+1}) + [\mathbf{b}]_i \gamma^{4L} R_i(\gamma), \quad \forall 0 \leq i < d, \quad \tilde{V}^{\pi[\mathbf{b}]}(s_d) = \gamma^L \tilde{V}^{\pi[\mathbf{b}]}(s_0). \quad (3.5)$$

We also obtain (3.5) with each s_i replaced by $\bar{s}_i$, since the corresponding selected paths have the same lengths and reward sequences (see Section 2.2). We obtain the common limit at s_0 and $\bar{s}_0$ by dividing (2.10) by $4\gamma F_0(\gamma)$ and taking $\gamma \rightarrow 1^-$ as in the derivation of (2.11). Since $\gamma^L \rightarrow 1$ and $R_i(\gamma) \rightarrow 0$, (3.5) gives this limit first at $s_d, \bar{s}_d$ and then at $s_i, \bar{s}_i$ for $i = d-1, \dots, 1$. Hence, we have

$$\lim_{\gamma \rightarrow 1^-} \tilde{V}^{\pi[\mathbf{b}]}(s_i) = 4 \sum_{j=0}^{d-1} \mathbf{g}_j [\mathbf{b}]_j = \lim_{\gamma \rightarrow 1^-} \tilde{V}^{\pi[\mathbf{b}]}(\bar{s}_i), \quad \forall 0 \leq i \leq d. \quad (3.6)$$

We express the policy iteration rule at states with two actions in terms of

$$\Delta^\pi(s) := \tilde{Q}^\pi(s, a_1) - \tilde{Q}^\pi(s, a_0). \quad (3.7)$$

By (1.2), the next policy selects a_1 if $\Delta^\pi(s) > 0$, selects a_0 if $\Delta^\pi(s) < 0$, and keeps $\pi(s)$ otherwise.

3.1 Iteration 1: identify the zero positions of $\mathbf{b}^+$

Before this iteration, $\pi_t = \pi[\mathbf{b}]$. We show that this iteration sets all bits at s_i to one, leaves the bits at $\bar{s}_i$ unchanged, and selects paths through $\mathcal{S}_2$ at the zero positions of $\mathbf{b}^+$.

Step 1.1. Determine the actions at s_i, s_d and $\{s_{i,j}\}_{j=1}^3$. For $0 \leq i < d$, (2.7) and (2.8) give

$$\pi_t(s_{i,j}) = \pi_0(s_{i,j}) = a_1, \quad \forall 1 \leq j \leq 3, \quad \pi_t(s_{i,j}) = \pi_0(s_{i,j}) = a_0, \quad \forall j \in \{4, 5\}.$$

Both actions at s_i , followed by π_t , reach s_{i+1} after $6L$ transitions. At s_d , the two L -transition paths end at s_0 and $\bar{s}_0$, whose values are equal by (2.10). By (2.4g), (2.9), and (3.7), we obtain

$$\Delta^{\pi_t}(s_i) = [\gamma^{4L}R_i(\gamma) + \gamma^{6L}\tilde{V}^{\pi_t}(s_{i+1})] - \gamma^{6L}\tilde{V}^{\pi_t}(s_{i+1}) = \gamma^{4L}R_i(\gamma) > 0, \quad \Delta^{\pi_t}(s_d) = -4f_{d-1}(\gamma) < 0.$$

At $s_{i,j}$, both actions followed by π_t reach s_{i+1} in $(6-j)L$ transitions, so the same comparison gives $\Delta^{\pi_t}(s_{i,j}) = \gamma^{(4-j)L}R_i(\gamma) > 0$ for $1 \leq j \leq 3$. We conclude that s_i and $\{s_{i,j}\}_{j=1}^3$ select a_1 for $0 \leq i < d$, while s_d switches to a_0 .

Step 1.2. Determine the actions at $\bar{s}_i, \{\bar{s}_{i,j}\}_{j=1}^4$ and in $\mathcal{S}_3$. For $s \in \{\bar{s}_{i,4} : 0 \leq i < d\} \cup \mathcal{S}_3$, (2.7) and (2.8) give $\pi_t(s) = a_0$. We compare the two action values after subtracting $\tilde{V}^{\pi_t}(\bar{s}_0)$.

- *Initial action a_0 .* By (2.4e), we have $\tilde{Q}^{\pi_t}(\bar{s}_{i,4}, a_0) = R_i(\gamma) + \gamma^{2L}\tilde{V}^{\pi_t}(\bar{s}_{i+1})$. By (3.6) and $R_i(\gamma) \rightarrow 0$, we have $\tilde{Q}^{\pi_t}(\bar{s}_{i,4}, a_0) - \tilde{V}^{\pi_t}(\bar{s}_0) \rightarrow 0$. By Table 1, the selected path from $s_{i,j}^r \in \mathcal{S}_3$ reaches $\bar{s}_0$ in $(2i+2)L$ transitions through either s_{-1}^c or s_{-1}^z . In the first case, the path has zero rewards until s_{-1}^c , whose path to $\bar{s}_0$ contributes $4f_{d-1}$ in the limit by (2.5). In the second case, each visited $s_{h,0}^r$ contributes f_h in the limit by (2.6a), giving a nonnegative sum bounded by $\sum_{h=0}^i f_h < 2f_{d-1}$ by (2.2). Combining these two cases with the preceding comparison at $\bar{s}_{i,4}$, we obtain from (3.4) and (3.6)

$$0 \leq \lim_{\gamma \rightarrow 1^-} (\tilde{Q}^{\pi_t}(s, a_0) - \tilde{V}^{\pi_t}(\bar{s}_0)) \leq 4f_{d-1}, \quad \forall s \in \{\bar{s}_{i,4} : 0 \leq i < d\} \cup \mathcal{S}_3. \quad (3.8)$$

- *Initial action a_1 .* By Table 1, the path from $\bar{s}_{i,4}$ ends at s_i . For the states in $\mathcal{S}_3$, the path from $s_{i,0}^r$ ends at s_{i+1} , while the paths from $s_{i,1}^r$ and $s_{i,2}^r$ both end at $s_{i,1}$. By (3.6), the values at s_i, s_{i+1} have the same limit as $\tilde{V}^{\pi_t}(\bar{s}_0)$. The identity $\tilde{V}^{\pi_t}(s_{i,1}) = \gamma^{3L}R_i(\gamma) + \gamma^{5L}\tilde{V}^{\pi_t}(s_{i+1})$ and $R_i(\gamma) \rightarrow 0$ give the same limit at $s_{i,1}$. Using the rewards in (2.4f) and (2.6b)–(2.6c), we obtain for $0 \leq i < d$

$$\lim_{\gamma \rightarrow 1^-} (\tilde{Q}^{\pi_t}(s, a_1) - \tilde{V}^{\pi_t}(\bar{s}_0)) = \begin{cases} -8f_{d-1}, & s = \bar{s}_{i,4}, \\ -8f_{d-1} - f_i, & s = s_{i,j}^r, \quad j \in \{0, 1, 2\}, \end{cases} \leq -8f_{d-1}. \quad (3.9)$$

Using (3.7), (3.8), and (3.9), we obtain

$$\lim_{\gamma \rightarrow 1^-} \Delta^{\pi_t}(s) \leq -8f_{d-1} < 0, \quad \forall s \in \{\bar{s}_{i,4} : 0 \leq i < d\} \cup \mathcal{S}_3.$$

We conclude that $\pi_{t+1}(s) = a_0$ for $s \in \{\bar{s}_{i,4} : 0 \leq i < d\} \cup \mathcal{S}_3$. We next show that each $\bar{s}_i$ keeps $a_{[\mathbf{b}]_i}$ and determine the actions selected at $\{\bar{s}_{i,j}\}_{j=1}^3$. By (2.8), π_t chooses $a_{[\mathbf{b}]_i}$ at $\bar{s}_i$ and $\{\bar{s}_{i,j}\}_{j=1}^3$.

- If $[\mathbf{b}]_i = 0$, action a_0 at $\bar{s}_i$ reaches $\bar{s}_{i+1}$ in $6L$ transitions. Action a_1 reaches $\bar{s}_{i,1}$ in L transitions, after which its current action a_0 reaches $\bar{s}_{i+1}$ in $5L$ transitions. Both paths have zero rewards by (2.4), and thus

$$\tilde{Q}^{\pi_t}(\bar{s}_i, a_1) = \gamma^L \tilde{V}^{\pi_t}(\bar{s}_{i,1}) = \gamma^{6L} \tilde{V}^{\pi_t}(\bar{s}_{i+1}) = \tilde{Q}^{\pi_t}(\bar{s}_i, a_0), \quad \Delta^{\pi_t}(\bar{s}_i) = 0.$$

At $\bar{s}_{i,3}$, both actions followed by π_t reach $\bar{s}_{i+1}$ in $3L$ transitions. The a_0 path has zero rewards, while the a_1 path passes through $\bar{s}_{i,4}$. For $j \in \{1, 2\}$, both actions at $\bar{s}_{i,j}$ followed by π_t reach $\bar{s}_{i+1}$ in $(6-j)L$ transitions with zero rewards. Using (2.4e) and (3.3), we obtain

$$\Delta^{\pi_t}(\bar{s}_{i,3}) = \gamma^L R_i(\gamma) > 0, \quad \Delta^{\pi_t}(\bar{s}_{i,j}) = 0, \quad \forall j \in \{1, 2\}.$$

We therefore find that $\bar{s}_{i,3}$ switches to a_1 , while $\bar{s}_i, \bar{s}_{i,1}, \bar{s}_{i,2}$ keep a_0 .

- If $[\mathbf{b}]_i = 1$, the a_1 path from $\bar{s}_i$ passes through $\bar{s}_{i,1}, \dots, \bar{s}_{i,4}$ and reaches $\bar{s}_{i+1}$ in $6L$ transitions. The a_0 path has the same length and endpoint but zero rewards. By (2.9) and (3.3), we obtain the action differences at $\bar{s}_i$ and $\{\bar{s}_{i,j}\}_{j=1}^3$:

$$\Delta^{\pi_t}(\bar{s}_i) = \gamma^{4L} R_i(\gamma) > 0, \quad \Delta^{\pi_t}(\bar{s}_{i,j}) = \gamma^{(4-j)L} R_i(\gamma) > 0, \quad \forall 1 \leq j \leq 3.$$

We conclude that $\bar{s}_i$ and $\{\bar{s}_{i,j}\}_{j=1}^3$ continue to choose a_1 .

If $[\mathbf{b}]_i = 0$, switching the action at $\bar{s}_{i,3}$ to a_1 leaves the selected a_0 path from $\bar{s}_i$ to $\bar{s}_{i+1}$ unchanged, because this path bypasses $\bar{s}_{i,3}$. Since the selected paths for $[\mathbf{b}]_i = 1$ also remain unchanged, we conclude that π_t and π_{t+1} generate the same trajectory and rewards from $\bar{s}_0$.

Step 1.3. Determine the actions at $s_{i,4}$ and $s_{i,5}$. By Table 1 and Figure 1, the actions at $s_{i,4}$ and $s_{i,5}$ determine which of s_{i+1}, s_i^z, s_i^c is reached from $s_{i,4}$ after $2L$ transitions. By (2.7) and (2.8), we have $\pi_t(s_j^c) = \pi_t(s_j^z) = a_{[\mathbf{b}]_j}$ for $0 \leq j < d$ and $\pi_t(s) = a_0$ for $s \in \mathcal{S}_3$. Under these choices, the paths from s_i^z and s_i^c reach $\bar{s}_0$ in $(2i+3)L$ transitions.

Determine $\pi_{t+1}(s_{i,5})$. By Table 1, actions a_0 and a_1 at $s_{i,5}$ reach s_{i+1} and s_i^z , respectively, in L transitions under any policy. By (2.5) and (2.6a), we have $\tilde{R}^{\pi_t}(s_j^z \rightsquigarrow s_{j-1}^z) = \gamma^L f_j(\gamma)(1 - [\mathbf{b}]_j)$ for $0 \leq j \leq i$. The final path from s_{-1}^z to $\bar{s}_0$ has zero rewards. Using (2.4c) and (2.4d), we obtain

$$\lim_{\gamma \rightarrow 1^-} \Delta^{\pi_t}(s_{i,5}) = \lim_{\gamma \rightarrow 1^-} \left[\gamma^L (\tilde{V}^{\pi_t}(s_i^z) - \tilde{V}^{\pi_t}(s_{i+1})) - f_i(\gamma) - \frac{1}{4} \right] \quad (3.10a)$$

$$= \lim_{\gamma \rightarrow 1^-} \left[\gamma^L (\tilde{R}^{\pi_t}(s_i^z \rightsquigarrow \bar{s}_0) + \gamma^{(2i+3)L} \tilde{V}^{\pi_t}(\bar{s}_0) - \tilde{V}^{\pi_t}(s_{i+1})) - f_i(\gamma) - \frac{1}{4} \right] \quad (3.10b)$$

$$= \sum_{j=0}^i f_j(1 - [\mathbf{b}]_j) - f_i - \frac{1}{4}. \quad (3.10c)$$

Here (3.10b) follows from (3.4), and (3.10c) uses the limits in (2.1) and (3.6). With k as defined in (3.1), the limit in (3.10c) equals $-1/4$ when $i = k$. If $i > k$ and $[\mathbf{b}]_i = 0$, it is at least $f_k - 1/4 > 0$. If $[\mathbf{b}]_i = 1$, it is at most $-5/4$ because $\sum_{j=0}^{i-1} f_j \leq f_i - 1$ by (2.2). We conclude that $\pi_{t+1}(s_{i,5}) = a_1$ exactly when $i > k$ and $[\mathbf{b}]_i = 0$.

Determine $\pi_{t+1}(s_{i,4})$. By (2.7) and (2.8), the current policy satisfies $\pi_t(s_{i,5}) = a_0$. For any policy π with $\pi(s_{i,5}) = a_0$, action a_0 at $s_{i,4}$ followed by π reaches s_{i+1} in $2L$ transitions with $\tilde{R}^\pi((s_{i,4}, a_0) \rightsquigarrow s_{i+1}) = R_i(\gamma)$. Action a_1 reaches s_i^c in the same number of transitions. For $i < k$, the path from s_i^c under π_t reaches s_{-1}^c with zero rewards because $[\mathbf{b}]_j = 1$ for $0 \leq j \leq i$. By (2.5), we obtain $\lim_{\gamma \rightarrow 1^-} \tilde{R}^{\pi_t}(s_i^c \rightsquigarrow \bar{s}_0) = 4f_{d-1}$. For $i \geq k$, let j be the largest index at most i with $[\mathbf{b}]_j = 0$. The paths from s_i^c and s_i^z reach $s_{j,0}^z$ after the same number of zero-reward transitions and then follow the same path to $\bar{s}_0$. Their discounted path rewards are therefore equal. Using (2.4a)–(2.4c), we obtain

$$\lim_{\gamma \rightarrow 1^-} \Delta^{\pi_t}(s_{i,4}) = \lim_{\gamma \rightarrow 1^-} \left[\gamma^{2L} (\tilde{V}^{\pi_t}(s_i^c) - \tilde{V}^{\pi_t}(s_{i+1})) + f_i(\gamma) - 4f_{d-1}(\gamma) - \frac{1}{2} - R_i(\gamma) \right] \quad (3.11a)$$

$$= \lim_{\gamma \rightarrow 1^-} \tilde{R}^{\pi_t}(s_i^c \rightsquigarrow \bar{s}_0) + f_i - 4f_{d-1} - \frac{1}{2} \quad (3.11b)$$

$$= \begin{cases} f_i - \frac{1}{2}, & i < k, \\ \sum_{j=0}^i f_j(1 - [\mathbf{b}]_j) + f_i - 4f_{d-1} - \frac{1}{2}, & i \geq k. \end{cases} \quad (3.11c)$$

Using (2.1), (3.4), and (3.6) with $R_i(\gamma) \rightarrow 0$, we obtain (3.11b). The limit in (3.11c) is positive for $i < k$. For $i \geq k$, it is less than $f_i - 2f_{d-1} - 1/2 < 0$, since $\sum_{j=0}^i f_j(1 - [\mathbf{b}]_j) < 2f_{d-1}$ by (2.2). We conclude that $\pi_{t+1}(s_{i,4}) = a_1$ for $i < k$ and a_0 otherwise.

Under π_{t+1} , the selected paths from $s_{i,4}$ reach s_i^c for $i < k$, s_i^z for $i > k$ with $[\mathbf{b}]_i = 0$, and s_{i+1} otherwise. They enter $\mathcal{S}_2$ exactly where $[\mathbf{b}^+]_i = 0$.

Step 1.4. Determine the actions at s_i^c and s_i^z . We have $\pi_t(s) = a_0$ for all $s \in \mathcal{S}_3$. We derive the comparison for any policy π with this property so that it can also be used in later iterations. At s_i^z , both actions followed by π reach s_{i-1}^z in $2L$ transitions. By (2.5), (2.6a), and (3.4), we obtain $\Delta^\pi(s_i^z) = 0 - \gamma^L f_i(\gamma) < 0$. By Table 1, either action at s_i^c followed by π reaches $\bar{s}_0$ in $(2i+3)L$ transitions through s_{-1}^c or s_{-1}^z , so (3.4) gives

$$\Delta^\pi(s_i^c) = \tilde{R}^\pi((s_i^c, a_1) \rightsquigarrow \bar{s}_0) - \tilde{R}^\pi((s_i^c, a_0) \rightsquigarrow \bar{s}_0). \quad (3.12)$$

- *Initial action a_0 at s_i^c .* By (2.5) and (2.6a), the first $2L$ -transition path reaches s_{i-1}^z with $\tilde{R}^\pi((s_i^c, a_0) \rightsquigarrow s_{i-1}^z) = \gamma^L f_i(\gamma)$. The path then passes through $s_{i-1}^z, \dots, s_{-1}^z$ to $\bar{s}_0$, collecting $\gamma^L f_j(\gamma)$ on the path from s_j^z to s_{j-1}^z whenever $\pi(s_j^z) = a_0$, and zero otherwise. Summing these discounted rewards and using (2.1), we

obtain

$$\lim_{\gamma \rightarrow 1^-} \tilde{R}^\pi((s_i^c, a_0) \rightsquigarrow \bar{s}_0) = f_i + \sum_{0 \leq j < i: \pi(s_j^z) = a_0} f_j \in [f_i, 2f_i - 1], \quad (3.13)$$

where the upper bound follows from $\sum_{j=0}^{i-1} f_j \leq f_i - 1$ in (2.2).

- *Initial action a_1 at s_i^c .* If $\pi(s_j^c) = a_1$ for all $0 \leq j < i$, the path reaches s_{-1}^c in $2(i+1)L$ zero-reward transitions, followed by the L -transition path to $\bar{s}_0$. By (2.5)–(2.6), we have $\tilde{R}^\pi((s_i^c, a_1) \rightsquigarrow \bar{s}_0) \rightarrow 4f_{d-1}$. Otherwise, let j be the first index encountered in $i-1, \dots, 0$ with $\pi(s_j^c) = a_0$. The path has zero rewards until s_j^c and then continues through $s_{j-1}^z, \dots, s_{-1}^z$ to $\bar{s}_0$. Using (2.2) and applying (3.13) at index j , we obtain

$$\lim_{\gamma \rightarrow 1^-} \tilde{R}^\pi((s_i^c, a_1) \rightsquigarrow \bar{s}_0) \leq \sum_{h=0}^j f_h \leq \sum_{h=0}^{i-1} f_h \leq f_i - 1. \quad (3.14)$$

Combining the two cases for initial action a_1 with (3.12)–(3.14), we obtain

$$\begin{cases} \lim_{\gamma \rightarrow 1^-} \Delta^\pi(s_i^c) \geq 4f_{d-1} - (2f_i - 1) > 0, & \text{if } \pi(s_j^c) = a_1 \text{ for all } 0 \leq j < i, \\ \lim_{\gamma \rightarrow 1^-} \Delta^\pi(s_i^c) \leq (f_i - 1) - f_i = -1 < 0, & \text{otherwise.} \end{cases} \quad (3.15)$$

By (2.8) and the definition of k , all s_j^c with $0 \leq j < i$ choose a_1 under π_t exactly when $i \leq k$. Using (3.15) and $\Delta^{\pi_t}(s_i^z) = -\gamma^L f_i(\gamma) < 0$, we obtain

$$\pi_{t+1}(s_i^c) = \begin{cases} a_1, & i \leq k, \\ a_0, & i > k, \end{cases} \quad \pi_{t+1}(s_i^z) = a_0, \quad \forall 0 \leq i < d.$$

3.2 Iteration 2: encode $\mathbf{b}^+$ in the first group

Under π_{t+1} , every s_i chooses a_1 , and the selected paths from $s_{i,4}$ enter $\mathcal{S}_2$ exactly where $[\mathbf{b}^+]_i = 0$. By Step 1.2 in Section 3.1, the paths from $\bar{s}_i$ remain those of $\pi[\mathbf{b}]$, so $\tilde{V}^{\pi_{t+1}}(\bar{s}_i) = \tilde{V}^{\pi[\mathbf{b}]}(\bar{s}_i)$ for $0 \leq i \leq d$.

Step 2.1. Determine the actions at s_i and $\{s_{i,j}\}_{j=1}^3$. For any policy π satisfying $\pi(s_{i,j}) = a_1$ for $1 \leq j \leq 3$, action a_0 at s_i reaches s_{i+1} in $6L$ transitions, while action a_1 followed by π reaches $s_{i,4}$ in $4L$ transitions. Both paths have zero rewards by (2.4), so we obtain

$$\Delta^\pi(s_i) = \gamma^{4L}(\tilde{V}^\pi(s_{i,4}) - \gamma^{2L}\tilde{V}^\pi(s_{i+1})), \quad \Delta^\pi(s_{i,j}) = \gamma^{-jL}\Delta^\pi(s_i), \quad \forall 1 \leq j \leq 3. \quad (3.16)$$

The same identities hold for $\bar{s}_i, \bar{s}_{i,j}$ when $\pi(\bar{s}_{i,j}) = a_1$ for $1 \leq j \leq 3$. By Step 1.1 in Section 3.1, we have $\pi_{t+1}(s_{i,j}) = a_1$ for $1 \leq j \leq 3$, so (3.16) applies. For $[\mathbf{b}^+]_i = 0$, we first compute $\tilde{V}^{\pi_{t+1}}(s_{i,4})$ from the selected paths. (I) If $i < k$, then $\pi_{t+1}(s_j^c) = a_1$ for $0 \leq j \leq i$, so the path continues through $s_i^c, \dots, s_{-1}^c$ to $\bar{s}_0$. By (2.4b) and (2.5), we have $\tilde{R}^{\pi_{t+1}}(s_{i,4} \rightsquigarrow \bar{s}_0) \rightarrow f_i - 4f_{d-1} - 1/2 + 4f_{d-1} = f_i - 1/2$. (II) If $i > k$, then $[\mathbf{b}]_i = 0$, and the path enters s_i^z . Since every s_j^z chooses a_0 , we obtain from (2.4a), (2.4d), and (2.6a) with $R_i(\gamma) \rightarrow 0$ that $\tilde{R}^{\pi_{t+1}}(s_{i,4} \rightsquigarrow \bar{s}_0) \rightarrow -f_i - 1/4 + \sum_{j=0}^i f_j = \sum_{j=0}^{i-1} f_j - 1/4$. The zero-reward path from s_i to $s_{i,4}$ gives $\tilde{V}^{\pi_{t+1}}(s_i) = \gamma^{4L}\tilde{V}^{\pi_{t+1}}(s_{i,4})$. By (3.4) and (3.6), we therefore obtain

$$\lim_{\gamma \rightarrow 1^-} (\tilde{V}^{\pi_{t+1}}(s_{i,4}) - \tilde{V}^{\pi_{t+1}}(\bar{s}_0)) = \lim_{\gamma \rightarrow 1^-} (\tilde{V}^{\pi_{t+1}}(s_i) - \tilde{V}^{\pi_{t+1}}(\bar{s}_0)) = \begin{cases} f_i - \frac{1}{2}, & i < k, \\ \sum_{j=0}^{i-1} f_j - \frac{1}{4}, & i > k, [\mathbf{b}]_i = 0. \end{cases} \quad (3.17)$$

To determine the sign of $\Delta^{\pi_{t+1}}(s_i)$ in (3.16), we next bound the limit of $\tilde{V}^{\pi_{t+1}}(s_{i+1}) - \tilde{V}^{\pi_{t+1}}(\bar{s}_0)$. Let j be the smallest index with $i < j < d$ and $[\mathbf{b}^+]_j = 0$, taking $j = d$ if no such index exists. By (3.3), the selected path from s_{i+1} to s_j under π_{t+1} has $6(j-i-1)L$ transitions and scaled reward $\sum_{h=i+1}^{j-1} \gamma^{(6(h-i-1)+4)L} R_h(\gamma) \rightarrow 0$. If $j < d$, the limit of $\tilde{V}^{\pi_{t+1}}(s_j) - \tilde{V}^{\pi_{t+1}}(\bar{s}_0)$ is at least $\sum_{h=0}^{j-1} f_h - 1/4 \geq f_i - 1/4$ and at most $4f_{d-1}$ by (2.2) and (3.17). If $j = d$, the difference tends to $4f_{d-1}$ by (2.4g). Thus, we obtain

$$f_i - \frac{1}{4} \leq \lim_{\gamma \rightarrow 1^-} (\tilde{V}^{\pi_{t+1}}(s_{i+1}) - \tilde{V}^{\pi_{t+1}}(\bar{s}_0)) \leq 4f_{d-1}, \quad \forall 0 \leq i < d. \quad (3.18)$$

For $[\mathbf{b}^+]_i = 0$, both cases in (3.17) are at most $\mathbf{f}_i - 1/2$ by (2.2). Combining (3.16) with (3.18), we obtain

$$\lim_{\gamma \rightarrow 1^-} \Delta^{\pi_{t+1}}(s_i) \leq (\mathbf{f}_i - \frac{1}{2}) - (\mathbf{f}_i - \frac{1}{4}) = -\frac{1}{4} < 0, \quad \pi_{t+2}(s_i) = a_0.$$

For $[\mathbf{b}^+]_i = 1$, the selected path from $s_{i,4}$ reaches s_{i+1} , so we obtain

$$\Delta^{\pi_{t+1}}(s_i) = \gamma^{4L} R_i(\gamma) > 0, \quad \pi_{t+2}(s_i) = a_1.$$

By (3.16), we also obtain $\pi_{t+2}(s_{i,j}) = a_{[\mathbf{b}^+]_i}$ for $1 \leq j \leq 3$.

Step 2.2. Restore a_0 at $s_{i,4}, s_{i,5}$. By Steps 1.2 and 1.4 in Section 3.1, π_{t+1} selects a_0 at every s_j^z and every state in $\mathcal{S}_3$. By Table 1 and (3.13), we have $\tilde{R}^{\pi_{t+1}}(s_i^z \rightsquigarrow \bar{s}_0) \rightarrow \sum_{j=0}^i \mathbf{f}_j$, and the path estimates in the proof of (3.15) give $\lim_{\gamma \rightarrow 1^-} \tilde{R}^{\pi_{t+1}}(s_i^c \rightsquigarrow \bar{s}_0) \leq 4\mathbf{f}_{d-1}$. Both paths reach $\bar{s}_0$ in $(2i+3)L$ transitions, so we obtain from (2.2), (3.4), and (3.6)

$$\lim_{\gamma \rightarrow 1^-} (\tilde{V}^{\pi_{t+1}}(s_i^c) - \tilde{V}^{\pi_{t+1}}(\bar{s}_0)) \leq 4\mathbf{f}_{d-1}, \quad \lim_{\gamma \rightarrow 1^-} (\tilde{V}^{\pi_{t+1}}(s_i^z) - \tilde{V}^{\pi_{t+1}}(\bar{s}_0)) = \sum_{j=0}^i \mathbf{f}_j \leq 2\mathbf{f}_i - 1. \quad (3.19)$$

Both actions at $s_{i,5}$ follow fixed L -transition paths, so the comparison in (3.10a) also applies under π_{t+1} . By (3.18) and (3.19), we obtain

$$\begin{aligned} \lim_{\gamma \rightarrow 1^-} \Delta^{\pi_{t+1}}(s_{i,5}) &= \lim_{\gamma \rightarrow 1^-} (\tilde{V}^{\pi_{t+1}}(s_i^z) - \tilde{V}^{\pi_{t+1}}(\bar{s}_0)) - \lim_{\gamma \rightarrow 1^-} (\tilde{V}^{\pi_{t+1}}(s_{i+1}) - \tilde{V}^{\pi_{t+1}}(\bar{s}_0)) - \mathbf{f}_i - \frac{1}{4} \\ &\leq (2\mathbf{f}_i - 1) - (\mathbf{f}_i - \frac{1}{4}) - \mathbf{f}_i - \frac{1}{4} = -1 < 0. \end{aligned}$$

If $\pi_{t+1}(s_{i,5}) = a_0$, the path comparison in (3.11a) applies under π_{t+1} . Using (3.18) and (3.19) with $R_i(\gamma) \rightarrow 0$, we obtain

$$\begin{aligned} \lim_{\gamma \rightarrow 1^-} \Delta^{\pi_{t+1}}(s_{i,4}) &= \lim_{\gamma \rightarrow 1^-} (\tilde{V}^{\pi_{t+1}}(s_i^c) - \tilde{V}^{\pi_{t+1}}(\bar{s}_0)) - \lim_{\gamma \rightarrow 1^-} (\tilde{V}^{\pi_{t+1}}(s_{i+1}) - \tilde{V}^{\pi_{t+1}}(\bar{s}_0)) + \mathbf{f}_i - 4\mathbf{f}_{d-1} - \frac{1}{2} \\ &\leq 4\mathbf{f}_{d-1} - (\mathbf{f}_i - \frac{1}{4}) + \mathbf{f}_i - 4\mathbf{f}_{d-1} - \frac{1}{2} = -\frac{1}{4} < 0. \end{aligned}$$

If $\pi_{t+1}(s_{i,5}) = a_1$, Step 1.3 in Section 3.1 gives $i > k$, and Step 1.4 gives $\pi_{t+1}(s_i^c) = \pi_{t+1}(s_i^z) = a_0$. By Table 1 and (2.5), the selected paths from s_i^c and s_i^z reach $s_{i,0}^r$ in L transitions with zero rewards, giving $\tilde{V}^{\pi_{t+1}}(s_i^c) = \gamma^L \tilde{V}^{\pi_{t+1}}(s_{i,0}^r) = \tilde{V}^{\pi_{t+1}}(s_i^z)$. From $s_{i,4}$, action a_1 reaches s_i^c in $2L$ transitions, whereas action a_0 followed by π_{t+1} reaches s_i^z through $s_{i,5}$ in $2L$ transitions. By (2.4a), (2.4b), and (2.4d), we obtain

$$\lim_{\gamma \rightarrow 1^-} \Delta^{\pi_{t+1}}(s_{i,4}) = \lim_{\gamma \rightarrow 1^-} \left\{ [\mathbf{f}_i(\gamma) - 4\mathbf{f}_{d-1}(\gamma) - \frac{1}{2}] - [R_i(\gamma) - \gamma^L(\mathbf{f}_i(\gamma) + \frac{1}{4})] \right\} = -4\mathbf{f}_{d-1} + 2\mathbf{f}_i - \frac{1}{4} < 0.$$

We conclude that $\pi_{t+2}(s_{i,4}) = \pi_{t+2}(s_{i,5}) = a_0$ for $0 \leq i < d$.

Step 2.3. Determine the actions at $s_d, \bar{s}_{i,4}$ and in $\mathcal{S}_3$. We first show that $\pi_{t+2}(s) = a_0$ for $s \in \{\bar{s}_{i,4} : 0 \leq i < d\} \cup \mathcal{S}_3$. By Table 1, the a_1 paths from $\bar{s}_{i,4}$ and the states in $\mathcal{S}_3$ end at $s_i, s_{i,1}, s_{i+1}$. By Step 1.1 in Section 3.1, we have $\tilde{V}^{\pi_{t+1}}(s_i) = \gamma^L \tilde{V}^{\pi_{t+1}}(s_{i,1}) = \gamma^{4L} \tilde{V}^{\pi_{t+1}}(s_{i,4})$. For $[\mathbf{b}^+]_i = 0$, (2.2) and (3.17) bound the limit of $\tilde{V}^{\pi_{t+1}}(s_{i,4}) - \tilde{V}^{\pi_{t+1}}(\bar{s}_0)$ by $\mathbf{f}_i - 1/2 < 4\mathbf{f}_{d-1}$. For $[\mathbf{b}^+]_i = 1$, the selected path gives $\tilde{V}^{\pi_{t+1}}(s_{i,4}) = R_i(\gamma) + \gamma^{2L} \tilde{V}^{\pi_{t+1}}(s_{i+1})$. Using (3.18) and $R_i(\gamma) \rightarrow 0$, we obtain

$$\lim_{\gamma \rightarrow 1^-} (\tilde{V}^{\pi_{t+1}}(s) - \tilde{V}^{\pi_{t+1}}(\bar{s}_0)) \leq 4\mathbf{f}_{d-1}, \quad \forall s \in \{s_i, s_{i,1}, s_{i+1}\}, \quad \forall 0 \leq i < d. \quad (3.20)$$

By Step 1.2 in Section 3.1, we have $\pi_{t+1}(s) = a_0$ for $s \in \{\bar{s}_{i,4} : 0 \leq i < d\} \cup \mathcal{S}_3$, and the trajectory from $\bar{s}_0$ is unchanged. The path reward estimates in the proofs of (3.8) and (3.9) therefore give $\lim_{\gamma \rightarrow 1^-} \tilde{R}^{\pi_{t+1}}(s \rightsquigarrow \bar{s}_0) \geq 0$ and $\lim_{\gamma \rightarrow 1^-} \tilde{R}^{\pi_{t+1}}((s, a_1) \rightsquigarrow s') \leq -8\mathbf{f}_{d-1}$, where $s' \in \{s_i, s_{i,1}, s_{i+1}\}$ for some $0 \leq i < d$. Using (3.4) and (3.20), we obtain

$$\lim_{\gamma \rightarrow 1^-} \Delta^{\pi_{t+1}}(s) \leq (-8\mathbf{f}_{d-1} + 4\mathbf{f}_{d-1}) - 0 = -4\mathbf{f}_{d-1} < 0, \quad \forall s \in \{\bar{s}_{i,4} : 0 \leq i < d\} \cup \mathcal{S}_3. \quad (3.21)$$

To determine $\pi_{t+2}(s_d)$, let $j = \min\{0 \leq i < d : [\mathbf{b}^+]_i = 0\}$. Under π_{t+1} , the selected path from s_0 to s_j takes $6jL$ transitions. By (3.3) and (3.4), we have

$$\tilde{V}^{\pi_{t+1}}(s_0) - \tilde{V}^{\pi_{t+1}}(s_j) = \sum_{i=0}^{j-1} \gamma^{(6i+4)L} R_i(\gamma) + (\gamma^{6jL} - 1) \tilde{V}^{\pi_{t+1}}(s_j) \longrightarrow 0,$$

where $R_i(\gamma) \rightarrow 0$ by (3.3), and $\tilde{V}^{\pi_{t+1}}(s_j)$ has a finite limit by (3.6) and (3.17). Combining this limit with (2.2), (2.4g), and (3.17), we obtain

$$\begin{aligned} \lim_{\gamma \rightarrow 1^-} \Delta^{\pi_{t+1}}(s_d) &= \lim_{\gamma \rightarrow 1^-} (\tilde{V}^{\pi_{t+1}}(s_j) - \tilde{V}^{\pi_{t+1}}(\bar{s}_0)) - 4f_{d-1} \\ &\leq f_j - \frac{1}{2} - 4f_{d-1} < 0. \end{aligned}$$

We conclude that $\pi_{t+2}(s) = a_0$ for $s \in \{s_d\} \cup \{\bar{s}_{i,4} : 0 \leq i < d\} \cup \mathcal{S}_3$.

Step 2.4. Determine the actions at $\bar{s}_i, \{\bar{s}_{i,j}\}_{j=1}^3, s_i^c, s_i^z$. For $[\mathbf{b}]_i = 0$, Step 1.2 in Section 3.1 gives $\pi_{t+1}(\bar{s}_i) = \pi_{t+1}(\bar{s}_{i,1}) = \pi_{t+1}(\bar{s}_{i,2}) = a_0$ and $\pi_{t+1}(\bar{s}_{i,3}) = a_1$. Both actions at $\bar{s}_i$, followed by π_{t+1} , reach $\bar{s}_{i+1}$ in $6L$ zero-reward transitions, and both actions at $\bar{s}_{i,1}$ do so in $5L$ transitions. At $\bar{s}_{i,2}$, the a_1 path now passes through $\bar{s}_{i,3}, \bar{s}_{i,4}$, so (2.4e) gives

$$\Delta^{\pi_{t+1}}(\bar{s}_i) = \Delta^{\pi_{t+1}}(\bar{s}_{i,1}) = 0, \quad \Delta^{\pi_{t+1}}(\bar{s}_{i,2}) = \gamma^{2L} R_i(\gamma) > 0.$$

The same path comparison gives $\Delta^{\pi_{t+1}}(\bar{s}_{i,3}) = \gamma^L R_i(\gamma) > 0$ for every i . When $[\mathbf{b}]_i = 1$, the positive action differences at $\bar{s}_i, \bar{s}_{i,1}, \bar{s}_{i,2}$ from Step 1.2 are unchanged. We therefore obtain

$$\pi_{t+2}(\bar{s}_i) = \pi_{t+2}(\bar{s}_{i,1}) = a_{[\mathbf{b}]_i}, \quad \pi_{t+2}(\bar{s}_{i,2}) = \pi_{t+2}(\bar{s}_{i,3}) = a_1, \quad \forall 0 \leq i < d.$$

If $[\mathbf{b}]_i = 0$, the unchanged a_0 path from $\bar{s}_i$ bypasses $\bar{s}_{i,2}$. By Step 2.3, $\bar{s}_{i,4}$ also keeps a_0 , so π_{t+1} and π_{t+2} generate the same trajectory and rewards from $\bar{s}_0$.

By Steps 1.2 and 1.4 in Section 3.1, π_{t+1} selects a_0 at every state in $\mathcal{S}_3$, and all s_j^c with $0 \leq j < i$ select a_1 exactly when $i \leq k+1$. Using (3.15) and $\Delta^{\pi_{t+1}}(s_i^z) = -\gamma^L f_i(\gamma) < 0$, we obtain $\pi_{t+2}(s_i^c) = a_1$ exactly when $i \leq k+1$, and $\pi_{t+2}(s_i^z) = a_0$ for $0 \leq i < d$.

3.3 Iteration 3: form the new cycle

Under π_{t+2} , the selected paths from s_i to s_{i+1} encode $\mathbf{b}^+$ and lead from s_0 to s_d , where a_0 still leads to $\bar{s}_0$. By Step 2.4 in Section 3.2, the paths from $\bar{s}_i$ remain those of $\pi[\mathbf{b}]$, so $\tilde{V}^{\pi_{t+2}}(\bar{s}_0) = \tilde{V}^{\pi[\mathbf{b}]}(\bar{s}_0)$.

Step 3.1. Select a_1 at s_d . Starting from s_d , action a_1 followed by π_{t+2} returns to s_d after $(6d+1)L$ transitions. This path has the same length and reward sequence as one cycle starting from s_d under $\pi[\mathbf{b}^+]$, so $\tilde{R}^{\pi_{t+2}}((s_d, a_1) \rightsquigarrow s_d) = (1 - \gamma^{(6d+1)L}) \tilde{V}^{\pi[\mathbf{b}^+]}(s_d)$. Since $\pi_{t+2}(s_d) = a_0$, we apply (3.4) to obtain

$$\begin{aligned} \Delta^{\pi_{t+2}}(s_d) &= [(1 - \gamma^{(6d+1)L}) \tilde{V}^{\pi[\mathbf{b}^+]}(s_d) + \gamma^{(6d+1)L} \tilde{V}^{\pi_{t+2}}(s_d)] - \tilde{V}^{\pi_{t+2}}(s_d) \\ &= (1 - \gamma^{(6d+1)L}) (\tilde{V}^{\pi[\mathbf{b}^+]}(s_d) - \tilde{V}^{\pi_{t+2}}(s_d)). \end{aligned} \tag{3.22}$$

By (2.4g), $\tilde{V}^{\pi_{t+2}}(s_d) = 4f_{d-1}(\gamma) + \gamma^L \tilde{V}^{\pi[\mathbf{b}]}(\bar{s}_0)$. We combine this identity with (2.11) and (3.6) to obtain

$$\lim_{\gamma \rightarrow 1^-} (\tilde{V}^{\pi[\mathbf{b}^+]}(s_d) - \tilde{V}^{\pi_{t+2}}(s_d)) = 4 \left(\mathbf{g}_k - \sum_{j=0}^{k-1} \mathbf{g}_j \right) - 4f_{d-1} \geq 12f_{d-1} > 0, \tag{3.23}$$

where we use $\sum_{j=0}^{k-1} \mathbf{g}_j \leq \mathbf{g}_k - \mathbf{g}_0$ and $\mathbf{g}_0 \geq 4f_{d-1}$ in (2.2). Combining (3.22) and (3.23), we obtain $\pi_{t+3}(s_d) = a_1$.

Step 3.2. Keep a_0 at $s_{i,4}, s_{i,5}, \bar{s}_{i,4}$ and in $\mathcal{S}_3$. By (3.3), the selected path from s_i to s_d satisfies

$$\tilde{R}^{\pi_{t+2}}(s_i \rightsquigarrow s_d) = \sum_{h=i}^{d-1} [\mathbf{b}^+]_h \gamma^{(6(h-i)+4)L} R_h(\gamma) \longrightarrow 0, \quad \forall 0 \leq i < d.$$

We also have $\tilde{V}^{\pi_{t+2}}(s_i) = \gamma^L \tilde{V}^{\pi_{t+2}}(s_{i,1})$. When $[\mathbf{b}^+]_i = 0$, this follows from the zero-reward paths of lengths $6L$ and $5L$ from s_i and $s_{i,1}$ to s_{i+1} . Using (3.4) and (3.6) with the value at s_d computed above, we obtain

$$\lim_{\gamma \rightarrow 1^-} (\tilde{V}^{\pi_{t+2}}(s) - \tilde{V}^{\pi_{t+2}}(\bar{s}_0)) = 4f_{d-1}, \quad \forall s \in \{s_i, s_{i,1}, s_{i+1}\}, \quad \forall 0 \leq i < d. \quad (3.24)$$

By Steps 2.3 and 2.4 in Section 3.2, π_{t+2} selects a_0 at $\bar{s}_{i,4}, s_i^z$ and every state in $\mathcal{S}_3$. The path reward bounds used in (3.19) and (3.21) therefore remain valid. Applying the proof of (3.21) with (3.24), we obtain

$$\lim_{\gamma \rightarrow 1^-} \Delta^{\pi_{t+2}}(s) \leq -4f_{d-1} < 0, \quad \forall s \in \{\bar{s}_{i,4} : 0 \leq i < d\} \cup \mathcal{S}_3.$$

We also use the bounds in (3.19) under π_{t+2} to compare the actions at $s_{i,5}$ and $s_{i,4}$. Since $\pi_{t+2}(s_{i,5}) = a_0$ by Step 2.2 in Section 3.2, the comparisons in (3.10a) and (3.11a) apply under π_{t+2} . Using (3.24), we obtain

$$\begin{aligned} \lim_{\gamma \rightarrow 1^-} \Delta^{\pi_{t+2}}(s_{i,5}) &\leq (2f_i - 1) - 4f_{d-1} - f_i - \frac{1}{4} = f_i - 4f_{d-1} - \frac{5}{4} < 0, \\ \lim_{\gamma \rightarrow 1^-} \Delta^{\pi_{t+2}}(s_{i,4}) &\leq 4f_{d-1} - 4f_{d-1} + f_i - 4f_{d-1} - \frac{1}{2} = f_i - 4f_{d-1} - \frac{1}{2} < 0. \end{aligned}$$

We conclude that $\pi_{t+3}(s_{i,4}) = \pi_{t+3}(s_{i,5}) = \pi_{t+3}(\bar{s}_{i,4}) = a_0$ for $0 \leq i < d$, and $\pi_{t+3}(s) = a_0$ for $s \in \mathcal{S}_3$.

Step 3.3. Determine the actions at $s_i, \bar{s}_i, \{s_{i,j}, \bar{s}_{i,j}\}_{j=1}^3, s_i^c, s_i^z$. The current policy satisfies $\pi_{t+2}(s_{i,4}) = \pi_{t+2}(s_{i,5}) = \pi_{t+2}(\bar{s}_{i,4}) = a_0$. Thus the selected $2L$ -transition paths from $s_{i,4}$ to s_{i+1} and from $\bar{s}_{i,4}$ to $\bar{s}_{i+1}$ have the same reward sequence, and the equal-length path comparison in Step 1.2 in Section 3.1 applies to both groups. For $[\mathbf{b}^+]_i = 0$, we obtain

$$\Delta^{\pi_{t+2}}(s_i) = \Delta^{\pi_{t+2}}(s_{i,1}) = \Delta^{\pi_{t+2}}(s_{i,2}) = 0, \quad \Delta^{\pi_{t+2}}(s_{i,3}) = \gamma^L R_i(\gamma) > 0.$$

We therefore obtain $\pi_{t+3}(s_{i,3}) = a_1$, while $s_i, s_{i,1}, s_{i,2}$ keep a_0 . For $[\mathbf{b}^+]_i = 1$, the positive action differences keep a_1 at s_i and $\{s_{i,j}\}_{j=1}^3$. Using the choices from Step 2.4 in Section 3.2, we obtain for $[\mathbf{b}]_i = 0$

$$\Delta^{\pi_{t+2}}(\bar{s}_i) = 0, \quad \Delta^{\pi_{t+2}}(\bar{s}_{i,1}) = \gamma^{3L} R_i(\gamma) > 0.$$

We obtain $\pi_{t+3}(\bar{s}_i) = a_0$ and $\pi_{t+3}(\bar{s}_{i,1}) = a_1$. The positive action differences keep a_1 at $\bar{s}_{i,2}, \bar{s}_{i,3}$ for every i , and at $\bar{s}_i, \bar{s}_{i,1}$ when $[\mathbf{b}]_i = 1$. The selected a_0 path from s_i when $[\mathbf{b}^+]_i = 0$ bypasses $s_{i,3}$, and that from $\bar{s}_i$ when $[\mathbf{b}]_i = 0$ bypasses $\bar{s}_{i,1}$, so these intermediate changes do not alter either encoded path.

By Step 2.4 in Section 3.2, all s_j^c with $0 \leq j < i$ select a_1 under π_{t+2} exactly when $i \leq k+2$. Applying (3.15) to π_{t+2} and using $\Delta^{\pi_{t+2}}(s_i^z) = -\gamma^L f_i(\gamma) < 0$, we obtain $\pi_{t+3}(s_i^c) = a_1$ exactly when $i \leq k+2$, and $\pi_{t+3}(s_i^z) = a_0$ for $0 \leq i < d$.

3.4 Iteration 4: connect to the new cycle

Under π_{t+3} , the cycles through s_0 and $\bar{s}_0$ encode $\mathbf{b}^+$ and $\mathbf{b}$, respectively. The selected paths therefore give

$$\tilde{V}^{\pi_{t+3}}(s_i) = \tilde{V}^{\pi[\mathbf{b}^+]}(s_i), \quad \tilde{V}^{\pi_{t+3}}(\bar{s}_i) = \tilde{V}^{\pi[\mathbf{b}]}(\bar{s}_i), \quad \forall 0 \leq i \leq d.$$

By (2.2), (2.11), and (3.6), we obtain

$$\lim_{\gamma \rightarrow 1^-} (\tilde{V}^{\pi_{t+3}}(s_i) - \tilde{V}^{\pi_{t+3}}(\bar{s}_0)) = 4 \left(\mathbf{g}_k - \sum_{j=0}^{k-1} \mathbf{g}_j \right) \geq 16f_{d-1}, \quad \forall 0 \leq i \leq d. \quad (3.25)$$

Step 4.1. Select a_1 at $\bar{s}_i, \{\bar{s}_{i,j}\}_{j=1}^4$ and in $\mathcal{S}_3$. For $0 \leq i < d$ and $s \in \{\bar{s}_{i,4}, s_{i,0}^r, s_{i,1}^r, s_{i,2}^r\}$, we have $\pi_{t+3}(s) = a_0$ by Step 3.2 in Section 3.3. By Table 1, action a_0 followed by π_{t+3} reaches $\bar{s}_0$, whereas the a_1 path ends at $s_i, s_{i,1}$, or s_{i+1} . The selected path from $s_{i,1}$ gives $\tilde{V}^{\pi_{t+3}}(s_{i,1}) = [\mathbf{b}^+]_i \gamma^{3L} R_i(\gamma) + \gamma^{5L} \tilde{V}^{\pi_{t+3}}(s_{i+1})$. Using $R_i(\gamma) \rightarrow 0$ and the path reward bounds from the proofs of (3.8) and (3.9), we obtain from (3.4) and (3.25)

$$\begin{cases} \lim_{\gamma \rightarrow 1^-} (\tilde{Q}^{\pi_{t+3}}(s, a_1) - \tilde{V}^{\pi_{t+3}}(\bar{s}_0)) \geq 16f_{d-1} - 8f_{d-1} - f_i, \\ \lim_{\gamma \rightarrow 1^-} (\tilde{Q}^{\pi_{t+3}}(s, a_0) - \tilde{V}^{\pi_{t+3}}(\bar{s}_0)) \leq 4f_{d-1}. \end{cases} \quad (3.26)$$

Using (3.7) and (3.26), we obtain

$$\lim_{\gamma \rightarrow 1^-} \Delta^{\pi_{t+3}}(s) \geq 16f_{d-1} - (8f_{d-1} + f_i) - 4f_{d-1} = 4f_{d-1} - f_i > 0.$$

By Steps 3.2 and 3.3 in Section 3.3, we have $\pi_{t+3}(\bar{s}_{i,4}) = a_0$ and $\pi_{t+3}(\bar{s}_{i,j}) = a_1$ for $1 \leq j \leq 3$. The path rewards in (3.3) and the comparisons in (3.16) give

$$\Delta^{\pi_{t+3}}(\bar{s}_i) = \gamma^{4L} R_i(\gamma) > 0, \quad \Delta^{\pi_{t+3}}(\bar{s}_{i,j}) = \gamma^{(4-j)L} R_i(\gamma) > 0, \quad \forall 1 \leq j \leq 3.$$

We conclude that π_{t+4} selects a_1 at $\bar{s}_i$, $\{\bar{s}_{i,j}\}_{j=1}^4$, and every state in $\mathcal{S}_3$.

Step 4.2. Determine the actions at $s_d, s_i, \{s_{i,j}\}_{j=1}^5, s_i^c, s_i^z$. By Steps 3.2 and 3.3 in Section 3.3, π_{t+3} selects a_0 at $s_{i,4}, s_{i,5}, s_i^z$ and every state in $\mathcal{S}_3$. The bounds in (3.19) therefore remain valid, and the comparisons in (3.10a) and (3.11a) apply under π_{t+3} . Using (3.25), we obtain for $0 \leq i < d$

$$\lim_{\gamma \rightarrow 1^-} \Delta^{\pi_{t+3}}(s) \leq f_i - \frac{1}{2} - 16f_{d-1} < 0, \quad \pi_{t+4}(s) = a_0, \quad \forall s \in \{s_{i,4}, s_{i,5}\}.$$

At s_d , we obtain from (2.4g) and (3.25)

$$\lim_{\gamma \rightarrow 1^-} \Delta^{\pi_{t+3}}(s_d) = \lim_{\gamma \rightarrow 1^-} (\tilde{V}^{\pi_{t+3}}(s_0) - \tilde{V}^{\pi_{t+3}}(\bar{s}_0)) - 4f_{d-1} \geq 12f_{d-1} > 0, \quad \pi_{t+4}(s_d) = a_1.$$

For $[\mathbf{b}^+]_i = 0$, Step 3.3 in Section 3.3 gives $\pi_{t+3}(s_i) = \pi_{t+3}(s_{i,1}) = \pi_{t+3}(s_{i,2}) = a_0$ and $\pi_{t+3}(s_{i,3}) = a_1$. The path comparison at $\bar{s}_i, \{\bar{s}_{i,j}\}_{j=1}^3$ in Step 2.4 in Section 3.2 therefore gives

$$\Delta^{\pi_{t+3}}(s_i) = \Delta^{\pi_{t+3}}(s_{i,1}) = 0, \quad \Delta^{\pi_{t+3}}(s_{i,2}) = \gamma^{2L} R_i(\gamma) > 0.$$

We conclude that $s_{i,2}$ switches to a_1 , while $s_i, s_{i,1}$ keep a_0 . The same comparison gives $\Delta^{\pi_{t+3}}(s_{i,3}) = \gamma^L R_i(\gamma) > 0$ for every i , and keeps a_1 at s_i and $\{s_{i,j}\}_{j=1}^3$ when $[\mathbf{b}^+]_i = 1$. When $[\mathbf{b}^+]_i = 0$, the selected a_0 path from s_i bypasses $s_{i,2}$, so the switch at $s_{i,2}$ does not alter the cycle through s_0 .

By Step 3.3 in Section 3.3, all s_j^c with $0 \leq j < i$ select a_1 under π_{t+3} exactly when $i \leq k+3$. Applying (3.15) and using $\Delta^{\pi_{t+3}}(s_i^z) = -\gamma^L f_i(\gamma) < 0$, we obtain $\pi_{t+4}(s_i^c) = a_1$ exactly when $i \leq k+3$, and $\pi_{t+4}(s_i^z) = a_0$ for $0 \leq i < d$.

3.5 Iteration 5: encode $\mathbf{b}^+$ in the second group

By Steps 4.1 and 4.2 in Section 3.4, the cycle through s_0 under π_{t+4} still encodes $\mathbf{b}^+$, and every $\bar{s}_{i,4}$ and every state in $\mathcal{S}_3$ chooses a_1 .

Step 5.1. Copy $\mathbf{b}^+$ to $\bar{s}_i, \{\bar{s}_{i,j}\}_{j=1}^3, s_i^c, s_i^z$. We express the action differences at $\bar{s}_i, s_i^c, s_i^z$ in terms of $\Delta^{\pi_{t+4}}(s_i)$. The unchanged cycle gives $\tilde{V}^{\pi_{t+4}}(s_i) = \tilde{V}^{\pi[\mathbf{b}^+]}(s_i)$ for $0 \leq i \leq d$. For $[\mathbf{b}^+]_i = 0$, Step 4.2 in Section 3.4 gives $\pi_{t+4}(s_{i,1}) = a_0$, so both actions at s_i followed by π_{t+4} reach s_{i+1} in $6L$ transitions with zero rewards. For $[\mathbf{b}^+]_i = 1$, action a_1 follows the full horizontal path. Since $\tilde{Q}^{\pi_{t+4}}(s_i, a_1) = \tilde{V}^{\pi_{t+4}}(s_i)$ in both cases, we obtain from (3.3)

$$\tilde{V}^{\pi_{t+4}}(s_i) - \gamma^{6L} \tilde{V}^{\pi_{t+4}}(s_{i+1}) = \Delta^{\pi_{t+4}}(s_i) = [\mathbf{b}^+]_i \gamma^{4L} R_i(\gamma), \quad \forall 0 \leq i < d. \quad (3.27)$$

By Step 4.1 in Section 3.4, π_{t+4} selects a_1 at $\bar{s}_i$ and $\{\bar{s}_{i,j}\}_{j=1}^4$. The path from $\bar{s}_i$ to s_i therefore has $4L$ zero-reward transitions followed by the L -transition path in (2.4f), giving

$$\tilde{V}^{\pi_{t+4}}(\bar{s}_i) = \gamma^{4L} (\gamma^L \tilde{V}^{\pi_{t+4}}(s_i) - 8f_{d-1}(\gamma)), \quad \forall 0 \leq i < d, \quad \tilde{V}^{\pi_{t+4}}(\bar{s}_d) = \gamma^L \tilde{V}^{\pi_{t+4}}(\bar{s}_0). \quad (3.28)$$

Using (3.27) and (3.28), we subtract the a_0 value $\gamma^{6L} \tilde{V}^{\pi_{t+4}}(\bar{s}_{i+1})$ from $\tilde{V}^{\pi_{t+4}}(\bar{s}_i)$ to obtain

$$\gamma^{-4L} \Delta^{\pi_{t+4}}(\bar{s}_i) = \gamma^L \Delta^{\pi_{t+4}}(s_i) - \begin{cases} 8f_{d-1}(\gamma)(1 - \gamma^{6L}), & 0 \leq i < d-1, \\ 8f_{d-1}(\gamma)(1 - \gamma^{7L}), & i = d-1. \end{cases} \quad (3.29)$$

For $i = d - 1$, the additional L transitions from $\bar{s}_d$ to $\bar{s}_0$ account for the exponent $7L$. By (3.16), we also have $\Delta^{\pi_{t+4}}(\bar{s}_{i,j}) = \gamma^{-jL} \Delta^{\pi_{t+4}}(\bar{s}_i)$ for $1 \leq j \leq 3$.

For $s \in \{s_i^c, s_i^z\}$, the a_0 path followed by π_{t+4} reaches s_{i+1} in $7L$ transitions, and the a_1 path reaches $s_{i,1}$ in $2L$ transitions. The identities $\tilde{Q}^{\pi_{t+4}}(s_i, a_0) = \gamma^{6L} \tilde{V}^{\pi_{t+4}}(s_{i+1})$ and $\tilde{Q}^{\pi_{t+4}}(s_i, a_1) = \gamma^L \tilde{V}^{\pi_{t+4}}(s_{i,1})$ let us express these action values in terms of those at s_i . By (2.6b) and (2.6c), we obtain

$$\begin{aligned}\tilde{Q}^{\pi_{t+4}}(s, a_0) &= \gamma^L [\tilde{Q}^{\pi_{t+4}}(s_i, a_0) - 8f_{d-1}(\gamma) - f_i(\gamma) + 1 - \gamma^L], \\ \tilde{Q}^{\pi_{t+4}}(s, a_1) &= \gamma^L [\tilde{Q}^{\pi_{t+4}}(s_i, a_1) - 8f_{d-1}(\gamma) - f_i(\gamma)].\end{aligned}\tag{3.30}$$

Subtracting the action values in (3.30), we obtain

$$\Delta^{\pi_{t+4}}(s) = \gamma^L (\Delta^{\pi_{t+4}}(s_i) - (1 - \gamma^L)), \quad \forall s \in \{s_i^c, s_i^z\}.\tag{3.31}$$

For $[\mathbf{b}^+]_i = 0$, (3.27) gives $\Delta^{\pi_{t+4}}(s_i) = 0$. Substituting this into (3.29) and (3.31), and using $f_{d-1}(\gamma) > 0$ for $\gamma < 1$ sufficiently close to one, we obtain $\Delta^{\pi_{t+4}}(\bar{s}_i), \Delta^{\pi_{t+4}}(s_i^c), \Delta^{\pi_{t+4}}(s_i^z) < 0$. For $[\mathbf{b}^+]_i = 1$, these action differences tend to zero. To determine their signs, we divide (3.29) and (3.31) by $1 - \gamma^L > 0$. Since $(1 - \gamma^{7L})/(1 - \gamma^L) \rightarrow 7$, the subtracted terms have limits at most $56f_{d-1}$ and 1, respectively. Using (2.2) and (3.3), we obtain

$$\lim_{\gamma \rightarrow 1^-} \frac{\Delta^{\pi_{t+4}}(s)}{1 - \gamma^L} \geq (24d + 4)g_i - 56f_{d-1} > 0, \quad \forall s \in \{\bar{s}_i, s_i^c, s_i^z\}.$$

We conclude that $\pi_{t+5}(s) = a_{[\mathbf{b}^+]_i}$ for $s \in \{\bar{s}_i, s_i^c, s_i^z\} \cup \{\bar{s}_{i,j}\}_{j=1}^3$, $0 \leq i < d$.

Step 5.2. Determine the actions at $s_i, s_d, \{s_{i,j}\}_{j=1}^5, \bar{s}_{i,4}$ and in $\mathcal{S}_3$. At $\bar{s}_{i,4}$, action a_0 has value $R_i(\gamma) + \gamma^{2L} \tilde{V}^{\pi_{t+4}}(\bar{s}_{i+1})$ by (2.4e). Combining (3.27) and (3.29), we obtain

$$\Delta^{\pi_{t+4}}(\bar{s}_{i,4}) = \gamma^{-4L} \Delta^{\pi_{t+4}}(\bar{s}_i) - R_i(\gamma) \leq (\gamma^{5L} - 1)R_i(\gamma) < 0, \quad \pi_{t+5}(\bar{s}_{i,4}) = a_0.$$

We next determine the actions in $\mathcal{S}_3$. By (3.27), we have $\tilde{Q}^{\pi_{t+4}}(s_i, a_1) = \tilde{V}^{\pi_{t+4}}(s_i)$ and $\Delta^{\pi_{t+4}}(s_i) \rightarrow 0$. Using (2.6b)–(2.6c) along the selected a_1 paths in $\mathcal{S}_3$, together with (3.6) and the action values at s_i^c, s_i^z in (3.30), we obtain

$$\lim_{\gamma \rightarrow 1^-} (\tilde{V}^{\pi_{t+4}}(s) - \tilde{V}^{\pi_{t+4}}(s_i)) = -8f_{d-1} - f_i, \quad \forall s \in \{s_i^c, s_i^z, s_{i,0}^r, s_{i,1}^r, s_{i,2}^r\}.\tag{3.32}$$

For $0 \leq i < d$, Table 1 gives the endpoints s_{i-1}^z for the a_0 paths from $s_{i,0}^r, s_{i,2}^r$, and s_{i-1}^c for the a_0 path from $s_{i,1}^r$. By (2.6a), we have $\tilde{R}^{\pi_{t+4}}((s_{i,0}^r, a_0) \rightsquigarrow s_{i-1}^z) = f_i(\gamma)$. The a_0 paths from $s_{i,1}^r, s_{i,2}^r$ have zero rewards. For $0 < i < d$, we subtract the a_0 value from the current a_1 value and use (3.6) and (3.32) to obtain

$$\lim_{\gamma \rightarrow 1^-} \Delta^{\pi_{t+4}}(s_{i,j}^r) \leq (-8f_{d-1} - f_i) - (-8f_{d-1} - f_{i-1}) = f_{i-1} - f_i \leq -1, \quad \forall 0 \leq j \leq 2.$$

For $i = 0$, we obtain from (2.5) and (3.28) that $\tilde{V}^{\pi_{t+4}}(s_{-1}^c) - \tilde{V}^{\pi_{t+4}}(s_0) \rightarrow -4f_{d-1}$ and $\tilde{V}^{\pi_{t+4}}(s_{-1}^z) - \tilde{V}^{\pi_{t+4}}(s_0) \rightarrow -8f_{d-1}$. Using (3.32), we therefore obtain

$$\lim_{\gamma \rightarrow 1^-} \Delta^{\pi_{t+4}}(s_{0,j}^r) \leq (-8f_{d-1} - f_0) - (-8f_{d-1}) = -f_0 = -1, \quad \forall 0 \leq j \leq 2.$$

We conclude that $\pi_{t+5}(s_{i,j}^r) = a_0$ for $0 \leq i < d$ and $0 \leq j \leq 2$. By Step 4.2 in Section 3.4, we have $\pi_{t+4}(s_{i,4}) = \pi_{t+4}(s_{i,5}) = a_0$. We use (3.10a), (3.11a), and (3.32) under π_{t+4} to obtain

$$\lim_{\gamma \rightarrow 1^-} \Delta^{\pi_{t+4}}(s_{i,4}) = -12f_{d-1} - \frac{1}{2} < 0, \quad \lim_{\gamma \rightarrow 1^-} \Delta^{\pi_{t+4}}(s_{i,5}) = -8f_{d-1} - 2f_i - \frac{1}{4} < 0.$$

We conclude that $s_{i,4}, s_{i,5}$ keep a_0 . At s_d , we use (2.4g) and (3.28) to obtain

$$\lim_{\gamma \rightarrow 1^-} \Delta^{\pi_{t+4}}(s_d) = 8f_{d-1} - 4f_{d-1} = 4f_{d-1} > 0, \quad \pi_{t+5}(s_d) = a_1.$$

For $[\mathbf{b}^+]_i = 0$, Step 4.2 in Section 3.4 gives $\pi_{t+4}(s_{i,1}) = a_0$ and $\pi_{t+4}(s_{i,2}) = \pi_{t+4}(s_{i,3}) = a_1$. Using (3.27) and the path comparison at $\bar{s}_{i,1}$ in Step 3.3 in Section 3.3, we obtain

$$\Delta^{\pi_{t+4}}(s_i) = 0, \quad \Delta^{\pi_{t+4}}(s_{i,1}) = \gamma^{3L} R_i(\gamma) > 0.$$

We conclude that s_i keeps a_0 and $s_{i,1}$ switches to a_1 . The same comparison keeps a_1 at $s_{i,2}, s_{i,3}$. For $[\mathbf{b}^+]_i = 1$, s_i and $\{s_{i,j}\}_{j=1}^3$ keep a_1 . All choices now agree with (2.8), so $\pi_{t+5} = \pi[\mathbf{b}^+]$.

3.6 Completing the proof

We verify that the action comparisons in Sections 3.1–3.5 hold at the discount in (2.7). Fix one of the five policies $\pi \in \{\pi_t, \dots, \pi_{t+4}\}$ specified in Table 2, and use the unscaled action values Q^π for the unshifted rewards. By Tables 1 and 2, every trajectory under π eventually reaches a cycle of length $(6d+1)L$. Thus, for any state s with two actions, we have

$$Q^\pi(s, a_1) - Q^\pi(s, a_0) = \frac{J(\gamma)}{1 - \gamma^{(6d+1)L}}, \quad J \in \mathbb{Z}[x]. \quad (3.33)$$

The transient path from either successor state and its eventual cycle contain at most N distinct states, so $\deg J \leq N$. Canceling the periodic terms in each action value leaves coefficients that are rewards or differences of two rewards, so every coefficient of J has magnitude at most $4r_{\max}$. For $J \neq 0$, we can write

$$J(\gamma) = (1 - \gamma)^{\ell_0} \left(c_0 + \sum_{\ell \geq 1} c_\ell (1 - \gamma)^\ell \right), \quad \sum_{\ell \geq 0} |c_\ell| \leq 4r_{\max} \sum_{j=0}^N 2^j < 2^{N+3} r_{\max},$$

where ℓ_0 is a nonnegative integer and $c_\ell \in \mathbb{Z}$ with $c_0 \neq 0$. The coefficient bound follows by expanding each $\gamma^j = (1 - (1 - \gamma))^j$, whose coefficients have absolute values summing to 2^j . The nonzero integer c_0 satisfies $|c_0| \geq 1$. At the discount in (2.7), we have

$$\left| \sum_{\ell \geq 1} c_\ell (1 - \gamma)^\ell \right| < (1 - \gamma) 2^{N+3} r_{\max} = 2^{3-N} < 1 \leq |c_0|. \quad (3.34)$$

Thus $J(\gamma)$ has the sign of c_0 both near one and at the chosen discount. Since the denominator in (3.33) is positive, all strict action comparisons are preserved, while $J = 0$ gives a tie for every discount. The bounds depend only on N and $r_{\max}$, so the same discount applies to every binary increment.

Thus $\pi_t = \pi[\mathbf{b}]$ implies $\pi_{t+5} = \pi[\mathbf{b}^+]$ for the specified discount. Induction from $\pi_0 = \pi[\mathbf{0}]$ proves the lemma and gives at least $5(2^d - 2) \geq 2^d$ iterations. $\square$

For $\mathbf{b} \neq \mathbf{1}$ and $s \in \{s_0, \bar{s}_0\}$, (2.10) expresses $V^{\pi[\mathbf{b}^+]}(s) - V^{\pi[\mathbf{b}]}(s)$ with denominator $1 - \gamma^{(6d+1)L}$ and an integer numerator of degree below N , with coefficients of magnitude at most $2r_{\max}$. Applying (3.34) to this numerator shows that the inequalities in Lemma 2.1 also hold at the discount in (2.7). The discount requires $O(N + \log r_{\max})$ bits.

Remark 3.2 (Comparison with Dantzig’s rule). Howard’s rule performs all improving switches using the same policy values, while the simplex method with Dantzig’s pivoting rule re-evaluates after each switch. Re-evaluation can change which switches are selected, so the two rules can produce different sequences of bit changes. For example, one can verify that, on the $d = 4$ instance in Section 4.1, Dantzig’s rule starting from $\pi[\mathbf{0}]$ can change the bits in each group in the order $0000 \rightarrow 1000 \rightarrow 1100 \rightarrow 1110 \rightarrow 1111$, with updates at other states omitted. Each bit changes only once along this trajectory, whereas Howard’s rule repeatedly changes bits to implement successive binary increments.

4 Proofs of the lower bounds

We choose the reward polynomials to prove Theorems 1.1 and 1.2. Lemma 3.1 supplies the bound of 2^d iterations. The choices below determine the number of states, reward magnitudes, and input length.

4.1 Proof of Theorem 1.1

Take $F_i(x) = 2^i$ and $G_i(x) = 2^{d+i+1}$. Conditions (2.1) and (2.2) hold with $\mathbf{f}_i = 2^i$ and $\mathbf{g}_i = 2^{d+i+1}$. Here $L = 2$, so (2.3) gives $N = \Theta(d)$. Lemma 3.1 gives at least 2^d iterations, which proves the $\exp(\Omega(N))$ lower bound. Since $r_{\max} = O(d4^d)$, each reward and the common discount use $O(d)$ bits, by (2.7). Each instance has encoding length $O(N^2)$ and can be constructed in time polynomial in N . $\square$

4.2 Proof of Theorem 1.2

A high-order zero at one allows us to obtain large limiting ratios by substituting powers of x , without increasing coefficient magnitudes. We first find a nonzero integer polynomial F_0 such that

$$\deg F_0 \leq 3d^2, \quad \max_{j \geq 0} |[x^j]F_0| \leq 3d^2, \quad F_0^{(r)}(1) = 0 \quad (0 \leq r < 2d). \quad (4.1)$$

Here $[x^j]F_0$ denotes the coefficient of x^j , and $F_0^{(r)}$ denotes its r -th derivative, with $F_0^{(0)} = F_0$. Let

$$\mathcal{F} = \left\{ \sum_{j=0}^{3d^2} a_j x^j : a_j \in \{0, \dots, 3d^2\} \right\} \quad \text{with} \quad |\mathcal{F}| = (3d^2 + 1)^{3d^2+1}.$$

For $\tilde{F} \in \mathcal{F}$, we define $\mathcal{T}(\tilde{F}) = (\tilde{F}(1), \tilde{F}'(1), \dots, \tilde{F}^{(2d-1)}(1))$. The entries of $\mathcal{T}(\tilde{F})$ are integers satisfying

$$0 \leq \tilde{F}^{(r)}(1) = \sum_{j=r}^{3d^2} a_j j(j-1) \cdots (j-r+1) \leq (3d^2 + 1)(3d^2)^{r+1}, \quad 0 \leq r < 2d.$$

Since each coordinate takes at most $(3d^2 + 1)(3d^2)^{r+1} + 1 < (3d^2 + 1)^{r+2}$ values and $d \geq 3$, we obtain

$$|\mathcal{T}(\mathcal{F})| < \prod_{r=0}^{2d-1} (3d^2 + 1)^{r+2} = (3d^2 + 1)^{2d^2+3d} < (3d^2 + 1)^{3d^2+1} = |\mathcal{F}|.$$

Hence, two distinct polynomials $\tilde{F}_1, \tilde{F}_2 \in \mathcal{F}$ satisfy $\mathcal{T}(\tilde{F}_1) = \mathcal{T}(\tilde{F}_2)$. Both $\tilde{F}_1 - \tilde{F}_2$ and $\tilde{F}_2 - \tilde{F}_1$ are nonzero and satisfy (4.1). Taylor expansion at $x = 1$ shows that one of them is positive on $(1 - \varepsilon, 1)$ for some $\varepsilon > 0$. Choose this difference as F_0 . Set

$$F_i(x) = F_0(x^{i+1}), \quad G_i(x) = F_0(x^{d+i+1}) \quad \forall 0 \leq i < d. \quad (4.2)$$

Let $r^* = \min\{r \geq 0 : F_0^{(r)}(1) \neq 0\} \geq 2d$. Taylor expansion at $x = 1$ gives $\lim_{x \rightarrow 1^-} F_0(x^q)/F_0(x) = q^{r^*}$ for every positive integer q . The limiting weights in (2.1) are therefore

$$\mathbf{f}_i = (i+1)^{r^*}, \quad \mathbf{g}_i = (d+i+1)^{r^*} \quad (0 \leq i < d), \quad \mathbf{f}_0 = 1. \quad (4.3)$$

For $1 \leq j \leq 2d-1$, the binomial theorem and $r^* \geq 2d$ give $(1 + 1/j)^{r^*} \geq (1 + 1/j)^j \geq 2$. It follows that

$$\mathbf{f}_i \geq 2\mathbf{f}_{i-1}, \quad \mathbf{g}_i \geq 2\mathbf{g}_{i-1} \quad (1 \leq i < d), \quad \frac{\mathbf{g}_0}{\mathbf{f}_{d-1}} = \left(1 + \frac{1}{d}\right)^{r^*} \geq \left[\left(1 + \frac{1}{d}\right)^d\right]^2 \geq 4. \quad (4.4)$$

Combining (4.3) and (4.4) yields (2.2). Lemma 3.1 then gives at least 2^d iterations.

We next bound the immediate rewards. Since $2d \leq r^* \leq \deg F_0 \leq 3d^2$, the definition of L and (4.2) give

$$4d^2 + 2 \leq L = 2d \deg F_0 + 2 \leq 6d^3 + 2. \quad (4.5)$$

By (4.1), (4.2), and (4.5), every coefficient of F_i, G_i has magnitude at most $3d^2 < L$. The reward formulas (2.4)–(2.6) therefore give $r_{\max} \leq (96d + 21)L$, with the largest bound coming from (2.4d). Combining this bound with $N = (67d + 5)L - 16d - 1$ from (2.3), we obtain

$$0 \leq r(s, a) \leq 2r_{\max} < 4N \quad \forall s \in \mathcal{S}, a \in \mathcal{A}_s. \quad (4.6)$$

Each reward and successor index uses $O(\log N)$ bits, while the common discount in (2.7) uses $O(N)$ bits. With at most two actions per state, the total encoding length is $O(N \log N)$. Finally, combining (4.5) with (2.3) gives $N = O(d^4)$. The lower bound of 2^d iterations therefore implies the claimed $\exp(\Omega(N^{1/4}))$ bound. $\square$

5 Further results and concluding remarks

We have established an exponential iteration lower bound for Howard's policy iteration on deterministic MDPs with at most two actions per state. A stretched-exponential lower bound holds even when each reward uses $O(\log N)$ bits.

The instances in Theorem 1.1 can be generated in time polynomial in N . The proof of Theorem 1.2 uses a pigeonhole argument to establish the existence of the reward polynomial. The following proposition gives an explicit construction under the same reward restriction, with a weaker iteration lower bound.

Proposition 5.1 (An explicit family). *There is a family of deterministic discounted MDPs with N states, at most two actions per state, and nonnegative integer rewards smaller than $4N$, on which Howard's policy iteration performs at least $N^{\Omega(\log N)}$ iterations from a specified initial policy. Each instance has encoding length $O(N \log N)$ and can be constructed in time polynomial in N .*

Proof. Take an integer $m \geq 2$ and set $d = m^2$. For $i = mj + r$, $0 \leq j, r < m$, set

$$F_i(x) = 2^r(1 - x^{2^j})^m, \quad G_i(x) = 2F_i(x^{2^m}). \quad (5.1)$$

Here $F_0(x) = (1 - x)^m > 0$ for $x < 1$. Since $(1 - x^q)/(1 - x) = 1 + x + \dots + x^{q-1} \rightarrow q$ as $x \rightarrow 1^-$ for every positive integer q , we obtain

$$f_i = 2^r(2^j)^m = 2^i, \quad g_i = 2^{r+1}(2^{j+m})^m = 2^{d+i+1}.$$

These weights satisfy (2.2). The coefficients in (5.1) have magnitude at most $2^{r+1}2^m \leq 4^m < L = m2^{2m-1} + 2$. Thus (4.6) and the $O(N \log N)$ encoding bound in Section 4.2 apply. The polynomial coefficients, transition paths, and discount in (2.7) can all be generated in time polynomial in N . By (2.3), we have

$$N = \Theta(m^3 4^m), \quad \log N = \Theta(m), \quad 2^{m^2} = N^{\Theta(\log N)}.$$

Lemma 3.1 gives at least $2^d = 2^{m^2}$ iterations, proving the claim. $\square$

The same iteration lower bounds apply to halving the initial optimality gap at s_0 .

Corollary 5.2. *Let V^* denote the optimal value function. For the families in Theorems 1.1 and 1.2 and Proposition 5.1, attaining*

$$V^*(s_0) - V^{\pi_t}(s_0) \leq \frac{1}{2}(V^*(s_0) - V^{\pi_0}(s_0)) \quad (5.2)$$

from the specified initial policies requires at least $\exp(\Omega(N))$, $\exp(\Omega(N^{1/4}))$, and $N^{\Omega(\log N)}$ iterations, respectively.

Proof. Omit the common reward shift, which preserves optimality gaps. Then $V^{\pi_0}(s_0) = V^{\pi^{[0]}}(s_0) = 0$. Let $\mathbf{b} \in \{0, 1\}^d$ have $\mathbf{b}_{d-2} = 1$ and all other entries zero. Lemma 3.1 gives policy $\pi[\mathbf{b}]$ after $5 \cdot 2^{d-2}$ iterations. We show that $V^{\pi^{[1]}}(s_0) > 2V^{\pi[\mathbf{b}]}(s_0)$. Using (2.1), (2.2), and (2.10), we obtain

$$\lim_{\gamma \rightarrow 1^-} \frac{V^{\pi^{[1]}}(s_0) - 2V^{\pi[\mathbf{b}]}(s_0)}{4\gamma F_0(\gamma)} = 4 \left(\sum_{i=0}^{d-1} g_i - 2g_{d-2} \right) \geq 4 \sum_{i=0}^{d-2} g_i > 0. \quad (5.3)$$

By (2.10), $V^{\pi^{[1]}}(s_0) - 2V^{\pi[\mathbf{b}]}(s_0)$ has denominator $1 - \gamma^{(6d+1)L}$ and an integer polynomial numerator of degree below N , with coefficients of magnitude at most $4r_{\max}$. Applying (3.34) to this numerator and using (5.3), we obtain $V^{\pi^{[1]}}(s_0) > 2V^{\pi[\mathbf{b}]}(s_0)$ for γ in (2.7). By monotonicity of policy iteration,

$$V^{\pi_t}(s_0) \leq V^{\pi[\mathbf{b}]}(s_0) < \frac{1}{2}V^{\pi^{[1]}}(s_0) \leq \frac{1}{2}V^*(s_0), \quad 0 \leq t \leq 5 \cdot 2^{d-2}.$$

Since $V^{\pi_0}(s_0) = 0$, the target in (5.2) has not been reached at any of these iterations. $\square$

The price of algorithmic anarchy also affects progress toward optimality. By Corollary 5.2, Howard's rule can require exponentially many iterations even to halve the initial optimality gap at s_0 , while Dantzig's rule reaches an optimal policy in polynomially many iterations.

AI use statement

We used generative AI to assist with some derivations, literature searches, proof checking, language editing, and L^AT_EX formatting. The authors carefully reviewed all AI-assisted material and independently verified the mathematical arguments. The authors take full responsibility for this paper.

References

- Asadi, A., Chatterjee, K. and de Raaij, J. (2025). Lower bound on Howard policy iteration for deterministic Markov decision processes. In *Proceedings of the Forty-first Conference on Uncertainty in Artificial Intelligence*, vol. 286 of *Proceedings of Machine Learning Research*. PMLR.
- Fearnley, J. (2010). Exponential lower bounds for policy iteration. In *Automata, Languages and Programming*, vol. 6199 of *Lecture Notes in Computer Science*. Springer.
- Goenka, R., Gupta, E., Khyalia, S. and Kalyanakrishnan, S. (2026). Upper bounds for all and max-gain policy iteration algorithms on deterministic MDPs. *Mathematics of Operations Research*, **51** 806–828.
- Hansen, T. D. (2012). *Worst-case Analysis of Strategy Iteration and the Simplex Method*. Ph.D. thesis, Aarhus University.
- Hansen, T. D. and Zwick, U. (2010). Lower bounds for Howard’s algorithm for finding minimum mean-cost cycles. In *Algorithms and Computation*, vol. 6506 of *Lecture Notes in Computer Science*. Springer.
- Hollanders, R., Delvenne, J.-C. and Jungers, R. M. (2012). The complexity of policy iteration is exponential for discounted Markov decision processes. In *Proceedings of the 51st IEEE Conference on Decision and Control*. IEEE.
- Howard, R. A. (1960). *Dynamic Programming and Markov Processes*. Technology Press of Massachusetts Institute of Technology, Cambridge, MA.
- Mukherjee, D. and Kalyanakrishnan, S. (2025). Howard’s policy iteration is subexponential for deterministic Markov decision problems with rewards of fixed bit-size and arbitrary discount factor. In *Proceedings of the International Conference on Automated Planning and Scheduling*, vol. 35. AAAI Press.
- Post, I. and Ye, Y. (2015). The simplex method is strongly polynomial for deterministic Markov decision processes. *Mathematics of Operations Research*, **40** 859–868.
- Puterman, M. L. (1994). *Markov Decision Processes: Discrete Stochastic Dynamic Programming*. Wiley, New York.
- Scherrer, B. (2016). Improved and generalized upper bounds on the complexity of policy iteration. *Mathematics of Operations Research*, **41** 758–774.
- Sutton, R. S. and Barto, A. G. (2018). *Reinforcement Learning: An Introduction*. 2nd ed. MIT Press, Cambridge, MA.
- Ye, Y. (2005). A new complexity result on solving the Markov decision problem. *Mathematics of Operations Research*, **30** 733–749.
- Ye, Y. (2011). The simplex and policy-iteration methods are strongly polynomial for the Markov decision problem with a fixed discount rate. *Mathematics of Operations Research*, **36** 593–603.