

STOCHASTIC GRADIENT DESCENT ASCENT IS SUBOPTIMAL FOR NONCONVEX-PŁ MIN-MAX GAMES

Junsoo Ha

junsoo.ha.contact@gmail.com

ABSTRACT

How far can stochastic gradient descent ascent (SGDA) go by tuning its timescale ratio and step sizes in nonconvex min-max games? We answer this question for nonconvex-PŁ (NC-PŁ) games by establishing the first tight complexity of two-timescale SGDA with a fixed timescale ratio and non-increasing step sizes. For ℓ -smooth games with an inner μ -PŁ inequality, we prove a complexity lower bound $\Omega(\kappa^2 \ell \varepsilon^{-2} + \kappa^4 \ell \sigma^2 \varepsilon^{-4})$, where $\kappa = \ell/\mu$ is the condition number, σ^2 is the gradient variance, and ε measures the outer gradient norm. This matches existing SGDA upper bounds and establishes a complexity separation from Smoothed-AGDA (Yang et al., 2022). In addition, we show that SGDA can fail to find a stationary point when its timescale ratio is as small as $o(\kappa^2)$. Our negative results highlight the fundamental limitation of SGDA in NC-PŁ games, and justify the development of alternative methods.

1 INTRODUCTION

Nonconvex min-max games have become a cornerstone of modern machine learning, underpinning generative adversarial networks (GANs) (Goodfellow et al., 2014), distributionally robust learning (Sinha et al., 2018), and language model alignment (Liu et al., 2026; Xu et al., 2025). Without additional structure, however, even finding a stationary point of a min-max game can be computationally intractable (Daskalakis et al., 2021; Bernasconi et al., 2026). For instance, games often exhibit limit cycles (Hsieh et al., 2021) and chaotic behavior (Sato et al., 2002). Such pathological examples motivate the study of *structured* nonconvex min-max games (Lin et al., 2020; 2025; Ha, 2026). In particular, nonconvex-PŁ (NC-PŁ) games assume a Polyak–Łojasiewicz (PŁ) inequality (Polyak, 1963) on the inner maximization problem and have been actively studied as one of the weakest forms of structure (Nouiehed et al., 2019; Yang et al., 2022; Cho and Yun, 2023; Ha, 2026; Cai et al., 2026).

Even in NC-PŁ games, finding a Nash equilibrium (Nash, 1951) can be still intractable; hence, we seek a stationary point of the outer function $\Phi(x) := \max_y f(x, y)$. Two-timescale SGDA is a direct analog of SGD for such min-max games: it uses a faster ascent step to track the outer gradient $\nabla \Phi(x)$ and performs a slower inexact descent on Φ . For ℓ -smooth nonconvex- μ -PŁ games with condition number $\kappa = \ell/\mu$ and gradient variance σ^2 , SGDA admits a complexity upper bound $O(\kappa^2 \ell \varepsilon^{-2} + \kappa^4 \ell \sigma^2 \varepsilon^{-4})$ for finding an ε -stationary point (Yang et al., 2022; Ha, 2026). Meanwhile, recent studies have developed more sophisticated techniques to attain sharper bounds (Yang et al., 2022; Huang et al., 2025; Cai et al., 2026). This raises the following natural question:

Can SGDA obtain better complexity by tuning its timescale ratio and step sizes?

Contributions. In this work, we answer this question negatively. Our contributions are twofold.

- (i) **Suboptimality of SGDA.** We prove matching lower bounds for SGDA with a fixed timescale ratio and nonincreasing step sizes, and establish its suboptimality in NC-PŁ games.
- (ii) **Optimal timescale ratio.** Additionally, we prove that SGDA with a small timescale ratio $o(\kappa^2)$ can fail to find a stationary point, and that existing $\Theta(\kappa^2)$ timescale ratio is optimal.

Sections 2–4 cover related work, the setup, and main results; Sections 5–6 prove the lower bounds; Section 7 discusses implications and future directions.

2 RELATED WORK

Nonconvex games are computationally intractable in general (Daskalakis et al., 2021), and even finding an approximate stationary point can be hard as well (Bernasconi et al., 2026). Such hardness results motivate structured games, including nonconvex-PŁ (NC-PŁ) (Nouiehed et al., 2019; Yang et al., 2020; 2022; Cho and Yun, 2023) and nonconvex-concave (NC-C) games (Thekumparampil et al., 2019; Lin et al., 2020; 2025; Ha, 2026), which assume a PŁ inequality and concavity on inner maximization problems, respectively. Notably, these structures make finding stationary points computationally tractable and admit inexact-descent interpretations (Nouiehed et al., 2019; Lin et al., 2020; 2025; Ha, 2026). In particular, Nouiehed et al. (2019) showed that taking multiple ascent steps per descent step suffices to find an approximate stationary point in NC-PŁ games, and interpreted these updates as inexact descent on $\Phi(x) = \max_y f(x, y)$. Lin et al. (2020; 2025) extended the same interpretation to two-timescale SGDA for nonconvex-strongly-concave (NC-SC) and NC-C games. On the other hand, weakly-convex-weakly-concave games (Liu et al., 2021; Grimmer et al., 2023; Hajizadeh et al., 2024) and nonmonotone variational inequalities (Mertikopoulos et al., 2019; Diakonikolas et al., 2021; Lee and Kim, 2021; Pethick et al., 2022; Böhm, 2023) have been studied as well, focusing on extragradient and optimistic gradient (OGDA), rather than SGDA.

NC-PŁ games. Yang et al. (2020) gave the first convergence analysis of two-timescale alternating gradient descent ascent (GDA) for NC-PŁ games. Subsequently, Yang et al. (2022) extended the analysis to stochastic settings and introduced Smoothed-AGDA, attaining $O(\kappa\ell\varepsilon^{-2} + \kappa^2\ell\sigma^2\varepsilon^{-4})$ under the comparison conditions in Appendix E. Meanwhile, Huang et al. (2025) used momentum to attain an improved leading stochastic complexity of $\tilde{O}((\kappa^3\ell\Delta\sigma + \kappa^4\sigma^3)\varepsilon^{-3})$ under Lipschitz sample gradient assumptions; Cai et al. (2026) obtained the same dependence on ε using Hessian-vector products to correct momentum bias. For finite-sum NC-PŁ games, Chen et al. (2022) and Cho and Yun (2023) obtained better complexities through variance reduction and sampling without replacement, respectively. Finally, recent studies also established high-probability guarantees for Smoothed-AGDA (Laguel et al., 2024) and SGDA (Ha, 2026).

Two-timescale SGDA exposes a timescale ratio hyperparameter ρ that controls the size of ascent steps relative to descent steps. Fiez and Ratliff (2021) showed that a sufficiently large timescale ratio ρ yields local linear convergence of simultaneous GDA to differential Stackelberg equilibria. For NC-SC games, Li et al. (2022) proved that $\rho = \Theta(\kappa)$ suffices for local convergence, whereas GDA with $\rho = o(\kappa)$ can fail to converge locally. On the other hand, existing global convergence guarantees for GDA and SGDA in NC-PŁ games rely on a timescale ratio $\rho = \Theta(\kappa^2)$ (Fiez et al., 2021; Yang et al., 2022; Ha, 2026). Cho and Yun (2023) showed that an excessively large timescale ratio can worsen convergence rates of GDA in strongly-convex-strongly-concave (SC-SC) games, and raised the question whether $\rho = \Theta(\kappa^2)$ is necessary for global convergence in NC-PŁ games.

Lower bounds for structured nonconvex min-max games have mainly addressed the minimax complexity of first-order methods. For instance, in NC-SC games, Zhang et al. (2021) proved a deterministic $\Omega(\sqrt{\kappa\ell\Delta\varepsilon^{-2}})$ lower bound, and Li et al. (2021) proved a stochastic $\Omega(\sqrt{\kappa\ell\Delta\varepsilon^{-2}} + \kappa^{1/3}\ell\Delta\sigma^2\varepsilon^{-4})$ bound. More recently, Pan and Li (2026) proved a sharper deterministic NC-PŁ lower bound $\Omega(\kappa\ell\Delta\varepsilon^{-2})$, establishing the minimax optimality of Smoothed-AGDA’s leading complexity. In this work, we derive SGDA-specific lower bounds that match existing upper bounds (Yang et al., 2022; Ha, 2026), and establish provable suboptimality of SGDA in NC-PŁ games.

3 PROBLEM SETUP

We study the following min-max optimization problem:

$$\min_{x \in \mathbb{R}^{d_x}} \max_{y \in \mathbb{R}^{d_y}} f(x, y), \quad (3.1)$$

where f satisfies a smoothness assumption and $f(x, \cdot)$ satisfies a PŁ inequality (Polyak, 1963).

Assumption 3.1 (Smoothness). The game $f : \mathbb{R}^{d_x} \times \mathbb{R}^{d_y} \rightarrow \mathbb{R}$ is differentiable and ℓ -smooth:

$$\begin{aligned} \|\nabla_x f(x_1, y_1) - \nabla_x f(x_2, y_2)\| &\leq \ell(\|x_1 - x_2\| + \|y_1 - y_2\|), \quad \forall x_1, x_2 \in \mathbb{R}^{d_x}, y_1, y_2 \in \mathbb{R}^{d_y}, \\ \|\nabla_y f(x_1, y_1) - \nabla_y f(x_2, y_2)\| &\leq \ell(\|x_1 - x_2\| + \|y_1 - y_2\|), \quad \forall x_1, x_2 \in \mathbb{R}^{d_x}, y_1, y_2 \in \mathbb{R}^{d_y}. \end{aligned}$$

We denote the outer objective of a game by $\Phi(x) := \max_{y \in \mathbb{R}^{d_y}} f(x, y)$, and let $\Phi^* := \inf_{x \in \mathbb{R}^{d_x}} \Phi(x)$. We define the condition number $\kappa := \ell/\mu \geq 1$, where $\mu > 0$ is the PL parameter.

Assumption 3.2 (NC-PL). The outer objective satisfies $\Phi(x) < \infty$ for every $x \in \mathbb{R}^{d_x}$, and $Y^*(x) := \arg \max_{y \in \mathbb{R}^{d_y}} f(x, y)$ is nonempty for every $x \in \mathbb{R}^{d_x}$. There exists $\mu > 0$ such that the following inequality holds for each $x \in \mathbb{R}^{d_x}$:

$$\frac{1}{2} \|\nabla_y f(x, y)\|^2 \geq \mu(\Phi(x) - f(x, y)), \quad \forall y \in \mathbb{R}^{d_y}. \quad (3.2)$$

Assumption 3.3 (Initialization). There exists $\Delta > 0$ and a universal constant $c_{\text{init}} \geq 1$ such that

$$0 \leq \Phi(x_0) - \Phi^* \leq \Delta, \quad 0 \leq \Phi(x_0) - f(x_0, y_0) \leq c_{\text{init}} \Delta. \quad (3.3)$$

Throughout the paper, we make the following standard stochastic gradient estimator assumptions.

Assumption 3.4 (Unbiasedness). The gradient estimators G_x, G_y are measurable and use fresh independent samples ξ from a fixed distribution. For every (x, y) ,

$$\mathbb{E} G_x(x, y; \xi) = \nabla_x f(x, y), \quad \mathbb{E} G_y(x, y; \xi) = \nabla_y f(x, y). \quad (3.4)$$

Assumption 3.5 (Bounded variance). For a fixed $\sigma \geq 0$ and every (x, y) ,

$$\mathbb{E} \|G_x(x, y; \xi) - \nabla_x f(x, y)\|^2 \leq \sigma^2, \quad \mathbb{E} \|G_y(x, y; \xi) - \nabla_y f(x, y)\|^2 \leq \sigma^2. \quad (3.5)$$

Two-timescale SGDA. SGDA with a fixed timescale ratio $\rho \geq 1$ and step sizes $\eta_t \geq 0$ follows:

$$x_{t+1} = x_t - \frac{\eta_t}{\rho} G_x(x_t, y_t; \xi_t^x), \quad y_{t+1} = y_t + \eta_t G_y(x_t', y_t; \xi_t^y), \quad (3.6)$$

where we use $x_t' = x_{t+1}$ for SGDA_{Alt}, and $x_t' = x_t$ for SGDA_{Sim}.

Minimax complexity. Let $\mathcal{C}(\ell, \kappa, \Delta, \sigma)$ collect all NC-PL game instances that satisfy Assumptions 3.1–3.5 with $d_x, d_y < \infty$. For each SGDA update order $\mathfrak{o} \in \{\text{Alt}, \text{Sim}\}$, we define the minimax convergence rate of SGDA with a fixed timescale ratio ρ and nonincreasing step sizes η_t as

$$\mathcal{R}_{T, \mathfrak{o}}(\rho) := \inf_{\eta_0 \geq \dots \geq \eta_{T-2} \geq 0} \sup_{I \in \mathcal{C}(\ell, \kappa, \Delta, \sigma)} \frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E} \|\nabla \Phi(x_t)\|^2. \quad (3.7)$$

We write the version optimized over the timescale ratio as $\mathcal{R}_{T, \mathfrak{o}}^* := \inf_{\rho \geq 1} \mathcal{R}_{T, \mathfrak{o}}(\rho)$. We emphasize that we allow parameter-dependent tuning of timescale ratio $\rho(\ell, \kappa, \Delta, \sigma, T)$ and step sizes $\eta_t(\ell, \kappa, \Delta, \sigma, T)$. Finally, we define the corresponding iteration complexities as

$$N_{\varepsilon, \mathfrak{o}}(\rho) := \inf \{T \in \mathbb{N}_{\geq 1} : \mathcal{R}_{T, \mathfrak{o}}(\rho) \leq \varepsilon^2\}, \quad N_{\varepsilon, \mathfrak{o}}^* := \inf \{T \in \mathbb{N}_{\geq 1} : \mathcal{R}_{T, \mathfrak{o}}^* \leq \varepsilon^2\}. \quad (3.8)$$

4 MAIN RESULTS

Our main result establishes a tight lower bound for two-timescale SGDA with a fixed timescale ratio $\rho \geq 1$ and non-increasing step size schedules η_t .

Theorem 4.1 (A lower bound for two-timescale SGDA in NC-PL games). *There are universal constants $c > 0$ and $\kappa_0 \geq 2$, depending on c_{init} , such that, for every $\kappa \geq \kappa_0$, $\ell > 0$, $\Delta > 0$, $\sigma \geq 0$, and integer $T \geq 1$,*

$$\mathcal{R}_{T, \text{Sim}}^*, \mathcal{R}_{T, \text{Alt}}^* \geq c \min \left\{ \ell \Delta, \max \left\{ \frac{\kappa^2 \ell \Delta}{T}, \sqrt{\frac{\kappa^4 \ell \Delta \sigma^2}{T}} \right\} \right\}. \quad (4.1)$$

Intuitively, the first term $\ell \Delta$ inside the minimum accounts for short horizons. Hence, in the usual small-error regime $\varepsilon^2 \leq c \ell \Delta$, Theorem 4.1 and the matching upper bounds for SGDA_{Alt} from Yang et al. (2022) and for SGDA_{Sim} in Theorem A.1 yield a tight complexity:

$$N_{\varepsilon, \text{Sim}}^*, N_{\varepsilon, \text{Alt}}^* \asymp \frac{\kappa^2 \ell \Delta}{\varepsilon^2} + \frac{\kappa^4 \ell \Delta \sigma^2}{\varepsilon^4}. \quad (4.2)$$

Table 1: **Upper and lower bounds in NC-PŁ games.** Our lower bounds match SGDA’s upper bounds after optimizing fixed timescale ratios and predetermined nonincreasing step sizes. With exact gradients, Theorem 6.3 rules out uniform convergence for $\rho = o(\kappa^2)$; $\Omega(1)$ denotes nonvanishing stationarity error. MSGDA and AdaMSGDA use $0 < \varepsilon \leq \sigma$. $\dagger$ AdaMSGDA measures the average expected squared norm of $A_t^{-1} \nabla \Phi(x_t)$, where A_t is its primal preconditioner. $\ddagger$ HCMM-2 shows only the variance-dependent cost for $\mathbb{E} \|\nabla \Phi\| \leq \varepsilon$, with Hessian variance at most σ^2 . Appendix F gives the complete bounds and assumptions.

Algorithm	Citation	Feedback	Complexity / obstruction
SGDA _{Alt}	Yang et al. (2022)	Bounded σ^2	$O(\kappa^2 \ell \varepsilon^{-2} + \kappa^4 \ell \sigma^2 \varepsilon^{-4})$
SGDA _{Sim}	Theorem A.1	Bounded σ^2	$O(\kappa^2 \ell \varepsilon^{-2} + \kappa^4 \ell \sigma^2 \varepsilon^{-4})$
Smoothed-AGDA	Yang et al. (2022)	Bounded σ^2	$\tilde{O}(\kappa \ell \varepsilon^{-2} + \kappa^2 \ell \sigma^2 \varepsilon^{-4})$
SPIDER-GDA	Chen et al. (2022)	Finite-sum	$O(n + \sqrt{n} \kappa^2 \ell \varepsilon^{-2})$
SGDA-RR	Cho and Yun (2023)	Finite-sum	$O(n \kappa^2 \ell \varepsilon^{-2} + \sqrt{n} \kappa^3 \ell \sigma \varepsilon^{-3})$
MSGDA	Huang et al. (2025)	Smooth samples	$\tilde{O}((\kappa^3 \ell \sigma + \kappa^4 \sigma^3) \varepsilon^{-3})$
AdaMSGDA †	Huang et al. (2025)	Smooth samples	$\tilde{O}((\kappa^3 \ell \sigma + \kappa^4 \sigma^3) \varepsilon^{-3})$
HCMM-2 ‡	Cai et al. (2026)	Hessian products	$O_\ell(\kappa^3 \sigma^3 \varepsilon^{-3})$
SGDA_{Alt}, SGDA_{Sim}	Theorem 4.1	Bounded σ^2	$\Theta(\kappa^2 \ell \varepsilon^{-2} + \kappa^4 \ell \sigma^2 \varepsilon^{-4})$
Linear-span (NC-SC)	Zhang et al. (2021)	Deterministic	$\Omega(\sqrt{\kappa} \ell \varepsilon^{-2})$
Linear-span (NC-SC)	Zhang et al. (2021)	Finite-sum	$\Omega(n + \sqrt{n} \kappa \ell \varepsilon^{-2})$
Zero-respecting (NC-SC)	Li et al. (2021)	Bounded σ^2	$\Omega(\sqrt{\kappa} \ell \varepsilon^{-2} + \kappa^{1/3} \ell \sigma^2 \varepsilon^{-4})$
First-order (NC-PŁ)	Pan and Li (2026)	Deterministic	$\Omega(\kappa \ell \varepsilon^{-2})$
SGDA _{Sim}	Li et al. (2022)	Deterministic	Local failure if $\rho = o(\kappa)$
SGDA_{Alt}, SGDA_{Sim}	Theorem 6.3	Deterministic	$\Omega(1)$ if $\rho = o(\kappa^2)$
SGDA_{Alt}, SGDA_{Sim}	Theorem 5.1	Bounded σ^2	$\Omega(\rho \ell \varepsilon^{-2} + \kappa^4 \ell \sigma^2 \varepsilon^{-4})$

Suboptimality of SGDA. Theorem 4.1 establishes a provable suboptimality of SGDA in NC-PŁ games: it yields a tight sample complexity $\Theta(\kappa^2 \ell \Delta \varepsilon^{-2} + \kappa^4 \ell \Delta \sigma^2 \varepsilon^{-4})$ for SGDA, which is suboptimal compared with the complexity upper bound $O(\kappa \ell \Delta \varepsilon^{-2} + \kappa^2 \ell \Delta \sigma^2 \varepsilon^{-4})$ of Smoothed-AGDA (Yang et al., 2022).¹ Hence, our result justifies the recent development of alternative methods such as Smoothed-AGDA (Yang et al., 2022) and momentum (Huang et al., 2025; Cai et al., 2026).

Optimal timescale ratio and step-size schedule. Notably, every existing analysis of two-timescale SGDA in NC-PŁ games relies on the choice of timescale ratio $\rho = \Theta(\kappa^2)$ (Fiez et al., 2021; Yang et al., 2022; Cho and Yun, 2023; Ha, 2026). However, the question whether it is actually optimal remained open. Together with the matching upper bounds (Yang et al., 2022; Ha, 2026), Theorem 4.1 shows that the timescale ratio $\rho = \Theta(\kappa^2)$ and the constant step size $\eta_t \equiv c_\eta \min\{1/\ell, \sqrt{\Delta/(\ell \sigma^2 T)}\}$ are optimal for two-timescale SGDA with a fixed timescale ratio and non-increasing step sizes.

In the following sections, we study two distinctive regimes of timescale ratios, namely the high-ratio regime $\rho = \Omega(\kappa^2)$ and low-ratio regime $\rho = O(\kappa^2)$, and explain the intuition behind the hard instances that establish Theorem 4.1. We defer the proof of Theorem 4.1 to Appendix A.1.

5 THE HIGH-RATIO LOWER BOUND

In this section, we establish the lower bound for large timescale ratios $\rho = \Omega(\kappa^2)$. In this high-ratio regime, smaller descent steps slow down the outer progress, whereas larger ascent steps increase noise-induced errors that accumulate on the outer variable. These complementary obstructions prevent SGDA from obtaining sharper convergence rates.

¹The Smoothed-AGDA bound uses the comparison conditions and output conversion in Appendix E.

Theorem 5.1 (High-ratio lower bound). *There are universal constants $c, c_{\text{high}} > 0$ and $\kappa_{\text{high}} \geq 2$ such that, for every choice of problem parameters $\ell, \Delta > 0$, $\sigma \geq 0$, and $\kappa \geq \kappa_{\text{high}}$, time horizon $T \in \mathbb{N}_{\geq 1}$, and timescale ratio $\rho \geq c_{\text{high}}\kappa^2$, both SGDA_{Sim} and SGDA_{Alt} satisfy*

$$\mathcal{R}_{T,\text{Sim}}(\rho), \mathcal{R}_{T,\text{Alt}}(\rho) \geq c \min \left\{ \ell\Delta, \max \left\{ \frac{\rho\ell\Delta}{T}, \sqrt{\frac{\kappa^4\ell\Delta\sigma^2}{T}} \right\} \right\}. \quad (5.1)$$

Together with the matching upper bounds from Yang et al. (2022) and Theorem A.1, Theorem 5.1 shows that the smallest possible timescale ratio $\rho = \Theta(\kappa^2)$ is optimal in the high-ratio regime, as larger ratios only increase the lower bound. This suggests that the standard choice $\rho = \Theta(\kappa^2)$ (Yang et al., 2022) might be optimal among all fixed ratios. We establish this optimality in Section 6 by ruling out a uniform convergence guarantee for timescale ratios of smaller order $\rho = o(\kappa^2)$.

In the following subsections, we construct two hard NC-PŁ game instances that yield Theorem 5.1.

5.1 THE INSTANCE FOR UNSTABLE STEP SIZES

Consider a simple linear quadratic game $f_{\text{unstable}} : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ and its outer objective:

$$f_{\text{unstable}}(x; y) = \frac{\ell + \mu}{4}(xy - y^2), \quad \Phi_{\text{unstable}}(x) = \frac{\ell + \mu}{16}x^2. \quad (5.2)$$

This game is ℓ -smooth and μ -strongly concave in y , and therefore, is an NC-PŁ game. For this game, an inner ascent step becomes unstable for $\eta_t \geq 8/\ell$. The following result shows that such m initial unstable steps can push x away from its minimizer.

Lemma 5.2 (A lower bound for f_{unstable}). *There is a universal $c > 0$ such that, for $\ell, \Delta > 0$, $\kappa \geq 2$, $\sigma \geq 0$, $\rho \geq 1$, integers $T \geq 1$ and $0 \leq m < T$, and predetermined nonnegative non-increasing step sizes with $\eta_t \geq 8/\ell$ for $t < m$, each of SGDA_{Sim} and SGDA_{Alt} admits an initialization (x_0, y_0) making f_{unstable} with $\mu = \ell/\kappa$ an exact-gradient instance in $\mathcal{C}(\ell, \kappa, \Delta, \sigma)$ with*

$$\frac{1}{T} \sum_{t=0}^{T-1} |\Phi'_{\text{unstable}}(x_t)|^2 \geq c \frac{m+1}{T} \ell\Delta. \quad (5.3)$$

For $m+1 \geq T/2$, Lemma 5.2 immediately proves Theorem 5.1 with the constant lower bound $c\ell\Delta$. We construct a distinct hard instance f_{stable} for the other case $m+1 < T/2$ in the next subsection.

5.2 THE INSTANCE FOR STABLE STEP SIZES

For the remaining case $m+1 < T/2$, we design an instance f_{stable} with outer coordinates $x = (u, v)$:

$$f_{\text{stable}}(u, v; y) = g_{\text{corridor}}(u) + g_{\sigma}(v, y), \quad \Phi_{\text{stable}}(u, v) = g_{\text{corridor}}(u) + \Phi_{\sigma}(v), \quad (5.4)$$

where we use a scalar coordinate $u \in \mathbb{R}$ for the deterministic function g_{corridor} , and define the outer objective $\Phi_{\sigma}(v) := \max_y g_{\sigma}(v, y)$ for the noisy function $g_{\sigma}(v, y)$. Intuitively, the corridor function g_{corridor} limits progress from small descent steps, and the noisy function g_{σ} transfers the inner stochastic perturbations to the outer variable. The stable ascent steps sum to $S = \sum_{t=m}^{T-2} \eta_t$.

5.2.1 THE CORRIDOR FUNCTION g_{corridor}

The following lower bound shows that the corridor function g_{corridor} keeps the remaining primal gradients constant and nonzero.

Lemma 5.3 (A lower bound for g_{corridor}). *For $\ell, \Delta > 0$, $\rho \geq 1$, integers $T \geq 1$, $0 \leq m < T$, and nonnegative steps with $S = \sum_{t=m}^{T-2} \eta_t$, there is an ℓ -smooth function $g_{\text{corridor}} : \mathbb{R} \rightarrow \mathbb{R}$ with initial gap $g_{\text{corridor}}(0) - \inf_u g_{\text{corridor}}(u) = \Delta$. Gradient descent from $u_0 = 0$ with steps η_t/ρ satisfies $g'_{\text{corridor}}(u_t) < 0$ for all $t < T$ and $g'_{\text{corridor}}(u_t) = -G$ for $m \leq t < T$, where $G > 0$ and $G^2 \asymp \min\{\ell\Delta, \rho\Delta/S\}$. Consequentially,*

$$\frac{1}{T} \sum_{t < T} |g'_{\text{corridor}}(u_t)|^2 \gtrsim \frac{T-m}{T} \min \left\{ \ell\Delta, \frac{\rho\Delta}{S} \right\}. \quad (5.5)$$




Figure 1: **The competing bounds in f_{stable} (schematic).** Let m count the initial large steps and $S = \sum_{t=m}^{T-2} \eta_t$. (a) The corridor keeps gradient magnitude G at $u_m, \dots, u_{T-1}$. (b) Gradient noise creates inner gap $\Phi_\sigma(v_t) - g_\sigma(v_t, y_t)$. (c) Pink bands show contributions to $t^{-1} \sum_{i < t} \|\nabla \Phi_\sigma(v_i)\|^2$; the initial increases are omitted. The dashed level is the bound's scale $\kappa^4 \ell \sigma^2 S / (\rho T)$ for $S \geq C \rho / (\ell \kappa)$.

We illustrate the corridor function g_{corridor} in Figure 1 (a). Intuitively, it consists of an entry curve and a constant-slope segment, followed by an exit curve. The entry curve brings the first m updates to the corridor, and the remaining iterates travel its length GS/ρ at slope $-G$, using gap $G^2 S/\rho$. The exit curve brings the slope to zero; with curvature of magnitude ℓ , the entry and exit curves together use gap of order G^2/ℓ . Hence $G^2(S/\rho + 1/\ell) \asymp \Delta$ gives the stated lower bound (Appendix C.2).

While Lemma 5.3 provides a lower bound for small cumulative steps S , it weakens as S grows. We complement this by constructing the function g_σ which induces a lower bound that grows with S .

5.2.2 THE LASTING EFFECT OF INNER PERTURBATION

To construct a noisy function g_σ whose lower bound grows with the cumulative steps S , we study an inner perturbation that causes a long-lasting outer error. To make our exposition clear, we represent the perturbation by a deterministic shift $\delta > 0$ in a simple linear-quadratic game $g_\delta : \mathbb{R} \times \mathbb{R}^2 \rightarrow \mathbb{R}$:

$$g_\delta(z; a, b) = \ell z a - \frac{\ell}{2\kappa} [a^2 + (a + b + \delta)^2] \quad (5.6)$$

with an outer variable z and inner variables a, b . Here, the inner maximizer $(\kappa z, -\kappa z - \delta)$ gives outer objective $\Phi_{g_\delta}(z) = \ell \kappa z^2 / 2$; hence, the inner perturbation δ does not affect the outer objective. From the origin, however, an ascent step follows the negative derivative $\partial_a g_\delta = -\ell \delta / \kappa$ and makes the inner variable a negative. A descent step then follows the derivative $-\partial_z g_\delta = -\ell a > 0$ and moves the outer variable z away from zero, which is the minimizer of the outer objective Φ_{g_δ} . Consequentially, the inner perturbation δ induces an outer error through GDA dynamics.

In the high-ratio regime $\rho = \Omega(\kappa^2)$, such an outer error takes longer to remove than to create. To see this clearly, let us measure the time by cumulative steps $\sum_{i < t} \eta_i$. The outer variable z moves slowly, so we may approximate its effect by fixing $z = 0$. Then, the inner gradients follow $\partial_a g_\delta = -(\ell/\kappa)(2a + b + \delta)$ and $\partial_b g_\delta = -(\ell/\kappa)(a + b + \delta)$. In particular, the gradients start at $-\ell \delta / \kappa$ and remain negative while $|a| + |b|$ is small relative to δ . Therefore, a reaches a negative constant fraction of δ on the cumulative timescale κ/ℓ ; then it returns to zero as b approaches $-\delta$. Substituting this response into $z_{i+1} - z_i = -\ell \eta_i a_i / \rho$ predicts the scale

$$\underbrace{\frac{\ell \delta}{\rho}}_{\text{outer speed}} \times \underbrace{\frac{\kappa}{\ell}}_{\text{formation time}} = \underbrace{\frac{\kappa \delta}{\rho}}_{\text{error size}}.$$

After the inner variables adapt enough to the perturbation δ , the inner maximizer gives the approximation $a \approx \kappa z$, so we obtain an outer contraction $z_{i+1} \approx (1 - \ell \kappa \eta_i / \rho) z_i$. This suggests that removing a fixed fraction of the error induced by an inner perturbation takes cumulative steps of order $\rho / (\ell \kappa)$, a factor ρ / κ^2 longer than its formation time. The following lemma captures this intuition and bounds the error on the outer variable z_t , and specifies the interval over which this error persists.

Lemma 5.4 (Persistent error under a deterministic shift). *There are universal constants $c, C, C_0 > 0$ such that, for $\ell, \delta > 0$, $\kappa \geq 1$, $\rho \geq C_0 \kappa^2$, and any finite step-size sequence with $0 \leq \eta_t \leq c\kappa$, both SGDA_{Sim} and SGDA_{Alt} with exact gradients on the linear quadratic game g_δ given by (5.6), initialized at $(0, 0, 0)$, satisfy $|z_t| \leq C\kappa\delta/\rho$ and $|a_t| + |b_t| \leq C\delta$ whenever $\sum_{i < t} \eta_i \leq c\rho/(\ell\kappa)$, and*

$$z_t \geq c \frac{\kappa\delta}{\rho} \quad \text{if} \quad C \frac{\kappa}{\ell} \leq \sum_{i < t} \eta_i \leq c \frac{\rho}{\ell\kappa}. \quad (5.7)$$

5.2.3 THE NOISY FUNCTION g_σ

We now construct the noisy function g_σ with its coordinate-wise components $g_{\sigma,j}$, each of which implements g_δ through a stochastic noise oracle. Specifically, we assign each update $j \in J := \{0, \dots, T-2\}$ a separate component $g_{\sigma,j} : \mathbb{R} \times \mathbb{R}^3 \rightarrow \mathbb{R}$ with $y = (a, b, s)$, initialized at the origin in coordinates $v[j], y[j] = (a[j], b[j], s[j])$. We denote its outer objective by $\Phi_{\sigma,j}$, choose the constants $K \asymp \kappa$ and $L \asymp \ell$ with fixed universal factors, and set $g_{\sigma,j} \equiv 0$ if $\eta_j \sigma = 0$.

$$g_\sigma(v, y) = \sum_{j \in J} g_{\sigma,j}(v[j]; y[j]), \quad (5.8)$$

$$g_{\sigma,j}(v; a, b, s) := Lva - \frac{L}{2K} \left[a^2 + (a + b + \chi_j(a + b)\delta_j(s))^2 \right], \quad (5.9)$$

$$\delta_j(s) := \sqrt{K} \eta_j \sigma \sin^2 \left(\frac{\pi s}{2\eta_j \sigma} \right). \quad (5.10)$$

The smooth cutoff $\chi_j : \mathbb{R} \rightarrow [0, 1]$ equals one along the trajectory segment used in Lemma 5.4, preserving g_δ with parameters L, K and inner perturbation $\delta_j(s)$. The cutoff function χ_j and the choice of constants K, L ensure ℓ -smoothness and the inner (ℓ/κ) -PL inequality (Appendix C.3.3).

Our stochastic oracle induces a gradient error $\pm \sigma$ in $s[j]$, and SGDA turns that error into $s[j] = \pm \eta_j \sigma$. Specifically, let $e(y)$ be the unit vector in the $s[j]$ -coordinate for the smallest $j \in J$ with $s[j] = 0$, or zero if none exists. With fresh independent uniform signs $\xi \in \{-1, 1\}$, we introduce a stochastic oracle

$$G_x = \nabla_x f_{\text{stable}}, \quad G_y = \nabla_y g_\sigma + \sigma \xi e(y). \quad (5.11)$$

Since the derivative of δ_j vanishes at $s = \pm \eta_j \sigma$, each $s[j]$ stays fixed once activated, while the remaining s -coordinates stay at the origin; only $s[j]$ receives noise. The oracle has variance $\sigma^2 \|e(y)\|^2 \leq \sigma^2$, and the estimators in (5.11) are pointwise unbiased, so they satisfy Assumptions 3.4–3.5.

For $j \geq m$ with enough remaining cumulative steps to cover $\asymp \rho/(\ell\kappa)$, Lemma 5.4 and curvature $\Phi''_{\sigma,j} \asymp \ell\kappa$ give an outer gradient $\Phi'_{\sigma,j}(v_t[j]) \gtrsim \ell\kappa^{5/2} \eta_j \sigma / \rho$. Subsequent steps are at most η_j due to the non-increasing step size schedule, hence, the following lower bound holds:

$$\sum_{t < T} \mathbb{E} |\Phi'_{\sigma,j}(v_t[j])|^2 \gtrsim \underbrace{\left(\frac{\ell\kappa^{5/2} \eta_j \sigma}{\rho} \right)^2}_{\text{squared outer gradient}} \times \underbrace{\frac{\rho}{\ell\kappa \eta_j}}_{\text{number of iterates}} = \frac{\ell\kappa^4 \sigma^2 \eta_j}{\rho}. \quad (5.12)$$

The next lemma captures this derivation, and lower bounds the outer gradient accumulation from a single inner perturbation.

Lemma 5.5 (Contribution of a single inner-gradient perturbation). *There are universal constants $c, C, c_{\text{step}}, c_{\text{high}} > 0$ and $\kappa_{\text{high}} \geq 2$ such that, for $\ell, \sigma > 0$, $\kappa \geq \kappa_{\text{high}}$, $\rho \geq \max\{1, c_{\text{high}} \kappa^2\}$, an integer $T \geq 2$, any predetermined step sizes $\eta_0 \geq \dots \geq \eta_{T-2} \geq 0$ and index $0 \leq j \leq T-2$ with $0 < \ell\eta_j \leq c_{\text{step}} \kappa$ admit an ℓ -smooth NC- (ℓ/κ) -PL component $g_{\sigma,j} : \mathbb{R} \times \mathbb{R}^3 \rightarrow \mathbb{R}$. Its outer objective $\Phi_{\sigma,j}(v) := \max_{y \in \mathbb{R}^3} g_{\sigma,j}(v; y)$ is quadratic with curvature $\Phi''_{\sigma,j} \asymp \ell\kappa$ and initial gaps $\Phi_{\sigma,j}(0) - \min_v \Phi_{\sigma,j}(v) = \Phi_{\sigma,j}(0) - g_{\sigma,j}(0; 0) = 0$. Starting from the origin with one equiprobable $\pm \sigma$ error in the third inner gradient at update j and exact gradients otherwise, both SGDA_{Sim} and SGDA_{Alt} generate outer iterates $v_t[j]$ satisfying*

$$\sum_{t < T} \mathbb{E} |\Phi'_{\sigma,j}(v_t[j])|^2 \geq c \frac{\ell\kappa^4 \sigma^2 \eta_j}{\rho} \quad \text{if} \quad \sum_{t=j+1}^{T-2} \eta_t \geq C \frac{\rho}{\ell\kappa}. \quad (5.13)$$

5.2.4 COMBINING THE CORRIDOR FUNCTION g_{corridor} AND NOISY FUNCTION g_{σ}

With enough remaining cumulative steps, $S \geq C\rho/(\ell\kappa)$, we can sum Lemma 5.5, omitting late perturbations with insufficient subsequent cumulative steps; their total step size is $O(\rho/(\ell\kappa))$. The corridor covers the other case of small remaining cumulative steps. The following lemma combines these bounds and captures a tradeoff in S (Appendix C.4).

Lemma 5.6 (Stable-step lower bound and admissibility). *There are universal $c, c_{\text{high}} > 0$ and $\kappa_{\text{high}} \geq 2$ such that, for $\ell, \Delta > 0$, $\sigma \geq 0$, $\kappa \geq \kappa_{\text{high}}$, $\rho \geq \max\{1, c_{\text{high}}\kappa^2\}$, an integer $T \geq 1$, and predetermined steps $\eta_0 \geq \dots \geq \eta_{T-2} \geq 0$, the following holds. Let m count the entries $\eta_t \geq 8/\ell$ and assume $m+1 < T/2$. For each of SGDA_{Sim} and SGDA_{Alt} , the game f_{stable} given by (5.4) and a fixed oracle can be chosen as an instance in $\mathcal{C}(\ell, \kappa, \Delta, \sigma)$ with initial outer gap Δ and inner gap zero at the origin. Writing $S = \sum_{t=m}^{T-2} \eta_t$ and $\rho\Delta/0 := +\infty$, its iterates satisfy*

$$\frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E} \|\nabla \Phi_{\text{stable}}(u_t, v_t)\|^2 \geq c \min \left\{ \ell\Delta, \max \left\{ \frac{\rho\Delta}{S}, \frac{\kappa^4 \ell \sigma^2 S}{\rho T} \right\} \right\}. \quad (5.14)$$

Figure 1 illustrates the tradeoff suggested by Lemma 5.6: increasing S weakens the corridor bound $\rho\Delta/S$ but strengthens the stochastic bound $\kappa^4 \ell \sigma^2 S/(\rho T)$. Applying the AM–GM inequality and $S < 8T/\ell$ to Lemma 5.6 proves Theorem 5.1 as follows:

$$\max \left\{ \frac{\rho\Delta}{S}, \frac{\ell\kappa^4 \sigma^2 S}{\rho T} \right\} \geq \max \left\{ \frac{\rho\ell\Delta}{8T}, \sqrt{\frac{\ell\kappa^4 \Delta \sigma^2}{T}} \right\}. \quad (5.15)$$

6 LOW-RATIO LOWER BOUND

In the low ratio regime $\rho = O(\kappa^2)$, inner variables may fail to track their maximizer due to relatively small inner step sizes. In particular, Mourrat (2026) recently constructed a PL–PL game that exhibits a periodic orbit of a continuous-time GDA flow, suggesting that a PL inequality can fail to prevent limit cycles of GDA. Below, we realize a similar construction for discrete-time GDA in NC–PL games.

6.1 INNER PL DOES NOT PREVENT A LIMIT CYCLE

We start with a simple bilinear game that creates a periodic orbit for GDA:

$$g(u, a) = \sqrt{2} u^T a, \quad u, a \in \mathbb{R}^2. \quad (6.1)$$

For timescale ratio $\rho = 1$, GDA flow oscillates in this game (Hsieh et al., 2021), but the infinite inner maximum for $u \neq 0$ violates NC–PL. Adding $-\|a\|^2$ gives the outer objective $\|u\|^2/2$ yet damps the flow: $\ddot{u} + 2\dot{u} + 2u = 0$. Hence, we introduce $b \in \mathbb{R}^2$ and a residual $R : \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow \mathbb{R}^2$ and define

$$g_{\text{cycle}}(u, a, b) = \sqrt{2} u^T a - \|a\|^2 - \frac{1}{2} \|R(a, b)\|^2. \quad (6.2)$$

Designing the residual $R(a, b)$. We want the residual $R(a, b)$ to restore a periodic motion while preserving an inner PL inequality and the quadratic outer objective $\|u\|^2/2$. If every a admits b that satisfies $R(a, b) = 0$, maximization over (a, b) can cancel the residual and leave the outer objective unchanged. On the other hand, the inner PL inequality follows when the map $(a, b) \mapsto (\sqrt{2}a, R(a, b))$ has a global smooth inverse with a uniformly bounded derivative (c.f. Lemma B.1). To specify the periodic motion we want g_{cycle} to induce, let J denote counterclockwise rotation by $\pi/2$, and define

$$u_*(t) = (\cos t, \sin t)^T, \quad a_*(t) = -Ju_*(t)/\sqrt{2}, \quad b_*(t) = u_*(t)/\sqrt{2}. \quad (6.3)$$

Our goal is to design the residual $R(a, b)$ for GDA to follow this path. Outer descent already has the required velocity along this path, since $\dot{u} = Ju = -\sqrt{2}a$. The inner variables must have velocities $\dot{a} = Ja$ and $\dot{b} = -a$. Hence, to obtain the desired inner motion, we can impose the restriction

$$\nabla_a \left(\frac{1}{2} \|R(a, b)\|^2 \right) = -2a + Ja, \quad \nabla_b \left(\frac{1}{2} \|R(a, b)\|^2 \right) = a, \quad (6.4)$$

on the circle $b = Ja$, $\|a\| = 1/\sqrt{2}$ traced by the motion (a_*, b_*) . With these identities, the GDA equations agree with the target path. We choose further derivatives so that nearby trajectories approach the orbit, and use cutoffs for global smoothness without changing these dynamics (c.f. Appendix D.1).

Lemma 6.1 (An attracting periodic orbit of GDA flow in g_{cycle}). *There are universal $C, \mu, h_\star > 0$ and a smooth residual R such that the base game $g_{\text{cycle}} : \mathbb{R}^2 \times \mathbb{R}^4 \rightarrow \mathbb{R}$ in (6.2) is jointly C -smooth and affine in u , has outer objective $\Phi_{g_{\text{cycle}}}(u) = \frac{1}{2} \|u\|^2$, and satisfies the inner μ -PŁ inequality. The path (6.3), with $y_\star = (a_\star, b_\star)$, traces a locally stable 2π -periodic orbit Γ of GDA flow. There is a compact neighborhood $\mathcal{U}$ of Γ such that both SGDA_{Sim} and SGDA_{Alt} , started in $\mathcal{U}$ with exact gradients and equal primal and dual steps $h_t \in [0, h_\star]$, satisfy $(u_t, y_t) \in \mathcal{U}$ and $\|\nabla \Phi_{g_{\text{cycle}}}(u_t)\| = \|u_t\| \geq \frac{1}{2}$ for all $t \geq 0$, where $y_t = (a_t, b_t)$.*

6.2 FROM THE CYCLE TO THE LOW-RATIO LOWER BOUND

Lemma 6.1 shows that continuous-time GDA can be trapped in a limit cycle in g_{cycle} . However, its discrete-time guarantee remains local and requires step sizes at most $h_\star$, which can be extremely small. The next lemma fills this gap by modifying the game, and shows that GDA can reach the orbit from a suitable initialization after any finite sequence of larger steps and remain nearby under subsequent small steps.

Lemma 6.2 (GDA can reach the orbit in a modified game $\tilde{g}_{\text{cycle}}$). *There are universal $C, \mu > 0$ with the following property. For any finite sequence of steps $h_0, \dots, h_{m-1} > h_\star$, including $m = 0$, we can modify the game g_{cycle} in Lemma 6.1 to obtain $\tilde{g}_{\text{cycle}} : \mathbb{R}^2 \times \mathbb{R}^5 \rightarrow \mathbb{R}$. This game is jointly C -smooth and affine in u , satisfies the inner μ -PŁ inequality, and has outer objective $\Phi_{\tilde{g}_{\text{cycle}}}(u) = \|u\|^2/2$. With $y = (a, b, r)$ and $e_1 = (1, 0)^\top$, there is an initialization (e_1, y_0) with inner gap at most C . From this initialization, both GDA_{Sim} and GDA_{Alt} , using exact gradients and equal primal and dual steps, keep $u_t = e_1$ for $0 \leq t \leq m$ and reach $y_m = (y_\star(0), 0)$. For every continuation with $h_t \in [0, h_\star]$, each algorithm follows its g_{cycle} dynamics in (u, a, b) , while r remains zero.*

We now rescale $\tilde{g}_{\text{cycle}}$ to satisfy ℓ -smoothness and the inner (ℓ/κ) -PŁ inequality. We choose $f_{\text{cycle}}(x, y) = A\tilde{g}_{\text{cycle}}(x/s_x, y/s_y)$ with positive scales satisfying $s_y^2 = \rho s_x^2$, which converts both updates at ratio ρ to equal-step dynamics. The scales in Appendix A.3 make f_{cycle} ℓ -smooth with inner PŁ constant of order $\ell/\sqrt{\rho}$, so $\rho \leq c_{\text{low}}\kappa^2$ ensures inner PŁ at least ℓ/κ . They also meet the initial-gap budgets and give $\|\nabla \Phi_{\text{cycle}}(x_t)\|^2 \gtrsim \ell\Delta$ at every iterate, proving the following theorem.

Theorem 6.3 (Necessity of an order- κ^2 fixed ratio). *There are universal $c, c_{\text{low}} > 0$ and $\kappa_{\text{low}} \geq 2$ such that, for every $\kappa \geq \kappa_{\text{low}}$, $\ell, \Delta > 0$, $\sigma \geq 0$, integer $T \geq 1$, $1 \leq \rho \leq c_{\text{low}}\kappa^2$, and predetermined nonnegative non-increasing step sizes, each of SGDA_{Sim} and SGDA_{Alt} admits a deterministic instance f_{cycle} in $\mathcal{C}(\ell, \kappa, \Delta, \sigma)$ with $d_x = 2$, $d_y = 5$, a strongly convex quadratic outer objective, and*

$$\min_{0 \leq t < T} \|\nabla \Phi(x_t)\|^2 \geq c\ell\Delta. \quad (6.5)$$

Crucially, the theorem answers the question raised by [Cho and Yun \(2023\)](#) affirmatively: *timescale ratio $\rho = \Omega(\kappa^2)$ is necessary for SGDA to converge in NC-PŁ games*. In particular, it shows that smaller timescale ratios $\rho = o(\kappa^2)$ cannot provide convergence guarantees in general, even with deterministic gradients and non-increasing step sizes. Together with Theorem 5.1, it yields our tight main lower bound in Theorem 4.1.

7 CONCLUSION

In this work, we established the first tight lower bounds of two-timescale SGDA with a fixed timescale ratio and nonincreasing step sizes in NC-PŁ games. The lower bounds explain the fundamental limitation of two-timescale SGDA: small ratios can keep outer gradients bounded away from zero, while large ratios leave a tradeoff between outer progress and errors induced by inner noise. Our negative result justifies the recent line of work developing alternative methods, such as Smoothed-AGDA ([Yang et al., 2022](#)) and methods based on momentum ([Huang et al., 2025](#); [Cai et al., 2026](#)).

On the other hand, our analysis only applies to SGDA with a fixed timescale ratio and nonincreasing step sizes; therefore, it does not rule out nonmonotone or signed step size schedules ([Altschuler and Parrilo, 2025](#); [Shugart and Altschuler, 2025](#)) that might improve SGDA’s complexity. An additional interesting direction to explore would be to extend the ideas behind our lower bounds to nonconvex-concave problems. We leave these two questions as intriguing future work.

AI USE STATEMENT

Generative AI tools were used to brainstorm and refine theoretical constructions, formulate and check proof strategies, survey relevant literature, draft and edit mathematical exposition, and prepare and check the $\LaTeX$ source. The authors are responsible for the final content, including all AI-assisted text, claims, proofs, and artifacts.

REPRODUCIBILITY STATEMENT

All assumptions and algorithms are stated in [Section 3](#). [Appendix A](#) contains the proofs of the main theorem, the high- and low-ratio lower bounds, and the matching SGDA_{Sim} upper bound. [Appendices C and D](#) prove the construction lemmas. Together, these proofs give the construction formulas, parameter choices, oracle definitions, initialization checks, and arguments for SGDA_{Sim} and SGDA_{Alt} .

REFERENCES

- Jason M. Altschuler and Pablo A. Parrilo. Acceleration by stepsize hedging: Silver Stepsize Schedule for smooth convex optimization. *Mathematical Programming*, 213:1105–1118, 2025. doi: 10.1007/s10107-024-02164-2. URL <https://doi.org/10.1007/s10107-024-02164-2>.
- Martino Bernasconi, Matteo Castiglioni, Andrea Celli, and Alexandros Hollender. The complexity of min-max optimization for quadratic polynomials. *arXiv preprint arXiv:2606.17000*, 2026. URL <https://arxiv.org/abs/2606.17000>.
- Axel Böhm. Solving nonconvex-nonconcave min-max problems exhibiting weak Minty solutions. *Transactions on Machine Learning Research*, 2023. URL <https://openreview.net/forum?id=Gp0pHyUyrb>.
- Haoyuan Cai, Sulaiman A. Alghunaim, and Ali H. Sayed. Accelerated stochastic min-max optimization based on bias-corrected momentum. *arXiv preprint arXiv:2406.13041v3*, 2026. URL <https://arxiv.org/abs/2406.13041v3>.
- Lesi Chen, Boyuan Yao, and Luo Luo. Faster stochastic algorithms for minimax optimization under Polyak–Łojasiewicz condition. In *Advances in Neural Information Processing Systems*, volume 35, 2022. URL <https://arxiv.org/abs/2307.15868>.
- Hanseul Cho and Chulhee Yun. SGDA with shuffling: Faster convergence for nonconvex-PL minimax optimization. In *International Conference on Learning Representations*, 2023. URL <https://iclr.cc/virtual/2023/poster/11501>.
- Constantinos Daskalakis, Stratis Skoulakis, and Manolis Zampetakis. The complexity of constrained min-max optimization. In *Proceedings of the 53rd Annual ACM SIGACT Symposium on Theory of Computing*, 2021. doi: 10.1145/3406325.3451125. URL <https://arxiv.org/abs/2009.09623>.
- Jelena Diakonikolas, Constantinos Daskalakis, and Michael I. Jordan. Efficient methods for structured nonconvex-nonconcave min-max optimization. In *Proceedings of the 24th International Conference on Artificial Intelligence and Statistics*, volume 130 of *Proceedings of Machine Learning Research*, pages 2746–2754. PMLR, 2021. URL <https://proceedings.mlr.press/v130/diakonikolas21a.html>.
- Tanner Fiez and Lillian J. Ratliff. Local convergence analysis of gradient descent ascent with finite timescale separation. In *International Conference on Learning Representations*, 2021. URL https://openreview.net/forum?id=AWOSz_mMAPx.
- Tanner Fiez, Lillian J. Ratliff, Eric Mazumdar, Evan Faulkner, and Adhyayan Narang. Global convergence to local minmax equilibrium in classes of nonconvex zero-sum games. In *Advances in Neural Information Processing Systems*, volume 34, 2021.

- Ian J. Goodfellow, Jean Pouget-Abadie, Mehdi Mirza, Bing Xu, David Warde-Farley, Sherjil Ozair, Aaron Courville, and Yoshua Bengio. Generative adversarial nets. In *Advances in Neural Information Processing Systems*, volume 27, 2014. URL <https://arxiv.org/abs/1406.2661>.
- Benjamin Grimmer, Haihao Lu, Pratik Worah, and Vahab Mirrokni. The landscape of the proximal point method for nonconvex-nonconcave minimax optimization. *Mathematical Programming*, 201(1–2):373–407, 2023. doi: 10.1007/s10107-022-01910-8. URL <https://doi.org/10.1007/s10107-022-01910-8>.
- Junsoo Ha. High probability convergence guarantees of stochastic gradient descent ascent in structured nonconvex min-max games. In *Proceedings of Thirty Ninth Conference on Learning Theory*, volume 336 of *Proceedings of Machine Learning Research*, pages 3005–3075, 2026. URL <https://proceedings.mlr.press/v336/ha26a.html>.
- Saeed Hajizadeh, Haihao Lu, and Benjamin Grimmer. On the linear convergence of extragradient methods for nonconvex-nonconcave minimax problems. *INFORMS Journal on Optimization*, 6(1):19–31, 2024. doi: 10.1287/ijoo.2022.0004. URL <https://doi.org/10.1287/ijoo.2022.0004>.
- Ya-Ping Hsieh, Panayotis Mertikopoulos, and Volkan Cevher. The limits of min-max optimization algorithms: Convergence to spurious non-critical sets. In *Proceedings of the 38th International Conference on Machine Learning*, volume 139 of *Proceedings of Machine Learning Research*, pages 4337–4348, 2021. URL <https://proceedings.mlr.press/v139/hsieh21a.html>.
- Feihu Huang, Chunyu Xuan, Xinrui Wang, Siqi Zhang, and Songcan Chen. Enhanced adaptive gradient algorithms for nonconvex-PL minimax optimization. In *Proceedings of the 28th International Conference on Artificial Intelligence and Statistics*, volume 258 of *Proceedings of Machine Learning Research*, pages 3439–3447, 2025. URL <https://proceedings.mlr.press/v258/huang25d.html>.
- Yassine Laguel, Yasa Syed, Necdet Serhat Aybat, and Mert Gürbüzbalaban. High-probability complexity bounds for stochastic non-convex minimax optimization. In *Advances in Neural Information Processing Systems*, volume 37, 2024. URL <https://arxiv.org/abs/2405.14130>.
- Sucheol Lee and Donghwan Kim. Fast extra gradient methods for smooth structured nonconvex-nonconcave minimax problems. In *Advances in Neural Information Processing Systems*, volume 34, pages 22588–22600, 2021. URL <https://proceedings.neurips.cc/paper/2021/hash/be767243ca8f574c740fb4c26cc6dceb-Abstract.html>.
- Haochuan Li, Yi Tian, Jingzhao Zhang, and Ali Jadbabaie. Complexity lower bounds for nonconvex-strongly-concave min-max optimization. In *Advances in Neural Information Processing Systems*, volume 34, 2021. URL https://proceedings.neurips.cc/paper_files/paper/2021/hash/0e105949d99a32ca1751703e94ece601-Abstract.html.
- Haochuan Li, Farzan Farnia, Subhro Das, and Ali Jadbabaie. On convergence of gradient descent ascent: A tight local analysis. In *Proceedings of the 39th International Conference on Machine Learning*, volume 162 of *Proceedings of Machine Learning Research*, pages 12717–12740, 2022. URL <https://proceedings.mlr.press/v162/li22e.html>.
- Tianyi Lin, Chi Jin, and Michael I. Jordan. On gradient descent ascent for nonconvex-concave minimax problems. In *Proceedings of the 37th International Conference on Machine Learning*, volume 119 of *Proceedings of Machine Learning Research*, pages 6083–6093, 2020. URL <https://proceedings.mlr.press/v119/lin20a.html>.
- Tianyi Lin, Chi Jin, and Michael I. Jordan. Two-timescale gradient descent ascent algorithms for nonconvex minimax optimization. *Journal of Machine Learning Research*, 26(11):1–45, 2025. URL <https://jmlr.org/papers/v26/22-0863.html>.

- Mingrui Liu, Hassan Rafique, Qihang Lin, and Tianbao Yang. First-order convergence theory for weakly-convex-weakly-concave min-max problems. *Journal of Machine Learning Research*, 22 (169):1–34, 2021. URL <https://jmlr.org/papers/v22/20-533.html>.
- Qihao Liu, Luoxin Ye, Wufei Ma, Yu-Cheng Chou, and Alan Yuille. Generative adversarial reasoner: Enhancing LLM reasoning with adversarial reinforcement learning. In *International Conference on Learning Representations*, 2026. URL <https://arxiv.org/abs/2512.16917>.
- Panayotis Mertikopoulos, Bruno Lecouat, Houssam Zenati, Chuan-Sheng Foo, Vijay Chandrasekhar, and Georgios Piliouras. Optimistic mirror descent in saddle-point problems: Going the extra (gradient) mile. In *International Conference on Learning Representations*, 2019. URL <https://openreview.net/forum?id=Bkg8jjC9KQ>.
- Jean-Christophe Mourrat. PL conditions do not guarantee convergence of gradient descent-ascent dynamics. *arXiv preprint arXiv:2602.16517*, 2026. URL <https://arxiv.org/abs/2602.16517>.
- John Nash. Non-cooperative games. *Annals of Mathematics*, 54(2):286–295, 1951. doi: 10.2307/1969529.
- Maher Nouiehed, Maziar Sanjabi, Tianjian Huang, Jason D. Lee, and Meisam Razaviyayn. Solving a class of non-convex min-max games using iterative first order methods. In *Advances in Neural Information Processing Systems*, volume 32, 2019.
- Siyu Pan and Jiajin Li. Lower bounds for nonconvex-PL minimax optimization. *arXiv preprint arXiv:2608.26799*, 2026. URL <https://arxiv.org/abs/2608.26799>.
- Thomas Pethick, Puya Latafat, Panagiotis Patrinos, Olivier Fercoq, and Volkan Cevher. Escaping limit cycles: Global convergence for constrained nonconvex-nonconcave minimax problems. In *International Conference on Learning Representations*, 2022. URL https://openreview.net/forum?id=2_vhkAMARK.
- Boris T. Polyak. Gradient methods for minimizing functionals. *USSR Computational Mathematics and Mathematical Physics*, 3(4):864–878, 1963. doi: 10.1016/0041-5553(63)90382-3.
- Yuzuru Sato, Eizo Akiyama, and J. Dooyne Farmer. Chaos in learning a simple two-person game. *Proceedings of the National Academy of Sciences*, 99(7):4748–4751, 2002. doi: 10.1073/pnas.032086299. URL <https://pmc.ncbi.nlm.nih.gov/articles/PMC123719/>.
- Henry Shugart and Jason M. Altschuler. Negative stepsizes make Gradient-Descent-Ascent converge. *arXiv preprint arXiv:2505.01423*, 2025. doi: 10.48550/arXiv.2505.01423. URL <https://arxiv.org/abs/2505.01423>.
- Aman Sinha, Hongseok Namkoong, and John Duchi. Certifying some distributional robustness with principled adversarial training. In *International Conference on Learning Representations*, 2018. URL <https://arxiv.org/abs/1710.10571v4>.
- Kiran K. Thekumparampil, Prateek Jain, Praneeth Netrapalli, and Sewoong Oh. Efficient algorithms for smooth minimax optimization. In *Advances in Neural Information Processing Systems*, volume 32, 2019. URL <https://arxiv.org/abs/1907.01543>.
- Zaiyan Xu, Sushil Vemuri, Kishan Panaganti, Dileep Kalathil, Rahul Jain, and Deepak Ramachandran. Robust LLM alignment via distributionally robust direct preference optimization. In *Advances in Neural Information Processing Systems*, volume 38, 2025. URL <https://arxiv.org/abs/2502.01930>.
- Junchi Yang, Negar Kiyavash, and Niao He. Global convergence and variance reduction for a class of nonconvex-nonconcave minimax problems. In *Advances in Neural Information Processing Systems*, volume 33, 2020. URL https://papers.neurips.cc/paper_files/paper/2020/hash/0cc6928e741d75e7a92396317522069e-Abstract.html.

Junchi Yang, Antonio Orvieto, Aurélien Lucchi, and Niao He. Faster single-loop algorithms for minimax optimization without strong concavity. In *Proceedings of the 25th International Conference on Artificial Intelligence and Statistics*, volume 151 of *Proceedings of Machine Learning Research*, pages 5485–5517, 2022. URL <https://proceedings.mlr.press/v151/yang22b.html>.

Siqi Zhang, Junchi Yang, Cristóbal Guzmán, Negar Kiyavash, and Niao He. The complexity of nonconvex-strongly-concave minimax optimization. In *Proceedings of the Thirty-Seventh Conference on Uncertainty in Artificial Intelligence*, volume 161 of *Proceedings of Machine Learning Research*, pages 482–492, 2021. URL <https://proceedings.mlr.press/v161/zhang21c.html>.

APPENDIX CONTENTS

A	Proof of main result	15
A.1	Proof of Theorem 4.1	15
A.2	Proof of Theorem 5.1	15
A.3	Proof of Theorem 6.3	16
A.4	Upper bound for SGDA_{Sim}	17
B	Preliminary lemmas	19
B.1	NC-PŁ games with a predetermined outer function	19
B.2	NC-PŁ geometry	19
C	Proofs for the high-ratio lower bound	20
C.1	Proof of the lower bound for f_{unstable}	20
C.2	Construction of g_{corridor}	21
C.3	Construction of g_{σ}	22
C.4	Proof of the lower bound for f_{stable}	28
D	Proofs for the low-ratio lower bound	29
D.1	Construction of g_{cycle}	29
D.2	Construction of $\tilde{g}_{\text{cycle}}$	32
E	Comparison with Smoothed-AGDA	34
E.1	Initialization budget of the hard instances	37
F	Complexity calculations for Table 1	37
F.1	SGDA and Smoothed-AGDA	38
F.2	Finite-sum upper bounds	38
F.3	MSGDA and AdaMSGDA	39
F.4	HCMM-2	40
F.5	Lower bounds and the nonconvergence entries	41

A PROOF OF MAIN RESULT

We first combine the two ratio regimes to prove Theorem 4.1. We then prove Theorems 5.1 and 6.3 using the construction lemmas from Sections 5 and 6, and establish the matching upper bound for SGDA_{Sim} .

A.1 PROOF OF THEOREM 4.1

Theorem 4.1 (A lower bound for two-timescale SGDA in NC-PŁ games). *There are universal constants $c > 0$ and $\kappa_0 \geq 2$, depending on c_{init} , such that, for every $\kappa \geq \kappa_0$, $\ell > 0$, $\Delta > 0$, $\sigma \geq 0$, and integer $T \geq 1$,*

$$\mathcal{R}_{T,\text{Sim}}^*, \mathcal{R}_{T,\text{Alt}}^* \geq c \min \left\{ \ell \Delta, \max \left\{ \frac{\kappa^2 \ell \Delta}{T}, \sqrt{\frac{\kappa^4 \ell \Delta \sigma^2}{T}} \right\} \right\}. \quad (4.1)$$

Proof. We first choose the construction constants so that the two ratio ranges overlap. The fixed cycle and its extension in Lemma 6.2 determine a universal inner-PŁ constant μ . Let C_{low} be the smoothness-normalization constant chosen in Appendix A.3; the low-ratio construction then applies up to $c_{\text{low}} = (\mu/C_{\text{low}})^2$. These constants are fixed before we calibrate the noise component. Let C_{resp} be the ratio constant C_0 in Lemma 5.4, enlarged as in Appendix C.3.3, let c_0, C_0 be the constants from Lemma C.1, and choose

$$0 < \gamma \leq \min \left\{ \sqrt{\frac{c_{\text{low}}}{C_{\text{resp}}}}, \frac{c_0}{2C_0} \right\}, \quad \theta := \frac{\gamma}{c_0}.$$

These choices calibrate only the high-ratio block: they give (C.23) and $c_{\text{high}} = \gamma^2 C_{\text{resp}} \leq c_{\text{low}}$. Thus its ratio range begins no later than the endpoint of the low-ratio range. The low-ratio constants do not depend on γ or θ .

Choose κ_0 above both theorem thresholds. In the high-ratio bound, $\rho \ell \Delta / T \geq c_{\text{high}} \kappa^2 \ell \Delta / T$; the stochastic term already matches (4.1). The low-ratio bound supplies the cap $c \ell \Delta$. Since the ranges cover every $\rho \geq 1$, taking the ratio infimum proves (4.1).

For $\varepsilon^2 \leq c \ell \Delta$ with sufficiently small universal $c > 0$, inversion gives the lower complexity bound in (4.2). The matching constant-step upper bounds cited in Section 4 complete the proof. $\square$

A.2 PROOF OF THEOREM 5.1

Theorem 5.1 (High-ratio lower bound). *There are universal constants $c, c_{\text{high}} > 0$ and $\kappa_{\text{high}} \geq 2$ such that, for every choice of problem parameters $\ell, \Delta > 0$, $\sigma \geq 0$, and $\kappa \geq \kappa_{\text{high}}$, time horizon $T \in \mathbb{N}_{\geq 1}$, and timescale ratio $\rho \geq c_{\text{high}} \kappa^2$, both SGDA_{Sim} and SGDA_{Alt} satisfy*

$$\mathcal{R}_{T,\text{Sim}}(\rho), \mathcal{R}_{T,\text{Alt}}(\rho) \geq c \min \left\{ \ell \Delta, \max \left\{ \frac{\rho \ell \Delta}{T}, \sqrt{\frac{\kappa^4 \ell \Delta \sigma^2}{T}} \right\} \right\}. \quad (5.1)$$

Proof. For each of SGDA_{Sim} and SGDA_{Alt} with predetermined non-increasing step sizes, let m count the initial excessively large steps $\ell \eta_t \geq 8$ and set $S = \sum_{t=m}^{T-2} \eta_t$. If $m+1 \geq T/2$, Lemma 5.2 gives an average squared outer gradient at least $c \ell \Delta$ on f_{unstable} , proving the capped bound. Otherwise, Lemma 5.6 applies to f_{stable} and also gives the cap when $S = 0$. For $S > 0$, the remaining steps satisfy $S \leq 8T/\ell$, while the maximum of two nonnegative numbers is at least their geometric mean. Hence

$$\max \left\{ \frac{\rho \Delta}{S}, \frac{\kappa^4 \ell \sigma^2 S}{\rho T} \right\} \geq \max \left\{ \frac{\rho \ell \Delta}{8T}, \sqrt{\frac{\kappa^4 \ell \Delta \sigma^2}{T}} \right\}. \quad (\text{A.1})$$

Substituting into (5.14) and adjusting the universal constant proves (5.1). In each case, the instance is chosen after fixing the schedule. Taking the instance supremum and then the schedule infimum proves the theorem. $\square$

A.3 PROOF OF THEOREM 6.3

Theorem 6.3 (Necessity of an order- κ^2 fixed ratio). *There are universal $c, c_{\text{low}} > 0$ and $\kappa_{\text{low}} \geq 2$ such that, for every $\kappa \geq \kappa_{\text{low}}, \ell, \Delta > 0, \sigma \geq 0$, integer $T \geq 1, 1 \leq \rho \leq c_{\text{low}}\kappa^2$, and predetermined nonnegative non-increasing step sizes, each of SGDA_{Sim} and SGDA_{Alt} admits a deterministic instance f_{cycle} in $\mathcal{C}(\ell, \kappa, \Delta, \sigma)$ with $d_x = 2, d_y = 5$, a strongly convex quadratic outer objective, and*

$$\min_{0 \leq t < T} \|\nabla \Phi(x_t)\|^2 \geq c\ell\Delta. \quad (6.5)$$

Proof. We rescale the fixed-dimensional game while retaining its strongly convex quadratic outer objective, then verify smoothness, inner PL, and the initial-gap budgets.

Let $C \geq 1$ bound the smoothness constant and initial inner gap in Lemma 6.2. Its construction also gives $\|\nabla_u \tilde{g}_{\text{cycle}}(e_1, y_0)\| \leq 1$, as verified in Appendix D.2. Choose a universal $C_{\text{low}} \geq 2C$, and fix $0 < \zeta \leq \min\{1/2, c_{\text{init}}/(3C)\}$. This choice also covers the regularized initial gap in Appendix E.1. For a ratio ρ , set

$$A := \frac{2\zeta\Delta}{\sqrt{\rho}}, \quad s_x^2 := \frac{2C_{\text{low}}\zeta\Delta}{\ell\rho}, \quad s_y^2 := \frac{2C_{\text{low}}\zeta\Delta}{\ell}. \quad (A.2)$$

For a fixed horizon and predetermined non-increasing step sizes (η_t) , put $h_t = \ell\eta_t/(C_{\text{low}}\sqrt{\rho})$. Let m count the initial steps with $h_t > h_*$. Apply Lemma 6.2 to those steps, writing $\tilde{g}_{\text{cycle}}$ for its game. For both SGDA_{Sim} and SGDA_{Alt} , the lemma keeps $u_t = e_1$ through time m and then identifies the continuation with the small-step dynamics of g_{cycle} . Hence Lemma 6.1 keeps all later states in $\mathcal{U} \times \{0\}$, so $\|u_t\| \geq 1/2$ at every iterate. This includes $m = 0, m = T - 1, T = 1$, and zero later steps.

Define the game $f_{\text{cycle}} : \mathbb{R}^2 \times \mathbb{R}^5 \rightarrow \mathbb{R}$ by

$$f_{\text{cycle}}(x, y) := A \tilde{g}_{\text{cycle}}\left(\frac{x}{s_x}, \frac{y}{s_y}\right), \quad (A.3)$$

initialized at $x_0 = s_x e_1$ and $y_0^{\text{cycle}} = s_y y_0$. Its outer objective and Hessian are

$$\Phi_{\text{cycle}}(x) = \frac{A}{2s_x^2} \|x\|^2 = \frac{\ell\sqrt{\rho}}{2C_{\text{low}}} \|x\|^2, \quad \nabla^2 \Phi_{\text{cycle}} = \frac{\ell\sqrt{\rho}}{C_{\text{low}}} I_2 \succ 0. \quad (A.4)$$

Thus the full outer objective, not merely one component, is strongly convex and quadratic.

After substituting $u = x/s_x$ and $v = y/s_y$ into the updates, both gradient coefficients equal

$$\frac{\eta_t A}{\rho s_x^2} = \frac{\eta_t A}{s_y^2} = \frac{\ell\eta_t}{C_{\text{low}}\sqrt{\rho}} = h_t. \quad (A.5)$$

Since f_{cycle} is affine in x , its Hessian with respect to x vanishes: $\nabla_{xx}^2 f_{\text{cycle}} = 0$. For every (x, y) , differentiating the partial gradients $\nabla_x f_{\text{cycle}}$ and $\nabla_y f_{\text{cycle}}$ with respect to y gives

$$\begin{aligned} \|\nabla_{xy}^2 f_{\text{cycle}}(x, y)\|_{\text{op}} &\leq C \frac{A}{s_x s_y} = C \frac{\ell}{C_{\text{low}}}, \\ \|\nabla_{yy}^2 f_{\text{cycle}}(x, y)\|_{\text{op}} &\leq C \frac{A}{s_y^2} = C \frac{\ell}{C_{\text{low}}\sqrt{\rho}}. \end{aligned}$$

The operator norm of the full Hessian is at most the sum of these two bounds, which is at most $2C\ell/C_{\text{low}} \leq \ell$ since $\rho \geq 1$. Thus the full gradient is ℓ -Lipschitz. If μ is the inner-PL constant of $\tilde{g}_{\text{cycle}}$, the scaled constant is $\mu A/s_y^2 = \mu\ell/(C_{\text{low}}\sqrt{\rho})$, which is at least ℓ/κ when

$$\rho \leq c_{\text{low}}\kappa^2, \quad c_{\text{low}} := \left(\frac{\mu}{C_{\text{low}}}\right)^2. \quad (A.6)$$

By Lemma 6.2, $\tilde{g}_{\text{cycle}}(x/s_x, \cdot)$ has a maximizer for every fixed x . Multiplying that maximizer by s_y gives a maximizer of $f_{\text{cycle}}(x, \cdot)$, since $A > 0$.

The initial outer gap $\Phi_{\text{cycle}}(x_0) - \inf_x \Phi_{\text{cycle}}(x)$ and inner gap $\Phi_{\text{cycle}}(x_0) - f_{\text{cycle}}(x_0, y_0^{\text{cycle}})$ satisfy the budgets in [Assumption 3.3](#):

$$\begin{aligned}\Phi_{\text{cycle}}(x_0) - \inf_x \Phi_{\text{cycle}}(x) &= \frac{A}{2} = \frac{\zeta \Delta}{\sqrt{\rho}} \leq \Delta, \\ \Phi_{\text{cycle}}(x_0) - f_{\text{cycle}}(x_0, y_0^{\text{cycle}}) &= A \left[\frac{1}{2} - \tilde{g}_{\text{cycle}}(e_1, y_0) \right] \leq CA \leq 2C\zeta\Delta \leq c_{\text{init}}\Delta.\end{aligned}\quad (\text{A.7})$$

The last inequalities use $\rho \geq 1$ and the chosen ζ . Finally, every iterate satisfies

$$\|\nabla \Phi_{\text{cycle}}(x_t)\|^2 = \frac{A^2}{s_x^2} \|u_t\|^2 = \frac{2\zeta\ell\Delta}{C_{\text{low}}} \|u_t\|^2 \geq \frac{\zeta}{2C_{\text{low}}} \ell\Delta. \quad (\text{A.8})$$

The instance therefore belongs to $\mathcal{C}(\ell, \kappa, \Delta, \sigma)$ for every $\sigma \geq 0$, using exact gradients. Its dimension is $(d_x, d_y) = (2, 5)$, independent of the horizon and schedule. Equation (A.8) proves the every-iterate bound (6.5). Averaging this inequality, taking the instance supremum, and then taking the schedule infimum in (3.7) gives $\mathcal{R}_{T,o}(\rho) \geq c\ell\Delta$ for both update orders. $\square$

A.4 UPPER BOUND FOR SGDA_{SIM}

We prove the matching upper bound for SGDA_{Sim} under the same assumptions and average squared-gradient criterion as the lower bound. The deterministic one-sided-PŁ counterpart is already established by [Chen et al. \(2022, Appendix B.2\)](#); the argument below includes unbiased stochastic gradients with bounded variance and our convention of averaging over $x_0, \dots, x_{T-1}$. The main issue is that $\nabla_x f(x_t, y_t)$ differs from $\nabla \Phi(x_t)$. Inner PŁ controls this discrepancy through the inner gradient by [Lemma B.2](#); the potential below uses a faster inner step to absorb it.

Theorem A.1 (Upper bound for SGDA_{Sim}). *There are constants $c_{\text{step}}, C > 0$, depending at most on c_{init} , such that the following holds for every $\kappa \geq 2, \ell, \Delta > 0, \sigma \geq 0$, and integer $T \geq 1$. On any instance in $\mathcal{C}(\ell, \kappa, \Delta, \sigma)$, run SGDA_{Sim} with*

$$\rho = 16\kappa^2, \quad \eta_x := c_{\text{step}} \min \left\{ \frac{1}{\kappa^2 \ell}, \sqrt{\frac{\Delta}{\kappa^4 \ell \sigma^2 T}} \right\}, \quad \eta_t \equiv \eta_y := 16\kappa^2 \eta_x, \quad (\text{A.9})$$

where the second term is $+\infty$ when $\sigma = 0$. Then

$$\frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E} \|\nabla \Phi(x_t)\|^2 \leq C \left(\frac{\kappa^2 \ell \Delta}{T} + \sqrt{\frac{\kappa^4 \ell \Delta \sigma^2}{T}} \right). \quad (\text{A.10})$$

Consequently, whenever $\varepsilon^2 \leq c\ell\Delta$ for a sufficiently small universal $c > 0$,

$$N_{\varepsilon, \text{Sim}}(16\kappa^2), N_{\varepsilon, \text{Sim}}^* \leq C \left(\frac{\kappa^2 \ell \Delta}{\varepsilon^2} + \frac{\kappa^4 \ell \Delta \sigma^2}{\varepsilon^4} \right). \quad (\text{A.11})$$

Proof of Theorem A.1. We track the outer objective together with the inner suboptimality through

$$\mathcal{V}_t := \Phi(x_t) + \frac{1}{8} [\Phi(x_t) - f(x_t, y_t)]. \quad (\text{A.12})$$

This potential is bounded below by Φ^* . Its decrease combines 9/8 times the outer-objective decrease with 1/8 times the game-value increase. The faster dual step makes the latter absorb the discrepancy between the primal and outer gradients.

Let $\mathcal{F}_t$ be the history before the two oracle calls at iteration t and $\mathbb{E}_t[\cdot] := \mathbb{E}[\cdot \mid \mathcal{F}_t]$. The constant primal and dual steps are η_x and $\eta_y = 16\kappa^2 \eta_x$ from (A.9). Choose the universal constant $c_{\text{step}} \leq 1/128$. Then

$$\ell_{\Phi} \eta_x \leq \frac{1}{4}, \quad \ell_{\eta_x} \leq \frac{1}{8}, \quad \ell_{\eta_y} \leq \frac{1}{8}. \quad (\text{A.13})$$

By ℓ_Φ -smoothness of Φ and the unbiased primal oracle,

$$\begin{aligned}\mathbb{E}_t[\Phi(x_t) - \Phi(x_{t+1})] &\geq \eta_x \langle \nabla \Phi(x_t), \nabla_x f(x_t, y_t) \rangle - \frac{\ell_\Phi \eta_x^2}{2} (\|\nabla_x f(x_t, y_t)\|^2 + \sigma^2) \\ &= \frac{\eta_x}{2} \|\nabla \Phi(x_t)\|^2 + \frac{\eta_x}{2} (1 - \ell_\Phi \eta_x) \|\nabla_x f(x_t, y_t)\|^2 \\ &\quad - \frac{\eta_x}{2} \|\nabla_x f(x_t, y_t) - \nabla \Phi(x_t)\|^2 - \frac{\ell_\Phi \eta_x^2}{2} \sigma^2.\end{aligned}\tag{A.14}$$

Assumption 3.1 implies that the full gradient is 2ℓ -Lipschitz in the Euclidean product norm. Since the primal and dual gradients in SGDA_{Sim} are evaluated at (x_t, y_t) , this gives

$$\begin{aligned}\mathbb{E}_t[f(x_{t+1}, y_{t+1}) - f(x_t, y_t)] &\geq -(\eta_x + \ell \eta_x^2) \|\nabla_x f(x_t, y_t)\|^2 + (\eta_y - \ell \eta_y^2) \|\nabla_y f(x_t, y_t)\|^2 \\ &\quad - \ell(\eta_x^2 + \eta_y^2) \sigma^2.\end{aligned}\tag{A.15}$$

Combining $\frac{9}{8}$ times (A.14) with $\frac{1}{8}$ times (A.15), using the gradient-bias bound (B.6), and then applying (A.13), yields

$$\mathbb{E}_t[\mathcal{V}_t - \mathcal{V}_{t+1}] \geq \frac{9\eta_x}{16} \|\nabla \Phi(x_t)\|^2 - 33\kappa^4 \ell \eta_x^2 \sigma^2.\tag{A.16}$$

Indeed, (A.13) makes the remaining squared-gradient coefficients nonnegative:

$$\begin{aligned}\frac{9\eta_x}{16} (1 - \ell_\Phi \eta_x) - \frac{\eta_x}{8} - \frac{\ell \eta_x^2}{8} &\geq 0, \\ \frac{\eta_y}{8} (1 - \ell \eta_y) - \frac{9\eta_x \kappa^2}{16} &\geq 0.\end{aligned}$$

The dual-noise contribution is $\ell \eta_y^2 / 8 = 32\kappa^4 \ell \eta_x^2$, which determines the fourth power of κ in the variance bound. The full noise coefficient satisfies

$$\frac{9\ell \eta_x^2}{16} + \frac{\ell}{8} (\eta_x^2 + \eta_y^2) \leq 33\kappa^4 \ell \eta_x^2 \quad (\kappa \geq 2).$$

We sum (A.16) over the $T - 1$ updates, retaining the terminal potential to control the last reported iterate. Smoothness and lower boundedness of Φ give $\|\nabla \Phi(x)\|^2 \leq 2\ell_\Phi [\Phi(x) - \Phi^*]$. Thus (A.13) implies

$$\frac{9\eta_x}{16} \mathbb{E} \|\nabla \Phi(x_{T-1})\|^2 \leq \frac{9}{32} \mathbb{E} [\mathcal{V}_{T-1} - \Phi^*].$$

Adding this bound to the sum and using both initial-gap bounds gives

$$\begin{aligned}\frac{9\eta_x}{16} \sum_{t=0}^{T-1} \mathbb{E} \|\nabla \Phi(x_t)\|^2 &\leq \mathcal{V}_0 - \Phi^* - \frac{23}{32} \mathbb{E} [\mathcal{V}_{T-1} - \Phi^*] \\ &\quad + 33\kappa^4 \ell \eta_x^2 \sigma^2 (T - 1) \\ &\leq \left(1 + \frac{c_{\text{init}}}{8}\right) \Delta + 33\kappa^4 \ell \eta_x^2 \sigma^2 T.\end{aligned}\tag{A.17}$$

With the empty-sum convention, the same argument applies when $T = 1$. Dividing by $\eta_x T$ and absorbing the numerical constants gives

$$\frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E} \|\nabla \Phi(x_t)\|^2 \leq C \left[\frac{\Delta}{\eta_x T} + \kappa^4 \ell \eta_x \sigma^2 \right].\tag{A.18}$$

Substitute (A.9). If the step-size bound $1/(\kappa^2 \ell)$ is active, then $\sigma^2 T \leq \ell \Delta$ and both terms in (A.18) are bounded by $C \kappa^2 \ell \Delta / T$. Otherwise the two terms are both bounded by $C \sqrt{\kappa^4 \ell \Delta \sigma^2 / T}$. This proves (A.10).

Finally, choosing

$$T \geq C \left(\frac{\kappa^2 \ell \Delta}{\varepsilon^2} + \frac{\kappa^4 \ell \Delta \sigma^2}{\varepsilon^4} \right)$$

makes the right-hand side of (A.10) at most ε^2 . The constant schedule is nonincreasing, so the minimax definitions in (3.7) give (A.11). $\square$

B PRELIMINARY LEMMAS

We first record the completion-of-squares construction used to preserve the outer function and inner PŁ. We then prove the consequences of inner PŁ used in the matching upper bound.

B.1 NC-PŁ GAMES WITH A PREDETERMINED OUTER FUNCTION

We use an onto inner map to determine the outer maximum exactly. A lower bound on its transposed derivative then converts the residual norm into an inner-gradient bound. This lemma proves inner PŁ, not global smoothness; the latter is checked for each constructed map.

Lemma B.1 (Outer function and inner PŁ for a completion-of-squares construction). *Let $b : \mathbb{R}^{d_x} \rightarrow \mathbb{R}^m$ and $T : \mathbb{R}^{d_y} \rightarrow \mathbb{R}^m$ be continuously differentiable, with T onto (surjective). Define*

$$g(x, y) := \langle b(x), T(y) \rangle - \frac{1}{2} \|T(y)\|^2. \quad (\text{B.1})$$

Then

$$\max_y g(x, y) = \frac{1}{2} \|b(x)\|^2, \quad \max_y g(x, y) - g(x, y) = \frac{1}{2} \|b(x) - T(y)\|^2. \quad (\text{B.2})$$

If, for some $\nu > 0$,

$$\|DT(y)^\top z\| \geq \nu \|z\| \quad \text{for all } y, z, \quad (\text{B.3})$$

then

$$\frac{1}{2} \|\nabla_y g(x, y)\|^2 \geq \nu^2 \left[\max_v g(x, v) - g(x, y) \right]. \quad (\text{B.4})$$

Proof. Since T is onto, maximizing (B.1) over y is equivalent to maximizing $\langle b(x), z \rangle - \|z\|^2/2$ over $z \in \mathbb{R}^m$. Completing the square gives (B.2). Moreover,

$$\nabla_y g(x, y) = DT(y)^\top [b(x) - T(y)],$$

so (B.3) and (B.2) imply (B.4). $\square$

B.2 NC-PŁ GEOMETRY

The following are standard consequences of smoothness and inner PŁ; see [Nouiehed et al. \(2019, Lemma A.5\)](#), [Chen et al. \(2022, Lemmas A.7–A.8\)](#), and [Ha \(2026, Lemmas A.2–A.3\)](#). We prove them under [Assumptions 3.1](#) and [3.2](#), without concavity or uniqueness of the maximizer.

Lemma B.2 (Inner error bound, gradient bias, and outer smoothness). *Under [Assumptions 3.1](#) and [3.2](#), Φ is differentiable with $\nabla \Phi(x) = \nabla_x f(x, y^*)$ for every $y^* \in Y^*(x)$. For every (x, y) ,*

$$\text{dist}(y, Y^*(x)) \leq \frac{\kappa}{\ell} \|\nabla_y f(x, y)\|, \quad (\text{B.5})$$

$$\|\nabla_x f(x, y) - \nabla \Phi(x)\| \leq \kappa \|\nabla_y f(x, y)\|. \quad (\text{B.6})$$

Moreover, Φ has ℓ_Φ -Lipschitz gradient with

$$\ell_\Phi \leq \ell(1 + \kappa). \quad (\text{B.7})$$

Proof. We first bound the inner gap after changing x while keeping a maximizer fixed; the resulting quadratic remainder proves differentiability of Φ . An inner gradient-flow argument then bounds the distance to a maximizer, giving the gradient-bias and outer-smoothness estimates.

Fix $x, y^* \in Y^*(x)$, and $h \in \mathbb{R}^{d_x}$. Since the inner domain is unconstrained, $\nabla_y f(x, y^*) = 0$. Inner PŁ at $(x + h, y^*)$ and [Assumption 3.1](#) give

$$0 \leq \Phi(x + h) - f(x + h, y^*) \leq \frac{\|\nabla_y f(x + h, y^*)\|^2}{2\mu} \leq \frac{\ell\kappa}{2} \|h\|^2.$$

Combining this with the Taylor bound for $f(\cdot, y^*)$ yields

$$|\Phi(x + h) - \Phi(x) - \langle \nabla_x f(x, y^*), h \rangle| \leq \frac{\ell(1 + \kappa)}{2} \|h\|^2.$$

Thus Φ is differentiable and $\nabla\Phi(x) = \nabla_x f(x, y^*)$ for every maximizer, including when the maximizer is not unique. If $d_y = 0$, this identity and all the claimed bounds follow directly from [Assumption 3.1](#); henceforth assume $d_y \geq 1$.

Fix x and define the inner objective gap $g_x(y) := \Phi(x) - f(x, y) \geq 0$. It is ℓ -smooth, attains its minimum zero, and satisfies $\|\nabla g_x(y)\|^2 \geq 2(\ell/\kappa)g_x(y)$. Since ∇g_x is globally Lipschitz, the gradient flow $\dot{y}(s) = -\nabla g_x(y(s))$, initialized at $y(0) = y$, exists for all $s \geq 0$. If it reaches $g_x = 0$ in finite time, the desired distance bound follows immediately by integrating up to that time. Otherwise $g_x(y(s)) > 0$ for every finite s , and

$$-\frac{d}{ds}\sqrt{g_x(y(s))} = \frac{\|\nabla g_x(y(s))\|^2}{2\sqrt{g_x(y(s))}} \geq \sqrt{\frac{\ell}{2\kappa}} \|\nabla g_x(y(s))\|.$$

Hence the total distance traveled by the gradient-flow solution is finite:

$$\int_0^\infty \|\dot{y}(s)\| ds \leq \sqrt{\frac{2\kappa g_x(y)}{\ell}}.$$

The finite path length makes $y(s)$ a Cauchy curve in $\mathbb{R}^{d_y}$, so it converges to a limit y^* . The limit must satisfy $g_x(y^*) = 0$: otherwise PŁ would keep $\|\nabla g_x\|$ bounded away from zero and the integral of its speed would be infinite. Thus $y^* \in Y^*(x)$, and therefore

$$\text{dist}(y, Y^*(x)) \leq \sqrt{\frac{2\kappa g_x(y)}{\ell}} \leq \frac{\kappa}{\ell} \|\nabla g_x(y)\|,$$

where the second inequality is the PŁ inequality. This proves [\(B.5\)](#).

Choose the maximizer supplied by [\(B.5\)](#). The gradient identity just proved and [Assumption 3.1](#) give

$$\|\nabla_x f(x, y) - \nabla\Phi(x)\| \leq \ell \|y - y^*\| \leq \kappa \|\nabla_y f(x, y)\|,$$

which is [\(B.6\)](#).

Finally, fix x, x' and $y^* \in Y^*(x)$. Since $\nabla_y f(x, y^*) = 0$, [Assumption 3.1](#) gives $\|\nabla_y f(x', y^*)\| \leq \ell \|x' - x\|$. Applying [\(B.5\)](#) to the inner problem at x' yields a maximizer $\bar{y} \in Y^*(x')$ such that $\|\bar{y} - y^*\| \leq \kappa \|x' - x\|$. Thus

$$\begin{aligned} \|\nabla\Phi(x') - \nabla\Phi(x)\| &= \|\nabla_x f(x', \bar{y}) - \nabla_x f(x, y^*)\| \\ &\leq \ell(\|x' - x\| + \|\bar{y} - y^*\|) \leq \ell(1 + \kappa) \|x' - x\|. \end{aligned}$$

This proves [\(B.7\)](#). □

C PROOFS FOR THE HIGH-RATIO LOWER BOUND

We prove the construction lemmas used in [Section 5](#). The lower bound for f_{unstable} accounts for initially large dual steps. On the other hand, the corridor function g_{corridor} controls the remaining primal movement, and the noisy function g_σ supplies the stochastic term in the lower bound. Finally, we combine the corridor function with the noisy function to obtain a single NC-PŁ game instance and prove its lower bound.

C.1 PROOF OF THE LOWER BOUND FOR f_{unstable}

Lemma 5.2 (A lower bound for f_{unstable}). *There is a universal $c > 0$ such that, for $\ell, \Delta > 0$, $\kappa \geq 2$, $\sigma \geq 0$, $\rho \geq 1$, integers $T \geq 1$ and $0 \leq m < T$, and predetermined nonnegative non-increasing step sizes with $\eta_t \geq 8/\ell$ for $t < m$, each of SGDA_{Sim} and SGDA_{Alt} admits an initialization (x_0, y_0) making f_{unstable} with $\mu = \ell/\kappa$ an exact-gradient instance in $\mathcal{C}(\ell, \kappa, \Delta, \sigma)$ with*

$$\frac{1}{T} \sum_{t=0}^{T-1} |\Phi'_{\text{unstable}}(x_t)|^2 \geq c \frac{m+1}{T} \ell \Delta. \quad (5.3)$$

Proof. We use the same initialization for both update rules. We show that two consecutive updates with dual step sizes at least $8/\ell$ take any state (x, y) satisfying $0 \leq x \leq y$ to a state (x'', y'') satisfying $0 \leq x \leq x'' \leq y''$. For $m \geq 1$, the first update gives $x_1 = x_0$ and $y_1 \geq 2x_0$. Repeating the two-update argument then gives $x_t \geq x_0$ for $t = 1, 3, 5, \dots$ with $t \leq m$.

Set $\mu = \ell/\kappa$, $x_0 = 4\sqrt{\Delta/(\ell + \mu)}$, and $y_0 = 0$. The Hessian is $(\ell + \mu) \begin{pmatrix} 0 & 1/4 \\ 1/4 & -1/2 \end{pmatrix}$, and the inner maximum is attained at $y = x/2$. Hence

$$\begin{aligned} \|\nabla^2 f_{\text{unstable}}\|_{\text{op}} &= \frac{1 + \sqrt{2}}{4}(\ell + \mu) \leq \frac{3(1 + \sqrt{2})}{8}\ell < \ell, \\ \Phi_{\text{unstable}}(x) - f_{\text{unstable}}(x, y) &= \frac{\ell + \mu}{4}(y - x/2)^2, \\ \frac{1}{2}|\nabla_y f_{\text{unstable}}(x, y)|^2 &= \frac{\ell + \mu}{2}(\Phi_{\text{unstable}}(x) - f_{\text{unstable}}(x, y)). \end{aligned}$$

Since $(\ell + \mu)/2 \geq \mu$, the game satisfies inner μ -PL. Its outer objective is differentiable and lower bounded. Both initial gaps equal Δ , so the initialization is admissible because $c_{\text{init}} \geq 1$.

Write $s_t = (\ell + \mu)\eta_t/2 \geq 4$ for $t < m$. A step of normalized size a sends (x, y) to

$$x' = x - \frac{a}{2\rho}y, \quad y' = \begin{cases} (1-a)y + \frac{a}{2}x, & \text{Sim,} \\ (1-a)y + \frac{a}{2}x', & \text{Alt.} \end{cases} \quad (\text{C.1})$$

If $m \geq 1$, the first step gives $x_1 = x_0$ and $y_1 = s_0 x_0/2 \geq 2x_0$. Consider two subsequent steps $a, b \geq 4$ starting from $0 \leq x \leq y$, with states (x', y') and (x'', y'') . Since $x' \leq x \leq y$, both update rules satisfy

$$\begin{aligned} y' &\leq (1 - a/2)y \leq 0, \\ x'' - x &= -\frac{ay + by'}{2\rho} \geq \frac{ab/2 - a - b}{2\rho}y \geq 0. \end{aligned}$$

The last coefficient is nonnegative because $(a-2)(b-2) \geq 4$. For SGDA_{Alt} , $y'' = (b/2)x'' + (1-b)y' \geq 2x''$, since $x'' \geq 0$ and $y' \leq 0$. For SGDA_{Sim} ,

$$\begin{aligned} x' - y' &= (1 - a/2)x + (a - 1 - a/(2\rho))y \geq \frac{a}{2}(1 - \rho^{-1})x \geq 0, \\ y'' - x'' &= (b/2 - 1)x' + (1 - b + b/(2\rho))y' \geq -\frac{b}{2}(1 - \rho^{-1})y' \geq 0. \end{aligned}$$

Here the first line uses $y \geq x$ and $a - 1 - a/(2\rho) \geq 0$; the second uses $x' \geq y'$ and $b/2 - 1 \geq 0$. Thus either pair of steps gives $0 \leq x \leq x'' \leq y''$.

Induction from (x_1, y_1) yields $x_t \geq x_0$ at every odd $t \leq m$. Including $t = 0$, we obtain

$$\frac{1}{T} \sum_{t=0}^{T-1} |\Phi'_{\text{unstable}}(x_t)|^2 \geq \frac{(\ell + \mu)^2}{64T} \sum_{t=0}^m x_t^2 \geq \frac{(\ell + \mu)^2 x_0^2}{64T} \left(1 + \left\lceil \frac{m}{2} \right\rceil\right) \geq \frac{m+1}{8T} \ell \Delta. \quad (\text{C.2})$$

The same bound holds directly for $m = 0$. Exact gradients are admissible for every $\sigma \geq 0$, completing the proof. $\square$

C.2 CONSTRUCTION OF g_{corridor}

Lemma 5.3 (A lower bound for g_{corridor}). *For $\ell, \Delta > 0$, $\rho \geq 1$, integers $T \geq 1$, $0 \leq m < T$, and nonnegative steps with $S = \sum_{t=m}^{T-2} \eta_t$, there is an ℓ -smooth function $g_{\text{corridor}} : \mathbb{R} \rightarrow \mathbb{R}$ with initial gap $g_{\text{corridor}}(0) - \inf_u g_{\text{corridor}}(u) = \Delta$. Gradient descent from $u_0 = 0$ with steps η_t/ρ satisfies $g'_{\text{corridor}}(u_t) < 0$ for all $t < T$ and $g'_{\text{corridor}}(u_t) = -G$ for $m \leq t < T$, where $G > 0$ and $G^2 \asymp \min\{\ell\Delta, \rho\Delta/S\}$. Consequentially,*

$$\frac{1}{T} \sum_{t < T} |g'_{\text{corridor}}(u_t)|^2 \gtrsim \frac{T-m}{T} \min \left\{ \ell\Delta, \frac{\rho\Delta}{S} \right\}. \quad (5.5)$$

Proof. We prescribe the derivative before choosing its magnitude. An initial linear segment makes the first m updates reach a constant-derivative interval. Its length is the distance traveled by the remaining steps; a final linear segment joins it to a minimum. Integrating the derivative then calibrates the exact initial gap.

Fix $m \in \{0, \dots, T-1\}$ and let $S = \sum_{t=m}^{T-2} \eta_t$. When $S = 0$, the minimum in the statement is $\ell\Delta$. Define

$$P_0 := 1, \quad P_t := \prod_{i=0}^{t-1} \left(1 + \frac{\ell\eta_i}{\rho}\right) \quad (1 \leq t \leq m), \quad (\text{C.3})$$

$$G^2 := \frac{\Delta}{S/\rho + \ell^{-1}(1 - 1/(2P_m^2))}, \quad G_0 := \frac{G}{P_m}, \quad b := \frac{G - G_0}{\ell} + \frac{GS}{\rho}. \quad (\text{C.4})$$

For $G > 0$, write $[z]_{[0,G]} := \min\{G, \max\{0, z\}\}$. Add a constant to g_{corridor} so that $\inf_u g_{\text{corridor}}(u) = 0$, and define

$$g'_{\text{corridor}}(u) := -[G_0 + \ell u]_{[0,G]} + [\ell(u - b)]_{[0,G]}. \quad (\text{C.5})$$

The two intervals on which the clipping terms vary are disjoint, so g'_{corridor} is continuous and ℓ -Lipschitz.

Starting from $u_0 = 0$, induction gives

$$u_t = \frac{P_t - 1}{\ell} G_0, \quad g'_{\text{corridor}}(u_t) = -P_t G_0, \quad 0 \leq t \leq m, \quad (\text{C.6})$$

$$u_t = \frac{G - G_0}{\ell} + G \sum_{j=m}^{t-1} \frac{\eta_j}{\rho}, \quad g'_{\text{corridor}}(u_t) = -G, \quad m \leq t \leq T-1. \quad (\text{C.7})$$

Direct integration gives

$$g_{\text{corridor}}(0) = \frac{G^2 - G_0^2}{2\ell} + \frac{G^2 S}{\rho} + \frac{G^2}{2\ell} = G^2 \left(\frac{S}{\rho} + \frac{1}{\ell} - \frac{1}{2\ell P_m^2} \right) = \Delta, \quad (\text{C.8})$$

and

$$G^2 \geq \frac{\Delta}{S/\rho + 1/\ell} \geq \frac{1}{2} \min \left\{ \ell\Delta, \frac{\rho\Delta}{S} \right\}. \quad (\text{C.9})$$

Since $P_m \geq 1$, the denominator in (C.4) is at least $S/\rho + 1/(2\ell)$, so also $G^2 \leq 2 \min\{\ell\Delta, \rho\Delta/S\}$. This proves the two-sided comparison in the statement.

The game is the single-player objective $f(u) = g_{\text{corridor}}(u)$, equivalently an instance with no inner variable. Hence the inner-PŁ condition is vacuous. Equations (C.7) and (C.9), summed over the last $T - m$ iterates, prove (5.5). Since $G_0, G > 0$ and $P_t > 0$, the derivative is negative at every iterate, so each positive update strictly increases the scalar iterate. The initial gap equals Δ by (C.8). Exact gradients are admissible for every σ . $\square$

C.3 CONSTRUCTION OF g_σ

We construct g_σ by giving each inner-gradient error its own component. The shifted quadratic game g_δ in Lemma 5.4 shows why this is useful: a shift in its inner gradients moves the outer variable away from its minimizer, and that displacement persists over a range of cumulative steps. We then build a smooth inner-PŁ component $g_{\sigma,j}$ in which either sign of the error at update j activates the same shift. The displacement estimate lets us count how many subsequent iterates have a large outer gradient. Summing the resulting squared-gradient bounds over components with enough remaining cumulative steps gives the lower bound on the average squared outer-gradient norm in Lemma C.2. The fixed oracle in (5.11) generates all of these errors.

C.3.1 PERSISTENT ERROR UNDER A DETERMINISTIC SHIFT

Lemma 5.4 (Persistent error under a deterministic shift). *There are universal constants $c, C, C_0 > 0$ such that, for $\ell, \delta > 0$, $\kappa \geq 1$, $\rho \geq C_0 \kappa^2$, and any finite step-size sequence with $0 \leq \ell \eta_t \leq c\kappa$, both SGDA_{Sim} and SGDA_{Alt} with exact gradients on the linear quadratic game g_δ given by (5.6), initialized at $(0, 0, 0)$, satisfy $|z_t| \leq C\kappa\delta/\rho$ and $|a_t| + |b_t| \leq C\delta$ whenever $\sum_{i < t} \eta_i \leq c\rho/(\ell\kappa)$, and*

$$z_t \geq c \frac{\kappa\delta}{\rho} \quad \text{if} \quad C \frac{\kappa}{\ell} \leq \sum_{i < t} \eta_i \leq c \frac{\rho}{\ell\kappa}. \quad (5.7)$$

Proof. Recall that $g_\delta(z; a, b) = \ell z a - \frac{\ell}{2\kappa} [a^2 + (a + b + \delta)^2]$ and $(z_0, a_0, b_0) = (0, 0, 0)$. We bound the inner coordinates in terms of the largest outer iterate, then use an exact identity to close the outer bound. Once the initial inner transient has decayed, the same identity gives a positive lower bound on the outer displacement. Normalize the coordinates and dual steps by setting

$$X_n = \frac{\kappa z_n}{\delta}, \quad Y_n = \frac{a_n}{\delta}, \quad Z_n = 1 + \frac{b_n}{\delta}, \quad h_n = \frac{\ell \eta_n}{\kappa}, \quad \alpha = \frac{\kappa^2}{\rho}.$$

The SGDA updates for g_δ become

$$\begin{aligned} X_{n+1} &= X_n - \alpha h_n Y_n, \\ Y_{n+1} &= Y_n + h_n (X'_n - 2Y_n - Z_n), \quad (X_0, Y_0, Z_0) = (0, 0, 1), \\ Z_{n+1} &= Z_n - h_n (Y_n + Z_n), \end{aligned} \quad (C.10)$$

where $X'_n = X_n$ for SGDA_{Sim} and $X'_n = X_{n+1}$ for SGDA_{Alt} . Write $\tau_n = \sum_{i < n} h_i$ for the cumulative normalized step. We choose universal positive constants $\alpha_*, \bar{h}, \tau_-, \tau_+$ below. For $\alpha \leq \alpha_*$ and $h_n \leq \bar{h}$, we will bound $|X_n|$ by a universal multiple of α when $\tau_n \leq \tau_+/\alpha$, and prove $X_n \geq \alpha/4$ when $\tau_- \leq \tau_n \leq \tau_+/\alpha$.

Subtracting the updates for Y_n and Z_n in (C.10) yields $Y_{n+1} - Z_{n+1} - (Y_n - Z_n) = h_n (X'_n - Y_n)$. Summing this equality over $0 \leq i < n$, using $Y_0 - Z_0 = -1$, and then summing the update for X_n gives

$$X_n = \alpha \left[1 + Y_n - Z_n - \sum_{i < n} h_i X'_i \right]. \quad (C.11)$$

The constant 1 records the initial offset $Y_0 - Z_0 = -1$. To obtain a positive outer displacement, we must make the inner-coordinate difference small while keeping the accumulated outer term small.

Let $q_n = (Y_n, Z_n)^\top \in \mathbb{R}^2$ collect the normalized inner coordinates in (C.10). Its update is $q_{n+1} = (I + h_n B)q_n + h_n X'_n (1, 0)^\top$, where $B = \begin{pmatrix} -2 & -1 \\ -1 & -1 \end{pmatrix}$. The eigenvalues of $-B$ are $(3 \pm \sqrt{5})/2$. Write $\lambda_0 = (3 - \sqrt{5})/2$ and choose $\bar{h} \leq 2/(3 + \sqrt{5})$. Then $I + hB$ is positive semidefinite and $\|I + hB\|_{\text{op}} = 1 - \lambda_0 h$ for $0 \leq h \leq \bar{h}$.

Fix an endpoint N , and let $M_N = \max_{0 \leq i \leq N} |X_i|$ be the largest magnitude of the normalized outer coordinate through iterate N . Since $|X'_n| \leq M_N$ for $n < N$, the formula for q_{n+1} and the norm bound on $I + h_n B$ give

$$\|q_{n+1}\| \leq (1 - \lambda_0 h_n) \|q_n\| + h_n M_N, \quad (C.12)$$

$$\|q_n\| \leq \prod_{i < n} (1 - \lambda_0 h_i) + \frac{M_N}{\lambda_0} \left[1 - \prod_{i < n} (1 - \lambda_0 h_i) \right] \leq e^{-\lambda_0 \tau_n} + \frac{M_N}{\lambda_0}. \quad (C.13)$$

Iterating the first inequality from $q_0 = (0, 1)^\top$ gives the product bound in the second line; $1 - a \leq e^{-a}$ gives its exponential bound.

For every $n \leq N$, $|Y_n - Z_n| \leq \sqrt{2} \|q_n\|$ and $\sum_{i < n} h_i |X'_i| \leq \tau_n M_N$. Substituting these bounds and (C.13) into (C.11), then taking the maximum over $0 \leq n \leq N$, gives

$$M_N \leq \alpha(1 + \sqrt{2}) + \alpha \left(\frac{\sqrt{2}}{\lambda_0} + \tau_N \right) M_N. \quad (C.14)$$

Put $C_X = 2(1 + \sqrt{2})$. If $\alpha_* \sqrt{2}/\lambda_0 \leq 1/4$ and $\tau_+ \leq 1/4$, then $\tau_N \leq \tau_+/\alpha$ makes the coefficient multiplying M_N in (C.14) at most $1/2$. Thus $M_N \leq C_X \alpha$, and (C.13) gives $\|q_n\| \leq e^{-\lambda_0 \tau_n} + C_X \alpha/\lambda_0$ for $0 \leq n \leq N$.

For a lower bound on X_N , use $Y_N - Z_N \geq -\sqrt{2}\|q_N\|$ in (C.11), together with (C.13) and $M_N \leq C_X \alpha$, to obtain

$$\frac{X_N}{\alpha} \geq 1 - \sqrt{2}e^{-\lambda_0 \tau_N} - \frac{\sqrt{2}}{\lambda_0} M_N - \tau_N M_N \geq 1 - \sqrt{2}e^{-\lambda_0 \tau_N} - \frac{\sqrt{2}C_X}{\lambda_0} \alpha - C_X \alpha \tau_N. \quad (\text{C.15})$$

Choose $\tau_- = \lambda_0^{-1} \log(4\sqrt{2})$, take $\tau_+ \leq \min\{1/4, 1/(4C_X)\}$, and reduce α_* so that

$$\alpha_* \leq \min \left\{ \frac{\lambda_0}{4\sqrt{2}C_X}, 1, \frac{\tau_+}{2(\tau_- + 2\bar{h})} \right\}.$$

For $\tau_- \leq \tau_N \leq \tau_+/\alpha$, the three subtracted terms $\sqrt{2}e^{-\lambda_0 \tau_N}$, $(\sqrt{2}C_X/\lambda_0)\alpha$, and $C_X \alpha \tau_N$ in (C.15) are each at most $1/4$, so $X_N \geq \alpha/4$. The choices also satisfy the conditions used in (C.14) and give $\tau_- < \tau_+/\alpha$. Since N was arbitrary, $|X_n| \leq C_X \alpha$ holds when $\tau_n \leq \tau_+/\alpha$, and $X_n \geq \alpha/4$ when $\tau_- \leq \tau_n \leq \tau_+/\alpha$. The argument permits $h_n = 0$ and both definitions of X'_n in (C.10).

To express the bounds in the original coordinates (z_n, a_n, b_n) , choose $c > 0$ no larger than $\bar{h}$, τ_+ , and $1/4$, and take $C \geq \tau_-$ large enough to bound $|X_n|/\alpha$ and $|Y_n| + |Z_n - 1|$. Set $C_0 = \alpha_*^{-1}$. These constants are fixed before choosing the cutoff in Appendix C.3.2 and the component scales in Appendix C.3.3.

If $\rho \geq C_0 \kappa^2$ and $\ell \eta_n \leq c\kappa$, then $\alpha \leq \alpha_*$ and $h_n \leq \bar{h}$. Substituting $z_n = \delta X_n/\kappa$, $a_n = \delta Y_n$, and $b_n = \delta(Z_n - 1)$ into the bounds on M_N and $\|q_n\|$ gives

$$|z_n| \leq C \frac{\kappa \delta}{\rho}, \quad |a_n| + |b_n| \leq C \delta \quad \text{when} \quad \sum_{i < n} \eta_i \leq c \frac{\rho}{\ell \kappa}.$$

If $C\kappa/\ell \leq \sum_{i < n} \eta_i \leq c\rho/(\ell\kappa)$, then $\tau_- \leq \tau_n \leq \tau_+/\alpha$, and (C.15) gives $z_n \geq c\kappa\delta/\rho$. No monotonicity of the steps was used. $\square$

C.3.2 A SMOOTH NC-PŁ GAME WITH THE SAME MECHANISM

The shifted quadratic starts with a nonzero inner gap at the origin. We introduce a third inner coordinate s so that a gradient error can activate the shift from an inner maximizer. We also cut off the shift when $|a + b|$ is large to bound the Hessian globally while preserving the quadratic updates along the trajectory segment in Lemma 5.4. Fix R_0 larger than the inner-coordinate bound in that lemma, and choose a smooth cutoff $\chi : \mathbb{R} \rightarrow [0, 1]$ equal to one on $[-R_0, R_0]$, zero outside $[-2R_0, 2R_0]$, with $|\chi'| \leq 1/4$ and bounded first two derivatives. The constant R_0 is fixed before all smoothness and PŁ constants below.

For a provisional condition parameter $K \geq 1$, define the residual $r_K : \mathbb{R}^3 \rightarrow \mathbb{R}$ and game $g_K : \mathbb{R} \times \mathbb{R}^3 \rightarrow \mathbb{R}$ by

$$r_K(a, b, s) = a + b + \sqrt{K} \chi\left(\frac{a+b}{\sqrt{K}}\right) \sin^2\left(\frac{\pi s}{2}\right), \quad g_K(u; a, b, s) = ua - \frac{a^2 + r_K(a, b, s)^2}{2K}. \quad (\text{C.16})$$

The profile vanishes at $s = 0$ and has zero derivative there. At $s = \pm 1$, it supplies the same shift and again has zero derivative. Thus either sign of a unit perturbation activates the shift, and the activation coordinate then stays fixed. Without the cutoff, the second derivative of g_K in s would grow linearly with $a + b$.

Lemma C.1 (Smoothness and inner PŁ of the single-perturbation game). *There are universal $c, C > 0$ such that, for every $K \geq 1$, g_K has a C -Lipschitz gradient,*

$$\max_{a,b,s} g_K(u; a, b, s) = \frac{K}{2} u^2, \quad \frac{1}{2} \|\nabla_{a,b,s} g_K\|^2 \geq \frac{c}{K} \left[\frac{K}{2} u^2 - g_K(u; a, b, s) \right]. \quad (\text{C.17})$$

The origin is stationary.

Proof. Completing the square gives

$$g_K(u; a, b, s) = \frac{K}{2}u^2 - \frac{(a - Ku)^2 + r_K(a, b, s)^2}{2K}. \quad (\text{C.18})$$

Both residuals vanish at $(a, b, s) = (Ku, -Ku, 0)$, proving the outer-objective formula and attainment. For the PŁ inequality, write $r = r_K(a, b, s)$, $e = a - Ku$, and

$$p = \partial_a r_K = \partial_b r_K = 1 + \chi' \left(\frac{a+b}{\sqrt{K}} \right) \sin^2 \left(\frac{\pi s}{2} \right) \in [3/4, 5/4].$$

The restriction on χ' keeps p bounded away from zero, so the gradients in a and b control both residuals e and r without using the gradient in s . Indeed, $\partial_a g_K = -(e+pr)/K$ and $\partial_b g_K = -pr/K$. Since $e = (e+pr) - pr$ and $r^2 \leq 16(pr)^2/9$,

$$e^2 + r^2 \leq 2(e+pr)^2 + \left(2 + \frac{16}{9}\right)(pr)^2 \leq 4[(e+pr)^2 + (pr)^2], \quad (\text{C.19})$$

$$\frac{1}{2} \|\nabla_{a,b,s} g_K\|^2 \geq \frac{e^2 + r^2}{8K^2} = \frac{1}{4K} \left[\frac{K}{2}u^2 - g_K(u; a, b, s) \right]. \quad (\text{C.20})$$

This proves the inner-PŁ inequality.

For smoothness, $|\partial_a r_K|, |\partial_b r_K| \leq C$ and $|\partial_s r_K| \leq C\sqrt{K}$. Wherever a second derivative is nonzero, the cutoff implies $|a+b| \leq 2R_0\sqrt{K}$ and $|r_K| \leq C\sqrt{K}$. Direct differentiation gives second-derivative bounds of order $K^{-1/2}$ in (a, b) , order one for mixed derivatives with s , and order $K^{1/2}$ for $\partial_{ss} r_K$. Hence, for $y = (a, b, s)$,

$$\|\nabla_y^2(r_K^2/(2K))\| \leq \frac{\|\nabla_y r_K\|^2 + |r_K| \|\nabla_y^2 r_K\|}{K} \leq C. \quad (\text{C.21})$$

Outside that region the residual is affine, so the same bound holds globally. The remaining terms $ua - a^2/(2K)$ also have bounded Hessian for $K \geq 1$. All first derivatives vanish at the origin. $\square$

At $s = \pm 1$ and $|a+b| \leq R_0\sqrt{K}$, the residual satisfies

$$r_K(a, b, s) = a + b + \sqrt{K}, \quad \partial_s r_K(a, b, s) = 0. \quad (\text{C.22})$$

Thus s stays fixed and the other coordinates follow the shifted quadratic of [Lemma 5.4](#) while the cutoff equals one. At $u = a = b = 0$, a unit perturbation of s creates inner gap $1/2$, since $g_K(0; 0, 0, \pm 1) = -1/2$. A unit perturbation of b , with $u = a = s = 0$, instead creates inner gap $1/(2K)$. The third coordinate therefore increases the gap by a factor K for the same perturbation magnitude. On the region where the cutoff equals one, the quadratic curvature in (a, b) remains of order $1/K$.

C.3.3 CHANGING VARIABLES AND THE CONTRIBUTION OF A SINGLE PERTURBATION

Write $c_r, C_r, C_{\text{resp}}$ for c, C, C_0 in [Lemma 5.4](#), and enlarge C_{resp} so that $c_r C_{\text{resp}} \geq 2(C_r + 2c_r)$. Let c_0, C_0 be the constants in [Lemma C.1](#). Choose universal $\gamma, \theta > 0$, fixed after the low-ratio construction, such that

$$\theta C_0 \leq 1, \quad \frac{\theta c_0}{\gamma} \geq 1. \quad (\text{C.23})$$

For condition number κ , set

$$K := \gamma\kappa, \quad L := \theta\ell. \quad (\text{C.24})$$

In [\(5.8\)](#), set $g_{\sigma,j} \equiv 0$ whenever $\eta_j \sigma = 0$; these components have zero gradients and gaps. Increase the universal lower bound on κ so that $K \geq 1$. For every block $j \in \{0, \dots, T-2\}$ with $\eta_j \sigma > 0$, write $y[j] = (a[j], b[j], s[j])$ and define

$$g_{\sigma,j}(v[j]; a[j], b[j], s[j]) := L(\eta_j \sigma)^2 g_K \left(\frac{v[j]}{\eta_j \sigma}; \frac{a[j]}{\eta_j \sigma}, \frac{b[j]}{\eta_j \sigma}, \frac{s[j]}{\eta_j \sigma} \right). \quad (\text{C.25})$$

The squared amplitude compensates exactly for the spatial rescaling. Writing $w \in \mathbb{R}^4$ for the component coordinates, we have

$$\nabla_w^2 \left[(\eta_j \sigma)^2 g_K \left(\frac{w}{\eta_j \sigma} \right) \right] = \nabla^2 g_K \left(\frac{w}{\eta_j \sigma} \right). \quad (\text{C.26})$$

Thus changing the positive amplitude $\eta_j \sigma$ does not change the Hessian bound. Define the shift as a function of the perturbed inner coordinate

$$\delta_j(s) = \sqrt{K} \eta_j \sigma \sin^2 \left(\frac{\pi s}{2 \eta_j \sigma} \right). \quad (\text{C.27})$$

The profile and its derivative vanish at zero, while $\delta_j(\pm \eta_j \sigma) = \sqrt{K} \eta_j \sigma$ and $\delta'_j(\pm \eta_j \sigma) = 0$. Thus either sign of the gradient error activates the same shift and leaves its derivative zero. Writing $\chi_j(q) := \chi(q/(\sqrt{K} \eta_j \sigma))$ gives exactly the component formula (5.9). Its displacement $\chi_j(a[j] + b[j])\delta_j(s[j])$ depends on the inner coordinates, and the cutoff keeps second derivatives bounded globally. Once $s[j]$ is fixed at either activated value, the updates of $(v[j], a[j], b[j])$ agree with those of g_δ in (5.6), with parameters $(L, K, \sqrt{K} \eta_j \sigma)$, wherever $|a[j] + b[j]| \leq R_0 \sqrt{K} \eta_j \sigma$.

Its full-gradient Lipschitz constant is at most $LC_0 \leq \ell$, and its inner-PŁ parameter is at least $Lc_0/K \geq \ell/\kappa$. Also $\Phi_{\sigma,j}(v[j]) = LK(v[j])^2/2$, and the origin is both an outer minimizer and an inner maximizer. These properties hold for every positive amplitude, including the initial large steps; the step-size restriction in Lemma 5.5 is needed only for its trajectory bound.

Lemma 5.5 (Contribution of a single inner-gradient perturbation). *There are universal constants $c, C, c_{\text{step}}, c_{\text{high}} > 0$ and $\kappa_{\text{high}} \geq 2$ such that, for $\ell, \sigma > 0$, $\kappa \geq \kappa_{\text{high}}$, $\rho \geq \max\{1, c_{\text{high}} \kappa^2\}$, an integer $T \geq 2$, any predetermined step sizes $\eta_0 \geq \dots \geq \eta_{T-2} \geq 0$ and index $0 \leq j \leq T-2$ with $0 < \ell \eta_j \leq c_{\text{step}} \kappa$ admit an ℓ -smooth NC- (ℓ/κ) -PŁ component $g_{\sigma,j} : \mathbb{R} \times \mathbb{R}^3 \rightarrow \mathbb{R}$. Its outer objective $\Phi_{\sigma,j}(v) := \max_{y \in \mathbb{R}^3} g_{\sigma,j}(v; y)$ is quadratic with curvature $\Phi''_{\sigma,j} \asymp \ell \kappa$ and initial gaps $\Phi_{\sigma,j}(0) - \min_v \Phi_{\sigma,j}(v) = \Phi_{\sigma,j}(0) - g_{\sigma,j}(0; 0) = 0$. Starting from the origin with one equiprobable $\pm \sigma$ error in the third inner gradient at update j and exact gradients otherwise, both SGDA_{Sim} and SGDA_{Alt} generate outer iterates $v_t[j]$ satisfying*

$$\sum_{t < T} \mathbb{E} |\Phi'_{\sigma,j}(v_t[j])|^2 \geq c \frac{\ell \kappa^4 \sigma^2 \eta_j}{\rho} \quad \text{if} \quad \sum_{t=j+1}^{T-2} \eta_t \geq C \frac{\rho}{\ell \kappa}. \quad (5.13)$$

Proof. Once the error activates a shift, the component agrees with the shifted quadratic as long as its inner coordinates stay inside the cutoff. We verify this agreement over the displacement window, then use the non-increasing steps to count the iterates in that window.

Let ξ take values ± 1 with equal probability. All coordinates remain zero until update j , which sends $s[j]$ to $\eta_j \sigma \xi$ and leaves the other coordinates zero. At either sign, $\delta_j(s[j]) = \sqrt{K} \eta_j \sigma$ and $\delta'_j(s[j]) = 0$. Thus $s[j]$ stays fixed and both signs activate the same shift.

Choose

$$c_{\text{high}} = \gamma^2 C_{\text{resp}}, \quad c_{\text{step}} = \frac{c_r \gamma}{\theta},$$

and take the constant in the subsequent-sum condition at least $c_r/(\theta \gamma)$. Assume that this condition holds. For $j+1 \leq t \leq T-1$, write

$$\tau_t = \frac{L}{K} \sum_{r=j+1}^{t-1} \eta_r.$$

By the hypotheses, $\rho \geq C_{\text{resp}} K^2$, each increment of τ_t is at most $L \eta_j / K \leq c_r$, and $\tau_{T-1} \geq c_r \rho / K^2$.

Consider the exact-gradient trajectory (v_t, a_t, b_t) of the uncut shifted quadratic in (5.6), with parameters $(L, K, \sqrt{K} \eta_j \sigma)$, starting at the origin at time $j+1$. Lemma 5.4 gives

$$\begin{aligned} v_t &\geq c_r \frac{K^{3/2} \eta_j \sigma}{\rho} && \text{when } C_r \leq \tau_t \leq \frac{c_r \rho}{K^2}, \\ |a_t| + |b_t| &\leq C_r \sqrt{K} \eta_j \sigma && \text{when } \tau_t \leq \frac{c_r \rho}{K^2}. \end{aligned}$$

Let i and k be the first indices with $\tau_i \geq C_r$ and $\tau_k \geq c_r \rho / K^2$. They exist because $\tau_{T-1} \geq c_r \rho / K^2$; moreover, $c_r \rho / K^2 \geq 2(C_r + 2c_r)$ and each increment is at most c_r , so $i < k \leq T - 1$.

For $j + 1 \leq t < k$, the quadratic trajectory satisfies $|a_t + b_t| \leq C_r \sqrt{K} \eta_j \sigma < R_0 \sqrt{K} \eta_j \sigma$. Starting from the common state at time $j + 1$, induction therefore identifies it with the actual component trajectory through iterate $k - 1$. Every gradient query needed for this induction lies where the cutoff is identically one. For SGDA_{Alt} , the inner-gradient query after the primal update has the same $a + b$, so the agreement also holds at that query. Thus $v_t[j] = v_t$ on the interval we use below.

Since $\eta_t \leq \eta_j$ for $t > j$ and the first crossing of C_r overshoots by at most c_r ,

$$(k - i) \frac{L \eta_j}{K} \geq \tau_k - \tau_i \geq \frac{c_r \rho}{K^2} - C_r - c_r \geq \frac{c_r \rho}{2K^2}. \quad (\text{C.28})$$

For every $i \leq t < k$, the displacement bound and $\Phi'_{\sigma,j}(v) = LKv$ give $\Phi'_{\sigma,j}(v_t[j]) \geq c_r LK^{5/2} \eta_j \sigma / \rho$. These are $k - i$ reported iterates, so

$$\sum_{t < T} \mathbb{E} |\Phi'_{\sigma,j}(v_t[j])|^2 \geq (k - i) \left(\frac{c_r LK^{5/2} \eta_j \sigma}{\rho} \right)^2 \geq \frac{c_r^3}{2} \frac{LK^4 \sigma^2 \eta_j}{\rho} \geq c \frac{\ell \kappa^4 \sigma^2 \eta_j}{\rho}. \quad (\text{C.29})$$

All constants are independent of the horizon and step-size sequence, and the same displacement estimate holds for either noise sign. In the aggregate game, the fixed oracle constructed in Lemma C.2 generates each component's prescribed perturbation at its designated update. $\square$

We now realize the prescribed perturbations through one fixed oracle and sum their contributions. Only components with enough subsequent cumulative steps contribute to the estimate; the omitted components account for a bounded total step size, leaving a constant fraction of S when S is sufficiently large.

Lemma C.2 (Lower bound from the noise components). *There are universal $c, C > 0$ with the following property. Fix $\ell, \sigma > 0$, $\kappa \geq \kappa_{\text{high}}$, $\rho \geq \max\{1, c_{\text{high}} \kappa^2\}$, an integer $T \geq 2$, a predetermined schedule $\eta_0 \geq \dots \geq \eta_{T-2} \geq 0$, and $m \in \{0, \dots, T - 1\}$ with $\ell \eta_t < 8$ for $m \leq t \leq T - 2$. Set $S = \sum_{t=m}^{T-2} \eta_t$. The components in (5.8) can be chosen ℓ -smooth and NC- (ℓ/κ) -PL, with $\Phi_{\sigma,j}(0) = \min \Phi_{\sigma,j} = g_{\sigma,j}(0, 0) = 0$. Both SGDA_{Sim} and SGDA_{Alt} on g_σ , initialized at the origin with exact outer gradients and the fixed inner oracle $G_y(v, y; \xi) = \nabla_y g_\sigma(v, y) + \sigma \xi e(y)$ from Section 5.2.3, satisfy*

$$\frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E} \|\nabla \Phi_\sigma(v_t)\|^2 \geq c \frac{\kappa^4 \ell \sigma^2 S}{\rho T} \quad \text{if } S \geq \frac{C \rho}{\kappa \ell}. \quad (\text{C.30})$$

Proof. Let $J = \{0, \dots, T - 2\}$. Set $g_{\sigma,j} \equiv 0$ for $\eta_j = 0$, and choose every positive-step block from (C.25). The preceding smoothness, inner-PL, and initialization estimates apply to all these blocks, with no restriction on the initial step sizes.

The selector $e(y)$ from Section 5.2.3 chooses the $s[j]$ -coordinate of the smallest $j \in J$ with $s[j] = 0$, or is zero if no such index exists. Its selection regions are finite intersections of coordinate hyperplanes and their complements, so it is measurable and has norm at most one. Fresh independent uniform signs make the inner oracle pointwise unbiased, with variance $\sigma^2 \|e(y)\|^2 \leq \sigma^2$.

We next verify that this oracle activates the intended component at each positive update. The induction invariant is that, at the start of a positive update j , all blocks with index at least j remain at the origin, while every earlier block has $s[i] = \pm \eta_i \sigma \neq 0$. Stationarity at the origin keeps the unactivated blocks fixed during the primal update. Thus $e(y)$ selects the third inner coordinate of block j for either update order, and update j sends $s[j]$ to $\pm \eta_j \sigma$ while leaving its other coordinates at zero.

At these peaks, $\partial_s r_K(a, b, \pm 1) = 0$ for every a, b , including outside the cutoff region. All subsequent exact $s[j]$ -gradients therefore vanish, so $s[j]$ stays fixed even if a large step changes the cutoff factor. The remaining unactivated blocks stay at the origin, completing the induction. Since the schedule is non-increasing, zero steps are trailing. The selector then repeatedly chooses the first zero-step block, but the zero step sizes leave every iterate unchanged. Consequently, each nonzero block receives one

equiprobable $\pm\sigma$ error at its assigned update and exact gradients otherwise, with independent signs across blocks.

Since the coordinates are disjoint,

$$\|\nabla\Phi_\sigma(v_t)\|^2 = \sum_{j \in J} \|\nabla\Phi_{\sigma,j}(v_t[j])\|^2.$$

Let C_{one} be the remaining-step constant in Lemma 5.5, and enlarge κ_{high} so that $8 \leq c_{\text{step}}\kappa$. Every positive step with index $t \geq m$ then satisfies its step-size condition. Define the tail indices $J_{\text{tail}} := \{m, \dots, T-2\}$ and $J_{\text{good}} := \{j \in J_{\text{tail}} : \sum_{t=j+1}^{T-2} \eta_t \geq C_{\text{one}}\rho/(\kappa\ell)\}$. Non-increasingness ensures $\eta_j > 0$ for every $j \in J_{\text{good}}$. If $J_{\text{tail}} \setminus J_{\text{good}}$ is nonempty, its first index j_0 satisfies

$$\sum_{j \in J_{\text{tail}} \setminus J_{\text{good}}} \eta_j \leq \sum_{t=j_0}^{T-2} \eta_t < \frac{8}{\ell} + \frac{C_{\text{one}}\rho}{\kappa\ell} \leq \frac{C_1\rho}{\kappa\ell}, \quad C_1 := C_{\text{one}} + \frac{8}{c_{\text{high}}},$$

where the last inequality uses $\rho \geq c_{\text{high}}\kappa^2$ and $\kappa \geq 1$. If the excluded set is empty, its total step size is zero. Taking $C := 2C_1$, the condition $S \geq C\rho/(\kappa\ell)$ implies $\sum_{j \in J_{\text{good}}} \eta_j \geq S/2$. Summing (5.13) over these indices gives

$$\frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E} \|\nabla\Phi_\sigma(v_t)\|^2 \geq c \frac{\kappa^4 \ell \sigma^2}{\rho T} \sum_{j \in J_{\text{good}}} \eta_j \geq \frac{c}{2} \frac{\kappa^4 \ell \sigma^2 S}{\rho T}.$$

Reducing the universal constant c proves the claim. $\square$

C.4 PROOF OF THE LOWER BOUND FOR f_{stable}

The aggregate construction already supplies a fixed oracle using only the inner coordinates. We add the deterministic corridor and combine its lower bound with the contributions of the noise components.

Lemma 5.6 (Stable-step lower bound and admissibility). *There are universal $c, c_{\text{high}} > 0$ and $\kappa_{\text{high}} \geq 2$ such that, for $\ell, \Delta > 0$, $\sigma \geq 0$, $\kappa \geq \kappa_{\text{high}}$, $\rho \geq \max\{1, c_{\text{high}}\kappa^2\}$, an integer $T \geq 1$, and predetermined steps $\eta_0 \geq \dots \geq \eta_{T-2} \geq 0$, the following holds. Let m count the entries $\eta_t \geq 8/\ell$ and assume $m+1 < T/2$. For each of SGDA_{Sim} and SGDA_{Alt} , the game f_{stable} given by (5.4) and a fixed oracle can be chosen as an instance in $\mathcal{C}(\ell, \kappa, \Delta, \sigma)$ with initial outer gap Δ and inner gap zero at the origin. Writing $S = \sum_{t=m}^{T-2} \eta_t$ and $\rho\Delta/0 := +\infty$, its iterates satisfy*

$$\frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E} \|\nabla\Phi_{\text{stable}}(u_t, v_t)\|^2 \geq c \min \left\{ \ell\Delta, \max \left\{ \frac{\rho\Delta}{S}, \frac{\kappa^4 \ell \sigma^2 S}{\rho T} \right\} \right\}. \quad (5.14)$$

Proof. For the chosen algorithm, SGDA_{Sim} or SGDA_{Alt} , use the predetermined non-increasing step sizes in the statement. Let $J = \{0, \dots, T-2\}$, and choose the corridor from Lemma 5.3. For $\sigma > 0$, choose the aggregate noise game g_σ and its fixed inner oracle from Lemma C.2; for $\sigma = 0$, set $g_{\sigma,j} \equiv 0$ for every $j \in J$ and use exact gradients. Use these games in (5.4), take exact outer gradients, and initialize every coordinate at zero. In either case, the dimensions are $d_x = 1 + |J| = T$ and $d_y = 3|J| = 3(T-1)$.

We first verify the game assumptions. Each component has full-gradient Lipschitz constant at most ℓ . Since the components use disjoint coordinates, summing the squared component inequalities gives

$$\|\nabla f_{\text{stable}}(u, v; y) - \nabla f_{\text{stable}}(\tilde{u}, \tilde{v}; \tilde{y})\|^2 \leq \ell^2 (\|u - \tilde{u}\|^2 + \|v - \tilde{v}\|^2 + \|y - \tilde{y}\|^2).$$

Thus the smoothness assumption holds without a dimension factor. Each inner maximum is finite and attained, so their finite product attains the maximum of f_{stable} . Its outer objective is

$$\Phi_{\text{stable}}(u, v) = g_{\text{corridor}}(u) + \Phi_\sigma(v) = g_{\text{corridor}}(u) + \frac{LK}{2} \sum_{\substack{j \in J \\ \eta_j \sigma > 0}} (v[j])^2.$$

It is differentiable and lower bounded. Summing the component inner-PŁ inequalities gives

$$\begin{aligned}
\frac{1}{2} \|\nabla_y f_{\text{stable}}(u, v; y)\|^2 &= \frac{1}{2} \sum_{j \in J} \|\nabla_{y[j]} g_{\sigma, j}(v[j]; y[j])\|^2 \\
&\geq \frac{\ell}{\kappa} \sum_{j \in J} (\Phi_{\sigma, j}(v[j]) - g_{\sigma, j}(v[j]; y[j])) \\
&= \frac{\ell}{\kappa} (\Phi_{\text{stable}}(u, v) - f_{\text{stable}}(u, v; y)).
\end{aligned}$$

The corridor supplies outer gap Δ , each perturbation component has zero outer gap at the origin, and every component is initially maximized in its inner variables. Hence the initial outer and inner gaps are Δ and zero. These conclusions also hold for the corridor alone, with a vacuous inner-PŁ inequality.

Adding the corridor does not change any noise-component trajectory, because its coordinates are disjoint and the inner oracle depends only on (v, y) . The oracle verification in the proof of [Lemma C.2](#) therefore also applies to f_{stable} . Together with the game assumptions above, this proves that the origin-initialized instance belongs to $\mathcal{C}(\ell, \kappa, \Delta, \sigma)$.

It remains to prove the lower bound. Let C be the threshold constant in [Lemma C.2](#), and increase κ_{high} so that $\kappa_{\text{high}} \geq C$. If $S < C\rho/(\kappa\ell)$, including $S = 0$, the corridor gives $c\ell\Delta$, since $(T - m)/T > 1/2$ and $S/\rho < 1/\ell$. This proves the capped bound.

Otherwise $S > 0$, and [Lemma C.2](#) applies when $\sigma > 0$; when $\sigma = 0$, the noise contribution is zero. Since the squared outer gradients of disjoint components add, their bounds give

$$\frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E} \|\nabla \Phi_{\text{stable}}(u_t, v_t)\|^2 \geq c \left[\min \left\{ \ell\Delta, \frac{\rho\Delta}{S} \right\} + \frac{\kappa^4 \ell \sigma^2 S}{\rho T} \right]. \quad (\text{C.31})$$

The bracket dominates the capped maximum in (5.14), proving the lemma. $\square$

D PROOFS FOR THE LOW-RATIO LOWER BOUND

We prove the two properties used by [Theorem 6.3](#): small steps preserve a region of nonzero outer gradient, and the initially larger steps can be made to end inside that region. The first requires a smooth game with a stable periodic solution; the second modifies that specific game away from the solution. The final rescaling and initialization checks appear in [Appendix A.3](#).

D.1 CONSTRUCTION OF g_{cycle}

Lemma 6.1 (An attracting periodic orbit of GDA flow in g_{cycle}). *There are universal $C, \mu, h_\star > 0$ and a smooth residual R such that the base game $g_{\text{cycle}} : \mathbb{R}^2 \times \mathbb{R}^4 \rightarrow \mathbb{R}$ in (6.2) is jointly C -smooth and affine in u , has outer objective $\Phi_{g_{\text{cycle}}}(u) = \frac{1}{2} \|u\|^2$, and satisfies the inner μ -PŁ inequality. The path (6.3), with $y_\star = (a_\star, b_\star)$, traces a locally stable 2π -periodic orbit Γ of GDA flow. There is a compact neighborhood $\mathcal{U}$ of Γ such that both SGDA_{sim} and SGDA_{Alt} , started in $\mathcal{U}$ with exact gradients and equal primal and dual steps $h_t \in [0, h_\star]$, satisfy $(u_t, y_t) \in \mathcal{U}$ and $\|\nabla \Phi_{g_{\text{cycle}}}(u_t)\| = \|u_t\| \geq \frac{1}{2}$ for all $t \geq 0$, where $y_t = (a_t, b_t)$.*

Proof. We first verify global smoothness and inner PŁ, then check that the prescribed curve solves the GDA equations. A rotating frame separates phase shifts along the orbit from departures that decay toward it. We use this decay to construct a Lyapunov function whose sublevel set is invariant under both discrete updates.

Global smoothness and inner PŁ. We construct $g_{\text{cycle}}(u, a, b) = \sqrt{2} u^\top a - \|a\|^2 - \|R(a, b)\|^2 / 2$ on $\mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{R}^2$. Its affine dependence on u will be essential to the ratio scaling. The residual has the form $R(a, b) = b + \chi_b(b)\psi(a)$: the values and first derivatives of ψ produce the periodic solution, its second derivatives control departures from that solution, and the cutoffs enforce global smoothness.

Let $J = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ be rotation by a right angle. The target path is $u_*(t) = (\cos t, \sin t)^\top$, with $a_* = -Ju_*/\sqrt{2}$ and $b_* = u_*/\sqrt{2}$. Since $\dot{u} = -\sqrt{2}a$ and, where the cutoffs equal one, $\dot{b} = -(b + \psi(a))$, the first and third equations require $\psi(a_*) = a_* - Ja_*$. The remaining equation requires $D\psi(a_*)^\top a_* = -2a_* + Ja_*$. We satisfy these value and derivative conditions first, then check contraction through the remaining derivatives.

Choose a smooth cutoff $\chi_a : [0, \infty) \rightarrow [0, 1]$ that equals one on a neighborhood of $\frac{1}{2}$ and vanishes outside a slightly larger compact interval. Define the polynomials

$$c(q) := 1 - 3(q - \tfrac{1}{2}) + \tfrac{7}{2}(q - \tfrac{1}{2})^2, \quad d(q) := -1 + (q - \tfrac{1}{2}), \quad (\text{D.1})$$

and set

$$\psi(a) := \chi_a(\|a\|^2) [c(\|a\|^2)a + d(\|a\|^2)Ja]. \quad (\text{D.2})$$

The orbit conditions require $c(1/2) = 1$, $d(1/2) = -1$, and $c'(1/2) = -3$. The remaining choices $c''(1/2) = 7$ and $d'(1/2) = 1$ make the transverse linearization stable, as verified below. The map ψ is smooth and compactly supported. Choose a smooth radial cutoff $\chi_b : \mathbb{R}^2 \rightarrow [0, 1]$ that equals one on a neighborhood of the circle $\|b\| = 1/\sqrt{2}$, vanishes outside a sufficiently large ball, and satisfies

$$\|\psi\|_\infty \|\nabla \chi_b\|_\infty \leq \frac{1}{2}. \quad (\text{D.3})$$

Define

$$R(a, b) := b + \chi_b(b)\psi(a), \quad (\text{D.4})$$

and

$$g_{\text{cycle}}(u, a, b) := \sqrt{2}u^\top a - \|a\|^2 - \frac{1}{2}\|R(a, b)\|^2. \quad (\text{D.5})$$

For every fixed a and target $w \in \mathbb{R}^2$, the map $b \mapsto w - \chi_b(b)\psi(a)$ is a contraction with factor at most $1/2$ by (D.3). Its unique fixed point solves $R(a, b) = w$. Thus $b \mapsto R(a, b)$ is bijective. Moreover,

$$D_b R = I + \psi(a)\nabla \chi_b(b)^\top, \quad \|D_b R - I\| \leq \tfrac{1}{2}, \quad \|(D_b R)^{-1}\| \leq 2.$$

The inverse function theorem gives a smooth inverse. Thus $T(a, b) := (\sqrt{2}a, R(a, b))$ is a smooth bijection. Its triangular derivative has diagonal blocks $\sqrt{2}I_2$ and $D_b R$; boundedness of $D_a R$ and $(D_b R)^{-1}$ gives $\|(DT)^{-1}\| \leq C$, and hence $\|DT^\top q\| \geq C^{-1}\|q\|$ for every (a, b) and $q \in \mathbb{R}^4$.

All derivatives of ψ are bounded. Whenever a second derivative of R is nonzero, both a and b lie in fixed compact sets; elsewhere R is affine. Hence DR and RD^2R are globally bounded, and so is $\nabla^2 \|R\|^2/2$. The full gradient of g_{cycle} is therefore universally Lipschitz.

For each a , maximizing over b cancels the residual. Maximizing over a then gives $\Phi_{g_{\text{cycle}}}(u) = \|u\|^2/2$ and

$$\Phi_{g_{\text{cycle}}}(u) - g_{\text{cycle}}(u, a, b) = \frac{1}{2}\|u - \sqrt{2}a\|^2 + \frac{1}{2}\|R(a, b)\|^2. \quad (\text{D.6})$$

The surjectivity and transposed-derivative bound for T verify the hypotheses of Lemma B.1 with outer vector $(u, 0)$, giving inner μ -PL for a universal $\mu > 0$.

The periodic orbit and its attraction. Along (6.3), $b_* = Ja_*$, $\|a_*\|^2 = \frac{1}{2}$, and both cutoffs are constant. Hence

$$\psi(a_*) = a_* - Ja_*, \quad R(a_*, b_*) = a_*.$$

On the neighborhood where both cutoff functions equal one, writing $q = \|a\|^2$,

$$D\psi(a)^\top a = [c(q) + 2qc'(q)]a - d(q)Ja. \quad (\text{D.7})$$

The values $c(\frac{1}{2}) = 1$, $c'(\frac{1}{2}) = -3$, and $d(\frac{1}{2}) = -1$ give

$$\dot{u}_* = -\sqrt{2}a_* = Ju_*, \quad \dot{a}_* = \sqrt{2}u_* - 2a_* - D\psi(a_*)^\top a_* = Ja_*, \quad \dot{b}_* = -R(a_*, b_*) = Jb_*.$$

Thus (6.3) solves the GDA differential equation and is 2π -periodic.

Along this solution, $-\nabla_u g_{\text{cycle}} = Ju_*$, $\nabla \Phi_{g_{\text{cycle}}} = u_*$, and $\|\nabla_{a,b} g_{\text{cycle}}\|^2 = \|\dot{a}_*\|^2 + \|\dot{b}_*\|^2 = 1$. By (D.6), the inner gap equals $5/4$. Inner ascent decreases this gap at rate one, while the motion of u increases it at the same rate: $\|\nabla_u g_{\text{cycle}}\|^2 = 1$ and $u_*^\top \dot{u}_* = 0$. These contributions cancel, so the gap remains positive despite global inner PL.

The definitions imply $\psi(e^{tJ}a) = e^{tJ}\psi(a)$ and $R(e^{tJ}a, e^{tJ}b) = e^{tJ}R(a, b)$. Thus the derivative calculation at any point of the periodic solution is the rotation of the calculation at $t = 0$. Direct differentiation at $t = 0$, using $c''(\frac{1}{2}) = 7$ and $d'(\frac{1}{2}) = 1$, gives

$$A_* := D\psi(a_*(0)) = \begin{pmatrix} 1 & 0 \\ -1 & -2 \end{pmatrix}, \quad S_* := D_a[D\psi(a)^\top R(a, b)]_{(a_*(0), b_*(0))} = \begin{pmatrix} -1 & 3 \\ 3 & 2 \end{pmatrix}. \quad (\text{D.8})$$

Write $z = (u, a, b)$, $z_*(t) = (u_*(t), a_*(t), b_*(t))$, and let $F = (-\nabla_u g_{\text{cycle}}, \nabla_{a,b} g_{\text{cycle}})$ be the GDA vector field. Set $\mathcal{J} = \text{diag}(J, J, J)$ and $\mathcal{Q}_t = \text{diag}(e^{tJ}, e^{tJ}, e^{tJ})$. Rotational equivariance gives $F(\mathcal{Q}_t z) = \mathcal{Q}_t F(z)$, so $\tilde{z}(t) = \mathcal{Q}_t^\top z(t)$ satisfies $\dot{\tilde{z}} = F(\tilde{z}) - \mathcal{J}\tilde{z}$. The periodic solution becomes the fixed point $z_*(0)$, whose Jacobian is

$$M_* = DF(z_*(0)) - \mathcal{J} = \begin{pmatrix} -J & -\sqrt{2}I_2 & 0 \\ \sqrt{2}I_2 & -2I_2 - S_* - J & -A_*^\top \\ 0 & -A_* & -I_2 - J \end{pmatrix}. \quad (\text{D.9})$$

A direct determinant calculation gives

$$\det(\lambda I - M_*) = \lambda(\lambda^5 + 7\lambda^4 + 7\lambda^3 + 21\lambda^2 + 10\lambda + 4). \quad (\text{D.10})$$

For the quintic factor, the first column of the Routh array is $1, 7, 4, 9/2, 370/63, 4$. It has no sign change, so the Routh–Hurwitz criterion excludes roots in the open right half-plane. The array has neither a zero first-column entry nor an all-zero row, excluding roots on the imaginary axis as well. Thus all five roots have strictly negative real parts. The remaining zero eigenvalue is simple and represents phase shifts $z_*(t) \mapsto z_*(t + s)$; its eigenvector is $\dot{z}_*(0) = \mathcal{J}z_*(0)$.

To control departures from the orbit, choose a basis $S \in \mathbb{R}^{6 \times 5}$ of the stable invariant subspace of M_* , and let A_- be its restriction in this basis.

For sufficiently small $\varepsilon > 0$, the map $\mathcal{X}(\vartheta, \xi) = \mathcal{Q}_\vartheta(z_*(0) + S\xi)$, with $\vartheta \in \mathbb{R}/(2\pi\mathbb{Z})$ and $\xi \in \mathbb{R}^5$, $\|\xi\| < \varepsilon$, gives smooth coordinates near Γ : the columns of S complement its tangent direction, and the orbit is compact and embedded. Using $M_* S = S A_-$ and Taylor expansion of F at $z_*(0)$, differentiation of these coordinates along GDA gives

$$(\dot{\vartheta} - 1)\mathcal{J}(z_*(0) + S\xi) + S(\dot{\xi} - A_- \xi) = O(\|\xi\|^2).$$

The coefficient matrix $[\mathcal{J}(z_*(0) + S\xi), S]$ remains uniformly invertible for small ξ . Hence $\dot{\vartheta} = 1 + O(\|\xi\|^2)$ and $\dot{\xi} = A_- \xi + O(\|\xi\|^2)$, uniformly in the phase ϑ . Since A_- is Hurwitz, choose $P \succ 0$ satisfying $A_-^\top P + P A_- = -I$, and define $V(\mathcal{X}(\vartheta, \xi)) = \xi^\top P \xi$. On a smaller neighborhood,

$$V(z) \asymp \text{dist}(z, \Gamma)^2, \quad \dot{V} = -\|\xi\|^2 + O(\|\xi\|^3) \leq -cV$$

for a universal $c > 0$. This proves local exponential orbital stability.

An invariant neighborhood for discrete GDA. We use $\dot{V} \leq -cV$ to obtain $V(z^+) \leq (1 - ch)V(z) + Ch^2$ for both update rules, then choose h_* so that a fixed sublevel set of V is invariant. Choose a fixed small $\delta > 0$ so that $\mathcal{U} := \{V \leq \delta\}$ is compactly contained in this neighborhood and in $\{\|u\| \geq 1/2\}$. Simultaneous GDA gives $z_{\text{sim}}^+ = z + hF(z)$. In the alternating update, the change in u is $-h\sqrt{2}a$, so its inner-gradient query changes only the a -gradient, by $-2ha$. Thus $z_{\text{Alt}}^+ - z_{\text{sim}}^+ = (0, -2h^2a, 0)$. For $z \in \mathcal{U}$ and sufficiently small h , both updates and the line segments joining them to z remain in a fixed compact neighborhood where V is smooth. Taylor expansion therefore gives, uniformly for either update,

$$V(z^+) \leq V(z) + h\nabla V(z)^\top F(z) + Ch^2 \leq (1 - ch)V(z) + Ch^2 \leq (1 - ch)\delta + Ch^2 \leq \delta,$$

where we choose $h_* > 0$ small enough that $ch_* \leq 1$ and $Ch_* \leq c\delta$. Thus $\mathcal{U}$ is invariant for every $h \in [0, h_*]$, including $h = 0$. Induction and $\nabla \Phi_{g_{\text{cycle}}}(u) = u$ give the discrete conclusion for every such step sequence. All constants depend only on the fixed game.

For the next lemma, fix a bounded open neighborhood $\mathcal{W} \subset \mathbb{R}^4$ of the inner projection of $\mathcal{U}$. We will preserve the game for every u whenever $(a, b) \in \mathcal{W}$. This also protects the alternating inner-gradient query after the primal update, since that update changes only u . $\square$

D.2 CONSTRUCTION OF $\tilde{g}_{\text{cycle}}$

To construct $\tilde{g}_{\text{cycle}}$, we extend the base game to accommodate finitely many initial steps $h_t > h_*$. At the initial query states y_t to be chosen, for $0 \leq t < m$ and $e_1 = (1, 0)^\top$, the required gradients are

$$\nabla_u \tilde{g}_{\text{cycle}}(e_1, y_t) = 0, \quad \nabla_y \tilde{g}_{\text{cycle}}(e_1, y_t) = \frac{y_{t+1} - y_t}{h_t}. \quad (\text{D.11})$$

They keep the outer variable fixed and send each inner state to the next, ending on the cycle. A local change of inner coordinates prescribes these gradients through its derivative. We preserve a to retain both the primal gradient $\sqrt{2}a$ and the bilinear term $\sqrt{2}u^\top a$. A nonlinear change of these coordinates would introduce inner-Hessian terms proportional to u , obstructing global joint smoothness. The following lemma realizes the prescribed derivative inside a ball; disjoint balls will keep its bounds independent of the number of steps.

Lemma D.1 (Local derivative modification that leaves the first two coordinates unchanged). *Fix $M < \infty$ and $\delta > 0$. There is $C = C(M, \delta)$ such that, for every $z \in \mathbb{R}^5$ and every*

$$N = \begin{pmatrix} I_2 & 0 \\ P & Q \end{pmatrix}, \quad \det N > 0, \quad \|N\| + \|N^{-1}\| \leq M, \quad (\text{D.12})$$

there is a smooth bijection $G : \mathbb{R}^5 \rightarrow \mathbb{R}^5$ with a smooth inverse. For every w , the first two coordinates of $G(w)$ equal those of w . Moreover, $G(z) = z$, $DG(z) = N$, G equals the identity outside $B(z, \delta)$, G maps this ball onto itself, and

$$\|DG\|_\infty + \|DG^{-1}\|_\infty + \|D^2G\|_\infty \leq C. \quad (\text{D.13})$$

Proof. We join the identity matrix to N through uniformly invertible matrices, then realize this matrix path as the derivative of a compactly supported flow. Since $\det N > 0$ and the first diagonal block is I_2 , the matrix Q is invertible and has positive determinant. Write $Q = US$, where $U^\top U = I$, $\det U = 1$, and S is symmetric positive definite. Choose a skew-symmetric matrix A with $e^A = U$ and $\|A\| \leq \pi$, and define

$$Q_\tau = e^{\tau A}((1 - \tau)I + \tau S), \quad N_\tau = \begin{pmatrix} I_2 & 0 \\ \tau P & Q_\tau \end{pmatrix}.$$

The matrices N_τ , their inverses, and their derivatives are uniformly bounded in terms of M . Put $A_\tau = \dot{N}_\tau N_\tau^{-1}$. Choose a smooth radial cutoff χ that equals one on $B(z, \delta/3)$ and vanishes outside $B(z, 2\delta/3)$, and set

$$X_\tau(w) = \chi(w)A_\tau(w - z).$$

This time-dependent differential equation has zero first two components and uniformly bounded first two derivatives. Let $G_\tau(w)$ denote its solution at time τ starting from w at time zero. Then G_τ fixes z , preserves the first two coordinates, and satisfies

$$\frac{d}{d\tau} DG_\tau(z) = A_\tau DG_\tau(z), \quad DG_0(z) = I,$$

so $DG_\tau(z) = N_\tau$. Differentiating this differential equation once and twice with respect to the initial point, and then applying Gronwall's inequality, gives (D.13) for $G = G_1$ and its inverse. Since the vector field vanishes in a neighborhood of the boundary, the ball is mapped onto itself. $\square$

Lemma 6.2 (GDA can reach the orbit in a modified game $\tilde{g}_{\text{cycle}}$). *There are universal $C, \mu > 0$ with the following property. For any finite sequence of steps $h_0, \dots, h_{m-1} > h_*$, including $m = 0$, we can modify the game g_{cycle} in Lemma 6.1 to obtain $\tilde{g}_{\text{cycle}} : \mathbb{R}^2 \times \mathbb{R}^5 \rightarrow \mathbb{R}$. This game is jointly C -smooth and affine in u , satisfies the inner μ -PL inequality, and has outer objective $\Phi_{\tilde{g}_{\text{cycle}}}(u) = \|u\|^2/2$. With $y = (a, b, r)$ and $e_1 = (1, 0)^\top$, there is an initialization (e_1, y_0) with inner gap at most C . From this initialization, both GDA_{Sim} and GDA_{Alb} , using exact gradients and equal primal and dual steps, keep $u_t = e_1$ for $0 \leq t \leq m$ and reach $y_m = (y_*(0), 0)$. For every continuation with $h_t \in [0, h_*]$, each algorithm follows its g_{cycle} dynamics in (u, a, b) , while r remains zero.*

Proof. We choose separated inner states whose required update vectors are uniformly bounded. At each state, a local coordinate modification realizes the desired gradient while preserving the first two inner coordinates. The residual representation then verifies inner PL, and the disjoint modification regions give uniform smoothness.

The case $m = 0$ uses $\tilde{g}_{\text{cycle}}(u, v, r) = g_{\text{cycle}}(u, v)$ and $y_0 = (y_*(0), 0)$. All initialization quantities are bounded on the fixed periodic solution, and the initial primal gradient is $\sqrt{2}a_*(0) = -Je_1$, of norm one. Suppose $m \geq 1$ and write $y = (a, b, r) \in \mathbb{R}^2 \times \mathbb{R}^2 \times \mathbb{R}$. The extra coordinate r separates the query states even when their first four coordinates agree. Its value is not itself penalized, so this separation need not increase the initial gap. Define

$$\begin{aligned} r_t &= -\sum_{j=t}^{m-1} h_j \quad (0 \leq t < m), \quad r_m = 0, \\ y_t &= (0, -Be_1, r_t) \quad (t < m), \quad y_m = (a_*(0), b_*(0), 0). \end{aligned} \quad (\text{D.14})$$

Here B is a fixed universal constant large enough that a neighborhood of $(a, b) = (0, -Be_1)$ is outside the support of the nonlinear part of R and separated from $\mathcal{W}$. We protect the lift $\mathcal{W} \times \mathbb{R}$ in the inner space (a, b, r) . Let $\tilde{H}(a, b, r) = ((a, R(a, b)), r)$. At all y_t with $t < m$, it is the identity in a fixed neighborhood.

Put $v_t = (y_{t+1} - y_t)/h_t = (v_t^a, v_t^b, 1)$ for $0 \leq t < m$. Its first four coordinates are zero except at $t = m - 1$; their magnitude at the last step is bounded by a universal constant since $h_{m-1} > h_*$ and the first four endpoint coordinates are fixed. Let $\pi_b : \mathbb{R}^5 \rightarrow \mathbb{R}^2$ project onto the two coordinates corresponding to b . We seek a smooth bijection $\tilde{H} : \mathbb{R}^5 \rightarrow \mathbb{R}^5$ that preserves a , fixes the initial query states $y_0, \dots, y_{m-1}$, and agrees with $\tilde{H}$ on $\mathcal{W} \times \mathbb{R}$. Define the game in terms of this map by

$$\tilde{g}_{\text{cycle}}(u, y) = \left\langle (u, 0), \tilde{T}(y) \right\rangle - \frac{1}{2} \|\tilde{T}(y)\|^2, \quad \tilde{T}(a, b, r) = (\sqrt{2}a, \pi_b \tilde{H}(a, b, r)) \in \mathbb{R}^4. \quad (\text{D.15})$$

At each initial query state y_t , $t < m$, we have $\tilde{T}(y_t) = (0, -Be_1)$, so $\nabla_u \tilde{g}_{\text{cycle}}(e_1, y_t) = 0$. The required inner gradient is therefore equivalent to

$$D\tilde{H}(y_t)^\top (\sqrt{2}e_1, Be_1, 0) = v_t. \quad (\text{D.16})$$

This identity determines the row corresponding to b_1 . We choose the remaining two rows to make the derivative uniformly invertible with positive determinant:

$$D\tilde{H}(y_t) = \begin{pmatrix} I_2 & 0 & 0 \\ (v_t^a - \sqrt{2}e_1)^\top/B & (v_t^b)^\top/B & 1/B \\ 0 & e_2^\top & 0 \\ 0 & -Be_1^\top & 0 \end{pmatrix}. \quad (\text{D.17})$$

The lower-right 3×3 block has determinant one, so the full derivative also has determinant one. It and its inverse are uniformly bounded, since B is fixed and v_t is uniformly bounded. Distinct y_t with $t < m$ are separated by at least h_* , because their scalar coordinates differ by sums of steps exceeding h_* . Choose a universal ball radius smaller than one quarter of this separation and small enough that $\tilde{H}$ is the identity on every ball and all balls avoid $\mathcal{W} \times \mathbb{R}$.

Apply Lemma D.1 in these disjoint balls. Each local map fixes its center, has the prescribed derivative, preserves the first two coordinates, and maps its ball onto itself. Let G be their composition and set $\tilde{H} = G \circ \tilde{H}$. Since $\tilde{H}$ is the identity near every center, (D.17) holds. At most one factor of G is nonidentity along the image of any point, so the derivative, inverse-derivative, and second-derivative

bounds do not grow with m . Since $\bar{H}$ is bijective and equals the identity on each modification ball, no point outside those balls is mapped into them. Thus $\bar{H} = \tilde{H}$ on $\mathcal{W} \times \mathbb{R}$.

Since $\tilde{H}$ is onto and preserves its first two coordinates, $\tilde{T}$ is onto. For $q = (q_a, q_b) \in \mathbb{R}^4$,

$$D\tilde{T}(y)^\top q = D\tilde{H}(y)^\top (\sqrt{2}q_a, q_b, 0), \quad \left\| D\tilde{T}(y)^\top q \right\| \geq c \|q\|,$$

where the last inequality uses the uniform inverse-derivative bound for $\tilde{H}$. Lemma B.1 now gives $\max_y \tilde{g}_{\text{cycle}}(u, y) = \|u\|^2/2$ and global inner PL with a universal constant.

To verify joint smoothness, write $K := \pi_b \tilde{H}$ for the last two residual coordinates. The derivative bounds above control DK and D^2K uniformly. Wherever D^2K is nonzero, K is bounded as well: outside the modification balls this follows from the fixed compact support in (a, b) of the original residual's nonlinear part, and inside a ball it follows because the ball maps onto itself. In particular, its b -coordinates stay near $-Be_1$, regardless of the scalar center r_t . Thus

$$\left\| \nabla^2 \left(\frac{1}{2} \|K\|^2 \right) \right\| \leq \|DK\|^2 + \|K\| \|D^2K\| \leq C.$$

Expanding (D.15) gives $\tilde{g}_{\text{cycle}}(u, a, b, r) = \sqrt{2}u^\top a - \|a\|^2 - \|K(a, b, r)\|^2/2$. Since a is unchanged, the outer-dependent term remains bilinear, so the full Hessian is uniformly bounded. On $\mathcal{W} \times \mathbb{R}$, we have $K(a, b, r) = R(a, b)$; hence $\tilde{g}_{\text{cycle}}$ agrees with g_{cycle} for every u and is independent of r there.

Finally, (D.16) gives the target gradients in (D.11). At each initial query, the zero primal gradient leaves $u = e_1$ unchanged, so the simultaneous and alternating methods use the same inner-gradient query and both move from y_t to y_{t+1} . The initial inner gap is $(1 + B^2)/2$ and the initial primal gradient is zero.

At time m , both methods reach $(e_1, y_*(0), 0)$. Agreement with the base game on $\mathcal{W} \times \mathbb{R}$ covers both update orders, including the alternating inner-gradient query after the primal update. Independence of r keeps $r_t = 0$, while Lemma 6.1 keeps $(u_t, a_t, b_t) \in \mathcal{U}$ under every small-step continuation. For either $m = 0$ or $m \geq 1$, the construction gives $\|\nabla_u \tilde{g}_{\text{cycle}}(e_1, y_0)\| \leq 1$. All constants depend only on the fixed small-step construction, not on the number or values of the prescribed steps. $\square$

E COMPARISON WITH SMOOTHED-AGDA

We compare the expected squared outer gradient at a uniformly selected SGDA iterate with that at a point computed from a Smoothed-AGDA iterate pair. Algorithm 1 states the updates of Yang et al. (2022, Algorithm 2), with an auxiliary center z_t initialized at x_0 and regularization coefficient 2ℓ .

Algorithm 1 Stochastic Smoothed-AGDA

Require: Initial pair (x_0, y_0) , integer $N \geq 1$, and step $0 < \alpha \leq 1/(3\ell)$.

- 1: Set $z_0 = x_0$ and $\beta = \alpha\ell/(76800\kappa)$.
 - 2: **for** $t = 0, \dots, N-1$ **do**
 - 3: Draw fresh independent samples ξ_t^x, ξ_t^y .
 - 4: $x_{t+1} = x_t - \alpha [G_x(x_t, y_t; \xi_t^x) + 2\ell(x_t - z_t)]$.
 - 5: $y_{t+1} = y_t + (\alpha/48)G_y(x_{t+1}, y_t; \xi_t^y)$.
 - 6: $z_{t+1} = z_t + \beta(x_{t+1} - z_t)$.
 - 7: **end for**
 - 8: Draw I uniformly from $\{0, \dots, N-1\}$, independently of the oracle samples.
 - 9: **return** $(\tilde{x}, \tilde{y}) = (x_I, y_I)$.
-

Smoothed-AGDA controls the squared game gradients, but directly bounding the squared outer gradient introduces an extra factor of κ through the inner-gradient error. The exact minimizer of the regularized outer objective avoids this loss, and stochastic updates approximate it. We apply the dual-first updates (E.7) to $f(x, y) + \ell \|x - \tilde{x}\|^2$, starting from $(\tilde{x}, \tilde{y})$. The guarantee below combines the mean-square estimates of Yang et al. (2022, Theorem 4.1, Proposition 2.1(b), and Corollary 4.1) with strong convexity in x .

Proposition E.1 (Outer stationarity for Smoothed-AGDA). *Suppose Assumptions 3.1–3.5 hold with $\kappa \geq 2$, and assume in addition that*

$$\Phi(x_0) - \inf_x \{f(x, y_0) + \ell \|x - x_0\|^2\} \leq c_{\text{init}} \Delta. \quad (\text{E.1})$$

For every $0 < \varepsilon^2 \leq \ell \Delta$, Smoothed-AGDA (Algorithm 1) followed by the dual-first conversion in (E.7) returns a point $\hat{x}$ with $\mathbb{E} \|\nabla \Phi(\hat{x})\|^2 \leq \varepsilon^2$, using at most

$$C \left[\frac{\kappa \ell \Delta}{\varepsilon^2} + \frac{\kappa^2 \ell \Delta \sigma^2}{\varepsilon^4} + \left(\kappa + \frac{\kappa^4 \sigma^2}{\varepsilon^2} \right) \log(2\kappa) \right] \quad (\text{E.2})$$

stochastic partial-gradient calls, where C depends at most on c_{init} . In particular, if $\varepsilon^2 \leq c \ell \Delta / (\kappa^2 \log(2\kappa))$ for a sufficiently small universal $c > 0$, this cost is

$$O\left(\frac{\kappa \ell \Delta}{\varepsilon^2} + \frac{\kappa^2 \ell \Delta \sigma^2}{\varepsilon^4}\right). \quad (\text{E.3})$$

Proof. All step sizes and iteration counts below depend only on the accuracy and problem parameters, not on the realized pair.

The iterate pair from Smoothed-AGDA. Initialize the auxiliary center of Yang et al. (2022, Algorithm 2) at $z_0 = x_0$. Their initial quantity b_0 is twice the left side of (E.1), so $b_0 \leq 2c_{\text{init}} \Delta$. Before tuning the primal step α , the proof of their Theorem 4.1, Eqs. (40)–(41), bounds the average squared game gradients by

$$\frac{1}{N} \sum_{t=0}^{N-1} \mathbb{E} \left[\|\nabla_x f(x_t, y_t)\|^2 + \kappa \|\nabla_y f(x_t, y_t)\|^2 \right] \leq C \frac{\kappa \Delta}{N \alpha} + C \kappa \ell \alpha \sigma^2, \quad 0 < \alpha \leq \frac{1}{3\ell}. \quad (\text{E.4})$$

Here the dual step is $\alpha/48$, the regularization coefficient is 2ℓ , and the center relaxation is $\alpha\ell/(76800\kappa)$, as in their proof. Their descent estimates use only the conditional means and variances of the two gradient calls; no smoothness of individual sampled gradients is required. They therefore apply to the oracle formulation in Assumptions 3.4 and 3.5.

Choose $\alpha = \min\{1/(3\ell), \sqrt{\Delta/(N\ell\sigma^2)}/2\}$, omitting the second term when $\sigma = 0$. With $N \geq C(\kappa\ell\Delta/\varepsilon^2 + \kappa^2\ell\Delta\sigma^2/\varepsilon^4)$, a uniformly selected pair $(\tilde{x}, \tilde{y})$ satisfies $\mathbb{E} e \leq \varepsilon^2/16$, where $e := \|\nabla_x f(\tilde{x}, \tilde{y})\|^2 + \kappa \|\nabla_y f(\tilde{x}, \tilde{y})\|^2$. Using the upper budget Δ in this choice, rather than the actual initial outer gap, also covers initializations with zero outer gap.

The exact proximal point. The gradient-bias bound in Lemma B.2 would only give $\|\nabla \Phi(\tilde{x})\|^2 \leq 2\kappa e$. Instead, condition on the selected pair and define the auxiliary game $\hat{f}(x, y) := f(x, y) + \ell \|x - \tilde{x}\|^2$. It is ℓ -strongly convex in x , retains inner PL with parameter ℓ/κ , and is 3ℓ -smooth in the partial-gradient convention of Assumption 3.1. Since $f(\cdot, y) + \ell \|\cdot\|^2/2$ is convex for every y , its supremum has the same property. Thus $\Phi(x) + \ell \|x - \tilde{x}\|^2$ is ℓ -strongly convex and has a unique minimizer $p := \arg \min_x \{\Phi(x) + \ell \|x - \tilde{x}\|^2\}$. Inner PL and strong convexity in x give

$$\begin{aligned} \Phi(\tilde{x}) - \min_x \hat{f}(x, \tilde{y}) &= \Phi(\tilde{x}) - f(\tilde{x}, \tilde{y}) + \hat{f}(\tilde{x}, \tilde{y}) - \min_x \hat{f}(x, \tilde{y}) \leq \frac{e}{2\ell}, \\ \frac{\ell}{2} \|p - \tilde{x}\|^2 &\leq \Phi(\tilde{x}) - \Phi(p) - \ell \|p - \tilde{x}\|^2 \leq \frac{e}{2\ell}. \end{aligned} \quad (\text{E.5})$$

The second line uses strong convexity of the auxiliary outer objective, then $\Phi(p) + \ell \|p - \tilde{x}\|^2 \geq \min_x \hat{f}(x, \tilde{y})$. Optimality of p now yields the pointwise bound

$$\nabla \Phi(p) = 2\ell(\tilde{x} - p), \quad \|\nabla \Phi(p)\|^2 \leq 4e. \quad (\text{E.6})$$

This is the squared-gradient estimate underlying Yang et al. (2022, Proposition 2.1(b), Eq. (7)); no conversion from a first-moment norm bound is used.

A finite stochastic conversion. Starting at $(x^0, y^0) = (\tilde{x}, \tilde{y})$, apply the dual-first alternating method used in that proposition to $\hat{f}$, with dual step τ and primal step 162τ . Explicitly, use fresh independent samples to update

$$\begin{aligned} y^{k+1} &= y^k + \tau G_y(x^k, y^k; \xi_k^y), \\ x^{k+1} &= x^k - 162\tau [G_x(x^k, y^{k+1}; \xi_k^x) + 2\ell(x^k - \tilde{x})], \quad 0 < \tau \leq \frac{1}{486\ell}. \end{aligned} \quad (\text{E.7})$$

These are an additional postprocessing procedure, not the primal-first recursion defining SGDA_{Alt} in Section 3.

We use the potential contraction of Yang et al. (2022, Theorem A.1 and the proof of Proposition 2.1(b)), then obtain the required distance bound directly from strong convexity. Write $\Psi(y) := \min_x \hat{f}(x, y)$ and $\Psi^* := \max_y \Psi(y)$. The auxiliary game has a saddle with outer coordinate p : any inner maximizer at p also minimizes $\hat{f}(\cdot, y)$ there, by Lemma B.2 and strong convexity. The potential combines the dual gap with the error in minimizing $\hat{f}(\cdot, y^k)$:

$$P_k := \mathbb{E}_{\text{conv}} \left[\Psi^* - \Psi(y^k) + \frac{1}{10} (\hat{f}(x^k, y^k) - \Psi(y^k)) \right],$$

where $\mathbb{E}_{\text{conv}}$ averages the fresh conversion samples conditional on $(\tilde{x}, \tilde{y})$. The initialization estimate and their contraction give

$$P_0 \leq \frac{11}{10} [\Phi(\tilde{x}) - \min_x \hat{f}(x, \tilde{y})] \leq \frac{11e}{20\ell}, \quad P_k \leq \left(1 - \frac{\ell\tau}{2\kappa}\right)^k P_0 + C\kappa\tau\sigma^2. \quad (\text{E.8})$$

To bound distance to p , let $x_y := \arg \min_x \hat{f}(x, y)$. Strong convexity at this minimizer and the saddle inequality $\hat{f}(p, y) \leq \Psi^*$ give

$$\|x - p\|^2 \leq 2\|x - x_y\|^2 + 2\|x_y - p\|^2 \leq \frac{4}{\ell} [\hat{f}(x, y) - \Psi(y) + \Psi^* - \Psi(y)]. \quad (\text{E.9})$$

Hence $\mathbb{E}_{\text{conv}} \|x^k - p\|^2 \leq 40P_k/\ell$. The step choices satisfy their conditions with smoothness 3ℓ and PL parameters ℓ and ℓ/κ . In particular, their variance term is $O(\kappa\tau\sigma^2)$. Substitution and expectation over the selected pair give

$$\mathbb{E} \|x^k - p\|^2 \leq \frac{C}{\ell^2} \left(1 - \frac{\ell\tau}{2\kappa}\right)^k \mathbb{E}e + \frac{C\kappa\tau\sigma^2}{\ell}. \quad (\text{E.10})$$

By Lemma B.2, the outer gradient is $2\kappa\ell$ -Lipschitz. To keep the conversion's contribution to the squared outer gradient below $\varepsilon^2/2$, it therefore suffices to make the squared distance at most $\varepsilon^2/(16\kappa^2\ell^2)$. Combining (E.6) and (E.10) gives

$$\begin{aligned} \mathbb{E} \|\nabla\Phi(x^k)\|^2 &\leq 8\mathbb{E}e + 8\kappa^2\ell^2\mathbb{E} \|x^k - p\|^2 \\ &\leq 8\mathbb{E}e + C\kappa^2 \left(1 - \frac{\ell\tau}{2\kappa}\right)^k \mathbb{E}e + C\kappa^3\ell\tau\sigma^2. \end{aligned} \quad (\text{E.11})$$

The last term determines the conversion step: for $\sigma > 0$, choose $\tau = \min\{1/(486\ell), c\varepsilon^2/(\kappa^3\ell\sigma^2)\}$ with a sufficiently small universal $c > 0$; use $\tau = 1/(486\ell)$ when $\sigma = 0$.

After

$$k = \left\lceil \frac{C\kappa}{\ell\tau} \log(2\kappa) \right\rceil = O\left(\left[\kappa + \frac{\kappa^4\sigma^2}{\varepsilon^2}\right] \log(2\kappa)\right)$$

updates, (E.10) meets this distance target. The factor κ^2 multiplying the contracting term in (E.11) accounts for the logarithm. This choice works in expectation over the random starting pair, without requiring its realized game gradients to satisfy a deterministic accuracy bound. With $\hat{x} = x^k$, the first line of (E.11) concludes

$$\mathbb{E} \|\nabla\Phi(\hat{x})\|^2 \leq 8\mathbb{E}e + \frac{\varepsilon^2}{2} \leq \varepsilon^2. \quad (\text{E.12})$$

There are two partial-gradient calls per update in both stages. Adding their costs proves (E.2). Under the stated small-error condition, the term $\kappa \log(2\kappa)$ is absorbed by $\kappa\ell\Delta/\varepsilon^2$, and $\kappa^4\sigma^2\varepsilon^{-2} \log(2\kappa)$ is absorbed by $\kappa^2\ell\Delta\sigma^2\varepsilon^{-4}$, giving (E.3). $\square$

E.1 INITIALIZATION BUDGET OF THE HARD INSTANCES

The additional budget (E.1) consists of the initial inner gap and the decrease obtainable by minimizing $f(x, y_0) + \ell \|x - x_0\|^2$ over x while keeping y_0 fixed. We verify that their sum is at most $c_{\text{init}}\Delta$ for every hard instance in Theorem 4.1, using the same declared budget Δ and its original initialization. The ordinary outer and inner gap checks remain in the construction proofs.

The unstable quadratic. For f_{unstable} in (5.2), Appendix C.1 chooses $x_0 = 4\sqrt{\Delta/(\ell + \mu)}$ and $y_0 = 0$. Thus $f_{\text{unstable}}(x, y_0) = 0$ for every x , and

$$\Phi_{\text{unstable}}(x_0) - \inf_x \{f_{\text{unstable}}(x, y_0) + \ell(x - x_0)^2\} = \Phi_{\text{unstable}}(x_0) = \Delta \leq c_{\text{init}}\Delta.$$

This verification includes the one-iterate case $T = 1$, for which the high-ratio proof uses this quadratic.

The corridor and noise components. The game f_{stable} in (5.4) is initialized at $(u_0, v_0; y_0) = (0, 0; 0)$. Equation (C.25) and the zero-component convention give $g_{\sigma,j}(v[j]; 0) = 0$ for every component, so $f_{\text{stable}}(u, v; 0) = g_{\text{corridor}}(u)$. Also $\Phi_{\text{stable}}(0, 0) = g_{\text{corridor}}(0)$, and (C.8) gives $g_{\text{corridor}}(0) - \inf_u g_{\text{corridor}}(u) = \Delta$. Therefore,

$$\begin{aligned} \Phi_{\text{stable}}(0, 0) - \inf_{u,v} \{f_{\text{stable}}(u, v; 0) + \ell(u^2 + \|v\|^2)\} \\ = g_{\text{corridor}}(0) - \inf_u \{g_{\text{corridor}}(u) + \ell u^2\} \leq \Delta \leq c_{\text{init}}\Delta. \end{aligned}$$

This also verifies the corridor alone, including $\sigma = 0$ and schedules with no positive noise-component steps, when all noise components vanish. Each noise component alone has regularized initial gap zero.

The low-ratio game. Retain the constants C, C_{low}, ζ and scales A, s_x, s_y from Appendix A.3, with f_{cycle} defined by (A.3) and initialized at $x_0 = s_x e_1, y_0^{\text{cycle}} = s_y y_0$. The construction in Appendix D.2 gives $\|\nabla_u \tilde{g}_{\text{cycle}}(e_1, y_0)\| \leq 1$, so $\|\nabla_x f_{\text{cycle}}(x_0, y_0^{\text{cycle}})\| \leq A/s_x$. Affinity in x , the gap bound (A.7), and the scales in (A.2) give

$$\begin{aligned} \Phi_{\text{cycle}}(x_0) - \inf_x \{f_{\text{cycle}}(x, y_0^{\text{cycle}}) + \ell \|x - x_0\|^2\} \\ = \Phi_{\text{cycle}}(x_0) - f_{\text{cycle}}(x_0, y_0^{\text{cycle}}) + \frac{\|\nabla_x f_{\text{cycle}}(x_0, y_0^{\text{cycle}})\|^2}{4\ell} \\ \leq \left(2C + \frac{1}{2C_{\text{low}}}\right) \zeta \Delta \leq 3C\zeta \Delta \leq c_{\text{init}}\Delta. \end{aligned}$$

The last inequalities use $C \geq 1, C_{\text{low}} \geq 2C$, and the choice $\zeta \leq c_{\text{init}}/(3C)$ already made in Appendix A.3. The check is uniform over the horizon and schedule, including $T = 1$. Both this game and the unstable quadratic use exact gradients, so their checks apply for every $\sigma \geq 0$.

Scope of the separation. All hard instances satisfy (E.1), so the SGDA lower bound persists on the subclass for which Proposition E.1 applies. Under its small-error condition, the complexity comparison uses the same declared gap budget Δ . SGDA returns a uniformly selected iterate, whereas $\hat{x}$ is computed by additional iterations.

F COMPLEXITY CALCULATIONS FOR TABLE 1

We derive the entries of Table 1, retaining initialization and additional source parameters until the final specialization. For the upper bounds, assume that the initial outer and inner gaps are $O(\Delta)$; the table fixes Δ . MSGDA and AdaMSGDA require every sampled gradient to be ℓ -Lipschitz, while SPIDER-GDA and SGDA-RR require ℓ -smooth finite-sum components. SGDA and Smoothed-AGDA require only population smoothness and bounded oracle variance. Expected-norm criteria use tolerance ε , and expected-squared-norm criteria use ε^2 . The complete complexities below specify the regimes for the MSGDA and AdaMSGDA specializations and distinguish HCMM-2's noise contribution from its full cost.

F.1 SGDA AND SMOOTHED-AGDA

Simultaneous and alternating SGDA. Under the setup of this paper, [Yang et al. \(2022, Theorem 3.1\)](#) for SGDA_{Alt} and [Theorem A.1](#) for SGDA_{Sim} give

$$\begin{aligned} \frac{1}{T} \sum_{t \leq T} \mathbb{E} \|\nabla \Phi(x_t)\|^2 &\leq C \left[\frac{\kappa^2 \ell \Delta}{T} + \kappa^2 \sigma \sqrt{\frac{\ell \Delta}{T}} \right], \\ T &= O \left(1 + \frac{\kappa^2 \ell \Delta}{\varepsilon^2} + \frac{\kappa^4 \ell \Delta \sigma^2}{\varepsilon^4} \right). \end{aligned} \quad (\text{F.1})$$

The second line makes each contribution in the first at most $\varepsilon^2/2$. Each iteration uses two fresh partial-gradient calls, so iteration and oracle complexities differ by a constant factor. Fixing Δ and taking the usual small-error regime $\varepsilon^2 \leq c\ell\Delta$ gives the two upper-bound entries. The matching lower bound in [Theorem 4.1](#) uses the same squared outer-gradient criterion and optimizes over fixed timescale ratios and predetermined nonincreasing step sizes.

Smoothed-AGDA. Proposition [E.1](#), derived from [Yang et al. \(2022, Theorem 4.1, Proposition 2.1\(b\), and Corollary 4.1\)](#), includes the conversion to a point $\hat{x}$ satisfying $\mathbb{E} \|\nabla \Phi(\hat{x})\|^2 \leq \varepsilon^2$. Under the additional regularized initialization condition ([E.1](#)), its total oracle cost is

$$O \left(\frac{\kappa \ell \Delta}{\varepsilon^2} + \frac{\kappa^2 \ell \Delta \sigma^2}{\varepsilon^4} + \left(\kappa + \frac{\kappa^4 \sigma^2}{\varepsilon^2} \right) \log(2\kappa) \right). \quad (\text{F.2})$$

If $\varepsilon^2 \leq c\ell\Delta/(\kappa^2 \log(2\kappa))$, then $\kappa^4 \sigma^2 \varepsilon^{-2} \log(2\kappa) \lesssim \kappa^2 \ell \Delta \sigma^2 \varepsilon^{-4}$ and $\kappa \log(2\kappa) \lesssim \kappa \ell \Delta \varepsilon^{-2}$. These absorptions give the table's $\tilde{O}(\kappa \ell \varepsilon^{-2} + \kappa^2 \ell \sigma^2 \varepsilon^{-4})$ entry with Δ fixed. The returned point is computed by the conversion in [Appendix E](#), rather than selected directly from the original trajectory.

F.2 FINITE-SUM UPPER BOUNDS

SPIDER-GDA. For $f = n^{-1} \sum_{i=1}^n f_i$ with ℓ -smooth components and attained inner μ -PL maximization, [Chen et al. \(2022, Theorem 6.1 and Lemma D.2\)](#) use $\kappa = \ell/\mu$ and initial potential $V_0 = \Phi(x_0) - \inf \Phi + (\Phi(x_0) - f(x_0, y_0))/24$. With primal step $\tau_x = \Theta((\kappa^2 \ell)^{-1})$, their proof gives a uniformly selected iterate $\hat{x}$ with

$$\begin{aligned} \mathbb{E} \|\nabla \Phi(\hat{x})\|^2 &\leq \frac{2V_0}{K\tau_x} \leq \frac{C\kappa^2 \ell V_0}{K}, \\ \text{component-gradient calls} &= O(n + K\sqrt{n}) = O \left(n + \frac{\sqrt{n} \kappa^2 \ell V_0}{\varepsilon^2} \right). \end{aligned} \quad (\text{F.3})$$

Here we choose $K = \max\{1, \lceil C\kappa^2 \ell V_0 / \varepsilon^2 \rceil\}$ updates, and the source chooses both the refresh period and recursive-estimator batch size of order $\sqrt{n}$. A full gradient costs n component calls, which also absorbs the rounding cost. The squared bound gives the source's stated expected-norm criterion by Jensen's inequality; $V_0 = O(\Delta)$ gives the table entry. No uniform component-variance bound is needed.

SGDA-RR. [Cho and Yun \(2023, Theorem 1\)](#) allow a partial-gradient component variance bound of the form $n^{-1} \sum_i \|\nabla_j f_i(z) - \nabla_j f(z)\|^2 \leq A \|\nabla_j f(z)\|^2 + B$, for $j \in \{x, y\}$. The table specializes to $A = 0$ and $B = \sigma^2$. With ratio $\rho \geq 14\kappa^2$, K epochs, and $V_0 = 4(\Phi(x_0) - \inf \Phi) + \Phi(x_0) - f(x_0, y_0)$, their bound gives

$$\begin{aligned} \frac{1}{K} \sum_{k=1}^K \mathbb{E} \|\nabla \Phi(x_0^k)\|^2 &\leq C \left[\frac{\rho \ell V_0}{K} + \rho \left(\frac{\ell^2 \sigma^2 V_0^2}{nK^2} \right)^{1/3} \right], \\ nK &= O \left(n + \frac{n\rho \ell V_0}{\varepsilon^2} + \frac{\sqrt{n} \rho^{3/2} \ell V_0 \sigma}{\varepsilon^3} \right). \end{aligned} \quad (\text{F.4})$$

The second line makes both error terms $O(\varepsilon^2)$ and counts n component calls per epoch, up to the constant factor for the two partial gradients; its first term covers rounding to a positive integer number of epochs. Substituting $\rho = \Theta(\kappa^2)$ and $V_0 = O(\Delta)$, then using $\varepsilon^2 \leq c\ell\Delta$ to absorb this rounding term, gives the table's two terms. The output is sampled from epoch starts. Component reuse and sampling without replacement differ from fresh-sample SGDA.

F.3 MSGDA AND ADAMSGDA

MSGDA. We specialize [Huang et al. \(2025, Theorem 1\)](#) to zero nonsmooth regularizer and unconstrained variables. Assume that each sampled partial gradient is unbiased with variance at most σ^2 , and that every sampled joint gradient is ℓ -Lipschitz. The source's full-gradient variance bound is then $2\sigma^2$, while its $L_f = \ell$ and $\kappa = \ell/\mu$. We use the valid outer-smoothness bound $L = \ell(1 + \kappa)$ from [Lemma B.2](#) in place of the smaller constant printed in the source. Each update draws one fresh sample and evaluates that same sampled function at two iterates, using a constant number of gradient calls.

The source's proof controls the primal estimator error, the inner gap, and the primal update direction. With w_t denoting its primal gradient estimator, the inner distance estimate in the proof of [Lemma B.2](#) gives

$$\|\nabla\Phi(x_t)\|^2 \leq 3\|w_t - \nabla_x f(x_t, y_t)\|^2 + 6\kappa\ell[\Phi(x_t) - f(x_t, y_t)] + 3\|w_t\|^2. \quad (\text{F.5})$$

Indeed, smoothness bounds the squared difference between the outer gradient and the game gradient by $2\kappa\ell$ times the inner gap. With no regularizer, the source's primal gradient mapping equals w_t . Thus its averaged squared estimates transfer directly to the squared outer-gradient criterion.

Write $D_x = \Phi(x_1) - \inf \Phi$, $D_y = \Phi(x_1) - f(x_1, y_1)$, and $D = D_x + D_y$. The accumulated estimate below bounds the initial gaps and initial estimator variance by a term proportional to $(\kappa\ell D + \kappa^2\sigma^2)/k$, and the accumulated estimator variance by a term proportional to $\kappa^4 k^2 \sigma^2$ up to logarithms. Balancing them for $\sigma > 0$ gives the following choice of the source's parameters:

$$\begin{aligned} k &= \left(\frac{\kappa\ell D + \kappa^2\sigma^2}{\kappa^4\sigma^2} \right)^{1/3}, & c_1 &= \frac{2}{3k^3} + 9\kappa^2, & c_2 &= \frac{2}{3k^3} + \frac{9}{4}, \\ m &= (c_1 k)^3, & \lambda &= \frac{\kappa}{8\ell}, & \gamma &= \frac{1}{256\kappa\ell}. \end{aligned} \quad (\text{F.6})$$

These choices satisfy the source proof's restrictions for every $\ell > 0$ and $\kappa \geq 1$: $k \geq \kappa^{-2/3}$, $9\kappa^2 \leq c_1 \leq (29/3)\kappa^2$, and $c_2 \leq c_1$ give all lower bounds on m . In particular, the restriction involving the corrected outer-smoothness constant holds because

$$\gamma = \frac{1}{256\kappa\ell} \leq \frac{1}{128\ell(1 + \kappa)} = \frac{\lambda\mu}{16L}.$$

The remaining primal and dual step restrictions are unchanged and are met by the displayed choices. The schedule is $\eta_t = k/(m + t)^{1/3}$, with momentum weights $c_1\eta_t^2$ and $c_2\eta_t^2$.

The source's coefficient H and the squared-gradient estimate obtained from its proof equations (59) and (61) satisfy

$$\begin{aligned} H &= \frac{D_x}{\gamma k} + \frac{9\kappa^2 D_y}{k\lambda} + \frac{4\sigma^2 m^{1/3}}{k^2} + 4k^2(c_1^2 + c_2^2)\sigma^2 \log(m + T) \\ &\leq C \left[\frac{\kappa\ell D + \kappa^2\sigma^2}{k} + \kappa^4 k^2 \sigma^2 \log(m + T) \right], \\ \frac{1}{T} \sum_{t=1}^T \mathbb{E} \|\nabla\Phi(x_t)\|^2 &\leq \frac{12H(m + T)^{1/3}}{T} \leq 12H \left(\frac{m^{1/3}}{T} + \frac{1}{T^{2/3}} \right). \end{aligned} \quad (\text{F.7})$$

The last line combines the source's averaged squared estimates with (F.5). Making its two contributions at most $\varepsilon^2/2$ requires $T = O(1 + Hm^{1/3}\varepsilon^{-2} + H^{3/2}\varepsilon^{-3})$. Substituting the balanced choice of k gives

$$\begin{aligned} Hm^{1/3} &= \tilde{O}(\kappa^3\ell D + \kappa^4\sigma^2), & H^{3/2} &= \tilde{O}(\kappa^3\ell D\sigma + \kappa^4\sigma^3), \\ T &= \tilde{O} \left(1 + \frac{\kappa^3\ell D + \kappa^4\sigma^2}{\varepsilon^2} + \frac{\kappa^3\ell D\sigma + \kappa^4\sigma^3}{\varepsilon^3} \right). \end{aligned} \quad (\text{F.8})$$

This is a feasible specialization derived from the cited theorem, rather than the source's accuracy-only statement. For $0 < \varepsilon \leq \sigma$ and fixed D , the ε^{-3} terms dominate the ε^{-2} terms, yielding the table's $\tilde{O}((\kappa^3\ell\sigma + \kappa^4\sigma^3)\varepsilon^{-3})$ entry. At $\sigma = 0$, instead take $k = \max\{\kappa^{-2/3}, (\ell D/(\kappa^3\varepsilon^2))^{1/3}\}$ with the remaining choices in (F.6); the same estimate gives $O(1 + \kappa^3\ell D\varepsilon^{-2})$.

AdaMSGDA. Theorem 2 of [Huang et al. \(2025\)](#) additionally assumes $A_t \succeq aI$ and $b_-I \preceq B_t \preceq b_+I$, where $a > 0$ and $0 < b_- \leq b_+$ are fixed spectral constants. To apply the source’s dual-update estimate, enlarge the upper bound to $\bar{b}_+ = \max\{b_+, b_-^{-1}\}$. Then $B_t \preceq \bar{b}_+I$ and $b_- \bar{b}_+ \geq 1$. For its dual gradient estimator v_t , the update $\tilde{y}_{t+1} = y_t + \lambda B_t^{-1} v_t$ satisfies

$$v_t^\top B_t^{-1} v_t \geq b_- \|B_t^{-1} v_t\|^2 \geq \frac{\|\tilde{y}_{t+1} - y_t\|^2}{\lambda^2 \bar{b}_+}.$$

This supplies the coefficient used in equation (66) of the source proof with $\bar{b}_+$ as the spectral upper bound.

Use the same k, c_2 as above, but take $c_1 = 2/(3k^3) + 9(\bar{b}_+/b_-)\kappa^2$ and $m = (c_1 k)^3$. For explicit admissible steps, set $d = 1/(8\bar{b}_+)$ and $g = \min\{d/(32\bar{b}_+), b_-/(16\bar{b}_+), a/8, a^2 d/(9\bar{b}_+)\}$, and choose $\lambda = d\kappa/\ell$ and $\gamma = g/(\kappa\ell)$. The outer-smoothness restriction again holds with $L = \ell(1 + \kappa)$, since $g \leq d/(32\bar{b}_+)$ gives

$$\gamma \leq \frac{d}{32\bar{b}_+\kappa\ell} \leq \frac{d}{16\bar{b}_+\ell(1 + \kappa)} = \frac{\lambda\mu}{16\bar{b}_+L}.$$

The other terms defining g satisfy the remaining primal step restrictions, while $m^{1/3}/k = c_1$ satisfies the schedule-dependent primal and dual restrictions. Its coefficient G obeys

$$\begin{aligned} G &= \frac{D_x}{\gamma k a} + \frac{9\bar{b}_+\kappa^2 D_y}{k\lambda a^2} + \frac{4\sigma^2 m^{1/3}}{k^2 a^2} + \frac{4k^2(c_1^2 + c_2^2)\sigma^2}{a^2} \log(m + T) \\ &\leq C_{a,b_-,b_+} \left[\frac{\kappa\ell D + \kappa^2\sigma^2}{k} + \kappa^4 k^2 \sigma^2 \log(m + T) \right]. \end{aligned} \quad (\text{F.9})$$

Equations (96)–(97) of the source proof give $T^{-1} \sum_{t=1}^T \mathbb{E} \|A_t^{-1} \nabla \Phi(x_t)\|^2 \leq 12G(m + T)^{1/3}/T$. Thus (F.8) and its table specialization hold for this adaptive gradient mapping, with constants depending on a, b_-, b_+ . A Euclidean outer-gradient guarantee additionally requires $A_t \preceq a_+I$: then $\|\nabla \Phi(x_t)\| \leq a_+ \|A_t^{-1} \nabla \Phi(x_t)\|$, so replace the mapping tolerance by ε/a_+ . The source’s lower bound on A_t alone does not supply this conversion. Both sampled-gradient methods therefore require stronger oracle information than the bounded-variance fresh-sample model of this paper.

F.4 HCMM-2

[Cai et al. \(2026, Theorem 2\)](#) control the average expected outer-gradient norm. Their assumptions require attained inner PL maximization, population-gradient smoothness ℓ , population-Hessian Lipschitzness M , and stochastic gradient and Hessian oracles based on fresh independent samples across iterations, with conditional unbiasedness and variances at most σ^2 and σ_h^2 , respectively. The parameters M and σ_h are not determined by ℓ, κ, σ . Initialization uses one stochastic gradient; each subsequent iteration uses a constant number of gradient and Hessian-vector operations.

Write $D_x = \Phi(x_0) - \inf \Phi$, $D_y = \Phi(x_0) - f(x_0, y_0)$, and $D = D_x + D_y$. For momentum weight $b \in (0, 1]$ and normalized step lengths $a_y = a > 0$ and $a_x = a/(6\kappa)$, the source’s conditions (27) hold. Its Appendix E, equation (134), gives

$$\begin{aligned} \frac{1}{T} \sum_{t \leq T} \mathbb{E} \|\nabla \Phi(x_t)\| &\leq C \left[\frac{\kappa D}{aT} + \frac{\kappa \sigma}{bT} + \frac{\kappa M a^2}{b} + \frac{\kappa \sigma_h a}{\sqrt{b}} \right. \\ &\quad \left. + \kappa \sigma \sqrt{b} + \kappa^2 \ell a + \frac{\sqrt{\kappa \ell D_y}}{T} \right]. \end{aligned} \quad (\text{F.10})$$

The last term uses the PL quadratic-growth bound $\text{dist}(y_0, \arg \max_y f(x_0, y)) \leq \sqrt{2\kappa D_y/\ell}$. We use the larger outer-gap coefficient obtained directly from the source’s descent inequality (46); the coefficient printed in (134) differs by a constant factor, which does not change these orders.

Choose $a = b = T^{-2/3}$. Substitution into (F.10), followed by making each contribution at most $\varepsilon/3$, yields

$$\begin{aligned} \frac{1}{T} \sum_{t < T} \mathbb{E} \|\nabla \Phi(x_t)\| &\leq C \left[\frac{\kappa(D + \sigma + \sigma_h)}{T^{1/3}} + \frac{\kappa M + \kappa^2 \ell}{T^{2/3}} + \frac{\sqrt{\kappa \ell D_y}}{T} \right], \\ T &= O \left(1 + \frac{\kappa^3(D^3 + \sigma^3 + \sigma_h^3)}{\varepsilon^3} + \frac{(\kappa M + \kappa^2 \ell)^{3/2}}{\varepsilon^{3/2}} + \frac{\sqrt{\kappa \ell D_y}}{\varepsilon} \right). \end{aligned} \quad (\text{F.11})$$

For a uniformly selected iterate, this is a sufficient complexity for expected norm at most ε , not expected squared norm at most ε^2 . Under $\sigma_h \leq \sigma$, its noise contribution at order ε^{-3} is $O(\kappa^3 \sigma^3 \varepsilon^{-3})$, giving the table's O_ℓ entry.

An additional sufficient regime in which this entry bounds the complete cost is $\sigma > 0$, $D = O(\sigma)$, $\sigma_h \leq \sigma$, and $0 < \varepsilon \leq c \min\{\sigma, \sigma^2/(\ell + M/\kappa)\}$. The second accuracy restriction absorbs the Hessian-regularity term, and the two restrictions together with $D_y \leq D$ absorb the terminal term. Hence (F.11) becomes $O(\kappa^3 \sigma^3 \varepsilon^{-3})$ throughout this regime. For general initialization, the full bound also contains $\kappa^3 D^3 \varepsilon^{-3}$; fixing the initial gaps does not make this term smaller in ε . The table then displays a noise contribution, while (F.11) retains the initialization and Hessian-regularity terms.

F.5 LOWER BOUNDS AND THE NONCONVERGENCE ENTRIES

NC-SC oracle lower bounds. For deterministic linear-span methods, Zhang et al. (2021, Theorem 3.1) prove $\Omega(\sqrt{\kappa} \ell \Delta \varepsilon^{-2})$ full-gradient calls to reach outer-gradient norm at most ε . Their finite-sum Theorem 3.2 gives $\Omega(n + \sqrt{n \kappa} \ell \Delta \varepsilon^{-2})$ expected component calls to reach an ε -stationary query for randomized linear-span methods. Its criterion is expected hitting time, as defined in their equations (4)–(6), rather than the expected gradient norm of a specified output. This result requires $\kappa \geq 2n$ and averaged component smoothness $n^{-1} \sum_i \|\nabla f_i(z) - \nabla f_i(z')\|^2 \leq \ell^2 \|z - z'\|^2$, which is weaker than smoothness of every component. Both rates apply in the stated small-error regime $\varepsilon^2 \leq c \ell \Delta$; fixing Δ gives the two table entries.

The first-order lower bound of Li et al. (2021, Theorem 2) is proved for zero-respecting algorithms on nonempty closed convex domains $\mathcal{X}, \mathcal{Y}$, with $0 \in \mathcal{X}$ and $\Phi(0) - \min_{x \in \mathcal{X}} \Phi(x) \leq \Delta$. Its projected mapping is $G_\Phi(x) := L_\Phi[x - P_{\mathcal{X}}(x - \nabla \Phi(x)/L_\Phi)]$, where $P_{\mathcal{X}}$ is Euclidean projection and $L_\Phi \leq (\kappa + 1)\ell$ is the source's value-function smoothness constant. The target is $\mathbb{E} \|G_\Phi(\hat{x})\| \leq \varepsilon$, and the lower bound is $\Omega(\sqrt{\kappa} \ell \Delta \varepsilon^{-2} + \kappa^{1/3} \ell \Delta \sigma^2 \varepsilon^{-4})$ calls to its bounded-variance stochastic first-order oracle. Fixing Δ gives the table entry; its criterion and algorithm class differ from the unconstrained SGDA recursion studied here.

NC-PŁ oracle lower bound. Pan and Li (2026, Theorem 3.2) prove $\Omega(\kappa \ell \Delta \varepsilon^{-2})$ exact first-order calls for arbitrary deterministic first-order methods, with only an initial outer-gap budget. The target is ordinary outer-gradient stationarity, with κ above a universal constant and $\varepsilon^2 \leq c \ell \Delta$. Fixing Δ gives the NC-PŁ first-order row.

Ratio obstructions and our matching complexity. Li et al. (2022, Theorem 4.1) construct an NC-SC quadratic on which the joint iterates of simultaneous GDA fail to converge locally for every positive constant step when $\rho \leq \kappa$. This implies the displayed local obstruction for $\rho = o(\kappa)$; it is not an averaged outer-gradient complexity bound.

For our update family, Theorems 4.1, 5.1 and 6.3 and the matching upper bounds give, respectively,

$$\begin{aligned} N_{\varepsilon, \text{Sim}}^*, N_{\varepsilon, \text{Alt}}^* &= \Theta \left(\frac{\kappa^2 \ell \Delta}{\varepsilon^2} + \frac{\kappa^4 \ell \Delta \sigma^2}{\varepsilon^4} \right), \\ N_{\varepsilon, \text{Sim}}(\rho), N_{\varepsilon, \text{Alt}}(\rho) &\geq c \left(\frac{\rho \ell \Delta}{\varepsilon^2} + \frac{\kappa^4 \ell \Delta \sigma^2}{\varepsilon^4} \right) \quad \text{if } \rho \geq c_{\text{high}} \kappa^2. \end{aligned} \quad (\text{F.12})$$

These complexity statements use $\varepsilon^2 \leq c \ell \Delta$, with κ above the respective theorem thresholds. Fixing Δ gives the optimized and high-ratio entries. For $1 \leq \rho \leq c_{\text{low}} \kappa^2$, Theorem 6.3 instead gives $\min_{t < T} \|\nabla \Phi(x_t)\|^2 \geq c \ell \Delta$ on a hard deterministic instance for every prescribed horizon and nonincreasing schedule. Thus the $\Omega(1)$ entry records a nonvanishing stationarity error, rather than an iteration complexity; the hard instance may depend on that horizon and schedule.