

Near-Linear Accuracy Bounds for Moreau–Yosida Unadjusted Langevin Sampling

Yuchen Xin^{*}, Zhihua Zhang[†]

Abstract

We establish near-linear accuracy bounds for the classical Moreau–Yosida unadjusted Langevin algorithm (MYULA). The target is $\pi \propto e^{-f-g}$, where $f \in C^2(\mathbb{R}^d)$ is m -strongly convex with Lipschitz gradient and g is convex and globally Lipschitz. Under an explicit parameter-dependent step-size condition, we bound the invariant-measure bias relative to the Moreau-smoothed target by $\tilde{O}(h)$, with only logarithmic dependence on the inverse smoothing parameter in the error coefficient. Combining this estimate with the Moreau approximation bias and Wasserstein contraction gives $\tilde{O}(\varepsilon^{-1})$ iterations to make the N th-iterate law μ_N satisfy $\sqrt{m} W_2(\mu_N, \pi) \leq \varepsilon$, for fixed model parameters and initialization. We bound the stationary error directly, without assuming third derivatives or a Lipschitz Hessian. Each iteration uses one gradient evaluation and one exact proximal evaluation. The key idea in our analysis is to convert a second-order stationary residual into a Wasserstein bound using a Poisson-based estimate.

Contents

1	Introduction	2
1.1	Contribution	3
1.2	Related Work	3
2	Preliminaries	4
2.1	Weak derivatives	5
2.2	Moreau regularization	5
2.3	Drift maps, pushforwards, and density evolution	5
2.4	Poisson equations and transport	7
2.5	Entropy and density-ratio moments	9
3	The Moreau–Yosida Unadjusted Langevin Sampling	9
3.1	Problem formulation and assumptions	9
3.2	Main results	10
4	Proof of the Main Results	11
4.1	The analytic estimates	11
4.2	The stationary residual	12
4.3	A residual-to-transport estimate	16
4.4	Proof of the fixed-smoothing theorem	18
4.5	Proof of the end-to-end complexity theorem	20

^{*}School of Mathematical Sciences, Peking University; email: 2301110087@pku.edu.cn

[†]School of Mathematical Sciences, Peking University; email: zhzhzhang@math.pku.edu.cn

5	Stationary Flux Estimates	20
5.1	The averaged drift flux	21
5.2	A finite heat expansion	21
5.3	Proof of the flux bounds	23
6	Conclusion	24
A	Target Moments and Normalized Operator Estimates	24
A.1	Target drift moments	24
A.2	A Gibbs change of variables for the heat operator	25
A.3	Gaussian derivatives without an extra dimension factor	26
A.4	Pushforward estimates by an integrated Jacobian	28
B	Finite-Order Stationary Density Ratios	29
B.1	Weighted energy for the density ratio	29
B.2	Gaussian integration by parts with the density-ratio weight	31
B.3	Proof of the density-ratio bound	33
C	Poisson Regularity and Negative Sobolev Norms	38
C.1	Existence and second-order energy	38
C.2	Approximation by compactly supported tests	39
C.3	Equivalent test classes for the negative Sobolev norm	40
D	Weak Regularity and Integrability Before Dissipation	41
D.1	Integration by parts and smooth approximation	41
D.2	Finiteness before the Rényi calculation	42
D.3	Removing the auxiliary mollification	50
	References	52

1 Introduction

We study a sampling problem from composite log-concave distributions

$$\pi(\mathrm{d}x) \propto e^{-f(x)-g(x)} \mathrm{d}x, \quad x \in \mathbb{R}^d,$$

where $f \in C^2(\mathbb{R}^d)$ is m -strongly convex with L_f -Lipschitz gradient, and g is convex and globally G -Lipschitz. The nonsmooth term can represent penalties such as ℓ_1 regularization. The Moreau–Yosida unadjusted Langevin algorithm (MYULA) of Durmus et al. [3] replaces g by its Moreau envelope g_λ and uses the update

$$\hat{X}_{k+1} = \hat{X}_k - h\nabla(f + g_\lambda)(\hat{X}_k) + \sqrt{2h}\xi_{k+1}, \quad \xi_{k+1} \stackrel{\text{i.i.d.}}{\sim} N(0, I_d).$$

The identity $\nabla g_\lambda = \lambda^{-1}(I - \text{prox}_{\lambda g})$ allows each iteration to use one evaluation of ∇f and one exact proximal evaluation of g , without an acceptance step.

The central issue is how smoothing affects discretization accuracy. Reducing λ improves the approximation of π by $\pi_\lambda \propto e^{-f-g_\lambda}$, but increases the global smoothness bound $L_\lambda = L_f + \lambda^{-1}$. Thus, a small-step bias estimate for a fixed λ does not by itself give a sharp complexity bound for the original target: its dependence on λ must also be controlled.

1.1 Contribution

In this work, we would address the above issue and present error bounds for MYULA. We denote

$$\tau_f := \sup_x \text{tr}[\nabla^2 f(x)], \quad R := \frac{G + \sqrt{G^2 + 4\tau_f}}{2},$$

and let $\ell_{h,\lambda} := 1 + \log(e + (mh)^{-1} + L_\lambda/m)$. Under the step-size condition (3.11), Theorem 3.4 proves that the MYULA invariant law satisfies

$$\sqrt{m} W_2(\hat{\pi}_{\lambda,h}, \pi_\lambda) \leq ChR^2 \ell_{h,\lambda}^2.$$

The error coefficient depends on λ^{-1} only through a logarithm, although the sufficient step-size restriction is stronger than $hL_\lambda \leq c$.

For $0 < \varepsilon \leq 1$, choose $\lambda = \varepsilon/G^2$ and the fixed step size in Theorem 3.5. Combining the invariant-measure estimate with the Moreau bias bound of Xin and Zhang [8, Proposition 5.9] gives the sufficient iteration complexity

$$N = \tilde{O} \left[\frac{L_f \tau_f}{m^2} + \frac{L_f R}{m^{3/2}} + \frac{1}{\varepsilon} \left(\frac{R^2}{m} + \frac{G^3 R}{m^2} \right) \right]$$

for $\sqrt{m} W_2(\mu_N, \pi) \leq \varepsilon$, where $\mu_N = \mathcal{L}(\hat{X}_N)$ and $\tilde{O}$ suppresses logarithmic factors. For fixed model parameters and initialization, this improves the $\tilde{O}(\varepsilon^{-4/3})$ guarantee of Xin and Zhang [9] to $\tilde{O}(\varepsilon^{-1})$. For fixed L_f/m and $G/\sqrt{m}$, the bound also gives $\tilde{O}(d/\varepsilon)$ iterations.

Proof strategy. The core strategy of our analysis is a Poisson-based estimate that converts a second-order stationary residual into a Wasserstein bound (Section 4.3). By testing the residual against a Poisson solution, we place derivatives on the test function rather than on the residual fields. Weighted energy estimates control the resulting terms while retaining the Hessian as a weight, instead of replacing it by its global bound. In particular, for MYULA, exact drift–heat identities express the residual through vector and matrix fluxes (Section 4.2). Finite-order density-ratio estimates and short-time heat estimates control these fluxes (Section 5), allowing the curvature contribution to be bounded through its average under π_λ . Together, these steps yield a near-linear invariant-measure bias with only logarithmic dependence on λ^{-1} in its coefficient, without additional higher-order smoothness assumptions (Section 4.4). Adding the smoothing and convergence errors gives the complexity bound in Section 4.5.

1.2 Related Work

MYULA and averaged curvature. Durmus et al. [3] introduced MYULA and established asymptotic and nonasymptotic convergence guarantees for nonsmooth log-concave targets. More recently, Xin and Zhang [8] analyzed the weak Moreau Hessian through its average along a reference heat path. For the structured penalties treated there, curvature localization yields $\tilde{O}(\varepsilon^{-2})$ end-to-end complexity. The closest result is Xin and Zhang [9], who consider the same assumptions and MYULA transition as this paper. Their discrete Poisson-corrector analysis gives an invariant-measure bias of $O(h) + \tilde{O}(h^{3/4})$, with only logarithmic smoothing dependence in its coefficients, and consequently $\tilde{O}(\varepsilon^{-4/3})$ complexity. Our stationary-density analysis replaces this bias bound by $\tilde{O}(h)$ under a stronger step-size restriction. The improvement concerns the fixed-parameter accuracy exponent; it does not assert uniformly better dependence on every model parameter.

First-order ULA bounds and the smoothing parameter. For a smooth, m -strongly convex potential with L -Lipschitz gradient, Pedrotti and Whalley [5, Theorem 1] prove

$$\sqrt{m} W_2(\hat{\pi}_h, \pi) \leq 6hL\sqrt{d}, \quad 0 < hL \leq 1.$$

Applying this estimate to $f + g_\lambda$, with the $C^{1,1}$ case obtained by mollification, and choosing $\lambda \asymp \varepsilon/G^2$ gives the sufficient MYULA complexity

$$\tilde{O}\left[\frac{\sqrt{d}}{m} \left(\frac{L_f}{\varepsilon} + \frac{G^2}{\varepsilon^2}\right)\right].$$

Thus, first-order bias for a fixed smooth target does not directly imply near-linear accuracy complexity after Moreau smoothing. The gain is in the joint smoothing–discretization analysis, not in establishing first-order ULA bias at fixed smoothness.

Other proximal samplers. Different transitions can achieve higher accuracy without a fixed Moreau approximation. For example, Fan et al. [4, Proposition 5] obtain polylogarithmic accuracy dependence in W_2 for strongly convex composite targets, under their initialization and subroutine assumptions. Their method uses a restricted Gaussian sampling step implemented by approximate rejection sampling, rather than a deterministic proximal update. These guarantees concern a different algorithm and computational model. Our result is an iteration bound for classical MYULA with one exact proximal evaluation per step.

2 Preliminaries

All measures are defined on $\mathbb{R}^d$, and $\mathcal{P}_2(\mathbb{R}^d)$ denotes the probability measures with finite second moment. For $\mu, \nu \in \mathcal{P}_2(\mathbb{R}^d)$,

$$W_2^2(\mu, \nu) := \inf_{\mathcal{L}(X)=\mu, \mathcal{L}(Y)=\nu} \mathbb{E} \|X - Y\|^2. \quad (2.1)$$

The same symbol denotes a measure and its Lebesgue density when a density exists. We use the Euclidean norm and inner product for vectors, the Frobenius norm $\|M\|_F^2 = \sum_{i,j} M_{ij}^2$ for matrices, and $\langle M, N \rangle = \text{tr}(M^\top N) = \sum_{i,j} M_{ij} N_{ij}$. For symmetric matrices, $M \preceq N$ means $v^\top M v \leq v^\top N v$ for every v . For a scalar, vector, or matrix field F , define

$$\|F\|_{L^s(\nu)} := \left(\int \|F(x)\|^s \nu(dx) \right)^{1/s}, \quad 1 \leq s < \infty. \quad (2.2)$$

The pointwise norm in (2.2) is the absolute value for scalars, the Euclidean norm for vectors, and the Frobenius norm for matrices. Thus, for a matrix field F ,

$$\|F\|_{L^2(\nu)}^2 = \int \|F(x)\|_F^2 \nu(dx) = \sum_{i,j} \int |F_{ij}(x)|^2 \nu(dx).$$

In particular, for a positive density ν and a scalar, vector, or matrix density F ,

$$\|F/\nu\|_{L^s(\nu)}^s = \int \|F(x)\|^s \nu(x)^{1-s} dx.$$

For $v, w \in \mathbb{R}^d$, the outer product is the matrix

$$(v \otimes w)_{ij} := v_i w_j, \quad \|v \otimes w\|_F = \|v\| \|w\|. \quad (2.3)$$

2.1 Weak derivatives

For a locally integrable function u , its distributional derivative is defined by

$$\langle \partial_i u, \phi \rangle = - \int u \partial_i \phi \, dx, \quad \phi \in C_c^\infty(\mathbb{R}^d),$$

where C_c^∞ denotes smooth functions with compact support. If this derivative is represented by a locally integrable function v , then v is the weak derivative of u . Lipschitz functions have almost-everywhere derivatives, meaning their weak derivatives. An equality of distributions implies an equality against every such test function. When both sides are locally integrable functions, this is equivalent to equality almost everywhere; it does not assert the existence of classical derivatives at every point.

For a vector field H and a matrix field E , our divergence convention is

$$\operatorname{div} H = \sum_i \partial_i H_i, \quad (\operatorname{div} E)_i = \sum_j \partial_j E_{ij}, \quad \operatorname{div} \operatorname{div} E = \sum_{i,j} \partial_i \partial_j E_{ij}. \quad (2.4)$$

In particular,

$$\langle \operatorname{div} H, \phi \rangle = - \int \langle \nabla \phi, H \rangle \, dx, \quad \langle \operatorname{div} \operatorname{div} E, \phi \rangle = \int \langle E, \nabla^2 \phi \rangle \, dx. \quad (2.5)$$

These formulas require only local integrability of the fields. They allow density equations to be used through integration by parts without differentiating the densities classically. Hessians are written explicitly as $\nabla^2 f$, $\nabla^2 g_\lambda$, or $\nabla^2 U_\lambda$; the letter H denotes a vector flux. Constants $c, C > 0$ are universal and may change from line to line. The notation $\hat{O}$ suppresses fixed powers of logarithms, but no polynomial parameter factors.

2.2 Moreau regularization

For a convex, globally G -Lipschitz function $g : \mathbb{R}^d \rightarrow \mathbb{R}$ and $\lambda > 0$, let

$$g_\lambda(x) := \inf_{y \in \mathbb{R}^d} \left\{ g(y) + \frac{\|x - y\|^2}{2\lambda} \right\}. \quad (2.6)$$

The unique minimizer is denoted by $\operatorname{prox}_{\lambda g}(x)$. The Moreau properties used below are

$$\nabla g_\lambda(x) = \lambda^{-1}(x - \operatorname{prox}_{\lambda g}(x)), \quad \|\nabla g_\lambda(x)\| \leq G, \quad 0 \preceq \nabla^2 g_\lambda \preceq \lambda^{-1}I \quad \text{a.e.} \quad (2.7)$$

In particular, $g_\lambda \in C^{1,1}$, meaning that its gradient is globally Lipschitz. The last inequality concerns its weak Hessian, not a continuous second derivative. We use these facts in the form recalled in Xin and Zhang [9, Section 3.2 and Appendix A].

2.3 Drift maps, pushforwards, and density evolution

The potential used for MYULA will be specified in [Section 3.1](#). For the remainder of this section, we use only its regularity and convexity properties: fix $\lambda > 0$ and let $U_\lambda \in C^{1,1}(\mathbb{R}^d)$ be m -strongly convex with L_λ -Lipschitz gradient, where $0 < m \leq L_\lambda < \infty$. Write π_λ for the probability density proportional to e^{-U_λ} . Its weak Hessian satisfies

$$mI \preceq \nabla^2 U_\lambda \preceq L_\lambda I \quad \text{almost everywhere.} \quad (2.8)$$

The deterministic drift step is

$$T_t(x) := x - t\nabla U_\lambda(x). \quad (2.9)$$

For an integrable scalar, vector, or matrix density F , the pushforward of $F dx$ assigns to each Borel set the integral of F over its inverse image under T_t . This defines a unique weighted measure, componentwise. If it has a Lebesgue density, we denote that density by $(T_t)_\# F$; equivalently,

$$\int \phi(y)(T_t)_\# F(y) dy = \int \phi(T_t x) F(x) dx \quad \text{for every } \phi \in C_c^\infty(\mathbb{R}^d). \quad (2.10)$$

For a probability density, the pushforward is the law of the transformed random variable $T_t X$. For vector and matrix densities, the pushforward is defined componentwise.

Proposition 2.1 (Drift pushforward). *If $0 \leq tL_\lambda < 1$, then T_t is a global bi-Lipschitz bijection and*

$$\text{Lip}(T_t^{-1}) \leq (1 - tL_\lambda)^{-1}.$$

For every integrable scalar, vector, or matrix density F , its pushforward has the density

$$(T_t)_\# F(y) = \frac{F(T_t^{-1}y)}{\det(I - t\nabla^2 U_\lambda(T_t^{-1}y))} \quad \text{for a.e. } y. \quad (2.11)$$

In particular, $\|(T_t)_\# F\|_{L^1(\mathbb{R}^d)} = \|F\|_{L^1(\mathbb{R}^d)}$.

Proof. For each y , the map $x \mapsto y + t\nabla U_\lambda(x)$ is a contraction on $\mathbb{R}^d$. Its unique fixed point is $T_t^{-1}y$. The fixed-point equations for two values of y give the stated inverse Lipschitz bound, and T_t is Lipschitz by definition. Its derivative is $DT_t = I - t\nabla^2 U_\lambda$ almost everywhere, with

$$\det(I - t\nabla^2 U_\lambda) \geq (1 - tL_\lambda)^d > 0.$$

The weighted area formula [7, Chapter 2, Theorem 3.3 and Remark 3.4(2)] applied to the Lipschitz bijection T_t gives

$$\int q(x) \det(I - t\nabla^2 U_\lambda(x)) dx = \int q(T_t^{-1}y) dy$$

for every nonnegative measurable scalar function q . Applying this formula with the integrand divided by the positive Jacobian, and then taking positive and negative parts componentwise, yields

$$\int \phi(T_t x) F(x) dx = \int \phi(y) \frac{F(T_t^{-1}y)}{\det(I - t\nabla^2 U_\lambda(T_t^{-1}y))} dy.$$

Both T_t and T_t^{-1} preserve Lebesgue null sets because they are Lipschitz, so the almost-everywhere derivatives suffice for these formulas. The last identity proves (2.11). Changing variables in its pointwise norm gives the stated L^1 identity. $\square$

The free heat semigroup is

$$\mathbf{H}_t F(x) := (4\pi t)^{-d/2} \int e^{-\|x-y\|^2/(4t)} F(y) dy, \quad t > 0, \quad \mathbf{H}_0 F = F. \quad (2.12)$$

It acts componentwise and adds an independent $N(0, 2tI_d)$ increment to a probability density. For $F \in L^1(\mathbb{R}^d)$, it preserves the integral and satisfies $\|\mathbf{H}_t F\|_{L^1} \leq \|F\|_{L^1}$. Moreover, $\mathbf{H}_t F \rightarrow F$ in L^1 as $t \downarrow 0$, since the Gaussian kernels form an approximate identity.

We consider a gradient bound and an integrated form of the heat equation. For scalar $F \in L^1(\mathbb{R}^d)$, differentiation of the Gaussian kernel, the triangle inequality, and Tonelli's theorem give

$$\begin{aligned} \|\nabla \mathbf{H}_t F\|_{L^1(\mathbb{R}^d)} &\leq \|F\|_{L^1(\mathbb{R}^d)} \int_{\mathbb{R}^d} \frac{\|z\|}{2t} (4\pi t)^{-d/2} e^{-\|z\|^2/(4t)} dz \\ &\leq \frac{C_d}{\sqrt{t}} \|F\|_{L^1(\mathbb{R}^d)}, \quad C_d := \sqrt{d/2}. \end{aligned} \quad (2.13)$$

The second inequality is Cauchy-Schwarz and the second moment $2td$ of this Gaussian density. In particular, $t \|\nabla \mathbf{H}_t F\|_{L^1}$ is integrable near $t = 0$.

For $\phi \in C_c^\infty(\mathbb{R}^d)$, symmetry of the Gaussian kernel gives

$$\int \phi \mathbf{H}_t F dx = \int F \mathbf{H}_t \phi dx.$$

For $t > 0$, differentiation of the kernel and integration by parts against the smooth function ϕ yield $\partial_t \mathbf{H}_t \phi = \mathbf{H}_t \Delta \phi$. Since

$$|F(x) \partial_t \mathbf{H}_t \phi(x)| \leq |F(x)| \|\Delta \phi\|_\infty,$$

differentiation under the integral is justified by dominated convergence. Hence

$$\frac{d}{dt} \int \phi \mathbf{H}_t F dx = \int (\Delta \phi) \mathbf{H}_t F dx, \quad t > 0. \quad (2.14)$$

The derivative is bounded by $\|F\|_{L^1} \|\Delta \phi\|_\infty$. Integrating first over a positive-time interval and then using $\mathbf{H}_t F \rightarrow F$ in L^1 at the left endpoint gives

$$\int \phi (\mathbf{H}_b F - \mathbf{H}_a F) dx = \int_a^b \int (\Delta \phi) \mathbf{H}_t F dx dt, \quad 0 \leq a \leq b < \infty. \quad (2.15)$$

Thus, for $F \in L^1(\mathbb{R}^d)$ and $\phi \in C_c^\infty(\mathbb{R}^d)$, the map $t \mapsto \int \phi \mathbf{H}_t F dx$ is absolutely continuous on $[0, T]$ for every $T > 0$.

The Langevin generator and its formal adjoint with respect to Lebesgue measure are defined by

$$\mathcal{L}_\lambda \phi = \Delta \phi - \langle \nabla U_\lambda, \nabla \phi \rangle, \quad \mathcal{L}_\lambda^* \rho = \Delta \rho + \operatorname{div}(\rho \nabla U_\lambda). \quad (2.16)$$

Here $\mathcal{L}_\lambda^* \rho$ is understood distributionally: $\langle \mathcal{L}_\lambda^* \rho, \phi \rangle = \int \rho \mathcal{L}_\lambda \phi dx$ for $\phi \in C_c^\infty(\mathbb{R}^d)$. Because $\nabla \pi_\lambda = -\pi_\lambda \nabla U_\lambda$, we have $\mathcal{L}_\lambda^* \pi_\lambda = 0$.

2.4 Poisson equations and transport

The Poisson equation will be used to convert a residual in the stationary density equation into a difference of expectations. The weighted Sobolev space $H^1(\pi_\lambda)$ consists of functions whose value and weak gradient belong to $L^2(\pi_\lambda)$. Strong convexity gives the Poincaré inequality

$$\int \left| v - \int v d\pi_\lambda \right|^2 d\pi_\lambda \leq \frac{1}{m} \int \|\nabla v\|^2 d\pi_\lambda; \quad (2.17)$$

see, for example, Chewi [1]. For $\phi \in L^2(\pi_\lambda)$, the mean-zero Poisson equation is

$$\mathcal{L}_\lambda \psi = \phi - \int \phi d\pi_\lambda, \quad \int \psi d\pi_\lambda = 0. \quad (2.18)$$

A weak solution is a function $\psi \in H^1(\pi_\lambda)$ satisfying

$$\int \langle \nabla \psi, \nabla v \rangle d\pi_\lambda = - \int \left(\phi - \int \phi d\pi_\lambda \right) v d\pi_\lambda, \quad v \in H^1(\pi_\lambda). \quad (2.19)$$

Here ϕ is prescribed and ψ is the unknown function; the zero-mean condition removes the freedom to add a constant. For $k = 1, 2$, the unweighted Sobolev space $H^k(\mathbb{R}^d)$ consists of functions whose weak derivatives of order at most k belong to $L^2(\mathbb{R}^d)$ with respect to Lebesgue measure. We write $u \in H_{\text{loc}}^2(\mathbb{R}^d)$ if $\eta u \in H^2(\mathbb{R}^d)$ for every $\eta \in C_c^\infty(\mathbb{R}^d)$. The next lemma shows the regularity and approximation needed below.

Lemma 2.2 (Poisson regularity and approximation). *Let $U_\lambda \in C^{1,1}(\mathbb{R}^d)$ be m -strongly convex with L_λ -Lipschitz gradient, where $0 < m \leq L_\lambda < \infty$. Let $\pi_\lambda \propto e^{-U_\lambda}$ and let $\mathcal{L}_\lambda$ be the generator in (2.16). For every $\phi \in L^2(\pi_\lambda)$, the Poisson equation (2.18) has a unique mean-zero weak solution $\psi \in H^1(\pi_\lambda)$. Moreover, $\psi \in H_{\text{loc}}^2(\mathbb{R}^d)$. If $\phi \in C_c^\infty(\mathbb{R}^d)$, then $\nabla^2 \psi \in L^2(\pi_\lambda)$ and*

$$\begin{aligned} & \int \|\nabla^2 \psi\|_F^2 d\pi_\lambda + \int \nabla \psi^\top (\nabla^2 U_\lambda) \nabla \psi d\pi_\lambda \\ & \leq \int \left| \phi - \int \phi d\pi_\lambda \right|^2 d\pi_\lambda \leq \frac{1}{m} \int \|\nabla \phi\|^2 d\pi_\lambda. \end{aligned} \quad (2.20)$$

For such ϕ , there exist $\psi_n \in C_c^\infty(\mathbb{R}^d)$ such that

$$\|\mathcal{L}_\lambda \psi_n - \mathcal{L}_\lambda \psi\|_{L^2(\pi_\lambda)} + \|\nabla^2 \psi_n - \nabla^2 \psi\|_{L^2(\pi_\lambda)} + \|\nabla \psi_n - \nabla \psi\|_{L^2(\pi_\lambda)} \longrightarrow 0. \quad (2.21)$$

The proof is given in [Appendix C](#). The estimate (2.20) controls the weak derivatives of the Poisson solution, whereas (2.21) allows it to be used in residual identities initially stated only for compactly supported smooth tests. Neither assertion requires a continuous Hessian or a third derivative of U_λ .

For an integrable signed density σ with $\int \sigma dx = 0$, define

$$\|\sigma\|_{\dot{H}^{-1}(\pi_\lambda)} := \sup_{\substack{\phi \in C_c^\infty(\mathbb{R}^d) \\ \|\nabla \phi\|_{L^2(\pi_\lambda)} \leq 1}} \left| \int \phi \sigma dx \right|. \quad (2.22)$$

Here σ represents the signed measure with Lebesgue density σ , not a density relative to π_λ . In particular, the function–measure pairing is

$$\langle \phi, \sigma(x) dx \rangle = \int \phi \sigma dx = \int \phi \frac{\sigma}{\pi_\lambda} d\pi_\lambda. \quad (2.23)$$

If $\sigma/\pi_\lambda \in L^2(\pi_\lambda)$, the same norm is obtained using either weighted Sobolev tests or C^1 tests:

$$\|\sigma\|_{\dot{H}^{-1}(\pi_\lambda)} = \sup_{\substack{\phi \in H^1(\pi_\lambda) \\ \|\nabla \phi\|_{L^2(\pi_\lambda)} \leq 1}} \left| \int \phi \sigma dx \right| = \sup_{\substack{\phi \in C^1(\mathbb{R}^d) \\ \|\nabla \phi\|_{L^2(\pi_\lambda)} \leq 1}} \left| \int \phi \sigma dx \right|. \quad (2.24)$$

The approximation and integrability needed for these equalities are proved in [Section C.3](#). Consequently, (2.22) agrees, in this setting, with the norm of the signed measure $\sigma(x) dx$ in Peyre [6, Section 1.1].

For a probability density $\nu \in \mathcal{P}_2(\mathbb{R}^d)$ with $\nu/\pi_\lambda \in L^2(\pi_\lambda)$, apply Peyre [6, Theorem 1] to the measures $\pi_\lambda(x) dx$ and $\nu(x) dx$. By (2.23) and (2.24), this gives

$$W_2(\nu, \pi_\lambda) \leq 2 \|\nu - \pi_\lambda\|_{\dot{H}^{-1}(\pi_\lambda)}. \quad (2.25)$$

2.5 Entropy and density-ratio moments

For $F \geq 0$, write

$$\text{Ent}_\nu F := \int F \log F \, d\nu - \left(\int F \, d\nu \right) \log \int F \, d\nu.$$

Strong convexity also gives the log-Sobolev inequality [1]

$$\text{Ent}_{\pi_\lambda}(w^2) \leq \frac{2}{m} \int \|\nabla w\|^2 \, d\pi_\lambda. \quad (2.26)$$

For $\alpha > 1$ and a probability density ν , the Rényi divergence is

$$\mathcal{R}_\alpha(\nu \| \pi_\lambda) := \frac{1}{\alpha - 1} \log \int \left(\frac{\nu}{\pi_\lambda} \right)^\alpha \, d\pi_\lambda. \quad (2.27)$$

Equivalently, $\|\nu/\pi_\lambda\|_{L^\alpha(\pi_\lambda)} = \exp\{(\alpha - 1)\mathcal{R}_\alpha(\nu \| \pi_\lambda)/\alpha\}$.

3 The Moreau–Yosida Unadjusted Langevin Sampling

In this section we study theoretical analysis for error bounds of the Moreau–Yosida unadjusted Langevin algorithm (MYULA). We first give some assumptions with the problem description, and then present our main theoretical results.

3.1 Problem formulation and assumptions

We consider the probability distribution

$$\pi(dx) = Z^{-1} \exp\{-f(x) - g(x)\} \, dx, \quad x \in \mathbb{R}^d. \quad (3.1)$$

The assumptions and normalization below follow those of Xin and Zhang [9].

Assumption 3.1 (Smooth component). *The function $f \in C^2(\mathbb{R}^d)$ satisfies*

$$mI \preceq \nabla^2 f(x) \preceq L_f I, \quad x \in \mathbb{R}^d, \quad 0 < m \leq L_f < \infty. \quad (3.2)$$

Write

$$\tau_f := \sup_{x \in \mathbb{R}^d} \text{tr}[\nabla^2 f(x)]. \quad (3.3)$$

In particular, $md \leq \tau_f \leq dL_f$.

Assumption 3.2 (Nonsmooth component). *The function $g: \mathbb{R}^d \rightarrow \mathbb{R}$ is convex and globally G -Lipschitz, for some $G > 0$.*

For the Moreau envelope in (2.6), set

$$U_\lambda := f + g_\lambda, \quad \pi_\lambda(dx) := Z_\lambda^{-1} e^{-U_\lambda(x)} \, dx, \quad L_\lambda := L_f + \lambda^{-1}. \quad (3.4)$$

By Assumptions 3.1 and 3.2 and (2.7), this choice of U_λ belongs to $C^{1,1}(\mathbb{R}^d)$ and satisfies (2.8). Thus the operator constructions and Poisson results in Sections 2.3 and 2.4 apply.

MYULA is the fixed-step recursion

$$\hat{X}_{k+1} = \hat{X}_k - h \nabla U_\lambda(\hat{X}_k) + \sqrt{2h} \xi_{k+1}, \quad \xi_{k+1} \stackrel{\text{i.i.d.}}{\sim} N(0, I_d). \quad (3.5)$$

Its transition kernel is $Q_{\lambda,h}$, and $\mu_k := \mathcal{L}(\widehat{X}_k)$. Under the step-size condition below, its invariant law is denoted by $\widehat{\pi}_{\lambda,h}$. The initial law μ_0 belongs to $\mathcal{P}_2(\mathbb{R}^d)$. We measure accuracy by $\sqrt{m} W_2(\mu_k, \pi)$. Each iteration uses one evaluation of ∇f and one exact proximal evaluation of g .

Let

$$R := \frac{G + \sqrt{G^2 + 4\tau_f}}{2}. \quad (3.6)$$

Thus, $R^2 = \tau_f + GR$, $R \geq G$, and $R^2 \geq \tau_f \geq md$.

Proposition 3.3 (Invariant law and contraction). *Under [Assumptions 3.1](#) and [3.2](#), if $0 < hL_\lambda < 1$, then $Q_{\lambda,h}$ has a unique invariant law in $\mathcal{P}_2(\mathbb{R}^d)$, with a smooth strictly positive density $\widehat{\pi}_{\lambda,h}$. For $\mu, \nu \in \mathcal{P}_2(\mathbb{R}^d)$,*

$$W_2(\mu Q_{\lambda,h}, \nu Q_{\lambda,h}) \leq (1 - mh) W_2(\mu, \nu). \quad (3.7)$$

Moreover,

$$\widehat{\pi}_{\lambda,h} = H_h((T_h)_\# \widehat{\pi}_{\lambda,h}). \quad (3.8)$$

Proof. For a smooth potential, integration of the Hessian along a segment gives $\|T_h x - T_h y\| \leq (1 - mh) \|x - y\|$. Mollification preserves the Hessian bounds and gives the same inequality for $U_\lambda \in C^{1,1}$ by uniform convergence of its gradient. Using the same Gaussian increment for two Euler updates proves (3.7). The linear growth of the drift implies that the kernel maps $\mathcal{P}_2$ into itself. The contraction theorem on the complete metric space $(\mathcal{P}_2, W_2)$ therefore gives the unique invariant law in that space. An Euler update consists of the pushforward by T_h followed by the heat operator H_h , which proves (3.8). Convolution of a probability measure with the Gaussian kernel has a smooth strictly positive density. $\square$

3.2 Main results

The first theorem controls the invariant-measure bias at fixed smoothing; the second gives a single step size for the original target. Set

$$\ell_{h,\lambda} := 1 + \log \left(e + \frac{1}{mh} + \frac{L_\lambda}{m} \right), \quad (3.9)$$

$$\ell_\varepsilon := 1 + \log \left(e + \frac{L_f}{m} + \frac{\tau_f}{m} + \frac{G^2}{m} + \frac{1}{\varepsilon} \right). \quad (3.10)$$

Theorem 3.4 (Invariant-measure bias at fixed smoothing). *Under [Assumptions 3.1](#) and [3.2](#), there exist universal constants $c, C > 0$ such that, for every $\lambda, h > 0$ satisfying*

$$h\ell_{h,\lambda}^2 \left[\frac{L_f \tau_f}{m} + \frac{L_f R}{\sqrt{m}} + \frac{GR}{m\lambda} + R^2 + \frac{1}{\lambda} \right] \leq c, \quad (3.11)$$

the kernel $Q_{\lambda,h}$ has a unique invariant law $\widehat{\pi}_{\lambda,h} \in \mathcal{P}_2(\mathbb{R}^d)$, and

$$\sqrt{m} W_2(\widehat{\pi}_{\lambda,h}, \pi_\lambda) \leq ChR^2 \ell_{h,\lambda}^2. \quad (3.12)$$

The proof of [Theorem 3.4](#) is given in [Section 4.4](#). The sufficient restriction (3.11) is stronger than $hL_\lambda \leq c$.

Theorem 3.5 (End-to-end complexity). *Under [Assumptions 3.1](#) and [3.2](#), let $0 < \varepsilon \leq 1$ and choose*

$$\lambda_\varepsilon := \frac{\varepsilon}{G^2}, \quad (3.13)$$

$$h_\varepsilon := \frac{c}{\ell_\varepsilon^2} \left[\frac{L_f \tau_f}{m} + \frac{L_f R}{\sqrt{m}} + \frac{R^2 + G^3 R/m}{\varepsilon} \right]^{-1}, \quad (3.14)$$

where $c > 0$ is a sufficiently small universal constant. For every $\mu_0 \in \mathcal{P}_2(\mathbb{R}^d)$, if

$$N \geq \frac{C}{mh_\varepsilon} \log \left(2 + \frac{\sqrt{m} W_2(\mu_0, \pi)}{\varepsilon} \right), \quad (3.15)$$

then

$$\sqrt{m} W_2(\mu_0 Q_{\lambda_\varepsilon, h_\varepsilon}^N, \pi) \leq \varepsilon. \quad (3.16)$$

Moreover, the smallest integer satisfying (3.15) obeys

$$N \leq 1 + \frac{C \ell_\varepsilon^2}{m} \left[\frac{L_f \tau_f}{m} + \frac{L_f R}{\sqrt{m}} + \frac{R^2 + G^3 R/m}{\varepsilon} \right] \log \left(2 + \frac{\sqrt{m} W_2(\mu_0, \pi)}{\varepsilon} \right). \quad (3.17)$$

In particular, for fixed model parameters and fixed initialization, $N(\varepsilon) = \tilde{O}(\varepsilon^{-1})$.

The proof of [Theorem 3.5](#) is given in [Section 4.5](#). The smoothing step uses the Moreau approximation estimate of Xin and Zhang [8, Proposition 5.9], also used in Xin and Zhang [9, Section 6.5]:

$$\sqrt{m} W_2(\pi_\lambda, \pi) \leq \frac{G^2 \lambda}{4}. \quad (3.18)$$

The choices (3.13)–(3.14) leave this bias budget unchanged.

Remark 3.6 (Parameter dependence). Using $R \leq G + \sqrt{\tau_f}$, the sufficient iteration bound (3.17) implies

$$N = \tilde{O} \left[\frac{L_f \tau_f}{m^2} + \frac{L_f G}{m^{3/2}} + \frac{1}{\varepsilon} \left(\frac{\tau_f}{m} + \frac{G^3 \sqrt{\tau_f}}{m^2} + \frac{G^4}{m^2} \right) \right], \quad (3.19)$$

where $\tilde{O}$ suppresses the logarithmic factors in (3.17). For fixed L_f/m and $G/\sqrt{m}$, the inequality $\tau_f \leq dL_f$ gives $N = \tilde{O}(d/\varepsilon)$.

4 Proof of the Main Results

This section proves [Theorems 3.4](#) and [3.5](#). We first use stationarity and a Poisson equation to bound $W_2(\hat{\pi}_{\lambda, h}, \pi_\lambda)$, using the flux estimates proved in [Section 5](#). We then combine this bound with the Moreau approximation error and contraction of the chain to obtain the iteration bound.

4.1 The analytic estimates

Lemma 4.1 (Target drift moments). *Under [Assumptions 3.1](#) and [3.2](#),*

$$\int \|\nabla U_\lambda\|^2 d\pi_\lambda = \int \text{tr}(\nabla^2 U_\lambda) d\pi_\lambda \leq R^2, \quad (4.1)$$

$$\|\nabla U_\lambda\|_{L^s(\pi_\lambda)} \leq 3\sqrt{s} R, \quad s \geq 2, \quad (4.2)$$

$$\int \exp \left(\frac{\|\nabla U_\lambda\|^2}{256 R^2} \right) d\pi_\lambda \leq 2. \quad (4.3)$$

The second-moment identity and bound follow from Xin and Zhang [9, Lemma 7.1 and Proposition A.4]. The higher-moment and exponential estimates are proved in [Section A.1](#). These estimates control the mean curvature and the drift moments using G and τ_f , uniformly in λ .

Lemma 4.2 (Finite-order stationary density ratios). *Let $\alpha \geq 2$. Suppose $hL_\lambda \leq c$ and*

$$h \frac{L_f \tau_f}{m} \leq \frac{c}{\alpha^2}, \quad h \frac{L_f R}{\sqrt{m}} \leq \frac{c}{\alpha}, \quad \frac{h GR}{\lambda m} \leq \frac{c}{\alpha^2}. \quad (4.4)$$

Then the stationary Euler interpolation

$$\nu_t := H_t((T_t)_\# \hat{\pi}_{\lambda,h}), \quad 0 \leq t \leq h, \quad \nu_0 = \nu_h = \hat{\pi}_{\lambda,h}, \quad (4.5)$$

satisfies

$$\sup_{0 \leq t \leq h} \|\nu_t / \pi_\lambda\|_{L^\alpha(\pi_\lambda)} \leq e^{1/4}. \quad (4.6)$$

The proof is given in [Appendix B](#). The prior integrability needed for differentiation is established independently in [Section D.2](#). The equal endpoint laws close a Rényi dissipation estimate over one Euler step using the log-Sobolev inequality for π_λ .

Lemma 4.3 (Normalized heat and pushforward estimates). *The estimates below apply to measurable scalar, vector, or matrix densities F , as indicated, whenever the norm on the right-hand side is finite. For $s \geq 2$, $t > 0$, and $ts^2 R^2 \leq c$,*

$$\|H_t F / \pi_\lambda\|_{L^s(\pi_\lambda)} \leq C \|F / \pi_\lambda\|_{L^{2s}(\pi_\lambda)}, \quad (4.7)$$

$$\|\nabla H_t F / \pi_\lambda\|_{L^s(\pi_\lambda)} \leq \frac{C}{\sqrt{t}} \|F / \pi_\lambda\|_{L^{2s}(\pi_\lambda)} \quad \text{for scalar } F, \quad (4.8)$$

$$\|\nabla \operatorname{div} H_t F / \pi_\lambda\|_{L^s(\pi_\lambda)} \leq \frac{C\sqrt{s}}{t} \|F / \pi_\lambda\|_{L^{4s}(\pi_\lambda)} \quad \text{for vector } F. \quad (4.9)$$

Moreover, if $st(L_\lambda + R^2) \leq c$, then

$$\|(T_t)_\# F / \pi_\lambda\|_{L^s(\pi_\lambda)} \leq C \|F / \pi_\lambda\|_{L^{2s}(\pi_\lambda)}. \quad (4.10)$$

The zero-order and pushforward estimates apply to scalar, vector, and matrix densities and extend to $t = 0$.

The proof is given in [Sections A.2](#) to [A.4](#).

4.2 The stationary residual

Assume $hL_\lambda < 1$ and let $\hat{\pi}_{\lambda,h}$ be the invariant density from [Proposition 3.3](#). Starting from $\hat{\pi}_{\lambda,h}$, a full drift step followed by heat evolution for time h returns to $\hat{\pi}_{\lambda,h}$. We compare this endpoint with the average along the heat path. Define

$$J := \frac{1}{h} \int_0^h (T_s)_\# (\nabla U_\lambda \hat{\pi}_{\lambda,h}) \, ds, \quad (4.11)$$

and

$$\rho_t := H_t((T_h)_\# \hat{\pi}_{\lambda,h}), \quad \bar{\rho} := \frac{1}{h} \int_0^h \rho_t \, dt, \quad 0 \leq t \leq h. \quad (4.12)$$

By stationarity,

$$\rho_0 = (T_h)_\# \widehat{\pi}_{\lambda,h}, \quad \rho_h = \widehat{\pi}_{\lambda,h}.$$

Unlike the Euler interpolation in (4.5), this path keeps T_h fixed as t varies. Define also

$$H := \frac{1}{h} \int_0^h t \nabla \rho_t \, dt, \quad (4.13)$$

$$E := -\frac{1}{h} \int_0^h (h-s)(T_s)_\# ((\nabla U_\lambda \otimes \nabla U_\lambda) \widehat{\pi}_{\lambda,h}) \, ds. \quad (4.14)$$

The gradient in (4.13) is used only for $t > 0$. The following lemma proves that these time integrals define L^1 fields and identifies their weak derivatives.

Lemma 4.4 (Exact stationary identities). *Under $hL_\lambda < 1$, the fields in (4.11)–(4.14) are well defined in $L^1(\mathbb{R}^d)$ and satisfy*

$$\begin{aligned} \widehat{\pi}_{\lambda,h} - \bar{\rho} &= \operatorname{div} H, \\ \nabla U_\lambda \widehat{\pi}_{\lambda,h} - J &= \operatorname{div} E, \\ \Delta \bar{\rho} + \operatorname{div} J &= 0. \end{aligned} \quad (4.15)$$

Consequently,

$$\mathcal{L}_\lambda^*(\bar{\rho} - \pi_\lambda) = \operatorname{div} \operatorname{div}(E - \nabla U_\lambda \otimes H) + \operatorname{div}((\nabla^2 U_\lambda)H). \quad (4.16)$$

All these identities hold in the sense of distributions. In particular, $\Delta \bar{\rho}$ denotes the distributional Laplacian; it has the L^1 representative $(\widehat{\pi}_{\lambda,h} - \rho_0)/h$.

Proof. Integrability of the fields. Since ∇U_λ has at most linear growth and $\widehat{\pi}_{\lambda,h} \in \mathcal{P}_2(\mathbb{R}^d)$,

$$\int (\|\nabla U_\lambda\| + \|\nabla U_\lambda\|^2) \widehat{\pi}_{\lambda,h} \, dx < \infty. \quad (4.17)$$

The pushforward and heat-kernel formulas give jointly measurable versions of the integrands in (4.11)–(4.14). Since ρ_0 is a probability density, so are ρ_t and $\bar{\rho}$. By Proposition 2.1 and (2.13),

$$\begin{aligned} \|J\|_{L^1} &\leq \int \|\nabla U_\lambda\| \widehat{\pi}_{\lambda,h} \, dx, \\ \|E\|_{L^1} &\leq \frac{h}{2} \int \|\nabla U_\lambda\|^2 \widehat{\pi}_{\lambda,h} \, dx, \\ \|H\|_{L^1} &\leq \frac{1}{h} \int_0^h t \|\nabla \rho_t\|_{L^1} \, dt \leq \frac{C_d}{h} \int_0^h \sqrt{t} \, dt = \frac{2C_d}{3} \sqrt{h}. \end{aligned}$$

All L^1 norms here are with respect to Lebesgue measure. These estimates justify the time integrals and their pairing with bounded test functions.

The drift balance. We first prove $\rho_0 - \widehat{\pi}_{\lambda,h} = h \operatorname{div} J$. Fix $\phi \in C_c^\infty(\mathbb{R}^d)$. For $\varepsilon \neq 0$ with $s, s + \varepsilon \in [0, h]$, the mean value theorem gives

$$\left| \frac{\phi(T_{s+\varepsilon}x) - \phi(T_sx)}{\varepsilon} \right| \widehat{\pi}_{\lambda,h}(x) \leq \|\nabla \phi\|_\infty \|\nabla U_\lambda(x)\| \widehat{\pi}_{\lambda,h}(x).$$

The right-hand side is integrable by (4.17) and independent of s, ε . Dominated convergence therefore permits differentiation under the integral in (2.10), giving

$$\begin{aligned} \frac{d}{ds} \int \phi(y)(T_s)_\# \widehat{\pi}_{\lambda,h}(y) \, dy &= - \int \langle \nabla \phi(T_sx), \nabla U_\lambda(x) \rangle \widehat{\pi}_{\lambda,h}(x) \, dx \\ &= - \int \langle \nabla \phi, (T_s)_\# (\nabla U_\lambda \widehat{\pi}_{\lambda,h}) \rangle \, dx. \end{aligned} \quad (4.18)$$

The same domination gives continuity of this derivative in s . Integrating (4.18) over $[0, h]$ yields

$$\int \phi(\rho_0 - \widehat{\pi}_{\lambda,h}) \, dx = -h \int \langle \nabla \phi, J \rangle \, dx = h \langle \operatorname{div} J, \phi \rangle.$$

Thus

$$\rho_0 - \widehat{\pi}_{\lambda,h} = (T_h)_\# \widehat{\pi}_{\lambda,h} - \widehat{\pi}_{\lambda,h} = h \operatorname{div} J \quad \text{in the sense of distributions.} \quad (4.19)$$

The heat balance and the endpoint difference. To prove the third identity in (4.15), we compute the distributional Laplacian through its definition. For $\phi \in C_c^\infty(\mathbb{R}^d)$, Fubini's theorem applies because

$$\int_0^h \int \rho_t |\Delta \phi| \, dx \, dt \leq h \|\Delta \phi\|_\infty < \infty.$$

The definition of the distributional Laplacian and (2.15) therefore give

$$\begin{aligned} \langle \Delta \bar{\rho}, \phi \rangle &:= \int \bar{\rho} \Delta \phi \, dx = \frac{1}{h} \int_0^h \int \rho_t \Delta \phi \, dx \, dt \\ &= \frac{1}{h} \int \phi(\rho_h - \rho_0) \, dx \\ &= \frac{1}{h} \int \phi(\widehat{\pi}_{\lambda,h} - \rho_0) \, dx = -\langle \operatorname{div} J, \phi \rangle. \end{aligned} \quad (4.20)$$

Here (2.15) is used with $F = \rho_0$, $a = 0$, and $b = h$, and the last equality is (4.19). This proves $\Delta \bar{\rho} + \operatorname{div} J = 0$ and the claimed L^1 representative of $\Delta \bar{\rho}$. No classical Laplacian of $\bar{\rho}$ was assumed.

We next prove $\widehat{\pi}_{\lambda,h} - \bar{\rho} = \operatorname{div} H$. The integrability already established for $t \nabla \rho_t$ permits Fubini's theorem. Spatial integration by parts for $t > 0$, followed by (2.14), gives

$$\begin{aligned} \langle \operatorname{div} H, \phi \rangle &= -\frac{1}{h} \int_0^h t \int \langle \nabla \phi, \nabla \rho_t \rangle \, dx \, dt \\ &= \frac{1}{h} \int_0^h t \int \rho_t \Delta \phi \, dx \, dt \\ &= \frac{1}{h} \int_0^h t \frac{d}{dt} \left(\int \phi \rho_t \, dx \right) \, dt \\ &= \frac{1}{h} \left[t \int \phi \rho_t \, dx \right]_0^h - \frac{1}{h} \int_0^h \int \phi \rho_t \, dx \, dt \\ &= \int \phi(\widehat{\pi}_{\lambda,h} - \bar{\rho}) \, dx. \end{aligned} \quad (4.21)$$

The time integration by parts is valid because $t \mapsto \int \phi \rho_t \, dx$ is absolutely continuous by (2.15). Its boundary term at zero vanishes since $|t \int \phi \rho_t \, dx| \leq t \|\phi\|_\infty$.

The matrix flux. We prove the second identity in (4.15) componentwise. For each fixed i , the difference quotients for the pairing in (2.10) with density $(\partial_i U_\lambda) \widehat{\pi}_{\lambda,h}$ are dominated by the integrable function $\|\nabla \phi\|_\infty \|\nabla U_\lambda\|^2 \widehat{\pi}_{\lambda,h}$. We may therefore differentiate under the integral. Keeping the initial weight $\partial_i U_\lambda(x)$ fixed gives

$$\begin{aligned} &\frac{d}{ds} \int \phi(y) [(T_s)_\# (\nabla U_\lambda \widehat{\pi}_{\lambda,h})]_i(y) \, dy \\ &= - \sum_j \int \partial_j \phi(T_s x) \partial_i U_\lambda(x) \partial_j U_\lambda(x) \widehat{\pi}_{\lambda,h}(x) \, dx \\ &= - \sum_j \int \partial_j \phi(y) [(T_s)_\# ((\nabla U_\lambda \otimes \nabla U_\lambda) \widehat{\pi}_{\lambda,h})]_{ij}(y) \, dy. \end{aligned} \quad (4.22)$$

The same second-moment bound justifies Fubini's theorem for the two time integrals below, since

$$\int_0^h \int_0^s \int \|\nabla U_\lambda\|^2 \hat{\pi}_{\lambda,h} dx du ds = \frac{h^2}{2} \int \|\nabla U_\lambda\|^2 \hat{\pi}_{\lambda,h} dx < \infty.$$

Integrating (4.22) first from 0 to s and then averaging over $s \in [0, h]$, we obtain

$$\begin{aligned} & \int \phi((\partial_i U_\lambda) \hat{\pi}_{\lambda,h} - J_i) dx \\ &= -\frac{1}{h} \int_0^h \int_0^s \frac{d}{du} \left(\int \phi(y) [(T_u)_\# (\nabla U_\lambda \hat{\pi}_{\lambda,h})]_i(y) dy \right) du ds \\ &= \frac{1}{h} \sum_j \int_0^h (h-u) \int \partial_j \phi(y) [(T_u)_\# ((\nabla U_\lambda \otimes \nabla U_\lambda) \hat{\pi}_{\lambda,h})]_{ij}(y) dy du \\ &= -\sum_j \int \partial_j \phi E_{ij} dx = \langle (\operatorname{div} E)_i, \phi \rangle. \end{aligned}$$

This proves $\nabla U_\lambda \hat{\pi}_{\lambda,h} - J = \operatorname{div} E$ and completes (4.15).

The residual equation. We use the product rule

$$\operatorname{div}(\nabla U_\lambda \otimes H) = \nabla U_\lambda (\hat{\pi}_{\lambda,h} - \bar{\rho}) + (\nabla^2 U_\lambda) H. \quad (4.23)$$

To justify it, test $\operatorname{div} H = \hat{\pi}_{\lambda,h} - \bar{\rho}$ against smooth approximations of $(\partial_i U_\lambda) \phi$. Mollifications of $\partial_i U_\lambda$ converge locally uniformly; their gradients are uniformly bounded and converge almost everywhere to its weak gradient. Since H and $\hat{\pi}_{\lambda,h} - \bar{\rho}$ are locally integrable, dominated convergence gives

$$-\int (\partial_i U_\lambda) \langle H, \nabla \phi \rangle dx = \int (\partial_i U_\lambda) (\hat{\pi}_{\lambda,h} - \bar{\rho}) \phi dx + \int \phi \sum_j (\partial_j \partial_i U_\lambda) H_j dx,$$

which is the i th component of (4.23). Now $\mathcal{L}_\lambda^* \pi_\lambda = 0$, (4.15), and (4.23) give

$$\begin{aligned} \mathcal{L}_\lambda^* (\bar{\rho} - \pi_\lambda) &= \Delta \bar{\rho} + \operatorname{div}(\nabla U_\lambda \bar{\rho}) \\ &= \operatorname{div}(\nabla U_\lambda \bar{\rho} - J) \\ &= \operatorname{div}(\operatorname{div} E - \nabla U_\lambda (\hat{\pi}_{\lambda,h} - \bar{\rho})) \\ &= \operatorname{div}(\operatorname{div} E - \operatorname{div}(\nabla U_\lambda \otimes H) + (\nabla^2 U_\lambda) H) \\ &= \operatorname{div} \operatorname{div}(E - \nabla U_\lambda \otimes H) + \operatorname{div}((\nabla^2 U_\lambda) H). \end{aligned}$$

All derivatives in this calculation are distributional. Equivalently, the resulting identity means that

$$\int (\bar{\rho} - \pi_\lambda) \mathcal{L}_\lambda \phi dx = \int \langle E - \nabla U_\lambda \otimes H, \nabla^2 \phi \rangle dx - \int \langle (\nabla^2 U_\lambda) H, \nabla \phi \rangle dx$$

for every $\phi \in C_c^\infty(\mathbb{R}^d)$. This proves (4.16). $\square$

For the quantitative flux bounds below and in Section 5, choose the moment order

$$r := 4 + 2 \log(1 + L_\lambda/m). \quad (4.24)$$

Lemma 4.5 (Stationary flux bounds). *Under (3.11), the fields in (4.11)–(4.14) satisfy*

$$\|H/\pi_\lambda\|_{L^r(\pi_\lambda)} \leq ChrR \log \frac{e}{mh}, \quad (4.25)$$

$$\|E/\pi_\lambda\|_{L^2(\pi_\lambda)} \leq ChR^2, \quad (4.26)$$

$$\|(E - \nabla U_\lambda \otimes H)/\pi_\lambda\|_{L^2(\pi_\lambda)} \leq ChrR^2 \log \frac{e}{mh}. \quad (4.27)$$

The proof is given in Section 5.

4.3 A residual-to-transport estimate

Proposition 4.6 (Second-order residual estimate). *Let $\nu, \rho \in \mathcal{P}_2(\mathbb{R}^d)$ be densities with $\nu/\pi_\lambda, \rho/\pi_\lambda \in L^2(\pi_\lambda)$. Suppose a vector density H and a matrix density E satisfy*

$$\nu - \rho = \operatorname{div} H, \quad (4.28)$$

$$\mathcal{L}_\lambda^*(\rho - \pi_\lambda) = \operatorname{div} \operatorname{div}(E - \nabla U_\lambda \otimes H) + \operatorname{div}((\nabla^2 U_\lambda)H) \quad (4.29)$$

in the sense of distributions. If the norms on the right below are finite, then

$$\begin{aligned} \sqrt{m} W_2(\nu, \pi_\lambda) &\leq 2\sqrt{m} \|H/\pi_\lambda\|_{L^2(\pi_\lambda)} + 2 \|(E - \nabla U_\lambda \otimes H)/\pi_\lambda\|_{L^2(\pi_\lambda)} \\ &\quad + 2 \left(\int \frac{H^\top (\nabla^2 U_\lambda) H}{\pi_\lambda} dx \right)^{1/2}. \end{aligned} \quad (4.30)$$

For clarity, (4.28) and (4.29) mean that, for every $v \in C_c^\infty(\mathbb{R}^d)$,

$$\begin{aligned} \int v(\nu - \rho) dx &= - \int \langle \nabla v, H \rangle dx, \\ \int (\rho - \pi_\lambda) \mathcal{L}_\lambda v dx &= \int \langle E - \nabla U_\lambda \otimes H, \nabla^2 v \rangle dx - \int \langle (\nabla^2 U_\lambda)H, \nabla v \rangle dx. \end{aligned}$$

All derivatives in these integral identities act on the test function. In particular, no classical derivatives of E or of the weak Hessian $\nabla^2 U_\lambda$ are required.

Proof of Proposition 4.6. Fix $\phi \in C_c^\infty(\mathbb{R}^d)$ and let ψ be its Poisson solution from Lemma 2.2. We first prove the pairing identity

$$\begin{aligned} \int \phi(\rho - \pi_\lambda) dx &= \int (\rho - \pi_\lambda) \mathcal{L}_\lambda \psi dx \\ &= \int \langle E - \nabla U_\lambda \otimes H, \nabla^2 \psi \rangle dx - \int \langle (\nabla^2 U_\lambda)H, \nabla \psi \rangle dx. \end{aligned} \quad (4.31)$$

The residual equation (4.29) is initially available only for compactly supported smooth tests. We therefore take the sequence $\psi_n \in C_c^\infty(\mathbb{R}^d)$ from (2.21). For every n , the weak formulation gives

$$\int (\rho - \pi_\lambda) \mathcal{L}_\lambda \psi_n dx = \int \langle E - \nabla U_\lambda \otimes H, \nabla^2 \psi_n \rangle dx - \int \langle (\nabla^2 U_\lambda)H, \nabla \psi_n \rangle dx. \quad (4.32)$$

We now justify passage to the limit in each integral of (4.32).

For the curvature integral, positivity of $\nabla^2 U_\lambda$ and Cauchy–Schwarz give, for every $v \in H^1(\pi_\lambda)$,

$$\left| \int \langle (\nabla^2 U_\lambda)H, \nabla v \rangle dx \right| \leq \left(\int \frac{H^\top (\nabla^2 U_\lambda) H}{\pi_\lambda} dx \right)^{1/2} \left(\int \nabla v^\top (\nabla^2 U_\lambda) \nabla v d\pi_\lambda \right)^{1/2}. \quad (4.33)$$

Using Cauchy–Schwarz for the density and matrix integrals, and (4.33) with $v = \psi_n - \psi$ for the curvature integral, we obtain

$$\begin{aligned} \left| \int (\rho - \pi_\lambda) (\mathcal{L}_\lambda \psi_n - \mathcal{L}_\lambda \psi) dx \right| &\leq \|\rho/\pi_\lambda - 1\|_{L^2(\pi_\lambda)} \|\mathcal{L}_\lambda \psi_n - \mathcal{L}_\lambda \psi\|_{L^2(\pi_\lambda)} \longrightarrow 0, \\ \left| \int \langle E - \nabla U_\lambda \otimes H, \nabla^2 \psi_n - \nabla^2 \psi \rangle dx \right| &\leq \|(E - \nabla U_\lambda \otimes H)/\pi_\lambda\|_{L^2(\pi_\lambda)} \|\nabla^2 \psi_n - \nabla^2 \psi\|_{L^2(\pi_\lambda)} \longrightarrow 0, \\ \left| \int \langle (\nabla^2 U_\lambda)H, \nabla \psi_n - \nabla \psi \rangle dx \right| &\leq \sqrt{L_\lambda} \left(\int \frac{H^\top (\nabla^2 U_\lambda) H}{\pi_\lambda} dx \right)^{1/2} \|\nabla \psi_n - \nabla \psi\|_{L^2(\pi_\lambda)} \longrightarrow 0. \end{aligned} \quad (4.34)$$

The coefficient norms are finite by the hypotheses of the proposition. The three approximation errors tend to zero by (2.21); the last estimate also uses $\nabla^2 U_\lambda \preceq L_\lambda I$. Thus (4.34) allows us to pass to the limit in (4.32), proving (4.31).

We next estimate the right-hand side of (4.31). For the matrix integral, Cauchy–Schwarz uses the Frobenius inner product; for the curvature integral, apply (4.33) with $v = \psi$. Consequently,

$$\begin{aligned} \left| \int \phi(\rho - \pi_\lambda) dx \right| &\leq \|(E - \nabla U_\lambda \otimes H)/\pi_\lambda\|_{L^2(\pi_\lambda)} \left(\int \|\nabla^2 \psi\|_F^2 d\pi_\lambda \right)^{1/2} \\ &\quad + \left(\int \frac{H^\top (\nabla^2 U_\lambda) H}{\pi_\lambda} dx \right)^{1/2} \left(\int \nabla \psi^\top (\nabla^2 U_\lambda) \nabla \psi d\pi_\lambda \right)^{1/2}. \end{aligned} \quad (4.35)$$

From (2.20),

$$\begin{aligned} \left(\int \|\nabla^2 \psi\|_F^2 d\pi_\lambda \right)^{1/2} &\leq \frac{1}{\sqrt{m}} \|\nabla \phi\|_{L^2(\pi_\lambda)}, \\ \left(\int \nabla \psi^\top (\nabla^2 U_\lambda) \nabla \psi d\pi_\lambda \right)^{1/2} &\leq \frac{1}{\sqrt{m}} \|\nabla \phi\|_{L^2(\pi_\lambda)}. \end{aligned} \quad (4.36)$$

Substituting (4.36) into (4.35) yields

$$\left| \int \phi(\rho - \pi_\lambda) dx \right| \leq \frac{\|\nabla \phi\|_{L^2(\pi_\lambda)}}{\sqrt{m}} \left[\|(E - \nabla U_\lambda \otimes H)/\pi_\lambda\|_{L^2(\pi_\lambda)} + \left(\int \frac{H^\top (\nabla^2 U_\lambda) H}{\pi_\lambda} dx \right)^{1/2} \right]. \quad (4.37)$$

The first residual identity (4.28), tested directly with ϕ , gives

$$\left| \int \phi(\nu - \rho) dx \right| = \left| \int \langle \nabla \phi, H \rangle dx \right| \leq \|H/\pi_\lambda\|_{L^2(\pi_\lambda)} \|\nabla \phi\|_{L^2(\pi_\lambda)}. \quad (4.38)$$

Since $\nu - \pi_\lambda = (\nu - \rho) + (\rho - \pi_\lambda)$, the triangle inequality, (4.38), and (4.37) imply

$$\begin{aligned} \left| \int \phi(\nu - \pi_\lambda) dx \right| &\leq \left| \int \phi(\nu - \rho) dx \right| + \left| \int \phi(\rho - \pi_\lambda) dx \right| \\ &\leq \|\nabla \phi\|_{L^2(\pi_\lambda)} \left[\|H/\pi_\lambda\|_{L^2(\pi_\lambda)} + \frac{1}{\sqrt{m}} \|(E - \nabla U_\lambda \otimes H)/\pi_\lambda\|_{L^2(\pi_\lambda)} \right. \\ &\quad \left. + \frac{1}{\sqrt{m}} \left(\int \frac{H^\top (\nabla^2 U_\lambda) H}{\pi_\lambda} dx \right)^{1/2} \right]. \end{aligned} \quad (4.39)$$

Taking the supremum in (4.39) over the test functions in (2.22) gives

$$\begin{aligned} \|\nu - \pi_\lambda\|_{\dot{H}^{-1}(\pi_\lambda)} &= \sup_{\substack{\phi \in C_c^\infty(\mathbb{R}^d) \\ \|\nabla \phi\|_{L^2(\pi_\lambda)} \leq 1}} \left| \int \phi(\nu - \pi_\lambda) dx \right| \\ &\leq \|H/\pi_\lambda\|_{L^2(\pi_\lambda)} + \frac{1}{\sqrt{m}} \|(E - \nabla U_\lambda \otimes H)/\pi_\lambda\|_{L^2(\pi_\lambda)} \\ &\quad + \frac{1}{\sqrt{m}} \left(\int \frac{H^\top (\nabla^2 U_\lambda) H}{\pi_\lambda} dx \right)^{1/2}. \end{aligned} \quad (4.40)$$

Finally, $\nu/\pi_\lambda \in L^2(\pi_\lambda)$ by assumption, so (2.25) applies. Using that comparison and then (4.40), we conclude that

$$\begin{aligned} \sqrt{m} W_2(\nu, \pi_\lambda) &\leq 2\sqrt{m} \|\nu - \pi_\lambda\|_{\dot{H}^{-1}(\pi_\lambda)} \\ &\leq 2\sqrt{m} \|H/\pi_\lambda\|_{L^2(\pi_\lambda)} + 2\|(E - \nabla U_\lambda \otimes H)/\pi_\lambda\|_{L^2(\pi_\lambda)} \\ &\quad + 2 \left(\int \frac{H^\top (\nabla^2 U_\lambda) H}{\pi_\lambda} dx \right)^{1/2}. \end{aligned}$$

This is (4.30). □

4.4 Proof of the fixed-smoothing theorem

Proof of Theorem 3.4. Assume (3.11). Since $\tau_f \geq md \geq m$ and $L_\lambda = L_f + \lambda^{-1}$, we have

$$h(L_\lambda + R^2) \leq h \left[\frac{L_f \tau_f}{m} + R^2 + \frac{1}{\lambda} \right] \leq \frac{c}{\ell_{h,\lambda}^2} \leq c. \quad (4.41)$$

Taking the universal constant in (3.11) sufficiently small gives $hL_\lambda \leq 1/2$. By Proposition 3.3, $Q_{\lambda,h}$ therefore has a unique invariant law $\hat{\pi}_{\lambda,h} \in \mathcal{P}_2(\mathbb{R}^d)$ with a strictly positive smooth density.

Let r be as in (4.24), and take $\bar{\rho}, H, E$ from (4.12)–(4.14). We will apply Proposition 4.6 with $\nu = \hat{\pi}_{\lambda,h}$ and $\rho = \bar{\rho}$. We first verify its density assumptions and then estimate the three terms in (4.30).

The density assumptions. (3.11) gives

$$64h \frac{L_f \tau_f}{m} + 8h \frac{L_f R}{\sqrt{m}} + 64h \frac{GR}{m\lambda} \leq 64h \left[\frac{L_f \tau_f}{m} + \frac{L_f R}{\sqrt{m}} + \frac{GR}{m\lambda} \right] \leq \frac{64c}{\ell_{h,\lambda}^2}. \quad (4.42)$$

Together with (4.41), these bounds imply the conditions of Lemma 4.2 with $\alpha = 8$, after decreasing the universal constant in (3.11). Evaluating (4.6) at $t = 0$ yields

$$\|\hat{\pi}_{\lambda,h}/\pi_\lambda\|_{L^2(\pi_\lambda)} \leq \|\hat{\pi}_{\lambda,h}/\pi_\lambda\|_{L^8(\pi_\lambda)} \leq e^{1/4}. \quad (4.43)$$

For $0 < t \leq h$, apply (4.7) with output order $s = 2$ and (4.10) with output order $s = 4$. Their time restrictions hold because

$$4tR^2 \leq 4hR^2 \leq 4c, \quad 4h(L_\lambda + R^2) \leq 4c$$

by (4.41). Thus, after the same choice of a sufficiently small constant,

$$\begin{aligned} \|\rho_t/\pi_\lambda\|_{L^2(\pi_\lambda)} &= \|\mathbf{H}_t((T_h)_\# \hat{\pi}_{\lambda,h})/\pi_\lambda\|_{L^2(\pi_\lambda)} \\ &\leq C \|(T_h)_\# \hat{\pi}_{\lambda,h}/\pi_\lambda\|_{L^4(\pi_\lambda)} \\ &\leq C \|\hat{\pi}_{\lambda,h}/\pi_\lambda\|_{L^8(\pi_\lambda)} \leq C. \end{aligned} \quad (4.44)$$

Jensen's inequality for the time average, followed by Tonelli's theorem and (4.44), gives

$$\begin{aligned} \|\bar{\rho}/\pi_\lambda\|_{L^2(\pi_\lambda)}^2 &= \int \left(\frac{1}{h} \int_0^h \frac{\rho_t(x)}{\pi_\lambda(x)} dt \right)^2 d\pi_\lambda(x) \\ &\leq \frac{1}{h} \int_0^h \|\rho_t/\pi_\lambda\|_{L^2(\pi_\lambda)}^2 dt < \infty. \end{aligned} \quad (4.45)$$

The heat-kernel representation also gives

$$\int \|y\|^2 \bar{\rho}(y) dy = \int \|T_h(x)\|^2 \hat{\pi}_{\lambda,h}(x) dx + dh < \infty.$$

Indeed, the Gaussian increment at time t contributes $2td$ to the second moment, and T_h has at most linear growth. Hence $\bar{\rho} \in \mathcal{P}_2(\mathbb{R}^d)$. Equations (4.43) and (4.45) verify the two L^2 density-ratio assumptions of Proposition 4.6.

The curvature term. The flux bounds (4.25) and (4.27) are supplied by Lemma 4.5. It remains to estimate the integral involving $\nabla^2 U_\lambda$ in (4.30). Since $\nabla^2 U_\lambda$ is positive semidefinite, its quadratic form is bounded by its trace times the squared Euclidean norm. Applying this pointwise and then using Hölder with conjugate exponents $r/2$ and $r/(r-2)$ gives

$$\begin{aligned} \int \frac{H^\top (\nabla^2 U_\lambda) H}{\pi_\lambda} dx &= \int \left(\frac{H}{\pi_\lambda} \right)^\top (\nabla^2 U_\lambda) \left(\frac{H}{\pi_\lambda} \right) d\pi_\lambda \\ &\leq \int \|H/\pi_\lambda\|^2 \operatorname{tr}(\nabla^2 U_\lambda) d\pi_\lambda \\ &\leq \|H/\pi_\lambda\|_{L^r(\pi_\lambda)}^2 \left(\int [\operatorname{tr}(\nabla^2 U_\lambda)]^{r/(r-2)} d\pi_\lambda \right)^{(r-2)/r}. \end{aligned} \quad (4.46)$$

To estimate the last integral, use $0 \leq \operatorname{tr}(\nabla^2 U_\lambda) \leq dL_\lambda$ almost everywhere and (4.1). Since $r/(r-2) = 1 + 2/(r-2)$,

$$\begin{aligned} \int [\operatorname{tr}(\nabla^2 U_\lambda)]^{r/(r-2)} d\pi_\lambda &= \int \operatorname{tr}(\nabla^2 U_\lambda) [\operatorname{tr}(\nabla^2 U_\lambda)]^{2/(r-2)} d\pi_\lambda \\ &\leq (dL_\lambda)^{2/(r-2)} \int \operatorname{tr}(\nabla^2 U_\lambda) d\pi_\lambda \\ &\leq (dL_\lambda)^{2/(r-2)} R^2. \end{aligned} \quad (4.47)$$

Moreover, $R^2 \geq md$ and the choice of r in (4.24) imply

$$\left(\frac{dL_\lambda}{R^2} \right)^{1/r} \leq \left(\frac{L_\lambda}{m} \right)^{1/r} = \exp \left\{ \frac{\log(L_\lambda/m)}{4 + 2\log(1 + L_\lambda/m)} \right\} \leq \sqrt{e}. \quad (4.48)$$

Substituting (4.47) into (4.46), taking square roots, and using (4.48), we obtain

$$\begin{aligned} \left(\int \frac{H^\top (\nabla^2 U_\lambda) H}{\pi_\lambda} dx \right)^{1/2} &\leq R \left(\frac{dL_\lambda}{R^2} \right)^{1/r} \|H/\pi_\lambda\|_{L^r(\pi_\lambda)} \\ &\leq \sqrt{e} R \|H/\pi_\lambda\|_{L^r(\pi_\lambda)}. \end{aligned} \quad (4.49)$$

In particular, this integral is finite by (4.25).

Application of the residual estimate. The identities (4.15) and (4.16) are precisely the weak identities required in Proposition 4.6 for $\nu = \hat{\pi}_{\lambda,h}$ and $\rho = \bar{\rho}$. Its density assumptions have been verified above, and its weighted norms are finite by (4.25), (4.27), and (4.49). Since $r \geq 4$, $\|H/\pi_\lambda\|_{L^2(\pi_\lambda)} \leq \|H/\pi_\lambda\|_{L^r(\pi_\lambda)}$. Applying Proposition 4.6 therefore gives

$$\begin{aligned} \sqrt{m} W_2(\hat{\pi}_{\lambda,h}, \pi_\lambda) &\leq 2\sqrt{m} \|H/\pi_\lambda\|_{L^2(\pi_\lambda)} + 2 \|(E - \nabla U_\lambda \otimes H)/\pi_\lambda\|_{L^2(\pi_\lambda)} \\ &\quad + 2 \left(\int \frac{H^\top (\nabla^2 U_\lambda) H}{\pi_\lambda} dx \right)^{1/2} \\ &\leq 2(\sqrt{m} + \sqrt{e} R) \|H/\pi_\lambda\|_{L^r(\pi_\lambda)} + 2 \|(E - \nabla U_\lambda \otimes H)/\pi_\lambda\|_{L^2(\pi_\lambda)} \\ &\leq C h r R^2 \log \frac{e}{mh}. \end{aligned} \quad (4.50)$$

The last inequality uses (4.25), (4.27), and $\sqrt{m} \leq R$.

Finally, the definitions (4.24) and (3.9) give

$$r \leq 4\ell_{h,\lambda}, \quad \log \frac{e}{mh} = 1 + \log \frac{1}{mh} \leq \ell_{h,\lambda}. \quad (4.51)$$

Combining (4.50) and (4.51) yields

$$\sqrt{m} W_2(\hat{\pi}_{\lambda,h}, \pi_\lambda) \leq ChR^2 \ell_{h,\lambda}^2.$$

This proves (3.12). $\square$

4.5 Proof of the end-to-end complexity theorem

Proof of Theorem 3.5. We allocate $\varepsilon/4$ to the Moreau bias, $\varepsilon/4$ to the invariant-measure bias, and $\varepsilon/2$ to convergence from the initial law. The choice (3.13) gives the first budget by (3.18). We verify that (3.14) gives the second and that (3.15) gives the third. Substitute $\lambda = \varepsilon/G^2$ into the bracket in (3.11). Because $R \geq G$ and $\varepsilon \leq 1$,

$$\frac{L_f \tau_f}{m} + \frac{L_f R}{\sqrt{m}} + \frac{GR}{m\lambda} + R^2 + \frac{1}{\lambda} \leq 2 \left[\frac{L_f \tau_f}{m} + \frac{L_f R}{\sqrt{m}} + \frac{R^2 + G^3 R/m}{\varepsilon} \right]. \quad (4.52)$$

Also $R/\sqrt{m} \leq G/\sqrt{m} + \sqrt{\tau_f/m}$. The reciprocal of mh_ε in (3.14) is consequently bounded above by a fixed polynomial in the dimensionless ratios appearing in (3.10), multiplied by ℓ_ε^2/c . It follows that

$$\ell_{h_\varepsilon, \lambda_\varepsilon} \leq C\ell_\varepsilon + \log(1/c). \quad (4.53)$$

Choose the absolute c small enough, using $c(1 + \log(1/c))^2 \rightarrow 0$. Equations (4.52) and (4.53) show that (3.11) holds and that

$$\sqrt{m} W_2(\hat{\pi}_{\lambda_\varepsilon, h_\varepsilon}, \pi_{\lambda_\varepsilon}) \leq \varepsilon/4. \quad (4.54)$$

Equation (3.18) gives another $\varepsilon/4$.

Synchronous coupling of (3.5) gives

$$W_2(\mu_0 Q_{\lambda,h}^N, \hat{\pi}_{\lambda,h}) \leq (1 - mh)^N W_2(\mu_0, \hat{\pi}_{\lambda,h}). \quad (4.55)$$

The triangle inequality and the two bias budgets imply

$$\sqrt{m} W_2(\mu_0, \hat{\pi}_{\lambda_\varepsilon, h_\varepsilon}) \leq \sqrt{m} W_2(\mu_0, \pi) + \varepsilon/2.$$

Thus (3.15) makes the contraction error at most $\varepsilon/2$. The triangle inequality gives

$$\begin{aligned} \sqrt{m} W_2(\mu_0 Q_{\lambda_\varepsilon, h_\varepsilon}^N, \pi) &\leq \sqrt{m} W_2(\mu_0 Q_{\lambda_\varepsilon, h_\varepsilon}^N, \hat{\pi}_{\lambda_\varepsilon, h_\varepsilon}) \\ &\quad + \sqrt{m} W_2(\hat{\pi}_{\lambda_\varepsilon, h_\varepsilon}, \pi_{\lambda_\varepsilon}) + \sqrt{m} W_2(\pi_{\lambda_\varepsilon}, \pi) \leq \varepsilon. \end{aligned}$$

This proves (3.16); substituting (3.14) into (3.15) proves (3.17).

This completes the proof of Theorem 3.5. $\square$

5 Stationary Flux Estimates

This section proves the bounds on H and E in Lemma 4.5; the fields are defined in Section 4.2. We first bound the averaged drift J , then use a finite heat expansion to estimate the gradient of the invariant density $\hat{\pi}_{\lambda,h}$ and obtain the bound on H . The bound on E follows from the drift moment and pushforward estimates.

5.1 The averaged drift flux

Assume (3.11). Since $\tau_f \geq m$, this condition implies $hL_\lambda \leq c$, so Proposition 3.3 supplies the invariant density $\hat{\pi}_{\lambda,h}$.

By (4.24) and (3.9), $r \leq 4\ell_{h,\lambda}$, and hence (3.11) implies

$$h(32r)^2 \left[\frac{L_f \tau_f}{m} + \frac{GR}{m\lambda} \right] + h(32r) \frac{L_f R}{\sqrt{m}} \leq Ch\ell_{h,\lambda}^2 \left[\frac{L_f \tau_f}{m} + \frac{L_f R}{\sqrt{m}} + \frac{GR}{m\lambda} \right] \leq Cc.$$

Taking the universal constant in (3.11) sufficiently small therefore ensures (4.4) with $\alpha = 32r$. By (4.6) with $\alpha = 32r$ and $t = 0$,

$$\|\hat{\pi}_{\lambda,h}/\pi_\lambda\|_{L^{32r}(\pi_\lambda)} \leq e^{1/4}. \quad (5.1)$$

For the next estimate, we use (4.10) with output order $s = 8r$ and times $0 \leq t \leq h$. Its restriction is satisfied after decreasing the constant in (3.11), since $L_\lambda \leq L_f \tau_f / m + \lambda^{-1}$ and $8rh(L_\lambda + R^2) \leq Cc$. Minkowski's inequality for the time average in (4.11), followed by (4.10) and Hölder's inequality, gives

$$\begin{aligned} \|J/\pi_\lambda\|_{L^{8r}(\pi_\lambda)} &\leq \frac{1}{h} \int_0^h \|(T_u)_\#(\nabla U_\lambda \hat{\pi}_{\lambda,h})/\pi_\lambda\|_{L^{8r}(\pi_\lambda)} du \\ &\leq C \|\nabla U_\lambda \hat{\pi}_{\lambda,h}/\pi_\lambda\|_{L^{16r}(\pi_\lambda)} \\ &\leq C \|\nabla U_\lambda\|_{L^{32r}(\pi_\lambda)} \|\hat{\pi}_{\lambda,h}/\pi_\lambda\|_{L^{32r}(\pi_\lambda)} \leq C\sqrt{r} R. \end{aligned} \quad (5.2)$$

The last step uses (4.2) at order $32r$ and (5.1).

5.2 A finite heat expansion

We next derive the finite heat expansion. By (4.19), the distributional divergence of J has the L^1 representative

$$\operatorname{div} J = \frac{(T_h)_\# \hat{\pi}_{\lambda,h} - \hat{\pi}_{\lambda,h}}{h}.$$

Since $J \in L^1(\mathbb{R}^d; \mathbb{R}^d)$, both J and $\operatorname{div} J$ can therefore be convolved with the heat kernel. For every $t > 0$,

$$\mathbf{H}_t(\operatorname{div} J) = \operatorname{div}(\mathbf{H}_t J). \quad (5.3)$$

To verify this identity, take $\phi \in C_c^\infty(\mathbb{R}^d)$. Symmetry of the heat kernel and $\nabla \mathbf{H}_t \phi = \mathbf{H}_t \nabla \phi$ give

$$\begin{aligned} \int \phi \mathbf{H}_t(\operatorname{div} J) dx &= \int (\mathbf{H}_t \phi) \operatorname{div} J dx \\ &= - \int \langle \nabla \mathbf{H}_t \phi, J \rangle dx \\ &= - \int \langle \nabla \phi, \mathbf{H}_t J \rangle dx = \int \phi \operatorname{div}(\mathbf{H}_t J) dx. \end{aligned}$$

For the integration by parts with $\mathbf{H}_t \phi$, first multiply it by smooth cutoffs χ_k equal to one on the ball of radius k , supported in the ball of radius $2k$, and satisfying $\|\nabla \chi_k\|_\infty \leq C/k$. The cutoff error is bounded by

$$\|\mathbf{H}_t \phi\|_\infty \|\nabla \chi_k\|_\infty \|J\|_{L^1} \longrightarrow 0.$$

The remaining terms converge by dominated convergence, since $J, \operatorname{div} J \in L^1$ and $\mathbf{H}_t \phi, \nabla \mathbf{H}_t \phi$ are bounded. This proves (5.3).

Applying H_h to (4.19), and using stationarity (3.8) and (5.3), yields

$$\begin{aligned}\widehat{\pi}_{\lambda,h} &= H_h((T_h)_\# \widehat{\pi}_{\lambda,h}) \\ &= H_h \widehat{\pi}_{\lambda,h} + h H_h(\operatorname{div} J) \\ &= H_h \widehat{\pi}_{\lambda,h} + h \operatorname{div} H_h J.\end{aligned}\tag{5.4}$$

For $k = 0, \dots, n-1$, the semigroup property and (5.3) then give

$$H_{kh} \widehat{\pi}_{\lambda,h} - H_{(k+1)h} \widehat{\pi}_{\lambda,h} = h \operatorname{div} H_{(k+1)h} J.$$

Summing these equalities gives the exact finite identity

$$\widehat{\pi}_{\lambda,h} = H_{nh} \widehat{\pi}_{\lambda,h} + h \sum_{j=1}^n \operatorname{div} H_{jh} J, \quad n \geq 1.\tag{5.5}$$

For positive time, convolution of an L^1 function with the Gaussian kernel is smooth: every spatial derivative can be placed on the kernel, whose derivatives are bounded and integrable. Thus the right-hand side of (5.5) is smooth. The identity, initially obtained distributionally, consequently holds pointwise for the smooth representative of $\widehat{\pi}_{\lambda,h}$.

Choose

$$\tau := \frac{c_1}{r^2 R^2}, \quad n := \left\lfloor \frac{\tau}{h} \right\rfloor,$$

where $0 < c_1 \leq 1$ is a sufficiently small universal constant. Since $r \leq 4\ell_{h,\lambda}$, (3.11) gives $hr^2 R^2 \leq 16c$. Decreasing the constant c in (3.11) relative to c_1 therefore ensures

$$h \leq \frac{\tau}{2}, \quad \frac{\tau}{2} \leq nh \leq \tau.$$

We will apply (4.8) and (4.9) with output order $s = 2r$, at times $t = nh$ and $t = jh$, respectively. Their common time restriction holds because, for $1 \leq j \leq n$,

$$(jh)(2r)^2 R^2 \leq \tau(2r)^2 R^2 = 4c_1.\tag{5.6}$$

All terms in (5.5) are smooth, and its sum is finite. We may therefore differentiate it to obtain

$$\nabla \widehat{\pi}_{\lambda,h} = \nabla H_{nh} \widehat{\pi}_{\lambda,h} + h \sum_{j=1}^n \nabla \operatorname{div} H_{jh} J.\tag{5.7}$$

Taking the weighted L^{2r} norm in (5.7), and using (4.8), (4.9), and $nh \geq \tau/2$, gives

$$\begin{aligned}\|\nabla \widehat{\pi}_{\lambda,h}/\pi_\lambda\|_{L^{2r}(\pi_\lambda)} &\leq \|\nabla H_{nh} \widehat{\pi}_{\lambda,h}/\pi_\lambda\|_{L^{2r}(\pi_\lambda)} + h \sum_{j=1}^n \|\nabla \operatorname{div} H_{jh} J/\pi_\lambda\|_{L^{2r}(\pi_\lambda)} \\ &\leq C\tau^{-1/2} \|\widehat{\pi}_{\lambda,h}/\pi_\lambda\|_{L^{4r}(\pi_\lambda)} + Ch \sum_{j=1}^n \frac{\sqrt{r}}{jh} \|J/\pi_\lambda\|_{L^{8r}(\pi_\lambda)} \\ &\leq CrR \left(1 + \sum_{j=1}^n \frac{1}{j}\right) \leq CrR(1 + \log n) \leq CrR \log \frac{e}{mh}.\end{aligned}\tag{5.8}$$

For the third inequality in (5.8), use (5.2), $\tau^{-1/2} = rR/\sqrt{c_1}$, and (5.1). The last two inequalities use $\sum_{j=1}^n j^{-1} \leq 1 + \log n$ and $n \leq \tau/h \leq 1/(mh)$, since $R^2 \geq m$ and $c_1 \leq 1$.

5.3 Proof of the flux bounds

Proof of Lemma 4.5. To estimate H , we express $\nabla \rho_t$ in terms of $\nabla \widehat{\pi}_{\lambda,h}$ and J . We first justify the commutation identity

$$\nabla(\mathbf{H}_t \widehat{\pi}_{\lambda,h}) = \mathbf{H}_t(\nabla \widehat{\pi}_{\lambda,h}), \quad t > 0. \quad (5.9)$$

By (5.8) and Hölder's inequality,

$$\int \|\nabla \widehat{\pi}_{\lambda,h}\| \, dx = \int \|\nabla \widehat{\pi}_{\lambda,h}/\pi_\lambda\| \, d\pi_\lambda \leq \|\nabla \widehat{\pi}_{\lambda,h}/\pi_\lambda\|_{L^{2r}(\pi_\lambda)} < \infty.$$

Thus both $\widehat{\pi}_{\lambda,h}$ and $\nabla \widehat{\pi}_{\lambda,h}$ belong to $L^1(\mathbb{R}^d)$. For each i , differentiation of the Gaussian kernel and integration by parts give

$$\begin{aligned} \partial_i(\mathbf{H}_t \widehat{\pi}_{\lambda,h})(x) &= \int \partial_{x_i} \left[\frac{e^{-\|x-y\|^2/(4t)}}{(4\pi t)^{d/2}} \right] \widehat{\pi}_{\lambda,h}(y) \, dy \\ &= - \int \partial_{y_i} \left[\frac{e^{-\|x-y\|^2/(4t)}}{(4\pi t)^{d/2}} \right] \widehat{\pi}_{\lambda,h}(y) \, dy \\ &= \int \frac{e^{-\|x-y\|^2/(4t)}}{(4\pi t)^{d/2}} \partial_i \widehat{\pi}_{\lambda,h}(y) \, dy = \mathbf{H}_t(\partial_i \widehat{\pi}_{\lambda,h})(x). \end{aligned}$$

For fixed $t > 0$, the kernel and its first derivatives are bounded. Differentiation under the integral therefore follows from $\widehat{\pi}_{\lambda,h} \in L^1$. The integration by parts is justified by the cutoff argument used for (5.3), now using $\widehat{\pi}_{\lambda,h}, \nabla \widehat{\pi}_{\lambda,h} \in L^1$. This proves (5.9).

For $0 < t \leq h$, the definition of ρ_t , (4.19), and (5.3) give

$$\rho_t = \mathbf{H}_t(\widehat{\pi}_{\lambda,h} + h \operatorname{div} J) = \mathbf{H}_t \widehat{\pi}_{\lambda,h} + h \operatorname{div} \mathbf{H}_t J.$$

Differentiating this identity between smooth functions and using (5.9), we obtain

$$\begin{aligned} \nabla \rho_t &= \nabla \mathbf{H}_t \widehat{\pi}_{\lambda,h} + h \nabla \operatorname{div} \mathbf{H}_t J \\ &= \mathbf{H}_t(\nabla \widehat{\pi}_{\lambda,h}) + h \nabla \operatorname{div} \mathbf{H}_t J, \quad 0 < t \leq h. \end{aligned} \quad (5.10)$$

Apply (4.7) to the vector field $\nabla \widehat{\pi}_{\lambda,h}$ and (4.9) to J , both with output order $s = r$. The time restrictions hold because $tr^2 R^2 \leq hr^2 R^2 \leq c_1/2$, as verified in Section 5.2. Using (5.2), (5.8), and $\|J/\pi_\lambda\|_{L^{4r}(\pi_\lambda)} \leq \|J/\pi_\lambda\|_{L^{8r}(\pi_\lambda)}$, we get

$$\begin{aligned} \|\nabla \rho_t/\pi_\lambda\|_{L^r(\pi_\lambda)} &\leq C \|\nabla \widehat{\pi}_{\lambda,h}/\pi_\lambda\|_{L^{2r}(\pi_\lambda)} + \frac{Ch\sqrt{r}}{t} \|J/\pi_\lambda\|_{L^{4r}(\pi_\lambda)} \\ &\leq CrR \log \frac{e}{mh} + \frac{ChrR}{t}, \quad 0 < t \leq h. \end{aligned}$$

The weight t in (4.13) cancels the apparent singularity at zero, proving (4.25). For (4.26), apply (4.10) with output order $s = 2$ and time $u \in [0, h]$. Its time restriction is weaker than the one already verified with output order $8r$ before (5.2). Using (2.3) and Hölder's inequality, we obtain

$$\begin{aligned} \|(T_u)_\#((\nabla U_\lambda \otimes \nabla U_\lambda) \widehat{\pi}_{\lambda,h})/\pi_\lambda\|_{L^2(\pi_\lambda)} &\leq C \|(\nabla U_\lambda \otimes \nabla U_\lambda) \widehat{\pi}_{\lambda,h}/\pi_\lambda\|_{L^4(\pi_\lambda)} \\ &\leq C \|\nabla U_\lambda\|_{L^{16}(\pi_\lambda)}^2 \|\widehat{\pi}_{\lambda,h}/\pi_\lambda\|_{L^8(\pi_\lambda)} \leq CR^2, \quad 0 \leq u \leq h. \end{aligned} \quad (5.11)$$

The last inequality follows from (4.2) at order 16 and (5.1). Minkowski's inequality in (4.14), followed by (5.11), now gives

$$\begin{aligned}\|E/\pi_\lambda\|_{L^2(\pi_\lambda)} &\leq \frac{1}{h} \int_0^h (h-u) \|(T_u)_\#((\nabla U_\lambda \otimes \nabla U_\lambda) \widehat{\pi}_{\lambda,h})/\pi_\lambda\|_{L^2(\pi_\lambda)} du \\ &\leq \frac{CR^2}{h} \int_0^h (h-u) du \leq ChR^2.\end{aligned}$$

This proves (4.26). Finally, $r \geq 4$ and (4.2) imply

$$\|\nabla U_\lambda \otimes H/\pi_\lambda\|_{L^2(\pi_\lambda)} \leq \|\nabla U_\lambda\|_{L^4(\pi_\lambda)} \|H/\pi_\lambda\|_{L^4(\pi_\lambda)} \leq CR \|H/\pi_\lambda\|_{L^r(\pi_\lambda)}.$$

Combining this with (4.26) proves (4.27). This proves (4.25), (4.26), and (4.27), completing the proof of Lemma 4.5. $\square$

6 Conclusion

We established a $\widetilde{O}(h)$ invariant-measure bias bound relative to the smoothed target under an explicit step-size condition, with only logarithmic dependence on λ^{-1} in the error coefficient. Combining this bound with Moreau approximation and Wasserstein contraction gives $\widetilde{O}(\varepsilon^{-1})$ iterations to achieve $\sqrt{m} W_2(\mu_N, \pi) \leq \varepsilon$ for fixed model parameters and initialization. These guarantees require no additional higher-order smoothness assumptions and leave the classical MYULA algorithm unchanged.

A Target Moments and Normalized Operator Estimates

A.1 Target drift moments

Proof of Lemma 4.1. The identity and bound in (4.1) follow from Xin and Zhang [9, Lemma 7.1 and Proposition A.4]. We prove the higher-moment and exponential estimates.

Since π_λ has Gaussian tails and ∇f has at most linear growth, all moments of ∇f are finite. Integration by parts with polynomially growing vector fields is justified by spatial cutoffs; see Section D.1.

For $s \geq 2$, apply integration by parts to the C^1 test field $\|\nabla f\|^{s-2} \nabla f$. Using $\nabla U_\lambda = \nabla f + \nabla g_\lambda$, we obtain

$$\begin{aligned}\int \|\nabla f\|^s d\pi_\lambda &= \int \|\nabla f\|^{s-2} \Delta f d\pi_\lambda \\ &\quad + (s-2) \int \|\nabla f\|^{s-4} (\nabla f)^\top \nabla^2 f \nabla f d\pi_\lambda \\ &\quad - \int \|\nabla f\|^{s-2} \langle \nabla f, \nabla g_\lambda \rangle d\pi_\lambda \\ &\leq [\tau_f + (s-2) \min(L_f, \tau_f)] \int \|\nabla f\|^{s-2} d\pi_\lambda \\ &\quad + G \int \|\nabla f\|^{s-1} d\pi_\lambda.\end{aligned}\tag{A.1}$$

The inequality uses $\Delta f \leq \tau_f$, $0 \preceq \nabla^2 f \preceq \min(L_f, \tau_f)I$, and $\|\nabla g_\lambda\| \leq G$. The term with coefficient $s-2$ is zero when $s=2$; for $s>2$, its integrand extends continuously by zero at $\nabla f=0$.

Hölder's inequality gives

$$\int \|\nabla f\|^{s-2} d\pi_\lambda \leq \|\nabla f\|_{L^s(\pi_\lambda)}^{s-2}, \quad \int \|\nabla f\|^{s-1} d\pi_\lambda \leq \|\nabla f\|_{L^s(\pi_\lambda)}^{s-1}.$$

If $\|\nabla f\|_{L^s(\pi_\lambda)} > 0$, substituting these bounds into (A.1) and dividing by $\|\nabla f\|_{L^s(\pi_\lambda)}^{s-2}$ yields

$$\|\nabla f\|_{L^s(\pi_\lambda)}^2 \leq \tau_f + (s-2) \min(L_f, \tau_f) + G \|\nabla f\|_{L^s(\pi_\lambda)}.$$

Solving this quadratic inequality gives

$$\|\nabla f\|_{L^s(\pi_\lambda)} \leq G + \sqrt{\tau_f + (s-2) \min(L_f, \tau_f)}. \quad (\text{A.2})$$

Since $\|\nabla g_\lambda\| \leq G$, $G \leq R$, and $\tau_f \leq R^2$, it follows that

$$\|\nabla U_\lambda\|_{L^s(\pi_\lambda)} \leq \|\nabla f\|_{L^s(\pi_\lambda)} + G \leq 2G + \sqrt{(s-1)\tau_f} \leq 3\sqrt{s}R,$$

which proves (4.2).

Finally, (4.2) gives $\int \|\nabla U_\lambda\|^{2k} d\pi_\lambda \leq (18kR^2)^k$. Expanding the exponential and using $k! \geq (k/e)^k$, we obtain

$$\int e^{\|\nabla U_\lambda\|^2/(256R^2)} d\pi_\lambda \leq 1 + \sum_{k \geq 1} (18e/256)^k \leq 2.$$

This proves (4.3). □

A.2 A Gibbs change of variables for the heat operator

This and the following two subsections prove Lemma 4.3. We use the exponential moment bound (4.3):

$$\int e^{\|\nabla U_\lambda\|^2/(256R^2)} d\pi_\lambda \leq 2. \quad (\text{A.3})$$

Let $X \sim \pi_\lambda$ and $Z_t \sim N(0, 2tI_d)$ be independent, and write φ_{2tI_d} for the density of Z_t . The Gibbs density satisfies

$$\pi_\lambda(x) = \pi_\lambda(y) e^{U_\lambda(y) - U_\lambda(x)}.$$

For every $a \geq 0$, independence and the substitution $y = x + z$ therefore give

$$\begin{aligned} & \mathbb{E} e^{a[U_\lambda(X+Z_t) - U_\lambda(X)]} \\ &= \iint \pi_\lambda(x) \varphi_{2tI_d}(y-x) e^{a[U_\lambda(y) - U_\lambda(x)]} dx dy \\ &= \iint \pi_\lambda(y) \varphi_{2tI_d}(y-x) e^{(a+1)[U_\lambda(y) - U_\lambda(x)]} dx dy \\ &\leq \int e^{(a+1)^2 t \|\nabla U_\lambda(y)\|^2} \pi_\lambda(y) dy. \end{aligned} \quad (\text{A.4})$$

For the last inequality, convexity gives

$$U_\lambda(y) - U_\lambda(x) \leq \langle \nabla U_\lambda(y), y - x \rangle,$$

and the Gaussian integral in x is

$$\int \varphi_{2tI_d}(y-x) e^{(a+1)\langle \nabla U_\lambda(y), y-x \rangle} dx = e^{(a+1)^2 t \|\nabla U_\lambda(y)\|^2}.$$

Thus, by (A.3), the expectation in (A.4) is at most 2 whenever $(a+1)^2 t R^2 \leq 1/256$.

Write $w = F/\pi_\lambda$. Since the heat kernel has total mass one, Jensen's inequality gives

$$\|\mathbf{H}_t F(y)\|^s = \left\| \int \varphi_{2tI_d}(y-x) F(x) dx \right\|^s \leq \int \varphi_{2tI_d}(y-x) \|F(x)\|^s dx.$$

Multiply by $\pi_\lambda(y)^{1-s}$ and integrate in y . Using Tonelli's theorem and $F(x) = \pi_\lambda(x)w(x)$, we obtain

$$\begin{aligned} \|\mathbf{H}_t F/\pi_\lambda\|_{L^s(\pi_\lambda)}^s &= \int \|\mathbf{H}_t F(y)\|^s \pi_\lambda(y)^{1-s} dy \\ &\leq \iint \varphi_{2tI_d}(y-x) \|F(x)\|^s \pi_\lambda(y)^{1-s} dx dy \\ &= \iint \pi_\lambda(x) \varphi_{2tI_d}(y-x) \|w(x)\|^s \left(\frac{\pi_\lambda(x)}{\pi_\lambda(y)} \right)^{s-1} dx dy \\ &= \mathbb{E} \left[\|w(X)\|^s e^{(s-1)[U_\lambda(X+Z_t)-U_\lambda(X)]} \right]. \end{aligned} \tag{A.5}$$

Cauchy–Schwarz then yields

$$\|\mathbf{H}_t F/\pi_\lambda\|_{L^s(\pi_\lambda)}^s \leq \|w\|_{L^{2s}(\pi_\lambda)}^s \left(\mathbb{E} e^{2(s-1)[U_\lambda(X+Z_t)-U_\lambda(X)]} \right)^{1/2}.$$

Apply (A.4) with $a = 2(s-1)$. Together with (A.3), this bounds the expectation by 2 whenever $(2s-1)^2 t R^2 \leq 1/256$. Since $2s-1 \leq 2s$, this condition follows from $ts^2 R^2 \leq c$ for a sufficiently small universal constant c . Taking s th roots proves (4.7). The argument applies to scalar, vector, and matrix densities, using the absolute value, Euclidean norm, and Frobenius norm, respectively.

A.3 Gaussian derivatives without an extra dimension factor

Let F be scalar with $\|F/\pi_\lambda\|_{L^{2s}(\pi_\lambda)} < \infty$. Since π_λ is a bounded probability density,

$$\int |F| dx \leq \|F/\pi_\lambda\|_{L^{2s}(\pi_\lambda)}, \quad \int |F|^2 dx \leq \|\pi_\lambda\|_\infty \|F/\pi_\lambda\|_{L^{2s}(\pi_\lambda)}^2.$$

For fixed $t > 0$, every derivative of the Gaussian kernel is bounded. Thus differentiation under the integral is justified by $F \in L^1(\mathbb{R}^d)$, and $\mathbf{H}_t F$ is smooth.

For a unit vector e , differentiating the kernel gives

$$\begin{aligned} e \cdot \nabla \mathbf{H}_t F(x) &= \int e \cdot \nabla_x \varphi_{2tI_d}(x-y) F(y) dy \\ &= - \int \frac{e \cdot (x-y)}{2t} \varphi_{2tI_d}(x-y) F(y) dy \\ &= - \mathbb{E} \left[\frac{e \cdot Z_t}{2t} F(x - Z_t) \right]. \end{aligned}$$

Using $\mathbb{E}(e \cdot Z_t)^2 = 2t$, Cauchy–Schwarz yields

$$\begin{aligned} \|\nabla \mathbf{H}_t F(x)\| &= \sup_{\|e\|=1} \left| \mathbb{E} \left[\frac{e \cdot Z_t}{2t} F(x - Z_t) \right] \right| \\ &\leq (2t)^{-1/2} (\mathbf{H}_t |F|^2(x))^{1/2}. \end{aligned}$$

For $s \geq 2$, Jensen then gives $\|\nabla \mathbf{H}_t F\|^s \leq (2t)^{-s/2} \mathbf{H}_t |F|^s$. Multiply by π_λ^{1-s} , integrate, and use the same endpoint exchange as in (A.5). This proves the scalar gradient estimate (4.8).

To estimate the divergence of a vector field, let $s' = s/(s-1)$ and $a = 2s/(2s-1)$ within this calculation. Initially take $\phi \in C_c^\infty(\mathbb{R}^d)$ with $\|\phi\|_{L^{s'}(\pi_\lambda)} \leq 1$. Integration by parts against the compactly supported ϕ and symmetry of the Gaussian kernel give

$$\begin{aligned} \int \phi \operatorname{div} \mathbf{H}_t F \, dx &= - \int \langle \nabla \phi, \mathbf{H}_t F \rangle \, dx \\ &= - \int \langle \mathbf{H}_t(\nabla \phi), F \rangle \, dx \\ &= - \int \langle \nabla \mathbf{H}_t \phi, F/\pi_\lambda \rangle \, d\pi_\lambda. \end{aligned}$$

The exchange of integrals is justified by $F \in L^1$ and the boundedness of $\nabla \phi$. The last equality also uses $\mathbf{H}_t(\nabla \phi) = \nabla \mathbf{H}_t \phi$. Since $1/(2s) + 1/a = 1$, Hölder's inequality with respect to π_λ yields

$$\left| \int \phi \operatorname{div} \mathbf{H}_t F \, dx \right| \leq \|F/\pi_\lambda\|_{L^{2s}(\pi_\lambda)} \|\nabla \mathbf{H}_t \phi\|_{L^a(\pi_\lambda)}. \quad (\text{A.6})$$

For every unit vector e , the scalar $e \cdot Z_t$ has distribution $N(0, 2t)$, and hence

$$(\mathbb{E}|e \cdot Z_t|^{2s})^{1/(2s)} \leq C\sqrt{st}.$$

Since $1/(2s) + 1/a = 1$, the derivative formula above and Hölder's inequality give

$$\begin{aligned} \|\nabla \mathbf{H}_t \phi(x)\| &= \sup_{\|e\|=1} \left| \mathbb{E} \left[\frac{e \cdot Z_t}{2t} \phi(x - Z_t) \right] \right| \\ &\leq \frac{1}{2t} \sup_{\|e\|=1} (\mathbb{E}|e \cdot Z_t|^{2s})^{1/(2s)} (\mathbb{E}|\phi(x - Z_t)|^a)^{1/a} \\ &\leq C\sqrt{s/t} (\mathbf{H}_t |\phi|^a(x))^{1/a}. \end{aligned}$$

To integrate this bound, use symmetry of the heat kernel and Hölder with conjugate exponents $s'/a = (2s-1)/(2s-2)$ and $2s-1$:

$$\begin{aligned} \int \mathbf{H}_t |\phi|^a \, d\pi_\lambda &= \int |\phi|^a \frac{\mathbf{H}_t \pi_\lambda}{\pi_\lambda} \, d\pi_\lambda \\ &\leq \|\phi\|_{L^{s'}(\pi_\lambda)}^a \|\mathbf{H}_t \pi_\lambda / \pi_\lambda\|_{L^{2s-1}(\pi_\lambda)}. \end{aligned}$$

Consequently,

$$\|\nabla \mathbf{H}_t \phi\|_{L^a(\pi_\lambda)} \leq C\sqrt{s/t} \|\phi\|_{L^{s'}(\pi_\lambda)} \|\mathbf{H}_t \pi_\lambda / \pi_\lambda\|_{L^{2s-1}(\pi_\lambda)}^{1/a} \leq C\sqrt{s/t}.$$

The last inequality uses $\|\phi\|_{L^{s'}(\pi_\lambda)} \leq 1$ and (4.7) with input π_λ and order $2s-1$. Since $2s-1 \leq 2s$, its time restriction follows from $ts^2 R^2 \leq c$ after decreasing the universal constant c . Duality and density of $C_c^\infty(\mathbb{R}^d)$ in $L^{s'}(\pi_\lambda)$ therefore yield

$$\|\operatorname{div} \mathbf{H}_t F / \pi_\lambda\|_{L^s(\pi_\lambda)} \leq C\sqrt{s/t} \|F/\pi_\lambda\|_{L^{2s}(\pi_\lambda)}. \quad (\text{A.7})$$

Finally, $\nabla \operatorname{div} \mathbf{H}_t = \nabla \mathbf{H}_{t/2} \circ \operatorname{div} \mathbf{H}_{t/2}$. Apply the scalar gradient bound and then (A.7), with order $2s$ in the latter. The result is (4.9).

A.4 Pushforward estimates by an integrated Jacobian

Recall that $0 \preceq \nabla^2 U_\lambda \preceq L_\lambda I$ almost everywhere. For $u \geq 0$, the expansion map $S_u(x) = x + u\nabla U_\lambda(x)$ is globally bi-Lipschitz. Indeed, $S_u^{-1}y$ is the unique minimizer of the strongly convex function $x \mapsto \frac{1}{2} \|x - y\|^2 + uU_\lambda(x)$; monotonicity of ∇U_λ gives a Lipschitz inverse. For $u > 0$, set $y = S_u(x) = x + u\nabla U_\lambda(x)$. Monotonicity of ∇U_λ gives

$$\|\nabla U_\lambda(x)\|^2 \leq \langle \nabla U_\lambda(y), \nabla U_\lambda(x) \rangle \leq \|\nabla U_\lambda(y)\| \|\nabla U_\lambda(x)\|,$$

so $\|\nabla U_\lambda(x)\| \leq \|\nabla U_\lambda(y)\|$. Convexity therefore implies

$$U_\lambda(y) - U_\lambda(x) \leq \langle \nabla U_\lambda(y), y - x \rangle = u \langle \nabla U_\lambda(y), \nabla U_\lambda(x) \rangle \leq u \|\nabla U_\lambda(y)\|^2.$$

Changing variables $y = S_u(x)$ and using the Gibbs density ratio, we obtain

$$\begin{aligned} \int \det(I + u\nabla^2 U_\lambda(x)) \pi_\lambda(x) dx &= \int \pi_\lambda(S_u^{-1}y) dy \\ &= \int e^{U_\lambda(y) - U_\lambda(S_u^{-1}y)} d\pi_\lambda(y) \\ &\leq \int e^{u\|\nabla U_\lambda(y)\|^2} d\pi_\lambda(y). \end{aligned} \tag{A.8}$$

The case $u = 0$ is immediate. By (A.3), the last integral is at most 2 whenever $uR^2 \leq 1/256$.

Under $st(L_\lambda + R^2) \leq c$, one has $tL_\lambda < 1$, so the contraction map $T_t = I - t\nabla U_\lambda$ is bi-Lipschitz by Proposition 2.1. Write $J_t(x) = \det(I - t\nabla^2 U_\lambda(x)) > 0$ almost everywhere. Using (2.11) and changing variables $y = T_t x$, we obtain

$$\begin{aligned} \|(T_t)_\# F / \pi_\lambda\|_{L^s(\pi_\lambda)}^s &= \int \left\| \frac{F(x)}{J_t(x)} \right\|^s \pi_\lambda(T_t x)^{1-s} J_t(x) dx \\ &= \int \left\| \frac{F(x)}{\pi_\lambda(x)} \right\|^s e^{(s-1)[U_\lambda(T_t x) - U_\lambda(x)]} J_t(x)^{1-s} d\pi_\lambda(x). \end{aligned} \tag{A.9}$$

Since $tL_\lambda < 1$, the smooth descent inequality gives

$$U_\lambda(T_t x) \leq U_\lambda(x) - t \left(1 - \frac{tL_\lambda}{2} \right) \|\nabla U_\lambda(x)\|^2 \leq U_\lambda(x).$$

Thus the exponential factor in (A.9) is at most one, and Cauchy–Schwarz yields

$$\|(T_t)_\# F / \pi_\lambda\|_{L^s(\pi_\lambda)}^s \leq \|F / \pi_\lambda\|_{L^{2s}(\pi_\lambda)}^s \left(\int J_t^{-2(s-1)} d\pi_\lambda \right)^{1/2}.$$

For $k \geq 1$ and $0 \leq z \leq 1/(4k)$, Bernoulli's inequality gives $(1 - z)^k \geq 1 - kz$, and hence

$$(1 - z)^{-k} \leq \frac{1}{1 - kz} = 1 + \frac{kz}{1 - kz} \leq 1 + 2kz.$$

Set $k = 2(s - 1)$. For almost every x , let $\mu_1, \dots, \mu_d$ be the eigenvalues of $\nabla^2 U_\lambda(x)$. If $ktL_\lambda \leq 1/4$, applying the preceding inequality to $z = t\mu_i$ gives

$$J_t(x)^{-k} = \prod_{i=1}^d (1 - t\mu_i)^{-k} \leq \prod_{i=1}^d (1 + 2kt\mu_i) = \det(I + 2kt\nabla^2 U_\lambda(x)).$$

Since $k \leq 2s$, the condition $st(L_\lambda + R^2) \leq c$ ensures both $ktL_\lambda \leq 1/4$ and $2ktR^2 \leq 1/256$ when c is sufficiently small. Applying (A.8) with $u = 2kt$ therefore yields

$$\int J_t^{-k} d\pi_\lambda \leq \int \det(I + 2kt\nabla^2 U_\lambda) d\pi_\lambda \leq 2.$$

Combining this with the preceding Cauchy–Schwarz bound gives

$$\|(T_t)_\# F/\pi_\lambda\|_{L^s(\pi_\lambda)} \leq 2^{1/(2s)} \|F/\pi_\lambda\|_{L^{2s}(\pi_\lambda)},$$

which proves (4.10). The changes of variables above are justified by the area formula for bi-Lipschitz maps; see Proposition 2.1. This completes the proof of Lemma 4.3.

B Finite-Order Stationary Density Ratios

This appendix proves the stationary density-ratio estimate in Lemma 4.2. The goal is to control

$$\sup_{0 \leq t \leq h} \|\nu_t/\pi_\lambda\|_{L^\alpha(\pi_\lambda)}$$

under the step restrictions (4.4).

The proof has three steps. In Section B.1, we establish the weighted drift and curvature moments needed for the calculation. In Section B.2, these moments give the drift-increment bound (B.7). In Section B.3, we insert that bound into the Rényi dissipation identity and use the stationary endpoint condition to obtain (B.12).

The integrability needed for differentiation and integration by parts is stated in Lemma B.1 and proved independently in Section D.2. The regularization and limiting arguments are supplied in Appendix D. The interpolation method follows the standard Rényi analysis of Langevin algorithms [2].

B.1 Weighted energy for the density ratio

Fix $\alpha \geq 2$ and λ, h satisfying the restrictions of Lemma 4.2. We first work with an auxiliary smooth regularization. Let $\varrho \in C_c^\infty(\mathbb{R}^d)$ be nonnegative and symmetric, with $\int \varrho dx = 1$, and set

$$\varrho_\delta(x) := \delta^{-d} \varrho(x/\delta), \quad f^{(\delta)} := f * \varrho_\delta, \quad g_\lambda^{(\delta)} := g_\lambda * \varrho_\delta, \quad \delta > 0.$$

Define

$$U_\lambda^{(\delta)} := f^{(\delta)} + g_\lambda^{(\delta)}, \quad \pi_\lambda^{(\delta)} \propto e^{-U_\lambda^{(\delta)}}, \quad T_t^{(\delta)}(x) := x - t\nabla U_\lambda^{(\delta)}(x).$$

The parameter δ is an auxiliary convolution scale, distinct from the Moreau parameter λ . It will tend to zero with λ and h fixed.

Let $Q_{\lambda,h}^{(\delta)}$ be the Euler kernel associated with $U_\lambda^{(\delta)}$, and let $\hat{\pi}_{\lambda,h}^{(\delta)}$ be its invariant density. The corresponding stationary interpolation is

$$\nu_t^{(\delta)} := H_t((T_t^{(\delta)})_\# \hat{\pi}_{\lambda,h}^{(\delta)}), \quad \nu_0^{(\delta)} = \nu_h^{(\delta)} = \hat{\pi}_{\lambda,h}^{(\delta)}.$$

The structural bounds needed below hold with the same parameters m, L_f, τ_f, G , and λ ; their preservation is verified in Section D.1.

For the calculation below, fix $\delta > 0$ and omit the superscript (δ) from the potentials and all associated densities, kernels, and maps. In particular, $\hat{\pi}_{\lambda,h}$ below denotes the invariant density of

the regularized Euler kernel. We first establish the integrability needed for the calculation at this fixed δ . We then obtain a quantitative estimate independent of δ , and pass to the original potential in [Section D.3](#).

Let $X \sim \hat{\pi}_{\lambda,h}$ and $Z \sim N(0, I_d)$ be independent, and, for $0 \leq t \leq h$, let

$$Y = X - t \nabla U_\lambda(X) + \sqrt{2t} Z.$$

The law of Y is ν_t . In this appendix only, suppressing the time argument, define

$$u := \frac{\nu_t}{\pi_\lambda}, \quad s := \nabla \log u, \\ e_t(y) := \nabla U_\lambda(y) - \mathbb{E}[\nabla U_\lambda(X) \mid Y = y].$$

The density ratio u is the quantity whose α -moment we seek to control. Its relative score $s = \nabla \log u$ appears in the dissipation of that moment. The field e_t measures the discrepancy between the current drift $\nabla U_\lambda(y)$ and the drift frozen at the starting point X . The mixed term involving e_t and s will be estimated in [Section B.3](#) using the drift-increment bound proved in [Section B.2](#).

Before carrying out these calculations, we record the integrability statement that makes them legitimate.

Lemma B.1 (Integrability before dissipation). *Consider the regularized interpolation defined above, with $\delta > 0$ fixed. Under the restrictions of [Lemma 4.2](#), ν_t is positive, locally $C^{1,2}$ in (t, y) for $0 < t < h$, and pointwise continuous at both time endpoints. Moreover,*

$$\sup_{0 \leq t \leq h} \int [u^\alpha (1 + \|s\|^2 + \|e_t\|^2) + u^{3\alpha/2}] d\pi_\lambda < \infty.$$

For every integer $k \geq 0$, one also has

$$\sup_{0 \leq t \leq h} \mathbb{E} \left[u(Y)^{\alpha-1} (1 + \|X\| + \|Y\|)^k \right] < \infty.$$

These bounds are uniform in $t \in [0, h]$. Their finite values may depend on the fixed parameters and on δ ; no uniform bound as $\delta \downarrow 0$ is asserted in this lemma.

The proof of [Lemma B.1](#) is given in [Section D.2](#). It establishes the required integrability independently of the dissipation calculation. The finite bounds from that proof justify the operations below; their numerical values do not enter the quantitative estimate.

We now prepare two moment bounds for the calculation in [Section B.2](#): an L^2 bound on ∇f and a bound on the mean of Δg_λ under the appropriate density-ratio weight. We define

$$\omega(dx) := \frac{u(x)^\alpha \pi_\lambda(x) dx}{\int u^\alpha d\pi_\lambda}, \quad I := \mathbb{E}_\omega \|s\|^2. \quad (\text{B.1})$$

A weighted moment of ∇f . Recall that $\nabla U_\lambda = \nabla f + \nabla g_\lambda$ and $\nabla \log \omega = \alpha s - \nabla U_\lambda$. Integration by parts gives

$$\begin{aligned} \mathbb{E}_\omega \Delta f &= -\mathbb{E}_\omega \langle \nabla f, \nabla \log \omega \rangle \\ &= \mathbb{E}_\omega \|\nabla f\|^2 + \mathbb{E}_\omega \langle \nabla f, \nabla g_\lambda \rangle - \alpha \mathbb{E}_\omega \langle \nabla f, s \rangle. \end{aligned}$$

The integration by parts is justified by spatial cutoffs and the moment and score bounds in [Lemma B.1](#). Rearranging the preceding identity, using $\Delta f \leq \tau_f$ and $\|\nabla g_\lambda\| \leq G$, and applying Cauchy–Schwarz, we obtain

$$\begin{aligned}\|\nabla f\|_{L^2(\omega)}^2 &= \mathbb{E}_\omega \Delta f - \mathbb{E}_\omega \langle \nabla f, \nabla g_\lambda \rangle + \alpha \mathbb{E}_\omega \langle \nabla f, s \rangle \\ &\leq \tau_f + G \mathbb{E}_\omega \|\nabla f\| + \alpha \|\nabla f\|_{L^2(\omega)} \sqrt{I} \\ &\leq \tau_f + (G + \alpha \sqrt{I}) \|\nabla f\|_{L^2(\omega)}.\end{aligned}$$

Using $R^2 = \tau_f + GR$, we solve this inequality and obtain

$$\|\nabla f\|_{L^2(\omega)} \leq R + \alpha \sqrt{I}. \quad (\text{B.2})$$

The weighted mean of the Moreau curvature. We next estimate $\mathbb{E}_\omega \Delta g_\lambda$. Applying integration by parts directly to ∇g_λ gives

$$\begin{aligned}\mathbb{E}_\omega \Delta g_\lambda &= -\mathbb{E}_\omega \langle \nabla g_\lambda, \nabla \log \omega \rangle \\ &= \mathbb{E}_\omega \langle \nabla g_\lambda, \nabla f + \nabla g_\lambda - \alpha s \rangle \\ &\leq G \|\nabla f\|_{L^2(\omega)} + G^2 + \alpha G \sqrt{I} \\ &\leq GR + G^2 + 2\alpha G \sqrt{I} \\ &\leq 2GR + 2\alpha G \sqrt{I}.\end{aligned} \quad (\text{B.3})$$

The first inequality uses $\|\nabla g_\lambda\| \leq G$ and Cauchy–Schwarz. The next line uses [\(B.2\)](#), and the last uses $G^2 \leq GR$.

B.2 Gaussian integration by parts with the density-ratio weight

This subsection proves [\(B.7\)](#).

For an integrable function $\Phi(X, Y)$, define the weighted joint expectation

$$\mathbb{E}_\alpha[\Phi(X, Y)] := \frac{\mathbb{E}[u(Y)^{\alpha-1} \Phi(X, Y)]}{\int u^\alpha d\pi_\lambda}.$$

Since Y has density $\nu_t = u\pi_\lambda$, the denominator normalizes this expectation, and its Y marginal is ω :

$$\mathbb{E}_\alpha[\phi(Y)] = \frac{\int \phi(y) u(y)^\alpha \pi_\lambda(y) dy}{\int u^\alpha d\pi_\lambda} = \mathbb{E}_\omega \phi$$

for every integrable test function ϕ .

Fix $0 < t \leq h$. Let w be the gradient of a convex function with L -Lipschitz gradient, and write

$$\Delta w := w(Y) - w(X).$$

Cocoercivity and $Y - X = -t \nabla U_\lambda(X) + \sqrt{2t} Z$ give

$$\mathbb{E}_\alpha \|\Delta w\|^2 \leq -tL \mathbb{E}_\alpha \langle \Delta w, \nabla U_\lambda(X) \rangle + L \mathbb{E}_\alpha \langle \Delta w, \sqrt{2t} Z \rangle.$$

To evaluate the Gaussian term, condition on X and integrate by parts under the original standard Gaussian law of Z , keeping $u(Y)^{\alpha-1}$ in the integrand. The identities

$$\partial_{Z_i} Y_j = \sqrt{2t} \delta_{ij}, \quad \nabla_y u(y)^{\alpha-1} = (\alpha-1)u(y)^{\alpha-1} s(y)$$

yield

$$\mathbb{E}_\alpha \left\langle \Delta w, \sqrt{2t} Z \right\rangle = 2t \mathbb{E}_\omega \operatorname{div} w + 2t(\alpha - 1) \mathbb{E}_\alpha \langle \Delta w, s(Y) \rangle.$$

The joint weighted moment bound with $k = 2$ in [Lemma B.1](#) and $\|\Delta w\| \leq L \|Y - X\|$ give $\mathbb{E}_\alpha \|\Delta w\|^2 < \infty$. The same lemma gives $I < \infty$. Since the Y marginal is ω , Cauchy–Schwarz yields

$$\mathbb{E}_\alpha |\langle \Delta w, s(Y) \rangle| \leq \left(\mathbb{E}_\alpha \|\Delta w\|^2 \right)^{1/2} \sqrt{I} < \infty.$$

The remaining terms are integrable by the joint moment bound and the bounded derivative of w . These bounds justify removing the cutoffs in the Gaussian integration by parts.

Substitution gives

$$\begin{aligned} \mathbb{E}_\alpha \|w(Y) - w(X)\|^2 &\leq tL \left[-\mathbb{E}_\alpha \langle w(Y) - w(X), \nabla U_\lambda(X) \rangle + 2\mathbb{E}_\omega \operatorname{div} w \right. \\ &\quad \left. + 2(\alpha - 1) \mathbb{E}_\alpha \langle w(Y) - w(X), s(Y) \rangle \right]. \end{aligned} \quad (\text{B.4})$$

The smooth component. We first estimate the squared increment of ∇f . Define

$$D_f := \mathbb{E}_\alpha \|\nabla f(Y) - \nabla f(X)\|^2.$$

To use [\(B.4\)](#), we also need a weighted second moment of $\nabla U_\lambda(X)$. The identity

$$\nabla U_\lambda(X) = \nabla f(Y) - (\nabla f(Y) - \nabla f(X)) + \nabla g_\lambda(X)$$

and the fact that the Y marginal of $\mathbb{E}_\alpha$ is ω give

$$\left(\mathbb{E}_\alpha \|\nabla U_\lambda(X)\|^2 \right)^{1/2} \leq \|\nabla f\|_{L^2(\omega)} + \sqrt{D_f} + G \leq 2R + \alpha\sqrt{I} + \sqrt{D_f},$$

where the last line uses [\(B.2\)](#) and $G \leq R$.

Apply [\(B.4\)](#) with $w = \nabla f$ and $L = L_f$. Using $\operatorname{div}(\nabla f) = \Delta f \leq \tau_f$ and Cauchy–Schwarz, we obtain

$$\begin{aligned} D_f &\leq tL_f \left[\sqrt{D_f}(2R + \alpha\sqrt{I} + \sqrt{D_f}) + 2\tau_f + 2(\alpha - 1)\sqrt{D_f}\sqrt{I} \right] \\ &\leq tL_f \left[D_f + (2R + 3\alpha\sqrt{I})\sqrt{D_f} + 2\tau_f \right]. \end{aligned}$$

Young’s inequality gives

$$tL_f(2R + 3\alpha\sqrt{I})\sqrt{D_f} \leq \frac{1}{4}D_f + 8t^2L_f^2R^2 + 18t^2L_f^2\alpha^2I.$$

For $tL_f \leq 1/4$, moving the terms involving D_f to the left therefore yields

$$\frac{1}{2}D_f \leq 2tL_f\tau_f + 8t^2L_f^2R^2 + 18t^2L_f^2\alpha^2I.$$

In particular, we may use

$$D_f \leq 6tL_f\tau_f + 16t^2L_f^2R^2 + 36t^2L_f^2\alpha^2I. \quad (\text{B.5})$$

Since $\tau_f \leq R^2$ and $tL_f \leq 1/4$, the preceding estimates also give

$$\left(\mathbb{E}_\alpha \|\nabla U_\lambda(X)\|^2 \right)^{1/2} \leq 6R + 3\alpha\sqrt{I}.$$

The Moreau component. We next estimate

$$D_g := \mathbb{E}_\alpha \|\nabla g_\lambda(Y) - \nabla g_\lambda(X)\|^2.$$

Apply (B.4) with $w = \nabla g_\lambda$ and $L = \lambda^{-1}$. The bounded-gradient property gives

$$\|\nabla g_\lambda(Y) - \nabla g_\lambda(X)\| \leq 2G.$$

It follows that

$$\begin{aligned} D_g &\leq \frac{t}{\lambda} \left[2G(\mathbb{E}_\alpha \|\nabla U_\lambda(X)\|^2)^{1/2} + 2\mathbb{E}_\omega \Delta g_\lambda + 4(\alpha - 1)G\sqrt{I} \right] \\ &\leq \frac{t}{\lambda} \left[2G(6R + 3\alpha\sqrt{I}) + 2(2GR + 2\alpha G\sqrt{I}) + 4(\alpha - 1)G\sqrt{I} \right]. \end{aligned}$$

The second line uses the bound above and (B.3). Collecting terms gives

$$D_g \leq \frac{t}{\lambda} (16GR + 14\alpha G\sqrt{I}). \quad (\text{B.6})$$

Combining the two components. Since $\nabla U_\lambda = \nabla f + \nabla g_\lambda$,

$$\mathbb{E}_\alpha \|\nabla U_\lambda(Y) - \nabla U_\lambda(X)\|^2 \leq 2D_f + 2D_g.$$

Equations (B.5) and (B.6) therefore yield

$$\begin{aligned} \mathbb{E}_\alpha \|\nabla U_\lambda(Y) - \nabla U_\lambda(X)\|^2 &\leq 12tL_f\tau_f + 32t^2L_f^2R^2 + 72t^2L_f^2\alpha^2I \\ &\quad + 32(t/\lambda)GR + 28(t/\lambda)\alpha G\sqrt{I}. \end{aligned} \quad (\text{B.7})$$

B.3 Proof of the density-ratio bound

This subsection proves (B.12), which implies (4.6).

Step 1. Evolution equation for ν_t : (B.8). Recall that

$$Y = X - t\nabla U_\lambda(X) + \sqrt{2t}Z.$$

For $\phi \in C_c^\infty(\mathbb{R}^d)$ and $0 < t < h$, differentiation under the expectation gives

$$\frac{d}{dt} \int \phi(y) \nu_t(y) dy = \mathbb{E} \left\langle \nabla \phi(Y), -\nabla U_\lambda(X) + Z/\sqrt{2t} \right\rangle.$$

This differentiation is valid on compact subintervals of $(0, h)$ because $\nabla \phi$ is bounded, the drift grows at most linearly, and X has finite moments by Lemma B.1.

For each coordinate i , Gaussian integration by parts, first conditional on X , gives

$$\mathbb{E}[Z_i \partial_i \phi(Y)] = \sqrt{2t} \mathbb{E}[\partial_{ii} \phi(Y)].$$

Consequently,

$$\begin{aligned} \frac{d}{dt} \int \phi(y) \nu_t(y) dy &= \mathbb{E}[\Delta \phi(Y) - \langle \nabla U_\lambda(X), \nabla \phi(Y) \rangle] \\ &= \int [\Delta \phi(y) - \langle \nabla U_\lambda(y) - e_t(y), \nabla \phi(y) \rangle] \nu_t(y) dy, \end{aligned}$$

where the second equality uses

$$\mathbb{E}[\nabla U_\lambda(X) \mid Y = y] = \nabla U_\lambda(y) - e_t(y).$$

Thus, in the weak sense,

$$\partial_t \nu_t = \Delta \nu_t + \operatorname{div}(\nu_t \nabla U_\lambda) - \operatorname{div}(\nu_t e_t).$$

Since $\nabla \log \pi_\lambda = -\nabla U_\lambda$, we have

$$\nabla \nu_t + \nu_t \nabla U_\lambda = \nu_t \nabla \log \frac{\nu_t}{\pi_\lambda}.$$

This proves

$$\partial_t \nu_t = \operatorname{div} \left(\nu_t \nabla \log \frac{\nu_t}{\pi_\lambda} \right) - \operatorname{div}(\nu_t e_t). \quad (\text{B.8})$$

Step 2. Derivative of the Rényi divergence: (B.9). By (2.27), with $u = \nu_t/\pi_\lambda$,

$$\mathcal{R}_\alpha(\nu_t \parallel \pi_\lambda) = \frac{1}{\alpha - 1} \log \int u^\alpha \, d\pi_\lambda.$$

We first compute the derivative of the integral on the right-hand side.

Choose $\chi_n \in C_c^\infty(\mathbb{R}^d)$ such that $0 \leq \chi_n \leq 1$, $\chi_n = 1$ on the ball of radius n , $\chi_n = 0$ outside the ball of radius $2n$, and $\|\nabla \chi_n\|_\infty \leq C/n$. Fix $0 < t_0 < t_1 < h$. For each fixed $t \in [t_0, t_1]$, Step 1 gives

$$\int \phi \partial_t \nu_t \, dx = - \int \langle \nabla \phi, s - e_t \rangle \nu_t \, dx, \quad \phi \in C_c^\infty(\mathbb{R}^d).$$

By Lemma B.1, u is positive and locally $C^{1,2}$, $\partial_t \nu_t$ is locally continuous, and $\nu_t(s - e_t)$ is locally integrable. Approximation in C^1 , with support in a fixed compact set, therefore extends this identity to $\phi \in C_c^1(\mathbb{R}^d)$.

At this fixed time, we may take $\phi = \chi_n u^{\alpha-1}$. Applying the chain rule first, and then the spatial identity above, gives

$$\begin{aligned} \frac{d}{dt} \int \chi_n u^\alpha \, d\pi_\lambda &= \alpha \int \chi_n u^{\alpha-1} \partial_t \nu_t \, dx \\ &= -\alpha \int \langle \nabla(\chi_n u^{\alpha-1}), s - e_t \rangle \nu_t \, dx. \end{aligned}$$

Since

$$\nabla(\chi_n u^{\alpha-1}) = u^{\alpha-1} \nabla \chi_n + (\alpha - 1) \chi_n u^{\alpha-2} \nabla u,$$

we obtain

$$\begin{aligned} \frac{d}{dt} \int \chi_n u^\alpha \, d\pi_\lambda &= -\alpha(\alpha - 1) \int \chi_n u^\alpha \|s\|^2 \, d\pi_\lambda \\ &\quad + \alpha(\alpha - 1) \int \chi_n u^\alpha \langle e_t, s \rangle \, d\pi_\lambda \\ &\quad - \alpha \int u^\alpha \langle s - e_t, \nabla \chi_n \rangle \, d\pi_\lambda. \end{aligned}$$

Keep t_0 and t_1 fixed and integrate the preceding identity in time. Writing $u(t_j) = \nu_{t_j}/\pi_\lambda$ at the endpoints gives

$$\begin{aligned} & \int \chi_n u(t_1)^\alpha d\pi_\lambda - \int \chi_n u(t_0)^\alpha d\pi_\lambda \\ &= -\alpha(\alpha-1) \int_{t_0}^{t_1} \int \chi_n u^\alpha \|s\|^2 d\pi_\lambda dt \\ &+ \alpha(\alpha-1) \int_{t_0}^{t_1} \int \chi_n u^\alpha \langle e_t, s \rangle d\pi_\lambda dt \\ &- \alpha \int_{t_0}^{t_1} \int u^\alpha \langle s - e_t, \nabla \chi_n \rangle d\pi_\lambda dt. \end{aligned}$$

We now remove the spatial cutoff while keeping t_0 and t_1 fixed. The last term satisfies

$$\alpha \left| \int_{t_0}^{t_1} \int u^\alpha \langle s - e_t, \nabla \chi_n \rangle d\pi_\lambda dt \right| \leq \frac{C\alpha}{n} \int_{t_0}^{t_1} \int u^\alpha (1 + \|s\|^2 + \|e_t\|^2) d\pi_\lambda dt.$$

The time integral on the right is finite and independent of n by [Lemma B.1](#). Moreover,

$$|u^\alpha \langle e_t, s \rangle| \leq \frac{1}{2} u^\alpha (\|e_t\|^2 + \|s\|^2),$$

so the same lemma provides an integrable bound for each of the other time integrands. We may therefore let $n \rightarrow \infty$ by dominated convergence, obtaining

$$\begin{aligned} & \int u(t_1)^\alpha d\pi_\lambda - \int u(t_0)^\alpha d\pi_\lambda \\ &= -\alpha(\alpha-1) \int_{t_0}^{t_1} \int u^\alpha \|s\|^2 d\pi_\lambda dt \\ &+ \alpha(\alpha-1) \int_{t_0}^{t_1} \int u^\alpha \langle e_t, s \rangle d\pi_\lambda dt. \end{aligned}$$

It remains to include the time endpoints. By [Lemma B.1](#), both time integrands in the last identity belong to $L^1(0, h)$. The endpoint limits [\(D.10\)](#), proved in [Section D.2](#), show that $t \mapsto \int u(t)^\alpha d\pi_\lambda$ is continuous at $t = 0$ and $t = h$. We can therefore let $t_0 \downarrow 0$ or $t_1 \uparrow h$ in this identity, and then include both endpoints. The same identity consequently holds for every $0 \leq t_0 < t_1 \leq h$.

Since its right-hand side is the time integral of an $L^1(0, h)$ function, this identity proves that $t \mapsto \int u(t)^\alpha d\pi_\lambda$ is absolutely continuous on $[0, h]$. Its derivative is therefore

$$\frac{d}{dt} \int u^\alpha d\pi_\lambda = -\alpha(\alpha-1) \int u^\alpha \|s\|^2 d\pi_\lambda + \alpha(\alpha-1) \int u^\alpha \langle e_t, s \rangle d\pi_\lambda$$

for almost every $t \in (0, h)$.

Because $\int u^\alpha d\pi_\lambda \geq 1$, the Rényi divergence is also absolutely continuous on $[0, h]$. The chain rule gives

$$\frac{d}{dt} \mathcal{R}_\alpha(\nu_t \| \pi_\lambda) = \frac{\frac{d}{dt} \int u^\alpha d\pi_\lambda}{(\alpha-1) \int u^\alpha d\pi_\lambda} = -\alpha I + \alpha \mathbb{E}_\omega \langle e_t, s \rangle.$$

The weight defining $\mathbb{E}_\alpha$ depends only on Y , so it leaves the conditional law of X given Y unchanged. Hence

$$e_t(Y) = \mathbb{E}_\alpha[\nabla U_\lambda(Y) - \nabla U_\lambda(X) \mid Y].$$

Conditional Jensen's inequality and the fact that the Y marginal is ω give

$$\mathbb{E}_\omega \|e_t\|^2 \leq \mathbb{E}_\alpha \|\nabla U_\lambda(Y) - \nabla U_\lambda(X)\|^2.$$

Using

$$\langle e_t, s \rangle \leq \frac{1}{2} \|s\|^2 + \frac{1}{2} \|e_t\|^2,$$

we conclude that, for almost every t ,

$$\begin{aligned} \frac{d}{dt} \mathcal{R}_\alpha(\nu_t \| \pi_\lambda) &= -\alpha I + \alpha \mathbb{E}_\omega \langle e_t, s \rangle \\ &\leq -\frac{\alpha}{2} I + \frac{\alpha}{2} \mathbb{E}_\omega \|e_t\|^2 \\ &\leq -\frac{\alpha}{2} I + \frac{\alpha}{2} \mathbb{E}_\alpha \|\nabla U_\lambda(Y) - \nabla U_\lambda(X)\|^2. \end{aligned} \tag{B.9}$$

Step 3. Applying the drift bound: (B.10). Substituting (B.7) into (B.9) gives

$$\begin{aligned} \frac{d}{dt} \mathcal{R}_\alpha(\nu_t \| \pi_\lambda) &\leq -\frac{\alpha}{2} I + 6\alpha t L_f \tau_f + 16\alpha t^2 L_f^2 R^2 \\ &\quad + 36\alpha^3 t^2 L_f^2 I + 16\alpha(t/\lambda)GR + 14\alpha^2(t/\lambda)G\sqrt{I}. \end{aligned}$$

Since $\tau_f \geq m$, the first condition in (4.4) implies $hL_f \leq c/\alpha^2$. Taking the universal constant c small enough gives, for $0 \leq t \leq h$,

$$36\alpha^3 t^2 L_f^2 I \leq \frac{\alpha}{8} I.$$

Also, Young's inequality gives

$$14\alpha^2(t/\lambda)G\sqrt{I} \leq \frac{\alpha}{8} I + 392\alpha^3(t/\lambda)^2 G^2.$$

Combining these estimates proves

$$\frac{d}{dt} \mathcal{R}_\alpha(\nu_t \| \pi_\lambda) \leq -\frac{\alpha}{4} I + C\alpha \left[tL_f \tau_f + t^2 L_f^2 R^2 + (t/\lambda)GR + \alpha^2(t/\lambda)^2 G^2 \right]. \tag{B.10}$$

Step 4. A lower bound for I : (B.11). By Lemma B.1,

$$\int u^\alpha d\pi_\lambda < \infty, \quad \int \left\| \nabla u^{\alpha/2} \right\|^2 d\pi_\lambda = \frac{\alpha^2}{4} \int u^\alpha \|s\|^2 d\pi_\lambda < \infty.$$

We may therefore apply (2.26) to $w = u^{\alpha/2}$. Dividing by $\int u^\alpha d\pi_\lambda$ gives

$$\frac{\text{Ent}_{\pi_\lambda}(u^\alpha)}{\int u^\alpha d\pi_\lambda} \leq \frac{2}{m} \frac{\int \left\| \nabla u^{\alpha/2} \right\|^2 d\pi_\lambda}{\int u^\alpha d\pi_\lambda} = \frac{\alpha^2}{2m} I.$$

We next bound the left-hand side from below. Since $\int u d\pi_\lambda = 1$, the definition of ω gives

$$\mathbb{E}_\omega[u^{1-\alpha}] = \frac{\int u d\pi_\lambda}{\int u^\alpha d\pi_\lambda} = \frac{1}{\int u^\alpha d\pi_\lambda}.$$

The $3\alpha/2$ moment in [Lemma B.1](#) ensures that $\mathbb{E}_\omega |\log u| < \infty$: $u^\alpha |\log u|$ is bounded when $0 < u \leq 1$, and is at most $C_\alpha u^{3\alpha/2}$ when $u \geq 1$. Jensen's inequality for the logarithm therefore gives

$$\begin{aligned} (1 - \alpha)\mathbb{E}_\omega \log u &= \mathbb{E}_\omega [\log(u^{1-\alpha})] \\ &\leq \log \mathbb{E}_\omega [u^{1-\alpha}] = -\log \int u^\alpha d\pi_\lambda. \end{aligned}$$

Because $\alpha > 1$, this implies

$$\mathbb{E}_\omega \log u \geq \frac{1}{\alpha - 1} \log \int u^\alpha d\pi_\lambda = \mathcal{R}_\alpha(\nu_t \| \pi_\lambda).$$

Consequently,

$$\begin{aligned} \frac{\text{Ent}_{\pi_\lambda}(u^\alpha)}{\int u^\alpha d\pi_\lambda} &= \alpha \mathbb{E}_\omega \log u - \log \int u^\alpha d\pi_\lambda \\ &\geq \alpha \mathcal{R}_\alpha(\nu_t \| \pi_\lambda) - (\alpha - 1) \mathcal{R}_\alpha(\nu_t \| \pi_\lambda) \\ &= \mathcal{R}_\alpha(\nu_t \| \pi_\lambda). \end{aligned}$$

Combining the upper and lower bounds proves

$$I \geq \frac{2m}{\alpha^2} \mathcal{R}_\alpha(\nu_t \| \pi_\lambda). \quad (\text{B.11})$$

Step 5. Uniform L^α bound for the density ratio: [\(B.12\)](#) and [\(4.6\)](#). In [\(B.10\)](#), use [\(B.11\)](#) and bound t by h in the four nonnegative terms. Multiplying the resulting inequality by $e^{mt/(2\alpha)}$ and integrating from 0 to t gives, after enlarging the universal constant C ,

$$\begin{aligned} \mathcal{R}_\alpha(\nu_t \| \pi_\lambda) &\leq e^{-mt/(2\alpha)} \mathcal{R}_\alpha(\nu_0 \| \pi_\lambda) \\ &\quad + \frac{C\alpha^2}{m} (1 - e^{-mt/(2\alpha)}) \left[hL_f\tau_f + h^2L_f^2R^2 + (h/\lambda)GR + \alpha^2(h/\lambda)^2G^2 \right]. \end{aligned}$$

Taking $t = h$ and using $\nu_h = \nu_0$, we move $e^{-mh/(2\alpha)} \mathcal{R}_\alpha(\nu_0 \| \pi_\lambda)$ to the left-hand side and cancel the positive factor $1 - e^{-mh/(2\alpha)}$. This bounds $\mathcal{R}_\alpha(\nu_0 \| \pi_\lambda)$ by $C\alpha^2/m$ times the bracket above. Substituting this bound back into the inequality for general t proves

$$\sup_{0 \leq t \leq h} \mathcal{R}_\alpha(\nu_t \| \pi_\lambda) \leq \frac{C\alpha^2}{m} \left[hL_f\tau_f + h^2L_f^2R^2 + (h/\lambda)GR + \alpha^2(h/\lambda)^2G^2 \right]. \quad (\text{B.12})$$

The conditions in [\(4.4\)](#), together with $R^2 \geq m$, give

$$\begin{aligned} \frac{\alpha^2}{m} hL_f\tau_f &\leq c, \\ \frac{\alpha^2}{m} h^2L_f^2R^2 &= \left(\frac{\alpha hL_fR}{\sqrt{m}} \right)^2 \leq c^2, \\ \frac{\alpha^2}{m} (h/\lambda)GR &\leq c, \\ \frac{\alpha^4}{m} (h/\lambda)^2G^2 &= \frac{m}{R^2} \left(\frac{\alpha^2}{m} (h/\lambda)GR \right)^2 \leq c^2. \end{aligned}$$

Taking the universal constant c small enough makes the right-hand side of (B.12) at most $1/4$. Finally, (2.27) gives

$$\sup_{0 \leq t \leq h} \|\nu_t / \pi_\lambda\|_{L^\alpha(\pi_\lambda)} = \exp \left\{ \frac{\alpha - 1}{\alpha} \sup_{0 \leq t \leq h} \mathcal{R}_\alpha(\nu_t \| \pi_\lambda) \right\} \leq e^{1/4}.$$

This is (4.6).

All quantitative constants above are independent of the auxiliary smoothing scale. Passing to the limit as described in Appendix D gives (4.6) for the original potential. This completes the proof of Lemma 4.2.

C Poisson Regularity and Negative Sobolev Norms

This appendix proves Lemma 2.2 and the test-function equivalence (2.24). Throughout, λ is fixed, $U_\lambda \in C^{1,1}(\mathbb{R}^d)$, and $mI \preceq \nabla^2 U_\lambda \preceq L_\lambda I$ almost everywhere.

C.1 Existence and second-order energy

Proof of the existence and energy assertions in Lemma 2.2. By (2.17), the gradient norm is a complete Hilbert norm on the mean-zero subspace of $H^1(\pi_\lambda)$. For $\phi \in L^2(\pi_\lambda)$, the functional

$$v \mapsto - \int \left(\phi - \int \phi d\pi_\lambda \right) v d\pi_\lambda$$

is continuous in that norm, with norm at most $m^{-1/2} \|\phi - \int \phi d\pi_\lambda\|_{L^2(\pi_\lambda)}$. The Riesz representation theorem gives a unique mean-zero $\psi \in H^1(\pi_\lambda)$ satisfying (2.19). The identity extends from mean-zero tests to every $v \in H^1(\pi_\lambda)$ because both sides are unchanged when a constant is added to v .

On compact sets, π_λ is bounded above and below by positive constants and ∇U_λ is bounded. The weak equation therefore gives, in the sense of distributions,

$$\Delta \psi = \phi - \int \phi d\pi_\lambda + \langle \nabla U_\lambda, \nabla \psi \rangle \in L^2_{\text{loc}}(\mathbb{R}^d).$$

For $\eta \in C_c^\infty(\mathbb{R}^d)$, the function $\eta\psi$ belongs to ordinary $H^1(\mathbb{R}^d)$ and

$$\Delta(\eta\psi) = \eta\Delta\psi + 2\langle \nabla\eta, \nabla\psi \rangle + \psi\Delta\eta \in L^2(\mathbb{R}^d).$$

Set $u = \eta\psi$ and let $u_\varepsilon = u * \varrho_\varepsilon$, where $\varrho_\varepsilon(x) = \varepsilon^{-d} \varrho(x/\varepsilon)$, $\varrho \in C_c^\infty(B(0,1))$, $\varrho \geq 0$, and $\int \varrho dx = 1$. Then $u_\varepsilon \in C_c^\infty(\mathbb{R}^d)$, $u_\varepsilon \rightarrow u$ in $H^1(\mathbb{R}^d)$, and

$$\Delta u_\varepsilon = (\Delta u) * \varrho_\varepsilon \longrightarrow \Delta u \quad \text{in } L^2(\mathbb{R}^d).$$

For every $w \in C_c^\infty(\mathbb{R}^d)$, two integrations by parts give

$$\int \|\nabla^2 w\|_F^2 dx = \sum_{i,j} \int (\partial_{ii} w)(\partial_{jj} w) dx = \int |\Delta w|^2 dx.$$

Applying this identity to $w = u_\varepsilon - u_{\varepsilon'}$ shows that $\nabla^2 u_\varepsilon$ is Cauchy in $L^2(\mathbb{R}^d)$. Its limit is the weak Hessian of u , as follows by passing to the limit in

$$\int \partial_{ij} u_\varepsilon \zeta dx = \int u_\varepsilon \partial_{ij} \zeta dx, \quad \zeta \in C_c^\infty(\mathbb{R}^d).$$

Thus $u \in H^2(\mathbb{R}^d)$. Since η was arbitrary, $\psi \in H_{\text{loc}}^2(\mathbb{R}^d)$.

Now let $\phi \in C_c^\infty(\mathbb{R}^d)$. For $u \in C_c^\infty(\mathbb{R}^d)$, differentiation of the generator in the weak sense gives

$$\nabla \mathcal{L}_\lambda u = \mathcal{L}_\lambda(\nabla u) - (\nabla^2 U_\lambda) \nabla u,$$

where $\mathcal{L}_\lambda$ acts componentwise on ∇u . Weighted integration by parts then yields

$$\begin{aligned} \int (\mathcal{L}_\lambda u)^2 d\pi_\lambda &= - \int \langle \nabla u, \nabla \mathcal{L}_\lambda u \rangle d\pi_\lambda \\ &= \int \|\nabla^2 u\|_F^2 d\pi_\lambda + \int \nabla u^\top (\nabla^2 U_\lambda) \nabla u d\pi_\lambda. \end{aligned} \quad (\text{C.1})$$

Only the bounded weak derivative of ∇U_λ is used. For a compactly supported $u \in H^2(\mathbb{R}^d)$, let $u_\varepsilon = u * \varrho_\varepsilon$. Mollification commutes with weak derivatives through order two, so $u_\varepsilon \rightarrow u$ in $H^2(\mathbb{R}^d)$. For $0 < \varepsilon < 1$, these functions are supported in a fixed compact set, on which π_λ and ∇U_λ are bounded. Consequently,

$$\|\mathcal{L}_\lambda u_\varepsilon - \mathcal{L}_\lambda u\|_{L^2(\pi_\lambda)} + \|\nabla^2 u_\varepsilon - \nabla^2 u\|_{L^2(\pi_\lambda)} + \|\nabla u_\varepsilon - \nabla u\|_{L^2(\pi_\lambda)} \rightarrow 0.$$

Together with the boundedness of $\nabla^2 U_\lambda$, these convergences justify passage to the limit in (C.1). The identity therefore holds for compactly supported $H^2(\mathbb{R}^d)$ functions.

Choose smooth cutoffs $0 \leq \chi_n \leq 1$, equal to one for $\|x\| \leq n$ and zero for $\|x\| \geq 2n$, such that $\|\nabla \chi_n\| \leq C/n$ and $\|\nabla^2 \chi_n\|_F \leq C_d/n^2$. Local regularity implies that $\chi_n \psi$ is a compactly supported H^2 function. Since ∇U_λ has at most linear growth, $\mathcal{L}_\lambda \chi_n$ is bounded uniformly in n on its annular support, with the model parameters fixed. Consequently,

$$\begin{aligned} \mathcal{L}_\lambda(\chi_n \psi) &= \chi_n \left(\phi - \int \phi d\pi_\lambda \right) + 2 \langle \nabla \chi_n, \nabla \psi \rangle + \psi \mathcal{L}_\lambda \chi_n \\ &\rightarrow \phi - \int \phi d\pi_\lambda = \mathcal{L}_\lambda \psi \quad \text{in } L^2(\pi_\lambda). \end{aligned}$$

The first term converges by dominated convergence; the other two tend to zero because $\psi, \nabla \psi \in L^2(\pi_\lambda)$ and the derivatives of χ_n are supported where $\|x\| \geq n$.

Apply (C.1) to $\chi_n \psi$. Since $\nabla^2 U_\lambda \geq 0$, both energy terms are nonnegative. Moreover, $\chi_n \psi = \psi$ on $B(0, n)$, so their derivatives agree there. Fatou's lemma and the convergence of $\mathcal{L}_\lambda(\chi_n \psi)$ give

$$\begin{aligned} &\int \|\nabla^2 \psi\|_F^2 d\pi_\lambda + \int \nabla \psi^\top \nabla^2 U_\lambda \nabla \psi d\pi_\lambda \\ &\leq \liminf_{n \rightarrow \infty} \int |\mathcal{L}_\lambda(\chi_n \psi)|^2 d\pi_\lambda \\ &= \int \left| \phi - \int \phi d\pi_\lambda \right|^2 d\pi_\lambda < \infty. \end{aligned}$$

In particular, $\nabla^2 \psi \in L^2(\pi_\lambda)$. This proves the first inequality in (2.20); the second follows from (2.17) applied to ϕ . $\square$

C.2 Approximation by compactly supported tests

Proof of the approximation assertion in Lemma 2.2. Let $\phi \in C_c^\infty(\mathbb{R}^d)$ and use the cutoffs from the preceding proof. The convergence of $\mathcal{L}_\lambda(\chi_n \psi)$ to $\mathcal{L}_\lambda \psi$ in $L^2(\pi_\lambda)$ has already been established. Now

that (2.20) gives global weighted integrability of $\nabla^2\psi$, the product formula

$$\begin{aligned}\nabla^2(\chi_n\psi) - \nabla^2\psi &= (\chi_n - 1)\nabla^2\psi + \nabla\chi_n \otimes \nabla\psi \\ &\quad + \nabla\psi \otimes \nabla\chi_n + \psi\nabla^2\chi_n\end{aligned}$$

implies convergence to zero in $L^2(\pi_\lambda)$. Likewise,

$$\nabla(\chi_n\psi) - \nabla\psi = (\chi_n - 1)\nabla\psi + \psi\nabla\chi_n \longrightarrow 0 \quad \text{in } L^2(\pi_\lambda).$$

For each fixed n , mollify $\chi_n\psi$. The mollifications converge in ordinary $H^2(\mathbb{R}^d)$ and remain supported in a fixed compact set. On that set, π_λ is bounded above and below by positive constants and ∇U_λ is bounded. Thus they also converge in each of the three norms in (2.21). Choose the mollification scale so that the sum of these three errors relative to $\chi_n\psi$ is at most $1/n$. The resulting sequence $\psi_n \in C_c^\infty(\mathbb{R}^d)$ satisfies (2.21). $\square$

C.3 Equivalent test classes for the negative Sobolev norm

Proof of (2.24). Let σ be an integrable signed density with $\int \sigma \, dx = 0$ and $\sigma/\pi_\lambda \in L^2(\pi_\lambda)$. We first compare compactly supported smooth tests with $H^1(\pi_\lambda)$ tests.

For $v \in H^1(\pi_\lambda)$, choose smooth cutoffs $0 \leq \chi_n \leq 1$, equal to one on the ball of radius n , supported in the ball of radius $2n$, and satisfying $\|\nabla\chi_n\| \leq C/n$. Then

$$\begin{aligned}\|\chi_nv - v\|_{L^2(\pi_\lambda)} &\longrightarrow 0, \\ \|\nabla(\chi_nv) - \nabla v\|_{L^2(\pi_\lambda)} &\leq \|(\chi_n - 1)\nabla v\|_{L^2(\pi_\lambda)} + \frac{C}{n} \|v\|_{L^2(\pi_\lambda)} \longrightarrow 0.\end{aligned}$$

On each compact set, π_λ is bounded above and below by positive constants. Mollifying χ_nv and choosing the smoothing scales diagonally therefore gives $v_n \in C_c^\infty(\mathbb{R}^d)$ with

$$\|v_n - v\|_{L^2(\pi_\lambda)} + \|\nabla v_n - \nabla v\|_{L^2(\pi_\lambda)} \longrightarrow 0.$$

The pairing is continuous under this convergence, since

$$\left| \int (v_n - v)\sigma \, dx \right| \leq \|\sigma/\pi_\lambda\|_{L^2(\pi_\lambda)} \|v_n - v\|_{L^2(\pi_\lambda)} \longrightarrow 0.$$

If $\|\nabla v\|_{L^2(\pi_\lambda)} \leq 1$, replace v_n by

$$\frac{v_n}{\max\{1, \|\nabla v_n\|_{L^2(\pi_\lambda)}\}}.$$

This preserves convergence to v in $H^1(\pi_\lambda)$ and makes every approximation admissible in (2.22). Thus the supremum over $H^1(\pi_\lambda)$ equals the supremum over $C_c^\infty(\mathbb{R}^d)$.

It remains to compare with C^1 tests. Let $v \in C^1(\mathbb{R}^d)$ satisfy $\|\nabla v\|_{L^2(\pi_\lambda)} < \infty$. To verify that $v \in H^1(\pi_\lambda)$, define the bounded truncations

$$v_k := \max\{-k, \min\{v, k\}\}.$$

They belong to $H^1(\pi_\lambda)$ and satisfy $\|\nabla v_k\| \leq \|\nabla v\|$ almost everywhere. The Poincaré inequality gives

$$\int \left| v_k - \int v_k \, d\pi_\lambda \right|^2 \, d\pi_\lambda \leq \frac{1}{m} \|\nabla v\|_{L^2(\pi_\lambda)}^2.$$

For $k > \sup_{\|x\| \leq 1} |v(x)|$, one has $v_k = v$ on the unit ball. Restricting the preceding variance bound to that ball gives

$$\left| \int v_k d\pi_\lambda \right| \leq \sup_{\|x\| \leq 1} |v(x)| + \frac{\|\nabla v\|_{L^2(\pi_\lambda)}}{\sqrt{m \pi_\lambda(\{x : \|x\| \leq 1\})}}.$$

The denominator is positive. Hence the means of v_k , and therefore their $L^2(\pi_\lambda)$ norms, are uniformly bounded. Since $v_k \rightarrow v$ pointwise, Fatou's lemma yields $v \in L^2(\pi_\lambda)$. Together with the assumed gradient bound, this proves $v \in H^1(\pi_\lambda)$.

Consequently, every admissible C^1 test is an admissible $H^1(\pi_\lambda)$ test. Since $C_c^\infty(\mathbb{R}^d) \subset C^1(\mathbb{R}^d)$ and the C_c^∞ and H^1 suprema have already been shown equal, the C^1 supremum has the same value. This proves (2.24). $\square$

D Weak Regularity and Integrability Before Dissipation

This appendix supplies the technical arguments used in [Appendix B](#). We first verify the properties of the auxiliary mollification. We then prove the integrability statement [Lemma B.1](#) independently of the dissipation calculation. Finally, [Section D.3](#) passes the quantitative estimate to the original potential.

D.1 Integration by parts and smooth approximation

We first justify the integrations by parts in [Section A.1](#). Let Φ be a locally Lipschitz vector field such that Φ and its weak divergence have at most polynomial growth. Choose $\chi_n \in C_c^\infty(\mathbb{R}^d)$ with $0 \leq \chi_n \leq 1$, equal to one on the ball of radius n , supported on the ball of radius $2n$, and satisfying $\|\nabla \chi_n\|_\infty \leq C/n$. Integration by parts with compact support gives

$$\int \chi_n \operatorname{div} \Phi d\pi_\lambda = \int \chi_n \langle \Phi, \nabla U_\lambda \rangle d\pi_\lambda - \int \langle \nabla \chi_n, \Phi \rangle d\pi_\lambda.$$

Strong convexity gives Gaussian tails for π_λ , and ∇U_λ grows at most linearly. The first two terms therefore converge by dominated convergence, while

$$\left| \int \langle \nabla \chi_n, \Phi \rangle d\pi_\lambda \right| \leq \frac{C}{n} \int \|\Phi\| d\pi_\lambda \rightarrow 0.$$

Thus

$$\int \operatorname{div} \Phi d\pi_\lambda = \int \langle \Phi, \nabla U_\lambda \rangle d\pi_\lambda.$$

The vector fields used in [Section A.1](#) satisfy these conditions.

Consider the mollified potentials defined in [Section B.1](#). Convolution with the nonnegative kernel ϱ_δ preserves the structural bounds:

$$\begin{aligned} mI &\preceq \nabla^2 f^{(\delta)} \preceq L_f I, & \operatorname{tr}(\nabla^2 f^{(\delta)}) &\leq \tau_f, \\ \|\nabla g_\lambda^{(\delta)}\| &\leq G, & 0 &\preceq \nabla^2 g_\lambda^{(\delta)} \preceq \lambda^{-1} I. \end{aligned}$$

Thus the same parameters and step restrictions apply for every $\delta > 0$. For each fixed δ , the mollified potentials are smooth and all their derivatives of order at least two are bounded. These derivative bounds may depend on δ .

The function $g_\lambda^{(\delta)}$ need not be the Moreau envelope of the original g . The estimates in [Appendix B](#) use the convexity, gradient, and Hessian bounds displayed above. The convergence of the regularized potentials and their associated laws is treated in [Section D.3](#).

D.2 Finiteness before the Rényi calculation

Proof of Lemma B.1. Fix $\alpha \geq 2$, λ , and h satisfying the restrictions of Lemma 4.2, and fix an auxiliary scale $\delta > 0$. Throughout this proof, the potentials, densities, kernels, and maps are the regularized objects defined in Section B.1; their superscripts (δ) are omitted. In particular, the function f and its minimizer introduced below belong to this regularized problem. In this subsection only, C and its subscripted versions may depend on the fixed problem data and on $\alpha, \lambda, h, \delta$. The step-size constant c remains universal.

We first establish an exponential moment of the invariant law, then derive density and gradient bounds uniform in $t \in [0, h]$, and finally control the weighted score and joint moments. The finite bounds obtained in this proof may depend on the fixed parameters and on δ . Their values are used only to justify the analytic operations in Appendix B and do not enter the quantitative step restrictions. For $hL_\lambda < c < 1$, the map T_h is $(1 - mh)$ -Lipschitz. The Euler kernel therefore contracts $\mathcal{P}_2(\mathbb{R}^d)$ in W_2 and has a unique invariant law.

An exponential moment of the invariant law. We prove the exponential moment bound (D.5) for $\hat{\pi}_{\lambda, h}$. Let $x_f = \arg \min f$, $F = f - f(x_f)$, and $\eta = 1 - 1/(8\alpha)$, only within this subsection. Strong convexity and smoothness give

$$F(x) \geq \frac{m}{2} \|x - x_f\|^2, \quad 2mF(x) \leq \|\nabla F(x)\|^2 \leq 2L_f F(x). \quad (\text{D.1})$$

For

$$Y = x - h\nabla U_\lambda(x) + \sqrt{2h} Z,$$

the L_f -smoothness of f gives

$$F(Y) - F(x) \leq \langle \nabla f(x), Y - x \rangle + \frac{L_f}{2} \|Y - x\|^2.$$

Substituting $Y - x = -h\nabla U_\lambda(x) + \sqrt{2h} Z$ and expanding the square yields

$$\begin{aligned} F(Y) - F(x) &\leq -h \langle \nabla f(x), \nabla U_\lambda(x) \rangle + \frac{h^2 L_f}{2} \|\nabla U_\lambda(x)\|^2 \\ &\quad + \sqrt{2h} \langle \nabla f(x) - hL_f \nabla U_\lambda(x), Z \rangle + hL_f \|Z\|^2. \end{aligned}$$

For $\xi \in \mathbb{R}^d$ and $\theta < 1/2$, completing the square gives

$$-\frac{1}{2} \|z\|^2 + \langle \xi, z \rangle + \theta \|z\|^2 = -\frac{1 - 2\theta}{2} \left\| z - \frac{\xi}{1 - 2\theta} \right\|^2 + \frac{\|\xi\|^2}{2(1 - 2\theta)}.$$

Integrating against the standard Gaussian density therefore gives

$$\mathbb{E} e^{\langle \xi, Z \rangle + \theta \|Z\|^2} = (1 - 2\theta)^{-d/2} \exp \left\{ \frac{\|\xi\|^2}{2(1 - 2\theta)} \right\}. \quad (\text{D.2})$$

The step-size restrictions ensure $2\eta h L_f < 1$. Apply (D.2) with

$$\theta = \eta h L_f, \quad \xi = \eta \sqrt{2h} (\nabla f(x) - hL_f \nabla U_\lambda(x)).$$

Since

$$\frac{Q_{\lambda, h} e^{\eta F(x)}}{e^{\eta F(x)}} = \mathbb{E} e^{\eta(F(Y) - F(x))},$$

we obtain

$$\begin{aligned}
\log \frac{Q_{\lambda,h} e^{\eta F}(x)}{e^{\eta F(x)}} &\leq -\frac{d}{2} \log(1 - 2\eta h L_f) - \eta h \langle \nabla f(x), \nabla U_\lambda(x) \rangle \\
&\quad + \frac{\eta h^2 L_f}{2} \|\nabla U_\lambda(x)\|^2 + \frac{\eta^2 h}{1 - 2\eta h L_f} \|\nabla f(x) - h L_f \nabla U_\lambda(x)\|^2 \\
&= -\frac{d}{2} \log(1 - 2\eta h L_f) \\
&\quad + \frac{\eta h}{1 - 2\eta h L_f} \left[\eta \|\nabla f(x)\|^2 - \langle \nabla f(x), \nabla U_\lambda(x) \rangle + \frac{h L_f}{2} \|\nabla U_\lambda(x)\|^2 \right].
\end{aligned}$$

Finally, substitute $\nabla U_\lambda = \nabla f + \nabla g_\lambda$ and use $\|\nabla g_\lambda\| \leq G$ to obtain

$$\begin{aligned}
\log \frac{Q_{\lambda,h} e^{\eta F}(x)}{e^{\eta F(x)}} &\leq -\frac{d}{2} \log(1 - 2\eta h L_f) \\
&\quad + \frac{\eta h}{1 - 2\eta h L_f} \left[(-(1 - \eta) + h L_f/2) \|\nabla f(x)\|^2 \right. \\
&\quad \left. - (1 - h L_f) \langle \nabla f(x), \nabla g_\lambda(x) \rangle + (h L_f/2) G^2 \right]. \tag{D.3}
\end{aligned}$$

The first condition in (4.4), together with $\tau_f \geq m$, gives $h L_f \leq c/\alpha^2$. Since $1 - \eta = 1/(8\alpha)$, taking the universal constant c small enough ensures

$$h L_f \leq \frac{1 - \eta}{2}, \quad 1 - 2\eta h L_f \geq \frac{1}{2}.$$

Hence

$$-(1 - \eta) + \frac{h L_f}{2} \leq -\frac{3}{4}(1 - \eta).$$

For the mixed term, Young's inequality gives

$$-(1 - h L_f) \langle \nabla f(x), \nabla g_\lambda(x) \rangle \leq G \|\nabla f(x)\| \leq \frac{1 - \eta}{4} \|\nabla f(x)\|^2 + \frac{G^2}{1 - \eta}.$$

The bracket in (D.3) is therefore at most

$$-\frac{1 - \eta}{2} \|\nabla f(x)\|^2 + \left(\frac{1}{1 - \eta} + \frac{h L_f}{2} \right) G^2.$$

Also, $-\log(1 - z) \leq 2z$ for $0 \leq z \leq 1/2$ gives

$$-\frac{d}{2} \log(1 - 2\eta h L_f) \leq 2\eta d h L_f.$$

Using $\|\nabla f(x)\|^2 \geq 2mF(x)$, we conclude that

$$\log \frac{Q_{\lambda,h} e^{\eta F}(x)}{e^{\eta F(x)}} \leq -\eta(1 - \eta)mhF(x) + C_\alpha h.$$

Thus, with $c_\alpha = \eta(1 - \eta) > 0$,

$$Q_{\lambda,h} e^{\eta F} \leq e^{\eta F} \exp\{-c_\alpha mhF + C_\alpha h\}. \tag{D.4}$$

Choose a finite $K > 0$ such that $c_\alpha m K \geq C_\alpha + 1$. For $F(x) \geq K$, (D.4) gives

$$Q_{\lambda,h} e^{\eta F}(x) \leq e^{-h} e^{\eta F(x)}.$$

For $F(x) < K$, it gives

$$Q_{\lambda,h} e^{\eta F}(x) \leq e^{\eta K + C_\alpha h}.$$

Combining the two bounds, we obtain, for every x ,

$$Q_{\lambda,h} e^{\eta F}(x) \leq e^{-h} e^{\eta F(x)} + e^{\eta K + C_\alpha h}.$$

Start the Euler chain at $X_0 = x_f$. Since $F(x_f) = 0$, taking expectations and iterating the preceding inequality gives

$$\mathbb{E} e^{\eta F(X_n)} \leq e^{-nh} + e^{\eta K + C_\alpha h} \sum_{j=0}^{n-1} e^{-jh} \leq 1 + \frac{e^{\eta K + C_\alpha h}}{1 - e^{-h}}.$$

Each expectation is finite by induction, and the bound is independent of n . The W_2 contraction gives convergence of the law of X_n to $\hat{\pi}_{\lambda,h}$. Since $e^{\eta F}$ is continuous and nonnegative, lower semicontinuity yields

$$\int e^{\eta F} \hat{\pi}_{\lambda,h} dx \leq \liminf_{n \rightarrow \infty} \mathbb{E} e^{\eta F(X_n)} < \infty. \quad (\text{D.5})$$

Bounds for the density and its gradient. We bound ν_t and its gradient uniformly for $0 \leq t \leq h$, as stated in (D.9).

We first prove the comparison (D.6) between $F(y)$ and $F(z)$. For $0 < \beta < \gamma$, Taylor's upper bound gives

$$\beta F(y) - \gamma F(z) \leq -(\gamma - \beta)F(z) + \beta \langle \nabla F(z), y - z \rangle + \frac{\beta L_f}{2} \|y - z\|^2.$$

By Young's inequality,

$$\begin{aligned} \beta \langle \nabla F(z), y - z \rangle &\leq \frac{\gamma - \beta}{2L_f} \|\nabla F(z)\|^2 + \frac{\beta^2 L_f}{2(\gamma - \beta)} \|y - z\|^2 \\ &\leq (\gamma - \beta)F(z) + \frac{\beta^2 L_f}{2(\gamma - \beta)} \|y - z\|^2, \end{aligned}$$

where the second inequality uses $\|\nabla F(z)\|^2 \leq 2L_f F(z)$. Substitution gives

$$\beta F(y) - \gamma F(z) \leq \frac{\beta \gamma L_f}{2(\gamma - \beta)} \|y - z\|^2. \quad (\text{D.6})$$

We also need an upper bound for $F(T_t x)$. For $0 \leq t \leq h$, Taylor's upper bound gives

$$\begin{aligned} F(T_t x) - F(x) &\leq -t \langle \nabla f(x), \nabla U_\lambda(x) \rangle + \frac{t^2 L_f}{2} \|\nabla U_\lambda(x)\|^2 \\ &= -\frac{t}{2} \|\nabla f(x)\|^2 + \frac{t}{2} \|\nabla g_\lambda(x)\|^2 - \frac{t(1 - tL_f)}{2} \|\nabla U_\lambda(x)\|^2 \\ &\leq -\frac{t}{2} \|\nabla f(x)\|^2 + \frac{t}{2} G^2 \leq \frac{t}{2} G^2, \end{aligned}$$

using $tL_f \leq 1$ and $\|\nabla g_\lambda\| \leq G$. Thus

$$F(T_t x) \leq F(x) + \frac{t}{2} G^2, \quad 0 \leq t \leq h. \quad (\text{D.7})$$

Choose the two intermediate exponents

$$\beta_1 := 1 - \frac{1}{4\alpha}, \quad \beta_2 := 1 - \frac{1}{2\alpha},$$

and decrease the universal step constant so that $h\alpha L_f \leq 1/32$.

The stationary identity and (2.12) give

$$\widehat{\pi}_{\lambda,h}(y) = (4\pi h)^{-d/2} \int e^{-\|y-T_h x\|^2/(4h)} \widehat{\pi}_{\lambda,h}(x) \, dx.$$

Differentiating the Gaussian kernel gives

$$\nabla \widehat{\pi}_{\lambda,h}(y) = -(4\pi h)^{-d/2} \int \frac{y - T_h x}{2h} e^{-\|y-T_h x\|^2/(4h)} \widehat{\pi}_{\lambda,h}(x) \, dx.$$

This differentiation is justified because the kernel and its spatial derivatives are bounded for fixed $h > 0$. Since

$$(4\pi h)^{-d/2} \left(1 + \frac{\|y - T_h x\|}{2h} \right) e^{-\|y-T_h x\|^2/(4h)} \leq C_h e^{-\|y-T_h x\|^2/(8h)},$$

the density and its gradient can be estimated using the same Gaussian bound. In (D.6), the choice $(\gamma, \beta) = (\eta, \beta_1)$ gives a coefficient at most $4\alpha L_f$. Thus

$$\begin{aligned} & e^{\beta_1 F(y)} (\widehat{\pi}_{\lambda,h}(y) + \|\nabla \widehat{\pi}_{\lambda,h}(y)\|) \\ & \leq C_h \int e^{\eta F(T_h x)} \exp \left\{ - \left(\frac{1}{8h} - 4\alpha L_f \right) \|y - T_h x\|^2 \right\} \widehat{\pi}_{\lambda,h}(x) \, dx \\ & \leq C_h e^{\eta h G^2/2} \int e^{\eta F(x)} \widehat{\pi}_{\lambda,h}(x) \, dx < \infty. \end{aligned}$$

The last inequality uses $h\alpha L_f \leq 1/32$, (D.7) with $t = h$, and (D.5). The bound is independent of y .

The change-of-variables formula gives

$$((T_t)_\# \widehat{\pi}_{\lambda,h})(y) = \widehat{\pi}_{\lambda,h}(T_t^{-1} y) |\det DT_t^{-1}(y)|.$$

By Proposition 2.1,

$$DT_t^{-1}(y) = (I - t\nabla^2 U_\lambda(T_t^{-1} y))^{-1}, \quad \|DT_t^{-1}(y)\|_{\text{op}} \leq (1 - hL_\lambda)^{-1}.$$

The determinant is positive. Differentiating the density formula gives

$$\begin{aligned} \nabla((T_t)_\# \widehat{\pi}_{\lambda,h})(y) &= \det DT_t^{-1}(y) [DT_t^{-1}(y)]^\top \nabla \widehat{\pi}_{\lambda,h}(T_t^{-1} y) \\ &\quad + \widehat{\pi}_{\lambda,h}(T_t^{-1} y) \nabla \det DT_t^{-1}(y). \end{aligned}$$

At the fixed smoothing scale, the third derivatives of U_λ are bounded. Differentiating the inverse-matrix formula above therefore shows that $\det DT_t^{-1}$ and $\nabla \det DT_t^{-1}$ are uniformly bounded for $0 \leq t \leq h$.

The preceding bounds on $\widehat{\pi}_{\lambda,h}$ and $\nabla\widehat{\pi}_{\lambda,h}$ now imply

$$((T_t)_\# \widehat{\pi}_{\lambda,h})(y) + \|\nabla((T_t)_\# \widehat{\pi}_{\lambda,h})(y)\| \leq C e^{-\beta_1 F(T_t^{-1}y)}.$$

Applying (D.7) with $x = T_t^{-1}y$ gives

$$F(y) \leq F(T_t^{-1}y) + tG^2/2.$$

Consequently,

$$((T_t)_\# \widehat{\pi}_{\lambda,h})(y) + \|\nabla((T_t)_\# \widehat{\pi}_{\lambda,h})(y)\| \leq C e^{-\beta_1 F(y)}, \quad 0 \leq t \leq h. \quad (\text{D.8})$$

with one constant for the whole time interval.

For the final heat convolution, apply (D.6) with $(\gamma, \beta) = (\beta_1, \beta_2)$, whose coefficient is at most $2\alpha L_f$. By (D.8) and (D.1), both the pushed density and its gradient are integrable. Spatial integration by parts therefore gives

$$\nabla H_t((T_t)_\# \widehat{\pi}_{\lambda,h}) = H_t \nabla((T_t)_\# \widehat{\pi}_{\lambda,h}).$$

For $t > 0$ it follows that

$$\begin{aligned} & e^{\beta_2 F(y)} (\nu_t(y) + \|\nabla \nu_t(y)\|) \\ & \leq C (4\pi t)^{-d/2} \int \exp\left\{-\left(\frac{1}{4t} - 2\alpha L_f\right) \|y - z\|^2\right\} dz \\ & = C (1 - 8\alpha t L_f)^{-d/2} \leq C'. \end{aligned}$$

The case $t = 0$ follows from (D.8), since T_0 is the identity, $F \geq 0$, and $\beta_1 > \beta_2$. Consequently, with a constant uniform for $0 \leq t \leq h$,

$$\nu_t(y) + \|\nabla \nu_t(y)\| \leq C_{\alpha,h} \exp\{-(1 - 1/(2\alpha))F(y)\}. \quad (\text{D.9})$$

Regularity and endpoint continuity. We show that the first time derivative and the spatial derivatives up to order two of ν_t exist and are continuous for $0 < t < h$. We then check continuity at the two time endpoints.

For $t > 0$, write the Gaussian representation as

$$\nu_t(y) = \int \kappa_t(y, x) \widehat{\pi}_{\lambda,h}(x) dx, \quad \kappa_t(y, x) = (4\pi t)^{-d/2} \exp\left\{-\frac{\|y - T_t x\|^2}{4t}\right\}.$$

Here $T_t x = x - t \nabla U_\lambda(x)$, while $\widehat{\pi}_{\lambda,h}$ is fixed throughout the interpolation.

The spatial derivatives of the kernel are

$$\begin{aligned} \nabla_y \kappa_t(y, x) &= -\frac{y - T_t x}{2t} \kappa_t(y, x), \\ \nabla_y^2 \kappa_t(y, x) &= \left[\frac{(y - T_t x)(y - T_t x)^\top}{4t^2} - \frac{I_d}{2t} \right] \kappa_t(y, x). \end{aligned}$$

For the time derivative, note that $\partial_t(y - T_t x) = \nabla U_\lambda(x)$. Differentiating both the prefactor and the exponent gives

$$\partial_t \kappa_t(y, x) = \left[-\frac{d}{2t} + \frac{\|y - T_t x\|^2}{4t^2} - \frac{\langle y - T_t x, \nabla U_\lambda(x) \rangle}{2t} \right] \kappa_t(y, x).$$

Fix $0 < \varepsilon < h$. The functions

$$z \mapsto (1 + \|z\| + \|z\|^2)e^{-\|z\|^2/4}$$

are bounded on $\mathbb{R}^d$. Applying this observation with $z = (y - T_t x)/\sqrt{t}$ to the displayed derivatives gives, for $\varepsilon \leq t \leq h$ and all x, y ,

$$\begin{aligned} & |\kappa_t(y, x)| + \|\nabla_y \kappa_t(y, x)\| + \|\nabla_y^2 \kappa_t(y, x)\| + |\partial_t \kappa_t(y, x)| \\ & \leq C_\varepsilon (1 + \|\nabla U_\lambda(x)\|). \end{aligned}$$

This bound is independent of t and y in the stated range.

The drift grows at most linearly, and the quadratic lower bound in (D.1) implies

$$1 + \|\nabla U_\lambda(x)\| \leq C(1 + \|x - x_f\|) \leq C' e^{\eta F(x)}.$$

Thus (D.5) makes the common bound integrable against $\hat{\pi}_{\lambda, h}(x) dx$.

We may therefore differentiate the integral once in time and twice in space:

$$\begin{aligned} \partial_t \nu_t(y) &= \int \partial_t \kappa_t(y, x) \hat{\pi}_{\lambda, h}(x) dx, \\ \nabla_y^j \nu_t(y) &= \int \nabla_y^j \kappa_t(y, x) \hat{\pi}_{\lambda, h}(x) dx, \quad j = 1, 2. \end{aligned}$$

For each fixed x , the kernel and all the displayed derivatives are continuous in (t, y) . Dominated convergence, using the same common bound, shows that their integrals are also continuous in (t, y) . Since $\varepsilon > 0$ is arbitrary, ν is locally $C^{1,2}$ on $(0, h) \times \mathbb{R}^d$.

Moreover, $\kappa_t(y, x) > 0$ for every $t > 0$ and every x, y . Since $\hat{\pi}_{\lambda, h}$ is a probability density, this gives $\nu_t(y) > 0$. The same dominated-convergence argument applies as $t \uparrow h$, so

$$\nu_t(y) \longrightarrow \nu_h(y) = \hat{\pi}_{\lambda, h}(y).$$

We now prove continuity at $t = 0$. First, the inverse-map bound in Proposition 2.1 gives

$$\begin{aligned} \|T_t^{-1}y - y\| &= \|T_t^{-1}y - T_t^{-1}(T_t y)\| \\ &\leq \frac{1}{1 - tL_\lambda} \|y - T_t y\| \\ &= \frac{t}{1 - tL_\lambda} \|\nabla U_\lambda(y)\| \longrightarrow 0. \end{aligned}$$

The inverse derivative also satisfies

$$DT_t^{-1}(y) = (I - t\nabla^2 U_\lambda(T_t^{-1}y))^{-1}, \quad \|DT_t^{-1}(y) - I_d\|_{\text{op}} \leq \frac{tL_\lambda}{1 - tL_\lambda} \longrightarrow 0.$$

Using the change-of-variables formula (2.11) and the continuity of $\hat{\pi}_{\lambda, h} = \nu_h$, we obtain

$$((T_t)_\# \hat{\pi}_{\lambda, h})(y) = \hat{\pi}_{\lambda, h}(T_t^{-1}y) \det DT_t^{-1}(y) \longrightarrow \hat{\pi}_{\lambda, h}(y).$$

We next estimate the effect of the Gaussian increment. Equation (D.8) and $F \geq 0$ give

$$\sup_{0 \leq r \leq h} \|\nabla((T_r)_\# \hat{\pi}_{\lambda, h})\|_\infty \leq C.$$

Thus these densities are Lipschitz with the same constant for all $0 \leq r \leq h$. By (2.12) and $\nu_t = \mathbf{H}_t((T_t)_\# \hat{\pi}_{\lambda,h})$,

$$\begin{aligned} & |\nu_t(y) - ((T_t)_\# \hat{\pi}_{\lambda,h})(y)| \\ &= \left| \mathbb{E} \left[((T_t)_\# \hat{\pi}_{\lambda,h})(y + \sqrt{2t} Z) - ((T_t)_\# \hat{\pi}_{\lambda,h})(y) \right] \right| \\ &\leq C\sqrt{2t} \mathbb{E} \|Z\| \longrightarrow 0. \end{aligned}$$

Together with $((T_t)_\# \hat{\pi}_{\lambda,h})(y) \rightarrow \hat{\pi}_{\lambda,h}(y)$, this proves

$$\nu_t(y) \longrightarrow \hat{\pi}_{\lambda,h}(y) = \nu_0(y) \quad \text{as } t \downarrow 0.$$

Density-ratio and score integrals. Since g_λ is G -Lipschitz,

$$\pi_\lambda(y) \geq C^{-1} \exp\{-F(y) - G\|y - x_f\|\}.$$

Combining this with (D.9) gives

$$\begin{aligned} u(y)^\alpha \pi_\lambda(y) &= \nu_t(y)^\alpha \pi_\lambda(y)^{1-\alpha} \\ &\leq C \exp \left\{ -\alpha \left(1 - \frac{1}{2\alpha} \right) F(y) + (\alpha - 1)(F(y) + G\|y - x_f\|) \right\} \\ &= C \exp \left\{ -\frac{1}{2} F(y) + (\alpha - 1)G\|y - x_f\| \right\}. \end{aligned}$$

Similarly,

$$u(y)^{3\alpha/2} \pi_\lambda(y) \leq C \exp \left\{ -\frac{1}{4} F(y) + \left(\frac{3\alpha}{2} - 1 \right) G\|y - x_f\| \right\}.$$

Both bounds are integrable because

$$F(y) \geq \frac{m}{2} \|y - x_f\|^2.$$

Their constants are uniform for $0 \leq t \leq h$.

For the score, use $s = \nabla \log \nu_t + \nabla U_\lambda$ to obtain

$$u^\alpha \pi_\lambda \|s\|^2 \leq 2\pi_\lambda^{1-\alpha} \nu_t^{\alpha-2} \|\nabla \nu_t\|^2 + 2\pi_\lambda^{1-\alpha} \nu_t^\alpha \|\nabla U_\lambda\|^2.$$

Since $\alpha \geq 2$, the factor $\nu_t^{\alpha-2}$ can be bounded using (D.9). The same equation bounds $\nabla \nu_t$, and ∇U_λ grows at most linearly. Thus the sum of the two terms is bounded by

$$C(1 + \|y - x_f\|^2) \exp \left\{ -\frac{1}{2} F(y) + (\alpha - 1)G\|y - x_f\| \right\}.$$

This bound is integrable and uniform in time. We have therefore proved

$$\sup_{0 \leq t \leq h} \int [u^\alpha(1 + \|s\|^2) + u^{3\alpha/2}] d\pi_\lambda < \infty.$$

The pointwise endpoint continuity of ν_t and the integrable bound for $u^\alpha \pi_\lambda$ give the following limits by dominated convergence:

$$\begin{aligned} \lim_{t \downarrow 0} \int u(t)^\alpha d\pi_\lambda &= \int \left(\frac{\hat{\pi}_{\lambda,h}}{\pi_\lambda} \right)^\alpha d\pi_\lambda, \\ \lim_{t \uparrow h} \int u(t)^\alpha d\pi_\lambda &= \int \left(\frac{\hat{\pi}_{\lambda,h}}{\pi_\lambda} \right)^\alpha d\pi_\lambda. \end{aligned} \tag{D.10}$$

Since $\nu_0 = \nu_h = \hat{\pi}_{\lambda,h}$, this proves continuity of $t \mapsto \int u(t)^\alpha d\pi_\lambda$ at both endpoints.

Joint weighted moments. We prove the joint moment bound in [Lemma B.1](#). The upper bound for ν_t and the lower bound for π_λ give

$$u(y)^{\alpha-1} \leq C \exp \left\{ \frac{\alpha-1}{2\alpha} F(y) + (\alpha-1)G \|y - x_f\| \right\}.$$

Strong convexity and Young's inequality give

$$(\alpha-1)G \|y - x_f\| \leq \frac{1}{8}F(y) + \frac{4(\alpha-1)^2 G^2}{m}.$$

Since $(\alpha-1)/(2\alpha) \leq 1/2$, it follows that

$$u(y)^{\alpha-1} \leq C e^{5F(y)/8}.$$

For every integer $k \geq 0$, the quadratic lower bound in [\(D.1\)](#) gives

$$(1 + \|Y\|)^k \leq C_k e^{F(Y)/32}.$$

Consequently,

$$\begin{aligned} u(Y)^{\alpha-1} (1 + \|X\| + \|Y\|)^k &\leq C_k (1 + \|X\|)^k e^{(5/8+1/32)F(Y)} \\ &= C_k (1 + \|X\|)^k e^{21F(Y)/32}. \end{aligned}$$

Conditional on X , Taylor's upper bound gives

$$F(Y) \leq F(T_t X) + \sqrt{2t} \langle \nabla F(T_t X), Z \rangle + tL_f \|Z\|^2.$$

Applying [\(D.2\)](#) yields

$$\begin{aligned} \mathbb{E} \left[e^{21F(Y)/32} \mid X \right] &\leq \left(1 - \frac{21}{16} tL_f \right)^{-d/2} \exp \left\{ \frac{21}{32} F(T_t X) + \frac{(21/32)^2 t}{1 - \frac{21}{16} tL_f} \|\nabla F(T_t X)\|^2 \right\} \\ &\leq \left(1 - \frac{21}{16} tL_f \right)^{-d/2} \exp \left\{ \frac{21F(T_t X)}{32(1 - \frac{21}{16} tL_f)} \right\}, \end{aligned}$$

where the second inequality uses $\|\nabla F\|^2 \leq 2L_f F$.

The condition $h\alpha L_f \leq 1/32$ and $\alpha \geq 2$ imply $tL_f \leq 1/64$. In particular,

$$\frac{21}{32(1 - \frac{21}{16} tL_f)} \leq \frac{3}{4},$$

and the Gaussian prefactor is bounded uniformly for $0 \leq t \leq h$. Using [\(D.7\)](#) with $x = X$, we obtain

$$\mathbb{E} \left[e^{21F(Y)/32} \mid X \right] \leq C e^{3F(X)/4}, \quad 0 \leq t \leq h.$$

Finally, $\eta = 1 - 1/(8\alpha) \geq 15/16$ gives $\eta - 3/4 \geq 3/16$. The quadratic lower bound in [\(D.1\)](#) therefore gives

$$(1 + \|X\|)^k e^{3F(X)/4} \leq C_k e^{\eta F(X)}.$$

Combining the preceding estimates gives

$$\begin{aligned} & \mathbb{E} \left[u(Y)^{\alpha-1} (1 + \|X\| + \|Y\|)^k \right] \\ & \leq C_k \mathbb{E} \left[(1 + \|X\|)^k e^{3F(X)/4} \right] \\ & \leq C'_k \int e^{\eta F(x)} \widehat{\pi}_{\lambda,h}(x) dx < \infty. \end{aligned}$$

The last inequality is (D.5), and the bound is uniform for $0 \leq t \leq h$.

It remains to bound the integral containing e_t . By its definition,

$$e_t(Y) = \mathbb{E}[\nabla U_\lambda(Y) - \nabla U_\lambda(X) \mid Y].$$

Conditional Jensen's inequality gives

$$\begin{aligned} \int u^\alpha \|e_t\|^2 d\pi_\lambda &= \mathbb{E} \left[u(Y)^{\alpha-1} \|e_t(Y)\|^2 \right] \\ &\leq \mathbb{E} \left[u(Y)^{\alpha-1} \|\nabla U_\lambda(Y) - \nabla U_\lambda(X)\|^2 \right] \\ &\leq L_\lambda^2 \mathbb{E} \left[u(Y)^{\alpha-1} \|Y - X\|^2 \right]. \end{aligned}$$

The last expression is uniformly finite by the joint moment bound with $k = 2$.

This completes the proof of [Lemma B.1](#). □

D.3 Removing the auxiliary mollification

We now restore the superscripts (δ) and let $\delta \downarrow 0$, with α, λ , and h fixed. The global Lipschitz bound on ∇U_λ gives

$$\varepsilon_\delta := \left\| \nabla U_\lambda^{(\delta)} - \nabla U_\lambda \right\|_\infty \longrightarrow 0.$$

By the definitions of $f^{(\delta)}$ and $g_\lambda^{(\delta)}$,

$$U_\lambda^{(\delta)}(x) = (U_\lambda * \varrho_\delta)(x) = \int U_\lambda(x - \delta z) \varrho(z) dz.$$

The symmetry of ϱ gives

$$\int z \varrho(z) dz = 0.$$

Since ∇U_λ is L_λ -Lipschitz, Taylor's remainder satisfies

$$|U_\lambda(x - \delta z) - U_\lambda(x) + \delta \langle \nabla U_\lambda(x), z \rangle| \leq \frac{L_\lambda \delta^2}{2} \|z\|^2.$$

Using the zero first moment, we can write

$$U_\lambda^{(\delta)}(x) - U_\lambda(x) = \int \left[U_\lambda(x - \delta z) - U_\lambda(x) + \delta \langle \nabla U_\lambda(x), z \rangle \right] \varrho(z) dz.$$

Integrating the remainder bound and taking the supremum over x therefore gives

$$\left\| U_\lambda^{(\delta)} - U_\lambda \right\|_\infty \leq \frac{L_\lambda \delta^2}{2} \int \|z\|^2 \varrho(z) dz = O(L_\lambda \delta^2).$$

The implicit constant depends only on the fixed kernel ϱ . Consequently, $\pi_\lambda^{(\delta)}/\pi_\lambda$ converges uniformly to one. In particular, $\pi_\lambda^{(\delta)} \leq 2\pi_\lambda$ for all sufficiently small δ . Since π_λ has a finite second moment, dominated convergence gives both weak convergence and convergence of the second moments. By the standard characterization of W_2 convergence,

$$W_2(\pi_\lambda^{(\delta)}, \pi_\lambda) \longrightarrow 0.$$

Recall that

$$\varepsilon_\delta := \left\| \nabla U_\lambda^{(\delta)} - \nabla U_\lambda \right\|_\infty, \quad T_h^{(\delta)} x := x - h \nabla U_\lambda^{(\delta)}(x).$$

The smoothed and original potentials satisfy the same lower and upper Hessian bounds. Hence the contraction argument in [Proposition 3.3](#) gives

$$\left\| T_h^{(\delta)} x - T_h^{(\delta)} y \right\| \leq (1 - mh) \|x - y\|.$$

Also,

$$\left\| T_h^{(\delta)} x - T_h x \right\| \leq h \varepsilon_\delta.$$

Choose an optimal coupling (X_δ, X) of $\hat{\pi}_{\lambda,h}^{(\delta)}$ and $\hat{\pi}_{\lambda,h}$, so that

$$\left(\mathbb{E} \|X_\delta - X\|^2 \right)^{1/2} = W_2(\hat{\pi}_{\lambda,h}^{(\delta)}, \hat{\pi}_{\lambda,h}).$$

Let $Z \sim N(0, I_d)$ be independent of this pair. By invariance,

$$T_h^{(\delta)} X_\delta + \sqrt{2h} Z \sim \hat{\pi}_{\lambda,h}^{(\delta)}, \quad T_h X + \sqrt{2h} Z \sim \hat{\pi}_{\lambda,h}.$$

These two updated random variables therefore give another coupling of the same invariant laws. Their Gaussian increments cancel in the difference. The definition of W_2 and the L^2 triangle inequality give

$$\begin{aligned} W_2(\hat{\pi}_{\lambda,h}^{(\delta)}, \hat{\pi}_{\lambda,h}) &\leq \left(\mathbb{E} \left\| T_h^{(\delta)} X_\delta - T_h X \right\|^2 \right)^{1/2} \\ &\leq \left(\mathbb{E} \left\| T_h^{(\delta)} X_\delta - T_h^{(\delta)} X \right\|^2 \right)^{1/2} + \left(\mathbb{E} \left\| T_h^{(\delta)} X - T_h X \right\|^2 \right)^{1/2} \\ &\leq (1 - mh) \left(\mathbb{E} \|X_\delta - X\|^2 \right)^{1/2} + h \varepsilon_\delta \\ &= (1 - mh) W_2(\hat{\pi}_{\lambda,h}^{(\delta)}, \hat{\pi}_{\lambda,h}) + h \varepsilon_\delta. \end{aligned}$$

Rearranging and using $h > 0$ gives

$$W_2(\hat{\pi}_{\lambda,h}^{(\delta)}, \hat{\pi}_{\lambda,h}) \leq \frac{\varepsilon_\delta}{m} \longrightarrow 0 \quad \text{as } \delta \downarrow 0.$$

The same coupling for the interpolations, together with $\text{Lip}(T_t^{(\delta)}) \leq 1 - mt$, yields

$$\begin{aligned} W_2(\nu_t^{(\delta)}, \nu_t) &\leq (1 - mt) W_2(\hat{\pi}_{\lambda,h}^{(\delta)}, \hat{\pi}_{\lambda,h}) + t \varepsilon_\delta \\ &\leq \frac{\varepsilon_\delta}{m}, \quad 0 \leq t \leq h. \end{aligned}$$

Thus the regularized interpolations converge to the original interpolation uniformly in time in W_2 .

Under the restrictions of [Lemma 4.2](#), the argument in [Appendix B](#) gives

$$\mathcal{R}_\alpha(\nu_t^{(\delta)} \|\pi_\lambda^{(\delta)}) \leq \frac{1}{4} \quad \text{for every } \delta > 0 \text{ and } 0 \leq t \leq h.$$

For each fixed $t \in [0, h]$, the W_2 convergence established above implies the weak convergence

$$\nu_t^{(\delta)} \Rightarrow \nu_t, \quad \pi_\lambda^{(\delta)} \Rightarrow \pi_\lambda.$$

Joint lower semicontinuity of the Rényi divergence, together with the uniform bound for the regularized problem, therefore gives

$$\mathcal{R}_\alpha(\nu_t \|\pi_\lambda) \leq \liminf_{\delta \downarrow 0} \mathcal{R}_\alpha(\nu_t^{(\delta)} \|\pi_\lambda^{(\delta)}) \leq \frac{1}{4}.$$

The same bound holds for every $t \in [0, h]$, so taking the supremum in time and using [\(2.27\)](#) proves [\(4.6\)](#). The same argument also preserves the full quantitative bound [\(B.12\)](#).

References

- [1] Sinho Chewi. Log-Concave Sampling. Book manuscript, 2026. URL <https://chewisinho.github.io/main.pdf>. Final draft, August 20, 2026.
- [2] Sinho Chewi, Murat A. Erdogdu, Mufan Bill Li, Ruoqi Shen, and Matthew Zhang. Analysis of Langevin Monte Carlo from Poincaré to Log-Sobolev. arXiv preprint arXiv:2112.12662v2, 2024. URL <https://arxiv.org/abs/2112.12662v2>.
- [3] Alain Durmus, Éric Moulines, and Marcelo Pereyra. Efficient Bayesian computation by proximal Markov chain Monte Carlo: When Langevin meets Moreau. *SIAM Journal on Imaging Sciences*, 11(1):473–506, 2018. doi: 10.1137/16M1108340.
- [4] Jiaojiao Fan, Bo Yuan, and Yongxin Chen. Improved dimension dependence of a proximal algorithm for sampling. In *Proceedings of Thirty Sixth Conference on Learning Theory*, volume 195 of *Proceedings of Machine Learning Research*, pages 1473–1521. PMLR, 2023. URL <https://proceedings.mlr.press/v195/fan23a.html>.
- [5] Francesco Pedrotti and Peter A. Whalley. Wasserstein mixing time of the unadjusted Langevin algorithm. arXiv preprint arXiv:2608.02430, 2026.
- [6] Rémi Peyre. Comparison between W_2 distance and $\dot{H}^{-1}$ norm, and localisation of Wasserstein distance. arXiv preprint arXiv:1104.4631v2, 2016. URL <https://arxiv.org/abs/1104.4631v2>.
- [7] Leon Simon. Introduction to geometric measure theory. Lecture notes, 2018. URL <https://math.stanford.edu/~lms/ntu-gmt-text.pdf>. NTU lectures, revised March 7, 2018.
- [8] Yuchen Xin and Zhihua Zhang. Active-trace complexity bounds for Moreau–Yosida unadjusted Langevin sampling. arXiv preprint arXiv:2608.13467v1, 2026. URL <https://arxiv.org/abs/2608.13467v1>.
- [9] Yuchen Xin and Zhihua Zhang. Poisson-corrector complexity bounds for Moreau–Yosida unadjusted Langevin sampling. arXiv preprint arXiv:2609.12594v1, 2026. URL <https://arxiv.org/abs/2609.12594v1>.