

Thin-Shell Stability of Gaussian Cooling: Logconcave Sampling with Sesteric Complexity from a Cold Start

Yunbum Kook
University of Michigan
ybkook@umich.edu

Santosh S. Vempala
Georgia Tech
vempala@gatech.edu

Abstract

We show that logconcave probability measures along the Gaussian cooling path have thin-shell stability, generalizing the thin-shell theorem. This result leads to improved complexity for the fundamental problem of sampling an arbitrary logconcave distribution from a cold start. For (near-)isotropic logconcave distributions, the complexity is nearly $n^{2.5}$, improving the previous bound of $n^{2.75}$, and matching the complexity of the abstract Speedy walk.

1 Introduction

The study of the complexity of logconcave sampling in high dimension has led to powerful algorithmic techniques and beautiful geometric phenomena. It is known that an arbitrary logconcave distribution in $\mathbb{R}^n$ with density proportional to $e^{-V(x)}$ for some convex function V can be sampled from a *warm* start¹ using $\tilde{O}(n^2 C_{\text{PI}} \text{polylog } \frac{1}{\varepsilon})$ evaluations of V [KV26b]. This is a tight bound for known methods and conjectured to be the best possible for any algorithm with only oracle query access to the target density. Here, the Poincaré constant C_{PI} is conjectured to be a universal constant for isotropic logconcave distributions (by the celebrated Kannan–Lovász–Simonovits (KLS) conjecture [KLS95]) and known to be bounded by $O(\log n)$ [Kla23].

In this paper, we focus on the problem of logconcave sampling from a *cold* start (i.e., no warm start), given access to an evaluation oracle for V . For the methods that achieve the best known bounds from a warm start, such as the Ball walk [KLS97] and In-and-Out [KVZ26, KV25b], initiating them from an arbitrary point could lead to very high complexity. This is due to the fact that for points near the boundary, these methods could make a large number of attempts (queries) before making a single proper step; moreover, this is unavoidable. Hit-and-Run has also been shown to have similar complexity from a warm start [LV06b, KV26a], and while it has the advantage of polynomial complexity from an arbitrary interior starting point, the known analysis incurs an additional factor of dimension, i.e., with complexity scaling as n^3 .

Sampling is a key ingredient in algorithms for volume computation [DFK91, LS93, KLS97, LV06c, CV18], integration, and rounding [LV06a, JLLV26, KV25b]. To give complete guarantees for the latter problems, researchers have addressed the challenge of generating a warm start efficiently, with Kannan, Lovász, and Simonovits [KLS97] giving an algorithm for convex bodies with complexity roughly n^5 . The main idea is annealing, i.e., a sequence of distributions, each one providing a warm start for the next. The starting distribution is easy to sample, and the final distribution is the target. Lovász and Vempala improved this complexity to n^4 [LV06c] and extended the approach to logconcave distributions [LV06a]. In fact, a key idea was to use a sequence of logconcave distributions

¹Roughly speaking, a warm start means that the initial distribution is already close to the target distribution.

in the annealing schedule, which provably achieves an improvement over using a sequence of uniform distributions even for the case of convex bodies [DFK91, KLS97].

A decade later, Cousins and Vempala [CV18] introduced **Gaussian cooling**, an accelerated adaptation of the previous annealing schemes, and combined it with the **Ball walk** : $R_\infty \rightarrow TV$ ² to obtain n^3 complexity for well-rounded distributions³. In particular, this implies that for an arbitrary near-isotropic distribution, sampling from a cold start has cubic complexity. The main idea in **Gaussian cooling** was an adaptive schedule, which efficiently accelerated the rate of progress towards the target, and this cubic complexity remained the state-of-the-art till 2025. We will discuss this in more detail presently.

The work of Lee and Vempala [LV18, LV24] suggests that further improvement might be possible. They showed that the **Speedy walk**, an abstract process introduced by [KLS97] and used only for analysis, has iteration complexity $n^{2.5}$ (not query complexity) and that this bound is tight. The **Speedy walk** counts only proper steps of the **Ball walk** and not wasted steps, which are needed by any known implementation; whether this $n^{2.5}$ iteration complexity can be matched by an actual algorithm in oracle-query complexity has remained an elusive question.

For uniform sampling, Kook, Vempala, and Zhang [KVZ24, KVZ26] developed the **In-and-Out** algorithm (also known as the proximal sampler [LST21]), using algorithmic diffusion and obtaining a $R_\infty \rightarrow R_q$ guarantee. Beyond its stronger output guarantee, the algorithmic connection to heat flow makes tools from Markov semigroup theory, functional inequalities, and convex geometry available for its analysis. Subsequently, Kook and Zhang [KZ25] refined the analysis of **Gaussian cooling** to obtain R_∞ -warmness with the same cubic complexity, but warm-start generation was still a bottleneck toward subcubic complexity from a cold start. **In-and-Out** and the annealing algorithm were extended to general logconcave distributions through the exponential lifting technique [KV25b], matching the complexity of uniform sampling.

These ideas were seminal for the first improvement in cold start sampling since [CV18]: using a more careful annealing schedule, refining the bound on the log-Sobolev constant of logconcave probability measures with compact support, and relying on the stronger and faster end-to-end sampling guarantee in q -Rényi divergence, [KV25a, KV25c] improved the complexity of cold start sampling to $n^{2.75}$, the first subcubic bound. They also gave an example showing a lower bound of $n^{8/3}$ for their general method. The motivation for the present paper is to understand the best possible complexity of sampling from a cold start.

Our main algorithmic result here is a refinement of **Gaussian cooling**, which, together with **In-and-Out**, achieves the complexity of nearly $n^{2.5}$ for arbitrary near-isotropic logconcave distributions. In other words, it matches the known iteration complexity of the abstract **Speedy walk** without any overhead. The core idea of the proof is a new stability phenomenon: while the Poincaré constant of logconcave distributions along the **Gaussian cooling** path can vary substantially (and thus using it, as in previous work, leads to higher complexity), the *thin-shell constant* is stable along this path, and this suffices! We will discuss this in more detail shortly, after presenting the main results precisely.

²Recall the q -Rényi divergence: $R_q(\mu \parallel \nu) := \frac{q}{q-1} \log \|\mathrm{d}\mu/\mathrm{d}\nu\|_{L^q(\nu)}$ for $q > 1$. It is monotonic in q , and satisfies $2\|\cdot\|_{TV}^2 \leq \text{KL} = \lim_{q \downarrow 1} R_q \leq R_2 = \log(1 + \chi^2) \leq R_\infty = \log \text{ess sup}_\nu \mathrm{d}\mu/\mathrm{d}\nu$. The notation $R_\infty \rightarrow TV$ means that the sampler requires an initial distribution with R_∞ -warmness and its output distribution is close to the target in total variation (TV) distance.

³A distribution is *well-rounded* if a level set of constant measure contains a ball of constant radius, and the trace of its covariance is $\tilde{O}(n)$. Any near-isotropic logconcave distribution, with covariance close to identity, is well-rounded.

1.1 Results

Setup. Let $V : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{\infty\}$ be a convex function such that $d\pi(x) \propto e^{-V(x)} dx$ is a full-dimensional probability measure in $\mathbb{R}^n$. Following the convention in [KV25b], let us denote its ground set by $L_g = \{x \in \mathbb{R}^n : V(x) - \inf V \leq 10n\}$, and standardize the setup by assuming $B(0, 1) \subset L_g$ (via translation and scaling). For the uniform distribution on a convex body $\mathcal{K} \subset \mathbb{R}^n$, this corresponds to $B(0, 1) \subseteq \mathcal{K}$. We assume access to its zeroth-order oracle (i.e., evaluation oracle for V telling us $V(x)$ for a queried point x).

We denote the barycenter by $m := \mathbb{E}_\pi X$, the covariance matrix by $\Sigma := \text{cov}_\pi X = \text{cov } \pi$, and its operator norm by $\Lambda := \|\text{cov } \pi\|_{\text{op}}$. Since the barycenter m is not necessarily at the origin, we also define $L := \|\Sigma + mm^\top\|_{\text{op}}$ and $R^2 = \mathbb{E}_\pi[\|\cdot\|^2]$. Note that $\Lambda \leq L, \text{tr } \Sigma \leq R^2$ in general. For $t > 0$, we define a Gaussian tilt of π as $\pi_t(dx) \propto \exp(-\frac{\|x\|^2}{2t}) \pi(dx)$, which can be thought of as the continuous version of **Gaussian cooling**. The stability proofs themselves use only logconcavity and these parameters, whereas the ground-set normalization is needed only for the sampling results.

Before we proceed, we define the *anisotropic thin-shell constant* Q_n of logconcave measures as the smallest number such that every isotropic logconcave distribution ν in $\mathbb{R}^n$ and every symmetric matrix M satisfy

$$\text{Var}_\nu(Z^\top M Z) \leq Q_n \text{tr}(M^2). \quad (1.1)$$

The ordinary thin-shell conjecture [ABP03, BK03], which was proven by Klartag and Lehec [KL25] in 2025, posits $Q_n = O(1)$ when $M = I_n$. Recently, Chen and Klartag showed that $Q_n = 8$ is sharp when $M = I_n$ [CK26]. For symmetric M , it holds that $Q_n \lesssim \log n$ due to Klartag’s KLS bound [Kla23].

Result 1: Thin-shell stability along Gaussian tilts. Consider the family of Gaussian tilts $\{\pi_t\}_{t \geq 0}$ that interpolates between $\pi_0 = \delta_0$ and $\pi_\infty = \pi$. The first result shows that the variance of $\|X\|^2$ can be bounded in terms of the barycenter and covariance matrix of the base measure π .

Theorem 1.1 (Stability of thin-shell estimates). *Let π be any logconcave probability measure in $\mathbb{R}^n$ with $m = \mathbb{E}_\pi X$ and $\Sigma = \text{cov } \pi$, and π_t be its Gaussian tilt for $t > 0$. Then,*

$$\sup_{t > 0} \text{Var}_{\pi_t}(\|X\|^2) \lesssim (1 + Q_n)^2 [\text{tr}(\Sigma^2) + \|m\|^2 (\text{tr}(\Sigma^2) + m^\top \Sigma m)^{1/2}] = \tilde{O}(R^2 L \wedge R^3 \Lambda^{1/2}).$$

In particular, when π is isotropic, this result implies that the thin-shell constant remains bounded: $\frac{1}{n} \text{Var}_{\pi_t}(\|X\|^2) \lesssim (1 + Q_n)^2$. Note that this extends the thin-shell bound [KL25, CK26] to *all* t (up to log factors). Its stronger version would be stability of the covariance operator norm, but this is false in general [Biz26, KV25c]; see §1.2.1 for more details.

Result 2: Faster warm-start generation for logconcave distributions. We start with warm-start generation for the uniform distribution on a convex body. Using the thin-shell stability result, our algorithm follows the **Gaussian cooling** scheme of [KV25c] with more aggressive updates to σ^2 . Roughly speaking, it updates σ^2 by a factor of $(1 + \sigma^2/V^{1/2})$ in each phase, where V is the thin-shell bound on $\sup_{t > 0} \text{Var}_{\pi_t}(\|X\|^2)$. Using **In-and-Out** for truncated Gaussians, whose query complexity is $n^2 \sigma^2$ from an $O(1)$ -warm start in R_q , doubling a given σ^2 requires $n^2 \sigma^2 \times V^{1/2}/\sigma^2 = n^2 V^{1/2}$ queries. Since there are logarithmically many doublings, the total query complexity for warm-start generation is $\tilde{O}(n^2 V^{1/2})$. This is simply $\tilde{O}(n^{2.5})$ for isotropic uniform distributions. Hence, we “algorithmically” achieve the iteration complexity of the **Speedy walk** in [LV24].

Theorem 1.2 (Faster warm start for uniform sampling). *Let $\mathcal{K} \subset \mathbb{R}^n$ be a convex body with $B(0, 1) \subset \mathcal{K}$ and π uniform on $\mathcal{K}$. Algorithm 3.5 outputs a law ν satisfying $R_q(\nu \parallel \pi) \leq 1$ for $q \geq 1$,*

using $\tilde{O}(q^{1/2}n^2V^{1/2}) = \tilde{O}(q^{1/2}n^2 \min\{RL^{1/2}, R^{3/2}\Lambda^{1/4}\})$ expected membership queries. When π is nearly isotropic (i.e., $\text{cov } \pi \approx I_n$), the total complexity is $\tilde{O}(q^{1/2}n^{2.5})$.

This improves the previously best complexity [KV25a, KV25c]. It is conceivable that $n^{2.5}$ is the best possible complexity for cold-start sampling in light of the $n^{2.5}$ lower bound for the iteration complexity of the Speedy walk [LV24].

For general logconcave distributions, we use the exponential lift [KV25b], streamline the tilted Gaussian annealing in [KV25c], and accelerate it using thin-shell stability.

Theorem 1.3 (Faster warm start for logconcave sampling). *Let $V : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{\infty\}$ be a convex function, and suppose that $\pi \propto e^{-V}$ satisfies $B(0, 1) \subseteq \mathcal{L}_g$. Given access to an evaluation oracle for V , Algorithm 4.5 outputs a law ν satisfying $R_q(\nu \parallel \pi) \leq 2$ for $q \geq 1$, using $\tilde{O}(n^{2.5} + q^{1/2}n^2V^{1/2})$ expected evaluation queries, which can be bounded as*

$$\tilde{O}(n^{2.5} + q^{1/2}n^2 \min\{RL^{1/2}, R^{3/2}\Lambda^{1/4}\}).$$

When π is nearly isotropic (i.e., $\text{cov } \pi \approx I_n$), the total complexity is $\tilde{O}(q^{1/2}n^{2.5})$.

1.2 Technical overview

Our proofs are organized around a direct connection between Gaussian cooling and thin-shell stability. We first recall the intuition behind Gaussian cooling [CV15, CV18], focusing on how thin-shell stability determines its annealing rate. We then outline the proof of thin-shell stability and explain how it leads to a faster annealing schedule for logconcave distributions.

Gaussian cooling and variance of thin shells. Let π be a logconcave distribution on $\mathbb{R}^n$, and consider the Gaussian-tilted family $\pi_{\sigma^2}(\text{d}x) \propto \exp(-\frac{\|x\|^2}{2\sigma^2}) \pi(\text{d}x)$. Gaussian cooling gradually increases σ^2 , thereby weakening the effect of Gaussian tilts and interpolating toward the target π . To understand the annealing rate, we take a geometric view of Gaussian cooling. The fluctuation scale of $\|X\|^2$ is $\text{Var}_{\pi_{\sigma^2}}(\|X\|^2)^{1/2}$. Geometrically, when we increase σ^2 , the next annealing distribution should overlap substantially with the thin shell of the previous annealing distribution π_{σ^2} so that consecutive distributions are “close” enough for a sampler to mix rapidly toward the next distribution. Hence, heuristically, $\mathbb{E}_{\pi_{\sigma_{\text{new}}^2}}[\|X\|^2]$ should lie in $\mathbb{E}_{\pi_{\sigma^2}}[\|X\|^2] \pm O(\text{Var}_{\pi_{\sigma^2}}(\|X\|^2)^{1/2})$.




Figure 1.1: Gaussian cooling viewed through the distribution of the squared radius. The schematic profiles show the laws of $\|X\|^2$ under consecutive Gaussian tilts. Increasing σ^2 to $(1 + \alpha)\sigma^2$ shifts the mean roughly by the amount indicated by the blue arrow. Comparing this displacement with the current fluctuation scale (black arrow) motivates the annealing step size.

To make this intuition concrete, let $\sigma_{\text{new}}^2 = (1 + \alpha) \sigma^2$ for a small increment α . Direct computation gives $\partial_{\sigma^2} \mathbb{E}_{\pi_{\sigma^2}}[\|X\|^2] = \text{Var}_{\pi_{\sigma^2}}(\|X\|^2)/(2\sigma^4)$. Thus, the center of the thin shell moves roughly by $\frac{\alpha}{2\sigma^2} \text{Var}_{\pi_{\sigma^2}}(\|X\|^2)$. Since we want this change to be smaller than the annulus width $O(\text{Var}_{\pi_{\sigma^2}}(\|X\|^2)^{1/2})$, this suggests the multiplicative step $\alpha \lesssim \sigma^2 / \text{Var}_{\pi_{\sigma^2}}(\|X\|^2)^{1/2}$. Note that a larger variance has two competing effects: it gives a larger annulus, but it moves its center even faster, so $\sigma^2 / \text{Var}_{\pi_{\sigma^2}}(\|X\|^2)^{1/2}$ is the ideal balance between these two effects.

This identifies $\text{Var}_{\pi_{\sigma^2}}(\|X\|^2)$ as a fundamental quantity underlying **Gaussian cooling** and motivates bounding it uniformly. Previous work bounds it by $\text{Var}_{\pi_{\sigma^2}}(\|X\|^2) \leq 4\sigma^2 \mathbb{E}_{\pi_{\sigma^2}}[\|X\|^2]$ using the strong logconcavity of π_{σ^2} and the Poincaré inequality (or Brascamp–Lieb). Since $\mathbb{E}_{\pi_{\sigma^2}}[\|X\|^2]$ can be made $O(R^2)$ by a suitable truncation, the previous bound yields $\alpha \asymp \sigma/R$. On the other hand, a uniform bound $\sup_{\sigma^2 > 0} \text{Var}_{\pi_{\sigma^2}}(\|X\|^2) \leq V$ suggests $\alpha \asymp \sigma^2/V^{1/2}$. Once $\sigma^2 \geq 1$, this suggests a faster annealing schedule than those in previous work. Theorem 1.1 establishes such a σ^2 -independent uniform bound along **Gaussian cooling**.

1.2.1 Thin-shell stability of Gaussian tilts

To build intuition, consider an isotropic logconcave probability measure ν on $\mathbb{R}^n$, and let $d\nu_s \propto e^{-s\|x\|^2} d\nu$ denote its Gaussian tilt for $s \in [0, \infty]$, where we use the inverse parameter $s = 1/(2t)$ for the sake of exposition. Prior work [KV25a] proposed a stronger version of the thin-shell stability conjecture as part of an attempt to design a faster warm-start algorithm, based on geometric intuition for stability of the largest eigenvalue of $\text{cov } \nu_s$. Indeed, by (PI),

$$\text{Var}_{\nu_s}(\|X\|^2) \leq 4C_{\text{PI}}(\nu_s) \mathbb{E}_{\nu_s}[\|X\|^2] \lesssim_{\log n} \|\text{cov } \nu_s\|_{\text{op}} \mathbb{E}_{\nu}[\|X\|^2] = n \|\text{cov } \nu_s\|_{\text{op}},$$

where the second inequality follows from $C_{\text{PI}}(\nu_s) \lesssim \|\text{cov } \nu_s\|_{\text{op}} \log n$ (i.e., the logarithmic KLS bound) and $\mathbb{E}_{\nu_s}[\|X\|^2] \leq \mathbb{E}_{\nu}[\|X\|^2]$ (which follows by differentiating $\mathbb{E}_{\nu_s}[\|X\|^2]$ with respect to s). Following the same reasoning, one might conjecture that $\lambda_s := \|\text{cov } \nu_s\|_{\text{op}}$ should be $O(\|\text{cov } \nu\|_{\text{op}}) = O(1)$. However, it was later disproved by [KV25c] using Bizeul’s example [Biz26].

A natural alternative is to directly control $V_s := \text{Var}_{\nu_s}(\|X\|^2)$ without going through (PI). Hence, we study the thin-shell variance V_s as s decreases from ∞ to 0. Two competing factors determine V_s : (i) the base measure ν and (ii) the variance $\frac{1}{2s}$ of the Gaussian factor. When s is large (so the Gaussian variance is small), the Gaussian factor pulls the ν -mass toward the origin. In this regime, the effect of the base measure ν should be limited. Hence, ν_s is concentrated near the origin, and intuitively, V_s should be small. As s decreases (so the Gaussian variance is larger), we expect the ν_s -measure to become more dispersed, which should in turn increase V_s . However, when s is sufficiently small, the Gaussian weight is $\Theta(1)$ throughout the effective support of ν , which contains most of the ν -measure (e.g., $\|x\| = O(n^{1/2})$). In this regime, the Gaussian tilt has little effect. Hence, ν_s should be distributed similar to the isotropic measure ν , and we expect $V_s \approx \text{Var}_{\nu}(\|X\|^2)$. By the thin-shell theorem, we expect $V_s = O(n)$. In summary, as s decreases from ∞ to 0, the thin-shell variance should initially increase as the Gaussian factor weakens. Once the effect of the original isotropic measure becomes dominant, however, V_s should remain $O(n)$. Thus, a plausible conjecture is that $V_s = O(n)$ for **all** $s \in [0, \infty]$.

Decomposition of variance. For a general logconcave measure π with $m = \mathbb{E}X$ and $\Sigma = \text{cov } \pi$, let $Z = \Sigma^{-1/2}(X - m)$ for $X \sim \pi$, and let ν be the law of Z , which is isotropic and logconcave. Under this change of variables, the tilted law π_t becomes

$$\nu_s(dz) \propto \exp(-s \|\Sigma^{1/2}z + m\|^2) \nu(dz) \quad \text{for } s = (2t)^{-1},$$

and $\|X\|^2 = \|\Sigma^{1/2}Z + m\|^2$. Hence, to bound $\text{Var}_{\pi_t}(\|X\|^2)$, we consider the following problem: for $M \succ 0$ and $u \in \mathbb{R}^n$, define $q_u(x) = \|M^{1/2}x + u\|^2$ under the tilt $\nu_s(dx) \propto e^{-sq_u(x)} \nu(dx)$ for $s \in [0, \infty]$ (later we can simply take $M = \Sigma$ and $u = m$). Write b_s and B_s for the mean and covariance of ν_s , and set $y_s = M^{1/2}b_s + u$ and $M_s = M^{1/2}B_sM^{1/2}$. Isotropizing ν_s and applying the definition of $\mathbf{Q}_n$ gives

$$\text{Var}_{\nu_s} q_u \lesssim \mathbf{Q}_n \text{tr}(M_s^2) + y_s^\top M_s y_s.$$

Hence, it suffices to control $\text{tr}(M_s^2)$ and the contribution of the moving mean y_s . The Brascamp–Lieb inequality yields $M_s \preceq (2s)^{-1} \text{Id}$, but applying this bound independently in every direction loses a factor of n and becomes vacuous as $s \downarrow 0$. As mentioned earlier, the plausible alternative of proving $\|B_s\|_{\text{op}} \lesssim \|B_0\|_{\text{op}}$ is also false: covariance operator-norm stability under Gaussian tilting is false in general. This suggests controlling the eigenvalues *collectively*, rather than the largest eigenvalue.

Collective spectral control of M_s . Consider first the centered case $u = 0$. Let $\lambda_i(s)$ be the eigenvalues of M_s , and let P_s and N_s denote the total increase and decrease of these eigenvalues relative to $M_0 = M$:

$$N_s := \sum_{\lambda_i(s) < \lambda_i(0)} (\lambda_i(0) - \lambda_i(s)) \quad \text{and} \quad P_s := \sum_{\lambda_i(s) > \lambda_i(0)} (\lambda_i(s) - \lambda_i(0)).$$

Note that $\text{tr } M_s - \text{tr } M_0 = P_s - N_s$. The score identity (Lemma 2.3) gives $\partial_s \mathbb{E}_s[X^\top M X] = -\text{Var}_{\nu_s} q_u$, which relates N_s , P_s , and $\|y_s\|^2$ as follows: $\Delta_s := \int_0^s \text{Var}_{\nu_r} q_u dr = N_s - P_s - \|y_s\|^2$, so $P_s + \|y_s\|^2 \leq N_s$. Also, differentiating $\lambda_i(s)$ along the same path gives $\partial_s \log \lambda_i(s) = -\text{cov}_s(Z_{s,i}^2, q_u)$, where Z_s is the random variable for isotropization of ν_s . The log-spectrum estimate yields $N_s^2 \leq s \mathbf{Q}_n \text{tr}(M^2) \Delta_s$. Since $\Delta_s \leq N_s$, both spectral gain and squared mean displacement are $O(s)$. Combining these relations yields $\text{tr}(M_s^2) \leq \text{tr}(M^2) (1 + \mathbf{Q}_n)$ and $y_s^\top M_s y_s \leq \frac{1}{2} \text{tr}(M^2) \mathbf{Q}_n$ with $\mathbf{Q}_n \lesssim \log n$ (Lemma 2.5). Note that we avoid the naïve factor- n loss: instead of using the same worst-case bound on every eigenvalue, we control the ℓ_2 -norm of the entire log-spectrum collectively. The detailed proof appears in §2.1.

Extension to the shifted case. When $u \neq 0$, the same argument starts from $y_0 = u$, and the identity becomes $\Delta_s = N_s - P_s + \|u\|^2 - \|y_s\|^2$. This change adds $O(\|u\|^2/s)$ to $\text{Var}_{\nu_s} q_u$, which is useful but vacuous near the untilted endpoint (i.e., $s = 0$). At that endpoint, the same decomposition and the quadratic-form estimate directly bound $\text{Var}_{\nu} q_u$ in terms of $G := \text{tr}(M^2) + u^\top M u$. Moreover, the score identity and reverse Hölder inequality for quadratic polynomials give $|\partial_s \text{Var}_{\nu_s} q_u| \lesssim (\text{Var}_{\nu_s} q_u)^{3/2}$. Solving this Grönwall-type inequality, we use the resulting bound up to $s_0 \asymp [(\mathbf{Q}_n + 4)G]^{-1/2}$ and the spectral estimate thereafter. Concatenating the two regimes yields the additional term $\|u\|^2 \sqrt{G}$ in Theorem 2.1. Putting these together, we obtain the bound $V = \tilde{O}(R^2 L \wedge R^3 \Lambda^{1/2})$.

1.2.2 From variance stability to faster Gaussian cooling

Rényi-divergence between annealing distributions. The heuristic calculation above suggests the natural scale of $\alpha \asymp \sigma^2/\sqrt{V}$. To make this quantitative, let $F(s)$ be the logarithm of the normalization constant of $\nu_s(dx) \propto e^{-s\|x\|^2} \pi(dx)$. Differentiation gives $F''(s) = \text{Var}_{\nu_s}(\|X\|^2)$. By the definition of q -Rényi divergence, one can check that $R_q(\nu_s \parallel \nu_{s'}) = [F(s + (q-1)\Delta) - qF(s) + (q-1)F(s-\Delta)]/(q-1)$ for $\Delta := s - s'$. Using Taylor expansion, for $I = [s', s' + q\Delta]$, we have

$$F(s + (q-1)\Delta) \leq F(s) + (q-1)\Delta F'(s) + \frac{V_I}{2} (q-1)^2 \Delta^2 \quad \text{and} \quad F(s-\Delta) \leq F(s) - \Delta F'(s) + \frac{V_I}{2} \Delta^2,$$

so

$$R_q(\nu_s \parallel \nu_{s'}) \leq \frac{q}{2} (s - s')^2 \sup_{t \in I} \text{Var}_{\nu_t}(\|X\|^2).$$

If $s = (2\sigma^2)^{-1}$, consecutive targets remain $O(1)$ -close in R_q under the update $\sigma^2 \leftarrow \sigma^2(1 + \Theta(\sigma^2/\sqrt{qV}))$. The same inequality shows that $\sigma^2 \gtrsim \sqrt{qV}$ is sufficient for warmness with respect to the target π , so we set $\sigma_{\text{last}}^2 \asymp \sqrt{qV}$. As discussed in Remark 3.3, the resulting bound recovers prior closeness bounds in [CV18, KV25a, KV25c]. Prior works use the following bound:

$$\text{Var}_{\nu_s}(\|X\|^2) \leq 4\sigma^2 \mathbb{E}_{\nu_s}[\|X\|^2] \lesssim \sigma^2 R^2$$

by (PI) (or Brascamp–Lieb); this σ^2 -dependent bound becomes loose as σ^2 increases.

Special case: uniform sampling. For a convex body containing the unit ball, we can sample the initial target at $\sigma_0^2 = 1/n$ exactly by Gaussian rejection sampling. We then follow this schedule using **ln-and-Out** (the proximal sampler) for truncated Gaussians [KV25c]. At scale σ^2 , each phase (i.e., sampling initialized at an $O(1)$ -warm start in R_q) requires $\tilde{O}(n^2\sigma^2)$ queries, and doubling the initial σ^2 requires $O(\sqrt{qV}/\sigma^2)$ phases. Thus, each doubling costs $\tilde{O}(n^2\sqrt{qV})$ queries. Since there are $O(\log(qnV))$ doublings, by the stability estimate, the total complexity of Gaussian annealing is $\tilde{O}(q^{1/2}n^2 \min\{R\sqrt{L}, R^{3/2}\Lambda^{1/4}\})$ queries; for a nearly isotropic target, this is $\tilde{O}(q^{1/2}n^{5/2})$. Using hypercontractivity of the proximal sampler [KV25c], this procedure propagates R_q -warmness from one target to the next. The details are given in §3.

General case: logconcave sampling. Given a convex potential V and an evaluation oracle for it, we use the lifting technique [KV25b]; instead of sampling directly from $\pi \propto e^{-V}$, we sample from a density proportional to $e^{-nt} \mathbb{1}_{\mathcal{K}}(x, t)$, where $\mathcal{K} = \{(x, t) : V(x) \leq nt\}$ is the epigraph of V/n . Its X -marginal is exactly proportional to $e^{-V(x)}$. Suitable truncation $\tilde{\mathcal{K}}$ of $\mathcal{K}$ contains at least half of the lifted measure and ensures a finite log-Sobolev constant. Since the truncated X -marginal has density at most twice that of π , its parameters R, L, Λ are bounded by universal multiples of those of π . Hence, the required thin-shell upper bound increases by at most a universal factor.

On the truncated lift, we use the two-parameter annealing distribution from [KV25b]:

$$\mu_{\sigma^2, \rho}(dx, dt) \propto \exp\left(-\frac{\|x\|^2}{2\sigma^2} - \rho t\right) \mathbb{1}_{\tilde{\mathcal{K}}}(x, t) dx dt.$$

Using the proximal sampler for this distribution [KV25c], we follow the path $(n^{-1}, 1) \rightarrow (n^{-1}, n) \rightarrow (\sigma_{\text{last}}^2, n)$ in the (σ^2, ρ) -plane, streamlining the annealing schedules proposed in [KV25a, KV25c]. The first segment changes only ρ while keeping the strongly logconcave Gaussian component fixed. Once $\rho = n$, the X -marginal is exactly the Gaussian tilt of the truncated target, so the second segment can reuse the schedule and theory developed for uniform sampling.

For the first phase $(n^{-1}, 1) \rightarrow (n^{-1}, n)$, we establish a new closeness bound. As in the analysis of the R_q bound between consecutive annealing distributions, we have to bound $\text{Var}_{(\sigma^2, \rho)} T := \text{Var}_{\mu_{\sigma^2, \rho}} T$. Varentropy of the joint logconcave density [FMW16], together with the Poincaré bound for its strongly logconcave X -marginal, gives $\text{Var}_{(n^{-1}, \rho)} T \lesssim n/\rho^2$ for $1 \leq \rho \leq n$. Hence, for $q_0 = \tilde{\Theta}(1)$, we multiply ρ by $1 + \Theta(1/\sqrt{q_0(q_0 + n)})$ at each phase. There are $\tilde{O}(\sqrt{n})$ such phases, and each costs $\tilde{O}(n^2)$ queries (since the proximal sampler costs $\tilde{O}(n^2(\sigma^2 \vee 1))$ from an $O(1)$ -warm start), using $\tilde{O}(n^{5/2})$ queries. We maintain the same base order q_0 while increasing σ^2 to 1. Thereafter, using the hypercontractivity of the proximal sampler again (to boost the Rényi order from q_0 to q), we use the thin-shell update up to σ_{last}^2 . These Gaussian-annealing phases cost $\tilde{O}(n^2\sqrt{qV})$ queries as in the uniform case. Finally, data processing under $(x, t) \mapsto x$ and the constant-mass truncation transfer the lifted warm start to the original target.

1.3 Preliminaries

We use the same symbol for a distribution and its density. Both $a \lesssim b$ and $a = O(b)$ mean $a \leq cb$ for a universal constant $c > 0$. $a = \Omega(b)$ means $a \gtrsim b$, and $a \asymp b$ means $a = O(b)$ and $a = \Omega(b)$. Lastly, $a = \tilde{O}(b)$ means $a = O(b \text{ polylog } b)$. For a positive semidefinite matrix Σ , $\|\Sigma\|_{\text{op}}$ denotes the operator norm of Σ . We use $a \vee b$ and $a \wedge b$ to denote their maximum and minimum, respectively. $B(x, r)$ denotes an ℓ_2 -ball of radius r centered at x .

Logconcavity. We call a function $f : \mathbb{R}^n \rightarrow [0, \infty)$ *logconcave* if $-\log f$ is convex in $\mathbb{R}^n$, and call a probability measure π (or distribution) *logconcave* if it has a logconcave density with respect to Lebesgue measure. We assume that all logconcave distributions considered in this work are full-dimensional⁴. For $t \geq 0$, π is called *t-strongly logconcave* if $-\log \pi$ is t -strongly convex (i.e., $-\log \pi - \frac{t}{2} \|\cdot\|^2$ is convex). Logconcavity is preserved under multiplication, and a classical result by Prékopa and Leindler ensures that convolution also preserves logconcavity. A distribution is called *isotropic* if its barycenter is at the origin and its covariance matrix is the identity. Logconcave distributions have exponentially decaying tails and hence finite moments of all orders.

Probability. Let μ and ν denote probability measures on $\mathbb{R}^n$ with $\mu \ll \nu$. For $q > 1$, we define the warmness convention $M_q = \|\mathrm{d}\mu/\mathrm{d}\nu\|_{L^q(\nu)}$. The q -Rényi divergence is defined as $R_q(\mu \parallel \nu) := \frac{1}{q-1} \log \|\frac{\mathrm{d}\mu}{\mathrm{d}\nu}\|_{L^q(\nu)}^q$, and the Rényi-infinity divergence is $R_\infty(\mu \parallel \nu) := \log \text{ess sup}_\nu \frac{\mathrm{d}\mu}{\mathrm{d}\nu}$. Recall monotonicity in the order: $R_q \leq R_{q'}$ for $q \leq q'$. A weak triangle inequality holds for Rényi divergence: for any $q > 1$ and probability measures μ, ν, π , it holds that

$$R_q(\mu \parallel \pi) \leq \frac{2q}{2q-1} R_{2q-1}(\mu \parallel \nu) + R_{2q}(\nu \parallel \pi).$$

We refer to [vEH14, Mir17] for more properties of Rényi divergence.

Functional inequalities. A probability measure π on $\mathbb{R}^n$ is said to satisfy a *Poincaré inequality* with constant C if for any locally Lipschitz function $f \in L^2(\pi)$,

$$\text{Var}_\pi f := \int \left(f - \int f \mathrm{d}\pi\right)^2 \mathrm{d}\pi \leq C \int |\nabla f|^2 \mathrm{d}\pi, \quad (\text{PI})$$

and the smallest such C is called the Poincaré constant $C_{\text{PI}}(\pi)$. A probability measure π on $\mathbb{R}^n$ is said to satisfy a *logarithmic Sobolev inequality* with constant C if for any locally Lipschitz function $f \in L^2(\pi)$,

$$\text{Ent}_\pi(f^2) := \int f^2 \log f^2 \mathrm{d}\pi - \int f^2 \mathrm{d}\pi \cdot \log \int f^2 \mathrm{d}\pi \leq 2C \int |\nabla f|^2 \mathrm{d}\pi, \quad (\text{LSI})$$

and the smallest such C is referred to as the log-Sobolev constant $C_{\text{LSI}}(\pi)$. In general, (LSI) is strictly stronger than (PI). By the Bakry–Émery criterion, every t -strongly logconcave probability measure π satisfies $C_{\text{LSI}}(\pi) \leq 1/t$ [BGL14].

Historical note and AI use. In early June, during the conference “Scalable MCMC Sampling” held at the FIM–Institute for Mathematical Research at ETH Zürich, the authors realized how to handle the $n^{8/3}$ lower bound example from [KV25a] in $n^{2.5}$ oracle complexity. They formulated and discussed the thin-shell stability conjecture and promising approaches to proving it. The first author then used GPT 5.5 and 5.6 Pro to obtain a preliminary version of the proof of Theorem 1.1 through guided prompting. The authors verified, streamlined, and wrote the full proof and its consequences.

⁴Otherwise, we may work on the affine subspace supporting the distribution.

2 Thin-shell stability of Gaussian tilts

In this section, we show that the thin-shell variance remains bounded for Gaussian tilts; motivated by (1.1), a natural anisotropic version of the conjecture would be $\text{Var}_{\nu_s}(X^\top M X) \lesssim_{\log n} \text{tr}(M^2)$ for every $s \geq 0$. In §2.1, we provide the proof of this conjecture, and in §2.2, we provide the following generalization to shifted quadratic functions:

Theorem 2.1 (Shifted anisotropic quadratic tilt). *Let ν be an isotropic logconcave probability measure on $\mathbb{R}^n$, $M \succeq 0$, and $u \in \mathbb{R}^n$. Set $q_u(x) = \|M^{1/2}x + u\|^2$ and define the probability measure $\nu_s(dx) \propto e^{-sq_u(x)} \nu(dx)$ for $s \geq 0$. Then, for every $s \geq 0$,*

$$\text{Var}_{\nu_s} q_u \lesssim (1 + Q_n)^2 (\text{tr}(M^2) + \|u\|^2 \sqrt{\text{tr}(M^2) + u^\top M u}).$$

Using this result, we readily obtain the thin-shell stability result (Theorem 1.1) needed for faster Gaussian cooling.

2.1 Proof idea: the centered case

We first present the proof in the centered case of Theorem 2.1 (i.e., $u = 0$).

Theorem 2.2 (Centered anisotropic quadratic tilt). *Let ν be an isotropic logconcave probability measure on $\mathbb{R}^n$, and $\nu_s(dx) \propto e^{-s\langle x, Mx \rangle} \nu(dx)$ for $s \geq 0$ and $M \succeq 0$. Then,*

$$\text{Var}_{\nu_s}(X^\top M X) \leq (2Q_n^2 + 6Q_n) \text{tr}(M^2) \quad \text{for every } s \geq 0. \quad (2.1)$$

We first prove the theorem for $M \succ 0$. Denote the barycenter and covariance of ν_s by $b_s = \mathbb{E}_s X$ and $B_s = \text{cov}_s X$. Under $x \mapsto y = T(x) := M^{1/2}x$, the pushforward $Y_s \sim T_\# \nu_s$ has barycenter $y_s = M^{1/2}b_s$ and covariance $M_s = M^{1/2}B_s M^{1/2}$. Isotropy of ν gives $b_0 = 0$ and $B_0 = \text{Id}$, so $y_0 = 0$ and $M_0 = M$. Although ν is centered, the quadratic tilt ν_s need not preserve its mean.

Controlling the covariance and the moving mean. Isotropize ν_s by $Z := M_s^{-1/2} M^{1/2}(X - b_s)$ for $X \sim \nu_s$. We now bound the variance of the target quadratic form $q(x) := x^\top M x$. We have the decomposition

$$q(x) = x^\top M x = (x - b_s)^\top M (x - b_s) + 2b_s^\top M x - b_s^\top M b_s = z^\top M_s z + 2y_s^\top M_s^{1/2} z + b_s^\top M b_s.$$

Hence, $\mathbb{E}_s q = \text{tr } M_s + \|y_s\|^2$ and $q - \mathbb{E}_s q = (Z^\top M_s Z - \text{tr } M_s) + 2y_s^\top M_s^{1/2} Z$. Since $\text{Var}_s q = \|q - \mathbb{E}_s q\|_{L^2(\nu_s)}^2$, the triangle inequality in L^2 gives

$$\sqrt{\text{Var}_s q} \leq \|z^\top M_s z - \text{tr } M_s\|_{L^2(\nu_s)} + 2\|y_s^\top M_s^{1/2} z\|_{L^2(\nu_s)} \leq \sqrt{Q_n \text{tr}(M_s^2)} + 2\sqrt{y_s^\top M_s y_s}. \quad (2.2)$$

Here, (1.1) bounds the first term by $[Q_n \text{tr}(M_s^2)]^{1/2}$, while $\|y_s^\top M_s^{1/2} z\|_{L^2(\nu_s)}^2 = \mathbb{E}_s[(y_s^\top M_s^{1/2} Z)^2] = y_s^\top M_s y_s$. Therefore, the proof reduces to controlling (i) the *covariance contribution* $\text{tr}(M_s^2)$ and (ii) the *moving-mean contribution* $y_s^\top M_s y_s$.

We now bound these two terms. By Brascamp–Lieb, $M_s \preceq (2s)^{-1} \text{Id}$, so each eigenvalue $(\lambda_i(s))_{i \in [n]}$ of M_s is bounded by $(2s)^{-1}$. However, applying this bound directly to $\text{tr}(M_s^2)$ would lose a factor of the dimension n and would blow up as $s \downarrow 0$. To refine the analysis, we track changes in the eigenvalues *collectively* relative to $M_0 = M$ and relate their total increase to their total decrease,

rather than applying the same worst-case bound to all n eigenvalues. To make this precise, let us define the total spectral “loss” and “gain” by

$$N_s := \sum_{\lambda_i(s) < \lambda_i(0)} (\lambda_i(0) - \lambda_i(s)) \quad \text{and} \quad P_s := \sum_{\lambda_i(s) > \lambda_i(0)} (\lambda_i(s) - \lambda_i(0)).$$

Note that $\text{tr } M_s - \text{tr } M_0 = P_s - N_s$.

We first track how much $\text{tr}(M_s^2)$ has changed from $\text{tr}(M_0^2)$: since $\text{tr}(M_s^2) - \text{tr}(M_0^2) = \sum_i (\lambda_i(s)^2 - \lambda_i(0)^2)$,

$$\text{tr}(M_s^2) - \text{tr}(M_0^2) \leq \sum_{\lambda_i(s) > \lambda_i(0)} (\lambda_i(s)^2 - \lambda_i(0)^2) \stackrel{(i)}{\leq} \frac{1}{s} \sum_{\lambda_i(s) > \lambda_i(0)} (\lambda_i(s) - \lambda_i(0)) = \frac{P_s}{s},$$

where (i) follows from $\lambda_i(s) + \lambda_i(0) \leq 2\lambda_i(s) \leq s^{-1}$ for an eigenvalue with $\lambda_i(s) > \lambda_i(0)$. Thus, the two target terms are bounded as

$$\text{tr}(M_s^2) \leq \text{tr}(M_0^2) + \frac{P_s}{s} \quad \text{and} \quad y_s^\top M_s y_s \leq \frac{\|y_s\|^2}{2s}. \quad (2.3)$$

Hence, it suffices to bound P_s and $\|y_s\|^2$ in terms of s . An elementary identity couples P_s , N_s , and $\|y_s\|^2$.

Coupling P_s , N_s , and $\|y_s\|^2$. We first recall the following identity.

Lemma 2.3. *Let ν be a probability measure, $q : \mathbb{R}^n \rightarrow \mathbb{R}$ measurable, and define an exponential tilt $\nu_s(dx) \propto e^{-sq(x)} \nu(dx)$ on an interval where the tilt is well-defined. If f_s is differentiable and differentiation can be passed under the defining integrals, then $\partial_s \mathbb{E}_s f_s = \mathbb{E}_s[\partial_s f_s] - \text{cov}_s(f_s, q)$, where the identity is interpreted entrywise when f_s is vector- or matrix-valued.*

Proof. Recall an elementary property of the score function: $\mathbb{E}_s[\partial_s \log \nu_s] = \int \partial_s \nu_s = 0$ due to $\partial_s \nu_s = \nu_s \partial_s \log \nu_s$. Differentiating the expectation under the integral gives

$$\partial_s \mathbb{E}_s f_s = \int (\partial_s f_s) \nu_s + \int f_s \partial_s \nu_s = \mathbb{E}_s[\partial_s f_s] + \mathbb{E}_s[f_s \partial_s \log \nu_s] = \mathbb{E}_s[\partial_s f_s] + \text{cov}_s(f_s, \partial_s \log \nu_s),$$

where the last equality follows from $\mathbb{E}_s[\partial_s \log \nu_s] = 0$. The claim follows from $\partial_s \log \nu_s = -(q - \mathbb{E}_s q)$. $\square$

Using this, $\partial_s \mathbb{E}_s q = -\text{Var}_{\nu_s} q$. Since $\mathbb{E}_s q = \text{tr } M_s + \|y_s\|^2$ and $\mathbb{E}_0 q = \text{tr } M_0$,

$$\Delta_s := \int_0^s \text{Var}_{\nu_r} q \, dr = \mathbb{E}_0 q - \mathbb{E}_s q = N_s - P_s - \|y_s\|^2. \quad (2.4)$$

Since $\Delta_s \geq 0$ (as $\text{Var}_{\nu_r} q \geq 0$), we have $P_s + \|y_s\|^2 \leq N_s$. It remains to bound N_s in terms of s .

For an index with $\lambda_i(s) < \lambda_i(0)$, the elementary inequality $1 - r \leq -\log r$ for $0 < r \leq 1$ gives $\lambda_i(0) - \lambda_i(s) \leq \lambda_i(0) |\log \frac{\lambda_i(s)}{\lambda_i(0)}|$. By Cauchy–Schwarz over the eigenvalues,

$$N_s \leq \left(\sum_i \lambda_i(0)^2 \right)^{1/2} \left(\sum_i \log^2 \frac{\lambda_i(s)}{\lambda_i(0)} \right)^{1/2} \leq (\text{tr}(M_0^2))^{1/2} \left(\sum_i \log^2 \frac{\lambda_i(s)}{\lambda_i(0)} \right)^{1/2}. \quad (2.5)$$

To obtain an explicit factor of s on the right-hand side, we apply Cauchy–Schwarz in time:

$$\sum_i \log^2 \frac{\lambda_i(s)}{\lambda_i(0)} = \sum_i \left(\int_0^s \partial_r \log \frac{\lambda_i(r)}{\lambda_i(0)} \, dr \right)^2 \leq s \int_0^s \sum_i (\partial_r \log \lambda_i(r))^2 \, dr. \quad (2.6)$$

Spectral analysis of M_s . We now analyze how fast each λ_i changes:

Lemma 2.4. *Let $u_i = u_i(s)$ be a suitable unit eigenvector of M_s such that $M_s u_i = \lambda_i(s) u_i$, and let $Z_i := u_i^\top Z$. Then,*

$$\partial_s \log \lambda_i(s) = -\text{cov}_s(Z_i^2, q). \quad (2.7)$$

Proof. By Lemma 2.3 for the centered covariance,

$$B'_s = \mathbb{E}_s[\partial_s((X - b_s)(X - b_s)^\top)] - \text{cov}_s((X - b_s)(X - b_s)^\top, q) = -\text{cov}_s((X - b_s)(X - b_s)^\top, q), \quad (2.8)$$

where the terms containing $\partial_s b_s$ vanish as $\mathbb{E}_s(X - b_s) = 0$. Since $M_s = M^{1/2} B_s M^{1/2}$, (2.8) implies

$$\partial_s M_s = -M^{1/2} \text{cov}_s((X - b_s)(X - b_s)^\top, q) M^{1/2}. \quad (2.9)$$

By perturbation theory for symmetric matrix paths [Kat95, Theorem 5.4 and Theorem 6.8], the eigenvalues $\lambda_i(s)$ of M_s may be labeled differentially, and at each s , one may choose an orthonormal eigenbasis $(u_i)_{i=1}^n$ of M_s such that $\partial_s \lambda_i(s) = u_i^\top (\partial_s M_s) u_i$. Hence, by (2.9),

$$\partial_s \lambda_i(s) = -\text{cov}_s\{(u_i^\top M^{1/2}(X - b_s))^2, q\}. \quad (2.10)$$

Using $M^{1/2}(X - b_s) = M_s^{1/2} Z$ and $M_s^{1/2} u_i = \sqrt{\lambda_i(s)} u_i$,

$$u_i^\top M^{1/2}(X - b_s) = u_i^\top M_s^{1/2} Z = \sqrt{\lambda_i(s)} Z_i.$$

Substituting this into (2.10) yields $\partial_s \lambda_i(s) = -\lambda_i(s) \text{cov}_s(Z_i^2, q)$, which completes the proof. $\square$

We use this identity to collect bounds on the key quantities (such as M_s and y_s) along the path indexed by s , from which the theorem will follow. Recall that $\nu_s \propto \exp(-sx^\top M x) \nu$ is a Gaussian tilt of an isotropic logconcave probability measure ν in $\mathbb{R}^n$ for $s \geq 0$, and that $b_s = \mathbb{E}_s X$ and $B_s = \text{cov}_s X$ are its barycenter and covariance, respectively.

Lemma 2.5 (Properties of M_s and y_s). *The barycenter $y_s = M^{1/2} b_s$ and covariance $M_s = M^{1/2} B_s M^{1/2}$ satisfy the following bounds for $s \geq 0$:*

- $P_s + \|y_s\|^2 \leq N_s \leq s \cdot \text{tr}(M_0^2) Q_n$
- $\text{tr}(M_s^2) \leq \text{tr}(M_0^2) (1 + Q_n)$
- $y_s^\top M_s y_s \leq \frac{1}{2} \text{tr}(M_0^2) Q_n$

Proof. For every unit vector $\alpha \in \mathbb{R}^n$, by Cauchy–Schwarz followed by the definition of Q_n for the quadratic form $\sum_i \alpha_i \alpha_i Z_i^2$,

$$\left| \sum_i \alpha_i \partial_s \log \lambda_i(s) \right| = \left| \text{cov}_s\left(\sum_i \alpha_i Z_i^2, q\right) \right| \leq \sqrt{Q_n} \|\alpha\| \sqrt{\text{Var}_{\nu_s} q}.$$

Hence, $\sum_i [\partial_s \log \lambda_i(s)]^2 \leq Q_n \text{Var}_{\nu_s} q$. Substituting this into (2.6),

$$\sum_i \log^2 \frac{\lambda_i(s)}{\lambda_i(0)} \leq s \int_0^s \sum_i (\partial_r \log \lambda_i(r))^2 dr \leq s Q_n \int_0^s \text{Var}_{\nu_r} q dr = s Q_n \Delta_s. \quad (2.11)$$

Using $\Delta_s \leq N_s$ and substituting this into (2.5), $N_s^2 \leq \text{tr}(M_0^2) \sum_i \log^2 \frac{\lambda_i(s)}{\lambda_i(0)} \leq s \text{tr}(M_0^2) Q_n N_s$. Hence,

$$(P_s + \|y_s\|^2 \leq) N_s \leq s \cdot \text{tr}(M_0^2) Q_n.$$

Thus, $P_s, N_s, \|y_s\|^2 = O(s)$ as desired. Substituting this into (2.3), we obtain $\text{tr}(M_s^2) \leq \text{tr}(M_0^2) (1 + Q_n)$ and $y_s^\top M_s y_s \leq \frac{1}{2} \text{tr}(M_0^2) Q_n$. $\square$

Substituting the second and third bounds into (2.2), we obtain, for every $s > 0$,

$$\text{Var}_s q \leq (2Q_n^2 + 6Q_n) \text{tr}(M^2),$$

which completes the proof of Theorem 2.2. When $M \succeq 0$, we apply the result to $M + \varepsilon I_n$ and let $\varepsilon \downarrow 0$.

Remark 2.6 (Collective control of eigenvalues). The proof controls the eigenvalues of M_s *collectively*, rather than individually. For example, considering the total continuous change in the trace (specifically, N_s and P_s terms), we obtain a useful bound. If we attempt to do this for each eigenvalue, then we may obtain some bound on each, but summing them up in a naïve way would lead to a factor of n . A further consequence is the second item of Lemma 2.5 (i.e., the stability of the Schatten 2-norm along the Gaussian tilt):

$$\text{tr}(M_s^2) \leq \text{tr}(M_0^2) (1 + Q_n).$$

As noted earlier, this type of bound does not hold for individual eigenvalues such as the largest eigenvalue (i.e., the operator norm).

2.2 Stability under shifted Gaussian tilts

We now consider the shifted quadratic: $q_u(x) = \|M^{1/2}x + u\|^2$. In this case, the initial transformed mean is $y_0 = u$, and we retain the same variance decomposition:

$$\sqrt{\text{Var}_s q} \leq \sqrt{Q_n \text{tr}(M_s^2)} + 2\sqrt{y_s^\top M_s y_s}. \quad (2.12)$$

The identity becomes $\Delta_s = N_s - P_s + \|u\|^2 - \|y_s\|^2$, so $P_s + \|y_s\|^2 \leq N_s$ is replaced by the weaker bound $P_s, \|y_s\|^2 \leq \|u\|^2 + N_s$. Applying Brascamp–Lieb again, this extra $\|u\|^2$ term yields $\|u\|^2/s$. The resulting bound is useful for moderate and large s but vacuous at the untilted endpoint (i.e., $s = 0$).

The proof follows the same steps and separately treats the regime near $s = 0$, solving the Grönwall-type inequality $|v'_s| \lesssim v_s^{3/2}$.

Proof of Theorem 2.1. Assume $M \succ 0$ and abbreviate $q = q_u$. Define $v_s := \text{Var}_s q$, $G := \text{tr}(M^2) + u^\top M u$, and

$$b_s := \mathbb{E}_s X, \quad B_s := \text{cov}_s X, \quad M_s := M^{1/2} B_s M^{1/2}, \quad y_s := M^{1/2} b_s + u.$$

By isotropy, we have $B_0 = \text{Id}$, $M_0 = M$, and $y_0 = u$.

By Lemma 2.3, we similarly have

$$(\mathbb{E}_s q)' = -v_s, \quad \Delta_s := \mathbb{E}_0 q - \mathbb{E}_s q = \int_0^s v_r dr, \quad B'_s = -\text{cov}_s((X - b_s)(X - b_s)^\top, q).$$

Let $\lambda_i(s)$ be an absolutely continuous labeling of the eigenvalues of M_s . The calculation in the preceding subsection gives the same identity: $\partial_s \log \lambda_i(s) = -\text{cov}_s(Z_i^2, q)$. Hence,

$$\sum_i (\partial_s \log \lambda_i(s))^2 \leq Q_n v_s \quad \text{and} \quad \sum_i \log^2 \frac{\lambda_i(s)}{\lambda_i(0)} \leq Q_n s \Delta_s.$$

Define N_s and P_s as before. Since $\mathbb{E}_s q = \text{tr} M_s + \|y_s\|^2$,

$$\Delta_s = N_s - P_s + \|u\|^2 - \|y_s\|^2 \quad \text{and} \quad \Delta_s, P_s, \|y_s\|^2 \leq \|u\|^2 + N_s.$$

Then, $N_s \leq \sqrt{\mathbf{Q}_n \operatorname{tr}(M^2) s \Delta_s} \leq \sqrt{\mathbf{Q}_n \operatorname{tr}(M^2) s (\|u\|^2 + N_s)}$. Solving this,

$$N_s \leq \mathbf{Q}_n \operatorname{tr}(M^2) s + \|u\| \sqrt{\mathbf{Q}_n \operatorname{tr}(M^2) s}.$$

Combining this with the bounds on P_s and y_s gives

$$\operatorname{tr}(M_s^2) \lesssim (1 + \mathbf{Q}_n) \operatorname{tr}(M^2) + \frac{\|u\|^2}{s} \quad \text{and} \quad y_s^\top M_s y_s \lesssim \mathbf{Q}_n \operatorname{tr}(M^2) + \frac{\|u\|^2}{s}.$$

Substituting these bounds into (2.12),

$$v_s \lesssim (\mathbf{Q}_n + \mathbf{Q}_n^2) \operatorname{tr}(M^2) + (1 + \mathbf{Q}_n) \frac{\|u\|^2}{s}. \quad (2.13)$$

The last term is the only singular term introduced by the shift, and it becomes vacuous as s goes to 0. However, at $s = 0$, the same decomposition gives $v_0 \leq (\mathbf{Q}_n + 4) G$. To combine the two regimes, we propagate the endpoint bound to small s .

Handling small s . By Lemma 2.3, $v'_s = -\mathbb{E}_s[(q - \mathbb{E}_s q)^3]$. By the polynomial reverse Hölder inequality [CW01, Theorem 7], $|v'_s| \lesssim v_s^{3/2}$. Let $\bar{v} := (\mathbf{Q}_n + 4) G$ and $s_0 = c\bar{v}^{-1/2}$ for sufficiently small $c > 0$. Since $M \succ 0$ and ν_s is full-dimensional, $v_s > 0$. Integrating the differential inequality, for $s \in [0, s_0]$, we have $v_s \leq 4\bar{v}$. Since $u^\top M u \leq \|u\|^2 \|M\|_{\text{op}} \leq \|u\|^2 \sqrt{G}$, we have $G = \operatorname{tr}(M^2) + u^\top M u \leq \operatorname{tr}(M^2) + \|u\|^2 \sqrt{G}$. Hence, $v_s \lesssim (1 + \mathbf{Q}_n) (\operatorname{tr}(M^2) + \|u\|^2 \sqrt{G})$.

For $s \geq s_0$, the pointwise bound (2.13) yields

$$v_s \lesssim (\mathbf{Q}_n + \mathbf{Q}_n^2) \operatorname{tr}(M^2) + (1 + \mathbf{Q}_n) \|u\|^2 \sqrt{(\mathbf{Q}_n + 4) G}.$$

Therefore,

$$v_s \lesssim (1 + \mathbf{Q}_n)^2 (\operatorname{tr}(M^2) + \|u\|^2 \sqrt{G}).$$

This proves the claim when $M \succ 0$. For $M \succeq 0$, apply the result to $M + \varepsilon I_n$ and let $\varepsilon \downarrow 0$; dominated convergence follows from the finite fourth moment of ν . $\square$

We now transfer the shifted stability bound to the original coordinates.

Proof of Theorem 1.1. Since π is full-dimensional, $\Sigma \succ 0$. Let $Z = \Sigma^{-1/2}(X - m)$ for $X \sim \pi$, and let ν be the law of Z , which is isotropic and logconcave. Under the same change of variables, the tilted law π_t becomes

$$\nu_s(dz) \propto \exp(-s \|\Sigma^{1/2} z + m\|^2) \nu(dz) \quad \text{for } s = (2t)^{-1},$$

and $\|X\|^2 = \|\Sigma^{1/2} Z + m\|^2$. Applying Theorem 2.1 with $M = \Sigma$ and $u = m$ proves that for $t > 0$,

$$\operatorname{Var}_{\pi_t}(\|X\|^2) \lesssim (1 + \mathbf{Q}_n)^2 (\operatorname{tr}(\Sigma^2) + \|m\|^2 \sqrt{\operatorname{tr}(\Sigma^2) + m^\top \Sigma m}).$$

To obtain the simpler scales, note that

$$\operatorname{tr}(\Sigma^2) \leq \Lambda \operatorname{tr} \Sigma \leq R^2 \Lambda, \quad \operatorname{tr}(\Sigma^2) + m^\top \Sigma m \leq \Lambda (\operatorname{tr} \Sigma + \|m\|^2) = R^2 \Lambda, \quad \|m\|^2 \leq L.$$

Consequently,

$$\operatorname{tr}(\Sigma^2) + \|m\|^2 \sqrt{\operatorname{tr}(\Sigma^2) + m^\top \Sigma m} \leq R^2 \Lambda + LR\sqrt{\Lambda} \leq 2 \min\{R^2 L, R^3 \sqrt{\Lambda}\}.$$

Here, for the $R^2 L$ bound, the last step uses $\Lambda \leq L$ and $\sqrt{\Lambda} \leq R$; for the $R^3 \sqrt{\Lambda}$ bound, it uses $\sqrt{\Lambda} \leq R$ and $L \leq R^2$. $\square$

3 Faster annealing for uniform sampling

In §3.1, we relate variance stability to Rényi divergence between consecutive Gaussian annealing distributions. This determines the step size in the Gaussian variance σ^2 in our annealing scheme. Then, in §3.2, we give a faster version of Gaussian cooling, proving Theorem 1.2. To go through the main calculations, the reader might find it convenient to think of q as a fixed constant, say 2.

3.1 Rényi closeness of annealing distributions

For a probability measure ν , let $\nu_s(dx) \propto e^{-s\|x\|^2} \nu(dx)$ for $s \geq 0$, and $F(s) := \log \int e^{-s\|x\|^2} \nu(dx)$. Differentiating gives $F'(s) = -\mathbb{E}_{\nu_s}[\|X\|^2]$ and $F''(s) = \text{Var}_{\nu_s}(\|X\|^2)$. Recall q -Rényi divergence: $R_q(\mu \parallel \nu) := \frac{1}{q-1} \log \int (\frac{d\mu}{d\nu})^q d\nu$ for $q > 1$.

Lemma 3.1. *Let $q > 1$ and $s \geq s' \geq 0$. Then,*

$$R_q(\nu_s \parallel \nu_{s'}) \leq \frac{q}{2} (s - s')^2 \sup_{t \in [s', qs - (q-1)s']} \text{Var}_{\nu_t}(\|X\|^2). \quad (3.1)$$

Proof. Let $Z(s) := \int e^{-s\|x\|^2} d\nu(x)$. For $\Delta := s - s'$, the density ratio is $\frac{d\nu_s}{d\nu_{s'}}(x) = \frac{Z(s')}{Z(s)} e^{-\Delta\|x\|^2}$, and

$$\int \left(\frac{d\nu_s}{d\nu_{s'}}\right)^q d\nu_{s'} = \frac{Z(s')^{q-1} Z(qs - (q-1)s')}{Z(s)^q}.$$

Thus,

$$R_q(\nu_s \parallel \nu_{s'}) = \frac{F(s + (q-1)\Delta) - qF(s) + (q-1)F(s - \Delta)}{q-1}.$$

Let $I := [s', s + (q-1)\Delta]$ and $V_I := \sup_{t \in I} \text{Var}_{\nu_t}(\|X\|^2) = \sup_{t \in I} F''(t)$. By Taylor expansion,

$$F(s + (q-1)\Delta) \leq F(s) + (q-1)\Delta F'(s) + \frac{V_I}{2} (q-1)^2 \Delta^2,$$

$$F(s - \Delta) \leq F(s) - \Delta F'(s) + \frac{V_I}{2} \Delta^2.$$

Substituting these estimates into R_q ,

$$\begin{aligned} R_q(\nu_s \parallel \nu_{s'}) &\leq \frac{F(s) + (q-1)\Delta F'(s) + \frac{V_I}{2} (q-1)^2 \Delta^2 - qF(s) + (q-1)(F(s) - \Delta F'(s) + \frac{V_I}{2} \Delta^2)}{q-1} \\ &= \frac{qV_I}{2} \Delta^2, \end{aligned}$$

which completes the proof. $\square$

This result does not require logconcavity of ν . For $\sigma^2 > 0$, define $\mu_{\sigma^2} := \nu_{1/(2\sigma^2)}$. The preceding lemma gives the following bounds; apply Lemma 3.1 with $s = (2\sigma^2)^{-1}$ and $s' = (2\sigma^2(1+\alpha))^{-1}$, and take $s' = 0$ for the second bound.

Corollary 3.2 (Closeness). *Let $V := \sup_{s \geq 0} \text{Var}_{\nu_s}(\|X\|^2)$. For every $q > 1$, $\sigma^2 > 0$, and $\alpha > 0$,*

$$R_q(\mu_{\sigma^2} \parallel \mu_{\sigma^2(1+\alpha)}) \leq \frac{qV}{8\sigma^4} \left(\frac{\alpha}{1+\alpha}\right)^2 \leq \frac{qV\alpha^2}{8\sigma^4} \quad \text{and} \quad R_q(\mu_{\sigma^2} \parallel \nu) \leq \frac{qV}{8\sigma^4}.$$

In particular, the update $\sigma^2 \mapsto \sigma^2(1+\alpha)$ with $\alpha \lesssim \sigma^2/\sqrt{qV}$ maintains $O(1)$ -closeness, and $\sigma^2 \gtrsim \sqrt{qV}$ ensures terminal $O(1)$ -warmness.

Remark 3.3 (Comparison with prior closeness results). [KV25a, Lemma 4.5] bounds $R_q(\mu_{\sigma^2} \parallel \mu_{\sigma^2(1+\alpha)})$ by $qD^2\alpha^2/\sigma^2$ for compactly supported logconcave ν with support diameter D . Later, [KV25c, Lemma 4.2] gives the bound $q^2R^2\alpha^2/(2\sigma^2)$ for the same quantity when $\mathbb{E}_\nu[\|X\|^2] = R^2 \leq D^2$.

Our estimate (3.1) yields an improved bound. For every $t > 0$, the measure ν_t is $2t$ -strongly logconcave. By Brascamp–Lieb or (PI) with $C_{\text{PI}}(\nu_t) \leq 1/(2t)$,

$$\text{Var}_{\nu_t}(\|X\|^2) \leq \frac{1}{2t} \mathbb{E}_{\nu_t}[4\|X\|^2] = \frac{2}{t} \mathbb{E}_{\nu_t}[\|X\|^2] \leq \frac{2R^2}{t},$$

where the last inequality follows from $\mathbb{E}_{\nu_t}[\|X\|^2] \leq \mathbb{E}_\nu[\|X\|^2]$. For $s = (2\sigma^2)^{-1}$ and $s' = [2\sigma^2(1+\alpha)]^{-1}$ with $t \geq s'$, we have

$$R_q(\mu_{\sigma^2} \parallel \mu_{\sigma^2(1+\alpha)}) \leq \frac{q}{2} \left(\frac{\alpha}{2\sigma^2(1+\alpha)} \right)^2 \frac{2R^2}{s'} = \frac{qR^2\alpha^2}{2\sigma^2(1+\alpha)}.$$

This improves the q^2 -dependence of the second bound to q . It also recovers the compact-support bound in the present convex-body setting (due to $R^2 \leq D^2$).

3.2 Gaussian cooling through thin-shell stability

Let $\mathcal{K} \subset \mathbb{R}^n$ be a convex body with $B(0, 1) \subset \mathcal{K}$, and let π be uniform on $\mathcal{K}$. Fix a Rényi order q and take $V = \tilde{O}(R^2L \wedge R^3\sqrt{\Lambda})$ by Theorem 1.1. We propose the following annealing schedule:

- Let $\sigma_{\text{start}}^2 := 1/n$ and $\sigma_{\text{last}}^2 := \sqrt{qV}$ be the initial and terminal variance parameters.
- At phase i , the annealing distribution is $\mu_{\sigma_i^2} \propto \exp\{-\|x\|^2/(2\sigma_i^2)\} \cdot \mathbb{1}_{\mathcal{K}}$. For $\sigma_i^2 \in [\sigma_{\text{start}}^2, \sigma_{\text{last}}^2]$, define σ_{i+1}^2 recursively by

$$\sigma_{i+1}^2 := \sigma_i^2 \left(1 + \frac{\sigma_i^2}{\sqrt{qV}} \right). \quad (3.2)$$

By Corollary 3.2, consecutive distributions are $1/4$ -close at R_{2q} .

Next, we recall a sampling algorithm for truncated Gaussian distributions [KV25c]. Starting from $x_0 \sim \mu_0$, with step size $h > 0$ and threshold τ , set $s := \sigma^2/(h + \sigma^2) \in (0, 1)$. One iteration of PS_{Gauss} consists of the following two steps:

- [Forward] Draw $y_{k+1} \sim \mathcal{N}(x_k, hI_n)$.
- [Backward] Repeatedly draw $x_{k+1} \sim \mathcal{N}(sy_{k+1}, shI_n)$ until $x_{k+1} \in \mathcal{K}$. If the number of rejection trials exceeds τ , declare failure.

This sampler follows the annealing path efficiently because the previous annealing distribution is an $O(1)$ -warm start for the next target distribution.

Proposition 3.4 (Truncated-Gaussian sampler). *Let $\mu_{\sigma^2} \propto e^{-\|x\|^2/(2\sigma^2)} \mathbb{1}_{\mathcal{K}}(x)$ for a convex body $\mathcal{K} \supset B(0, 1)$ given by a membership oracle. Given $\varepsilon > 0$, $\eta \in (0, 1/2)$, $q \geq q_0 \geq 2$, and an initial distribution μ_0 with $M_{q_0} := \|\text{d}\mu_0/\text{d}\mu_{\sigma^2}\|_{L^{q_0}(\mu_{\sigma^2})} \leq 10$, initialize PS_{Gauss} with μ_0 and iterate it $N = \tilde{O}(n^2\sigma^2 \log \frac{q}{\varepsilon} \log \frac{1}{\eta})$ times with $h = (10n^2 \log Z)^{-1}$ and $\tau = Z^2 \log^3 Z$, where $Z = 16NM_2/\eta$. Assume $M_{q_0} \leq 10$ and $q_0 \geq 1 + \log \tau$.*

- The latter condition is ensured by $q_0 \geq 2 \vee O(\log \frac{N}{\eta})$.

- With probability at least $1-\eta$, PS_{Gauss} completes all N iterations without failure. Conditioned on this event, its output has law ν satisfying $R_q(\nu \parallel \mu_{\sigma^2}) \leq \varepsilon + 2 \log \frac{1}{1-\eta}$, using $\tilde{O}(n^2 \sigma^2 \log \frac{q}{\varepsilon} \log^5 \frac{1}{\eta})$ membership queries in expectation. The corresponding uncapped chain (i.e., $\tau = \infty$) has output law $\bar{\nu}$ satisfying $R_q(\bar{\nu} \parallel \mu_{\sigma^2}) \leq \varepsilon$.

Thus, we need $\tilde{O}(n^2 \sigma_i^2 \log q)$ membership queries at stage i . Let J be the first stage with $\sigma_J^2 \geq \sigma_{\text{last}}^2$.

Algorithm 3.5 (Thin-shell Gaussian cooling). **Input:** membership oracle for $\mathcal{K}$ with $B(0,1) \subset \mathcal{K}$, Rényi-order q , and $V = \tilde{O}(R^2 L \wedge R^3 \Lambda^{1/2})$.

1. Initialize $i = 0$, and draw $X_0 \sim \mu_0 := \mu_{\sigma_{\text{start}}^2}$ by rejection sampling from $N(0, \sigma_{\text{start}}^2 I_n)$.

2. While $\sigma_i^2 < \sigma_{\text{last}}^2$,

(a) compute σ_{i+1}^2 by (3.2): $\sigma_{i+1}^2 := \sigma_i^2 (1 + \frac{\sigma_i^2}{\sqrt{qV}})$.

(b) Iterate $\text{PS}_{\text{Gauss}} : R_q \rightarrow R_q$ for $N_{i+1} = \tilde{O}(n^2 \sigma_{i+1}^2 \log q \log J)$ iterations with target $\mu_{\sigma_{i+1}^2}$, accuracy $\varepsilon_{i+1} = 1/20$, and failure $\eta_{i+1} = 1/(100J)$. Use the parameters of the Truncated Gaussian Sampler (Proposition 3.4) with $Z_{i+1} := 160N_{i+1}/\eta_{i+1}$, restarting if a cap is hit. After each successful phase, replace i by $i+1$, and return the current point if $\sigma_i^2 \geq \sigma_{\text{last}}^2$.

Proof of Theorem 1.2. Let $q_{\min} := 2 \vee \Theta(\log(en^3 V))$. For $q < q_{\min}$, run the algorithm at order $q_{\min}$ and use monotonicity of Rényi divergence; this increases the cost by only a polylogarithmic factor. Thus, we may assume $q \geq q_{\min}$. Define $t_i := \sigma_i^2$. The initial rejection sampler produces $X_0 \sim \mu_{t_0}$ exactly and takes $\Theta(1)$ expected trials, since $t_0 = n^{-1}$ and $B(0,1) \subset \mathcal{K}$. As in [KV25c, High-level idea of analysis], we first consider the *ideal* truncated-Gaussian sampler where every threshold is removed (i.e., $\tau = \infty$) and then bound the bias introduced by the cap. Let $\bar{\nu}_i$ be its output law after phase i (so $\bar{\nu}_0 = \mu_{t_0}$).

The uncapped mixing estimate in Proposition 3.4, applied with input order q , output order $2q$, and accuracy $1/20$, ensures $R_{2q}(\bar{\nu}_i \parallel \mu_{t_i}) \leq \frac{1}{20}$ for $0 \leq i \leq J$. For $i < J$, the update is $\alpha_i = t_i/\sqrt{qV}$, so Corollary 3.2 gives $R_{2q}(\mu_{t_i} \parallel \mu_{t_{i+1}}) \leq 1/4$. By the weak triangle inequality,

$$R_q(\bar{\nu}_i \parallel \mu_{t_{i+1}}) \leq \frac{q-1/2}{q-1} R_{2q}(\bar{\nu}_i \parallel \mu_{t_i}) + R_{2q-1}(\mu_{t_i} \parallel \mu_{t_{i+1}}) \leq \frac{13}{40}.$$

Hence, the warmness satisfies $M_q < e$. The schedule gives $J = O(n\sqrt{qV})$ and $\sigma_i^2 \leq 2\sqrt{qV}$, so $\log \tau_i \lesssim \log(en^3 q(1 \vee V))$. Since $q \geq q_{\min}$, choosing the hidden constant in $q_{\min}$ sufficiently large ensures $q \geq 1 + \log \tau_i$ at every phase, as required by Proposition 3.4.

We now return to the capped algorithm with threshold τ . Couple a capped warm-start algorithm to the ideal warm-start algorithm, using the same randomness until a cap is hit, and let S be the event that all J phases succeed. The per-phase failure estimate in Proposition 3.4 is taken under the ideal input law. As in [KV25c, High-level idea of analysis], the union bound gives $\mathbb{P}(S) \geq 1 - \sum_{i=1}^J \eta_i \geq 99/100$. On S , the capped and ideal paths coincide, so restarting independent full attempts until success returns $\nu = \text{law}(\bar{X}_J \mid S)$. At the terminal index, $R_{2q}(\mu_{t_J} \parallel \pi) \leq 1/4$ by Corollary 3.2, and applying the weak triangle inequality gives $R_q(\bar{\nu}_J \parallel \pi) \leq 13/40$. Finally,

$$\frac{d\nu}{d\nu_J}(x) = \frac{\mathbb{P}(S|\bar{X}_J=x)}{\mathbb{P}(S)} \leq \frac{1}{\mathbb{P}(S)}, \text{ so}$$

$$R_q(\nu \parallel \pi) \leq R_q(\bar{\nu}_J \parallel \pi) + \frac{q}{q-1} \log \frac{1}{\mathbb{P}(S)} < 1.$$

It remains to bound the query complexity. Starting from a given σ_i^2 , doubling this value requires $O(\sqrt{qV}/\sigma_i^2)$ phases. Since the query complexity in each phase during doubling is $\tilde{O}(n^2\sigma_i^2 \log q \log^5 J)$, the total query cost of a doubling is $\tilde{O}(q^{1/2}n^2V^{1/2})$. Since there are only logarithmically many doublings, the total query complexity is

$$\tilde{O}(q^{1/2}n^2V^{1/2}) \times O(\log_2(n\sqrt{qV})) = \tilde{O}(q^{1/2}n^2V^{1/2}).$$

If C denotes the cost of one full attempt, then the restart wrapper has expected cost $\mathbb{E}C_{\text{restart}} = \frac{\mathbb{E}C}{\mathbb{P}(S)} \leq \frac{100}{99} \mathbb{E}C$. Thus, restarting increases the cost by only a universal factor. $\square$

4 Extension to logconcave sampling

Let $V : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$ be a convex function such that $d\pi(x) \propto e^{-V(x)} dx$ is a full-dimensional logconcave probability measure in $\mathbb{R}^n$. Assume that V is given by an evaluation oracle, which returns the value of V at the queried point. Denote $v_* := \inf V$, but neither v_* nor a minimizer of V is assumed known. The ground set is $L_g := \{x \in \mathbb{R}^n : V(x) - \inf V \leq 10n\}$, and we impose the normalization $B(0, 1) \subset L_g$ following [KV25b].

We first recall the preprocessing steps from [KV25b].

Reduction. Define the epigraph⁵

$$\mathcal{K} := \{(x, t) \in \mathbb{R}^n \times \mathbb{R} : V(x) \leq nt\}. \quad (4.1)$$

Let $a := V(0)/n$, which is known from one oracle query. Since $0 \leq V(0) - v_* \leq 10n$, every point of $\mathcal{K}$ has $t \geq a - 10$. Moreover, $V(x) \leq V(0) + 10n$ for $x \in B(0, 1)$, so $B_{n+1}((0, a + 11), 1) \subset \mathcal{K}$.

Instead of sampling directly from π , we use the exponential-lifting technique [KV25b] and define the lifted distribution by

$$\pi^{X,T}(dx, dt) \propto e^{-nt} \mathbf{1}_{\mathcal{K}}(x, t) dx dt,$$

whose X -marginal is the original target π , since integrating over t gives a density proportional to $e^{-V(x)}$.

Constant-mass truncation. To ensure that the lifted distribution has a finite log-Sobolev constant, let $\ell := \log(4e)$, $D := 1 \vee R\ell$, and $b := 13\ell + 5$. We then truncate the epigraph $\mathcal{K}$ as follows:

$$\bar{\mathcal{K}} := \mathcal{K} \cap (B(0, D) \times [a - 10, a + b]).$$

Since $D \geq 1$ and $b > 12$, the ball $B_{n+1}((0, a + 11), 1)$ remains inside $\bar{\mathcal{K}}$. We now verify that this truncation has constant mass. If $E \sim \text{Exp}(1)$ is independent of $X \sim \pi$, then the conditional law of the lift gives

$$T = \frac{V(X) + E}{n}.$$

By [FMW16, Theorem 2.3 and Corollary 4.5], $\mathbb{E}_\pi[V(X) - v_*] \leq n$ and $\text{Var}_\pi V \leq n$. Hence,

$$\mathbb{E}T \leq a + 2 \quad \text{and} \quad \text{Var} T = \frac{\text{Var}_\pi V + 1}{n^2} \leq 2.$$

⁵We retain the original t -coordinate and do not normalize $\inf V$.

By Markov's and Chebyshev's inequalities,

$$\pi^{X,T}(\bar{\mathcal{K}}^c) \leq \mathbb{P}(\|X\| > R\ell) + \mathbb{P}(T > a + b) \leq \frac{1}{\ell^2} + \frac{2}{(b-2)^2} < \frac{1}{2},$$

so $p := \pi^{X,T}(\bar{\mathcal{K}}) \geq 1/2$. Let $\bar{\pi}^{X,T} \propto \pi^{X,T} \mathbb{1}_{\bar{\mathcal{K}}}$ be the truncated distribution, and write $\bar{\pi}$ for its X -marginal. Clearly, $d\bar{\pi}/d\pi \leq p^{-1} \leq 2$. Note that the truncated lift $\bar{\pi}^{X,T}$ is logconcave.

The truncated X -marginal need not have the same mean as π . The following comparison shows that the relevant parameters change by only a constant factor.

Lemma 4.1. *Let π and $\bar{\pi}$ be probability measures on $\mathbb{R}^n$ such that $\bar{\pi}$ is logconcave, and $d\bar{\pi}/d\pi \leq p_0^{-1}$ for some $p_0 > 0$. Let $\bar{R}, \bar{L}, \bar{\Lambda}$ denote the analogues of R, L, Λ for $\bar{\pi}$. Then:*

- $\mathbb{E}_{\bar{\pi}}[XX^\top] \preceq p_0^{-1} \mathbb{E}_{\pi}[XX^\top]$ and $\text{cov}_{\bar{\pi}} X \preceq p_0^{-1} \Sigma$;
- $\bar{R}^2 \leq p_0^{-1} R^2$, $\bar{L} \leq p_0^{-1} L$, and $\bar{\Lambda} \leq p_0^{-1} \Lambda$;
- $\sup_{\sigma^2 > 0} \text{Var}_{\bar{\pi}_{\sigma^2}}(\|X\|^2) \lesssim p_0^{-2} (1 + \mathbf{Q}_n)^2 (R^2 L \wedge R^3 \sqrt{\Lambda})$.

Proof. For every $u \in \mathbb{R}^n$, we have $\mathbb{E}_{\bar{\pi}}[\langle u, X \rangle^2] \leq p_0^{-1} \mathbb{E}_{\pi}[\langle u, X \rangle^2]$. For $m = \mathbb{E}_{\pi} X$,

$$\text{Var}_{\bar{\pi}}(\langle u, X \rangle) \leq \mathbb{E}_{\bar{\pi}}[\langle u, X - m \rangle^2] \leq p_0^{-1} \mathbb{E}_{\pi}[\langle u, X - m \rangle^2] = p_0^{-1} \text{Var}_{\pi}(\langle u, X \rangle),$$

which proves the first item. The second item follows by taking traces and operator norms. Applying Theorem 1.1 to $\bar{\pi}$ and using $\bar{R}^2 \bar{L} \leq p_0^{-2} R^2 L$ and $\bar{R}^3 \sqrt{\bar{\Lambda}} \leq p_0^{-2} R^3 \sqrt{\Lambda}$ proves the third item. $\square$

4.1 Rényi closeness of lifted annealing distributions

As in [KV25a, KV25c], we use the following annealing distribution with parameters $\rho \geq 0$ and $\sigma^2 > 0$:

$$\mu_{\sigma^2, \rho}(dx, dt) \propto \exp\left(-\frac{\|x\|^2}{2\sigma^2} - \rho t\right) \mathbb{1}_{\bar{\mathcal{K}}}(x, t) dx dt, \quad (4.2)$$

and follow a two-phase annealing schedule, which simplifies the schedules in [KV25a, KV25c]. In the (σ^2, ρ) -plane, we first follow $(n^{-1}, 1) \rightarrow (n^{-1}, n)$ and then increase σ^2 with $\rho = n$ fixed. As in §3, we bound the Rényi divergence between neighboring targets, follow the annealing path with a logconcave sampler, and sum the per-phase query costs.

Rényi closeness. In the first phase $(n^{-1}, 1) \rightarrow (n^{-1}, n)$, we use the multiplicative update $\rho \leftarrow \rho(1 + O(\frac{1}{[q(q+n)]^{1/2}}))$. To justify this update, we bound the divergence incurred by changing the t -tilt:

Lemma 4.2. *Let $\nu_s := \mu_{n^{-1}, s}$ for $s > 0$. Then, for $q > 1$ and $0 \leq \alpha < (q-1)^{-1}$ such that $(1+\alpha)s \leq n$,*

$$\mathbf{R}_q(\nu_s \parallel \nu_{(1+\alpha)s}) \lesssim \frac{qn\alpha^2}{(1 - (q-1)\alpha)^2}.$$

Proof. Let $\Phi(s) := \log \int_{\bar{\mathcal{K}}} \exp(-n\|x\|^2/2 - st) dx dt$. As in the proof of Lemma 3.1, we obtain

$$\mathbf{R}_q(\nu_s \parallel \nu_{(1+\alpha)s}) = \frac{\Phi(s(1 - (q-1)\alpha)) - q\Phi(s) + (q-1)\Phi((1+\alpha)s)}{q-1}.$$

Set $s_- := s(1-(q-1)\alpha) > 0$, $s_+ := s(1+\alpha)$, $I := [s_-, s_+]$, and $V_I := \sup_{u \in I} \Phi''(u) = \sup_{u \in I} \text{Var}_{\nu_u} T$. By Taylor expansion,

$$\begin{aligned}\Phi(s_-) &\leq \Phi(s) - (q-1)\alpha s \Phi'(s) + \frac{(q-1)^2 \alpha^2 s^2}{2} V_I, \\ \Phi(s_+) &\leq \Phi(s) + \alpha s \Phi'(s) + \frac{\alpha^2 s^2}{2} V_I.\end{aligned}$$

Substituting these estimates into the R_q -identity yields

$$R_q(\nu_s \parallel \nu_{(1+\alpha)s}) \leq \frac{(q-1)^2 \alpha^2 s^2 V_I + (q-1) \alpha^2 s^2 V_I}{2(q-1)} = \frac{q \alpha^2 s^2}{2} V_I.$$

We now bound V_I . To this end, we first bound the second moment of ν_u as $\mathbb{E}_{\nu_u}[\|X\|^2] \lesssim 1$ uniformly over $u \in [0, n]$. Note that the X -marginal of ν_u is proportional to $e^{-n\|x\|^2/2} h_u(x)$, where $h_u(x) := \int e^{-ut} \mathbf{1}_{\bar{\mathcal{K}}}(x, t) dt$ is logconcave. By the reduction above, $\{0\} \times [a, a+1] \subset \bar{\mathcal{K}} \subset \mathbb{R}^n \times [a-10, a+b]$. Hence, for $u \in [0, n]$,

$$\begin{aligned}h_u(0) &\geq \int_a^{a+1} e^{-ut} dt \geq e^{-u(a+1)}, \\ h_u(x) &\leq \int_{a-10}^{a+b} e^{-ut} dt \leq (b+10) e^{-u(a-10)}.\end{aligned}$$

Let x_u be a mode of this marginal. Since $\nu_u^X(x_u)/\nu_u^X(0) \geq 1$,

$$\frac{n}{2} \|x_u\|^2 \leq \log \frac{h_u(x_u)}{h_u(0)} \leq \log(b+10) + 11u \lesssim n.$$

Also, since the X -marginal of ν_u is n -strongly logconcave, the standard mode-moment estimate gives $\mathbb{E}_{\nu_u}[\|X - x_u\|^2] \leq 1$ (see [Che25, Lemma 4.0.1]). Combining these bounds gives $\mathbb{E}_{\nu_u}[\|X\|^2] \lesssim 1$. Now, by (PI), for $Q(x) = n\|x\|^2/2$,

$$\text{Var}_{\nu_u} Q \leq \frac{1}{n} \mathbb{E}_{\nu_u}[\|\nabla Q\|^2] = n \mathbb{E}_{\nu_u}[\|X\|^2] \lesssim n.$$

By the varentropy inequality [FMW16], we have $\text{Var}_{\nu_u}(Q(X) + uT) \leq n + 1$. Using the L^2 -triangle inequality,

$$u\sqrt{\text{Var}_{\nu_u} T} \leq \sqrt{\text{Var}_{\nu_u}(Q(X) + uT)} + \sqrt{\text{Var}_{\nu_u} Q} \lesssim \sqrt{n}.$$

Hence, $\Phi''(u) = \text{Var}_{\nu_u} T \lesssim n/u^2 \leq n/s_-^2$ for every $u \in I$, which gives $V_I \lesssim n/[s_-^2(1-(q-1)\alpha)^2]$. Therefore,

$$R_q(\nu_s \parallel \nu_{(1+\alpha)s}) \lesssim \frac{qn\alpha^2}{(1-(q-1)\alpha)^2}.$$

This completes the proof. $\square$

In the second phase, we keep $\rho = n$ fixed and increase σ^2 . While $\sigma^2 < 1$, we use the universal Gaussian-annealing update $\sigma^2 \leftarrow \sigma^2(1 + O(\frac{1}{(qn)^{1/2}}))$. Once $\sigma^2 \geq 1$, we use the thin-shell update $\sigma^2 \leftarrow \sigma^2(1 + O(\frac{\sigma^2}{(qV)^{1/2}}))$ as in §3. Let $\mu_{\sigma^2} := \mu_{\sigma^2, n}$, and note that its X -marginal is $\bar{\pi}_{\sigma^2}$. The conditional distribution $T|X = x$ is independent of σ^2 , so

$$R_q(\mu_{\sigma^2} \parallel \mu_{\tilde{\sigma}^2}) = R_q(\bar{\pi}_{\sigma^2} \parallel \bar{\pi}_{\tilde{\sigma}^2}) \quad \text{for all } q > 1 \text{ and } \sigma^2, \tilde{\sigma}^2 > 0.$$

Hence, the closeness bound in Corollary 3.2 applies in $\sigma^2 \geq 1$ and at the terminal point, but the regime of $\sigma^2 \leq 1$ needs an additional argument.

Lemma 4.3. *Let $\mu_s := \mu_{s,n}$ for $s \in (0, 1)$. Then, for $q > 1$ and $\alpha > 0$,*

$$R_q(\mu_s \parallel \mu_{(1+\alpha)s}) \lesssim qn\alpha^2.$$

Proof. Let x_σ be a mode of $\bar{\pi}_{\sigma^2}$. The bounds on h_n in the preceding proof give $\|x_\sigma\|^2/(2\sigma^2) \leq \log(h_n(x_\sigma)/h_n(0)) \lesssim n$. Since $\bar{\pi}_{\sigma^2}$ is σ^{-2} -strongly logconcave, the same mode-moment estimate gives $\mathbb{E}_{\bar{\pi}_{\sigma^2}}[\|X - x_\sigma\|^2] \leq n\sigma^2$. Thus, $\mathbb{E}_{\bar{\pi}_{\sigma^2}}[\|X\|^2] \lesssim n\sigma^2$, so by (PI)

$$\text{Var}_{\bar{\pi}_{\sigma^2}}(\|X\|^2) \leq 4\sigma^2 \mathbb{E}_{\bar{\pi}_{\sigma^2}}[\|X\|^2] \lesssim n\sigma^4.$$

By Lemma 3.1 with $s = (2\sigma^2)^{-1}$ and $s' = s/(1 + \alpha)$,

$$R_q(\mu_{\sigma^2} \parallel \mu_{(1+\alpha)\sigma^2}) \lesssim qn \left(\frac{s - s'}{s'} \right)^2 = qn\alpha^2,$$

which justifies the update when $\sigma^2 \leq 1$. □

4.2 Gaussian cooling in the lifted space

Fix any target Rényi order $q \geq q_0$. We use PS_{ann} to follow the annealing scheme described above [KV25a, KV25c]. Starting from $v_0 \sim \mu_0$, with step size $h > 0$ and threshold τ , write $v_k = (x_k, t_k)$, $w_{k+1} = (y_{k+1}, s_{k+1})$, $r := \sigma^2/(h + \sigma^2)$, and $h_r := rh$. One iteration of PS_{ann} consists of the following two steps.

- [Forward] Draw $w_{k+1} \sim \mathbf{N}(v_k, hI_{n+1})$.
- [Backward] Repeatedly draw $v_{k+1} \sim \mathbf{N}(ry_{k+1}, h_r I_n) \otimes \mathbf{N}(s_{k+1} - \rho h, h)$ until $v_{k+1} \in \bar{\mathcal{K}}$. If the rejection loop exceeds τ trials, declare failure.

We use the following per-target guarantee from [KV25c, Proposition 5.2]. For the cited guarantee, translate t by $-(a + 11)$. The shifted body contains $B_{n+1}(0, 1)$ and lies in $\mathbb{R}^n \times [-21, b - 11]$, while the normalized targets and both Gaussian steps transform by the same translation. Thus, rejection counts and failure events are unchanged, and the guarantee has no dependence on a .

Proposition 4.4 (Lifted annealing sampler). *Let $\mu = \mu_{\sigma^2, \rho}$ be as in (4.2) with $\sigma^2 \geq 1/n$ and $1 \leq \rho \leq n$. Given access to an evaluation oracle for V , let $\varepsilon > 0$, $\eta \in (0, 1/2)$, $q \geq q_0 \geq 2$, and let μ_0 be an initial distribution with $M_{q_0} := \|\frac{d\mu_0}{d\mu}\|_{L^{q_0}(\mu)}$. Initialize PS_{ann} from μ_0 and iterate it $N = \tilde{O}(n^2(\sigma^2 \vee 1) \log \frac{q}{\varepsilon} \log \frac{1}{\eta})$ times with $h = (24^2 n^2 \log S)^{-1}$ and $\tau = 2S^2 \log^2 S$, where $S = 16NM_2/\eta$. Assume $M_{q_0} \leq 10$ and $q_0 \geq 1 + \log \tau$.*

- The latter condition is ensured by $q_0 \geq 2 \vee O(\log \frac{N}{\eta})$.
- With probability at least $1 - \eta$, PS_{ann} completes all N iterations without failure. Conditioned on this event, its output law ν satisfies $R_q(\nu \parallel \mu) \leq \varepsilon + 2 \log \frac{1}{1-\eta}$, using $\tilde{O}(n^2(\sigma^2 \vee 1) \log \frac{q}{\varepsilon} \log \frac{1}{\eta})$ evaluation queries in expectation. The corresponding uncapped chain (i.e., $\tau = \infty$) has output law $\bar{\nu}$ satisfying $R_q(\bar{\nu} \parallel \mu) \leq \varepsilon$.

Let J be the total number of phases. Note that with $\varepsilon_i = 1/20$ and $\eta_i = (100J)^{-1}$, each phase requires $\tilde{O}(n^2(\sigma^2 \vee 1))$ evaluation queries, and its desired output is $1/20$ -close to its target.

Algorithm 4.5 (Thin-shell Gaussian cooling). **Input:** evaluation oracle for V , a target order q , $V \lesssim_{\log n} R^2 L \wedge R^3 \sqrt{\Lambda}$, $\sigma_{\text{last}}^2 = \sqrt{qV}$, and the base order $q_0 = 2 \vee \tilde{\Theta}(\log(qn^3V))$, and a sufficiently small constant $c > 0$.

Every sampler call below uses the indicated input and output Rényi orders, accuracy $\varepsilon_{i+1} = 1/20$, failure probability $\eta_{i+1} = (100J)^{-1}$, and the parameters of Proposition 4.4, using $S_{i+1} := 160N_{i+1}/\eta_{i+1}$. After each successful phase, denote its output by Z_{i+1} and replace i by $i+1$. Restart the entire construction if a cap is hit. The steps are as follows:

1. Independently draw $G \sim \mathcal{N}(0, n^{-1}I_n)$ conditioned on $\|G\| \leq D$ and $U \sim \text{Exp}(1)$ conditioned on $0 \leq U \leq b+10$, set $T := a - 10 + U$, and repeat until $V(G) \leq nT$. Set $Z_0 := (G, T) \sim \mu_0 := \mu_{n^{-1}, 1}$, $(\sigma_{\text{start}}^2, \rho_0) := (n^{-1}, 1)$, $\sigma_0^2 := \sigma_{\text{start}}^2$, and $i = 0$.

2. [Phase I] While $\rho_i < n$ and $\sigma_i^2 = n^{-1}$, set $\sigma_{i+1}^2 \leftarrow n^{-1}$ and

$$\rho_{i+1} \leftarrow \min\left\{n, \left(1 + \frac{c}{\sqrt{q_0(q_0 + n)}}\right) \rho_i\right\}, \quad (4.3)$$

and sample from $\mu_{i+1} := \mu_{n^{-1}, \rho_{i+1}}$ using $\text{PS}_{\text{ann}} : R_{q_0} \rightarrow R_{2q_0}$.

3. [Phase II] First, while $\sigma_i^2 < 1$ and $\rho_i = n$, set $\rho_{i+1} \leftarrow n$ and

$$\sigma_{i+1}^2 \leftarrow \left(1 + \frac{c}{\sqrt{q_0 n}}\right) \sigma_i^2,$$

and sample from $\mu_{i+1} := \mu_{\sigma_{i+1}^2, n}$ using $\text{PS}_{\text{ann}} : R_{q_0} \rightarrow R_{2q_0}$. After this loop, set $\sigma_{i+1}^2 := \sigma_i^2$, $\rho_{i+1} := \rho_i$, and $\mu_{i+1} := \mu_i$, and make one call at this target with $\text{PS}_{\text{ann}} : R_{q_0} \rightarrow R_{2q}$. Then, while $\sigma_i^2 < \sigma_{\text{last}}^2$, set $\rho_{i+1} \leftarrow n$ and

$$\sigma_{i+1}^2 \leftarrow \left(1 + \frac{\sigma_i^2}{\sqrt{qV}}\right) \sigma_i^2,$$

and sample from $\mu_{i+1} := \mu_{\sigma_{i+1}^2, n}$ using $\text{PS}_{\text{ann}} : R_q \rightarrow R_{2q}$. After the second loop, return the X -coordinate of Z_i .

For Phase I updates, Lemma 4.2 ensures $R_{2q_0}(\mu_{n^{-1}, \rho_i} \parallel \mu_{n^{-1}, \rho_{i+1}}) \leq \frac{1}{18}$. In the first Phase II loop, Lemma 4.3 ensures 1/4-closeness in R_{2q_0} -divergence. In the second loop, as in §3, Corollary 3.2 ensures 1/4-closeness in R_{2q} .

Proof of Theorem 1.3. We may assume $q \geq q_0$; otherwise, we can proceed with $q = q_0$, and use monotonicity of R_q . Let $\mu_0, \dots, \mu_J$ denote the annealing distributions in Algorithm 4.5, and set $\bar{\nu}_0 := \mu_0$. For $i \geq 1$, let $Z_i \sim \bar{\nu}_i$ be the output of the ideal PS_{ann} at phase i .

Let $p = q_0$ before the order-boost and $p = q$ after it. By the weak triangle inequality,

$$R_p(\bar{\nu}_{i-1} \parallel \mu_i) \leq \frac{p-1/2}{p-1} R_{2p}(\bar{\nu}_{i-1} \parallel \mu_{i-1}) + R_{2p-1}(\mu_{i-1} \parallel \mu_i) \leq \frac{3}{40} + \frac{1}{4} = \frac{13}{40}.$$

Hence, Proposition 4.4 with $(p, r) = (q_0, 2q_0)$ before the boost and $(p, r) = (q, 2q)$ afterward preserves M_p -warmness.

As in the proof of Theorem 1.2, we couple the capped and ideal paths until a cap is hit; the success event S satisfies $\mathbb{P}(S) \geq 99/100$. Write $\nu^{X,T} = \text{law}(Z_J \mid S)$. The terminal bound and the

weak triangle inequality yield $R_q(\bar{\nu}_J \parallel \bar{\pi}^{X,T}) \leq 13/40$. Since $d\nu^{X,T}/d\bar{\nu}_J \leq 1/\mathbb{P}(S)$,

$$R_q(\nu^{X,T} \parallel \bar{\pi}^{X,T}) \leq \frac{13}{40} + \frac{q}{q-1} \log \frac{100}{99} < 1.$$

Since $\bar{\pi}^{X,T}(dx, dt) = p^{-1} \mathbb{1}_{\bar{\mathcal{K}}}(x, t) \pi^{X,T}(dx, dt)$ for $p = \pi^{X,T}(\bar{\mathcal{K}}) \geq 1/2$, data processing under $(x, t) \mapsto x$ gives

$$R_q(\nu^X \parallel \pi) \stackrel{\text{DPI}}{\leq} R_q(\nu^{X,T} \parallel \pi^{X,T}) = R_q(\nu^{X,T} \parallel \bar{\pi}^{X,T}) + \log \frac{1}{p} \leq 1 + \log 2 < 2.$$

It remains to bound the query complexity. Rejection sampling gives an exact initial sample and uses $O(1)$ expected evaluation queries. In Phase I, the update in (4.3) requires $O(\sqrt{q_0(q_0 + n)} \log n)$ phases. Each phase requires $\tilde{O}(n^2)$ queries, so

$$\text{Phase I cost} \leq \tilde{O}(n^2) \times O(\sqrt{q_0(q_0 + n)} \log n) = \tilde{O}(n^{5/2}).$$

For Phase II, the universal updates below $\sigma^2 = 1$ use $O(\sqrt{q_0 n} \log n)$ phases, and each phase needs $\tilde{O}(n^2)$ queries. After $\sigma^2 \geq 1$, since the doubling of an initial σ^2 requires $O(\sqrt{qV}/\sigma^2)$ phases, and each phase needs $\tilde{O}(n^2 \sigma^2)$ queries, the doubling requires $\tilde{O}(n^2 \sqrt{qV})$ queries. Summing the query costs of the two loops,

$$\text{Phase II cost} \leq \tilde{O}(n^{2.5} + n^2 \sqrt{qV}) = \tilde{O}(n^{5/2} + n^2 \sqrt{q} (R\sqrt{L} \wedge R^{3/2} \Lambda^{1/4})).$$

Finally, since $\mathbb{P}(S) \geq 99/100$, the restart wrapper multiplies the expected cost by at most $100/99$, which completes the proof. $\square$

Acknowledgments. This work was supported in part by NSF Awards CCF-2504995, CCF-2236669, CCF-2504994, and a Simons Investigator award.

References

- [ABP03] Milla Anttila, Keith Ball, and Irini Perissinaki. The central limit problem for convex bodies. *Transactions of the American Mathematical Society*, 355(12):4723–4735, 2003.
- [BGL14] Dominique Bakry, Ivan Gentil, and Michel Ledoux. *Analysis and geometry of Markov diffusion operators*, volume 348. Springer, Cham, 2014.
- [Biz26] Pierre Bizeul. On the log-Sobolev constant of log-concave vectors. *Journal of Functional Analysis*, 290(9):111368, May 2026.
- [BK03] Sergey G. Bobkov and Alexander Koldobsky. On the central limit property of convex bodies. In *Geometric Aspects of Functional Analysis*, volume 1807 of *Lecture Notes in Math.*, pages 44–52. Springer, Berlin, 2003.
- [Che25] Sinho Chewi. *Log-concave sampling*. Book draft available at <https://chewisinho.github.io>, 2025.
- [CK26] Yuansi Chen and Bo’az Klartag. Digesting the proof of the sharp thin-shell inequality. *arXiv preprint arXiv:2607.23307*, 2026.
- [CV15] Ben Cousins and Santosh S. Vempala. Bypassing KLS: Gaussian cooling and an $O^*(n^3)$ volume algorithm. In *Symposium on Theory of Computing*, pages 539–548. ACM, 2015.
- [CV18] Ben Cousins and Santosh S. Vempala. Gaussian cooling and $O^*(n^3)$ algorithms for volume and Gaussian volume. *SIAM Journal on Computing*, 47(3):1237–1273, 2018.

- [CW01] Anthony Carbery and James Wright. Distributional and L^q norm inequalities for polynomials over convex bodies in $\mathbb{R}^n$. *Mathematical Research Letters*, 8(3):233–248, 2001.
- [DFK91] Martin Dyer, Alan Frieze, and Ravi Kannan. A random polynomial-time algorithm for approximating the volume of convex bodies. *Journal of the ACM*, 38(1):1–17, 1991.
- [FMW16] Matthieu Fradelizi, Mokshay Madiman, and Liyao Wang. Optimal concentration of information content for log-concave densities. In *High Dimensional Probability VII*, volume 71 of *Progress in Probability*, pages 45–60. Birkhäuser, Cham, 2016.
- [JLLV26] He Jia, Aditi Laddha, Yin Tat Lee, and Santosh Vempala. Reducing isotropy and volume to kls: Faster rounding and volume algorithms. *J. ACM*, 73(2), April 2026.
- [Kat95] Tosio Kato. *Perturbation Theory for Linear Operators*. Classics in Mathematics. Springer, Berlin, Heidelberg, 2 edition, 1995.
- [KL25] Bo’az Klartag and Joseph Lehec. Thin-shell bounds via parallel coupling. *arXiv preprint arXiv:2507.15495*, 2025.
- [Kla23] Bo’az Klartag. Logarithmic bounds for isoperimetry and slices of convex sets. *Ars Inveniendi Analytica*, 2023. Paper No. 4, 17 pp.
- [KLS95] Ravi Kannan, László Lovász, and Miklós Simonovits. Isoperimetric problems for convex bodies and a localization lemma. *Discrete & Computational Geometry*, 13(3-4):541–559, 1995.
- [KLS97] Ravi Kannan, László Lovász, and Miklós Simonovits. Random walks and an $O^*(n^5)$ volume algorithm for convex bodies. *Random Structures & Algorithms*, 11(1):1–50, 1997.
- [KV25a] Yunbum Kook and Santosh S. Vempala. Faster logconcave sampling from a cold start in high dimension. In *Symposium on Foundations of Computer Science*, pages 997–1006. IEEE, 2025.
- [KV25b] Yunbum Kook and Santosh S. Vempala. Sampling and integration of logconcave functions by algorithmic diffusion. In *Symposium on Theory of Computing*, pages 924–932. ACM, 2025.
- [KV25c] Yunbum Kook and Santosh S. Vempala. Zeroth-order logconcave sampling. *arXiv preprint arXiv:2507.18021*, 2025.
- [KV26a] Yunbum Kook and Santosh S. Vempala. Spectral gaps of Hit-and-Run and Coordinate Hit-and-Run. *arXiv preprint arXiv:2608.16878*, 2026.
- [KV26b] Yunbum Kook and Santosh S. Vempala. A unified complexity bound for logconcave sampling. *arXiv preprint arXiv:2606.12694*, 2026.
- [KVZ24] Yunbum Kook, Santosh S. Vempala, and Matthew S. Zhang. In-and-Out: algorithmic diffusion for sampling convex bodies. In *Advances in Neural Information Processing Systems*, volume 37, pages 108354–108388. Curran Associates, Inc., 2024.
- [KVZ26] Yunbum Kook, Santosh S. Vempala, and Matthew S. Zhang. In-and-Out: algorithmic diffusion for sampling convex bodies. *Random Structures & Algorithms*, 68(3):e70061, 2026.
- [KZ25] Yunbum Kook and Matthew S. Zhang. Rényi-infinity constrained sampling with d^3 membership queries. In *Symposium on Discrete Algorithms*, pages 5278–5306. SIAM, 2025.
- [LS93] László Lovász and Miklós Simonovits. Random walks in a convex body and an improved volume algorithm. *Random Structures & Algorithms*, 4(4):359–412, 1993.
- [LST21] Yin Tat Lee, Ruochi Shen, and Kevin Tian. Structured logconcave sampling with a restricted Gaussian oracle. In *Conference on Learning Theory*, volume 134, pages 2993–3050. PMLR, 2021.
- [LV06a] László Lovász and Santosh S. Vempala. Fast algorithms for logconcave functions: sampling, rounding, integration and optimization. In *Symposium on Foundations of Computer Science*, pages 57–68. IEEE, 2006.
- [LV06b] László Lovász and Santosh S. Vempala. Hit-and-run from a corner. *SIAM Journal on Computing*, 35(4):985–1005, 2006.
- [LV06c] László Lovász and Santosh S. Vempala. Simulated annealing in convex bodies and an $O^*(n^4)$ volume algorithm. *Journal of Computer and System Sciences*, 72(2):392–417, 2006.

- [LV18] Yin Tat Lee and Santosh S. Vempala. Stochastic localization + Stieltjes barrier = tight bound for log-Sobolev. In *Symposium on Theory of Computing*, STOC 2018, pages 1122–1129, New York, NY, USA, 2018. ACM.
- [LV24] Yin Tat Lee and Santosh S. Vempala. Eldan’s stochastic localization and the KLS conjecture: isoperimetry, concentration and mixing. *Annals of Mathematics*, 199(3):1043–1092, 2024.
- [Mir17] Ilya Mironov. Rényi differential privacy. In *Computer Security Foundations Symposium*, pages 263–275. IEEE, 2017.
- [vEH14] Tim van Erven and Peter Harremoës. Rényi divergence and Kullback-Leibler divergence. *IEEE Transactions on Information Theory*, 60(7):3797–3820, 2014.