

Variational Outlier-Robust Gaussian Process Regression with Generative Modeling

Arslan Majal and Aamir Hussain Chughtai

Abstract—Outliers can substantially distort Gaussian process regression (GPR) due to its conventional Gaussian observation likelihood, leading to inaccurate model learning and prediction. To address this limitation, this article introduces a generative GPR model that captures observation-specific contamination and adaptively mitigates the influence of outliers. Subsequently, a variational generalized expectation-maximization procedure is used to learn the latent variables and GPR model parameters. Experiments on synthetic and real datasets under different contamination settings demonstrate that the proposed method remains competitive with—and in several cases outperforms—robust GPR baselines in prediction accuracy. Moreover, the proposed method shares the cubic computational scaling of the compared GPR methods.

Index Terms—Gaussian process regression, outlier-robust regression, variational Bayesian inference, robust signal processing, uncertainty quantification, expectation-maximization

I. INTRODUCTION

GAUSSIAN process regression (GPR) [1] is a Bayesian nonparametric regression method that provides flexible nonlinear modeling and principled uncertainty quantification, while remaining effective in data-limited settings. These properties make GPR useful in numerous signal-processing and learning applications.

Standard GPR assumes Gaussian observation noise, making it sensitive to outliers caused by sensor faults, calibration errors, transmission noise, or unexpected disturbances. Under this model, large residuals are penalized quadratically, allowing even a few corrupted observations to substantially distort the estimated latent function [2]. This motivates robust GPR methods that reduce the influence of unreliable observations.

Outliers in GPR are commonly handled by modifying the observation likelihood or the corresponding loss function. Existing robust approaches often rely on fixed likelihood modifications or robust losses with user-defined tuning parameters, including Huber- and Hampel-type penalties, heavy-tailed Student- t likelihoods, and generalized-Bayes likelihood-induced losses [3]–[5]. Although tractable, these approaches can be sensitive to prescribed thresholds, loss parameters, or distributional shape parameters that control the influence of atypical observations. To overcome these limitations, more expressive adaptive models have been proposed. Instead of fixing the robust cost or likelihood parameters a priori, these methods adapt them from the observed data, thereby providing data-driven robustness.

The main challenge, however, is to make the outlier model sufficiently expressive while retaining tractable inference. To this end, existing approaches introduce latent variables, hierarchical likelihoods, mixture-based noise models, or optimization-based robust formulations, including graduated non-convexity (GNC), to account for corrupted measurements [6]–[13].

A relevant class of adaptive methods employs generative models with variational inference to identify and mitigate outliers [7], [11]. Building on this idea, adaptive selective observation rejecting (ASOR) [10] learns outlier statistics rather than completely discarding suspected observations, thereby adapting their influence according to the learned contamination characteristics. ASOR has demonstrated favorable estimation accuracy and computational efficiency relative to state-of-the-art GNC methods in robotic perception problems [13]. Its conjugate hierarchical structure enables closed-form variational updates for the model parameters [10], with related ideas extended to robust filtering in [12].

Motivated by the adaptive outlier treatment in ASOR, we propose a formulation tailored to outlier-robust GPR. The key development is to integrate observation-specific precision modeling with the latent GP function and its associated noise, mean, and kernel parameters within a unified model. Since exact Bayesian inference is intractable, we employ a variational generalized expectation-maximization (EM) procedure to learn the latent variables and model parameters. Closed-form updates are obtained for all quantities except the kernel hyperparameters, which are optimized using the gradient-based method [9].

To evaluate performance, the proposed method is extensively compared with representative GPR baselines on synthetic and real-data regression problems under diverse contamination settings.

II. PROPOSED METHODOLOGY

Notation: Nonbold symbols denote scalars or scalar-valued functions, bold lowercase symbols denote vectors, and bold uppercase symbols denote matrices. Moreover, $(\cdot)^\top$ denotes transpose, $[\mathbf{A}]_{ij}$ denotes the (i, j) th entry of $\mathbf{A}$, and $\text{diag}(\cdot)$ and $\text{tr}(\cdot)$ denote the diagonal and trace operators, respectively.

Consider n input–output samples arranged as

$$\mathbf{X} = \begin{bmatrix} x_{11} & \cdots & x_{1d_x} \\ \vdots & & \vdots \\ x_{i1} & \cdots & x_{id_x} \\ \vdots & & \vdots \\ x_{n1} & \cdots & x_{nd_x} \end{bmatrix}, \quad \mathbf{Y} = \begin{bmatrix} y_{11} & \cdots & y_{1j} & \cdots & y_{1d_y} \\ \vdots & & \vdots & & \vdots \\ y_{i1} & \cdots & y_{ij} & \cdots & y_{id_y} \\ \vdots & & \vdots & & \vdots \\ y_{n1} & \cdots & y_{nj} & \cdots & y_{nd_y} \end{bmatrix} \quad (\mathbf{y}^i)^\top$$

$\mathbf{y}_j$

A. Majal is with Zunitech, LLC, Houston, USA.

A. H. Chughtai is with the Institute of Data Science, University of Engineering and Technology, Lahore, Pakistan.

where $\mathbf{X} \in \mathbb{R}^{n \times d_x}$ contains the input samples and $\mathbf{Y} \in \mathbb{R}^{n \times d_y}$ contains the corresponding outputs. The i th rows are $(\mathbf{x}^i)^\top$ and $(\mathbf{y}^i)^\top$, respectively, while $\mathbf{y}_j$ denotes the j th output column.

Assuming independent output components and contamination processes, the problem is decomposed into d_y scalar-output GPR models, where the j th model is trained using $\mathbf{X}$ and $\mathbf{y}_j$.

A. Scalar Observation Model

For output dimension j , let $f_{ij} \triangleq f_j(\mathbf{x}^i)$ denote the latent function value at the i th input. The observation model is

$$y_{ij} = f_{ij} + \varepsilon_{ij}, \quad i = 1, \dots, n,$$

where ε_{ij} denotes the observation error and σ_j^2 denotes its nominal variance. Following the ASOR construction, a positive latent precision multiplier $\mathcal{I}_{ij} > 0$ is introduced such that

$$p(\varepsilon_{ij} | \mathcal{I}_{ij}, \sigma_j^2) = \mathcal{N}(\varepsilon_{ij} | 0, \sigma_j^2 / \mathcal{I}_{ij}),$$

where $\mathcal{N}(x | \mu, v)$ denotes a Gaussian density in x with mean μ and variance v . Accordingly, the observation likelihood is

$$p(y_{ij} | f_{ij}, \mathcal{I}_{ij}, \sigma_j^2) = (2\pi\sigma_j^2)^{-1/2} \mathcal{I}_{ij}^{1/2} \exp\left[-\frac{\mathcal{I}_{ij}}{2\sigma_j^2} (y_{ij} - f_{ij})^2\right]. \quad (1)$$

The value $\mathcal{I}_{ij} = 1$ recovers the nominal observation model, whereas $\mathcal{I}_{ij} \neq 1$ adaptively modifies the observation variance to account for outliers.

Let $\mathbf{f}_j \triangleq [f_{1j}, \dots, f_{nj}]^\top$. The GP prior is

$$p(\mathbf{f}_j | \mathbf{X}, m_j, \boldsymbol{\kappa}_j) = \mathcal{N}(\mathbf{f}_j | m_j \mathbf{1}_n, \mathbf{K}_j), \quad (2)$$

$$[\mathbf{K}_j]_{i\ell} = k_j(\mathbf{x}^i, \mathbf{x}^\ell; \boldsymbol{\kappa}_j).$$

where $\mathbf{1}_n \in \mathbb{R}^n$ denotes the vector of ones and, for the multivariate case, $\mathcal{N}(\mathbf{x} | \boldsymbol{\mu}, \boldsymbol{\Sigma})$ denotes a Gaussian density with mean $\boldsymbol{\mu}$ and covariance matrix $\boldsymbol{\Sigma}$. The covariance matrix is constructed using the automatic relevance determination (ARD) squared-exponential kernel

$$k_j(\mathbf{x}^i, \mathbf{x}^\ell; \boldsymbol{\kappa}_j) = \sigma_{f,j}^2 e^{-0.5 \sum_{r=1}^{d_x} (x_{ir} - x_{\ell r})^2 / \ell_{j,r}^2}, \quad (3)$$

$$\boldsymbol{\kappa}_j \triangleq [\sigma_{f,j}^2, \ell_{j,1}, \dots, \ell_{j,d_x}]^\top.$$

Here, $\sigma_{f,j}^2$ is the signal variance and $\ell_{j,r}$ is the length scale associated with the r th input dimension.

B. Hierarchical Priors

$\mathcal{I}_{ij}$ is assigned the unit-spike-and-Gamma prior

$$p(\mathcal{I}_{ij} | b_j) = (1 - \theta_{ij}) \mathcal{G}(\mathcal{I}_{ij} | a, b_j) + \theta_{ij} \delta(\mathcal{I}_{ij} - 1), \quad (4)$$

where $\delta(\cdot)$ denotes the Dirac delta function and $\mathcal{G}(\mathcal{I} | a, b)$ denotes the Gamma density with shape $a > 0$ and rate $b > 0$:

$$\mathcal{G}(\mathcal{I} | a, b) \triangleq \frac{b^a}{\Gamma(a)} \mathcal{I}^{a-1} e^{-b\mathcal{I}}. \quad (5)$$

where $\Gamma(\cdot)$ denotes the Gamma function. The parameter $\theta_{ij} \in (0, 1)$ is the prior probability that y_{ij} belongs to the nominal branch.

To retain Bayesian conjugacy, the output-specific Gamma rate b_j is assigned the Gamma hyperprior

$$p(b_j) = \mathcal{G}(b_j | A, B), \quad (6)$$

where $A > 1$ and $B > 0$ are fixed hyperparameters.

Similarly, the nominal observation-noise variance is assigned the inverse-Gamma prior

$$p(\sigma_j^2) = \mathcal{IG}\left(\sigma_j^2 \left| \frac{\nu_0}{2}, \frac{s_{0j}}{2} \right.\right) \propto (\sigma_j^2)^{-(\nu_0+2)/2} e^{-s_{0j}/2\sigma_j^2}. \quad (7)$$

where $\nu_0 > 0$ and $s_{0j} > 0$ are fixed hyperparameters.

The fixed prior hyperparameters associated with output dimension j are collected as

$$\mathcal{P}_j \triangleq \{a, \{\theta_{ij}\}_{i=1}^n, A, B, \nu_0, s_{0j}\}. \quad (8)$$

C. Variational Generalized-EM Inference

For output dimension j , define

$$\mathcal{I}_j \triangleq [\mathcal{I}_{1j}, \dots, \mathcal{I}_{nj}]^\top, \quad \mathbf{Z}_j \triangleq \{\mathbf{f}_j, \mathcal{I}_j\}, \quad \boldsymbol{\Theta}_j \triangleq \{m_j, \boldsymbol{\kappa}_j, \sigma_j^2, b_j\}. \quad (9)$$

For $j = 1, \dots, d_y$, the joint posterior satisfies

$$p(\mathbf{Z}_j, \boldsymbol{\Theta}_j | \mathbf{y}_j, \mathbf{X}, \mathcal{P}_j) \propto p(\mathbf{f}_j | \mathbf{X}, m_j, \boldsymbol{\kappa}_j) p(\sigma_j^2) p(b_j) \\ \times \prod_{i=1}^n p(\mathcal{I}_{ij} | b_j) p(y_{ij} | f_{ij}, \mathcal{I}_{ij}, \sigma_j^2). \quad (10)$$

We use flat non-informative priors for m_j and $\boldsymbol{\kappa}_j$, whose constant terms are omitted from (10). Since the exact joint posterior does not admit closed form, we employ variational generalized EM [14].

The E-step uses the mean-field factorization

$$q_j(\mathbf{Z}_j) = q_j(\mathbf{f}_j) \prod_{i=1}^n q_j(\mathcal{I}_{ij}). \quad (11)$$

For $\xi \in \{\mathbf{f}_j, \mathcal{I}_{1j}, \dots, \mathcal{I}_{nj}\}$, the coordinate-optimal update is

$$\log q_j^*(\xi) = \mathbb{E}_{q_j(\mathbf{Z}_j \setminus \xi)} [\log p(\mathbf{y}_j, \mathbf{Z}_j | \mathbf{X}, \boldsymbol{\Theta}_j, \mathcal{P}_j)] + c, \quad (12)$$

where c is independent of ξ . Applying (12) to $\mathbf{f}_j$ and each $\mathcal{I}_{ij}$ gives the closed-form updates below. At each generalized-EM iteration, these factors are updated sequentially, and their product gives $q_j(\mathbf{Z}_j)$.

The M-step aims to solve

$$\boldsymbol{\Theta}_j^* \in \arg \max_{\boldsymbol{\Theta}_j} \mathcal{Q}_j(\boldsymbol{\Theta}_j). \quad (13)$$

where $\mathcal{Q}_j(\boldsymbol{\Theta}_j) \triangleq \mathbb{E}_{q_j(\mathbf{Z}_j)} [\log p(\mathbf{y}_j, \mathbf{Z}_j, \boldsymbol{\Theta}_j | \mathbf{X}, \mathcal{P}_j)]$. Closed-form updates are obtained for m_j , σ_j^2 , and b_j , whereas $\boldsymbol{\kappa}_j$ is updated using gradient descent with backtracking. The priors on σ_j^2 and b_j yield maximum a posteriori (MAP) updates, whereas the uninformative priors on m_j and $\boldsymbol{\kappa}_j$ yield maximum-likelihood updates.

D. Variational E-Step: Latent-Function Update

For fixed output j , define

$$w_{ij} \triangleq \mathbb{E}_{q_j(\mathcal{I}_{ij})} [\mathcal{I}_{ij}], \quad \boldsymbol{\Lambda}_j \triangleq \text{diag}(w_{1j}, \dots, w_{nj}) / \sigma_j^2. \quad (14)$$

Applying (12) to $\mathbf{f}_j$ gives

$$q_j(\mathbf{f}_j) = \mathcal{N}(\mathbf{f}_j | \boldsymbol{\mu}_{f,j}, \mathbf{C}_{f,j}), \\ \mathbf{C}_{f,j} = (\mathbf{K}_j^{-1} + \boldsymbol{\Lambda}_j)^{-1}, \quad (15)$$

$$\boldsymbol{\mu}_{f,j} = \mathbf{C}_{f,j} (\mathbf{K}_j^{-1} m_j \mathbf{1}_n + \boldsymbol{\Lambda}_j \mathbf{y}_j).$$

E. Variational E-Step: Precision-Multiplier Update

Using the moments of $q_j(\mathbf{f}_j)$, define

$$S_{ij} \triangleq (y_{ij} - \mu_{f,ij})^2 + [\mathbf{C}_{f,j}]_{ii}, \quad R_{ij} \triangleq S_{ij}/\sigma_j^2, \quad (16)$$

where $\mu_{f,ij}$ is the i th entry of $\boldsymbol{\mu}_{f,j}$. Let

$$\tilde{a} \triangleq a + 1/2, \quad \tilde{b}_{ij} \triangleq b_j + R_{ij}/2. \quad (17)$$

Applying (12) to $\mathcal{I}_{ij}$ gives

$$q_j(\mathcal{I}_{ij}) = (1 - \Omega_{ij})\mathcal{G}(\mathcal{I}_{ij} | \tilde{a}, \tilde{b}_{ij}) + \Omega_{ij}\delta(\mathcal{I}_{ij} - 1), \quad (18)$$

where

$$\Omega_{ij} = \left[1 + \frac{1 - \theta_{ij}}{\theta_{ij}} \frac{\Gamma(\tilde{a})}{\Gamma(a)} \frac{b_j^a}{\tilde{b}_{ij}^a} \exp\left(\frac{R_{ij}}{2}\right) \right]^{-1}. \quad (19)$$

The corresponding robustness weight is

$$w_{ij} = \Omega_{ij} + (1 - \Omega_{ij})\tilde{a}/\tilde{b}_{ij}. \quad (20)$$

F. Generalized M-Step: Parameter Updates

With the variational quantities fixed, $\mathcal{Q}_j(\boldsymbol{\Theta}_j)$ separates into terms involving σ_j^2 , b_j , and $(m_j, \boldsymbol{\kappa}_j)$.

Collecting the terms that depend on σ_j^2 and setting their derivative to zero gives the MAP update

$$\sigma_j^2 \leftarrow \frac{S_j + s_{0j}}{n + \nu_0 + 2}. \quad (21)$$

where $S_j \triangleq \sum_{i=1}^n w_{ij} S_{ij}$. Similarly, the MAP update of the Gamma rate parameter is

$$b_j \leftarrow \frac{A - 1 + a \sum_{i=1}^n (1 - \Omega_{ij})}{B + \sum_{i=1}^n (1 - \Omega_{ij})\tilde{a}/\tilde{b}_{ij}}. \quad (22)$$

For fixed $\boldsymbol{\kappa}_j$, maximizing the GP-dependent terms of $\mathcal{Q}_j(\boldsymbol{\Theta}_j)$ with respect to m_j gives

$$m_j \leftarrow (\mathbf{1}_n^\top \mathbf{K}_j^{-1} \boldsymbol{\mu}_{f,j}) / (\mathbf{1}_n^\top \mathbf{K}_j^{-1} \mathbf{1}_n). \quad (23)$$

For the kernel update, define

$$\begin{aligned} \phi_j &\triangleq [\log \sigma_{f,j} \quad \log \ell_{j1} \quad \cdots \quad \log \ell_{jd_x}]^\top, \\ \boldsymbol{\kappa}_j(\phi_j) &\triangleq [\exp(2\phi_{j1}) \quad \exp(\phi_{j2}) \quad \cdots \quad \exp(\phi_{j,d_x+1})]^\top. \end{aligned} \quad (24)$$

This logarithmic parameterization enforces positivity of the signal variance and length scales while allowing optimization over ϕ_j .

For fixed m_j , maximizing the terms of $\mathcal{Q}_j(\boldsymbol{\Theta}_j)$ that depend on $\boldsymbol{\kappa}_j$ through $\mathbf{K}_j$ is equivalent to minimizing

$$\mathcal{J}_j(\phi_j) \triangleq \frac{1}{2} \text{tr}(\mathbf{K}_j^{-1} \mathbf{D}_j) + \frac{1}{2} \log |\mathbf{K}_j|. \quad (25)$$

where

$$\mathbf{D}_j \triangleq \mathbf{C}_{f,j} + (\boldsymbol{\mu}_{f,j} - m_j \mathbf{1}_n)(\boldsymbol{\mu}_{f,j} - m_j \mathbf{1}_n)^\top. \quad (26)$$

To obtain a descent direction for minimizing $\mathcal{J}_j(\phi_j)$, its gradient components are

$$\frac{\partial \mathcal{J}_j}{\partial \phi_{jr}} = \frac{1}{2} \text{tr} \left[(\mathbf{K}_j^{-1} - \mathbf{K}_j^{-1} \mathbf{D}_j \mathbf{K}_j^{-1}) \frac{\partial \mathbf{K}_j}{\partial \phi_{jr}} \right]. \quad (27)$$

Algorithm 1: ASOR-GPR

Input: Training data $(\mathbf{X}, \mathbf{Y})$; prior hyperparameters $\{\mathcal{P}_j\}_{j=1}^{d_y}$; initial parameters $\{\boldsymbol{\Theta}_j\}_{j=1}^{d_y}$.

Output: Converged independent-output ASOR-GPR models.

for $j \leftarrow 1$ **to** d_y **do**

Extract $\mathbf{y}_j$, initialize $\boldsymbol{\Theta}_j$, set ϕ_j from $\boldsymbol{\kappa}_j$ using (24), set $w_{ij} \leftarrow 1$, $i = 1, \dots, n$, and construct $\mathbf{K}_j$

repeat

Variational E-step:

Update $q_j(\mathbf{f}_j)$ using (14) and (15)

Update $\{q_j(\mathcal{I}_{ij}), \Omega_{ij}, w_{ij}\}_{i=1}^n$ using (16)–(20)

Generalized M-step:

Update σ_j^2 , b_j , and m_j using (21)–(23)

Update ϕ_j by gradient descent with backtracking using (25)–(30)

Update $\boldsymbol{\kappa}_j$ using (24) and reconstruct $\mathbf{K}_j$ using (3)

until convergence

Perform a final variational E-step and store the converged quantities

Return The converged ASOR-GPR models

The required kernel derivatives are

$$\frac{\partial \mathbf{K}_j}{\partial \phi_{j1}} = 2\mathbf{K}_j, \quad \left[\frac{\partial \mathbf{K}_j}{\partial \phi_{j,r+1}} \right]_{il} = [\mathbf{K}_j]_{il} \frac{(x_{ir} - x_{lr})^2}{\ell_{jr}^2}. \quad (28)$$

For a trial step size $\alpha_j > 0$, the normalized-gradient candidate is

$$\phi_j^{\text{cand}} = \phi_j^{\text{old}} - \alpha_j \frac{\mathbf{g}_j}{\max\{1, \|\mathbf{g}_j\|_2\}}, \quad (29)$$

where $\mathbf{g}_j \triangleq \nabla_{\phi_j} \mathcal{J}_j$. A backtracking line search halves α_j until the Armijo condition

$$\mathcal{J}_j(\phi_j^{\text{cand}}) \leq \mathcal{J}_j(\phi_j^{\text{old}}) - c_1 \alpha_j \frac{\|\mathbf{g}_j\|_2^2}{\max\{1, \|\mathbf{g}_j\|_2\}} \quad (30)$$

is satisfied, where c_1 is the Armijo constant. The accepted value is $\phi_j \leftarrow \phi_j^{\text{cand}}$, followed by $\boldsymbol{\kappa}_j \leftarrow \boldsymbol{\kappa}_j(\phi_j)$ and $\mathbf{K}_j \leftarrow \mathbf{K}_j(\boldsymbol{\kappa}_j)$.

Iterations terminate when the relative change in the weighted residual diagnostic S_j falls below a prescribed tolerance or the maximum iteration count is reached.

G. Posterior Prediction

Let $\hat{m}_j$, $\hat{\boldsymbol{\kappa}}_j$, and $\hat{\sigma}_j^2$ denote the learned parameters, and let

$$q_j(\mathbf{f}_j) = \mathcal{N}(\mathbf{f}_j | \hat{\boldsymbol{\mu}}_{f,j}, \hat{\mathbf{C}}_{f,j}). \quad (31)$$

For a test input $\mathbf{x}^*$, let $f_{*j} \triangleq f_j(\mathbf{x}^*)$ and define

$$\begin{aligned} \hat{\mathbf{K}}_j &\triangleq \mathbf{K}_j(\hat{\boldsymbol{\kappa}}_j), \quad \hat{\mathbf{k}}_{*j} \triangleq [\mathbf{k}_j(\mathbf{x}^i, \mathbf{x}^*; \hat{\boldsymbol{\kappa}}_j)]_{i=1}^n, \\ \hat{k}_{**j} &\triangleq k_j(\mathbf{x}^*, \mathbf{x}^*; \hat{\boldsymbol{\kappa}}_j). \end{aligned} \quad (32)$$

Marginalizing the conditional GP distribution over (31) gives

$$\begin{aligned} q_j(f_{*j}) &= \mathcal{N}(f_{*j} | \mu_{*j}, v_{*j}), \\ \mu_{*j} &= \hat{m}_j + \hat{\mathbf{k}}_{*j}^\top \hat{\mathbf{K}}_j^{-1} (\hat{\boldsymbol{\mu}}_{f,j} - \hat{m}_j \mathbf{1}_n), \\ v_{*j} &= \hat{k}_{**j} - \hat{\mathbf{k}}_{*j}^\top \hat{\mathbf{K}}_j^{-1} \hat{\mathbf{k}}_{*j} + \hat{\mathbf{k}}_{*j}^\top \hat{\mathbf{K}}_j^{-1} \hat{\mathbf{C}}_{f,j} \hat{\mathbf{K}}_j^{-1} \hat{\mathbf{k}}_{*j}. \end{aligned} \quad (33)$$

For a nominal future observation y_{*j} , the predictive distribution is obtained as

$$q_j(y_{*j}) = \mathcal{N}(y_{*j} | \mu_{*j}, v_{*j} + \hat{\sigma}_j^2). \quad (34)$$




Fig. 1. Prediction performance under different outlier settings: (a) synthetic data with uniform outliers over $[0, 10]$; (b) Air Quality data with uniform outliers over $[-5, 15]$; and (c) Energy Efficiency data with zero-mean Gaussian outliers of variance 25.

III. NUMERICAL EXPERIMENTS

All experiments were conducted in MATLAB R2026a on an AMD Ryzen 9 9950X desktop with 64 GB RAM. We compare ASOR-GPR with standard GPR [1], robust conjugate GPR (RCGPR) [5], Student- t GPR [4], Gaussian-mixture-likelihood GPR (GMM-GPR) [9], and an Oracle GPR supplied with all available ground-truth contamination information.

All methods use independent scalar-output models with the ARD squared-exponential kernel, initialized with $\sigma_{f,j}^{(0)} = 1$ and $\ell_{jr}^{(0)} = 0.2d_{\text{med}}$, where d_{med} is the median pairwise training-input distance. All non-oracle methods are initialized with nominal-noise variance 0.5. For synthetic data, this is deliberately misspecified relative to the true variance 0.25. For the real datasets, the true nominal-noise variance is unknown. Predictive experiments use $n_{\text{train}} = n_{\text{test}} = 100$.

Outliers are injected entry-wise only into the training targets over 30 Monte Carlo trials. Root mean square error (RMSE) for synthetic data is evaluated against the known latent function at the test inputs. For real data, the original observations are treated as nominal, with outliers injected only into the training targets and RMSE evaluated against the unmodified test targets, which retain their inherent observation noise.

Baseline-specific parameters follow their respective references unless otherwise stated. ASOR-GPR uses $a = 1$, $\theta_{ij} = 0.5$, $A = 10$, $B = 0.45$, $\nu_0 = 3$, $b_j^{(0)} = 20$, and $s_{0j} = 1.25$ in all real-data and synthetic experiments. Up to 1000 variational iterations are run, terminating when relative change in S_j falls below 10^{-5} . Kernel hyperparameters are optimized by normalized gradient descent with Armijo backtracking using initial step $\alpha_j = 0.05$, Armijo constant $c_1 = 10^{-4}$, minimum step $\alpha_{\min} = 10^{-7}$, and at most 25 inner iterations.

For the synthetic experiment, $d_x = 2$ and $d_y = 4$. Inputs are Latin-hypercube sampled uniformly over $[-3, 3]^2$. Defining $z_{ir} = (x_{ir} + 3)/6$, $r = 1, 2$, the clean outputs are generated as

$$\begin{aligned} g_j(\mathbf{x}^i) &= 10 \sin(\pi z_{i1} z_{i2}) + 5 + 0.75 \sin((0.8 + 0.1j)x_{i1}) \\ &\quad + 0.50 \cos((0.6 + 0.05j)x_{i2}) + 0.10j x_{i1}, \\ \tilde{\mathbf{g}}(\mathbf{x}^i) &= \mathbf{C}_y^\top [g_1(\mathbf{x}^i) \quad \cdots \quad g_{d_y}(\mathbf{x}^i)]^\top, \\ \mathbf{C}_y &= \mathbf{I}_{d_y} + 0.1 (\mathbf{1}_{d_y} \mathbf{1}_{d_y}^\top - \mathbf{I}_{d_y}). \end{aligned} \quad (35)$$

Here, $\tilde{\mathbf{g}}(\mathbf{x}^i)$ denotes the mixed clean output vector. The outputs are standardized to zero mean and unit variance before adding nominal Gaussian noise and entry-wise contamination, with $p_{\text{out}} \in \{0, 0.1, \dots, 0.8\}$.

Fig. 1a considers outliers drawn uniformly from $[0, 10]$. ASOR-GPR remains competitive across the contamination range and achieves the lowest RMSE among the compared non-oracle methods at most contamination levels. For real-data evaluation, we use the Air Quality [15] and Energy Efficiency [16] datasets, standardized using training-set statistics. Using the same probability sweep, Fig. 1b considers uniform outliers over $[-5, 15]$, while Fig. 1c considers zero-mean Gaussian outliers with variance 25. ASOR-GPR provides clear gains under asymmetric contamination in Fig. 1b and remains competitive with robust likelihood-based methods under symmetric Gaussian outliers in Fig. 1c. Similar trends are observed over a wider range of contamination intensity. This reflects the effectiveness of the proposed adaptive inference of observation-specific precisions and nominal/outlier associations.

TABLE I
MEDIAN SERIAL MODEL-FITTING TIME (S) OVER 30 MONTE CARLO TRIALS.

Method	n_{train}				
	200	400	600	800	1000
Oracle GPR	0.10	0.70	1.99	4.37	6.76
Std. GPR	0.19	1.46	4.24	9.43	15.05
RCGPR	0.50	4.74	15.58	36.45	62.73
Student- t GPR	3.25	16.23	41.39	80.75	123.53
ASOR-GPR	24.60	31.13	85.52	212.41	335.96
GMM-GPR	31.96	151.23	376.66	644.76	963.38

For runtime evaluation, Table I reports median model-fitting times with increasing training-set size at $p_{\text{out}} = 0.4$ under zero-mean Gaussian contamination with variance 10. Oracle GPR and standard GPR have the lowest runtimes, with the Oracle benefiting from known contamination information. RCGPR and Student- t GPR incur moderate computational cost. ASOR-GPR and GMM-GPR exhibit the highest runtimes, with ASOR-GPR consistently faster than GMM-GPR over the reported training sizes. This relative computational behavior remains broadly consistent across different contamination probabilities. All methods involve dense covariance-matrix operations and retain cubic complexity, while runtime differences arise from their method-specific inference procedures.

IV. CONCLUSION

We proposed an outlier-robust GPR approach that embeds adaptive observation-specific precision modeling within a hierarchical generative framework. Variational generalized-EM inference enables observation-specific contamination effects and the underlying GP model to be learned jointly. Results on synthetic and real datasets show that ASOR-GPR achieves competitive predictive performance with enhanced robustness across a range of contamination settings, while preserving the cubic computational scaling of the standard GPR.

REFERENCES

- [1] C. E. Rasmussen and C. K. I. Williams, *Gaussian Processes for Machine Learning*. MIT Press, 2006.
- [2] P. J. Huber, "Robust estimation of a location parameter," *The Annals of Mathematical Statistics*, vol. 35, no. 1, pp. 73–101, 1964.
- [3] L. Chang, K. Li, and B. Hu, "Huber's m-estimation-based process uncertainty robust filter for integrated INS/GPS," *IEEE Sensors Journal*, vol. 15, no. 6, pp. 3367–3374, 2015.
- [4] P. Jylänki, J. Vanhatalo, and A. Vehtari, "Robust Gaussian process regression with a Student- t likelihood," *JMLR*, 2011.
- [5] M. Altamirano, F.-X. Briol, and J. Knoblauch, "Robust and conjugate Gaussian process regression," *ICML*, 2024.
- [6] H. Wang, H. Li, J. Fang, and H. Wang, "Robust Gaussian Kalman filter with outlier detection," *IEEE Signal Processing Letters*, vol. 25, no. 8, pp. 1236–1240, 2018.
- [7] A. H. Chughtai, M. Tahir, and M. Uppal, "Outlier-robust filtering for nonlinear systems with selective observations rejection," *IEEE Sensors Journal*, vol. 22, no. 7, pp. 6887–6897, 2022.
- [8] A. Nakabayashi and G. Ueno, "Nonlinear filtering method using a switching error model for outlier-contaminated observations," *IEEE Transactions on Automatic Control*, vol. 65, no. 7, pp. 3150–3156, 2019.
- [9] A. Daemi, H. Kodamana, and B. Huang, "Gaussian process modelling with Gaussian mixture likelihood," *Journal of Process Control*, vol. 81, pp. 209–220, 2019.
- [10] A. H. Chughtai, M. Tahir, and M. Uppal, "Bayesian heuristics for robust spatial perception," *IEEE Transactions on Instrumentation and Measurement*, vol. 73, pp. 1–11, 2024.
- [11] —, "EMORF/S: EM-based outlier-robust filtering and smoothing with correlated measurement noise," *IEEE Transactions on Signal Processing*, vol. 72, pp. 4318–4331, 2024.
- [12] A. Majal, A. H. Chughtai, and M. Tahir, "EMORF-II: Adaptive EM-based outlier-robust filtering with correlated measurement noise," in *2025 IEEE 35th International Workshop on Machine Learning for Signal Processing (MLSP)*, 2025, pp. 1–6.
- [13] H. Yang, P. Antonante, V. Tzoumas, and L. Carlone, "Graduated non-convexity for robust spatial perception: From non-minimal solvers to global outlier rejection," *IEEE Robotics and Automation Letters (RA-L)*, vol. 5, no. 2, pp. 1127–1134, 2020.
- [14] V. Šmídl and A. Quinn, *The Variational Bayes Method in Signal Processing*. Springer, 2006.
- [15] S. Vito, "Air Quality," UCI Machine Learning Repository, 2008, DOI: <https://doi.org/10.24432/C59K5F>.
- [16] A. Tsanas and A. Xifara, "Energy Efficiency," UCI Machine Learning Repository, 2012, DOI: <https://doi.org/10.24432/C51307>.