arXiv:2610.39575v1 [cond-mat.soft] 29 Oct 2026

Hidden Phase Transition in Single-Polymer Stretching under Elongational Flow

Aiqing Zhu[1], Fushuai Wang[2,3], Eleonore Vissol-Gaudin[4], Qianxiao Li[*1,3], and Beatrice W. Soh[†2,3]

[1]Department of Mathematics, National University of Singapore, 10 Lower Kent Ridge Road, 119076, Singapore
[2]Department of Chemical and Biomolecular Engineering, National University of Singapore, 4 Engineering Drive 4, 117585, Singapore
[3]Institute for Functional Intelligent Materials, National University of Singapore, 4 Science Drive 2, 117544, Singapore
[4]Fujitsu Research of Europe, 3–9 Albert Street, Slough SL1 2BE, United Kingdom

Abstract

Using machine learning, we identify a second transition in polymer chain stretching driven by planar elongational flow, beyond the well-known coil-stretch transition. We achieve this by developing a conditional extension of Universal and Identifiable Dissipative Dynamics (UIDD) that jointly learns reduced stochastic dynamics from microscopic trajectories across flow strengths and yields the corresponding stationary nonequilibrium free energy landscapes. As the flow strength increases, the evolving landscape exhibits the expected coil-stretch transition, and then a previously unrecognized dynamical transition. In the latter, the stretched states persist, but the distribution along a folding coordinate becomes bimodal when conditioned on small end-to-end separation. Large-deviation analysis of a simplified analytical chain model reproduces both transitions and shows that they are phase transitions in the long-chain limit. Single-molecule DNA experiments provide qualitative support for the dynamical transition through the predicted reversal of the folding response.

Introduction— The coil-stretch transition in elongational flow is a paradigmatic problem in nonequilibrium polymer dynamics, with important consequences for the rheology of polymer solutions [1, 2]. As predicted by de Gennes [3], the competition between flow-induced deformation and entropic relaxation drives a sharp change from an entropically coiled state to a highly extended state above a critical flow strength, while conformation-dependent hydrodynamic drag gives rise to hysteresis. Single-molecule experiments have resolved this transition directly [4, 5] and observed hysteresis for sufficiently long chains [6]. Minimal analytical models capture this behavior with a single extension variable [3, 7, 8]. Although a single chain is a finite system, the transition sharpens with increasing chain length, so coil-stretch behaves as a phase transition in the long-chain limit [3, 6]. These models leave open the question of whether coil-stretch fully defines the transition structure of the microscopic chain, or whether additional transitions emerge along conformational degrees of freedom not resolved by extension.

Experiments and simulations suggest that the underlying dynamics are richer than extension alone. Single-molecule experiments reveal "molecular individualism" during stretching, in which chains with the same extension adopt qualitatively distinct conformations and evolve along different pathways [4, 5]. Brownian dynamics simulations reproduce these conformational classes and show that populations shift with flow strength [9]. This motivates examining how the overall conformational distribution reorganizes with flow strength, without restricting the description to extension alone.

The direct construction of such a description from microscopic dynamics remains difficult, due to the high dimensionality of the underlying polymer system and its many coupled degrees of freedom. Conventional tools each address only part of the problem. Histograms in chosen coordinates presuppose which coordinates matter, while dimensionality reduction methods such as principal component analysis and diffusion maps[10] can identify relevant coordinates, but do not yield dynamics or an interpretable potential. Deep learning offers a way to learn reduced coordinates and their dynamics jointly [11–14]. Stochastic OnsagerNet demonstrated that effective landscapes can be learned directly from microscopic polymer trajectories [15]. Universal and Identifiable Dissipative Dynamics (UIDD) further established the identifiability of the learned potential [16]. In previous work, landscapes were learned separately at each flow strength, so the reduced coordinates were not necessarily consistent across conditions, making it difficult to track how the landscape reorganizes and to identify new transitions. Since UIDD identifies the landscape uniquely, extending it to a single model with varying driving conditions would place all landscapes in shared coordinates. Their reorganization with flow strength can then be tracked directly. This turns landscape learning into a systematic tool for detecting transitions, including

*Contact author: qianxiao@nus.edu.sg
†Contact author: bwysoh@nus.edu.sg

those invisible to conventional order parameters.

In this Letter, we develop this approach to revisit the transition structure of a single polymer chain in planar elongational flow (Fig. 1) and show that, beyond coil-stretch, the chain undergoes a second dynamical transition. At higher flow strength, oppositely backfolded saddle states emerge among chains with small end-to-end separation, while the dominant stretched states persist, which is a change invisible in extension. We then develop an independent analytical model and derive its nonequilibrium free energy landscapes using large-deviation theory. This simplified model reproduces both transitions and traces them to a competition between conformational entropy and flow-driven axial spreading. In the long-chain limit, the second becomes a phase transition between unfolded and oppositely folded states. Our single-molecule DNA experiments further provide qualitative support for the dynamical transition.

Kinetic model— The microscopic kinetic model we use is Brownian dynamics for linear, touching-bead polymer chains subjected to an external planar elongational flow of strain rate $\dot{\epsilon}$ [17], where $\dot{\epsilon}$ serves as the control parameter, as illustrated in Fig. 1(a). Each chain consists of $N+1=300$ beads, connected by N rigid rods. The governing stochastic dynamics accounts for excluded volume and Brownian fluctuations, and neglects hydrodynamic interactions (Supplemental Material). For each microscopic conformation, we record the three-dimensional coordinates of all beads, from which the chain extension Z_1 along the elongational axis x is computed. To construct an effective reduced description, we augment Z_1 with two additional coordinates [15], Z_2 and Z_3, using a symmetry-preserving principal component analysis (PCA). Correlation analysis associates Z_2 and Z_3 with the signed end-to-end extension and the signed foldedness, respectively (End Matter). Each microscopic trajectory at strain rate $\dot{\epsilon}$ thus yields a time series of (Z_1, Z_2, Z_3).

To learn how dynamics and stationary statistics vary with external conditions, we introduce conditional UIDD, an extension of UIDD [16]. For a d-dimensional state $\mathbf{Z} \in \mathbb{R}^d$ and condition $c \in \mathcal{C}$, shared neural functions define a family of Itô SDEs of the form

$$\begin{aligned} \mathrm{d}\mathbf{Z}_t = \Big[&-\big(M(\mathbf{Z}_t;c) + W(\mathbf{Z}_t;c)\big)\nabla_{\mathbf{Z}} V(\mathbf{Z}_t;c) \\ &+ \nabla_{\mathbf{Z}}\cdot M(\mathbf{Z}_t;c) + \nabla_{\mathbf{Z}}\cdot W(\mathbf{Z}_t;c)\Big]\,\mathrm{d}t + \sigma(\mathbf{Z}_t;c)\,\mathrm{d}\mathbf{B}_t, \end{aligned} \tag{1}$$

where V is a scalar potential and M, W, σ are $d \times d$ matrices depending on state $\mathbf{Z}$ and condition c. Here, the diffusion σ and Brownian motion $\mathbf{B}_t$ model the thermal fluctuations, and the symmetric matrix $M = \sigma\sigma^\top/2$ characterizes the dissipative response satisfying the fluctuation-dissipation relation. The conservative parts are encoded by a banded antisymmetric matrix W with nonzero entries only on the first sub- and super-diagonals. The fields depend on $(\mathbf{Z}, c)$, while all derivatives act only on $\mathbf{Z}$.

This construction possesses desirable structural properties linking the dynamics to its stationary statistics under fixed condition. It is proved that the dynamics retains full expressivity, and the resulting density

$$\rho(\mathbf{Z};c) = \frac{1}{\mathcal{Z}(c)}\exp[-V(\mathbf{Z};c)] \tag{2}$$

satisfies the stationary Fokker-Planck equation, constituting a key advantage over other neural dynamics models [13–15, 18, 19] (End Matter and Supplemental Material). Consequently, V is identifiable and coincides, up to an additive constant, with $F = -\ln\rho$. We refer to V (and F) as the nonequilibrium free energy; in a driven steady state, its minima and barriers encode the stationary statistics of the chain. Therefore, changes in the critical point structure of the learned landscape $V(\mathbf{Z};c)$ as c varies provide finite-size signatures of transitions.

Here, we express the flow strength in dimensionless form as the Weissenberg number $\mathrm{Wi} = \dot{\varepsilon}\tau$, where τ is the longest relaxation time of the chain. We take the condition parameter to be $c = \mathrm{Wi}$ and the reduced state to be $\mathbf{Z} = (Z_1, Z_2, Z_3)$. V, W and σ are each parameterized as trainable deep neural networks, and their parameters are fitted by jointly maximizing the conditional transition likelihood (End Matter). The learned dynamics, validated on various initial chain conformations, accurately predict the evolution of chain extension Z_1 in all cases (Supplemental Material).

To examine end-separation statistics and the associated folding structure, we introduce the dimensionless marginal free energies

$$F_{23}(Z_2, Z_3; c) = -\ln\int \exp[-V(\mathbf{Z};c)]\,\mathrm{d}Z_1,$$
$$F_2(Z_2; c) = -\ln\iint \exp[-V(\mathbf{Z};c)]\,\mathrm{d}Z_1\,\mathrm{d}Z_3,$$

and the section $F_{23}(0, Z_3; c)$, which probes the landscape along $Z_2 = 0$ (small-end-separation) and is distinct from a marginal free energy obtained by integrating over Z_2. The results are presented in Fig. 2 (a-d).

At $\mathrm{Wi} = 0.01$, F_{23} has a single minimum at the origin and F_2 exhibits a single well, consistent with a chain in the coiled state. By $\mathrm{Wi} = 0.66$, the origin becomes a saddle of F_{23}, with two dominant minima along Z_2; F_2 becomes double-welled while $F_{23}(0, Z_3)$ retains a single minimum at $Z_3 = 0$. Since Z_2 is strongly correlated with the signed end-to-end separation, the two minima correspond to the two orientations of a stretched chain. Hence, the coil-stretch transition appears as a symmetric splitting of the coiled minimum, consistent with the absence of coil-stretch bistability in a model without hydrodynamic interactions [20]. This reorganization recovers the onset of coil-stretch near the critical value $\mathrm{Wi}_t = 0.5$ [3, 4] commonly identified through chain extension, here as a change in the nonequilibrium free energy landscape learned from the microscopic kinetic model.



Figure 1: Overview of the polymer system and transition analysis. (**a**) A single polymer chain is represented by a bead-rod model, subjected to planar elongational flow, with chain extension measured along the elongational axis. (**b**) Plot of chain extension as a function of time across varying flow strengths. (**c**) Microscopic trajectories are reduced by PCA and fitted with conditional UIDD to identify phase transitions; large-deviation theory independently supports both transitions, while single-molecule DNA experiments provide qualitative support for the second transition.



Figure 2: Free energy landscapes and transition diagrams. (**a-d**) Learned marginal free energies $F_2(Z_2)$ and sections $F_{23}(0, Z_3)$ at the indicated Wi. Insets show the corresponding landscapes $F_{23}(Z_2, Z_3)$. (**e,f**) Near-minimizing sets of the learned F_2 ($\eta = 0.6$) and $F_{23}(0, \cdot)$ ($\eta = 1.0$), respectively, defined by $\{q : F(q; \mathrm{Wi}) - \min_{q'} F(q'; \mathrm{Wi}) \le \eta\}$. Color intensity indicates the number of the five independently trained models satisfying this condition. For each model, the transition point Wi_t is determined by the first crossing of $F(0; \mathrm{Wi}) - \min_q F(q; \mathrm{Wi})$ above η. Orange dashed lines and bands indicate the mean and range of Wi_t across models, respectively, with sensitivities shown in the insets. (**g-j**) Free energy profiles per bond of the analytical model, $F_{2,n}(z_2)$ and $F_{23,n}(0, z_3)$. (**k,l**) Minimizing sets of $F_{2,\infty}(z_2)$ and $F_{23,\infty}(0, z_3)$, respectively. Beyond the second transition, the gray region in (**l**) denotes the minimizing interval, with diamonds marking its endpoints.

At $\mathrm{Wi} = 2.64$, the origin changes from a saddle to a local maximum of F_{23}, while two symmetry-related saddles emerge along Z_3, appearing as off-center minima of $F_{23}(0, Z_3)$. This marks a second transition, which splits the small-end-separation population into two oppositely oriented backfolded conformations, while the dominant stretched minima persist. Unlike coil-stretch, this transition changes the unstable states of the landscape instead of the dominant stable states. Since saddles govern the pathways and timescales of unfolding, we refer to it as a dynamical transition, and show below that it becomes a phase transition in the long-chain limit. Previous work [15] identified prolonged residence near folded state saddles as a key driver of heterogeneous unfolding times [4–6]. This transition thus indicates the flow-induced emergence of the unstable states underlying this dynamical heterogeneity, consistent with the growing prevalence of folded unraveling pathways at strong flows. To our knowledge, this transition has not been identified previously, and it is invisible to analyses restricted to stable states

Fig. 2 (e,f) locate the two transitions through a systematic scan of Wi in the learned landscapes. Learning errors introduce seed-dependent variations in the free energy profiles, so we track near-minimizing sets, defined as the regions where the free energy lies within a tolerance η of its minimum, across five independently trained models. The near-minimizing set first separates into stretched branches along Z_2, followed by a separation in the section $F_{23}(0, Z_3)$ at higher flow strength. The stretched branches persist through the dynamical transition, distinguishing the reorganization of the small-end-separation population from the initial coil-stretch transition. At the selected tolerances, the mean transition estimates are $\mathrm{Wi}_t^{(1)} \simeq 0.42$ and $\mathrm{Wi}_t^{(2)} \simeq 1.46$, with respective ranges $[0.37, 0.46]$ and $[0.89, 1.92]$. The two ranges do not overlap, confirming that the transitions are distinct.

Analytical model— To assess the theoretical plausibility of the two transitions and explore their mechanism, we consider a simplified two-dimensional model. It consists of $n+1$ beads at positions $r_0, \ldots, r_n \in \mathbb{R}^2$, subject to $\sum_{k=0}^{n} r_k = 0$ and $|r_k - r_{k-1}| = 1$ for $k = 1, \ldots, n$, with no additional inter-bead interactions. The dimensionless effective flow potential for a planar elongational flow is

$$U_{n,\lambda}(\{r_k\}_{k=0}^n) = -\frac{\lambda}{n^2} \sum_{k=0}^{n} \left(r_{k,x}^2 - r_{k,y}^2\right), \qquad (3)$$

where $\lambda \geq 0$ controls the elongational strength ($\mathrm{Wi} = 0.096\lambda$, see Sec. S3.C). The interpretation of the learned closure coordinates motivates the reduced observables

$$\hat{z}_2 = (r_{n,x} - r_{0,x})/n, \quad \hat{z}_3 = (r_{n,x} + r_{0,x})/n.$$

To compare with the learned estimates, we introduce

$$\mathcal{Z}_{2,n}(z_2; \lambda) = \int e^{-U_{n,\lambda}} \delta(z_2 - \hat{z}_2)\, \mathrm{d}\nu_n,$$
$$\mathcal{Z}_{23,n}(0, z_3; \lambda) = \int e^{-U_{n,\lambda}} \delta(\hat{z}_2)\delta(z_3 - \hat{z}_3)\, \mathrm{d}\nu_n,$$

where δ denotes the Dirac distribution and ν_n is the reference configurational measure induced by independent, uniformly distributed bond directions. The corresponding dimensionless free energies per bond are

$$F_{2,n} = -\ln \mathcal{Z}_{2,n}/n, \quad F_{23,n} = -\ln \mathcal{Z}_{23,n}/n. \qquad (4)$$

In the long-chain limit, the scaled interpolated chain belongs to the following subset of the absolutely continuous path space $\mathrm{AC}([0,1]; \mathbb{R}^2)$:

$$\mathcal{A} = \left\{ m \in \mathrm{AC}([0,1]; \mathbb{R}^2) : m(0) = 0,\ |m'(s)| \leq 1 \text{ a.e.} \right\}.$$

Writing $\bar{m} = \int_0^1 m(s), \mathrm{d}s$, we define the effective energy and conformational entropy by

$$\mathcal{E}[m] = -\int_0^1 \left[(m_x - \bar{m}_x)^2 - (m_y - \bar{m}_y)^2\right] \mathrm{d}s,$$
$$\mathcal{S}[m] = -\int_0^1 \sup_{h \in \mathbb{R}} \left\{h|m'| - \ln I_0(h)\right\} \mathrm{d}s, \qquad (5)$$

where $I_0(h) = (2\pi)^{-1} \int_0^{2\pi} e^{h\cos\theta} \mathrm{d}\theta$ is the modified Bessel function of the first kind of order zero. By Mogulskii's theorem and Varadhan's lemma [21–23], the limiting free energies are given by (End Matter)

$$F_{2,\infty}(z_2; \lambda) = \inf_{\substack{m \in \mathcal{A} \\ m_x(1) = z_2}} \left\{\lambda \mathcal{E}[m] - \mathcal{S}[m]\right\},$$
$$F_{23,\infty}(0, z_3; \lambda) = \inf_{\substack{m \in \mathcal{A},\ m_x(1) = 0 \\ \bar{m}_x = -z_3/2}} \left\{\lambda \mathcal{E}[m] - \mathcal{S}[m]\right\}. \qquad (6)$$

These variational problems show that the topological changes of free energy minima arise from the competition between the energetic preference for axial spreading and the entropic cost of local alignment.

Fig. 2 (g-j) presents the variational solutions of (6) and Monte Carlo estimates of finite chain free energies (4). The analytical model qualitatively reproduces both the coil-stretch transition, marked by the splitting of the extension minimum, and the subsequent dynamical transition at zero axial end separation. This sequence arises as increasing Wi (and hence λ) strengthens the elongational energy contribution relative to entropy, reshaping the free energy landscape. At $\mathrm{Wi} = 5.28$, $F_{23,\infty}(0, z_3)$ is minimized on a symmetric interval containing the origin (Theorem S3), which finite-size effects turn into the two isolated minima seen for $n = 16, 32$ and in the machine learning results. This flat interval signals phase coexistence between oppositely folded states, making the second transition a phase transition in the long-chain limit. Scans of the variational free energies locate the two transitions at $\mathrm{Wi}_t^{(1)} \simeq 0.95$ and $\mathrm{Wi}_t^{(2)} \simeq 3.79$ [Fig. 2 (k,l)], in



Figure 3: (**a**) Definition of the image-based folding coordinate $\widetilde{Z}_3$. (**b**) Schematic of the single-well to double-well transition in the free energy landscape and the corresponding increments near the origin. (**c**) Experimental signatures of folding reorganization. Symbols show the odd component of the conditional finite-lag mean increment of $\widetilde{Z}_3$. Error bars indicate uncertainty in the estimated response. Curves show polynomial fits; triangles mark zero crossings interpolated from the binned data at 40 V. Insets show apparent potentials obtained as negative antiderivatives of the fitted polynomials.

qualitative agreement with the learned estimates for the full kinetic model.

Experimental signatures— To test whether the predicted dynamical transition appears in experiment, we performed single-molecule experiments [24] on DNA molecules stretched in a planar elongational electrokinetic field that is analogous to a planar elongational flow, with the field strength controlled by the applied voltage (Supplemental Material). The applied voltages of 10-50 V correspond to Wi $\approx$ [0.7,4.8]. The transition is characterized by changes in the free energy landscape along Z_3, but this coordinate is not directly accessible from molecular images. Motivated by the association of Z_3 with the axial displacement of the chain-end midpoint from the center of mass, we define its image-based surrogate $\widetilde{Z}_3$ as twice the axial displacement of the midpoint between the molecule's axial edges from the intensity-weighted centroid, centered by its trajectory median [Fig. 3(a)].

Across the second transition, the origin of $F_{23}(0, Z_3)$ changes from a local minimum to a local maximum. Since the drift near the origin follows the local slope of the landscape, small deviations of Z_3 from zero should be restored below the transition and amplified above it, reversing the slope of the mean response at the origin from negative to positive [Fig. 3(b)]. To test this, we measure the mean change in $\widetilde{Z}_3$ over a fixed time lag, conditioned on its current value, for molecules within a common window of normalized extension during initial stretching, and retain its odd component. As seen in Fig. 3(c), at 10 and 20 V, the response is restoring near $\widetilde{Z}_3 = 0$, whereas at 40 and 50 V it is repelling, with outer zero crossings at 40 V marking the two folded states, see End Matter for details. This reversal agrees with the predicted change of the landscape and provides qualitative experimental support for the dynamical transition.

Discussion— Beyond the coil-stretch transition, a polymer chain in planar elongational flow undergoes a second dynamical transition, in which backfolded saddle states emerge among chains with small end-to-end separation, while the stretched states persist. To identify this, we developed conditional UIDD to learn stochastic dynamics and free energy landscapes jointly across flow strengths. Large-deviation analysis on a simplified model reproduces both transitions and shows that they are phase transitions in the long-chain limit. Since saddles divide the pathways by which chains stretch, sending a fraction of chains through folded intermediates, this transition offers a landscape-level explanation for why folds become common during unraveling at strong flows [5, 9]. As this transition is invisible in the end-to-end vector, it also identifies a folding coordinate that dumbbell-based constitutive models omit. The landscapes further specify what to measure in experiment, namely a reversal of the folding response from restoring to repelling, which our single-molecule DNA experiments observe.

More broadly, our workflow applies to any system observed through trajectories under a tunable control, such as biomolecules under force or colloidal assembly at varying interaction strength. Since its identifiability does not require detailed balance and it accommodates several conditions at once, it can map transitions across nonequilibrium phase diagrams by tracking how unstable states change across multi-parameter spaces, including those that change unstable states while the most probable states stay the same.

Methods

Closure coordinates— In our previous study of the same polymer system [15, 16], two closure coordinates are identified and associated with the axial end-to-end distance and degree of foldedness, respectively. These coordinates were obtained from a nonlinear PCA-ResNet encoder and interpreted a posteriori through geometric correlations [15]. To compare conformational statistics and free energy landscapes across flow strength without ambiguity from changes in the representation itself, we construct a common symmetry-preserving PCA representation with prescribed parity under the physical symmetries of the chain. Here $\mathbf{r}_i = (r_{i,x}, r_{i,y}, r_{i,z})$ is the position of bead $i = 0, \ldots, N$, with 0 and N labeling the chain ends. The x axis is the elongational direction, and $r_{\mathrm{com},x} = (N+1)^{-1} \sum_{i=0}^{N} r_{i,x}$ is the axial center-of-mass coordinate. Specifically, we impose chain-end exchange

$R : \mathbf{r}_i \mapsto \mathbf{r}_{N-i}$ and the three Cartesian coordinate reflections $T_\alpha : r_{i,\alpha} \mapsto -r_{i,\alpha}$, $\alpha = x, y, z$. The coordinate Z_2 is odd under R and T_x and even under T_y and T_z, whereas Z_3 is even under R, odd under T_x, and even under T_y and T_z. A single projection and normalization are used over all data. Details are given in Supplemental Material.



Figure 4: Correlations of the PCA coordinates. Top: Z_2 versus the signed axial endpoint difference, $r_{0,x} - r_{N,x}$. Bottom: Z_3 versus the signed foldedness, $(r_{0,x} + r_{N,x})/2 - r_{\mathrm{com},x}$. Correlation coefficients are computed separately for each Wi interval.

We then test whether these symmetry-adapted coordinates retain the geometric interpretations across flow strengths by evaluating the corresponding correlations separately. As shown in Fig. 4, Z_2 is strongly correlated with the signed axial endpoint difference, whereas Z_3 is strongly correlated with the signed foldedness.

Conditional UIDD— Conditional UIDD learns a family of stochastic dynamics on $\mathbb{R}^d$ from trajectories labeled by a fixed external condition $c \in \mathcal{C} \subset \mathbb{R}^p$. It couples the inference problems at different conditions through shared state-and-condition maps V, W, and σ. For each fixed c, normalizability and dissipation conditions [25] yield the existence and uniqueness of the stationary density for Eq. (1). Substituting ρ from Eq. (2) into the Fokker-Planck current [26] we have

$$\mathbf{J} = f\rho - \nabla_{\mathbf{Z}} \cdot (M\rho) = \nabla_{\mathbf{Z}} \cdot (W\rho),$$

where $(\nabla_{\mathbf{Z}} \cdot A)_i = \sum_j \partial_{Z_j} A_{ij}$. The tensor $W\rho$ is antisymmetric, so its double divergence vanishes and thus

$$\nabla_{\mathbf{Z}} \cdot \mathbf{J} = 0.$$

Consequently, the energy potential within each condition are identifiable, although the scalar reference $C(c)$ need not be unique.

We first split the time series into multiple transitions. For each transition a, the state $\mathbf{Z}_a$, condition c_a, increment $\Delta\mathbf{Z}_a$, and observation interval Δt_a enter a joint conditional likelihood. With f denoting the drift of Eq. (1), define $e_a = \Delta\mathbf{Z}_a - f(\mathbf{Z}_a; c_a)\Delta t_a$ and the positive-definite covariance $C_a = 2M(\mathbf{Z}_a; c_a)\Delta t_a$. The Euler-Maruyama approximation gives, up to parameter-independent terms, the objective [26–28]

$$\mathcal{L}(\theta) = \frac{1}{2} \sum_a \left[e_a^\top C_a^{-1} e_a + \ln \det C_a \right]. \tag{7}$$

Minimizing this objective over all condition-labeled transitions fits the parameters θ. Training requires neither stationary initial samples nor evaluation of the normalization $\mathcal{Z}(c)$. The proof of universality of the conditional UIDD representation, the network construction, training settings, and landscape evaluation are given in Supplemental Material.

Large-deviation derivation— Fix $\lambda \geq 0$ and linearly interpolate $m_n(k/n) = (r_k - r_0)/n$. These paths lie in $\mathcal{A}$; write $\mathbb{P}_n$ for their distribution under ν_n. The unit bond directions are independent and uniform, with log moment-generating function

$$\Lambda(\mathbf{h}) = \ln \langle e^{\mathbf{h}\cdot(r_1 - r_0)} \rangle_{\nu_n} = \ln I_0(|\mathbf{h}|).$$

Its Legendre transform is therefore radial:

$$\Lambda^*(v) = \ell(|v|), \qquad \ell(u) = \sup_{h \in \mathbb{R}} \{hu - \ln I_0(h)\}. \tag{8}$$

Because Λ is finite on $\mathbb{R}^2$, Mogulskii's theorem [21] gives a path large-deviation principle (LDP) in the uniform topology, with speed n and conformational entropy (5):

$$\mathcal{S}[m] = -\int_0^1 \ell(|m'(s)|)\, \mathrm{d}s, \qquad m \in \mathcal{A},$$

with $\mathcal{S}[m] = -\infty$ outside $\mathcal{A}$.

The zero center-of-mass condition gives the exact relation

$$\frac{r_k}{n} = m_n(k/n) - \frac{1}{n+1} \sum_{j=0}^{n} m_n(j/n).$$

Substitution into Eq. (3) gives

$$-U_{n,\lambda} = -\lambda n \mathcal{E}[m_n] + o(n), \qquad \hat{z}_2 = m_{n,x}(1),$$
$$\mathcal{E}[m] = -\int_0^1 \left[(m_x - \bar{m}_x)^2 - (m_y - \bar{m}_y)^2 \right] \mathrm{d}s.$$

The error is uniform because the paths are 1-Lipschitz. Thus the n^{-2} prefactor makes the flow exponent scale with the LDP speed n. For $|z_2| < 1$, integrating the partition density over a window of fixed width 2ε gives

$$W_{2,n,\varepsilon}(z_2; \lambda) = \int e^{-U_{n,\lambda}} \mathbf{1}_{\{|\hat{z}_2 - z_2| < \varepsilon\}}\, \mathrm{d}\nu_n$$
$$\asymp \int_{\substack{m \in \mathcal{A} \\ |m_x(1) - z_2| < \varepsilon}} e^{-n\lambda\mathcal{E}[m]}\, \mathrm{d}\mathbb{P}_n(m),$$

where $\asymp$ means that the logarithms differ by $o(n)$. The energy $\mathcal{E}$ is bounded and continuous on $\mathcal{A}$. The constrained Laplace bounds [22] therefore give

$$-\lim_{n\to\infty} \frac{1}{n} \ln W_{2,n,\varepsilon} = \inf_{\substack{m \in \mathcal{A} \\ |m_x(1) - z_2| < \varepsilon}} \{\lambda\mathcal{E}[m] - \mathcal{S}[m]\}.$$

At each sufficiently large fixed n, the continuous density satisfies $W_{2,n,\varepsilon}/(2\varepsilon) \to \mathcal{Z}_{2,n}(z_2;\lambda)$ as $\varepsilon \downarrow 0$. Theorem S2 in the Supplemental Material proves that the ε and n limits commute at the logarithmic scale. Consequently,

$$\begin{aligned} F_{2,\infty}(z_2;\lambda) &= -\lim_{\varepsilon\downarrow 0}\lim_{n\to\infty}\frac{1}{n}\ln\left[\frac{W_{2,n,\varepsilon}(z_2;\lambda)}{2\varepsilon}\right] \\ &= \inf_{\substack{m\in\mathcal{A}\\ m_x(1)=z_2}}\{\lambda\mathcal{E}[m]-\mathcal{S}[m]\}. \end{aligned} \tag{9}$$

The section $F_{23,\infty}(0,z_3;\lambda)$ is obtained in the same way.

Conditional response of image asymmetry— We examine how the mean response of image asymmetry changes with increasing voltage (Fig. 3). To compare similar stages of initial stretching, reference frames are selected within a common window of normalized image span. The raw dataset comprises 48 trajectories (47,339 frames) at 10 V, 201 (34,716 frames) at 20 V, 335 (27,922 frames) at 40 V, and 448 (33,726 frames) at 50 V.

For each image, let $\mathcal{M}$ be the retained molecular mask, $x_{\min}, x_{\max}$ its axial extrema, and I_p the background-corrected intensity at pixel coordinate x_p. Define

$$\begin{aligned} x_{\mathrm{com}} &= \frac{\sum_{p\in\mathcal{M}} I_p x_p}{\sum_{p\in\mathcal{M}} I_p}, \quad \frac{q}{2} = \frac{x_{\min}+x_{\max}}{2} - x_{\mathrm{com}}, \\ \widetilde{Z}_3^{(i)}(t) &= q_i(t) - \operatorname*{median}_{s\in\mathcal{Q}_i} q_i(s). \end{aligned} \tag{10}$$

Here $\mathcal{Q}_i$ contains all quality-filtered frames of trajectory i (Sec. S4.A); coordinates are in pixels. Such image-based coordinate $\tilde{Z}_3$ correlates with the microscopic folding asymmetry $(r_{0,x}+r_{N,x})/2 - r_{\mathrm{com},x}$, as shown in Fig. 5. For conditioning, set $z(t) = \widetilde{Z}_3(t) - \mu$, where μ is the mean reference value at each voltage. This shift leaves increments unchanged. For two later frames, $0 < \delta_1 < \delta_2$, we define

$$\begin{aligned} \Delta\widetilde{Z}_3(t) &= \widetilde{Z}_3(t+\delta_2) - \widetilde{Z}_3(t+\delta_1), \\ f(z) &= \left\langle \Delta\widetilde{Z}_3(t) \,|\, z(t) = z \right\rangle, \\ g(z) &= \frac{f(z) - f(-z)}{2}. \end{aligned} \tag{11}$$

The use of two later frames separates the conditioning measurement from the finite-lag increment. The antisymmetrization extracts the component odd under $z \mapsto -z$, thereby focusing the analysis on the restoring or amplifying tendency shared by positive and negative deviations from the chosen center.

We estimate g in symmetric bins and quantify uncertainty by bootstrap resampling entire trajectories within each voltage ensemble, with the bin edges held fixed. The error bars in Fig. 3 denote pointwise 16th-84th percentile intervals from 2000 bootstrap resamples. The resampling specification and its remaining implementation details are discussed in Sec. S4.B. The marked zero crossings at 40 V are obtained by linear interpolation between adjacent odd-response bins.



Figure 5: Correlations of the image-based coordinates. The image-based coordinate $\tilde{Z}_3$ is extracted from images generated from simulated polymer conformations, whereas the microscopic folding asymmetry $x_1 + x_N - 2x_{\mathrm{com}}$ are computed directly from the simulated configurations.

The smooth curves in Fig. 3 are unweighted least-squares fits of

$$g_{\mathrm{fit}}(z) = c_1 z + c_3 z^3$$

to the retained bins, with no sign constraints on the coefficients. The corresponding apparent potentials are defined by

$$U_{\mathrm{app}}(z) = -\int_0^z g_{\mathrm{fit}}(u)\,\mathrm{d}u + C = -\frac{c_1 z^2}{2} - \frac{c_3 z^4}{4} + C.$$

Each inset is shifted to zero minimum over the displayed range and rescaled independently, so the comparison concerns the evolution of potential shape across voltages. These apparent potentials provide a compact representation of the fitted finite-lag responses. Experimental details, selection and lag parameters, and sensitivity analyses are given in Supplemental Material.

References

[1] Ronald G Larson. The rheology of dilute solutions of flexible polymers: Progress and problems. *Journal of Rheology*, 49(1):1–70, 2005.

[2] CM Schroeder. Single polymer dynamics for molecular rheology. *Journal of Rheology*, 62(1):371–403, 2018.

[3] PG De Gennes. Coil-stretch transition of dilute flexible polymers under ultrahigh velocity gradients. *The Journal of Chemical Physics*, 60(12): 5030–5042, 1974.

[4] Thomas T Perkins, Douglas E Smith, and Steven Chu. Single polymer dynamics in an elongational flow. *Science*, 276(5321):2016–2021, 1997.

[5] Douglas E Smith and Steven Chu. Response of flexible polymers to a sudden elongational flow. *Science*, 281(5381):1335–1340, 1998.

[6] Charles M Schroeder, Hazen P Babcock, Eric SG Shaqfeh, and Steven Chu. Observation of polymer conformation hysteresis in extensional flow. *Science*, 301(5639):1515–1519, 2003.

[7] EJ Hinch. Mechanical models of dilute polymer solutions in strong flows. *The Physics of Fluids*, 20 (10):S22–S30, 1977.

[8] RG Larson and JJ Magda. Coil-stretch transitions in mixed shear and extensional flows of dilute polymer solutions. *Macromolecules*, 22(7):3004–3010, 1989.

[9] RG Larson, Hua Hu, DE Smith, and S Chu. Brownian dynamics simulations of a dna molecule in an extensional flow field. *Journal of Rheology*, 43(2): 267–304, 1999.

[10] Ronald R Coifman and Stéphane Lafon. Diffusion maps. *Applied and computational harmonic analysis*, 21(1):5–30, 2006.

[11] Ricky TQ Chen, Yulia Rubanova, Jesse Bettencourt, and David K Duvenaud. Neural ordinary differential equations. In *Advances in neural information processing systems*, volume 31, 2018.

[12] Pinchen Xie, Roberto Car, and Weinan E. Ab initio generalized langevin equation. *Proceedings of the National Academy of Sciences*, 121(14): e2308668121, 2024.

[13] Haijun Yu, Xinyuan Tian, Weinan E, and Qianxiao Li. Onsagernet: Learning stable and interpretable dynamics using a generalized onsager principle. *Physical Review Fluids*, 6(11):114402, 2021.

[14] Zhen Zhang, Yeonjong Shin, and George Em Karniadakis. GFINNs: GENERIC formalism informed neural networks for deterministic and stochastic dynamical systems. *Philosophical Transactions of the Royal Society A*, 380(2229):20210207, 2022.

[15] Xiaoli Chen, Beatrice W Soh, Zi-En Ooi, Eleonore Vissol-Gaudin, Haijun Yu, Kostya S Novoselov, Kedar Hippalgaonkar, and Qianxiao Li. Constructing custom thermodynamics using deep learning. *Nature Computational Science*, 4(1):66–85, 2024.

[16] Aiqing Zhu, Beatrice W Soh, Grigorios A Pavliotis, and Qianxiao Li. Identifiable learning of dissipative dynamics. *Nature Communications*, 17(1): 9460, 2026. doi: 10.1038/s41467-026-76255-3.

[17] Beatrice W Soh, Alexander R Klotz, and Patrick S Doyle. Untying of complex knots on stretched polymers in elongational fields. *Macromolecules*, 51(23): 9562–9571, 2018.

[18] Shaan A Desai, Marios Mattheakis, David Sondak, Pavlos Protopapas, and Stephen J Roberts. Port-hamiltonian neural networks for learning explicit time-dependent dynamical systems. *Physical Review E*, 104(3):034312, 2021.

[19] Kookjin Lee, Nathaniel Trask, and Panos Stinis. Machine learning structure preserving brackets for forecasting irreversible processes. *Advances in Neural Information Processing Systems*, 34:5696–5707, 2021.

[20] US Agarwal, Rohit Bhargava, and RA Mashelkar. Brownian dynamics simulation of a polymer molecule in solution under elongational flow. *The Journal of chemical physics*, 108(4):1610–1617, 1998.

[21] Anatolii A Mogul'skii. Large deviations for trajectories of multi-dimensional random walks. *Theory of Probability & Its Applications*, 21(2):300–315, 1977.

[22] SR Srinivasa Varadhan. *Large deviations*, volume 27. American Mathematical Soc., 2016.

[23] Arsen Akopyan and Vladislav Vysotsky. Large deviations of convex hulls of planar random walks and Brownian motions. *Annales Henri Lebesgue*, 4:1163–1201, 2021. doi: 10.5802/ahl.100.

[24] Beatrice W Soh, Zi-En Ooi, Eleonore Vissol-Gaudin, Chang Jie Leong, and Kedar Hippalgaonkar. Automated electrokinetic stretcher for manipulating nanomaterials. *Lab on a Chip*, 23(16): 3716–3726, 2023.

[25] Wen Huang, Min Ji, Zhenxin Liu, and Yingfei Yi. Steady states of fokker–planck equations: I. existence. *Journal of Dynamics and Differential Equations*, 27(3):721–742, 2015.

[26] Grigorios A Pavliotis. Stochastic processes and applications. *Texts in applied mathematics*, 60:41–43, 2014.

[27] Felix Dietrich, Alexei Makeev, George Kevrekidis, Nikolaos Evangelou, Tom Bertalan, Sebastian Reich, and Ioannis G Kevrekidis. Learning effective stochastic differential equations from microscopic simulations: Linking stochastic numerics to deep learning. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 33(2), 2023.

[28] Aiqing Zhu and Qianxiao Li. DynGMA: A robust approach for learning stochastic differential equations from data. *Journal of Computational Physics*, 513:113200, 2024.

[29] Alexander Vologodskii. Brownian dynamics simulation of knot diffusion along a stretched dna molecule. *Biophysical journal*, 90(5):1594–1597, 2006.

[30] Tony W Liu. Flexible polymer chain dynamics and rheological properties in steady flows. *The Journal of chemical physics*, 90(10):5826–5842, 1989.

[31] Madan Somasi, Bamin Khomami, Nathanael J Woo, Joe S Hur, and Eric SG Shaqfeh. Brownian dynamics simulations of bead-rod and bead-spring chains: numerical algorithms and coarse-graining issues. *Journal of non-newtonian fluid mechanics*, 108(1-3):227–255, 2002.

[32] Kang Feng and Zaijiu Shang. Volume-preserving algorithms for source-free dynamical systems. *Numerische Mathematik*, 71(4):451–463, 1995.

[33] Kurt Hornik, Maxwell Stinchcombe, and Halbert White. Universal approximation of an unknown mapping and its derivatives using multilayer feedforward networks. *Neural networks*, 3(5):551–560, 1990.

[34] Milton Abramowitz. Handbook of mathematical functions, national bureau of standards. *Applied Mathematics Series*, (55), 1964.

Supplemental Material for Hidden Phase Transition in Single-Polymer Stretching under Elongational Flow

S1 Microscopic dynamics and reduced coordinates

S1.A Bead-rod chain and its numerical integration

We generate the microscopic trajectories using a Brownian dynamics model of a linear chain of $N+1=300$ touching beads in three dimensions [17]. Each bead has diameter $b = 10$nm, and adjacent beads are connected by rigid rods of the same length:

$$|\mathbf{r}_{i+1} - \mathbf{r}_i| = b, \qquad i = 0, \dots, N-1. \tag{S1}$$

where the beads are indexed by $i = 0, \dots, N$, $\mathbf{r}_i(t)$ is the position of bead i and $\mathbf{r}_{\rm cm}(t) = (N+1)^{-1}\sum_{i=0}^{N}\mathbf{r}_i(t)$ is the instantaneous center of mass. The imposed planar elongational flow is evaluated relative to the center of mass:

$$\mathbf{u}_i(t) = \boldsymbol{\kappa}[\mathbf{r}_i(t) - \mathbf{r}_{\rm cm}(t)], \qquad \boldsymbol{\kappa} = \dot{\epsilon}\begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

The stagnation point follows the instantaneous center of mass, with extension along the x-axis, compression along the y-axis and no imposed velocity in the z-axis.

Let $\zeta = 3\pi\eta b$ denote the Stokes drag coefficient of a single bead in a solvent of viscosity η. We neglect hydrodynamic interactions between beads, such that the solvent acts only through the imposed flow and the bead-wise drag coefficient. Neglecting inertia, the force balance is

$$\zeta\dot{\mathbf{r}}_i = \zeta\mathbf{u}_i + \mathbf{F}_i^{\rm ev} + \mathbf{F}_i^{\rm c} + \mathbf{F}_i^{\rm B},$$

where $\mathbf{F}_i^{\rm ev}$ is the excluded volume force, $\mathbf{F}_i^{\rm c}$ enforces Eq. (S1) and $\mathbf{F}_i^{\rm B}$ is the random Brownian force. With $r_{ij} = |\mathbf{r}_i - \mathbf{r}_j|$, the short-range contact repulsion is given by

$$\mathbf{F}_i^{\rm ev} = \mu \sum_{j\neq i,\ r_{ij}<b} [\mathbf{r}_i - \mathbf{r}_j]/r_{ij},$$

where $\mu = 35$ pN is chosen to suppress chain crossings [29]. For $r_{ij} \neq b$, this force follows from the continuous pair potential $E_{\rm ev} = \mu\sum_{i<j}(b - r_{ij})_+$, where $(a)_+ = \max(a, 0)$. Defining the bond unit vector $\widehat{\mathbf{b}}_i = (\mathbf{r}_{i+1} - \mathbf{r}_i)/b$ and setting $T_{-1} = T_N = 0$ (with the boundary terms omitted), the constraint force on bead i is

$$\mathbf{F}_i^{\rm c} = T_i\widehat{\mathbf{b}}_i - T_{i-1}\widehat{\mathbf{b}}_{i-1}.$$

The tensions T_i enforce the fixed length constraints. The Brownian force $\mathbf{F}_i^{\rm B}$ has zero mean and satisfies the fluctuation-dissipation relation

$$\langle\mathbf{F}_i^{\rm B}(t)\rangle = 0, \quad \langle\mathbf{F}_i^{\rm B}(t)\mathbf{F}_j^{\rm B}(t')^{\top}\rangle = 2\zeta k_{\rm B}T\delta_{ij}\mathbf{I}\delta(t-t').$$

We integrate the force balance equation with a predictor-corrector scheme [30] and solve the nonlinear bond-tension equations by Newton's method at each step [31]. We use a time step $\Delta t = 5\times10^{-4}\tau_d$, where $\tau_d = b^2\zeta/(k_{\rm B}T)$; the dimensionless contact repulsion strength is $\mu b/(k_{\rm B}T) = 85.0$. Each chain is equilibrated for $10^4\tau_d$ (without any flow) before turning on the flow. A single-exponential fit to the final 30% extension during relaxation of an initially stretched chain yields the longest relaxation time $\tau = 4321\tau_d$, defining the Weissenberg number $\mathrm{Wi} = \dot{\epsilon}\tau$. In simulation units, $\widetilde{\mathbf{r}} = \mathbf{r}/b$, $\widetilde{t} = t/\tau_d$, and $\widetilde{\mathbf{F}} = b\mathbf{F}/(k_{\rm B}T)$, such that

$$\widetilde{\dot{\epsilon}} = \dot{\epsilon}\tau_d, \qquad \mathrm{Wi} = \widetilde{\dot{\epsilon}}\frac{\tau}{\tau_d} = 4321\widetilde{\dot{\epsilon}}.$$

S1.B Symmetry-adapted dimensionality reduction

We represent each conformation by the stacked bead positions $\mathbf{r} = (\mathbf{r}_0^\top, \ldots, \mathbf{r}_N^\top)^\top \in \mathbb{R}^{900}$, retaining the instantaneous center-of-mass position. The first reduced coordinate is the normalized chain extension,

$$Z_1 = \frac{\max_i r_{i,x} - \min_i r_{i,x}}{(N+1)b}.$$

To construct the two additional coordinates, we sample K training conformations across chain-extension bins within each batch, supplemented by random draws without replacement.

We constrain the projection directions by chain reversal R and coordinate reflections T_x, T_y, T_z. These operations commute and each squares to the identity. Their 16 combinations form the symmetry group $G = \{R^{a_R} T_x^{a_x} T_y^{a_y} T_z^{a_z} : a_R, a_x, a_y, a_z \in \{0,1\}\}$. For each $g \in G$, an orthogonal matrix $U_g \in \mathbb{R}^{900\times 900}$ represents its action on $\mathbf{r}$. Specifically, $(U_R\mathbf{r})_i = \mathbf{r}_{N-i}$, where i labels a three-component bead block, and U_{T_α} is diagonal with -1 on the α-coordinate entries of all beads and $+1$ elsewhere ($\alpha = x, y, z$). Composed operations satisfy $U_{gh} = U_g U_h$ for $g, h \in G$. A one-dimensional character $\chi : G \to \{+1, -1\}$ satisfies $\chi(gh) = \chi(g)\chi(h)$ and is labeled by the generator signs $(\chi(R), \chi(T_x), \chi(T_y), \chi(T_z))$. Each character defines a symmetry subspace of projection vectors $\mathbf{v} \in \mathbb{R}^{900}$ satisfying $U_g\mathbf{v} = \chi(g)\mathbf{v}$ for all $g \in G$, where $\chi(g) = +1$ or -1 indicates that g preserves or reverses $\mathbf{v}$, respectively. The orthogonal projector onto this subspace is

$$P_\chi = \frac{1}{16}\sum_{g\in G} \chi(g) U_g.$$

We select $\chi_2 = (-1, -1, +1, +1)$ and $\chi_3 = (+1, -1, +1, +1)$ for the two PCA directions.

The group-averaged sample mean vanishes under the coordinate reflections. Accordingly, for the sampled conformations $\mathbf{r}^{(m)}$, $m = 1, \ldots, K$, we form the second-moment matrices

$$C_0 = \frac{1}{K-1}\sum_{m=1}^{K} \mathbf{r}^{(m)}\mathbf{r}^{(m)\top}, \qquad C = \frac{1}{16}\sum_{g\in G} U_g C_0 U_g^\top.$$

This enforces a zero mean by symmetry without removing the center of mass from individual conformations. Within each subspace, the unit vector $\mathbf{v}_j$ maximizing $\mathbf{v}^\top C\mathbf{v}$ is obtained as the normalized leading eigenvector of $P_{\chi_j} C P_{\chi_j}$. For the projections $a_j^{(m)} = \mathbf{v}_j^\top \mathbf{r}^{(m)}$ on the original fitting samples, define

$$\bar{a}_j = \frac{1}{K}\sum_m a_j^{(m)}, \quad s_j^2 = \frac{1}{K}\sum_m (a_j^{(m)} - \bar{a}_j)^2, \quad Z_j = \frac{\mathbf{v}_j^\top \mathbf{r}}{s_j}, \quad j = 2, 3.$$

Thus, each coordinate is scaled by its sample standard deviation without mean subtraction. The resulting map $\mathbf{r} \mapsto \mathbf{Z} = (Z_1, Z_2, Z_3)$, including direction signs and scales, is fixed across all flow strengths and training and test trajectories.

S2 Conditional UIDD: formulation, learning, and landscape analysis

S2.A Expressivity and identifiability

We next establish the expressivity and identifiability of conditional UIDD. Specifically, we show that any diffusion family satisfying the assumptions below admits a conditional UIDD representation whose potential is unique up to a c-dependent additive constant. Consider a family of Itô SDEs on $\mathbb{R}^d$ ($d \geq 1$) indexed by a condition $c \in \mathcal{U} \subset \mathbb{R}^p$:

$$\mathrm{d}\mathbf{Z}_t = b(\mathbf{Z}_t; c)\mathrm{d}t + \sigma(\mathbf{Z}_t; c)\mathrm{d}\mathbf{B}_t. \tag{S2}$$

Here $\mathbf{B}_t$ is a standard d-dimensional Brownian motion, $b(\mathbf{Z}; c) \in \mathbb{R}^d$ is the drift, and $\sigma(\mathbf{Z}; c) \in \mathbb{R}^{d\times d}$ is the noise coefficient. The condition domain $\mathcal{U}$ is an open set containing the conditions $\mathcal{C}$, and c is fixed along each trajectory. We assume a positive-definite diffusion matrix $M = \sigma\sigma^\top/2$. We write $\mathbf{e}_i$ for the standard basis vectors of $\mathbb{R}^d$.

Assumption S1. *For each fixed $c \in \mathcal{U}$, the coefficients $b(\cdot; c) \in C^1$ and $\sigma(\cdot; c) \in C^2$ of Eq. (S2) satisfy global linear growth and Lipschitz bounds in $\mathbf{Z}$ (with c-dependent constants), and the process admits a unique invariant probability density $\pi(\cdot; c) \in C^2 \cap L^1$.*

Assumption S2. *For all $i, j = 1, \dots, d$, assume*

$$\begin{aligned} b,\ \partial_{Z_i} b &\in C^0(\mathbb{R}^d \times \mathcal{U}), \\ M,\ \partial_{Z_i} M,\ \partial_{Z_i}\partial_{Z_j} M &\in C^0(\mathbb{R}^d \times \mathcal{U}), \\ \pi,\ \partial_{Z_i}\pi,\ \partial_{Z_i}\partial_{Z_j}\pi &\in C^0(\mathbb{R}^d \times \mathcal{U}). \end{aligned}$$

Here continuity is joint in $(\mathbf{Z}, c)$ and is understood componentwise for vector and matrix fields.

For Eq. (S2), the stationary Fokker-Planck equation requires a divergence-free probability current:

$$\mathbf{J} = b\pi - \nabla_{\mathbf{Z}} \cdot (M\pi), \qquad \nabla_{\mathbf{Z}} \cdot \mathbf{J} = 0. \tag{S3}$$

Here $\mathbf{J}(\cdot; c) \in C^1$ for each c. Under Assumption S2, $\mathbf{J}$ and its first $\mathbf{Z}$-derivatives are jointly continuous in $(\mathbf{Z}, c)$.

Lemma S1. *Let $d \geq 2$ and $\mathbf{J} : \mathbb{R}^d \times \mathcal{U} \to \mathbb{R}^d$ satisfy $\nabla_{\mathbf{Z}} \cdot \mathbf{J} = 0$ and $\mathbf{J}, \partial_{Z_i}\mathbf{J} \in C^0(\mathbb{R}^d \times \mathcal{U})$ for $i = 1, \dots, d$. There exists a skew-symmetric matrix field $\Psi : \mathbb{R}^d \times \mathcal{U} \to \mathbb{R}^{d\times d}$, with $\Psi_{ij} = 0$ for $|i - j| > 1$, such that $\nabla_{\mathbf{Z}} \cdot \Psi = \mathbf{J}$ and $\Psi, \partial_{Z_i}\Psi_{i,i+1}, \partial_{Z_{i+1}}\Psi_{i,i+1} \in C^0(\mathbb{R}^d \times \mathcal{U})$ for $i = 1, \dots, d-1$.*

Proof. We adapt the coordinate-integration construction in Ref. [32], using the reference hyperplanes $Z_j = 0$ for every c. For a scalar field $f = f(\mathbf{Z}; c)$, define

$$(I_j f)(\mathbf{Z}; c) = \int_0^{Z_j} f(Z_1, \dots, s, \dots, Z_d; c)\mathrm{d}s, \quad (E_j f)(\mathbf{Z}; c) = f(Z_1, \dots, 0, \dots, Z_d; c),$$

where s or 0 replaces the jth coordinate of $\mathbf{Z}$ at fixed c. Both operators preserve joint continuity in $(\mathbf{Z}, c)$, since E_j is a coordinate evaluation and

$$(I_j f)(\mathbf{Z}; c) = Z_j \int_0^1 f(Z_1, \dots, uZ_j, \dots, Z_d; c)\mathrm{d}u$$

is continuous in $(\mathbf{Z}, c)$ by uniform continuity on compact sets. Define $S_i(\mathbf{Z}; c) = \sum_{k=1}^{i} \partial_{Z_k} J_k(\mathbf{Z}; c)$ and $S_0 = 0$. For $d = 2$, set

$$A_1(\mathbf{Z}; c) = (I_2 J_1 - I_1 E_2 J_2)(\mathbf{Z}; c).$$

At each c, the divergence-free condition $\nabla_{\mathbf{Z}} \cdot \mathbf{J} = 0$ implies $\partial_{Z_2} A_1(\mathbf{Z}; c) = J_1(\mathbf{Z}; c)$ and $\partial_{Z_1} A_1(\mathbf{Z}; c) = -J_2(\mathbf{Z}; c)$. For $d \geq 3$, we take

$$\begin{aligned} A_1(\mathbf{Z}; c) &= (I_2 J_1)(\mathbf{Z}; c), \\ A_i(\mathbf{Z}; c) &= (I_{i+1} J_i + I_{i+1} I_i S_{i-1})(\mathbf{Z}; c), \quad 2 \leq i \leq d-2, \\ A_{d-1}(\mathbf{Z}; c) &= (I_d E_{d-1} J_{d-1} - I_{d-1} J_d)(\mathbf{Z}; c). \end{aligned}$$

Differentiating at fixed c and using $\partial_{Z_i}(I_i S_{i-1}) = S_{i-1}$, we obtain that, for $1 \leq i \leq d-2$,

$$\begin{aligned} \partial_{Z_i} A_i(\mathbf{Z}; c) &= [I_{i+1}(\partial_{Z_i} J_i + S_{i-1})](\mathbf{Z}; c) = (I_{i+1} S_i)(\mathbf{Z}; c), \\ \partial_{Z_{i+1}} A_i(\mathbf{Z}; c) &= J_i(\mathbf{Z}; c) + (I_i S_{i-1})(\mathbf{Z}; c). \end{aligned}$$

For the last entry, $\nabla_{\mathbf{Z}} \cdot \mathbf{J}(\mathbf{Z}; c) = 0$ yields

$$\partial_{Z_{d-1}} A_{d-1}(\mathbf{Z}; c) = -J_d(\mathbf{Z}; c), \quad \partial_{Z_d} A_{d-1}(\mathbf{Z}; c) = J_{d-1}(\mathbf{Z}; c) + (I_{d-1} S_{d-2})(\mathbf{Z}; c).$$

Set $\Psi_{i,i+1}(\mathbf{Z}; c) = A_i(\mathbf{Z}; c) = -\Psi_{i+1,i}(\mathbf{Z}; c)$ and all other entries to zero. Then

$$\begin{aligned} (\nabla_{\mathbf{Z}} \cdot \Psi)_1(\mathbf{Z}; c) &= J_1(\mathbf{Z}; c), \\ (\nabla_{\mathbf{Z}} \cdot \Psi)_i(\mathbf{Z}; c) &= -\partial_{Z_{i-1}} A_{i-1}(\mathbf{Z}; c) + \partial_{Z_{i+1}} A_i(\mathbf{Z}; c) = J_i(\mathbf{Z}; c), \quad 2 \leq i \leq d-1, \\ (\nabla_{\mathbf{Z}} \cdot \Psi)_d(\mathbf{Z}; c) &= -\partial_{Z_{d-1}} A_{d-1}(\mathbf{Z}; c) = J_d(\mathbf{Z}; c). \end{aligned}$$

Since J_i and S_i are jointly continuous in $(\mathbf{Z}, c)$ and I_j, E_j preserve continuity, Ψ and its stated derivatives are jointly continuous. The construction requires no c-derivatives and needs only $\mathbf{J}(\cdot; c) \in C^1$ at each fixed c. For $d = 2, 3$, all first $\mathbf{Z}$-derivatives of Ψ are jointly continuous; for $d \geq 4$, this holds if all second $\mathbf{Z}$-derivatives of $\mathbf{J}$ are jointly continuous. □

Theorem S1 (Expressivity and identifiability)**.** *For the family of SDEs in Eq. (S2) satisfying Assumption S1, there exist a potential $V(\cdot;c) \in C^2(\mathbb{R}^d)$ and an antisymmetric matrix $W(\mathbf{Z};c)$, with $W_{ij} = 0$ for $|i-j| > 1$, such that $e^{-V(\cdot;c)} \in L^1(\mathbb{R}^d)$ and*

$$b(\mathbf{Z};c) = -[M(\mathbf{Z};c) + W(\mathbf{Z};c)]\nabla_{\mathbf{Z}} V(\mathbf{Z};c) + \nabla_{\mathbf{Z}} \cdot M(\mathbf{Z};c) + \nabla_{\mathbf{Z}} \cdot W(\mathbf{Z};c) \tag{S4}$$

for every $(\mathbf{Z},c) \in \mathbb{R}^d \times \mathcal{U}$, with $W = 0$ for $d = 1$. For each fixed c, the entries $H_i = W_{i,i+1}$ and their derivatives $\partial_{Z_i} H_i$, $\partial_{Z_{i+1}} H_i$ are continuous in $\mathbf{Z}$. The potential can be chosen as $V = -\ln \pi$. Among representations with continuous potentials whose normalized exponentials are invariant for Eq. (S2), it is unique up to an additive function $C(c)$ independent of $\mathbf{Z}$.

If Assumption S2 also holds, V (up to second $\mathbf{Z}$-derivatives) and $H_i, \partial_{Z_i} H_i, \partial_{Z_{i+1}} H_i$ can be chosen jointly continuous in $(\mathbf{Z},c)$.

Proof. We follow the UIDD decomposition [16], first fixing c. On any bounded ball, the stationary Fokker-Planck operator is uniformly elliptic. Bounding its zeroth-order coefficient below by a nonpositive constant makes $\pi \geq 0$ a supersolution, so the strong minimum principle and normalization imply $\pi > 0$ on $\mathbb{R}^d$. Using the stationary current in Eq. (S3), decompose the drift as

$$b = \frac{\nabla_{\mathbf{Z}} \cdot (M\pi)}{\pi} + \frac{\mathbf{J}}{\pi}, \qquad \nabla_{\mathbf{Z}} \cdot \mathbf{J} = 0.$$

For $d \geq 2$, the construction in Lemma S1 yields a banded antisymmetric matrix Ψ with $\nabla_{\mathbf{Z}} \cdot \Psi = \mathbf{J}$. Define

$$V(\mathbf{Z};c) = -\ln \pi(\mathbf{Z};c), \qquad W(\mathbf{Z};c) = \frac{\Psi(\mathbf{Z};c)}{\pi(\mathbf{Z};c)}.$$

The matrix W has the same banded antisymmetric structure as Ψ. Since $\nabla_{\mathbf{Z}} \pi = -\pi \nabla_{\mathbf{Z}} V$, the two terms in the drift become

$$\frac{\nabla_{\mathbf{Z}} \cdot (M\pi)}{\pi} = -M\nabla_{\mathbf{Z}} V + \nabla_{\mathbf{Z}} \cdot M, \quad \frac{\mathbf{J}}{\pi} = \frac{\nabla_{\mathbf{Z}} \cdot (W\pi)}{\pi} = -W\nabla_{\mathbf{Z}} V + \nabla_{\mathbf{Z}} \cdot W.$$

Adding these identities yields Eq. (S4), and $e^{-V} = \pi$ is normalized and invariant.

Under Assumption S2, $\mathbf{J}$ and its first $\mathbf{Z}$-derivatives are jointly continuous, so Lemma S1 yields a $\Psi(\mathbf{Z};c)$ with the stated joint continuity. Dividing by $\pi > 0$ and applying the chain rule gives the claimed regularity of W and V without differentiating in c.

Finally, if $\widetilde{V}$ is another continuous potential whose normalized exponential is invariant, uniqueness of the continuous invariant density implies

$$\frac{e^{-\widetilde{V}(\mathbf{Z};c)}}{\widetilde{\mathcal{Z}}(c)} = \pi(\mathbf{Z};c), \quad \widetilde{\mathcal{Z}}(c) = \int_{\mathbb{R}^d} e^{-\widetilde{V}(\mathbf{Z};c)} \mathrm{d}\mathbf{Z}.$$

Thus $\widetilde{V}(\mathbf{Z};c) = -\ln \pi(\mathbf{Z};c) + C(c)$ with $C(c) = -\ln \widetilde{\mathcal{Z}}(c)$, proving identifiability. □

S2.B Neural parameterization

We extend the UIDD network modules [16] by supplying the joint input $(\mathbf{Z},c)$ and sharing all parameters θ across conditions:

$$\sigma_\theta(\mathbf{Z};c) = \sqrt{2}\,[\alpha_\sigma T_\theta(\mathbf{Z};c) + \operatorname{diag}\phi(s_\theta(\mathbf{Z};c))]\,,$$

$$W_\theta(\mathbf{Z};c) = \begin{pmatrix} 0 & H_{\theta,1}(\mathbf{Z};c) & 0 & \cdots & 0 & 0 \\ -H_{\theta,1}(\mathbf{Z};c) & 0 & H_{\theta,2}(\mathbf{Z};c) & \cdots & 0 & 0 \\ 0 & -H_{\theta,2}(\mathbf{Z};c) & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 0 & H_{\theta,d-1}(\mathbf{Z};c) \\ 0 & 0 & 0 & \cdots & -H_{\theta,d-1}(\mathbf{Z};c) & 0 \end{pmatrix},$$

where $T_\theta \in \mathbb{R}^{d\times d}$ is strictly lower triangular, $s_\theta \in \mathbb{R}^d$, $H_\theta \in \mathbb{R}^{d-1}$, and $\phi(s) = (\sqrt{1+s^2}+s)/2$ is applied componentwise. Thus $M_\theta = \sigma_\theta \sigma_\theta^\top/2 = L_\theta L_\theta^\top$, where $L_\theta = \alpha_\sigma T_\theta + \operatorname{diag}\phi(s_\theta)$ has positive diagonal. The off-diagonal scale factor is fixed at $\alpha_\sigma = 0.1$. The potential is

$$V_\theta(\mathbf{Z};c) = \frac{\beta_1}{2}\|U_\theta(\mathbf{Z};c) + \beta_2 \Gamma(\mathbf{Z}^\top, c^\top)^\top\|^2 + \beta_3\|\mathbf{Z}\|^2,$$

with $U_\theta \in \mathbb{R}^m$, $m \geq 1$, trainable $\Gamma \in \mathbb{R}^{m\times(d+p)}$, and fixed $\beta_1, \beta_2, \beta_3 > 0$. Thus M_θ is positive definite, W_θ is banded and antisymmetric, and $e^{-V_\theta} \leq e^{-\beta_3\|\mathbf{Z}\|^2}$ is integrable.

To establish expressivity, consider tanh networks. Under Assumptions S1 and S2, let V, W be the fields in Theorem S1 and let $Q = K \times \mathcal{C}_0$, where $K \subset \mathbb{R}^d$ and $\mathcal{C}_0 \subset \mathcal{U}$ are nonempty compact sets. For every $\epsilon > 0$, there exist shared parameters θ and a constant C_Q, independent of c, such that

$$\sup_{(\mathbf{Z},c)\in Q} [|V_\theta - V - C_Q| + \|f_\theta - b\| + \|M_\theta - M\|] < \epsilon, \tag{S5}$$

where f_θ is the drift in Eq. (1) assembled from the neural fields.

To see this, let $M = LL^\top$ be the Cholesky factorization, with L lower triangular and $L_{ii} > 0$. We approximate $10L_{ij}$ for $i > j$ and $\phi^{-1}(L_{ii})$ by the corresponding components of T_θ and s_θ, respectively. For the potential, we choose C_Q such that $V + C_Q - \beta_3\|\mathbf{Z}\|^2 > 0$ on a neighborhood of Q, set $\Gamma = 0$, and approximate

$$q(\mathbf{Z}; c) = \sqrt{\frac{2}{\beta_1}\left[V(\mathbf{Z}; c) + C_Q - \beta_3\|\mathbf{Z}\|^2\right]}$$

by one component of U_θ, with the remaining components set to zero. We approximate these targets together with their required $\mathbf{Z}$-derivatives (in particular, $H_i = W_{i,i+1}$ along with $\partial_{Z_i} H_i$ and $\partial_{Z_{i+1}} H_i$). By Assumption S2 and Lemma S1, all targets and required derivatives are jointly continuous in $(\mathbf{Z}, c)$. Multiplying by a smooth cutoff equal to one near Q and mollifying in $(\mathbf{Z}, c)$ yields smooth functions that converge uniformly on Q together with these derivatives. Simultaneous neural approximation of functions and derivatives [33], combined with the product and chain rules, then yields Eq. (S5).

Polymer implementation. We symmetrize the potential as

$$V_\theta^{\mathrm{sym}}(\mathbf{Z}; c) = \frac{1}{4}\sum_{g\in G_Z} V_\theta(g\mathbf{Z}; c), \qquad G_Z = \{\mathrm{diag}(1, \pm 1, \pm 1)\}.$$

The generators $\mathrm{diag}(1, -1, 1)$ and $\mathrm{diag}(1, -1, -1)$ are the reduced actions of R and T_x, respectively (Sec. S1.B). Thus the potential is even in Z_2 and Z_3 and retains the Gaussian bound; σ_θ and W_θ are not symmetrized. For a G_Z-invariant target potential and a G_Z-invariant K, averaging does not increase the uniform errors in the potential and its state gradient. The preceding approximation argument therefore also applies to this symmetry-restricted class.

Network hyperparameters are summarized in the table below.

Output	Layer widths	Hidden activation
U_θ	$4 \to 128 \to 32$	r
s_θ	$4 \to 32 \to 32 \to 3$	r
T_θ (before triangular projection)	$4 \to 32 \to 32 \to 9$	r
H_θ	$4 \to 32 \to 32 \to 2$	tanh

Here $r(x) = \max(0, x)^2 - \max(0, x - \frac{1}{2})^2$. For T_θ, the nine outputs are reshaped into a 3×3 matrix before retaining its strictly lower-triangular part. Here Γ is a bias-free linear layer with $m = 32$, and $(\beta_1, \beta_2, \beta_3) = (100, 1, 0.01)$.

S2.C Training and testing

We fit the conditional model to 11145 trajectories by minimizing the loss in Eq. (7). Of these, 4945 trajectories start from a pool of coiled conformations and another 4945 from a single folded conformation. Both ensembles span $\mathrm{Wi} \in (0.005, 10.8]$ with closely matched condition distributions. An additional 1255 trajectories sample $\mathrm{Wi} \in (0, 1)$ from the same pool of coiled conformations. The condition is held fixed along each trajectory.

Training uses adjacent transitions from 500-frame segments with $\Delta t_{\mathrm{train}} = 5 \times 10^{-4}$. Minibatches contain at most 256 segments and are processed in a fixed order. For each minibatch, a single transformation is drawn from the identity, R, and T_x with probabilities 1/2, 1/4, and 1/4, respectively. The model is trained for 300 epochs with Adam, starting from a learning rate of 10^{-3}, which is reduced when the loss plateaus. We retain the parameters from the final epoch and train five independent models from different random initializations.

We evaluate transient stretching from three initial microscopic conformations (labeled fast, medium, and slow) excluded from both PCA fitting and model training. For each conformation and each of ten test Wi values, we compare 500 microscopic trajectories against 500 reduced trajectories simulated from the corresponding encoded

initial state. To quantify unfolding times, let $Z_1^{(j)}(t)$ denote the normalized span of trajectory j, and define the unfolding step $n_{\mathrm{u}}^{(j)}$ as

$$n_{\mathrm{u}}^{(j)} = \min\left\{n \in \{0,\dots,n_{\max}\} : Z_1^{(j)}(n\Delta t_{\mathrm{test}}) > 0.95 Z_1^{(j)}(t_{\max})\right\}, \tag{S6}$$

where $n_{\max} = t_{\max}/\Delta t_{\mathrm{test}}$. This threshold is defined relative to each trajectory's final extension and is recorded at discrete output steps without interpolation ($n_{\mathrm{u}}^{(j)} = 0$ if exceeded initially). Figure S1 shows that the conditional UIDD model accurately reproduces the ensemble mean and standard deviation of $Z_1(t)$ as well as the normalized histograms of n_{u} across all three initial conformations and flow strengths.

S2.D Landscape evaluation and transition estimates

We evaluate the marginal integrals using uniform-grid Riemann sums (via log-sum-exp) with equal grid-spacing weights on the following grids:

Profile	Integration grids (nodes)	Tabulation grid (nodes; spacing)
	$Z_1 \in [-0.5, 2.5]$ (512),	
F_2	$Z_3 \in [-15, 15]$ (769)	$Z_2 \in [-3, 3]$ (601; 0.01)
$F_{23}(0,\cdot)$	$Z_1 \in [-0.5, 2.5]$ (512)	$Z_3 \in [-15, 15]$ (2401; 0.0125)

All profiles and insets use the same 512-node quadrature over $Z_1 \in [-0.5, 2.5]$.

For $k = 1,\dots,5$, let $\mathcal{F}_k(q;\mathrm{Wi})$ denote either $F_2^{(k)}(q;\mathrm{Wi})$ or $F_{23}^{(k)}(0,q;\mathrm{Wi})$, and let $\mathcal{Q}$ be the corresponding tabulation grid, which includes zero. For $\eta > 0$, define

$$\mathcal{A}_{\eta,k}(\mathrm{Wi}) = \left\{q \in \mathcal{Q} : \mathcal{F}_k(q;\mathrm{Wi}) - \min_{q'\in\mathcal{Q}} \mathcal{F}_k(q';\mathrm{Wi}) \le \eta\right\},$$
$$D_k(\mathrm{Wi}) = \mathcal{F}_k(0;\mathrm{Wi}) - \min_{q\in\mathcal{Q}} \mathcal{F}_k(q;\mathrm{Wi}).$$

The condition $0 \notin \mathcal{A}_{\eta,k}$ is equivalent to $D_k > \eta$. Using $\eta = 0.6$ for F_2 and $\eta = 1.0$ for $F_{23}(0,\cdot)$ [Fig. 2(i,l)], we estimate each model's transition point $\mathrm{Wi}_{t,k}$ (denoted Wi_t in the main text) by linearly interpolating $D_k(\mathrm{Wi}) - \eta$ across its first upward zero crossing. The F_2 scan contains 146 values of Wi: spacing 0.02 on $[0.02, 1]$, refined to 0.005 on $[0.15, 0.55]$; spacing 0.05 on $[1.05, 2]$; and 15 logarithmically spaced values on $[2.2, 10]$. The $F_{23}(0,\cdot)$ scan uses 439 values: spacing 0.01 on $[0.02, 3]$ and 0.05 on $[3.05, 10]$. At finite η, $\mathrm{Wi}_{t,k}$ marks where the central barrier reaches η, rather than the exact onset of peak splitting.

The tolerance-sensitivity panels in Fig. 2(j,m) show the individual model estimates, their mean, and error bars of one sample standard deviation over five initializations, with

$$F_2: \quad \eta \in \{0.02, 0.05, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.8, 1.0, 1.2, 1.5, 2.0\},$$
$$F_{23}(0,\cdot): \quad \eta \in \{0.05, 0.1, 0.2, 0.3, 0.4, 0.6, 0.8, 1.0, 1.2, 1.5, 2.0, 3.0\}.$$

The spread across models reflects initialization sensitivity on the fixed training set. For visibility in Fig. 2(i,l), near-minimizing regions are plotted with minimum widths of four Z_2 and nine Z_3 grid cells; all $\mathrm{Wi}_{t,k}$ estimates use the unbroadened sets.

The numerical quadratures are stable under grid refinement. Increasing the F_2 quadrature to 1023 nodes in Z_1 and 1537 nodes in Z_3, and the section quadrature to 2048 nodes in Z_1, changes D_k by less than 10^{-3} at the tested conditions with $\mathrm{Wi} \le 3$ and by less than 4×10^{-2} at those up to $\mathrm{Wi} = 10$. All tabulated minima of $F_{23}(0, Z_3)$ satisfy $|Z_3| < 9$. To test domain truncation, we compare the integral I_{win} on the default window with I_{ref} on an expanded window ($Z_1 \in [-1.50, 3.50]$, $Z_3 \in [-20, 20]$) at matched spacings (3/511 in Z_1, 10/256 in Z_3). Across all five models, evaluating $1 - I_{\mathrm{win}}/I_{\mathrm{ref}}$ at $Z_2 = 0$ and at the grid minimizer for five representative controls (the point nearest $\mathrm{Wi}_{t,k}^{(1)}$, namely $0.385, 0.455, 0.435, 0.370, 0.460$, plus $\mathrm{Wi} = 1, 2.73, 5.23, 10$) yields values below 10^{-12}, confirming negligible truncation error.

S3 Large-deviation analysis of the analytical model

We prove the limit interchange in Eq. (9) and characterize the minimizing set of the joint free energy. We use the potential, observables, and reference measure of the main-text analytical model, with $\mathcal{E}$ and $\mathcal{S}$ defined in Eq. (5), and ℓ in Eq. (8). For $\alpha = 2, 23$, write $d_\alpha = 1, 2$, respectively, and set

$$\hat{Q}_2 = \hat{z}_2, \quad \hat{Q}_{23} = (\hat{z}_2, \hat{z}_3), \qquad Q_2(m) = m_x(1), \quad Q_{23}(m) = (m_x(1), m_x(1) - 2\bar{m}_x).$$



Figure S1: Transient stretching from the fast, medium, and slow initial conformations (column groups) at different Wi (rows). Each group compares the mean and standard deviation of Z_1 and the distribution of n_u defined in Eq. (S6) obtained from microscopic simulations (blue) and the conditional UIDD model (orange), using 500 realizations per ensemble. Horizontal axes are given in output-step units.

Let $\mathcal{Z}_{\alpha,n}$ be the unnormalized Boltzmann-weighted density of $\hat{Q}_\alpha$, defined by the Dirac δ integrals in the main text. We use its continuous version for all sufficiently large n and define

$$W_{\alpha,n,\varepsilon}(q;\lambda) = \int e^{-U_{n,\lambda}} \mathbf{1}_{\{|\hat{Q}_\alpha - q|_\infty < \varepsilon\}} \mathrm{d}\nu_n, \quad A_{\alpha,n,\varepsilon}(q;\lambda) = \frac{W_{\alpha,n,\varepsilon}(q;\lambda)}{(2\varepsilon)^{d_\alpha}}.$$

Here $|\cdot|_\infty$ denotes the maximum norm, unsubscripted norms are Euclidean, and we set $\log 0 = -\infty$ and $\inf \varnothing = +\infty$. We now state our two main results.

Theorem S2 (Exchange of the two limits in Eq. (9))**.** *Fix $\lambda \geq 0$. For $\alpha = 2$, let $q = z_2 \in (-1,1)$; for $\alpha = 23$, let $q = (0, z_3)$ with $|z_3| < 1/2$. The pointwise limits $F_{\alpha,\infty}(q;\lambda) := \lim_{n\to\infty} F_{\alpha,n}(q;\lambda)$, where $F_{\alpha,n} = -n^{-1}\log \mathcal{Z}_{\alpha,n}$, exist and satisfy*

$$F_{\alpha,\infty}(q;\lambda) = -\lim_{n\to\infty}\lim_{\varepsilon\downarrow 0}\frac{1}{n}\log A_{\alpha,n,\varepsilon}(q;\lambda) = -\lim_{\varepsilon\downarrow 0}\lim_{n\to\infty}\frac{1}{n}\log A_{\alpha,n,\varepsilon}(q;\lambda) = \inf_{\substack{m\in\mathcal{A}\\ Q_\alpha(m)=q}} \{\lambda\mathcal{E}[m] - \mathcal{S}[m]\}.$$

Theorem S3 (The minimizing set is an interval)**.** *For every finite $\lambda \geq 0$, there is $a_\lambda \geq 0$ such that*

$$\operatorname*{argmin}_{c\in\mathbb{R}} F_{23,\infty}(0,c;\lambda) = [-a_\lambda, a_\lambda]. \tag{S7}$$

The interval opens at the exact threshold $\lambda_2 = 4\pi^2$. That is, $a_\lambda = 0$ $(0 \leq \lambda \leq 4\pi^2)$, $a_\lambda > 0$ $(\lambda > 4\pi^2)$.

The two-dimensional variational problem is able to be reduced to a scalar one. Let $\bar{u} = \int_0^1 u(s)\mathrm{d}s$, and take

$$\mathcal{A}_1 = \{u \in \mathrm{AC}[0,1] : u(0) = 0,\ |u'| \leq 1 \text{ a.e.}\}, \quad \mathcal{F}_\lambda^{(1)}[u] = \int_0^1 \ell(u')\mathrm{d}s - \lambda\int_0^1 (u-\bar{u})^2\mathrm{d}s.$$

Since ℓ is even, convex, and nondecreasing on $[0,1]$ and $\lambda \geq 0$, projecting $m = (u,v)$ onto $(u,0)$ preserves the constraints, drops the nonnegative transverse energy, and does not increase the entropy cost:

$$\lambda\mathcal{E}[(u,v)] - \mathcal{S}[(u,v)] \geq \lambda\mathcal{E}[(u,0)] - \mathcal{S}[(u,0)] = \mathcal{F}_\lambda^{(1)}[u].$$

Theorem S2 therefore yields the scalar variational formulas used in the proof of Theorem S3 and in the numerical evaluation:

$$F_{2,\infty}(z_2;\lambda) = \inf_{\substack{u\in\mathcal{A}_1\\ u(1)=z_2}} \mathcal{F}_\lambda^{(1)}[u], \quad F_{23,\infty}(0,z_3;\lambda) = \inf_{\substack{u\in\mathcal{A}_1\\ u(1)=0,\ \bar{u}=-z_3/2}} \mathcal{F}_\lambda^{(1)}[u]. \tag{S8}$$

S3.A Proof of the limit-interchange theorem

Window limit. We use the path large-deviation principle and uniform energy approximation from the End Matter. With $\bar{m}_{n,x} = \int_0^1 m_{n,x}(s)\mathrm{d}s$ denoting the continuum average of the piecewise-linear path m_n, trapezoidal integration yields that the observables satisfy $\hat{z}_2 = m_{n,x}(1)$ and

$$\hat{z}_3 = \frac{n}{n+1}\left[m_{n,x}(1) - 2\bar{m}_{n,x}\right] = \frac{1}{n}\sum_{i=1}^{n}\frac{2i-n-1}{n+1}\cos\theta_i.$$

Thus $\hat{Q}_{23} = Q_{23}(m_n) + O(n^{-1})$ uniformly, so windows of widths $\varepsilon - \eta$ and $\varepsilon + \eta$ give the constrained Laplace bounds for any fixed $0 < \eta < \varepsilon$.

For q as in Theorem S2, define

$$m_{\mathrm{ref}}(s) = \begin{cases} (z_2 s, 0), & \alpha = 2, \quad q = z_2, \\ (-2z_3 \min(s, 1-s), 0), & \alpha = 23, \quad q = (0, z_3). \end{cases}$$

Each path lies in $\mathcal{A}$ and satisfies $Q_\alpha(m_{\mathrm{ref}}) = q$. Its y-component is zero, so $\mathcal{E}[m_{\mathrm{ref}}] \leq 0$. The derivative is

$$m'_{\mathrm{ref}}(s) = \begin{cases} (z_2, 0), & \alpha = 2, \\ (-2z_3, 0), & \alpha = 23, \quad 0 < s < 1/2, \\ (2z_3, 0), & \alpha = 23, \quad 1/2 < s < 1. \end{cases}$$

Substituting into the entropy integral and using $\ell(-a) = \ell(a)$ gives

$$J(q) := -\mathcal{S}[m_{\mathrm{ref}}] = \int_0^1 \ell(|m'_{\mathrm{ref}}(s)|)\,\mathrm{d}s = \begin{cases} \ell(z_2), & \alpha = 2, \\ \ell(2z_3), & \alpha = 23. \end{cases}$$

These entropy costs are finite because $|z_2| < 1$ for $\alpha = 2$ and $2|z_3| < 1$ for $\alpha = 23$.

For any finite-entropy path m in the closed window $|Q_\alpha(m) - q|_\infty \le \varepsilon$, linearity of Q_α and $Q_\alpha(m_{\mathrm{ref}}) = q$ imply that $(1-\delta)m + \delta m_{\mathrm{ref}}$ lies in the open window $|Q_\alpha(\cdot) - q|_\infty < \varepsilon$ for every $\delta \in (0, 1]$. Convexity of $-\mathcal{S}$ and continuity of $\mathcal{E}$ ensure that its free energy approaches that of m as $\delta \downarrow 0$. The two window infima therefore agree. Letting $\eta \downarrow 0$, then $\varepsilon \downarrow 0$, compactness of $\mathcal{A}$ and lower semicontinuity of $\lambda\mathcal{E} - \mathcal{S}$ give

$$\begin{aligned} S_{\alpha,\varepsilon}(q) &:= \lim_{n\to\infty} \frac{1}{n} \log A_{\alpha,n,\varepsilon}(q;\lambda) = -\inf_{|Q_\alpha(m)-q|_\infty < \varepsilon} \{\lambda\mathcal{E}[m] - \mathcal{S}[m]\}, \\ S_{\alpha,\varepsilon}(q) &\to S_\alpha(q) := -\inf_{Q_\alpha(m)=q} \{\lambda\mathcal{E}[m] - \mathcal{S}[m]\} \quad (\varepsilon \downarrow 0). \end{aligned} \tag{S9}$$

All infima are over $\mathcal{A}$; the fixed volume factor $(2\varepsilon)^{d_\alpha}$ does not affect the exponential rate.

Continuous densities. Under $\nu_n = \prod_i \mathrm{d}\theta_i/(2\pi)$, set $H_n = -U_{n,\lambda}$ and $d = d_\alpha$. Write

$$T_n = \frac{1}{n}\sum_{i=1}^n w_i \cos\theta_i = \hat{Q}_\alpha, \qquad G = \frac{1}{n}\sum_{i=1}^n \sin^2\theta_i w_i w_i^T,$$
$$w_i = 1 \ (d=1), \qquad w_i = (1, c_i)^T, \quad c_i = \frac{2i-n-1}{n+1} \ (d=2).$$

The matrix G measures the sensitivity of the observables to bond rotations. For a continuous periodic weight Ω, write

$$f_\Omega(q) = \int \Omega(\theta)\,\delta(T_n(\theta) - q)\,\mathrm{d}\nu_n(\theta), \qquad f_n = f_{e^{H_n}}.$$

In ordinary integral terms, $\int_D f_\Omega(q)\,\mathrm{d}q$ is the total weight of configurations with $T_n(\theta) \in D$, for any box D in observable space. We write $\mathbb{P}(B) = \int \mathbf{1}_B\,\mathrm{d}\nu_n$ for the probability of an angular event B.

Lemma S2 (Smoothing by a few bonds)**.** *For all sufficiently large n, $q \mapsto f_\Omega(q)$ is continuous for every continuous periodic angular weight $\Omega(\theta)$. Moreover, there is a choice of J_n containing three indices for $d = 1$, or six for $d = 2$, such that, with $\theta_{\mathrm{out}} := (\theta_i)_{i\notin J_n}$,*

$$0 \le \Omega(\theta) \le K\mathbf{1}_B(\theta_{\mathrm{out}}) \quad \Longrightarrow \quad 0 \le f_\Omega(q) \le KCn^d\mathbb{P}(B) \quad (q \in \mathbb{R}^d).$$

Here $K > 0$, the event B depends only on θ_{out}, and C is independent of n, q, Ω, K, B.

Proof. Choose J_n as any three indices for $d = 1$, or the six indices nearest $nj/7$, $j = 1, \dots, 6$, for $d = 2$. With θ_{out} fixed, write

$$q_{\mathrm{out}} := \frac{1}{n}\sum_{i\notin J_n} w_i \cos\theta_i, \qquad T_n = q_{\mathrm{out}} + \frac{1}{n}\sum_{i\in J_n} w_i\cos\theta_i.$$

Let only θ_i, $i \in J_n$, be random, independently and uniformly distributed on $[0, 2\pi)$. Then T_n has probability density

$$g_{J_n}(q \mid \theta_{\mathrm{out}}) := \int_{[0,2\pi)^{|J_n|}} \delta\left(q_{\mathrm{out}} + \frac{1}{n}\sum_{i\in J_n} w_i\cos\theta_i - q\right) \prod_{i\in J_n} \frac{\mathrm{d}\theta_i}{2\pi}.$$

A single projection $X_i = \cos\theta_i$ has the arcsine law and its Fourier transform φ is the Bessel function of the first kind of order zero. The Bessel asymptotics give $|\varphi(t)| \le C(1+|t|)^{-1/2}$; the same decay holds with any polynomial amplitude $P(\cos\theta)$, by expansion in $\cos(k\theta)$ and the fixed-order Bessel asymptotics [34].

Let $Y := \sum_{i\in J_n} w_i X_i$. Independence gives its Fourier transform explicitly as

$$\mathbb{E}[e^{\mathrm{i}s\cdot Y}] = \prod_{i\in J_n} \varphi(s\cdot w_i).$$

For $d = 1$, this is $\varphi(s)^3$. For $d = 2$, the six selected coefficients satisfy $|c_i - c_j| \geq \delta_c > 0$ for $i \neq j$, with δ_c independent of n for large n. At most one index can satisfy $|s_1 + c_i s_2| < b|s|$ for a sufficiently small fixed $b > 0$. Indeed, if two indices $i \neq j$ satisfied this inequality, subtracting the two expressions and using $|c_i| \leq 1$ would imply

$$|s_2| < \frac{2b}{\delta_c}|s|, \quad |s_1| \leq |s_1 + c_i s_2| + |c_i||s_2| < \left(1 + \frac{2}{\delta_c}\right) b|s|,$$

which contradicts $|s|^2 = s_1^2 + s_2^2$. Thus at least five factors satisfy

$$|\varphi(s_1 + c_i s_2)| \leq C(1 + |s|)^{-1/2},$$

while any remaining factor is bounded by one. Consequently,

$$\left|\mathbb{E}[e^{\mathrm{i}s\cdot Y}]\right| \leq C \begin{cases} (1+|s|)^{-3/2}, & d = 1, \\ (1+|s|)^{-5/2}, & d = 2. \end{cases}$$

The first bound is integrable on $\mathbb{R}$; the second is integrable on $\mathbb{R}^2$. Fourier inversion therefore gives Y a continuous bounded density. Rescaling by $1/n$ and shifting by q_{out} shows that g_{J_n} is continuous and

$$0 \leq g_{J_n}(q \mid \theta_{\mathrm{out}}) \leq Cn^d \qquad \text{uniformly in } q, \theta_{\mathrm{out}}.$$

For $X = (X_i) \in [-1, 1]^n$, define the sign-averaged weight

$$\overline{\Omega}(X) := 2^{-n} \sum_{\varepsilon \in \{-1,1\}^n} \Omega(\varepsilon_1 \arccos X_1, \ldots, \varepsilon_n \arccos X_n),$$

where the angles are interpreted periodically. This is continuous on $[-1, 1]^n$. Since the changes of variables $\theta_i \mapsto -\theta_i$ preserve the uniform angular average and T_n, for every box D, we have

$$\int_{\{T_n \in D\}} \Omega(\theta)\, \mathrm{d}\nu_n = \int_{\{T_n \in D\}} \overline{\Omega}(X(\theta))\, \mathrm{d}\nu_n.$$

Choose polynomials P_m converging uniformly to $\overline{\Omega}$ on $[-1, 1]^n$, and set $\Omega_m(\theta) := P_m(X(\theta))$. Each P_m is a finite linear combination of products $\prod_i p_i(X_i)$, so it suffices to prove continuity for one such product. For each one-variable polynomial p, write $\varphi_p(t) := \frac{1}{2\pi} \int_0^{2\pi} p(\cos\theta) e^{\mathrm{i}t\cos\theta}\, \mathrm{d}\theta$. The polynomial-amplitude estimate above gives $|\varphi_p(t)| \leq C_p(1 + |t|)^{-1/2}$. For a product $P(X) = \prod_i p_i(X_i)$, set $\Omega_P(\theta) := P(X(\theta))$ and define its weighted density with θ_{out} fixed by

$$h_P(q \mid \theta_{\mathrm{out}}) := \int_{[0,2\pi)^{|J_n|}} \Omega_P(\theta) \delta(T_n(\theta) - q) \prod_{i \in J_n} \frac{\mathrm{d}\theta_i}{2\pi}.$$

Its Fourier transform in q can be expressed as

$$\widehat{h}_P(s \mid \theta_{\mathrm{out}}) := \int_{\mathbb{R}^d} e^{\mathrm{i}s\cdot q} h_P(q \mid \theta_{\mathrm{out}})\, \mathrm{d}q = e^{\mathrm{i}s\cdot q_{\mathrm{out}}} \left[\prod_{i \notin J_n} p_i(\cos\theta_i)\right] \prod_{i \in J_n} \varphi_{p_i}\Big(\frac{s \cdot w_i}{n}\Big).$$

The bracketed product is uniformly bounded because every p_i is bounded on $[-1, 1]$. Using the same three or six reserved factors as above therefore yields

$$|\widehat{h}_P(s \mid \theta_{\mathrm{out}})| \leq M_{P,n}(s) := C_{P,n} \begin{cases} (1+|s|)^{-3/2}, & d = 1, \\ (1+|s|)^{-5/2}, & d = 2. \end{cases}$$

$M_{P,n}$ is integrable, and its Fourier inversion implies h_P continuous in q, and the continuity estimate

$$|h_P(q + u \mid \theta_{\mathrm{out}}) - h_P(q \mid \theta_{\mathrm{out}})| \leq \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} |e^{-\mathrm{i}s\cdot u} - 1| M_{P,n}(s)\, \mathrm{d}s \longrightarrow 0 \qquad (u \to 0).$$

Since the right-hand side is independent of θ_{out}, it also bounds the change in the full density

$$f_{\Omega_P}(q) = \int h_P(q \mid \theta_{\mathrm{out}}) \prod_{i \notin J_n} \frac{\mathrm{d}\theta_i}{2\pi}.$$

Hence f_{Ω_P} is continuous. By linearity in the weight, the same holds for each f_{Ω_m}.

The bound on g_{J_n} also gives the unweighted density bound

$$f_1(q) = \int g_{J_n}(q \mid \theta_{\text{out}}) \prod_{i \notin J_n} \frac{\mathrm{d}\theta_i}{2\pi} \le Cn^d.$$

For any box D, the defining integral identities for f_{Ω_m} and f_{Ω_k} give

$$\left| \int_D (f_{\Omega_m} - f_{\Omega_k})(p)\,\mathrm{d}p \right| = \left| \int_{\{T_n \in D\}} [P_m(X(\theta)) - P_k(X(\theta))]\,\mathrm{d}\nu_n \right| \le \|P_m - P_k\|_\infty \int_D f_1(p)\,\mathrm{d}p.$$

Take $D = D_r(q) := q + (-r, r)^d$, divide by its volume $(2r)^d$, and let $r \downarrow 0$. Since f_{Ω_m}, f_{Ω_k}, and f_1 are continuous, their averages over $D_r(q)$ converge to their values at q. Hence

$$|f_{\Omega_m}(q) - f_{\Omega_k}(q)| \le \|P_m - P_k\|_\infty f_1(q) \le Cn^d \|P_m - P_k\|_\infty.$$

At fixed n, uniform convergence of P_m makes the right-hand side tend to zero as $m, k \to \infty$, uniformly in q. Thus the densities have a uniform limit, which we denote by $F(q) := \lim_{m\to\infty} f_{\Omega_m}(q)$. As a uniform limit of continuous functions, F is continuous. For each m and each fixed bounded box D,

$$\int_D f_{\Omega_m}(q)\,\mathrm{d}q = \int_{\{T_n \in D\}} P_m(X(\theta))\,\mathrm{d}\nu_n.$$

Both sides have limits as $m \to \infty$. Passing to the limit in the preceding identity, and then using the sign-averaging identity, we obtain

$$\int_D F(q)\,\mathrm{d}q = \int_{\{T_n \in D\}} \overline{\Omega}(X(\theta))\,\mathrm{d}\nu_n = \int_{\{T_n \in D\}} \Omega(\theta)\,\mathrm{d}\nu_n.$$

Thus F is the continuous density f_Ω.

Finally, assume $0 \le \Omega(\theta) \le K\mathbf{1}_B(\theta_{\text{out}})$.Averaging first over the angles in J_n gives, we obtain for any box D,

$$\int_D f_\Omega(q)\,\mathrm{d}q \le K \int \mathbf{1}_B(\theta_{\text{out}}) \left[\int_D g_{J_n}(q \mid \theta_{\text{out}})\,\mathrm{d}q \right] \prod_{i \notin J_n} \frac{\mathrm{d}\theta_i}{2\pi} \le KCn^d \mathbb{P}(B)|D|.$$

Taking $D = D_r(q)$, dividing by $(2r)^d$, and letting $r \downarrow 0$ concludes that

$$0 \le f_\Omega(q) \le KCn^d \mathbb{P}(B).$$

□

Negligible degenerate configurations. We first show that configurations with small $\det G$ contribute an arbitrarily small exponential remainder to the density. Choose a smooth function $\chi : [0, \infty) \to [0, 1]$ that is zero on $[0, 1]$ and one on $[2, \infty)$, and set

$$\chi_\kappa(\theta) = \chi\left(\frac{\det G(\theta)}{\kappa} \right), \qquad b_{n,\kappa} = f_{(1-\chi_\kappa)e^{H_n}}.$$

Lemma S3 (An arbitrarily small exponential remainder)**.** *For every $M > 0$, there exist $\kappa \in (0, 1/2]$ and $C_M > 0$ such that, for all sufficiently large n and all q,*

$$0 \le b_{n,\kappa}(q) \le B_n := C_M n^d e^{-Mn}. \tag{S10}$$

Proof. We need an upper bound on H_n to control the Boltzmann weight. We also record its angular derivative bound for the transport argument below. The centered bead positions satisfy

$$r_k = \sum_{i=1}^n b_{ki}(\cos\theta_i, \sin\theta_i), \qquad b_{ki} = \mathbf{1}_{\{i \le k\}} - \frac{n+1-i}{n+1}, \qquad k = 0, \dots, n.$$

Since $|b_{ki}| \le 1$, we have $|r_k| \le n$ and $|\partial_{\theta_i} r_k| \le 1$. Thus

$$H_n = \frac{\lambda}{n^2} \sum_{k=0}^n (r_{k,x}^2 - r_{k,y}^2) \le \lambda(n+1), \quad |\partial_{\theta_i} H_n| \le \frac{2\lambda}{n^2} \sum_{k=0}^n |r_k|\,|\partial_{\theta_i} r_k| \le \frac{2\lambda(n+1)}{n}.$$

We then have

$$H_n \le 2\lambda n, \qquad |\partial_{\theta_i} H_n| \le 4\lambda. \tag{S11}$$

We first construct an event depending only on θ_{out} so that Lemma S2 applies to the weight $(1-\chi_\kappa)e^{H_n}$. Define

$$G_{\text{out}} = \frac{1}{n}\sum_{i\notin J_n} \sin^2\theta_i w_i w_i^T.$$

Here matrix norms are operator norms. The definitions give

$$0 \le G_{\text{out}} \le G, \qquad \|G_{\text{out}}\| \le \operatorname{tr} G \le 2. \tag{S12}$$

For any $\kappa \in (0, 1/2]$, set $\eta = \sqrt{2\kappa}$. Then $\det G \le 2\kappa$ yields $\lambda_{\min}(G) \le \eta$. Since $G_{\text{out}} \le G$ by Eq. (S12), we have

$$\{\det G \le 2\kappa\} \subset \mathcal{B}_\eta, \qquad \mathcal{B}_\eta := \{\lambda_{\min}(G_{\text{out}}) \le \eta\}. \tag{S13}$$

Since $1-\chi_\kappa$ vanishes for $\det G \ge 2\kappa$, combining estimates (S13) and (S11) we obtain

$$0 \le (1-\chi_\kappa)e^{H_n} \le e^{2\lambda n}\mathbf{1}_{\mathcal{B}_\eta}(\theta_{\text{out}}).$$

The event $\mathcal{B}_\eta$ depends only on θ_{out}. Using Lemma S2, with taking $K = e^{2\lambda n}$ and $B = \mathcal{B}_\eta$, we obtain

$$b_{n,\kappa}(q) \le Cn^d e^{2\lambda n}\mathbb{P}(\mathcal{B}_\eta). \tag{S14}$$

It remains to make this probability decay fast enough to compensate for $e^{2\lambda n}$.

To estimate $\mathbb{P}(\mathcal{B}_\eta)$, we will show that a small eigenvalue forces the sine squares of many independent angles to have a small sum. We first ensure that every direction has enough indices with a fixed lower bound on $|a\cdot w_i|^2$. For this purpose, define the deterministic matrix

$$K_{\text{out}} = \frac{1}{n}\sum_{i\notin J_n} w_i w_i^T.$$

Direct summation of the coefficients w_i gives

$$\frac{1}{n}\sum_{i=1}^{n} w_i w_i^T = \begin{cases} 1, & d=1, \\ \operatorname{diag}\left(1, \dfrac{n-1}{3(n+1)}\right), & d=2. \end{cases}$$

Deleting $|J_n| \le 6$ terms changes this matrix by at most $12/n$ since $|w_i|^2 \le 2$. Therefore, for sufficiently large n,

$$K_{\text{out}} \ge I_d/12. \tag{S15}$$

For any unit vector a, we set

$$I_a = \{i \notin J_n : |a\cdot w_i|^2 \ge 1/24\}. \tag{S16}$$

Using Eq. (S15), the definition in Eq. (S16), and $|a\cdot w_i|^2 \le 2$, we obtain

$$\frac{n}{12} \le \sum_{i\notin J_n} |a\cdot w_i|^2 \le \frac{n}{24} + 2|I_a|, \qquad |I_a| \ge \frac{n}{48}. \tag{S17}$$

A unit eigenvector corresponding to $\lambda_{\min}(G_{\text{out}})$ may depend on the angles. Using this vector as a in Eq. (S16) could therefore make the index set depend on the angles. Instead, we first choose a finite set $\mathcal{N}_\eta$ of unit vectors, independently of the angles and of n, such that

$$\text{for every } v\in\mathbb{R}^d \text{ with } |v|=1, \quad \text{there is } a\in\mathcal{N}_\eta \text{ with } |a-v| \le \eta/4.$$

For $d=1$, take $\mathcal{N}_\eta = \{-1, 1\}$; for $d=2$, take sufficiently many equally spaced points on the unit circle. For each fixed $a\in\mathcal{N}_\eta$, the set I_a in Eq. (S16) depends only on a and the given coefficients w_i, and hence does not depend on the angles.

Now consider any angle configuration in $\mathcal{B}_\eta$. Choose a unit eigenvector v for the smallest eigenvalue of G_{out}, and then choose $a\in\mathcal{N}_\eta$ with $|a-v| \le \eta/4$. Using Eq. (S12), we get

$$a^T G_{\text{out}} a \le v^T G_{\text{out}} v + 2\|G_{\text{out}}\|\,|a-v| \le \eta + 4\frac{\eta}{4} = 2\eta.$$

For this a, the definition in Eq. (S16) implies

$$\frac{1}{24n}\sum_{i\in I_a}\sin^2\theta_i \le a^T G_{\text{out}} a \le 2\eta.$$

Thus, for every configuration in $\mathcal{B}_\eta$, this inequality holds for at least one $a \in \mathcal{N}_\eta$. The choice of a may vary with the configuration, so we take the union over all these fixed directions:

$$\mathcal{B}_\eta \subset \bigcup_{a\in\mathcal{N}_\eta}\left\{\sum_{i\in I_a}\sin^2\theta_i \le 48\eta n\right\}. \tag{S18}$$

The bound $|I_a| \ge n/48$ from Eq. (S17) applies to every fixed index set in this union.

We next bound the probability of each scalar-sum event in Eq. (S18). For $t > 0$, define

$$\psi(t) = \frac{1}{2\pi}\int_0^{2\pi} e^{-t\sin^2\theta}\,\mathrm{d}\theta.$$

Splitting this integral into $|\sin\theta| \le r$ and its complement gives

$$\psi(t) \le \frac{2}{\pi}\arcsin r + e^{-tr^2}, \qquad 0 < r < 1.$$

Letting $t \to \infty$ and then $r \downarrow 0$ shows that $\psi(t) \to 0$. The exponential Markov inequality and independence now give, for each fixed $a \in \mathcal{N}_\eta$,

$$\mathbb{P}\left\{\sum_{i\in I_a}\sin^2\theta_i \le 48\eta n\right\} \le e^{48t\eta n}\mathbb{E}\left[e^{-t\sum_{i\in I_a}\sin^2\theta_i}\right] = e^{48t\eta n}\psi(t)^{|I_a|} \le \exp\left\{n\left[48t\eta + \frac{1}{48}\log\psi(t)\right]\right\}. \tag{S19}$$

The last inequality uses $0 < \psi(t) \le 1$ and the cardinality bound in Eq. (S17). Combining the event inclusion in Eq. (S18) with Eq. (S19), and summing over the finite set of directions, yields

$$\mathbb{P}(\mathcal{B}_\eta) \le |\mathcal{N}_\eta|\exp\left\{n\left[48t\eta + \frac{1}{48}\log\psi(t)\right]\right\}. \tag{S20}$$

To compensate for the factor $e^{2\lambda n}$ in Eq. (S14) and obtain the prescribed rate M, first choose t so that $(\log\psi(t))/48 < -(M + 2\lambda + 2)$, and then choose κ so that $48t\sqrt{2\kappa} < 1$. With these choices, Eq. (S20) gives

$$\mathbb{P}(\mathcal{B}_\eta) \le |\mathcal{N}_\eta| e^{-n(M+2\lambda+1)}, \tag{S21}$$

and both κ and $|\mathcal{N}_\eta|$ are independent of n.

Substituting Eq. (S21) into the density bound in Eq. (S14), we conclude that

$$b_{n,\kappa}(q) \le Cn^d e^{2\lambda n}|\mathcal{N}_\eta|e^{-n(M+2\lambda+1)} = C|\mathcal{N}_\eta|n^d e^{-n(M+1)} \le C_M n^d e^{-Mn},$$

where the fixed factor $C|\mathcal{N}_\eta|$ is absorbed into C_M. □

Density comparison by bond rotations. For configurations with $\chi_\kappa > 0$, we use bond rotations to translate T_n from q to a nearby p. We bound the changes in the Boltzmann weight and angular volume to compare the densities at these points. Lemma S3 bounds the remaining contribution to the density.

Lemma S4 (Density comparison by transport)**.** *Let κ and B_n be as in Lemma S3, and set $\delta_\kappa = \kappa/(16\sqrt{2})$ and $L_\kappa = 8\lambda/\sqrt{\kappa} + 8\sqrt{2}/\kappa$. For all sufficiently large n and all $p, q \in \mathbb{R}^d$ with $|p - q| < \delta_\kappa$, we have*

$$f_n(q) \le e^{nL_\kappa|p-q|} f_n(p) + B_n. \tag{S22}$$

Proof. For a unit vector $a \in \mathbb{R}^d$ and $\det G > 0$, define

$$b_i = w_i^T G^{-1} a, \qquad d_i = w_i^T G^{-1} w_i, \qquad \mathcal{V}_i = -\sin\theta_i b_i.$$

We write $\mathcal{V}A = \sum_i \mathcal{V}_i \partial_{\theta_i} A$ for the derivative of an angular function A along the vector field $\mathcal{V}$. Since $\partial_{\theta_i} T_n = -n^{-1} \sin\theta_i w_i$,

$$\mathcal{V}T_n = \frac{1}{n}\sum_i \sin^2\theta_i w_i w_i^T G^{-1} a = GG^{-1}a = a.$$

Thus moving the angles with velocity $\mathcal{V}$ translates T_n at unit speed in direction a.

To bound the change in $\det G$ and the divergence of $\mathcal{V}$, differentiate G and its inverse:

$$\partial_{\theta_i} G = \frac{2}{n}\sin\theta_i\cos\theta_i w_i w_i^T, \qquad \partial_{\theta_i} G^{-1} = -G^{-1}(\partial_{\theta_i} G)G^{-1}.$$

Hence $\partial_{\theta_i}\log\det G = (2/n)\sin\theta_i\cos\theta_i d_i$ and $\partial_{\theta_i} b_i = -(2/n)\sin\theta_i\cos\theta_i d_i b_i$. Substituting into $\mathcal{V}\log\det G$ and $\operatorname{div}\mathcal{V} := \sum_i \partial_{\theta_i}\mathcal{V}_i$ yields

$$\begin{aligned} \mathcal{V}\log\det G &= -\frac{2}{n}\sum_i \sin^2\theta_i\cos\theta_i b_i d_i, \\ \operatorname{div}\mathcal{V} &= -\sum_i \cos\theta_i b_i + \frac{2}{n}\sum_i \sin^2\theta_i\cos\theta_i d_i b_i, \\ \frac{1}{n}\sum_i \sin^2\theta_i d_i &= \operatorname{tr}(G^{-1}G) = d. \end{aligned} \tag{S23}$$

Set $\mu = \lambda_{\min}(G)$. The first and third identities in Eq. (S23), together with $d_i \geq 0$ and $|b_i| \leq \sqrt{2}/\mu$, give

$$|\mathcal{V}\det G| \leq \frac{2\det G}{n}\max_i |b_i| \sum_i \sin^2\theta_i d_i \leq 2d\sqrt{2}\,\frac{\det G}{\mu} \leq 8\sqrt{2}. \tag{S24}$$

Here $\det G/\mu = 1$ for $d = 1$, while for $d = 2$ it equals $\lambda_{\max}(G) \leq \operatorname{tr} G \leq 2$.

To obtain a smooth invertible flow for the angular change of variables, we extend $\mathcal{V}$ to all angles. Choose a smooth $\rho : [0,\infty) \to [0,1]$ that is zero on $[0,1/4]$ and one on $[1/2,\infty)$. Define

$$\widetilde{\mathcal{V}} = \rho(\det G/\kappa)\mathcal{V} \quad (\det G > 0), \qquad \widetilde{\mathcal{V}} = 0 \quad (\det G = 0),$$

and let $\Phi_t(\theta)$ solve

$$\frac{\mathrm{d}}{\mathrm{d}t}\Phi_t(\theta) = \widetilde{\mathcal{V}}(\Phi_t(\theta)), \qquad \Phi_0(\theta) = \theta.$$

All angles are interpreted periodically. The extended field is smooth and bounded, so its flow exists for all t and has inverse Φ_{-t}.

Consider a trajectory starting at θ with $\chi_\kappa(\theta) > 0$, so $\det G(\theta) > \kappa$. Along this trajectory, the cutoff $\rho(\det G/\kappa)$ equals one until $\det G$ first reaches $\kappa/2$. Equation (S24) then gives

$$\det G(\Phi_t\theta) \geq \det G(\theta) - 8\sqrt{2}\,t > \frac{\kappa}{2}, \qquad 0 \leq t < \delta_\kappa = \frac{\kappa}{16\sqrt{2}}.$$

Thus $\det G$ remains above $\kappa/2$ for $0 \leq t < \delta_\kappa$. On this interval,

$$T_n(\Phi_t\theta) = T_n(\theta) + ta, \qquad \mu \geq \frac{\det G}{2} > \frac{\kappa}{4}.$$

The eigenvalue bound uses $\det G/\mu \leq 2$.

To bound the change in H_n along these trajectories, substitute the angular velocities and use $\sum_i \sin^2\theta_i w_i w_i^T = nG$:

$$\sum_i \mathcal{V}_i^2 = \sum_i \sin^2\theta_i (w_i^T G^{-1} a)^2 = a^T G^{-1}\left(\sum_i \sin^2\theta_i w_i w_i^T\right) G^{-1} a = n a^T G^{-1} G G^{-1} a = n a^T G^{-1} a \leq \frac{n}{\mu} \leq \frac{4n}{\kappa}.$$

Cauchy–Schwarz yields $\sum_i |\mathcal{V}_i| \leq 2n/\sqrt{\kappa}$, so Eq. (S11) implies

$$|\mathcal{V}H_n| \leq 4\lambda \sum_i |\mathcal{V}_i| \leq \frac{8\lambda n}{\sqrt{\kappa}}.$$

For the angular volume change, we bound $\operatorname{div}\mathcal{V}$. Using $|b_i| \leq 4\sqrt{2}/\kappa$ in Eq. (S23) we obtain

$$|\operatorname{div}\mathcal{V}| \leq n\max_i |b_i| + \frac{2}{n}\max_i |b_i| \sum_i \sin^2\theta_i d_i \leq \frac{4\sqrt{2}n + 16\sqrt{2}}{\kappa} \leq \frac{8\sqrt{2}n}{\kappa} \qquad (n \geq 4).$$

The Jacobian of the angular change of variables is

$$j_t(\theta) = |\det D_\theta \Phi_t(\theta)|.$$

Differentiating the flow equation with respect to the initial angles gives

$$\frac{\mathrm{d}}{\mathrm{d}t} D_\theta \Phi_t(\theta) = D\widetilde{\mathcal{V}}(\Phi_t\theta)\, D_\theta \Phi_t(\theta),$$

where $D\widetilde{\mathcal{V}}$ is the matrix with entries $\partial_{\theta_j} \widetilde{\mathcal{V}}_i$. The matrix $D_\theta \Phi_t$ is invertible because Φ_t has the smooth inverse Φ_{-t}. Jacobi's formula and cyclic invariance of the trace give

$$\frac{\mathrm{d}}{\mathrm{d}t} \log j_t(\theta) = \operatorname{tr}\left[(D_\theta \Phi_t)^{-1} \frac{\mathrm{d}}{\mathrm{d}t} D_\theta \Phi_t\right] = \operatorname{tr}\left[(D_\theta \Phi_t)^{-1} D\widetilde{\mathcal{V}}(\Phi_t\theta) D_\theta \Phi_t\right] = \operatorname{tr} D\widetilde{\mathcal{V}}(\Phi_t\theta) = (\operatorname{div} \widetilde{\mathcal{V}})(\Phi_t\theta).$$

For trajectories starting at θ with $\chi_\kappa(\theta) > 0$, the cutoff $\rho(\det G/\kappa)$ equals one in a neighborhood of every trajectory point for $0 \le t < \delta_\kappa$. Hence $\widetilde{\mathcal{V}}$ and its derivatives agree with $\mathcal{V}$ there. Combining the bounds on $\mathcal{V}H_n$ and $\operatorname{div} \mathcal{V}$ gives

$$\left|\frac{\mathrm{d}}{\mathrm{d}t} \log\left[e^{H_n(\Phi_t\theta)} j_t(\theta)\right]\right| = |(\mathcal{V}H_n + \operatorname{div} \mathcal{V})(\Phi_t\theta)| \le nL_\kappa.$$

Integrating from 0 to t, with $j_0 = 1$, yields

$$e^{H_n(\theta)} \le e^{nL_\kappa t} e^{H_n(\Phi_t\theta)} j_t(\theta), \qquad \chi_\kappa(\theta) > 0, \quad 0 \le t < \delta_\kappa. \tag{S25}$$

For any box $D \subset \mathbb{R}^d$, the identity $T_n(\Phi_t\theta) = T_n(\theta) + ta$ shows that Φ_t maps $\{T_n \in D, \chi_\kappa > 0\}$ into $\{T_n \in D + ta\}$. Integrating Eq. (S25), using $0 \le \chi_\kappa \le 1$, and changing angular variables by Φ_t gives

$$\int_D f_{\chi_\kappa e^{H_n}}(x)\,\mathrm{d}x = \int_{\{T_n \in D\}} \chi_\kappa e^{H_n}\,\mathrm{d}\nu_n \le e^{nL_\kappa t} \int_{\{T_n \in D, \chi_\kappa > 0\}} e^{H_n(\Phi_t\theta)} j_t(\theta)\,\mathrm{d}\nu_n(\theta) \le e^{nL_\kappa t} \int_{D+ta} f_n(x)\,\mathrm{d}x.$$

Take $D = D_r(q) = q + (-r, r)^d$, divide by $(2r)^d$, and let $r \downarrow 0$. The densities $f_{\chi_\kappa e^{H_n}}$ and f_n are continuous by Lemma S2, so

$$f_n(q) - b_{n,\kappa}(q) = f_{\chi_\kappa e^{H_n}}(q) \le e^{nL_\kappa t} f_n(q + ta).$$

For $p \ne q$, choose $a = (p - q)/|p - q|$ and $t = |p - q|$. The bound $b_{n,\kappa}(q) \le B_n$ from Lemma S3 then concludes Eq. (S22). □

From window averages to pointwise free energies.

Proof of Theorem S2. Fix q as in Theorem S2 and abbreviate $A_{n,\varepsilon} = A_{\alpha,n,\varepsilon}(q;\lambda)$. For m_{ref} and $J(q)$ defined above, we have $Q_\alpha(m_{\mathrm{ref}}) = q$, $\mathcal{E}[m_{\mathrm{ref}}] \le 0$, and $J(q) < \infty$. Evaluating the variational problems at m_{ref} gives

$$S_{\alpha,\varepsilon}(q) \ge S_\alpha(q) \ge -J(q) > -\infty.$$

Choose $M > J(q) + 5$ and the corresponding κ, B_n from Lemma S3. These choices are independent of ε.

For $|p - q|_\infty < \varepsilon < \delta_\kappa/\sqrt{d}$, we have $|p - q| \le \sqrt{d}\,\varepsilon$. Applying Lemma S4 in both directions and using $f_n \ge 0$ gives

$$f_n(q) \le e^{nC_\kappa \varepsilon} f_n(p) + B_n, \qquad f_n(p) \le e^{nC_\kappa \varepsilon} f_n(q) + B_n, \qquad C_\kappa = \sqrt{d}\, L_\kappa.$$

Averaging over $|p - q|_\infty < \varepsilon$ and rearranging yields

$$e^{-nC_\kappa \varepsilon}[f_n(q) - B_n] \le A_{n,\varepsilon} \le e^{nC_\kappa \varepsilon} f_n(q) + B_n, \qquad 0 < \varepsilon < \delta_\kappa/\sqrt{d}. \tag{S26}$$

To show that $B_n/f_n(q) \to 0$ independently of ε, we first bound $f_n(q)$ from below. Fix $0 < \varepsilon_0 < \delta_\kappa/\sqrt{d}$ with $C_\kappa \varepsilon_0 \le 1$. Equation (S9) and $S_{\alpha,\varepsilon_0}(q) \ge -J(q)$ give

$$A_{n,\varepsilon_0} \ge e^{-n[J(q)+1]} \qquad \text{for sufficiently large } n.$$

Since $M > J(q) + 5$, we also have $B_n \le \frac{1}{2} e^{-n[J(q)+1]}$ for sufficiently large n. The upper bound in Eq. (S26) at $\varepsilon = \varepsilon_0$ therefore implies

$$f_n(q) \ge e^{-nC_\kappa \varepsilon_0}[A_{n,\varepsilon_0} - B_n] \ge \tfrac{1}{2} e^{-n[J(q)+2]} > 0. \tag{S27}$$

Combining Eqs. (S27) and (S10) gives

$$\mathcal{R}_n := \frac{B_n}{f_n(q)} \le 2C_M n^d e^{-n[M-J(q)-2]} \longrightarrow 0,$$

This bound is independent of ε. Dividing Eq. (S26) by $f_n(q)$ and using $e^{nC_\kappa\varepsilon} \ge 1$ we obtain

$$e^{-nC_\kappa\varepsilon}(1-\mathcal{R}_n) \le \frac{A_{n,\varepsilon}}{f_n(q)} \le e^{nC_\kappa\varepsilon}(1+\mathcal{R}_n). \tag{S28}$$

Since $\mathcal{R}_n < 1$ for sufficiently large n, the lower bound in Eq. (S28) is positive. Taking logarithms gives

$$-C_\kappa\varepsilon + \frac{1}{n}\log(1-\mathcal{R}_n) \le \frac{1}{n}\log A_{n,\varepsilon} - \frac{1}{n}\log f_n(q) \le C_\kappa\varepsilon + \frac{1}{n}\log(1+\mathcal{R}_n).$$

Since $\log(1+\mathcal{R}_n) \le -\log(1-\mathcal{R}_n)$, this yields

$$\left|\frac{1}{n}\log A_{n,\varepsilon} - \frac{1}{n}\log f_n(q)\right| \le C_\kappa\varepsilon + \xi_n, \qquad \xi_n := -\frac{1}{n}\log(1-\mathcal{R}_n) \longrightarrow 0. \tag{S29}$$

The error ξ_n is independent of ε. For each fixed $0 < \varepsilon < \delta_\kappa/\sqrt{d}$, combining estimates (S29) and (S9), we obtain

$$S_{\alpha,\varepsilon}(q) - C_\kappa\varepsilon \le \liminf_{n\to\infty}\frac{1}{n}\log f_n(q) \le \limsup_{n\to\infty}\frac{1}{n}\log f_n(q) \le S_{\alpha,\varepsilon}(q) + C_\kappa\varepsilon.$$

Letting $\varepsilon \downarrow 0$ and using Eq. (S9) shows that

$$\lim_{n\to\infty}\frac{1}{n}\log f_n(q) = S_\alpha(q). \tag{S30}$$

To take $\varepsilon \downarrow 0$ before $n \to \infty$, fix n large enough for Eq. (S27). The continuity of f_n from Lemma S2 and the bound $f_n(q) > 0$ imply

$$\lim_{\varepsilon\downarrow 0}\frac{1}{n}\log A_{n,\varepsilon} = \frac{1}{n}\log f_n(q).$$

Taking $n \to \infty$ and using Eq. (S30), we obtain $\lim_{n\to\infty}\lim_{\varepsilon\downarrow 0} n^{-1}\log A_{n,\varepsilon} = S_\alpha(q)$. Equation (S9) yields $\lim_{\varepsilon\downarrow 0}\lim_{n\to\infty} n^{-1}\log A_{n,\varepsilon} = S_\alpha(q)$ for the opposite order of limits. Thus both iterated limits agree,which concludes Theorem S2. □

S3.B Proof of the minimizing-interval theorem

Proof. We begin with the scalar variational problem in Eq. (S8). Write its constrained value as

$$G_\lambda(c) := \inf_{\substack{u\in\mathcal{A}_1 \\ u(1)=0,\ -2\bar{u}=c}} \mathcal{F}_\lambda^{(1)}[u], \qquad c \in \mathbb{R}.$$

Equation (S8) implies $G_\lambda(c) = F_{23,\infty}(0,c;\lambda)$ for $|c| < 1/2$; we verify the boundary and exterior values below. First, drop the mean constraint and define

$$\mathcal{M}_\lambda = \operatorname*{argmin}_{\substack{u\in\mathcal{A}_1 \\ u(1)=0}} \mathcal{F}_\lambda^{(1)}[u].$$

The admissible set is compact in the uniform topology by the Arzelà–Ascoli theorem. On this set, the convex entropy integral is lower semicontinuous, while the quadratic energy term is continuous and bounded. Since the entropy term is nonnegative and $\mathcal{F}_\lambda^{(1)}[0] = 0$, the minimum $m_\lambda := \min_{u\in\mathcal{A}_1,\, u(1)=0}\mathcal{F}_\lambda^{(1)}[u] \le 0$ is finite, $\mathcal{M}_\lambda$ is nonempty and compact, and its elements have finite entropy cost. Fixing the mean restricts the admissible set, so $G_\lambda(c) \ge m_\lambda$ for every $c \in \mathbb{R}$. If $u \in \mathcal{M}_\lambda$ and $c = -2\bar{u}$, then

$$m_\lambda \le G_\lambda(c) \le \mathcal{F}_\lambda^{(1)}[u] = m_\lambda.$$

Thus each such c minimizes G_λ, with $\min_c G_\lambda(c) = m_\lambda$. Conversely, if c minimizes G_λ, the fixed-mean set is compact with finite infimum $G_\lambda(c) = m_\lambda$, which is attained by a path $u_c \in \mathcal{M}_\lambda$ satisfying $c = -2\bar{u}_c$. Therefore,

$$\operatorname*{argmin}_{c\in\mathbb{R}} G_\lambda(c) = \{-2\bar{u} : u \in \mathcal{M}_\lambda\}.$$

For $u \in \mathcal{M}_\lambda$, let $\widetilde{u}$ be its 1-periodic extension and shift the starting point by $a \in [0,1]$. Subtracting $u(a)$ preserves the zero endpoint values, and the shifted paths satisfy

$$u_a(s) = \widetilde{u}(s+a) - u(a), \qquad u_a(0) = u_a(1) = 0, \qquad \mathcal{F}_\lambda^{(1)}[u_a] \;= \mathcal{F}_\lambda^{(1)}[u], \qquad -2\bar{u}_a = 2\big(u(a) - \bar{u}\big). \tag{S31}$$

Indeed, the derivative is shifted over one period, and subtracting a constant leaves the variance unchanged. Thus $u_a \in \mathcal{M}_\lambda$. As a varies over $[0,1]$, continuity of u ensures that $-2\bar{u}_a$ sweeps out $[2(\min u - \bar{u}), 2(\max u - \bar{u})]$, which contains 0. Every such interval consists of minimizing values of c, and every minimizing value belongs to one of them by taking $a = 0$ (since $u_0 = u$). Their union over $u \in \mathcal{M}_\lambda$ is therefore an interval containing 0. It is compact because the mean map $u \mapsto \bar{u}$ is continuous on $\mathcal{M}_\lambda$, and symmetric because $u \mapsto -u$ preserves the functional. Consequently,

$$\operatorname*{argmin}_{c \in \mathbb{R}} G_\lambda(c) = [-a_\lambda, a_\lambda],$$

where

$$a_\lambda = 2 \max_{u \in \mathcal{M}_\lambda} |\bar{u}|.$$

We next determine when the interval opens. The series $I_0(h) = \sum_{k \geq 0} (h^2/4)^k/(k!)^2$ gives $I_0(h) < e^{h^2/4}$ for $h \neq 0$. Taking $h = 2v$ in the Legendre supremum gives $\ell(v) > v^2$ for $v \neq 0$. Set $K(h) = \log I_0(h)$. Since K is strictly convex and $K''(0) = 1/2$, the Legendre maximizer near $v = 0$ solves $K'(h) = v$. The series gives

$$K(h) = \frac{h^2}{4} - \frac{h^4}{64} + O(h^6), \qquad K'(h) = \frac{h}{2} - \frac{h^3}{16} + O(h^5),$$
$$h(v) = 2v + v^3 + O(v^5).$$

Substitution into $\ell(v) = vh(v) - K(h(v))$ yields

$$\ell(0) = 0, \qquad \ell(v) > v^2 \quad (v \neq 0), \qquad \ell(v) = v^2 + \tfrac{1}{4}v^4 + O(v^6) \quad (v \to 0). \tag{S32}$$

For nonzero $u \in \mathcal{A}_1$ with $u(1) = 0$, the periodic Wirtinger inequality gives $\int_0^1 |u'|^2 \, \mathrm{d}s \geq 4\pi^2 \int_0^1 (u - \bar{u})^2 \, \mathrm{d}s$. Together with Eq. (S32), this implies $\mathcal{F}_\lambda^{(1)}[u] > 0$ when $\lambda \leq 4\pi^2$. Hence $\mathcal{M}_\lambda = \{0\}$ and $a_\lambda = 0$ in this range. For $\lambda > 4\pi^2$, the trial path $u_\varepsilon(s) = \varepsilon \sin(2\pi s)$ is admissible for $|\varepsilon| \leq (2\pi)^{-1}$ and satisfies

$$\mathcal{F}_\lambda^{(1)}[u_\varepsilon] = \frac{\varepsilon^2}{2}(4\pi^2 - \lambda) + O(\varepsilon^4) < 0$$

for sufficiently small $\varepsilon \neq 0$. Thus $m_\lambda < 0$, and every $u \in \mathcal{M}_\lambda$ is nonconstant. Choosing $a \in [0,1]$ with $u(a) \neq \bar{u}$ in Eq. (S31) then gives a minimizing path $u_a \in \mathcal{M}_\lambda$ with nonzero mean. Consequently, $a_\lambda > 0$.

Finally, we identify G_λ with the density-defined free energy on the remaining values of c. For $h > 0$ and $0 < r < \pi$, splitting the angular integral gives $e^{-h} I_0(h) \leq r/\pi + e^{-h(1-\cos r)}$. Letting $h \to \infty$ and then $r \downarrow 0$ gives $e^{-h} I_0(h) \to 0$, so the Legendre definition yields $\ell(\pm 1) = +\infty$. Any finite-entropy $u \in \mathcal{A}_1$ with $u(1) = 0$ then has $|u'| < 1$ almost everywhere; integration from either endpoint gives $|u(s)| < \min(s, 1-s)$ for $0 < s < 1$, and hence $|\bar{u}| < 1/4$. Thus $G_\lambda(c) = +\infty$ for $|c| \geq 1/2$. Since $|\hat{z}_3| \leq n^{-1} \sum_i |c_i| \leq n/[2(n+1)] < 1/2$, the density-defined free energy is also $+\infty$ there. Together with Eq. (S8) on $|c| < 1/2$, this gives $G_\lambda(c) = F_{23,\infty}(0, c; \lambda)$ for all $c \in \mathbb{R}$, proving Eq. (S7) and the stated threshold. □

First transition. For the endpoint observable, $|\hat{z}_2| \leq 1$, so its continuous weighted density vanishes for $|z_2| > 1$ and, by continuity, at $z_2 = \pm 1$. In the scalar problem, $u(1) = \pm 1$ forces $u' = \pm 1$ almost everywhere and hence infinite entropy cost, while $|u(1)| > 1$ is impossible. Thus Eq. (S8) extends to

$$F_{2,\infty}(z_2; \lambda) = \inf_{\substack{u \in \mathcal{A}_1 \\ u(1) = z_2}} \mathcal{F}_\lambda^{(1)}[u], \qquad z_2 \in \mathbb{R}.$$

A similar argument determines the first transition threshold $\lambda_1 = \pi^2$. Since the admissible set $\mathcal{A}_1$ is compact in the uniform topology and $\mathcal{F}_\lambda^{(1)}$ is lower semicontinuous, $\mathcal{F}_\lambda^{(1)}$ attains its global minimum on $\mathcal{A}_1$, and $\operatorname{argmin}_{z_2 \in \mathbb{R}} F_{2,\infty}(z_2; \lambda) = \{u(1) : u \in \operatorname{argmin}_{\mathcal{A}_1} \mathcal{F}_\lambda^{(1)}\}$. For $w = u - \bar{u}$ without the periodic boundary condition, expanding in the cosine basis $\{\cos(\pi k s)\}_{k \geq 1}$ on $[0,1]$ yields the sharp Poincaré inequality $\int_0^1 |w'|^2 \, \mathrm{d}s \geq \pi^2 \int_0^1 w^2 \, \mathrm{d}s$. By Eq. (S32), $u = 0$ is therefore the unique global minimizer of $\mathcal{F}_\lambda^{(1)}$ for all $\lambda \leq \pi^2$. For $\lambda > \pi^2$, the trial path

$u_\varepsilon(s) = \varepsilon[\cos(\pi s) - 1]$ satisfies $\mathcal{F}_\lambda^{(1)}[u_\varepsilon] = \frac{1}{2}\varepsilon^2(\pi^2 - \lambda) + O(\varepsilon^4) < 0$ for sufficiently small $\varepsilon \neq 0$, so any global minimizer is nonconstant. Moreover, every global minimizer u must be monotone. Indeed, replacing u with $v(t) = \int_0^t |u'(s)|\mathrm{d}s$ preserves the entropy cost while increasing the variance:

$$\int_0^1 (u - \bar{u})^2 \mathrm{d}t = \frac{1}{2}\int_0^1 \int_0^1 [u(t) - u(s)]^2 \mathrm{d}s\mathrm{d}t \leq \frac{1}{2}\int_0^1 \int_0^1 [v(t) - v(s)]^2 \mathrm{d}s\mathrm{d}t = \int_0^1 (v - \bar{v})^2 \mathrm{d}t. \tag{S33}$$

If u' takes both positive and negative values on sets of positive measure, then $|u(1) - u(0)| < v(1) - v(0)$. By continuity, $|u(t) - u(s)| < |v(t) - v(s)|$ holds on a neighborhood of $(s, t) = (0, 1)$, making the inequality in Eq. (S33) strict and thus strictly decreasing $\mathcal{F}_\lambda^{(1)}$. Consequently, for $\lambda > \pi^2$, every global minimizer is nonconstant and monotone, and thus has a nonzero endpoint value $u(1)$, with opposite signs occurring in pairs by the reflection symmetry $u \mapsto -u$.

S3.C Numerical evaluation

The scalar variational problems in Eq. (S8) are evaluated by a piecewise-linear discretization on M uniform intervals of length $h = 1/M$. Writing v_j for the slope on interval j, we set

$$u_0 = 0, \qquad u_j = h\sum_{i=1}^{j} v_i, \qquad |v_j| \leq 1 - 10^{-6}, \qquad j = 1, \ldots, M.$$

With trapezoidal weights $\omega_0 = \omega_M = 1/2$ and $\omega_j = 1$ otherwise, the discrete functional is

$$\bar{u}_M = h\sum_{j=0}^{M} \omega_j u_j, \qquad \mathcal{F}_{\lambda,M}^{(1)}(v) = h\sum_{j=1}^{M} \ell(v_j) - \lambda h \sum_{j=0}^{M} \omega_j (u_j - \bar{u}_M)^2.$$

Both problems impose $u_M = z_2$. The joint problem also imposes the mean constraint

$$h\sum_{j=1}^{M} (1 - s_{j-1/2}) v_j = \frac{z_2 - z_3}{2}, \qquad s_{j-1/2} = (j - \tfrac{1}{2})h,$$

and the section used in the main text is obtained by setting $z_2 = 0$.

For $0 < |v| < 1$, we evaluate the entropy density from

$$\ell(v) = h_v|v| - \log I_0(h_v), \qquad \frac{I_1(h_v)}{I_0(h_v)} = |v|,$$

with $h_v \geq 0$ determined numerically and $\ell(0) = 0$. Here I_0 and I_1 are modified Bessel functions. The constrained minimization is performed with SLSQP to a tolerance of 10^{-9}. The endpoint solutions are continued in $|z_2|$ from constant-slope initializations. For the joint problem, continuation is supplemented by multiple projected initializations; solutions with residuals above 2×10^{-7} are discarded, and the smallest converged objective is retained. Reflection symmetry supplies the negative half of each coordinate grid.

At fixed λ, let $\widehat{\mathcal{F}}(q)$ denote the retained minimum on the coordinate grid $\mathcal{Q}$. The corresponding numerical free energy profile, shifted to zero minimum, is $\widehat{F}(q) = \widehat{\mathcal{F}}(q) - \min_{q' \in \mathcal{Q}} \widehat{\mathcal{F}}(q')$. For the joint section, the minimizing interval is estimated from the connected grid points whose objective lies within 10^{-6} of the grid minimum, with adjacent exterior points bracketing its endpoints.

Finally, the analytical-model parameter can be related to the flow strength used for the kinetic model. For beadwise mobility $1/\zeta$ and physical bond length b, matching the quadratic planar-flow potential in units of $k_\mathrm{B}T$ to Eq. (3) gives

$$\lambda = \frac{\zeta \dot{\epsilon} b^2 n^2}{2k_\mathrm{B}T} = \frac{\zeta b^2 n^2}{2k_\mathrm{B}T\tau}\mathrm{Wi}, \qquad \mathrm{Wi} = \dot{\epsilon}\tau.$$

For the 300-bead reference chain, $\widetilde{\dot{\epsilon}} = \dot{\epsilon}\tau_d$ and $\tau/\tau_d = 4321$, so we have $\frac{\lambda}{n^2} = \frac{\widetilde{\dot{\epsilon}}}{2}$, $\mathrm{Wi} = \frac{8640}{n^2}\lambda$. Using the approximation $n \simeq 300$ for this reference chain, we express the analytical results in Fig. 2 in terms of $\mathrm{Wi} = \frac{8640}{300^2}\lambda = 0.096\lambda$.

S4 Single-molecule experiments and image-response analysis

S4.A Experimental protocol and image observables

We acquired single-molecule DNA fluorescence trajectories using the automated electrokinetic stretcher of Ref. [24], which controls field strength via the applied voltages and keeps the molecule near the stagnation point via active feedback control (see Ref. [24] for experimental details).

Image coordinates and trajectory centering follow Eq. (10). We threshold pixels at $I_p \geq 20$, discard frames with fewer than 40 foreground pixels, and remove connected components smaller than 10% of the largest component to extract the projected span $L = x_{\max} - x_{\min}$. Since $x_{\min}$ and $x_{\max}$ can lie on folded interior segments rather than the true chain ends, these image observables serve as surrogates, instead of exact matches to the microscopic coordinates.

We define the reference span $L_{\max}$ as the 99th percentile of L under matched optics. We retain frames whose mask stays clear of the image boundary and changes in L by less than $0.25L_{\max}$ between consecutive frames. Reference frames are further restricted to $0.30 \leq L(t)/L_{\max} < 0.55$ prior to the first crossing of $L > 0.72L_{\max}$, isolating a common stage of initial stretching (rather than imposing $Z_2 = 0$).

S4.B Lag selection and response estimation

Both future frames in each finite-lag increment (End Matter) must also pass these quality checks. At each voltage, we select the lag pair (δ_1, δ_2) such that the mean relative span growth

$$r_L(\delta_1, \delta_2) = \left\langle \frac{L(t+\delta_2) - L(t+\delta_1)}{L(t)} \right\rangle .$$

is closest to 5%. The table below lists the resulting lag pairs, relative growth r_L, and number of contributing trajectories N:

Voltage (V)	N	δ_1 (frames)	δ_2 (frames)	r_L
10	44	20	59	0.006
20	201	2	5	0.047
40	335	1	2	0.045
50	444	1	2	0.058

The conditioning coordinate $z(t)$, increment $\Delta\widetilde{Z}_3(t)$, raw response f, and odd response g follow Eq. (11), with all valid pairs weighted equally. By default, we use eight equal-width bins over $[-2.6\sigma_z, 2.6\sigma_z]$ (where σ_z is the standard deviation of z), requiring at least 800 valid pairs overall and 150 per retained bin. Figure S2(a,b) compares f and g at 20 and 40 V, showing that the central positive slope and outer downturn at 40 V are already evident in the raw response f prior to odd symmetrization.

Odd-response estimates require both symmetric bins in a pair to be valid. We compute error bars from the 16th–84th percentiles across 2000 bootstrap resamples of the N trajectories (with replacement and fixed bin edges, omitting bins with fewer than 150 pairs). We estimate the central slope s_0 by a linear fit to the odd response over bins with $|z| \leq \sigma_z$ (requiring at least three valid bins; four innermost bins by default), which provides a local alternative to the full-range cubic coefficient c_1 in the End Matter.

To test robustness across analysis parameters [Fig. S2(c)], we vary the span window ($[0.28, 0.52)$, $[0.30, 0.55)$, $[0.32, 0.58)$), target growth r_L (0.040, 0.051, 0.060), bin count, and binning range ($\pm 2.3\sigma_z$, $\pm 2.6\sigma_z$, $\pm 2.9\sigma_z$). Among the 108 combinations, all 72 admissible settings yield $s_0 < 0$ at 20 V and $s_0 > 0$ at 40 V, confirming a robust sign reversal of the central slope despite sampling uncertainty at 40 V; the outer zero crossing is resolved in 54 of the 72 settings.



Figure S2: Experimental response checks and sensitivity to analysis choices. (a,b) Raw conditional response f (gray circles) and odd component g (red squares) at 20 and 40 V. (c) Central slopes s_0 at 20 and 40 V for the 72 admissible analysis settings.