

Finite-Time Concentration and Convergence Rates for Projected Two-Time-Scale Stochastic Approximation with Markov Noise

Rahul Singh¹, Vivek S. Borkar², and Eric Moulines^{1,3}

¹Laboratoire de Recherche de l'EPITA, Paris, France

²Indian Institute of Technology Bombay

³Mohamed bin Zayed University of Artificial Intelligence, Abu Dhabi, United Arab Emirates

Abstract

We study projected two-time-scale stochastic approximation algorithms driven by a controlled Markov chain. The fast recursion is governed by an averaged contractive map, while the slow recursion is constrained to a compact convex polyhedron through a projection step. The limiting slow dynamics are described by a projected differential equation, whose vector field may be discontinuous on the boundary of the constraint set. This feature prevents a direct application of standard finite-time stochastic approximation arguments based on Lipschitz limiting vector fields.

We establish explicit blockwise high-probability bounds showing that the fast iterate tracks the moving fixed point $x^*(\theta_n)$ of the averaged fast recursion and that the step embedding of the slow iterate tracks the associated projected ODE. The bounds separate fast and slow martingale deviations, Poisson residuals generated by Markovian sampling, and the deterministic two-time-scale bias. Under a Lipschitz Lyapunov function satisfying a uniform one-block decrease condition for the projected ODE, these estimates yield almost-sure convergence of the last-iterate Lyapunov error and explicit rates when the decrease modulus has a power lower bound near zero. Uniform one-block contraction is the linear-decrease special case.

A uniform decrease condition for a Lipschitz Lyapunov function over fixed time intervals yields almost-sure convergence of the slow Lyapunov error, with explicit last-iterate rates when the decrease admits a power lower bound near zero. Under uniform Lyapunov contraction, suitably chosen polynomial step sizes give joint fast-tracking and slow Lyapunov-error exponents arbitrarily close to $1/3$, while a slow step size of order $1/n$ attains the fast-tracking exponent $1/3$. When the reduced slow update map is also a Euclidean contraction, logarithmically separated step sizes improve the joint rate to $O_{a.s.}(n^{-1/2} \log n)$, including when the equilibrium lies on the boundary. The same rate holds under a distinct geometric condition: the slow equilibrium set is a strictly attracting face of a box, and the fast equilibrium is constant on that face.

We apply the results to an actor-critic algorithm, obtaining an almost-sure value-gap rate of $O(n^{-1} \log n)$ relative to the optimum within the constrained policy class. Further applications include projected TD(0) and projected stochastic gradient descent. We also extend the analysis to projection of the fast recursion under Euclidean contractivity.

Keywords: Stochastic approximation, two-time-scale methods, projected dynamical systems, controlled Markov chains, Skorokhod map, finite-time concentration, reinforcement learning

1 Introduction

Stochastic approximation algorithms are iterative schemes for solving fixed-point, root-finding, and optimization problems when only noisy observations of the underlying operator are available. They play a central role in stochastic control, reinforcement learning, simulation-based optimization, and adaptive systems. Many modern applications involve coupled recursions evolving on different time scales, with one component adapting rapidly and another changing more slowly. In addition, constraints on the iterates are often enforced through projection steps. This paper studies finite-time behavior of a projected two-time-scale stochastic approximation scheme driven by a controlled Markov chain.

The algorithm consists of a fast recursion x_n and a slow projected recursion θ_n . The slow iterate is projected onto a compact convex polyhedron Θ , while the transition probabilities of the controlled Markov chain are determined by the current value of θ_n . Because the slow variable changes on a smaller time scale, the fast recursion is expected to equilibrate near the fixed point $x^*(\theta_n)$ of the averaged fast-time-scale map. We make this intuition quantitative by proving high-probability bounds on the tracking error $x_n - x^*(\theta_n)$.

The slow recursion is then compared with the solution of an associated projected ODE. This comparison is nonstandard because the projection may make the limiting vector field discontinuous on the boundary of Θ .

We overcome this difficulty by representing each projected path through the Skorokhod map. The Lipschitz property of this map then allows us to control differences between projected paths by differences between their unconstrained driving paths. This yields finite-time concentration bounds for the slow iterate around the projected ODE trajectory. If the associated projected ODE admits a Lipschitz Lyapunov function satisfying a uniform one-block decrease condition along its trajectories, these bounds imply almost-sure convergence of the last-iterate Lyapunov error to zero. Explicit rates follow when the decrease modulus has a power lower bound near zero, with uniform one-block contraction as the linear-decrease special case.

Consider the coupled stochastic recursions

$$x_{n+1} = x_n + \alpha_n [h(Y_n, x_n, \theta_n) - x_n + M_{n+1}], \quad (1.1)$$

$$\theta_{n+1} = \text{proj}_\Theta \{ \theta_n + \beta_n [f(Y_n, x_n, \theta_n) + M'_{n+1}] \}. \quad (1.2)$$

where the step sizes satisfy $\beta_n/\alpha_n \rightarrow 0$ as $n \rightarrow \infty$, Θ is a compact convex polyhedron, and proj_Θ denotes projection onto Θ . This separation of step sizes gives the two-time-scale structure: the x_n -recursion evolves on the faster time scale, while the θ -recursion evolves on the slower time scale. The processes $\{M_n\}$ and $\{M'_n\}$ are martingale-difference noise terms. The sequence $\{Y_n\}$ is a controlled Markov chain taking values in the finite set $\mathcal{Y}$, and its transition kernel depends on the current slow iterate θ_n .

For each $\theta \in \Theta$, let $\mu^{(\theta)}$ be the stationary distribution of the controlled Markov chain induced by θ , and define

$$h^{(\text{av.})}(x; \theta) := \sum_{y \in \mathcal{Y}} \mu^{(\theta)}(y) h(y, x, \theta).$$

The unique fixed point of $h^{(\text{av.})}(\cdot; \theta)$ is denoted by $x^*(\theta)$. We prove that the fast iterate tracks the moving fixed point $x^*(\theta_n)$ with high probability. We then show that the right-continuous step embedding of the slow iterate tracks the solution of the projected ODE

$$\dot{z}(t) = \Pi_\Theta \left(z(t), \sum_{y \in \mathcal{Y}} \mu^{(z(t))}(y) f(y, x^*(z(t)), z(t)) \right).$$

Here $\Pi_\Theta(z, v)$ denotes the projected vector at z in direction v ; its formal definition is given in (4.1).

Using these concentration bounds, we obtain iteration-wise almost-sure last-iterate rates for $\alpha_n = (N_0 + n)^{-\mathfrak{a}}$ and $\beta_n = (N_0 + n)^{-\mathfrak{b}}$, where $1/2 < \mathfrak{a} < \mathfrak{b} \leq 1$. Under uniform one-block contraction and $\mathfrak{b} < 1$, Corollary 8.1 gives joint fast-tracking and slow Lyapunov-error rates with polynomial exponent arbitrarily close to $1/3$, up to a $\sqrt{\log(N_0 + n)}$ factor. At the harmonic endpoint $\mathfrak{b} = 1$, for which $\beta_n = (N_0 + n)^{-1}$, taking $\mathfrak{a} = 2/3$ gives $\|x_n - x^*(\theta_n)\| = O_{\text{a.s.}}((N_0 + n)^{-1/3} \sqrt{\log(N_0 + n)})$. The polynomial exponent governing the slow Lyapunov error is then $\min\{\lambda_T, 1/3\}$, with the precise logarithmic factors given in the same corollary; here λ_T is defined in (8.7).

More generally, suppose that the decrease in the Lyapunov function over one fixed block of projected-ODE time is bounded below by a positive constant times its value at the start of the block raised to a power $p_J > 1$, whenever that value is sufficiently small. Corollary 8.2 gives the optimized $\mathfrak{b} < 1$ rate

$$J(\theta_n) = O_{\text{a.s.}}\left((N_0 + n)^{-1/(4p_J-1)} \{\log(N_0 + n)\}^{1/(2p_J)}\right),$$

whereas its harmonic-endpoint rate is $O_{\text{a.s.}}(\{\log(N_0 + n)\}^{-1/(p_J-1)})$. In Appendices S.1–S.3 of the Online Companion, we specialize the framework to the case where the fast recursion is absent, obtain sharper rate bounds for the remaining projected recursion, and give applications to projected TD(0) and projected SGD.

Additional equilibrium geometry improves these polynomial exponents. If the slow equilibrium set is a strictly attracting box face and the fast equilibrium is constant on that face, Proposition 8.1 gives a joint rate $O_{\text{a.s.}}(n^{-1/2} \log n)$ with $\alpha_n = A \log(N_0 + n)/(N_0 + n)$ and $\beta_n = B/(N_0 + n)$. Actor–critic algorithm from reinforcement learning satisfies these conditions and also achieves a value-gap rate of $O_{\text{a.s.}}(n^{-1} \log n)$.

The projection step introduces a difficulty that is absent in standard finite-time analyses of stochastic approximation. The limiting dynamics are described by a projected differential equation, whose vector field need not be continuous at the boundary of the constraint set. Consequently, standard perturbation arguments for stochastic approximation with Lipschitz limiting vector fields do not directly apply. We therefore follow the ODE method (Borkar [3]): on each finite block of accumulated slow time, we compare the slow step embedding with the projected-ODE solution initialized at the block entrance. We make this projected-ODE comparison quantitative: the Lipschitz continuity of the Skorokhod map on finite time intervals yields explicit path-tracking concentration bounds, and a separate one-block Lyapunov decrease condition converts those bounds into last-iterate rates, with contraction as its linear special case.

1.1 Contributions

Our main contributions are as follows.

- (i) We establish explicit finite-time high-probability concentration bounds showing that the fast iterate x_n tracks the moving fixed point $x^*(\theta_n)$ in projected two-time-scale stochastic approximation driven by an iterate-controlled finite-state Markov chain. Only the averaged fast map is assumed contractive; no contraction is imposed on the slow recursion, which is projected onto the compact convex polyhedron Θ . The bounds separate the effects of martingale noise, Markovian sampling, movement of the fast equilibrium, and time-scale separation; see Proposition 6.1.
- (ii) We prove a finite-horizon supremum-norm concentration bound for the slow iterates around the associated projected-ODE trajectory. We use the Skorokhod map to handle the possible discontinuity of the projected vector field at the boundary; see Theorem 7.1.

- (iii) For the slow iterates, we prove almost-sure convergence to the equilibrium set of the associated projected ODE, assuming that this ODE admits a Lipschitz Lyapunov function that is nonincreasing along trajectories and satisfies a uniform decrease condition over a fixed time horizon $T > 0$. The horizon can be chosen to establish this decrease, without requiring contraction or nonexpansiveness of the reduced slow update map. This goes beyond the reduced slow-map contraction assumptions of Chandak [6], Chandak et al. [9] and the nonexpansiveness assumption of Chandak [5]. The actor-critic application in Section 9 illustrates this distinction: its reduced slow update map need not be nonexpansive, while its projected ODE reaches the restricted-optimal equilibrium face in uniformly bounded time and therefore satisfies one-block Lyapunov contraction for a sufficiently large T . To quantify this convergence, we assume that, for sufficiently small Lyapunov errors, the decrease over time T is bounded below by a positive constant times a power of the current Lyapunov error. This yields explicit optimized rates, with one-block Lyapunov contraction as the linear-decrease special case. Under this contraction condition and polynomial step sizes, the joint fast-tracking and slow Lyapunov-error exponent can be taken arbitrarily close to $1/3$; see Theorems 8.2 and 8.4 and Corollaries 8.1 and 8.2. When the slow equilibrium set is a face of a box constraint, the averaged slow drift points strictly outward in each coordinate fixed on that face, and $x^*(\theta)$ is constant on the face, Proposition 8.1 gives the joint rate $O_{\text{a.s.}}(n^{-1/2} \log n)$ with logarithmically separated step sizes.
- (iv) We apply our results to an actor-critic algorithm in which the current policy controls the Markov sampling process. With logarithmically separated step sizes, the actor value gap relative to the optimum within the constrained policy class is $O_{\text{a.s.}}(n^{-1} \log n)$, and the critic error is $O_{\text{a.s.}}(n^{-1/2} \log n)$. The Online Companion contains the proof and applications to projected TD(0) and projected SGD. It also gives a saddle-point application in Appendix S.18, including examples with vanishing curvature and rotational coupling from a bilinear term. These examples fail both nonexpansiveness of the reduced forward map in any fixed norm and uniform quadratic dissipation, but their projected ODEs satisfy a power Lyapunov decrease condition; see Remark S.18.1. Finally, we extend the analysis to algorithms that project the fast iterate onto a compact convex polyhedron at every step. This makes the fast iterates bounded by construction, removing Assumption 3.7, while retaining contraction of the averaged fast map in the Euclidean norm and bounded noise before projection.

2 Related work

Representative finite-time results include mean-square, and high-probability concentration bounds due to Borkar [2], Chandak et al. [8], Chen et al. [10, 11]. In two-time-scale stochastic approximation (TTSA), one first studies the fast averaged dynamics with the slow variable frozen. The corresponding reduced slow dynamics are then obtained by replacing the fast variable with its equilibrium $x^*(\theta)$. Existing analyses often impose stability or contraction both on this frozen-slow fast subsystem and on the resulting equilibrium-reduced slow subsystem; see, for example, Chandak [6]. In the linear setting with i.i.d. noise, Konda and Tsitsiklis [17] characterize the asymptotic covariance of two-time-scale stochastic approximation, and establish asymptotic normality. Dalal et al. [12] derive a concentration bound for linear two-time-scale stochastic approximation, with applications to reinforcement-learning algorithms. Doan [13] prove almost-sure convergence and an $O(k^{-2/3})$ mean-square bound for nonlinear TTSA under strong-monotonicity assumptions on the frozen-slow fast operator and the equilibrium-reduced slow operator. Chandak et al. [9] obtain an $O(n^{-2/3})$ mean-square rate under arbitrary-norm contractions and Markov noise. Chandak

[6] obtains $O(k^{-a})$ mean-square error for the two-time-scale schedule $\alpha_k \asymp k^{-a}$, $\beta_k \asymp k^{-1}$, for $a \in (1/2, 1)$. These rates use contraction on both reduced operators. Chandak [5] allows a non-expansive slow map and a projected fast recursion, with a residual mean-square rate $O(k^{-1/4+\varepsilon})$, where $\varepsilon > 0$ is arbitrarily small. Classical works on projected stochastic approximation focus on asymptotic convergence, weak convergence, or large-deviation principles; see, for example, Dupuis and Kushner [15], Kushner and Yin [19]. Borowski and Miasojedow [4] repair the gap in Kushner and Yin [19, Theorem 5.2.1] for single-time-scale SA with projection onto a hyperrectangle, proving almost-sure convergence to the stationary set of its projected ODE.

Notation Throughout, $\|\cdot\|$ denotes the Euclidean norm for vectors and the induced Euclidean operator norm for matrices. For vectors, $\|\cdot\|_\infty$ denotes the sup norm. For matrices,

$$\|A\|_\infty := \max_i \sum_j |A(i, j)|$$

denotes the operator norm induced by the vector sup norm, equivalently the maximum absolute row-sum norm. Boldface $\mathbf{0}$ and $\mathbf{1}$ denote, respectively, the zero and all-ones vectors of the appropriate dimension. If $\mathcal{E}$ is an event, then $\mathbf{1}\{\mathcal{E}\}$ is its indicator. If $\mathcal{B}$ is a finite nonempty set, then $\Delta(\mathcal{B})$ denotes the probability simplex on $\mathcal{B}$, viewed as a subset of $\mathbb{R}^{|\mathcal{B}|}$. For a set X and point x , $\text{dist}(x, X)$ denotes the distance from x to X . For a positive deterministic sequence a_n , we write $Z_n = O_{\text{a.s.}}(a_n)$ if, on an event of probability one, there exist a finite nonnegative random constant C and a finite integer-valued random index N such that $|Z_n| \leq Ca_n$ for every $n \geq N$, with absolute value replaced by the Euclidean norm for vector-valued Z_n . For positive deterministic sequences $\{a_n\}$ and $\{b_n\}$, we write $a_n \asymp b_n$ when they are comparable up to positive multiplicative constants for all sufficiently large n ; that is, when $cb_n \leq a_n \leq Cb_n$ eventually for some $0 < c \leq C < \infty$. We write $\mathbb{N} := \{1, 2, \dots\}$ and write $\mathbb{R}_+$ for the set of nonnegative reals. For $d \geq 1$, define

$$\kappa_d := \sup_{\|u\|_\infty \leq 1, u \in \mathbb{R}^d} \|u\|.$$

Thus $\|z\| \leq \kappa_d \|z\|_\infty$ for every $z \in \mathbb{R}^d$.

Sections 3–5 give the model, notation, and proof strategy; Sections 6–8 establish concentration and last-iterate rates; Section 9 presents the actor–critic application; and Section 10 concludes. Selected proofs appear in Appendix A; the remaining proofs, the one-time-scale specialization and applications, the fast-projection extension, and auxiliary results are collected in the Online Companion.

3 Problem Formulation

We study the recursions (1.1)–(1.2) introduced in Section 1. Here $x_n \in \mathbb{R}^{d_x}$ is the fast iterate, $\theta_n \in \Theta \subset \mathbb{R}^{d_\theta}$ is the slow iterate, and Θ is a nonempty compact convex polyhedron. The map $\text{proj}_\Theta(\cdot)$ denotes Euclidean projection onto Θ . The initial slow iterate satisfies $\theta_1 \in \Theta$ almost surely. The process $\{Y_n\}$ is a controlled Markov chain taking values in the finite state space $\mathcal{Y}$, and its transition law depends on the current slow iterate θ_n . The functions $h : \mathcal{Y} \times \mathbb{R}^{d_x} \times \Theta \rightarrow \mathbb{R}^{d_x}$ and $f : \mathcal{Y} \times \mathbb{R}^{d_x} \times \Theta \rightarrow \mathbb{R}^{d_\theta}$ define the fast and slow updates, respectively. The processes $\{M_n\}$ in $\mathbb{R}^{d_x}$ and $\{M'_n\}$ in $\mathbb{R}^{d_\theta}$ are noise processes.

In addition to the time-scale separation stated in Section 1, the step sizes satisfy

$$\sum_n \alpha_n = \sum_n \beta_n = \infty, \quad \sum_n (\alpha_n^2 + \beta_n^2) < \infty.$$

We impose the following standing assumptions.

Assumption 3.1 (Controlled Markov chain). The process $\{Y_n\}$ is a controlled Markov process taking values in the finite state space $\mathcal{Y}$. Let

$$\mathcal{F}_n = \sigma(x_1, \theta_1, \{Y_i\}_{i=1}^n, \{M_i\}_{i=1}^n, \{M'_i\}_{i=1}^n), n = 1, 2, \dots$$

Then

$$\mathbb{P}(Y_{n+1} = y' | \mathcal{F}_n) = p^{(\theta_n)}(Y_n, y'), \quad n \in \mathbb{N}, \quad \text{a.s.},$$

where $\{p^{(\theta)}(y, y')\}_{y, y' \in \mathcal{Y}}$ are the controlled transition probabilities of the Markov chain induced by θ . For each $\theta \in \Theta$, the Markov chain with transition probabilities $p^{(\theta)}$ has a unique stationary distribution, denoted by $\mu^{(\theta)}$.

Assumption 3.2 (Lipschitz transition kernel). There exists $L_p > 0$ such that, for all $i \in \mathcal{Y}$ and all $\theta, \theta' \in \Theta$,

$$\sum_{j \in \mathcal{Y}} |p^{(\theta)}(i, j) - p^{(\theta')}(i, j)| \leq L_p \|\theta - \theta'\|.$$

Thus, the controlled transition kernel is Lipschitz in θ .

This regularity condition is standard in finite-time analyses of stochastic approximation; see, e.g., Chandak et al. [8], Metivier and Priouret [20].

Assumption 3.3 (Martingale-difference noise). The sequences $\{M_n\}$ and $\{M'_n\}$ are martingale-difference sequences with respect to $\{\mathcal{F}_n\}$. Thus,

$$\mathbb{E}(M_{n+1} | \mathcal{F}_n) = \mathbf{0}, \quad \mathbb{E}(M'_{n+1} | \mathcal{F}_n) = \mathbf{0}, \quad \text{a.s.}$$

Moreover, there exist constants $\mathbf{c}_1, \mathbf{c}_2 > 0$ such that, almost surely,

$$\|M_{n+1}\| \leq \mathbf{c}_1 + \mathbf{c}_2 \|x_n\|, \quad \|M'_{n+1}\| \leq \mathbf{c}_1 + \mathbf{c}_2 \|x_n\|, \quad \forall n \in \mathbb{N}. \quad (3.1)$$

Martingale-difference noise assumptions of this form are standard in stochastic approximation; see, e.g., Borkar [3], Kushner and Yin [19].

Assumption 3.4 (Uniform contraction of the averaged fast map). For each $\theta \in \Theta$ define the averaged fast-time-scale map

$$h^{(\text{av.})}(x; \theta) := \sum_{y \in \mathcal{Y}} \mu^{(\theta)}(y) h(y, x, \theta), \quad x \in \mathbb{R}^{d_x}. \quad (3.2)$$

For a fixed θ , $h^{(\text{av.})}(x; \theta)$ is the mean fast-update map obtained by freezing the slow parameter and averaging $h(y, x, \theta)$ with respect to the stationary distribution $\mu^{(\theta)}$ of the resulting Markov chain. Thus, $h^{(\text{av.})}(x; \theta) - x$ is the corresponding averaged drift of the fast recursion. There exists $\gamma \in [0, 1)$ such that

$$\|h^{(\text{av.})}(x; \theta) - h^{(\text{av.})}(x'; \theta)\| \leq \gamma \|x - x'\|, \quad x, x' \in \mathbb{R}^{d_x}, \theta \in \Theta.$$

The unique fixed point of $h^{(\text{av.})}(\cdot; \theta)$ is denoted by $x^*(\theta)$, so that

$$h^{(\text{av.})}(x^*(\theta); \theta) = x^*(\theta).$$

Remark 3.1 (Fast-scale contraction). Assumption 3.4 imposes a contraction condition only on the averaged fast-time-scale map. No contractivity assumption on a discrete slow update map is needed for the concentration and path-tracking results; the slow dynamics are instead characterized through the projected ODE studied later. The constant γ may be replaced by a parameter-dependent contraction factor $\gamma(\theta)$, provided $\sup_{\theta \in \Theta} \gamma(\theta) < 1$.

The extension to an arbitrary fixed norm for the fast coordinate is recorded in Remark S.5.1 of the Online Companion.

Assumption 3.5 (Lipschitz update functions on bounded fast-state sets). For every $0 \leq R < \infty$, there exist finite constants $L_f(R)$, $L_{h,x}(R)$, and $L_{h,\theta}(R)$ such that, for all $y \in \mathcal{Y}$, $\theta, \tilde{\theta} \in \Theta$, and $x, \tilde{x} \in \mathbb{R}^{d_x}$ satisfying $\|x\|, \|\tilde{x}\| \leq R$,

$$\|f(y, x, \theta) - f(y, \tilde{x}, \tilde{\theta})\| \leq L_f(R) (\|x - \tilde{x}\| + \|\theta - \tilde{\theta}\|),$$

and

$$\|h(y, x, \theta) - h(y, \tilde{x}, \tilde{\theta})\| \leq L_{h,x}(R) \|x - \tilde{x}\| + L_{h,\theta}(R) \|\theta - \tilde{\theta}\|.$$

The constants may depend on R , but not on y , x , $\tilde{x}$, θ , or $\tilde{\theta}$ within the displayed sets. The moduli may be chosen nondecreasing in R .

For $0 \leq R < \infty$, Assumption 3.5, finiteness of $\mathcal{Y}$, and compactness of Θ imply

$$B(R) := \sup_{\substack{y \in \mathcal{Y}, \theta \in \Theta \\ \|x\| \leq R}} \max\{\|h(y, x, \theta)\|, \|f(y, x, \theta)\|\} < \infty.$$

Write $B_0 := B(0)$.

Assumption 3.6 (Uniform hitting-time bound). There exists a reference state $y^* \in \mathcal{Y}$ and $\mathfrak{c}_5 < \infty$ such that, for every $\theta \in \Theta$ and every initial state $y \in \mathcal{Y}$,

$$\mathbb{E}_y^{(\theta)}(\tau_{y^*}) \leq \mathfrak{c}_5,$$

where $\tau_{y^*} := \inf\{n \geq 1 : Y_n = y^*\}$ for the Markov chain with transition probabilities $p^{(\theta)}$. The superscript (θ) on $\mathbb{E}_y^{(\theta)}$ indicates that the chain uses the fixed transition kernel $p^{(\theta)}$ at every step, while the subscript y specifies the initial state $Y_0 = y$.

This condition provides uniform control of the Poisson-equation solutions used to decompose the Markov noise; see Lemmas 6.1 and 7.1. The next lemma records two regularity consequences of the preceding controlled-Markov assumptions. Together, Assumptions 3.2 and 3.6 imply Lipschitz continuity of the stationary-distribution map. Combined with Assumptions 3.5 and 3.4, this also gives Lipschitz continuity of $x^*(\cdot)$.

Lemma 3.1. *The stationary distribution map $\theta \mapsto \mu^{(\theta)}$ and the fixed-point map $\theta \mapsto x^*(\theta)$ are Lipschitz on Θ . That is, there exist constants $L_\mu, L_{fp} \geq 0$ such that*

$$\|\mu^{(\theta)} - \mu^{(\theta')}\|_1 \leq L_\mu \|\theta - \theta'\|, \quad \|x^*(\theta) - x^*(\theta')\| \leq L_{fp} \|\theta - \theta'\|, \quad \theta, \theta' \in \Theta.$$

Here, for any signed measure ν on $\mathcal{Y}$,

$$\|\nu\|_1 := \sum_{y \in \mathcal{Y}} |\nu(y)|.$$

In particular, $\sup_{\theta \in \Theta} \|x^*(\theta)\| < \infty$.

The proof is given in Appendix S.4 of the Online Companion. We henceforth write

$$\mathfrak{c}_8 := \sup_{\theta \in \Theta} \|x^\star(\theta)\| \leq \frac{B_0}{1 - \gamma} < \infty.$$

Assumption 3.7 (Stability of the fast iterate). There exists a deterministic constant $\mathfrak{c}_6 < \infty$ such that

$$\sup_{n \in \mathbb{N}} \|x_n\| \leq \mathfrak{c}_6 \text{ a.s.}$$

Contraction of the averaged fast map does not by itself keep the fast iterates bounded: individual sampled updates may still produce large excursions. Assumption 3.7 ensures that the martingale increments, Poisson-equation solutions, and drift perturbations used throughout the concentration proof are uniformly bounded. It can be verified by finding a deterministic bounded set that contains the initial iterate and is preserved by the recursion, as illustrated in the Online Companion. Alternatively, projecting the fast iterate onto a compact convex polyhedron at every step ensures boundedness by construction. Appendix S.9 develops this extension, removing the need for Assumption 3.7.

Set

$$B_6 := B(\mathfrak{c}_6), \quad B_8 := B(\mathfrak{c}_8), \quad R_{\text{loc}} := \max\{\mathfrak{c}_6, \mathfrak{c}_8\}.$$

Hereafter, L_f , $L_{h,x}$, and $L_{h,\theta}$ denote, respectively, the constants $L_f(R_{\text{loc}})$, $L_{h,x}(R_{\text{loc}})$, and $L_{h,\theta}(R_{\text{loc}})$ from Assumption 3.5. The ball of radius R_{loc} contains every fast iterate and every fixed point $x^\star(\theta)$.

Unless an alternative schedule is stated explicitly, we use polynomial step sizes with a positive-integer shift $N_0 \geq 1$:

$$\alpha_n = \frac{1}{(n + N_0)^{\mathfrak{a}}}, \beta_n = \frac{1}{(n + N_0)^{\mathfrak{b}}}, n \in \mathbb{N},$$

where $\mathfrak{a} \in (0, 1)$ and $\mathfrak{b} \in (0, 1]$. The exponent $\mathfrak{a}$ corresponds to the fast recursion and $\mathfrak{b}$ to the slow recursion. Since the slow step size must decay faster, we require $\mathfrak{a} < \mathfrak{b}$. The hyperparameters $\mathfrak{a}, \mathfrak{b}$ satisfy the following.

Condition 3.1 (Step-size exponents). The step-size exponents satisfy

$$\frac{1}{2} < \mathfrak{a} < \mathfrak{b} \leq 1.$$

The shift N_0 is part of the algorithm. For the polynomial two-time-scale schedule above, we assume throughout, unless explicitly stated otherwise, that

$$(N_0 + 1)^{1-\mathfrak{a}} \geq 4. \tag{3.3}$$

This choice ensures that the step-size bounds needed in the analysis hold from index $n = 1$; see Appendix S.16 of the Online Companion.

Alternatively, we can begin the analysis after a finite number of iterations. In this case, we take $N_0 = 1$ and run the recursion from $n = 1$, but begin the block construction at the first index $n_0 \geq 1$ satisfying

$$(n_0 + 1)^{1-\mathfrak{a}} \geq 4.$$

Appendix S.16 of the Online Companion verifies that the analysis applies from $n = n_0$, with convergence rates expressed in terms of $n + 1$.

4 Continuous-time step embeddings and projected dynamical systems

We first introduce the cumulative time grid and the right-continuous step paths used in the projected-ODE analysis. For $n \geq m$, write

$$a(m, n) := \sum_{k=m}^{n-1} \beta_k, \quad b(n) := \sum_{k \geq n} \beta_k^2,$$

and

$$\chi(m, n) := \begin{cases} \prod_{k=m}^n (1 - \alpha_k), & m \leq n, \\ 1, & m > n. \end{cases}$$

For $\ell \geq 1$, define the cumulative time grid associated with the θ -recursion by

$$t(\ell) := \sum_{k=1}^{\ell-1} \beta_k.$$

The empty sum is zero, so $t(1) = 0$. Moreover,

$$t(n+1) - t(n) = \beta_n, \quad t(n) - t(m) = a(m, n), \quad n \geq m \geq 1.$$

Define the right-continuous step embedding of the slow iterate by

$$\theta^\circ(r) := \theta_n, \quad r \in [t(n), t(n+1)).$$

Thus,

$$\theta^\circ(t(n)) = \theta_n.$$

By construction, the path θ° is right-continuous with left limits (càdlàg) and piecewise constant. Fix a slow-time block horizon $T > 0$. Let n_0 be the starting index of the analysis specified in Section 3, with $n_0 = 1$ under the default choice of N_0 . Define recursively

$$n_m := \min\{n > n_{m-1} : t(n) \geq t(n_{m-1}) + T\}, \quad m \geq 1.$$

For $m \geq 0$, set

$$T_m := t(n_m), \quad \mathcal{K}_m := \{n_m, n_m + 1, \dots, n_{m+1} - 1\},$$

and write

$$K_m := n_{m+1} - n_m, \quad H_m := T_{m+1} - T_m,$$

We call $\mathcal{K}_m$ the m th discrete-time block; its associated slow-time interval is $[T_m, T_{m+1}]$. A local index k measures displacement from the block entrance n_m , with $k = K_m$ corresponding to the right endpoint n_{m+1} .

For $0 \leq \ell \leq K_m$, define the shifted slow-time grid by

$$t^{(m)}(\ell) := t(n_m + \ell) - t(n_m).$$

Then

$$t^{(m)}(0) = 0, \quad t^{(m)}(K_m) = H_m,$$

and

$$t^{(m)}(\ell + 1) - t^{(m)}(\ell) = \beta_{n_m + \ell}.$$

For any discrete-time sequence $\{u_n\}_{n \geq 1}$, write

$$u_\ell^{(m)} := u_{n_m + \ell}.$$

For any sequence $\{q_k^{(m)}\}_{0 \leq k \leq K_m}$ indexed by the local position k relative to the entrance of block m , define its right-continuous block step path by

$$q^{(m), \circ}(r) := q_k^{(m)}, \quad r \in [t^{(m)}(k), t^{(m)}(k+1)), \quad 0 \leq k < K_m,$$

and set

$$q^{(m), \circ}(H_m) := q_{K_m}^{(m)}.$$

Let Γ denote the normal-reflection Skorokhod map on Θ , and let L_{Sk} denote a finite Lipschitz constant for this map, as stated in Theorem S.13.1 of the Online Companion. When a block path is passed to the Skorokhod map, it is extended constantly beyond H_m ; this does not affect any block estimate. In particular,

$$\theta^{(m), \circ}(r) = \theta^\circ(T_m + r), \quad 0 \leq r \leq H_m.$$

We use parenthesized superscripts to label shifted objects. Thus, $u_\ell^{(m)}$ denotes a discrete-time sequence reindexed at the block entrance n_m , whereas $z^{(T_m)}$ denotes the projected-ODE trajectory initialized at the slow-time anchor T_m .

4.1 Projected dynamical systems

Following Dupuis and Nagurney [16], let $C \subset \mathbb{R}^d$ be a nonempty closed convex set, and let proj_C denote Euclidean projection onto C . For $z \in C$ and $v \in \mathbb{R}^d$, define

$$\Pi_C(z, v) := \lim_{\varepsilon \downarrow 0} \frac{\text{proj}_C(z + \varepsilon v) - z}{\varepsilon}. \quad (4.1)$$

For $\varepsilon > 0$, also define

$$\Pi_{C, \varepsilon}(z, v) := \frac{\text{proj}_C(z + \varepsilon v) - z}{\varepsilon}. \quad (4.2)$$

We omit the set subscript when it is clear from the context.

Definition 4.1 (Dupuis and Nagurney [16]). Let $F : C \rightarrow \mathbb{R}^d$. A function $z : [0, \infty) \rightarrow C$ is a solution of $\dot{z} = \Pi_C(z, F(z))$ if it is absolutely continuous and

$$\dot{z}(t) = \Pi_C(z(t), F(z(t)))$$

for almost every $t \geq 0$.

For the slow recursion, we apply these definitions with $C = \Theta$ and $d = d_\theta$.

5 Concentration proof sketch and overview

The concentration arguments are developed in Sections 6 and 7; selected proofs appear in Appendix A and supporting proofs in the Online Companion. Since the fast recursion is contractive after averaging, the first step of the analysis is to show that x_n tracks the moving fixed point $x^*(\theta_n)$ with high probability. This allows the slow recursion to be compared with an auxiliary projected recursion in which x_n is replaced by $x^*(\theta_n)$. After averaging over the controlled Markov chain, the limiting slow dynamics are described by the differential equation

$$\dot{z} = \Pi_{\Theta}(z(t), f^{(\text{av.})}(z(t); z(t))), \quad (5.1)$$

where, for $\theta, \theta' \in \Theta$, we define

$$f^{(\text{av.})}(\theta; \theta') := \sum_{y \in \mathcal{Y}} \mu^{(\theta')}(y) f(y, x^*(\theta), \theta). \quad (5.2)$$

In $f^{(\text{av.})}(\theta; \theta')$, the first argument θ is substituted into both the fixed-point map $x^*(\cdot)$ and the parameter coordinate of f , while the second argument θ' determines the stationary distribution used to average over y . We refer to $f^{(\text{av.})}$ as the two-parameter averaged field. Its evaluation is called diagonal when the two arguments agree and off-diagonal otherwise.

We first establish a high-probability bound on the fast tracking error $\|x_n - x^*(\theta_n)\|$; see Proposition 6.1. Using this estimate, we then obtain a concentration bound on the maximum deviation of the embedding of the slow iterates from the projected-ODE trajectory over each finite slow-time block; see Theorem 7.1.

For $\tau \geq 0$, let $z^{(\tau)}(\cdot)$ denote the solution to (5.1) satisfying

$$z^{(\tau)}(0) = \theta^\circ(\tau).$$

In particular, on the m th block, we compare

$$\theta^{(m), \circ}(r) = \theta^\circ(T_m + r)$$

with $z^{(T_m)}(r)$ for $0 \leq r \leq H_m$.

Our analysis has three main components. First, Section 6 establishes a high-probability tracking bound for the fast iterate x_n around the moving equilibrium $x^*(\theta_n)$. Second, Section 7 compares the step embedding θ° with the projected ODE over finite time intervals. Third, Section 8 combines this finite-time tracking estimate with a Lipschitz Lyapunov function satisfying a uniform one-block decrease condition for the projected ODE to obtain almost-sure convergence. Explicit rates follow when, for sufficiently small Lyapunov errors, the decrease over a fixed time horizon is bounded below by a positive constant times a power of the current error. One-block Lyapunov contraction is the linear-decrease special case.

The main technical obstacle is the projection. The projected vector field in the limiting ODE may be discontinuous on the boundary of Θ , so standard stochastic-approximation concentration arguments based on Lipschitz ODE fields do not apply directly. We instead use right-continuous step embeddings of the discrete projected recursions and their unconstrained drivers. More precisely, each projected step path is exactly the image of its corresponding unconstrained driving path under the Skorokhod map. The finite-horizon supremum-norm Lipschitz property of that map then controls the difference between two projected paths through the difference between their unconstrained drivers.

6 Blockwise concentration of the fast iterate

We analyze the fast recursion on the slow-time blocks introduced in Section 4. On block m , our goal is to bound the *fast tracking error*, the distance between the fast iterate and its moving equilibrium:

$$\|x_{n_m+k} - x^*(\theta_{n_m+k})\|, \quad 0 \leq k \leq K_m.$$

We bound this error through two comparisons: first, between the original fast recursion and a block-restarted averaged recursion; second, between this averaged recursion and the moving equilibrium.

Fix $m \geq 0$. For $0 \leq k \leq K_m$, write

$$x_k^{(m)} := x_{n_m+k}, \quad \theta_k^{(m)} := \theta_{n_m+k}, \quad \alpha_k^{(m)} := \alpha_{n_m+k}, \quad \beta_k^{(m)} := \beta_{n_m+k}.$$

On block m , we compare the original fast iterate with an averaged recursion initialized at the same value x_{n_m} and driven by the same slow-parameter sequence $\{\theta_k^{(m)}\}_{0 \leq k \leq K_m}$. Consequently, the difference between these two recursions vanishes at the block entrance and records only the explicit martingale increments and the Markov noise terms

$$h(Y_{n_m+k}, x_{n_m+k}, \theta_k^{(m)}) - h^{(\text{av.})}(x_{n_m+k}; \theta_k^{(m)})$$

accumulated within the block. The discrepancy already present at time n_m enters through the distance between the averaged comparison recursion and the moving fixed point $x^*(\theta_k^{(m)})$.

6.1 Block-restarted averaged recursion

Define the block-restarted averaged recursion $\{\tilde{x}_k^{(m)}\}_{0 \leq k \leq K_m}$ by

$$\tilde{x}_0^{(m)} := x_{n_m},$$

and, for $0 \leq k < K_m$,

$$\tilde{x}_{k+1}^{(m)} = \tilde{x}_k^{(m)} + \alpha_k^{(m)} \left[h^{(\text{av.})}(\tilde{x}_k^{(m)}; \theta_k^{(m)}) - \tilde{x}_k^{(m)} \right]. \quad (6.1)$$

Since $\|h^{(\text{av.})}(0; \theta)\| \leq B_0$, contraction gives

$$\|\tilde{x}_{k+1}^{(m)}\| \leq \{1 - (1 - \gamma)\alpha_k^{(m)}\} \|\tilde{x}_k^{(m)}\| + \alpha_k^{(m)} B_0.$$

Since $\tilde{x}_0^{(m)} = x_{n_m}$, induction gives

$$\|\tilde{x}_k^{(m)}\| \leq R_{\text{aux}} := \max \left\{ \mathfrak{c}_6, \frac{B_0}{1 - \gamma} \right\}.$$

The auxiliary iterates need not lie in the ball of radius R_{loc} . At these iterates, we use the global contraction of the averaged map; no bounded-set estimate for the unaveraged maps h or f is needed. The recursion (6.1) applies the averaged map $h^{(\text{av.})}$ from (3.2) to its own state $\tilde{x}_k^{(m)}$, while using the same slow-parameter sequence $\{\theta_k^{(m)}\}_{0 \leq k \leq K_m}$. Thus it contains neither the explicit noise increment M_{n_m+k+1} nor the sampled Markov state Y_{n_m+k} . It remains random only through its block initial value and the random sequence $\{\theta_k^{(m)}\}$.

To quantify the first comparison—the discrepancy between the original fast recursion and this averaged recursion—we define for $0 \leq k \leq K_m$,

$$D_k^{(m)} := x_k^{(m)} - \tilde{x}_k^{(m)}.$$

By construction, $D_0^{(m)} = 0$.

6.2 Poisson decomposition and local fast fluctuations

The Markov noise

$$h(Y_n, x_n, \theta_n) - h^{(\text{av.})}(x_n; \theta_n)$$

is generally not a martingale difference because Y_n is Markovian. The Poisson equation is used to decompose this error into a martingale-difference term and a difference between two successive-state evaluations of the Poisson solution. Since the sampling error enters the fast update multiplied by α_n , iterating the recursion produces weighted partial sums of the two terms. We control the martingale contribution using a concentration inequality and bound the remaining Poisson residual by summation by parts, using the slow variation of (x_n, θ_n) .

Definition 6.1 (Poisson equation for the fast recursion). Fix $x \in \mathbb{R}^{d_x}$ and $\theta \in \Theta$. For the Markov chain on $\mathcal{Y}$ with transition matrix $p^{(\theta)}$, consider the Poisson equation

$$u(y) - \sum_{i \in \mathcal{Y}} p^{(\theta)}(y, i) u(i) = h(y, x, \theta) - h^{(\text{av.})}(x; \theta), \quad y \in \mathcal{Y}. \quad (6.2)$$

We use the particular solution, as in Chandak et al. [8]

$$u_h^{(x, \theta)}(y) := \mathbb{E}^{(\theta)} \left[\sum_{r=0}^{\tau_{y^*}-1} (h(Y_r, x, \theta) - h^{(\text{av.})}(x; \theta)) \mid Y_0 = y \right], \quad (6.3)$$

where τ_{y^*} is the hitting time of the reference state y^* from Assumption 3.6. The superscript (θ) on $\mathbb{E}^{(\theta)}$ indicates that the controlled Markov chain (Y_r) is run with transition matrix $p^{(\theta)}$; the arguments x and θ in the summand are held fixed as the chain evolves.

The expectation in (6.3) is finite by Assumption 3.6. Moreover, by the definition of $h^{(\text{av.})}$,

$$\sum_{y \in \mathcal{Y}} \mu^{(\theta)}(y) \{h(y, x, \theta) - h^{(\text{av.})}(x; \theta)\} = \mathbf{0}.$$

Lemma S.15.7 of the Online Companion, applied coordinatewise, therefore gives $u_h^{(x, \theta)}(y^*) = \mathbf{0}$. Conditioning on the first transition then shows that (6.3) solves (6.2).

Let $C_{h, \text{Pois.}} < \infty$ denote the Poisson-sensitivity constant from Lemma S.10.1. In particular, whenever $\|x\|, \|x'\| \leq \mathfrak{c}_6$, $\theta, \theta' \in \Theta$, and $y \in \mathcal{Y}$,

$$\left\| u_h^{(x, \theta)}(y) - u_h^{(x', \theta')}(y) \right\| \leq C_{h, \text{Pois.}} (\|x - x'\| + \|\theta - \theta'\|).$$

Lemma 6.1 (Poisson decomposition). *Define*

$$M''_{n+1} := u_h^{(x_n, \theta_n)}(Y_{n+1}) - \sum_{y \in \mathcal{Y}} p^{(\theta_n)}(Y_n, y) u_h^{(x_n, \theta_n)}(y), \quad n \geq 1. \quad (6.4)$$

Then $\{M''_{n+1}\}_{n \geq 1}$ is a martingale-difference sequence with respect to $\{\mathcal{F}_n\}$, and, almost surely,

$$h(Y_n, x_n, \theta_n) - h^{(\text{av.})}(x_n; \theta_n) = M''_{n+1} + u_h^{(x_n, \theta_n)}(Y_n) - u_h^{(x_n, \theta_n)}(Y_{n+1}). \quad (6.5)$$

The sequence $\{M''_{n+1}\}$ is defined on the original time axis and therefore does not carry a block index. On block m , the relevant increments are simply

$$M''_{n_m+k+1}, \quad 0 \leq k < K_m.$$

For later use, define

$$\begin{aligned}\Delta_{n+1}^h &:= u_h^{(x_n, \theta_n)}(Y_n) - u_h^{(x_n, \theta_n)}(Y_{n+1}), \\ \Xi_{n+1} &:= M_{n+1} + M_{n+1}'', \quad n \geq 1.\end{aligned}$$

Then (6.5) becomes

$$h(Y_n, x_n, \theta_n) - h^{(\text{av.})}(x_n; \theta_n) + M_{n+1} = \Xi_{n+1} + \Delta_{n+1}^h.$$

Set

$$C_{\text{mg}}^{(x)} := \mathbf{c}_1 + \mathbf{c}_2 \mathbf{c}_6 + 4 \mathbf{c}_5 \mathbf{B}_6, \quad c_{\text{mg}}^x := \frac{1}{4 \kappa_{d_x}^2 (C_{\text{mg}}^{(x)})^2}.$$

For $0 \leq k \leq K_m$, define the local weighted martingale sum using $\alpha_j^{(m)} = \alpha_{n_m+j}$, by

$$\mathcal{M}_{m,k}^x := \sum_{j=0}^{k-1} \alpha_j^{(m)} \chi(n_m + j + 1, n_m + k - 1) \Xi_{n_m+j+1}, \quad (6.6)$$

with $\mathcal{M}_{m,0}^x := \mathbf{0}$. For $\delta > 0$, define the blockwise fast-martingale event

$$\mathcal{G}_x^{(m)}(\delta) := \left\{ \max_{0 \leq k \leq K_m} \kappa_{d_x} \left\| \mathcal{M}_{m,k}^x \right\|_{\infty} < \delta \right\}. \quad (6.7)$$

Lemma 6.2 (Blockwise fast-martingale tail bound). *For every $m \geq 0$ and $\delta > 0$,*

$$\mathbb{P} \left(\left[\mathcal{G}_x^{(m)}(\delta) \right]^c \right) \leq 2 d_x K_m \exp \left\{ - \frac{c_{\text{mg}}^x \delta^2}{\alpha_{n_m}} \right\}. \quad (6.8)$$

Equivalently,

$$\mathbb{P} \left(\left[\mathcal{G}_x^{(m)}(\delta) \right]^c \right) \leq 2 d_x K_m \exp \left\{ - c_{\text{mg}}^x \delta^2 (N_0 + n_m)^a \right\}.$$

6.3 Comparison with the block-restarted averaged recursion

The successive-state term in the Poisson decomposition produces a stepsize-weighted telescoping residual. Summation by parts and the slow variation of (x_n, θ_n) bound this residual by $O(\alpha_{n_m})$. Contraction then converts the martingale and residual terms into the following comparison. The residual calculation remains in Appendix S.5; see Sections A.2 and A.3 for the comparison proofs.

Lemma 6.3 (Local comparison with the averaged recursion). *There exists $C_{\text{aux}} < \infty$, depending only on the problem parameters, such that, on $\mathcal{G}_x^{(m)}(\delta)$,*

$$\left\| x_k^{(m)} - \tilde{x}_k^{(m)} \right\| \leq C_{\text{aux}} (\delta + \alpha_{n_m}), \quad 0 \leq k \leq K_m. \quad (6.9)$$

6.4 Tracking the moving fixed point

We next compare $\tilde{x}_k^{(m)}$ with the moving fixed point $x^*(\theta_k^{(m)})$. Set $q := 1 - \gamma$ and, for $0 \leq k \leq K_m$, define

$$\mathcal{W}_m(k) := \prod_{\ell=0}^{k-1} \left(1 - q \alpha_{\ell}^{(m)} \right),$$

with the empty product equal to one. Thus $\mathcal{W}_m(k)$ is the contraction accumulated from the block entrance to local index k .

Lemma 6.4 (Moving-equilibrium comparison on a block). *There exists $C_{\text{mov}} < \infty$, depending only on the problem parameters, such that, almost surely, for every $m \geq 0$ and $0 \leq k \leq K_m$,*

$$\begin{aligned} \left\| \tilde{x}_k^{(m)} - x^*(\theta_k^{(m)}) \right\| &\leq \mathcal{W}_m(k) \|x_{n_m} - x^*(\theta_{n_m})\| \\ &\quad + \frac{C_{\text{mov}}}{q} \frac{\beta_{n_m}}{\alpha_{n_m}}. \end{aligned} \quad (6.10)$$

The two terms in (6.10) are, respectively, the contracted block-entry error and the deterministic tracking lag of order β_{n_m}/α_{n_m} .

6.5 Blockwise fast tracking

Finally, we combine the comparisons in Lemmas 6.3 and 6.4 using the triangle inequality. This gives pointwise, stepsize-weighted, and block-endpoint bounds for the fast tracking error. The block-endpoint estimate can then be iterated over successive blocks.

Proposition 6.1 (Blockwise fast tracking). *Let*

$$\varepsilon_n^{(x)} := \|x_n - x^*(\theta_n)\|.$$

There exists $C_x < \infty$, depending only on the problem parameters, such that, on $\mathcal{G}_x^{(m)}(\delta)$,

$$\varepsilon_{n_m+k}^{(x)} \leq \mathcal{W}_m(k) \varepsilon_{n_m}^{(x)} + C_x \left(\delta + \alpha_{n_m} + \frac{\beta_{n_m}}{\alpha_{n_m}} \right), \quad 0 \leq k \leq K_m. \quad (6.11)$$

Moreover, there exists $C_{x,T} < \infty$, depending only on T and the problem parameters, such that, on the same event,

$$\sum_{r=n_m}^{n_{m+1}-1} \beta_r \|x_r - x^*(\theta_r)\| \leq C_{x,T} \left(\delta + \alpha_{n_m} + \frac{\beta_{n_m}}{\alpha_{n_m}} \right). \quad (6.12)$$

At the block endpoint,

$$\varepsilon_{n_{m+1}}^{(x)} \leq \mathcal{W}_m(K_m) \varepsilon_{n_m}^{(x)} + C_x \left(\delta + \alpha_{n_m} + \frac{\beta_{n_m}}{\alpha_{n_m}} \right). \quad (6.13)$$

Proof. See Appendix A.1. □

For $m \geq 0$ and $\delta > 0$, set

$$g_m^{(x)}(\delta) := \delta + \alpha_{n_m} + \frac{\beta_{n_m}}{\alpha_{n_m}}. \quad (6.14)$$

Corollary 6.1 (Fast tracking across successive blocks). *For every $m \geq 1$ and every choice of positive real numbers $\delta_0, \dots, \delta_{m-1}$, on $\bigcap_{j=0}^{m-1} \mathcal{G}_x^{(j)}(\delta_j)$,*

$$\varepsilon_{n_m}^{(x)} \leq \left(\prod_{\ell=0}^{m-1} \mathcal{W}_\ell(K_\ell) \right) \varepsilon_{n_0}^{(x)} + C_x \sum_{j=0}^{m-1} \left(\prod_{\ell=j+1}^{m-1} \mathcal{W}_\ell(K_\ell) \right) g_j^{(x)}(\delta_j). \quad (6.15)$$

Furthermore,

$$\mathbb{P} \left(\bigcup_{j=0}^{m-1} \left[\mathcal{G}_x^{(j)}(\delta_j) \right]^c \right) \leq \sum_{j=0}^{m-1} 2d_x K_j \exp \left\{ -\frac{c_{\text{mg}}^x \delta_j^2}{\alpha_{n_j}} \right\}. \quad (6.16)$$

7 Concentration of the slow iterate

We now establish a blockwise high-probability comparison between the step embedding of the slow iterate and the projected ODE (5.1). On each block, we introduce an auxiliary projected recursion and decompose the comparison into two stages. First, we compare the original slow recursion with the auxiliary recursion, thereby isolating the fast-tracking error, the martingale noise in the slow recursion, and the Markov noise. Second, we compare the auxiliary recursion with the projected ODE. This second comparison controls the time-grid discretization error and the mismatch arising because the auxiliary recursion evaluates the first and second arguments of the averaged field at its own current state and the original slow state, respectively, whereas the ODE evaluates both arguments at its current state.

Throughout this section, “auxiliary recursion” refers to the slow projected recursion introduced below. It is distinct from the block-restarted averaged fast recursion of Section 6.

Since the projection operator is nonlinear, both comparisons are carried out through the corresponding unconstrained driving paths. Bounds on the driving paths are then transferred to the projected paths using the Lipschitz property of the Skorokhod map.

Fix $m \geq 0$ and use the block notation introduced in Section 4. In particular, for $0 \leq n \leq K_m$, write

$$\theta_n^{(m)} := \theta_{n_m+n}, \quad x_n^{(m)} := x_{n_m+n}, \quad Y_n^{(m)} := Y_{n_m+n}.$$

For $0 \leq n \leq K_m - 1$, define

$$\beta_n^{(m)} := \beta_{n_m+n}, \quad (M')_{n+1}^{(m)} := M'_{n_m+n+1}.$$

For $0 \leq n \leq K_m$, also define the block-shifted filtration and the fast-tracking error by

$$\mathcal{F}_n^{(m)} := \mathcal{F}_{n_m+n}, \quad e_n^{(m)} := x_n^{(m)} - x^\star(\theta_n^{(m)}).$$

Throughout this section, the parenthesized superscript (m) denotes reindexing relative to the m th block; it is neither an exponent nor an abbreviation for “martingale.”

Our goal is to control

$$\sup_{0 \leq t \leq H_m} \left\| \theta^{(m),\circ}(t) - z^{(T_m)}(t) \right\|,$$

where $z^{(T_m)}$ solves (5.1) with

$$z^{(T_m)}(0) = \theta^\circ(T_m) = \theta_{n_m}.$$

7.1 Auxiliary projected recursion

The shifted slow recursion can be written as

$$\theta_{n+1}^{(m)} = \theta_n^{(m)} + \beta_n^{(m)} \Pi_{\Theta, \beta_n^{(m)}} \left(\theta_n^{(m)}, f(Y_n^{(m)}, x^\star(\theta_n^{(m)}), \theta_n^{(m)}) + \mathcal{E}_{f,n}^{(m)} + (M')_{n+1}^{(m)} \right),$$

where

$$\mathcal{E}_{f,n}^{(m)} := f(Y_n^{(m)}, x_n^{(m)}, \theta_n^{(m)}) - f(Y_n^{(m)}, x^\star(\theta_n^{(m)}), \theta_n^{(m)}).$$

Thus, $\mathcal{E}_{f,n}^{(m)}$ is the drift error caused by the fast iterate not being at its moving fixed point.

Define the auxiliary projected recursion by setting

$$\tilde{\theta}_0^{(m)} = \theta_0^{(m)}$$

and, for $0 \leq n \leq K_m - 1$,

$$\theta_{n+1}^{(m)} = \tilde{\theta}_n^{(m)} + \beta_n^{(m)} \Pi_{\Theta, \beta_n^{(m)}} \left(\tilde{\theta}_n^{(m)}, f^{(\text{av.})}(\tilde{\theta}_n^{(m)}; \theta_n^{(m)}) \right). \quad (7.1)$$

Equivalently,

$$\tilde{\theta}_{n+1}^{(m)} = \text{proj}_{\Theta} \left(\tilde{\theta}_n^{(m)} + \beta_n^{(m)} f^{(\text{av.})}(\tilde{\theta}_n^{(m)}; \theta_n^{(m)}) \right).$$

In (7.1), the two-parameter averaged field is evaluated off diagonal:

$$f^{(\text{av.})}(\tilde{\theta}_n^{(m)}; \theta_n^{(m)}) = \sum_{y \in \mathcal{Y}} \mu^{(\theta_n^{(m)})}(y) f(y, x^*(\tilde{\theta}_n^{(m)}), \tilde{\theta}_n^{(m)}).$$

The auxiliary iterate $\tilde{\theta}_n^{(m)}$ determines where the drift is evaluated. The original slow iterate $\theta_n^{(m)}$ determines the averaging distribution $\mu^{(\theta_n^{(m)})}$, since the Markov chain uses the transition kernel $p^{(\theta_n^{(m)})}$ at this iteration. Replacing the second argument by $\tilde{\theta}_n^{(m)}$ gives the diagonal field used by the projected ODE. The error introduced by this replacement is bounded by a constant times $\|\tilde{\theta}_n^{(m)} - \theta_n^{(m)}\|$, which we control by comparing the original and auxiliary recursions.

Before comparing the projected paths, we compare the unprojected update directions that drive their Skorokhod representations. The difference between the original and auxiliary update directions has the exact decomposition

$$\begin{aligned} & f(Y_n^{(m)}, x_n^{(m)}, \theta_n^{(m)}) + (M')_{n+1}^{(m)} - f^{(\text{av.})}(\tilde{\theta}_n^{(m)}; \theta_n^{(m)}) \\ &= [f(Y_n^{(m)}, x_n^{(m)}, \theta_n^{(m)}) - f(Y_n^{(m)}, x^*(\theta_n^{(m)}), \theta_n^{(m)})] \\ &+ [f(Y_n^{(m)}, x^*(\theta_n^{(m)}), \theta_n^{(m)}) - f^{(\text{av.})}(\theta_n^{(m)}; \theta_n^{(m)})] \\ &+ [f^{(\text{av.})}(\theta_n^{(m)}; \theta_n^{(m)}) - f^{(\text{av.})}(\tilde{\theta}_n^{(m)}; \theta_n^{(m)})] + (M')_{n+1}^{(m)}. \end{aligned}$$

The four terms are, respectively, the fast-tracking error, the centered Markovian sampling error, the auxiliary-state mismatch, and the explicit slow martingale noise.

To compare the projected recursions, introduce their unconstrained driving sequences. Set

$$v_0^{(m)} = \theta_0^{(m)}, \quad \tilde{v}_0^{(m)} = \tilde{\theta}_0^{(m)},$$

and, for $0 \leq n \leq K_m - 1$, define

$$v_{n+1}^{(m)} = v_n^{(m)} + \beta_n^{(m)} \left(f(Y_n^{(m)}, x_n^{(m)}, \theta_n^{(m)}) + (M')_{n+1}^{(m)} \right), \quad (7.2)$$

and

$$\tilde{v}_{n+1}^{(m)} = \tilde{v}_n^{(m)} + \beta_n^{(m)} f^{(\text{av.})}(\tilde{\theta}_n^{(m)}; \theta_n^{(m)}). \quad (7.3)$$

The step-path representation and Lipschitz comparison are proved in Lemma S.13.1 of the Online Companion's Skorokhod section. In particular, for every $0 \leq n \leq K_m$,

$$\max_{0 \leq k \leq n} \|\theta_k^{(m)} - \tilde{\theta}_k^{(m)}\| \leq L_{\text{Sk}} \max_{0 \leq k \leq n} \|v_k^{(m)} - \tilde{v}_k^{(m)}\|.$$

Subtracting (7.3) from (7.2) gives

$$\begin{aligned} v_{n+1}^{(m)} - \tilde{v}_{n+1}^{(m)} &= v_n^{(m)} - \tilde{v}_n^{(m)} + \beta_n^{(m)} \left\{ f^{(\text{av.})}(\theta_n^{(m)}; \theta_n^{(m)}) - f^{(\text{av.})}(\tilde{\theta}_n^{(m)}; \theta_n^{(m)}) \right\} \\ &+ \beta_n^{(m)} \left\{ f(Y_n^{(m)}, x^*(\theta_n^{(m)}), \theta_n^{(m)}) - f^{(\text{av.})}(\theta_n^{(m)}; \theta_n^{(m)}) \right\} \\ &+ \beta_n^{(m)} \left\{ f(Y_n^{(m)}, x_n^{(m)}, \theta_n^{(m)}) - f(Y_n^{(m)}, x^*(\theta_n^{(m)}), \theta_n^{(m)}) \right\} \\ &+ \beta_n^{(m)} (M')_{n+1}^{(m)}. \end{aligned} \quad (7.4)$$

The second term on the right-hand side of (7.4), namely,

$$f(Y_n^{(m)}, x^*(\theta_n^{(m)}), \theta_n^{(m)}) - f^{(\text{av.})}(\theta_n^{(m)}; \theta_n^{(m)}). \quad (7.5)$$

is the Markov noise obtained by subtracting the stationary average under $\mu^{(\theta_n^{(m)})}$. To control it, we use a Poisson-equation decomposition for the controlled Markov chain.

Definition 7.1 (Poisson equation for the slow drift). For $\theta \in \Theta$, consider the Poisson equation

$$u(y) - \sum_{j \in \mathcal{Y}} p^{(\theta)}(y, j)u(j) = f(y, x^*(\theta), \theta) - f^{(\text{av.})}(\theta; \theta), \quad y \in \mathcal{Y}. \quad (7.6)$$

Let $\{Y_n\}$ be the Markov chain with transition matrix $p^{(\theta)}$. For $y \in \mathcal{Y}$, define

$$u_f^{(\theta)}(y) := \mathbb{E}^{(\theta)} \left[\sum_{n=0}^{\tau-1} \left(f(Y_n, x^*(\theta), \theta) - f^{(\text{av.})}(\theta; \theta) \right) \mid Y_0 = y \right], \quad (7.7)$$

where τ is the hitting time of the reference state y^* in Assumption 3.6. The expectation $\mathbb{E}^{(\theta)}$ is taken under a Markov chain that uses the transition kernel $p^{(\theta)}$ at every step. The parameter θ in the summand is held fixed as the chain evolves.

The same argument as in Definition 6.1 shows that (7.7) solves (7.6), with $u_f^{(\theta)}(y^*) = \mathbf{0}$.

Next we define a martingale-difference sequence.

Lemma 7.1 (Poisson decomposition). Define, for $0 \leq n \leq K_m - 1$,

$$(M''')_{n+1}^{(m)} := u_f^{(\theta_n^{(m)})}(Y_{n+1}^{(m)}) - \sum_{y \in \mathcal{Y}} p^{(\theta_n^{(m)})}(Y_n^{(m)}, y) u_f^{(\theta_n^{(m)})}(y). \quad (7.8)$$

Then $\{(M''')_{n+1}^{(m)}\}_{0 \leq n \leq K_m - 1}$ is a martingale-difference sequence with respect to $\{\mathcal{F}_n^{(m)}\}_{0 \leq n \leq K_m}$; more precisely,

$$(M''')_{n+1}^{(m)} \text{ is } \mathcal{F}_{n+1}^{(m)}\text{-measurable,} \quad \mathbb{E}[(M''')_{n+1}^{(m)} \mid \mathcal{F}_n^{(m)}] = \mathbf{0},$$

for every $0 \leq n \leq K_m - 1$. Moreover, for every $0 \leq n \leq K_m - 1$, almost surely,

$$\begin{aligned} f(Y_n^{(m)}, x^*(\theta_n^{(m)}), \theta_n^{(m)}) - f^{(\text{av.})}(\theta_n^{(m)}; \theta_n^{(m)}) \\ = u_f^{(\theta_n^{(m)})}(Y_n^{(m)}) - u_f^{(\theta_n^{(m)})}(Y_{n+1}^{(m)}) + (M''')_{n+1}^{(m)}. \end{aligned} \quad (7.9)$$

Let C_P and C_L denote the constants in Lemma S.11.3, and set

$$R_m := 3C_P \beta_{n_m} + C_L \{b(n_m) - b(n_{m+1})\}.$$

To handle the Markov noise (7.5), we use the Poisson decomposition. Its martingale-difference term is controlled later on the slow-martingale good event defined below, and its Poisson boundary term is controlled by the deterministic residual bound R_m .

Substituting (7.9) into the Markov noise term in (7.4) gives

$$\begin{aligned} v_{n+1}^{(m)} - \tilde{v}_{n+1}^{(m)} &= v_n^{(m)} - \tilde{v}_n^{(m)} + \beta_n^{(m)} \left\{ f^{(\text{av.})}(\theta_n^{(m)}; \theta_n^{(m)}) - f^{(\text{av.})}(\tilde{\theta}_n^{(m)}; \theta_n^{(m)}) \right\} \\ &\quad + \beta_n^{(m)} \left\{ u_f^{(\theta_n^{(m)})}(Y_n^{(m)}) - u_f^{(\theta_n^{(m)})}(Y_{n+1}^{(m)}) \right\} \\ &\quad + \beta_n^{(m)} \left\{ f(Y_n^{(m)}, x_n^{(m)}, \theta_n^{(m)}) - f(Y_n^{(m)}, x^*(\theta_n^{(m)}), \theta_n^{(m)}) \right\} \\ &\quad + \beta_n^{(m)} \left((M''')_{n+1}^{(m)} + (M')_{n+1}^{(m)} \right). \end{aligned} \quad (7.10)$$

The final term in (7.10) is the combined slow martingale: $(M')_{n+1}^{(m)}$ is the explicit martingale noise in the slow recursion, while $(M''')_{n+1}^{(m)}$ arises from the Poisson decomposition of the slow Markov noise. For each block m and each slow-martingale deviation threshold $\delta_s > 0$, define

$$\mathcal{G}_\theta^{(m)}(\delta_s) := \left\{ \max_{1 \leq k \leq K_m} \left\| \sum_{\ell=0}^{k-1} \beta_\ell^{(m)} \left[(M')_{\ell+1}^{(m)} + (M''')_{\ell+1}^{(m)} \right] \right\| < \delta_s \right\}. \quad (7.11)$$

Thus, on $\mathcal{G}_\theta^{(m)}(\delta_s)$, the cumulative slow-martingale perturbation remains below δ_s throughout block m . The subscript θ identifies this as the slow-coordinate martingale event, while m is the block index. Define

$$C_{\text{mg}}^{(\theta)} := \mathbf{c}_1 + \mathbf{c}_2 \mathbf{c}_6 + 4\mathbf{c}_5 \mathbf{B}_8.$$

Lemma 7.2. *For every $m \geq 0$ and $\delta_s > 0$,*

$$\mathbb{P} \left(\left(\mathcal{G}_\theta^{(m)}(\delta_s) \right)^c \right) \leq 2d_\theta \sum_{k=1}^{K_m} \exp \left(- \frac{\delta_s^2}{2\kappa_{d_\theta}^2 (C_{\text{mg}}^{(\theta)})^2 \{b(n_m) - b(n_m + k)\}} \right).$$

Proposition 7.1 (Comparison between $\theta^{(m)}$ and $\tilde{\theta}^{(m)}$). *Fix $m \geq 0$ and $\delta_s > 0$. On the event $\mathcal{G}_\theta^{(m)}(\delta_s)$, for every $0 \leq n \leq K_m$,*

$$\begin{aligned} \sup_{0 \leq k \leq n} \left\| \theta_k^{(m)} - \tilde{\theta}_k^{(m)} \right\| &\leq L_{\text{Sk}} \left[L_f \sum_{\ell=0}^{n-1} \beta_\ell^{(m)} \left\| e_\ell^{(m)} \right\| + \delta_s + R_m \right] \\ &\quad \times \exp \left(L_f (L_{fp} + 1) L_{\text{Sk}} a(n_m, n_m + n) \right), \end{aligned} \quad (7.12)$$

Here R_m is the deterministic Poisson-residual bound proved in Lemma S.11.3.

Proof. See Appendix A.4. □

Proposition 7.1 leaves the weighted fast-tracking term

$$F_m^x := \sum_{\ell=0}^{K_m-1} \beta_\ell^{(m)} \left\| e_\ell^{(m)} \right\| = \sum_{r=n_m}^{n_{m+1}-1} \beta_r \left\| x_r - x^*(\theta_r) \right\|$$

explicit. For a fast-martingale deviation level $\delta > 0$, define

$$S_m(\delta) := C_{x,T} \left(\delta + \alpha_{n_m} + \frac{\beta_{n_m}}{\alpha_{n_m}} \right), \quad (7.13)$$

where $C_{x,T}$ is the constant in Proposition 6.1. By (6.12), on $\mathcal{G}_x^{(m)}(\delta)$,

$$F_m^x \leq S_m(\delta). \quad (7.14)$$

For $\delta_f, \delta_s > 0$, set

$$\begin{aligned} E_m^{(\theta, \tilde{\theta})}(\delta_f, \delta_s) &:= L_{\text{Sk}} [L_f S_m(\delta_f) + \delta_s + R_m] \\ &\quad \times \exp \{ L_f (L_{fp} + 1) L_{\text{Sk}} a(n_m, n_{m+1}) \}. \end{aligned} \quad (7.15)$$

Corollary 7.1 (Original–auxiliary slow comparison). *Fix $m \geq 0$ and $\delta_f, \delta_s > 0$. On*

$$\mathcal{G}_x^{(m)}(\delta_f) \cap \mathcal{G}_\theta^{(m)}(\delta_s),$$

we have

$$\max_{0 \leq k \leq K_m} \left\| \theta_k^{(m)} - \tilde{\theta}_k^{(m)} \right\| \leq E_m^{(\theta, \tilde{\theta})}(\delta_f, \delta_s).$$

Equivalently,

$$\sup_{0 \leq t \leq H_m} \left\| \theta^{(m), \circ}(t) - \tilde{\theta}^{(m), \circ}(t) \right\| \leq E_m^{(\theta, \tilde{\theta})}(\delta_f, \delta_s).$$

7.2 Comparison of the auxiliary recursion with the projected ODE

We next compare the auxiliary recursion with the projected ODE

$$\dot{z}(t) = \Pi_{\Theta} \left(z(t), f^{(\text{av.})}(z(t); z(t)) \right).$$

Recall that $z^{(T_m)}(0) = \theta_{n_m}$. The auxiliary recursion uses the off-diagonal evaluation $f^{(\text{av.})}(\tilde{\theta}_n^{(m)}; \theta_n^{(m)})$, whereas the ODE uses the diagonal evaluation $f^{(\text{av.})}(z; z)$. For any $s \in [0, H_m]$ within block m , add and subtract $f^{(\text{av.})}(\tilde{\theta}^{(m),\circ}(s); \tilde{\theta}^{(m),\circ}(s))$ to write

$$\begin{aligned} & f^{(\text{av.})}(z^{(T_m)}(s); z^{(T_m)}(s)) - f^{(\text{av.})}(\tilde{\theta}^{(m),\circ}(s); \theta^{(m),\circ}(s)) \\ &= \left[f^{(\text{av.})}(z^{(T_m)}(s); z^{(T_m)}(s)) - f^{(\text{av.})}(\tilde{\theta}^{(m),\circ}(s); \tilde{\theta}^{(m),\circ}(s)) \right] \\ &+ \left[f^{(\text{av.})}(\tilde{\theta}^{(m),\circ}(s); \tilde{\theta}^{(m),\circ}(s)) - f^{(\text{av.})}(\tilde{\theta}^{(m),\circ}(s); \theta^{(m),\circ}(s)) \right]. \end{aligned}$$

The first bracket is controlled by the difference between the ODE state $z^{(T_m)}(s)$ and the auxiliary state $\tilde{\theta}^{(m),\circ}(s)$. The second is the off-diagonal mismatch between the stationary laws indexed by $\tilde{\theta}^{(m),\circ}(s)$ and $\theta^{(m),\circ}(s)$. To compare the auxiliary path $\tilde{\theta}^{(m),\circ}$ with the projected-ODE trajectory $z^{(T_m)}$, we first bound the difference between their unconstrained driving paths. The Lipschitz property of the Skorokhod map then bounds the difference between these two projected paths. Lemma S.13.2 of the Online Companion's Skorokhod section provides an absolutely continuous driver $w^{(\tau)}$ such that

$$w^{(\tau)}(0) = z^{(\tau)}(0) = \theta^\circ(\tau), \quad \dot{w}^{(\tau)}(t) = f^{(\text{av.})}(z^{(\tau)}(t); z^{(\tau)}(t)) \text{ a.e.}, \quad z^{(\tau)} = \Gamma(w^{(\tau)}).$$

We compare $w^{(T_m)}$ with the step driver $\tilde{v}^{(m),\circ}$, and Lemma S.13.1 bounds the difference between the projected paths $z^{(T_m)}$ and $\tilde{\theta}^{(m),\circ}$ in terms of the difference between their unconstrained driving paths.

Let $L_{f,\text{avg.}}$ denote the diagonal-field Lipschitz constant defined in Lemma S.11.2(ii).

Proposition 7.2 (Auxiliary-recursion–ODE comparison). *Fix $m \geq 0$ and $\delta_f, \delta_s > 0$, and recall from Section 4 that $H_m = t^{(m)}(K_m) = a(n_m, n_{m+1})$. On*

$$\mathcal{G}_x^{(m)}(\delta_f) \cap \mathcal{G}_\theta^{(m)}(\delta_s),$$

define, for $0 \leq t \leq H_m$,

$$\mathcal{A}_m(t) := \sup_{0 \leq s \leq t} \|w^{(T_m)}(s) - \tilde{v}^{(m),\circ}(s)\|.$$

Then

$$\mathcal{A}_m(t) \leq \exp\{L_{f,\text{avg.}} L_{\text{Sk}} t\} \left[\mathbf{B}_8 L_\mu t E_m^{(\theta, \tilde{\theta})}(\delta_f, \delta_s) + \mathbf{B}_8 \beta_{n_m} \right]. \quad (7.16)$$

Consequently,

$$\begin{aligned} & \sup_{0 \leq s \leq t} \|z^{(T_m)}(s) - \tilde{\theta}^{(m),\circ}(s)\| \\ & \leq L_{\text{Sk}} \exp\{L_{f,\text{avg.}} L_{\text{Sk}} t\} \left[\mathbf{B}_8 L_\mu t E_m^{(\theta, \tilde{\theta})}(\delta_f, \delta_s) + \mathbf{B}_8 \beta_{n_m} \right]. \end{aligned}$$

Proof. See Appendix A.5. □

7.3 Putting the estimates together

For later use, define

$$p_{\text{det}} := \mathbf{b} - \mathbf{a},$$

which is the decay exponent of the step-size ratio β_n/α_n . For $m \geq 0$, a fast-martingale deviation threshold $\delta_f > 0$, and a slow-martingale deviation threshold $\delta_s > 0$, define

$$\begin{aligned} B_m(\delta_f, \delta_s) &:= E_m^{(\theta, \tilde{\theta})}(\delta_f, \delta_s) \\ &\quad + L_{\text{Sk}} \exp\{L_{f, \text{avg}} L_{\text{Sk}} a(n_m, n_{m+1})\} \left[\mathbf{B}_8 L_{\mu} a(n_m, n_{m+1}) E_m^{(\theta, \tilde{\theta})}(\delta_f, \delta_s) \right. \\ &\quad \left. + \mathbf{B}_8 \beta_{n_m} \right]. \end{aligned} \quad (7.17)$$

Theorem 7.1 (Localized blockwise tracking of the projected ODE). *The deviation thresholds δ_f and δ_s may be chosen arbitrarily for each block m . Section 8 later substitutes the rate-specific choices $\delta_{\text{dev}}^{(x)}(m)$ and $\delta_{\text{dev}}^{(\theta)}(m)$. Here dev denotes a deviation threshold; x and θ denote the fast and slow martingales, respectively. Fix $m \geq 0$ and $\delta_f, \delta_s > 0$. On*

$$\mathcal{G}_x^{(m)}(\delta_f) \cap \mathcal{G}_\theta^{(m)}(\delta_s),$$

we have

$$\sup_{0 \leq t \leq H_m} \|\theta^{(m), \circ}(t) - z^{(T_m)}(t)\| \leq B_m(\delta_f, \delta_s). \quad (7.18)$$

Moreover, there exists $C_{B,T} < \infty$, depending only on T and the problem parameters, such that

$$B_m(\delta_f, \delta_s) \leq C_{B,T} \left\{ \delta_f + \delta_s + \alpha_{n_m} + \frac{\beta_{n_m}}{\alpha_{n_m}} \right\}. \quad (7.19)$$

Consequently,

$$B_m(\delta_f, \delta_s) \leq C_{B,T} \left\{ \delta_f + \delta_s + (N_0 + n_m)^{-p_{\text{det}}} \right\}. \quad (7.20)$$

Proof. See Appendix A.6. $\square$

8 Almost-sure convergence and last-iterate rates

We now convert the localized blockwise projected-ODE tracking estimate of Theorem 7.1 into iteration-wise almost-sure rates. The argument has three steps. First, we choose block-dependent fast- and slow-martingale deviation thresholds whose failure probabilities are summable over the blocks. The Borel–Cantelli lemma then implies that the blockwise tracking estimate holds on every sufficiently large block almost surely. Second, a uniform one-block Lyapunov decrease condition yields a perturbed nonlinear recursion at the block endpoints. Third, we analyze this recursion and use the within-block tracking estimate to transfer the endpoint bound to every discrete iterate. Uniform one-block contraction is the special case in which the Lyapunov value decreases by at least a fixed proportion over each block. Throughout this section, the standing assumptions and Condition 3.1 of Section 3, together with (3.3), remain in force.

For $s \geq 0$, let $\mathcal{S}_s$ denote the time- s flow map of the projected ODE (5.1). The well-posedness of (5.1) is established in Lemma S.15.1. Let $\mathcal{EQ}_\Theta$ denote the set of equilibrium points of the ODE (5.1):

$$\mathcal{EQ}_\Theta := \left\{ z \in \Theta : \Pi_\Theta(z, f^{(\text{av.})}(z; z)) = \mathbf{0} \right\}.$$

Definition 8.1 (Lyapunov function). Consider the projected ODE (5.1), repeated here for convenience:

$$\dot{z}(t) = \Pi_{\Theta} \left(z(t), f^{(\text{av.})}(z(t); z(t)) \right), \quad z(0) = z_0 \in \Theta.$$

A continuous function $J : \Theta \rightarrow \mathbb{R}_+$ is called a Lyapunov function for (5.1) if $J(z) = 0$ if and only if $z \in \mathcal{EQ}_{\Theta}$, and if, for every solution $z(\cdot)$ of (5.1),

$$J(z(t)) = J(\mathbf{S}_{t-s}(z(s))) \leq J(z(s)), \quad 0 \leq s \leq t < \infty.$$

Thus J is nonincreasing along trajectories of the projected ODE.

For any Lyapunov function J as in Definition 8.1, write

$$J_{\max} := \max_{\theta \in \Theta} J(\theta) < \infty,$$

where finiteness follows from the continuity of J and the compactness of Θ . We use a Lyapunov function to measure convergence to the equilibrium set.

Assumption 8.1. The projected ODE (5.1) admits a Lyapunov function in the sense of Definition 8.1.

Theorem 7.1 bounds, on each slow-time block, the gap between the path of the slow iterates $\theta^{(m),\circ}$ and the projected-ODE trajectory $z^{(T_m)}$ started at θ_{n_m} . We now choose block-dependent thresholds δ_f and δ_s that vanish as $m \rightarrow \infty$, while their block-failure probabilities remain summable. Smaller deviation levels sharpen this bound but increase the block-failure probabilities; the logarithmic choices below make both levels vanish while keeping those probabilities summable.

For $m \geq 0$, set

$$D_m := b(n_m) - b(n_{m+1}), \quad L_m^x := \log(4d_x K_m(m+2)^2), \quad L_m^\theta := \log(4d_\theta K_m(m+2)^2).$$

Define

$$\delta_{\text{dev}}^{(x)}(m) := \left(\frac{\alpha_{n_m} L_m^x}{c_{\text{mg}}^x} \right)^{1/2},$$

and

$$\delta_{\text{dev}}^{(\theta)}(m) := \kappa_{d_\theta} C_{\text{mg}}^{(\theta)} \sqrt{2D_m L_m^\theta}.$$

Define the blockwise rate event

$$\mathcal{G}_m^{\text{rate}} := \mathcal{G}_x^{(m)}(\delta_{\text{dev}}^{(x)}(m)) \cap \mathcal{G}_\theta^{(m)}(\delta_{\text{dev}}^{(\theta)}(m)).$$

On $\mathcal{G}_m^{\text{rate}}$, both martingale perturbations stay below their rate-calibrated levels, allowing the blockwise tracking estimate below to be applied. By Lemmas 6.2 and 7.2, the chosen thresholds satisfy

$$\mathbb{P}\left([\mathcal{G}_x^{(m)}(\delta_{\text{dev}}^{(x)}(m))]^c\right) \leq \frac{1}{2(m+2)^2}, \quad \mathbb{P}\left([\mathcal{G}_\theta^{(m)}(\delta_{\text{dev}}^{(\theta)}(m))]^c\right) \leq \frac{1}{2(m+2)^2}.$$

By the union bound, $\sum_{m \geq 1} \mathbb{P}((\mathcal{G}_m^{\text{rate}})^c) < \infty$, and the Borel–Cantelli lemma implies that $\mathcal{G}_m^{\text{rate}}$ holds eventually almost surely.

Define

$$d_m^{\text{rate}} := \delta_{\text{dev}}^{(x)}(m) + \delta_{\text{dev}}^{(\theta)}(m) + \alpha_{n_m} + \frac{\beta_{n_m}}{\alpha_{n_m}}.$$

By Theorem 7.1, applied with

$$\delta_f = \delta_{\text{dev}}^{(x)}(m), \quad \delta_s = \delta_{\text{dev}}^{(\theta)}(m),$$

on $\mathcal{G}_m^{\text{rate}}$,

$$\sup_{0 \leq r \leq H_m} \left\| \theta^{(m), \circ}(r) - z^{(T_m)}(r) \right\| \leq C_{B,T} d_m^{\text{rate}}. \quad (8.1)$$

The estimate (8.1) controls the stochastic perturbation accumulated within one block. To propagate this estimate over successive blocks, we impose a Lyapunov decrease condition over time T along the projected-ODE flow, uniformly over initial states $\theta \in \Theta$.

Assumption 8.2 (Uniform one-block Lyapunov decrease). Let J be the Lyapunov function in Assumption 8.1. Let $T > 0$ be the block horizon used in the definition of $\{n_m\}$. There exists a continuous function

$$\mathcal{D}_J : [0, J_{\max}] \rightarrow \mathbb{R}_+$$

such that

$$\mathcal{D}_J(0) = 0, \quad 0 < \mathcal{D}_J(u) \leq u, \quad 0 < u \leq J_{\max},$$

and

$$J(S_T(\theta)) \leq J(\theta) - \mathcal{D}_J(J(\theta)), \quad \theta \in \Theta.$$

Since J is nonincreasing along the projected-ODE flow, for every $s \geq T$,

$$J(S_s(\theta)) \leq J(S_T(\theta)) \leq J(\theta) - \mathcal{D}_J(J(\theta)), \quad \theta \in \Theta.$$

A decrease proportional to the current Lyapunov value gives the following contraction condition.

Assumption 8.3 (Uniform one-block Lyapunov contraction). Let J be the Lyapunov function in Assumption 8.1, and let $T > 0$ be the block horizon used in the definition of $\{n_m\}$. There exists $\rho_T \in (0, 1)$ such that

$$J(S_T(\theta)) \leq \rho_T J(\theta), \quad \theta \in \Theta.$$

Assumption 8.3 is the special case of Assumption 8.2 obtained by taking

$$\mathcal{D}_J(u) := (1 - \rho_T)u.$$

Define

$$m(n) := \max\{m \geq 0 : n_m \leq n\},$$

so that $n_{m(n)} \leq n < n_{m(n)+1}$.

Theorem 8.1 (Fast tracking rate). *Under Assumptions 3.1, 3.2, 3.3, 3.4, 3.5, 3.6, and 3.7, the step-size exponent requirement in Condition 3.1, and Condition (3.3),*

$$\|x_n - x^*(\theta_n)\| = O_{\text{a.s.}} \left((N_0 + n)^{-r_x(\mathbf{a}, \mathbf{b})} \sqrt{\log(N_0 + n)} \right), \quad r_x(\mathbf{a}, \mathbf{b}) := \min \left\{ \frac{\mathbf{a}}{2}, \mathbf{b} - \mathbf{a} \right\}.$$

The exponent $r_x(\mathbf{a}, \mathbf{b})$ furnished by this bound is maximized over $1/2 < \mathbf{a} < \mathbf{b} \leq 1$ at $\mathbf{b} = 1$ and $\mathbf{a} = 2/3$, where $r_x(2/3, 1) = 1/3$.

Theorem 8.2 (Almost-sure convergence under one-block Lyapunov decrease). *Fix a Lyapunov function J as in Assumption 8.1. Suppose that J is L_J -Lipschitz on Θ , and that Assumption 8.2 holds for the block horizon T used to define $\{n_m\}$. Define*

$$U_m := J(\theta_{n_m}), \quad e_m^J := L_J C_{B,T} d_m^{\text{rate}}.$$

Then, almost surely, there exists a finite random index $m_0 = m_0(\omega)$ such that

$$U_{m+1} \leq U_m - \mathcal{D}_J(U_m) + e_m^J, \quad m \geq m_0. \quad (8.2)$$

Whenever $m(n) \geq m_0$,

$$J(\theta_n) \leq U_{m(n)} + e_{m(n)}^J. \quad (8.3)$$

Consequently,

$$J(\theta_n) \longrightarrow 0, \quad \text{dist}(\theta_n, \mathcal{EQ}_\Theta) \longrightarrow 0 \quad \text{almost surely.}$$

Proof. See Appendix A.7. □

When $\mathfrak{b} = 1$, $\beta_n = (N_0 + n)^{-1}$, and both the accumulated slow time and the number of completed blocks $m(n)$ grow as $\log(N_0 + n)$; for $\mathfrak{b} < 1$, they grow as $(N_0 + n)^{1-\mathfrak{b}}$. Theorem 8.2 gives almost-sure convergence in both cases. Explicit rates require a quantitative lower bound on $\mathcal{D}_J$, as imposed in the next result.

Theorem 8.3 (Rates under a power-law decrease condition). *Assume the hypotheses of Theorem 8.2. Suppose, in addition, that there exist constants $p_J > 1$, $\kappa_J > 0$, and $u_J > 0$ such that*

$$\mathcal{D}_J(u) \geq \kappa_J u^{p_J}, \quad 0 \leq u \leq u_J. \quad (8.4)$$

If, for some $r > 0$ and $C_r < \infty$,

$$d_m^{\text{rate}} \leq C_r (N_0 + n_m)^{-r} \sqrt{\log(N_0 + n_m)}$$

for all sufficiently large m , the following conclusions hold. If $\mathfrak{b} < 1$, then

$$J(\theta_n) = O_{\text{a.s.}} \left((N_0 + n)^{-\frac{1-\mathfrak{b}}{p_J-1}} + (N_0 + n)^{-\frac{r}{p_J}} (\log(N_0 + n))^{\frac{1}{2p_J}} \right). \quad (8.5)$$

If $\mathfrak{b} = 1$, then

$$J(\theta_n) = O_{\text{a.s.}} \left(\{\log(N_0 + n)\}^{-\frac{1}{p_J-1}} \right). \quad (8.6)$$

The rate (8.6) for $\mathfrak{b} = 1$ is sharp under the power-decrease hypothesis; see Remark S.7.1 in the Online Companion.

Theorem 8.2 allows a general lower bound on the Lyapunov decrease over one block. In the contraction case, the block recursion can be unrolled explicitly, giving the sharper estimate below.

Theorem 8.4 (Slow-iterate Lyapunov error rates under one-block contraction). *Fix a Lyapunov function J as in Assumption 8.1. Suppose that J is L_J -Lipschitz on Θ , and that Assumption 8.3 holds for the block horizon T used to define $\{n_m\}$. Then, almost surely, there exists a finite random index $m_0 = m_0(\omega)$ such that, with d_j^{rate} as defined above, the following inequality holds for every n satisfying $m(n) \geq m_0$:*

$$J(\theta_n) \leq \rho_T^{m(n)-m_0} J_{\max} + L_J C_{B,T} \sum_{j=m_0}^{m(n)-1} \rho_T^{m(n)-1-j} d_j^{\text{rate}} + L_J C_{B,T} d_{m(n)}^{\text{rate}}.$$

Consequently, suppose that, for some $r > 0$ and $C_r < \infty$,

$$d_m^{\text{rate}} \leq C_r (N_0 + n_m)^{-r} \sqrt{\log(N_0 + n_m)}, \quad m \geq m_0.$$

If $\mathfrak{b} < 1$, then

$$J(\theta_n) = O_{\text{a.s.}} \left((N_0 + n)^{-r} \sqrt{\log(N_0 + n)} \right).$$

If $\mathfrak{b} = 1$, define

$$\lambda_T := -\frac{\log \rho_T}{T} > 0. \quad (8.7)$$

Then

$$J(\theta_n) = \begin{cases} O_{\text{a.s.}} \left((N_0 + n)^{-r} \sqrt{\log(N_0 + n)} \right), & \lambda_T > r, \\ O_{\text{a.s.}} \left((N_0 + n)^{-r} \{\log(N_0 + n)\}^{3/2} \right), & \lambda_T = r, \\ O_{\text{a.s.}} \left((N_0 + n)^{-\lambda_T} \right), & 0 < \lambda_T < r. \end{cases} \quad (8.8)$$

The block estimates in Lemmas S.15.4 and S.15.5 of the Online Companion imply that, for the polynomial step sizes above under Condition 3.1,

$$d_m^{\text{rate}} \leq C_T (N_0 + n_m)^{-r_x(\mathfrak{a}, \mathfrak{b})} \sqrt{\log(N_0 + n_m)}$$

for all sufficiently large m . Thus the rate-envelope conditions in Theorem 8.3 and in the rate part of Theorem 8.4 hold with exponent $r(\mathfrak{a}, \mathfrak{b})$; see the proof of Corollary 8.1 in Appendix S.7 of the Online Companion.

Corollary 8.1 (Optimized two-time-scale rate under contraction). *Assume the hypotheses of Theorem 8.4. For $\mathfrak{b} < 1$,*

$$J(\theta_n) + \|x_n - x^*(\theta_n)\| = O_{\text{a.s.}} \left((N_0 + n)^{-r_x(\mathfrak{a}, \mathfrak{b})} \sqrt{\log(N_0 + n)} \right).$$

Moreover,

$$\sup_{\frac{1}{2} < \mathfrak{a} < \mathfrak{b} < 1} r_x(\mathfrak{a}, \mathfrak{b}) = \frac{1}{3}.$$

More precisely, for every

$$0 < \varepsilon_{\text{rate}} < \frac{1}{12},$$

choose

$$\mathfrak{b} = 1 - 3\varepsilon_{\text{rate}}, \quad \mathfrak{a} = \frac{2\mathfrak{b}}{3} = \frac{2}{3} - 2\varepsilon_{\text{rate}}.$$

Then the step-size conditions hold and

$$J(\theta_n) + \|x_n - x^*(\theta_n)\| = O_{\text{a.s.}} \left((N_0 + n)^{-1/3 + \varepsilon_{\text{rate}}} \sqrt{\log(N_0 + n)} \right).$$

At the harmonic endpoint, take $\mathfrak{b} = 1$ and $\mathfrak{a} = 2/3$. The slow rate is (8.8) with $r = 1/3$, and Theorem 8.1 gives fast-tracking rate $O_{\text{a.s.}}((N_0 + n)^{-1/3} \sqrt{\log(N_0 + n)})$. If $J(\theta) = \|\theta - \theta^*\|$ for a fixed equilibrium θ^* of the projected ODE (5.1), each rate stated for $J(\theta_n)$ is the corresponding slow-iterate distance rate.

Corollary 8.2 (Optimized polynomial exponent under power Lyapunov decrease). *Assume the hypotheses of Theorem 8.2, together with (8.4) for some $p_J > 1$. For the case $\mathfrak{b} < 1$, choose*

$$\mathfrak{a} = \frac{2p_J}{4p_J - 1}, \quad \mathfrak{b} = \frac{3p_J}{4p_J - 1}.$$

Then $1/2 < \mathfrak{a} < \mathfrak{b} < 1$, and

$$J(\theta_n) = O_{\text{a.s.}} \left((N_0 + n)^{-\frac{1}{4p_J - 1}} (\log(N_0 + n))^{\frac{1}{2p_J}} \right),$$

and

$$\|x_n - x^*(\theta_n)\| = O_{\text{a.s.}} \left((N_0 + n)^{-\frac{p_J}{4p_J - 1}} \sqrt{\log(N_0 + n)} \right).$$

If $J(\theta) = \|\theta - \theta^\|$ for a fixed equilibrium θ^* of the projected ODE (5.1), the first display is also the corresponding slow-iterate distance bound.*

At the harmonic endpoint, take $\mathfrak{b} = 1$ and $\mathfrak{a} = 2/3$. The slow rate is (8.6), and Theorem 8.1 gives fast-tracking rate $O_{\text{a.s.}}((N_0 + n)^{-1/3} \sqrt{\log(N_0 + n)})$.

8.1 Faster rates at an attracting equilibrium face

The general fast-tracking bound accounts for movement of $x^*(\theta_n)$ as the slow iterate evolves. This contribution can be reduced when the slow equilibrium set is a strictly attracting face of a box and x^* has the same value at every point of that face. The resulting rates are given next.

Proposition 8.1 (Rates near an equilibrium face). *For (1.1)–(1.2), retain the standing assumptions of Section 3 on the chain, update maps, noise, and fast-iterate stability, and the assumptions on the Lipschitz Lyapunov function J in Theorem 8.2. Write $t_n = N_0 + n$. In place of Condition 3.1 and (3.3), allow either*

$$\begin{aligned} \alpha_n &= t_n^{-\mathfrak{a}}, \quad \beta_n = t_n^{-\mathfrak{b}}, \quad \frac{1}{2} < \mathfrak{a} < \mathfrak{b} \leq 1, \quad t_1^{1-\mathfrak{a}} \geq 4; \\ \alpha_n &= A \log(t_n)/t_n, \quad \beta_n = B/t_n, \quad A, B > 0, \quad N_0 \geq 3, \quad \alpha_1 \leq 1, \quad A \log t_1 \geq \max\{4, B\}. \end{aligned} \quad (8.9)$$

Suppose that $\Theta = \prod_{i=1}^{d_\theta} [\ell_i, u_i]$, with $\ell_i < u_i$, and that, for some $I \subseteq \{1, \dots, d_\theta\}$, its projected-ODE equilibrium set is

$$\mathcal{EQ}_\Theta = \{\theta \in \Theta : \theta_i = v_i \ (i \in I)\}, \quad v_i \in \{\ell_i, u_i\}.$$

For $i \in I$, set $s_i = -1$ at a lower endpoint and $s_i = 1$ at an upper endpoint. Assume

$$\inf_{\theta \in \mathcal{EQ}_\Theta} s_i f_i^{(\text{av.})}(\theta; \theta) > 0, \quad i \in I. \quad (8.10)$$

Then

$$\text{dist}(\theta_n, \mathcal{EQ}_\Theta) + J(\theta_n) = O_{\text{a.s.}}(\beta_n \log t_n). \quad (8.11)$$

If, additionally, $x^(\theta) = x^\dagger$ for every $\theta \in \mathcal{EQ}_\Theta$, for some fixed $x^\dagger$, then*

$$\|x_n - x^\dagger\| + \|x_n - x^*(\theta_n)\| = O_{\text{a.s.}}(\sqrt{\alpha_n \log t_n}). \quad (8.12)$$

Thus $\mathfrak{b} = 1$, $\mathfrak{a} = 1 - 2\varepsilon$, with $0 < \varepsilon < 1/4$, gives a joint rate $O_{\text{a.s.}}(n^{-1/2+\varepsilon} \sqrt{\log n})$. The second schedule in (8.9) gives $O_{\text{a.s.}}(n^{-1/2} \log n)$.

Proof. See Appendix S.7.6. □

8.2 Improved rates under contraction of the reduced slow map

A different mechanism yields the $n^{-1/2}$ polynomial exponent: assume that the reduced slow update map is a Euclidean contraction. Together with logarithmic separation of the two step sizes, this controls the slow error directly, even when its equilibrium lies on the boundary. Unlike Proposition 8.1, the next result permits an arbitrary compact convex polyhedron and does not require an attracting box face or a common fast target along a face. Its contraction assumption is stronger than the one-block Lyapunov condition.

Proposition 8.2 (Contraction rate with boundary equilibria). *For (1.1)–(1.2), retain the standing assumptions of Section 3 on the chain, update maps, noise, fast contraction, and fast-iterate stability. Suppose that*

$$\overline{G}(\theta) := \theta + f^{(\text{av.})}(\theta; \theta), \quad \|\overline{G}(\theta) - \overline{G}(\vartheta)\| \leq \rho_s \|\theta - \vartheta\|, \quad \theta, \vartheta \in \Theta, \quad (8.13)$$

for some $\rho_s \in [0, 1)$, in the Euclidean slow-coordinate norm. Let θ^* be the unique fixed point of $\text{proj}_\Theta \circ \overline{G}$. In place of Condition 3.1 and (3.3), take

$$t_n = N_0 + n, \quad \alpha_n = \frac{A \log t_n}{t_n}, \quad \beta_n = \frac{B}{t_n}, \quad A > 0, \quad B \geq \frac{2}{1 - \rho_s}, \quad (8.14)$$

where $N_0 \geq 3$, $\alpha_1 < 1$, and $A \log t_1 \geq \max\{4, B\}$. Assumption 3.7 is required for this selected schedule. Then

$$\|x_n - x^*(\theta_n)\| + \|\theta_n - \theta^*\| = O_{\text{a.s.}}\left(\frac{\log t_n}{\sqrt{t_n}}\right). \quad (8.15)$$

The same bound holds for $\|x_n - x^*(\theta^*)\|$ and for $J(\theta_n)$ whenever J is Lipschitz and $J(\theta^*) = 0$.

Proof. See Appendix S.17. □

The equilibrium drift may be nonzero and outward; neither interiority nor strict complementarity is assumed. Condition (8.13) implies contraction of distance along the projected ODE and is stronger than the one-block Lyapunov contraction used in Theorem 8.4.

In the terminology of Chandak [6], “true” two-time-scale stochastic approximation requires $\beta_n/\alpha_n \rightarrow 0$, not specifically polynomial separation. The schedule (8.14) satisfies this condition since $\beta_n/\alpha_n = B/(A \log t_n)$. Under the additional contraction condition (8.13), the bound (8.15) has polynomial exponent $1/2$ for both iterate errors, with only logarithmic separation of the step sizes. This improvement uses the stronger contraction assumption together with the revised schedule and proof; it does not follow from changing the schedule alone. The proof adapts the auxiliary noise-averaging idea of that work to the projected recursion and controlled Markov noise, using a feasible auxiliary sequence to handle boundary equilibria.

One setting in which the reduced-map condition can be checked directly is projected gradient descent for a smooth strongly convex objective. If $\overline{G}(\theta) = \theta - \eta \nabla F(\theta)$ and $mI \preceq \nabla^2 F(\theta) \preceq LI$ on Θ , then, for $0 < \eta < 2/L$, the mean-value theorem gives

$$\|\overline{G}(\theta) - \overline{G}(\vartheta)\| \leq \max\{|1 - \eta m|, |1 - \eta L|\} \|\theta - \vartheta\|.$$

The displayed factor is less than one. Projection onto a compact convex polyhedron preserves this contraction bound, including when the constrained minimizer lies on a boundary face.

9 Actor–Critic Application

Actor–critic provides an example in which the slow iterate controls both the sampling law and the projected update. We first specify the single-trajectory algorithm and its constrained policy class, then state the rate obtained from the attracting-face result. The verification of the assumptions is in the Online Companion; standalone projected TD(0) and projected-SGD applications appear in Appendices S.1 and S.2 there.

Consider a finite-state, finite-action Markov decision process (MDP); see, for example, Puterman [21]. Let $\mathcal{S}$ and $\mathcal{A}$ be its state and action spaces, $P(s' | s, a)$ its transition probabilities, and $c(s, a)$ its expected one-step cost. At time n , the current actor parameter determines a policy from which A_n is drawn at state S_n . The environment then produces S_{n+1} and a realized one-step cost C_{n+1} . Let $\mathcal{F}_n^{\text{MDP}}$ be the history immediately before A_n is drawn. For every $s' \in \mathcal{S}$,

$$\mathbb{P}(S_{n+1} = s' | \mathcal{F}_n^{\text{MDP}}, A_n) = P(s' | S_n, A_n), \quad \mathbb{E}[C_{n+1} | \mathcal{F}_n^{\text{MDP}}, A_n] = c(S_n, A_n).$$

The Online Companion, Appendix S.8, expresses the critic and actor updates in the drift-plus-martingale form of (1.1)–(1.2) and checks the assumptions used in the rate results.

Fix $0 < \gamma_{\text{disc}} < 1$. For a stationary policy π , its discounted total-cost value is

$$V_\pi(s) := \mathbb{E}_s^\pi \left[\sum_{k=0}^{\infty} \gamma_{\text{disc}}^k C_{k+1} \right], \quad s \in \mathcal{S}.$$

We analyze an online actor–critic recursion using one observed state–action transition per iteration, motivated by Konda and Borkar [18, Algorithm 3]. In the original asynchronous form, an update associated with a state–action pair uses a step size indexed by that pair’s visit count. Here the critic coordinate for the observed state and the actor coordinate for the observed state–action pair are updated using the global step sizes α_n and β_n , indexed by the number n of environment transitions. The rate below applies to this global-clock recursion.

Assume $|C_{n+1}| \leq C_{\max}$. The actor parameter $\vartheta = (\vartheta(s, a))_{s,a}$ consists of state–action logits, also called preference parameters. Fix $B_{\text{ac}} > 0$ and set

$$\Theta_{\text{ac}} := [-B_{\text{ac}}, B_{\text{ac}}]^{|\mathcal{S}||\mathcal{A}|}, \quad \pi_\vartheta(a | s) := \frac{e^{\vartheta(s,a)}}{\sum_{b \in \mathcal{A}} e^{\vartheta(s,b)}},$$

and draw $A_n \sim \pi_{\vartheta_n}(\cdot | S_n)$ before the time- n update. For $s \in \mathcal{S}$ and $a \in \mathcal{A}$, let $e_s \in \mathbb{R}^{|\mathcal{S}|}$ and $e_{s,a} \in \mathbb{R}^{|\mathcal{S}||\mathcal{A}|}$ denote the corresponding standard coordinate vectors. With

$$\delta_{n+1}^{\text{ac}} := C_{n+1} + \gamma_{\text{disc}} V_n(S_{n+1}) - V_n(S_n),$$

the recursion is

$$\begin{aligned} V_{n+1} &= V_n + \alpha_n e_{S_n} \delta_{n+1}^{\text{ac}}, \\ \vartheta_{n+1} &= \text{proj}_{\Theta_{\text{ac}}} (\vartheta_n - \beta_n e_{S_n, A_n} \delta_{n+1}^{\text{ac}}). \end{aligned} \tag{9.1}$$

For $\vartheta \in \Theta_{\text{ac}}$, let d_ϑ and V_ϑ be the stationary state distribution and discounted cost value under π_ϑ . Assume that the controlled state process induced by the policy family $\{\pi_\vartheta : \vartheta \in \Theta_{\text{ac}}\}$ satisfies Assumptions 3.1, 3.2, and 3.6 uniformly over $\vartheta \in \Theta_{\text{ac}}$. Assume also that the directed graph on $\mathcal{S}$, with an edge $s \rightarrow s'$ whenever $P(s' | s, a) > 0$ for some $a \in \mathcal{A}$, is strongly connected. As shown in Appendix S.8 of the Online Companion, the full support of the bounded-logit softmax policies and compactness of Θ_{ac} then imply

$$d_{\min}^{\text{ac}} := \inf_{\vartheta \in \Theta_{\text{ac}}} \min_{s \in \mathcal{S}} d_\vartheta(s) > 0. \tag{9.2}$$

We refer to (9.2) as uniform stationary state coverage. Set

$$\pi_{\min}^{\text{ac}} := \frac{e^{-2B_{\text{ac}}}}{|\mathcal{A}|}, \quad q_{\min}^{\text{ac}} := d_{\min}^{\text{ac}} \pi_{\min}^{\text{ac}}.$$

Initialize $\vartheta_1 \in \Theta_{\text{ac}}$ and choose $\|V_1\|_{\infty} \leq C_{\max}/(1 - \gamma_{\text{disc}})$. The critic recursion preserves this cube, which also contains every discounted value vector V_{ϑ} . On that bounded set, the finite MDP and the softmax policy on the compact logit box give the localized Lipschitz bounds required for the sampled updates. In the fast-coordinate norm $\|\cdot\|_{\infty}$, the averaged critic map is contractive on $\mathbb{R}^{|\mathcal{S}|}$; cube invariance supplies fast-iterate stability. The Online Companion verifies these claims.

Let $V_{B_{\text{ac}}}^*$ be the optimal value among policies representable, state by state, by logits in $[-B_{\text{ac}}, B_{\text{ac}}]^{|\mathcal{A}|}$. We measure the actor's aggregate value gap by

$$J_{B_{\text{ac}}}(\vartheta) := \sum_{s \in \mathcal{S}} \{V_{\vartheta}(s) - V_{B_{\text{ac}}}^*(s)\}.$$

For a fixed actor parameter ϑ , the stationary frequency of the state–action pair (s, a) is $q_{\vartheta}(s, a) := d_{\vartheta}(s) \pi_{\vartheta}(a | s)$. These frequencies weight the coordinates of the averaged actor drift. Appendix S.8 of the Online Companion uses their uniform positive lower bound to establish Lyapunov decrease, identify the restricted-optimal equilibrium face, and show that the projected actor ODE reaches that face in uniformly bounded time. On the face, the critic equilibrium is $V_{B_{\text{ac}}}^*$, and each coordinate fixed at an endpoint has strictly outward drift. Hence Proposition 8.1 applies.

Corollary 9.1 (Actor–critic rate). *Suppose that the preceding MDP, policy, and initialization conditions hold, and choose either step-size schedule in Proposition 8.1. Then*

$$J_{B_{\text{ac}}}(\vartheta_n) = O_{\text{a.s.}}(\beta_n \log(N_0 + n)), \quad (9.3)$$

and

$$\|V_n - V_{\vartheta_n}\|_{\infty} + \|V_n - V_{B_{\text{ac}}}^*\|_{\infty} = O_{\text{a.s.}}\left(\sqrt{\alpha_n \log(N_0 + n)}\right). \quad (9.4)$$

For $\alpha_n = A \log(N_0 + n)/(N_0 + n)$ and $\beta_n = B/(N_0 + n)$, with the shift specified in that proposition, the actor value gap is $O_{\text{a.s.}}(n^{-1} \log n)$, and

$$J_{B_{\text{ac}}}(\vartheta_n) + \|V_n - V_{\vartheta_n}\|_{\infty} = O_{\text{a.s.}}\left(\frac{\log(N_0 + n)}{\sqrt{N_0 + n}}\right). \quad (9.5)$$

This controls the aggregate value Lyapunov gap and the critic tracking error. If V^* is the unrestricted optimal value, then

$$0 \leq \sum_{s \in \mathcal{S}} \{V_{B_{\text{ac}}}^*(s) - V^*(s)\} \leq \frac{2|\mathcal{S}|(|\mathcal{A}| - 1)C_{\max}e^{-2B_{\text{ac}}}}{(1 - \gamma_{\text{disc}})^2}.$$

Consequently, for any prescribed target tolerance $\varepsilon_{\text{pol}} > 0$, a fixed B_{ac} can be chosen sufficiently large that the displayed restriction bias is at most ε_{pol} . The unrestricted aggregate value gap is then at most ε_{pol} plus the vanishing term in (9.3).

10 Conclusion

We obtained finite-time concentration bounds for projected two-time-scale stochastic approximation driven by a controlled Markov chain. On each slow-time block, the bounds compare the fast

iterate with its moving equilibrium and the projected slow path with its ODE trajectory. The Skorokhod-map comparison handles boundary discontinuities, while the estimates identify separately the martingale, Markovian, and time-scale contributions.

A one-block Lyapunov decrease gives almost-sure convergence to the slow equilibrium set; power decrease and contraction give explicit last-iterate rates. For polynomial step sizes, the optimized joint exponent under one-block contraction approaches $1/3$. With a strictly attracting equilibrium face and a common fast target, logarithmically separated step sizes give the joint $O_{\text{a.s.}}(n^{-1/2} \log n)$ rate. Euclidean contraction of the reduced slow map gives the same joint order on a general compact convex polyhedron, including boundary equilibria. In actor–critic, the attracting-face result further gives an $O_{\text{a.s.}}(n^{-1} \log n)$ constrained-policy value gap. The Online Companion develops the actor–critic proof, projected TD(0), projected SGD, and fast-coordinate projection.

A Selected Proofs of the Main Results

We give the path comparisons underlying the main concentration bounds and the argument that carries Lyapunov control from block endpoints to every iteration. These comprise the two comparisons for the fast recursion, their fast-tracking consequence, the original–auxiliary and auxiliary–ODE comparisons for the slow recursion, and the localized projected-ODE comparison. The step-path Skorokhod representation, supporting Poisson-equation and martingale estimates, and the remaining proofs are collected in the Online Companion.

A.1 Proof of Proposition 6.1

Proof. The triangle inequality and Lemmas 6.3 and 6.4 give, on $\mathcal{G}_x^{(m)}(\delta)$ and for $0 \leq k \leq K_m$,

$$\varepsilon_{n_m+k}^{(x)} \leq \mathcal{W}_m(k) \varepsilon_{n_m}^{(x)} + C_{\text{aux}}(\delta + \alpha_{n_m}) + \frac{C_{\text{mov}}}{q} \frac{\beta_{n_m}}{\alpha_{n_m}}.$$

This proves (6.11), for instance with

$$C_x := \max \left\{ C_{\text{aux}}, \frac{C_{\text{mov}}}{q} \right\}.$$

Assumption 3.7 and the definition of $\mathbf{c}_8$ give $\varepsilon_{n_m}^{(x)} \leq \mathbf{c}_6 + \mathbf{c}_8$. Moreover, monotonicity of β_n/α_n and $\mathcal{W}_m(k+1) = \mathcal{W}_m(k)(1 - q\alpha_{n_m+k})$ yield

$$\sum_{k=0}^{K_m-1} \beta_{n_m+k} \mathcal{W}_m(k) \leq \frac{\beta_{n_m}}{q\alpha_{n_m}} [1 - \mathcal{W}_m(K_m)] \leq \frac{\beta_{n_m}}{q\alpha_{n_m}}.$$

Since $\sum_{r=n_m}^{n_{m+1}-1} \beta_r \leq T+1$, summing the pointwise bound proves (6.12) with $C_{x,T} = (\mathbf{c}_6 + \mathbf{c}_8)/q + (T+1)C_x$. Equation (6.13) is (6.11) at $k = K_m$. $\square$

A.2 Proof of Lemma 6.3

We first control the discrepancy between the original block-reindexed fast iterate $x_k^{(m)}$ and the block-restarted averaged recursion $\tilde{x}_k^{(m)}$. Both recursions use the same slow-parameter path and are unprojected, so their comparison is controlled directly by contraction in the fast state; no parameter-path or Skorokhod-map comparison is required. Subtracting their updates and using the Poisson decomposition separates this discrepancy into a martingale transform, a Poisson residual, and a contractive propagation term. The first term is controlled on $\mathcal{G}_x^{(m)}(\delta)$, while the second is bounded deterministically.

Proof. Fix $m \geq 0$, write $s := n_m$, and recall $q = 1 - \gamma$. Let $\mathcal{R}_{m,k}^x$ and C_{res}^x be as in (S.5.2) and Proposition S.5.1, respectively, and set

$$c := \delta + C_{\text{res}}^x \alpha_s.$$

Subtracting the averaged recursion (6.1) from the original update (1.1), applying the Poisson decomposition (6.5), and iterating from $D_0^{(m)} = 0$ gives

$$\begin{aligned} D_k^{(m)} &= \mathcal{M}_{m,k}^x + \mathcal{R}_{m,k}^x \\ &\quad + \sum_{j=0}^{k-1} \alpha_{s+j} \chi(s+j+1, s+k-1) [h^{(\text{av.})}(x_j^{(m)}; \theta_j^{(m)}) - h^{(\text{av.})}(\tilde{x}_j^{(m)}; \theta_j^{(m)})]. \end{aligned}$$

Define

$$H_k^{(m)} := \sum_{j=0}^{k-1} \alpha_{s+j} \chi(s+j+1, s+k-1) \|D_j^{(m)}\|, \quad H_0^{(m)} := 0.$$

The polynomial step sizes satisfy $0 < \alpha_n < 1$, so the weights $\alpha_{s+j} \chi(s+j+1, s+k-1)$ are nonnegative. Hence the triangle inequality and Assumption 3.4, which makes the averaged fast map γ -contractive in its state argument, give

$$\begin{aligned} &\left\| \sum_{j=0}^{k-1} \alpha_{s+j} \chi(s+j+1, s+k-1) [h^{(\text{av.})}(x_j^{(m)}; \theta_j^{(m)}) - h^{(\text{av.})}(\tilde{x}_j^{(m)}; \theta_j^{(m)})] \right\| \\ &\leq \gamma \sum_{j=0}^{k-1} \alpha_{s+j} \chi(s+j+1, s+k-1) \|D_j^{(m)}\| = \gamma H_k^{(m)}. \end{aligned}$$

On $\mathcal{G}_x^{(m)}(\delta)$, the definition of κ_{d_x} and (6.7) give $\|\mathcal{M}_{m,k}^x\| \leq \kappa_{d_x} \|\mathcal{M}_{m,k}^x\|_\infty < \delta$, while Proposition S.5.1 gives $\|\mathcal{R}_{m,k}^x\| \leq C_{\text{res}}^x \alpha_{n_m} = C_{\text{res}}^x \alpha_s$. Combining these estimates and weakening the strict martingale bound to a non-strict one gives

$$\|D_k^{(m)}\| \leq c + \gamma H_k^{(m)}. \tag{A.1}$$

Moreover,

$$\begin{aligned} H_{k+1}^{(m)} &= (1 - \alpha_{s+k}) H_k^{(m)} + \alpha_{s+k} \|D_k^{(m)}\| \\ &\leq (1 - q \alpha_{s+k}) H_k^{(m)} + \alpha_{s+k} c. \end{aligned}$$

Iteration and telescoping give

$$\begin{aligned} H_k^{(m)} &\leq c \sum_{j=0}^{k-1} \alpha_{s+j} \prod_{\ell=j+1}^{k-1} (1 - q \alpha_{s+\ell}) \\ &= \frac{c}{q} \left[1 - \prod_{\ell=0}^{k-1} (1 - q \alpha_{s+\ell}) \right] \leq \frac{c}{q}. \end{aligned}$$

Substitution into (A.1), together with $1 + \gamma/q = 1/q$, yields

$$\|D_k^{(m)}\| \leq \frac{c}{q} = \frac{\delta + C_{\text{res}}^x \alpha_{n_m}}{q} \leq C_{\text{aux}} (\delta + \alpha_{n_m}),$$

where

$$C_{\text{aux}} := \frac{1}{q} \max\{1, C_{\text{res}}^x\}.$$

This proves (6.9). □

A.3 Proof of Lemma 6.4

Proof. Set $C_{\text{mov}} := L_{fp}(\mathfrak{c}_1 + \mathfrak{c}_2\mathfrak{c}_6 + \mathbf{B}_6)$, and define

$$d_k^{(m)} := \left\| \tilde{x}_k^{(m)} - x^*(\theta_k^{(m)}) \right\|.$$

The fixed-point identity $h^{(\text{av.})}(x^*(\theta_k^{(m)}); \theta_k^{(m)}) = x^*(\theta_k^{(m)})$, Assumption 3.4, and (6.1) give

$$d_{k+1}^{(m)} \leq (1 - q\alpha_{n_m+k})d_k^{(m)} + \left\| x^*(\theta_{k+1}^{(m)}) - x^*(\theta_k^{(m)}) \right\|.$$

By Lemma 3.1, nonexpansiveness of the projection in (1.2), and Assumptions 3.3 and 3.7,

$$\left\| x^*(\theta_{k+1}^{(m)}) - x^*(\theta_k^{(m)}) \right\| \leq C_{\text{mov}}\beta_{n_m+k}.$$

Iteration of the scalar recursion yields

$$\begin{aligned} d_k^{(m)} &\leq \mathcal{W}_m(k)d_0^{(m)} \\ &\quad + C_{\text{mov}} \sum_{j=0}^{k-1} \beta_{n_m+j} \prod_{\ell=j+1}^{k-1} (1 - q\alpha_{n_m+\ell}). \end{aligned}$$

Since β_n/α_n is nonincreasing,

$$\begin{aligned} &\sum_{j=0}^{k-1} \beta_{n_m+j} \prod_{\ell=j+1}^{k-1} (1 - q\alpha_{n_m+\ell}) \\ &\leq \frac{\beta_{n_m}}{\alpha_{n_m}} \sum_{j=0}^{k-1} \alpha_{n_m+j} \prod_{\ell=j+1}^{k-1} (1 - q\alpha_{n_m+\ell}) \\ &= \frac{\beta_{n_m}}{q\alpha_{n_m}} [1 - \mathcal{W}_m(k)] \\ &\leq \frac{\beta_{n_m}}{q\alpha_{n_m}}. \end{aligned}$$

Indeed,

$$\begin{aligned} &q \sum_{j=0}^{k-1} \alpha_{n_m+j} \prod_{\ell=j+1}^{k-1} (1 - q\alpha_{n_m+\ell}) \\ &= \sum_{j=0}^{k-1} \left[\prod_{\ell=j+1}^{k-1} (1 - q\alpha_{n_m+\ell}) - \prod_{\ell=j}^{k-1} (1 - q\alpha_{n_m+\ell}) \right] \\ &= 1 - \prod_{\ell=0}^{k-1} (1 - q\alpha_{n_m+\ell}) = 1 - \mathcal{W}_m(k). \end{aligned}$$

In the first step, we have used

$$\beta_{n_m+j} \leq \frac{\beta_{n_m}}{\alpha_{n_m}} \alpha_{n_m+j}.$$

Finally,

$$d_0^{(m)} = \|x_{n_m} - x^*(\theta_{n_m})\|,$$

which proves (6.10). □

A.4 Proof of Proposition 7.1

Proof. For $0 \leq n \leq K_m$, let

$$A_n := \sup_{0 \leq k \leq n} \|v_k^{(m)} - \tilde{v}_k^{(m)}\|, \quad L_0 := L_f(L_{fp} + 1)L_{\text{Sk}}.$$

Lemma S.11.2 and the Skorokhod comparison in Lemma S.13.1 give

$$\|f^{(\text{av.})}(\theta_\ell^{(m)}; \theta_\ell^{(m)}) - f^{(\text{av.})}(\tilde{\theta}_\ell^{(m)}; \theta_\ell^{(m)})\| \leq L_0 A_\ell.$$

Moreover, since $e_\ell^{(m)} = x_\ell^{(m)} - x^\star(\theta_\ell^{(m)})$, Assumption 3.5 gives

$$\|f(Y_\ell^{(m)}, x_\ell^{(m)}, \theta_\ell^{(m)}) - f(Y_\ell^{(m)}, x^\star(\theta_\ell^{(m)}), \theta_\ell^{(m)})\| \leq L_f \|e_\ell^{(m)}\|.$$

Iterating (7.10), maximizing over terminal indices, and using $\mathcal{G}_\theta^{(m)}(\delta_s)$ and Lemma S.11.3 therefore gives

$$A_n \leq L_0 \sum_{\ell=0}^{n-1} \beta_\ell^{(m)} A_\ell + B_n, \quad B_n := L_f \sum_{\ell=0}^{n-1} \beta_\ell^{(m)} \|e_\ell^{(m)}\| + \delta_s + R_m.$$

Because B_n is nondecreasing, discrete Gronwall yields

$$A_n \leq B_n \exp\{L_0 a(n_m, n_m + n)\}.$$

A final application of Lemma S.13.1 gives $\sup_{0 \leq k \leq n} \|\theta_k^{(m)} - \tilde{\theta}_k^{(m)}\| \leq L_{\text{Sk}} A_n$, which is (7.12). $\square$

A.5 Proof of Proposition 7.2

Proof. Write $F(\vartheta) := f^{(\text{av.})}(\vartheta; \vartheta)$ and $C_\star := B_8$. On $[t^{(m)}(\ell), t^{(m)}(\ell+1))$, let $G_m(s) := f^{(\text{av.})}(\tilde{\theta}_\ell^{(m)}; \theta_\ell^{(m)})$, and set $G_m(H_m) := \mathbf{0}$. The ODE driver satisfies

$$w^{(T_m)}(t) = \theta_{n_m} + \int_0^t F(z^{(T_m)}(s)) ds.$$

The auxiliary step-path driver accumulates only completed grid increments. Consequently, its difference from $\theta_{n_m} + \int_0^t G_m(s) ds$ has norm at most $C_\star \beta_{n_m}$: the difference is the contribution of the single grid interval containing t . Therefore

$$\|w^{(T_m)}(t) - \tilde{v}^{(m), \circ}(t)\| \leq \int_0^t \|F(z^{(T_m)}(s)) - G_m(s)\| ds + C_\star \beta_{n_m}.$$

For almost every $s \in [0, H_m]$, Lemma S.11.2, the identities $z^{(T_m)} = \Gamma(w^{(T_m)})$ and $\tilde{\theta}^{(m), \circ} = \Gamma(\tilde{v}^{(m), \circ})$ from Lemmas S.13.2 and S.13.1, and Corollary 7.1 give

$$\|F(z^{(T_m)}(s)) - G_m(s)\| \leq L_{f, \text{avg.}} L_{\text{Sk}} \mathcal{A}_m(s) + C_\star L_\mu E_m^{(\theta, \tilde{\theta})}(\delta_f, \delta_s).$$

Taking the supremum up to t therefore yields

$$\mathcal{A}_m(t) \leq L_{f, \text{avg.}} L_{\text{Sk}} \int_0^t \mathcal{A}_m(s) ds + C_\star L_\mu t E_m^{(\theta, \tilde{\theta})}(\delta_f, \delta_s) + C_\star \beta_{n_m}.$$

Gronwall proves (7.16). Applying the same finite-horizon Skorokhod estimate to these drivers proves the final assertion. $\square$

A.6 Proof of Theorem 7.1

Proof. By the triangle inequality,

$$\begin{aligned} & \sup_{0 \leq t \leq H_m} \left\| \theta^{(m),\circ}(t) - z^{(T_m)}(t) \right\| \\ & \leq \sup_{0 \leq t \leq H_m} \left\| \theta^{(m),\circ}(t) - \tilde{\theta}^{(m),\circ}(t) \right\| + \sup_{0 \leq t \leq H_m} \left\| \tilde{\theta}^{(m),\circ}(t) - z^{(T_m)}(t) \right\|. \end{aligned}$$

Corollary 7.1 bounds the first term by $E_m^{(\theta, \tilde{\theta})}(\delta_f, \delta_s)$. Proposition 7.2, applied with $t = H_m = a(n_m, n_{m+1})$, bounds the second term by

$$L_{\text{Sk}} e^{L_{f, \text{avg}} \cdot L_{\text{Sk}} a(n_m, n_{m+1})} \left[\mathbf{B}_8 L_\mu a(n_m, n_{m+1}) E_m^{(\theta, \tilde{\theta})}(\delta_f, \delta_s) + \mathbf{B}_8 \beta_{n_m} \right].$$

This proves (7.18) with $B_m(\delta_f, \delta_s)$ defined in (7.17).

For the rate bounds, let $C_{B,T}$ increase from line to line. Since $a(n_m, n_{m+1}) \leq T + 1$, the definitions of B_m and $E_m^{(\theta, \tilde{\theta})}$ imply

$$B_m(\delta_f, \delta_s) \leq C_{B,T} \{S_m(\delta_f) + \delta_s + R_m + \beta_{n_m}\}.$$

Using $R_m \leq C_{B,T} \beta_{n_m}$, (7.13), and $\beta_{n_m} \leq \alpha_{n_m}$ proves (7.19). Finally,

$$\alpha_{n_m} + \frac{\beta_{n_m}}{\alpha_{n_m}} \leq 2(N_0 + n_m)^{-p_{\text{det}}},$$

which gives (7.20). $\square$

A.7 Proof of Theorem 8.2

Proof. By the Borel–Cantelli argument in Section 8, almost surely there exists $m_0 < \infty$ such that $\mathcal{G}_m^{\text{rate}}$ holds for every $m \geq m_0$. Fix such a sample path and write $U_m := J(\theta_{n_m})$ and $e_m^J := L_J C_{B,T} d_m^{\text{rate}}$. At the block endpoint, (8.1) and the Lipschitz continuity of J , followed by nonincrease of J along the ODE flow, $H_m \geq T$, and Assumption 8.2, give

$$U_{m+1} \leq J(z^{(T_m)}(H_m)) + e_m^J \leq U_m - \mathcal{D}_J(U_m) + e_m^J.$$

This proves (8.2).

For $n_m \leq n < n_{m+1}$, let $r_n := t(n) - T_m \in [0, H_m]$. Then $\theta^{(m),\circ}(r_n) = \theta_n$, and the within-block tracking estimate gives

$$J(\theta_n) \leq J(z^{(T_m)}(r_n)) + e_m^J \leq U_m + e_m^J.$$

This proves (8.3).

It remains to prove convergence. If $J_{\text{max}} = 0$, the conclusion is immediate. Otherwise, fix $0 < \eta \leq 2J_{\text{max}}$ and set $c_\eta := \min_{u \in [\eta/2, J_{\text{max}}]} \mathcal{D}_J(u) > 0$. The definition of d_m^{rate} and Lemmas S.15.4 and S.15.5, in their respective cases, give $d_m^{\text{rate}} \rightarrow 0$, and hence $e_m^J \rightarrow 0$. Thus, for all sufficiently large m ,

$$e_m^J \leq \min\{c_\eta/2, \eta/2\}, \quad U_m \geq \eta/2 \implies U_{m+1} \leq U_m - c_\eta/2.$$

The sequence must therefore eventually enter $[0, \eta/2)$. Once it does, the two preceding inequalities show that it remains in $[0, \eta]$: below $\eta/2$, its next increase is at most e_m^J , while on $[\eta/2, \eta]$ it decreases by at least $c_\eta/2$. Since $\eta > 0$ is arbitrary, $U_m \rightarrow 0$. The within-block estimate and $e_m^J \rightarrow 0$ give $J(\theta_n) \rightarrow 0$. Finally, compactness of Θ , continuity of J , and $J^{-1}(\{0\}) = \mathcal{E}\mathcal{Q}_\Theta$ imply $\text{dist}(\theta_n, \mathcal{E}\mathcal{Q}_\Theta) \rightarrow 0$. $\square$

Acknowledgments

During the preparation of this manuscript, the authors used large language models (LLMs) to improve its presentation and clarity.

References

- [1] Jalaaj Bhandari, Daniel Russo, and Raghav Singal. A finite time analysis of temporal difference learning with linear function approximation. *Operations Research*, 69(3):950–973, 2021. doi: 10.1287/opre.2020.2024.
- [2] Vivek S. Borkar. A concentration bound for contractive stochastic approximation. *Systems & Control Letters*, 153:104947, 2021. ISSN 0167-6911. doi: 10.1016/j.sysconle.2021.104947.
- [3] Vivek S Borkar. *Stochastic approximation: a dynamical systems viewpoint*. Springer, 2 edition, 2023.
- [4] Michał Borowski and Błażej Miasojedow. Convergence of projected stochastic approximation algorithm. *arXiv preprint arXiv:2501.08256*, 2025.
- [5] Siddharth Chandak. Nonexpansive mappings in two-time-scale stochastic approximation: Finite-time analysis. *SIAM Journal on Control and Optimization*, 64(4):3024–3049, 2026.
- [6] Siddharth Chandak. $O(1/k)$ finite-time bound for nonlinear two-time-scale stochastic approximation. *IEEE Transactions on Automatic Control*, 2026. URL <https://ieeexplore.ieee.org/document/11523503>.
- [7] Siddharth Chandak and Vivek S. Borkar. A concentration bound for TD(0) with function approximation. *Stochastic Systems*, 16(1):44–60, 2026. doi: 10.1287/stsy.2023.0055.
- [8] Siddharth Chandak, Vivek S Borkar, and Parth Dodhia. Concentration of contractive stochastic approximation and reinforcement learning. *Stochastic Systems*, 12(4):411–430, 2022.
- [9] Siddharth Chandak, Shaan Ul Haque, and Nicholas Bambos. Finite-time bounds for two-time-scale stochastic approximation with arbitrary norm contractions and Markovian noise, 2025. Proceedings of the 64th IEEE Conference on Decision and Control.
- [10] Zaiwei Chen, Siva Theja Maguluri, Sanjay Shakkottai, and Karthikeyan Shanmugam. Finite-sample analysis of contractive stochastic approximation using smooth convex envelopes. *Advances in Neural Information Processing Systems*, 33:8223–8234, 2020.
- [11] Zaiwei Chen, Sheng Zhang, Thinh T. Doan, John-Paul Clarke, and Siva Theja Maguluri. Finite-sample analysis of nonlinear stochastic approximation with applications in reinforcement learning. *Automatica*, 146:110623, 2022.
- [12] Gal Dalal, Balázs Szörényi, Gugu Thoppe, and Shie Mannor. Finite sample analysis of two-timescale stochastic approximation with applications to reinforcement learning. In *Proceedings of the 31st Conference on Learning Theory*, volume 75 of *Proceedings of Machine Learning Research*, pages 1199–1233. PMLR, 2018.
- [13] Thinh T. Doan. Nonlinear two-time-scale stochastic approximation: Convergence and finite-time performance. *IEEE Transactions on Automatic Control*, 68(8):4695–4705, 2022.
- [14] Thinh T Doan. Fast nonlinear two-time-scale stochastic approximation: Achieving $O(1/k)$ finite-sample complexity. *IEEE Transactions on Automatic Control*, 71(1):230–242, 2025.
- [15] Paul Dupuis and Harold J Kushner. Asymptotic behavior of constrained stochastic approximations via the theory of large deviations. *Probability theory and related fields*, 75(2):223–244, 1987.
- [16] Paul Dupuis and Anna Nagurney. Dynamical systems and variational inequalities. *Annals of Operations Research*, 44:7–42, 1993.
- [17] Vijay R Konda and John N Tsitsiklis. Convergence rate of linear two-time-scale stochastic approximation. *The Annals of Applied Probability*, 14(2):796–819, 2004.
- [18] Vijaymohan R. Konda and Vivek S. Borkar. Actor-critic-type learning algorithms for Markov decision processes. *SIAM Journal on control and Optimization*, 38(1):94–123, 1999.

- [19] Harold J. Kushner and G George Yin. *Stochastic approximation and recursive algorithms and applications*. Springer, 2003.
- [20] M. Metivier and P. Priouret. Applications of a Kushner and Clark lemma to general classes of stochastic algorithms. *IEEE Transactions on Information Theory*, 30(2):140–151, 1984. doi: 10.1109/TIT.1984.1056894.
- [21] Martin L. Puterman. *Markov Decision Processes: Discrete Stochastic Dynamic Programming*. John Wiley & Sons, 1994. doi: 10.1002/9780470316887.
- [22] John N. Tsitsiklis and Benjamin Van Roy. An analysis of temporal-difference learning with function approximation. *IEEE Transactions on Automatic Control*, 42(5):674–690, 1997. doi: 10.1109/9.580874.

Supplementary Appendix: Proofs and Additional Results

Contents

S.1	Projected On-Policy Linear TD(0)	38
S.2	Optimization: Projected Stochastic Gradient Descent	40
S.3	One-time-scale specialization	41
S.4	Proofs from Main-Paper Section 3	43
S.5	Supporting Results and Proofs for Main-Paper Section 6	44
S.6	Proofs from Main-Paper Section 7	47
S.7	Proofs from Main-Paper Section 8	48
S.8	Proofs from Main-Paper Section 9 (Actor-Critic)	61
S.9	Extension: Projection of the Fast-Scale Iterate	68
S.10	Auxiliary Proofs Used for Main-Paper Section 6	80
S.11	Auxiliary Results Used for Main-Paper Section 7	82
S.12	Strong monotonicity and exponential stability of projected ODEs	84
S.13	Skorokhod problem	86
S.14	Step-size-related results	88
S.15	Miscellaneous results	90
S.16	Step-size shift and the starting index of the analysis	95
S.17	Rates under contraction of the reduced slow map (Proof of Proposition 8.2)	98
S.18	Stochastic saddle-point learning without strong convexity–concavity	105

Guide to the Supplementary Appendix. Appendix S.1 gives a standalone projected on-policy linear TD(0) application and its last-iterate rate. Appendix S.2 gives the projected-SGD application, including distance and objective-gap rates, and Appendix S.3 states the general one-time-scale specializations under contraction and power Lyapunov decrease.

Appendix S.4 establishes regularity of the stationary law and fast equilibrium. Appendices S.5–S.7 contain, respectively, the remaining fast-scale proofs and supporting results; the remaining slow-scale martingale, Poisson, and projected-path arguments; and the remaining convergence and rate proofs, including those for the one-time-scale specializations. Appendix S.7 also proves Proposition 8.1 and records the harmonic-endpoint sharpness example. The seven selected proofs of the main results appear in Sections A.1–A.7 of the main-paper appendix. Appendix S.5 also records, in Remark S.5.1, the extension of the unprojected fast-scale analysis to an arbitrary fixed fast-coordinate norm. Appendix S.8 verifies the actor–critic application, including controlled-chain coverage, critic contraction, Lyapunov geometry, restriction bias, and improved actor and critic rates. Appendix S.9 gives the optional extension with every-step compact-polyhedral Euclidean projection of the fast iterate, including high-probability tracking and almost-sure rates. Appendices S.10–S.12 collect, in order, the auxiliary fast-scale Poisson estimates, the auxiliary slow-scale Poisson estimates, and the strong-monotonicity consequences for projected ODEs. Appendix S.13 contains the Skorokhod-map tools used for projection. Appendix S.14 derives auxiliary results related to step-sizes. Appendix S.15 contains several miscellaneous results, while Appendix S.16 discusses the shifted-step-size and finite-burn-in alternatives.

For $1/2 < \mathfrak{b} \leq 1$ and $\lambda > 0$, define the rate envelope used in the one-time-scale contraction results below (including the projected TD(0) and projected-SGD applications) by

$$\mathfrak{R}_n(\mathfrak{b}, \lambda) := \begin{cases} (N_0 + n)^{-\mathfrak{b}/2} \sqrt{\log(N_0 + n)}, & \frac{1}{2} < \mathfrak{b} < 1, \\ (N_0 + n)^{-1/2} \sqrt{\log(N_0 + n)}, & \mathfrak{b} = 1, \lambda > \frac{1}{2}, \\ (N_0 + n)^{-1/2} \{\log(N_0 + n)\}^{3/2}, & \mathfrak{b} = 1, \lambda = \frac{1}{2}, \\ (N_0 + n)^{-\lambda}, & \mathfrak{b} = 1, 0 < \lambda < \frac{1}{2}. \end{cases}$$

S.1 Projected On-Policy Linear TD(0)

Projected linear TD(0) approximates a fixed policy's discounted value from one Markov trajectory. Fix a policy π in a finite discounted Markov reward process. Let P be the resulting irreducible transition matrix on $\mathcal{S}$, let d be its stationary distribution, and set $D := \text{diag}(d)$. Given the natural pre-transition history $\mathcal{F}_n^{\text{td}}$, suppose that (R_{n+1}, S_{n+1}) is drawn from a fixed joint reward–transition kernel depending only on S_n , with transition marginal P , and that

$$r(s) := \mathbb{E}[R_{n+1} \mid S_n = s], \quad |R_{n+1}| \leq R_{\max} \quad \text{a.s.}$$

Fix $0 < \gamma_{\text{td}} < 1$, and let V^π denote the fixed policy's value function, so that $V^\pi = r + \gamma_{\text{td}} P V^\pi$. Let $\Psi = (\psi(s)^\top)_{s \in \mathcal{S}} \in \mathbb{R}^{|\mathcal{S}| \times d_w}$ be a full-column-rank feature matrix; its columns span the linear class $V_w := \Psi w$, so $V_w(s) = \psi(s)^\top w$. Let W be a nonempty compact convex polyhedron, assume $w_1 \in W$, and consider

$$\begin{aligned} w_{n+1} &= \text{proj}_W \left[w_n + \eta_n \psi(S_n) \left\{ R_{n+1} + \gamma_{\text{td}} \psi(S_{n+1})^\top w_n - \psi(S_n)^\top w_n \right\} \right], \\ \eta_n &= c_{\text{td}} (N_0 + n)^{-\mathbf{b}}, \quad c_{\text{td}} > 0, \quad \frac{1}{2} < \mathbf{b} \leq 1. \end{aligned} \tag{S.1.1}$$

Here $c_{\text{td}} > 0$ is the multiplicative gain in the TD step size. Define

$$\begin{aligned} A_{\text{td}} &:= \Psi^\top D (I - \gamma_{\text{td}} P) \Psi, \quad q_{\text{td}} := \Psi^\top D r, \\ \mu_{\text{td}} &:= (1 - \gamma_{\text{td}}) \lambda_{\min}(\Psi^\top D \Psi), \quad w_{\text{td}}^* := A_{\text{td}}^{-1} q_{\text{td}}. \end{aligned} \tag{S.1.2}$$

The proof below verifies explicitly that A_{td} is nonsingular and $\mu_{\text{td}} > 0$. Assume that $w_{\text{td}}^* \in W$.

Corollary S.1.1 (Projected linear TD(0)). *Under the preceding conditions, with $\lambda_{\text{td}} := c_{\text{td}} \mu_{\text{td}}$,*

$$\|w_n - w_{\text{td}}^*\| = O_{\text{a.s.}}(\mathfrak{R}_n(\mathbf{b}, \lambda_{\text{td}})).$$

Thus, if $1/2 < \mathbf{b} < 1$,

$$\|w_n - w_{\text{td}}^*\| = O_{\text{a.s.}} \left((N_0 + n)^{-\mathbf{b}/2} \sqrt{\log(N_0 + n)} \right).$$

At $\mathbf{b} = 1$, equivalently,

$$\|w_n - w_{\text{td}}^*\| = \begin{cases} O_{\text{a.s.}} \left((N_0 + n)^{-1/2} \sqrt{\log(N_0 + n)} \right), & \lambda_{\text{td}} > 1/2, \\ O_{\text{a.s.}} \left((N_0 + n)^{-1/2} \{\log(N_0 + n)\}^{3/2} \right), & \lambda_{\text{td}} = 1/2, \\ O_{\text{a.s.}} \left((N_0 + n)^{-\lambda_{\text{td}}} \right), & 0 < \lambda_{\text{td}} < 1/2. \end{cases}$$

When $\mathbf{b} = 1$ and $\lambda_{\text{td}} > 1/2$,

$$\|\Psi(w_n - w_{\text{td}}^*)\|_D^2 = O_{\text{a.s.}} \left(\frac{\log(N_0 + n)}{N_0 + n} \right), \quad \|v\|_D^2 := v^\top D v.$$

Proof. Write $\beta_n = (N_0 + n)^{-\mathbf{b}}$ and define

$$g_{\text{td}}(s, w) := c_{\text{td}} \psi(s) \left\{ r(s) + \gamma_{\text{td}} \sum_{s'} P(s, s') \psi(s')^\top w - \psi(s)^\top w \right\}.$$

Set

$$N_{n+1}^{\text{td}} := c_{\text{td}} \psi(S_n) \left\{ R_{n+1} + \gamma_{\text{td}} \psi(S_{n+1})^\top w_n - \psi(S_n)^\top w_n \right\} - g_{\text{td}}(S_n, w_n).$$

Then (S.1.1) has the form

$$w_{n+1} = \text{proj}_W \left[w_n + \beta_n \{ g_{\text{td}}(S_n, w_n) + N_{n+1}^{\text{td}} \} \right],$$

and

$$\mathbb{E}[N_{n+1}^{\text{td}} \mid \mathcal{F}_n^{\text{td}}] = \mathbf{0}.$$

Because W is compact, $\mathcal{S}$ is finite, and the rewards and features are bounded, $\{N_{n+1}^{\text{td}}\}$ is uniformly bounded. Because the transition kernel P does not depend on w , the required Lipschitz condition on the kernel holds with constant zero. Finite-state irreducibility gives the unique stationary law and the required hitting-time bound, and $g_{\text{td}}(s, \cdot)$ is uniformly Lipschitz on W . Moreover,

$$\sum_s d(s) g_{\text{td}}(s, w) = c_{\text{td}} \{ q_{\text{td}} - A_{\text{td}} w \}.$$

It remains to verify contraction of the projected mean ODE. For $u \in \mathbb{R}^{d_w}$, put $v = \Psi u$. Stationarity of d and Jensen's inequality give

$$\|Pv\|_D^2 = \sum_s d(s) \left(\sum_{s'} P(s, s') v(s') \right)^2 \leq \sum_{s, s'} d(s) P(s, s') v(s')^2 = \|v\|_D^2.$$

Consequently,

$$\begin{aligned} u^\top A_{\text{td}} u &= \|v\|_D^2 - \gamma_{\text{td}} \langle v, Pv \rangle_D \\ &\geq (1 - \gamma_{\text{td}}) \|v\|_D^2 \geq \mu_{\text{td}} \|u\|^2. \end{aligned} \tag{S.1.3}$$

Here the first inequality also uses Cauchy–Schwarz and $\|Pv\|_D \leq \|v\|_D$. Finite-state irreducibility gives $d(s) > 0$ for every s , and full column rank of Ψ therefore makes $\Psi^\top D \Psi$ positive definite. Hence $\mu_{\text{td}} > 0$. If $A_{\text{td}} u = 0$, then (S.1.3) gives

$$0 = u^\top A_{\text{td}} u \geq \mu_{\text{td}} \|u\|^2,$$

so $u = 0$. Thus A_{td} is nonsingular.

Consider

$$\dot{z} = \Pi_W(z, c_{\text{td}} \{ q_{\text{td}} - A_{\text{td}} z \}).$$

For every $z, \bar{w} \in W$ and $h \in \mathbb{R}^{d_w}$, the tangent-cone projection satisfies

$$\langle z - \bar{w}, \Pi_W(z, h) - h \rangle \leq 0.$$

Apply this inequality with $\bar{w} = w_{\text{td}}^*$ and $h = c_{\text{td}} \{ q_{\text{td}} - A_{\text{td}} z \}$. Since $q_{\text{td}} = A_{\text{td}} w_{\text{td}}^*$, equation (S.1.3) gives, for almost every t ,

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|z(t) - w_{\text{td}}^*\|^2 &= \langle z(t) - w_{\text{td}}^*, \Pi_W(z(t), c_{\text{td}} \{ q_{\text{td}} - A_{\text{td}} z(t) \}) \rangle \\ &\leq -c_{\text{td}} (z(t) - w_{\text{td}}^*)^\top A_{\text{td}} (z(t) - w_{\text{td}}^*) \\ &\leq -c_{\text{td}} \mu_{\text{td}} \|z(t) - w_{\text{td}}^*\|^2. \end{aligned}$$

Thus its flow satisfies

$$\left\| \mathcal{S}_t^{(1)}(w) - w_{\text{td}}^* \right\| \leq e^{-c_{\text{td}} \mu_{\text{td}} t} \|w - w_{\text{td}}^*\|.$$

Proposition S.3.1, with $J(w) = \|w - w_{\text{td}}^*\|$, $\rho_T = e^{-c_{\text{td}} \mu_{\text{td}} T}$, and $\lambda_T^{(1)} = c_{\text{td}} \mu_{\text{td}}$, gives the stated rates. The squared value-error conclusion follows from norm equivalence in finite dimensions. $\square$

Let $T_{\text{td}}v := r + \gamma_{\text{td}}Pv$, and let

$$\Pi_D := \Psi(\Psi^\top D\Psi)^{-1}\Psi^\top D$$

be the D -orthogonal projector onto $\text{range}(\Psi)$. Then $\Psi w_{\text{td}}^* = \Pi_D T_{\text{td}}(\Psi w_{\text{td}}^*)$, so Ψw_{td}^* is the usual on-policy projected-Bellman solution; see Tsitsiklis and Van Roy [22]. This Bellman projection differs from proj_W , which safeguards the iterate. If W excludes w_{td}^* , the target becomes the constrained variational-inequality solution for $A_{\text{td}}w - q_{\text{td}}$; assuming $w_{\text{td}}^* \in W$ preserves the usual TD target.

Finite-time TD results differ in sampling model, projection, output, and guarantee. Bhandari et al. [1] study expectation bounds for projected TD driven by a Markov trajectory, whereas Chandak and Borkar [7] obtain uniform-in-time concentration for raw unprojected online TD iterates. The result above instead gives an eventual almost-sure envelope for the raw iterate of explicitly projected TD as a direct specialization of Proposition S.3.1.

S.2 Optimization: Projected Stochastic Gradient Descent

We consider the constrained convex optimization problem $\Phi^* := \min_{\theta \in \Theta} \Phi(\theta)$, where Θ is a compact convex polyhedron and Φ is continuously differentiable on an open neighborhood of Θ , convex on Θ , and has an L_Φ -Lipschitz gradient there. Projected SGD for this problem is

$$\theta_{n+1} = \text{proj}_\Theta [\theta_n - \beta_n G_{n+1}], \quad \beta_n = (N_0 + n)^{-\mathfrak{b}}, \quad \frac{1}{2} < \mathfrak{b} \leq 1, \quad (\text{S.2.1})$$

Let $\mathcal{F}_n^{\text{opt}}$ denote the natural optimization history through time n , including the current iterate and all oracle samples revealed before G_{n+1} is sampled. Assume that, for some deterministic $B_{\text{sgd}} < \infty$,

$$\mathbb{E}[G_{n+1} \mid \mathcal{F}_n^{\text{opt}}] = \nabla \Phi(\theta_n), \quad \|G_{n+1} - \nabla \Phi(\theta_n)\| \leq B_{\text{sgd}} \quad \text{a.s.}$$

This is a one-time-scale singleton-state application, so Markov noise is absent. Its limiting projected gradient flow is

$$\dot{z} = \Pi_\Theta(z, -\nabla \Phi(z)). \quad (\text{S.2.2})$$

Corollary S.2.1 (Almost-sure rates for projected SGD). *Suppose that $1/2 < \mathfrak{b} \leq 1$ and that Φ is m_Φ -strongly convex on an open neighborhood of Θ , and let θ^* be its unique minimizer over Θ . Then,*

$$\|\theta_n - \theta^*\| = O_{\text{a.s.}}(\mathfrak{R}_n(\mathfrak{b}, m_\Phi)), \quad (\text{S.2.3})$$

$$\Phi(\theta_n) - \Phi^* = O_{\text{a.s.}}(\mathfrak{R}_n(\mathfrak{b}, m_\Phi)). \quad (\text{S.2.4})$$

If, in addition, $\theta^* \in \text{int}(\Theta)$, then

$$\Phi(\theta_n) - \Phi^* = O_{\text{a.s.}}(\mathfrak{R}_n(\mathfrak{b}, m_\Phi)^2). \quad (\text{S.2.5})$$

Proof. Write

$$-G_{n+1} = -\nabla \Phi(\theta_n) + \{\nabla \Phi(\theta_n) - G_{n+1}\}.$$

By the oracle assumptions following (S.2.1), the term in braces is a uniformly bounded martingale difference with respect to $\{\mathcal{F}_n^{\text{opt}}\}$. Hence (S.2.1) has the one-time-scale form in Proposition S.3.1, with drift $-\nabla \Phi$. Taking the Markov state to be a singleton makes Assumptions 3.1, 3.2, and 3.6 automatic. The oracle condition gives the bounded martingale difference required by Assumption 3.3,

while $g = -\nabla\Phi$ is L_Φ -Lipschitz on Θ , as required by Assumption 3.5; compactness of Θ also makes this drift bounded. The required flow contraction is verified next.

Under strong convexity,

$$\langle -\nabla\Phi(\theta) + \nabla\Phi(\theta^*), \theta - \theta^* \rangle \leq -m_\Phi \|\theta - \theta^*\|^2.$$

The first-order optimality condition $\langle \nabla\Phi(\theta^*), \theta - \theta^* \rangle \geq 0$ for every $\theta \in \Theta$ is equivalent to $\Pi_\Theta(\theta^*, -\nabla\Phi(\theta^*)) = 0$; see Dupuis and Nagurney [16, Lemma 1]. Thus θ^* is an equilibrium of (S.2.2).

Hence Corollary S.12.1 gives, with $S_t^{(1)}$ denoting the flow in (S.2.2),

$$\|S_t^{(1)}(\theta) - \theta^*\| \leq e^{-m_\Phi t} \|\theta - \theta^*\|, \quad t \geq 0.$$

Thus, for $J(\theta) = \|\theta - \theta^*\|$, one may take $\rho_T = e^{-m_\Phi T}$, so $\lambda_T^{(1)} = m_\Phi$. Proposition S.3.1 then gives $J(\theta_n) = O_{\text{a.s.}}(\mathfrak{R}_n(\mathfrak{b}, m_\Phi))$, proving (S.2.3). Since $\nabla\Phi$ is bounded on the compact set Θ , the fundamental theorem of calculus along the segment from θ^* to θ , followed by Cauchy–Schwarz, gives

$$0 \leq \Phi(\theta) - \Phi^* \leq \left(\sup_{u \in \Theta} \|\nabla\Phi(u)\| \right) \|\theta - \theta^*\|.$$

Thus the objective-function gap is bounded by a finite deterministic multiple of the distance to θ^* . Combining this inequality with the distance rate (S.2.3) proves the objective-gap rate (S.2.4). If $\theta^* \in \text{int}(\Theta)$, then $\nabla\Phi(\theta^*) = 0$, and the descent lemma gives

$$0 \leq \Phi(\theta_n) - \Phi^* \leq \frac{L_\Phi}{2} \|\theta_n - \theta^*\|^2.$$

Thus, in the interior case, the objective-function gap is controlled by a finite deterministic multiple of the squared distance. Squaring the distance rate in (S.2.3) therefore proves (S.2.5). $\square$

S.3 One-time-scale specialization

Recursions without a fast variable can be embedded in the general scheme by taking the fast coordinate, its update, and its noise identically zero. We state the resulting one-time-scale guarantee as follows.

Proposition S.3.1 (One-time-scale contraction specialization). *Consider*

$$\theta_{n+1} = \text{proj}_\Theta(\theta_n + \beta_n \{g(Y_n, \theta_n) + N_{n+1}\}), \quad \beta_n = (N_0 + n)^{-\mathfrak{b}}, \quad (\text{S.3.1})$$

where $1/2 < \mathfrak{b} \leq 1$. Suppose that $\{Y_n\}$ is a controlled Markov chain on the finite state space $\mathcal{Y}$ whose conditional transition kernel, given $\mathcal{F}_n$, is $p^{(\theta_n)}(Y_n, \cdot)$. Assume that each $p^{(\theta)}$ has a unique stationary distribution $\mu^{(\theta)}$, that the kernels depend Lipschitz-continuously on θ , and that they satisfy the uniform hitting-time bound. Suppose also that $g(y, \cdot)$ is Lipschitz on Θ , uniformly in $y \in \mathcal{Y}$, and that $\{N_{n+1}\}_{n \geq 1}$ is a uniformly bounded martingale-difference sequence with respect to $\{\mathcal{F}_n\}$. Define

$$\bar{g}(\theta) := \sum_{y \in \mathcal{Y}} \mu^{(\theta)}(y) g(y, \theta)$$

and consider the projected ODE

$$\dot{z} = \Pi_\Theta(z, \bar{g}(z)).$$

Let $S_s^{(1)}$ denote the time- s flow of this ODE. Assume that a Lipschitz function $J : \Theta \rightarrow \mathbb{R}_+$ is nonincreasing along this ODE and that there exist $T > 0$ and $\rho_T \in (0, 1)$ such that

$$J(S_s^{(1)}(\theta)) \leq \rho_T J(\theta), \quad \theta \in \Theta, \quad s \geq T.$$

If $\mathfrak{b} < 1$, then, almost surely, there are finite random variables C and N such that

$$J(\theta_n) \leq C(N_0 + n)^{-\mathfrak{b}/2} \sqrt{\log(N_0 + n)}, \quad n \geq N.$$

If $\mathfrak{b} = 1$, set

$$\lambda_T^{(1)} := -\frac{\log \rho_T}{T}.$$

Then, almost surely, there are finite random variables C_1 and N_1 such that, for $n \geq N_1$,

$$J(\theta_n) \leq C_1 \begin{cases} (N_0 + n)^{-1/2} \sqrt{\log(N_0 + n)}, & \lambda_T^{(1)} > 1/2, \\ (N_0 + n)^{-1/2} \{\log(N_0 + n)\}^{3/2}, & \lambda_T^{(1)} = 1/2, \\ (N_0 + n)^{-\lambda_T^{(1)}}, & 0 < \lambda_T^{(1)} < 1/2. \end{cases}$$

Proposition S.3.2 (One-time-scale specialization under power Lyapunov decrease). *Consider (S.3.1) under the stochastic regularity and boundedness hypotheses of Proposition S.3.1, but replace its one-block contraction hypothesis as follows. Let $S_s^{(1)}$ denote the flow of the limiting projected ODE, and suppose that $J : \Theta \rightarrow \mathbb{R}_+$ is Lipschitz and nonincreasing along this flow. Let*

$$J_{\max} := \sup_{\theta \in \Theta} J(\theta).$$

Assume that there exist $T > 0$ and a continuous function $\mathcal{D}_J : [0, J_{\max}] \rightarrow \mathbb{R}_+$ such that

$$\mathcal{D}_J(0) = 0, \quad 0 < \mathcal{D}_J(u) \leq u \quad (0 < u \leq J_{\max}),$$

and

$$J(S_T^{(1)}(\theta)) \leq J(\theta) - \mathcal{D}_J(J(\theta)), \quad \theta \in \Theta.$$

Then $J(\theta_n) \rightarrow 0$ almost surely.

If, in addition, for some $p_J > 1$, $\kappa_J > 0$, and $u_J > 0$,

$$\mathcal{D}_J(u) \geq \kappa_J u^{p_J}, \quad 0 \leq u \leq u_J,$$

then the following quantitative conclusions hold. If $\mathfrak{b} < 1$,

$$J(\theta_n) = O_{\text{a.s.}} \left((N_0 + n)^{-\frac{1-\mathfrak{b}}{p_J-1}} + (N_0 + n)^{-\frac{\mathfrak{b}}{2p_J}} (\log(N_0 + n))^{\frac{1}{2p_J}} \right).$$

The optimized choice within the branch $1/2 < \mathfrak{b} < 1$ is

$$\mathfrak{b} = \frac{2p_J}{3p_J - 1},$$

for which

$$J(\theta_n) = O_{\text{a.s.}} \left((N_0 + n)^{-\frac{1}{3p_J-1}} (\log(N_0 + n))^{\frac{1}{2p_J}} \right).$$

At the harmonic endpoint $\mathfrak{b} = 1$, the same power-decrease hypothesis gives instead

$$J(\theta_n) = O_{\text{a.s.}} \left(\{\log(N_0 + n)\}^{-\frac{1}{p_J-1}} \right).$$

S.4 Proofs from Main-Paper Section 3

Proof of Lemma 3.1. The fixed-point identity, Assumption 3.4, and $\|h^{(\text{av.})}(0; \theta)\| \leq B_0$ give

$$\|x^*(\theta)\| = \|h^{(\text{av.})}(x^*(\theta); \theta)\| \leq \gamma \|x^*(\theta)\| + B_0.$$

Hence $\|x^*(\theta)\| \leq B_0/(1 - \gamma)$ uniformly on Θ , so $\mathfrak{c}_8 = \sup_{\theta \in \Theta} \|x^*(\theta)\| < \infty$.

We next prove Lipschitz continuity of $\theta \mapsto \mu^{(\theta)}$. Let $\mathcal{Y}_0 := \mathcal{Y} \setminus \{y^*\}$. If $\mathcal{Y}_0$ is empty, then $\mu^{(\theta)} = \delta_{y^*}$ for every θ ; take $L_\mu = 0$ and proceed directly to the fixed-point argument. Otherwise, let $Q^{(\theta)}$ be the restriction of $p^{(\theta)}$ to $\mathcal{Y}_0 \times \mathcal{Y}_0$. We regard distributions on $\mathcal{Y}_0$ as row vectors. Under the global matrix-norm convention,

$$\|w\|_1 := \sum_{y \in \mathcal{Y}_0} |w(y)|, \quad \|A\|_\infty = \max_{i \in \mathcal{Y}_0} \sum_{j \in \mathcal{Y}_0} |A(i, j)| = \sup_{w \neq \mathbf{0}} \frac{\|wA\|_1}{\|w\|_1}.$$

In particular, $\|\cdot\|_\infty$ is submultiplicative. For $i, j \in \mathcal{Y}_0$,

$$(Q^{(\theta)})^n(i, j) = \mathbb{P}_i^{(\theta)}(Y_n = j, \tau_{y^*} > n).$$

Consequently,

$$\sum_{n \geq 0} \sum_{j \in \mathcal{Y}_0} (Q^{(\theta)})^n(i, j) = \mathbb{E}_i^{(\theta)}[\tau_{y^*}] \leq \mathfrak{c}_5,$$

and therefore

$$Z^{(\theta)} := \sum_{n \geq 0} (Q^{(\theta)})^n = (I - Q^{(\theta)})^{-1}, \quad \|Z^{(\theta)}\|_\infty \leq \mathfrak{c}_5.$$

Order the states as y^* followed by $\mathcal{Y}_0$ and write

$$p^{(\theta)} = \begin{pmatrix} p^{(\theta)}(y^*, y^*) & r^{(\theta)} \\ b^{(\theta)} & Q^{(\theta)} \end{pmatrix}.$$

Set $v^{(\theta)} := r^{(\theta)} Z^{(\theta)}$ and $d_{\text{cyc}}^{(\theta)} := 1 + v^{(\theta)} \mathbf{1}$, where $\mathbf{1}$ is the column vector of ones indexed by $\mathcal{Y}_0$. In this state ordering, $(1, v^{(\theta)})$ is a nonnegative row vector with total mass $d_{\text{cyc}}^{(\theta)}$. Since $v^{(\theta)}(I - Q^{(\theta)}) = r^{(\theta)}$,

$$r^{(\theta)} + v^{(\theta)} Q^{(\theta)} = v^{(\theta)}.$$

Row stochasticity also gives $b^{(\theta)} = \mathbf{1} - Q^{(\theta)} \mathbf{1}$ and hence

$$v^{(\theta)} b^{(\theta)} = r^{(\theta)} \mathbf{1} = 1 - p^{(\theta)}(y^*, y^*).$$

Thus $(1, v^{(\theta)})/d_{\text{cyc}}^{(\theta)}$ is a stationary probability vector. Uniqueness in Assumption 3.1 gives

$$\mu^{(\theta)} = \frac{(1, v^{(\theta)})}{d_{\text{cyc}}^{(\theta)}}.$$

Put $\Delta_\theta := \|\theta - \theta'\|$. Applying Assumption 3.2 to the rows indexed by $\mathcal{Y}_0$ and to the reference-state row gives

$$\|Q^{(\theta)} - Q^{(\theta')}\|_\infty \leq L_p \Delta_\theta, \quad \|r^{(\theta)} - r^{(\theta')}\|_1 \leq L_p \Delta_\theta.$$

The resolvent identity yields

$$Z^{(\theta)} - Z^{(\theta')} = Z^{(\theta)}(Q^{(\theta)} - Q^{(\theta')})Z^{(\theta')},$$

so submultiplicativity and $\|Z^{(\theta)}\|_\infty \leq \mathfrak{c}_5$ give

$$\|Z^{(\theta)} - Z^{(\theta')}\|_\infty \leq \mathfrak{c}_5^2 L_p \Delta_\theta.$$

Since $\|r^{(\theta')}\|_1 \leq 1$,

$$\begin{aligned} \|v^{(\theta)} - v^{(\theta')}\|_1 &\leq \|r^{(\theta)} - r^{(\theta')}\|_1 \|Z^{(\theta)}\|_\infty + \|r^{(\theta')}\|_1 \|Z^{(\theta)} - Z^{(\theta')}\|_\infty \\ &\leq L_p \mathfrak{c}_5 (1 + \mathfrak{c}_5) \Delta_\theta. \end{aligned}$$

Also, $|d_{\text{cyc}}^{(\theta)} - d_{\text{cyc}}^{(\theta')}| \leq \|v^{(\theta)} - v^{(\theta')}\|_1$ and $d_{\text{cyc}}^{(\theta)} \geq 1$. Therefore,

$$\begin{aligned} \|\mu^{(\theta)} - \mu^{(\theta')}\|_1 &\leq \frac{\|v^{(\theta)} - v^{(\theta')}\|_1 + |d_{\text{cyc}}^{(\theta)} - d_{\text{cyc}}^{(\theta')}|}{d_{\text{cyc}}^{(\theta)}} \\ &\leq 2L_p \mathfrak{c}_5 (1 + \mathfrak{c}_5) \|\theta - \theta'\|. \end{aligned}$$

Thus one may take $L_\mu := 2L_p \mathfrak{c}_5 (1 + \mathfrak{c}_5)$.

Finally, contraction of $h^{(\text{av.})}(\cdot; \theta)$ gives

$$\begin{aligned} \|x^*(\theta) - x^*(\theta')\| &\leq \gamma \|x^*(\theta) - x^*(\theta')\| \\ &\quad + \|h^{(\text{av.})}(x^*(\theta'); \theta) - h^{(\text{av.})}(x^*(\theta'); \theta')\|. \end{aligned}$$

By Assumption 3.5, the preceding stationary-distribution estimate, and $\|x^*(\theta')\| \leq \mathfrak{c}_8$, the last term is at most

$$[L_{h,\theta}(\mathfrak{c}_8) + L_\mu \mathfrak{B}_8] \|\theta - \theta'\|.$$

Rearranging proves

$$\|x^*(\theta) - x^*(\theta')\| \leq \frac{L_{h,\theta}(\mathfrak{c}_8) + L_\mu \mathfrak{B}_8}{1 - \gamma} \|\theta - \theta'\|.$$

Thus the displayed coefficient is an admissible choice of L_{fp} . □

S.5 Supporting Results and Proofs for Main-Paper Section 6

S.5.1 Proof of Lemma 6.1

Proof. The controlled Markov property gives

$$\mathbb{E} \left[u_h^{(x_n, \theta_n)}(Y_{n+1}) \mid \mathcal{F}_n \right] = \sum_{y \in \mathcal{Y}} p^{(\theta_n)}(Y_n, y) u_h^{(x_n, \theta_n)}(y).$$

Hence $\mathbb{E}[M''_{n+1} \mid \mathcal{F}_n] = \mathbf{0}$ almost surely. Since $u_h^{(x_n, \theta_n)}$ solves (6.2),

$$\begin{aligned} h(Y_n, x_n, \theta_n) - h^{(\text{av.})}(x_n; \theta_n) \\ = u_h^{(x_n, \theta_n)}(Y_n) - \sum_{y \in \mathcal{Y}} p^{(\theta_n)}(Y_n, y) u_h^{(x_n, \theta_n)}(y). \end{aligned}$$

Adding and subtracting $u_h^{(x_n, \theta_n)}(Y_{n+1})$, and using (6.4), proves (6.5). □

S.5.2 Proof of Lemma 6.2

Proof. Fix $m \geq 0$, $1 \leq k \leq K_m$, and $i \in \{1, \dots, d_x\}$. For these fixed indices, the partial sums in j defining $\mathcal{M}_{m,k}^x(i)$ form a scalar martingale. By Assumptions 3.3 and 3.7, and by part (iii) of Lemma S.10.1, its j th increment is bounded in absolute value by $\alpha_{n_m+j}\chi(n_m+j+1, n_m+k-1)C_{\text{mg}}^{(x)}$. Lemma S.14.1 bounds the corresponding squared-weight sum by $2\alpha_{n_m}$. Applying the Azuma–Hoeffding inequality in Theorem S.15.1, using the coordinate threshold δ/κ_{d_x} , and taking a union bound over $i = 1, \dots, d_x$ and $k = 1, \dots, K_m$ proves (6.8) with the stated constant c_{mg}^x .

Under the arbitrary-norm formulation of Remark S.5.1, let $\|\cdot\|_x$ be the fixed fast-coordinate norm and set $\kappa_x := \sup_{\|u\|_\infty \leq 1} \|u\|_x$ and $\lambda_x := \sup_{\|u\|_x \leq 1} \|u\|_\infty$. For every $r \geq 1$, $\|\Xi_{r+1}\|_x \leq C_{\text{mg}}^{(x)}$, and hence $\|\Xi_{r+1}\|_\infty \leq \lambda_x C_{\text{mg}}^{(x)}$. The good event uses the coordinate threshold δ/κ_x . The same calculation therefore replaces c_{mg}^x by

$$\frac{1}{4\kappa_x^2 \lambda_x^2 (C_{\text{mg}}^{(x)})^2},$$

which is the asserted norm-robust constant. $\square$

S.5.3 Fast Poisson residual

For the residual calculation, set

$$\begin{aligned} A_x &:= \mathbf{c}_1 + \mathbf{B}_6 + \mathbf{c}_6(1 + \mathbf{c}_2), & A_\theta &:= \mathbf{c}_1 + \mathbf{c}_2 \mathbf{c}_6 + \mathbf{B}_6, \\ U_h &:= 2\mathbf{c}_5 \mathbf{B}_6, & C_\theta &:= C_{h,\text{Pois.}} A_\theta, & C_{\text{res}}^x &:= 8C_{h,\text{Pois.}} A_x + 3U_h. \end{aligned}$$

Here A_x and A_θ control the one-step changes of the fast and slow iterates, respectively, while U_h bounds the fast Poisson solution.

The definitions above and Condition 3.1 imply

$$C_\theta \beta_1 \leq C_{h,\text{Pois.}} A_x \alpha_1. \quad (\text{S.5.1})$$

Indeed, $A_x = A_\theta + \mathbf{c}_6 \geq A_\theta$, $C_\theta = C_{h,\text{Pois.}} A_\theta$, and $\beta_1 \leq \alpha_1$. Together with the monotonicity of β_n/α_n , this is the domination used below to control changes in the Poisson solution.

Recall that

$$\Delta_{n+1}^h = u_h^{(x_n, \theta_n)}(Y_n) - u_h^{(x_n, \theta_n)}(Y_{n+1}).$$

For $m \geq 0$ and $0 \leq k \leq K_m$, define the associated local Poisson residual by

$$\mathcal{R}_{m,k}^x := \sum_{j=0}^{k-1} \alpha_{n_m+j} \chi(n_m+j+1, n_m+k-1) \Delta_{n_m+j+1}^h, \quad (\text{S.5.2})$$

with $\mathcal{R}_{m,0}^x := \mathbf{0}$.

Proposition S.5.1 (Local fast Poisson residual). *Under (3.3), with (S.5.1) as established above,*

$$\|\mathcal{R}_{m,k}^x\| \leq C_{\text{res}}^x \alpha_{n_m}, \quad m \geq 0, \quad 0 \leq k \leq K_m. \quad (\text{S.5.3})$$

Proof. The case $k = 0$ is immediate. Fix $k \geq 1$. For the remainder of this proof, write

$$s := n_m, \quad t := n_m + k, \quad u_r := u_h^{(x_r, \theta_r)}, \quad A_r := \alpha_r \chi(r+1, t-1).$$

Summation by parts in (S.5.2) gives

$$\begin{aligned}\mathcal{R}_{m,k}^x &= A_s u_s(Y_s) - A_{t-1} u_{t-1}(Y_t) \\ &\quad + \sum_{r=s+1}^{t-1} A_r [u_r(Y_r) - u_{r-1}(Y_r)] \\ &\quad + \sum_{r=s+1}^{t-1} (A_r - A_{r-1}) u_{r-1}(Y_r),\end{aligned}$$

with empty sums interpreted as zero. By part (iv) of Lemma S.10.1, (S.5.1), and the monotonicity of β_n/α_n ,

$$\|u_r(Y_r) - u_{r-1}(Y_r)\| \leq C_{h,\text{Pois}} A_x \alpha_{r-1} + C_{\theta} \beta_{r-1} \leq 2C_{h,\text{Pois}} A_x \alpha_{r-1}.$$

Since $\alpha_{r-1} \leq 2\alpha_r$, Lemma S.14.1 yields

$$\left\| \sum_{r=s+1}^{t-1} A_r [u_r(Y_r) - u_{r-1}(Y_r)] \right\| \leq 8C_{h,\text{Pois}} A_x \alpha_{t-1}.$$

The subadditivity argument used in Lemma S.14.2 shows that A_r is nondecreasing. Hence

$$\sum_{r=s+1}^{t-1} |A_r - A_{r-1}| = A_{t-1} - A_s \leq \alpha_{t-1}.$$

Part (ii) of Lemma S.10.1 bounds the second interior sum by $U_h \alpha_{t-1}$, and the two boundary terms by $2U_h \alpha_s$. Since $\alpha_{t-1} \leq \alpha_s$,

$$\|\mathcal{R}_{m,k}^x\| \leq (8C_{h,\text{Pois}} A_x + 3U_h) \alpha_s = C_{\text{res}}^x \alpha_{n_m},$$

which proves (S.5.3). □

S.5.4 Proof of Corollary 6.1

Proof. On $\mathcal{G}_x^{(j)}(\delta_j)$, (6.13) gives

$$\varepsilon_{n_{j+1}}^{(x)} \leq \mathcal{W}_j(K_j) \varepsilon_{n_j}^{(x)} + C_x g_j^{(x)}(\delta_j).$$

Iterating this recursion for $j = 0, \dots, m-1$ proves (6.15). The probability estimate (6.16) follows from the union bound and Lemma 6.2. □

S.5.5 Arbitrary fast-coordinate norms

Remark S.5.1 (Choice of norm for the fast coordinate). The fast-scale arguments in Sections 6–8 remain valid if the Euclidean norm on $\mathbb{R}^{d_x}$ is replaced by any fixed norm $\|\cdot\|_x$, provided that the contraction, local Lipschitz, bounded-set, stability, and noise bounds involving the fast coordinate are formulated in that norm. In $\mathcal{B}(R)$, the fast-state radius and the norm of h are then interpreted using $\|\cdot\|_x$, while the norm of f remains Euclidean. Recall the two norm-equivalence constants introduced in the proof of Lemma 6.2:

$$\kappa_x = \sup_{\|u\|_\infty \leq 1} \|u\|_x, \quad \lambda_x = \sup_{\|u\|_x \leq 1} \|u\|_\infty.$$

Thus $\|z\|_x \leq \kappa_x \|z\|_\infty$ and $\|z\|_\infty \leq \lambda_x \|z\|_x$. The first inequality converts the coordinatewise martingale estimate to $\|\cdot\|_x$, while the second converts an x -norm increment bound to a coordinatewise one. The corresponding fast-martingale event and tail constant are recorded in the proof of Lemma 6.2 in Section S.5.2 of the Online Companion. The slow projection and Skorokhod-map arguments remain Euclidean.

S.6 Proofs from Main-Paper Section 7

S.6.1 Proof of Lemma 7.1

Proof. The martingale-difference property follows from

$$\mathbb{E} \left[u_f^{(\theta_n^{(m)})}(Y_{n+1}^{(m)}) \mid \mathcal{F}_n^{(m)} \right] = \sum_{y \in \mathcal{Y}} p^{(\theta_n^{(m)})}(Y_n^{(m)}, y) u_f^{(\theta_n^{(m)})}(y).$$

Since $u_f^{(\theta_k^{(m)})}$ solves (7.6) with the parameter θ equal to $\theta_k^{(m)}$, we have

$$f(Y_k^{(m)}, x^*(\theta_k^{(m)}), \theta_k^{(m)}) - f^{(\text{av.})}(\theta_k^{(m)}; \theta_k^{(m)}) = u_f^{(\theta_k^{(m)})}(Y_k^{(m)}) - \sum_{y \in \mathcal{Y}} p^{(\theta_k^{(m)})}(Y_k^{(m)}, y) u_f^{(\theta_k^{(m)})}(y).$$

Adding and subtracting $u_f^{(\theta_k^{(m)})}(Y_{k+1}^{(m)})$ gives (7.9). $\square$

S.6.2 Proof of Lemma 7.2

Proof. By Assumptions 3.3 and 3.7, and part (iii) of Lemma S.11.1, the coordinate increments of the weighted martingale differences $\beta_\ell^{(m)}[(M')_{\ell+1}^{(m)} + (M'')_{\ell+1}^{(m)}]$ are bounded by $\beta_\ell^{(m)} C_{\text{mg}}^{(\theta)}$. Apply the Azuma–Hoeffding inequality in Theorem S.15.1 coordinatewise, use $\sum_{\ell=0}^{k-1} (\beta_\ell^{(m)})^2 = b(n_m) - b(n_m + k)$, and take a union bound over $i = 1, \dots, d_\theta$ and $k = 1, \dots, K_m$ to obtain the stated bound. $\square$

S.6.3 Proof of Corollary 7.1

Proof. On $\mathcal{G}_x^{(m)}(\delta_f)$, Proposition 6.1, specifically (6.12), gives

$$\sum_{\ell=0}^{K_m-1} \beta_\ell^{(m)} \|e_\ell^{(m)}\| = \sum_{r=n_m}^{n_{m+1}-1} \beta_r \|x_r - x^*(\theta_r)\| \leq S_m(\delta_f).$$

On $\mathcal{G}_\theta^{(m)}(\delta_s)$, Proposition 7.1 with $n = K_m$ therefore yields

$$\sup_{0 \leq k \leq K_m} \|\theta_k^{(m)} - \tilde{\theta}_k^{(m)}\| \leq E_m^{(\theta, \tilde{\theta})}(\delta_f, \delta_s)$$

by (7.15). Since both paths are right-continuous step embeddings on the same slow-time grid,

$$\sup_{0 \leq t \leq H_m} \|\theta^{(m), \circ}(t) - \tilde{\theta}^{(m), \circ}(t)\| = \max_{0 \leq k \leq K_m} \|\theta_k^{(m)} - \tilde{\theta}_k^{(m)}\|.$$

The asserted step-path bound follows. $\square$

S.7 Proofs from Main-Paper Section 8

S.7.1 Proof of Theorem 8.3

Proof. Theorem 8.2 first gives $U_m \rightarrow 0$. Consequently, the local power condition (8.4) and (8.2) imply, eventually almost surely,

$$U_{m+1} \leq U_m - \kappa_J U_m^{p_J} + e_m^J.$$

Suppose first that $\mathfrak{b} < 1$. By (S.15.5),

$$N_0 + n_m \asymp (m+1)^{1/(1-\mathfrak{b})}.$$

The assumed rate envelope for d_m^{rate} therefore gives

$$e_m^J \leq C(m+2)^{-\frac{r}{1-\mathfrak{b}}} \{\log(m+2)\}^{1/2}.$$

Applying Lemma S.15.2 with $\zeta = r/(1-\mathfrak{b})$ and $\ell = 1/2$ yields

$$U_m \leq C(\omega) \left\{ (m+2)^{-1/(p_J-1)} + (m+2)^{-\frac{r}{p_J(1-\mathfrak{b})}} \{\log(m+2)\}^{1/(2p_J)} \right\}.$$

Using (S.15.5) once more gives

$$U_m \leq C(\omega) \left\{ (N_0 + n_m)^{-\frac{1-\mathfrak{b}}{p_J-1}} + (N_0 + n_m)^{-\frac{r}{p_J}} \{\log(N_0 + n_m)\}^{\frac{1}{2p_J}} \right\}.$$

For $n_m \leq n < n_{m+1}$, the within-block contribution in (8.3) is of order $(N_0 + n_m)^{-r} \sqrt{\log(N_0 + n_m)}$, which is eventually bounded by its p_J th root. The comparison between $N_0 + n$ and $N_0 + n_m$ used in the proof of Theorem 8.4 then proves (8.5).

Suppose now that $\mathfrak{b} = 1$. By Lemma S.15.5, $N_0 + n_m \asymp e^{mT}$. Hence the assumed envelope gives

$$e_m^J \leq C e^{-rTm} (m+1)^{1/2}.$$

For every $Q > 0$, the right-hand side is eventually at most $C_Q(m+2)^{-Q}$. Choose $Q > p_J/(p_J-1)$ and apply Lemma S.15.2 with $\zeta = Q$ and $\ell = 0$. Then

$$U_m = O_{\text{a.s.}} \left((m+2)^{-1/(p_J-1)} \right).$$

Here and below, the implicit constants in $O_T(\cdot)$ may depend on the fixed block horizon T . Since $m = T^{-1} \log(N_0 + n_m) + O_T(1)$, and since Lemma S.15.5 makes $N_0 + n$ comparable with $N_0 + n_m$ throughout block m , while e_m^J is exponentially smaller, the within-block estimate yields

$$J(\theta_n) = O_{\text{a.s.}} \left(\{\log(N_0 + n)\}^{-1/(p_J-1)} \right),$$

which is (8.6). □

Remark S.7.1 (Sharpness of the harmonic-endpoint power rate). The logarithmic rate in (8.6) is intrinsic to the assumed power decrease. Indeed, for $u_1 > 0$ and N_0 sufficiently large, the deterministic scalar recursion

$$u_{n+1} = u_n - \frac{\kappa_J}{N_0 + n} u_n^{p_J}, \quad p_J > 1,$$

remains positive and satisfies

$$\lim_{n \rightarrow \infty} u_n [(p_J - 1)\kappa_J \log n]^{1/(p_J - 1)} = 1.$$

To see this, set $v_n := u_n^{-(p_J - 1)}$. A Taylor expansion gives

$$v_{n+1} - v_n = \frac{(p_J - 1)\kappa_J}{N_0 + n} + O((N_0 + n)^{-2}),$$

and summation yields $v_n = (p_J - 1)\kappa_J \log n + O(1)$. Thus a polynomial iteration-wise rate cannot be guaranteed at $\mathbf{b} = 1$ from the power-decrease condition alone. In particular, the rate is $O_{\text{a.s.}}(1/\log n)$ when $p_J = 2$.

S.7.2 Proof of Theorem 8.4

Proof. Since

$$\sum_{m \geq 1} \mathbb{P}((\mathcal{G}_m^{\text{rate}})^c) < \infty,$$

the Borel–Cantelli lemma implies that, almost surely, there exists $m_0 < \infty$ such that $\mathcal{G}_m^{\text{rate}}$ holds for every $m \geq m_0$. Fix such a sample path and write

$$U_m := J(\theta^{(m), \circ}(0)).$$

From (8.1), the L_J -Lipschitz continuity of J , $H_m \geq T$, the ODE contraction assumption, and $\theta^{(m), \circ}(H_m) = \theta^{(m+1), \circ}(0)$, we obtain

$$U_{m+1} \leq \rho_T U_m + L_J C_{B,T} d_m^{\text{rate}}, \quad m \geq m_0.$$

Iteration gives

$$U_{m(n)} \leq \rho_T^{m(n) - m_0} J_{\max} + L_J C_{B,T} \sum_{j=m_0}^{m(n)-1} \rho_T^{m(n)-1-j} d_j^{\text{rate}}.$$

If n lies in block $m = m(n)$, then

$$t(n) - T_m \in [0, H_m], \quad \theta^{(m), \circ}(t(n) - T_m) = \theta_n.$$

Hence (8.1), the Lipschitz continuity of J , and the nonincrease of J along the ODE trajectory imply

$$\begin{aligned} J(\theta_n) &\leq J(z^{(T_m)}(t(n) - T_m)) + L_J C_{B,T} d_m^{\text{rate}} \\ &\leq U_m + L_J C_{B,T} d_m^{\text{rate}}. \end{aligned}$$

Substituting the preceding bound for U_m proves the first claim. For the rate assertion, suppose first that $\mathbf{b} < 1$. Applying Lemma S.15.3, with the present value of r , to the assumed bound on d_m^{rate} shows that both the geometric convolution and the initial transient are bounded by a constant multiple of

$$(N_0 + n_{m(n)})^{-r} \sqrt{\log(N_0 + n_{m(n)})}.$$

Moreover, Lemma S.15.4 gives

$$N_0 + n_{m(n)} \leq N_0 + n \leq N_0 + n_{m(n)} + C_T (N_0 + n_{m(n)})^{\mathbf{b}} \leq 2(N_0 + n_{m(n)})$$

for all sufficiently large $m(n)$. Thus $N_0 + n_{m(n)}$ and $N_0 + n$, as well as their logarithms, are comparable. This proves the $\mathfrak{b} < 1$ assertion.

Suppose $\mathfrak{b} = 1$, and recall λ_T from (8.7). Lemma S.15.6, with $\ell = 1/2$ and $\rho = \rho_T$, bounds the endpoint term U_m and its initial transient. For $n_m \leq n < n_{m+1}$, the within-block estimate established above gives

$$J(\theta_n) \leq U_m + L_J C_{B,T} d_m^{\text{rate}}.$$

Writing $X_m := N_0 + n_m$, Lemma S.15.5 gives

$$X_m \leq N_0 + n < (e^T + 1)X_m.$$

Thus $N_0 + n \asymp_T X_m$, and their logarithms are comparable. Moreover, the current-block term $d_m^{\text{rate}} = O(X_m^{-r} \sqrt{\log X_m})$ is bounded by the corresponding endpoint order in each of the three cases below. Consequently,

$$J(\theta_n) = \begin{cases} O_{\text{a.s.}}((N_0 + n)^{-r} \sqrt{\log(N_0 + n)}), & \lambda_T > r, \\ O_{\text{a.s.}}((N_0 + n)^{-r} \{\log(N_0 + n)\}^{3/2}), & \lambda_T = r, \\ O_{\text{a.s.}}((N_0 + n)^{-\lambda_T}), & 0 < \lambda_T < r, \end{cases}$$

which proves (8.8). □

S.7.3 Proof of Theorem 8.1

Proof. Let

$$\varepsilon_n^{(x)} := \|x_n - x^*(\theta_n)\|,$$

and recall $q = 1 - \gamma$. Lemma 6.2 and the definition of $\delta_{\text{dev}}^{(x)}(m)$ give

$$\mathbb{P}\left(\left[\mathcal{G}_x^{(m)}(\delta_{\text{dev}}^{(x)}(m))\right]^c\right) \leq \frac{1}{2(m+2)^2}.$$

Thus, by Borel–Cantelli, these events hold for all sufficiently large m , almost surely. On those blocks, (6.13) gives

$$\varepsilon_{n_{m+1}}^{(x)} \leq \mathcal{W}_m(K_m) \varepsilon_{n_m}^{(x)} + C_x g_m^{(x)}(\delta_{\text{dev}}^{(x)}(m)).$$

Moreover,

$$\sum_{r=n_m}^{n_{m+1}-1} \alpha_r \geq \frac{\alpha_{n_m}}{\beta_{n_m}} a(n_m, n_{m+1}) \geq T(N_0 + n_m)^{\mathfrak{b}-\mathfrak{a}},$$

so

$$\mathcal{W}_m(K_m) \leq \exp\{-qT(N_0 + n_m)^{\mathfrak{b}-\mathfrak{a}}\}.$$

Suppose first that $\mathfrak{b} < 1$. The preceding factor is at most $1/2$ for all sufficiently large m . Hence, eventually almost surely,

$$\varepsilon_{n_{m+1}}^{(x)} \leq \frac{1}{2} \varepsilon_{n_m}^{(x)} + C_x g_m^{(x)}(\delta_{\text{dev}}^{(x)}(m)).$$

Lemma S.15.4 gives

$$g_m^{(x)}(\delta_{\text{dev}}^{(x)}(m)) \leq C_T (N_0 + n_m)^{-r_x(\mathfrak{a}, \mathfrak{b})} \sqrt{\log(N_0 + n_m)}.$$

Applying Lemma S.15.3 to the endpoint recursion yields

$$\varepsilon_{n_m}^{(x)} \leq C(\omega) (N_0 + n_m)^{-r_x(\mathfrak{a}, \mathfrak{b})} \sqrt{\log(N_0 + n_m)}$$

for all sufficiently large m . If $n_m \leq n < n_{m+1}$, then (6.11) and $\mathcal{W}_m(n - n_m) \leq 1$ give

$$\varepsilon_n^{(x)} \leq \varepsilon_{n_m}^{(x)} + C_x g_m^{(x)} \left(\delta_{\text{dev}}^{(x)}(m) \right).$$

Finally, Lemma S.15.4 gives $K_m \leq C_T(N_0 + n_m)^{\mathfrak{b}}$, and $\mathfrak{b} < 1$ implies $N_0 + n \leq 2(N_0 + n_m)$ for all sufficiently large m . The endpoint rate therefore holds at every iterate n .

Suppose now that $\mathfrak{b} = 1$, and put

$$\varphi_m^x := X_m^{-r_x(\mathfrak{a}, 1)} \sqrt{\log X_m}, \quad X_m := N_0 + n_m.$$

Lemma S.15.5 gives $g_m^{(x)} \left(\delta_{\text{dev}}^{(x)}(m) \right) \leq C_T \varphi_m^x$. Choose any $\eta \in (0, e^{-r_x(\mathfrak{a}, 1)T})$. Since

$$\mathcal{W}_m(K_m) \leq \exp\{-qT X_m^{1-\mathfrak{a}}\},$$

eventually $\mathcal{W}_m(K_m) \leq \eta$. The endpoint recursion and Lemma S.15.6, with $\rho = \eta$ and $\ell = 1/2$, therefore give

$$\varepsilon_{n_m}^{(x)} = O_{\text{a.s.}}(X_m^{-r_x(\mathfrak{a}, 1)} \sqrt{\log X_m}).$$

The pointwise block estimate and Lemma S.15.5 transfer this bound to every n , proving the harmonic-endpoint assertion. $\square$

S.7.4 Proof of Corollary 8.1

Proof. Suppose first that $\mathfrak{b} < 1$. We identify the rate of d_m^{rate} . By Lemma S.15.4, for all sufficiently large m ,

$$D_m \leq (T + 1)\beta_{n_m}, \quad K_m \leq C_T(N_0 + n_m)^{\mathfrak{b}},$$

and

$$L_m^x + L_m^\theta \leq C_T \log(N_0 + n_m).$$

Consequently,

$$\delta_{\text{dev}}^{(x)}(m) \leq C_T(N_0 + n_m)^{-\mathfrak{a}/2} \sqrt{\log(N_0 + n_m)},$$

and

$$\delta_{\text{dev}}^{(\theta)}(m) \leq C_T(N_0 + n_m)^{-\mathfrak{b}/2} \sqrt{\log(N_0 + n_m)}.$$

Moreover,

$$\alpha_{n_m} + \frac{\beta_{n_m}}{\alpha_{n_m}} \leq 2(N_0 + n_m)^{-p_{\text{det}}}.$$

Since

$$r_x(\mathfrak{a}, \mathfrak{b}) \leq p_{\text{det}},$$

all deterministic terms are covered by the same envelope. Hence

$$d_m^{\text{rate}} \leq C_T(N_0 + n_m)^{-r_x(\mathfrak{a}, \mathfrak{b})} \sqrt{\log(N_0 + n_m)}.$$

Theorem 8.4 then gives the same iteration-wise exponent for $J(\theta_n)$, and Theorem 8.1 gives the fast tracking bound.

It remains to optimize $r_x(\mathfrak{a}, \mathfrak{b})$. For fixed $\mathfrak{b} > 3/4$, its maximum is obtained by balancing $\mathfrak{a}/2 = \mathfrak{b} - \mathfrak{a}$, that is, $\mathfrak{a} = 2\mathfrak{b}/3$, and equals $\mathfrak{b}/3$. For $1/2 < \mathfrak{b} \leq 3/4$, its supremum is $\mathfrak{b} - 1/2$,

approached as $\mathfrak{a} \downarrow 1/2$ (and not attained under the strict constraint). In either case, the supremum over $1/2 < \mathfrak{a} < \mathfrak{b} < 1$ is at most $1/3$. For the displayed choice

$$\mathfrak{b} = 1 - 3\varepsilon_{\text{rate}}, \quad \mathfrak{a} = \frac{2\mathfrak{b}}{3} = \frac{2}{3} - 2\varepsilon_{\text{rate}},$$

the restriction $0 < \varepsilon_{\text{rate}} < 1/12$ gives $\mathfrak{b} > 3/4$, so the balancing branch above applies, and

$$\frac{1}{2} < \mathfrak{a} < \mathfrak{b} < 1,$$

and

$$r_x(\mathfrak{a}, \mathfrak{b}) = \min \left\{ \frac{1}{3} - \varepsilon_{\text{rate}}, \frac{1}{3} - \varepsilon_{\text{rate}} \right\} = \frac{1}{3} - \varepsilon_{\text{rate}}.$$

This proves the stated $\mathfrak{b} < 1$ slow and fast optimized rates.

For the harmonic endpoint $\mathfrak{b} = 1$, set $X_m := N_0 + n_m$. Lemma S.15.5 gives

$$D_m \leq C_T X_m^{-1}, \quad K_m \leq C_T X_m, \quad L_m^x + L_m^\theta \leq C_T \log X_m.$$

Consequently,

$$d_m^{\text{rate}} \leq C_T X_m^{-r_x(\mathfrak{a}, 1)} \sqrt{\log X_m}.$$

The slow assertions follow from Theorem 8.4, and the fast assertion follows from Theorem 8.1. Finally, $r_x(\mathfrak{a}, 1)$ is maximized by $\mathfrak{a}/2 = 1 - \mathfrak{a}$, namely $\mathfrak{a} = 2/3$, and its maximum is $1/3$. This proves the harmonic-endpoint claims. $\square$

Remark S.7.2 (Comparison with recent mean-square bounds). For every fixed $0 < \varepsilon_{\text{rate}} < 1/12$, Corollary 8.1 gives an almost-sure distance envelope of order $n^{-1/3+\varepsilon_{\text{rate}}} \sqrt{\log n}$, whose square has trajectory-wise scale $n^{-2/3+2\varepsilon_{\text{rate}}} \log n$. This observation does *not* imply a mean-square bound: passing from an almost-sure envelope with a random prefactor to an expectation requires an additional integrability argument. At the level of exponents, the trajectory-wise squared scale agrees, up to logarithmic and arbitrarily small polynomial losses, with the $O(n^{-2/3})$ mean-square rate for contractive two-time-scale SA with arbitrary-norm contractions and Markov noise in Chandak et al. [9]. Other $O(k^{-a})$ or $O(k^{-1})$ mean-square results have been obtained by Chandak [6], Doan [14] under different assumptions or for modified recursions. Because their algorithms, assumptions, and modes of convergence differ, those results are not directly comparable with the almost-sure envelope above. Our result instead provides blockwise high-probability tracking for projected-ODE slow dynamics without assuming a contractive discrete slow update map. The comparison in this remark concerns the linear-decrease branch, which uses uniform one-block contraction of a Lipschitz Lyapunov function. The nonlinear decrease result gives different exponents and is not being compared here with the cited contractive mean-square bounds.

S.7.5 Proof of Corollary 8.2

Proof. For the case $\mathfrak{b} < 1$, the proof of Corollary 8.1 shows that

$$d_m^{\text{rate}} \leq C_T (N_0 + n_m)^{-r_x(\mathfrak{a}, \mathfrak{b})} \sqrt{\log(N_0 + n_m)}.$$

Theorem 8.3 therefore gives the slow polynomial exponent

$$s(\mathfrak{a}, \mathfrak{b}) := \min \left\{ \frac{1 - \mathfrak{b}}{p_J - 1}, \frac{r_x(\mathfrak{a}, \mathfrak{b})}{p_J} \right\}.$$

Fix $\sigma > 0$. If $s(\mathfrak{a}, \mathfrak{b}) \geq \sigma$, then necessarily

$$\mathfrak{a} \geq 2p_J\sigma, \quad \mathfrak{b} - \mathfrak{a} \geq p_J\sigma, \quad \mathfrak{b} \leq 1 - (p_J - 1)\sigma.$$

Hence

$$3p_J\sigma \leq \mathfrak{b} \leq 1 - (p_J - 1)\sigma,$$

which implies

$$\sigma \leq \frac{1}{4p_J - 1}.$$

Equality is attained at

$$\mathfrak{a} = \frac{2p_J}{4p_J - 1}, \quad \mathfrak{b} = \frac{3p_J}{4p_J - 1}.$$

For this choice,

$$r_x(\mathfrak{a}, \mathfrak{b}) = \frac{p_J}{4p_J - 1}.$$

The slow bound follows from Theorem 8.3, while the fast bound follows from Theorem 8.1.

At the harmonic endpoint $\mathfrak{b} = 1$, take $\mathfrak{a} = 2/3$. The harmonic branch of Theorem 8.3 gives

$$J(\theta_n) = O_{\text{a.s.}}\left(\{\log(N_0 + n)\}^{-1/(p_J-1)}\right),$$

while Theorem 8.1 gives

$$\|x_n - x^*(\theta_n)\| = O_{\text{a.s.}}\left((N_0 + n)^{-1/3} \sqrt{\log(N_0 + n)}\right).$$

□

S.7.6 Proof of Proposition 8.1

Proof. All bounds below hold for the iterates generated by the selected schedule; in particular, the deterministic stability assumption is retained for those iterates. We first establish convergence for both schedules. This step is needed because the second schedule is outside Condition 3.1.

A comparison on fixed fast-time intervals. For either schedule in (8.9), the sequences α_n , β_n , and β_n/α_n are nonincreasing, with

$$0 < \beta_n \leq \alpha_n \leq 1, \quad \frac{\beta_n}{\alpha_n} \longrightarrow 0, \quad t_n \alpha_n \longrightarrow \infty.$$

Both step-size sums diverge. For the second schedule these facts follow directly from

$$\frac{\beta_n}{\alpha_n} = \frac{B}{A \log t_n}, \quad \sum_{n < k} \alpha_n = \frac{A}{2} (\log t_k)^2 + O(1), \quad \sum_{n < k} \beta_n = B \log t_k + O(1).$$

The shift conditions ensure that α_n , β_n , and β_n/α_n are nonincreasing and that $0 < \beta_n \leq \alpha_n \leq 1$ for every $n \geq 1$.

Fix a fast-time block horizon $T_x > 0$, and define deterministic block endpoints

$$k_0 = 1, \quad k_{j+1} = \min \left\{ k > k_j : \sum_{n=k_j}^{k-1} \alpha_n \geq T_x \right\}.$$

For either schedule,

$$k_{j+1} - k_j = O(\alpha_{k_j}^{-1}), \quad \frac{t_{k_{j+1}}}{t_{k_j}} \longrightarrow 1, \quad \sum_{n=k_j}^{k_{j+1}-1} \alpha_n \in [T_x, T_x + \alpha_{k_j}]. \quad (\text{S.7.1})$$

Indeed, $t\alpha(t) \rightarrow \infty$, and each displayed step-size function has logarithmic derivative of magnitude at most $1/t$ for $t \geq t_1$. On intervals of length $O(1/\alpha_{k_j})$, its ratio to its initial value therefore tends to one. This proves (S.7.1), as well as uniform comparability of the steps within each such interval.

Put

$$\xi_{n+1} = h(Y_n, x_n, \theta_n) - h^{(\text{av.})}(x_n; \theta_n) + M_{n+1}.$$

The Poisson equation (6.3) decomposes this term as a bounded martingale difference plus $u_n(Y_n) - u_n(Y_{n+1})$, where $u_n = u_h^{(x_n, \theta_n)}$. The increment and Poisson-sensitivity bounds of Lemma S.10.1 use only the standing assumptions and $\beta_n \leq \alpha_n$; they therefore hold for both schedules. Summation by parts gives, uniformly for $k_j < r \leq k_{j+1}$,

$$\begin{aligned} & \left\| \sum_{n=k_j}^{r-1} \alpha_n \{u_n(Y_n) - u_n(Y_{n+1})\} \right\| \\ & \leq C \left\{ \alpha_{k_j} + \sum_{n=k_j+1}^{r-1} |\alpha_n - \alpha_{n-1}| + \sum_{n=k_j+1}^{r-1} \alpha_{n-1} (\|x_n - x_{n-1}\| + \|\theta_n - \theta_{n-1}\|) \right\} \leq C_{T_x} \alpha_{k_j}. \end{aligned}$$

Here and below deterministic constants may depend on T_x . The martingale part has bounded increments of order α_n , and

$$\sum_{n=k_j}^{k_{j+1}-1} \alpha_n^2 \leq (T_x + 1) \alpha_{k_j}.$$

The maximal martingale inequality, applied coordinatewise, consequently bounds its maximum partial sum above u with probability at most $C \exp\{-cu^2/\alpha_{k_j}\}$. Choosing a sufficiently large multiple of $\sqrt{\alpha_{k_j} \log t_{k_j}}$, the failure probabilities are at most $C t_{k_j}^{-3}$, and are summable since $k_j \geq j+1$. Thus, almost surely,

$$\max_{k_j \leq r \leq k_{j+1}} \left\| \sum_{n=k_j}^{r-1} \alpha_n \xi_{n+1} \right\| = O\left(\sqrt{\alpha_{k_j} \log t_{k_j}}\right). \quad (\text{S.7.2})$$

Restart an averaged recursion at each k_j :

$$\hat{x}_{k_j}^{(j)} = x_{k_j}, \quad \hat{x}_{n+1}^{(j)} = (1 - \alpha_n) \hat{x}_n^{(j)} + \alpha_n h^{(\text{av.})}(\hat{x}_n^{(j)}; \theta_n).$$

The parameter sequence in this recursion is the actual θ_n . The averaged drift $h^{(\text{av.})}(x; \theta) - x$ is $(1 + \gamma)$ -Lipschitz in x , uniformly in θ . Subtracting the two recursions, taking partial sums, and applying discrete Gronwall over a fast-time interval of length at most $T_x + 1$ to (S.7.2) yields

$$\max_{k_j \leq n \leq k_{j+1}} \|x_n - \hat{x}_n^{(j)}\| = O_{\text{a.s.}}\left(\sqrt{\alpha_{k_j} \log t_{k_j}}\right). \quad (\text{S.7.3})$$

This argument estimates martingale partial sums before making a pathwise drift comparison; it introduces no martingale weights containing future slow iterates.

Initial convergence. Let $e_j := \|x_{k_j} - x^*(\theta_{k_j})\|$, and recall $q = 1 - \gamma > 0$. One step of the averaged recursion contracts the distance to the current fast target by $1 - q\alpha_n$, while the target moves by at most $L_{fp}A_\theta\beta_n$. Using (S.7.3) and monotonicity of β_n/α_n , we obtain

$$e_{j+1} \leq e^{-qT_x} e_j + O_{\text{a.s.}} \left(\sqrt{\alpha_{k_j} \log t_{k_j}} + \frac{\beta_{k_j}}{\alpha_{k_j}} \right). \quad (\text{S.7.4})$$

The deterministic envelope on the right tends to zero and has consecutive ratio tending to one by (S.7.1). Geometric comparison gives the same bound for e_j , and the within-interval comparison gives

$$\|x_n - x^*(\theta_n)\| = O_{\text{a.s.}} \left(\sqrt{\alpha_n \log t_n} + \frac{\beta_n}{\alpha_n} \right) \rightarrow 0. \quad (\text{S.7.5})$$

For clarity, the geometric comparison follows by dividing (S.7.4) by its envelope at k_{j+1} . The coefficient of the normalized previous error tends to $e^{-qT_x} < 1$, so the normalized errors are eventually bounded.

Now use the deterministic slow-time blocks n_m with horizon T from Assumption 8.2. Their lengths in slow time lie in $[T, T + \beta_{n_m}]$. For either schedule, $t_{n_{m+1}}/t_{n_m}$ is bounded. In the second schedule it tends to $e^{T/B}$. Apply the slow Poisson decomposition (Lemma 7.1) to $f(y, x^*(\theta), \theta) - f^{(\text{av.})}(\theta; \theta)$. The same partial-sum argument, now with weights β_n , gives almost-sure martingale errors of order $\sqrt{\beta_{n_m} \log t_{n_m}}$ and Poisson residuals of order β_{n_m} . Their failure probabilities can again be chosen at most $Ct_{n_m}^{-3}$. The forcing due to the fast error is bounded by $(T + 1)L_f \sup_{n_m \leq n < n_{m+1}} \|x_n - x^*(\theta_n)\|$. Consequently, the deterministic Skorokhod-map and Gronwall comparison used in the proof of Theorem 7.1 gives a uniform slow-path error on block m tending to zero. This uses only the driver estimates just obtained and the vanishing largest step in the block. In particular, it applies to the second schedule without invoking any pure-power rate formula.

Lipschitz continuity of J and its one-block decrease therefore give

$$J(\theta_{n_{m+1}}) \leq J(\theta_{n_m}) - \mathcal{D}_J(J(\theta_{n_m})) + o(1) \quad \text{almost surely,}$$

with a within-block increase bounded by $o(1)$. Since $\mathcal{D}_J$ is continuous and strictly positive away from zero, the scalar convergence argument in Theorem 8.2 gives $J(\theta_n) \rightarrow 0$. Compactness of Θ and $J^{-1}(\{0\}) = \mathcal{EQ}_\Theta$ then imply $\text{dist}(\theta_n, \mathcal{EQ}_\Theta) \rightarrow 0$ almost surely. These consistency statements are the only inputs from this step needed for the sharper rates.

Distance from a required endpoint. Fix $i \in I$, choose $0 < \eta_i \leq \inf_{\theta \in \mathcal{EQ}_\Theta} s_i f_i^{(\text{av.})}(\theta; \theta)$, and write

$$y_n := s_i(v_i - \theta_{n,i}), \quad X_n := s_i\{f_i(Y_n, x_n, \theta_n) + M'_{n+1,i}\}.$$

Thus $0 \leq y_n \leq u_i - \ell_i$, and coordinatewise box projection gives the exact recursion

$$y_{n+1} = \min\{u_i - \ell_i, [y_n - \beta_n X_n]_+\}. \quad (\text{S.7.6})$$

The standing boundedness assumptions give a deterministic constant $D < \infty$ with $|X_n| \leq D$ for every n .

We next obtain a positive conditional mean over a fixed number of steps from the averaged condition (8.10). Let $\Pi_{\mathcal{EQ}_\Theta}$ denote Euclidean projection onto $\mathcal{EQ}_\Theta$, and define

$$A_i := \sup_{y \in \mathcal{Y}, \zeta \in \mathcal{EQ}_\Theta} |f_i(y, x^*(\zeta), \zeta)|, \quad C_i := L_\mu A_i, \\ g_i(y, \theta) := s_i f_i(y, x^*(\Pi_{\mathcal{EQ}_\Theta} \theta), \Pi_{\mathcal{EQ}_\Theta} \theta) + C_i \text{dist}(\theta, \mathcal{EQ}_\Theta).$$

These functions are bounded and Lipschitz: the maps x^* , $\Pi_{\mathcal{E}\mathcal{Q}_\Theta}$, and the distance function $\text{dist}(\cdot, \mathcal{E}\mathcal{Q}_\Theta)$ are Lipschitz, and the arguments of f remain in a fixed bounded set. The stationary-distribution bound in Lemma 3.1 gives

$$\begin{aligned}\bar{g}_i(\theta) &:= \sum_y \mu^{(\theta)}(y) g_i(y, \theta) \\ &\geq s_i f_i^{(\text{av.})}(\Pi_{\mathcal{E}\mathcal{Q}_\Theta} \theta; \Pi_{\mathcal{E}\mathcal{Q}_\Theta} \theta) - L_\mu A_i \text{dist}(\theta, \mathcal{E}\mathcal{Q}_\Theta) + C_i \text{dist}(\theta, \mathcal{E}\mathcal{Q}_\Theta) \geq \eta_i, \quad \theta \in \Theta.\end{aligned}$$

The added term makes this lower bound valid throughout the box. It vanishes as the iterates approach the face.

Put

$$X_n^0 := g_i(Y_n, \theta_n) + s_i M'_{n+1, i}.$$

For the bounded-set Lipschitz constant L_f from the standing assumptions,

$$|X_n - X_n^0| \leq L_f \|x_n - x^*(\theta_n)\| + \{L_f(1 + L_{fp}) + C_i\} \text{dist}(\theta_n, \mathcal{E}\mathcal{Q}_\Theta) \longrightarrow 0 \quad \text{almost surely.} \quad (\text{S.7.7})$$

The variables X_n^0 are uniformly bounded.

A conditional exponential estimate. The same finite-state Poisson argument as in Lemma S.10.1, applied to the bounded Lipschitz scalar observable g_i , gives solutions

$$u_i^{(\theta)} - P^{(\theta)} u_i^{(\theta)} = g_i(\cdot, \theta) - \bar{g}_i(\theta)$$

with a uniform bound U_i and a uniform Lipschitz constant $L_{u, i}$ in θ . The standing bounds and nonexpansiveness of projection give $\|\theta_{n+1} - \theta_n\| \leq A_\theta \beta_n$. Telescoping the Poisson equation over $k, \dots, k+L-1$, and taking conditional expectations, therefore gives

$$\mathbb{E} \left[\sum_{j=k}^{k+L-1} X_j^0 \middle| \mathcal{F}_k \right] \geq L\eta_i - 2U_i - L_{u, i} A_\theta L \beta_k. \quad (\text{S.7.8})$$

Indeed, the Poisson martingale and the original martingale increments have zero conditional mean. The two endpoint terms contribute at most $2U_i$, and changing the Poisson parameter between consecutive indices contributes at most $L_{u, i} \sum_{j=k}^{k+L-2} \|\theta_{j+1} - \theta_j\|$.

Choose a positive integer L such that $2U_i \leq \eta_i L/4$. For all sufficiently large deterministic k , the last term in (S.7.8) is at most $\eta_i L/4$, and hence

$$\mathbb{E} \left[\sum_{j=k}^{k+L-1} X_j^0 \middle| \mathcal{F}_k \right] \geq \eta_i L/2.$$

Set $W_n := X_n^0 - \eta_i/4$, and let H_i be a deterministic bound on $|W_n|$. For sufficiently small $\lambda > 0$, Taylor's inequality gives, uniformly over all such k ,

$$\begin{aligned}\mathbb{E} \left[\exp \left\{ -\lambda \sum_{j=k}^{k+L-1} W_j \right\} \middle| \mathcal{F}_k \right] \\ \leq 1 - \lambda \eta_i L/4 + \frac{1}{2} \lambda^2 L^2 H_i^2 e^{\lambda L H_i} \leq \rho < 1.\end{aligned}$$

Iterating over consecutive full blocks and bounding the last incomplete block by $(L-1)H_i$ yields constants $C, c > 0$ such that

$$\mathbb{E} \left[\exp \left\{ -\lambda \sum_{j=k}^{n-1} W_j \right\} \middle| \mathcal{F}_k \right] \leq C e^{-c(n-k)}, \quad n > k, \quad (\text{S.7.9})$$

whenever k exceeds a fixed deterministic index.

Rescaling the endpoint distance. Let $t_n = N_0 + n$, $z_n = y_n/\beta_n$, and $r_n = \beta_n/\beta_{n+1}$. Both schedules satisfy $r_n - 1 \leq 1/t_n$. Equation (S.7.6) implies

$$z_{n+1} \leq [r_n z_n - r_n X_n]_+ \leq [z_n - X_n + e_n]_+,$$

where

$$0 \leq e_n := (r_n - 1)(z_n + D) \leq \frac{y_n}{t_n \beta_n} + D/t_n \longrightarrow 0 \quad \text{almost surely.}$$

Here $y_n \rightarrow 0$ and $1/(t_n \beta_n)$ is bounded: it equals t_n^{b-1} for the power schedule and $1/B$ for the second schedule. Together with (S.7.7), this proves that, almost surely, for every sufficiently large n ,

$$z_{n+1} \leq [z_n - W_n]_+. \quad (\text{S.7.10})$$

To handle this random starting index, fix a deterministic N beyond the starting index in (S.7.9), and define

$$Q_N = z_N, \quad Q_{n+1} = [Q_n - W_n]_+, \quad n \geq N.$$

On the event that (S.7.10) holds for every $n \geq N$, monotonicity gives $z_n \leq Q_n$. The union of these events over deterministic N has probability one. Direct iteration of the last recursion gives

$$Q_n = \max \left\{ z_N - \sum_{j=N}^{n-1} W_j, \max_{N < k \leq n} \left(- \sum_{j=k}^{n-1} W_j \right) \right\}.$$

Since $z_N \leq (u_i - \ell_i)/\beta_N$ deterministically, Markov's inequality, (S.7.9), and a union bound over the suffixes give, for every $u > 0$,

$$\begin{aligned} \mathbb{P}(Q_n > u) &\leq C e^{-\lambda u + \lambda(u_i - \ell_i)/\beta_N} e^{-c(n-N)} + C e^{-\lambda u} \sum_{m \geq 0} e^{-cm} \\ &\leq C_N e^{-\lambda u}. \end{aligned}$$

Take $u = K \log(N_0 + n)$, with $\lambda K > 2$. Borel–Cantelli yields $Q_n = O_{\text{a.s.}}(\log n)$ for every fixed N . Intersecting these probability-one events over the countable set of N , and using eventual domination, gives

$$y_n = O_{\text{a.s.}}(\beta_n \log n).$$

There are finitely many constrained coordinates, and $\text{dist}(\theta_n, \mathcal{E}Q_\Theta)^2 = \sum_{i \in I} y_{n,i}^2$. This proves the distance bound. Lipschitz continuity of J and $J = 0$ on $\mathcal{E}Q_\Theta$ imply $J(\theta_n) \leq L_J \text{dist}(\theta_n, \mathcal{E}Q_\Theta)$, proving (8.11). If I is empty, this conclusion is immediate because $\mathcal{E}Q_\Theta = \Theta$.

The fast rate under a common target. Suppose $x^*(\theta) = x^\dagger$ on $\mathcal{E}Q_\Theta$. The fixed-point identity, contraction, and Lipschitz continuity of x^* give

$$\left\| h^{(\text{av.})}(x^\dagger; \theta) - x^\dagger \right\| \leq (1 + \gamma) \left\| x^*(\theta) - x^\dagger \right\| \leq (1 + \gamma) L_{fp} \text{dist}(\theta, \mathcal{E}Q_\Theta). \quad (\text{S.7.11})$$

Return to the fixed fast-time intervals (S.7.1) and their averaged recursions. One averaged step contracts the distance to $x^\dagger$ by $1 - q\alpha_n$ and adds at most $\alpha_n(1 + \gamma)L_{fp} \text{dist}(\theta_n, \mathcal{E}Q_\Theta)$. The sum of the resulting contraction weights is at most $1/q$. Combining this with (S.7.3) and the face bound already proved yields

$$\begin{aligned} \left\| x_{k_{j+1}} - x^\dagger \right\| &\leq e^{-qT_x} \left\| x_{k_j} - x^\dagger \right\| \\ &\quad + O_{\text{a.s.}} \left(\sqrt{\alpha_{k_j} \log t_{k_j}} + \beta_{k_j} \log t_{k_j} \right). \end{aligned}$$

The deterministic envelope again has consecutive ratio tending to one. Geometric comparison and the within-interval bound therefore give

$$\|x_n - x^\dagger\| = O_{\text{a.s.}}\left(\sqrt{\alpha_n \log t_n} + \beta_n \log t_n\right).$$

For both schedules, $\beta_n \log t_n = o(\sqrt{\alpha_n \log t_n})$. Moreover,

$$\|x_n - x^\star(\theta_n)\| \leq \|x_n - x^\dagger\| + L_{fp} \text{dist}(\theta_n, \mathcal{EQ}_\Theta).$$

This proves (8.12). The two stated polynomial exponents follow by substituting the corresponding schedules. No lower bound on A relative to q is needed: each fixed fast-time interval has a fixed contraction factor, and its forcing envelope has consecutive ratio tending to one. $\square$

S.7.7 Proof of Proposition S.3.1

Proof. Repeat the proof of Theorem 7.1 with the fast component absent. The term corresponding to F_m^x is absent.

For $\vartheta, \theta' \in \Theta$, define the two-parameter one-time-scale averaged field

$$g^{(\text{av.})}(\vartheta; \theta') := \sum_{y \in \mathcal{Y}} \mu^{(\theta')}(y) g(y, \vartheta).$$

Thus

$$g^{(\text{av.})}(\theta; \theta) = \bar{g}(\theta).$$

For $\tau \geq 0$, let $z_g^{(\tau)}(\cdot)$ solve

$$\begin{aligned} \dot{z}_g^{(\tau)}(s) &= \Pi_\Theta\left(z_g^{(\tau)}(s), g^{(\text{av.})}(z_g^{(\tau)}(s); z_g^{(\tau)}(s))\right) \\ &= \Pi_\Theta\left(z_g^{(\tau)}(s), \bar{g}(z_g^{(\tau)}(s))\right), \quad z_g^{(\tau)}(0) = \theta^\circ(\tau). \end{aligned}$$

Equivalently,

$$z_g^{(\tau)}(s) = S_s^{(1)}(\theta^\circ(\tau)).$$

Let $u_g^{(\theta)}$ be the hitting-time-normalized Poisson solution of

$$u_g^{(\theta)}(y) - \sum_{j \in \mathcal{Y}} p^{(\theta)}(y, j) u_g^{(\theta)}(j) = g(y, \theta) - \bar{g}(\theta), \quad y \in \mathcal{Y},$$

and define

$$M_{n+1}^g := u_g^{(\theta_n)}(Y_{n+1}) - \sum_{j \in \mathcal{Y}} p^{(\theta_n)}(Y_n, j) u_g^{(\theta_n)}(j).$$

Then $\{M_{n+1}^g\}$ is a martingale-difference sequence and

$$g(Y_n, \theta_n) - \bar{g}(\theta_n) = u_g^{(\theta_n)}(Y_n) - u_g^{(\theta_n)}(Y_{n+1}) + M_{n+1}^g.$$

The one-time-scale analogues of the standing bounded-noise and uniform hitting-time assumptions, together with compactness of Θ , give constants $C_N, C_P^{(1)}, C_L^{(1)} < \infty$ such that

$$\|N_{n+1}\| \leq C_N \quad \text{almost surely for every } n \geq 1, \quad C_P^{(1)} := \sup_{\theta \in \Theta, y \in \mathcal{Y}} \|u_g^{(\theta)}(y)\| < \infty,$$

and

$$\sup_{y \in \mathcal{Y}} \|u_g^{(\theta)}(y) - u_g^{(\theta')}(y)\| \leq C_L^{(1)} \|\theta - \theta'\|, \quad \theta, \theta' \in \Theta.$$

Let

$$A_g := \sup_{y \in \mathcal{Y}, \theta \in \Theta} \|g(y, \theta)\| + C_N < \infty.$$

Because $\theta_r \in \Theta$ and Euclidean projection is nonexpansive,

$$\|\theta_{r+1} - \theta_r\| \leq A_g \beta_r \quad \text{almost surely,} \quad r \geq 1.$$

For $m \geq 1$ and $1 \leq k \leq K_m$, define

$$\mathcal{R}_{m,k}^{(1)} := \sum_{\ell=0}^{k-1} \beta_{n_m+\ell} \left[u_g^{(\theta_{n_m+\ell})}(Y_{n_m+\ell}) - u_g^{(\theta_{n_m+\ell})}(Y_{n_m+\ell+1}) \right].$$

The summation-by-parts argument used in the proof of Lemma S.11.3 gives

$$\max_{1 \leq k \leq K_m} \|\mathcal{R}_{m,k}^{(1)}\| \leq R_m^{(1)},$$

where

$$R_m^{(1)} := 3C_P^{(1)} \beta_{n_m} + C_L^{(1)} A_g D_m = O_T(\beta_{n_m}).$$

Hence, with

$$C_{\text{mg}}^{(\theta,1)} := C_N + 2C_P^{(1)},$$

we have, for every $n \geq 1$,

$$\|N_{n+1} + M_{n+1}^g\| \leq C_{\text{mg}}^{(\theta,1)} \quad \text{almost surely.}$$

For $m \geq 1$ and $\delta > 0$, define the one-time-scale good event

$$\mathcal{G}_{\text{one}}^{(m)}(\delta) := \left\{ \max_{1 \leq k \leq K_m} \left\| \sum_{\ell=0}^{k-1} \beta_{n_m+\ell} (N_{n_m+\ell+1} + M_{n_m+\ell+1}^g) \right\| < \delta \right\}.$$

Coordinatewise Azuma–Hoeffding and a union bound give

$$\mathbb{P}\left([\mathcal{G}_{\text{one}}^{(m)}(\delta)]^c\right) \leq 2d_\theta \sum_{k=1}^{K_m} \exp\left\{-\frac{\delta^2}{2\kappa_{d_\theta}^2 (C_{\text{mg}}^{(\theta,1)})^2 \{b(n_m) - b(n_m + k)\}}\right\}.$$

Set

$$\delta_{\text{dev}}^{(1)}(m) := \kappa_{d_\theta} C_{\text{mg}}^{(\theta,1)} \sqrt{2D_m L_m^\theta}.$$

Since $b(n_m) - b(n_m + k) \leq D_m$, the definition of L_m^θ implies

$$\mathbb{P}\left([\mathcal{G}_{\text{one}}^{(m)}(\delta_{\text{dev}}^{(1)}(m))]^c\right) \leq \frac{1}{2(m+2)^2}.$$

Thus the failure probabilities are summable. These are exactly the substitutions

$$f(y, x^*(\theta), \theta) \mapsto g(y, \theta), \quad f^{(\text{av.})}(\theta; \theta') \mapsto g^{(\text{av.})}(\theta; \theta'), \quad M'_{n+1} \mapsto N_{n+1}, \quad M'''_{n+1} \mapsto M_{n+1}^g$$

in (7.6)–(7.11), with every fast term omitted. The resulting version of Lemma 7.2 and the comparison proof of Theorem 7.1 show that there exist $C_T, C'_T < \infty$ such that, on $\mathcal{G}_{\text{one}}^{(m)}(\delta_{\text{dev}}^{(1)}(m))$,

$$\begin{aligned} \sup_{0 \leq r \leq H_m} \left\| \theta^{(m), \circ}(r) - z_g^{(T_m)}(r) \right\| &\leq C_T \left\{ \delta_{\text{dev}}^{(1)}(m) + R_m^{(1)} + \beta_{n_m} \right\} \\ &\leq C'_T \left\{ \delta_{\text{dev}}^{(1)}(m) + \beta_{n_m} \right\}. \end{aligned}$$

As before,

$$D_m = b(n_m) - b(n_{m+1}) \leq (T+1)\beta_{n_m},$$

and Lemmas S.15.4 and S.15.5, in their respective cases, show that the logarithmic union-bound factor is $O(\log(N_0 + n_m))$. Therefore,

$$\delta_{\text{dev}}^{(1)}(m) \leq C_T(N_0 + n_m)^{-\mathfrak{b}/2} \sqrt{\log(N_0 + n_m)}.$$

Since $\beta_{n_m} = (N_0 + n_m)^{-\mathfrak{b}}$ is of smaller order, the Borel–Cantelli lemma yields, eventually almost surely,

$$\sup_{0 \leq r \leq H_m} \left\| \theta^{(m), \circ}(r) - z_g^{(T_m)}(r) \right\| \leq C_T(N_0 + n_m)^{-\mathfrak{b}/2} \sqrt{\log(N_0 + n_m)}.$$

If $\mathfrak{b} < 1$, the block-contraction argument in the proof of Theorem 8.4, together with Lemma S.15.3, gives the claimed iteration-wise rate. If $\mathfrak{b} = 1$, applying Lemma S.15.6 with $r = \ell = 1/2$ and $\rho = \rho_T$, and then using the within-block comparison in Lemma S.15.5, gives the three cases stated in Proposition S.3.1. $\square$

S.7.8 Proof of Proposition S.3.2

Proof. The proof of Proposition S.3.1 gives, eventually almost surely,

$$\sup_{0 \leq r \leq H_m} \left\| \theta^{(m), \circ}(r) - z_g^{(T_m)}(r) \right\| \leq C_T(N_0 + n_m)^{-\mathfrak{b}/2} \sqrt{\log(N_0 + n_m)}.$$

Set $U_m := J(\theta_{n_m})$. Repeating the endpoint and within-block argument in the proof of Theorem 8.2 gives qualitative convergence and, under the power-decrease condition, the eventual recursion

$$U_{m+1} \leq U_m - \kappa_J U_m^{p_J} + C_T(N_0 + n_m)^{-\mathfrak{b}/2} \sqrt{\log(N_0 + n_m)}.$$

Suppose first that $\mathfrak{b} < 1$. By (S.15.5), the perturbation on the right-hand side is bounded by

$$C(m+2)^{-\frac{\mathfrak{b}}{2(1-\mathfrak{b})}} \{\log(m+2)\}^{1/2}.$$

Applying Lemma S.15.2 with $\zeta = \mathfrak{b}/\{2(1-\mathfrak{b})\}$ and $\ell = 1/2$, and then converting block number back to iteration number gives

$$J(\theta_n) = O_{\text{a.s.}} \left((N_0 + n)^{-\frac{1-\mathfrak{b}}{p_J-1}} + (N_0 + n)^{-\frac{\mathfrak{b}}{2p_J}} \{\log(N_0 + n)\}^{\frac{1}{2p_J}} \right).$$

To optimize the polynomial exponent, consider

$$\min \left\{ \frac{1-\mathfrak{b}}{p_J-1}, \frac{\mathfrak{b}}{2p_J} \right\}.$$

The two terms are equal when

$$\mathfrak{b} = \frac{2p_J}{3p_J - 1},$$

and their common value is $1/(3p_J - 1)$, proving the final claim.

Suppose now that $\mathfrak{b} = 1$. By Lemma S.15.5, the perturbation is

$$O\left(e^{-Tm/2}(m+1)^{1/2}\right),$$

and hence is $O((m+2)^{-Q})$ for every $Q > 0$.

Taking $Q > p_J/(p_J - 1)$ and applying Lemma S.15.2 with $\zeta = Q$ and $\ell = 0$ gives

$$U_m = O_{\text{a.s.}}\left((m+2)^{-1/(p_J-1)}\right).$$

Since $m = T^{-1} \log(N_0 + n_m) + O_T(1)$, the within-block comparison in Lemma S.15.5 yields

$$J(\theta_n) = O_{\text{a.s.}}\left(\{\log(N_0 + n)\}^{-1/(p_J-1)}\right),$$

as claimed. □

S.8 Proofs from Main-Paper Section 9 (Actor-Critic)

We begin by discussing the filtration and sampling conventions used in the proof. For $n \geq 1$, let

$$\mathcal{F}_n^{\text{MDP}} := \sigma(V_1, \vartheta_1, S_1, \{(A_k, C_{k+1}, S_{k+1})\}_{1 \leq k < n}),$$

the history immediately before A_n is drawn. Then V_n , ϑ_n , and $Y_n = S_n$ are $\mathcal{F}_n^{\text{MDP}}$ -measurable. The filtration is taken before A_n is drawn from the current policy; the bounded one-step cost and finite-valued next state are then observed before the update. We now prove Corollary 9.1.

Proof of Corollary 9.1. For $\vartheta \in \Theta_{\text{ac}}$, define

$$P_\vartheta(s, s') := \sum_{a \in \mathcal{A}} \pi_\vartheta(a | s) P(s' | s, a).$$

With $c(s, a)$ as in the finite-MDP description in the main paper, define

$$c_\vartheta(s) := \sum_{a \in \mathcal{A}} \pi_\vartheta(a | s) c(s, a).$$

Take $Y_n = S_n$, and take $\mathcal{F}_n^{\text{MDP}}$ before A_n is drawn. Conditional on this history, $A_n \sim \pi_{\vartheta_n}(\cdot | S_n)$; thereafter, for every $s' \in \mathcal{S}$,

$$\mathbb{E}[C_{n+1} | \mathcal{F}_n^{\text{MDP}}, A_n] = c(S_n, A_n), \quad \mathbb{P}(S_{n+1} = s' | \mathcal{F}_n^{\text{MDP}}, A_n) = P(s' | S_n, A_n).$$

Thus the controlled kernel for Y_n is $P^{(\vartheta)} = P_\vartheta$.

For $s \in \mathcal{S}$, $V \in \mathbb{R}^{|\mathcal{S}|}$, and $\vartheta \in \Theta_{\text{ac}}$, set

$$\bar{\delta}_{\text{ac}}(s, V, \vartheta) := c_\vartheta(s) + \gamma_{\text{disc}} \sum_{s'} P_\vartheta(s, s') V(s') - V(s),$$

$$h_{\text{ac}}(s, V, \vartheta) := V + e_s \bar{\delta}_{\text{ac}}(s, V, \vartheta),$$

and

$$f_{\text{ac}}(s, V, \vartheta) := - \sum_{a \in \mathcal{A}} \pi_{\vartheta}(a \mid s) e_{s,a} \left\{ c(s, a) + \gamma_{\text{disc}} \sum_{s'} P(s' \mid s, a) V(s') - V(s) \right\}.$$

Define

$$M_{n+1}^V := e_{S_n} \left\{ \delta_{n+1}^{\text{ac}} - \bar{\delta}_{\text{ac}}(S_n, V_n, \vartheta_n) \right\}$$

and

$$M_{n+1}^{\vartheta} := -e_{S_n, A_n} \delta_{n+1}^{\text{ac}} - f_{\text{ac}}(S_n, V_n, \vartheta_n).$$

The conditional cost and transition identities above imply

$$\mathbb{E}[\delta_{n+1}^{\text{ac}} \mid \mathcal{F}_n^{\text{MDP}}, A_n] = c(S_n, A_n) + \gamma_{\text{disc}} \sum_{s' \in \mathcal{S}} P(s' \mid S_n, A_n) V_n(s') - V_n(S_n).$$

Averaging this identity over $A_n \sim \pi_{\vartheta_n}(\cdot \mid S_n)$, using the tower property, gives

$$\mathbb{E}[M_{n+1}^V \mid \mathcal{F}_n^{\text{MDP}}] = \mathbb{E}[M_{n+1}^{\vartheta} \mid \mathcal{F}_n^{\text{MDP}}] = \mathbf{0}.$$

Hence (9.1) is precisely (1.1)–(1.2) under the identifications

$$x_n = V_n, \quad \theta_n = \vartheta_n, \quad Y_n = S_n, \quad h = h_{\text{ac}}, \quad f = f_{\text{ac}}.$$

Let $D_{\vartheta} := \text{diag}((d_{\vartheta}(s))_{s \in \mathcal{S}})$ and define

$$T_{\vartheta} V := c_{\vartheta} + \gamma_{\text{disc}} P_{\vartheta} V.$$

The averaged fast map is

$$h_{\text{ac}}^{(\text{av.})}(V; \vartheta) = V + D_{\vartheta}(T_{\vartheta} V - V),$$

whose unique fixed point is V_{ϑ} . For all V, V' ,

$$\left\| h_{\text{ac}}^{(\text{av.})}(V; \vartheta) - h_{\text{ac}}^{(\text{av.})}(V'; \vartheta) \right\|_{\infty} \leq \{1 - (1 - \gamma_{\text{disc}}) d_{\min}^{\text{ac}}\} \|V - V'\|_{\infty}. \quad (\text{S.8.1})$$

Thus $h_{\text{ac}}^{(\text{av.})}(\cdot; \vartheta)$ is a contraction in $\|\cdot\|_{\infty}$ with the common modulus $1 - (1 - \gamma_{\text{disc}}) d_{\min}^{\text{ac}} < 1$, uniformly over $\vartheta \in \Theta_{\text{ac}}$. The critic is bounded: if $B_V := C_{\max}/(1 - \gamma_{\text{disc}})$ and $\|V_n\|_{\infty} \leq B_V$, then its updated coordinate is a convex combination of $V_n(S_n)$ and $C_{n+1} + \gamma_{\text{disc}} V_n(S_{n+1})$, both in $[-B_V, B_V]$, since $0 < \alpha_n \leq 1$. Induction therefore gives

$$\sup_{n \geq 1} \|V_n\|_{\infty} \leq B_V.$$

We apply Remark S.5.1 with $\|\cdot\|_x = \|\cdot\|_{\infty}$. Thus $\kappa_x = \lambda_x = 1$, and the fast contraction, local Lipschitz, bounded-set, stability, and martingale bounds are interpreted in this norm; the slow coordinate and its projection remain Euclidean.

We next verify the localized Lipschitz condition directly for the original update maps. Since the softmax map $\vartheta \mapsto \pi_{\vartheta}$ is continuously differentiable and its Jacobian is continuous, its Jacobian is uniformly bounded on the compact logit box. Moreover, c_{ϑ} and P_{ϑ} are linear functions of π_{ϑ} . Hence, by the line-segment form of the mean-value theorem, there exist finite constants $L_{\pi}, L_c, L_P > 0$ such that, for all $\vartheta, \vartheta' \in \Theta_{\text{ac}}$,

$$\begin{aligned} \max_{s \in \mathcal{S}} \|\pi_{\vartheta}(\cdot \mid s) - \pi_{\vartheta'}(\cdot \mid s)\|_1 &\leq L_{\pi} \|\vartheta - \vartheta'\|, \\ \|c_{\vartheta} - c_{\vartheta'}\|_{\infty} &\leq L_c \|\vartheta - \vartheta'\|, \\ \|P_{\vartheta} - P_{\vartheta'}\|_{\infty} &\leq L_P \|\vartheta - \vartheta'\|. \end{aligned}$$

Recall that $\|\cdot\|_\infty$ denotes the induced matrix infinity norm, equivalently the maximum absolute row-sum norm. Consequently, when $\|V\|_\infty, \|V'\|_\infty \leq R$,

$$\|P_\vartheta V - P_{\vartheta'} V'\|_\infty \leq \|V - V'\|_\infty + L_P R \|\vartheta - \vartheta'\|.$$

Substitution in the definitions of h_{ac} and f_{ac} therefore gives finite constants $L_h(R), L_f(R)$, independent of s , such that

$$\begin{aligned} \|h_{ac}(s, V, \vartheta) - h_{ac}(s, V', \vartheta')\|_\infty &\leq L_h(R) \{\|V - V'\|_\infty + \|\vartheta - \vartheta'\|\}, \\ \|f_{ac}(s, V, \vartheta) - f_{ac}(s, V', \vartheta')\| &\leq L_f(R) \{\|V - V'\|_\infty + \|\vartheta - \vartheta'\|\}. \end{aligned}$$

Thus Assumption 3.5 holds on every bounded critic set; in particular, the invariant cube contains every actual V_n , while the Bellman resolvent gives $\sup_\vartheta \|V_\vartheta\|_\infty \leq B_V$.

On the invariant cube $\|V\|_\infty \leq B_V$, the definitions give

$$\|h_{ac}(s, V, \vartheta)\|_\infty \leq B_V, \quad \|f_{ac}(s, V, \vartheta)\| \leq C_{\max} + (1 + \gamma_{\text{disc}})B_V = 2B_V,$$

uniformly over $s \in \mathcal{S}$ and $\vartheta \in \Theta_{ac}$. These are the required deterministic bounded-set drift envelopes. The definitions of M_{n+1}^V and M_{n+1}^ϑ , together with $|C_{n+1}| \leq C_{\max}$, separately give the bounded martingale-noise estimates in Assumption 3.3. The remaining boundedness and martingale hypotheses follow from the invariant cube and the stated controlled-kernel assumptions. To prove (9.2), observe that full support of the softmax policy gives

$$P_\vartheta(s, s') > 0 \iff P(s' | s, a) > 0 \text{ for some } a \in \mathcal{A}.$$

Thus the strongly connected graph assumed in the main paper is the support graph of every P_ϑ . Hence P_ϑ is irreducible and $d_\vartheta(s) > 0$ for every $s \in \mathcal{S}$ and $\vartheta \in \Theta_{ac}$. If $\vartheta_k \rightarrow \vartheta$, every subsequential limit of d_{ϑ_k} is stationary for P_ϑ , by continuity of $\vartheta \mapsto P_\vartheta$. Uniqueness of the stationary distribution therefore gives $d_{\vartheta_k} \rightarrow d_\vartheta$. Thus $\vartheta \mapsto \min_{s \in \mathcal{S}} d_\vartheta(s)$ is continuous and strictly positive. Compactness of Θ_{ac} proves (9.2). The separately assumed uniform hitting-time bound provides the uniform Poisson-equation bounds required by the abstract theorem. For the actor calculation, define the reduced cost

$$K_{s,a}(\vartheta) := c(s, a) + \gamma_{\text{disc}} \sum_{s'} P(s' | s, a) V_\vartheta(s') - V_\vartheta(s).$$

It satisfies $\sum_a \pi_\vartheta(a | s) K_{s,a}(\vartheta) = 0$. At the fast fixed point, the averaged slow field is

$$F_{ac}(\vartheta) = - \sum_{s,a} q_\vartheta(s, a) e_{s,a} K_{s,a}(\vartheta), \quad q_\vartheta(s, a) := d_\vartheta(s) \pi_\vartheta(a | s). \quad (\text{S.8.2})$$

Here $q_\vartheta(s, a)$ is the stationary activation frequency of coordinate (s, a) . Put

$$\pi_{\min}^{\text{ac}} := \frac{e^{-2B_{ac}}}{|\mathcal{A}|}, \quad q_{\min}^{\text{ac}} := d_{\min}^{\text{ac}} \pi_{\min}^{\text{ac}}.$$

Then $\pi_\vartheta(a | s) \geq \pi_{\min}^{\text{ac}}$ and $q_\vartheta(s, a) \geq q_{\min}^{\text{ac}}$.

Lyapunov decrease. Let

$$L_{ac}(\vartheta) := \sum_{s \in \mathcal{S}} V_\vartheta(s).$$

Encode the coordinates blocked by the box projection by setting $\chi_{s,a}(\vartheta) = 0$ when either $\vartheta(s, a) = B_{ac}$ and $K_{s,a}(\vartheta) < 0$, or $\vartheta(s, a) = -B_{ac}$ and $K_{s,a}(\vartheta) > 0$, and let $\chi_{s,a}(\vartheta) = 1$ otherwise. Thus, almost everywhere along the projected actor ODE,

$$\dot{\vartheta}(s, a) = -q_\vartheta(s, a) \chi_{s,a}(\vartheta) K_{s,a}(\vartheta).$$

Along any absolutely continuous solution of this ODE, differentiating the softmax map gives

$$\dot{\pi}_{\vartheta}(a \mid s) = \pi_{\vartheta}(a \mid s) \left\{ \dot{\vartheta}(s, a) - \sum_b \pi_{\vartheta}(b \mid s) \dot{\vartheta}(s, b) \right\}.$$

Consequently, using $\sum_a \pi_{\vartheta}(a \mid s) K_{s,a}(\vartheta) = 0$, we obtain

$$\begin{aligned} \sum_a K_{s,a}(\vartheta) \dot{\pi}_{\vartheta}(a \mid s) &= \sum_a \pi_{\vartheta}(a \mid s) K_{s,a}(\vartheta) \dot{\vartheta}(s, a) \\ &= - \sum_a \pi_{\vartheta}(a \mid s) q_{\vartheta}(s, a) \chi_{s,a}(\vartheta) K_{s,a}(\vartheta)^2 \leq 0. \end{aligned}$$

Applying Konda and Borkar [18, Lemma 5.4] to the induced policy-space vector field, as in their proof of Lemma 5.10, and summing over s therefore gives

$$\begin{aligned} \frac{d}{dt} J_{B_{\text{ac}}}(\vartheta(t)) &= \frac{d}{dt} L_{\text{ac}}(\vartheta(t)) \\ &\leq - \sum_{s,a} \pi_{\vartheta}(a \mid s) q_{\vartheta}(s, a) \chi_{s,a}(\vartheta) K_{s,a}(\vartheta)^2 \\ &\leq - \pi_{\min}^{\text{ac}} q_{\min}^{\text{ac}} \sum_{s,a} \chi_{s,a}(\vartheta) K_{s,a}(\vartheta)^2. \end{aligned} \tag{S.8.3}$$

Control through the Bellman residual. To express the preceding Lyapunov-decrease estimate in terms of $J_{B_{\text{ac}}}$, we first bound $J_{B_{\text{ac}}}$ by the Bellman residual $\|V_{\vartheta} - T_{B_{\text{ac}}} V_{\vartheta}\|_{\infty}$, and then bound that residual by the square root of the active squared-gap sum. The set of policies generated by the logit box is given by

$$\mathcal{P}_{B_{\text{ac}}} := \left\{ \mu \in \Delta(\mathcal{A}) : \mu(a) \leq e^{2B_{\text{ac}}} \mu(b), \ a, b \in \mathcal{A} \right\}.$$

This is exactly the softmax image of the logit box. The forward inclusion follows from the logit ratios. Conversely, the ratio constraints force every $\mu(a) > 0$ and make the range of $\{\log \mu(a) : a \in \mathcal{A}\}$ at most $2B_{\text{ac}}$. Hence

$$\eta(a) := \log \mu(a) - \frac{1}{2} \left\{ \max_b \log \mu(b) + \min_b \log \mu(b) \right\} \in [-B_{\text{ac}}, B_{\text{ac}}]$$

and $e^{\eta(a)} / \sum_{b \in \mathcal{A}} e^{\eta(b)} = \mu(a)$ for every $a \in \mathcal{A}$, which proves the reverse inclusion. The restricted Bellman operator is

$$[T_{B_{\text{ac}}} V](s) := \min_{\mu \in \mathcal{P}_{B_{\text{ac}}}} \sum_a \mu(a) \left\{ c(s, a) + \gamma_{\text{disc}} \sum_{s'} P(s' \mid s, a) V(s') \right\},$$

and $V_{B_{\text{ac}}}^*$ is its fixed point. Fix s , and for $\eta \in [-B_{\text{ac}}, B_{\text{ac}}]^{|\mathcal{A}|}$ define

$$N_s(\eta) := \sum_a e^{\eta(a)} K_{s,a}(\vartheta), \quad Z_s(\eta) := \sum_a e^{\eta(a)}.$$

Because $N_s(\vartheta(s, \cdot)) = 0$, and every coordinate blocked by the projection is already at its beneficial endpoint,

$$N_s(\eta) \geq -(e^{B_{\text{ac}}} - e^{-B_{\text{ac}}}) \sum_a \chi_{s,a}(\vartheta) |K_{s,a}(\vartheta)|.$$

Indeed, in the difference $N_s(\eta) - N_s(\vartheta(s, \cdot))$, every blocked coordinate contributes nonnegatively, while the sum of the remaining contributions is bounded below by the right-hand side above. Because the softmax image of the logit box is exactly the restricted policy class,

$$0 \leq V_\vartheta(s) - [T_{B_{ac}} V_\vartheta](s) = - \min_{\eta \in [-B_{ac}, B_{ac}]^{|\mathcal{A}|}} \frac{N_s(\eta)}{Z_s(\eta)} \leq \frac{e^{2B_{ac}} - 1}{|\mathcal{A}|} \sum_a \chi_{s,a}(\vartheta) |K_{s,a}(\vartheta)|.$$

Indeed, $N_s(\vartheta(s, \cdot)) = 0$, so the minimum is nonpositive, which gives the left inequality. The preceding lower bound on $N_s(\eta)$, together with $Z_s(\eta) \geq |\mathcal{A}|e^{-B_{ac}}$, gives the right inequality. Since $T_{B_{ac}}$ is a γ_{disc} -contraction in the sup norm,

$$\|V_\vartheta - V_{B_{ac}}^*\|_\infty \leq \frac{1}{1 - \gamma_{\text{disc}}} \|V_\vartheta - T_{B_{ac}} V_\vartheta\|_\infty.$$

Using the componentwise inequality $V_\vartheta \geq V_{B_{ac}}^*$, we therefore have

$$\begin{aligned} J_{B_{ac}}(\vartheta) &= \sum_{s \in \mathcal{S}} \{V_\vartheta(s) - V_{B_{ac}}^*(s)\} \\ &\leq |\mathcal{S}| \|V_\vartheta - V_{B_{ac}}^*\|_\infty \\ &\leq \frac{|\mathcal{S}|}{1 - \gamma_{\text{disc}}} \|V_\vartheta - T_{B_{ac}} V_\vartheta\|_\infty \\ &\leq \frac{|\mathcal{S}|(e^{2B_{ac}} - 1)}{|\mathcal{A}|(1 - \gamma_{\text{disc}})} \max_{s \in \mathcal{S}} \sum_a \chi_{s,a}(\vartheta) |K_{s,a}(\vartheta)|. \end{aligned}$$

Finally, Cauchy–Schwarz and $\chi_{s,a} \in \{0, 1\}$ bound the preceding maximum by $\sqrt{|\mathcal{A}|} \{\sum_{s,a} \chi_{s,a}(\vartheta) K_{s,a}(\vartheta)^2\}^{1/2}$. Therefore,

$$J_{B_{ac}}(\vartheta) \leq A_{B_{ac}} \left\{ \sum_{s,a} \chi_{s,a}(\vartheta) K_{s,a}(\vartheta)^2 \right\}^{1/2}, \quad A_{B_{ac}} := \frac{|\mathcal{S}|(e^{2B_{ac}} - 1)}{(1 - \gamma_{\text{disc}})\sqrt{|\mathcal{A}|}}. \quad (\text{S.8.4})$$

Combining (S.8.3) and (S.8.4) yields

$$\frac{d}{dt} J_{B_{ac}}(\vartheta(t)) \leq -\kappa_{B_{ac}}^{\text{ac}} J_{B_{ac}}(\vartheta(t))^2 \quad \text{for almost every } t, \quad \kappa_{B_{ac}}^{\text{ac}} := \frac{\pi_{\min}^{\text{ac}} q_{\min}^{\text{ac}}}{A_{B_{ac}}^2} > 0. \quad (\text{S.8.5})$$

The map $t \mapsto J_{B_{ac}}(\vartheta(t))$ is absolutely continuous. If its initial value is positive, applying the chain rule to its reciprocal while it remains positive and integrating (S.8.5) gives the following bound; if it reaches zero, nonnegativity and monotonicity keep it there. Thus,

$$J_{B_{ac}}(\vartheta(t)) \leq \frac{J_{B_{ac}}(\vartheta(0))}{1 + \kappa_{B_{ac}}^{\text{ac}} t J_{B_{ac}}(\vartheta(0))}. \quad (\text{S.8.6})$$

Equilibrium set and finite-time convergence of the actor ODE. To verify the uniform one-block Lyapunov contraction condition required by Theorem 8.4, we first identify the equilibrium set of the projected actor ODE and then show that every trajectory reaches this set within a uniform finite time. Define

$$K_{s,a}^* := c(s, a) + \gamma_{\text{disc}} \sum_{s'} P(s' \mid s, a) V_{B_{ac}}^*(s') - V_{B_{ac}}^*(s)$$

and

$$\mathcal{E}_{B_{ac}} := \{\vartheta \in \Theta_{ac} : \vartheta(s, a) = B_{ac} \text{ if } K_{s,a}^* < 0, \\ \vartheta(s, a) = -B_{ac} \text{ if } K_{s,a}^* > 0\}.$$

Coordinates with $K_{s,a}^* = 0$ are unrestricted. For each state s and $\eta \in [-B_{ac}, B_{ac}]^{|\mathcal{A}|}$, define

$$N_s^*(\eta) := \sum_a e^{\eta(a)} K_{s,a}^*.$$

Because $V_{B_{ac}}^*$ is the fixed point of the restricted Bellman operator and the softmax image of the logit box is exactly $\mathcal{P}_{B_{ac}}$,

$$0 = [T_{B_{ac}} V_{B_{ac}}^*](s) - V_{B_{ac}}^*(s) = \min_{\eta \in [-B_{ac}, B_{ac}]^{|\mathcal{A}|}} \frac{N_s^*(\eta)}{Z_s(\eta)}.$$

The claim $J_{B_{ac}}(\vartheta) = 0 \iff \vartheta \in \mathcal{E}_{B_{ac}}$ identifies both the hitting target and, as shown below, the actor ODE equilibria. First suppose $J_{B_{ac}}(\vartheta) = 0$. Since $V_\vartheta - V_{B_{ac}}^*$ is componentwise nonnegative, this implies $V_\vartheta = V_{B_{ac}}^*$. The policy-evaluation fixed-point equation then gives

$$N_s^*(\vartheta(s, \cdot)) = Z_s(\vartheta(s, \cdot)) \{[T_\vartheta V_{B_{ac}}^*](s) - V_{B_{ac}}^*(s)\} = 0.$$

If $K_{s,a}^* > 0$ but $\vartheta(s, a) > -B_{ac}$, lowering only that logit makes N_s^* strictly negative. Since $Z_s > 0$, this contradicts the preceding restricted Bellman minimum. Likewise, if $K_{s,a}^* < 0$ but $\vartheta(s, a) < B_{ac}$, raising only that logit makes N_s^* strictly negative. Hence $\vartheta \in \mathcal{E}_{B_{ac}}$.

Conversely, suppose $\vartheta \in \mathcal{E}_{B_{ac}}$. The endpoint signs imply, for every feasible η ,

$$N_s^*(\eta) - N_s^*(\vartheta(s, \cdot)) = \sum_a (e^{\eta(a)} - e^{\vartheta(s,a)}) K_{s,a}^* \geq 0.$$

The restricted Bellman minimum is zero, so every feasible numerator is nonnegative. Moreover, compactness gives a minimizer $\bar{\eta}_s$ with $N_s^*(\bar{\eta}_s) = 0$. The preceding comparison, together with the feasibility of $\vartheta(s, \cdot)$, therefore gives $N_s^*(\vartheta(s, \cdot)) = 0$. Thus

$$[T_\vartheta V_{B_{ac}}^*](s) - V_{B_{ac}}^*(s) = \frac{N_s^*(\vartheta(s, \cdot))}{Z_s(\vartheta(s, \cdot))} = 0.$$

Since this holds for every state, $V_{B_{ac}}^*$ is the fixed point of T_ϑ . Uniqueness of that fixed point gives $V_\vartheta = V_{B_{ac}}^*$, and hence

$$J_{B_{ac}}(\vartheta) = 0 \iff \vartheta \in \mathcal{E}_{B_{ac}}. \quad (\text{S.8.7})$$

If $\vartheta \in \mathcal{E}_{B_{ac}}$, the equality $V_\vartheta = V_{B_{ac}}^*$ gives $K_{s,a}(\vartheta) = K_{s,a}^*$, and the endpoint conditions give $\chi_{s,a}(\vartheta) K_{s,a}(\vartheta) = 0$ for every (s, a) . Conversely, if these products vanish, the residual bound gives $J_{B_{ac}}(\vartheta) = 0$, so (S.8.7) yields $\vartheta \in \mathcal{E}_{B_{ac}}$. Since $q_\vartheta(s, a) > 0$, the products vanish exactly when the projected actor velocity vanishes. Thus $\mathcal{E}_{B_{ac}}$, equivalently the zero set of $J_{B_{ac}}$, is precisely the equilibrium set of the projected actor ODE.

If $K_{s,a}^* = 0$ for every (s, a) , then $\mathcal{E}_{B_{ac}} = \Theta_{ac}$, $J_{B_{ac}}$ is identically zero, and we set $T_{\text{hit}} := 0$. The finite-hitting property is then immediate. Henceforth suppose that the nonzero restricted-gap set is nonempty, and let

$$\Delta_{B_{ac}} := \min_{K_{s,a}^* \neq 0} |K_{s,a}^*| > 0, \quad \eta_{B_{ac}} := \frac{\Delta_{B_{ac}}}{2(1 + \gamma_{\text{disc}})}.$$

Since $V_{\vartheta} - V_{B_{ac}}^*$ is componentwise nonnegative,

$$\|V_{\vartheta} - V_{B_{ac}}^*\|_{\infty} \leq J_{B_{ac}}(\vartheta),$$

and hence

$$|K_{s,a}(\vartheta) - K_{s,a}^*| \leq (1 + \gamma_{\text{disc}})J_{B_{ac}}(\vartheta).$$

By (S.8.6), every ODE trajectory enters the sublevel $J_{B_{ac}} \leq \eta_{B_{ac}}$ within time at most

$$T_{\text{ent}} := \frac{1}{\kappa_{B_{ac}}^{\text{ac}} \eta_{B_{ac}}},$$

with entry time zero if it starts in that sublevel. This sublevel is invariant because $J_{B_{ac}}$ is nonincreasing. Thereafter every nonzero $K_{s,a}^*$ has its final sign and $|K_{s,a}(\vartheta)| \geq \Delta_{B_{ac}}/2$. Each corresponding logit therefore moves toward its required endpoint at speed at least $q_{\min}^{\text{ac}} \Delta_{B_{ac}}/2$, and reaches it within an additional time at most

$$\frac{4B_{ac}}{q_{\min}^{\text{ac}} \Delta_{B_{ac}}}.$$

It remains there because the sign is preserved. In the nondegenerate case one may therefore take

$$T_{\text{hit}} := T_{\text{ent}} + \frac{4B_{ac}}{q_{\min}^{\text{ac}} \Delta_{B_{ac}}} < \infty.$$

Together with the choice $T_{\text{hit}} := 0$ in the degenerate case, in either case

$$J_{B_{ac}}(S_t(\vartheta)) = 0, \quad \vartheta \in \Theta_{ac}, \quad t \geq T_{\text{hit}}. \quad (\text{S.8.8})$$

For any block horizon $T \geq T_{\text{hit}}$ and any $\rho_T \in (0, 1)$, the uniform one-block contraction condition follows.

Approximation error from the bounded-logit restriction. It remains to bound the difference between the optimal value $V_{B_{ac}}^*$ over policies representable by logits in $[-B_{ac}, B_{ac}]^{|\mathcal{A}|}$ and the unrestricted optimal value V^* . For each state s , choose an unrestricted optimal action $a^*(s)$, and let $\hat{\vartheta} \in \Theta_{ac}$ assign logit B_{ac} to $a^*(s)$ and logit $-B_{ac}$ to every other action. Under $\pi_{\hat{\vartheta}}$, the total probability of the other actions is at most $(|\mathcal{A}| - 1)e^{-2B_{ac}}$. Bellman optimality and $\|V^*\|_{\infty} \leq C_{\max}/(1 - \gamma_{\text{disc}})$ give, for every $s \in \mathcal{S}$ and $a \in \mathcal{A}$,

$$0 \leq c(s, a) + \gamma_{\text{disc}} \sum_{s' \in \mathcal{S}} P(s' | s, a) V^*(s') - V^*(s) \leq \frac{2C_{\max}}{1 - \gamma_{\text{disc}}},$$

with equality on the left when $a = a^*(s)$. Consequently,

$$\|T_{\hat{\vartheta}} V^* - V^*\|_{\infty} \leq \frac{2(|\mathcal{A}| - 1)C_{\max}e^{-2B_{ac}}}{1 - \gamma_{\text{disc}}}.$$

Since $T_{\hat{\vartheta}}$ is a γ_{disc} -contraction, the fixed-point residual inequality gives

$$\|V_{\hat{\vartheta}} - V^*\|_{\infty} \leq \frac{1}{1 - \gamma_{\text{disc}}} \|T_{\hat{\vartheta}} V^* - V^*\|_{\infty}.$$

Finally, optimality over the unrestricted and restricted policy classes gives $V^* \leq V_{B_{ac}}^* \leq V_{\hat{\vartheta}}$ componentwise, and therefore

$$\|V_{B_{ac}}^* - V^*\|_{\infty} \leq \frac{2(|\mathcal{A}| - 1)C_{\max}e^{-2B_{ac}}}{(1 - \gamma_{\text{disc}})^2}.$$

Summing over states proves the restriction-bias bound in the corollary.

It remains to apply Proposition 8.1. The Bellman resolvent and smoothness of the softmax map on the compact box show that $J_{B_{ac}}$ is Lipschitz. Taking $T = \max\{1, T_{\text{hit}}\}$ and $\rho_T = e^{-T}$, (S.8.8) verifies the one-block Lyapunov decrease condition needed for initial convergence. The equilibrium set $\mathcal{E}_{B_{ac}}$ is a box face, and $V_{\vartheta} = V_{B_{ac}}^*$ throughout that face. For a coordinate fixed on the face, the outward sign is $-\text{sgn}(K_{s,a}^*)$, so its signed averaged drift satisfies

$$q_{\vartheta}(s, a)|K_{s,a}^*| \geq q_{\min}^{\text{ac}} \Delta_{B_{ac}} > 0.$$

If no coordinate is fixed, the outward-drift condition is vacuous and $J_{B_{ac}} \equiv 0$. All other model, Lipschitz, and contraction assumptions were verified above. Both schedules in (8.9) have $0 < \alpha_n \leq 1$, so the critic cube remains invariant for either choice. Proposition 8.1, with common fast target $V_{B_{ac}}^*$, gives the component bounds and the joint rate (9.5). Together with the restriction-bias estimate already proved, this establishes Corollary 9.1. $\square$

S.9 Extension: Projection of the Fast-Scale Iterate

This section gives an optional extension of the framework in which the fast iterate is projected at every iteration. The main-paper recursion and the results proved for it are not changed. Instead, throughout this section we replace (1.1) by

$$x_{n+1} = \text{proj}_{\mathcal{X}}(x_n + \alpha_n\{h(Y_n, x_n, \theta_n) - x_n + M_{n+1}\}), \quad x_1 \in \mathcal{X}, \quad (\text{S.9.1})$$

while retaining the projected slow recursion (1.2). Here $\mathcal{X} \subset \mathbb{R}^{d_x}$ is fixed: it does not depend on n or on θ . Consequently, $x_n \in \mathcal{X}$ for every n , and the fast-stability Assumption 3.7 is no longer needed.

This extension is useful when boundedness of the unprojected fast iterates is not available a priori or would require a separate application-specific stability proof. The main paper assumes such boundedness, whereas the modified recursion enforces it by projecting the fast iterate at every step onto a prescribed compact feasible region. Subject to the Euclidean contraction condition on $\mathcal{X}$ and the bounded-preprojection-noise condition stated below, the extension then gives the corresponding high-probability path-tracking and almost-sure last-iterate guarantees.

S.9.1 Euclidean projection and contraction

Throughout this section, the fast-coordinate norm is the Euclidean norm. Let $\mathcal{X} \subset \mathbb{R}^{d_x}$ be a nonempty compact convex polyhedron, and write $P_{\mathcal{X}} := \text{proj}_{\mathcal{X}}$ for its Euclidean projection. We impose the averaged-contraction condition in the Euclidean norm and only on $\mathcal{X}$:

$$\left\| h^{(\text{av.})}(x; \theta) - h^{(\text{av.})}(x'; \theta) \right\| \leq \gamma \|x - x'\|, \quad x, x' \in \mathcal{X}, \quad \theta \in \Theta. \quad (\text{S.9.2})$$

Throughout this section, (S.9.2) replaces the global contraction requirement in Assumption 3.4. Assumption 3.5 is retained with both fast-state arguments restricted to $\mathcal{X}$: there exist finite constants $L_f^{\mathcal{X}}$, $L_{h,x}^{\mathcal{X}}$, and $L_{h,\theta}^{\mathcal{X}}$ such that, uniformly over $y \in \mathcal{Y}$, $x, \tilde{x} \in \mathcal{X}$, and $\theta, \tilde{\theta} \in \Theta$,

$$\left\| f(y, x, \theta) - f(y, \tilde{x}, \tilde{\theta}) \right\| \leq L_f^{\mathcal{X}} (\|x - \tilde{x}\| + \|\theta - \tilde{\theta}\|),$$

and

$$\left\| h(y, x, \theta) - h(y, \tilde{x}, \tilde{\theta}) \right\| \leq L_{h,x}^{\mathcal{X}} \|x - \tilde{x}\| + L_{h,\theta}^{\mathcal{X}} \|\theta - \tilde{\theta}\|.$$

For every $v \in D_{BV}([0, \infty); \mathbb{R}^{d_x})$ with $v(0) \in \mathcal{X}$, let $\Gamma_{\mathcal{X}}(v)$ denote the projected component of the Skorokhod problem on $\mathcal{X}$, as in Definition S.13.1, with (Θ, d) replaced by $(\mathcal{X}, d_x)$. Theorem S.13.1 ensures that $\Gamma_{\mathcal{X}}$ is well defined.

Proposition S.9.1 (Projection properties and contraction of $\mathcal{T}_{s,\theta}$). *Under (S.9.2), the following properties hold.*

(i) *The Euclidean projection is nonexpansive:*

$$\|P_{\mathcal{X}}(u) - P_{\mathcal{X}}(v)\|^2 \leq \langle P_{\mathcal{X}}(u) - P_{\mathcal{X}}(v), u - v \rangle, \quad u, v \in \mathbb{R}^{d_x}.$$

Consequently, $\|P_{\mathcal{X}}(u) - P_{\mathcal{X}}(v)\| \leq \|u - v\|$.

(ii) *The Skorokhod map $\Gamma_{\mathcal{X}}$ is Lipschitz: there exists $L_{\text{Sk}}^x < \infty$, independent of the time horizon, such that*

$$\sup_{0 \leq t \leq H} \|\Gamma_{\mathcal{X}}(v)(t) - \Gamma_{\mathcal{X}}(\tilde{v})(t)\| \leq L_{\text{Sk}}^x \sup_{0 \leq t \leq H} \|v(t) - \tilde{v}(t)\|$$

for every $H < \infty$ and every

$$v, \tilde{v} \in D_{BV}([0, \infty); \mathbb{R}^{d_x}) \quad \text{with} \quad v(0), \tilde{v}(0) \in \mathcal{X}.$$

(iii) *For $s \in [0, 1]$, define*

$$\mathcal{T}_{s,\theta}(x) := P_{\mathcal{X}}((1-s)x + sh^{(\text{av.})}(x; \theta)).$$

Then

$$\|\mathcal{T}_{s,\theta}(x) - \mathcal{T}_{s,\theta}(x')\| \leq \{1 - (1-\gamma)s\} \|x - x'\|, \quad x, x' \in \mathcal{X}. \quad (\text{S.9.3})$$

In particular, $P_{\mathcal{X}} \circ h^{(\text{av.})}(\cdot; \theta)$ is a γ -contraction on $\mathcal{X}$.

Proof. Let $p = P_{\mathcal{X}}(u)$ and $q = P_{\mathcal{X}}(v)$. The two projection variational inequalities give

$$\langle u - p, q - p \rangle \leq 0, \quad \langle v - q, p - q \rangle \leq 0.$$

Adding them yields $\|p - q\|^2 \leq \langle p - q, u - v \rangle$, which proves part (i). Part (ii) follows by applying Theorem S.13.1 to the polyhedron $\mathcal{X}$. Finally, part (i) and (S.9.2) give

$$\begin{aligned} \|\mathcal{T}_{s,\theta}(x) - \mathcal{T}_{s,\theta}(x')\| &\leq \|(1-s)(x - x') + s\{h^{(\text{av.})}(x; \theta) - h^{(\text{av.})}(x'; \theta)\}\| \\ &\leq \{1 - (1-\gamma)s\} \|x - x'\|, \end{aligned}$$

which proves part (iii). $\square$

All remaining standing assumptions are imposed on $\mathcal{Y} \times \mathcal{X} \times \Theta$. In particular, we retain Assumptions 3.1, 3.2, 3.3, 3.5, and 3.6, with the fast-coordinate bounds interpreted in the Euclidean norm; Assumption 3.4 is replaced by (S.9.2). Let $R_{\mathcal{X}} := \sup_{x \in \mathcal{X}} \|x\| < \infty$. Finiteness of $\mathcal{Y}$, compactness, and Assumption 3.5 give a finite $\mathbf{B}_{\mathcal{X}}$ bounding both drift maps on $\mathcal{Y} \times \mathcal{X} \times \Theta$, while Assumption 3.3 gives the preprojection-noise bound $N_{\text{noise}}^{\mathcal{X}} := \mathbf{c}_1 + \mathbf{c}_2 R_{\mathcal{X}}$. Set

$$A_x^{\mathcal{X}} := R_{\mathcal{X}} + \mathbf{B}_{\mathcal{X}} + N_{\text{noise}}^{\mathcal{X}}, \quad A_{\theta}^{\mathcal{X}} := \mathbf{B}_{\mathcal{X}} + N_{\text{noise}}^{\mathcal{X}}. \quad (\text{S.9.4})$$

Then, almost surely,

$$\begin{aligned} \sup_{x \in \mathcal{X}, \theta \in \Theta, y \in \mathcal{Y}} \{\|x\| + \|h(y, x, \theta)\|\} + \sup_n \|M_{n+1}\| &\leq A_x^{\mathcal{X}}, \\ \|x_{n+1} - x_n\| &\leq A_x^{\mathcal{X}} \alpha_n, \quad \|\theta_{n+1} - \theta_n\| \leq A_{\theta}^{\mathcal{X}} \beta_n. \end{aligned} \quad (\text{S.9.5})$$

The first increment bound follows from $x_n = P_{\mathcal{X}}(x_n)$ and Proposition S.9.1(i). Thus fast projection bounds the state but does not by itself bound the preprojection noise. The pathwise bounded-noise part of Assumption 3.3 is still required.

S.9.2 Constrained equilibrium and projected fast ODE

For $x \in \mathcal{X}$ and $v \in \mathbb{R}^{d_x}$, define the fast-set analogue of (4.1) by

$$\Pi_{\mathcal{X}}(x, v) := \lim_{\eta \downarrow 0} \frac{P_{\mathcal{X}}(x + \eta v) - x}{\eta}. \quad (\text{S.9.6})$$

Let $L_{\bar{h}, \theta}$ be a uniform Lipschitz constant of $h^{(\text{av.})}(x; \cdot)$ on Θ , uniform in $x \in \mathcal{X}$. Such a constant follows from Assumptions 3.2, 3.5, and 3.6, exactly as in Lemma 3.1, because $\mathcal{X}$ is compact.

Proposition S.9.2 (Constrained fast equilibrium and exponential stability). *Under the contraction condition (S.9.2), for every $\theta \in \Theta$, the map $P_{\mathcal{X}} \circ h^{(\text{av.})}(\cdot; \theta)$ has a unique fixed point in $\mathcal{X}$. Denote it by $x_{\mathcal{X}}^*(\theta)$; equivalently,*

$$x_{\mathcal{X}}^*(\theta) = P_{\mathcal{X}}\left(h^{(\text{av.})}(x_{\mathcal{X}}^*(\theta); \theta)\right). \quad (\text{S.9.7})$$

Moreover,

$$\|x_{\mathcal{X}}^*(\theta) - x_{\mathcal{X}}^*(\theta')\| \leq \frac{L_{\bar{h}, \theta}}{q} \|\theta - \theta'\|, \quad \theta, \theta' \in \Theta. \quad (\text{S.9.8})$$

For frozen θ , the projected fast ODE

$$\dot{z}(t) = \Pi_{\mathcal{X}}(z(t), h^{(\text{av.})}(z(t); \theta) - z(t)) \quad (\text{S.9.9})$$

has $x_{\mathcal{X}}^*(\theta)$ as its unique equilibrium, and its flow $S_t^{x, \theta}$ satisfies

$$\|S_t^{x, \theta}(z) - S_t^{x, \theta}(z')\| \leq e^{-qt} \|z - z'\|, \quad z, z' \in \mathcal{X}, \quad t \geq 0. \quad (\text{S.9.10})$$

In particular,

$$\|S_t^{x, \theta}(z) - x_{\mathcal{X}}^*(\theta)\| \leq e^{-qt} \|z - x_{\mathcal{X}}^*(\theta)\|.$$

Proof. Proposition S.9.1(iii) and Banach's fixed-point theorem prove existence and uniqueness. Comparing the two fixed-point identities at θ and θ' gives

$$\|x_{\mathcal{X}}^*(\theta) - x_{\mathcal{X}}^*(\theta')\| \leq \gamma \|x_{\mathcal{X}}^*(\theta) - x_{\mathcal{X}}^*(\theta')\| + L_{\bar{h}, \theta} \|\theta - \theta'\|,$$

which proves (S.9.8).

The projection characterization also shows that $x_{\mathcal{X}}^*(\theta)$ is a fixed point of $\mathcal{T}_{s, \theta}$ for every $s \in (0, 1]$: indeed, $x = P_{\mathcal{X}}(h^{(\text{av.})}(x; \theta))$ is equivalent to $h^{(\text{av.})}(x; \theta) - x$ belonging to the normal cone at x , and that cone is positively homogeneous.

Let

$$N_{\mathcal{X}}(z) := \left\{ \eta \in \mathbb{R}^{d_x} : \langle \eta, y - z \rangle \leq 0 \text{ for every } y \in \mathcal{X} \right\}$$

be the Euclidean outward normal cone. For fixed θ , the map $x \mapsto h^{(\text{av.})}(x; \theta) - x$ is Lipschitz on $\mathcal{X}$ with constant at most $1 + \gamma$. Hence the projected ODE is well posed by the same projected-dynamical-systems argument as in Lemma S.15.1. The projection characterization shows that the fixed point $x_{\mathcal{X}}^*(\theta)$ is an equilibrium of the ODE. For two solutions, the tangent-normal decomposition gives, for almost every t ,

$$\dot{z} = h^{(\text{av.})}(z; \theta) - z - \eta, \quad \dot{z}' = h^{(\text{av.})}(z'; \theta) - z' - \eta',$$

where $\eta \in N_{\mathcal{X}}(z)$ and $\eta' \in N_{\mathcal{X}}(z')$. Here the normals are outward, so $-\eta$ and $-\eta'$ are the corresponding inward correction terms. Since the normal-cone map is monotone,

$$\langle \eta - \eta', z - z' \rangle \geq 0.$$

Therefore,

$$\begin{aligned} \frac{d}{dt} \frac{1}{2} \|z - z'\|^2 &= \left\langle z - z', h^{(\text{av.})}(z; \theta) - h^{(\text{av.})}(z'; \theta) \right\rangle \\ &\quad - \|z - z'\|^2 - \langle z - z', \eta - \eta' \rangle \\ &\leq -(1 - \gamma) \|z - z'\|^2 = -q \|z - z'\|^2. \end{aligned}$$

Gronwall's inequality proves (S.9.10). It also proves uniqueness of the equilibrium and the final displayed bound in the proposition. $\square$

The constrained target need not equal the fixed point used in the unprojected framework. For comparison with that framework, suppose additionally that the original global Assumption 3.4 holds, and let $x^*(\theta)$ denote its unique fixed point. If

$$x^*(\theta) \in \mathcal{X}, \quad \theta \in \Theta, \quad (\text{S.9.11})$$

then $x_{\mathcal{X}}^*(\theta) = x^*(\theta)$, so the projection is only a safeguard and the reduced slow ODE is unchanged. Without this additional global assumption and compatibility condition, the extension instead analyzes the constrained reduced field

$$f_{\mathcal{X}}^{(\text{av.})}(\theta; \theta') := \sum_{y \in \mathcal{Y}} \mu^{(\theta')}(y) f(y, x_{\mathcal{X}}^*(\theta), \theta), \quad (\text{S.9.12})$$

and the slow projected ODE

$$\dot{z} = \Pi_{\Theta}(z, f_{\mathcal{X}}^{(\text{av.})}(z; z)). \quad (\text{S.9.13})$$

The field $\theta \mapsto f_{\mathcal{X}}^{(\text{av.})}(\theta; \theta)$ is Lipschitz by (S.9.8) and the standing assumptions; hence the same projected-dynamical-systems argument as in Lemma S.15.1 makes this ODE well posed. Let $\mathbf{S}_t^{\mathcal{X}}$ denote the flow of (S.9.13), and let

$$\mathcal{EQ}_{\Theta}^{\mathcal{X}} := \left\{ z \in \Theta : \Pi_{\Theta}(z, f_{\mathcal{X}}^{(\text{av.})}(z; z)) = \mathbf{0} \right\} \quad (\text{S.9.14})$$

be its equilibrium set. In this section, the constrained analogues of Assumptions 8.1, 8.2, and 8.3 mean those assumptions with $\mathbf{S}_s$, $\mathcal{EQ}_{\Theta}$, and $f^{(\text{av.})}$ replaced, respectively, by $\mathbf{S}_s^{\mathcal{X}}$, $\mathcal{EQ}_{\Theta}^{\mathcal{X}}$, and $f_{\mathcal{X}}^{(\text{av.})}$. When the constrained analogue of Assumption 8.3 is invoked, denote its one-block contraction factor by $\rho_T^{\mathcal{X}} \in (0, 1)$. If $\mathcal{X}$ fails to contain $x^*(\theta)$ for some $\theta \in \Theta$, the projected recursion targets $x_{\mathcal{X}}^*(\theta)$ instead, and the resulting reduced slow field, limiting ODE, and equilibrium set may differ from those of the unprojected recursion.

S.9.3 Fixed fast-time blocks and the moving-parameter comparison

The projected-fast comparison cannot be run over an entire slow-time block by a direct Gronwall argument: the fast-time duration of such a block is of order α_{n_m}/β_{n_m} , which diverges. We therefore partition each slow block into subblocks having a fixed duration in the fast clock.

Fix a fast-time horizon $T_x > 0$. On the slow block $\mathcal{K}_m$, put $\nu_{m,0} := n_m$, and, while $\nu_{m,j} < n_{m+1}$, define

$$\nu_{m,j+1} := \min \left\{ n_{m+1}, \inf \left\{ n > \nu_{m,j} : \sum_{r=\nu_{m,j}}^{n-1} \alpha_r \geq T_x \right\} \right\}. \quad (\text{S.9.15})$$

Here, m indexes the slow-time blocks and j indexes the fast-time subblocks within $\mathcal{K}_m$. The index $\nu_{m,j}$ is the left endpoint of the j -th subblock, while $\nu_{m,j+1}$ is its right endpoint, chosen when the

accumulated fast-clock time first reaches T_x , or as n_{m+1} if the slow block ends first. Let N_m^{sub} be the number of resulting subblocks. For $j = 0, \dots, N_m^{\text{sub}} - 1$, define the discrete index set

$$I_{m,j} := \{r \in \mathbb{N} : \nu_{m,j} \leq r < \nu_{m,j+1}\}, \quad H_{m,j}^x := \sum_{r \in I_{m,j}} \alpha_r.$$

Every subblock except possibly the last one satisfies

$$T_x \leq H_{m,j}^x < T_x + \alpha_{\nu_{m,j}}, \quad (\text{S.9.16})$$

and the last subblock satisfies the corresponding upper bound. The arguments below use only the bounds in (S.9.16) on the accumulated fast time of each subblock. For integers $r < s$, call the index set $\{r, r+1, \dots, s-1\}$ admissible if

$$\sum_{i=r}^{s-1} \alpha_i \leq T_x + \alpha_r,$$

and call it full if the same sum is also at least T_x . Thus every local subblock is admissible, and every one except possibly the last is full. The terms admissible and full will also be used for the discrete index sets $\{\nu_j, \nu_j + 1, \dots, \nu_{j+1} - 1\}$ in the trajectory-wide fast-clock partition defined later in (S.9.44). Unlike the local subblocks $I_{m,j}$, these blocks partition the entire iteration suffix and are not restarted or truncated at the slow-block endpoints n_{m+1} .

For each admissible block with integer endpoints $r < s$, define a fresh block-restarted, moving-parameter averaged recursion $\{\tilde{x}_k\}_{k=r}^s$ by

$$\begin{aligned} \tilde{x}_r &:= x_r, \\ \tilde{x}_{k+1} &:= P_{\mathcal{X}} \left(\tilde{x}_k + \alpha_k \{h^{(\text{av.})}(\tilde{x}_k; \theta_k) - \tilde{x}_k\} \right), \quad r \leq k < s. \end{aligned} \quad (\text{S.9.17})$$

The dependence of $\tilde{x}_k$ on the selected block endpoints r, s is suppressed. In particular, for the local subblock $I_{m,j}$, we take $r = \nu_{m,j}$ and $s = \nu_{m,j+1}$, and restart the auxiliary recursion from $x_{\nu_{m,j}}$. The auxiliary recursion uses the actual moving parameter θ_k .

We retain the fast Poisson solution from Definition 6.1, now restricted to $(x, \theta) \in \mathcal{X} \times \Theta$, rather than $(x, \theta) \in \mathbb{R}^{d_x} \times \Theta$. With M_{n+1}'' as in (6.4), set

$$\Xi_{n+1} := M_{n+1} + M_{n+1}'', \quad \Delta_{n+1}^h := u_h^{(x_n, \theta_n)}(Y_n) - u_h^{(x_n, \theta_n)}(Y_{n+1}).$$

Adding and subtracting $u_h^{(x_n, \theta_n)}(Y_{n+1})$ in the Poisson equation, as in Lemma 6.1, gives, pathwise,

$$h(Y_n, x_n, \theta_n) - h^{(\text{av.})}(x_n; \theta_n) + M_{n+1} = \Xi_{n+1} + \Delta_{n+1}^h. \quad (\text{S.9.18})$$

The sequence $\{\Xi_{n+1}\}$ is a martingale-difference sequence with respect to $\{\mathcal{F}_n\}$. Since $\mathcal{Y}$ is finite and $\mathcal{X} \times \Theta$ is compact, the retained bounded-noise and Lipschitz assumptions, together with Assumption 3.6, give deterministic constants $C_{\Xi}, U_h^{\mathcal{X}}, C_u < \infty$ such that the first bound below holds almost surely for every $n \geq 1$, whereas the remaining two bounds hold for every $(x, \theta), (x', \theta') \in \mathcal{X} \times \Theta$:

$$\begin{aligned} \|\Xi_{n+1}\| &\leq C_{\Xi}, \\ \sup_{y \in \mathcal{Y}} \|u_h^{(x, \theta)}(y)\| &\leq U_h^{\mathcal{X}}, \\ \sup_{y \in \mathcal{Y}} \|u_h^{(x, \theta)}(y) - u_h^{(x', \theta')}(y)\| &\leq C_u \{\|x - x'\| + \|\theta - \theta'\|\}. \end{aligned} \quad (\text{S.9.19})$$

For $r \leq k \leq s$, define

$$\mathcal{N}_{r,k}^x := \sum_{i=r}^{k-1} \alpha_i \Xi_{i+1}, \quad (\text{S.9.20})$$

$$\mathcal{R}_{r,k}^x := \sum_{i=r}^{k-1} \alpha_i \Delta_{i+1}^h, \quad (\text{S.9.21})$$

and, for $\delta > 0$, let

$$\mathcal{G}_{r,s}^{x,\text{proj}}(\delta) := \left\{ \max_{r \leq k \leq s} \|\mathcal{N}_{r,k}^x\| < \delta \right\}. \quad (\text{S.9.22})$$

Lemma S.9.1 (Fast martingale and Poisson residual on a fast-time block). *There are constants $c_x, C_x^{\text{res}} \in (0, \infty)$, independent of r, s , such that every admissible block with endpoints $r < s$ satisfies*

$$\mathbb{P} \left([\mathcal{G}_{r,s}^{x,\text{proj}}(\delta)]^c \right) \leq 2d_x(s-r) \exp \left\{ -\frac{c_x \delta^2}{\alpha_r} \right\}, \quad (\text{S.9.23})$$

$$\max_{r \leq k \leq s} \|\mathcal{R}_{r,k}^x\| \leq C_x^{\text{res}} \alpha_r. \quad (\text{S.9.24})$$

Proof. Coordinatewise Azuma–Hoeffding, followed by a union bound over the coordinates and the at most $s-r$ partial sums, gives

$$\mathbb{P} \left(\max_{r \leq k \leq s} \|\mathcal{N}_{r,k}^x\| \geq \delta \right) \leq 2d_x(s-r) \exp \left\{ -\frac{\delta^2}{2\kappa_{d_x}^2 C_{\Xi}^2 \sum_{i=r}^{s-1} \alpha_i^2} \right\}.$$

By admissibility and monotonicity of the step sizes,

$$\sum_{i=r}^{s-1} \alpha_i^2 \leq \alpha_r \sum_{i=r}^{s-1} \alpha_i \leq (T_x + \alpha_1) \alpha_r.$$

This proves (S.9.23).

For the residual, abbreviate $U_i(y) := u_h^{(x_i, \theta_i)}(y)$. Summation by parts gives, for $r < k \leq s$,

$$\begin{aligned} \mathcal{R}_{r,k}^x &= \alpha_r U_r(Y_r) - \alpha_{k-1} U_{k-1}(Y_k) \\ &\quad + \sum_{i=r+1}^{k-1} \{\alpha_i U_i(Y_i) - \alpha_{i-1} U_{i-1}(Y_i)\}. \end{aligned}$$

Using (S.9.5) and (S.9.19), its norm is at most a constant times

$$\alpha_r + \sum_{i=r+1}^{s-1} |\alpha_i - \alpha_{i-1}| + \sum_{i=r}^{s-1} \alpha_i^2 + \sum_{i=r}^{s-1} \alpha_i \beta_i.$$

The first two terms are $O(\alpha_r)$, because α_i is decreasing. The third term was bounded above. Finally, $\beta_i \leq \alpha_i$ under Condition 3.1, so the last term is also $O(\alpha_r)$. This proves (S.9.24). $\square$

Lemma S.9.2 (Projected comparison on a fast-time block). *There is $C_{T_x} < \infty$, independent of r, s , such that, for every $r < s$ satisfying the admissibility condition above, on $\mathcal{G}_{r,s}^{x,\text{proj}}(\delta)$,*

$$\max_{r \leq k \leq s} \|x_k - \tilde{x}_k\| \leq C_{T_x}(\delta + \alpha_r). \quad (\text{S.9.25})$$

Proof. Let $F_x(x, \theta) := h^{(\text{av.})}(x; \theta) - x$. On the selected block, embed the projected stochastic recursion (S.9.1) and the block-restarted averaged recursion (S.9.17) as right-continuous step paths on the local fast-time grid $t_{r,k} := \sum_{i=r}^{k-1} \alpha_i$. Their unconstrained drivers are

$$\begin{aligned} v_{r,k} &:= x_r + \sum_{i=r}^{k-1} \alpha_i \{F_x(x_i, \theta_i) + \Xi_{i+1} + \Delta_{i+1}^h\}, \\ \tilde{v}_{r,k} &:= x_r + \sum_{i=r}^{k-1} \alpha_i F_x(\tilde{x}_i, \theta_i). \end{aligned}$$

For $r \leq k < s$ and $t \in [t_{r,k}, t_{r,k+1})$, set

$$x^\circ(t) := x_k, \quad \tilde{x}^\circ(t) := \tilde{x}_k, \quad v^\circ(t) := v_{r,k}, \quad \tilde{v}^\circ(t) := \tilde{v}_{r,k},$$

and extend all four paths constantly with their corresponding index- s values for $t \geq t_{r,s}$. At each jump, the normal correction of the Skorokhod problem is precisely the correction made by Euclidean projection. Uniqueness of the Skorokhod problem therefore gives $x^\circ = \Gamma_{\mathcal{X}}(v^\circ)$ and $\tilde{x}^\circ = \Gamma_{\mathcal{X}}(\tilde{v}^\circ)$.

Let $L_{F,x}$ be the fast-state Lipschitz constant of F_x , and write $D_k := \max_{r \leq i \leq k} \|x_i - \tilde{x}_i\|$. Proposition S.9.1(ii), Lemma S.9.1, and the good event give

$$D_k \leq L_{\text{Sk}}^x \left\{ L_{F,x} \sum_{i=r}^{k-1} \alpha_i D_i + \delta + C_x^{\text{res}} \alpha_r \right\}.$$

Discrete Gronwall and $\sum_{i=r}^{s-1} \alpha_i \leq T_x + \alpha_1$ prove (S.9.25). $\square$

For $r \leq k \leq s$, set

$$\mathcal{W}_{r,k}^x := \prod_{i=r}^{k-1} (1 - q\alpha_i), \quad \mathcal{W}_{r,r}^x := 1.$$

Lemma S.9.3 (Moving constrained equilibrium). *Let $L_{fp}^{\mathcal{X}} := L_{h,\theta}/q$. On every admissible fast-time interval,*

$$\|\tilde{x}_k - x_{\mathcal{X}}^*(\theta_k)\| \leq \mathcal{W}_{r,k}^x \|x_r - x_{\mathcal{X}}^*(\theta_r)\| + \frac{L_{fp}^{\mathcal{X}} A_{\theta}^{\mathcal{X}}}{q} \frac{\beta_r}{\alpha_r}, \quad r \leq k \leq s. \quad (\text{S.9.26})$$

Proof. For $r \leq i < s$, the auxiliary recursion (S.9.17) can be written as

$$\tilde{x}_{i+1} = \mathcal{T}_{\alpha_i, \theta_i}(\tilde{x}_i).$$

As observed in the proof of Proposition S.9.2, $x_{\mathcal{X}}^*(\theta_i)$ is a fixed point of every relaxed map $\mathcal{T}_{a, \theta_i}$, $a \in (0, 1]$. Hence

$$\mathcal{T}_{\alpha_i, \theta_i}(x_{\mathcal{X}}^*(\theta_i)) = x_{\mathcal{X}}^*(\theta_i).$$

Adding and subtracting $x_{\mathcal{X}}^*(\theta_i)$, and then using the relaxed contraction (S.9.3), the equilibrium Lipschitz bound (S.9.8), and the slow-increment bound (S.9.5), gives

$$\begin{aligned} \|\tilde{x}_{i+1} - x_{\mathcal{X}}^*(\theta_{i+1})\| &\leq \|\mathcal{T}_{\alpha_i, \theta_i}(\tilde{x}_i) - \mathcal{T}_{\alpha_i, \theta_i}(x_{\mathcal{X}}^*(\theta_i))\| + \|x_{\mathcal{X}}^*(\theta_i) - x_{\mathcal{X}}^*(\theta_{i+1})\| \\ &\leq (1 - q\alpha_i) \|\tilde{x}_i - x_{\mathcal{X}}^*(\theta_i)\| + L_{fp}^{\mathcal{X}} \|\theta_{i+1} - \theta_i\| \\ &\leq (1 - q\alpha_i) \|\tilde{x}_i - x_{\mathcal{X}}^*(\theta_i)\| + L_{fp}^{\mathcal{X}} A_{\theta}^{\mathcal{X}} \beta_i. \end{aligned}$$

Iterate this inequality. Since $\beta_i/\alpha_i \leq \beta_r/\alpha_r$ for $i \geq r$, the forcing sum is at most

$$L_{fp}^{\mathcal{X}} A_{\theta}^{\mathcal{X}} \frac{\beta_r}{\alpha_r} \sum_{i=r}^{k-1} \alpha_i \prod_{\ell=i+1}^{k-1} (1 - q\alpha_{\ell}) \leq \frac{L_{fp}^{\mathcal{X}} A_{\theta}^{\mathcal{X}}}{q} \frac{\beta_r}{\alpha_r},$$

where the last sum telescopes after multiplication by q . $\square$

Combining the preceding two lemmas gives, on $\mathcal{G}_{r,s}^{x,\text{proj}}(\delta)$,

$$\begin{aligned} \|x_k - x_{\mathcal{X}}^*(\theta_k)\| &\leq \mathcal{W}_{r,k}^x \|x_r - x_{\mathcal{X}}^*(\theta_r)\| \\ &\quad + C_{T_x} \left(\delta + \alpha_r + \frac{\beta_r}{\alpha_r} \right), \quad r \leq k \leq s. \end{aligned} \quad (\text{S.9.27})$$

If the interval is full, then

$$\mathcal{W}_{r,s}^x \leq \exp \left\{ -q \sum_{i=r}^{s-1} \alpha_i \right\} \leq e^{-qT_x} =: \rho_x < 1. \quad (\text{S.9.28})$$

S.9.4 Slow-block interface and high-probability event

For the remainder of this section, define the projected-fast tracking error by

$$e_n^{(x)} := \|x_n - x_{\mathcal{X}}^*(\theta_n)\|, \quad n \geq 1.$$

Define the projected-fast event on the m th slow block by

$$\mathcal{H}_{x,m}^{\text{proj}}(\delta) := \bigcap_{j=0}^{N_m^{\text{sub}}-1} \mathcal{G}_{\nu_{m,j}, \nu_{m,j+1}}^{x,\text{proj}}(\delta). \quad (\text{S.9.29})$$

Thus, $\mathcal{H}_{x,m}^{\text{proj}}(\delta)$ is the event that, on every fast-time subblock of the m th slow block, the maximum accumulated martingale fluctuation is strictly less than δ , with the accumulation restarted at each subblock entrance. On this event, (S.9.27) can be applied successively across all fast-time subblocks of the slow block.

Proposition S.9.3 (Projected-fast input to the slow proof). *For every fixed slow-block horizon T , there are constants $c_{x,T}, C_{x,T}^{\text{proj}} \in (0, \infty)$ such that, for all $m \geq 0$ and $\delta > 0$,*

$$\mathbb{P} \left([\mathcal{H}_{x,m}^{\text{proj}}(\delta)]^c \right) \leq 2d_x K_m \exp \left\{ -\frac{c_{x,T} \delta^2}{\alpha_{n_m}} \right\}. \quad (\text{S.9.30})$$

On $\mathcal{H}_{x,m}^{\text{proj}}(\delta)$,

$$\sum_{r=n_m}^{n_{m+1}-1} \beta_r \|x_r - x_{\mathcal{X}}^*(\theta_r)\| \leq C_{x,T}^{\text{proj}} \left(\delta + \alpha_{n_m} + \frac{\beta_{n_m}}{\alpha_{n_m}} \right). \quad (\text{S.9.31})$$

Proof. Applying the union bound to the subblock failure events in (S.9.29) and using (S.9.23), together with

$$\sum_{j=0}^{N_m^{\text{sub}}-1} (\nu_{m,j+1} - \nu_{m,j}) = K_m, \quad \alpha_{\nu_{m,j}} \leq \alpha_{n_m},$$

gives (S.9.30), after decreasing the constant in the exponent if necessary. We next prove (S.9.31), which bounds the cumulative fast-tracking error weighted by the slow step sizes. Recall $g_m^{(x)}(\delta)$ from (6.14); its three terms dominate the corresponding terms on later subblocks. Iterating the endpoint form of (S.9.27) over the full fast subblocks and using (S.9.28) gives

$$e_{\nu_{m,j}}^{(x)} \leq \rho_x^j e_{n_m}^{(x)} + C_{T_x} g_m^{(x)}(\delta),$$

up to a change in C_{T_x} . Since $x_{n_m}, x_{\mathcal{X}}^*(\theta_{n_m}) \in \mathcal{X}$, compactness of $\mathcal{X}$ gives a deterministic bound on $e_{n_m}^{(x)}$.

On an admissible subblock with integer endpoints $r < s$, monotonicity of β_i/α_i and the telescoping identity used in the proof of Lemma S.9.3 yield

$$\sum_{i=r}^{s-1} \beta_i \mathcal{W}_{r,i}^x \leq \frac{\beta_r}{q\alpha_r}.$$

Also,

$$\sum_{i=r}^{s-1} \beta_i \leq \frac{\beta_r}{\alpha_r} H_{m,j}^x \leq C_{T_x} \frac{\beta_r}{\alpha_r}.$$

Apply these bounds to (S.9.27) and sum over j . By the preceding entrance-error bound, the part of $e_{\nu_{m,j}}^{(x)}$ inherited from the initial slow-block error is bounded by $\rho_x^j e_{n_m}^{(x)}$, and hence decays geometrically across the fast subblocks. The remaining terms are controlled using

$$\sum_{j=0}^{N_m^{\text{sub}}-1} \frac{\beta_{\nu_{m,j}}}{\alpha_{\nu_{m,j}}} \leq C_T$$

because each full subblock has fast duration at least T_x and the total slow duration is at most $T + \beta_{n_m}$. We verify this bound explicitly. Fixed fast duration and the power-step formulas make $(N_0 + i)/(N_0 + r)$ uniformly bounded on each admissible block with endpoints $r < s$, so β_i/α_i and β_r/α_r are uniformly comparable there. Consequently, on each full subblock,

$$\beta_r/\alpha_r \leq \frac{C}{T_x} \sum_{i=r}^{s-1} \beta_i.$$

Summing over the full subblocks gives a constant times the slow duration; for the possibly short final subblock, monotonicity gives $\beta_r/\alpha_r \leq \beta_{n_m}/\alpha_{n_m}$. Moreover, $\sum_{j=0}^{N_m^{\text{sub}}-1} \rho_x^j \frac{\beta_{\nu_{m,j}}}{\alpha_{\nu_{m,j}}} \leq \frac{\beta_{n_m}}{\alpha_{n_m}(1-\rho_x)}$. Thus the initial transient is multiplied by β_{n_m}/α_{n_m} , already a term in $g_m^{(x)}(\delta)$, and no factor N_m^{sub} is lost. This proves (S.9.31). $\square$

Equation (S.9.31) is the constrained analogue of (7.14), which controls the weighted fast-tracking term in the original-auxiliary comparison (7.12). After establishing the constrained slow Poisson and martingale bounds below, the proofs of Propositions 7.1 and 7.2 carry over under the following substitutions:

$$x^*(\theta) \mapsto x_{\mathcal{X}}^*(\theta), \quad f^{(\text{av.})} \mapsto f_{\mathcal{X}}^{(\text{av.})}, \quad \mathcal{G}_x^{(m)}(\delta) \mapsto \mathcal{H}_{x,m}^{\text{proj}}(\delta). \quad (\text{S.9.32})$$

To make the slow-coordinate transfer explicit, define the function describing the slow Markov noise

$$g_{f,\mathcal{X}}^{(\theta)}(y) := f(y, x_{\mathcal{X}}^*(\theta), \theta) - f_{\mathcal{X}}^{(\text{av.})}(\theta; \theta), \quad y \in \mathcal{Y}, \quad (\text{S.9.33})$$

and its return-time Poisson solution

$$u_{f,\mathcal{X}}^{(\theta)}(y) := \mathbb{E}^{(\theta)} \left[\sum_{r=0}^{\tau-1} g_{f,\mathcal{X}}^{(\theta)}(Y_r) \mid Y_0 = y \right], \quad (\text{S.9.34})$$

where τ is the return time used in Definition 7.1. For $n \geq 1$, set

$$M_{n+1}'''^{\mathcal{X}} := u_{f,\mathcal{X}}^{(\theta_n)}(Y_{n+1}) - \sum_{y \in \mathcal{Y}} p^{(\theta_n)}(Y_n, y) u_{f,\mathcal{X}}^{(\theta_n)}(y). \quad (\text{S.9.35})$$

Then $\{M_{n+1}'''^{\mathcal{X}}\}$ is a martingale-difference sequence and

$$g_{f,\mathcal{X}}^{(\theta_n)}(Y_n) = u_{f,\mathcal{X}}^{(\theta_n)}(Y_n) - u_{f,\mathcal{X}}^{(\theta_n)}(Y_{n+1}) + M_{n+1}'''^{\mathcal{X}}. \quad (\text{S.9.36})$$

By compactness of $\mathcal{X}$ and Θ , (S.9.8), and the same killed-resolvent argument used for Lemma S.11.1, there are finite deterministic constants $U_{f,\mathcal{X}}$, $C_{f,\mathcal{X}}$, and $C_{\text{mg}}^{(\theta,\mathcal{X})}$ such that, for every $\theta, \theta' \in \Theta$,

$$\sup_{\vartheta \in \Theta, y \in \mathcal{Y}} \|u_{f,\mathcal{X}}^{(\vartheta)}(y)\| \leq U_{f,\mathcal{X}}, \quad \sup_{y \in \mathcal{Y}} \|u_{f,\mathcal{X}}^{(\theta)}(y) - u_{f,\mathcal{X}}^{(\theta')}(y)\| \leq C_{f,\mathcal{X}} \|\theta - \theta'\|, \quad (\text{S.9.37})$$

and, for every $n \geq 1$, $\|M'_{n+1} + M_{n+1}'''^{\mathcal{X}}\| \leq C_{\text{mg}}^{(\theta,\mathcal{X})}$ almost surely. Define

$$\mathcal{G}_{\theta,\mathcal{X}}^{(m)}(\delta_s) := \left\{ \max_{1 \leq k \leq K_m} \left\| \sum_{\ell=0}^{k-1} \beta_{n_m+\ell} (M'_{n_m+\ell+1} + M_{n_m+\ell+1}'''^{\mathcal{X}}) \right\| < \delta_s \right\}. \quad (\text{S.9.38})$$

The coordinatewise Azuma–Hoeffding argument gives

$$\mathbb{P} \left([\mathcal{G}_{\theta,\mathcal{X}}^{(m)}(\delta_s)]^c \right) \leq 2d_\theta \sum_{k=1}^{K_m} \exp \left\{ - \frac{\delta_s^2}{2\kappa_{d_\theta}^2 (C_{\text{mg}}^{(\theta,\mathcal{X})})^2 \{b(n_m) - b(n_m + k)\}} \right\}. \quad (\text{S.9.39})$$

Proposition S.9.4 (Localized slow-ODE tracking with both projections). *Fix a slow-block horizon T . There is $C_{B,T}^{\mathcal{X}} < \infty$ such that, for every $m \geq 0$ and $\delta_f, \delta_s > 0$, on*

$$\mathcal{H}_{x,m}^{\text{proj}}(\delta_f) \cap \mathcal{G}_{\theta,\mathcal{X}}^{(m)}(\delta_s), \quad \sup_{0 \leq t \leq H_m} \left\| \theta^{(m),\circ}(t) - z_{\mathcal{X}}^{(T_m)}(t) \right\| \leq C_{B,T}^{\mathcal{X}} \left(\delta_f + \delta_s + \alpha_{n_m} + \frac{\beta_{n_m}}{\alpha_{n_m}} \right), \quad (\text{S.9.40})$$

where $z_{\mathcal{X}}^{(T_m)}$ solves (S.9.13) from θ_{n_m} . Moreover, the failure probability of this event is at most the sum of the right-hand sides of (S.9.30) and (S.9.39).

Proof. Use (S.9.36) in the original slow-driver decomposition. Bounds (S.9.8) and (S.9.37) give the same Poisson residual, auxiliary-recursion, and Skorokhod estimates as in Section 7, with x^* , $f^{(\text{av.})}$, and the stepsize-weighted fast input replaced according to (S.9.32). Proposition S.9.3 supplies that stepsize-weighted input. The same summation-by-parts calculation gives a deterministic constrained slow-Poisson residual bounded by $C\{\beta_{n_m} + b(n_m) - b(n_{m+1})\} = O(\beta_{n_m})$. The union bound gives the stated failure probability. $\square$

For a rate-specific choice, set

$$L_m^{x,\text{proj}} := \log\{4d_x K_m(m+2)^2\}, \quad \delta_{\text{dev}}^{(x,\text{proj})}(m) := \left(\frac{\alpha_{n_m} L_m^{x,\text{proj}}}{c_{x,T}} \right)^{1/2}. \quad (\text{S.9.41})$$

Then (S.9.30) gives

$$\mathbb{P} \left([\mathcal{H}_{x,m}^{\text{proj}}(\delta_{\text{dev}}^{(x,\text{proj})}(m))]^c \right) \leq \frac{1}{2(m+2)^2}.$$

Combine this event with $\mathcal{G}_{\theta, \mathcal{X}}^{(m)}(\delta_{\text{dev}}^{(\theta, \mathcal{X})}(m))$, where

$$\delta_{\text{dev}}^{(\theta, \mathcal{X})}(m) := \kappa_{d_\theta} C_{\text{mg}}^{(\theta, \mathcal{X})} \sqrt{2D_m L_m^\theta}.$$

Equations (S.9.39) and (S.9.30), followed by Borel–Cantelli, give the same eventual pathwise slow-block event as in Section 8, now with

$$d_{m, \mathcal{X}}^{\text{rate}} := \delta_{\text{dev}}^{(x, \text{proj})}(m) + \delta_{\text{dev}}^{(\theta, \mathcal{X})}(m) + \alpha_{n_m} + \frac{\beta_{n_m}}{\alpha_{n_m}}. \quad (\text{S.9.42})$$

On the resulting event, Proposition S.9.4 gives

$$\sup_{0 \leq t \leq H_m} \left\| \theta^{(m), \circ}(t) - z_{\mathcal{X}}^{(T_m)}(t) \right\| \leq C_{B, T}^{\mathcal{X}} d_{m, \mathcal{X}}^{\text{rate}}, \quad (\text{S.9.43})$$

S.9.5 Almost-sure rates, including the harmonic slow step

The preceding local construction is enough for the slow proof. For completeness, a global partition of the fast clock gives the iteration-wise fast rate. Under the shifted-step condition (3.3), put $\nu_0 := 1$; under the finite-prefix alternative, put $\nu_0 := n_0$, where n_0 is the suffix index defined in Appendix S.16. Define

$$\nu_{j+1} := \inf \left\{ n > \nu_j : \sum_{r=\nu_j}^{n-1} \alpha_r \geq T_x \right\}. \quad (\text{S.9.44})$$

Apply (S.9.27) on these successive blocks with

$$\delta_j = C_\delta \sqrt{\alpha_{\nu_j} \log(N_0 + \nu_j)},$$

where C_δ is chosen so that $c_x C_\delta^2 > 1$. Since $N_0 + \nu_j \asymp (j+1)^{1/(1-a)}$, integral comparison for the power steps gives $\nu_{j+1} - \nu_j = O((N_0 + \nu_j)^a)$. Hence the right-hand side of (S.9.23) is at most

$$C(N_0 + \nu_j)^{a - c_x C_\delta^2},$$

which is summable in j . Therefore, eventually almost surely,

$$e_{\nu_{j+1}}^{(x)} \leq \rho_x e_{\nu_j}^{(x)} + C_{T_x} \left\{ \sqrt{\alpha_{\nu_j} \log(N_0 + \nu_j)} + \alpha_{\nu_j} + \frac{\beta_{\nu_j}}{\alpha_{\nu_j}} \right\}. \quad (\text{S.9.45})$$

A geometric convolution of this recursion has the same order as its right-hand envelope. On a global fast block, $(N_0 + n)/(N_0 + \nu_j)$ is uniformly bounded and tends to one; applying the admissible-interval version of (S.9.27) to the final partial block therefore gives

$$\|x_n - x_{\mathcal{X}}^*(\theta_n)\| = O_{\text{a.s.}} \left((N_0 + n)^{-r_x(a, b)} \sqrt{\log(N_0 + n)} \right). \quad (\text{S.9.46})$$

The next theorem records the complete transfer to the slow results.

Theorem S.9.1 (Rates with projection on both time scales). *Let $\mathcal{X}$ be a nonempty compact convex polyhedron and equip the fast coordinate with the Euclidean norm. Suppose Condition 3.1 and the standing controlled-Markov, Euclidean contraction, Lipschitz, bounded-noise, and hitting-time assumptions hold on $\mathcal{Y} \times \mathcal{X} \times \Theta$. Use either the shifted-step condition (3.3) or the finite-prefix alternative of Appendix S.16. Let J be a Lipschitz Lyapunov function satisfying the constrained analogue of Assumption 8.1 for (S.9.13). Assumption 3.7 is not required. Then (S.9.46) holds. The following slow conclusions also hold.*

(i) Under the constrained analogue of the general one-block decrease condition in Assumption 8.2,

$$J(\theta_n) \longrightarrow 0, \quad \text{dist}(\theta_n, \mathcal{EQ}_\Theta^\chi) \longrightarrow 0 \quad \text{almost surely,}$$

where $\mathcal{EQ}_\Theta^\chi$ is defined in (S.9.14).

(ii) Under the hypotheses of part (i), suppose additionally that there exist $p_J > 1$, $\kappa_J > 0$, and $u_J > 0$ such that

$$\mathcal{D}_J(u) \geq \kappa_J u^{p_J}, \quad 0 \leq u \leq u_J.$$

If $\mathfrak{b} < 1$, then

$$J(\theta_n) = O_{\text{a.s.}} \left((N_0 + n)^{-(1-\mathfrak{b})/(p_J-1)} + (N_0 + n)^{-r_x(\mathfrak{a}, \mathfrak{b})/p_J} \{\log(N_0 + n)\}^{1/(2p_J)} \right). \quad (\text{S.9.47})$$

If $\mathfrak{b} = 1$, then

$$J(\theta_n) = O_{\text{a.s.}} \left(\{\log(N_0 + n)\}^{-1/(p_J-1)} \right). \quad (\text{S.9.48})$$

Thus the harmonic rate is $O_{\text{a.s.}}(1/\log n)$ when $p_J = 2$.

(iii) Under the constrained analogue of the linear one-block contraction condition in Assumption 8.3, if $\mathfrak{b} < 1$, then

$$J(\theta_n) = O_{\text{a.s.}} \left((N_0 + n)^{-r_x(\mathfrak{a}, \mathfrak{b})} \sqrt{\log(N_0 + n)} \right). \quad (\text{S.9.49})$$

If $\mathfrak{b} = 1$, put

$$\lambda_T^\chi := -\frac{\log \rho_T^\chi}{T} > 0.$$

Then

$$J(\theta_n) = \begin{cases} O_{\text{a.s.}} \left((N_0 + n)^{-r_x(\mathfrak{a}, 1)} \sqrt{\log(N_0 + n)} \right), & \lambda_T^\chi > r_x(\mathfrak{a}, 1), \\ O_{\text{a.s.}} \left((N_0 + n)^{-r_x(\mathfrak{a}, 1)} \{\log(N_0 + n)\}^{3/2} \right), & \lambda_T^\chi = r_x(\mathfrak{a}, 1), \\ O_{\text{a.s.}} \left((N_0 + n)^{-\lambda_T^\chi} \right), & 0 < \lambda_T^\chi < r_x(\mathfrak{a}, 1). \end{cases} \quad (\text{S.9.50})$$

At $\mathfrak{b} = 1$ and $\mathfrak{a} = 2/3$,

$$\|x_n - x_\chi^*(\theta_n)\| = O_{\text{a.s.}} \left((N_0 + n)^{-1/3} \sqrt{\log(N_0 + n)} \right), \quad (\text{S.9.51})$$

and the slow rate is given by (S.9.50) with $r_x(2/3, 1) = 1/3$.

If, in addition, the original global Assumption 3.4 holds and (S.9.11) holds, then $\mathcal{EQ}_\Theta^\chi = \mathcal{EQ}_\Theta$, and every target and reduced ODE in the theorem is the same as in the main paper. The shifted-step-size and finite-prefix alternatives of Appendix S.16 carry over without change: the global fast-time construction begins at 1 in the former case and at n_0 in the latter.

Proof. By the block-geometry lemmas, (S.9.41) and the constrained slow threshold satisfy

$$\delta_{\text{dev}}^{(x, \text{proj})}(m) = O \left((N_0 + n_m)^{-\mathfrak{a}/2} \sqrt{\log(N_0 + n_m)} \right),$$

and

$$\delta_{\text{dev}}^{(\theta, \chi)}(m) = O \left((N_0 + n_m)^{-\mathfrak{b}/2} \sqrt{\log(N_0 + n_m)} \right).$$

Therefore

$$d_{m,\mathcal{X}}^{\text{rate}} = O\left((N_0 + n_m)^{-r_x(\mathbf{a},\mathbf{b})} \sqrt{\log(N_0 + n_m)}\right).$$

Propositions S.9.3 and S.9.4 provide, respectively, the stepsize-weighted fast input and the localized slow-ODE tracking estimate used in the block-endpoint proofs. Those deterministic proofs use only these estimates, the Lipschitz property of J , and the stated Lyapunov decrease or contraction of the relevant flow. Repeating them with the constrained analogues defined above gives all the stated slow conclusions, including the harmonic branch. Finally, (S.9.46) follows from (S.9.45); setting $(\mathbf{a}, \mathbf{b}) = (2/3, 1)$ gives (S.9.51). $\square$

S.10 Auxiliary Proofs Used for Main-Paper Section 6

Lemma S.10.1. *The following bounds hold.*

(i) For every $k \geq 1$,

$$\|x_{k+1} - x_k\| \leq A_x \alpha_k,$$

and

$$\|\theta_{k+1} - \theta_k\| \leq A_\theta \beta_k.$$

(ii) The solution (6.3) of the fast Poisson equation satisfies

$$\|u_h^{(x,\theta)}(y)\| \leq 2\mathbf{c}_5 \mathbf{B}_6, \quad x \in \mathbb{R}^{d_x}, \theta \in \Theta, y \in \mathcal{Y},$$

whenever $\|x\| \leq \mathbf{c}_6$.

(iii) The martingale difference sequence defined in (6.4) satisfies

$$\|M''_{n+1}\| \leq 4\mathbf{c}_5 \mathbf{B}_6, \quad n \geq 1.$$

(iv) Define

$$C_{h,\text{Pois.}} := \max\{2\mathbf{c}_5 L_{h,x}, \mathbf{c}_5 [2L_{h,\theta} + (L_\mu + 2L_p \mathbf{c}_5) \mathbf{B}_6]\}.$$

For all $x, x' \in \mathbb{R}^{d_x}$ with $\|x\|, \|x'\| \leq \mathbf{c}_6$, all $\theta, \theta' \in \Theta$, and all $y \in \mathcal{Y}$,

$$\|u_h^{(x,\theta)}(y) - u_h^{(x',\theta')}(y)\| \leq C_{h,\text{Pois.}} (\|x - x'\| + \|\theta - \theta'\|).$$

Proof. For part (i), the update equations, Assumptions 3.3 and 3.7, and the definition of $\mathbf{B}_6$ give

$$\begin{aligned} \|x_{k+1} - x_k\| &\leq A_x \alpha_k, \\ \|\theta_{k+1} - \theta_k\| &\leq A_\theta \beta_k, \end{aligned}$$

where the second inequality also uses nonexpansiveness of projection.

For parts (ii)–(iv), write

$$g_{x,\theta}(y) := h(y, x, \theta) - h^{(\text{av.})}(x; \theta).$$

When $\|x\| \leq \mathbf{c}_6$, $\sup_y \|g_{x,\theta}(y)\| \leq 2\mathbf{B}_6$. The return-time representation (6.3) and Assumption 3.6 therefore imply

$$\|u_h^{(x,\theta)}(y)\| \leq 2\mathbf{c}_5 \mathbf{B}_6,$$

proving part (ii). Part (iii) follows from (6.4), since

$$\|M''_{n+1}\| \leq 2 \sup_y \|u_h^{(x,\theta)}(y)\| \leq 4\mathfrak{c}_5 \mathbf{B}_6.$$

It remains to prove part (iv). Let $\mathcal{Y}_0 := \mathcal{Y} \setminus \{y^\star\}$. The claim is immediate if $\mathcal{Y}_0$ is empty. For vector-valued functions on $\mathcal{Y}_0$, use $\|v\|_{\infty, \text{vec}} := \max_y \|v(y)\|$; then

$$\|Av\|_{\infty, \text{vec}} \leq \|A\|_{\infty} \|v\|_{\infty, \text{vec}}$$

for every scalar matrix A , with no dimension-dependent factor.

Let $Q^{(\theta)}$ be the restriction of $p^{(\theta)}$ to $\mathcal{Y}_0 \times \mathcal{Y}_0$. The hitting-time identity gives

$$\begin{aligned} Z^{(\theta)} &:= \sum_{n \geq 0} (Q^{(\theta)})^n = (I - Q^{(\theta)})^{-1}, \\ \sup_{\theta \in \Theta} \|Z^{(\theta)}\|_{\infty} &\leq \mathfrak{c}_5. \end{aligned} \tag{S.10.1}$$

Assumption 3.6 makes $y^\star$ positive recurrent and gives $\mu^{(\theta)}(y^\star) > 0$. Since $\sum_y \mu^{(\theta)}(y) g_{x,\theta}(y) = 0$, Lemma S.15.7, applied coordinatewise to the return-time representation (6.3), gives $u_h^{(x,\theta)}(y^\star) = 0$. First-step decomposition then yields

$$u_h^{(x,\theta)} = Z^{(\theta)} g_{x,\theta} \quad \text{on } \mathcal{Y}_0. \tag{S.10.2}$$

Set $\Delta_x := \|x - x'\|$ and $\Delta_\theta := \|\theta - \theta'\|$. Assumption 3.5, the bound $\mathbf{B}_6$, and Lemma 3.1 give

$$\|h^{(\text{av.})}(x; \theta) - h^{(\text{av.})}(x'; \theta')\| \leq L_{h,x} \Delta_x + (L_{h,\theta} + L_\mu \mathbf{B}_6) \Delta_\theta,$$

and hence

$$\|g_{x,\theta} - g_{x',\theta'}\|_{\infty, \text{vec}} \leq 2L_{h,x} \Delta_x + (2L_{h,\theta} + L_\mu \mathbf{B}_6) \Delta_\theta. \tag{S.10.3}$$

Also, $\|g_{x',\theta'}\|_{\infty, \text{vec}} \leq 2\mathbf{B}_6$. Since

$$\|Q^{(\theta)} - Q^{(\theta')}\|_{\infty} \leq L_p \Delta_\theta,$$

the resolvent identity and (S.10.1) yield

$$\|Z^{(\theta)} - Z^{(\theta')}\|_{\infty} \leq \mathfrak{c}_5^2 L_p \Delta_\theta.$$

Using (S.10.2)–(S.10.3),

$$\begin{aligned} \sup_{y \in \mathcal{Y}_0} \|u_h^{(x,\theta)}(y) - u_h^{(x',\theta')}(y)\| &\leq 2\mathfrak{c}_5 L_{h,x} \Delta_x \\ &\quad + \mathfrak{c}_5 [2L_{h,\theta} + (L_\mu + 2L_p \mathfrak{c}_5) \mathbf{B}_6] \Delta_\theta. \end{aligned}$$

Both solutions vanish at $y^\star$, so this bound holds on all of $\mathcal{Y}$. Taking the larger coefficient gives $C_{h, \text{Pois.}}$, proving part (iv). $\square$

S.11 Auxiliary Results Used for Main-Paper Section 7

Lemma S.11.1. *Consider the Poisson equation (7.6). The following bounds hold.*

(i) *For every $\theta \in \Theta$ and $y \in \mathcal{Y}$,*

$$\left\| u_f^{(\theta)}(y) \right\| \leq 2\mathfrak{c}_5 \mathbf{B}_8.$$

(ii) *For every $\theta, \theta' \in \Theta$ and $y \in \mathcal{Y}$,*

$$\left\| u_f^{(\theta)}(y) - u_f^{(\theta')}(y) \right\| \leq C_{f,\text{Pois.}} \|\theta - \theta'\|,$$

where

$$C_{f,\text{Pois.}} := \mathfrak{c}_5 [2L_f(L_{fp} + 1) + (L_\mu + 2L_p\mathfrak{c}_5)\mathbf{B}_8].$$

(iii) *The martingale difference sequence defined in (7.8) satisfies*

$$\left\| (M''')_{n+1}^{(m)} \right\| \leq 4\mathfrak{c}_5 \mathbf{B}_8.$$

Proof. Set

$$g_\theta(y) := f(y, x^\star(\theta), \theta) - f^{(\text{av.})}(\theta; \theta).$$

The definition of $\mathbf{B}_8$ gives $\sup_y \|g_\theta(y)\| \leq 2\mathbf{B}_8$. Hence the return-time representation and Assumption 3.6 imply

$$\left\| u_f^{(\theta)}(y) \right\| \leq 2\mathfrak{c}_5 \mathbf{B}_8,$$

which proves part (i).

For part (ii), let $\Delta_\theta := \|\theta - \theta'\|$. Assumption 3.5 and Lemma 3.1 give

$$\sup_y \|f(y, x^\star(\theta), \theta) - f(y, x^\star(\theta'), \theta')\| \leq L_f(L_{fp} + 1)\Delta_\theta.$$

Adding and subtracting the stationary average with measure $\mu^{(\theta)}$ gives

$$\left\| f^{(\text{av.})}(\theta; \theta) - f^{(\text{av.})}(\theta'; \theta') \right\| \leq [L_f(L_{fp} + 1) + L_\mu \mathbf{B}_8] \Delta_\theta.$$

Use the vector-valued sup norm introduced in the proof of Lemma S.10.1. The bounded-set estimate above gives

$$\|g_{\theta'}\|_{\infty, \text{vec}} \leq 2\mathbf{B}_8. \tag{S.11.1}$$

Combining the preceding two Lipschitz bounds gives

$$\|g_\theta - g_{\theta'}\|_{\infty, \text{vec}} \leq [2L_f(L_{fp} + 1) + L_\mu \mathbf{B}_8] \Delta_\theta. \tag{S.11.2}$$

These are the only bounds on this function needed.

If $\mathcal{Y}_0 := \mathcal{Y} \setminus \{y^\star\}$ is empty, the claim is immediate. Otherwise, use the killed kernels $Q^{(\theta)}$ and resolvents $Z^{(\theta)}$ constructed in the proof of Lemma S.10.1. Assumption 3.6 and Lemma S.15.7, applied coordinatewise to the return-time representation (7.7), give $u_f^{(\theta)}(y^\star) = 0$; first-step decomposition then gives

$$u_f^{(\theta)} = Z^{(\theta)} g_\theta \quad \text{on } \mathcal{Y}_0, \quad \left\| Z^{(\theta)} \right\|_\infty \leq \mathfrak{c}_5.$$

As in that proof,

$$\left\| Z^{(\theta)} - Z^{(\theta')} \right\|_\infty \leq \mathfrak{c}_5^2 L_p \Delta_\theta.$$

Therefore, adding and subtracting $Z^{(\theta)}g_{\theta'}$ and using (S.11.1)–(S.11.2),

$$\sup_{y \in \mathcal{Y}} \left\| u_f^{(\theta)}(y) - u_f^{(\theta')}(y) \right\| \leq \mathfrak{c}_5 \left[2L_f(L_{fp} + 1) + (L_\mu + 2L_p\mathfrak{c}_5)\mathbf{B}_8 \right] \Delta_\theta.$$

The coefficient is $C_{f,\text{Pois.}}$, proving part (ii). Finally, (7.8) and part (i) give

$$\left\| (M''')_{n+1}^{(m)} \right\| \leq 2 \sup_y \left\| u_f^{(\theta)}(y) \right\| \leq 4\mathfrak{c}_5\mathbf{B}_8,$$

which proves part (iii). □

Lemma S.11.2. *For all $\theta, \theta', \tilde{\theta} \in \Theta$, the following hold:*

(i)

$$\left\| f^{(\text{av.})}(\theta'; \theta) - f^{(\text{av.})}(\tilde{\theta}; \theta) \right\| \leq L_f(L_{fp} + 1) \left\| \theta' - \tilde{\theta} \right\|.$$

(ii)

$$\left\| f^{(\text{av.})}(\theta; \theta) - f^{(\text{av.})}(\theta'; \theta') \right\| \leq L_{f,\text{avg.}} \left\| \theta - \theta' \right\|,$$

where

$$L_{f,\text{avg.}} := L_f(L_{fp} + 1) + L_\mu\mathbf{B}_8.$$

(iii)

$$\left\| f^{(\text{av.})}(\theta; \theta') - f^{(\text{av.})}(\theta; \tilde{\theta}) \right\| \leq L_\mu\mathbf{B}_8 \left\| \theta' - \tilde{\theta} \right\|.$$

Proof. For part (i),

$$\begin{aligned} \left\| f^{(\text{av.})}(\theta'; \theta) - f^{(\text{av.})}(\tilde{\theta}; \theta) \right\| &\leq \sum_{y \in \mathcal{Y}} \mu^{(\theta)}(y) \left\| f(y, x^*(\theta'), \theta') - f(y, x^*(\tilde{\theta}), \tilde{\theta}) \right\| \\ &\leq L_f \left(\left\| x^*(\theta') - x^*(\tilde{\theta}) \right\| + \left\| \theta' - \tilde{\theta} \right\| \right) \\ &\leq L_f(L_{fp} + 1) \left\| \theta' - \tilde{\theta} \right\|. \end{aligned}$$

For part (ii), add and subtract $\sum_{y \in \mathcal{Y}} \mu^{(\theta)}(y) f(y, x^*(\theta'), \theta')$ and use part (i), Lemma 3.1, and the bounded-set estimate

$$\left\| f(y, x^*(\theta'), \theta') \right\| \leq \mathbf{B}_8.$$

Part (iii) follows directly from Lemma 3.1 and the same bounded-set estimate. □

Lemma S.11.3 (Residual term in the Poisson decomposition). *For every $1 \leq n \leq K_m$,*

$$\left\| \sum_{k=0}^{n-1} \beta_k^{(m)} \left(u_f^{(\theta_k^{(m)})}(Y_k^{(m)}) - u_f^{(\theta_{k+1}^{(m)})}(Y_{k+1}^{(m)}) \right) \right\| \leq R_m,$$

where

$$R_m = 3C_P\beta_{n_m} + C_L(b(n_m) - b(n_{m+1})), \tag{S.11.3}$$

with

$$C_P := 2\mathfrak{c}_5\mathbf{B}_8, \quad C_L := C_{f,\text{Pois.}}(\mathfrak{c}_1 + \mathfrak{c}_2\mathfrak{c}_6 + \mathbf{B}_6) = C_{f,\text{Pois.}}A_\theta.$$

Here $C_{f,\text{Pois.}}$ is the constant defined in part (ii) of Lemma S.11.1.

Proof of Lemma S.11.3. The sum can be written as follows:

$$\begin{aligned} & \sum_{k=0}^{n-1} \beta_k^{(m)} \left(u_f^{(\theta_k^{(m)})}(Y_k^{(m)}) - u_f^{(\theta_{k+1}^{(m)})}(Y_{k+1}^{(m)}) \right) \\ &= \beta_0^{(m)} u_f^{(\theta_0^{(m)})}(Y_0^{(m)}) - \beta_{n-1}^{(m)} u_f^{(\theta_{n-1}^{(m)})}(Y_n^{(m)}) \\ &+ \sum_{k=1}^{n-1} \left(\beta_k^{(m)} u_f^{(\theta_k^{(m)})}(Y_k^{(m)}) - \beta_{k-1}^{(m)} u_f^{(\theta_{k-1}^{(m)})}(Y_k^{(m)}) \right). \end{aligned}$$

The two boundary terms are bounded by

$$\beta_0^{(m)} C_P + \beta_{n-1}^{(m)} C_P \leq 2C_P \beta_{n_m}.$$

For the summation term, write

$$\begin{aligned} & \beta_k^{(m)} u_f^{(\theta_k^{(m)})}(Y_k^{(m)}) - \beta_{k-1}^{(m)} u_f^{(\theta_{k-1}^{(m)})}(Y_k^{(m)}) \\ &= \beta_{k-1}^{(m)} \{ u_f^{(\theta_k^{(m)})}(Y_k^{(m)}) - u_f^{(\theta_{k-1}^{(m)})}(Y_k^{(m)}) \} \\ &+ (\beta_k^{(m)} - \beta_{k-1}^{(m)}) u_f^{(\theta_k^{(m)})}(Y_k^{(m)}). \end{aligned}$$

By part (ii) of Lemma S.11.1 and the update bound for $\theta_k^{(m)}$,

$$\left\| u_f^{(\theta_k^{(m)})}(Y_k^{(m)}) - u_f^{(\theta_{k-1}^{(m)})}(Y_k^{(m)}) \right\| \leq \beta_{k-1}^{(m)} C_L.$$

Also, $\left\| u_f^{(\theta_k^{(m)})}(Y_k^{(m)}) \right\| \leq C_P$ and

$$\sum_{k=1}^{n-1} |\beta_k^{(m)} - \beta_{k-1}^{(m)}| \leq \beta_0^{(m)} = \beta_{n_m}.$$

Combining these estimates gives

$$\left\| \sum_{k=0}^{n-1} \beta_k^{(m)} \left(u_f^{(\theta_k^{(m)})}(Y_k^{(m)}) - u_f^{(\theta_{k+1}^{(m)})}(Y_{k+1}^{(m)}) \right) \right\| \leq 3C_P \beta_{n_m} + C_L \{b(n_m) - b(n_{m+1})\} = R_m.$$

□

S.12 Strong monotonicity and exponential stability of projected ODEs

Definition S.12.1. Let $K \subset \mathbb{R}^d$ be a nonempty closed convex set and let $F : K \rightarrow \mathbb{R}^d$. Define

$$\Pi_K(x, v) := \lim_{h \downarrow 0} \frac{\text{proj}_K(x + hv) - x}{h}, \quad x \in K, \quad v \in \mathbb{R}^d,$$

and consider the projected dynamical system

$$\dot{x} = \Pi_K(x, F(x)).$$

The vector field F is strongly monotone at $x^* \in K$ if there exists $\mu_{\text{sm}} > 0$ such that

$$\langle F(x) - F(x^*), x - x^* \rangle \leq -\mu_{\text{sm}} \|x - x^*\|^2, \quad x \in K.$$

Theorem S.12.1. *Let K be a nonempty compact convex polyhedron and let $F : K \rightarrow \mathbb{R}^d$ be Lipschitz. Suppose that $x^\star \in K$ is an equilibrium of*

$$\dot{x} = \Pi_K(x, F(x))$$

and that F is strongly monotone at $x^\star$, in the sense of Definition S.12.1, with constant $\mu_{\text{sm}} > 0$. Then the projected ODE is well posed and every solution satisfies

$$\|x(t) - x^\star\| \leq e^{-\mu_{\text{sm}} t} \|x(0) - x^\star\|, \quad t \geq 0.$$

In particular, $x^\star$ is the unique globally exponentially stable equilibrium.

Proof. Set

$$F_{\text{ext}}(z) := F(\text{proj}_K z), \quad z \in \mathbb{R}^d.$$

Since Euclidean projection is nonexpansive, F_{ext} is globally Lipschitz; since K is compact, it is also bounded. Thus Assumption 1 of Dupuis and Nagurney [16] holds. Taking $x_0^n = x_0$, $b_i^n = F_{\text{ext}}$, and $a_i^n = 1/n$ in their Theorem 3 gives existence, while their Theorem 2 gives uniqueness.

Since $x^\star$ is an equilibrium, it is a constant solution. Repeating the comparison calculation leading to equation (15) in the proof of Dupuis and Nagurney [16, Theorem 2], with their drift $b = F_{\text{ext}}$ and with the two solutions $x(\cdot)$ and $x^\star$, gives, for almost every $t \geq 0$,

$$\frac{1}{2} \frac{d}{dt} \|x(t) - x^\star\|^2 \leq \langle F(x(t)) - F(x^\star), x(t) - x^\star \rangle \leq -\mu_{\text{sm}} \|x(t) - x^\star\|^2.$$

Gronwall's inequality gives

$$\|x(t) - x^\star\| \leq e^{-\mu_{\text{sm}} t} \|x(0) - x^\star\|, \quad t \geq 0.$$

Applying this estimate to any constant equilibrium trajectory also proves that $x^\star$ is the unique equilibrium. $\square$

Corollary S.12.1 (Lyapunov and one-block consequences). *Under the hypotheses of Theorem S.12.1, let S_t^F denote the projected-ODE flow and define*

$$J(x) := \|x - x^\star\|.$$

Then J is a 1-Lipschitz Lyapunov function and, for every $T > 0$,

$$J(S_T^F(x)) \leq \rho_T J(x), \quad \rho_T := e^{-\mu_{\text{sm}} T} \in (0, 1), \quad x \in K.$$

Thus the uniform one-block Lyapunov contraction condition holds for every $T > 0$.

Lemma S.12.1. *Let K be a nonempty compact convex polyhedron. Consider the projected ODE*

$$\dot{x} = \Pi_K(x, \mathcal{T}_{\text{ctr}}(x) - x),$$

where $\mathcal{T}_{\text{ctr}} : K \rightarrow K$ is a contraction with contraction factor $\gamma \in [0, 1)$. If $x^\star$ is an equilibrium, then $\mathcal{T}_{\text{ctr}}(x^\star) = x^\star$, the vector field

$$F_{\text{ctr}}(x) := \mathcal{T}_{\text{ctr}}(x) - x$$

is strongly monotone at $x^\star$, and Corollary S.12.1 applies.

Proof. Since $\mathcal{T}_{\text{ctr}}(x^*) \in K$, the vector

$$\mathcal{T}_{\text{ctr}}(x^*) - x^*$$

belongs to the tangent cone of K at x^* . Because x^* is an equilibrium,

$$\Pi_K(x^*, \mathcal{T}_{\text{ctr}}(x^*) - x^*) = 0.$$

It follows that

$$\mathcal{T}_{\text{ctr}}(x^*) = x^*.$$

For every $x \in K$,

$$\begin{aligned} \langle F_{\text{ctr}}(x) - F_{\text{ctr}}(x^*), x - x^* \rangle &= \langle \mathcal{T}_{\text{ctr}}(x) - \mathcal{T}_{\text{ctr}}(x^*), x - x^* \rangle - \|x - x^*\|^2 \\ &\leq \|\mathcal{T}_{\text{ctr}}(x) - \mathcal{T}_{\text{ctr}}(x^*)\| \|x - x^*\| - \|x - x^*\|^2 \\ &\leq -(1 - \gamma) \|x - x^*\|^2. \end{aligned}$$

Thus F_{ctr} is strongly monotone at x^* , with constant $1 - \gamma$. The conclusion now follows from Corollary S.12.1. $\square$

S.13 Skorokhod problem

We recall the Skorokhod problem and the Lipschitz property of its solution map. This property is used to convert bounds between unconstrained driving paths into bounds between the corresponding projected paths.

Let $D([0, \infty); \mathbb{R}^d)$ denote the space of right-continuous functions with left limits from $[0, \infty)$ to $\mathbb{R}^d$. If $v : [0, \infty) \rightarrow \mathbb{R}^d$, let $|v|(T)$ denote its total variation on $[0, T]$. Let $D_{BV}([0, \infty); \mathbb{R}^d)$ be the set of all $v \in D([0, \infty); \mathbb{R}^d)$ such that v has bounded variation on every compact interval. Let $\partial\Theta$ denote the boundary of Θ . For $x \in \partial\Theta$, define the set of unit inward normals

$$n_{\Theta}(x) := \left\{ \nu \in \mathbb{R}^d : \|\nu\| = 1, \langle \nu, y - x \rangle \geq 0 \text{ for all } y \in \Theta \right\}.$$

For $x \in \text{int}(\Theta)$, set $n_{\Theta}(x) := \emptyset$.

Definition S.13.1 (Skorokhod problem). Let $v \in D([0, \infty); \mathbb{R}^d)$ with $v(0) \in \Theta$ be given. A pair (ϕ, η_{sk}) solves the Skorokhod problem with respect to Θ if, for all $t \geq 0$,

- (i) $\phi(t) = v(t) + \eta_{sk}(t)$ and $\phi(0) = v(0)$;
- (ii) $\phi(t) \in \Theta$;
- (iii) $|\eta_{sk}|(t) < \infty$;
- (iv) $|\eta_{sk}|(t) = \int_{[0, t]} \mathbf{1}_{\partial\Theta}(\phi(s)) d|\eta_{sk}|(s)$;
- (v) there exists a measurable function $\nu : [0, \infty) \rightarrow \mathbb{R}^d$ such that $\nu(s) \in n_{\Theta}(\phi(s))$ for $d|\eta_{sk}|$ -a.e. s , and

$$\eta_{sk}(t) = \int_{(0, t]} \nu(s) d|\eta_{sk}|(s).$$

The path v is called the unconstrained driving path, while ϕ is the projected path. When the solution is unique, write $\Gamma(v) := \phi$ for its projected component. The following is Dupuis and Nagurney [16, Theorem 1] in the present notation.

Theorem S.13.1 (Lipschitz property of the Skorokhod map). *Let $\Theta \subset \mathbb{R}^d$ be a convex polyhedron. For every $v \in D_{BV}([0, \infty); \mathbb{R}^d)$ with $v(0) \in \Theta$, there exists a unique solution to the Skorokhod problem. Moreover, the corresponding map Γ is Lipschitz, i.e., there exists a constant $L_{\text{Sk}} > 0$ such that, for every $T < \infty$ and every $v_1, v_2 \in D_{BV}([0, \infty); \mathbb{R}^d)$ with $v_1(0), v_2(0) \in \Theta$,*

$$\sup_{0 \leq t \leq T} \|\Gamma(v_1)(t) - \Gamma(v_2)(t)\| \leq L_{\text{Sk}} \sup_{0 \leq t \leq T} \|v_1(t) - v_2(t)\|.$$

Whenever a block step path $q^{(m), \circ}$, defined in Section 4, is used as an argument of Γ , extend it constantly beyond the block endpoint:

$$q^{(m), \circ}(r) := q_{K_m}^{(m)}, \quad r \geq H_m.$$

This convention does not affect any estimate on $[0, H_m]$.

Lemma S.13.1 (Step-path Skorokhod comparison on a block). *Fix $m \geq 0$. Let*

$$\theta^{(m), \circ}, \quad \tilde{\theta}^{(m), \circ}, \quad v^{(m), \circ}, \quad \tilde{v}^{(m), \circ}$$

be the right-continuous block step embeddings of the original slow sequence and of the sequences in (7.1)–(7.3), using the convention of Section 4. Then

$$\theta^{(m), \circ} = \Gamma(v^{(m), \circ}), \quad \tilde{\theta}^{(m), \circ} = \Gamma(\tilde{v}^{(m), \circ}).$$

The Lipschitz constant L_{Sk} depends only on the geometry of Θ , not on m or the driving paths. Consequently, for every $0 \leq n \leq K_m$,

$$\max_{0 \leq k \leq n} \|\theta_k^{(m)} - \tilde{\theta}_k^{(m)}\| \leq L_{\text{Sk}} \max_{0 \leq k \leq n} \|v_k^{(m)} - \tilde{v}_k^{(m)}\|.$$

Proof. For the first recursion, set $\eta_0^{(m)} := 0$ and $\eta_k^{(m)} := \theta_k^{(m)} - v_k^{(m)}$. For $0 \leq k < K_m$, write

$$\Delta v_{k+1}^{(m)} := v_{k+1}^{(m)} - v_k^{(m)}, \quad \Delta \eta_{k+1}^{(m)} := \eta_{k+1}^{(m)} - \eta_k^{(m)}.$$

Then, at the grid time $t^{(m)}(k+1)$,

$$\theta_{k+1}^{(m)} = \text{proj}_{\Theta}(\theta_k^{(m)} + \Delta v_{k+1}^{(m)}), \quad \Delta \eta_{k+1}^{(m)} = \theta_{k+1}^{(m)} - (\theta_k^{(m)} + \Delta v_{k+1}^{(m)}).$$

Because Θ is closed and convex and $\theta_{k+1}^{(m)} = \text{proj}_{\Theta}(\theta_k^{(m)} + \Delta v_{k+1}^{(m)})$, the first-order optimality condition for the Euclidean projection yields, for every $y \in \Theta$,

$$\langle \theta_{k+1}^{(m)} - (\theta_k^{(m)} + \Delta v_{k+1}^{(m)}), y - \theta_{k+1}^{(m)} \rangle \geq 0.$$

The first argument is $\Delta \eta_{k+1}^{(m)}$. Hence

$$\langle \Delta \eta_{k+1}^{(m)}, y - \theta_{k+1}^{(m)} \rangle \geq 0, \quad y \in \Theta.$$

Thus every nonzero regulator jump is an inward normal at $\theta_{k+1}^{(m)}$, which must then lie on $\partial\Theta$; its normalized direction belongs to $n_{\Theta}(\theta_{k+1}^{(m)})$. Because the block has only finitely many jumps and

both step embeddings are constant between grid points, the step regulator $\eta^{(m),\circ} := \theta^{(m),\circ} - v^{(m),\circ}$ has finite variation supported on times at which $\theta^{(m),\circ} \in \partial\Theta$. Hence

$$\nu_{k+1}^{(m)} := \frac{\Delta\eta_{k+1}^{(m)}}{\|\Delta\eta_{k+1}^{(m)}\|}$$

at each nonzero jump defines a measurable direction process with

$$\eta^{(m),\circ}(t) = \int_{(0,t]} \nu^{(m)}(s) d|\eta^{(m),\circ}|(s).$$

Consequently,

$$\left(\theta^{(m),\circ}, \theta^{(m),\circ} - v^{(m),\circ}\right)$$

solves the càdlàg Skorokhod problem driven by $v^{(m),\circ}$, and uniqueness gives $\theta^{(m),\circ} = \Gamma(v^{(m),\circ})$. The same argument yields $\tilde{\theta}^{(m),\circ} = \Gamma(\tilde{v}^{(m),\circ})$. Applying Theorem S.13.1 on $[0, t^{(m)}(n)]$ therefore gives

$$\begin{aligned} \max_{0 \leq k \leq n} \|\theta_k^{(m)} - \tilde{\theta}_k^{(m)}\| &= \sup_{0 \leq t \leq t^{(m)}(n)} \|\theta^{(m),\circ}(t) - \tilde{\theta}^{(m),\circ}(t)\| \\ &\leq L_{\text{Sk}} \sup_{0 \leq t \leq t^{(m)}(n)} \|v^{(m),\circ}(t) - \tilde{v}^{(m),\circ}(t)\| \\ &= L_{\text{Sk}} \max_{0 \leq k \leq n} \|v_k^{(m)} - \tilde{v}_k^{(m)}\|. \end{aligned}$$

□

Lemma S.13.2 (Unconstrained path for the projected ODE). *Fix $\tau \in \mathbb{R}_+$. There exists an absolutely continuous path $w^{(\tau)} : [0, \infty) \rightarrow \mathbb{R}^{d_\theta}$ satisfying*

$$w^{(\tau)}(0) = z^{(\tau)}(0) = \theta^\circ(\tau)$$

and

$$\dot{w}^{(\tau)}(t) = f^{(\text{av.})}(z^{(\tau)}(t); z^{(\tau)}(t))$$

for Lebesgue-a.e. $t \geq 0$, such that

$$z^{(\tau)} = \Gamma(w^{(\tau)}).$$

Proof. This follows from the Skorokhod representation of projected dynamical systems in Dupuis and Nagurney [16, Theorem 2]. □

S.14 Step-size-related results

Lemma S.14.1. *Let*

$$\alpha_n = (n + N_0)^{-\mathfrak{a}}, \quad 0 < \mathfrak{a} < 1,$$

and define

$$\chi(m, n) := \begin{cases} \prod_{\ell=m}^n (1 - \alpha_\ell), & m \leq n, \\ 1, & m > n. \end{cases}$$

Suppose that

$$(N_0 + 1)^{1-\mathfrak{a}} \geq 4.$$

Then, for every $n \geq 1$,

$$\sum_{k=1}^n \alpha_k^2 \chi(k+1, n) \leq 2\alpha_n,$$

and

$$\sum_{k=1}^n \alpha_k^2 \chi^2(k+1, n) \leq 2\alpha_n.$$

Proof. Let

$$S_n := \sum_{k=1}^n \alpha_k^2 \chi(k+1, n).$$

Then $S_1 = \alpha_1^2 \leq 2\alpha_1$ and, for $n \geq 2$,

$$S_n = \alpha_n^2 + (1 - \alpha_n)S_{n-1}.$$

Assume inductively that $S_{n-1} \leq 2\alpha_{n-1}$ and set $s_n := n + N_0$. Since $0 < \mathfrak{a} < 1$,

$$\frac{\alpha_{n-1}}{\alpha_n} = \left(\frac{s_n}{s_n - 1} \right)^{\mathfrak{a}} \leq \frac{s_n}{s_n - 1} \leq 1 + \frac{2}{s_n}.$$

Therefore,

$$\begin{aligned} S_n &\leq \alpha_n^2 + 2(1 - \alpha_n)\alpha_n \left(1 + \frac{2}{s_n} \right) \\ &\leq 2\alpha_n - \alpha_n^2 + \frac{4\alpha_n}{s_n} \leq 2\alpha_n, \end{aligned}$$

where the last inequality uses $4/s_n \leq s_n^{-\mathfrak{a}} = \alpha_n$, which follows from $(N_0 + 1)^{1-\mathfrak{a}} \geq 4$. This proves the first bound. Because $0 \leq \chi(k+1, n) \leq 1$,

$$\sum_{k=1}^n \alpha_k^2 \chi^2(k+1, n) \leq \sum_{k=1}^n \alpha_k^2 \chi(k+1, n) \leq 2\alpha_n,$$

proving the second. □

Lemma S.14.2. For $\alpha_n = (n + N_0)^{-\mathfrak{a}}$ and $0 < \mathfrak{a} < 1$, for every $n \geq 2$,

$$\sum_{k=2}^n |\alpha_k \chi(k+1, n) - \alpha_{k-1} \chi(k, n)| \leq \alpha_n.$$

Consequently,

$$\left\| \sum_{k=2}^n \{ \alpha_k \chi(k+1, n) - \alpha_{k-1} \chi(k, n) \} u_h^{(x_{k-1}, \theta_{k-1})}(Y_k) \right\| \leq 2\mathfrak{c}_5 \mathbf{B}_6 \alpha_n,$$

almost surely.

Proof. Let $A_k := \alpha_k \chi(k+1, n)$. Since $\chi(k, n) = (1 - \alpha_k) \chi(k+1, n)$,

$$A_k - A_{k-1} = \{ \alpha_k - \alpha_{k-1}(1 - \alpha_k) \} \chi(k+1, n).$$

With $s = k + N_0$, the inequality $\alpha_k \geq \alpha_{k-1}(1 - \alpha_k)$ is equivalent to

$$\left(\frac{s-1}{s} \right)^{\mathfrak{a}} \geq 1 - s^{-\mathfrak{a}},$$

which follows from subadditivity of $x \mapsto x^{\mathfrak{a}}$ on $\mathbb{R}_+$. Thus $A_k - A_{k-1} \geq 0$, and

$$\sum_{k=2}^n |A_k - A_{k-1}| = A_n - A_1 \leq \alpha_n.$$

The final bound follows from part (ii) of Lemma S.10.1. □

S.15 Miscellaneous results

Theorem S.15.1 (Azuma–Hoeffding). *Suppose $\{X_k\}_{k \geq 0}$ is a scalar martingale and $\{c_k\}_{k \geq 1}$ is a deterministic nonnegative sequence such that $|X_k - X_{k-1}| \leq c_k$ almost surely. Then, for every $N \in \mathbb{N}$ and every $\epsilon > 0$,*

$$\mathbb{P}(|X_N - X_0| \geq \epsilon) \leq 2 \exp \left(-\frac{\epsilon^2}{2 \sum_{k=1}^N c_k^2} \right).$$

Lemma S.15.1 (Well-posedness of the projected ODE). *For every $z_0 \in \Theta$, the projected ODE*

$$\dot{z}(t) = \Pi_{\Theta} \left(z(t), f^{(\text{av.})}(z(t); z(t)) \right), \quad z(0) = z_0,$$

has a unique solution, and this solution remains in Θ for all $t \geq 0$.

Proof. Lemma 3.1 gives $\sup_{\theta \in \Theta} \|x^*(\theta)\| \leq \mathfrak{c}_8$. Hence Assumption 3.5, applied on the ball of radius $\mathfrak{c}_8$, makes the diagonal field $F(\theta) := f^{(\text{av.})}(\theta; \theta)$ Lipschitz on Θ . Define $F_{\text{ext}}(z) := F(\text{proj}_{\Theta} z)$. Nonexpansiveness of projection makes F_{ext} globally Lipschitz; compactness of Θ makes it bounded, and it agrees with F on Θ . Existence follows from Dupuis and Nagurney [16, Theorem 3], while the Skorokhod representation and uniqueness follow from Dupuis and Nagurney [16, Theorem 2]. Hence the projected ODE is well posed and its solution remains in Θ . $\square$

Lemma S.15.2 (Perturbed power-drift recursion). *Let $p > 1$, $\kappa > 0$, $\zeta > 0$, $\ell \geq 0$, and $C_e \in [0, \infty)$. Let $\{u_m\}$ and $\{e_m\}$ be nonnegative sequences such that $u_m \rightarrow 0$ and, for all sufficiently large m ,*

$$u_{m+1} \leq u_m - \kappa u_m^p + e_m,$$

and

$$e_m \leq C_e (m+2)^{-\zeta} \{\log(m+2)\}^{\ell}.$$

Then there exists $C < \infty$ such that, for all sufficiently large m ,

$$u_m \leq C \left\{ (m+2)^{-1/(p-1)} + (m+2)^{-\zeta/p} \{\log(m+2)\}^{\ell/p} \right\}.$$

Proof. Set

$$w_m := (m+2)^{-1/(p-1)}, \quad \eta_m := (m+2)^{-\zeta/p} \{\log(m+2)\}^{\ell/p},$$

and $h_m := w_m + \eta_m$. These sequences are eventually decreasing. Because

$$\frac{1}{p-1} + 1 = \frac{p}{p-1},$$

the mean-value theorem gives, eventually,

$$0 \leq w_m - w_{m+1} \leq C_w w_m^p.$$

It also gives

$$0 \leq \eta_m - \eta_{m+1} \leq C_{\eta} \frac{\eta_m}{m+2}.$$

If $\zeta \leq p/(p-1)$, then

$$\frac{\eta_m}{m+2} = O(\eta_m^p),$$

where the logarithmic factor at equality is harmless because $\ell \geq 0$. If $\zeta > p/(p-1)$, then

$$\frac{\eta_m}{m+2} = O(w_m^p).$$

Consequently, for some $C_h < \infty$,

$$0 \leq h_m - h_{m+1} \leq C_h(w_m^p + \eta_m^p) \quad (\text{S.15.1})$$

eventually.

Choose $R > 0$ so that $\kappa p R^{p-1} \leq 1/2$. Since $u_m \rightarrow 0$ and $h_m \rightarrow 0$, choose $A_0 \geq 1$ and then a sufficiently large index M such that

$$\kappa A_0^{p-1} \geq 2C_h, \quad \kappa A_0^p \geq 2C_e, \quad u_M \leq R/2, \quad A_0 h_M \leq R/2.$$

Set

$$A := \max \left\{ A_0, \frac{u_M}{h_M} \right\}, \quad v_m := Ah_m.$$

Then $u_M \leq v_M \leq R/2$, and $v_m \leq R/2$ for every $m \geq M$. Since $(w_m + \eta_m)^p \geq w_m^p + \eta_m^p$, (S.15.1) and the choice of A give

$$\begin{aligned} v_m - \kappa v_m^p + e_m &\leq Ah_m - \kappa A^p(w_m^p + \eta_m^p) + C_e \eta_m^p \\ &\leq Ah_m - AC_h(w_m^p + \eta_m^p) \\ &\leq Ah_{m+1} = v_{m+1}. \end{aligned}$$

The map $F(u) := u - \kappa u^p$ is nondecreasing on $[0, R]$. Therefore, if $u_m \leq v_m$, then

$$u_{m+1} \leq F(u_m) + e_m \leq F(v_m) + e_m \leq v_{m+1}.$$

Induction proves $u_m \leq v_m$ for every $m \geq M$, which is the claimed bound. $\square$

Lemma S.15.3 (Geometric convolution of the blockwise rate envelope). *Suppose $1/2 < \mathfrak{b} < 1$. Fix $\rho \in (0, 1)$ and $r > 0$, and set*

$$\varphi_m := (N_0 + n_m)^{-r} \sqrt{\log(N_0 + n_m)}.$$

Then, for every $m_0 \geq 0$, there exist $m_1 \geq m_0$ and $C_{\text{geo}} < \infty$ such that, for every $m \geq m_1$,

$$\sum_{j=m_0}^{m-1} \rho^{m-1-j} \varphi_j \leq C_{\text{geo}} \varphi_m. \quad (\text{S.15.2})$$

Moreover,

$$\rho^{m-m_0} \leq C_{\text{geo}} \varphi_m, \quad m \geq m_1. \quad (\text{S.15.3})$$

Consequently, if

$$d_j \leq C_r \varphi_j, \quad j \geq m_0,$$

then

$$\sum_{j=m_0}^{m-1} \rho^{m-1-j} d_j \leq C_r C_{\text{geo}} \varphi_m, \quad m \geq m_1.$$

Proof. By the minimality of n_m ,

$$T \leq t(n_m) - t(n_{m-1}) < T + \beta_{n_m-1} \leq T + \beta_{n_0}.$$

This holds for $m \geq 1$. Summing over the first m blocks gives

$$mT \leq t(n_m) - t(n_0) \leq m(T + \beta_{n_0}), \quad m \geq 1. \quad (\text{S.15.4})$$

Integral comparison for $t(n) = \sum_{k=1}^{n-1} (N_0 + k)^{-\mathfrak{b}}$, together with (S.15.4), yields constants $c_{T,-}, c_{T,+} > 0$ and $M < \infty$ such that

$$c_{T,-}(m+1)^{1/(1-\mathfrak{b})} \leq N_0 + n_m \leq c_{T,+}(m+1)^{1/(1-\mathfrak{b})}, \quad m \geq M. \quad (\text{S.15.5})$$

Set

$$p := \frac{r}{1-\mathfrak{b}}, \quad \psi_m := (m+2)^{-p} \sqrt{\log(m+2)}.$$

Equation (S.15.5) gives constants $c_-, c_+ > 0$ such that

$$c_- \psi_m \leq \varphi_m \leq c_+ \psi_m, \quad m \geq M. \quad (\text{S.15.6})$$

For $m \geq j \geq M$,

$$\frac{\psi_j}{\psi_m} = \left(\frac{m+2}{j+2} \right)^p \sqrt{\frac{\log(j+2)}{\log(m+2)}} \leq \left(\frac{m+2}{j+2} \right)^p \leq (m-j+1)^p.$$

Hence, by (S.15.6),

$$\varphi_j \leq C(m-j+1)^p \varphi_m, \quad m \geq j \geq M. \quad (\text{S.15.7})$$

Putting $k = m-1-j$ in (S.15.7) gives

$$\sum_{j=M}^{m-1} \rho^{m-1-j} \varphi_j \leq C \varphi_m \sum_{k=0}^{\infty} \rho^k (k+2)^p \leq C_{\text{geo}} \varphi_m,$$

because $\rho \in (0, 1)$. The finitely many terms with $m_0 \leq j < M$ are $O(\rho^{m-M}) = O(\varphi_m)$, since an exponential sequence decays faster than a polynomial-logarithmic one. This proves (S.15.2). The same comparison gives $\rho^{m-m_0} = O(\varphi_m)$, proving (S.15.3). Finally, if $d_j \leq C_r \varphi_j$, multiply the first bound by C_r to obtain the stated consequence. $\square$

Lemma S.15.4 (Block estimates for nonharmonic slow steps). *Suppose $1/2 < \mathfrak{b} < 1$. There exist $C_T < \infty$ and $m_T < \infty$ such that, for every $m \geq m_T$,*

$$D_m \leq (T+1)\beta_{n_m}, \quad K_m \leq C_T(N_0 + n_m)^{\mathfrak{b}},$$

and

$$L_m^x + L_m^\theta \leq C_T \log(N_0 + n_m).$$

Proof. Set $X_m := N_0 + n_m$. Minimality of n_{m+1} gives

$$a(n_m, n_{m+1}) < T + \beta_{n_{m+1}-1} \leq T + \beta_{n_0} \leq T + 1,$$

and hence

$$D_m \leq \beta_{n_m} \sum_{k=n_m}^{n_{m+1}-1} \beta_k \leq (T+1)\beta_{n_m}.$$

Let

$$\overline{K}_m := \left\lceil 2^{\mathfrak{b}} T X_m^{\mathfrak{b}} \right\rceil.$$

Because $\mathfrak{b} < 1$, eventually $\overline{K}_m \leq X_m$, and then

$$t(n_m + \overline{K}_m) - t(n_m) \geq \overline{K}_m (X_m + \overline{K}_m)^{-\mathfrak{b}} \geq 2^{-\mathfrak{b}} \overline{K}_m X_m^{-\mathfrak{b}} \geq T.$$

Minimality therefore yields

$$K_m = n_{m+1} - n_m \leq \overline{K}_m \leq C_T X_m^{\mathfrak{b}}.$$

Finally, (S.15.5) implies $m + 2 \leq C_T X_m^{1-\mathfrak{b}}$ eventually, so

$$\begin{aligned} L_m^x + L_m^\theta &= \log\{4d_x K_m(m+2)^2\} + \log\{4d_\theta K_m(m+2)^2\} \\ &\leq C + 2 \log K_m + 4 \log(m+2) \leq C_T \log X_m. \end{aligned}$$

Increasing m_T and C_T , if necessary, proves the result. □

Lemma S.15.5 (Block estimates for harmonic slow steps). *Suppose $\mathfrak{b} = 1$, and set*

$$X_m := N_0 + n_m, \quad A_T := e^T.$$

Then, for every $m \geq 0$,

$$(A_T - 1)(X_m - 1) \leq K_m < (A_T - 1)X_m + 1.$$

Moreover, there exists $\xi_T > 0$ such that

$$X_m = \xi_T e^{mT} + O_T(1) \quad \text{as } m \rightarrow \infty.$$

Finally, there exist $C_T < \infty$ and a deterministic integer $m_T \geq 0$ such that, for every $m \geq m_T$,

$$D_m \leq (T + 1)X_m^{-1}, \quad L_m^x + L_m^\theta \leq C_T \log X_m,$$

and, whenever $n_m \leq n < n_{m+1}$,

$$X_m \leq N_0 + n < X_{m+1} < e^T X_m + 1 \leq (e^T + 1)X_m.$$

Proof. Minimality of n_{m+1} gives

$$T \leq \sum_{r=0}^{K_m-1} \frac{1}{X_m + r} < T + \frac{1}{X_{m+1} - 1}.$$

Integral comparison, applied also to the same sum with its last term removed, gives

$$T \leq \log \frac{X_m + K_m - 1}{X_m - 1}, \quad \log \frac{X_m + K_m - 1}{X_m} < T,$$

which proves the asserted bounds on K_m . Hence

$$X_{m+1} = e^T X_m + \eta_m, \quad -(e^T - 1) \leq \eta_m < 1.$$

Thus

$$e^{-mT} X_m = X_0 + \sum_{j=0}^{m-1} e^{-(j+1)T} \eta_j \longrightarrow \xi_T,$$

and the tail of this absolutely convergent series gives $X_m = \xi_T e^{mT} + O_T(1)$. Also, $X_{m+1} - 1 \geq e^T(X_m - 1)$, so $\xi_T \geq X_0 - 1 > 0$.

Since $\beta_k \leq X_m^{-1}$ on block m ,

$$D_m \leq X_m^{-1} \sum_{k=n_m}^{n_{m+1}-1} \beta_k \leq (T+1)X_m^{-1}.$$

The preceding estimates give $K_m = O_T(X_m)$ and $m = O_T(\log X_m)$. Substitution in the definitions of L_m^x and L_m^θ proves the logarithmic bound. The final within-block comparison follows from the upper bound on K_m . $\square$

Lemma S.15.6 (Geometric convolution for harmonic slow steps). *Suppose $\mathfrak{b} = 1$. Fix $\rho \in (0, 1)$, $r > 0$, and $\ell \geq 0$, and set*

$$\varphi_m := (N_0 + n_m)^{-r} \{\log(N_0 + n_m)\}^\ell, \quad \lambda_{\rho,T} := -\frac{\log \rho}{T}.$$

When $\rho = \rho_T$, the quantity $\lambda_{\rho,T}$ coincides with λ_T in (8.7). For every fixed m_0 , as $m \rightarrow \infty$,

$$\rho^{m-m_0} + \sum_{j=m_0}^{m-1} \rho^{m-1-j} \varphi_j = \begin{cases} O(\varphi_m), & \lambda_{\rho,T} > r, \\ O\left((N_0 + n_m)^{-r} \{\log(N_0 + n_m)\}^{\ell+1}\right), & \lambda_{\rho,T} = r, \\ O\left((N_0 + n_m)^{-\lambda_{\rho,T}}\right), & 0 < \lambda_{\rho,T} < r. \end{cases}$$

The same bounds hold with φ_j replaced by any nonnegative sequence satisfying $d_j \leq C\varphi_j$.

Proof. Lemma S.15.5 gives $N_0 + n_m \asymp e^{mT}$ and $\log(N_0 + n_m) \asymp m + 1$. Therefore,

$$\varphi_m \asymp e^{-rTm} (m+1)^\ell.$$

Up to constant factors, the convolution is

$$\sum_{j=m_0}^{m-1} \rho^{m-1-j} e^{-rTj} (j+1)^\ell.$$

If $\lambda_{\rho,T} > r$, equivalently $\rho < e^{-rT}$, factor out $e^{-rT(m-1)}$ and sum the convergent geometric series in

$$(\rho e^{rT})^{m-1-j}.$$

If $\lambda_{\rho,T} = r$, equivalently $\rho = e^{-rT}$, sum $(j+1)^\ell$. If $0 < \lambda_{\rho,T} < r$, equivalently $\rho > e^{-rT}$, factor out ρ^{m-1} and sum the convergent series in

$$\left(\frac{e^{-rT}}{\rho}\right)^j (j+1)^\ell.$$

These give, respectively,

$$O(e^{-rTm} m^\ell), \quad O(e^{-rTm} m^{\ell+1}), \quad O(\rho^m).$$

The initial transient has the same or smaller order, and converting back from m proves the result. $\square$

Lemma S.15.7 (Finite-state Kac cycle identity). *Let P be a transition matrix on a finite state space $\mathcal{Y}$, with unique stationary distribution μ . Fix $y^* \in \mathcal{Y}$, and let $(Y_n)_{n \geq 0}$ be the homogeneous Markov chain with transition matrix P , started from $Y_0 = y^*$. Let $\mathbb{E}_{y^*}$ denote expectation under this law, and set*

$$\tau_{y^*} := \inf\{n \geq 1 : Y_n = y^*\}.$$

If $\mathbb{E}_{y^}[\tau_{y^*}] < \infty$, then, for every integer $d_g \geq 1$ and every $g : \mathcal{Y} \rightarrow \mathbb{R}^{d_g}$,*

$$\mathbb{E}_{y^*} \left[\sum_{n=0}^{\tau_{y^*}-1} g(Y_n) \right] = \frac{1}{\mu(y^*)} \sum_{y \in \mathcal{Y}} \mu(y) g(y).$$

Proof. For $j \in \mathcal{Y}$, define the finite cycle occupation measure

$$\nu(j) := \mathbb{E}_{y^*} \left[\sum_{n=0}^{\tau_{y^*}-1} \mathbf{1}\{Y_n = j\} \right].$$

It is finite because

$$\sum_{j \in \mathcal{Y}} \nu(j) = \mathbb{E}_{y^*}[\tau_{y^*}] < \infty.$$

The Markov property and $Y_0 = Y_{\tau_{y^*}} = y^*$ give, for every j ,

$$\sum_{i \in \mathcal{Y}} \nu(i) P(i, j) = \mathbb{E}_{y^*} \left[\sum_{n=1}^{\tau_{y^*}} \mathbf{1}\{Y_n = j\} \right] = \nu(j).$$

Thus ν is a finite invariant measure. Uniqueness of the stationary distribution implies $\nu = c\mu$, while the first-return property gives $\nu(y^*) = 1$, and hence $c = 1/\mu(y^*)$. Therefore

$$\mathbb{E}_{y^*} \left[\sum_{n=0}^{\tau_{y^*}-1} g(Y_n) \right] = \sum_{j \in \mathcal{Y}} \nu(j) g(j) = \frac{1}{\mu(y^*)} \sum_{j \in \mathcal{Y}} \mu(j) g(j),$$

as claimed. □

S.16 Step-size shift and the starting index of the analysis

This appendix describes two ways to ensure that the step-size conditions required by the analysis hold. We can choose N_0 sufficiently large and begin the analysis at $n = 1$, or take $N_0 = 1$ and begin the analysis at a sufficiently large index n_0 . In both cases, the algorithm runs from $n = 1$.

S.16.1 Choosing N_0 to begin the analysis at $n = 1$

This is the convention already adopted throughout the main paper. We first give a sufficient lower bound on N_0 for the analysis to apply from $n = 1$. For the blockwise fast analysis in Section 6, choose N_0 so that

$$(N_0 + 1)^{1-\alpha} \geq 4.$$

Equivalently, it is sufficient to take

$$N_0 \geq \left\lceil 4^{1/(1-\alpha)} \right\rceil - 1.$$

The named domination inequality (S.5.1) then holds automatically because $A_x \geq A_\theta$, $C_\theta = C_{h,\text{Pois.}}A_\theta$, and $\beta_1 \leq \alpha_1$. The small-step condition is used in the local martingale and Poisson-residual estimates. No additional infinite-horizon condition involving a fixed fast-martingale threshold is needed: the block-dependent thresholds will be chosen later so that their failure probabilities are summable.

S.16.2 Beginning the analysis at $n = n_0$ when $N_0 = 1$

Alternatively, take $N_0 = 1$ and define

$$n_0 := \min\{n \geq 1 : (n+1)^{1-a} \geq 4\}.$$

The algorithm runs from $n = 1$, while the blockwise analysis begins at $n = n_0$, using the iterates generated by the algorithm up to that point. We retain the original iteration index and slow-time grid. For every $n \geq n_0$,

$$\alpha_n = (n+1)^{-a}, \quad \beta_n = (n+1)^{-b}, \quad (n+1)^{1-a} \geq 4.$$

We now establish the analogue of Lemma S.14.1 for sums beginning at n_0 . For every $n \geq n_0$,

$$\sum_{k=n_0}^n \alpha_k^2 \chi(k+1, n) \leq 2\alpha_n, \quad \sum_{k=n_0}^n \alpha_k^2 \chi^2(k+1, n) \leq 2\alpha_n. \quad (\text{S.16.1})$$

Indeed, at $n = n_0$, the first sum is $\alpha_{n_0}^2 \leq 2\alpha_{n_0}$. If the first inequality holds with terminal index $n-1$, the product structure of χ gives

$$\begin{aligned} \sum_{k=n_0}^n \alpha_k^2 \chi(k+1, n) &= \alpha_n^2 + (1 - \alpha_n) \sum_{k=n_0}^{n-1} \alpha_k^2 \chi(k+1, n-1) \\ &\leq \alpha_n^2 + 2(1 - \alpha_n)\alpha_{n-1}. \end{aligned}$$

Moreover,

$$\frac{\alpha_{n-1}}{\alpha_n} = \left(\frac{n+1}{n}\right)^a \leq \frac{n+1}{n} \leq 1 + \frac{2}{n+1},$$

and $4/(n+1) \leq \alpha_n$ because $(n+1)^{1-a} \geq 4$. Consequently,

$$\sum_{k=n_0}^n \alpha_k^2 \chi(k+1, n) \leq 2\alpha_n - \alpha_n^2 + \frac{4\alpha_n}{n+1} \leq 2\alpha_n.$$

This proves the first inequality in (S.16.1) by induction. The second follows from $0 \leq \chi^2(k+1, n) \leq \chi(k+1, n)$.

Every block entrance satisfies $n_m \geq n_0$. Therefore, whenever the fast-tail or Poisson-residual proof invokes Lemma S.14.1, its relevant block-local sum is bounded instead by the corresponding suffix sum in (S.16.1). For example, for $n \geq n_m$,

$$\sum_{k=n_m}^n \alpha_k^2 \chi^2(k+1, n) \leq \sum_{k=n_0}^n \alpha_k^2 \chi^2(k+1, n) \leq 2\alpha_n \leq 2\alpha_{n_m},$$

and the same argument applies without the square on χ . The weight-variation bound from Lemma S.14.2 is likewise valid on the suffix.

Under the large-shift convention used in the main paper, inequality (S.5.1), together with the monotonicity of β_n/α_n , is used in Appendix S.5, in the proof of Proposition S.5.1, immediately after the summation-by-parts decomposition of (S.5.2), to bound the Poisson-solution increments. When $N_0 = 1$, the same inequality holds for every $n \geq n_0$, since

$$C_\theta \beta_n = C_{h,\text{Pois.}} A_\theta \beta_n \leq C_{h,\text{Pois.}} A_x \alpha_n, \quad n \geq n_0,$$

using $A_\theta \leq A_x$ and $\beta_n \leq \alpha_n$.

Finally, the original slow-time grid is retained. The block construction gives

$$T \leq t(n_m) - t(n_{m-1}) < T + \beta_{n_m-1} \leq T + \beta_{n_0},$$

and hence

$$mT \leq t(n_m) - t(n_0) \leq m(T + \beta_{n_0}).$$

Beginning the block construction at $t(n_0)$ changes only the constants in the block-geometry and geometric-convolution bounds, for both $\mathfrak{b} < 1$ and $\mathfrak{b} = 1$. Thus, every blockwise result assuming (3.3) applies to the blocks beginning at n_0 , with (S.16.1) providing the required step-size estimate. The corresponding almost-sure rates hold for all sufficiently large original indices n . Since $N_0 = 1$, these rates are expressed in terms of $n + 1$, with the same polynomial and logarithmic exponents.

S.17 Rates under contraction of the reduced slow map (Proof of Proposition 8.2)

This appendix proves Proposition 8.2. We first establish Lemma S.17.1, which bounds the distance between a noisy projected recursion and a projected auxiliary recursion driven by the same input.

The next lemma assumes that the noise ξ_{n+1} admits the decomposition (S.17.1) into a martingale-difference term m_{n+1} , a difference $R_n - R_{n+1}$ of bounded correction terms, and a small remainder r_n . These auxiliary sequences are introduced only for the analysis and are not computed by the algorithm. When applying the lemma to the slow recursion in the proof of Proposition 8.2, we take

$$\xi_{n+1} = f(Y_n, x_n, \theta_n) - \sum_{y \in \mathcal{Y}} \mu^{(\theta_n)}(y) f(y, x_n, \theta_n) + M'_{n+1}.$$

Here ξ_{n+1} combines the Markov noise with the original martingale-difference noise M'_{n+1} . We construct m_{n+1} , R_n , and r_n using the Poisson equation, as shown below; the remainder r_n accounts for changes in the current pair (x_n, θ_n) .

Lemma S.17.1. *Let Θ be compact and convex, and consider adapted sequences*

$$\theta_{n+1} = \text{proj}_\Theta[(1 - \beta_n)\theta_n + \beta_n G_n + \beta_n \xi_{n+1}], \quad z_{n+1} = \text{proj}_\Theta[(1 - \beta_n)z_n + \beta_n G_n],$$

where G_n is $\mathcal{F}_n$ -measurable, $\theta_1, z_1 \in \Theta$, and $\beta_n = B/t_n$, with $t_n = N_0 + n$, $B > 1/2$, and $0 < \beta_n \leq 1$. Suppose that G_n and ξ_{n+1} are uniformly bounded by deterministic constants and

$$\xi_{n+1} = m_{n+1} + R_n - R_{n+1} + r_n, \quad \mathbb{E}[m_{n+1} | \mathcal{F}_n] = 0, \quad \|m_{n+1}\| \leq K_{\text{md}}, \quad (\text{S.17.1})$$

where R_n is adapted and, for finite deterministic constants $P, C_r \geq 0$, $\|R_n\| \leq P$ and $\|r_n\|^2 \leq C_r \beta_n$ almost surely for every n . Here $K_{\text{md}} := K_\xi + 2P + \sqrt{C_r}$, where $K_\xi < \infty$ is a deterministic uniform bound on $\|\xi_{n+1}\|$. The bound on m_{n+1} follows from

$$\begin{aligned} \|m_{n+1}\| &= \|\xi_{n+1} - R_n + R_{n+1} - r_n\| \\ &\leq K_\xi + 2P + \sqrt{C_r \beta_n} \leq K_{\text{md}}, \end{aligned}$$

using $\beta_n \leq 1$. In particular, K_{md} is independent of n and of the block index. Then

$$\|\theta_n - z_n\| = O_{\text{a.s.}}\left(\sqrt{\frac{\log t_n}{t_n}}\right). \quad (\text{S.17.2})$$

All norms in this lemma are Euclidean.

Proof. Nonexpansiveness of projection and boundedness of ξ_{n+1} give

$$\begin{aligned} \|\theta_{n+1} - z_{n+1}\|^2 &\leq (1 - \beta_n)^2 \|\theta_n - z_n\|^2 \\ &\quad + 2\beta_n(1 - \beta_n)\langle \theta_n - z_n, \xi_{n+1} \rangle + C\beta_n^2. \end{aligned} \quad (\text{S.17.3})$$

Here and below, C denotes a finite deterministic constant whose value may increase between inequalities. Since projection fixes every point of Θ , boundedness of the inputs also yields

$$\|\theta_{n+1} - \theta_n\| + \|z_{n+1} - z_n\| \leq C\beta_n.$$

To handle the difference $R_n - R_{n+1}$ in (S.17.1), we work directly with the squared shifted distance $\|\theta_n - z_n + \beta_n R_n\|^2$. Expand this square at indices n and $n + 1$, and substitute (S.17.1) into (S.17.3). The terms involving $R_n - R_{n+1}$ cancel up to an error bounded by $C\beta_n^2$, giving

$$\begin{aligned} & \|\theta_{n+1} - z_{n+1} + \beta_{n+1} R_{n+1}\|^2 \\ & \leq (1 - \beta_n)^2 \|\theta_n - z_n + \beta_n R_n\|^2 \\ & \quad + 2\beta_n(1 - \beta_n)\langle\theta_n - z_n, m_{n+1}\rangle \\ & \quad + 2\beta_n(1 - \beta_n)\langle\theta_n - z_n, r_n\rangle + C\beta_n^2. \end{aligned} \tag{S.17.4}$$

Indeed, the uncanceled terms are

$$\begin{aligned} & 2(1 - \beta_n)\beta_n^2\langle\theta_n - z_n, R_n\rangle \\ & \quad + 2\beta_{n+1}\langle(\theta_{n+1} - \theta_n) - (z_{n+1} - z_n), R_{n+1}\rangle \\ & \quad + 2\{\beta_{n+1} - (1 - \beta_n)\beta_n\}\langle\theta_n - z_n, R_{n+1}\rangle \\ & \quad + \beta_{n+1}^2\|R_{n+1}\|^2 - (1 - \beta_n)^2\beta_n^2\|R_n\|^2. \end{aligned}$$

Their total absolute value is $O(\beta_n^2)$, by the preceding increment bound, compactness of Θ , boundedness of R_n , and $\beta_{n+1} - (1 - \beta_n)\beta_n = O(\beta_n^2)$.

Fix c with $1 < c < 2B$. Using

$$\|\theta_n - z_n\| \leq \|\theta_n - z_n + \beta_n R_n\| + P\beta_n$$

and $\|r_n\|^2 \leq C_r\beta_n$, Young's inequality gives

$$\begin{aligned} & 2\beta_n(1 - \beta_n)|\langle\theta_n - z_n, r_n\rangle| \\ & \leq (2 - c/B)\beta_n\|\theta_n - z_n + \beta_n R_n\|^2 + C\beta_n^2. \end{aligned}$$

Since $\beta_n = B/t_n$, inequality (S.17.4) therefore becomes

$$\begin{aligned} & \|\theta_{n+1} - z_{n+1} + \beta_{n+1} R_{n+1}\|^2 \\ & \leq \left(1 - \frac{c}{t_n} + \frac{B^2}{t_n^2}\right) \|\theta_n - z_n + \beta_n R_n\|^2 \\ & \quad + 2\beta_n(1 - \beta_n)\langle\theta_n - z_n, m_{n+1}\rangle + \frac{C}{t_n^2}. \end{aligned} \tag{S.17.5}$$

Conditional Hoeffding's inequality, together with $\|\theta_n - z_n\|^2 \leq 2\|\theta_n - z_n + \beta_n R_n\|^2 + 2P^2\beta_n^2$, gives

$$\begin{aligned} & \mathbb{E}[\exp\{2s\beta_n(1 - \beta_n)\langle\theta_n - z_n, m_{n+1}\rangle\} \mid \mathcal{F}_n] \\ & \leq \exp\left\{4K_{\text{md}}^2 s^2 \beta_n^2 (\|\theta_n - z_n + \beta_n R_n\|^2 + P^2\beta_n^2)\right\}, \quad s \in \mathbb{R}. \end{aligned} \tag{S.17.6}$$

Choose $d > 0$ so that $4B^2K_{\text{md}}^2d < c - 1$. Exponentiate (S.17.5) after multiplying by dt_{n+1} , and apply (S.17.6) with $s = dt_{n+1} = d(t_n + 1)$. The coefficient of $dt_n\|\theta_n - z_n + \beta_n R_n\|^2$ in the resulting exponent is

$$\begin{aligned} & \frac{t_n + 1}{t_n} \left(1 - \frac{c}{t_n} + \frac{B^2}{t_n^2}\right) + 4B^2K_{\text{md}}^2d \frac{(t_n + 1)^2}{t_n^3} \\ & = 1 - \frac{c - 1 - 4B^2K_{\text{md}}^2d}{t_n} + O(t_n^{-2}). \end{aligned}$$

The remaining terms in that exponent are bounded by C/t_n . Consequently, there exists $\delta > 0$ such that, for all sufficiently large n ,

$$\begin{aligned} & \mathbb{E} \left[e^{dt_{n+1} \|\theta_{n+1} - z_{n+1} + \beta_{n+1} R_{n+1}\|^2} \middle| \mathcal{F}_n \right] \\ & \leq e^{C/t_n} \left(e^{dt_n \|\theta_n - z_n + \beta_n R_n\|^2} \right)^{1-\delta/t_n}, \quad 0 < 1 - \delta/t_n < 1. \end{aligned}$$

Taking expectations and applying Jensen's inequality to the concave power $u \mapsto u^{1-\delta/t_n}$ yields

$$\begin{aligned} & \log \mathbb{E} \left[e^{dt_{n+1} \|\theta_{n+1} - z_{n+1} + \beta_{n+1} R_{n+1}\|^2} \right] \\ & \leq \left(1 - \frac{\delta}{t_n} \right) \log \mathbb{E} \left[e^{dt_n \|\theta_n - z_n + \beta_n R_n\|^2} \right] + \frac{C}{t_n}. \end{aligned}$$

The squared shifted distance is bounded by a deterministic constant, so its exponential moment is finite at every fixed index. Starting at an index where the preceding recursion holds, induction bounds the logarithm of the exponential moment by the larger of its initial value and C/δ . Including the finite prefix therefore gives

$$\sup_{n \geq 1} \mathbb{E} \left[e^{dt_n \|\theta_n - z_n + \beta_n R_n\|^2} \right] < \infty.$$

Markov's inequality now implies

$$\mathbb{P} \left(\|\theta_n - z_n + \beta_n R_n\|^2 \geq K \frac{\log t_n}{t_n} \right) \leq C t_n^{-dK}.$$

Choose $K > 1/d$ and apply Borel–Cantelli. Finally,

$$\|\theta_n - z_n\| \leq \|\theta_n - z_n + \beta_n R_n\| + \frac{PB}{t_n} = O_{\text{a.s.}} \left(\sqrt{\frac{\log t_n}{t_n}} \right),$$

which proves (S.17.2). □

Proof of Proposition 8.2. The proof proceeds in four steps:

- (i) We identify the projected equilibrium and establish almost-sure convergence of the original iterates under the logarithmically separated step sizes.
- (ii) We construct Poisson decompositions for the fast and slow noise. For the fast recursion, we form the recursive weighted average U_n of the fast noise and analyze $p_n = x_n - U_n$. For the slow recursion, we construct the projected auxiliary sequence z_n and apply Lemma S.17.1 to bound $\|\theta_n - z_n\|$. These auxiliary sequences are defined in (S.17.10) and are used only in the proof.
- (iii) We show that the auxiliary slow iterate eventually remains in the face of Θ maximizing the inner product with the averaged drift at the equilibrium. On this face, that drift drops out of the projection objective, yielding a sharper bound on the auxiliary iterate's increments.
- (iv) We use fast contraction and reduced slow-map contraction to derive coupled error bounds. An induction gives the rate $O_{\text{a.s.}}(\log(N_0 + n)/\sqrt{N_0 + n})$, which we then transfer from the auxiliary sequences to the original iterates.

We use the original fast norm and Euclidean slow norm. Define the full averaged slow drift and its associated map by

$$F(x, \theta) = \sum_{y \in \mathcal{Y}} \mu^{(\theta)}(y) f(y, x, \theta), \quad G(x, \theta) = \theta + F(x, \theta).$$

Thus $F(x^*(\theta), \theta) = f^{(\text{av.})}(\theta; \theta)$ and $G(x^*(\theta), \theta) = \overline{G}(\theta)$. Constants below are deterministic unless stated otherwise; their values may increase from one estimate to the next.

Projected equilibrium and initial convergence. The map $\text{proj}_\Theta \circ \overline{G} : \Theta \rightarrow \Theta$ is a contraction, so it has a unique fixed point θ^* . Put $c = 1 - \rho_s > 0$ and $v = F(x^*(\theta^*), \theta^*)$. With the outward normal cone convention

$$N_\Theta(\theta) = \{w : \langle w, y - \theta \rangle \leq 0 \text{ for every } y \in \Theta\},$$

the projection characterization gives

$$v \in N_\Theta(\theta^*), \quad \theta^* = \text{proj}_\Theta[\theta^* + \beta v] \quad (\beta \geq 0). \quad (\text{S.17.7})$$

At a boundary equilibrium, the averaged slow drift v may be nonzero and point outward. Projection nevertheless leaves θ^* fixed, as expressed in (S.17.7). Contraction of $\overline{G}$ implies

$$\langle \theta - \theta^*, f^{(\text{av.})}(\theta; \theta) - v \rangle \leq -c \|\theta - \theta^*\|^2.$$

Both v and the normal correction in the projected ODE have nonpositive contributions to the derivative of squared distance to θ^* . Hence its flow satisfies

$$\|\mathbb{S}_t \theta - \theta^*\| \leq e^{-ct} \|\theta - \theta^*\|, \quad t \geq 0. \quad (\text{S.17.8})$$

In particular, the equilibrium set is $\mathcal{EQ}_\Theta = \{\theta^*\}$, and distance to θ^* is a global Lipschitz Lyapunov function with a uniform one-block contraction.

The initial-convergence argument in Section S.7.6, through (S.7.5) and the subsequent slow-block comparison, applies to the present schedule. That part of the argument uses the standing assumptions, the step-size relations, and Lyapunov decrease; it does not use the box-face or common-target conditions. In the present case $\beta_n/\alpha_n = B/(A \log t_n) \rightarrow 0$, $t_n \alpha_n \rightarrow \infty$, and both step-size sums diverge. The endpoint ratios on fixed fast-time blocks tend to one, while those on fixed slow-time blocks remain bounded. Thus the same Poisson and martingale comparisons, together with (S.17.8), give

$$\|x_n - x^*(\theta_n)\| = O_{\text{a.s.}} \left(\sqrt{\alpha_n \log t_n} + \frac{\beta_n}{\alpha_n} \right) \rightarrow 0, \quad \theta_n \rightarrow \theta^* \quad \text{almost surely.} \quad (\text{S.17.9})$$

The bounds in (S.17.9) follow from the block comparison estimates applied with the step sizes in (8.14).

Noise averaging within the feasible set. Center both noise streams at the actual pair (x_n, θ_n) :

$$\begin{aligned} \xi_{n+1}^x &= h(Y_n, x_n, \theta_n) - h^{(\text{av.})}(x_n; \theta_n) + M_{n+1}, \\ \xi_{n+1}^\theta &= f(Y_n, x_n, \theta_n) - F(x_n, \theta_n) + M'_{n+1}. \end{aligned}$$

Both admit (S.17.1), with bounded martingale differences and correctors and with $\|r_n\| \leq C\alpha_n$. To see this for either observation, let $u_{x,\theta}$ solve its Poisson equation centered at its full average, and put $u_n = u_{x_n, \theta_n}$. Then take

$$\begin{aligned} R_n &= u_n(Y_n), \\ m_{n+1} &= M_{n+1}^a + u_n(Y_{n+1}) - \sum_{y \in \mathcal{Y}} p^{(\theta_n)}(Y_n, y) u_n(y), \\ r_n &= u_{n+1}(Y_{n+1}) - u_n(Y_{n+1}), \end{aligned}$$

where M^a denotes M or M' , respectively. The transition rule for Y_{n+1} makes m_{n+1} a martingale difference with respect to the original filtration. The Poisson equation verifies the decomposition exactly, including its index Y_n . Lemma S.10.1 supplies the boundedness and sensitivity estimates for h . To obtain them for f , put $g_{x,\theta}(y) = f(y, x, \theta) - F(x, \theta)$. On a bounded fast-state set this function is bounded and satisfies

$$\sup_y \|g_{x,\theta}(y) - g_{x',\theta'}(y)\| \leq C(\|x - x'\| + \|\theta - \theta'\|),$$

by update Lipschitzness and Lemma 3.1. It is centered under $\mu^{(\theta)}$. The return-time solution therefore vanishes at the reference state and equals $Z^{(\theta)} g_{x,\theta}$ off that state, where $Z^{(\theta)}$ is the killed resolvent in (S.10.1). Its uniform bound and the resolvent identity give

$$\sup_y \|u_{x,\theta}(y)\| \leq C, \quad \sup_y \|u_{x,\theta}(y) - u_{x',\theta'}(y)\| \leq C(\|x - x'\| + \|\theta - \theta'\|).$$

The actual increments are $O(\alpha_n)$ and $O(\beta_n)$, so the sensitivity estimate gives the stated remainder bound. No independence between M_{n+1}^a and Y_{n+1} is needed.

Define the recursive weighted average U_n of the fast noise and the projected slow auxiliary sequence z_n by

$$\begin{aligned} U_1 &= 0, & U_{n+1} &= (1 - \alpha_n)U_n + \alpha_n \xi_{n+1}^x, & p_n &= x_n - U_n, \\ z_1 &= \theta_1, & z_{n+1} &= \text{proj}_\Theta[(1 - \beta_n)z_n + \beta_n G(x_n, \theta_n)], & \Delta_n &= \theta_n - z_n. \end{aligned} \tag{S.17.10}$$

For completeness, the deterministic averaging weights

$$w_{n,i} = \alpha_i \prod_{j=i+1}^{n-1} (1 - \alpha_j), \quad 1 \leq i < n,$$

satisfy $\sum_{i < n} w_{n,i}^2 = O(\alpha_n)$ and $\sum_{i < n} w_{n,i} \alpha_i = O(\alpha_n)$. These follow by applying a multiple of α_n as a supersolution to the respective recursions $v_{n+1} = (1 - \alpha_n)^2 v_n + \alpha_n^2$ and $b_{n+1} = (1 - \alpha_n)b_n + \alpha_n^2$, since $\alpha_n - \alpha_{n+1} = o(\alpha_n^2)$. The weights are eventually nondecreasing in i , because $1/\alpha_{i+1} - 1/\alpha_i \rightarrow 0$; each fixed initial weight decays faster than every inverse power of t_n . Summation by parts therefore bounds the coboundary contribution by $O(\alpha_n)$, as does the remainder contribution. Applying the Azuma–Hoeffding inequality coordinatewise to $\sum_{i < n} w_{n,i} m_{i+1}$, using $\sum_{i < n} w_{n,i}^2 = O(\alpha_n)$, gives

$$\mathbb{P}\left(\left\|\sum_{i < n} w_{n,i} m_{i+1}\right\|_2 > K\sqrt{\alpha_n \log t_n}\right) \leq C t_n^{-2}$$

for sufficiently large deterministic K . Since these probabilities are summable, the Borel–Cantelli lemma bounds the martingale contribution by $O_{\text{a.s.}}(\sqrt{\alpha_n \log t_n})$. Norm equivalence gives the same order in the original fast norm.

The original slow recursion for θ_n and the recursion for the projected auxiliary sequence z_n in (S.17.10) have the shared-input form required by Lemma S.17.1. Its remainder condition holds because $\alpha_n^2 = O(\beta_n)$. Consequently,

$$\|U_n\| = O_{\text{a.s.}}\left(\frac{\log t_n}{\sqrt{t_n}}\right), \quad \|\Delta_n\| = O_{\text{a.s.}}\left(\sqrt{\frac{\log t_n}{t_n}}\right). \tag{S.17.11}$$

The inputs and actual iterates, together with the auxiliary sequences U_n , p_n , and z_n , are bounded by deterministic constants. Thus all subsequent Lipschitz estimates may be taken on one enlarged deterministic fast-state ball.

The face reached by the auxiliary. Set

$$S = \operatorname{argmax}_{y \in \Theta} \langle v, y \rangle, \quad e_n = F(x_n, \theta_n) - v + \Delta_n.$$

Equations (S.17.9) and (S.17.11) give $z_n \rightarrow \theta^*$, $e_n \rightarrow 0$, and

$$z_{n+1} = \operatorname{proj}_{\Theta}[z_n + \beta_n(v + e_n)].$$

Suppose first that $S \neq \Theta$. A polyhedron has only finitely many distinct normal cones, all closed. Therefore

$$\delta = \min\{\operatorname{dist}(v, C) : C = N_{\Theta}(y) \text{ for some } y \in \Theta, v \notin C\} > 0.$$

Work after $\|e_n\| \leq \delta/4$. If $z_{n+1} \notin S$, let $r = (z_{n+1} - z_n)/\beta_n$. Projection optimality gives $v + e_n - r \in N_{\Theta}(z_{n+1})$, a cone excluding v , so $\|r\| \geq 3\delta/4$. Testing the same normal inequality against z_n yields

$$\langle v, r \rangle \geq \|r\|^2 - \|e_n\| \|r\| \geq 3\delta^2/8.$$

Every step ending outside S therefore increases $\langle v, z_n \rangle$ by at least $3\delta^2\beta_n/8$. Since $z_n \in \Theta$ and Θ is compact, $\langle v, z_n \rangle$ is bounded above by $\max_{y \in \Theta} \langle v, y \rangle < \infty$. If z_n remained outside S , the preceding increment bound and $\sum_n \beta_n = \infty$ would imply $\langle v, z_n \rangle \rightarrow \infty$, a contradiction. Thus z_n enters S after finitely many iterations. After entry, a step leaving S would require a strict increase from its maximum, which is impossible. Hence the auxiliary eventually remains in S .

For $z_n, z_{n+1} \in S$, restrict the projection objective to S . The term involving v is constant there, giving

$$z_{n+1} = \operatorname{proj}_S[z_n + \beta_n e_n], \quad \|z_{n+1} - z_n\| \leq \beta_n \|e_n\|. \quad (\text{S.17.12})$$

If $S = \Theta$, the above relations hold from the start. The argument showing that z_n eventually enters and remains in S also applies when Θ is lower-dimensional. If $v \neq 0$ is orthogonal to every difference $y - y'$, with $y, y' \in \Theta$, then $\langle v, y \rangle$ is constant over Θ , so $S = \Theta$. Only the auxiliary sequence z_n must eventually remain in S ; the noisy iterates θ_n need not do so. No strict complementarity is required: θ^* may lie on the relative boundary of S . Moreover, the fast equilibrium $x^*(\theta)$ need not be constant as θ varies over S .

Coupled contraction estimates. Recall $q = 1 - \gamma > 0$, and let K be a Lipschitz constant of x^* , and let L_g bound the Lipschitz dependence of G on its fast argument. Write

$$e_n^{(x)} := \|p_n - x^*(z_n)\|, \quad e_n^{(\theta)} := \|z_n - \theta^*\|, \quad \rho_n = \|U_n\| + \|\Delta_n\|, \quad s_n = \frac{\log t_n}{\sqrt{t_n}}.$$

By (S.17.11), $\rho_n = O_{\text{a.s.}}(s_n)$. Subtracting the update for U_{n+1} in (S.17.10) from the update for x_{n+1} cancels the noise term $\alpha_n \xi_{n+1}^x$. Using $p_n = x_n - U_n$, we obtain

$$p_{n+1} = (1 - \alpha_n)p_n + \alpha_n h^{(\text{av.})}(x_n; \theta_n).$$

Since $x_n - p_n = U_n$ and $\theta_n - z_n = \Delta_n$, Lipschitz continuity gives, for a deterministic constant L ,

$$\begin{aligned} \|h^{(\text{av.})}(x_n; \theta_n) - h^{(\text{av.})}(p_n; z_n)\| &\leq L(\|U_n\| + \|\Delta_n\|) = L\rho_n, \\ \|G(x_n, \theta_n) - G(p_n, z_n)\| &\leq L(\|U_n\| + \|\Delta_n\|) = L\rho_n. \end{aligned}$$

Also,

$$\|e_n\| \leq L_g e_n^{(x)} + (1 + \rho_s) e_n^{(\theta)} + L\rho_n,$$

because $f^{(\text{av.})}(z_n; z_n) - v = \overline{G}(z_n) - \overline{G}(\theta^*) - (z_n - \theta^*)$. Fast contraction and (S.17.12) consequently give, eventually almost surely,

$$e_{n+1}^{(x)} \leq (1 - q\alpha_n + KL_g\beta_n)e_n^{(x)} + K(1 + \rho_s)\beta_n e_n^{(\theta)} + (L\alpha_n + KL\beta_n)\rho_n. \quad (\text{S.17.13})$$

For the slow error, compare the auxiliary update with $\text{proj}_\Theta[(1 - \beta_n)\theta^* + \beta_n\overline{G}(\theta^*)] = \theta^*$, using (S.17.7). Nonexpansiveness and contraction of $\overline{G}$ give

$$e_{n+1}^{(\theta)} \leq (1 - c\beta_n)e_n^{(\theta)} + L_g\beta_n e_n^{(x)} + L\beta_n\rho_n. \quad (\text{S.17.14})$$

Thus the comparison retains the equilibrium normal v without assuming that the averaged drift vanishes at θ^* .

To close these inequalities, note that

$$\frac{s_{n+1}}{s_n} = 1 - \frac{1}{2t_n} + \frac{1}{t_n \log t_n} + O(t_n^{-2}).$$

Choose $K_x > 2L/q$, and then choose K_θ so large that

$$c - \frac{L_g K_x + L}{K_\theta} > \frac{1}{2B};$$

this is possible since $Bc \geq 2$. On each almost-sure path, choose a sufficiently large index N after face identification and all eventual estimates. A finite random C can be chosen so that $\rho_n \leq Cs_n$ for $n \geq N$, $e_N^{(x)} \leq CK_x s_N$, and $e_N^{(\theta)} \leq CK_\theta s_N$. Assuming the latter two bounds at index n , (S.17.14) proves the slow bound at $n + 1$. For (S.17.13), the decrement below $CK_x s_n$, divided by Cs_n , is at least

$$\alpha_n(qK_x - L) - \beta_n\{KL_g K_x + K(1 + \rho_s)K_\theta + KL\}.$$

Since $\beta_n/\alpha_n \rightarrow 0$ and $t_n\alpha_n \rightarrow \infty$, this eventually exceeds $K_x(1 - s_{n+1}/s_n)$, proving the fast bound at $n + 1$. Induction gives $e_n^{(x)} + e_n^{(\theta)} = O_{\text{a.s.}}(s_n)$. Finally,

$$\|\theta_n - \theta^*\| \leq e_n^{(\theta)} + \|\Delta_n\|, \quad \|x_n - x^*(\theta_n)\| \leq e_n^{(x)} + \|U_n\| + K\|\Delta_n\|.$$

Together with (S.17.11), these inequalities prove (8.15). Lipschitzness of x^* gives the same bound for $\|x_n - x^*(\theta^*)\|$; Lipschitzness of J and $J(\theta^*) = 0$ give it for $|J(\theta_n)|$. $\square$

S.18 Stochastic saddle-point learning without strong convexity–concavity

Consider the problem

$$\min_{u \in U} \max_{v \in V} \mathcal{L}(u, v), \quad \mathcal{L}(u, v) = \varphi(u) + u^\top A v - \psi(v), \quad (\text{S.18.1})$$

where $U \subset \mathbb{R}^{d_u}$ and $V \subset \mathbb{R}^{d_v}$ are nonempty compact convex polyhedra, $A \in \mathbb{R}^{d_u \times d_v}$, and φ and ψ are convex differentiable functions with Lipschitz gradients on neighborhoods of U and V . Write $\theta = (u, v)$ and $\Theta = U \times V$. Let $\theta^* = (u^*, v^*)$ be a saddle point, so that

$$\mathcal{L}(u^*, v) \leq \mathcal{L}(u^*, v^*) \leq \mathcal{L}(u, v^*), \quad u \in U, v \in V.$$

Assumption S.18.1 (Radial power curvature). There exist $q > 2$ and $c_{\text{sp}} > 0$ such that

$$\begin{aligned} \langle u - u^*, \nabla \varphi(u) - \nabla \varphi(u^*) \rangle + \langle v - v^*, \nabla \psi(v) - \nabla \psi(v^*) \rangle \\ \geq c_{\text{sp}} \|\theta - \theta^*\|^q, \quad \theta = (u, v) \in \Theta. \end{aligned} \quad (\text{S.18.2})$$

All norms in this section are Euclidean. Condition (S.18.2) permits curvature to vanish at the saddle and does not require strong convexity–concavity. Convexity–concavity alone, however, does not guarantee convergence of the descent–ascent ODE, which can cycle even in bilinear games.

Define the descent–ascent field

$$F_{\text{sp}}(u, v) := (-\nabla \varphi(u) - A v, A^\top u - \nabla \psi(v)). \quad (\text{S.18.3})$$

Suppose the finite controlled Markov chain satisfies Assumptions 3.1, 3.2, and 3.6. Let $\hat{F}_{\text{sp}}(y, \theta)$ be bounded and Lipschitz in θ , uniformly in y , with stationary mean

$$\sum_{y \in \mathcal{Y}} \mu^{(\theta)}(y) \hat{F}_{\text{sp}}(y, \theta) = F_{\text{sp}}(\theta), \quad \theta \in \Theta. \quad (\text{S.18.4})$$

Consider

$$\begin{aligned} x_{n+1} &= (1 - \alpha_n) x_n + \alpha_n \hat{F}_{\text{sp}}(Y_n, \theta_n), \\ \theta_{n+1} &= \text{proj}_\Theta(\theta_n + \beta_n x_n). \end{aligned} \quad (\text{S.18.5})$$

The fast variable estimates the entire saddle field; both primal and dual variables evolve on the slow time scale. Initialize $\theta_1 \in \Theta$ and $\|x_1\| \leq R$ almost surely, where R is a deterministic bound also satisfying $\sup_{y, \theta} \|\hat{F}_{\text{sp}}(y, \theta)\| \leq R$.

Proposition S.18.1 (Saddle-point learning under power curvature). *Under the preceding conditions, the reduced projected ODE has the unique equilibrium θ^* , and $J(\theta) := \|\theta - \theta^*\|$ satisfies Assumption 8.2 with power $p_J = q - 1$. For recursion (S.18.5), choose*

$$\alpha_n = (N_0 + n)^{-\mathfrak{a}}, \quad \beta_n = (N_0 + n)^{-\mathfrak{b}}, \quad \mathfrak{a} = \frac{2(q-1)}{4q-5}, \quad \mathfrak{b} = \frac{3(q-1)}{4q-5}, \quad (\text{S.18.6})$$

with N_0 satisfying (3.3). Then

$$\begin{aligned} \|\theta_n - \theta^*\| &= O_{\text{a.s.}} \left((N_0 + n)^{-\frac{1}{4q-5}} \{\log(N_0 + n)\}^{\frac{1}{2(q-1)}} \right), \\ \|x_n - F_{\text{sp}}(\theta_n)\| &= O_{\text{a.s.}} \left((N_0 + n)^{-\frac{q-1}{4q-5}} \sqrt{\log(N_0 + n)} \right). \end{aligned} \quad (\text{S.18.7})$$

The saddle may lie on the boundary of Θ ; no strict-complementarity condition is imposed.

Proof. In (1.1)–(1.2), take $h(y, x, \theta) = \widehat{F}_{\text{sp}}(y, \theta)$, $f(y, x, \theta) = x$, and $M_{n+1} = M'_{n+1} = 0$. By (S.18.4), the averaged fast map is constant in x :

$$h^{(\text{av.})}(x; \theta) = F_{\text{sp}}(\theta), \quad x^*(\theta) = F_{\text{sp}}(\theta).$$

Thus Assumption 3.4 holds with contraction factor zero. The oracle regularity and $f(y, x, \theta) = x$ verify Assumption 3.5. Since $0 < \alpha_n \leq 1$,

$$\|x_{n+1}\| \leq (1 - \alpha_n) \|x_n\| + \alpha_n R \leq R$$

whenever $\|x_n\| \leq R$. This proves Assumption 3.7; the martingale assumption holds trivially. The controlled-chain assumptions were imposed above.

The reduced slow field is F_{sp} . Let $\theta(t)$ solve its projected ODE, and use the outward normal cone

$$N_{\Theta}(\theta) := \{w : \langle w, z - \theta \rangle \leq 0 \text{ for every } z \in \Theta\}.$$

The saddle inequalities imply $F_{\text{sp}}(\theta^*) \in N_{\Theta}(\theta^*)$. Almost everywhere,

$$\dot{\theta}(t) = F_{\text{sp}}(\theta(t)) - \eta(t), \quad \eta(t) \in N_{\Theta}(\theta(t)).$$

For $d = \theta(t) - \theta^*$, monotonicity of the normal cone gives $\langle d, \eta(t) - F_{\text{sp}}(\theta^*) \rangle \geq 0$. Consequently, cancellation of the bilinear terms yields

$$\begin{aligned} \frac{d}{dt} \frac{\|d\|^2}{2} &\leq \langle d, F_{\text{sp}}(\theta(t)) - F_{\text{sp}}(\theta^*) \rangle \\ &= -\langle u - u^*, \nabla \varphi(u) - \nabla \varphi(u^*) \rangle - \langle v - v^*, \nabla \psi(v) - \nabla \psi(v^*) \rangle \\ &\leq -c_{\text{sp}} \|d\|^q. \end{aligned} \tag{S.18.8}$$

Applying the same inequality to a constant equilibrium trajectory shows that every equilibrium equals θ^* . The function J is therefore 1-Lipschitz and has exactly the required zero set. Wherever $J(\theta(t)) > 0$, (S.18.8) gives

$$\frac{d}{dt} J(\theta(t)) \leq -c_{\text{sp}} J(\theta(t))^{q-1}.$$

If a trajectory reaches θ^* , it remains there by uniqueness of the projected flow. Integrating the preceding inequality gives, for each fixed $T > 0$,

$$J(S_T(\theta)) \leq J(\theta) [1 + c_{\text{sp}}(q-2)TJ(\theta)^{q-2}]^{-1/(q-2)}. \tag{S.18.9}$$

In particular, J is a Lyapunov function, and

$$\mathcal{D}_J(u) := u - u[1 + c_{\text{sp}}(q-2)Tu^{q-2}]^{-1/(q-2)}$$

is continuous, vanishes at zero, and satisfies $0 < \mathcal{D}_J(u) \leq u$ for $u > 0$. Moreover,

$$\lim_{u \downarrow 0} \frac{\mathcal{D}_J(u)}{u^{q-1}} = c_{\text{sp}} T > 0.$$

This verifies the uniform one-block decrease and its power lower bound with $p_J = q - 1$. Since $q > 2$, the exponents in (S.18.6) satisfy $1/2 < \mathfrak{a} < \mathfrak{b} < 1$. Corollary 8.2 now gives (S.18.7). $\square$

A family with vanishing curvature. Let $0 \in U \times V$ and take

$$\varphi(u) = \frac{\|u\|^q}{q}, \quad \psi(v) = \frac{\|v\|^q}{q}, \quad q > 2, \quad (\text{S.18.10})$$

with arbitrary dimensions and arbitrary A . The origin is a saddle, and

$$\langle u, \nabla \varphi(u) \rangle + \langle v, \nabla \psi(v) \rangle = \|u\|^q + \|v\|^q \geq 2^{1-q/2} \|(u, v)\|^q.$$

Thus (S.18.2) holds. The Hessians of both penalties vanish at zero, so on sets containing a neighborhood of zero neither strong convexity nor strong concavity is available. The proposition also applies when zero lies on the boundary. For example, $q = 4$ gives $\mathfrak{a} = 6/11$, $\mathfrak{b} = 9/11$, and slow-iterate rate $O_{\text{a.s.}}((N_0 + n)^{-1/11} \{\log(N_0 + n)\}^{1/6})$. These are the rates supplied by the general power-decrease theorem; no optimality claim is made.

Remark S.18.1 (Comparison with existing frameworks). The obstruction in these applications is in the reduced slow dynamics; the fast averaged map is contractive. The comparisons below concern the hypotheses of the cited results, rather than possible extensions of their proof techniques.

For (S.18.10), suppose 0 is interior to Θ and $A \neq 0$. Then

$$DF_{\text{sp}}(0) = R_A := \begin{pmatrix} 0 & -A \\ A^\top & 0 \end{pmatrix}.$$

If $\sigma > 0$ is a nonzero singular value of A , the derivative of the forward map $\theta \mapsto \theta + hF_{\text{sp}}(\theta)$ has eigenvalues $1 \pm ih\sigma$, whose moduli exceed one for every $h > 0$. Projection is inactive in a neighborhood of zero for each fixed h . Hence even the projected forward map is not locally nonexpansive in any fixed norm. This rules out the reduced-map contraction hypotheses of Chandak [6], Chandak et al. [9] and the nonexpansive-map hypothesis of Chandak [5] for this family. The nonconvex-gradient extension in Chandak [5] does not apply either: R_A is nonsymmetric, so F_{sp} cannot be the negative Euclidean gradient of a scalar objective near zero. Its minimax specialization, which uses strong concavity in the fast player, also differs from (S.18.5), where the fast variable estimates the field and neither player requires strong curvature.

The contractive-map analysis of Chen et al. [10] likewise does not cover this reduced map. The more general drift condition in Chen et al. [11, Assumption 2.2] requires uniform quadratic dissipation toward an unprojected root. Here

$$\langle \theta, F_{\text{sp}}(\theta) \rangle = -\|u\|^q - \|v\|^q,$$

which cannot be bounded above by $-c\|\theta\|^2$ with $c > 0$ near zero. Nor can a positive-definite quadratic change of Lyapunov function restore local exponential dissipation: the linearization R_A has nonzero purely imaginary eigenvalues. The power-decrease condition (S.18.9) is sufficient for our result.

For the actor–critic recursion in Section 9, the lack of a nonexpansive slow map is already visible in a one-state, two-action MDP. Take costs 0 and 1, deterministic return to the same state, $B_{\text{ac}} = 1$, and write $p = (1 + e^{-(\vartheta_1 - \vartheta_2)})^{-1}$. At the exact critic, the averaged actor field is

$$F_{\text{ac}}(\vartheta) = (p(1 - p), -p(1 - p)).$$

Its Jacobian has eigenvalue $2p(1 - p)(1 - 2p) > 0$ in direction $(1, -1)$ when $p < 1/2$. At $\vartheta = (-1/2, 1/2)$, both this point and $\vartheta + F_{\text{ac}}(\vartheta)$ are interior to the logit box. Thus $\vartheta \mapsto \text{proj}_{\Theta_{\text{ac}}}(\vartheta + F_{\text{ac}}(\vartheta))$ is not locally nonexpansive in any fixed norm. The same obstruction persists for sufficiently

small positive forward steps. At the constrained optimum $(1, -1)$, the actor field is nonzero and points outward, although the projected velocity is zero. Consequently, neither a reduced-map contraction/nonexpansiveness condition nor an unprojected-root condition follows from the hypotheses of our actor-critic application. The unprojected slow-gradient result in Chandak [5] does not provide a rate to this constrained equilibrium. In the full application, the controlled Markov sampling and value-based Lyapunov function must also be retained.

The actor-critic result concerns precisely the global-clock recursion stated in Section 9, motivated by Konda and Borkar [18, Algorithm 3]; it does not extend the present rate claim to the original visit-count or delayed implementation. Neither projection nor Markov noise alone distinguishes our analysis from all earlier work. The point of these two applications is that projected slow-ODE Lyapunov decrease can hold when the slow-map assumptions in the comparison results fail.