

Sharp Stationary Gaussian Approximation for Constant-Stepsize SGD

Junghoon Seo

PIT IN Corp., South Korea

Abstract

We prove a sharp Gaussian approximation for the invariant law of constant-stepsizes SGD with bounded additive noise generated by an exogenous uniformly ergodic Markov chain. For a smooth, strongly convex objective with a Lipschitz Hessian and nondegenerate long-run noise covariance, the centered iterate normalized by the square root of the stepsize is $O(\sqrt{\alpha})$ -close in 1-Wasserstein distance to its limiting Gaussian. The proof combines blockwise Gaussian comparison with long-run contraction. A four-state example gives a matching lower bound although the one-time noise marginal is symmetric and every nonzero-lag autocovariance vanishes. In this example, an adjacent third-order mixed moment produces the leading correction.

1. Introduction

Constant-stepsizes stochastic gradient descent (SGD) is naturally studied through its invariant distribution when the deterministic drift is stable and the gradient noise persists (Pflug, 1986; Dieuleveut et al., 2020; Chen et al., 2022). We consider

$$X_{k+1} = X_k - \alpha \nabla f(X_k) + \alpha \xi(Z_k), \quad k \geq 0, \quad (1)$$

where $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is smooth and strongly convex, (Z_k) is an exogenous Markov chain with invariant distribution π , and $\xi : \mathcal{Z} \rightarrow \mathbb{R}^d$ is centered under π . Exogeneity means that the transition kernel of (Z_k) is independent of the iterate. At stationarity, X_k and Z_k are generally dependent because the iterate contains the accumulated effect of earlier Markov states.

Let $x_\star$ be the unique minimizer of f and set $H = \nabla^2 f(x_\star)$. The scale $\sqrt{\alpha}$ is suggested by a simple variance calculation. Over $O(\alpha^{-1})$ effective steps, the recursion accumulates noise increments of size α , so their total standard deviation is of order $\sqrt{\alpha^{-1}}\alpha = \sqrt{\alpha}$. For a stationary chain, temporal dependence changes the covariance through the long-run covariance

$$\Sigma_M = \mathbb{E}[\xi(Z_0)\xi(Z_0)^\top] + \sum_{\ell \geq 1} \mathbb{E}[\xi(Z_0)\xi(Z_\ell)^\top + \xi(Z_\ell)\xi(Z_0)^\top]. \quad (2)$$

Under the assumptions below, this series converges absolutely. The Gaussian benchmark is $\gamma = \mathcal{N}(0, \Sigma)$, where

$$H\Sigma + \Sigma H = \Sigma_M. \quad (3)$$

This is the covariance equation for the Ornstein–Uhlenbeck approximation obtained by linearizing the drift at $x_\star$ (Pflug, 1986; Chen et al., 2022).

Main contribution. Wang et al. (2026) give an $O(\sqrt{\alpha} \log(1/\alpha))$ upper bound for the W_1 distance between the $\alpha^{-1/2}$ -scaled centered stationary law and its Gaussian limit. We improve this to $O(\sqrt{\alpha})$ (Theorem 1) and give a four-state example attaining the matching lower bound (Theorem 2).

Proof architecture. The proof separates a finite-block Gaussian approximation from long-run contractivity. One block contains $n = \lceil T/\alpha \rceil$ updates, so its deterministic contraction factor is approximately e^{-mT} . Within a block, we linearize the drift and use a Poisson equation to decompose the Markovian noise into a martingale difference and a coboundary. The propagated Gaussian initial condition smooths Lipschitz test functions, which permits a second-order Lindeberg expansion. A second Poisson equation telescopes the state-dependent conditional covariance. A coalescent coupling and strong convexity then transfer the block estimate to the invariant law.

Related work. Stationary and diffusion descriptions of constant-stepsizes stochastic approximation appear in Pflug (1986); Mandt et al. (2017); Dieuleveut et al. (2020), and general stationary weak limits are developed by Chen et al. (2022). Poisson-equation methods for Markovian stochastic approximation are classical (Benveniste et al., 2012; Kushner and Yin, 2003; Fort, 2015). Quantitative analyses use concentration, Lyapunov methods, coupling, and linear stochastic-approximation techniques (Thoppe and Borkar, 2019; Chen et al., 2024; Mou et al., 2024; Durmus et al., 2025). Stationary bias and prelimit coupling under Markovian noise are studied in Huo et al. (2026, 2024); Allmeier and Gast (2024); Merad and Gaïffas (2025); Zhang et al. (2026). Dependent-data and finite-horizon Gaussian approximations provide complementary benchmarks (Zhang and Xie, 2026a; Wei et al., 2025; Haque et al., 2026; Kong and Srikanth, 2026). Appendix A discusses connections with recent Gaussian approximation results Zhang and Xie (2026a); Haque et al. (2026); Zhang and Xie (2026b,c).

2. Problem setup and main results

Fix $d \geq 1$. Vector norms and $\langle \cdot, \cdot \rangle$ are Euclidean, matrix inequalities use the Loewner order, and $\text{Lip}(h)$ denotes the Lipschitz constant of h . For matrices, $\|\cdot\|_{\text{op}}$, $\|\cdot\|_F$, and $\|\cdot\|_*$ denote the operator, Frobenius, and nuclear norms. We write I_d for the identity matrix and, for $B, C \in \mathbb{R}^{d \times d}$, $\langle B, C \rangle_F = \text{tr}(B^\top C)$. The notation $\mathbf{1}_A$ denotes the indicator of A . For a bounded function g taking values in a normed space, let $\|g\|_\infty = \sup_z \|g(z)\|$, where the pointwise norm is understood from context and is displayed explicitly for matrices when needed. Unquantified constants denoted by C or C_T are independent of α and may change from line to line. Explicitly quantified constants remain fixed. For probability measures on $\mathbb{R}^d$ with finite first moments, we use the Kantorovich–Rubinstein dual representation as the definition

$$W_1(\mu, \nu) = \sup_{\text{Lip}(h) \leq 1} \left| \int h \, d\mu - \int h \, d\nu \right|. \quad (4)$$

Every coupling (U, V) of (μ, ν) satisfies $W_1(\mu, \nu) \leq \mathbb{E}\|U - V\|$ by weak duality (Villani et al., 2009).

Objective assumptions. Assume that f is twice continuously differentiable and that there are constants $0 < m \leq L$ and $0 \leq M < \infty$ such that

$$mI_d \preceq \nabla^2 f(x) \preceq LI_d, \quad \|\nabla^2 f(x) - \nabla^2 f(y)\|_{\text{op}} \leq M\|x - y\| \quad (5)$$

for all $x, y \in \mathbb{R}^d$. Thus f is m -strongly convex and L -smooth, its Hessian is M -Lipschitz, and $H = \nabla^2 f(x_*)$ satisfies $mI_d \preceq H \preceq LI_d$.

Markovian noise assumptions. Let P be a Markov kernel on a Polish space $\mathbf{Z}$ with invariant probability measure π . We also use P for the associated Markov operator, $Pg(z) = \int g(u)P(z, du)$. Assume that, for constants $C_Z < \infty$ and $\rho \in (0, 1)$,

$$\sup_{z \in \mathbf{Z}} \|P^k(z, \cdot) - \pi\|_{\text{TV}} \leq C_Z \rho^k, \quad k \geq 0. \quad (6)$$

Here $\|\mu - \nu\|_{\text{TV}} = \sup_B |\mu(B) - \nu(B)|$, with the supremum over Borel sets. Assume also that the measurable observable $\xi : \mathbf{Z} \rightarrow \mathbb{R}^d$ satisfies

$$\int \xi(z) \pi(dz) = 0, \quad \sup_z \|\xi(z)\| \leq b < \infty. \quad (7)$$

We write $\mathbb{E}_\pi$ for expectation under a stationary P -chain. An unsubscripted $\mathbb{E}$ refers to the probability space currently under consideration. For a stationary P -chain, define

$$\Gamma_\ell = \mathbb{E}_\pi[\xi(Z_0)\xi(Z_\ell)^\top], \quad \ell \geq 0, \quad \Gamma_{-\ell} = \Gamma_\ell^\top, \quad \ell \geq 1. \quad (8)$$

Appendix C proves

$$C_\Gamma := \sum_{\ell \in \mathbb{Z}} \|\Gamma_\ell\|_* < \infty. \quad (9)$$

Hence $\Sigma_M = \sum_{\ell \in \mathbb{Z}} \Gamma_\ell$ agrees with (2). We assume $\Sigma_M \succ 0$. The Lyapunov equation (3) then has the unique positive-definite solution

$$\Sigma = \int_0^\infty e^{-tH} \Sigma_M e^{-tH} dt, \quad \gamma = \mathcal{N}(0, \Sigma). \quad (10)$$

Moreover, $\lambda_{\min}(\Sigma) \geq \lambda_{\min}(\Sigma_M)/(2L)$ and $\|\Sigma\|_{\text{op}} \leq \|\Sigma_M\|_{\text{op}}/(2m)$. Thus the constants below can be chosen with the dependence stated in Theorem 1.

Theorem 1 (Gaussian approximation for the invariant law) *Under (5)–(7) and $\Sigma_M \succ 0$, there are constants $\alpha_* > 0$ and $C < \infty$, depending only on d, m, L, M, b, C_Z, ρ , and Σ_M , such that the joint recursion*

$$Z_{k+1} \mid (X_j, Z_j)_{j \leq k} \sim P(Z_k, \cdot), \quad X_{k+1} = X_k - \alpha \nabla f(X_k) + \alpha \xi(Z_k)$$

has a unique invariant probability measure Π_α for every $0 < \alpha \leq \alpha_$. Its Z -marginal is π . If $(X_\alpha, Z_\alpha) \sim \Pi_\alpha$, then*

$$W_1\left(\mathcal{L}\left(\frac{X_\alpha - x_*}{\sqrt{\alpha}}\right), \mathcal{N}(0, \Sigma)\right) \leq C\sqrt{\alpha}. \quad (11)$$

Consequently,

$$W_1(\mathcal{L}(X_\alpha), \mathcal{N}(x_*, \alpha\Sigma)) \leq C\alpha. \quad (12)$$

Theorem 2 (Matching lower bound from temporal dependence) *There are constants $c_0, \alpha_0 > 0$ and a scalar instance satisfying (5)–(7) such that, for the stationary scaled iterate $Y_\alpha = (X_\alpha - x_\star)/\sqrt{\alpha}$,*

$$W_1(\mathcal{L}(Y_\alpha), \mathcal{N}(0, 5/4)) \geq c_0\sqrt{\alpha}, \quad 0 < \alpha \leq \alpha_0. \quad (13)$$

The stationary noise marginal is symmetric, $\mathbb{E}[\xi(Z_0)^3] = 0$, and $\mathbb{E}[\xi(Z_0)\xi(Z_\ell)] = 0$ for every $\ell \in \mathbb{Z} \setminus \{0\}$. The leading correction is generated by the mixed moment $\mathbb{E}[\xi(Z_0)\xi(Z_1)^2] = -9/4$. Independent sampling from the same marginal gives zero expectation for the bounded 1-Lipschitz test function $h(x) = \sin x$.

The construction takes $Z_k = (\eta_{k-1}, \eta_k)$ for independent Rademacher innovations and $\xi(Z_k) = \eta_k(3 - \eta_{k-1})/2$. Adjacent noise variables share η_k and are dependent. Appendix G derives the stationary weighted series, computes its exact second and third moments, and obtains $\mathbb{E}[\sin(Y_\alpha)] = \frac{3}{8}e^{-5/8}\sqrt{\alpha} + O(\alpha)$ through an elementary two-state characteristic-function recursion. Since $\sin$ is 1-Lipschitz and every centered Gaussian law is symmetric, this expansion proves (13).

3. Block approximation and stationary transfer

Define the scaled iterate and its deterministic update map by

$$Y_k = \frac{X_k - x_\star}{\sqrt{\alpha}}, \quad \mathcal{T}_\alpha(y) = y - \sqrt{\alpha}\nabla f(x_\star + \sqrt{\alpha}y), \quad (14)$$

$$Y_{k+1} = \mathcal{T}_\alpha(Y_k) + \sqrt{\alpha}\xi(Z_k). \quad (15)$$

For $\alpha \leq L^{-1}$, the mean-value theorem and (5) give

$$\|\mathcal{T}_\alpha(y) - \mathcal{T}_\alpha(y')\| \leq (1 - m\alpha)\|y - y'\|. \quad (16)$$

Thus one step has a contraction gap of order α . Taking $n = \lceil T/\alpha \rceil$ steps turns this into a fixed contraction factor of order e^{-mT} .

Let P_α be the transition kernel of (Y_k, Z_k) , also use P_α for its Markov operator on functions, write λP_α^n for the law after n steps from λ , and let $\nu = \gamma \otimes \pi$. The next propositions isolate the block approximation and joint stability. Appendix B gives a proof guide, followed by the formal arguments in Appendices D and E.

Proposition 3 (One-block Gaussian approximation) *For every fixed $T > 0$, there are $C_T < \infty$ and $\alpha_T > 0$ such that, if $(Y_0, Z_0) \sim \gamma \otimes \pi$ and $n = \lceil T/\alpha \rceil$, then*

$$W_1(\mathcal{L}(Y_n), \gamma) \leq C_T\sqrt{\alpha}, \quad 0 < \alpha \leq \alpha_T. \quad (17)$$

For $\kappa > 0$, equip $\mathbb{R}^d \times \mathbb{Z}$ with the cost

$$d_{\alpha, \kappa}((y, z), (y', z')) = \|y - y'\| + \kappa\sqrt{\alpha}\mathbf{1}_{\{z \neq z'\}}. \quad (18)$$

For product-space laws λ and λ' with finite first Y -moments, $W_{d_{\alpha, \kappa}}(\lambda, \lambda')$ is the supremum of $|\int h d\lambda - \int h d\lambda'|$ over measurable functions $h : \mathbb{R}^d \times \mathbb{Z} \rightarrow \mathbb{R}$ satisfying $|h(y, z) - h(y', z')| \leq d_{\alpha, \kappa}((y, z), (y', z'))$. Every coupling upper-bounds this distance by its expected cost. The factor $\sqrt{\alpha}$ matches the size of one noise increment in (15).

Proposition 4 (One-block contraction of the joint process) *There are $T > 0$, $0 < \kappa < \infty$, $q \in (0, 1)$, and $\alpha_c > 0$ such that, with $n = \lceil T/\alpha \rceil$,*

$$W_{d_{\alpha, \kappa}}(\lambda P_{\alpha}^n, \lambda' P_{\alpha}^n) \leq q W_{d_{\alpha, \kappa}}(\lambda, \lambda') \quad (19)$$

for all probability measures with finite first Y -moments and every $0 < \alpha \leq \alpha_c$.

The one-block approximation uses two Poisson equations. We state them here because they identify the two places where serial dependence enters the proof.

Lemma 5 (Noise and covariance correctors) *Let $\mathcal{F}_k = \sigma(Z_0, \dots, Z_k)$ and define*

$$V = \sum_{j \geq 0} P^j \xi, \quad (20)$$

$$D_{k+1} = V(Z_{k+1}) - PV(Z_k), \quad (21)$$

$$\mathcal{C}(z) = P(VV^{\top})(z) - (PV)(z)(PV)(z)^{\top}, \quad (22)$$

$$W = \sum_{j \geq 0} P^j (\mathcal{C} - \Sigma_M). \quad (23)$$

Both series converge uniformly, with nuclear-norm convergence for W . Moreover, $V - PV = \xi$, (D_{k+1}) is a bounded martingale difference sequence, and

$$\mathbb{E}[D_{k+1} D_{k+1}^{\top} \mid \mathcal{F}_k] = \mathcal{C}(Z_k), \quad \int \mathcal{C} \, d\pi = \Sigma_M. \quad (24)$$

The exact identities

$$\xi(Z_k) = D_{k+1} + V(Z_k) - V(Z_{k+1}), \quad W - PW = \mathcal{C} - \Sigma_M \quad (25)$$

will be used for summation by parts and covariance telescoping.

The first corrector turns the Markov sum into a martingale sum plus a bounded coboundary remainder. The second turns predictable covariance fluctuations into an exact telescoping sum. These are the two points at which temporal dependence changes the Gaussian replacement. Appendix E constructs the invariant law and gives the complete stationary-transfer argument.

4. Proof of Theorem 1

Fix the block duration from Proposition 4, let $n = \lceil T/\alpha \rceil$, and start from $(Y_0, Z_0) \sim \gamma \otimes \pi$. We follow the comparison path and record the order of each error.

Step 1: linearize the drift. Set $A = I_d - \alpha H$. This matrix is symmetric. Let

$$\bar{Y}_{k+1} = A\bar{Y}_k + \sqrt{\alpha}\xi(Z_k), \quad \bar{Y}_0 = Y_0.$$

The Lipschitz Hessian condition and Taylor's formula give

$$\|\mathcal{T}_{\alpha}(y) - Ay\| \leq \frac{M}{2} \alpha^{3/2} \|y\|^2. \quad (26)$$

Stationarity of the Markov chain and absolute summability of (Γ_ℓ) imply

$$\sup_{0 \leq k \leq n} \mathbb{E} \|\bar{Y}_k\|^2 \leq \text{tr}(\Sigma) + (T+1)C_\Gamma.$$

Coupling Y_k and $\bar{Y}_k$ with the same Markov path, then iterating (16) and (26), yields

$$\mathbb{E} \|Y_n - \bar{Y}_n\| \leq C_T \sqrt{\alpha}. \quad (27)$$

Step 2: reduce the Markov sum to a martingale sum. For $r = 1, \dots, n$, write $B_r = \sqrt{\alpha} A^{n-r}$. Substituting the first identity in (25) into the explicit linear recursion and applying weighted summation by parts gives

$$\bar{Y}_n = A^n Y_0 + \sum_{r=1}^n B_r D_r + R_{V,n}, \quad (28)$$

where

$$R_{V,n} = \sqrt{\alpha} \left[A^{n-1} V(Z_0) - V(Z_n) + \alpha \sum_{j=1}^{n-1} A^{n-1-j} H V(Z_j) \right].$$

The first two terms are endpoints, and the sum records the variation of the matrix weights. Since V is bounded, $\|A^j\|_{\text{op}} \leq 1$, and $\alpha n \leq T+1$,

$$\|R_{V,n}\| \leq C_T \sqrt{\alpha}. \quad (29)$$

Step 3: replace the martingale increments by Gaussian increments. Let $N_r \stackrel{\text{iid}}{\sim} \mathcal{N}(0, \Sigma_M)$ be independent of $(Y_0, (Z_k))$. After restricting $\alpha \leq (2L)^{-1}$, $\text{Cov}(A^n Y_0) \succeq c_T I_d$ for some $c_T > 0$ independent of α . Hence the function $\psi(u) = \mathbb{E}[h(A^n Y_0 + u)]$ has a bounded and Lipschitz Hessian for every 1-Lipschitz h .

For the r th replacement, let $U_r = \sum_{1 \leq i < r} B_i D_i + \sum_{r < i \leq n} B_i N_i$. Conditional centering cancels the linear terms, while the Taylor remainders total $O(n\alpha^{3/2}) = O(\sqrt{\alpha})$. With $K_r = B_r^\top \nabla^2 \psi(U_r) B_r$, the second Poisson equation gives

$$\begin{aligned} \sum_{r=1}^n \mathbb{E} \langle K_r, \mathcal{C}(Z_{r-1}) - \Sigma_M \rangle_{\text{F}} &= \mathbb{E} \langle K_1, W(Z_0) \rangle_{\text{F}} - \mathbb{E} \langle K_n, W(Z_n) \rangle_{\text{F}} \\ &\quad + \sum_{r=1}^{n-1} \mathbb{E} \langle K_{r+1} - K_r, W(Z_r) \rangle_{\text{F}}. \end{aligned}$$

The boundary terms are $O(\alpha)$, while $\mathbb{E} \|K_{r+1} - K_r\|_{\text{op}} = O(\alpha^{3/2} + \alpha^2)$. Thus the variation sum is $O(\sqrt{\alpha})$, and

$$W_1 \left(\mathcal{L} \left(A^n Y_0 + \sum_{r=1}^n B_r D_r \right), \mathcal{L} \left(A^n Y_0 + \sum_{r=1}^n B_r N_r \right) \right) \leq C_T \sqrt{\alpha}. \quad (30)$$

Step 4: identify the target covariance. The Gaussian vector in (30) has covariance $\Sigma_{\alpha,n} = A^n \Sigma A^n + \alpha \sum_{j=0}^{n-1} A^j \Sigma_M A^j$. The identities $\Sigma_M = H \Sigma + \Sigma H$ and $A = I_d - \alpha H$ give

$$\Sigma_{\alpha,n} - \Sigma = \alpha^2 \sum_{j=0}^{n-1} A^j H \Sigma H A^j, \quad \|\Sigma_{\alpha,n} - \Sigma\|_F \leq C_T \alpha. \quad (31)$$

Appendix F converts this covariance error into an $O(\alpha)$ Wasserstein error. Equations (27), (29), and (30) then prove Proposition 3.

Step 5: contract the joint process and transfer to stationarity. Uniform ergodicity gives a coupling with coalescence time τ satisfying $\mathbb{P}(\tau > k) \leq C_{\text{cpl}} \rho_{\text{cpl}}^k$. Under this coupling, strong convexity contracts the iterate distance by $1 - m\alpha$ after the Markov paths meet, while bounded noise adds at most $2b\sqrt{\alpha}$ before meeting. Choosing T and κ therefore makes the n -step joint kernel contract by a fixed factor $q < 1$. Appendix E constructs its unique invariant law ϖ_α and proves $(1 - q)W_{d_{\alpha,\kappa}}(\varpi_\alpha, \nu) \leq (C_T + 2\kappa)\sqrt{\alpha}$. Projection gives (11), and undoing the scaling gives (12).

References

- Sebastian Allmeier and Nicolas Gast. Computing the bias of constant-step stochastic approximation with Markovian noise. *Advances in Neural Information Processing Systems*, 2024.
- Albert Benveniste, Michel Métivier, and Pierre Priouret. *Adaptive algorithms and stochastic approximations*. Springer Science & Business Media, 2012.
- Rajendra Bhatia. *Positive definite matrices*. Princeton University Press, 2009.
- Sourav Chatterjee. A generalization of the Lindeberg principle. *Annals of Probability*, 2006.
- Zaiwei Chen, Shancong Mou, and Siva Theja Maguluri. Stationary behavior of constant stepsize SGD-type algorithms: An asymptotic characterization. *Proceedings of the ACM on Measurement and Analysis of Computing Systems*, 2022.
- Zaiwei Chen, Siva T Maguluri, Sanjay Shakkottai, and Karthikeyan Shanmugam. A Lyapunov theory for finite-sample guarantees of Markovian stochastic approximation. *Operations Research*, 2024.
- Persi Diaconis and David Freedman. Iterated random functions. *SIAM review*, 1999.
- Aymeric Dieuleveut, Alain Durmus, and Francis Bach. Bridging the gap between constant step size stochastic gradient descent and Markov chains. *Annals of Statistics*, 2020.
- Randal Douc, Eric Moulines, Pierre Priouret, and Philippe Soulier. *Markov Chains*. Springer, 2018.
- Alain Durmus, Eric Moulines, Alexey Naumov, and Sergey Samsonov. Finite-time high-probability bounds for Polyak–Ruppert averaged iterates of linear stochastic approximation. *Mathematics of Operations Research*, 2025.

- Arnold M Faden. The existence of regular conditional probabilities: necessary and sufficient conditions. *The Annals of Probability*, 1985.
- Gersende Fort. Central limit theorems for stochastic approximation with controlled Markov chain dynamics. *ESAIM: Probability and Statistics*, 2015.
- David Griffeath. A maximal coupling for Markov chains. *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete*, 1975.
- Abhishek Gupta and William B Haskell. Convergence of recursive stochastic algorithms using Wasserstein divergence. *SIAM Journal on Mathematics of Data Science*, 2021.
- Shaan Ul Haque, Zedong Wang, Zixuan Zhang, and Siva Theja Maguluri. How accurately can a Gaussian approximate stochastic approximation iterates? *arXiv preprint arXiv:2602.13906*, 2026.
- Dongyan Huo, Yudong Chen, and Qiaomin Xie. Bias and extrapolation in Markovian linear stochastic approximation with constant step sizes. *Mathematics of Operations Research*, 2026.
- Dongyan Lucy Huo, Yixuan Zhang, Yudong Chen, and Qiaomin Xie. The collusion of memory and nonlinearity in stochastic approximation with constant stepsize. *Advances in Neural Information Processing Systems*, 2024.
- Seo Taek Kong and R Srikant. Finite-sample Wasserstein error bounds and concentration inequalities for nonlinear stochastic approximation. *arXiv preprint arXiv:2602.02445*, 2026.
- Harold J Kushner and G George Yin. *Stochastic approximation and recursive algorithms and applications*. Springer, 2003.
- Stephan Mandt, Matthew D Hoffman, and David M Blei. Stochastic gradient descent as approximate Bayesian inference. *Journal of Machine Learning Research*, 2017.
- Ibrahim Merad and Stéphane Gaïffas. Convergence and concentration properties of constant step-size SGD through markov chains. *Electronic Journal of Statistics*, 2025.
- Sean P Meyn and Richard L Tweedie. *Markov chains and stochastic stability*. Springer Science & Business Media, 2012.
- Wenlong Mou, Ashwin Pananjady, Martin J Wainwright, and Peter L Bartlett. Optimal and instance-dependent guarantees for Markovian linear stochastic approximation. *Mathematical Statistics and Learning*, 2024.
- Georg Ch Pflug. Stochastic minimization with constant step-size: asymptotic laws. *SIAM Journal on Control and Optimization*, 1986.
- Bernhard A Schmitt. Perturbation bounds for matrix square roots and Pythagorean sums. *Linear algebra and its applications*, 1992.

- Gugan Thoppe and Vivek Borkar. A concentration bound for stochastic approximation via Alekseev’s formula. *Stochastic Systems*, 2019.
- Cédric Villani et al. *Optimal transport: old and new*. Springer, 2009.
- Yuyang Wang, Felix Wang, Zedong Wang, Ijay Narang, Yuzhou Wang, and Siva Theja Maguluri. Steady-state behavior of constant-stepsizes stochastic approximation: Gaussian approximation and tail bounds. In *Forty-third International Conference on Machine Learning*, 2026.
- Ziyang Wei, Jiaqi Li, Zhipeng Lou, and Wei Biao Wu. Gaussian approximation and concentration of constant learning-rate stochastic gradient descent. In *Advances in Neural Information Processing Systems*, 2025.
- Yixuan Zhang and Qiaomin Xie. Wasserstein- p central limit theorem rates: From local dependence to Markov chains. In *Abstracts of the 2026 ACM SIGMETRICS International Conference on Measurement and Modeling of Computer Systems*, pages 216–218. Association for Computing Machinery, 2026a. Full version: arXiv:2601.08184v3.
- Yixuan Zhang and Qiaomin Xie. Gaussian approximation for multivariate Martingale sums from uniformly ergodic Markov chains. *arXiv preprint arXiv:2609.09480*, 2026b.
- Yixuan Zhang and Qiaomin Xie. Steady-state convergence of stochastic approximation. *arXiv preprint arXiv:2609.14922*, 2026c.
- Yixuan Zhang, Dongyan Huo, Yudong Chen, and Qiaomin Xie. Prelimit coupling and steady-state convergence of constant-step-size nonsmooth contractive stochastic approximation. *Operations Research*, 2026.

Appendix A. Connections with recent Gaussian approximation results

The CLT for time-varying observables in [Zhang and Xie \(2026a\)](#), Theorem 3) offers an alternative to our direct Gaussian comparison for the linearized block. Fix $T > 0$, let $n = \lceil T/\alpha \rceil$, and set $A = I_d - \alpha H$. The noise contribution to this block is

$$S_{\alpha,n} = \sqrt{\alpha} \sum_{k=0}^{n-1} A^{n-1-k} \xi(Z_k) = \frac{1}{\sqrt{n}} \sum_{k=0}^{n-1} h_{\alpha,k}(Z_k),$$

$$h_{\alpha,k}(z) = \sqrt{n\alpha} A^{n-1-k} \xi(z).$$

For $0 < \alpha \leq \min\{1, L^{-1}\}$, these centered observables satisfy $\sup_{0 \leq k < n} \|h_{\alpha,k}\|_{\infty} \leq \sqrt{T+1} b$. Under stationarity, geometric covariance decay gives

$$\text{Cov}(S_{\alpha,n}) = \alpha \sum_{j=0}^{n-1} A^j \Sigma_M A^j + O(\alpha) \longrightarrow \int_0^T e^{-tH} \Sigma_M e^{-tH} dt \succ 0 \quad \text{as } \alpha \downarrow 0,$$

where the error is measured in operator norm. For the fixed chain and horizon, these bounds make the CLT constants uniform for sufficiently small α , yielding an $O(n^{-1/2}) = O(\sqrt{\alpha})$ Wasserstein-1 comparison with $\mathcal{N}(0, \text{Cov}(S_{\alpha,n}))$. The Gaussian comparator for $A^n Y_0 + S_{\alpha,n}$, with $Y_0 \sim \gamma$ independent of the chain, has covariance $\Sigma + O(\alpha)$ and is $O(\alpha)$ -close to γ in Wasserstein-1 distance. The nonlinear comparison and joint contraction then yield our stationary bound. Our proof establishes the block comparison directly by Poisson–Lindeberg interpolation.

[Haque et al. \(2026, Proposition 4.5\)](#) obtain a finite-time lower bound for linear stochastic approximation from a nonzero third moment of an i.i.d. noise projection. Our matching stationary lower bound instead isolates temporal dependence: the noise marginal is symmetric and all nonzero-lag autocovariances vanish, yet $\mathbb{E}[\xi(Z_0)\xi(Z_1)^2] = -9/4$ produces the leading order- $\sqrt{\alpha}$ correction.

Concurrent work by [Zhang and Xie \(2026b,c\)](#) develops sharp Gaussian bounds for martingale sums from uniformly ergodic Markov chains and for contractive stochastic approximation with multiplicative Markov noise. Under local quadratic linearization, their stationary result gives the optimal $O(\sqrt{\alpha})$ rate in Wasserstein-2 distance for this broader class of recursions. By Hölder’s inequality, $W_1(\mu, \nu) \leq W_2(\mu, \nu)$ for the same pair of laws, so their stationary result also implies the $O(\sqrt{\alpha})$ rate in Wasserstein-1 distance. For SGD with bounded additive Markov noise, our direct Wasserstein-1 proof uses a Gaussian initial condition for smoothing and Poisson correctors for the noise and covariance fluctuations. The four-state example above gives a matching $\Omega(\sqrt{\alpha})$ lower bound driven by higher-order temporal dependence, even with zero stationary mean bias and vanishing nonzero-lag noise autocovariances.

Appendix B. Reader’s guide to the upper-bound proof

This section gives a calculation-level roadmap for the upper bound. It explains the block length, the two Poisson equations, the Lindeberg interpolation, and the transfer from one block to stationarity. The formal estimates appear in Appendices [C–F](#).

B.1. Block length and the linear reference recursion

After centering and scaling, the update is

$$Y_{k+1} = \mathcal{T}_\alpha(Y_k) + \sqrt{\alpha} \xi(Z_k).$$

For two points y, y' , the fundamental theorem of calculus gives

$$\mathcal{T}_\alpha(y) - \mathcal{T}_\alpha(y') = \left[I_d - \alpha \int_0^1 \nabla^2 f(x_\star + \sqrt{\alpha}(y' + t(y - y'))) dt \right] (y - y').$$

The averaged Hessian in brackets has eigenvalues in $[m, L]$. When $\alpha \leq L^{-1}$, the bracketed matrix therefore has operator norm at most $1 - m\alpha$. This proves the one-step contraction in (16).

Linearizing the drift at $x_\star$ gives

$$A = I_d - \alpha H, \quad \bar{Y}_{k+1} = A\bar{Y}_k + \sqrt{\alpha} \xi(Z_k).$$

The eigenvalues of A lie in $[0, 1 - m\alpha]$, so

$$\|A^j\|_{\text{op}} \leq (1 - m\alpha)^j \leq e^{-m\alpha j}.$$

One update has only an order- α contraction gap. A block of $n = \lceil T/\alpha \rceil$ updates satisfies

$$\|A^n\|_{\text{op}} \leq e^{-m\alpha n} \leq e^{-mT},$$

which is a fixed contraction factor once T is fixed.

Iterating the linear recursion gives

$$\bar{Y}_n = A^n Y_0 + \sqrt{\alpha} \sum_{k=0}^{n-1} A^{n-1-k} \xi(Z_k).$$

After restricting $\alpha \leq (2L)^{-1}$, the first term remains a nondegenerate Gaussian vector over a fixed block when $Y_0 \sim \gamma$. The second term is a matrix-weighted additive functional of the Markov chain.

B.2. The first Poisson equation removes the conditional mean

For a centered observable,

$$P^j \xi(z) = \mathbb{E}[\xi(Z_j) \mid Z_0 = z].$$

Uniform ergodicity makes this conditional mean decay geometrically. Hence

$$V = \sum_{j \geq 0} P^j \xi$$

converges uniformly and satisfies $V - PV = \xi$. Define

$$D_{k+1} = V(Z_{k+1}) - PV(Z_k).$$

The Markov property yields $\mathbb{E}[D_{k+1} \mid \mathcal{F}_k] = 0$, and the Poisson equation gives

$$\xi(Z_k) = D_{k+1} + V(Z_k) - V(Z_{k+1}).$$

Thus the additive functional is a martingale sum plus a coboundary.

For the weights

$$B_r = \sqrt{\alpha} A^{n-r},$$

write $V_r = V(Z_r)$. Expanding the weighted coboundary sum and collecting the coefficient of each interior V_r gives

$$\sum_{r=1}^n B_r (V_{r-1} - V_r) = B_1 V_0 - B_n V_n + \sum_{r=1}^{n-1} (B_{r+1} - B_r) V_r.$$

This is the discrete summation-by-parts identity used in the proof. The endpoint weights have operator norm at most $\sqrt{\alpha}$, while

$$B_{r+1} - B_r = \alpha^{3/2} A^{n-r-1} H.$$

The variation of one weight is therefore $O(\alpha^{3/2})$. There are $O(\alpha^{-1})$ weights in a block, so boundedness of V makes the entire coboundary correction $O(\sqrt{\alpha})$.

B.3. Gaussian smoothing and Lindeberg interpolation

After the first Poisson equation, the leading variable is

$$A^n Y_0 + \sum_{r=1}^n B_r D_r.$$

Let N_r be independent $\mathcal{N}(0, \Sigma_M)$ vectors. The Lindeberg interpolation replaces $B_r D_r$ by $B_r N_r$ one index at a time.

The propagated initial condition provides the smoothing needed for a second-order Taylor expansion. Under the restriction $\alpha \leq (2L)^{-1}$, its covariance is $A^n \Sigma A^n$, which is bounded below by $c_T I_d$ for a constant $c_T > 0$ independent of α . For a 1-Lipschitz test function h , define

$$\psi(u) = \mathbb{E}[h(A^n Y_0 + u)].$$

Gaussian convolution gives a bounded Hessian and a Lipschitz Hessian for ψ .

At position r , set

$$U_r = \sum_{i < r} B_i D_i + \sum_{i > r} B_i N_i, \quad \mathcal{G}_r = \mathcal{F}_{r-1} \vee \sigma(N_i : i > r).$$

Then U_r is $\mathcal{G}_r$ -measurable. The future Gaussian variables are independent of the Markov chain, so

$$\mathbb{E}[D_r \mid \mathcal{G}_r] = 0, \quad \mathbb{E}[D_r D_r^\top \mid \mathcal{G}_r] = \mathcal{C}(Z_{r-1}).$$

The Gaussian candidate N_r is conditionally centered with covariance Σ_M . The linear terms in the two Taylor expansions cancel after conditioning. Since $\|B_r\|_{\text{op}} \leq \sqrt{\alpha}$, each cubic remainder is $O(\alpha^{3/2})$. Summing over $n = O(\alpha^{-1})$ positions gives an $O(\sqrt{\alpha})$ contribution.

B.4. The second Poisson equation removes covariance fluctuations

The quadratic Taylor term contains

$$\mathcal{C}(Z_{r-1}) = \mathbb{E}[D_r D_r^\top \mid \mathcal{F}_{r-1}].$$

The first Poisson equation is constructed so that

$$\int \mathcal{C} \, d\pi = \Sigma_M.$$

Hence $\mathcal{C} - \Sigma_M$ is centered under π . Uniform ergodicity gives the bounded matrix corrector

$$W = \sum_{j \geq 0} P^j (\mathcal{C} - \Sigma_M), \quad W - PW = \mathcal{C} - \Sigma_M.$$

Let

$$K_r = B_r^\top \nabla^2 \psi(U_r) B_r.$$

Because K_r is $\mathcal{G}_r$ -measurable, the Markov property gives

$$\mathbb{E}\langle K_r, PW(Z_{r-1}) \rangle_F = \mathbb{E}\langle K_r, W(Z_r) \rangle_F.$$

Therefore

$$\begin{aligned} & \sum_{r=1}^n \mathbb{E}\langle K_r, \mathcal{C}(Z_{r-1}) - \Sigma_M \rangle_F \\ &= \sum_{r=1}^n \mathbb{E}\langle K_r, W(Z_{r-1}) \rangle_F - \sum_{r=1}^n \mathbb{E}\langle K_r, W(Z_r) \rangle_F \\ &= \mathbb{E}\langle K_1, W(Z_0) \rangle_F - \mathbb{E}\langle K_n, W(Z_n) \rangle_F + \sum_{r=1}^{n-1} \mathbb{E}\langle K_{r+1} - K_r, W(Z_r) \rangle_F. \end{aligned}$$

The second equality follows by shifting the index in the second sum. This is the stochastic summation-by-parts step.

The bounded Hessian gives $\|K_r\|_{\text{op}} = O(\alpha)$, so the two boundary terms are $O(\alpha)$. To control the variation, write $J_r = \nabla^2 \psi(U_r)$ and decompose

$$\begin{aligned} K_{r+1} - K_r &= (B_{r+1} - B_r)^\top J_{r+1} B_{r+1} + B_r^\top (J_{r+1} - J_r) B_{r+1} \\ &\quad + B_r^\top J_r (B_{r+1} - B_r). \end{aligned}$$

The first and third terms are $O(\alpha^2)$. The middle term is $O(\alpha^{3/2})$ in expectation because the Hessian is Lipschitz and

$$\mathbb{E}\|U_{r+1} - U_r\| = O(\sqrt{\alpha}).$$

Summing $O(\alpha^{-1})$ variation terms gives $O(\sqrt{\alpha})$.

B.5. Target covariance and transfer to stationarity

After all replacements, the comparison variable is Gaussian with covariance

$$\Sigma_{\alpha,n} = A^n \Sigma A^n + \alpha \sum_{j=0}^{n-1} A^j \Sigma_M A^j.$$

The continuous Lyapunov equation and $A = I_d - \alpha H$ give the one-step identity

$$A \Sigma A + \alpha \Sigma_M = \Sigma + \alpha^2 H \Sigma H.$$

Multiplying by A^j on the left and right and summing over j makes the left-hand side telescope. Hence

$$\Sigma_{\alpha,n} - \Sigma = \alpha^2 \sum_{j=0}^{n-1} A^j H \Sigma H A^j.$$

Since $n = O(\alpha^{-1})$, the covariance error is $O(\alpha)$.

Let ϖ_α be the invariant law of the joint scaled process and let $\nu = \gamma \otimes \pi$. The block coupling gives a fixed contraction factor $q < 1$ in the joint transportation distance. The Z -marginal of both νP_α^n and ν is π . For an admissible joint test, replacing the state coordinate by a fixed state changes each expectation by at most $\kappa\sqrt{\alpha}$. Proposition 3 therefore implies

$$W_{d_{\alpha,\kappa}}(\nu P_\alpha^n, \nu) \leq W_1(\mathcal{L}(Y_n), \gamma) + 2\kappa\sqrt{\alpha} = O(\sqrt{\alpha}).$$

Invariance and the triangle inequality now give

$$W_{d_{\alpha,\kappa}}(\varpi_\alpha, \nu) \leq q W_{d_{\alpha,\kappa}}(\varpi_\alpha, \nu) + W_{d_{\alpha,\kappa}}(\nu P_\alpha^n, \nu).$$

Because $1 - q$ is a fixed positive number, rearranging preserves the $O(\sqrt{\alpha})$ order.

For reference, the estimates used in one block are summarized below.

Source	Scale calculation	Block error
Nonlinear drift	$\alpha^{3/2} \sum_{k=0}^{n-1} \mathbb{E} \ \tilde{Y}_k\ ^2$	$O(\sqrt{\alpha})$
First Poisson correction	$\sqrt{\alpha} + n\alpha^{3/2}$	$O(\sqrt{\alpha})$
Lindeberg remainders	$n\alpha^{3/2}$	$O(\sqrt{\alpha})$
Conditional covariance	$\alpha + n(\alpha^{3/2} + \alpha^2)$	$O(\sqrt{\alpha})$
Discrete covariance	$n\alpha^2$	$O(\alpha)$

Appendix C. Consequences of uniform ergodicity

This appendix develops the Markov-chain estimates used throughout the proof. For a centered function g , the quantity $P^k g(z)$ is the conditional mean of $g(Z_k)$ when the chain starts from z . Uniform ergodicity makes this conditional mean decay geometrically. That estimate yields three consequences: a coalescent coupling with a geometric tail, absolute summability of the stationary covariance sequence, and bounded solutions of the two Poisson equations in Lemma 5. For a bounded function g taking values in a finite-dimensional normed space, recall the supremum-norm convention from Section 2 and write

$$\pi g = \int g(z) \pi(dz).$$

Lemma 6 (Uniform decay of centered observables) *Let $g : \mathbf{Z} \rightarrow E$ be bounded and measurable, where E is a finite-dimensional normed space, and suppose $\pi g = 0$. Under (6),*

$$\sup_{z \in \mathbf{Z}} \|P^k g(z)\| \leq 2C_Z \rho^k \|g\|_\infty, \quad k \geq 0. \quad (32)$$

Proof Fix $z \in \mathbf{Z}$. The centering condition $\pi g = 0$ gives

$$P^k g(z) = \int g(u) [P^k(z, du) - \pi(du)].$$

Let E^* be the dual space of E , with dual norm denoted by $\|\cdot\|_{E^*}$. The dual representation of the norm yields

$$\|P^k g(z)\| = \sup_{\substack{\ell \in E^* \\ \|\ell\|_{E^*} \leq 1}} \left| \int \ell(g(u)) [P^k(z, du) - \pi(du)] \right|.$$

For every functional in this supremum, $|\ell(g(u))| \leq \|g\|_\infty$. With the total-variation convention fixed above, integration against a bounded scalar function satisfies

$$\left| \int q d(\mu - \nu) \right| \leq 2\|q\|_\infty \|\mu - \nu\|_{\text{TV}}.$$

Consequently,

$$\|P^k g(z)\| \leq 2\|g\|_\infty \|P^k(z, \cdot) - \pi\|_{\text{TV}}.$$

Equation (6) now gives the stated bound uniformly in z . ■

Lemma 7 (Geometric coalescent coupling from uniform ergodicity) *Under (6), there exist constants $C_{\text{cpl}} < \infty$ and $\rho_{\text{cpl}} \in (0, 1)$, depending only on C_Z and ρ , and a coupling kernel measurable in the initial pair $(z, z') \in \mathbf{Z}^2$. Under this kernel, the two P -chains have coalescence time*

$$\tau = \inf\{k \geq 0 : Z_j = Z'_j \text{ for every } j \geq k\}$$

satisfying

$$\mathbb{P}_{z, z'}(\tau > k) \leq C_{\text{cpl}} \rho_{\text{cpl}}^k \mathbf{1}_{\{z \neq z'\}}, \quad k \geq 0. \quad (33)$$

Here $\mathbb{P}_{z, z'}$ denotes probability under the coupling kernel started from (z, z') . The kernel can be chosen Markovian at the boundaries of fixed-length blocks, which is the structure used in the joint contraction argument.

Proof The construction uses the r -step skeleton of the chain. Choose an integer $r \geq 1$ such that

$$2C_Z \rho^r \leq \frac{1}{2}.$$

For any two initial states $z, z' \in \mathbf{Z}$, the triangle inequality and uniform ergodicity imply

$$\begin{aligned} \|P^r(z, \cdot) - P^r(z', \cdot)\|_{\text{TV}} &\leq \|P^r(z, \cdot) - \pi\|_{\text{TV}} + \|P^r(z', \cdot) - \pi\|_{\text{TV}} \\ &\leq 2C_Z \rho^r \leq \frac{1}{2}. \end{aligned}$$

Thus the two r -step transition laws overlap uniformly. At each block boundary, use a maximal coupling for the two endpoints at the next boundary. Their conditional disagreement probability is at most $1/2$. On a Polish space, the maximal coupling can be selected measurably in the initial pair (Griffeath, 1975; Douc et al., 2018; Meyn and Tweedie, 2012).

Apply this endpoint coupling successively at times $0, r, 2r, \dots$. The iterate recursion also uses the intermediate states inside each block. To retain the correct path law, sample those states from the regular conditional law of a Markov path given its initial and terminal states. Such bridge laws exist on a Polish space (Faden, 1985). Once the two endpoints agree at a block boundary, use the same transitions in both coordinates thereafter. The two paths then remain equal.

If $z = z'$, this synchronous construction starts at time zero. If $z \neq z'$, every block before coalescence has conditional meeting probability at least $1/2$. Therefore

$$\mathbb{P}_{z,z'}(\tau > jr) \leq 2^{-j}, \quad j \geq 0.$$

For an arbitrary integer $k \geq 0$, the event $\{\tau > k\}$ is contained in the event that the chains have not coalesced by the preceding block boundary. Hence

$$\mathbb{P}_{z,z'}(\tau > k) \leq 2^{-\lfloor k/r \rfloor} \leq 2(2^{-1/r})^k.$$

This proves (33) with $C_{\text{cpl}} = 2$ and $\rho_{\text{cpl}} = 2^{-1/r}$. ■

Lemma 8 (Absolute covariance summability) *Let Γ_ℓ be the stationary covariance matrices defined in (8). Under (6)–(7),*

$$\|\Gamma_\ell\|_* \leq 2b^2 C_Z \rho^\ell, \quad \ell \geq 1. \tag{34}$$

Consequently, the constant C_Γ in (9) is finite, and the long-run covariance series in (2) converges absolutely in nuclear norm.

Proof By the Markov property and stationarity,

$$\Gamma_\ell = \mathbb{E}_\pi \left[\xi(Z_0) (P^\ell \xi)(Z_0)^\top \right].$$

For a rank-one matrix $uv^\top$, the nuclear norm is $\|u\| \|v\|$. Lemma 6, applied to the centered function ξ , therefore gives

$$\begin{aligned} \|\Gamma_\ell\|_* &\leq \mathbb{E}_\pi \left[\|\xi(Z_0)\| \|P^\ell \xi(Z_0)\| \right] \\ &\leq b \cdot 2C_Z \rho^\ell b = 2b^2 C_Z \rho^\ell. \end{aligned}$$

The term Γ_0 is finite because ξ is bounded. Since $\Gamma_{-\ell} = \Gamma_\ell^\top$ and the nuclear norm is invariant under transposition, the same geometric estimate holds for negative lags. Summing over $\ell \in \mathbb{Z}$ proves (9). ■

Proof [Proof of Lemma 5] The proof follows the two Poisson equations in the order in which they are used later.

Step 1: construct the noise corrector. Apply Lemma 6 to the centered function ξ . Since $P^j \xi(z) = \mathbb{E}[\xi(Z_j) \mid Z_0 = z]$, the j th term represents the expected noise j steps into the future. The series

$$V(z) = \sum_{j=0}^{\infty} P^j \xi(z)$$

converges uniformly because the j th term is bounded by $2bC_Z \rho^j$. In particular,

$$\|V\|_{\infty} \leq \frac{2bC_Z}{1-\rho} =: v_V. \quad (35)$$

Uniform convergence permits termwise application of P , so

$$V - PV = \sum_{j=0}^{\infty} P^j \xi - \sum_{j=1}^{\infty} P^j \xi = \xi.$$

Step 2: obtain the martingale-coboundary decomposition. Define

$$D_{k+1} = V(Z_{k+1}) - PV(Z_k).$$

The Markov property gives

$$\mathbb{E}[V(Z_{k+1}) \mid \mathcal{F}_k] = PV(Z_k),$$

and hence $\mathbb{E}[D_{k+1} \mid \mathcal{F}_k] = 0$. Also,

$$D_{k+1} + V(Z_k) - V(Z_{k+1}) = V(Z_k) - PV(Z_k) = \xi(Z_k).$$

Thus the first identity in (25) holds. Equation (35) gives the uniform increment bound

$$\|D_{k+1}\| \leq 2v_V \quad \text{almost surely.} \quad (36)$$

A direct conditional second-moment calculation yields

$$\begin{aligned} \mathbb{E}[D_{k+1} D_{k+1}^{\top} \mid \mathcal{F}_k] &= P(VV^{\top})(Z_k) - (PV)(Z_k)(PV)(Z_k)^{\top} \\ &= \mathcal{C}(Z_k). \end{aligned}$$

Step 3: identify the mean conditional covariance. The absolute convergence in (9) allows us to sum the lag covariances inside expectation. Since $V = \sum_{j \geq 0} P^j \xi$, the Markov property gives

$$\mathbb{E}_{\pi}[\xi(Z_0) V(Z_0)^{\top}] = \sum_{j \geq 0} \mathbb{E}_{\pi}[\xi(Z_0) (P^j \xi)(Z_0)^{\top}] = \sum_{j \geq 0} \Gamma_j, \quad \mathbb{E}_{\pi}[V(Z_0) \xi(Z_0)^{\top}] = \sum_{j \geq 0} \Gamma_j^{\top}.$$

The term Γ_0 appears in both sums. Therefore

$$\Sigma_M = \mathbb{E}_{\pi} \left[\xi V^{\top} + V \xi^{\top} - \xi \xi^{\top} \right], \quad (37)$$

where all functions in the expectation are evaluated at Z_0 . Substitute $\xi = V - PV$ into the matrix inside the expectation. The cross terms cancel and give

$$\xi V^\top + V \xi^\top - \xi \xi^\top = V V^\top - (PV)(PV)^\top.$$

Finally, invariance of π implies

$$\mathbb{E}_\pi[V V^\top] = \mathbb{E}_\pi[P(V V^\top)].$$

Combining these identities with the definition of $\mathcal{C}$ proves

$$\int \mathcal{C}(z) \pi(\mathrm{d}z) = \Sigma_M. \quad (38)$$

This direct calculation is the reason the covariance in the Gaussian replacement is exactly the long-run covariance.

Step 4: construct the covariance corrector. The matrix $\mathcal{C}(z)$ is positive semidefinite. By (36),

$$\sup_z \|\mathcal{C}(z)\|_* = \sup_z \mathrm{tr}(\mathcal{C}(z)) \leq 4v_V^2.$$

Thus $g := \mathcal{C} - \Sigma_M$ is a bounded matrix-valued function, and (38) shows that $\pi g = 0$. Lemma 6, applied to symmetric matrices with the nuclear norm, shows that

$$W = \sum_{j=0}^{\infty} P^j g$$

converges uniformly. Termwise subtraction gives $W - PW = g$, and

$$\|W\|_{\infty,*} \leq \frac{2C_Z}{1-\rho} \|\mathcal{C} - \Sigma_M\|_{\infty,*} < \infty.$$

This proves all claims. ■

Appendix D. Proof of the one-block Gaussian approximation

This appendix proves Proposition 3. Fix $T > 0$, restrict to $0 < \alpha \leq 1$, let $n = \lceil T/\alpha \rceil$, and assume $(Y_0, Z_0) \sim \gamma \otimes \pi$. The comparison proceeds from the nonlinear recursion to its linearization, then to a weighted martingale sum, then to a Gaussian weighted sum, and finally to the target Gaussian law. Appendix B gives an informal derivation of the same path. Here every approximation is stated with the estimate used in the final triangle inequality.

D.1. Linearization and finite-block moment bounds

Set

$$A := I_d - \alpha H, \quad \bar{Y}_{k+1} = A \bar{Y}_k + \sqrt{\alpha} \xi(Z_k), \quad \bar{Y}_0 = Y_0. \quad (39)$$

The matrix A is the explicit Euler update for the linearized drift $-Hy$. To compare the two recursions, observe that

$$\mathcal{T}_\alpha(y) - Ay = -\sqrt{\alpha} [\nabla f(x_\star + \sqrt{\alpha} y) - H \sqrt{\alpha} y].$$

The integral Taylor formula gives

$$\nabla f(x_\star + \sqrt{\alpha}y) - H\sqrt{\alpha}y = \int_0^1 [\nabla^2 f(x_\star + t\sqrt{\alpha}y) - H] \sqrt{\alpha}y \, dt.$$

Using the Hessian Lipschitz condition and integrating t over $[0, 1]$ yields

$$\|\mathcal{T}_\alpha(y) - Ay\| \leq \frac{M}{2} \alpha^{3/2} \|y\|^2. \quad (40)$$

The linear recursion has the explicit representation

$$\bar{Y}_k = A^k Y_0 + \sqrt{\alpha} \sum_{i=0}^{k-1} A^{k-1-i} \xi(Z_i).$$

The two terms are centered and independent because Y_0 is independent of the stationary Markov chain. For $\alpha \leq L^{-1}$, the symmetric matrix A has spectrum in $[0, 1]$, so $\|A^j\|_{\text{op}} \leq 1$. The next calculation shows that the weighted Markov sum has a second moment bounded uniformly over a fixed block. Using (8), stationarity gives

$$\mathbb{E}\|\bar{Y}_k\|^2 = \text{tr}(A^k \Sigma A^k) + \alpha \sum_{i,j=0}^{k-1} \text{tr}(A^{k-1-i} \Gamma_{j-i} A^{k-1-j}). \quad (41)$$

Stationarity expresses all cross terms through the lag covariances Γ_{j-i} , whose absolute summability controls their total contribution. The first term is at most $\text{tr}(\Sigma)$. For the second term, cyclicity of the trace and operator–nuclear norm duality give

$$\left| \text{tr}(A^{k-1-i} \Gamma_{j-i} A^{k-1-j}) \right| \leq \|\Gamma_{j-i}\|_*.$$

For each i , the sum over j is bounded by C_Γ . Therefore

$$\sup_{0 \leq k \leq n} \mathbb{E}\|\bar{Y}_k\|^2 \leq \text{tr}(\Sigma) + (T+1)C_\Gamma =: C_{\text{mom}}(T), \quad (42)$$

where we used $\alpha n \leq T+1$.

Construct Y_k and $\bar{Y}_k$ with the same initial condition and the same Markov path. Add and subtract $\mathcal{T}_\alpha(\bar{Y}_k)$, use the contraction (16), and apply (40) at $\bar{Y}_k$:

$$\|Y_{k+1} - \bar{Y}_{k+1}\| \leq (1 - m\alpha) \|Y_k - \bar{Y}_k\| + \frac{M}{2} \alpha^{3/2} \|\bar{Y}_k\|^2.$$

Iteration from $Y_0 = \bar{Y}_0$ gives

$$\begin{aligned} \mathbb{E}\|Y_n - \bar{Y}_n\| &\leq \frac{M}{2} \alpha^{3/2} \sum_{k=0}^{n-1} (1 - m\alpha)^{n-1-k} \mathbb{E}\|\bar{Y}_k\|^2 \\ &\leq \frac{MC_{\text{mom}}(T)}{2m} \sqrt{\alpha}. \end{aligned} \quad (43)$$

The final bound uses (42) and the geometric-series estimate $\sum_{j \geq 0} (1 - m\alpha)^j = (m\alpha)^{-1}$. Thus the nonlinear-to-linear comparison contributes $O(\sqrt{\alpha})$ over one block.

D.2. Weighted martingale–coboundary decomposition

Insert (25) into the explicit solution of the linear recursion. With the reindexing $r = i + 1$, define

$$B_r = \sqrt{\alpha} A^{n-r}, \quad r = 1, \dots, n. \quad (44)$$

The coboundary terms can be summed exactly. This is the matrix-weighted version of the elementary identity $\sum_r (v_{r-1} - v_r) = v_0 - v_n$. Indeed,

$$\begin{aligned} \sum_{r=1}^n B_r [V(Z_{r-1}) - V(Z_r)] \\ = B_1 V(Z_0) - B_n V(Z_n) + \sum_{r=1}^{n-1} (B_{r+1} - B_r) V(Z_r). \end{aligned} \quad (45)$$

Here $B_1 = \sqrt{\alpha} A^{n-1}$, $B_n = \sqrt{\alpha} I_d$, and

$$B_{r+1} - B_r = \alpha^{3/2} A^{n-r-1} H.$$

Substitution into the linear recursion gives the exact decomposition

$$\bar{Y}_n = A^n Y_0 + \sum_{r=1}^n B_r D_r + R_{V,n}, \quad (46)$$

where

$$R_{V,n} = \sqrt{\alpha} \left[A^{n-1} V(Z_0) - V(Z_n) + \alpha \sum_{j=1}^{n-1} A^{n-1-j} H V(Z_j) \right]. \quad (47)$$

The first two terms are the endpoints of the coboundary sum. The final term is the accumulated variation of the matrix weights in (45). Since $\|A^j\|_{\text{op}} \leq 1$, $\|H\|_{\text{op}} \leq L$, and $\alpha n \leq T + 1$, we obtain

$$\|R_{V,n}\| \leq v_V (2 + L(T + 1)) \sqrt{\alpha}. \quad (48)$$

Consequently, replacing $\bar{Y}_n$ by the weighted martingale sum in (46) costs $O(\sqrt{\alpha})$ in Wasserstein distance.

D.3. Gaussian smoothing for Lipschitz test functions

Let $N_1, \dots, N_n$ be independent $\mathcal{N}(0, \Sigma_M)$ vectors, independent of Y_0 and the Markov chain. Set $\beta_1 = \mathbb{E}\|N_1\|$ and $\beta_3 = \mathbb{E}\|N_1\|^3$. Both quantities are finite and depend only on Σ_M . We compare

$$A^n Y_0 + \sum_{r=1}^n B_r D_r \quad \text{with} \quad A^n Y_0 + \sum_{r=1}^n B_r N_r.$$

The propagated initial condition provides the smoothing needed for this comparison. Since $Y_0 \sim \mathcal{N}(0, \Sigma)$, the covariance of $A^n Y_0$ is $A^n \Sigma A^n$. Restrict $\alpha \leq (2L)^{-1}$. Then $\lambda_{\min}(A) \geq 1 - \alpha L$ and $\log(1 - \alpha L) \geq -2\alpha L$. Since $\alpha n \leq T + 1$,

$$A^n \Sigma A^n \succeq \lambda_{\min}(\Sigma) e^{-4L(T+1)} I_d =: c_T I_d. \quad (49)$$

Thus the smoothing covariance remains uniformly nondegenerate over the block. For a 1-Lipschitz test function h , define

$$\psi(u) = \mathbb{E}[h(A^n Y_0 + u)].$$

Gaussian convolution makes ψ twice differentiable with a Lipschitz Hessian. Appendix F proves the uniform bounds

$$\sup_u \|\nabla^2 \psi(u)\|_{\text{op}} \leq a_{2,T}, \quad \|\nabla^2 \psi(u) - \nabla^2 \psi(v)\|_{\text{op}} \leq a_{3,T} \|u - v\|, \quad (50)$$

where one may take $a_{2,T} = a_2(c_T)$ and $a_{3,T} = a_3(c_T)$ with the functions defined in Appendix F. These constants are independent of α .

D.4. Lindeberg replacement

For $r = 1, \dots, n$, define the Lindeberg hybrid

$$U_r = \sum_{i=1}^{r-1} B_i D_i + \sum_{i=r+1}^n B_i N_i. \quad (51)$$

Thus the increments before position r are martingale increments, those after position r are Gaussian, and the r th position is left open. Empty sums are zero. Replacing the open position and summing over r gives the Lindeberg telescoping identity (Chatterjee, 2006)

$$\begin{aligned} & \mathbb{E}\psi\left(\sum_{i=1}^n B_i D_i\right) - \mathbb{E}\psi\left(\sum_{i=1}^n B_i N_i\right) \\ &= \sum_{r=1}^n [\mathbb{E}\psi(U_r + B_r D_r) - \mathbb{E}\psi(U_r + B_r N_r)]. \end{aligned} \quad (52)$$

Indeed, $U_r + B_r D_r$ is the hybrid with martingale increments through position r , while $U_r + B_r N_r$ is the hybrid with martingale increments only through position $r-1$. Consecutive hybrids therefore cancel in the sum. For the r th summand, condition on

$$\mathcal{G}_r = \mathcal{F}_{r-1} \vee \sigma(N_i : r < i \leq n).$$

This σ -field contains every random variable already present in U_r , so U_r is $\mathcal{G}_r$ -measurable. Adding the independent future Gaussian variables does not change conditional moments of D_r . The martingale property therefore gives

$$\mathbb{E}[D_r \mid \mathcal{G}_r] = 0, \quad \mathbb{E}[D_r D_r^\top \mid \mathcal{G}_r] = \mathcal{C}(Z_{r-1}),$$

while N_r is conditionally centered with covariance Σ_M .

Apply the second-order Taylor formula for ψ at U_r . For any increment $x \in \mathbb{R}^d$, the Hessian-Lipschitz bound in (50) gives

$$\left| \psi(U_r + x) - \psi(U_r) - \langle \nabla \psi(U_r), x \rangle - \frac{1}{2} x^\top \nabla^2 \psi(U_r) x \right| \leq \frac{a_{3,T}}{6} \|x\|^3. \quad (53)$$

Use (53) first with $x = B_r D_r$ and then with $x = B_r N_r$. Conditional on $\mathcal{G}_r$, both increments are centered, so the linear terms vanish. Writing

$$K_r = B_r^\top \nabla^2 \psi(U_r) B_r,$$

the difference of the quadratic terms is

$$\begin{aligned} & \mathbb{E}\psi(U_r + B_r D_r) - \mathbb{E}\psi(U_r + B_r N_r) \\ &= \frac{1}{2} \mathbb{E}\langle K_r, \mathcal{C}(Z_{r-1}) - \Sigma_M \rangle_F + \mathcal{R}_r. \end{aligned} \quad (54)$$

By (21) and boundedness of V , $\|D_r\| \leq 2v_V$, while (44) gives $\|B_r\|_{\text{op}} \leq \sqrt{\alpha}$. The two Taylor remainders therefore satisfy the explicit bound

$$|\mathcal{R}_r| \leq \frac{a_{3,T}}{6} \alpha^{3/2} ((2v_V)^3 + \beta_3). \quad (55)$$

Since $n \leq T/\alpha + 1$, summing (55) gives

$$\sum_{r=1}^n |\mathcal{R}_r| \leq C_T n \alpha^{3/2} \leq C_T \sqrt{\alpha}. \quad (56)$$

D.5. Predictable covariance correction

We next sum the predictable covariance errors. By (25),

$$\mathcal{C}(Z_{r-1}) - \Sigma_M = W(Z_{r-1}) - PW(Z_{r-1}).$$

The key point is that K_r is $\mathcal{G}_r$ -measurable even though it remains correlated with Z_{r-1} . Thus K_r is predictable for the transition from Z_{r-1} to Z_r . The Markov property and conditional expectation give

$$\begin{aligned} \mathbb{E}\langle K_r, W(Z_r) \rangle_F &= \mathbb{E}[\mathbb{E}[\langle K_r, W(Z_r) \rangle_F \mid \mathcal{G}_r]] \\ &= \mathbb{E}\langle K_r, PW(Z_{r-1}) \rangle_F. \end{aligned}$$

Thus the correlation between K_r and the current Markov state is retained inside an exact telescoping identity. More precisely, each summand equals

$$\mathbb{E}\langle K_r, W(Z_{r-1}) \rangle_F - \mathbb{E}\langle K_r, W(Z_r) \rangle_F.$$

After summing over r , the first sum is indexed by $Z_0, \dots, Z_{n-1}$ and the second by $Z_1, \dots, Z_n$. Pairing the common interior states gives

$$\begin{aligned} \sum_{r=1}^n \mathbb{E}\langle K_r, \mathcal{C}(Z_{r-1}) - \Sigma_M \rangle_F &= \mathbb{E}\langle K_1, W(Z_0) \rangle_F - \mathbb{E}\langle K_n, W(Z_n) \rangle_F \\ &\quad + \sum_{r=1}^{n-1} \mathbb{E}\langle K_{r+1} - K_r, W(Z_r) \rangle_F. \end{aligned} \quad (57)$$

The boundary terms can be bounded directly. By (50) and (44),

$$\|K_r\|_{\text{op}} \leq a_{2,T} \|B_r\|_{\text{op}}^2 \leq a_{2,T} \alpha.$$

Operator–nuclear norm duality therefore gives

$$|\mathbb{E}\langle K_1, W(Z_0) \rangle_{\text{F}}| + |\mathbb{E}\langle K_n, W(Z_n) \rangle_{\text{F}}| \leq 2a_{2,T} \|W\|_{\infty,*} \alpha. \quad (58)$$

For the variation terms, the weights and hybrids satisfy, for $r = 1, \dots, n-1$,

$$\|B_{r+1} - B_r\|_{\text{op}} \leq L\alpha^{3/2}, \quad U_{r+1} - U_r = B_r D_r - B_{r+1} N_{r+1}. \quad (59)$$

Write $J_r = \nabla^2 \psi(U_r)$. Adding and subtracting $B_r^\top J_{r+1} B_{r+1}$ and $B_r^\top J_r B_{r+1}$ gives

$$\begin{aligned} K_{r+1} - K_r &= (B_{r+1} - B_r)^\top J_{r+1} B_{r+1} \\ &\quad + B_r^\top (J_{r+1} - J_r) B_{r+1} \\ &\quad + B_r^\top J_r (B_{r+1} - B_r). \end{aligned}$$

The first and third terms measure the change in the weights. The middle term measures the change in the smoothed Hessian along the hybrid increment. Using (50) and (59),

$$\begin{aligned} \mathbb{E}\|K_{r+1} - K_r\|_{\text{op}} &\leq 2a_{2,T} L\alpha^2 + a_{3,T} \alpha \mathbb{E}\|U_{r+1} - U_r\| \\ &\leq 2a_{2,T} L\alpha^2 + a_{3,T} (2v_V + \beta_1) \alpha^{3/2}. \end{aligned}$$

Summing over $r = 1, \dots, n-1$ gives

$$\sum_{r=1}^{n-1} \mathbb{E}\|K_{r+1} - K_r\|_{\text{op}} \leq C_T \sqrt{\alpha}. \quad (60)$$

Multiplying the variation bound by $\|W\|_{\infty,*}$ and combining it with the two boundary terms shows that the right-hand side of (57) is $O(\sqrt{\alpha})$. Summing (54), using (56), and taking the supremum over all 1-Lipschitz h yields

$$W_1 \left(\mathcal{L} \left(A^n Y_0 + \sum_{r=1}^n B_r D_r \right), \mathcal{L} \left(A^n Y_0 + \sum_{r=1}^n B_r N_r \right) \right) \leq C_T \sqrt{\alpha}. \quad (61)$$

D.6. Identification of the target covariance

The Gaussian vector in the second law has covariance

$$\Sigma_{\alpha,n} = A^n \Sigma A^n + \alpha \sum_{j=0}^{n-1} A^j \Sigma_M A^j. \quad (62)$$

Using $\Sigma_M = H\Sigma + \Sigma H$ and $A = I_d - \alpha H$ gives

$$A\Sigma A + \alpha \Sigma_M = \Sigma + \alpha^2 H\Sigma H.$$

Rearrange this identity as $A\Sigma A - \Sigma = -\alpha\Sigma_M + \alpha^2 H\Sigma H$. Multiplying on both sides by A^j and summing over $j = 0, \dots, n-1$ makes the covariance terms telescope and gives

$$\Sigma_{\alpha,n} - \Sigma = \alpha^2 \sum_{j=0}^{n-1} A^j H \Sigma H A^j. \quad (63)$$

Since $\|A^j\|_{\text{op}} \leq 1$ and $n\alpha \leq T+1$,

$$\|\Sigma_{\alpha,n} - \Sigma\|_F \leq (T+1)\alpha \|H\Sigma H\|_F.$$

Moreover, (63) shows that $\Sigma_{\alpha,n} \succeq \Sigma$. Hence both covariance matrices are bounded below by $\lambda_{\min}(\Sigma)I_d$. Coupling the two Gaussian vectors through the same standard normal vector and applying Lemma 10 give

$$W_1(\mathcal{N}(0, \Sigma_{\alpha,n}), \mathcal{N}(0, \Sigma)) \leq C_T \alpha. \quad (64)$$

Finally, apply the triangle inequality along the comparison chain displayed at the beginning of the section. The nonlinear-to-linear term is bounded by (43), the coboundary term by (48), the martingale-to-Gaussian term by (61), and the covariance term by (64). Therefore

$$\begin{aligned} W_1(\mathcal{L}(Y_n), \gamma) &\leq C_T \sqrt{\alpha} + C_T \alpha \\ &\leq C'_T \sqrt{\alpha}, \end{aligned}$$

where the last inequality uses $\alpha \leq 1$. This proves Proposition 3.

Appendix E. Joint contraction and construction of the invariant law

This section proves Proposition 4, constructs the invariant law, and completes the proof of Theorem 1. The coupling first brings the two Markov paths together. Strong convexity then contracts the iterate coordinates because both recursions receive the same noise after coalescence.

E.1. Contraction of the joint block kernel

Fix two initial states (y, z) and (y', z') . Lemma 7 supplies a coupling of the two P -chains with coalescence time τ . Write $\mathbb{P}_{z,z'}$ and $\mathbb{E}_{z,z'}$ for probability and expectation under this coupling when it starts from (z, z') . Expectations in this subsection are taken under that coupling. Then

$$\mathbb{P}_{z,z'}(\tau > k) \leq C_{\text{cpl}} \rho_{\text{cpl}}^k \mathbf{1}_{\{z \neq z'\}}.$$

Use the coupled Markov paths to drive the two scaled recursions. Set

$$a_\alpha = 1 - m\alpha, \quad \Delta_k = \|Y_k - Y'_k\|.$$

Before coalescence, the two scaled noise increments differ by at most $2b\sqrt{\alpha}$. After coalescence, the increments agree. Combining this observation with (16) gives the pathwise inequality

$$\Delta_{k+1} \leq a_\alpha \Delta_k + 2b\sqrt{\alpha} \mathbf{1}_{\{\tau > k\}}. \quad (65)$$

Iterating (65) and taking expectations yields

$$\mathbb{E}\Delta_n \leq a_\alpha^n \Delta_0 + 2bC_{\text{cpl}}\sqrt{\alpha} \sum_{k=0}^{n-1} a_\alpha^{n-1-k} \rho_{\text{cpl}}^k \mathbf{1}_{\{z \neq z'\}}.$$

Since $a_\alpha \rightarrow 1$ and $\rho_{\text{cpl}} < 1$, restrict α further so that $a_\alpha - \rho_{\text{cpl}} \geq (1 - \rho_{\text{cpl}})/2$. The geometric convolution can then be evaluated exactly:

$$\sum_{k=0}^{n-1} a_\alpha^{n-1-k} \rho_{\text{cpl}}^k = \frac{a_\alpha^n - \rho_{\text{cpl}}^n}{a_\alpha - \rho_{\text{cpl}}} \leq \frac{2a_\alpha^n}{1 - \rho_{\text{cpl}}}.$$

Consequently,

$$\mathbb{E}\Delta_n \leq a_\alpha^n \Delta_0 + \frac{4bC_{\text{cpl}}}{1 - \rho_{\text{cpl}}} a_\alpha^n \sqrt{\alpha} \mathbf{1}_{\{z \neq z'\}}. \quad (66)$$

The same coupling controls the terminal Markov-state discrepancy:

$$\mathbb{P}(Z_n \neq Z'_n) \leq C_{\text{cpl}} \rho_{\text{cpl}}^n \mathbf{1}_{\{z \neq z'\}}. \quad (67)$$

Choose $T > 0$ so that $e^{-mT} \leq 1/4$ and take $n = \lceil T/\alpha \rceil$. Since $n\alpha \geq T$,

$$a_\alpha^n \leq e^{-m\alpha n} \leq e^{-mT} \leq \frac{1}{4}.$$

Next choose

$$\kappa \geq \frac{4bC_{\text{cpl}}}{1 - \rho_{\text{cpl}}},$$

and restrict α so that $C_{\text{cpl}} \rho_{\text{cpl}}^{\lceil T/\alpha \rceil} \leq 1/4$. The initial iterate discrepancy is then multiplied by at most $1/4$. If the initial Markov states differ, the noise-mismatch term contributes at most $\kappa\sqrt{\alpha}/4$, and the terminal state mismatch contributes at most another $\kappa\sqrt{\alpha}/4$. Therefore (66) and (67) give

$$\mathbb{E}d_{\alpha,\kappa}((Y_n, Z_n), (Y'_n, Z'_n)) \leq \frac{1}{2} d_{\alpha,\kappa}((Y_0, Z_0), (Y'_0, Z'_0)). \quad (68)$$

To pass from point states to probability measures, let h be an admissible joint Kantorovich–Rubinstein test. Coupling the two copies from point states p, p' and using (68) gives

$$|(P_\alpha^n h)(p) - (P_\alpha^n h)(p')| \leq \frac{1}{2} d_{\alpha,\kappa}(p, p').$$

For a fixed base point p_0 , the function $2[P_\alpha^n h - (P_\alpha^n h)(p_0)]$ is therefore admissible. The subtraction of the constant does not change its integral against $\lambda - \lambda'$. Integrating and taking the supremum over h proves Proposition 4 with $q = 1/2$.

E.2. Construction and uniqueness of the invariant law

Fix $0 < \alpha \leq L^{-1}$. Because Z is Polish and π is invariant, a stationary P -chain with marginal π has a two-sided realization $(Z_k)_{k \in \mathbb{Z}}$. Since $\mathcal{T}_\alpha(0) = 0$, the contraction bound gives

$$\|\mathcal{T}_\alpha(y) + \sqrt{\alpha}\xi(z)\| \leq (1 - m\alpha)\|y\| + b\sqrt{\alpha}. \quad (69)$$

For each $N \geq 1$, initialize the scaled recursion at time $-N$ from zero and denote its value at time zero by $Y_0^{(N)}$. Iterating (69) shows that every such trajectory stays in the deterministic ball of radius $b/(m\sqrt{\alpha})$.

If $N' > N$, the two trajectories use the same noise path from time $-N$ onward. At time $-N$, the trajectory started there equals zero, while the older trajectory lies in the absorbing ball. Their distance at that time is at most $b/(m\sqrt{\alpha})$. Contractivity over the next N updates gives

$$\|Y_0^{(N')} - Y_0^{(N)}\| \leq \frac{b}{m\sqrt{\alpha}}(1 - m\alpha)^N.$$

The right-hand side tends to zero, so $(Y_0^{(N)})$ is almost surely Cauchy. Let Y_0 denote its limit. This limit is measurable with respect to the infinite past of the noise chain.

Apply the same pullback construction after shifting the time origin. This defines Y_k for every $k \in \mathbb{Z}$, and the shift invariance of the two-sided noise chain makes $(Y_k, Z_k)_{k \in \mathbb{Z}}$ stationary. Passing to the limit in the recursion shows that it satisfies (15) (Diaconis and Freedman, 1999; Gupta and Haskell, 2021). Hence $\varpi_\alpha := \mathcal{L}(Y_0, Z_0)$ is invariant for P_α and has Z -marginal π .

The pullback law just constructed is supported in the deterministic ball

$$\|Y\| \leq \frac{b}{m\sqrt{\alpha}} \quad \text{almost surely.} \quad (70)$$

Indeed, the finite pullback trajectories stay in this ball by (69), and the ball is closed, so their almost-sure limit has the same bound.

Now restrict further to $\alpha \leq \alpha_c$, where α_c is supplied by Proposition 4. Every invariant law has a finite first Y -moment. Indeed, iterating (69) gives

$$\|Y_k\| \leq (1 - m\alpha)^k \|Y_0\| + \frac{b}{m\sqrt{\alpha}}. \quad (71)$$

If the law of Y_0 is invariant, then Y_k has the same law as Y_0 . For every $\varepsilon > 0$, the preceding pathwise bound gives

$$\begin{aligned} \mathbb{P}\left(\|Y_0\| > \frac{b}{m\sqrt{\alpha}} + \varepsilon\right) &= \mathbb{P}\left(\|Y_k\| > \frac{b}{m\sqrt{\alpha}} + \varepsilon\right) \\ &\leq \mathbb{P}\left(\|Y_0\| > \varepsilon(1 - m\alpha)^{-k}\right). \end{aligned}$$

The threshold on the right tends to infinity, so the probability tends to zero. Thus every invariant law is supported in the same deterministic ball as (70). Its Z -marginal is π because uniform ergodicity makes π the unique invariant law of P .

Let μ_1 and μ_2 be two invariant laws of the joint recursion. Both have finite first Y -moments, so Proposition 4 applies:

$$W_{d_{\alpha,\kappa}}(\mu_1, \mu_2) = W_{d_{\alpha,\kappa}}(\mu_1 P_{\alpha}^n, \mu_2 P_{\alpha}^n) \leq q W_{d_{\alpha,\kappa}}(\mu_1, \mu_2).$$

The dual class contains suitably rescaled products of bounded Lipschitz functions of y and bounded measurable functions of z , so it determines the joint law. Hence $q < 1$ forces $\mu_1 = \mu_2$, and the pullback law ϖ_{α} is the unique invariant law of the scaled recursion.

E.3. Transfer of the block estimate to stationarity

Recall $\nu = \gamma \otimes \pi$. Let h be an admissible joint test and fix an arbitrary $z_0 \in Z$. The function $y \mapsto h(y, z_0)$ is 1-Lipschitz. Moreover,

$$|h(y, z) - h(y, z_0)| \leq \kappa \sqrt{\alpha} \quad \text{for every } y, z.$$

Both νP_{α}^n and ν have Z -marginal π . Replacing $h(y, z)$ by $h(y, z_0)$ in each of the two expectations therefore costs at most $\kappa \sqrt{\alpha}$. Proposition 3 now gives

$$W_{d_{\alpha,\kappa}}(\nu P_{\alpha}^n, \nu) \leq (C_T + 2\kappa) \sqrt{\alpha}. \quad (72)$$

Set $w_{\alpha} = W_{d_{\alpha,\kappa}}(\varpi_{\alpha}, \nu)$. Invariance of ϖ_{α} , the triangle inequality, and Proposition 4 yield

$$\begin{aligned} w_{\alpha} &= W_{d_{\alpha,\kappa}}(\varpi_{\alpha} P_{\alpha}^n, \nu) \\ &\leq W_{d_{\alpha,\kappa}}(\varpi_{\alpha} P_{\alpha}^n, \nu P_{\alpha}^n) + W_{d_{\alpha,\kappa}}(\nu P_{\alpha}^n, \nu) \\ &\leq q w_{\alpha} + (C_T + 2\kappa) \sqrt{\alpha}. \end{aligned}$$

Therefore

$$w_{\alpha} \leq \frac{C_T + 2\kappa}{1 - q} \sqrt{\alpha}. \quad (73)$$

Projection onto the first coordinate can only decrease Wasserstein distance. Hence the Y -marginal of ϖ_{α} is within $C \sqrt{\alpha}$ of γ . The pushforward of ϖ_{α} under $(y, z) \mapsto (x_{\star} + \sqrt{\alpha} y, z)$ is the unique invariant probability measure Π_{α} of the original recursion. Scaling the iterate distance by $\sqrt{\alpha}$ also gives the unscaled $O(\alpha)$ bound. Taking $\alpha_{\star}$ to be the minimum of the finitely many positive thresholds introduced above completes the proof of Theorem 1.

Appendix F. Gaussian smoothing and covariance perturbation

This appendix records the two analytic estimates used in the Lindeberg argument. Gaussian convolution turns a Lipschitz test function into a function of class C^2 with a quantitatively Lipschitz Hessian. A separate matrix perturbation estimate converts a covariance error into a Wasserstein-1 error between centered Gaussian laws.

Lemma 9 (Derivative bounds for Gaussian convolution) *Let $S \succeq cI_d$ with $c > 0$, let $G \sim \mathcal{N}(0, S)$, and let $h : \mathbb{R}^d \rightarrow \mathbb{R}$ be 1-Lipschitz. Define $\psi(x) = \mathbb{E}[h(x + G)]$. Then ψ is*

of class C^2 and

$$\sup_x \|\nabla^2 \psi(x)\|_{\text{op}} \leq a_2(c) := \sqrt{\frac{2}{\pi}} c^{-1/2}, \quad (74)$$

$$\|\nabla^2 \psi(x) - \nabla^2 \psi(y)\|_{\text{op}} \leq a_3(c) \|x - y\|, \quad a_3(c) := \sqrt{2} c^{-1}. \quad (75)$$

Consequently, for all $x, u \in \mathbb{R}^d$,

$$\left| \psi(x+u) - \psi(x) - \langle \nabla \psi(x), u \rangle - \frac{1}{2} u^\top \nabla^2 \psi(x) u \right| \leq \frac{a_3(c)}{6} \|u\|^3. \quad (76)$$

Proof For a 1-Lipschitz function h , begin with a standard mollification h_δ . Mollification preserves the Lipschitz constant, so $\|\nabla h_\delta\|_\infty \leq 1$, and the estimates below are uniform in δ . After subtracting the constant $h(0)$, these functions have at most linear growth. Dominated convergence in the Gaussian formulas permits $\delta \downarrow 0$. It therefore suffices to prove the estimates for smooth h with $\|\nabla h\|_\infty \leq 1$.

Fix unit vectors $u, v \in \mathbb{R}^d$. Differentiating in direction v and applying Gaussian integration by parts in direction u gives

$$u^\top \nabla^2 \psi(x) v = \mathbb{E} \left[\partial_v h(x+G) u^\top S^{-1} G \right]. \quad (77)$$

The scalar $u^\top S^{-1} G$ is centered Gaussian with variance $u^\top S^{-1} u \leq c^{-1}$. Since $|\partial_v h| \leq 1$,

$$|u^\top \nabla^2 \psi(x) v| \leq \mathbb{E} |u^\top S^{-1} G| \leq \sqrt{\frac{2}{\pi}} c^{-1/2}.$$

Taking the supremum over unit u and v proves (74).

A second Gaussian integration by parts gives

$$u^\top \nabla^2 \psi(x) v = \mathbb{E} [h(x+G) Q_{u,v}(G)], \quad (78)$$

where

$$Q_{u,v}(G) = (u^\top S^{-1} G)(v^\top S^{-1} G) - u^\top S^{-1} v.$$

The random variable $Q_{u,v}(G)$ is centered. Subtracting (78) at x and y , then using the Lipschitz property of h , yields

$$\begin{aligned} |u^\top (\nabla^2 \psi(x) - \nabla^2 \psi(y)) v| &\leq \|x - y\| \mathbb{E} |Q_{u,v}(G)| \\ &\leq \|x - y\| (\mathbb{E} Q_{u,v}(G)^2)^{1/2}. \end{aligned}$$

Write $G = S^{1/2} G_0$ with $G_0 \sim \mathcal{N}(0, I_d)$. The Gaussian fourth-moment identity gives

$$\mathbb{E} Q_{u,v}(G)^2 = (u^\top S^{-1} u)(v^\top S^{-1} v) + (u^\top S^{-1} v)^2 \leq 2c^{-2}.$$

Taking the supremum over unit u, v proves (75).

Finally, for arbitrary $x, w \in \mathbb{R}^d$, Taylor's formula with integral remainder gives

$$\begin{aligned} \psi(x+w) - \psi(x) - \langle \nabla \psi(x), w \rangle - \frac{1}{2} w^\top \nabla^2 \psi(x) w \\ = \int_0^1 (1-t) w^\top [\nabla^2 \psi(x+tw) - \nabla^2 \psi(x)] w \, dt. \end{aligned}$$

Using (75) and $\int_0^1 t(1-t) \, dt = 1/6$ proves (76). ■

Lemma 10 (Perturbation of centered Gaussian laws) *Let $S_1, S_2 \succ 0$ and suppose $S_1, S_2 \succeq \lambda I_d$ for some $\lambda > 0$. Then*

$$W_1(\mathcal{N}(0, S_1), \mathcal{N}(0, S_2)) \leq \frac{\|S_1 - S_2\|_F}{2\sqrt{\lambda}}. \quad (79)$$

Proof Let $G_0 \sim \mathcal{N}(0, I_d)$ and use the common-randomness coupling $S_1^{1/2}G_0$ and $S_2^{1/2}G_0$. Jensen's inequality gives

$$\begin{aligned} W_1(\mathcal{N}(0, S_1), \mathcal{N}(0, S_2)) &\leq \mathbb{E}\|(S_1^{1/2} - S_2^{1/2})G_0\| \\ &\leq (\mathbb{E}\|(S_1^{1/2} - S_2^{1/2})G_0\|^2)^{1/2} \\ &= \|S_1^{1/2} - S_2^{1/2}\|_F. \end{aligned}$$

We next control the difference of the matrix square roots. Set $\Delta_S = S_1^{1/2} - S_2^{1/2}$. From this definition, we obtain the Sylvester identity

$$S_1 - S_2 = S_1^{1/2}\Delta_S + \Delta_S S_2^{1/2}.$$

Cauchy–Schwarz for the Frobenius inner product then gives

$$\|S_1 - S_2\|_F \|\Delta_S\|_F \geq |\langle \Delta_S, S_1 - S_2 \rangle_F| = \langle \Delta_S, S_1^{1/2}\Delta_S + \Delta_S S_2^{1/2} \rangle_F.$$

The two terms on the right are nonnegative. More explicitly,

$$\begin{aligned} \langle \Delta_S, S_1^{1/2}\Delta_S \rangle_F &= \text{tr}(\Delta_S^\top S_1^{1/2}\Delta_S) \geq \sqrt{\lambda}\|\Delta_S\|_F^2, \\ \langle \Delta_S, \Delta_S S_2^{1/2} \rangle_F &= \text{tr}(\Delta_S^\top \Delta_S S_2^{1/2}) \geq \sqrt{\lambda}\|\Delta_S\|_F^2. \end{aligned}$$

Here we used $S_i^{1/2} \succeq \sqrt{\lambda}I_d$ and the fact that $\Delta_S^\top \Delta_S$ is positive semidefinite. Therefore

$$\langle \Delta_S, S_1^{1/2}\Delta_S + \Delta_S S_2^{1/2} \rangle_F \geq 2\sqrt{\lambda}\|\Delta_S\|_F^2.$$

For $\Delta_S \neq 0$, division by $2\sqrt{\lambda}\|\Delta_S\|_F$ yields $\|\Delta_S\|_F \leq \|S_1 - S_2\|_F / (2\sqrt{\lambda})$. The same inequality is immediate when $\Delta_S = 0$. Combining this estimate with the coupling bound proves the result. Related matrix square-root perturbation inequalities are discussed by [Schmitt \(1992\)](#); [Bhatia \(2009\)](#). ■

Appendix G. Matching lower bound from temporal dependence

This appendix proves Theorem 2. The calculation has four steps. We first verify that the finite-state noise has a symmetric marginal distribution and zero autocovariance at every nonzero lag. We then identify the third-order temporal moment that remains and compute the exact second and third moments of the stationary weighted series. Kantorovich–Rubinstein duality requires a Lipschitz test function, whereas $x \mapsto x^3$ is not Lipschitz. We therefore evaluate the bounded 1-Lipschitz test function $\sin x$ through an explicit two-state characteristic-function recursion.

G.1. Markov model and third-order temporal dependence

Take $f(x) = x^2/2$. Then $x_\star = 0$, $H = 1$, and the objective assumptions in (5) hold with $m = L = 1$ and $M = 0$. Let $(\eta_k)_{k \in \mathbb{Z}}$ be independent Rademacher random variables, so

$$\mathbb{P}(\eta_k = 1) = \mathbb{P}(\eta_k = -1) = \frac{1}{2}. \quad (80)$$

Define

$$Z_k = (\eta_{k-1}, \eta_k), \quad \xi(Z_k) = \frac{\eta_k(3 - \eta_{k-1})}{2}. \quad (81)$$

The four possible noise values are displayed by

	$\eta_k = 1$	$\eta_k = -1$
$\eta_{k-1} = 1$	1	-1
$\eta_{k-1} = -1$	2	-2

The state space is $\{-1, 1\}^2$. From a state (a, b) , the chain moves to (b, c) , where c is a fresh Rademacher variable. Thus

$$P((a, b), \{(b, c)\}) = \frac{1}{2}, \quad c \in \{-1, 1\}.$$

Its invariant probability measure π is uniform on the four states. After two transitions, both coordinates are fresh and independent of the initial state, which gives

$$P^2(z, \cdot) = \pi \quad \text{for every } z. \quad (82)$$

Equation (82) verifies (6), for example with $C_Z = 1$ and $\rho = 1/2$. The observable satisfies (7) with $b = 2$.

The four equally likely values of $\xi(Z_0)$ are $-2, -1, 1, 2$. Hence the marginal distribution is symmetric and

$$\mathbb{E}\xi(Z_0) = 0, \quad \mathbb{E}\xi(Z_0)^2 = \frac{5}{2}, \quad \mathbb{E}\xi(Z_0)^3 = 0. \quad (83)$$

For every $\ell \geq 1$, the variable $\xi(Z_\ell)$ contains the fresh factor η_ℓ to the first power. Conditional on all earlier innovations, this factor has mean zero. Therefore

$$\mathbb{E}[\xi(Z_0)\xi(Z_\ell)] = 0, \quad \ell \geq 1. \quad (84)$$

Stationarity gives the same conclusion at negative lags. The long-run covariance is therefore

$$\Sigma_M = \mathbb{E}\xi(Z_0)^2 = \frac{5}{2}. \quad (85)$$

Since $H = 1$, the Lyapunov equation $2\Sigma = \Sigma_M$ gives $\Sigma = 5/4$.

At second order, this process agrees with independent draws having the same marginal distribution because every nonzero-lag covariance vanishes. Third-order moments separate the two models. Direct conditioning gives

$$\begin{aligned} \mathbb{E}[\xi(Z_0)\xi(Z_1)^2] &= \mathbb{E}\left[\frac{\eta_0(3 - \eta_{-1})}{2} \frac{(3 - \eta_0)^2}{4}\right] \\ &= \frac{1}{8}\mathbb{E}[3 - \eta_{-1}]\mathbb{E}[\eta_0(3 - \eta_0)^2] = -\frac{9}{4}. \end{aligned} \quad (86)$$

Here $\mathbb{E}[3 - \eta_{-1}] = 3$ and $\mathbb{E}[\eta_0(3 - \eta_0)^2] = -6$. By contrast, for independent centered variables $\tilde{\xi}_0$ and $\tilde{\xi}_1$ with this same marginal law,

$$\mathbb{E}[\tilde{\xi}_0 \tilde{\xi}_1^2] = \mathbb{E}\tilde{\xi}_0 \mathbb{E}\tilde{\xi}_1^2 = 0.$$

Thus (86) is generated by temporal dependence.

G.2. Stationary weighted series and exact low-order moments

Fix $\alpha \in (0, 1)$ and set $r_\alpha = 1 - \alpha$. The recursion is

$$X_{k+1} = r_\alpha X_k + \alpha \xi(Z_k).$$

Iterating backward N steps in a two-sided stationary realization gives

$$X_0 = r_\alpha^N X_{-N} + \alpha \sum_{j=0}^{N-1} r_\alpha^j \xi(Z_{-1-j}).$$

Iterating $|X_{k+1}| \leq r_\alpha |X_k| + 2\alpha$ gives $|X_0| \leq r_\alpha^N |X_{-N}| + 2(1 - r_\alpha^N)$. For every $\delta > 0$, stationarity then gives

$$\mathbb{P}(|X_0| > 2 + \delta) \leq \mathbb{P}(|X_0| > 2 + \delta r_\alpha^{-N}) \longrightarrow 0.$$

Letting $N \rightarrow \infty$ shows that $|X_0| \leq 2$ almost surely. By stationarity, the same bound holds for every X_{-N} . Hence $r_\alpha^N X_{-N} \rightarrow 0$ almost surely. The remaining series converges absolutely, and

$$Y_\alpha = \frac{X_0}{\sqrt{\alpha}} = \sqrt{\alpha} \sum_{j=0}^{\infty} r_\alpha^j \zeta_j \quad \text{almost surely,} \quad (87)$$

where we set $\varepsilon_j = \eta_{-1-j}$. The reversed variables remain independent Rademacher variables, and

$$\xi(Z_{-1-j}) = \frac{\eta_{-1-j}(3 - \eta_{-2-j})}{2} = \frac{\varepsilon_j(3 - \varepsilon_{j+1})}{2} =: \zeta_j. \quad (88)$$

The sequence (ζ_j) is 1-dependent because ζ_j only uses $(\varepsilon_j, \varepsilon_{j+1})$.

Time reversal changes the orientation of the mixed moment. The original quantity

$$\mathbb{E}[\xi(Z_0)\xi(Z_1)^2]$$

appears as $\mathbb{E}[\zeta_j^2 \zeta_{j+1}]$ in the backward stationary series. This is the nonzero orientation in the third-moment calculation below.

The marginal moments in (83) also hold for ζ_j . If $j \neq k$, then $\mathbb{E}[\zeta_j \zeta_k] = 0$. For $|j - k| \geq 2$, this follows from independence and centering. For adjacent indices, the innovation appearing only in the variable at the left endpoint has mean zero. The variance of (87) is consequently

$$\mathbb{E}[Y_\alpha^2] = \alpha \sum_{j=0}^{\infty} r_\alpha^{2j} \mathbb{E}\zeta_j^2 = \frac{5}{2} \frac{\alpha}{1 - r_\alpha^2} = \frac{5}{2(2 - \alpha)} = \frac{5}{4} + O(\alpha). \quad (89)$$

For the third moment, only configurations with two equal indices and one adjacent index can contribute. There are two orientations. Direct calculation gives

$$\mathbb{E}[\zeta_j^2 \zeta_{j+1}] = -\frac{9}{4}, \quad \mathbb{E}[\zeta_j \zeta_{j+1}^2] = 0. \quad (90)$$

For the first identity, independence separates the expectation as

$$\frac{1}{8} \mathbb{E}[\varepsilon_{j+1}(3 - \varepsilon_{j+1})^2] \mathbb{E}[3 - \varepsilon_{j+2}] = -\frac{9}{4}.$$

The second identity is zero because ε_j appears to the first power and is independent of the remaining factors. A product involving three distinct indices also has expectation zero. If the three indices are not consecutive, independence separates off a centered factor. If they are consecutive, the leftmost innovation appears only to the first power and again has conditional mean zero.

Expanding the cube in (87) and grouping the two possible adjacent repeated-index patterns gives

$$\begin{aligned} \mathbb{E}[Y_\alpha^3] &= 3\alpha^{3/2} \sum_{j=0}^{\infty} [r_\alpha^{3j+1} \mathbb{E}(\zeta_j^2 \zeta_{j+1}) + r_\alpha^{3j+2} \mathbb{E}(\zeta_j \zeta_{j+1}^2)] \\ &= -\frac{27}{4} \alpha^{3/2} \sum_{j=0}^{\infty} r_\alpha^{3j+1} = -\frac{27}{4} \frac{\alpha^{3/2} r_\alpha}{1 - r_\alpha^3}. \end{aligned} \quad (91)$$

Since $1 - r_\alpha^3 = 3\alpha - 3\alpha^2 + \alpha^3$,

$$\mathbb{E}[Y_\alpha^3] = -\frac{9}{4} \sqrt{\alpha} + O(\alpha^{3/2}). \quad (92)$$

The filtered sum therefore has an order- $\sqrt{\alpha}$ third moment while every individual observation has a symmetric law.

G.3. Elementary characteristic-function recursion

Each ζ_j depends on two adjacent Rademacher variables. After the variables up to ε_j have been processed, only the value of ε_j is needed for the next factor. This two-state memory yields an explicit recursion for the characteristic function.

Lemma 11 (Two-state characteristic-function expansion) *For the stationary series in (87), the characteristic function is nonzero at frequency one for all sufficiently small α . Here $\mathbf{i} = \sqrt{-1}$. With the logarithm defined by the near-identity product in the proof,*

$$\log \mathbb{E}[e^{\mathbf{i}Y_\alpha}] = -\frac{1}{2} \mathbb{E}[Y_\alpha^2] - \frac{\mathbf{i}}{6} \mathbb{E}[Y_\alpha^3] + O(\alpha) \quad \text{as } \alpha \downarrow 0. \quad (93)$$

Proof Set

$$\theta_j = \sqrt{\alpha} r_\alpha^j, \quad Y_{\alpha,N} = \sum_{j=0}^{N-1} \theta_j \zeta_j.$$

We integrate the innovations from left to right while retaining the terminal sign. Define

$$\begin{aligned} u_j^+ &= \mathbb{E} \left[e^{i \sum_{k=0}^{j-1} \theta_k \zeta_k} \mathbf{1}_{\{\varepsilon_j=1\}} \right], \\ u_j^- &= \mathbb{E} \left[e^{i \sum_{k=0}^{j-1} \theta_k \zeta_k} \mathbf{1}_{\{\varepsilon_j=-1\}} \right], \end{aligned}$$

and set

$$s_j = u_j^+ + u_j^-, \quad d_j = u_j^+ - u_j^-.$$

Since $u_0^+ = u_0^- = 1/2$, we have $s_0 = 1$ and $d_0 = 0$. Also, $s_N = \mathbb{E}[e^{iY_{\alpha,N}}]$.

Condition on ε_j and then average over the fresh sign ε_{j+1} . If $\varepsilon_{j+1} = 1$, then $\zeta_j = \varepsilon_j$ and the new factor is $e^{i\theta_j \varepsilon_j}$. If $\varepsilon_{j+1} = -1$, then $\zeta_j = 2\varepsilon_j$ and the new factor is $e^{2i\theta_j \varepsilon_j}$. Therefore

$$\begin{aligned} u_{j+1}^+ &= \frac{1}{2} \left(e^{i\theta_j} u_j^+ + e^{-i\theta_j} u_j^- \right), \\ u_{j+1}^- &= \frac{1}{2} \left(e^{2i\theta_j} u_j^+ + e^{-2i\theta_j} u_j^- \right). \end{aligned}$$

Adding and subtracting gives

$$s_{j+1} = a(\theta_j) s_j + i b(\theta_j) d_j, \tag{94}$$

$$d_{j+1} = c(\theta_j) s_j + i e(\theta_j) d_j, \tag{95}$$

where

$$\begin{aligned} a(t) &= \frac{\cos t + \cos(2t)}{2}, & b(t) &= \frac{\sin t + \sin(2t)}{2}, \\ c(t) &= \frac{\cos t - \cos(2t)}{2}, & e(t) &= \frac{\sin t - \sin(2t)}{2}. \end{aligned}$$

For $|t| \leq 1/2$, Taylor's theorem gives uniform expansions

$$a(t) = 1 - \frac{5}{4}t^2 + O(t^4), \quad b(t) = \frac{3}{2}t + O(t^3), \tag{96}$$

$$c(t) = \frac{3}{4}t^2 + O(t^4), \quad e(t) = -\frac{1}{2}t + O(t^3). \tag{97}$$

All $O(\cdot)$ constants below are uniform in j and in sufficiently small α .

We now control the normalized terminal-state imbalance. Whenever $s_j \neq 0$, let

$$\chi_j = \frac{d_j}{s_j}.$$

Dividing (95) by (94) gives

$$\chi_{j+1} = \frac{c(\theta_j) + i e(\theta_j) \chi_j}{a(\theta_j) + i b(\theta_j) \chi_j}. \tag{98}$$

We first establish a coarse bound that justifies every division. Choose α small enough that $\theta_j \leq \sqrt{\alpha} \leq 1/2$ and $r_\alpha^{-1} \leq 2$. The Taylor bounds imply that, for a numerical constant C_2 and every $|t| \leq 1/2$,

$$|a(t) - 1| \leq C_2 t^2, \quad |b(t)| + |e(t)| \leq C_2 |t|, \quad |c(t)| \leq C_2 t^2.$$

Set $C_\chi = 4C_2$. We claim that

$$|\chi_j| \leq C_\chi \theta_{j-1}^2, \quad |a(\theta_j) + ib(\theta_j)\chi_j| \geq \frac{1}{2}, \quad j \geq 1. \quad (99)$$

For the base case, $\chi_0 = 0$, so $|a(\theta_0)| \geq 1 - C_2\theta_0^2$. Thus χ_1 is well defined and $|\chi_1| \leq 2C_2\theta_0^2 \leq C_\chi\theta_0^2$ once α is small enough.

It remains to verify the induction implication. Suppose $|\chi_j| \leq C_\chi\theta_{j-1}^2$ and put $t = \theta_j$. Since $\theta_{j-1} \leq 2t$,

$$\begin{aligned} |c(t) + ie(t)\chi_j| &\leq C_2t^2 + 4C_2C_\chi t^3, \\ |a(t) + ib(t)\chi_j| &\geq 1 - C_2t^2 - 4C_2C_\chi t^3. \end{aligned}$$

After decreasing the upper bound on α , the second quantity is at least $1/2$, and twice the first is at most $C_\chi t^2$. Equation (98) then gives $|\chi_{j+1}| \leq C_\chi\theta_j^2$. Applying this implication first at $j = 1$ and then successively proves (99). Since $s_{j+1} = s_j[a(\theta_j) + ib(\theta_j)\chi_j]$ and $s_0 = 1$, it also proves that every s_j is nonzero.

The coarse estimate allows us to sharpen the real and imaginary parts. We claim that

$$\Re\chi_j = \frac{3}{4}\theta_{j-1}^2 + O(\theta_{j-1}^4), \quad \Im\chi_j = O(\theta_{j-1}^3), \quad j \geq 1. \quad (100)$$

For $j = 1$, the ratio $\chi_1 = c(\theta_0)/a(\theta_0)$ is real, and (100) follows from (96)–(97).

Assume (100) at index j and write $\chi_j = x_j + iy_j$. Put $t = \theta_j$. Since $\theta_{j-1} = t/r_\alpha$ and $r_\alpha^{-1} \leq 2$,

$$x_j = O(t^2), \quad y_j = O(t^3).$$

The numerator and denominator of (98) become

$$\begin{aligned} c(t) + ie(t)\chi_j &= c(t) - e(t)y_j + ie(t)x_j \\ &= \frac{3}{4}t^2 + O(t^4) + iO(t^3), \\ a(t) + ib(t)\chi_j &= a(t) - b(t)y_j + ib(t)x_j \\ &= 1 + O(t^2) + iO(t^3). \end{aligned}$$

The reciprocal of the denominator is $1 + O(t^2) + iO(t^3)$. Multiplication gives

$$\Re\chi_{j+1} = \frac{3}{4}t^2 + O(t^4), \quad \Im\chi_{j+1} = O(t^3),$$

which proves (100).

Define the one-step characteristic-function factor

$$g_j = \frac{s_{j+1}}{s_j} = a(\theta_j) + ib(\theta_j)\chi_j.$$

For $j = 0$, use $\chi_0 = 0$. For $j \geq 1$, substitute (100) into (96). Since $ib\chi = -b\Im\chi + ib\Re\chi$, we obtain

$$g_j - 1 = -\frac{5}{4}\theta_j^2 + \frac{9i}{8}\mathbf{1}_{\{j \geq 1\}}\theta_j\theta_{j-1}^2 + O(\theta_j^4). \quad (101)$$

The real correction from $-b\Im\chi$ is $O(\theta_j^4)$. The leading imaginary term is $(3\theta_j/2)(3\theta_{j-1}^2/4)$.

For small α , (101) gives $|g_j - 1| \leq 1/2$ uniformly in j . Let $\log g_j$ denote the principal logarithm on this disk. Since $\log(1+w) = w + O(|w|^2)$,

$$\log g_j = -\frac{5}{4}\theta_j^2 + \frac{9i}{8}\mathbf{1}_{\{j \geq 1\}}\theta_j\theta_{j-1}^2 + O(\theta_j^4).$$

Because $s_0 = 1$ and $s_N = \prod_{j=0}^{N-1} g_j$, define

$$\log s_N = \sum_{j=0}^{N-1} \log g_j.$$

The series of logarithms converges absolutely because $\sum_j \theta_j^2 < \infty$. Its limit defines the logarithm in the statement of the lemma, and the corresponding infinite product is nonzero. Also, $Y_{\alpha,N} \rightarrow Y_\alpha$ almost surely. Bounded convergence gives $s_N \rightarrow \mathbb{E}[e^{iY_\alpha}]$. Letting $N \rightarrow \infty$ yields

$$\log \mathbb{E}[e^{iY_\alpha}] = -\frac{5}{4} \sum_{j \geq 0} \theta_j^2 + \frac{9i}{8} \sum_{j \geq 1} \theta_j \theta_{j-1}^2 + O\left(\sum_{j \geq 0} \theta_j^4\right). \quad (102)$$

The geometric sums are

$$\sum_{j \geq 0} \theta_j^2 = \frac{1}{2 - \alpha}, \quad \sum_{j \geq 1} \theta_j \theta_{j-1}^2 = \frac{\alpha^{3/2} r_\alpha}{1 - r_\alpha^3}, \quad \sum_{j \geq 0} \theta_j^4 = O(\alpha). \quad (103)$$

Equation (89) identifies the first term in (102) as $-\mathbb{E}[Y_\alpha^2]/2$. Equation (91) identifies the second as $-i\mathbb{E}[Y_\alpha^3]/6$. This proves (93). $\blacksquare$

Apply Lemma 11 and substitute the exact moments (89) and (91). We obtain

$$\begin{aligned} \log \mathbb{E}[e^{iY_\alpha}] &= -\frac{1}{2} \frac{5}{2(2-\alpha)} + \frac{9i}{8} \frac{\alpha^{3/2} r_\alpha}{1 - r_\alpha^3} + O(\alpha) \\ &= -\frac{5}{8} + \frac{3i}{8} \sqrt{\alpha} + O(\alpha). \end{aligned} \quad (104)$$

Exponentiating gives

$$\mathbb{E}[e^{iY_\alpha}] = e^{-5/8} \left(1 + \frac{3i}{8} \sqrt{\alpha} + O(\alpha)\right). \quad (105)$$

Taking imaginary parts yields

$$\mathbb{E}[\sin(Y_\alpha)] = \frac{3}{8} e^{-5/8} \sqrt{\alpha} + O(\alpha). \quad (106)$$

G.4. Wasserstein test function and independent-sampling comparison

Let $G \sim \mathcal{N}(0, 5/4)$. The function $h(x) = \sin x$ is 1-Lipschitz, and symmetry gives $\mathbb{E}h(G) = 0$. Kantorovich–Rubinstein duality and (106) imply

$$W_1(\mathcal{L}(Y_\alpha), \mathcal{L}(G)) \geq |\mathbb{E} \sin(Y_\alpha)|.$$

Thus there exists $\alpha_0 > 0$ such that, for every $0 < \alpha \leq \alpha_0$,

$$|\mathbb{E} \sin(Y_\alpha)| \geq \frac{3}{16} e^{-5/8} \sqrt{\alpha}.$$

This proves Theorem 2 with $c_0 = \frac{3}{16} e^{-5/8}$.

It remains to make the dependence comparison explicit. Let $(\tilde{\zeta}_j)_{j \geq 0}$ be independent with the same marginal law as ζ_0 , and define

$$\tilde{Y}_\alpha = \sqrt{\alpha} \sum_{j=0}^{\infty} r_\alpha^j \tilde{\zeta}_j.$$

Independence and symmetry make the full sequence jointly invariant under global sign reversal. Hence $\tilde{Y}_\alpha$ is symmetric and $\mathbb{E} \sin(\tilde{Y}_\alpha) = 0$ for every $\alpha \in (0, 1)$. Moreover, (84) shows that Y_α and $\tilde{Y}_\alpha$ have the same variance, so they share the same limiting Gaussian comparator. The Markov construction has the order- $\sqrt{\alpha}$ sine term generated by (86), while any jointly sign-symmetric construction has zero sine expectation.

G.5. Interpretation of the lower bound

The quadratic objective $f(x) = x^2/2$ has exactly linear drift, so the nonlinear Taylor remainder from the upper-bound proof is absent. The order- $\sqrt{\alpha}$ discrepancy therefore comes entirely from temporal dependence. Nevertheless, the Markovian and independent models agree in their one-time noise law and second-order structure. Both have the symmetric marginal distribution on $\{-2, -1, 1, 2\}$ and long-run covariance $\Sigma_M = 5/2$, and every nonzero-lag covariance in the Markovian model is zero. Thus both have the limiting Gaussian law $\mathcal{N}(0, 5/4)$.

The distinction appears in the adjacent mixed moment

$$\mathbb{E}[\xi(Z_0)\xi(Z_1)^2] = -\frac{9}{4}.$$

This moment survives the geometric filtering in the stationary recursion and produces an order- $\sqrt{\alpha}$ third cumulant. The characteristic-function recursion converts that local temporal asymmetry into the following expectation:

$$\mathbb{E}[\sin(Y_\alpha)] = \frac{3}{8} e^{-5/8} \sqrt{\alpha} + O(\alpha).$$

Independent sampling has the same covariance calculation, while joint sign symmetry makes its sine expectation exactly zero.

The scale can also be read directly from the exact third-moment sum. Each adjacent repeated-index pattern has weight of order $\alpha^{3/2}r_\alpha^{3j}$, and the number of effectively contributing indices is of order α^{-1} . Equivalently,

$$\alpha^{3/2} \sum_{j \geq 0} r_\alpha^{3j} = \frac{\alpha^{3/2}}{1 - r_\alpha^3} = O(\sqrt{\alpha}).$$

The two-state recursion transfers this scale from an unbounded third-moment calculation to the bounded sine test function required by Wasserstein duality.

The coefficient $3e^{-5/8}/8$ in (106) combines the order- $\sqrt{\alpha}$ imaginary correction $3/8$ with the Gaussian damping factor $e^{-5/8}$.

The two proofs separate three effects. Strong convexity gives block contraction, Hessian regularity controls nonlinear drift, and the Poisson correctors encode temporal dependence. Under independent sampling, $\mathcal{C}(z) = \Sigma_M$ and $W = 0$. The lower example shows that higher-order dependence can still affect the order- $\sqrt{\alpha}$ correction. Thus long-run covariance determines the Gaussian limit, while the first correction may also reflect objective curvature and higher-order temporal structure.