

Subspace Uncertainty and Sharp Sampling Thresholds on the Boolean Cube

Thomas Weinberger

School of Computer and Communication Sciences, EPFL, Switzerland
thomas.weinberger@epfl.ch

October 9, 2026

Abstract

We study Gaussian regression under squared population L_2 loss in a known m -dimensional subspace of degree-at-most- k functions on the d -dimensional Boolean cube. Random inputs can undersample regions essential for prediction, delaying the parametric rate even when the model is known.

For fixed $q_0 < 1/2$, $1 \leq k \leq q_0 d$, and sufficiently large fixed A , the worst-subspace sample threshold for minimax error $A\sigma^2(m+t)/n$ with confidence $1 - e^{-t}$, $t \geq \log 4$, is

$$N = (m+t) \exp\{E_{d,k} + O(k^{1/3})\}, \quad E_{d,k} = d\Psi(k/d),$$

where $\Psi(q) = \log 2 - H(\frac{1}{2} - \sqrt{q(1-q)})$ and H is binary entropy with natural logarithms. The upper bound holds for every feasible m ; the matching lower bound holds when $m \leq \binom{d}{\lfloor k^{1/3} \rfloor}$ or $t \geq m$.

We sharpen the Polyanskiy–Samorodnitsky uncertainty principle in two respects. First, for fixed leakage $\rho \in (0, 1)$, the smallest set carrying a fraction $1 - \rho$ of a nonzero degree-at-most- k polynomial's energy has probability $\exp\{-E_{d,k} + O_{\rho,q_0}(k^{1/3})\}$. An Airy-kernel construction proves that the remainder cannot be $o(k^{1/3})$ in general. Second, we construct a subspace of dimension $\binom{d}{\lfloor k^{1/3} \rfloor}$ such that every function in the subspace has at least a fraction $1 - \rho$ of its energy on the same set, whose probability is at most $\exp\{-E_{d,k} + C_{\rho,q_0}k^{1/3}\}$. For sufficiently large k , this set is a Hamming ball.

A striking consequence is an exponential cost of noise: the parametric rate can require $(m+t)4^k \exp\{-O(k^{1/3})\}$ samples, whereas $O((m+t)2^k)$ suffice for noiseless identification. As $k \rightarrow \infty$ with $k/d \rightarrow 0$, the noisy threshold is $(m+t) \exp\{2k + o(k)\}$.

Keywords: minimax linear regression, uncertainty principle, sample complexity, boolean cube, low-degree polynomials.

Contents

1	Introduction	1
1.1	Preliminaries	2
1.2	Main results	4
1.3	Related work	6
2	Constructing a polynomial concentrated on a Hamming ball	7
2.1	Designing the polynomial in the Krawchouk basis	7
2.2	Choosing the coefficients and bounding the error	9
2.3	From an approximate eigenvector to a rare event	10
3	From moment bounds to stable sampling	11
3.1	The Kirshner–Samorodnitsky moment bound	12
3.2	Cancellation at the second moment	12
3.3	From moments to clipping	12
3.4	Uniform empirical control after clipping	13
4	Many directions on one Hamming ball	14
4.1	Separating radial concentration from variation within slices	14
4.2	Uniform energy control across harmonic directions	15
4.3	Proof of Theorem 1.8	17
5	From concentration to minimax regression	17
5.1	From shared concentration to small empirical eigenvalues	17
5.2	From empirical eigenvalues to minimax risk	18
5.3	Proof of Theorem 1.5	19
6	Sharpness of the uniform uncertainty remainder	19
6.1	Convergence and limits	19
6.2	A Gaussian polynomial with a fixed amount of tail energy	20
6.3	Transferring the construction to a Boolean subspace	21
7	Discussion	23
	References	23
A	The entropy curve and the cubic remainder	26
A.1	The entropy curve	26
A.2	Uniform cubic remainder	26
B	Classical analytic and probabilistic tools	28
B.1	Bonami–Beckner hypercontractivity	28
B.2	A bound independent of the ambient dimension	29
B.3	A bounded empirical-process inequality	29

B.4	Symmetrization and contraction	29
B.5	Elementary Gaussian probability bounds	30
C	The radial-weight identity	31
D	Probability constants for the sampling lower bounds	31
E	Comparison to the Polyanskiy–Samorodnitsky uncertainty boundary	33
F	Further bounds and model examples	33
F.1	Ambient dimension and model dimension	34
F.2	Known Walsh coordinates and finite-space caps	35

1 Introduction

A familiar promise of linear regression is that prediction error eventually scales like the number of unknown coefficients divided by the number of observations. How long must one wait before this parametric rate becomes valid? For a known model and random inputs, the answer depends on how well the observations reveal the model’s population geometry. A small model can still be difficult if much of its predictive variation occurs on rarely sampled inputs. This issue connects statistical estimation to the stability of random least squares and sampling discretization [CDL13; KKLT22; Mou22; EHE24].

We study this question for low-degree functions on the Boolean cube. The learner is given a model (in the form of a linear subspace) before observing noisy evaluations at uniformly random inputs. A model has three parameters: the ambient dimension counts input coordinates; the degree bounds the order of interactions; the subspace dimension counts unknown coefficients. The model dimension determines the number of coefficients to estimate, but not how well random observations reveal them. Stricter bounds on degree and ambient dimension restrict the class of possible models and can rule out models that are difficult to estimate. Knowing the model specifies the basis functions, but leaves their coefficients unrestricted. Some choices may therefore remain difficult to estimate from random observations, which delays the point at which the parametric rate can be guaranteed uniformly over all targets in the model. This raises the question of how these three parameters jointly determine the sample size needed for accurate prediction.

Statistical question. Among known subspaces with a prescribed ambient dimension, degree, and model dimension, how many random examples are needed before the usual parametric prediction rate is guaranteed?

A central distinction is between identification and accurate prediction. Without noise, the sampled evaluations determine the coefficients exactly when the evaluation map from the model subspace to the observed values has a trivial null space, leaving no indistinguishable targets. With noise, merely distinguishing targets in principle is insufficient: their observed responses must differ enough to be distinguished reliably.

The distribution of energy determines this separation. Throughout this work, we focus on differences between possible targets and denote these functions by h . Using a basis orthonormal under the uniform input distribution, the squared coefficient distance between two targets equals $\mathbb{E}[h(X)^2]$. On the observed inputs $X_1, \dots, X_n$, however, their squared separation is $\sum_{i=1}^n h(X_i)^2$. If this quantity is small relative to the noise variance, the data cannot reliably distinguish the targets, even when their coefficients are far apart. Concentration of the energy of h on a rarely sampled set can create precisely this discrepancy. Exact Gaussian minimax results formalize this connection through the spectrum of the empirical covariance matrix [Mou22; EHE24].

The number of poorly observed directions also matters. For noise variance σ^2 , the parametric rate allows error of order σ^2/n per coefficient, hence total error of order $\sigma^2 m/n$ in a model of dimension m . An increase confined to one direction may still fit within this total allowance. To exceed it, a single direction would have to be exceptionally difficult, or many directions would have to be difficult simultaneously. Our approach makes many directions difficult at once, so their error contributions accumulate. Concentrating their energy on one shared set ensures that samples containing few points from that set leave many directions poorly observed. Each observation inside the set supplies at most one linear constraint, while observations outside it carry little energy in these directions.

The relevant geometric tool is an uncertainty principle. Low Fourier degree restricts how much

of a function's energy can concentrate on a small set of inputs. Polyanskiy and Samorodnitsky identify a sharp asymptotic boundary for this tradeoff [PS19]. Our statistical lower bound calls for a quantitative version that also accounts for how many directions can concentrate on the same set.

Geometric question. How small can the probability of one set of Boolean inputs be while that set still carries most of the energy of every function in a large low-degree subspace?

We provide an answer via a quantitative subspace uncertainty principle, with a remainder of uniformly optimal order. It yields matching upper and lower sample thresholds in a substantial range of model dimensions. For unrestricted ambient dimension, the leading exponential factor is e^{2k} , compared with 2^k for noiseless identification. Our refinement also establishes exponential savings at finite ambient dimension.

1.1 Preliminaries

Let μ_d be uniform measure on $\{-1, 1\}^d$. The Walsh characters $\chi_S(x) = \prod_{i \in S} x_i$ form an orthonormal basis, and

$$\mathcal{P}_{d,k} = \text{span}\{\chi_S : |S| \leq k\}, \quad D_{d,k} = \dim \mathcal{P}_{d,k} = \sum_{j=0}^{\min(k,d)} \binom{d}{j}.$$

Thus d counts coordinates and k bounds interaction order. The learner receives a fixed m -dimensional $V \subseteq \mathcal{P}_{d,k}$, including an evaluable basis. There are no computational restrictions. Choose a population-orthonormal basis $\Phi_V = (\phi_1, \dots, \phi_m)^\top$.

We write $\mathbb{E}[f] = \int f(x) d\mu_d(x)$ and $\mathbb{E}_n[f] = n^{-1} \sum_i f(X_i)$. The random experiment is

$$X_i \stackrel{\text{iid}}{\sim} \mu_d, \quad Y_i = f_\theta(X_i) + \xi_i, \quad f_\theta = \theta^\top \Phi_V, \quad \xi_i \stackrel{\text{iid}}{\sim} N(0, \sigma^2), \quad (1)$$

with independent inputs and noise, $\sigma > 0$, and unknown $\theta \in \mathbb{R}^m$. The learner knows V and μ_d and may know σ . V and θ are fixed before sampling. Given the observations $\{(X_i, Y_i)\}_{i=1}^n$, it outputs an estimate $\hat{\theta} \in \mathbb{R}^m$, defining $\hat{f} = \hat{\theta}^\top \Phi_V$. By orthonormality, the population prediction loss is $\mathbb{E}[(\hat{f} - f_\theta)^2] = \|\hat{\theta} - \theta\|_2^2$.

The empirical covariance and its population counterpart are

$$\hat{\Sigma}_V = \frac{1}{n} \sum_i \Phi_V(X_i) \Phi_V(X_i)^\top, \quad \Sigma_V = \mathbb{E}[\Phi_V(X) \Phi_V(X)^\top] = I_m.$$

For a nonnegative, possibly infinite random variable Z , define $Q_{1-\delta}(Z) = \inf\{r \geq 0 : \mathbb{P}(Z \leq r) \geq 1 - \delta\}$, with $\inf \emptyset = +\infty$. Throughout, we write $t = \log(1/\delta)$ for the confidence parameter. The supplied-model minimax risk is

$$\mathcal{R}_{n,\delta}^*(V) = \inf_{\hat{\theta}_V} \sup_{\theta \in \mathbb{R}^m} Q_{1-\delta,\theta}(\|\hat{\theta}_V - \theta\|_2^2).$$

Probabilities include inputs, noise, and any estimator randomness.

Our main object of interest is the minimax prediction risk for the worst admissible subspace, with the learner allowed to adapt its estimator to the supplied V .

Definition 1.1 (Worst-subspace minimax risk). *Define*

$$\mathfrak{R}_{n,\delta}(d, k, m) := \sup_{\substack{V \subseteq \mathcal{P}_{d,k} \\ \dim V = m}} \mathcal{R}_{n,\delta}^*(V) = \sup_{\substack{V \subseteq \mathcal{P}_{d,k} \\ \dim V = m}} \inf_{\hat{\theta}_V} \sup_{\theta \in \mathbb{R}^m} Q_{1-\delta,\theta}(\|\hat{\theta}_V - \theta\|_2^2).$$

Remark 1.2 (Why the subspace is known). *Giving V to the learner separates coefficient estimation from discovering the relevant low-degree directions. Without this information, even $m = 1$ permits every degree-at-most- k target, since each lies in its own one-dimensional span. Knowing V fixes the number of unknown coefficients at m , allowing us to isolate how a degree bound k restricts the complexity of the known model and the resulting difficulty of estimation.*

We ask how many samples are needed before this risk reaches the parametric rate $\sigma^2(m + \log(1/\delta))/n$. The following threshold is the central quantity we determine.

Definition 1.3 (Parametric threshold). *For a fixed benchmark constant $A \geq 16$ and $t = \log(1/\delta)$, let $N_{\text{par}}(d, k, m, A, \delta)$ be the least integer $N \geq 1$ such that*

$$\mathfrak{R}_{n,\delta}(d, k, m) \leq \frac{A\sigma^2(m + t)}{n} \quad \text{for every } n \geq N.$$

We take $A \geq 16$ to simplify the proof constants; the value 16 has no special significance. Our results have the same leading exponent for every fixed $A \geq 16$.

We also track when empirical squared norms uniformly preserve a fixed fraction of population squared norms.

Definition 1.4 (Stable-sampling threshold). *For fixed $c \in (0, 1)$, let $N_{\text{stab}}(d, k, m, c, \delta)$ be the least integer $N \geq 1$ such that, for every $n \geq N$ and every fixed eligible V ,*

$$\mathbb{P}\{\mathbb{E}_n[h^2] \geq c\mathbb{E}[h^2] \text{ for all } h \in V\} \geq 1 - \delta.$$

Equivalently, $\mathbb{P}\{\hat{\Sigma}_V \succeq cI_m\} \geq 1 - \delta$.

Our proof of [Theorem 1.5](#) will reveal that, in its stated parameter range, N_{stab} has the same worst-subspace exponential scale as N_{par} , with constants depending on q_0 and c . Sufficiently strong stable sampling guarantees the parametric regression rate, but need not be necessary for it.

Throughout, all logarithms are natural. We define the binary entropy and the uncertainty curve by

$$\mathsf{H}(u) = -u \log u - (1 - u) \log(1 - u), \quad \Psi(q) = \log 2 - \mathsf{H}\left(\frac{1}{2} - \sqrt{q(1 - q)}\right),$$

with $0 \log 0 = 0$, $u \in [0, 1]$, and $q \in [0, 1/2]$. Set

$$E_{d,k} = d\Psi(k/d), \quad M_{d,k} = \binom{d}{\lfloor k^{1/3} \rfloor}.$$

Further notation. For real numbers, $a \vee b = \max\{a, b\}$, $a \wedge b = \min\{a, b\}$, and $(a)_+ = \max\{a, 0\}$. For symmetric matrices, $A \succeq B$ means $a^\top(A - B)a \geq 0$ for every a . We use $\|a\|_2^2 = \sum_j a_j^2$ for vectors and $\|f\|_2^2 = \mathbb{E}[f^2]$, $\langle f, g \rangle = \mathbb{E}[fg]$ for functions on the cube under discussion. The indicator of E is $\mathbf{1}_E$. We abbreviate the feature map $\Phi_V = (\phi_1, \dots, \phi_m)^\top$ by Φ when V is clear, and write $\hat{f}(S) = \mathbb{E}[f\chi_S]$ for Fourier coefficients. The Rayleigh quotient of a symmetric matrix B is $\mathcal{R}_B(a) = a^\top B a / \|a\|_2^2$ for $a \neq 0$. We write $a \lesssim b$ (equivalently $b \gtrsim a$) for an inequality up to a positive constant, and $a \asymp b$ for both inequalities; allowed parameter dependence is stated locally. Finally, $\chi_r^2 = \sum_{j=1}^r G_j^2$ for independent $G_j \sim N(0, 1)$ has the chi-squared distribution with r degrees of freedom, and $Z \succeq_{\text{st}} Y$ means $\mathbb{P}(Z > u) \geq \mathbb{P}(Y > u)$ for every real u .

1.2 Main results

Throughout, $1 \leq k \leq q_0 d$ for fixed $q_0 < 1/2$. Constants may depend on q_0 and fixed target parameters, but not on d, k, m, n, δ .

1.2.1 Sharp sample thresholds for regression

Our first result determines when the worst-subspace minimax risk reaches the parametric rate, with the dependence on degree and ambient dimension captured by $E_{d,k}$.

Theorem 1.5 (Minimax sample threshold). *Fix $q_0 \in (0, 1/2)$ and $A \geq 16$. For $1 \leq k \leq q_0 d$, $1 \leq m \leq D_{d,k}$ and $0 < \delta \leq 1/4$, put $t = \log(1/\delta)$. If $m \leq M_{d,k} \vee t$, then*

$$\left| \log \frac{N_{\text{par}}(d, k, m, A, \delta)}{m + t} - E_{d,k} \right| \leq C_{q_0, A} k^{1/3}. \quad (2)$$

The upper bound holds for every feasible m and is attained by ordinary least squares.

Remark 1.6 (Polynomial Range). *In particular, for every fixed $B > 0$, equation (2) holds for every feasible $m \leq d^B$, with constants depending additionally on B : for sufficiently large k depending on B , $M_{d,k} \geq d^B$, while for bounded k , the estimate follows from $N_{\text{par}} \gtrsim m + t$ and $E_{d,k} \leq 2k$.*

Taking the worst case over all ambient dimensions gives a particularly simple threshold, valid for every model dimension m .

Corollary 1.7 (Unrestricted ambient dimension). *For integers $k, m \geq 1$, fixed $A \geq 16$, and $0 < \delta \leq 1/4$, put $t = \log(1/\delta)$ and define*

$$N_{\text{par}}^\infty(k, m, A, \delta) := \sup_{\substack{d \geq 1 \\ m \leq D_{d,k}}} N_{\text{par}}(d, k, m, A, \delta).$$

Then

$$\left| \log \frac{N_{\text{par}}^\infty(k, m, A, \delta)}{m + t} - 2k \right| \leq C_A k^{1/3}.$$

The analogous statement holds for the stable-sampling threshold, with a constant depending on its fixed conditioning level c .

The upper bound is uniform over all ambient dimensions. For the lower bound, choosing d sufficiently large makes $m \leq M_{d,k}$ and $E_{d,k} = 2k + O(k^{1/3})$. See Appendix F.1 for details.

The finite-dimensional correction. The dependence on d can substantially reduce the threshold. As $k \rightarrow \infty$ with $k/d \rightarrow 0$, provided that $m \leq M_{d,k}$ or $t \geq m$,

$$\log \frac{N_{\text{par}}(d, k, m, A, \delta)}{m + t} = 2k - \frac{2k^2}{3d} + O_A \left(\frac{k^3}{d^2} + k^{1/3} \right).$$

When $k \ll d \ll k^{5/3}$, the correction $2k^2/(3d)$ dominates both error terms. Consequently, the threshold is smaller than its unrestricted-dimension counterpart by a factor

$$\frac{N_{\text{par}}^\infty(k, m, A, \delta)}{N_{\text{par}}(d, k, m, A, \delta)} = \exp \left\{ \left(\frac{2}{3} + o(1) \right) \frac{k^2}{d} \right\}.$$

This follows from the expansion of Ψ , see Appendix F.1.

1.2.2 The exponential cost of noise

A direct statistical consequence of Theorem 1.5 is an exponential separation between noiseless identification and noisy prediction.

Indeed, concavity of Ψ , together with $\Psi(0) = 0$ and $\Psi(1/2) = \log 2$, gives $\Psi(q) \geq 2q \log 2$ for $0 \leq q \leq \frac{1}{2}$ and hence $E_{d,k} \geq k \log 4$. Thus, for every $1 \leq k \leq q_0 d$ with fixed $q_0 < 1/2$, whenever $m \leq M_{d,k}$ or $t \geq m$,

$$N_{\text{par}}(d, k, m, A, \delta) \geq (m + t) 4^k \exp\{-C_{q_0, A} k^{1/3}\}.$$

In contrast, $C2^k(m + t)$ samples suffice for noiseless identification in every admissible subspace. Indeed, a nonzero degree-at-most- k polynomial is nonzero on at least a 2^{-k} fraction of the cube. To see this, choose a monomial of maximal degree $r \leq k$ with nonzero coefficient. For every assignment of the other coordinates, this coefficient remains unchanged, so the resulting polynomial is nonzero on at least one of the 2^r assignments to those r coordinates. Whenever the evaluation matrix has rank below m , its null space therefore contains a function that the next independent input detects with probability at least 2^{-k} , increasing the rank. Consequently,

$$\mathbb{P}(\text{rank} < m) \leq \mathbb{P}\{\text{Bin}(n, 2^{-k}) < m\} \leq e^{-t} \quad \text{if } n \geq 8 \cdot 2^k(m + t),$$

by a standard binomial lower-tail bound. Full rank permits exact recovery of the coefficients.

The separation is even larger as $k \rightarrow \infty$ with $k/d \rightarrow 0$: the expansion in Appendix F.1 gives $E_{d,k} = 2k + o(k)$. In the range $m \leq M_{d,k}$ or $t \geq m$, where our upper and lower bounds agree up to an $O(k^{1/3})$ error in the exponent, the parametric threshold is therefore $(m + t) \exp\{2k + o(k)\}$, compared with the noiseless upper bound $O((m + t)2^k)$.

1.2.3 A quantitative subspace uncertainty principle

Our geometric result quantifies energy concentration jointly over a subspace.

Theorem 1.8 (Subspace uncertainty principle). *Fix $\rho \in (0, 1)$. For every set E , if some nonzero $h \in \mathcal{P}_{d,k}$ satisfies $\mathbb{E}[h^2 \mathbf{1}_{E^c}] \leq \rho \mathbb{E}[h^2]$, then*

$$\mu_d(E) \geq \exp\{-E_{d,k} - C_{\rho, q_0} k^{1/3}\}.$$

Conversely, there exist $W \subseteq \mathcal{P}_{d,k}$ of dimension $M_{d,k}$ and a permutation-invariant set E such that

$$\mu_d(E) \leq \exp\{-E_{d,k} + C_{\rho, q_0} k^{1/3}\}, \quad \mathbb{E}[h^2 \mathbf{1}_{E^c}] \leq \rho \mathbb{E}[h^2] \quad (h \in W). \quad (3)$$

For sufficiently large k , E is a Hamming ball.

We also refer to the parameter ρ as *leakage*. Let $p_\rho(d, k, m)$ be the smallest probability of a set admitting such an m -dimensional concentrating subspace. The theorem implies

$$\left| \log \frac{1}{p_\rho(d, k, m)} - E_{d,k} \right| \leq C_{\rho, q_0} k^{1/3}, \quad 1 \leq m \leq M_{d,k}. \quad (4)$$

It gives both a one-dimensional obstruction to concentration and a subspace construction at the same precision, uniformly even as $k/d \rightarrow 0$. The next result shows that the order of this uniform remainder is optimal and cannot be improved to $o(k^{1/3})$ in general.

Proposition 1.9 (Optimal order of the uniform remainder). *There are a fixed $\rho_* \in (0, 1)$, integers $d_k \geq k^3$, and $C < \infty$ such that, for all sufficiently large k and all $1 \leq m \leq \binom{d_k}{\lfloor k^{1/3} \rfloor}$,*

$$2k^{1/3} \leq \log \frac{1}{p_{\rho_*}(d_k, k, m)} - E_{d_k, k} \leq Ck^{1/3}. \quad (5)$$

1.3 Related work

Minimax regression and low-degree learning. Mourtada relates expected minimax least-squares risk to inverse sample covariance [Mou22]. El Hanchi, Maddison, and Erdogdu give the exact quantile identity used here [EHE24]. We do not prove any new Gaussian minimax theorem. Rather, we determine its consequences for the worst known low-degree subspace by controlling the empirical spectrum sharply. Learning bounded low-degree targets without a supplied subspace [EI22; EIS23] is a different problem: target normalization and model selection change both the relevant error guarantees and the role of dimension. Noiseless interpolation and Reed–Muller erasure recovery [ASW15; BHSS22] concern rank rather than noise amplification. These distinctions explain why knowing the model removes model selection but leaves a substantial statistical question about random observation.

Random least squares and sampling stability. Cohen, Davenport, and Leviatan control random least squares through the pointwise envelope $\sup_x \|\Phi_V(x)\|_2^2$ [CDL13]. Cohen and Migliorati show that choosing a model-dependent sampling measure and suitable weights permits stability with sample size of order $m \log m$ [CM17]. In contrast, our input law is fixed and uniform: the learner can adapt its estimator to V , but cannot redirect observations toward informative regions. We seek the worst geometry compatible with Fourier degree, and obtain a uniform lower bound on empirical energy even when the feature vectors can have very large norms. This is a one-sided sampling-discretization problem [KKLT22], closely related to lower bounds on empirical covariance matrices [Yas14; Oli16] and the small-ball method [Men14; Men21].

Uncertainty and log-Sobolev inequalities. Log-Sobolev inequalities relate concentration of a function’s energy to its Fourier complexity. With the normalization $\mathcal{D}(f, f) = \sum_S |S| \hat{f}(S)^2$, the classical logarithmic Sobolev inequality on the cube [BLM13, Chapter 5], for $\mathbb{E}[f^2] = 1$, reads

$$\text{Ent}_{\mu_d}(f^2) := \mathbb{E}[f^2 \log f^2] \leq 2\mathcal{D}(f, f) \leq 2k \quad (f \in \mathcal{P}_{d,k}).$$

For an exactly supported unit function, Jensen’s inequality gives $\text{Ent}(f^2) \geq \log(1/\mu_d(\text{supp } f))$. Thus entropy inequalities already constrain spatial concentration. For approximate concentration, the retained fraction $1 - \rho$ also enters and a sharp fixed-leakage boundary requires more than this elementary bound. Samorodnitsky’s modified log-Sobolev inequality strengthens the entropy–energy relation and connects it to small-set geometry [Sam08]. Polyanskiy and Samorodnitsky develop nonlinear log-Sobolev inequalities and improved hypercontractivity to obtain their uncertainty curve [PS19]. We use the same uncertainty curve Ψ , but sharpen the energy-concentration bounds and extend them jointly to every function in a large subspace.

Hypercontractivity and sharp moment bounds. The classical Bonami–Beckner inequality [Bon70; Bec75] bounds higher moments of low-degree polynomials in terms of their second moment; see Lemma B.1. Its well-known fourth-moment specialization gives $\mathbb{E}[h^4] \leq 9^k$ for a unit-norm degree-at-most- k polynomial, yielding a loose upper bound of order $9^k(m+t)$ on the worst-subspace parametric threshold, for fixed target constants. However, using moments approaching two already gives $\exp 2k + O(k^{1/3})$ from classical hypercontractivity (Proposition B.2), which is sharp for unrestricted ambient dimension. For our dimension-dependent results, classical hypercontractivity, even when tuned for the optimal moment parameter, is not sharp enough. Instead, to match the sharper finite-dimensional exponent $E_{d,k} = d\Psi(k/d)$, we optimize the Kirshner–Samorodnitsky moment bound [KS21].

Harmonic analysis on Boolean slices. Harmonic multilinear polynomials provide the additional directions in our construction. Filmus constructs an explicit orthogonal basis for functions on a

Boolean slice, organized by harmonic degree [Fil16]. Filmus–Mossel establish the dimension formula for each harmonic degree and describe how inner products behave under exchangeable measures [FM19]. We use these latter results directly: harmonic polynomials let us turn one concentrated polynomial into many orthogonal functions whose energy concentrates on the same set.

Krawchouk polynomials and concentration. Krawchouk polynomials connect Fourier analysis on the cube with the geometry of Hamming distance. Levenshtein uses them to derive universal bounds for codes and designs in Hamming spaces [Lev95]. Their extreme zeros also enter the analysis of sum-of-squares bounds for binary polynomial optimization [SL23]. The extreme zeros indicate where the polynomial in our subsequent construction concentrates its energy, providing a geometric connection with these classical results.

Clipping and empirical quadratic processes. Combining truncation with empirical-process bounds is an established method for controlling empirical quadratic forms from below. Mendelson uses bounded Lipschitz approximations in the small-ball method [Men14], and clipped squares together with contraction and concentration in [Men21]. We follow this approach, using Bousquet’s inequality [Bou02] to track the dependence on the clipping level explicitly. Our contribution to this approach is a sharper degree- and dimension-dependent clipping scale, derived from the Kirshner–Samorodnitsky moment bound.

Airy asymptotics and sharpness. Airy functions arise in the asymptotic behavior of classical orthogonal polynomials near the outer edges of their oscillatory region. For Krawchouk polynomials, Dai and Wong obtain Airy asymptotics as part of their global analysis at a fixed positive degree-to-size ratio [DW07]. Our sharpness argument uses the convergence of the Hermite kernel to the Airy kernel [DG07; SX22]. From this, we construct a polynomial with fixed positive energy on a Gaussian tail interval, then transfer it to the cube and use harmonic polynomials to construct many orthogonal directions concentrating on the same set. The resulting tail probability establishes the optimal order $k^{1/3}$ of the uniform uncertainty remainder.

2 Constructing a polynomial concentrated on a Hamming ball

We first construct one unit polynomial whose energy is concentrated on a small set of inputs. Any such set could serve the statistical lower bound, but we aim for a Hamming ball. The motivation behind this choice is threefold: (i) its probability follows a binomial tail, naturally giving rise to the small sets we are looking for; (ii) radial polynomials admit a convenient three-term recurrence in the Krawchouk basis; and (iii) its permutation symmetry allows the later extension to many directions within V . In particular, consider $S_D = x_1 + \dots + x_D$, for which $E_u = \{S_D \geq u\}$ is the ball around the all-plus vertex with radius $\lfloor (D - u)/2 \rfloor$. Our aim is to place most of a polynomial’s energy where S_D is well above its typical scale $\sqrt{D}$ under the uniform measure.

2.1 Designing the polynomial in the Krawchouk basis

We seek a polynomial whose energy is concentrated where $S_D(x) = x_1 + \dots + x_D$ is unusually large. The set $E_u = \{S_D \geq u\}$ is a Hamming ball, and increasing u shrinks it. Since E_u depends only on S_D , we look for a polynomial in S_D .

Krawchouk polynomials form an orthonormal basis for such functions under the uniform measure, and their three-term recurrence lets us express concentration near a chosen value of S_D as a condition on the polynomial’s coefficients.

Our polynomial must satisfy four constraints:

- **Bounded degree:** $\deg p \leq \ell$.
- **Unit norm:** $\mathbb{E}[p^2] = 1$, so $p^2 d\mu_D$ is a probability measure.
- **Radial symmetry:** p depends only on S_D , allowing the harmonic lift in [Section 4](#).
- **Concentration at a high threshold:** for a prescribed leakage ρ ,

$$\mathbb{E}[p^2 \mathbf{1}_{\{S_D < u\}}] \leq \rho,$$

with u large enough that E_u has the required small probability.

The Krawchouk basis makes the first three constraints explicit and gives a useful algebraic description of the fourth. Define

$$e_j(x) = \binom{D}{j}^{-1/2} \sum_{|S|=j} \chi_S(x), \quad 0 \leq j \leq D.$$

These are orthonormal radial functions of degree j that form a basis for all functions of S_D . Thus $p = \sum_{j=0}^{\ell} c_j e_j$ with $\sum_{j=0}^{\ell} c_j^2 = 1$ automatically satisfies the degree, symmetry, and normalization constraints.

There is also a connection to heavy tails: Krawchouk polynomials nearly attain sharp higher-moment bounds among polynomials of the same degree and L_2 norm. Kirshner and Samorodnitsky [KS21] establish this near extremality. However, large moments alone do not give our prescribed concentration on one ball. Indeed, $e_j(-x) = (-1)^j e_j(x)$, so a single e_j has equally heavy upper and lower tails. We will combine several degrees to concentrate energy in the upper tail.

Multiplication measures concentration. For any unit p ,

$$\varepsilon^2 := \|(S_D - \tau)p\|_2^2 = \mathbb{E}[(S_D - \tau)^2 p^2]$$

describes the mean squared distance of S_D from τ under $p^2 d\mu_D$. Hence

$$\mathbb{E}[p^2 \mathbf{1}_{\{S_D < \tau - a\}}] \leq \frac{\varepsilon^2}{a^2} \quad (a > 0).$$

For $\varepsilon > 0$, taking $a = \varepsilon/\sqrt{\rho}$ gives the desired leakage on the ball with cutoff $u = \tau - \varepsilon/\sqrt{\rho}$.

If $\varepsilon = 0$, all energy is within $S_D = \tau$, so $u = \tau$ works. We therefore want a large center τ and a small multiplication residual ε . Both quantities matter: a large residual would force us to lower the cutoff, making the ball larger.

The following recurrence allows us to express this objective in terms of the coefficients c_j .

Lemma 2.1 (Radial recurrence). *Put $a_j = \sqrt{(j+1)(D-j)}$, with $a_{-1} = a_D = 0$ and $e_{-1} = e_{D+1} = 0$. Then*

$$S_D e_j = a_j e_{j+1} + a_{j-1} e_{j-1}. \quad (6)$$

Each e_j is a univariate polynomial of degree j in S_D .

Proof. Multiplying a Walsh character by x_i adds i to its index set if absent and removes it if present. Each resulting set of size $j+1$ occurs $j+1$ times, and each set of size $j-1$ occurs $D-j+1$ times. Normalizing gives (6). Starting with $e_0 = 1$, solving successively for e_{j+1} proves the degree assertion. $\square$

Consequently, multiplying p by S_D transforms its coefficient vector according to the linear map

$$J_D = \begin{pmatrix} 0 & a_0 & 0 & \cdots & 0 \\ a_0 & 0 & a_1 & \ddots & \vdots \\ 0 & a_1 & 0 & \ddots & 0 \\ \vdots & \ddots & \ddots & \ddots & a_{D-1} \\ 0 & \cdots & 0 & a_{D-1} & 0 \end{pmatrix}, \quad (J_D c)_j = a_{j-1}c_{j-1} + a_j c_{j+1}.$$

Here $c_j = 0$ outside $0, \dots, \ell$. Orthonormality gives the exact identity

$$\|(S_D - \tau)p\|_2 = \|(J_D - \tau I)c\|_2.$$

Choosing the coefficients c_j . We now seek a unit vector supported on degrees at most ℓ for which $\|(J_D - \tau I)c\|_2$ is small: in other words, an *approximate eigenvector* of J_D with approximate eigenvalue τ . Let L denote the number of nonzero coefficients. Three constraints guide our construction:

- **Use consecutive degrees.** J_D couples only adjacent indices. Consecutive coefficients allow the two neighboring contributions to reinforce each other while gaps interrupt this reinforcement.
- **Place them near ℓ .** The weights a_j increase through the relevant range below $D/2$. Degrees near the largest permitted degree therefore give large weights. Over a short interval near ℓ , both neighboring weights are close to $a_* = \sqrt{\ell(D - \ell)}$. If $c_{j-1} + c_{j+1} \approx 2c_j$, then $(J_D c)_j \approx 2a_* c_j$. This suggests $\tau = 2a_*$. When $\ell \ll D$, this center is approximately $2\sqrt{D\ell}$, well above the uniform scale $\sqrt{D}$ for large ℓ . A cutoff close to it therefore defines a rare event.
- **Taper them to zero.** The degree restriction requires $c_{\ell+1} = 0$, but $((J_D - \tau I)c)_{\ell+1} = a_\ell c_\ell$. An abrupt jump from $c_\ell = L^{-1/2}$ to zero would therefore contribute $a_\ell/\sqrt{L}$ to the residual:

$$\|(J_D - \tau I)c\|_2 \geq \frac{a_\ell}{\sqrt{L}}.$$

We instead make the coefficients rise gradually from zero and fall gradually back to zero, keeping $\Delta^2 c_j = c_{j-1} - 2c_j + c_{j+1}$ small, including at the ends of the selected interval.

We are hence looking to construct a *smooth window*: a coefficient sequence supported on L consecutive degrees and tapered to zero at both ends, with small second differences. The next section constructs such a sequence. Increasing L permits smaller second differences, but includes matrix weights farther from a_* . Balancing these two errors determines L .

2.2 Choosing the coefficients and bounding the error

The next lemma makes this plan quantitative.

Lemma 2.2 (A concentrating polynomial). *Fix $q_0 < 1/2$. For all sufficiently large $\ell \leq q_0 D$, there is a unit radial polynomial p of exact degree ℓ such that*

$$\|(S_D - 2\sqrt{\ell(D - \ell)})p\|_2 \leq C_{q_0} \sqrt{D} \ell^{-1/6}. \quad (7)$$

Proof. We use the $L = \lfloor \ell^{1/3} \rfloor$ degrees $\ell - L + 1, \dots, \ell$. To make the coefficients small near the two ends of this interval, define

$$b(x) = \begin{cases} x^2(1-x)^2, & 0 \leq x \leq 1, \\ 0, & \text{otherwise,} \end{cases} \quad c_{\ell-L+r} = Z_L^{-1/2} b\left(\frac{r}{L+1}\right) \quad (1 \leq r \leq L),$$

where $Z_L = \sum_{r=1}^L b(r/(L+1))^2$ normalizes $\sum_j c_j^2$ to one. All other coefficients are zero, including at integer indices outside $0, \dots, D$. For large ℓ , $2 \leq L \leq \ell/4$.

We first bound the second differences of this sequence. The function b is bounded above and below by positive constants on $[1/4, 3/4]$. A fixed positive fraction of the L sampling points lie in that interval, so $cL \leq Z_L \leq CL$ for absolute positive constants c, C . In particular, the normalization contributes a factor of order $L^{-1/2}$.

Both b and its first derivative vanish at 0 and 1. Extending b by zero therefore leaves its derivative continuous, with $|b'(u) - b'(v)| \leq C|u - v|$ for all u, v : inside $[0, 1]$ its second derivative is bounded, and outside it the derivative is zero. For the sampling step $\delta_L = 1/(L+1)$, it follows that

$$|b(x + \delta_L) - 2b(x) + b(x - \delta_L)| = \left| \int_0^{\delta_L} (b'(x+t) - b'(x - \delta_L + t)) dt \right| \leq C\delta_L^2.$$

This estimate also holds when one of the sampled points is outside $[0, 1]$. Consequently, for $\Delta^2 c_j = c_{j+1} - 2c_j + c_{j-1}$, every second difference has magnitude at most $CZ_L^{-1/2}\delta_L^2 \leq CL^{-5/2}$. Only the $L+2$ indices $\ell-L, \dots, \ell+1$ can contribute, since at all other indices the three coefficients are zero. Hence

$$\|\Delta^2 c\|_{\ell_2(\mathbb{Z})} = \left(\sum_{j \in \mathbb{Z}} |\Delta^2 c_j|^2 \right)^{1/2} \leq \sqrt{L+2} CL^{-5/2} \leq CL^{-2}.$$

Set $p = \sum_j c_j e_j$, so $\|p\|_2 = 1$ and $c_\ell > 0$.

For $a_* = \sqrt{\ell(D-\ell)}$, the coefficient of $(S_D - 2a_*)p$ at j is

$$((J_D - 2a_* I)c)_j = a_* \Delta^2 c_j + (a_j - a_*)c_{j+1} + (a_{j-1} - a_*)c_{j-1}.$$

Whenever a shifted coefficient is nonzero, $|j - \ell| \leq L + 2$ and j is comparable to ℓ . Since

$$|a_j^2 - a_*^2| \leq CD(|j - \ell| + 1), \quad a_j + a_* \geq c_{q_0} \sqrt{D\ell},$$

we have $|a_j - a_*| \leq C_{q_0} \sqrt{D/\ell}(L+3)$. Thus

$$\|(J_D - 2a_* I)c\|_2 \leq C_{q_0} \left(\frac{\sqrt{D\ell}}{L^2} + \sqrt{\frac{D}{\ell}}(L+3) \right) \leq C_{q_0} \sqrt{D} \ell^{-1/6}.$$

The estimate includes the two indices $\ell - L$ and $\ell + 1$, just outside the chosen interval of degrees. Orthogonality of the e_j proves (7). $\square$

2.3 From an approximate eigenvector to a rare event

Lemma 2.2 places the energy of p near $\tau = 2\sqrt{\ell(D-\ell)}$, but does not determine how much lies above τ . The ball $\{S_D \geq \tau\}$ could therefore miss substantial energy. Instead, put its cutoff at $u = \tau - a$, with $a > 0$. Every excluded point then has distance greater than a from τ , so

$$\mathbb{E}[p^2 \mathbf{1}_{\{S_D < u\}}] \leq \frac{\|(S_D - \tau)p\|_2^2}{a^2} \leq \frac{C_{q_0}^2 D \ell^{-1/3}}{a^2}.$$

Choosing $a = M_{\rho,q_0} \sqrt{D} \ell^{-1/6}$ with a sufficiently large constant makes this leakage at most ρ . This displacement is only $a/\tau = O_{\rho,q_0}(\ell^{-2/3})$ relative to the center. The next lemma also shows that enlarging the ball in this way costs only $O_{\rho,q_0}(\ell^{1/3})$ in its probability exponent.

Lemma 2.3 (One-dimensional concentration witness). *Fix $\rho \in (0, 1)$ and $q_0 \in (0, 1/2)$. There are constants $M_{\rho,q_0}, C_{\rho,q_0} > 0$ such that, for all sufficiently large $\ell \leq q_0 D$, there is a unit radial polynomial p of exact degree ℓ for which the cutoff $\tau = 2\sqrt{\ell(D-\ell)}$, $u = \tau - M_{\rho,q_0} \sqrt{D} \ell^{-1/6} > 0$ satisfies*

$$\mathbb{E}[p^2 \mathbf{1}_{\{S_D < u\}}] \leq \rho, \quad \mu_D(S_D \geq u) \leq \exp\{-D\Psi(\ell/D) + C_{\rho,q_0} \ell^{1/3}\}.$$

Proof. Step 1: retaining the energy. Take p from Lemma 2.2 and set $a = M_{\rho,q_0} \sqrt{D} \ell^{-1/6}$, so $u = \tau - a$. On $\{S_D < u\}$, the distance from S_D to τ exceeds a . Consequently,

$$\mathbb{E}[p^2 \mathbf{1}_{\{S_D < u\}}] \leq \frac{\mathbb{E}[(S_D - \tau)^2 p^2]}{a^2} \leq \frac{C_{q_0}^2 D \ell^{-1/3}}{M_{\rho,q_0}^2 D \ell^{-1/3}} \leq \rho$$

if $M_{\rho,q_0} \geq C_{q_0}/\sqrt{\rho}$. Also $\tau \geq 2\sqrt{(1-q_0)D\ell}$, so $a/\tau = O_{\rho,q_0}(\ell^{-2/3})$. Thus $0 < u < \tau < D$ for large ℓ .

Step 2: bounding the uniform probability. For $0 \leq z < 1$, write $I(z) = \log 2 - H((1-z)/2)$, $I'(z) = \operatorname{arctanh} z$. The binomial Chernoff bound gives $\mu_D(S_D \geq u) \leq \exp\{-DI(u/D)\}$. At the center τ , the rate is exactly

$$DI(\tau/D) = D\Psi(\ell/D).$$

We therefore need to bound the loss in this rate when the cutoff moves from τ down to u . Since I' is increasing,

$$D\Psi(\ell/D) - DI(u/D) = D \int_{u/D}^{\tau/D} I'(z) dz \leq (\tau - u) I'(\tau/D). \quad (8)$$

The assumption $\ell/D \leq q_0 < 1/2$ gives $\tau/D \leq 2\sqrt{q_0(1-q_0)} < 1$. On this interval, $\operatorname{arctanh} z \leq C_{q_0} z$, and therefore

$$I'(\tau/D) \leq C_{q_0} \sqrt{\ell/D}.$$

Combining the last two bounds yields $D\Psi(\ell/D) - DI(u/D) \leq C_{\rho,q_0} \sqrt{D} \ell^{-1/6} \sqrt{\ell/D} = C_{\rho,q_0} \ell^{1/3}$. Substituting into the Chernoff bound proves the second assertion. $\square$

3 From moment bounds to stable sampling

The preceding construction shows how a low-degree polynomial can “hide” its energy on rarely sampled inputs. For an upper sampling bound, we need the converse: a level B at which clipping h^2 to $h^2 \wedge B$ preserves a fixed amount of energy for every unit polynomial h . These clipped functions are bounded, so their empirical averages can be controlled uniformly over a known subspace. Since $\mathbb{E}_n[h^2] \geq \mathbb{E}_n[h^2 \wedge B]$, this gives the required lower bound on the empirical covariance without an upper spectral bound. The construction also bounds the necessary clipping level from below. Optimizing a moment bound will give the matching upper estimate.

For unit-norm h , define

$$b_\eta(h) = \inf\{B > 0 : \mathbb{E}[h^2 \wedge B] \geq \eta\}, \quad T_{d,k}(\eta) = \sup_{\substack{h \in \mathcal{P}_{d,k} \\ \mathbb{E}[h^2]=1}} b_\eta(h).$$

3.1 The Kirshner–Samorodnitsky moment bound

For $p \geq 2$ and $0 < q < 1/2$, let $z \in (0, 1)$ be the unique solution of

$$q = \frac{z((1+z)^{p-1} - (1-z)^{p-1})}{(1+z)^p + (1-z)^p}. \quad (9)$$

Define

$$F(p, q) = \log \frac{(1+z)^p + (1-z)^p}{2} - pq \log z - \frac{p}{2} \mathbf{H}(q). \quad (10)$$

At the endpoints, set $F(p, 0) = 0$ and $F(p, 1/2) = (p/2 - 1) \log 2$, the respective limits of the formula.

The moment bound stated below is precisely [KS21, Corollary 1.4]. We verify uniqueness in Section A.2.

Theorem 3.1 (Kirshner–Samorodnitsky [KS21]). *For $p \geq 2$, $0 \leq k \leq d/2$, and nonzero $f \in \mathcal{P}_{d,k}$,*

$$\frac{\mathbb{E}[|f|^p]}{(\mathbb{E}[f^2])^{p/2}} \leq \exp(dF(p, k/d)),$$

with F defined in (10).

3.2 Cancellation at the second moment

The next lemma identifies $\Psi(q)$ as the linear term in the moment bound near $p = 2$, with no quadratic error. The cubic remainder here will be responsible for the $k^{1/3}$ correction in the subsequent clipping bound.

Lemma 3.2 (Cancellation at the second moment). *For every $q_0 < 1/2$, there is $C_0 < \infty$ such that, for $0 < q \leq q_0$ and $0 \leq s \leq 1/2$,*

$$F(2 + 2s, q) \leq s\Psi(q) + C_0 q s^3. \quad (11)$$

Consequently, if $\mathbb{E}[h^2] = 1$ and $\deg h \leq k \leq q_0 d$, then

$$\log \mathbb{E}[|h|^{2+2s}] \leq sE_{d,k} + C_0 k s^3. \quad (12)$$

The proof uses an elementary variational comparison and Taylor’s theorem. Evaluating the variational formula at a carefully chosen function of s gives an upper bound whose second derivative vanishes at $s = 0$, leaving only a cubic remainder. See Appendix A.2 for the details.

3.3 From moments to clipping

The remaining step is the standard moment-to-tail argument, which exploits that a moment slightly above two bounds the energy removed by truncation. We give the parameter choice explicitly because it determines the $k^{1/3}$ correction.

Proposition 3.3 (One-dimensional clipping). *For every fixed $\eta \in (0, 1)$,*

$$|\log T_{d,k}(\eta) - E_{d,k}| \leq C_{\eta, q_0} k^{1/3}.$$

Upper bound. Step 1: bounding the energy removed by clipping. Let $\mathbb{E}[h^2] = 1$, $L_\eta = \log(1/(1-\eta))$, and $0 < s \leq 1/2$. For $z \geq 0$,

$$(z - B)_+ \leq z \mathbf{1}_{\{z > B\}} \leq z \left(\frac{z}{B}\right)^s = z^{1+s} B^{-s}.$$

Using [Lemma 3.2](#),

$$\mathbb{E}[(h^2 - B)_+] \leq B^{-s} \mathbb{E}[|h|^{2+2s}] \leq \exp(-s(\log B - E_{d,k}) + C_0 k s^3).$$

Step 2: balancing the two error terms. To make the last exponent at most $-L_\eta$, it suffices to take $\log B - E_{d,k} = C_0 k s^2 + L_\eta/s$. Balancing these terms suggests s of order $k^{-1/3}$. Choose

$$s = \frac{1}{2} \wedge \left(\frac{L_\eta}{2C_0 k}\right)^{1/3}, \quad \log B = E_{d,k} + C_0 k s^2 + \frac{L_\eta}{s}. \quad (13)$$

Then $\mathbb{E}[(h^2 - B)_+] \leq 1 - \eta$, so $\mathbb{E}[h^2 \wedge B] \geq \eta$. If the second value in the minimum in (13) is selected, the two error terms are each $O_{q_0, \eta}(k^{1/3})$. Otherwise, k is bounded in terms of q_0, η , and the same conclusion holds after increasing the constant. Taking the supremum proves the claim. $\square$

Lower bound. Fix the retained fraction η . For large k , apply [Lemma 2.3](#) with $D = d$, $\ell = k$, and leakage $\rho = \eta/2$. Let $E = \{S_d \geq u\}$ be its small ball. Outside E , the polynomial has at most $\eta/2$ energy; inside E , clipping at B leaves at most B at each input. Hence

$$\mathbb{E}[p^2 \wedge B] \leq \mathbb{E}[p^2 \mathbf{1}_{E^c}] + B\mu_d(E) \leq \eta/2 + B\mu_d(E).$$

Every $B < \eta/(2\mu_d(E))$ therefore retains less than η energy. By the definitions of b_η and $T_{d,k}(\eta)$,

$$T_{d,k}(\eta) \geq b_\eta(p) \geq \frac{\eta}{2\mu_d(E)} \geq \frac{\eta}{2} \exp\{E_{d,k} - C_{\eta, q_0} k^{1/3}\}.$$

For bounded k , the constant polynomial gives $T_{d,k}(\eta) \geq \eta$. Since $E_{d,k} \leq 2k$, increasing C_{η, q_0} ensures $\log T_{d,k}(\eta) \geq E_{d,k} - C_{\eta, q_0} k^{1/3}$ in this case as well. $\square$

3.4 Uniform empirical control after clipping

We now convert the clipping estimate into a uniform lower bound on empirical energy. The argument combines three standard techniques: symmetrization, contraction, and Bousquet's inequality; see [Appendix B](#).

Proposition 3.4 (Clipping gives stable sampling). *Let V have dimension m and an orthonormal basis. Suppose $\mathbb{E}[h^2 \wedge B] \geq \alpha$ for every unit $h \in V$, where $B \geq 1$ and $0 < c < \alpha < 1$. Then*

$$n \geq C_{\alpha, c} B(m + t) \implies \mathbb{P}(\widehat{\Sigma}_V \succeq cI_m) \geq 1 - e^{-t}.$$

Proof. Let $\psi_B(u) = u^2 \wedge B$, and let $\mathcal{S} = \{h \in V : \mathbb{E}[h^2] = 1\}$. This function is $2\sqrt{B}$ -Lipschitz and vanishes at zero. Define

$$D_B = \sup_{h \in \mathcal{S}} (\mathbb{E}[\psi_B(h)] - \mathbb{E}_n[\psi_B(h)]), \quad \overline{D}_B = \max\{D_B, 0\}.$$

Since $\overline{D}_B$ is bounded by the corresponding absolute supremum, [Theorem B.4\(i\)](#) (symmetrization), followed by part (ii) (contraction) conditionally on the inputs with $\varphi = \psi_B$ and $L = 2\sqrt{B}$, gives

$$\mathbb{E}\overline{D}_B \leq 8\sqrt{B} \mathbb{E} \sup_{h \in \mathcal{S}} \left| \frac{1}{n} \sum_i \varepsilon_i h(X_i) \right| = 8\sqrt{B} \mathbb{E} \left\| \frac{1}{n} \sum_i \varepsilon_i \Phi_V(X_i) \right\|_2 \leq 8\sqrt{\frac{Bm}{n}}.$$

The last step uses Jensen's inequality and $\mathbb{E} \left\| \frac{1}{n} \sum_i \varepsilon_i \Phi_V(X_i) \right\|_2^2 = \frac{1}{n} \mathbb{E} \|\Phi_V(X)\|_2^2 = \frac{m}{n}$. For a unit h , the centered function

$$g_h = \frac{\mathbb{E}[\psi_B(h)] - \psi_B(h)}{B}$$

has absolute value at most one and variance at most $1/B$ since $\mathbb{E}[\psi_B(h)^2] \leq B \mathbb{E}[\psi_B(h)] \leq B \mathbb{E}[h^2] = B$.

Apply [Theorem B.3](#) (Bousquet's inequality) to these functions and the zero function. Its supremum is $Z = n\overline{D}_B/B$. After rescaling, with probability at least $1 - e^{-t}$,

$$\overline{D}_B \leq \mathbb{E}\overline{D}_B + \sqrt{\frac{2Bt}{n}(1 + 2\mathbb{E}\overline{D}_B)} + \frac{Bt}{3n}.$$

When $n \geq CB(m+t)$ with C large, the expectation is at most one. The right side is then bounded by

$$C' \left(\sqrt{\frac{B(m+t)}{n}} + \frac{B(m+t)}{n} \right).$$

Choose the sample-size constant so this is at most $\alpha - c$. On that event, for every $h \in \mathcal{S}$,

$$\mathbb{E}_n[h^2] \geq \mathbb{E}_n[\psi_B(h)] \geq \mathbb{E}[\psi_B(h)] - \overline{D}_B \geq \alpha - (\alpha - c) = c.$$

Finally, homogeneity gives the claim for all $h \in V$. □

4 Many directions on one Hamming ball

The one-dimensional construction gives one polynomial concentrated on a small Hamming ball. We now seek a whole subspace for which every function concentrates on the same ball.

4.1 Separating radial concentration from variation within slices

A radial polynomial $p(S_d)$ is constant on each slice $\{x : S_d(x) = s\}$. The Krawchouk basis lets us choose its values across these slices so that its energy concentrates at large s . To obtain more than one direction, we multiply a shared radial factor by polynomials that can distinguish points within a slice:

$$h_f(x) = f(x)p_r(S_d(x)), \quad f \in \mathcal{H}_{d,r}.$$

Here $\mathcal{H}_{d,r}$ is the space of homogeneous multilinear degree- r *harmonic polynomials*, defined by the polynomial identity

$$\sum_{i=1}^d \partial_i f = 0.$$

For example, $\mathcal{H}_{d,0}$ consists of constants, while $\mathcal{H}_{d,1} = \{\sum_i v_i x_i : \sum_i v_i = 0\}$.

We choose $\deg p_r \leq k - r$, so each product has Boolean degree at most k . The available number of harmonic directions is

$$\dim \mathcal{H}_{d,r} = \binom{d}{r} - \binom{d}{r-1}, \quad \binom{d}{-1} = 0,$$

see [\[FM19, Corollary 3.9\(b\)\]](#).

4.2 Uniform energy control across harmonic directions

All unit polynomials in $\mathcal{H}_{d,r}$ have the same distribution of squared energy across slices. A shared radial factor therefore changes this distribution identically in every direction. Moreover, different harmonic degrees remain orthogonal under radial weighting. These facts follow from the identity below, a direct specialization of [FM19, Theorem 3.10]. See Appendix C for the proof.

Lemma 4.1 (Radial-weight identity). *For integers $0 \leq r, s \leq \lfloor d/2 \rfloor$, $f \in \mathcal{H}_{d,r}$, $g \in \mathcal{H}_{d,s}$, and any real function H of the coordinate sum,*

$$\mathbb{E}_{\mu_d}[fgH(S_d)] = \begin{cases} 0, & r \neq s, \\ \langle f, g \rangle_d \mathbb{E}_{\mu_{d-2r}}[H(S_{d-2r})], & r = s. \end{cases} \quad (14)$$

Here $\langle f, g \rangle_d = \mathbb{E}_{\mu_d}[fg]$ and $S_0 = 0$.

To see why $d - 2r$ appears, consider the unit harmonic polynomial

$$Q_r(x) = \prod_{j=1}^r \frac{x_{2j-1} - x_{2j}}{\sqrt{2}}.$$

Taking $f = g = Q_r$ introduces the weight Q_r^2 in the uniform expectation. This weight vanishes unless $x_{2j-1} = -x_{2j}$ for every $j \leq r$. Each pair contributes zero to S_d , leaving the sum of $d - 2r$ independent signs. The lemma says that the same law of the coordinate sum results from weighting by $f^2/\|f\|_2^2$ for every nonzero $f \in \mathcal{H}_{d,r}$.

In particular, choose p_r with unit norm under S_{d-2r} and energy at most ρ below a cutoff u . Apply the identity with $H(s) = p_r(s)^2$ and with $H(s) = p_r(s)^2 \mathbf{1}_{\{s < u\}}$. For every nonzero $h_f = fp_r(S_d)$, this gives

$$\|h_f\|_2^2 = \|f\|_2^2, \quad \frac{\mathbb{E}[h_f^2 \mathbf{1}_{\{S_d < u\}}]}{\|h_f\|_2^2} = \mathbb{E}[p_r(S_{d-2r})^2 \mathbf{1}_{\{S_{d-2r} < u\}}] \leq \rho.$$

Thus multiplication by the radial factor preserves dimension and transfers the one-dimensional concentration bound to the entire space $W_r = \{fp_r(S_d) : f \in \mathcal{H}_{d,r}\}$, with no dimension factor in the leakage. Different harmonic degrees also remain orthogonal after radial weighting, including restriction to a Hamming ball. This lets us combine the spaces W_r and their corresponding concentration bounds once we choose a cutoff valid for all of them. We call this a *harmonic lift*: a single one-dimensional witness p_r is turned into a whole space of witnesses by the map $f \mapsto fp_r(S_d)$ on $\mathcal{H}_{d,r}$.

Combining harmonic degrees $0 \leq r \leq R$ supplies $\binom{d}{R}$ concentrated directions, but leaves only $k - R$ degrees for the radial factor in sector r . Its effective coordinate sum has $d - 2R$ terms, so the one-dimensional construction has center $2\sqrt{(k - R)(d - k - R)}$. This center decreases with R . The common cutoff must therefore lie below the smallest center, attained at R . We define that center and its binomial tail exponent as

$$\tau_R = 2\sqrt{(k - R)(d - k - R)}, \quad \mathcal{E}_{d,k}(R) = dI(\tau_R/d), \quad 0 \leq R \leq k.$$

Thus $\mathcal{E}_{d,k}(R)$ determines the leading exponential decay of the ball's probability after reserving degree R for the harmonic factors, before adjusting the cutoff to ensure energy concentration. At $R = 0$ it equals $E_{d,k}$, and increasing R trades a larger subspace for a smaller concentration exponent. Explicitly parametrizing this tradeoff will allow for the subsequent optimized choice $R = \lfloor k^{1/3} \rfloor$.

Proposition 4.2 (Harmonic lift). *Fix $\rho \in (0, 1)$ and $q_0 \in (0, 1/2)$. For integers d, k, R with $1 \leq k \leq q_0 d$ and $0 \leq R \leq k$, there are $W_{\leq R} \subseteq \mathcal{P}_{d,k}$ of dimension $\binom{d}{R}$ and a permutation-invariant E such that, with $K = k - R$,*

$$\mu_d(E) \leq e^{-\mathcal{E}_{d,k}(R) + C_{\rho,q_0} K^{1/3}}, \quad \mathbb{E}[h^2 \mathbf{1}_{E^c}] \leq \rho \mathbb{E}[h^2] \quad (h \in W_{\leq R}).$$

Consequently,

$$\mu_d(E) \leq e^{-E_{d,k} + C_{\rho,q_0}(R + k^{1/3})}. \quad (15)$$

Proof. Write $M = M_{\rho,q_0}$ from Lemma 2.3. Fix an integer $K_0 = K_0(\rho, q_0) \geq 1$ above that lemma's degree threshold and large enough that $M K_0^{-2/3} < 2\sqrt{1 - 2q_0}$.

Case (i): $K \geq K_0$.

Step 1: choosing a common ball. For $0 \leq r \leq R$, put $D_r = d - 2r$ and $K_r = k - r$. Since $K_r \geq K_0$ and $K_r/D_r \leq k/d \leq q_0$, Lemma 2.3 provides a polynomial p_r of degree K_r , with unit norm under S_{D_r} and leakage at most ρ below

$$u_r = 2\sqrt{(k-r)(d-k-r)} - M\sqrt{D_r} K_r^{-1/6}.$$

Set $u = \tau_R - M\sqrt{d} K^{-1/6}$ and $E = \{S_d \geq u\}$. The centers decrease with r , while $\sqrt{D_r} K_r^{-1/6} \leq \sqrt{d} K^{-1/6}$, so $u \leq u_r$ for all r . Moreover, $\tau_R \geq 2\sqrt{(1 - 2q_0)dK}$; our choice of K_0 therefore ensures $0 < u < \tau_R < d$.

Step 2: combining directions without increasing leakage. Define $W_r = \{f p_r(S_d) : f \in \mathcal{H}_{d,r}\}$, whose elements have degree at most $r + K_r = k$. By Lemma 4.1, multiplication by $p_r(S_d)$ preserves norms, and

$$\mathbb{E}[h_r^2 \mathbf{1}_{E^c}] \leq \rho \|h_r\|_2^2 \quad (h_r \in W_r).$$

To extend this bound to $h = \sum_r h_r$, we must also control the cross terms in both $\|h\|_2^2$ and $\mathbb{E}[h^2 \mathbf{1}_{E^c}]$. For $r \neq s$, apply Lemma 4.1 with the weights $H(t) = p_r(t)p_s(t)$ and $H(t) = p_r(t)p_s(t)\mathbf{1}_{\{t < u\}}$, respectively. This yields

$$\mathbb{E}[h_r h_s] = \mathbb{E}[h_r h_s \mathbf{1}_{E^c}] = 0.$$

Hence $W_{\leq R} = \bigoplus_{r=0}^R W_r$ has dimension $\sum_{r=0}^R [\binom{d}{r} - \binom{d}{r-1}] = \binom{d}{R}$, and

$$\mathbb{E}[h^2 \mathbf{1}_{E^c}] = \sum_r \mathbb{E}[h_r^2 \mathbf{1}_{E^c}] \leq \rho \sum_r \|h_r\|_2^2 = \rho \|h\|_2^2.$$

Step 3: bounding the probability of the ball. Use the integral estimate in (8), now with $D = d$ and center τ_R . Since $\tau_R/d \leq 2\sqrt{q_0(1 - q_0)} < 1$ and $I'(\tau_R/d) \leq C_{q_0} \sqrt{K/d}$, it gives

$$\mathcal{E}_{d,k}(R) - dI(u/d) \leq (\tau_R - u)I'(\tau_R/d) \leq C_{\rho,q_0} \sqrt{d} K^{-1/6} \sqrt{K/d} = C_{\rho,q_0} K^{1/3}.$$

The binomial Chernoff bound $\mu_d(E) \leq e^{-dI(u/d)}$ now proves the asserted probability bound.

Case (ii): $0 \leq K < K_0$. Take E to be the whole cube and $W_{\leq R}$ to be the span of all degree- R Walsh characters. This space has dimension $\binom{d}{R}$, lies in $\mathcal{P}_{d,k}$, and has zero leakage. To check the probability bound, note that $\tau_R \leq 2\sqrt{K(d - K)}$ and $\Psi(q) \leq 2q$ give $0 \leq \mathcal{E}_{d,k}(R) \leq d\Psi(K/d) \leq 2K$. For $1 \leq K < K_0$, choosing $C_{\rho,q_0} \geq 2K_0^{2/3}$ makes $-\mathcal{E}_{d,k}(R) + C_{\rho,q_0} K^{1/3} \geq 0$, so the claimed upper bound is at least $\mu_d(E) = 1$. For $K = 0$, both exponents are zero.

Finally, for $z_R = \tau_R/d$ and by continuity at $R = k$,

$$\left| \frac{d}{dR} \mathcal{E}_{d,k}(R) \right| = 2(1 - 2R/d) \frac{\operatorname{arctanh} z_R}{z_R} \leq C_{q_0}.$$

Integrating gives $\mathcal{E}_{d,k}(R) \geq E_{d,k} - C_{q_0} R$, which proves (15). $\square$

4.3 Proof of Theorem 1.8

Proof of Theorem 1.8. Step 1: the universal lower bound on $\mu_d(E)$. Let $h \neq 0$ satisfy $\mathbb{E}[h^2 \mathbf{1}_{E^c}] \leq \rho \|h\|_2^2$ and rescale it to $\|h\|_2 = 1$. Set $\eta = (1 + \rho)/2 > \rho$. By Proposition 3.3, clipping at $B = \exp\{E_{d,k} + C_{\eta,q_0} k^{1/3}\}$ retains at least η energy. On E^c the clipped energy is at most ρ , while on E the integrand is at most B . Hence

$$\eta \leq \mathbb{E}[h^2 \wedge B] \leq \mathbb{E}[h^2 \mathbf{1}_{E^c}] + B\mu_d(E) \leq \rho + B\mu_d(E).$$

It follows that $\mu_d(E) \geq (1 - \rho)/(2B)$, giving the asserted bound after absorbing the fixed prefactor into C_{ρ,q_0} .

Step 2: a subspace attaining the bound up to the remainder. Apply Proposition 4.2 with $R = \lfloor k^{1/3} \rfloor$. It supplies a space of dimension $\binom{d}{R} = M_{d,k}$ and one event E with leakage at most ρ for every function in that space. Moreover, (15) gives

$$\mu_d(E) \leq \exp\{-E_{d,k} + C_{\rho,q_0}(R + k^{1/3})\} \leq \exp\{-E_{d,k} + C'_{\rho,q_0} k^{1/3}\}.$$

□

Remark 4.3 (Rare event is a Hamming ball). *For sufficiently large k , the choice $R = \lfloor k^{1/3} \rfloor$ gives $K = k - R \geq K_0(\rho, q_0)$. We therefore land in case (i) of the proof of Proposition 4.2, which constructs $E = \{S_d \geq u\}$ with $u > 0$, rather than the whole-cube fallback of case (ii). Since $S_d(x) = d - 2 \text{dist}_H(x, (1, \dots, 1))$, this event is a Hamming ball, establishing the final assertion of Theorem 1.8.*

5 From concentration to minimax regression

We now convert the geometric bounds into regression guarantees. These reductions apply to any isotropic feature distribution.

5.1 From shared concentration to small empirical eigenvalues

To lower-bound the sample size required for the parametric regression rate, we use a subspace whose functions concentrate on a common set E . In an orthonormal basis, this gives

$$\mathbb{E}[\Phi(X)\Phi(X)^\top \mathbf{1}_{\{X \notin E\}}] \preceq \rho I_m, \quad \mathbb{E}[\Phi(X)\Phi(X)^\top] = I_m. \quad (16)$$

When $n\mu_d(E)$ is small relative to m , few observations fall inside E , while those outside contribute little total energy. Together, these properties force many eigenvalues of the empirical covariance matrix to be small.

Lemma 5.1 (Many small empirical eigenvalues). *Let $\mathbb{P}(E) = p \in (0, 1/2]$. (i) If (16) holds, $m \geq 2$, and $n \leq m/(128p)$, then with probability at least $15/16$, at least $\lfloor m/2 \rfloor$ eigenvalues of $\hat{\Sigma}$ are at most 128ρ . (ii) If $t \geq \log 4$, $n \leq t/(512p)$, and a fixed unit vector v satisfies $\mathbb{E}[(v^\top \Phi)^2 \mathbf{1}_{E^c}] \leq \rho$, then with probability at least $(15/16)e^{-t/256}$ the smallest eigenvalue of $\hat{\Sigma}$ is at most 32ρ .*

Proof. Split the empirical covariance according to where each observation falls:

$$\hat{\Sigma} = \hat{\Sigma}_E + \hat{\Sigma}_{E^c}, \quad \hat{\Sigma}_A = \frac{1}{n} \sum_{i=1}^n \Phi(X_i)\Phi(X_i)^\top \mathbf{1}_{\{X_i \in A\}}, \quad A \in \{E, E^c\}.$$

Both matrices are positive semidefinite.

Step 1: many small eigenvalues (part (i)). Let $N_E = \sum_i \mathbf{1}_{\{X_i \in E\}}$. Then $\text{rank}(\hat{\Sigma}_E) \leq N_E$, $\mathbb{E}[N_E] = np \leq m/128$, and $\mathbb{E}[\text{tr} \hat{\Sigma}_{E^c}] \leq m\rho$ by (16). Markov's inequality and a union bound show that, with probability at least $15/16$, both $N_E \leq m/4$ and $\text{tr} \hat{\Sigma}_{E^c} \leq 32m\rho$ hold. We condition on this event and set $L = \ker \hat{\Sigma}_E$, so that $\dim L \geq m - \lfloor m/4 \rfloor$. Let P_L be orthogonal projection onto L . The operator

$$B_L = (P_L \hat{\Sigma}_{E^c} P_L)|_L : L \rightarrow L$$

describes the *compression* to L : it has the same quadratic form as $\hat{\Sigma}_{E^c}$ on vectors in L , and $\text{tr} B_L \leq \text{tr} \hat{\Sigma}_{E^c} \leq 32m\rho$. Since its eigenvalues are nonnegative, at most $\lfloor m/4 \rfloor$ can exceed 128ρ . Let $U \subseteq L$ be the span of the eigenvectors of B_L with eigenvalues at most 128ρ . Then $\dim U \geq \dim L - \lfloor m/4 \rfloor \geq \lfloor m/2 \rfloor$.

For $0 \neq w \in U$,

$$\mathcal{R}_{\hat{\Sigma}}(w) = \frac{w^\top \hat{\Sigma}_{E^c} w}{\|w\|_2^2} = \frac{w^\top B_L w}{\|w\|_2^2} \leq 128\rho,$$

because $\hat{\Sigma}_E w = 0$. The min-max principle now gives the claimed number of small eigenvalues of $\hat{\Sigma}$.

Step 2: one small eigenvalue (part (ii)). Fix a unit vector v satisfying the assumption in (ii) and condition on $N_E = 0$. Independence gives

$$\Pr(N_E = 0) = (1-p)^n \geq e^{-2np} \geq e^{-t/256}, \quad \mathbb{E}[v^\top \hat{\Sigma} v \mid N_E = 0] = \frac{\mathbb{E}[(v^\top \Phi)^2 \mathbf{1}_{E^c}]}{1-p} \leq 2\rho.$$

Markov's inequality gives $v^\top \hat{\Sigma} v \leq 32\rho$ with conditional probability at least $15/16$. Since $\lambda_{\min}(\hat{\Sigma}) \leq \mathcal{R}_{\hat{\Sigma}}(v) = v^\top \hat{\Sigma} v$, the unconditional probability is at least $(15/16)e^{-t/256}$. $\square$

5.2 From empirical eigenvalues to minimax risk

For the design matrix $\mathbf{X}$ with rows $\Phi_V(X_i)^\top$, a full-rank sample gives

$$(\hat{\theta}_{\text{OLS}} - \theta) \mid \mathbf{X} \sim N\left(0, \frac{\sigma^2}{n} \hat{\Sigma}_V^{-1}\right).$$

Thus an empirical eigenvalue λ produces variance $\sigma^2/(n\lambda)$ in a unit population direction. The following identity shows that the corresponding quantile is minimax over all estimators and is a direct corollary of the results in [EHE24].

Theorem 5.2 (El Hanchi–Maddison–Erdogdu). *For a fixed known subspace V in (1), let $G = (G_1, \dots, G_m) \sim N(0, I_m)$ be independent of the sampled inputs, and define*

$$Z_{n,V} = \begin{cases} (\sigma^2/n) G^\top \hat{\Sigma}_V^{-1} G, & \hat{\Sigma}_V \text{ invertible,} \\ +\infty, & \hat{\Sigma}_V \text{ singular.} \end{cases}$$

Then $\mathcal{R}_{n,\delta}^(V) = Q_{1-\delta}(Z_{n,V})$, and any measurable least-squares minimizer attains this minimax value. Moreover,*

$$\mathcal{R}_{n,\delta}^*(V) = +\infty \iff \Pr(\hat{\Sigma}_V \text{ singular}) \geq \delta.$$

Conditionally on any nonsingular design with at least r eigenvalues at most $\lambda > 0$, the theorem's comparison variable satisfies

$$Z_{n,V} \succeq_{\text{st}} \frac{\sigma^2}{n\lambda} \chi_r^2.$$

The comparison also holds on singular designs. If $t < m$, the many eigenvalues in [Lemma 5.1\(i\)](#) provide the factor m . If $t \geq m$, part (ii) and a Gaussian tail provide the factor t . Choosing ρ sufficiently small depending only on A , the benchmark is unattainable whenever

$$n \leq \frac{m+t}{1024p}. \quad (17)$$

[Appendix D](#) applies the Gaussian tail estimates from [Lemma B.5](#) to give the complete probability calculation, including the cases of bounded m, t . It also proves the stable-sampling lower bound, which uses only [Lemma 5.1](#), with ρ chosen depending on c .

5.3 Proof of Theorem 1.5

Proof of Theorem 1.5. Write $N = N_{\text{par}}(d, k, m, A, \delta)$. For the upper bound, use [Proposition 3.3](#) with a fixed retained fraction greater than $1/2$, then [Proposition 3.4](#) at covariance level $1/2$ and failure probability $\delta/2$. Conditional on this event, the error of ordinary least squares is bounded by $(2\sigma^2/n)\chi_m^2$. [Corollary B.6](#) gives the benchmark with constant $12 \leq A$. This proves $N_{\text{par}} \leq C(m+t)e^{E_{d,k}+Ck^{1/3}}$ for every feasible m .

For the lower bound choose the leakage required in [Proposition D.3](#). For large k , [Theorem 1.8](#) gives a common set with $0 < p \leq e^{-E_{d,k}+Ck^{1/3}} \leq 1/2$. If $m \leq M_{d,k}$, take an m -dimensional subspace of the concentrating space. If $t \geq m$, extend one concentrated unit vector to any feasible m -dimensional space. The common-set or one-direction version of (17) gives $N \geq (m+t)/(1024p)$: failure at every positive integer up to this quantity forces the eventual threshold above its integer part, otherwise, if the quantity is below one, use $N \geq 1$. For bounded k , the baseline $N \geq c_0(m+t)$ in [Lemma D.1](#), together with $E_{d,k} \leq 2k$, suffices. Taking logarithms absorbs fixed factors into $Ck^{1/3}$.

For stable sampling the upper proof uses retained fraction $(1+c)/2$ and [Proposition 3.4](#). The lower bound is proved by [Proposition D.2](#). $\square$

6 Sharpness of the uniform uncertainty remainder

We prove [Proposition 1.9](#) by constructing a polynomial with a fixed fraction $\beta > 0$ of its squared L_2 -norm concentrated on the chosen set, a sequence of dimensions $d_k \geq k^3$, and spaces of dimension $\binom{d_k}{\lfloor k^{1/3} \rfloor}$ concentrated on events of probability at most $e^{-E_{d_k,k}-2k^{1/3}}$. In other words, here the leakage is $\rho_* = 1 - \beta$, and only a small fixed fraction of energy needs to be retained. This differs from the construction of [Section 2](#), which allows any prescribed leakage.

The proof has three stages. We first construct a polynomial of degree at most N whose energy under the standard Gaussian measure has a fixed positive fraction concentrated on $\{x > 2\sqrt{N}\}$. For each sufficiently large degree k , we transfer that polynomial to a sufficiently large Boolean cube and apply the harmonic lift. Finally, a tail bound shows that the common event has the required probability.

6.1 Convergence and limits

Two different limits underlie the construction. We describe them below.

From the cube to Gaussian polynomials: fix the degree, let the ambient dimension grow. Write the

normalized radial Krawchouk polynomial as

$$e_{D,j}(x) = \binom{D}{j}^{-1/2} \sum_{|A|=j} \prod_{i \in A} x_i = P_{D,j}(S_D(x)/\sqrt{D}).$$

For every fixed j , the coefficients of the univariate polynomial $P_{D,j}$ converge as $D \rightarrow \infty$ to those of $H_j(t) = \text{He}_j(t)/\sqrt{j!}$, the degree- j orthonormal Hermite polynomial for the standard Gaussian distribution. In particular, $P_{D,j}(t) \rightarrow H_j(t)$ uniformly on bounded intervals of t . This explains why Gaussian polynomials are useful models for radial polynomials on large cubes. Our transfer below only needs the central limit theorem and moment convergence for a fixed polynomial.

From Hermite polynomials to the Airy kernel as the degree cutoff increases. We combine Hermite degrees $0, \dots, N-1$ into the kernel K_N defined below. The external asymptotic theorem we use evaluates this kernel at points

$$x_N(s) = 2\sqrt{N} + sN^{-1/6},$$

where s remains in a fixed bounded interval as $N \rightarrow \infty$. After multiplying the kernel by $N^{-1/6}$, its limit is the *Airy kernel* in (18). Here it describes the limiting behavior of the Hermite kernel near the edge of its concentration region.

6.2 A Gaussian polynomial with a fixed amount of tail energy

Step 1: use the Hermite kernel to combine degrees. Let $\gamma = N(0,1)$ have density $\varphi(x) = (2\pi)^{-1/2}e^{-x^2/2}$, and define

$$H_j(x) = \frac{\text{He}_j(x)}{\sqrt{j!}}, \quad \text{He}_j(x) = (-1)^j e^{x^2/2} \frac{d^j}{dx^j} e^{-x^2/2}.$$

The H_j are orthonormal in $L_2(\gamma)$. Set $\psi_j = H_j\sqrt{\varphi}$ and define the Hermite kernel

$$K_N(x, y) = \sum_{j=0}^{N-1} \psi_j(x)\psi_j(y).$$

The Airy function is defined for real s by the oscillatory integral

$$\text{Ai}(s) = \frac{1}{\pi} \lim_{R \rightarrow \infty} \int_0^R \cos\left(\frac{u^3}{3} + su\right) du.$$

The associated Airy kernel is

$$K_{\text{Ai}}(s, t) = \int_0^\infty \text{Ai}(s+u) \text{Ai}(t+u) du.$$

To describe the Hermite kernel near $2\sqrt{N}$, put $x_N(s) = 2\sqrt{N} + sN^{-1/6}$. Then

$$N^{-1/6} K_N(x_N(s), x_N(t)) \longrightarrow K_{\text{Ai}}(s, t) \tag{18}$$

as $N \rightarrow \infty$, uniformly for (s, t) in compact subsets of $\mathbb{R}^2$. This convergence follows from [SX22, Theorem 2.3], which in turn restates the result of [DG07].

We center a kernel vector at $y_N = x_N(3)$, slightly beyond $2\sqrt{N}$. This is the standard reproducing-kernel construction: the coefficients are the basis values at the point where we want concentration. For $K = N - 1$, define

$$q_K(x) = \frac{\sum_{j=0}^K H_j(x) H_j(y_N)}{(\sum_{j=0}^K H_j(y_N)^2)^{1/2}}.$$

Orthonormality gives $\deg q_K \leq K$ and $\int q_K^2 d\gamma = 1$. Moreover,

$$q_K(x)^2 \varphi(x) = \frac{K_N(x, y_N)^2}{K_N(y_N, y_N)}.$$

Step 2: retain positive energy on a fixed scaled interval. We estimate the energy on the interval $[2\sqrt{N} + 3N^{-1/6}, 2\sqrt{N} + 4N^{-1/6}]$. Under the change of variables $x = 2\sqrt{N} + sN^{-1/6}$, this becomes $s \in [3, 4]$, where the uniform convergence in (18) lets us pass to the limit inside the integral. The left endpoint is chosen far enough above $2\sqrt{N}$ to accommodate the degree reserved for the harmonic lift below. Any fixed right endpoint larger than 3 would work. A positive lower bound on the energy in this interval suffices, since the interval lies within the tail event we will use.

To track the scaling, write $\widetilde{K}_N(s, t) = N^{-1/6} K_N(x_N(s), x_N(t))$. Since $dx = N^{-1/6} ds$, the kernel factors cancel and yield

$$\int_{x_N(3)}^{x_N(4)} q_{N-1}(x)^2 d\gamma(x) = \frac{\int_3^4 \widetilde{K}_N(s, 3)^2 ds}{\widetilde{K}_N(3, 3)} \xrightarrow{N \rightarrow \infty} \alpha := \frac{\int_3^4 K_{\text{Ai}}(s, 3)^2 ds}{K_{\text{Ai}}(3, 3)} > 0. \quad (19)$$

Indeed, $\text{Ai}(s) > 0$ for $s \geq 0$, so the denominator and integrand in the limit are positive. Also $\alpha \leq 1$, since each q_{N-1} has unit norm. Fix $\beta = \alpha/8 > 0$, independently of all subsequent degrees and dimensions. By convergence to α , there is N_0 such that for every $N \geq N_0$ the left side of (19) is at least $\alpha/2 = 4\beta$. The factor four leaves room for approximation errors in passing to the cube and comparing norms.

6.3 Transferring the construction to a Boolean subspace

Proof of Proposition 1.9. Step 1: reserving enough degree for the harmonic directions. For each sufficiently large integer k , set

$$R = \lfloor k^{1/3} \rfloor, \quad K = k - R, \quad N = K + 1, \quad a_k = 2\sqrt{k} + k^{-1/6}.$$

As $k \rightarrow \infty$, $N = k - \lfloor k^{1/3} \rfloor + 1 \rightarrow \infty$, so eventually $N \geq N_0$. We need the interval used above to lie beyond a_k , which is measured relative to the larger degree k . For such k , $N \leq k$ and

$$2(\sqrt{k} - \sqrt{N}) = \frac{2(R - 1)}{\sqrt{k} + \sqrt{N}} \leq 2k^{-1/6}, \quad 3N^{-1/6} \geq 3k^{-1/6}.$$

Hence $x_N(3) \geq a_k$, and (19) implies that $\int q_K^2 \mathbf{1}_{[a_k, \infty)} d\gamma \geq 4\beta$.

Step 2: transfer to the cube with k fixed. Fix one such k , and hence fix R, K, N, a_k and the polynomial q_K . We now let the ambient dimension d tend to infinity. For each $r \in \{0, \dots, R\}$, the radial-weight identity (Lemma 4.1) expresses energy in harmonic degree r as an expectation over a cube with $d - 2r$ coordinates. Accordingly, define the total energy and the energy above a_k by

$$c_{r,d} = \mathbb{E}_{\mu_{d-2r}}[q_K(S_{d-2r}/\sqrt{d})^2], \quad b_{r,d} = \mathbb{E}_{\mu_{d-2r}}[q_K(S_{d-2r}/\sqrt{d})^2 \mathbf{1}_{\{S_{d-2r}/\sqrt{d} \geq a_k\}}].$$

For each fixed r , the central limit theorem gives $S_{d-2r}/\sqrt{d} \rightarrow G$ in distribution, where $G \sim N(0, 1)$. To obtain convergence of the expectations above, we must also control the growth of q_K^2 . Since q_K has fixed degree K and

$$\sup_{d > 2r} \mathbb{E}[|S_{d-2r}/\sqrt{d}|^{2K+2}] < \infty,$$

the contribution to either expectation from $|S_{d-2r}/\sqrt{d}| > M$ tends to zero uniformly in d as $M \rightarrow \infty$. Together with $\mathbb{P}(G = a_k) = 0$, this gives

$$c_{r,d} \xrightarrow{d \rightarrow \infty} \mathbb{E}[q_K(G)^2] = 1, \quad b_{r,d} \xrightarrow{d \rightarrow \infty} \mathbb{E}[q_K(G)^2 \mathbf{1}_{\{G \geq a_k\}}] \geq 4\beta.$$

These limits hold with k fixed; no uniform convergence as k grows is needed. Since there are only $R + 1$ values of r , we may choose $d = d_k \geq k^3$ sufficiently large that

$$\frac{1}{2} \leq c_{r,d} \leq 2, \quad b_{r,d} \geq 2\beta \quad \text{for every } r \in \{0, \dots, R\}.$$

Step 3: lift the one-dimensional witness to many directions. On the chosen cube, let $E = \{S_d/\sqrt{d} \geq a_k\}$ and

$$W_r = q_K(S_d/\sqrt{d}) \mathcal{H}_{d,r}, \quad W = \bigoplus_{r=0}^R W_r.$$

The degree of each product is at most $K + r \leq k$. By [Lemma 4.1](#), multiplication on $\mathcal{H}_{d,r}$ changes the squared norm by the positive factor $c_{r,d}$, so it is injective. For a nonzero $h_r = f q_K(S_d/\sqrt{d}) \in W_r$, the same identity gives

$$\frac{\mathbb{E}[h_r^2 \mathbf{1}_E]}{\mathbb{E}[h_r^2]} = \frac{b_{r,d} \|f\|_2^2}{c_{r,d} \|f\|_2^2} \geq \frac{2\beta}{2} = \beta.$$

Different W_r are orthogonal both globally and on E , by the same radial-weight identity. Thus

$$\dim W = \sum_{r=0}^R \left(\binom{d}{r} - \binom{d}{r-1} \right) = \binom{d}{R}, \quad \mathbb{E}[h^2 \mathbf{1}_E] \geq \beta \mathbb{E}[h^2] \quad (h \in W). \quad (20)$$

This proves the common-event assertion for every linear combination.

Step 4: bound the common event's probability. The standard Chernoff argument uses $\mathbb{E}[e^{\lambda S_d/\sqrt{d}}] = \cosh(\lambda/\sqrt{d})^d \leq e^{\lambda^2/2}$. Applying Markov's inequality to $e^{\lambda S_d/\sqrt{d}}$ with $\lambda = a_k$ gives

$$\mu_d(E) \leq e^{-a_k^2/2} = \exp\left(-2k - 2k^{1/3} - \frac{1}{2}k^{-1/3}\right) \leq e^{-E_{d,k} - 2k^{1/3}},$$

using $\Psi(q) \leq 2q$ from [Lemma A.1](#). The cross term in $a_k^2/2 = (2\sqrt{k} + k^{-1/6})^2/2$ is $2k^{1/3}$: this is why moving the cutoff above $2\sqrt{k}$ yields the required correction. Take $\rho_* = 1 - \beta$. Restricting W to any m -dimensional subspace in (20) proves the lower inequality in (5). The upper inequality is (4), applied, for example, with $q_0 = 1/4$. Since $d_k \geq k^3$, this range holds eventually. $\square$

For each sufficiently large k , the transfer works once d is large enough; choose one such dimension $d_k \geq k^3$. The argument does not give an explicit upper bound on d_k .

This proves that a positive correction of order $k^{1/3}$ is necessary at one fixed leakage level, for all subspace dimensions in the stated range. The smooth-window construction remains essential: it works for every $k \leq q_0 d$ and allows any prescribed leakage $\rho > 0$. By contrast, the Airy construction guarantees only that a fixed fraction $\beta > 0$ of the energy lies in E . The remaining fraction $1 - \beta$ need not be small enough to establish the regression lower bound for a prescribed benchmark constant $A \geq 16$.

7 Discussion

Our subspace uncertainty principle determines the worst-subspace parametric threshold, throughout the matching range, to an $O(k^{1/3})$ error in the exponent. It explains the exponential cost of noise relative to identification. The uncertainty remainder is optimal in general.

Section F derives the finite-dimensional correction, the worst-case threshold over ambient dimensions, and lower bounds for larger model dimensions, including the leading exponent when $\log m = o(d)$ and $k/d \rightarrow q \in (0, 1/2)$.

Limitations and open questions. At fixed confidence, determining the threshold beyond $m \leq M_{d,k}$ remains open, particularly when $\log m$ is comparable to d . How does increasing the model dimension change the exponent $E_{d,k}$? A necessary constraint is $\mu_d(E) \geq (1 - \rho)m / \dim \mathcal{P}_{d,k}$ whenever an m -dimensional subspace concentrates on E with leakage ρ .

Our sharpness result concerns one fixed leakage level and sufficiently large ambient dimensions. Optimality of the remainder for every prescribed leakage or regression benchmark, and uniform estimates as $k/d \uparrow 1/2$, remain open. Another question is to identify geometric properties of a given subspace V that predict its sample threshold before sampling. Such criteria should distinguish models requiring exponentially many samples in k from models, such as spans of Walsh characters, that attain the parametric rate after $O(m(\log m + t))$ samples, see Section F.2.

AI methodologies

During the preparation of this paper, the author used OpenAI’s ChatGPT 5.6 Sol and 6.0 Astra for the tasks described below. Through an extended exchange with ChatGPT 6.0 Astra, the author explored a matching converse establishing uniform sharpness of the uncertainty remainder. ChatGPT eventually suggested the Airy approach presented in Section 6, whose proof is substantially based on the argument it supplied. ChatGPT 5.6 Sol assisted with proofreading and minor copy-editing. The author independently re-derived and verified the proofs in Section 6 and takes full responsibility for the paper’s content.

References

- [ASW15] Emmanuel Abbe, Amir Shpilka, and Avi Wigderson. *Reed–Muller codes for random erasures and errors*. *Proceedings of STOC*, pp. 297–306, 2015. Journal version: *IEEE Transactions on Information Theory* 61(10):5229–5252, 2015. [arXiv:1411.4590](#).
- [Bec75] William Beckner. *Inequalities in Fourier analysis*. *Annals of Mathematics* 102(1):159–182, 1975. [doi:10.2307/1970980](#).
- [BHSS22] Siddharth Bhandari, Prahladh Harsha, Ramprasad Saptharishi, and Srikanth Srinivasan. *Vanishing spaces of random sets and applications to Reed–Muller codes*. *Computational Complexity Conference, LIPIcs* 234, Article 31, 2022. [doi:10.4230/LIPIcs.CCC.2022.31](#).
- [BLM13] Stéphane Boucheron, Gábor Lugosi, and Pascal Massart. *Concentration Inequalities: A Nonasymptotic Theory of Independence*. Oxford University Press, 2013. [doi:10.1093/acprof:oso/9780199535255.001.0001](#).
- [Bon70] Aline Bonami. *Étude des coefficients de Fourier des fonctions de $L^p(G)$* . *Annales de l’Institut Fourier* 20(2):335–402, 1970. [doi:10.5802/aif.357](#).

- [Bou02] Olivier Bousquet. *A Bennett concentration inequality and its application to suprema of empirical processes*. *Comptes Rendus Mathématique* 334(6):495–500, 2002. doi:10.1016/S1631-073X(02)02292-6.
- [CDL13] Albert Cohen, Mark A. Davenport, and Dany Leviatan. *On the stability and accuracy of least squares approximations*. *Foundations of Computational Mathematics* 13:819–834, 2013. arXiv:1111.4422.
- [CM17] Albert Cohen and Giovanni Migliorati. *Optimal weighted least-squares methods*. *SMAI Journal of Computational Mathematics* 3:181–203, 2017. doi:10.5802/smai-jcm.24.
- [DG07] Percy Deift and Dimitri Gioev. *Universality at the edge of the spectrum for unitary, orthogonal, and symplectic ensembles of random matrices*. *Communications on Pure and Applied Mathematics* 60(6):867–910, 2007. doi:10.1002/cpa.20164. arXiv:math-ph/0507023.
- [DW07] Dan Dai and Roderick Wong. *Global asymptotics of Krawtchouk polynomials: a Riemann–Hilbert approach*. *Chinese Annals of Mathematics, Series B* 28(1):1–34, 2007. doi:10.1007/s11401-006-0195-3.
- [EHE24] Ayoub El Hanchi, Chris J. Maddison, and Murat A. Erdogdu. *Minimax linear regression under the quantile risk*. *Proceedings of COLT*, PMLR 247:1516–1572, 2024. PMLR proceedings.
- [EI22] Alexandros Eskenazis and Paata Ivanisvili. *Learning low-degree functions from a logarithmic number of random queries*. *Proceedings of STOC*, pp. 203–207, 2022. arXiv:2109.10162.
- [EIS23] Alexandros Eskenazis, Paata Ivanisvili, and Lauritz Streck. *Low-degree learning and the metric entropy of polynomials*. *Discrete Analysis*, Article 17, 2023. arXiv:2203.09659.
- [Fil16] Yuval Filmus. *An orthogonal basis for functions over a slice of the Boolean hypercube*. *Electronic Journal of Combinatorics* 23(1), Paper P1.23, 2016. doi:10.37236/4567. arXiv:1406.0142v2.
- [FM19] Yuval Filmus and Elchanan Mossel. *Harmonicity and invariance on slices of the Boolean cube*. *Probability Theory and Related Fields* 175:721–782, 2019. doi:10.1007/s00440-019-00900-w. arXiv:1507.02713v5.
- [KKLT22] Boris Kashin, Egor Kosov, Irina Limonova, and Vladimir Temlyakov. *Sampling discretization and related problems*. *Journal of Complexity* 71:101653, 2022. arXiv:2109.07567.
- [KS21] Naomi Kirshner and Alex Samorodnitsky. *A moment ratio bound for polynomials and some extremal properties of Krawchouk polynomials and Hamming spheres*. *IEEE Transactions on Information Theory* 67(6):3509–3541, 2021. arXiv:1909.11929v1.
- [Lev95] Vladimir I. Levenshtein. *Krawtchouk polynomials and universal bounds for codes and designs in Hamming spaces*. *IEEE Transactions on Information Theory* 41(5):1303–1321, 1995. doi:10.1109/18.412678.
- [Men14] Shahar Mendelson. *Learning without concentration*. *Proceedings of COLT*, PMLR 35:25–39, 2014. PMLR proceedings.
- [Men21] Shahar Mendelson. *Extending the scope of the small-ball method*. *Studia Mathematica* 256(2):147–167, 2021. doi:10.4064/sm190420-21-11. arXiv:1709.00843.
- [Mou22] Jaouad Mourtada. *Exact minimax risk for linear least squares, and the lower tail of sample covariance matrices*. *Annals of Statistics* 50(4):2157–2178, 2022. arXiv:1912.10754.

- [Oli16] Roberto Imbuzeiro Oliveira. *The lower tail of random quadratic forms with applications to ordinary least squares*. *Probability Theory and Related Fields* 166:1175–1194, 2016. [arXiv:1312.2903](#).
- [PS19] Yury Polyanskiy and Alex Samorodnitsky. *Improved log-Sobolev inequalities, hypercontractivity and uncertainty principle on the hypercube*. *Journal of Functional Analysis* 277(11):108280, 2019. [arXiv:1606.07491v3](#).
- [Sam08] Alex Samorodnitsky. *A modified logarithmic Sobolev inequality for the Hamming cube and some applications*. *arXiv preprint*, 2008. [arXiv:0807.1679](#).
- [SL23] Lucas Slot and Monique Laurent. *Sum-of-squares hierarchies for binary polynomial optimization*. *Mathematical Programming* 197:621–660, 2023 (published online in 2022). [doi:10.1007/s10107-021-01745-9](#). [arXiv:2011.04027v3](#).
- [SX22] Kevin Schnelli and Yuanyuan Xu. *Convergence rate to the Tracy–Widom laws for the largest eigenvalue of Wigner matrices*. *Communications in Mathematical Physics* 393:839–907, 2022. [doi:10.1007/s00220-022-04377-y](#).
- [Tropp12] Joel A. Tropp. *User-friendly tail bounds for sums of random matrices*. *Foundations of Computational Mathematics* 12(4):389–434, 2012. [arXiv:1004.4389v7](#).
- [Yas14] Pavel Yaskov. *Lower bounds on the smallest eigenvalue of a sample covariance matrix*. *Electronic Communications in Probability* 19, Article 83, pp. 1–10, 2014. [doi:10.1214/ECP.v19-3807](#). [arXiv:1409.6188](#).

A The entropy curve and the cubic remainder

We collect the elementary properties of the entropy curve and the proof of [Lemma 3.2](#), whose statement and use in clipping appear in [Section 3](#).

A.1 The entropy curve

Recall that all logarithms are natural and

$$H(x) = -x \log x - (1-x) \log(1-x), \quad \Psi(q) = \log 2 - H\left(\frac{1}{2} - \sqrt{q(1-q)}\right), \quad E_{d,k} = d\Psi(k/d),$$

with $0 \log 0 = 0$. The following properties follow by elementary differentiation; we include the details used later.

Lemma A.1 (Elementary calculus). *For $q \in [0, 1/2]$, the function Ψ is increasing and concave, with*

$$\Psi(0) = 0, \quad \Psi(1/2) = \log 2, \quad \Psi'(0) = 2.$$

For any $q_0 < 1/2$, uniformly on $0 \leq q \leq q_0$,

$$\Psi(q) = 2q - \frac{2}{3}q^2 - \frac{8}{15}q^3 + O_{q_0}(q^4), \quad \frac{\Psi(q_0)}{q_0}q \leq \Psi(q) \leq 2q.$$

Proof. Step 1: monotonicity and concavity. Let $z = 2\sqrt{q(1-q)}$ and $w = 1 - 2q$, so $z^2 + w^2 = 1$. Then

$$\begin{aligned} \Psi(q) &= \frac{1+z}{2} \log(1+z) + \frac{1-z}{2} \log(1-z), \\ \Psi'(q) &= \frac{2w}{z} \operatorname{arctanh} z, \\ \Psi''(q) &= \frac{4(z - \operatorname{arctanh} z)}{z^3} < 0. \end{aligned}$$

Here $\Psi'(q) > 0$ and $\Psi''(q) < 0$ for $0 < q < 1/2$. The endpoint values and $\Psi'(0) = 2$ follow by continuity. Concavity places Ψ above its chord from 0 to q_0 and below its tangent at 0, proving the linear bounds.

Step 2: expansion at zero. Integrating the power series for $\operatorname{arctanh} z$ gives

$$\frac{1+z}{2} \log(1+z) + \frac{1-z}{2} \log(1-z) = \sum_{j \geq 1} \frac{z^{2j}}{2j(2j-1)}.$$

Substitute $z^2 = 4q(1-q)$ and collect the first three powers of q . The omitted term is uniformly $O_{q_0}(q^4)$ by analyticity at zero and smoothness on a compact interval ending below $1/2$. $\square$

A.2 Uniform cubic remainder

Proof of Lemma 3.2. Step 1: express the known rate as a minimum. This is an algebraic reformulation of the rate in [\(10\)](#). Take $z = \tanh v$. Then

$$\begin{aligned} \log \frac{(1+z)^p + (1-z)^p}{2} &= \log \cosh(pv) - p \log \cosh v, \\ F(p, q) &= \min_{v > 0} \{\log \cosh(pv) - pg_q(v)\}, \\ g_q(v) &= \log \cosh v + q \log \tanh v + \frac{1}{2} H(q). \end{aligned}$$

Differentiating the above expression in braces gives

$$\begin{aligned} \frac{\partial}{\partial v} \{\log \cosh(pv) - pg_q(v)\} &= \frac{p}{\sinh v \cosh v} (q_p(v) - q), \\ q_p(v) &= \sinh v \cosh v (\tanh(pv) - \tanh v) = \frac{1}{2} \left(1 - \frac{\cosh((p-2)v)}{\cosh(pv)} \right). \end{aligned} \quad (21)$$

For $p \geq 2$, the ratio on the last line decreases strictly from one to zero. This can be seen by inspecting its logarithmic derivative $(p-2)\tanh((p-2)v) - p\tanh(pv) < 0$.

Thus q_p increases strictly from zero to $1/2$. The derivative in (21) changes sign exactly once, from negative to positive. This proves the variational form and uniqueness. It also shows that evaluation at any value of v gives an upper bound on F .

Step 2: choose a test point v . Fix $q \in (0, q_0]$ and set

$$a = \operatorname{arctanh} \sqrt{\frac{q}{1-q}}, \quad q = \frac{\sinh^2 a}{\cosh(2a)}. \quad (22)$$

At $p = 2$, the minimizer is $v = a$. For $p = 2 + 2s$, we can use $v = a/(1+s)$ since this keeps $pv = 2a$ fixed, making the first term of the variational expression constant. Define

$$M_q(s) = \log \cosh(2a) - 2(1+s)g_q\left(\frac{a}{1+s}\right).$$

Then $F(2+2s, q) \leq M_q(s)$.

Step 3: verify cancellation up to the second order. Let $a_0 = \operatorname{arctanh} \sqrt{q_0/(1-q_0)}$. Basic algebra using (22) yields

$$\begin{aligned} 2g_q(a) &= \log \cosh(2a), \\ g'_q(a) &= \tanh(2a), \\ g''_q(a) &= 0, \\ 2a \tanh(2a) - \log \cosh(2a) &= \Psi(q). \end{aligned} \quad (23)$$

To verify these identities, one can simply take $z = \tanh a$. Then $q = z^2/(1+z^2)$ and

$$\begin{aligned} H(q) &= \log(1+z^2) - 2q \log z, \\ g_q(a) &= \log \cosh a + \frac{1}{2} \log(1+z^2) = \frac{1}{2} \log \cosh(2a), \\ g''_q(a) &= \operatorname{sech}^2 a - q \frac{\cosh(2a)}{\sinh^2 a \cosh^2 a} = 0. \end{aligned}$$

The first derivative follows from $g'_q(a) = \tanh a + q/(\sinh a \cosh a) = \tanh(2a)$. Also $\tanh(2a) = 2\sqrt{q(1-q)}$. For $u = \tanh(2a)$, $\log 2 - H((1-u)/2) = 2au - \log \cosh(2a)$, which proves the last line of (23).

Write $w = 1+s$ and $v = a/w$. Differentiation yields

$$\begin{aligned} M'_q(s) &= 2\{vg'_q(v) - g_q(v)\}, \\ M''_q(s) &= -\frac{2v^2}{w} g''_q(v), \\ M'''_q(s) &= \frac{2}{w^2} \{3v^2 g''_q(v) + v^3 g'''_q(v)\}. \end{aligned} \quad (24)$$

It follows from (23) that

$$M_q(0) = 0, \quad M'_q(0) = \Psi(q), \quad M''_q(0) = 0.$$

Step 4: bound the remainder uniformly as $q \downarrow 0$. Taylor's theorem concludes the proof once $|M_q'''(s)| \leq C_{q_0} q$.

Let $\ell(v) = \log \tanh v$. Its derivatives satisfy

$$\begin{aligned}\ell''(v) &= -\frac{4 \cosh(2v)}{\sinh^2(2v)}, \\ \ell'''(v) &= -\frac{8}{\sinh(2v)} + \frac{16 \cosh^2(2v)}{\sinh^3(2v)}.\end{aligned}$$

For $0 < v \leq a_0$, the quantities $v^2|\ell''(v)|$ and $v^3|\ell'''(v)|$ are bounded by constants depending only on a_0 . This can be seen directly from $\sinh(2v) \geq 2v$, since their limits at zero are 1 and 2. Moreover,

$$v^2 \leq a^2 \leq q \cosh(2a_0),$$

because $q = \sinh^2 a / \cosh(2a)$ and $\sinh a \geq a$. The derivatives of $\log \cosh v$ are bounded on $[0, a_0]$. In $g_q(v) = \log \cosh v + q\ell(v) + H(q)/2$, the first term therefore contributes $O_{q_0}(v^2)$ and $O_{q_0}(v^3)$ to the weighted second and third derivatives respectively, while the ℓ term contributes $O_{q_0}(q)$. Since $v \leq a_0$ and $v^2 = O_{q_0}(q)$, this gives

$$|v^2 g_q''(v)| \leq C_{q_0} q, \quad |v^3 g_q'''(v)| \leq C_{q_0} q.$$

For $0 \leq s \leq 1/2$, (24) yields

$$|M_q'''(s)| \leq C_{q_0} q.$$

Taylor's theorem and the bound on the third derivative give

$$M_q(s) \leq s\Psi(q) + C_0 q s^3.$$

Since $F(2 + 2s, q) \leq M_q(s)$, this proves (11). Finally, we apply [Theorem 3.1](#) and multiply by d , using $dq = k$, to obtain (12). $\square$

B Classical analytic and probabilistic tools

This section collects the hypercontractive, empirical-process, and Gaussian probability bounds used in the sampling arguments.

B.1 Bonami–Beckner hypercontractivity

We use the classical degree form of the Bonami–Beckner inequality [[Bon70](#); [Bec75](#)].

Lemma B.1 (Bonami–Beckner inequality). *For $p \geq 2$ and every $h \in \mathcal{P}_{d,k}$,*

$$(\mathbb{E}[|h|^p])^{1/p} \leq (p-1)^{k/2} (\mathbb{E}[h^2])^{1/2}.$$

In particular, for $\mathbb{E}[h^2] = 1$,

$$\mathbb{E}[h^4] \leq 9^k, \quad \log \mathbb{E}[|h|^{2+2s}] \leq k(1+s) \log(1+2s) \quad (s \geq 0).$$

B.2 A bound independent of the ambient dimension

For our unrestricted-dimension result we also need the classical Bonami–Beckner hypercontractive estimate. The clipping step is the same as in [Proposition 3.3](#).

Proposition B.2 (Classical hypercontractive comparison). *For $k \geq 1$, $\eta \in (0, 1)$, and $T_k(\eta) = \sup_{d \geq 1} T_{d,k}(\eta)$,*

$$\log T_k(\eta) \leq 2k + \left(\frac{1}{6} + 2 \log \frac{1}{1-\eta} \right) k^{1/3}.$$

Proof. If $\mathbb{E}[h^2] = 1$ and $\deg h \leq k$, [Lemma B.1](#) with $p = 2 + 2s$ gives

$$\log \mathbb{E}[|h|^{2+2s}] \leq k(1+s) \log(1+2s) \leq 2ks + \frac{2}{3}ks^3 \quad (0 \leq s \leq 1/2).$$

Step 1: a Taylor bound. Indeed, $f(s) = (1+s) \log(1+2s)$ satisfies $f(0) = f''(0) = 0$, $f'(0) = 2$, and $f'''(s) = 4(1-2s)/(1+2s)^3 \leq 4$ on this interval.

Step 2: clipping. We take $L = \log(1/(1-\eta))$, $s = k^{-1/3}/2$, and $\log B = 2k + (2/3)ks^2 + L/s$. Then

$$\mathbb{E}[(h^2 - B)_+] \leq B^{-s} \mathbb{E}[|h|^{2+2s}] \leq e^{-L} = 1 - \eta,$$

so $\mathbb{E}[h^2 \wedge B] \geq \eta$. Substituting s proves the bound uniformly over d and h . $\square$

B.3 A bounded empirical-process inequality

We use the following result in its bounded, centered form. For convenience, we include the zero function in the class, ensuring that $Z \geq 0$.

Theorem B.3 (Bousquet, Theorem 2.3 of [\[Bou02\]](#)). *Let $X_1, \dots, X_n$ be independent and identically distributed, and let $\mathcal{G}$ be a countable class of measurable real functions such that*

$$\mathbb{E}[g] = 0, \quad |g| \leq 1, \quad \mathbb{E}[g^2] \leq v \quad (g \in \mathcal{G}).$$

Assume $0 \in \mathcal{G}$, and put $Z = \sup_{g \in \mathcal{G}} \sum_{i=1}^n g(X_i)$. Then, for every $t > 0$,

$$\mathbb{P} \left(Z > \mathbb{E}Z + \sqrt{2(nv + 2\mathbb{E}Z)t} + \frac{t}{3} \right) \leq e^{-t}.$$

By continuity in the coefficient vector, all suprema may be taken over a countable dense subset of the unit sphere, so Bousquet’s countability assumption is satisfied.

B.4 Symmetrization and contraction

The following are two standard results, see [\[BLM13, Lemma 11.4 and Theorem 11.6\]](#).

Lemma B.4 (Symmetrization and contraction). *Let $\varepsilon_1, \dots, \varepsilon_n$ be independent random signs, each uniform on $\{-1, 1\}$.*

(i) Symmetrization. *Let $X_1, \dots, X_n$ be independent and identically distributed, independently of the signs. For a countable class $\mathcal{F}$ of integrable functions,*

$$\mathbb{E} \sup_{f \in \mathcal{F}} |\mathbb{E}_n[f] - \mathbb{E}[f]| \leq 2 \mathbb{E} \sup_{f \in \mathcal{F}} \left| \frac{1}{n} \sum_{i=1}^n \varepsilon_i f(X_i) \right|.$$

The expectation on the right includes both the sample and the signs.

(ii) Contraction. For a bounded set $\mathcal{A} \subseteq \mathbb{R}^n$ and an L -Lipschitz function $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ satisfying $\varphi(0) = 0$,

$$\mathbb{E}_\varepsilon \sup_{a \in \mathcal{A}} \left| \sum_{i=1}^n \varepsilon_i \varphi(a_i) \right| \leq 2L \mathbb{E}_\varepsilon \sup_{a \in \mathcal{A}} \left| \sum_{i=1}^n \varepsilon_i a_i \right|.$$

B.5 Elementary Gaussian probability bounds

Lemma B.5 (Bounds with all ranges explicit). For $r \geq 1$, the chi-squared variable defined in Section 1.1 satisfies the following bounds. For $u \geq 0$ and $0 < a < 1$,

$$\mathbb{P}(\chi_r^2 \geq 4(r+u)) \leq e^{-u}, \quad (25)$$

$$\mathbb{P}(\chi_r^2 \leq ar) \leq (ae^{1-a})^{r/2}. \quad (26)$$

For a one-dimensional standard Gaussian G and every $t \geq \log 4$,

$$\frac{15}{16} e^{-t/256} \mathbb{P}(G^2 \geq t/64) > e^{-t}. \quad (27)$$

Proof. Completing the square in the Gaussian integral gives $\mathbb{E}[e^{v\chi_r^2}] = (1-2v)^{-r/2}$ for $v < 1/2$. Exponential Markov with $v = 1/4$ gives

$$\mathbb{P}(\chi_r^2 \geq 4(r+u)) \leq e^{-r-u} 2^{r/2} \leq e^{-u}.$$

For the lower tail use $v = -(1-a)/(2a) < 0$:

$$\mathbb{P}(\chi_r^2 \leq ar) \leq e^{-var} (1-2v)^{-r/2} = (ae^{1-a})^{r/2}.$$

To prove (27), first suppose $\log 4 \leq t \leq 16$. Then $\sqrt{t/64} \leq 1/2$. The Gaussian density is bounded by $(2\pi)^{-1/2}$, so

$$\mathbb{P}(|G| \geq 1/2) \geq 1 - (2\pi)^{-1/2} > 3/5.$$

The left side of (27) is then greater than $(15/16)e^{-1/16}(3/5) > 1/2$, while $e^{-t} \leq 1/4$.

For $t \geq 16$, put $b = \sqrt{t}/8$. Integrate over $[b, b+1]$ and $[-b-1, -b]$:

$$\mathbb{P}(G^2 \geq t/64) \geq \sqrt{2/\pi} e^{-(b+1)^2/2} = \sqrt{2/\pi} e^{-1/2} e^{-t/128 - \sqrt{t}/8} > \frac{1}{4} e^{-5t/128},$$

since $\sqrt{t}/8 \leq t/32$. Thus the left side of (27) exceeds $(15/64)e^{-11t/256}$. This is larger than e^{-t} for every $t \geq 16$. $\square$

Corollary B.6 (OLS after lower conditioning). If $\mathbb{P}(\hat{\Sigma}_V \succeq \frac{1}{2}I_m) \geq 1 - \delta/2$, where $\delta \leq 1/4$, then

$$\mathcal{R}_{n,\delta}^*(V) \leq 12\sigma^2 \frac{m + \log(1/\delta)}{n}.$$

OLS satisfies the same bound uniformly over θ .

Proof. Conditionally on a design with $\hat{\Sigma}_V \succeq I_m/2$,

$$\|\hat{\theta}_{\text{OLS}} - \theta\|_2^2 \preceq_{\text{st}} \frac{2\sigma^2}{n} \chi_m^2.$$

Use (25) with $u = t + \log 2$, where $t = \log(1/\delta)$, and add the design failure probability. Since $t \geq \log 4$, $\log 2 \leq t/2$, and $8(m+t+\log 2) \leq 12(m+t)$.

The minimax assertion follows because OLS is an admissible estimator (or from Theorem 5.2). $\square$

C The radial-weight identity

We derive [Lemma 4.1](#) directly from [\[FM19, Theorem 3.10\]](#). We first apply that theorem to the uniform cube to identify its measure-independent constant, and then to the cube weighted by $H(S_d)$.

Proof. Filmus–Mossel’s theorem states that, for homogeneous harmonic multilinear f, g of degrees r, s and any exchangeable probability measure α , $\mathbb{E}_\alpha[fg] = 0$ if $r \neq s$. If $r = s$, there is a constant $C_{f,g}$ independent of α such that

$$\mathbb{E}_\alpha[fg] = C_{f,g} \mathbb{E}_\alpha \left[\prod_{j=1}^r (x_{2j-1} - x_{2j})^2 \right].$$

Apply this first with $\alpha = \mu_d$. Each squared difference has expectation 2, and the pairs are independent, hence

$$C_{f,g} = 2^{-r} \langle f, g \rangle_d.$$

For $H \geq 0$ with $Z = \mathbb{E}_{\mu_d}[H(S_d)] > 0$, now choose

$$\alpha(x) = \frac{H(S_d(x))}{Z} \mu_d(x).$$

This probability measure is exchangeable because both S_d and μ_d are invariant under coordinate permutations. Substituting it into the theorem and multiplying by Z gives

$$\mathbb{E}_{\mu_d}[fgH(S_d)] = \langle f, g \rangle_d \mathbb{E}_{\mu_d}[Q_r^2 H(S_d)], \quad Q_r = \prod_{j=1}^r \frac{x_{2j-1} - x_{2j}}{\sqrt{2}}.$$

On the other hand, for unequal degrees, the same substitution yields zero.

To evaluate the remaining expectation, let A_r be the event that the signs differ in each of the first r pairs. Then

$$\Pr_{\mu_d}(A_r) = 2^{-r}, \quad Q_r^2 = 2^r \mathbf{1}_{A_r}.$$

Conditional on A_r , those pairs contribute zero to S_d , while the remaining $d - 2r$ coordinates are independent uniform signs. Thus

$$\mathbb{E}_{\mu_d}[Q_r^2 H(S_d)] = \mathbb{E}_{\mu_d}[H(S_d) \mid A_r] = \mathbb{E}_{\mu_{d-2r}}[H(S_{d-2r})],$$

which is the equal-degree case of [\(14\)](#). The empty product for $r = 0$ is one. If $H \geq 0$ and $Z = 0$, the weighted identities are immediate since $H(S_d) = 0$ on the cube. Finally, decomposing a real H into its positive and negative parts extends the result to signed weights by linearity. $\square$

D Probability constants for the sampling lower bounds

We give the complete probability bounds for the stable-sampling and Gaussian-risk obstructions from [Lemma 5.1](#). Throughout, $\mathbb{P}(E) = p \in (0, 1/2]$. The shared-set case uses [\(16\)](#). For $t \geq m$, the directional assumption in [Lemma 5.1\(ii\)](#) suffices.

Lemma D.1 (Baseline sample complexity). *For $d, k \geq 1$, $1 \leq m \leq D_{d,k}$, $0 < \delta \leq 1/4$, $0 < c < 1$, and $A \geq 16$, there is a universal $c_0 > 0$ such that*

$$N_{\text{stab}}(d, k, m, c, \delta) \wedge N_{\text{par}}(d, k, m, A, \delta) \geq c_0(m + \log(1/\delta)).$$

Proof of Lemma D.1. If $n < m$, every feature matrix is singular, so both guarantees fail. This gives the lower bound m .

For the confidence term, include $h(x) = (1 + x_1)/\sqrt{2}$ in the model and extend it to an orthonormal basis of any feasible size m . If $t = \log(1/\delta)$, then

$$n \leq \frac{t}{2 \log 2} \implies \mathbb{P}(X_{i1} = -1 \text{ for every } i) = 2^{-n} \geq e^{-t/2} > \delta.$$

On this event the feature column for h is zero, so both guarantees fail. Taking an integer sample size below the displayed bound yields a universal constant times t . If that range contains no positive integer, use $N \geq 1$. Finally, $m \vee t \geq (m + t)/2$. $\square$

Proposition D.2 (Stable-sampling obstruction). *Fix $0 < c < 1$ and choose $0 < \rho < c/256$. Under (16), or under the directional assumption in Lemma 5.1(ii) when $t \geq m$, for every $t \geq \log 4$ and every positive integer $n \leq (m + t)/(1024p)$, we have*

$$\mathbb{P}(\widehat{\Sigma} \not\preceq cI_m) > e^{-t}.$$

Proof. If $t < m$, then $m \geq 2$ and $n \leq m/(512p)$, so Lemma 5.1(i) applies. Its small eigenvalues are strictly below c , and its probability $15/16$ exceeds $e^{-t} \leq 1/4$. If $t \geq m$, then $n \leq t/(512p)$, and part (ii) applies. Now $32\rho < c$, and

$$\frac{15}{16}e^{-t/256} > e^{-t} \quad (t \geq \log 4).$$

This proves the strict failure bound in both regimes. $\square$

Proposition D.3 (Estimator-independent parametric obstruction). *Fix $A \geq 16$. There is $\rho_A > 0$ such that, under (16), or under the directional assumption in Lemma 5.1(ii) when $t \geq m$, with $\rho \leq \rho_A$, for every $t \geq \log 4$ and every positive integer $n \leq (m + t)/(1024p)$, the Gaussian model satisfies*

$$\mathcal{R}_{n,e^{-t}}^*(V) > A\sigma^2 \frac{m + t}{n}.$$

Proof. Take $b_0 = 1/(1024e)$ and choose $0 < \rho_A < \min \left\{ \frac{1}{8}, \frac{b_0}{3072A}, \frac{1}{16384A} \right\}$. Use the random variable $Z_{n,V}$ in Theorem 5.2. A singular design makes it infinite, so it is enough to focus on nonsingular designs.

Suppose first that $t < m$. Then $m \geq 2$. With probability at least $15/16$, at least $r = \lfloor m/2 \rfloor \geq m/3$ eigenvalues of $\widehat{\Sigma}$ are at most 128ρ . Conditional on any such nonsingular design,

$$Z_{n,V} \succeq_{\text{st}} \frac{\sigma^2}{128\rho n} \chi_r^2.$$

By (26), $\mathbb{P}(\chi_r^2 < b_0 r) \leq (eb_0)^{r/2} \leq 1/32$. Thus, with unconditional probability at least $(15/16)(31/32) > 1/4$,

$$Z_{n,V} \geq \frac{\sigma^2 b_0 r}{128\rho n} \geq \frac{\sigma^2 b_0 m}{384\rho n} > 8A\sigma^2 \frac{m}{n} > A\sigma^2 \frac{m + t}{n}.$$

If $t \geq m$, Lemma 5.1(ii) gives an event D of probability at least $(15/16)e^{-t/256}$ on which $\lambda_1(\widehat{\Sigma}) \leq 32\rho$. Conditional on such a design, one coordinate of the independent standard Gaussian vector yields

$$Z_{n,V} \succeq_{\text{st}} \frac{\sigma^2}{32\rho n} G^2.$$

By (27), the event

$$Z_{n,V} \geq \frac{\sigma^2 t}{2048 \rho n}$$

has probability strictly greater than e^{-t} . Its displayed threshold is greater than $8A\sigma^2 t/n$, hence greater than the benchmark because $m \leq t$. This proves that the lower $(1 - e^{-t})$ -quantile of $Z_{n,V}$ strictly exceeds the benchmark. Lastly, apply Theorem 5.2. $\square$

E Comparison to the Polyanskiy–Samorodnitsky uncertainty boundary

We restate the Polyanskiy–Samorodnitsky uncertainty principle in our notation and explain how it yields the leading exponent of our one-dimensional clipping threshold when k/d tends to a fixed value in $(0, 1/2)$.

Let $B_{d,r}$ denote a Hamming ball of radius $\lfloor rd \rfloor$ and put $r_*(q) = \frac{1}{2} - \sqrt{q(1-q)}$.

Let $\Pi_{\leq k}$ be orthogonal projection onto $\mathcal{P}_{d,k}$ and $C_E = \Pi_{\leq k} \mathbf{1}_E \Pi_{\leq k}$ its concentration operator, where $\mathbf{1}_E$ acts by multiplication. Its eigenvalues are denoted $\kappa_1(C_E) \geq \kappa_2(C_E) \geq \dots$.

Then, Theorem 9 of [PS19] implies the following statements for fixed $q \in (0, 1/2)$ and fixed $r \in (0, 1/2)$, with $k = \lfloor qd \rfloor$: for some $c = c(q, r) > 0$ and all sufficiently large d ,

$$\begin{aligned} r < r_*(q), \quad |E| \leq e^{dH(r)} &\implies \kappa_1(C_E) \leq e^{-cd}, \\ r > r_*(q), \quad E = B_{d,r} &\implies \kappa_1(C_E) \geq 1 - e^{-cd}. \end{aligned}$$

The cosine of their angle between the spatial and Fourier subspaces is the square root of $\kappa_1(C_E)$; squaring only changes the constants in these bounds. The boundary equation is $(1-2r)^2 + (1-2q)^2 = 1$, and $\mu_d(B_{d,r}) = \exp\{-d(\log 2 - H(r)) + o(d)\}$. Thus the critical probability exponent is precisely $E_{d,k}$.

For each fixed $\eta \in (0, 1)$, these statements imply

$$\log T_{d, \lfloor qd \rfloor}(\eta) = d\Psi(q) + o(d).$$

For the upper bound, choose $B = \exp\{d(\Psi(q) + \varepsilon)\}$ with fixed $\varepsilon > 0$. For every unit polynomial h of degree at most $\lfloor qd \rfloor$, the set $E = \{h^2 > B\}$ has probability at most $1/B$. The first implication above therefore gives $\mathbb{E}[h^2 \mathbf{1}_E] = o(1)$, uniformly in h , and hence $\mathbb{E}[h^2 \wedge B] \geq 1 - o(1) \geq \eta$ for sufficiently large d .

For the lower bound, the second implication provides a unit polynomial h with $\mathbb{E}[h^2 \mathbf{1}_{E^c}] = o(1)$ on a Hamming ball E whose probability exponent is arbitrarily close to $\Psi(q)$. Choosing this ball so that $\mu_d(E) \leq \exp\{-d(\Psi(q) - \varepsilon/2)\}$ and setting $B = \exp\{d(\Psi(q) - \varepsilon)\}$ gives

$$\mathbb{E}[h^2 \wedge B] \leq \mathbb{E}[h^2 \mathbf{1}_{E^c}] + B\mu_d(E) = o(1) < \eta.$$

Thus $\log T_{d, \lfloor qd \rfloor}(\eta)/d \rightarrow \Psi(q)$.

F Further bounds and model examples

We derive the additional bounds and justify the model examples discussed in Section 7.

F.1 Ambient dimension and model dimension

The finite-dimensional correction. By Lemma A.1,

$$\Psi(q) = 2q - \frac{2}{3}q^2 - \frac{8}{15}q^3 + O(q^4).$$

Substituting this expansion into Theorem 1.5 gives, throughout the matching range and as $k/d \rightarrow 0$,

$$\log \frac{N_{\text{par}}}{m+t} = 2k - \frac{2k^2}{3d} + O\left(\frac{k^3}{d^2} + k^{1/3}\right).$$

Thus the first correction to the unrestricted-dimension exponent $2k$ is $-2k^2/(3d)$. Relative to its magnitude, the two error terms have orders k/d and $d/k^{5/3}$, respectively. Both vanish when $k \ll d \ll k^{5/3}$. Alternatively, if $k/d \rightarrow q \in (0, 1/2)$, then

$$\frac{1}{d} \log \frac{N_{\text{par}}}{m+t} \longrightarrow \Psi(q)$$

throughout the matching range.

Worst case over the ambient dimension. Let N_{par}^∞ be the least threshold valid on every cube admitting a model of dimension m , and define N_{stab}^∞ analogously. For either threshold N^∞ , we show that

$$\log \frac{N^\infty}{m+t} = 2k + O(k^{1/3}),$$

with constants depending on the fixed target parameters.

For the upper bound, Propositions B.2 and 3.4 and Corollary B.6 give

$$N^\infty \leq C(m+t) \exp\{2k + Ck^{1/3}\},$$

uniformly over d , including $k > d$. For the lower bound, it suffices to exhibit one suitable ambient dimension. Choose

$$d = \lceil \max\{m+1, 10k, k^{5/3}\} \rceil.$$

Then $m < d \leq M_{d,k}$ and $k/d \leq 1/10$, so the matching lower bound applies. Moreover,

$$E_{d,k} \geq 2k - Ck^2/d \geq 2k - Ck^{1/3}.$$

The lower bound of Theorem 1.5, and its stable-sampling counterpart, therefore prove the claim. The same choice of d , together with the classical upper bound and Proposition 3.3, also gives

$$\log \sup_d T_{d,k}(\eta) = 2k + O_\eta(k^{1/3}).$$

A lower bound for larger model dimensions. By allocating more degree to the harmonic directions, we can extend the lower bound beyond $m \leq M_{d,k}$, at the cost of a larger error. Specifically, for any integer $0 \leq R \leq k$ with $m \leq \binom{d}{R}$, we have

$$\log \frac{N_{\text{par}}}{m+t} \geq E_{d,k} - C(R + k^{1/3}).$$

The same bound holds for N_{stab} , with constants depending on its fixed conditioning level.

To prove this, [Proposition 4.2](#) supplies a common concentration set of probability at most e^{-a_R} , where

$$a_R = \mathcal{E}_{d,k}(R) - C(k - R)^{1/3}.$$

If $a_R \geq \log 2$, this probability is at most $1/2$, so [Proposition D.3](#) gives

$$\log \frac{N_{\text{par}}}{m+t} \geq a_R - O(1) \geq E_{d,k} - C'(R + k^{1/3}),$$

where the last inequality uses (15). If $a_R < \log 2$, the same entropy comparison gives $E_{d,k} \leq C'(R + k^{1/3}) + O(1)$. In this case, the baseline bound $N_{\text{par}} \geq c_0(m+t)$ from [Lemma D.1](#) proves the claim after increasing the constant. For stable sampling, use [Proposition D.2](#) in place of [Proposition D.3](#).

The leading exponent for subexponential model dimensions. Suppose $k/d \rightarrow q \in (0, 1/2)$ and $\log m = o(d)$. Then, even outside the cube-root matching range,

$$\log \frac{N_{\text{par}}}{m+t} = E_{d,k} + o(d),$$

and likewise for N_{stab} .

Indeed, choose $R = \lceil \log_2 m \rceil$, with $R = 0$ when $m = 1$. We have $R = o(d)$, so eventually $R \leq k/2$ and $d \geq 2R$. For $R > 0$,

$$\binom{d}{R} \geq (d/R)^R \geq 2^R \geq m.$$

The preceding lower bound therefore applies, with error $O(R + k^{1/3}) = o(d)$. The upper bound of [Theorem 1.5](#) also has error $o(d)$ and holds for every feasible m . Together these establish the leading exponent, uniformly in $t \geq \log 4$, without asserting an $O(k^{1/3})$ remainder throughout this larger dimension range.

F.2 Known Walsh coordinates and finite-space caps

For an isotropic feature vector with $\|\Phi(x)\|_2^2 \leq L$, matrix Chernoff [[Tropp12](#), Corollary 5.2] gives, for $0 < c < 1$,

$$\mathbb{P}(\lambda_{\min}(\widehat{\Sigma}) \leq c) \leq m \exp\left(-\frac{(1-c)^2 n}{2L}\right). \quad (28)$$

Indeed, its matrix summands are positive semidefinite, have norm at most L , and their expectations sum to nI_m . The standard Chernoff factor is $[e^{-u}/(1-u)^{1-u}]^{n/L}$ with $u = 1 - c$. The inequality $-u - (1-u)\log(1-u) \leq -u^2/2$ follows by differentiation.

For a known span of m Walsh characters, $\|\Phi(x)\|_2^2 = m$ exactly. Hence $n \geq C_c m(\log m + t)$ gives stable sampling. Choosing $c = 1/2$, failure probability $\delta/2$, and using [Corollary B.6](#) gives parametric prediction. Character values need not be mutually independent; only independent sample rows are used.

Let $\chi_{\leq k}(x) = (\chi_S(x))_{|S| \leq \min(k,d)} \in \mathbb{R}^{D_{d,k}}$ be the column vector of Walsh characters. Expanding each basis function of V in this basis gives

$$\Phi_V(x) = U \chi_{\leq k}(x),$$

where $U \in \mathbb{R}^{m \times D_{d,k}}$ contains the expansion coefficients.

Since U has orthonormal rows, $\|\Phi_V(x)\|_2^2 \leq \|\chi_{\leq k}(x)\|_2^2 = D_{d,k}$. Thus (28) gives

$$N \leq CD_{d,k}(\log m + t)$$

for either the stable-sampling or parametric threshold, with C depending on the fixed target parameters.

Likewise, every unit polynomial $h \in \mathcal{P}_{d,k}$ satisfies $h(x)^2 \leq D_{d,k}$. The inequality $z \wedge (\eta D_{d,k}) \geq \eta z$ for $0 \leq z \leq D_{d,k}$ therefore gives

$$\mathbb{E}[h^2 \wedge (\eta D_{d,k})] \geq \eta, \quad T_{d,k}(\eta) \leq \eta D_{d,k}.$$

When $k \geq d$, equality follows by taking $h = 2^{d/2} \mathbf{1}_{\{x=x_0\}}$ for any cube vertex x_0 . These bounds reflect the finite size of the cube: once $k \geq d$, the polynomial space no longer grows, so the threshold cannot continue to scale as e^{2k} .