

Shared Gaussianization: What Gaussian Regularizers Certify About Contrastive Learning, and What They Miss

Ruoyu Zhao^{1,*}, Yuting Chen^{2,*}, Jinheng Zhang³, Zhehao Zou⁴, and Tong Che^{5,†}

¹City University of Hong Kong, ruoyuzhao8-c@my.cityu.edu.hk

²Georgia Institute of Technology, yuting3123@gmail.com

³University of Pennsylvania, jinhengz@seas.upenn.edu

⁴The Chinese University of Hong Kong, 1155257174@link.cuhk.edu.hk

⁵NVIDIA Research, tongc@nvidia.com

Abstract

What can a distribution-matching regularizer such as SIGReg in LeJEPa certify about contrastive learning? We study shared Gaussianization (SG), a characteristic-function Gaussianity test on the average of two normalized views, scaled by an independent χ_d radius. Because disagreeing views shorten the average, one test detects both misalignment and non-uniformity. SG vanishes exactly at the aligned, uniform minimizers of population InfoNCE, and under equal marginals it bounds the InfoNCE excess by $4 \cdot 3^{3/4} \beta \sqrt{\mathcal{J}}$ plus an $O(\mathcal{J})$ term. The rate and the dimension-free leading constant are sharp, and no squared mean-embedding distance on view pairs achieves a faster rate. With an explicit alignment term, a rotation-invariant uniformity test gives a linear bound if and only if its spectrum dominates that of $e^{\beta u^T v}$; SG's own test does, Gaussian kernels $e^{-\gamma \|u-v\|^2}$ qualify exactly when $\gamma \geq \beta/2$, and moment matching never does. Away from the optimum, the objectives differ. Along an isotropic nuisance channel, pure SG lowers its loss by adding per-view nuisance whenever the shared code is non-uniform. An alignment weight above the channel's gain makes the nuisance-free solution a strict local minimizer; for LeJEPa, the same rule gives a critical SIGReg weight that decreases with the batch size. At finite batch size, an off-diagonal U-statistic removes a plug-in bias toward misalignment. In controlled latent-variable models, pure SG retains per-view style, an alignment weight above the measured gain removes it, and for LeJEPa at three batch sizes the measured gain separates the encoders that retain style from those that do not. InfoNCE training also reaches a lower $SG_{0.2}$ loss than $SG_{0.2}$ training from scratch, which points to an optimization gap.

1. Introduction

Self-supervised representation learning commonly uses two families of joint-embedding objectives: contrastive methods and distribution-matching methods. Contrastive methods such as InfoNCE and SimCLR (van den Oord et al., 2018; Chen et al., 2020) pull positive pairs together and push the rest of the batch away; population InfoNCE is *alignment plus uniformity* on the sphere (Wang and Isola, 2020). Distribution-matching methods drop the negatives and prevent collapse with a test of the embedding distribution: variance–covariance penalties (Bardes et al., 2022; Zbontar et al., 2021), spherical uniformity tests (Zheng et al., 2023; Nicollier et al., 2026a), and the sketched isotropic Gaussian regularizer (SIGReg) of LeJEPa (Balestriero and LeCun, 2025), recently recast as a kernel maximum mean discrepancy (MMD) (Zimmermann et al., 2025). Such a test needs no pairwise similarity matrix; LeJEPa enforces view agreement with a separate alignment (prediction) term. Tuned carefully, the families achieve

comparable accuracy (Garrido et al., 2023), which leaves open what a distribution test certifies about contrastive learning and whether the two learn the same representation.

We ask whether a single distribution-matching loss can enforce both alignment and uniformity, and study *shared Gaussianization* (SG). For normalized embeddings $U, V \in \mathbb{S}^{d-1}$ of two augmented views, SG tests the law of $Z = RC$, with centroid $C = \frac{1}{2}(U + V)$ and an independent radius $R \sim \chi_d$, against $\mathcal{N}(0, I_d)$ using SIGReg's characteristic-function loss $\mathcal{J}$. A uniform direction times an independent χ_d radius is exactly Gaussian, while $\|C\|^2 = 1 - \frac{1}{4}\|U - V\|^2$, so disagreement shortens the centroid, and the same test sees both global non-uniformity and pairwise misalignment.

To compare SG with population InfoNCE, we separate three questions: whether two objectives share the same ideal solution, whether one quantitatively controls the other, and whether they favor the same representations away from that solution. SG answers

*Equal contribution. †Project lead.

yes to the first two and no to the third. Away from the optimum, SG can keep per-view nuisance that InfoNCE rejects, unless an explicit alignment price exceeds the gain of the nuisance channel. Finite-batch estimation adds a further difference. Our main contributions are as follows.

- **Population guarantees** (Section 4). SG vanishes exactly when the views agree almost surely and their common law is uniform. Under equal marginals it controls excess InfoNCE at a square-root rate whose dimension-free leading constant is attained by an explicit extremal family (Theorem 1), and no mean-embedding (Hilbertian) pair discrepancy with this target can do better (Proposition 3). With an explicit alignment term the certificate becomes linear, and a spectral criterion, condition (U), characterizes which uniformity tests allow this (Theorem 2).
- **Finite-batch estimation** (Section 5). The plug-in estimator inherits SIGReg’s finite-sample bias, and the centroid turns it into a preference for imperfect alignment. An off-diagonal U-statistic removes the bias at $O(BdM)$ cost without a pairwise similarity matrix (Propositions 4 and 5).
- **Nuisance preferences and the alignment price** (Section 6). Population agreement does not fix preferences away from the optimum. Along an isotropic channel, per-view nuisance lowers SG at first order whenever the shared code is non-uniform, while population InfoNCE rejects it near uniformity (Theorem 3). An alignment price (SG_α), $\mathcal{J}_\alpha = \mathcal{J} + \alpha\Delta$ with Δ the expected misalignment, makes the nuisance-free solution a strict local minimizer when α exceeds the channel’s gain and not when it falls below; the same rule applies to two-term objectives such as LeJEPa (Proposition 6).
- **Experiments** (Section 7). In controlled latent models, SG_α comes close to InfoNCE without per-view nuisance, and Gaussian uniformity tests that satisfy (U) reach InfoNCE’s uniformity in training while moment matching does not. With nuisance, pure SG retains or increases it, and an alignment price above the gain measured on the trained encoders removes it; for LeJEPa at three batch sizes the measured gain separates the encoders that keep style from those that do not. InfoNCE training also reaches lower values of the $SG_{0.2}$ objective than $SG_{0.2}$ trained from scratch, which points to an *optimization gap*.

2. Related Work

Alignment and uniformity. Our comparison builds on the decomposition of InfoNCE, in the limit of infinitely many negatives, into alignment and uniformity (Wang and Isola, 2020); see Wang and Liu (2021) for the role of temperature. Downstream guarantees (Saunshi et al., 2019; HaoChen et al., 2021) and identifiability results for contrastive (Zimmermann et al.,

2021; von Kügelgen et al., 2021; Rusak et al., 2025) and LeJEPa-type objectives (Klindt et al., 2026; Nicollier et al., 2026c) characterize what the optimum represents. Betser et al. (2026) show that, under alignment and concentration assumptions, low-dimensional projections of InfoNCE representations become Gaussian as the dimension grows. We instead compare a distribution test with InfoNCE away from the optimum, through a sharp bound on InfoNCE’s excess that holds for every law with equal marginals and through local preferences.

Gaussian and kernel regularizers. SIGReg (Balestriero and LeCun, 2025) matches sliced characteristic functions of each view to $\mathcal{N}(0, I)$ (Epps and Pulley, 1983; Baringhaus and Henze, 1988) and adds a predictive alignment term; KerJEPa (Zimmermann et al., 2025) recasts it as a sliced MMD; we use its kernel form and SIGReg’s bias identity directly. SFRIK (Zheng et al., 2023) and SPHERE-JEPa (Nicollier et al., 2026a,b) use zonal or sliced uniformity tests on the sphere, which are Sobolev tests of uniformity (Giné, 1975; García-Portugués and Verdebout, 2018). Reported limitations of SIGReg-type regularizers include vanishing gradients (Yu et al., 2026; Wu et al., 2026) and failure to prevent spurious attributes from structuring the latent space (Nicollier et al., 2026d), and one-view marginals do not identify the coupling of the views (Jenkinson Alvarez, 2026); our gain–price rule (Section 6.3) gives the invariance weight at which a two-term objective nevertheless rejects a given nuisance channel. A sphere–interval decomposition of the Gaussian also appears in Parakhin et al. (2026). These regularizers test marginal embedding distributions, whereas SG tests a statistic of the pair. Testing the pair gives InfoNCE’s exact target and a sharp comparison with it, and it turns SIGReg’s finite-sample bias into a bias toward misalignment.

Contrastive and non-contrastive. Garrido et al. (2023) relate covariance criteria to sample-Gram criteria, both second-moment objects, as is the whitening of W-MSE (Ermolov et al., 2021); SSL-HSIC maximizes a kernel dependence (Li et al., 2021), and Fang et al. (2024) propose a Wasserstein uniformity metric; MMCR (Yerxa et al., 2023; Schaeffer et al., 2024) also uses the norm of the view centroid, with a nuclear-norm objective. Mikulasch and Zenke (2026) unify contrastive, non-contrastive and predictive objectives as latent distribution matching, Chen et al. (2024) recast InfoNCE as optimal-transport alignment, and Dufumier et al. (2023) replace its uniformity term with a kernel-based decoupled uniformity. We compare targets, certificates and local preferences quantitatively, and show that second-moment matching cannot control InfoNCE’s uniformity term (Corollary 1).

3. Shared Gaussianization

Let $d \geq 2$, $\mathbb{S}^{d-1} \subset \mathbb{R}^d$ be the unit sphere, and σ be its uniform probability measure. A *paired law* P is the joint law of $(U, V) \in \mathbb{S}^{d-1} \times \mathbb{S}^{d-1}$, with marginals q_U, q_V and mixture marginal $\bar{q} = \frac{1}{2}(q_U + q_V)$; P has *equal marginals* if $q_U = q_V =: q$. We measure misalignment by

$$\Delta(P) = \frac{1}{4} \mathbb{E} \|U - V\|^2 = \frac{1}{2} (1 - \mathbb{E} U^\top V) \in [0, 1]. \quad (1)$$

The shared loss. Given $(U, V) \sim P$, let

$$C = \frac{1}{2}(U + V), \quad Z = RC, \quad R \sim \chi_d, \quad (2)$$

with R independent of (U, V) , and write $\varphi_Z(\omega) = \mathbb{E} e^{i\omega^\top Z}$. SG matches the law of Z to $\mathcal{N}(0, I_d)$ through

$$\mathcal{J}(P) = \mathbb{E}_{a \sim \sigma, t \sim \mathcal{N}(0,1)} |\varphi_Z(ta) - e^{-t^2/2}|^2, \quad (3)$$

where $e^{-t^2/2}$ is the standard Gaussian characteristic function at ta ($\|a\| = 1$). This is the sliced Epps–Pulley statistic of SIGReg (Epps and Pulley, 1983; Balestriero and LeCun, 2025), applied to a statistic of the pair instead of to each view; equivalently, $\mathcal{J}$ is the squared MMD (Gretton et al., 2012) between the law of Z and $\mathcal{N}(0, I_d)$ for the kernel ${}_1F_1(\frac{1}{2}; \frac{d}{2}; -\|x - y\|^2/2)$ (Zimmermann et al., 2025, Thm. 7). If $U = V \sim \sigma$, then $Z = RU \sim \mathcal{N}(0, I_d)$; otherwise $\|C\| < 1$ on disagreeing pairs. The centroid is not renormalized since its norm carries the alignment signal. SG itself is defined for arbitrary paired laws; equal marginals are needed only for the comparison with InfoNCE.

Population InfoNCE. Let P have equal marginals q and let $\beta = 1/\tau > 0$. InfoNCE with N negatives, minus $\log N$, converges as $N \rightarrow \infty$ to (Wang and Isola, 2020)

$$\mathcal{L}_\beta(P) = -\beta \mathbb{E} U^\top V + \mathbb{E}_{U \sim q} \log \mathbb{E}_{W \sim q} e^{\beta U^\top W}. \quad (4)$$

With $\kappa_d(\beta) = \mathbb{E}_{W \sim \sigma} e^{\beta u^\top W}$, which does not depend on $u \in \mathbb{S}^{d-1}$, and $\mathcal{L}_\beta^* = -\beta + \log \kappa_d(\beta)$, the excess separates into a misalignment cost and a uniformity excess (Lemma 3):

$$\mathcal{L}_\beta(P) - \mathcal{L}_\beta^* = 2\beta \Delta(P) + \mathcal{U}_\beta(q), \quad (5)$$

where $\mathcal{U}_\beta(q) = \mathbb{E}_{U \sim q} \log \mathbb{E}_{W \sim q} e^{\beta U^\top W} - \log \kappa_d(\beta) \geq 0$, with equality only at $q = \sigma$ (Lemma 4). Our question is whether the single discrepancy $\mathcal{J}(P)$ controls both terms. For NT-Xent, whose negatives are pooled from both views, the population limit is $\mathcal{L}_\beta(\tilde{P})$, where $\tilde{P}$ swaps the coordinates with probability $\frac{1}{2}$; $\tilde{P}$ has equal marginals $\bar{q}$ and the same Δ and $\mathcal{J}$, so equal marginals suffice for this symmetrized comparison.

4. Population Guarantees for SG

Because view disagreement shortens the centroid (Section 3), we can identify the exact population minimizers of SG and obtain a sharp comparison with population InfoNCE, at a rate that no Hilbertian pair discrepancy with InfoNCE’s target can improve. Once

misalignment is charged linearly, a spectral condition decides which uniformity tests give a linear certificate. Proofs are in Appendix A.

4.1. Alignment from the Centroid Radius

Let $H = \frac{1}{2}(U - V)$ and $Q = \|H\|^2$. Then $\|C\|^2 + Q = 1$ and $\mathbb{E} Q = \Delta(P)$, so disagreement appears as a deficit in the centroid radius. For a fixed centroid, averaging over the uniform direction a and the radius turns Ra into a standard Gaussian vector, hence

$$\mathbb{E}_{a \sim \sigma} \varphi_Z(ta) = \mathbb{E} e^{-t^2 \|C\|^2/2} = e^{-t^2/2} \mathbb{E} e^{t^2 Q/2}. \quad (6)$$

Lemma 1 (Alignment control). *Let $j(\Delta) = (3 - 2\Delta)^{-1/2} - 2(3 - \Delta)^{-1/2} + 3^{-1/2}$, which is increasing on $[0, 1]$. For every paired law P ,*

$$\mathcal{J}(P) \geq j(\Delta(P)) \geq \frac{\Delta(P)^2}{12\sqrt{3}},$$

hence $\Delta(P) \leq k_0 \sqrt{\mathcal{J}(P)}$ with $k_0 = 2 \cdot 3^{3/4} \approx 4.56$. The first inequality is an equality on an extremal family of pairs at a fixed angle about a uniform centroid direction (Section 4.3).

Proof sketch. Jensen’s inequality over a inside the modulus, (6), and convexity of $x \mapsto e^{t^2 x/2}$ with $\mathbb{E} Q = \Delta$ (both sides are ≥ 1 , so squaring preserves the order) give

$$\begin{aligned} \mathcal{J}(P) &\geq \mathbb{E}_t [e^{-t^2} (\mathbb{E} e^{t^2 Q/2} - 1)^2] \\ &\geq \mathbb{E}_t [e^{-t^2} (e^{t^2 \Delta/2} - 1)^2] = j(\Delta) \\ &\geq \frac{\Delta^2}{4} \mathbb{E}_t [t^4 e^{-t^2}] = \frac{\Delta^2}{12\sqrt{3}}. \end{aligned}$$

Since $j(\Delta) = \Delta^2/(12\sqrt{3}) + O(\Delta^3)$, the alignment floor rises only quadratically, and on the extremal family, which attains it, so does $\mathcal{J}$; this is the source of the square root below.

4.2. Exact Population Minimizers

Proposition 1 (Exact population minimizers). *For every paired law P , $\mathcal{J}(P) = 0$ if and only if $U = V$ almost surely and $U \sim \sigma$.*

The alignment bound already forces agreement; the Gaussian target then determines the distribution of their common direction.

Proof. If $\mathcal{J} = 0$, Lemma 1 gives $U = V$ almost surely. Since $\omega = ta$ covers $\mathbb{R}^d$ and characteristic functions are continuous, $\varphi_Z(\omega) = e^{-\|\omega\|^2/2}$ for all ω , so $Z = RU \sim \mathcal{N}(0, I_d)$; as $R > 0$, $U = Z/\|Z\| \sim \sigma$. The converse is the polar decomposition. $\square$

Thus SG and population InfoNCE have the same minimizers, the aligned, uniform laws. For two-view objectives that test each view separately, the target




Figure 1 | **Theory on controlled families** (Appendix B). (a) InfoNCE excess relative to the leading term $2\beta k_0 \sqrt{\mathcal{J}}$ of Theorem 1 ($d = 16, \beta = 2$): misaligned families attain the sharp constant, the mixture path of Proposition 3 stays at 0.77, and the aligned von Mises–Fisher (vMF) path has excess $\Theta(\mathcal{J})$. (b) The plug-in estimator is minimized away from alignment (dotted), whereas the U-statistic follows $\mathcal{J}$ (extremal family, $d = 16$; markers: Monte Carlo). (c) Along the isotropic channel, pure SG’s optimal misalignment for residual non-uniformity of degree ℓ follows $\rho_\ell \mathcal{J}_{\text{unif}}$ (Theorem 3, $d = 8$; ρ_ℓ as after (11)); InfoNCE’s is zero. (d) With $\mathcal{J}_\alpha$, the rescaled misalignment collapses onto $[1 - \alpha/\alpha_c]_+$, where $\alpha_c = g$ is the channel’s gain.

depends on the test. Let $X_1, X_2 \in \mathbb{R}^d$ be unnormalized view embeddings. Write $I(X_1, X_2)$ for a nonnegative population invariance loss that vanishes exactly when $X_1 = X_2$ almost surely, and $S(X)$ for a nonnegative characteristic population Gaussian discrepancy, integrated over all directions and a full-support frequency law, that vanishes exactly when $X \sim \mathcal{N}(0, I_d)$. By population VICReg we mean a positive weighted sum of the two-view invariance loss, coordinatewise variance hinges with target standard deviation one, and squared off-diagonal covariance penalties, all evaluated at population moments.

Proposition 2 (Directional targets).

- (i) For $\lambda \in (0, 1)$, the idealized population objective $(1-\lambda)I(X_1, X_2) + \frac{\lambda}{2}\{S(X_1) + S(X_2)\}$ is zero if and only if $X_1 = X_2$ almost surely and the common embedding is $\mathcal{N}(0, I_d)$. Its normalized directions therefore have the aligned, uniform target of Proposition 1.
- (ii) Population VICReg’s zero set also contains aligned embeddings uniform on the cross-polytope $\{\pm\sqrt{d}e_i : i \leq d\}$. These embeddings have mean zero and covariance I_d , while their directions are non-uniform.

Thus an idealized population LeJEPa objective (exact invariance, no predictor) has the same normalized target as SG and InfoNCE, whereas population VICReg, which constrains only the first two moments, also admits aligned embeddings with non-uniform directions. Section 7 finds the same contrast in trained encoders.

4.3. A Sharp Comparison with Population InfoNCE

Sharing minimizers gives only qualitative agreement. Under equal marginals, SG also controls the excess population InfoNCE loss.

Theorem 1 (Sharp comparison with population InfoNCE). *Let P have equal marginals and $\beta > 0$. Then*

$$0 \leq \mathcal{L}_\beta(P) - \mathcal{L}_\beta^* \leq 2\beta k_0 \sqrt{\mathcal{J}(P)} + \log(1 + \bar{c}_{d,\beta} \mathcal{J}(P)), \quad (7)$$

where $\bar{c}_{d,\beta} < \infty$ depends only on d and β . Moreover, there is a family of paired laws with uniform marginals such that $\mathcal{J}(P) \rightarrow 0$ and

$$\mathcal{L}_\beta(P) - \mathcal{L}_\beta^* = 2\beta k_0 \sqrt{\mathcal{J}(P)} + o(\sqrt{\mathcal{J}(P)}). \quad (8)$$

Hence both the square-root order and the dimension-free leading constant $2\beta k_0$ are sharp.

Moreover, for laws with equal marginals, $\mathcal{J}(P_n) \rightarrow 0$ if and only if $\mathcal{L}_\beta(P_n) - \mathcal{L}_\beta^* \rightarrow 0$ (Appendix A.11). Explicitly, $\bar{c}_{d,\beta} = (1 + \frac{3}{2}k_0)^2 C_{d,\beta}^* / \kappa_d(\beta)$ with the kernel-comparison constant $C_{d,\beta}^* \leq \sqrt{6} e^{4\beta}$ of Proposition 7. Unlike $2\beta k_0$, $\bar{c}_{d,\beta}$ depends on d and grows at least like $e^{\beta/4 + o(\beta)}$ as $\beta \rightarrow \infty$ at fixed d ; at $d = 8, \beta = 5$ it is about 4.9×10^4 . Replacing $2\beta k_0 \sqrt{\mathcal{J}}$ by $2\beta \Delta^*(\mathcal{J})$, with $\Delta^*(\mathcal{J}) = j^{-1}(\mathcal{J})$ for $\mathcal{J} \leq j(1)$ and 1 otherwise, gives a bound whose first term is exact on the extremal family; aligned laws have $\Delta = 0$ and $\mathcal{J} = D_{\text{lift}}^2(q, \sigma)$, so their excess is $\mathcal{U}_\beta(q) \leq (C_{d,\beta}^* / \kappa_d(\beta)) \mathcal{J}$ (Lemma 3 and Proposition 7; Fig. 1a).

Proof sketch. By (5), it suffices to bound $2\beta \Delta$ and $\mathcal{U}_\beta(q)$. *Alignment:* Lemma 1 gives $\Delta \leq k_0 \sqrt{\mathcal{J}}$. *Transfer:* let $D_{\text{lift}}^2(q, \sigma)$, the lifted discrepancy, be the SG loss of the aligned law with marginal q . Since $\frac{1}{2}(e^{ix^T U} + e^{ix^T V}) = e^{ix^T C} \cos(x^T H)$, the pair test controls the marginal (Theorem 4):

$$|\sqrt{\mathcal{J}(P)} - D_{\text{lift}}(\bar{q}, \sigma)| \leq \frac{3}{2} \Delta(P). \quad (9)$$

Under equal marginals, $\bar{q} = q$. *Kernel comparison:* D_{lift} is an MMD on the sphere whose zonal kernel, like $e^{\beta u^T v}$, diagonalizes in spherical harmonics; comparing eigenvalues degree by degree gives $D_\beta^2 \leq C_{d,\beta}^* D_{\text{lift}}^2$

for the MMD D_β of $e^{\beta u^\top v}$ (Proposition 7). *Uniformity:* by Jensen's inequality and $\mathbb{E}_{W \sim \sigma} e^{\beta u^\top W} = \kappa_d(\beta)$, $\mathbb{E}_q \log \mathbb{E}_q e^{\beta U^\top W} \leq \log \mathbb{E}_{q \otimes q} e^{\beta U^\top W} = \log(\kappa_d(\beta) + D_\beta^2(q, \sigma))$, so $\mathcal{U}_\beta(q) \leq \log(1 + D_\beta^2/\kappa_d(\beta))$ (Lemma 3).

Sharpness. Let (n, h) be a Haar-distributed orthonormal pair in $\mathbb{R}^d$ and, for $s \in [0, 1]$,

$$U_s = \sqrt{s}n + \sqrt{1-s}h, \quad V_s = \sqrt{s}n - \sqrt{1-s}h.$$

Both marginals are uniform, $C_s = \sqrt{s}n$ and $\Delta = 1 - s$, so $Z_s \sim \mathcal{N}(0, sI_d)$ and $\mathcal{J} = j(\Delta)$, so this extremal family attains Lemma 1. Its uniformity excess vanishes, so $\mathcal{L}_\beta - \mathcal{L}_\beta^* = 2\beta\Delta$, and $j(\Delta) = \Delta^2/(12\sqrt{3}) + O(\Delta^3)$ gives (8). The same family is used in Proposition 4 and Fig. 1a,b.

4.4. Optimality of the Square-Root Rate

The square-root rate is optimal for a broad class of pair discrepancies. Call a pair discrepancy F *Hilbertian* (a squared mean-embedding distance) if there exist a bounded feature map Ψ into a Hilbert space $\mathcal{H}$, a target $m^* \in \mathcal{H}$ and independent auxiliary randomness ξ such that $F(P) = \|\mathbb{E}_P \Psi(U, V, \xi) - m^*\|_{\mathcal{H}}^2$. This class contains $\mathcal{J}$, for any radius law and frequency weight, as well as every bounded-kernel squared MMD of a statistic of the pair.

Proposition 3 (Square-root barrier). *Let F be Hilbertian with $F(P_0) = 0$ for an aligned law P_0 with uniform marginals. Let P_1 have uniform marginals and $\Delta(P_1) > 0$, and let $P_p = (1-p)P_0 + pP_1$. Then $F(P_p) = p^2 F(P_1)$, while $\mathcal{L}_\beta(P_p) - \mathcal{L}_\beta^* = 2\beta p \Delta(P_1)$. Hence either F vanishes on a misaligned law, or no bound $\mathcal{L}_\beta - \mathcal{L}_\beta^* \leq \psi(F)$ with $\psi(x) = o(\sqrt{x})$ holds.*

Proof. $F(P_0) = 0$ gives $\mathbb{E}_{P_0} \Psi = m^*$, so $\mathbb{E}_{P_p} \Psi - m^* = p(\mathbb{E}_{P_1} \Psi - m^*)$ and $F(P_p) = p^2 F(P_1)$. The marginals of P_p stay uniform, so $\mathcal{U}_\beta = 0$, and since Δ is linear in P with $\Delta(P_0) = 0$, the excess is $2\beta p \Delta(P_1)$ by (5). If $F(P_1) > 0$, the excess equals $(2\beta \Delta(P_1)/\sqrt{F(P_1)})\sqrt{F(P_p)}$ for all p , which no $\psi(x) = o(\sqrt{x})$ dominates. $\square$

A mean embedding sees a fraction p of misaligned pairs at order p^2 , whereas InfoNCE sees it at order p ; along the mixture path only the coupling changes. Thus no Hilbertian pair discrepancy controls the InfoNCE excess faster than $\sqrt{F}$, and SG attains this rate with the sharp constant of Theorem 1; for SG the order comes from Lemma 1, since $\mathcal{J}$ responds to misalignment only at second order.

4.5. Which Uniformity Tests Control InfoNCE Linearly

Charging misalignment directly removes the barrier. With the alignment price $\mathcal{J}_\alpha = \mathcal{J} + \alpha\Delta$ of Section 6.3, under equal marginals, $\mathcal{L}_\beta - \mathcal{L}_\beta^* \leq c_{\beta,\alpha} \mathcal{J}_\alpha$ for every




Figure 2 | **Condition (U)** ($d = 8, \beta = 5$). (a) Eigenvalues μ_ℓ of zonal squared MMDs relative to those of $e^{\beta u^\top v}$, normalized at $\ell = 1$: the ratio is bounded below for SG's lift and Gaussian kernels with $\gamma \geq \beta/2$, and not for $\gamma = 2$ or moment matching ($\mu_\ell = 0$ for $\ell \geq 3$). (b) Exact $\mathcal{U}_\beta$ over the squared MMD of $e^{-\gamma \|u-v\|^2}$ for a law perturbed in degree ℓ ; bounded in ℓ if and only if $\gamma \geq \beta/2$ (Corollary 1).

$\alpha > 0$, with explicit $c_{\beta,\alpha}$ (Proposition 8). More generally, once misalignment is paid for linearly, a linear certificate needs a test of the marginal that controls InfoNCE's uniformity term $\mathcal{U}_\beta$ linearly. Which uniformity tests do?

Squared MMDs to σ with zonal (rotation-invariant) kernels are classical Sobolev statistics of uniformity; they diagonalize in spherical harmonics, and their weights decide which departures from uniformity they detect (Giné, 1975; García-Portugués and Verdebout, 2018; García-Portugués et al., 2023). $\mathcal{U}_\beta(q) = \int \log(\int k_\beta(\cdot, w) dq(w)) dq - \log \kappa_d(\beta)$, with $k_\beta(u, v) = e^{\beta u^\top v}$, is not of this form, because the logarithm sits inside the outer expectation. Let λ_ℓ^β be the degree- ℓ eigenvalues of k_β , so $\lambda_0^\beta = \kappa_d(\beta)$, and let D_β be the MMD of k_β . For a zonal kernel K with eigenvalues (μ_ℓ) , $\text{MMD}_K^2(q, \sigma) = \sum_{\ell \geq 1} \mu_\ell \|\hat{q}_\ell\|^2$, where $\hat{q}_\ell$ are the harmonic coefficients of q (Appendix A.8).

Theorem 2 (InfoNCE uniformity is spectral). *Fix $d \geq 2$, $\beta > 0$, and write $\kappa = \kappa_d(\beta)$.*

(i) *Let $\delta := \|dq/d\sigma - 1\|_\infty \leq \frac{1}{2}$ and*

$$Q_\beta(q) = \sum_{\ell \geq 1} \frac{\lambda_\ell^\beta}{\kappa} \left(1 - \frac{\lambda_\ell^\beta}{2\kappa}\right) \|\hat{q}_\ell\|^2.$$

Then $\frac{1}{2} D_\beta^2/\kappa \leq Q_\beta \leq D_\beta^2/\kappa$ and $|\mathcal{U}_\beta(q) - Q_\beta(q)| \leq \frac{3}{2} \delta D_\beta^2(q, \sigma)/\kappa$.

- (ii) For a continuous positive-definite zonal kernel K with eigenvalues (μ_ℓ) , the best constant c_K in $\mathcal{U}_\beta \leq c_K \text{MMD}_K^2$ satisfies

$$\sup_{\ell \geq 1} \frac{\lambda_\ell^\beta (1 - \lambda_\ell^\beta / 2\kappa)}{\kappa \mu_\ell} \leq c_K \leq \sup_{\ell \geq 1} \frac{\lambda_\ell^\beta}{\kappa \mu_\ell} =: \frac{S_K}{\kappa}.$$

with the convention $x/0 = \infty$ for $x > 0$.

Part (i) expands the log-partition functional to second order, with a remainder uniform in d and β ; near σ , $\mathcal{U}_\beta$ lies within a factor 2 of the squared MMD with kernel $e^{\beta u^\top v} / \kappa$, up to an $O(\delta)$ error. Part (ii) combines this local necessity with the global bound $\mathcal{U}_\beta \leq \log(1 + D_\beta^2 / \kappa)$ (Lemma 3). A zonal test controls InfoNCE's uniformity term at temperature $1/\beta$ linearly if and only if its spectrum dominates that of $e^{\beta u^\top v}$, that is, $S_K < \infty$. We call this condition (U).

Corollary 1 (Which tests satisfy (U)). *For every $d \geq 2$ and $\beta > 0$:*

- (a) The Gaussian lift of SG: $S_K = C_{d,\beta}^* < \infty$, with $C_{d,\beta}^* / \kappa_d(\beta) \geq e^{\beta/4 + o(\beta)}$ as $\beta \rightarrow \infty$ with d fixed.
- (b) Gaussian kernels $e^{-\gamma \|u-v\|^2}$: $S_K < \infty$ if and only if $\gamma \geq \beta/2$. The same threshold decides linear control of $\mathcal{U}_\beta$ by the excess of the uniformity loss $\log \mathbb{E}_{q \otimes q} e^{-\gamma \|U-U'\|^2}$ (U' an independent copy) of Wang and Isola (2020).
- (c) Heat kernels, band-limited kernels, and first- and second-moment matching: $S_K = \infty$.

Characteristic Gaussian and heat kernels still identify σ when (U) fails, unlike finite-moment tests, but they control $\mathcal{U}_\beta$ only sublinearly, with a defect that grows with the harmonic degree (Fig. 2). The squared MMD with the default $\gamma = 2$ kernel of Wang and Isola (2020), whose uniformity loss it matches to first order near σ up to a constant factor, therefore satisfies (U) exactly for $\tau \geq 1/4$. Since $e^{-\gamma \|u-v\|^2} = e^{-2\gamma} e^{2\gamma u^\top v}$, the threshold says that the kernel must be at least as concentrated as InfoNCE's own kernel $e^{\beta u^\top v}$ (inverse temperature $2\gamma \geq \beta$). SG's lift passes at every temperature, with a constant that grows exponentially in β (at $d = 8$, $\beta = 5$ the supremum in S_K is attained at $\ell = 7$, with $S_K / \kappa \approx 790$), whereas the matched Gaussian kernel $\gamma = \beta/2$ has $S_K / \kappa = e^{\beta} / \kappa_d(\beta) = O(\beta^{(d-1)/2})$; a Gaussian kernel certifies only $\beta \leq 2\gamma$, so the lift trades a larger constant for validity at every temperature. With an alignment term, condition (U) is necessary and sufficient for a linear certificate.

Corollary 2 (Linear certificates). *Let $\alpha > 0$ and let K be a continuous positive-definite zonal kernel. Under equal marginals,*

$$\mathcal{L}_\beta - \mathcal{L}_\beta^* \leq \max \left\{ \frac{2\beta}{\alpha}, \frac{S_K}{\kappa_d(\beta)} \right\} (\alpha \Delta + \text{MMD}_K^2(q, \sigma)),$$

and no bound of this form holds if $S_K = \infty$.

Section 7 tests this distinction in trained encoders.

5. Finite-Batch Estimation

In training, we estimate $\mathcal{J}$ from a batch of B i.i.d. pairs with independent radii, $Z_j = R_j C_j$, and M fresh frequencies $\omega_m = t_m a_m$ with $a_m \sim \sigma$, $t_m \sim \mathcal{N}(0, 1)$. Let $\hat{\phi}_m = \frac{1}{B} \sum_j e^{i\omega_m^\top Z_j}$ and $g_m = e^{-t_m^2/2}$. The plug-in estimator inherits SIGReg's finite-sample bias (Balestriero and LeCun, 2025, Thm. 6); through the centroid, this bias changes the alignment incentive. Proofs are in Appendix A.16.

5.1. Plug-in Bias Toward Misalignment

Proposition 4 (Plug-in bias favors misalignment). *The plug-in estimator $\hat{\mathcal{J}}_V = \frac{1}{M} \sum_m |\hat{\phi}_m - g_m|^2$ satisfies*

$$\mathbb{E} \hat{\mathcal{J}}_V = \mathcal{J}(P) + \frac{1}{B} \mathbb{E}_{a,t} [1 - |\varphi_Z(ta)|^2].$$

For every $B \geq 2$, no aligned uniform law minimizes $\mathbb{E} \hat{\mathcal{J}}_V$: along the extremal family, $\mathbb{E} \hat{\mathcal{J}}_V$ has a unique minimizer, with pair cosine $\mathbb{E} U^\top V = 1 - 4/B + O(B^{-2})$.

The bias comes from the diagonal terms: $\mathbb{E} \hat{\phi}_m = \varphi_Z(\omega_m)$ and $\mathbb{E} |\hat{\phi}_m|^2 = (1 - \frac{1}{B}) |\varphi_Z(\omega_m)|^2 + \frac{1}{B}$, and expanding $|\hat{\phi}_m - g_m|^2$ gives the identity. Shortening the centroid concentrates Z , which raises $|\varphi_Z|$ and lowers the bias term. On the extremal family, $|\varphi_Z(ta)|^2 = e^{-t^2(1-\Delta)}$, so the bias is $\frac{1}{B} [1 - (3 - 2\Delta)^{-1/2}]$, with slope $-3^{-3/2}/B$ at $\Delta = 0$, against $j'(\Delta) \approx \Delta/(6\sqrt{3})$; balancing the two gives $\Delta \approx 2/B$, that is, pair cosine $1 - 4/B$ (Fig. 1b).

5.2. An Unbiased U-Statistic Estimator

Since $\mathbb{E}(1 - |\hat{\phi}_m|^2) = (1 - \frac{1}{B})(1 - |\varphi_Z(\omega_m)|^2)$, removing the diagonal terms gives

$$\hat{\mathcal{J}}_U = \hat{\mathcal{J}}_V - \frac{1}{M(B-1)} \sum_{m=1}^M (1 - |\hat{\phi}_m|^2). \quad (10)$$

Proposition 5 (Unbiased SG estimator). *For every paired law P , $B \geq 2$ and $M \geq 1$, $\mathbb{E} \hat{\mathcal{J}}_U = \mathcal{J}(P)$, where the expectation is over the batch, the radii and the frequencies. $\hat{\mathcal{J}}_U$ is a U-statistic over distinct pairs $j \neq k$ and is computed from the sums $\hat{\phi}_m$ in $O(BdM)$ time and $O(BM + dM)$ memory, without a $B \times B$ similarity matrix.*

The correction is the standard U-statistic; for SG, the bias it removes also shifts the alignment optimum. $\hat{\mathcal{J}}_U$ can be negative on a batch; clipping it would reintroduce bias, so we optimize it directly. In our experiments ($d = 8$, $B = 256$), SG_α (Section 6.3, $\alpha = 0.2$) with $M = B/2$ frequencies reaches the CKA of $M = 256$ and $M = 1024$ in all runs but one vMF seed, while $M = 16$ is too few (Appendix B).

6. Nuisance Preferences and the Alignment Price

SG and InfoNCE share their minimizers but can favor different representations away from the optimum. We exhibit a population failure mode that persists with exact expectations. When the shared code is non-uniform, pure SG can prefer independent per-view nuisance, because the nuisance improves the marginal distributional match faster than SG penalizes the resulting misalignment. Proofs are in [Appendices A.13](#) and [A.15](#).

6.1. An Isotropic Nuisance Channel

Let $W \sim q$ be a shared code on $\mathbb{S}^{d-1}$ and, given W , draw U_ϑ, V_ϑ independently from the heat kernel on $\mathbb{S}^{d-1}$ at time $\vartheta \geq 0$. At $\vartheta = 0$ the views coincide; for $\vartheta > 0$ they carry independent perturbations, with $\Delta(\vartheta) = \frac{1}{2}(1 - e^{-2(d-1)\vartheta})$, and their marginal is smoothed toward σ . Let $\mathcal{J}_q(\vartheta)$ be the SG loss along this channel and $\mathcal{J}_{\text{unif}}(q) = \mathcal{J}_q(0) = D_{\text{lift}}^2(q, \sigma)$.

6.2. Why Pure SG Can Prefer Nuisance

The channel creates two competing effects. Independent perturbations shorten the centroid and hurt alignment, but they also smooth a non-uniform marginal toward the uniform distribution. For pure SG, the latter effect dominates to first order.

Theorem 3 (Nuisance preference of pure SG). *Along the isotropic nuisance channel:*

- (i) *For every $q \neq \sigma$, the right derivative satisfies $\partial_\vartheta \mathcal{J}_q|_{\vartheta=0} = -\sum_{\ell \geq 1} G_\ell \|\hat{q}_\ell\|^2 < 0$, where $\hat{q}_\ell$ is the degree- ℓ harmonic component of $q - \sigma$ and $G_\ell > 0$ is explicit. Thus a small amount of nuisance strictly decreases $\mathcal{J}$, and, for this fixed code q , the aligned law is not a local minimizer of $\mathcal{J}$ along the channel.*
- (ii) *Let $q_\varepsilon = (1 + \varepsilon f)\sigma$ with $f \neq 0$ smooth and $\int f d\sigma = 0$. There is $\vartheta_0(d) > 0$ such that, for all small ε , $\mathcal{J}_{q_\varepsilon}$ has a unique minimizer on $[0, \vartheta_0]$, which is interior, and its misalignment Δ_{eq} there satisfies $\Delta_{\text{eq}}/\mathcal{J}_{\text{unif}}(q_\varepsilon) \rightarrow \rho(f)$, an explicit constant with $\rho(f) > 6\sqrt{3}$.*
- (iii) *In the same setting, $\vartheta = 0$ is the unique minimizer of $\mathcal{L}_\beta - \mathcal{L}_\beta^*$ on $[0, \vartheta_0]$ for all small ε .*

Nuisance helps SG at first order because the centroid direction diffuses at half the speed of the views, which smooths the non-uniformity, and radial contraction reduces the remaining harmonic contributions. Misalignment is cheap, because along the channel the coefficient of Δ^2 is $1/(12\sqrt{3})$ to leading order, as in [Lemma 1](#). In the regime of (ii), therefore,

$$\mathcal{J} \approx \mathcal{J}_{\text{unif}} - g \Delta + \frac{\Delta^2}{12\sqrt{3}}, \quad g = \frac{\rho(f) \mathcal{J}_{\text{unif}}}{6\sqrt{3}}, \quad (11)$$

where $g = -d\mathcal{J}/d\Delta|_{\Delta=0}$ is the channel's gain. InfoNCE pays $2\beta\Delta$ at first order, while its uniformity improvement is of order ε^2 ; near uniformity InfoNCE therefore keeps no nuisance. For f of pure degree ℓ write $\rho_\ell = \rho(f)$; at $d = 8$, $\rho_1, \dots, \rho_4 = 10.8, 28.5, 49.5, 73.8$; at equal $\mathcal{J}_{\text{unif}}$, finer residuals induce more nuisance ([Fig. 1c](#)). The certificate still applies: $\Delta \leq k_0\sqrt{\mathcal{J}}$ bounds the amount of nuisance that pure SG keeps.

6.3. Adding an Alignment Price

This mismatch suggests a minimal modification, SG_α :

$$\mathcal{J}_\alpha(P) = \mathcal{J}(P) + \alpha\Delta(P), \quad \alpha \geq 0. \quad (12)$$

With (11), $\mathcal{J}_\alpha \approx \mathcal{J}_{\text{unif}} + (\alpha - g)\Delta + \Delta^2/(12\sqrt{3})$. If $\alpha > g$, infinitesimal nuisance increases the objective and alignment is locally preferred; if $\alpha < g$, the incentive remains. In the regime of (ii), with α of order ε^2 , the minimizer is $\Delta = [\rho(f)\mathcal{J}_{\text{unif}} - 6\sqrt{3}\alpha]_+ + o(\varepsilon^2)$, so the equilibrium nuisance vanishes for $\alpha \geq g$ ([Corollary 3](#), [Fig. 1d](#)). The price keeps SG's minimizers ([Proposition 8](#)).

The rule applies beyond the isotropic model. For a family $(P_\vartheta)_{\vartheta \in [0, \vartheta_0]}$ of paired laws with $\Delta(P_0) = 0$, along which Δ and an objective F are right-differentiable at 0 with $\Delta'(0) > 0$, call $g = -(dF/d\vartheta)/(d\Delta/d\vartheta)|_0$ the gain of the family, and $\alpha > g$ condition (A). LeJEPa minimizes $\lambda S + (1 - \lambda)I$ where S is the average of the per-view SIGReg terms and I is an invariance term; its reference implementation multiplies S by the batch size B .

Proposition 6 (Condition (A): gain against price). *Let g be the gain of a family as above.*

- (i) *If $\alpha > g$, P_0 is a strict local minimizer of $F + \alpha\Delta$ along the family; if $\alpha < g$, it is not.*
- (ii) *For $\lambda S + (1 - \lambda)I$ with $\lambda \in (0, 1)$, along a family with $I(P_0) = 0$, on which S and I are right-differentiable at 0 with $dI/d\vartheta|_0 > 0$, (i) holds with F replaced by S , Δ by I , α by $(1 - \lambda)/\lambda$, and g by $g_S = -dS/d\vartheta|_0$. Equivalently, for $g_S > 0$, P_0 is a strict local minimizer if $\lambda < \lambda_c = 1/(1 + g_S)$ and is not one if $\lambda > \lambda_c$. Multiplying S by B replaces g_S by Bg_S .*

At $\alpha = g$, higher-order terms decide. For LeJEPa, the price of misalignment is $(1 - \lambda)/\lambda$, set against the gain Bg_S of its batch-scaled regularizer; at fixed per-sample gain, the critical SIGReg weight λ_c therefore decreases with B . The gain depends on the channel and can be measured on a trained encoder ([Section 7](#), [Fig. 3](#)).

Beyond the isotropic channel. If the two views are conditionally independent given a latent, with common conditional mean $c(\Lambda)W$, $c(\Lambda) \in [0, 1]$, for a mean code $W \sim q$ on $\mathbb{S}^{d-1}$, then $|\sqrt{\mathcal{J}(P)} - \sqrt{\mathcal{J}_{\text{unif}}(q)}| \leq 6.5 \Delta(P)$. Hence every such family with fixed q that starts at the

aligned law $U = V = W$ has gain at most $13\sqrt{\mathcal{J}_{\text{unif}}(q)}$ (Proposition 10 and Corollary 4), and heat channels whose diffusion time varies with the code point attain this order (Proposition 9). Condition (A) thus becomes easier to meet as the code becomes uniform.

7. Experiments

We test Sections 4 to 6 in trained encoders. We ask whether SG reaches InfoNCE’s representation when the target is the only difference, whether uniformity tests that satisfy (U) reach InfoNCE’s uniformity, whether pure SG prefers per-view nuisance, and whether the measured gain decides which SG_α and LeJEPa encoders keep it.

Setup. We use latent-variable models with known latent content (Zimmermann et al., 2021; von Kügelgen et al., 2021). In the *content–style* model, a fixed random injective network maps a *content* $c \sim \sigma$ on $\mathbb{S}^7$, shared by the two views, and an independent per-view *style* $s \sim \mathcal{N}(0, I_4)$ to an observation $x = m(c, s)$. An encoder that recovers c up to a measure-preserving map and ignores s is aligned and uniform, so style is a per-view nuisance channel and the content plays the role of the shared code. In the von Mises–Fisher (vMF) model of Zimmermann et al. (2021) there is no style, and the second view’s content is $\tilde{c} \sim \text{vMF}(c, 10)$. We train the same MLP encoder ($d = 8$, $B = 256$, 8k steps, 3 seeds; Appendix B) with InfoNCE ($\beta = 5$), SG and SG_α (with $\hat{\mathcal{J}}_U$), LeJEPa (SIGReg weight λ) and VICReg. To separate the objective from the optimization path, we also switch InfoNCE encoders to SG_α halfway through training (warm start). The *gain* of a trained encoder is measured on its own style channel. We rescale the style input by η and fit $F - F(0) \approx -g\Delta + c_2\Delta^2$ for small η , with $F = \hat{\mathcal{J}}_U$ (for LeJEPa, the per-sample SIGReg term against I), averaging over independent repetitions. Style is *detected* when a held-out probe from (U, c) to the style reaches $R^2 > 0.02$; CKA is linear centered kernel alignment (Kornblith et al., 2019) to InfoNCE encoders of other seeds.

Without per-view nuisance. With noisy positives and no per-view style (vMF), $\text{SG}_{0.2}$ and LeJEPa learn representations close to InfoNCE’s: CKA 0.993 and 0.981, against 0.997 between InfoNCE seeds (Table 1), with content R_c^2 of 0.99 and 0.97 (Table 3). Pure SG follows at 0.938 (0.935 with $\hat{\mathcal{J}}_V$, as the bias of Proposition 4 is $O(1/B)$), and VICReg at 0.811. On content–style, VICReg aligns the views (pair cosine 0.979, Table 2) but remains far from uniform ($\mathcal{U}_\beta = 1.38$ against 0.02), in line with its larger target set (Proposition 2). LeJEPa and VICReg were selected on small grids (Appendix B), so their rows are not fully tuned.

Condition (U) in training. Section 4.5 separates uniformity tests that permit a linear certificate from those

Table 1 | Controlled study ($d = 8$, $\beta = 5$; mean over 3 seeds). CKA to InfoNCE encoders of other seeds; style: held-out probe R^2 (detected if > 0.02); $\mathcal{J}$ and $\mathcal{J}_{0.2} = \mathcal{J} + 0.2\Delta$, in units of 10^{-3} , are evaluated on held-out pairs for every row, whatever the training loss. Warm starts switch InfoNCE to SG or $\text{SG}_{0.2}$ after 4k of 8k steps. Standard deviations: Tables 2 and 3.

Method	vMF	content-style				
	CKA	CKA	style	$\mathcal{U}_\beta$	$\mathcal{J}$	$\mathcal{J}_{0.2}$
<i>from scratch</i>						
InfoNCE	0.997	0.982	−0.02	0.02	0.17	1.7
SG	0.938	0.646	0.11	0.05	0.85	20.6
SG _{0.2}	0.993	0.828	−0.02	0.16	0.98	3.6
LeJEPa, $\lambda = 0.01$	0.981	0.696	−0.02	0.26	1.57	6.0
VICReg	0.811	0.597	−0.02	1.38	95	97
<i>warm start from InfoNCE (4k steps)</i>						
switch point	–	0.969	0.03	0.05	1.27	3.9
→ SG	–	0.962	0.14	0.02	0.12	4.0
→ SG _{0.2}	–	0.975	−0.02	0.04	0.27	2.2

that do not, and we check whether the distinction persists in training. We train the Gaussian uniformity loss of Wang and Isola (2020), with alignment term $\frac{\beta}{2}\mathbb{E}\|U - V\|^2 = 2\beta\Delta$, at $\gamma \in \{1, 2, 2.5, 5\}$ (2 seeds), and first- and second-moment matching with alignment term Δ (3 seeds; Appendix B). All of them align the views (pair cosine 0.97–0.99). With $\gamma \geq \beta/2 = 2.5$ the uniformity excess reaches InfoNCE’s ($\mathcal{U}_\beta = 0.022$ and 0.010, against 0.021 for InfoNCE in these runs), at $\gamma = 2$ it is 0.037, and at $\gamma = 1$ it is 0.18. Moment matching, which fails (U) at every temperature, keeps $\mathcal{U}_\beta = 0.29$, and its residual non-uniformity is 7–63 times InfoNCE’s at every degree $\ell = 3, \dots, 8$ (Fig. 4).

Pure SG prefers nuisance. From scratch, pure SG keeps style ($R^2 = 0.11$; pair cosine 0.80). The warm start isolates the preference. The half-trained InfoNCE encoders carry weak style ($R^2 = 0.03$; two of three seeds), which continuing with InfoNCE or with $\text{SG}_{0.2}$ removes (−0.02); continuing with SG raises it to 0.14. With the same start and budget, the SG branch reaches the lowest $\mathcal{J}$ in Table 1, 0.12×10^{-3} , against 0.27×10^{-3} for the style-free $\text{SG}_{0.2}$ branch and 0.17×10^{-3} for fully trained InfoNCE (Table 1), in line with Theorem 3(i).

The alignment price. Across 38 SG and SG_α encoders, every encoder with detected style has a gain above its price (Fig. 3a). Style is detected in some seeds up to $\alpha = 0.015$ and in none from $\alpha = 0.02$, so the transition lies just below the crossing of the mean gain with α ($\alpha \approx 0.023$). Warm starts compare prices from a common representation. Style is still detected at $\alpha = 0.003$ (mean $R^2 = 0.027$) and is removed for $\alpha \geq 0.01$, so the transition overlaps the range of the warm-started encoders’ own gains, [0.002, 0.008] (Fig. 3c).

Beyond SG: LeJEPa. For LeJEPa, Proposition 6(ii) gives the critical weight $\lambda_c = 1/(1 + Bg_S)$. Across 34 encoders and three batch sizes, style is retained by ex-




Figure 3 | **Gain against price in trained encoders** (content–style model, $d = 8$; filled: style detected). (a) SG_α: style is detected only when the own-channel gain exceeds α . (b) LeJEPa, three batch sizes: style is detected exactly when Bg_S exceeds the price $(1 - \lambda)/\lambda$. (c) Warm starts stay close to InfoNCE; the style added by pure SG vanishes once α exceeds their gains (shaded; dashed: detection threshold).




Figure 4 | **Condition (U) in trained encoders** (alignment term + uniformity term; content–style model, $d = 8$, $\beta = 5$). (a) $\mathcal{U}_\beta$ of encoders trained with the Gaussian uniformity loss at scale γ (one point per seed; blue: $\gamma \geq \beta/2$; amber: $\gamma < \beta/2$; gray line: mean over seeds) and mean over 3 seeds for moment matching (dashed) and InfoNCE (dotted). (b) Residual non-uniformity $\|\hat{q}_\ell\|^2$ per harmonic degree, relative to InfoNCE’s (mean over seeds; dotted: equal to InfoNCE).

actly those encoders whose measured gain Bg_S exceeds the price $(1 - \lambda)/\lambda$ (Fig. 3b). The observed boundary moves from $\lambda \in (0.03, 0.05)$ at $B = 256$ to $(0.1, 0.2)$ at $B = 64$ (two seeds) and to $(0.01, 0.02)$ at $B = 1024$ (one seed). A single weight trades nuisance rejection against uniformity. At $\lambda = 0.01$, LeJEPa discards style but has $\mathcal{U}_\beta = 0.26$ (InfoNCE: 0.02), and at $\lambda = 0.1$ it keeps style ($R^2 = 0.52$; Table 2). The finite-batch bias of the plug-in SIGReg used in training (Proposition 4) changes no classification (Appendix B).

An optimization gap. With per-view nuisance, SG_{0.2} from scratch stays farther from InfoNCE (CKA 0.83) than without nuisance (0.993), although it retains no style, so this gap is separate from the preference of Theorem 3; sharing InfoNCE’s minimizers does not ensure that training reaches InfoNCE’s representation. With a shared optimizer, schedule, budget and peak

learning rate, InfoNCE training reaches a lower value of the SG_{0.2} objective than SG_{0.2} training itself. In units of 10^{-3} , $\mathcal{J}_{0.2} = 1.7$ for InfoNCE against 3.6 for SG_{0.2} from scratch; from the common switch point (3.9), continuing SG_{0.2} reaches 2.2 and keeps InfoNCE’s representation (CKA 0.975; InfoNCE seeds 0.982). Pure SG likewise ends above InfoNCE on its own objective ($\mathcal{J} = 0.85$ against 0.17). More frequencies do not move SG_{0.2} toward InfoNCE (CKA 0.825 at $M = 1024$; Appendix B), and without nuisance the gap nearly closes (vMF, SG_{0.2} against InfoNCE: $\mathcal{J} = 1.42$ and 1.37, $\mathcal{J}_{0.2} = 32.0$ and 31.6; Table 3). SG_{0.2} training thus stops at representations that its own objective scores worse than InfoNCE’s, which suggests that part of the difference lies in optimization.

8. Discussion

A single centroid test shares InfoNCE’s minimizers and controls its excess at the optimal rate among Hilbertian pair discrepancies, yet away from the optimum it can favor per-view nuisance, as our controlled experiments illustrate. Three practical consequences follow. First, a linear guarantee needs two ingredients, an explicit alignment term and a uniformity test whose spectrum dominates that of $e^{\beta u^T v}$; Gaussian kernels with $\gamma \geq \beta/2$ and SG’s lift qualify and moment matching does not; within the Gaussian family, and against moment matching, the difference is visible in trained encoders. Second, the gain of a nuisance channel can be measured on a trained encoder, and comparing it with the alignment weight checks whether that weight rejects the channel. Third, because LeJEPa’s reference implementation scales SIGReg by the batch size, its critical SIGReg weight falls with B , so a weight that rejects nuisance at one batch size can admit it at a larger one.

Limitations and next steps. Our guarantees are population statements (equal marginals, infinitely many

negatives), and the nuisance results are local to a channel. The certificate’s second term has a constant that grows exponentially in β ; at $d = 8$, $\beta = 5$ it is about 4.9×10^4 , so the second term exceeds the leading one for $10^{-6} \lesssim \mathcal{J} \lesssim 2 \times 10^{-2}$, a range that contains the $\mathcal{J}$ of trained encoders. The gains in Section 7 are measured on the trained encoders, so the comparisons with the price check condition (A) at the end of training; predicting retention from a gain measured before training, and testing the rule on natural images and at scale, are next steps, as is identifying the cause of the optimization gap.

References

- Donald E. Amos. Computation of modified Bessel functions and their ratios. *Mathematics of Computation*, 28(125), 1974. doi: 10.1090/S0025-5718-1974-0333287-7.
- Kendall Atkinson and Weimin Han. *Spherical Harmonics and Approximations on the Unit Sphere: An Introduction*, volume 2044 of *Lecture Notes in Mathematics*. Springer, Berlin, Heidelberg, 2012. doi: 10.1007/978-3-642-25983-8.
- Randall Balestriero and Yann LeCun. LeJEPa: Provable and scalable self-supervised learning without the heuristics. *arXiv preprint arXiv:2511.08544*, 2025.
- Adrien Bardes, Jean Ponce, and Yann LeCun. VICReg: Variance-invariance-covariance regularization for self-supervised learning. In *International Conference on Learning Representations (ICLR)*, 2022.
- Ludwig Baringhaus and Norbert Henze. A consistent test for multivariate normality based on the empirical characteristic function. *Metrika*, 35(1), 1988. doi: 10.1007/BF02613322.
- Roy Betser, Eyal Gofer, Meir Yossef Levi, and Guy Gilboa. InfoNCE induces Gaussian distribution. In *International Conference on Learning Representations (ICLR)*, 2026. arXiv:2602.24012.
- Ting Chen, Simon Kornblith, Mohammad Norouzi, and Geoffrey Hinton. A simple framework for contrastive learning of visual representations. In *Proceedings of the 37th International Conference on Machine Learning (ICML)*, volume 119 of *Proceedings of Machine Learning Research*, 2020.
- Zihao Chen, Chi-Heng Lin, Ran Liu, Jingyun Xiao, and Eva L. Dyer. Your contrastive learning problem is secretly a distribution alignment problem. In *Advances in Neural Information Processing Systems (NeurIPS)*, volume 37, 2024. doi: 10.52202/079017-2907.
- Feng Dai and Yuan Xu. *Approximation Theory and Harmonic Analysis on Spheres and Balls*. Springer Monographs in Mathematics. Springer, New York, 2013. doi: 10.1007/978-1-4614-6660-4.
- DLMF. NIST digital library of mathematical functions. <https://dlmf.nist.gov/>, 2026. F. W. J. Olver et al., eds.
- Benoit Dufumier, Carlo Alberto Barbano, Robin Louiset, Edouard Duchesnay, and Pietro Gori. Integrating prior knowledge in contrastive learning with kernel. In *Proceedings of the 40th International Conference on Machine Learning (ICML)*, volume 202 of *Proceedings of Machine Learning Research*, 2023.
- T. W. Epps and Lawrence B. Pulley. A test for normality based on the empirical characteristic function. *Biometrika*, 70(3), 1983. doi: 10.1093/biomet/70.3.723.
- Aleksandr Ermolov, Aliaksandr Siarohin, Enver Sangineto, and Nicu Sebe. Whitening for self-supervised representation learning. In *Proceedings of the 38th International Conference on Machine Learning (ICML)*, volume 139 of *Proceedings of Machine Learning Research*, 2021.
- Xianghong Fang, Jian Li, Qiang Sun, and Benyou Wang. Rethinking the uniformity metric in self-supervised learning. In *International Conference on Learning Representations (ICLR)*, 2024.
- Eduardo García-Portugués and Thomas Verdebout. An overview of uniformity tests on the hypersphere. *arXiv preprint arXiv:1804.00286*, 2018.
- Eduardo García-Portugués, Paula Navarro-Esteban, and Juan A. Cuesta-Albertos. On a projection-based class of uniformity tests on the hypersphere. *Bernoulli*, 29(1), 2023. doi: 10.3150/21-BEJ1454.
- Quentin Garrido, Yubei Chen, Adrien Bardes, Laurent Najman, and Yann LeCun. On the duality between contrastive and non-contrastive self-supervised learning. In *International Conference on Learning Representations (ICLR)*, 2023.
- Evarist M. Giné. Invariant tests for uniformity on compact Riemannian manifolds based on Sobolev norms. *The Annals of Statistics*, 3(6), 1975. doi: 10.1214/aos/1176343283.
- Arthur Gretton, Karsten M. Borgwardt, Malte J. Rasch, Bernhard Schölkopf, and Alexander Smola. A kernel two-sample test. *Journal of Machine Learning Research*, 13(25), 2012.
- Jeff Z. HaoChen, Colin Wei, Adrien Gaidon, and Tengyu Ma. Provable guarantees for self-supervised deep learning with spectral contrastive loss. In *Advances in Neural Information Processing Systems (NeurIPS)*, volume 34, 2021.
- Robert Jenkinson Alvarez. Beyond isotropy in JEPAs: Hamiltonian geometry and symplectic prediction. *arXiv preprint arXiv:2605.20107*, 2026.
- David Klindt, Yann LeCun, and Randall Balestriero. When does LeJEPa learn a world model? *arXiv preprint arXiv:2605.26379*, 2026.
- Simon Kornblith, Mohammad Norouzi, Honglak Lee, and Geoffrey Hinton. Similarity of neural network representations revisited. In *Proceedings of the 36th International Conference on Machine Learning (ICML)*, volume 97 of *Proceedings of Machine Learning Research*, 2019.
- Yazhe Li, Roman Pogodin, Danica J. Sutherland, and Arthur Gretton. Self-supervised learning with kernel dependence maximization. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2021.

- Fabian A. Mikulasch and Friedemann Zenke. Understanding self-supervised learning via latent distribution matching. In *Proceedings of the 43rd International Conference on Machine Learning (ICML)*, 2026. arXiv:2605.03517.
- Léo Nicollier, Max Dunitz, Marc Pic, Pablo Musé, Enric Meinhardt-Llopis, and Gabriele Facciolo. SPHERE-JEPA: Spherical prediction with homogeneous embeddings. *arXiv preprint arXiv:2605.26900*, 2026a.
- Léo Nicollier, Enric Meinhardt-Llopis, Max Dunitz, Marc Pic, Pablo Musé, and Gabriele Facciolo. Expanding SPHERE-JEPA: A family of statistical regularizers for the hypersphere. *arXiv preprint arXiv:2606.17603*, 2026b.
- Léo Nicollier, Enric Meinhardt-Llopis, Marc Pic, Pablo Musé, and Gabriele Facciolo. Beyond Gaussian worlds: Latent geometry matters for JEPAs. *arXiv preprint arXiv:2609.21656*, 2026c.
- Léo Nicollier, Marc Pic, Pablo Musé, Enric Meinhardt-Llopis, and Gabriele Facciolo. Unbiased open world regularization for fair self-supervised learning. *arXiv preprint arXiv:2607.22149*, 2026d.
- Mikhail Parakhin, André M. Carvalho, and Patrick Haluptzok. The wristband Gaussian loss: Deterministic, composable latents via a sphere–interval decomposition. *arXiv preprint arXiv:2605.08749*, 2026.
- Evgenia Rusak, Patrik Reizinger, Attila Juhos, Oliver Bringmann, Roland S. Zimmermann, and Wieland Brendel. InfoNCE: Identifying the gap between theory and practice. In *Proceedings of the 28th International Conference on Artificial Intelligence and Statistics (AISTATS)*, volume 258 of *Proceedings of Machine Learning Research*, 2025.
- Nikunj Saunshi, Orestis Plevrakis, Sanjeev Arora, Mikhail Khodak, and Hrishikesh Khandeparkar. A theoretical analysis of contrastive unsupervised representation learning. In *Proceedings of the 36th International Conference on Machine Learning (ICML)*, volume 97 of *Proceedings of Machine Learning Research*, 2019.
- Rylan Schaeffer, Victor Lecomte, Dhruv Bhandarkar Pai, Andres Carranza, Berivan Isik, Alyssa Unell, Mikail Khona, Thomas Yerxa, Yann LeCun, SueYeon Chung, Andrey Gromov, Ravid Shwartz-Ziv, and Sanmi Koyejo. Towards an improved understanding and utilization of maximum manifold capacity representations. *arXiv preprint arXiv:2406.09366*, 2024.
- I. J. Schoenberg. Positive definite functions on spheres. *Duke Mathematical Journal*, 9(1), 1942. doi: 10.1215/S0012-7094-42-00908-6.
- Aaron van den Oord, Yazhe Li, and Oriol Vinyals. Representation learning with contrastive predictive coding. *arXiv preprint arXiv:1807.03748*, 2018.
- Julius von Kügelgen, Yash Sharma, Luigi Gresele, Wieland Brendel, Bernhard Schölkopf, Michel Besserve, and Francesco Locatello. Self-supervised learning with data augmentations provably isolates content from style. In *Advances in Neural Information Processing Systems (NeurIPS)*, volume 34, 2021.
- Feng Wang and Huaping Liu. Understanding the behaviour of contrastive loss. In *IEEE/CVF Conference on Computer Vision and Pattern Recognition (CVPR)*, 2021. doi: 10.1109/CVPR46437.2021.00252.
- Tongzhou Wang and Phillip Isola. Understanding contrastive representation learning through alignment and uniformity on the hypersphere. In *Proceedings of the 37th International Conference on Machine Learning (ICML)*, volume 119 of *Proceedings of Machine Learning Research*, 2020.
- Haiyu Wu, Randall Balestriero, and Morgan Levine. VISReg: Variance-invariance-sketching regularization for JEPA training. *arXiv preprint arXiv:2606.02572*, 2026.
- Thomas Yerxa, Yilun Kuang, Eero P. Simoncelli, and SueYeon Chung. Learning efficient coding of natural images with maximum manifold capacity representations. In *Advances in Neural Information Processing Systems (NeurIPS)*, volume 36, 2023. doi: 10.52202/075280-1047.
- Zhoushun Yu, Xiaoyu Hu, and Xiangyu Xu. QQWorld: Quantile-quantile matching for world model regularization. *arXiv preprint arXiv:2607.28415*, 2026.
- Jure Zbontar, Li Jing, Ishan Misra, Yann LeCun, and Stéphane Deny. Barlow Twins: Self-supervised learning via redundancy reduction. In *Proceedings of the 38th International Conference on Machine Learning (ICML)*, volume 139 of *Proceedings of Machine Learning Research*, 2021.
- Léon Zheng, Gilles Puy, Elisa Riccietti, Patrick Pérez, and Rémi Gribonval. Self-supervised learning with rotation-invariant kernels. In *International Conference on Learning Representations (ICLR)*, 2023.
- Eric Zimmermann, Harley Wiltzer, Justin Szeto, David Alvarez-Melis, and Lester Mackey. KerJEPA: Kernel discrepancies for Euclidean self-supervised learning. *arXiv preprint arXiv:2512.19605*, 2025.
- Roland S. Zimmermann, Yash Sharma, Steffen Schneider, Matthias Bethge, and Wieland Brendel. Contrastive learning inverts the data generating process. In *Proceedings of the 38th International Conference on Machine Learning (ICML)*, volume 139 of *Proceedings of Machine Learning Research*, 2021.

Appendix

A. Proofs

A.1. Auxiliary statements

The following statements are cited in the main text; their proofs are given below.

Theorem 4 (Structure). *For every paired law, $|\sqrt{\mathcal{J}(P)} - D_{\text{lift}}(\bar{q}, \sigma)| \leq \frac{3}{2}\Delta(P)$.*

Proof in [Appendix A.6](#).

Proposition 7 (Lifted kernel). *Let $\lambda_\ell^{\text{lift}}$ be the eigenvalues of K_{lift} and $C_{d,\beta}^* = \sup_{\ell \geq 1} \lambda_\ell^\beta / \lambda_\ell^{\text{lift}}$. Then $D_\beta^2 \leq C_{d,\beta}^* D_{\text{lift}}^2$, no smaller constant works, and $C_{d,\beta}^* \leq \sqrt{6} e^\beta \max_{\ell \geq 1} (3\beta)^\ell / \ell! \leq \sqrt{6} e^{4\beta}$. Moreover, for fixed d , $\liminf_{\beta \rightarrow \infty} \beta^{-1} \log(C_{d,\beta}^* / \kappa_d(\beta)) \geq \frac{1}{4}$.*

Proof in [Appendix A.10](#). For $d = 8$ and $\beta = 5$, the setting of [Section 7](#), the supremum is attained at $\ell = 7$ and $C_{d,\beta}^* / \kappa_d(\beta) \approx 790$, so $\bar{c}_{d,\beta} \approx 4.9 \times 10^4$ in (7); the leading term then dominates for $\mathcal{J} \lesssim 10^{-6}$.

Proposition 8 (Calibrated shared loss). *For $\alpha > 0$, $\mathcal{J}_\alpha = 0$ if and only if $\mathcal{J} = 0$, and under equal marginals $\mathcal{L}_\beta(P) - \mathcal{L}_\beta^* \leq c_{\beta,\alpha} \mathcal{J}_\alpha(P)$ with $c_{\beta,\alpha} = \max\{(2\beta + \frac{9}{2}c_\beta)/\alpha, 2c_\beta\}$ and $c_\beta = C_{d,\beta}^* / \kappa_d(\beta)$.*

Corollary 3 (Calibration threshold). *In the setting of [Theorem 3\(ii\)](#), let $\mathcal{J}_\alpha = \mathcal{J} + \alpha\Delta$ with $\alpha = a\varepsilon^2$ for a fixed $a \geq 0$. Then $\mathcal{J}_\alpha$ has a unique minimizer on $[0, \vartheta_0]$ for small ε , and*

$$\Delta(\vartheta_\alpha^*) = [\rho(f) \mathcal{J}_{\text{unif}}(q_\varepsilon) - 6\sqrt{3}\alpha]_+ + o(\varepsilon^2).$$

In particular $\vartheta = 0$ is the unique minimizer if and only if $\alpha \geq \alpha_c := \rho(f) \mathcal{J}_{\text{unif}}(q_\varepsilon) / (6\sqrt{3})$; for any fixed $\alpha > 0$ this holds for all small ε .

Proofs in [Appendices A.13](#) and [A.14](#).

A.2. Preliminaries

Throughout, $D = d/2$, $a \sim \sigma$, $t \sim \mathcal{N}(0, 1)$, $R \sim \chi_d$ are mutually independent and independent of the pair (U, V) , and $g(t) = e^{-t^2/2}$. We write $\|F\|$ for the norm of $L^2(\sigma \otimes \mathcal{N}(0, 1))$ applied to a function $F(a, t)$, so that $\mathcal{J} = \|\varphi_Z(t \cdot) - g\|^2$. For a probability measure μ on $\mathbb{S}^{d-1}$, $T_\mu = RW$ with $W \sim \mu$ independent of R , so $T_\sigma \sim \mathcal{N}(0, I_d)$ and $Z = T_q$ when $U = V \sim q$. With the feature map $\psi_u(a, t) = \mathbb{E}_R e^{itRa^\top u}$ we have $\varphi_{T_\mu}(ta) = \int \psi_u(a, t) d\mu(u)$; the Gaussian lift is the kernel $K_{\text{lift}}(u, v) = \langle \psi_u, \psi_v \rangle$ (Hermitian inner product; K_{lift} is real since $\psi_u(-a, t) = \overline{\psi_u(a, t)}$), and $D_{\text{lift}}(\mu, \sigma) = \|\varphi_{T_\mu}(t \cdot) - g\|$ is the corresponding MMD ([Lemma 5](#)). Symbols are collected in [Appendix C](#). We use the following standard facts.

- (F1) *Gaussian polar decomposition.* $Ra \sim \mathcal{N}(0, I_d)$. Conversely, if $G \sim \mathcal{N}(0, I_d)$ then $\|G\| \sim \chi_d$ and $G/\|G\| \sim \sigma$ are independent.
- (F2) For $c \in \mathbb{R}^d$ and $t \in \mathbb{R}$: $\mathbb{E}_{a,R} e^{itRa^\top c} = e^{-t^2 \|c\|^2/2}$ (by (F1)).
- (F3) For $\theta \geq 0$ and integer $\ell \geq 0$: $\mathbb{E} e^{-\theta t^2} = (1 + 2\theta)^{-1/2}$ and $\mathbb{E} t^{2\ell} e^{-\theta t^2} = (2\ell - 1)!! (1 + 2\theta)^{-\ell-1/2}$, with $(-1)!! = 1$. In particular $\mathbb{E} t^4 e^{-t^2} = 3^{-3/2}$ and $\mathbb{E} t^4 = 3$.
- (F4) For symmetric $A \in \mathbb{R}^{d \times d}$: $\mathbb{E}_a (a^\top A a)^2 = \frac{(\text{tr} A)^2 + 2 \text{tr}(A^2)}{d(d+2)}$.
- (F5) $\mathbb{E} R^p = 2^{p/2} \Gamma(D + p/2) / \Gamma(D)$ for $p > -d$; in particular $\mathbb{E} R^2 = d$.

We also use $H = \frac{1}{2}(U - V)$, $C = \frac{1}{2}(U + V)$, for which $U = C + H$, $V = C - H$, $C^\top H = 0$, $\|C\|^2 + \|H\|^2 = 1$, and $\mathbb{E} \|H\|^2 = \Delta$.

A.3. Proof of Lemma 1 (alignment control)

Let $Q = \|H\|^2 = 1 - \|C\|^2 \in [0, 1]$, so $\mathbb{E}Q = \Delta$. Fix t . Conditioning on (U, V) and applying (F2),

$$\mathbb{E}_a \varphi_Z(ta) = \mathbb{E}_{U,V} \mathbb{E}_{a,R} e^{itRa^\top C} = \mathbb{E} e^{-t^2 \|C\|^2 / 2} = g(t) \mathbb{E} e^{t^2 Q / 2} \geq g(t) e^{t^2 \Delta / 2},$$

by Jensen's inequality for the convex map $x \mapsto e^{t^2 x / 2}$. Hence $\mathbb{E}_a \varphi_Z(ta) - g(t) \geq g(t)(e^{t^2 \Delta / 2} - 1) \geq 0$. By Jensen's inequality for $w \mapsto |w|^2$ on $\mathbb{C}$,

$$\mathbb{E}_a |\varphi_Z(ta) - g(t)|^2 \geq |\mathbb{E}_a \varphi_Z(ta) - g(t)|^2 \geq e^{-t^2} (e^{t^2 \Delta / 2} - 1)^2 = e^{-(1-\Delta)t^2} - 2e^{-(1-\Delta/2)t^2} + e^{-t^2}.$$

Taking $\mathbb{E}_t$ with (F3) gives $\mathcal{J} \geq \mathbb{E}_t e^{-t^2} (e^{t^2 \Delta / 2} - 1)^2 = (3 - 2\Delta)^{-1/2} - 2(3 - \Delta)^{-1/2} + 3^{-1/2} = j(\Delta)$. Since $e^x - 1 \geq x$, $j(\Delta) \geq \frac{\Delta^2}{4} \mathbb{E} t^4 e^{-t^2} = \frac{\Delta^2}{12\sqrt{3}}$ by (F3). Moreover $j'(\Delta) = (3 - 2\Delta)^{-3/2} - (3 - \Delta)^{-3/2} > 0$ on $(0, 1]$, so j is increasing, with $j(1) = 1 - \sqrt{2} + 3^{-1/2} \approx 0.163$. If $\mathcal{J} \leq j(1)$, this gives $\Delta \leq j^{-1}(\mathcal{J}) = \Delta^*(\mathcal{J})$; if $\mathcal{J} > j(1)$, then $\Delta \leq 1 = \Delta^*(\mathcal{J})$. In both cases $\Delta^*(\mathcal{J}) \leq k_0 \sqrt{\mathcal{J}}$, because $j(\Delta) \geq \Delta^2 / k_0^2$ and $k_0 \sqrt{j(1)} > 1$. On the family of Section 4.3, $Q \equiv \Delta$, so the first Jensen step is an equality. Also $\varphi_Z(ta) = e^{-(1-\Delta)t^2/2}$ does not depend on a , so the second one is too. $\square$

A.4. Proof of Proposition 1

($\Rightarrow$) If $\mathcal{J} = 0$, Lemma 1 gives $\Delta = 0$, i.e., $\mathbb{E}\|U - V\|^2 = 0$, so $U = V$ a.s. The function $(a, t) \mapsto \varphi_Z(ta) - g(t)$ is continuous on $\mathbb{S}^{d-1} \times \mathbb{R}$, and its squared modulus integrates to zero against $\sigma \otimes \mathcal{N}(0, 1)$, a measure with full support. Hence it vanishes identically. Since $\{ta\} = \mathbb{R}^d$, $\varphi_Z(\omega) = e^{-\|\omega\|^2/2}$ for all ω , and $Z \sim \mathcal{N}(0, I_d)$ by uniqueness of characteristic functions. As $U = V$, $Z = RU$ with $R > 0$ a.s., so $U = Z/\|Z\| \sim \sigma$ by (F1). ($\Leftarrow$) If $U = V \sim \sigma$, then $Z = RU \sim \mathcal{N}(0, I_d)$ by (F1), and $\mathcal{J} = 0$. $\square$

A.5. Proof of Proposition 2 (directional targets)

Proof. (i) Both terms are nonnegative and both weights are positive. A zero value therefore gives $I(X_1, X_2) = 0$ and $S(X_j) = 0$ for $j = 1, 2$. By the defining properties of I and the characteristic population discrepancy S , $X_1 = X_2$ almost surely and their common law is $\mathcal{N}(0, I_d)$. The converse is immediate. The Gaussian polar decomposition then gives aligned, uniform normalized directions.

(ii) Let $X_1 = X_2 = X$ and draw X uniformly from $\{\pm\sqrt{d} e_i : i \leq d\}$. Then $\mathbb{E}X = 0$, $\mathbb{E}XX^\top = I_d$, and the two views coincide. Hence the invariance, variance-hinge (target standard deviation one), and off-diagonal covariance terms all vanish. The direction $X/\|X\|$ has finite support and is not σ . $\square$

A.6. The centroid identity and the proof of Theorem 4

Lemma 2 (Centroid versus mixture). *For every paired law, with $T = T_{\tilde{q}}$,*

$$\|\varphi_Z(t \cdot) - \varphi_T(t \cdot)\| \leq \frac{3}{2} \sqrt{\frac{d}{d+2}} \Delta \leq \frac{3}{2} \Delta.$$

Proof. For every $x \in \mathbb{R}^d$, $\frac{1}{2}(e^{ix^\top U} + e^{ix^\top V}) = e^{ix^\top C} \cos(x^\top H)$. Put $x = tRa$ and take expectations over (U, V, R) . Since $T = RW$ with $W \sim \tilde{q} = \frac{1}{2}(q_U + q_V)$,

$$\varphi_T(ta) = \frac{1}{2}(\mathbb{E}e^{itRa^\top U} + \mathbb{E}e^{itRa^\top V}) = \mathbb{E}[e^{itRa^\top C} \cos(tRa^\top H)].$$

Hence $\varphi_T(ta) - \varphi_Z(ta) = \mathbb{E}[e^{itRa^\top C}(\cos(tRa^\top H) - 1)]$. Using $0 \leq 1 - \cos y \leq y^2/2$, independence of R , and (F5),

$$|\varphi_T(ta) - \varphi_Z(ta)| \leq \frac{1}{2} t^2 \mathbb{E}[R^2(a^\top H)^2] = \frac{d}{2} t^2 a^\top \Sigma_H a, \quad \Sigma_H = \mathbb{E}HH^\top \succeq 0, \quad \text{tr } \Sigma_H = \Delta.$$

Squaring, integrating and using (F3)–(F4),

$$\|\varphi_T - \varphi_Z\|^2 \leq \frac{d^2}{4} \cdot 3 \cdot \frac{(\text{tr } \Sigma_H)^2 + 2 \text{tr}(\Sigma_H^2)}{d(d+2)} \leq \frac{9d}{4(d+2)} \Delta^2,$$

because $\text{tr}(\Sigma_H^2) \leq (\text{tr } \Sigma_H)^2$ for $\Sigma_H \succeq 0$. $\square$

Proof of Theorem 4. By the reverse triangle inequality in L^2 and Lemma 2, $|\sqrt{\mathcal{J}} - D_{\text{lift}}(\tilde{q}, \sigma)| = \|\varphi_Z - g\| - \|\varphi_T - g\| \leq \|\varphi_Z - \varphi_T\| \leq \frac{3}{2} \Delta$. $\square$

A.7. InfoNCE: decomposition and uniqueness of the uniform minimizer

For a probability measure q on $\mathbb{S}^{d-1}$ let $g_q(u) = \int e^{\beta u^\top w} dq(w) \in [e^{-\beta}, e^\beta]$ and $\Phi(q) = \int \log g_q dq$, so that $\mathcal{U}_\beta(q) = \Phi(q) - \log \kappa_d(\beta)$.

Lemma 3 (Decomposition). *If P has equal marginals q , then $\mathcal{L}_\beta(P) - \mathcal{L}_\beta^* = 2\beta\Delta + \mathcal{U}_\beta(q)$ and $\mathcal{U}_\beta(q) \leq \log(1 + D_\beta^2(q, \sigma)/\kappa_d(\beta))$.*

Proof. $-\beta \mathbb{E} U^\top V + \beta = \beta(1 - \mathbb{E} U^\top V) = 2\beta\Delta$, which gives the identity. Since $\int e^{\beta u^\top w} d\sigma(w) = \kappa_d(\beta)$ for every u ,

$$D_\beta^2(q, \sigma) = \iint k_\beta dq dq - 2 \iint k_\beta dq d\sigma + \iint k_\beta d\sigma d\sigma = \int g_q dq - \kappa_d(\beta). \quad (13)$$

By Jensen's inequality for the concave logarithm, $\Phi(q) \leq \log \int g_q dq = \log(\kappa_d(\beta) + D_\beta^2(q, \sigma))$. $\square$

Wang and Isola (2020, Thm. 1) show that perfectly uniform encoders minimize $\mathcal{U}_\beta$; the following lemma gives a self-contained proof that σ is its unique minimizer over all laws.

Lemma 4 (Uniform is the unique minimizer). *Φ is continuous for the weak topology, and $\mathcal{U}_\beta(q) \geq 0$ for every q , with equality if and only if $q = \sigma$.*

Proof. Continuity. Since $|e^{\beta u^\top w} - e^{\beta u'^\top w}| \leq \beta e^\beta \|u - u'\|$, the family $\{g_q\}$ is uniformly equicontinuous and uniformly bounded. If $q_n \rightarrow q$, then $g_{q_n} \rightarrow g_q$ pointwise and hence uniformly on the compact sphere (Arzelà–Ascoli). As $\log$ is Lipschitz on $[e^{-\beta}, e^\beta]$, $|\Phi(q_n) - \Phi(q)| \leq e^\beta \|g_{q_n} - g_q\|_\infty + \int |\log g_q d(q_n - q)| \rightarrow 0$.

Existence and structure of minimizers. The set of probability measures on $\mathbb{S}^{d-1}$ is weakly compact, so Φ attains its minimum at some q . Let $m = \min_{u \in \text{supp } q} g_q(u)$, which exists by continuity. Fix $\varepsilon > 0$ and let $S_\varepsilon = \{u : g_q(u) < m + \varepsilon\}$, an open set with $q(S_\varepsilon) > 0$. Let $q_S = q(\cdot \cap S_\varepsilon)/q(S_\varepsilon)$ and $q_\eta = (1 - \eta)q + \eta q_S$ for $\eta \in [0, 1]$. Then $g_{q_\eta} = g_q + \eta(g_{q_S} - g_q)$ takes values in $[e^{-\beta}, e^\beta]$, and $\eta \mapsto \Phi(q_\eta) = \int \log g_{q_\eta} dq_\eta$ is differentiable, with right derivative at 0

$$\frac{d}{d\eta} \Phi(q_\eta) \Big|_{\eta=0^+} = \int \frac{g_{q_S}}{g_q} dq - 1 + \int \log g_q dq_S - \Phi(q).$$

Minimality of q forces this derivative to be ≥ 0 . Now $g_q \geq m$ q -a.e., so by Fubini and the symmetry of k_β ,

$$\int \frac{g_{q_S}}{g_q} dq = \iint \frac{k_\beta(u, v)}{g_q(u)} dq(u) dq_S(v) \leq \frac{1}{m} \int g_q dq_S \leq \frac{m + \varepsilon}{m},$$

and $\int \log g_q dq_S \leq \log(m + \varepsilon)$. Therefore $\Phi(q) \leq \frac{\varepsilon}{m} + \log(m + \varepsilon)$ for every $\varepsilon > 0$, so $\Phi(q) \leq \log m$. But $\Phi(q) = \int \log g_q dq \geq \log m$. Hence $\int (\log g_q - \log m) dq = 0$ with a nonnegative integrand, and $g_q = m$ q -a.e. Integrating and using (13), $m = \int g_q dq = \kappa_d(\beta) + D_\beta^2(q, \sigma)$, so

$$\min \Phi = \Phi(q) = \log(\kappa_d(\beta) + D_\beta^2(q, \sigma)) \geq \log \kappa_d(\beta) = \Phi(\sigma),$$

where $\Phi(\sigma) = \log \kappa_d(\beta)$ because $g_\sigma \equiv \kappa_d(\beta)$. Thus $\min \Phi = \log \kappa_d(\beta)$, i.e., $\mathcal{U}_\beta \geq 0$. Moreover, every minimizer satisfies $D_\beta(q, \sigma) = 0$, hence $q = \sigma$, because k_β is characteristic on $\mathbb{S}^{d-1}$ (all its eigenvalues λ_ℓ^β with $\ell \geq 1$ are positive, Lemma 6). $\square$

A.8. Spectral analysis of the two kernels

Let $\mathcal{H}_\ell$ be the space of degree- ℓ spherical harmonics on $\mathbb{S}^{d-1}$, of dimension $N_\ell = \frac{(2\ell+d-2)\Gamma(\ell+d-2)}{\Gamma(\ell+1)\Gamma(d-1)}$ ($N_0 = 1$), with an $L^2(\sigma)$ -orthonormal basis $\{Y_{\ell m}\}_{m \leq N_\ell}$. Let P_ℓ be the Gegenbauer polynomial normalized by $P_\ell(1) = 1$ (for $d = 2$, use $P_\ell(\cos \theta) = \cos(\ell \theta)$ and $N_\ell = 2$ for $\ell \geq 1$); then $|P_\ell| \leq 1$ on $[-1, 1]$ and the addition theorem reads $\sum_m Y_{\ell m}(u) Y_{\ell m}(v) = N_\ell P_\ell(u^\top v)$ (Dai and Xu, 2013; Atkinson and Han, 2012). Put $\nu = D - 1$.

Zonal kernels and MMD. By Schoenberg's theorem (Schoenberg, 1942), a continuous positive-definite zonal kernel has an expansion $K(u, v) = \sum_{\ell \geq 0} \lambda_\ell N_\ell P_\ell(u^\top v)$ with $\lambda_\ell \geq 0$ and $\sum_\ell \lambda_\ell N_\ell = K(u, u) < \infty$, converging absolutely and uniformly. Here $\lambda_\ell = \iint K(u, v) Y(u) Y(v) d\sigma d\sigma$ for any unit-norm $Y \in \mathcal{H}_\ell$. By uniform convergence, for every probability measure q ,

$$\text{MMD}_K^2(q, \sigma) = \sum_{\ell \geq 1} \lambda_\ell \|\hat{q}_\ell\|^2, \quad \|\hat{q}_\ell\|^2 := \sum_m \left| \int Y_{\ell m} dq \right|^2, \quad (14)$$

where the $\ell = 0$ term drops because $q - \sigma$ has zero mass. Since a finite measure on $\mathbb{S}^{d-1}$ is determined by its harmonic coefficients, K is characteristic if and only if $\lambda_\ell > 0$ for all $\ell \geq 1$.

Funk–Hecke formulas. For $Y \in \mathcal{H}_\ell$, $a \in \mathbb{S}^{d-1}$, $s \in \mathbb{R}$ and $\beta > 0$ (see, e.g., Dai and Xu, 2013; Atkinson and Han, 2012),

$$\int e^{is^\top u} Y(u) d\sigma(u) = i^\ell b_\ell(s) Y(a), \quad b_\ell(s) = \Gamma(D) \sum_{k \geq 0} \frac{(-1)^k (s/2)^{2k+\ell}}{k! \Gamma(k + \ell + D)}, \quad (15)$$

$$\int e^{\beta a^\top u} Y(u) d\sigma(u) = \lambda_\ell^\beta Y(a), \quad \lambda_\ell^\beta = \Gamma(D) \sum_{k \geq 0} \frac{(\beta/2)^{2k+\ell}}{k! \Gamma(k + \ell + D)} = \Gamma(D) \left(\frac{2}{\beta}\right)^\nu I_{\ell+\nu}(\beta). \quad (16)$$

For $s > 0$, $b_\ell(s) = \Gamma(D)(2/s)^\nu J_{\ell+\nu}(s)$. In (15), b_ℓ is an entire function with parity $(-1)^\ell$, and J denotes a Bessel function, unrelated to the loss $\mathcal{J}$. Formula (16) identifies the eigenvalues of k_β , with $\lambda_0^\beta = \kappa_d(\beta)$.

Lemma 5 (Lifted eigenvalues). K_{lift} is a continuous positive-definite zonal kernel with $\text{MMD}_{K_{\text{lift}}}(q, \sigma) = D_{\text{lift}}(q, \sigma)$. Its eigenvalues are $\lambda_\ell^{\text{lift}} = \mathbb{E}_t h_\ell(t)^2$, where

$$h_\ell(t) = \mathbb{E}_R b_\ell(tR) = c_\ell t^\ell {}_1F_1(D + \frac{\ell}{2}; D + \ell; -\frac{t^2}{2}) = c_\ell t^\ell e^{-t^2/2} {}_1F_1(\frac{\ell}{2}; D + \ell; \frac{t^2}{2}), \quad c_\ell = \frac{\Gamma(D + \ell/2)}{2^{\ell/2} \Gamma(D + \ell)}.$$

Proof. Continuity follows from dominated convergence and positive-definiteness from the feature-map form $K_{\text{lift}}(u, v) = \langle \psi_u, \psi_v \rangle_{L^2}$. Zonality holds because $\psi_{Ou}(a, t) = \psi_u(O^\top a, t)$ for orthogonal O and σ is rotation invariant. Since $\varphi_{T_q}(ta) = \int \psi_u(a, t) dq(u)$ and $\varphi_{T_\sigma}(ta) = g(t)$, expanding the square gives $D_{\text{lift}}^2(q, \sigma) = \iint K_{\text{lift}} d(q - \sigma)^{\otimes 2}$. For unit-norm $Y \in \mathcal{H}_\ell$, Fubini (all integrands are bounded) and (15) give

$$\int \psi_u(a, t) Y(u) d\sigma(u) = \mathbb{E}_R \int e^{itRa^\top u} Y(u) d\sigma(u) = i^\ell \mathbb{E}_R b_\ell(tR) Y(a) = i^\ell h_\ell(t) Y(a),$$

so $\lambda_\ell^{\text{lift}} = \mathbb{E}_{a,t} |h_\ell(t) Y(a)|^2 = \mathbb{E}_t h_\ell(t)^2$. For the series, integrate (15) termwise against the χ_d law using (F5). This is legitimate because $\sum_k \mathbb{E} R^{2k+\ell} (|t|/2)^{2k+\ell} / (k! \Gamma(k + \ell + D)) < \infty$, the ratio of consecutive terms tending to zero. It gives

$$h_\ell(t) = \frac{t^\ell}{2^{\ell/2}} \sum_{k \geq 0} \frac{\Gamma(D + \ell/2 + k)}{k! \Gamma(D + \ell + k)} \left(-\frac{t^2}{2}\right)^k = c_\ell t^\ell {}_1F_1(D + \frac{\ell}{2}; D + \ell; -\frac{t^2}{2}).$$

The last expression follows from Kummer’s transformation ${}_1F_1(a; b; -z) = e^{-z} {}_1F_1(b - a; b; z)$ (DLMF, 2026, Eq. 13.2.39). $\square$

Lemma 6 (Eigenvalue bounds). For all $\ell \geq 1$, $d \geq 2$ and $\beta > 0$:

- (i) $\frac{\Gamma(D)}{\Gamma(D + \ell)} \left(\frac{\beta}{2}\right)^\ell \leq \lambda_\ell^\beta \leq \frac{\Gamma(D)}{\Gamma(D + \ell)} \left(\frac{\beta}{2}\right)^\ell \exp\left(\min\left\{\beta, \frac{\beta^2}{4(D + \ell)}\right\}\right)$;
- (ii) $c_\ell^2 (2\ell - 1)!! 3^{-\ell-1/2} \leq \lambda_\ell^{\text{lift}} \leq c_\ell^2 (2\ell - 1)!!$;
- (iii) $A(D, \ell) := \frac{\Gamma(D)\Gamma(D+\ell)}{\Gamma(D+\ell/2)^2}$ satisfies $1 \leq A(D, \ell) \leq A(1, \ell) \leq \sqrt{2} (2\ell - 1)!!/\ell!$.

Proof. (i) The lower bound is the $k = 0$ term of the positive series (16). For the first upper bound, use $I_\mu(\beta) = \frac{(\beta/2)^\mu}{\sqrt{\pi} \Gamma(\mu+1/2)} \int_{-1}^1 e^{\beta x} (1-x^2)^{\mu-1/2} dx$ for $\mu > -\frac{1}{2}$ (DLMF, 2026, Eq. 10.32.2), with $\mu = \ell + \nu \geq \ell \geq 1$. Bounding $e^{\beta x} \leq e^\beta$ and using $\int_{-1}^1 (1-x^2)^{\mu-1/2} dx = \sqrt{\pi} \Gamma(\mu + \frac{1}{2})/\Gamma(\mu + 1)$ gives $I_\mu(\beta) \leq (\beta/2)^\mu e^\beta/\Gamma(\mu + 1)$. For the second, $\Gamma(k + \ell + D) = \Gamma(\ell + D)(\ell + D)_k \geq \Gamma(\ell + D)(\ell + D)^k$ in (16).

(ii) Lower bound: in the Kummer form of Lemma 5, ${}_1F_1(\ell/2; D + \ell; t^2/2) \geq 1$ since all its terms are nonnegative, so $h_\ell(t)^2 \geq c_\ell^2 t^{2\ell} e^{-t^2}$; apply (F3) with $\theta = 1$. Upper bound: $|J_\mu(s)| \leq (s/2)^\mu/\Gamma(\mu+1)$ for $s \geq 0$ and $\mu \geq -\frac{1}{2}$ (DLMF, 2026, Eq. 10.14.4) gives $|b_\ell(s)| \leq \Gamma(D)(s/2)^\ell/\Gamma(D + \ell)$ for $s \geq 0$, and by the parity of b_ℓ also $|b_\ell(s)| \leq \Gamma(D)|s/2|^\ell/\Gamma(D + \ell)$ for all real s . Hence, by (F5), $|h_\ell(t)| \leq \Gamma(D)(|t|/2)^\ell \mathbb{E} R^\ell/\Gamma(D + \ell) = c_\ell |t|^\ell$, and $\lambda_\ell^{\text{lift}} \leq c_\ell^2 \mathbb{E} t^{2\ell} = c_\ell^2 (2\ell - 1)!!$.

(iii) $A \geq 1$ is log-convexity of Γ . Next, $\partial_D \log A(D, \ell) = \psi(D) + \psi(D + \ell) - 2\psi(D + \ell/2) \leq 0$ because the digamma function ψ is concave on $(0, \infty)$. Hence $A(D, \ell) \leq A(1, \ell) = \ell!/\Gamma(1 + \ell/2)^2$ for $D \geq 1$. Let $x = (\ell + 1)/2$. The duplication formula $\Gamma(2z) = \pi^{-1/2} 2^{2z-1} \Gamma(z) \Gamma(z + \frac{1}{2})$ (DLMF, 2026, Eq. 5.5.5), applied with $2z = \ell + 1$ and with $2z = \ell + \frac{1}{2}$, gives $\ell!/\Gamma(1 + \ell/2) = 2^\ell \Gamma(x)/\sqrt{\pi}$ and $(2\ell - 1)!! = 2^\ell \Gamma(\ell + \frac{1}{2})/\sqrt{\pi} = 2^{2\ell-1/2} \Gamma(x - \frac{1}{4}) \Gamma(x + \frac{1}{4})/\pi$. Therefore

$$\frac{\ell! A(1, \ell)}{(2\ell - 1)!!} = \sqrt{2} \frac{\Gamma(x)^2}{\Gamma(x - \frac{1}{4}) \Gamma(x + \frac{1}{4})} \leq \sqrt{2},$$

again by log-convexity of Γ . $\square$

A.9. Proof of Theorem 2 and Corollary 1

Lemma 7 (Eigenvalues of k_β lie in $(0, \kappa_d(\beta))$). For $\ell \geq 1$, $0 < \lambda_\ell^\beta < \kappa_d(\beta) = \lambda_0^\beta$.

Proof. Positivity follows from the positive series (16). By the Funk–Hecke formula, $\lambda_\ell^\beta = \mathbb{E}[e^{\beta W_1} P_\ell(W_1)]$ and $\kappa_d(\beta) = \mathbb{E}[e^{\beta W_1}]$ for $W \sim \sigma$. Since $|P_\ell| \leq 1$, with $|P_\ell(x)| < 1$ for all but finitely many $x \in (-1, 1)$ when $\ell \geq 1$, and W_1 has a density, the inequality is strict. $\square$

Proof of Theorem 2. Global bound. $0 \leq \mathcal{U}_\beta \leq \log(1 + D_\beta^2/\kappa)$ follows from Lemmas 3 and 4.

(i) Let $\kappa = \kappa_d(\beta)$, $u = dq/d\sigma - 1$ (so $\int u d\sigma = 0$ and $\|u\|_\infty = \delta$), and let $\mathcal{K}$ be the integral operator of k_β on $L^2(\sigma)$, which acts on degree ℓ as multiplication by $\lambda_\ell := \lambda_\ell^\beta$. Then $g_q = \kappa + \mathcal{K}u$, and $h := \mathcal{K}u/\kappa$ satisfies $\int h d\sigma = 0$ and $\|h\|_\infty \leq \delta \leq \frac{1}{2}$, because $|\mathcal{K}u(v)| \leq \delta \int e^{\beta v^\top w} d\sigma(w) = \delta \kappa$ for every v . For $|x| \leq \frac{1}{2}$, $\log(1+x) = x - \frac{1}{2}x^2 + r(x)$ with $|r(x)| \leq \frac{|x|^3}{3(1-|x|)} \leq \frac{2}{3}|x|^3$. Hence

$$\mathcal{U}_\beta(q) = \int \log(1+h)(1+u) d\sigma = \underbrace{\int hu d\sigma}_{=:Q_\beta(q)} - \frac{1}{2} \int h^2 d\sigma - \underbrace{\frac{1}{2} \int h^2 u d\sigma}_{=:R} + \int r(h)(1+u) d\sigma.$$

By orthogonality of harmonics, $\int hu = \sum_\ell \lambda_\ell \|u_\ell\|^2/\kappa$ and $\int h^2 = \sum_\ell \lambda_\ell^2 \|u_\ell\|^2/\kappa^2 \leq D_\beta^2/\kappa$, using $\lambda_\ell < \kappa$ (Lemma 7) and $u_\ell = \hat{q}_\ell$ for $\ell \geq 1$. Therefore $|R| \leq (\frac{\delta}{2} + \frac{2}{3}(1+\delta)\|h\|_\infty) \int h^2 \leq (\frac{1}{2} + \frac{2}{3} \cdot \frac{3}{2}) \delta D_\beta^2/\kappa = \frac{3}{2} \delta D_\beta^2/\kappa$. Finally, each weight $\frac{\lambda_\ell}{\kappa} (1 - \frac{\lambda_\ell}{2\kappa})$ lies in $[\frac{\lambda_\ell}{2\kappa}, \frac{\lambda_\ell}{\kappa}]$, and (14) gives the bounds on Q_β .

(ii) Upper bound: by the global bound, $\log(1+x) \leq x$ and (14), $\mathcal{U}_\beta(q) \leq D_\beta^2/\kappa = \kappa^{-1} \sum_{\ell \geq 1} \lambda_\ell \|\hat{q}_\ell\|^2 \leq (S_K/\kappa) \sum_{\ell \geq 1} \mu_\ell \|\hat{q}_\ell\|^2 = (S_K/\kappa) \text{MMD}_K^2$. Lower bound: fix $\ell \geq 1$ and $e \in \mathbb{S}^{d-1}$, and apply (i) to $q_e = (1 + \varepsilon P_\ell(e^\top \cdot))\sigma$, for which $\delta \leq \varepsilon$ and only the degree- ℓ component of $q_e - \sigma$ is nonzero. If $\mu_\ell = 0$, then $\text{MMD}_K(q_e, \sigma) = 0 < \mathcal{U}_\beta(q_e)$ for small $\varepsilon > 0$ by uniqueness (Lemma 4), so $c_K = \infty = S_K$. Otherwise

$$\frac{\mathcal{U}_\beta(q_e)}{\text{MMD}_K^2(q_e, \sigma)} = \frac{\lambda_\ell(1 - \lambda_\ell/(2\kappa))}{\kappa \mu_\ell} + O(\varepsilon) \geq \frac{\lambda_\ell}{2\kappa \mu_\ell} + O(\varepsilon),$$

using $\lambda_\ell < \kappa$ (Lemma 7). Let $\varepsilon \rightarrow 0$ and take the supremum over ℓ . $\square$

Proof of Corollary 1. (a) is Proposition 7. (b) Since $\|u - v\|^2 = 2 - 2u^\top v$, we have $e^{-\gamma\|u-v\|^2} = e^{-2\gamma} k_{2\gamma}(u, v)$, so $\mu_\ell = e^{-2\gamma} \lambda_\ell^{2\gamma}$. By (16), $\lambda_\ell^\beta = \Gamma(D)(x/2)^\ell S_\ell(x)$ with $S_\ell(x) = \sum_k (x/2)^{2k}/(k!\Gamma(k+\ell+D))$, which is increasing in $x > 0$. If $2\gamma \geq \beta$, then $\lambda_\ell^\beta/\mu_\ell = e^{2\gamma}(\beta/2\gamma)^\ell S_\ell(\beta)/S_\ell(2\gamma) \leq e^{2\gamma}$. At $2\gamma = \beta$, $\mu_\ell = e^{-\beta} \lambda_\ell^\beta$ for every ℓ , so $S_K = e^\beta$; by (16), $\kappa_d(\beta) = \Gamma(D)(2/\beta)^{D-1} I_{D-1}(\beta)$, and $I_\nu(\beta) = e^\beta (2\pi\beta)^{-1/2} (1 + O(\beta^{-1}))$ (DLMF, 2026, Eq. 10.40.1) gives $e^\beta/\kappa_d(\beta) = O(\beta^{(d-1)/2})$. If $2\gamma < \beta$, the same ratio is at least $e^{2\gamma}(\beta/2\gamma)^\ell \rightarrow \infty$. For the uniformity loss, $\log \mathbb{E}_{q \otimes q} e^{-\gamma\|u-v\|^2} = -2\gamma + \log \iint k_{2\gamma} dq dq = -2\gamma + \log(\kappa_d(2\gamma) + D_{2\gamma}^2(q, \sigma))$ by (13). Hence $D_{2\gamma}^2 = \kappa_d(2\gamma)(e^{\Delta\mathcal{L}} - 1)$, where $\Delta\mathcal{L}$ is the excess over σ . When $2\gamma \geq \beta$, $\lambda_\ell^\beta \leq \lambda_\ell^{2\gamma}$ for all $\ell \geq 1$, so $\mathcal{U}_\beta \leq D_{2\gamma}^2/\kappa_d(\beta)$; with $\iint k_{2\gamma} dq dq \leq e^{2\gamma}$ this gives $D_{2\gamma}^2 = \kappa_d(2\gamma)(e^{\Delta\mathcal{L}} - 1) \leq e^{2\gamma} \Delta\mathcal{L}$ and $\mathcal{U}_\beta \leq e^{2\gamma} \Delta\mathcal{L}/\kappa_d(\beta)$. Conversely, $\Delta\mathcal{L} \leq D_{2\gamma}^2/\kappa_d(2\gamma)$, so linear control of $\mathcal{U}_\beta$ by $\Delta\mathcal{L}$ fails whenever $2\gamma < \beta$. With $\gamma = 2$, linear control holds exactly when $\tau = 1/\beta \geq 1/4$. (c) For the heat kernel, $\mu_\ell = e^{-s\ell(\ell+d-2)}$ decays faster than $\lambda_\ell^\beta \geq \Gamma(D)(\beta/2)^\ell/\Gamma(D+\ell)$ (Lemma 6), so $S_K = \infty$. A band-limited kernel has $\mu_\ell = 0$ for some $\ell \geq 1$. First- and second-moment matching is MMD_K for $K(u, v) = u^\top v + (u^\top v)^2$, whose eigenvalues vanish for $\ell \geq 3$. $\square$

A.10. Proof of Proposition 7 (lifted kernel)

Comparison and optimality. By (14) for both kernels, $D_\beta^2 = \sum_{\ell \geq 1} \lambda_\ell^\beta \|\hat{q}_\ell\|^2 \leq C_{d,\beta}^* \sum_{\ell \geq 1} \lambda_\ell^{\text{lift}} \|\hat{q}_\ell\|^2 = C_{d,\beta}^* D_{\text{lift}}^2$. Note that $\lambda_\ell^{\text{lift}} > 0$ by Lemma 6(ii). Conversely, fix $e \in \mathbb{S}^{d-1}$, $\ell \geq 1$, and let $q = (1 + \frac{1}{2} P_\ell(e^\top u)) \sigma(du)$, a probability measure because $|P_\ell| \leq 1$ and $\int P_\ell(e^\top u) d\sigma = 0$. Then $q - \sigma$ has only a degree- ℓ component, so $D_\beta^2/D_{\text{lift}}^2 = \lambda_\ell^\beta/\lambda_\ell^{\text{lift}}$. Hence no constant smaller than $C_{d,\beta}^*$ works.

Upper bound. Writing $c_\ell^2 = \Gamma(D)/(2^\ell A(D, \ell)\Gamma(D+\ell))$ and combining Lemma 6(i)–(iii),

$$\frac{\lambda_\ell^\beta}{\lambda_\ell^{\text{lift}}} \leq \frac{\Gamma(D)(\beta/2)^\ell e^\beta}{\Gamma(D+\ell)} \cdot \frac{2^\ell A(D, \ell)\Gamma(D+\ell) 3^{\ell+1/2}}{\Gamma(D)(2\ell-1)!!} = \sqrt{3} e^\beta (3\beta)^\ell \frac{A(D, \ell)}{(2\ell-1)!!} \leq \sqrt{6} e^\beta \frac{(3\beta)^\ell}{\ell!} \leq \sqrt{6} e^{4\beta}.$$

Growth of $C_{d,\beta}^*/\kappa_d(\beta)$. The constant that enters (7) is $C_{d,\beta}^*/\kappa_d(\beta)$. For its lower bound, fix $\ell \geq 1$. By Lemma 6(ii), $C_{d,\beta}^* \geq \lambda_\ell^\beta / (c_\ell^2(2\ell - 1)!!)$, and by (16),

$$\frac{\lambda_\ell^\beta}{\kappa_d(\beta)} = \frac{I_{\nu+\ell}(\beta)}{I_\nu(\beta)} = \prod_{k=0}^{\ell-1} r_{\nu+k}(\beta), \quad r_\mu := \frac{I_{\mu+1}}{I_\mu} \geq \frac{\beta}{\mu + 1 + \sqrt{\beta^2 + (\mu + 1)^2}} = e^{-\operatorname{asinh}((\mu+1)/\beta)},$$

where the inequality is Amos' bound, valid for $\mu \geq 0$ (Amos, 1974); here $\nu = D - 1 \geq 0$. Since asinh is increasing, $\sum_{k=0}^{\ell-1} \operatorname{asinh} \frac{\nu+k+1}{\beta} \leq \beta \int_0^{(\nu+\ell+1)/\beta} \operatorname{asinh}(x) dx$. On the other side, $c_\ell^2(2\ell - 1)!! = \Gamma(D + \frac{\ell}{2})^2 \Gamma(\ell + \frac{1}{2}) / (\sqrt{\pi} \Gamma(D + \ell)^2)$, and Stirling's formula gives $\frac{1}{\ell} \log(c_\ell^2(2\ell - 1)!!) \rightarrow -\log 2$ as $\ell \rightarrow \infty$ for fixed d . Take $\ell = \lfloor c\beta \rfloor$ and let $\beta \rightarrow \infty$:

$$\liminf_{\beta \rightarrow \infty} \frac{1}{\beta} \log \frac{C_{d,\beta}^*}{\kappa_d(\beta)} \geq c \log 2 - \int_0^c \operatorname{asinh}(x) dx = c \log 2 - c \operatorname{asinh} c + \sqrt{1 + c^2} - 1.$$

The right-hand side is maximized where $\operatorname{asinh} c = \log 2$, i.e., $c = \frac{3}{4}$, where it equals $\frac{5}{4} - 1 = \frac{1}{4}$. This proves the last claim of Proposition 7. By Theorem 2(ii), $C_{d,\beta}^*/\kappa_d(\beta)$ is optimal up to a factor of 2 among bounds of the form $\mathcal{U}_\beta \leq c D_{\text{lift}}^2$, so the exponential growth is intrinsic to the lifted kernel. $\square$

A.11. Proof of Theorem 1 (sharp comparison with population InfoNCE)

Upper bound. With equal marginals, $\bar{q} = q$. By Lemma 3, $\mathcal{L}_\beta - \mathcal{L}_\beta^* \leq 2\beta\Delta + \log(1 + D_\beta^2(q, \sigma)/\kappa_d(\beta))$. By Lemma 1, $\Delta \leq k_0\sqrt{\mathcal{J}}$. By Proposition 7, Theorem 4, and Lemma 1,

$$D_\beta^2(q, \sigma) \leq C_{d,\beta}^* D_{\text{lift}}^2(q, \sigma) \leq C_{d,\beta}^* (\sqrt{\mathcal{J}} + \frac{3}{2}\Delta)^2 \leq C_{d,\beta}^* (1 + \frac{3}{2}k_0)^2 \mathcal{J}.$$

This gives (7). The lower bound $\mathcal{L}_\beta \geq \mathcal{L}_\beta^*$ is Lemmas 3 and 4.

Minimizing sequences. “ $\Rightarrow$ ” follows from (7). For “ $\Leftarrow$ ”, suppose $\mathcal{L}_\beta(P_n) - \mathcal{L}_\beta^* \rightarrow 0$. By Lemmas 3 and 4, both nonnegative terms $2\beta\Delta_n$ and $\mathcal{U}_\beta(q_n)$ tend to zero. We claim $q_n \rightarrow \sigma$. Otherwise, by compactness, some subsequence converges weakly to $q' \neq \sigma$, and continuity of Φ (Lemma 4) gives $\mathcal{U}_\beta(q') = 0$, which contradicts uniqueness. Since K_{lift} is bounded and continuous, $q_n \otimes q_n \rightarrow \sigma \otimes \sigma$ implies $D_{\text{lift}}^2(q_n, \sigma) = \iint K_{\text{lift}} d(q_n - \sigma)^{\otimes 2} \rightarrow 0$. By Theorem 4, $\sqrt{\mathcal{J}(P_n)} \leq D_{\text{lift}}(q_n, \sigma) + \frac{3}{2}\Delta_n \rightarrow 0$.

Refined alignment term and sharpness. In the upper bound we may use $\Delta \leq \Delta^*(\mathcal{J})$ (Lemma 1) in place of $\Delta \leq k_0\sqrt{\mathcal{J}}$, which gives the refined form stated after Theorem 1. Consider the family of Section 4.3. Let (n, h) be a Haar-distributed orthonormal pair and, for $s \in [0, 1]$, let $U = \sqrt{s}n + \sqrt{1-s}h$ and $V = \sqrt{s}n - \sqrt{1-s}h$. Both are unit vectors, and for every orthogonal O the pair (On, Oh) has the law of (n, h) ; hence U and V are both σ -distributed. Moreover $C = \sqrt{s}n$, $\Delta = 1 - s$, and $Z = \sqrt{s}Rn \sim \mathcal{N}(0, sI_d)$ by (F1), so by (F3)

$$\mathcal{J} = \mathbb{E}_t(e^{-st^2/2} - e^{-t^2/2})^2 = (1 + 2s)^{-1/2} - 2(2 + s)^{-1/2} + 3^{-1/2} = j(1 - s) = j(\Delta). \quad (17)$$

Since $q = \sigma$, $\mathcal{U}_\beta(q) = 0$ and $\mathcal{L}_\beta - \mathcal{L}_\beta^* = 2\beta\Delta = 2\beta\Delta^*(\mathcal{J})$ exactly. For the limit, the upper bound gives $(\mathcal{L}_\beta - \mathcal{L}_\beta^*)/\sqrt{\mathcal{J}} \leq 2\beta k_0 + c\sqrt{\varepsilon}$ when $0 < \mathcal{J} \leq \varepsilon$, with $c = \bar{c}_{d,\beta}$. Along the family, $j(\Delta) = \Delta^2/(12\sqrt{3}) + O(\Delta^3)$ (indeed $j(0) = j'(0) = 0$ and $j''(0) = \frac{3}{2}3^{-5/2}$), so the ratio tends to $2\beta k_0$ as $\Delta \downarrow 0$, with $\mathcal{J} > 0$ for $\Delta > 0$. $\square$

A.12. Proof of Proposition 3 (square-root barrier)

Write $F(P) = \|\mathcal{A}(P) - m^*\|_{\mathcal{H}}^2$ with $\mathcal{A}(P) = \mathbb{E}_P \Psi(U, V, \xi)$, which is affine in P : $\mathcal{A}(P_p) = (1 - p)\mathcal{A}(P_0) + p\mathcal{A}(P_1)$. Since $F(P_0) = 0$, $\mathcal{A}(P_0) = m^*$, so $\mathcal{A}(P_p) - m^* = p(\mathcal{A}(P_1) - m^*)$ and $F(P_p) = p^2 F(P_1)$. Both marginals of P_0 and of P_1 are σ , hence so are those of P_p , and $\mathcal{U}_\beta(\sigma) = 0$. Since Δ is linear in the law and $\Delta(P_0) = 0$, Lemma 3 gives $\mathcal{L}_\beta(P_p) - \mathcal{L}_\beta^* = 2\beta\Delta(P_p) = 2\beta p\Delta(P_1)$.

For the consequence, let P_1 be any law with uniform marginals and $\Delta(P_1) > 0$ (for instance $\sigma \otimes \sigma$, or a member of the family of Section 4.3). If $F(P_1) = 0$, then F vanishes on a law with $\mathcal{L}_\beta > \mathcal{L}_\beta^*$. Otherwise, along P_p we have $\mathcal{L}_\beta(P_p) - \mathcal{L}_\beta^* = c\sqrt{F(P_p)}$ with $c = 2\beta\Delta(P_1)/\sqrt{F(P_1)} > 0$ and $F(P_p) \downarrow 0$ as $p \downarrow 0$, which contradicts $\mathcal{L}_\beta - \mathcal{L}_\beta^* \leq \psi(F)$ with $\psi(x)/\sqrt{x} \rightarrow 0$. Coverage. $\mathcal{J}$ is Hilbertian with $\mathcal{H} = L^2(\sigma \otimes \mathcal{N}(0, 1))$, $\xi = R$, $\Psi(U, V, R) = [(a, t) \mapsto e^{itRa^\top C}]$ (of norm 1), and $m^* = g$; changing the radius law or the frequency weight changes only Ψ and $\mathcal{H}$. Every squared MMD with a bounded kernel is Hilbertian through its kernel mean embedding. $\square$

A.13. Proof of Theorem 3 (nuisance preference of pure SG) and Corollary 3

Setting. Let $(P_\vartheta)_{\vartheta \geq 0} = (e^{\vartheta \nabla_{\mathbb{S}}^2})$ be the heat semigroup of the Laplace–Beltrami operator $\nabla_{\mathbb{S}}^2$ on $\mathbb{S}^{d-1}$, with kernel $p_\vartheta(w, \cdot)$. Given $W \sim q$, the views U, V are drawn independently from $p_\vartheta(W, \cdot)$; write P_ϑ^q for the resulting paired law. Since $\nabla_{\mathbb{S}}^2$ acts on $\mathcal{H}_\ell$ as $-\ell(\ell + d - 2)$, $\mathbb{E}[U | W] = e^{-(d-1)\vartheta} W$, so $\mathbb{E}U^\top V = e^{-2(d-1)\vartheta}$ and $\Delta(\vartheta) = \frac{1}{2}(1 - e^{-2(d-1)\vartheta})$, independently of q . The marginal of each view is $P_\vartheta q$, with harmonic coefficients $e^{-\ell(\ell+d-2)\vartheta} \hat{q}_\ell$.

Step 1: the pair generator. For a smooth $F : \mathbb{R}^d \rightarrow \mathbb{C}$ let $G(u, v) = F(\frac{u+v}{2})$. For a function ϕ on $\mathbb{R}^d$ restricted to the sphere, $\nabla_{\mathbb{S}}^2 \phi(u) = \text{tr}(\Pi_u \nabla^2 \phi(u)) - (d-1)u^\top \nabla \phi(u)$ with $\Pi_u = I - uu^\top$. Applying this in u and in v and evaluating at $u = v = w$ gives

$$(\nabla_{\mathbb{S},u}^2 + \nabla_{\mathbb{S},v}^2)G|_{u=v=w} = \frac{1}{2}\text{tr}(\Pi_w \nabla^2 F(w)) - (d-1)w^\top \nabla F(w) = \frac{1}{2}\nabla_{\mathbb{S}}^2 F(w) - \frac{d-1}{2}w^\top \nabla F(w) =: \mathcal{G}F(w).$$

The first term is the heat generator at half speed; the second is the generator of the radial contraction $c \mapsto e^{-(d-1)\vartheta/2}c$. Since G is smooth on $\mathbb{S}^{d-1} \times \mathbb{S}^{d-1}$ and the product semigroup $P_\vartheta \otimes P_\vartheta$ has generator $\nabla_{\mathbb{S},u}^2 + \nabla_{\mathbb{S},v}^2$,

$$\frac{d}{d\vartheta} \mathbb{E}F(C)|_{\vartheta=0} = \int \mathcal{G}F dq, \quad C = \frac{1}{2}(U + V). \quad (18)$$

We apply this to $F_{a,t}(c) = \mathbb{E}_R e^{itRa^\top c}$, for which $c^\top \nabla F_{a,t}(c) = t \partial_t F_{a,t}(c)$. All derivatives of $F_{a,t}$ are bounded by polynomials in $|t|$ (moments of R , (F5)). Since $\frac{d}{d\vartheta} \mathbb{E}F_{a,t}(C) = \mathbb{E}[(\mathcal{L}_u + \mathcal{L}_v)^k G(U, V)]$ with $\mathcal{L} = \nabla_{\mathbb{S}}^2$, the ϑ -derivatives of every order are bounded by polynomials in $|t|$ uniformly in $\vartheta \in [0, 1]$. This justifies differentiating under $\mathbb{E}_{a,t}$ below by dominated convergence and shows that $\vartheta \mapsto \mathcal{J}$ is C^∞ on $[0, 1]$.

Step 2: part (i). Write $\varphi_\vartheta(ta) = \mathbb{E}F_{a,t}(C)$ under P_ϑ^q and $u = dq/d\sigma - 1 = \sum_{\ell \geq 1} u_\ell$ (for a general probability measure q , read $\int \cdot u d\sigma$ as integration against $q - \sigma$ and u_ℓ as the degree- ℓ projection of $q - \sigma$; the expansions below converge because $F_{a,t}$ and $\mathcal{G}F_{a,t}$ are smooth on the sphere). At $\vartheta = 0$, $C = W$, and by Lemma 5

$$D(a, t) := \varphi_0(ta) - g(t) = \int F_{a,t} d(q - \sigma) = \sum_{\ell \geq 1} i^\ell h_\ell(t) u_\ell(a).$$

By (18), $\partial_\vartheta \varphi_\vartheta|_0 = \int \mathcal{G}F_{a,t} d\sigma + \int \mathcal{G}F_{a,t} u d\sigma$. The first integral equals $-\frac{d-1}{2}t \partial_t \int F_{a,t} d\sigma = \frac{d-1}{2}t^2 g(t)$, a function of t alone. For the second, self-adjointness of $\nabla_{\mathbb{S}}^2$ and $\int F_{a,t} u d\sigma = D(a, t)$ give

$$\int \mathcal{G}F_{a,t} u d\sigma = -\frac{1}{2} \sum_{\ell \geq 1} \ell(\ell + d - 2) i^\ell h_\ell(t) u_\ell(a) - \frac{d-1}{2} \sum_{\ell \geq 1} i^\ell t h'_\ell(t) u_\ell(a).$$

Now $\partial_\vartheta \mathcal{J}|_0 = 2 \text{Re } \mathbb{E}_{a,t}[\bar{D} \partial_\vartheta \varphi_\vartheta|_0]$. The contribution of $\frac{d-1}{2}t^2 g(t)$ vanishes, because $\mathbb{E}_a D(a, t) = 0$ (every u_ℓ with $\ell \geq 1$ has mean zero under σ). By orthogonality of harmonics of different degrees, and since h_ℓ is real,

$$\partial_\vartheta \mathcal{J}|_{\vartheta=0} = - \sum_{\ell \geq 1} \left(\ell(\ell + d - 2) \mathbb{E}_t h_\ell^2 + (d-1) \mathbb{E}_t [t h_\ell h'_\ell] \right) \|u_\ell\|^2.$$

Gaussian integration by parts, $\mathbb{E}[t \phi'(t)] = \mathbb{E}[(t^2 - 1)\phi(t)]$ for $t \sim \mathcal{N}(0, 1)$, applied to $\phi = h_\ell^2$ gives $(d-1)\mathbb{E}[t h_\ell h'_\ell] = \frac{d-1}{2} \mathbb{E}[(t^2 - 1)h_\ell^2]$, which proves part (i) with $G_\ell = \ell(\ell + d - 2)\lambda_\ell^{\text{lift}} + \frac{d-1}{2} \mathbb{E}_t[(t^2 - 1)h_\ell(t)^2]$ and $\|u_\ell\|^2 = \|\hat{q}_\ell\|^2$. For the bounds on G_ℓ we show

$$\frac{2\ell + 1}{3} < \frac{\mathbb{E}_t[t^2 h_\ell(t)^2]}{\mathbb{E}_t[h_\ell(t)^2]} < 2\ell + 1. \quad (19)$$

By Kummer's Euler integral, ${}_1F_1(a; b; -z) = \mathbb{E} e^{-zS}$ with $S \sim \text{Beta}(a, b-a)$ for $b > a > 0$ (DLMF, 2026, Eq. 13.4.1), so Lemma 5 gives $h_\ell(t) = c_\ell t^\ell \mathbb{E} e^{-St^2/2}$ with $S \sim \text{Beta}(D + \frac{\ell}{2}, \frac{\ell}{2})$. Let S' be an independent copy and $A = 1 + S + S' \in (1, 3)$ almost surely. By Fubini and (F3),

$$\mathbb{E}_t[t^{2m} h_\ell^2] = c_\ell^2 \mathbb{E}_{S, S'} \mathbb{E}_t[t^{2\ell+2m} e^{-(S+S')t^2/2}] = c_\ell^2 (2\ell + 2m - 1)!! \mathbb{E} A^{-\ell-m-1/2},$$

so the ratio in (19) equals $(2\ell + 1) \mathbb{E} A^{-\ell-3/2} / \mathbb{E} A^{-\ell-1/2}$, and $1/3 < A^{-1} < 1$ gives (19). Hence $\frac{2(\ell-1)}{3} \lambda_\ell^{\text{lift}} < \mathbb{E}[(t^2 - 1)h_\ell^2] < 2\ell \lambda_\ell^{\text{lift}}$, which, combined with the expression for G_ℓ above, gives

$$\frac{(d-1)(\ell-1)}{3} < \frac{G_\ell}{\lambda_\ell^{\text{lift}}} - \ell(\ell + d - 2) < (d-1)\ell; \quad (20)$$

in particular the radial term is strictly positive for every $\ell \geq 1$. Finally $\lambda_\ell^{\text{lift}} > 0$ (Lemma 6(ii)).

Step 3: part (ii). The map $q \mapsto P_g^q$ is linear, so for $q_\varepsilon = (1 + \varepsilon f)\sigma$ we have $\varphi_g - g = A_g + \varepsilon B_g$, where $A_g(t) = \varphi_g^\sigma(ta) - g(t)$ depends on t only (rotation invariance) and $B_g(a, t) = \int \mathbb{E}[F_{a,t}(C) \mid W = w]f(w) d\sigma(w)$. Rotation equivariance of the channel makes $\mathbb{E}_a \mathbb{E}[F_{a,t}(C) \mid W = w]$ independent of w , so $\mathbb{E}_a B_g(a, t) = 0$ and the cross term vanishes:

$$\mathcal{J}(\vartheta, \varepsilon) = \mathcal{A}(\vartheta) + \varepsilon^2 C(\vartheta), \quad \mathcal{A}(\vartheta) = \mathbb{E}_t A_g^2, \quad C(\vartheta) = \mathbb{E}_{a,t} |B_g|^2.$$

Here $\mathcal{A}, C \in C^\infty[0, 1]$ by Step 1; $C(0) = \sum_\ell \lambda_\ell^{\text{lift}} \|\hat{f}_\ell\|^2 = \mathcal{J}_{\text{unif}}(q_\varepsilon)/\varepsilon^2$ and $C'(0) = -\sum_\ell G_\ell \|\hat{f}_\ell\|^2 =: -G < 0$ by part (i). By Step 2 with $q = \sigma$, $A_g(t) = \frac{d-1}{2} \vartheta t^2 g(t) + O(\vartheta^2)$ with a polynomially bounded remainder, so by (F3)

$$\mathcal{A}(\vartheta) = a_2 \vartheta^2 + O(\vartheta^3), \quad a_2 = \frac{(d-1)^2}{4} \mathbb{E} t^4 e^{-t^2} = \frac{(d-1)^2}{12\sqrt{3}}.$$

(This matches Lemma 1: to leading order the channel is the extremal family with $\Delta = (d-1)\vartheta$.) Choose $\vartheta_0 \in (0, 1]$ with $\mathcal{A}'' \geq a_2$ on $[0, \vartheta_0]$, and let $M = \max_{[0,1]} |C''|$. Then $\mathcal{A}'(\vartheta) \geq a_2 \vartheta$ and $C'(\vartheta) \geq -G - M\vartheta$ on $[0, \vartheta_0]$, so $\partial_\vartheta^2 \mathcal{J} \geq a_2 - \varepsilon^2 M \geq a_2/2$ on $[0, \vartheta_0]$ once $\varepsilon^2 M \leq a_2/2$, and $\mathcal{J}$ is strictly convex there. Moreover $\partial_\vartheta \mathcal{J} \geq (a_2 - \varepsilon^2 M)\vartheta - \varepsilon^2 G > 0$ for $\vartheta > 2\varepsilon^2 G/a_2$; take ε also so small that $2\varepsilon^2 G/a_2 < \vartheta_0$. Since $\partial_\vartheta \mathcal{J}(0, \varepsilon) = -\varepsilon^2 G < 0$, the unique minimizer ϑ_ε^* over $[0, \vartheta_0]$ is interior and satisfies $\vartheta_\varepsilon^* \leq 2\varepsilon^2 G/a_2$. At an interior minimizer, $0 = \mathcal{A}'(\vartheta^*) + \varepsilon^2 C'(\vartheta^*) = 2a_2 \vartheta^* - \varepsilon^2 G + O(\vartheta^{*2} + \varepsilon^2 \vartheta^*)$, so $\vartheta_\varepsilon^* = \frac{\varepsilon^2 G}{2a_2} (1 + O(\varepsilon^2))$. With $\Delta(\vartheta) = (d-1)\vartheta(1 + O(\vartheta))$,

$$\frac{\Delta(\vartheta_\varepsilon^*)}{\mathcal{J}_{\text{unif}}(q_\varepsilon)} \rightarrow \frac{(d-1)G}{2a_2 C(0)} = \frac{6\sqrt{3} G}{(d-1)C(0)} = \rho(f) > 6\sqrt{3},$$

using $G > (d-1)C(0)$, which follows from part (i) since $\ell(\ell+d-2) \geq d-1$. For $f \in \mathcal{H}_\ell$, part (i) gives $6\sqrt{3} \left[\frac{\ell(\ell+d-2)}{d-1} + \frac{\ell-1}{3} \right] < \rho(f) < 6\sqrt{3} \left[\frac{\ell(\ell+d-2)}{d-1} + \ell \right]$.

Step 4: part (iii). *InfoNCE.* The paired law $P_g^{q_\varepsilon}$ has equal marginals $r_g \sigma$ with $r_g = 1 + \varepsilon P_g f$, so by Lemma 3 $\mathcal{L}_\beta - \mathcal{L}_\beta^* = 2\beta \Delta(\vartheta) + \mathcal{U}_\beta(r_g \sigma)$. Let $g_g = \kappa + \varepsilon \mathcal{K} P_g f$ and $\dot{r} = \varepsilon \nabla_\Sigma^2 P_g f$, so $\|\dot{r}\|_\infty \leq \varepsilon \|\nabla_\Sigma^2 f\|_\infty$ and $\int \dot{r} d\sigma = 0$. Differentiating $\mathcal{U}_\beta = \int \log g_g r_g d\sigma - \log \kappa$,

$$\frac{d}{d\vartheta} \mathcal{U}_\beta = \int \log \frac{g_g}{\kappa} \dot{r} d\sigma + \int (\mathcal{K} \dot{r}) \left(\frac{r_g}{g_g} - \frac{1}{\kappa} \right) d\sigma,$$

where we subtracted $\int \dot{r} d\sigma \cdot \log \kappa = 0$ and $\kappa^{-1} \int \mathcal{K} \dot{r} d\sigma = \int \dot{r} d\sigma = 0$. For $\varepsilon \|f\|_\infty \leq \frac{1}{4}$, $|\log(g_g/\kappa)| \leq 2\varepsilon \|f\|_\infty$, $|r_g/g_g - 1/\kappa| \leq 4\varepsilon \|f\|_\infty/\kappa$ and $|\mathcal{K} \dot{r}| \leq \kappa \|\dot{r}\|_\infty$. Hence $|\frac{d}{d\vartheta} \mathcal{U}_\beta| \leq 6\varepsilon^2 \|f\|_\infty \|\nabla_\Sigma^2 f\|_\infty$, while $\frac{d}{d\vartheta} 2\beta \Delta = 2\beta(d-1)e^{-2(d-1)\vartheta} \geq 2\beta(d-1)e^{-2(d-1)}$ on $[0, 1]$. For small ε the objective is strictly increasing on $[0, \vartheta_0]$, and $\vartheta = 0$ is its unique minimizer. $\square$

Proof of Corollary 3. With $\alpha = a\varepsilon^2$, $\mathcal{J}_\alpha(\vartheta) = \mathcal{A}(\vartheta) + \varepsilon^2 C(\vartheta) + a\varepsilon^2 \Delta(\vartheta)$, and $|\Delta''| \leq 2(d-1)^2$, so $\partial_\vartheta^2 \mathcal{J}_\alpha \geq a_2 - \varepsilon^2(M + 2a(d-1)^2) > 0$ on $[0, \vartheta_0]$ for small ε : $\mathcal{J}_\alpha$ is strictly convex there and has a unique minimizer. Its derivative at 0 is $\varepsilon^2(a(d-1) - G)$. If $a(d-1) \geq G$, the minimizer is $\vartheta = 0$. Otherwise it is interior, and as in Step 3, $0 = 2a_2 \vartheta^* - \varepsilon^2(G - a(d-1)) + O(\vartheta^{*2} + \varepsilon^2 \vartheta^*)$ gives $\vartheta^* = \varepsilon^2(G - a(d-1))/(2a_2) (1 + O(\varepsilon^2))$. Hence $\Delta(\vartheta^*) = \varepsilon^2 \left[(d-1)G - a(d-1)^2 \right]_+ / (2a_2) + o(\varepsilon^2)$. Since $(d-1)\varepsilon^2 G/(2a_2) = \rho(f) \mathcal{J}_{\text{unif}}(q_\varepsilon)$ and $(d-1)^2/(2a_2) = 6\sqrt{3}$, this is the claim. More generally, for every $\alpha \geq \varepsilon^2 G/(d-1)$ and $\vartheta \in (0, \vartheta_0]$, $\partial_\vartheta \mathcal{J}_\alpha \geq (a_2 - \varepsilon^2 M)\vartheta - \varepsilon^2 G(1 - e^{-2(d-1)\vartheta}) \geq \vartheta(a_2 - \varepsilon^2 M - 2(d-1)\varepsilon^2 G) > 0$ for small ε , so $\vartheta = 0$ is the unique minimizer; for $\alpha < \varepsilon^2 G/(d-1)$ the derivative at 0 is negative and the minimizer is positive. In particular every fixed $\alpha > 0$ gives $\vartheta = 0$ for small ε . $\square$

Numerical constants. The values ρ_ℓ quoted after (11) are $\rho(f)$ for $f \in \mathcal{H}_\ell$, i.e., $\rho_\ell = 6\sqrt{3} G_\ell / ((d-1)\lambda_\ell^{\text{lift}})$, computed by quadrature from Lemma 5.

A.14. Proof of Proposition 8 and Corollary 2 (linear certificates)

If $\mathcal{J} = 0$ then $\Delta = 0$ by Lemma 1, so $\mathcal{J}_\alpha = 0$; the converse is trivial. By Lemma 3, $\log(1+x) \leq x$, Proposition 7 and Theorem 4,

$$\mathcal{L}_\beta - \mathcal{L}_\beta^* \leq 2\beta \Delta + \frac{C_{d,\beta}^*}{\kappa_d(\beta)} D_{\text{lift}}^2(q, \sigma) \leq 2\beta \Delta + \frac{C_{d,\beta}^*}{\kappa_d(\beta)} (\sqrt{\mathcal{J}} + \frac{3}{2} \Delta)^2.$$

Using $(x+y)^2 \leq 2x^2 + 2y^2$ and $\Delta^2 \leq \Delta$, the right-hand side is at most $(2\beta + \frac{9}{2} c_\beta) \Delta + 2c_\beta \mathcal{J} \leq c_{\beta,\alpha} (\alpha \Delta + \mathcal{J})$ with $c_\beta = C_{d,\beta}^* / \kappa_d(\beta)$. $\square$

Proof of Corollary 2. By Lemma 3, $\log(1+x) \leq x$ and (14), $\mathcal{L}_\beta - \mathcal{L}_\beta^* \leq 2\beta\Delta + D_\beta^2/\kappa \leq 2\beta\Delta + (S_K/\kappa) \text{MMD}_K^2(q, \sigma)$, which is at most $\max\{2\beta/\alpha, S_K/\kappa\}(\alpha\Delta + \text{MMD}_K^2)$. If $S_K = \infty$, restrict to aligned laws, where $\Delta = 0$ and the bound would read $\mathcal{U}_\beta \leq c \text{MMD}_K^2$; this contradicts the lower bound of Theorem 2(ii), which gives $c_K \geq \sup_\ell \lambda_\ell^\beta / (2\kappa\mu_\ell) = \infty$. $\square$

A.15. General channels

This section proves Proposition 6 and the gain bounds stated after it: Proposition 10 bounds the gain of every centered channel, and Proposition 9 shows that heteroscedastic heat channels attain the order $\sqrt{\mathcal{J}_{\text{unif}}}$. We use the notation and facts of Appendices A.2, A.8 and A.13.

Notation. For a probability measure q on $\mathbb{S}^{d-1}$ and $\ell \geq 1$ let

$$u_\ell(x) = \int N_\ell P_\ell(x^\top y) dq(y) = \sum_{m \leq N_\ell} Y_{\ell m}(x) \int Y_{\ell m} dq,$$

the degree- ℓ component of $q - \sigma$ (for $q = (1+u)\sigma$ this is the projection of u onto $\mathcal{H}_\ell$), so that $\|u_\ell\|_{L^2(\sigma)} = \|\hat{q}_\ell\|$ and $\mathcal{J}_{\text{unif}}(q) = D_{\text{lift}}^2(q, \sigma) = \sum_{\ell \geq 1} \lambda_\ell^{\text{lift}} \|\hat{q}_\ell\|^2$ by (14) and Lemma 5. By the Cauchy–Schwarz inequality and the addition theorem,

$$|u_\ell(x)| \leq \sqrt{N_\ell} \|\hat{q}_\ell\|, \quad \|\hat{q}_\ell\|^2 = \iint N_\ell P_\ell(x^\top y) dq(x) dq(y) \leq N_\ell. \quad (21)$$

$G_\ell > 0$ are the constants of Theorem 3(i) and (20). A family is a map $\vartheta \mapsto P_\vartheta$, $\vartheta \in [0, \vartheta_0]$, into paired laws; for an objective F we write $f(\vartheta) = F(P_\vartheta)$ and $\Delta(\vartheta) = \Delta(P_\vartheta)$.

Lemma 8 (Summability). $G_\ell < \ell(\ell + 2d - 3) \lambda_\ell^{\text{lift}}$ for $\ell \geq 1$, and $\sum_{\ell \geq 1} N_\ell \ell^k \lambda_\ell^{\text{lift}} < \infty$ for every $k \geq 0$. Consequently

$$K_d := \left(\sum_{\ell \geq 1} \frac{N_\ell G_\ell^2}{\lambda_\ell^{\text{lift}}} \right)^{1/2} < \infty, \quad \Gamma_q(x) := \sum_{\ell \geq 1} G_\ell u_\ell(x)$$

converges absolutely and uniformly on $\mathbb{S}^{d-1}$, and Γ_q is continuous with $\int \Gamma_q d\sigma = 0$.

Proof. The first claim is the upper bound in (20). By Lemma 6(ii), $\lambda_\ell^{\text{lift}} \leq c_\ell^2 (2\ell - 1)!!$, and $\frac{1}{\ell} \log(c_\ell^2 (2\ell - 1)!!) \rightarrow -\log 2$ (proof of Proposition 7), so $\lambda_\ell^{\text{lift}} \leq C_d (3/5)^\ell$ for all ℓ ; $N_\ell \leq c_d \ell^{d-2}$. This gives the series bound, and $K_d^2 \leq \sum_\ell N_\ell \ell^2 (\ell + 2d - 3)^2 \lambda_\ell^{\text{lift}} < \infty$. By (21), $\sum_\ell G_\ell \|u_\ell\|_\infty \leq \sum_\ell G_\ell N_\ell < \infty$. Each u_ℓ is continuous with σ -mean zero. $\square$

Proof of Proposition 6. Write $f(\vartheta) = F(P_\vartheta)$ and $f_\alpha = f + \alpha\Delta$. By definition of g , the right derivative is $f'_\alpha(0) = f'(0) + \alpha\Delta'(0) = \Delta'(0)(\alpha - g)$, so $f_\alpha(\vartheta) - f_\alpha(0) = \Delta'(0)(\alpha - g)\vartheta + o(\vartheta)$, which gives (i). For (ii), $\lambda S + (1 - \lambda)I = \lambda(S + \frac{1-\lambda}{\lambda}I)$; dividing by $\lambda > 0$ does not change minimizers, and (i) applies with I as the misalignment coordinate. Multiplying S by B multiplies its derivative, and hence g_S , by B . $\square$

For the isotropic heat channel, $g = \rho(f) \mathcal{J}_{\text{unif}}(q_\varepsilon) / (6\sqrt{3})$, and Corollary 3 gives the global version of (i) on $[0, \vartheta_0]$.

Heteroscedastic heat channels

Let $w : \mathbb{S}^{d-1} \rightarrow [0, \infty)$ be measurable with $0 < \mathbb{E}_q w < \infty$. The heteroscedastic heat channel $P_\vartheta^{q,w}$ draws $W \sim q$ and then U, V independently from $p_{\vartheta w(W)}(W, \cdot)$; $w \equiv 1$ is the channel of Theorem 3. It is a centered channel in the sense of Proposition 10 below, with $c(W) = e^{-(d-1)\vartheta w(W)}$.

Proposition 9 (Gain of heteroscedastic heat channels). *Let $q \neq \sigma$, and let w be as above.*

- (i) $\Delta(\vartheta) = \frac{1}{2}(1 - \mathbb{E}_q e^{-2(d-1)\vartheta w(W)})$ and $\vartheta \mapsto \mathcal{J}(P_\vartheta^{q,w})$ are continuous on $[0, \infty)$ and right-differentiable at 0, Δ is strictly increasing with $\Delta'(0) = (d-1)\mathbb{E}_q w$, and the gain of the family is

$$g(w) = \frac{\mathbb{E}_q[w \Gamma_q]}{(d-1) \mathbb{E}_q w}.$$

If w is bounded, both functions are C^∞ on $[0, \infty)$.

- (ii) For $w \equiv 1$, $g = \frac{1}{d-1} \sum_{\ell \geq 1} G_\ell \|\hat{q}_\ell\|^2 = \frac{\rho(q)}{6\sqrt{3}} \mathcal{J}_{\text{unif}}(q)$, with $\rho(q) := \frac{6\sqrt{3} \sum_\ell G_\ell \|\hat{q}_\ell\|^2}{(d-1) \sum_\ell \lambda_\ell^{\text{lift}} \|\hat{q}_\ell\|^2} > 6\sqrt{3}$.
- (iii) $\sup_w g(w) = \frac{1}{d-1} \max_{\text{supp } q} \Gamma_q$, the supremum being over bounded w or over all admissible w . It is attained if and only if $q(\{x \in \text{supp } q : \Gamma_q(x) = \max_{\text{supp } q} \Gamma_q\}) > 0$.
- (iv) $\|\Gamma_q\|_\infty \leq K_d \sqrt{\mathcal{J}_{\text{unif}}(q)}$; hence $\sup_w g(w) \leq \frac{K_d}{d-1} \sqrt{\mathcal{J}_{\text{unif}}(q)}$. The constant is sharp: $\sup\{g(w)/\sqrt{\mathcal{J}_{\text{unif}}(q)}\} = K_d/(d-1)$ over all $q \neq \sigma$ and w , and codes with $\sup_w g(w) \geq (K_d/(d-1) - \eta) \sqrt{\mathcal{J}_{\text{unif}}(q)}$ exist for every $\eta > 0$ and with $\mathcal{J}_{\text{unif}}(q)$ arbitrarily small.

Proof. Write $Q_s = P_s \otimes P_s$ for the product heat semigroup on $\mathbb{S}^{d-1} \times \mathbb{S}^{d-1}$, with generator $L_2 = \nabla_{\mathbb{S},u}^2 + \nabla_{\mathbb{S},v}^2$, and, as in Step 1 of [Appendix A.13](#), $G_{a,t}(u, v) = F_{a,t}(\frac{u+v}{2})$ with $F_{a,t}(c) = \mathbb{E}_R e^{itRa^\top c}$. Throughout, p_k denotes a polynomial, depending only on d and k , with $\|L_2^k G_{a,t}\|_\infty \leq p_k(|t|)$ (Step 1 of [Appendix A.13](#)).

Semigroup bounds. For smooth G , $s \mapsto Q_s G$ is C^∞ into $C(\mathbb{S}^{d-1} \times \mathbb{S}^{d-1})$ with $\partial_s^k Q_s G = Q_s L_2^k G$, and Q_s is a sup-norm contraction. Hence, for all $s \geq 0$ and $x \in \mathbb{S}^{d-1}$,

$$|Q_s G(x, x) - G(x, x)| \leq s \|L_2 G\|_\infty, \quad |Q_s G(x, x) - G(x, x) - s L_2 G(x, x)| \leq \frac{s^2}{2} \|L_2^2 G\|_\infty, \quad (22)$$

and $L_2 G_{a,t}(x, x) = \mathcal{G} F_{a,t}(x)$ by Step 1 of [Appendix A.13](#).

(i), *misalignment.* The coordinates are degree-one harmonics, so $\mathbb{E}[U | W] = e^{-(d-1)\vartheta w(W)} W$, and conditional independence gives $\mathbb{E}[U^\top V | W] = e^{-2(d-1)\vartheta w(W)}$, which is the formula for Δ . It is continuous and strictly increasing because $q(w > 0) > 0$. Since $0 \leq (1 - e^{-2(d-1)\vartheta w})/\vartheta \leq 2(d-1)w \in L^1(q)$, dominated convergence gives $\Delta'(0) = (d-1)\mathbb{E}_q w$.

(i), *first variation of $\mathcal{J}$.* Conditionally on $W = x$, $\mathbb{E}[F_{a,t}(C) | W = x] = Q_{\vartheta w(x)} G_{a,t}(x, x)$, so $\varphi_\vartheta(ta) := \varphi_Z(ta) = \int Q_{\vartheta w(x)} G_{a,t}(x, x) dq(x)$. Put $D = \varphi_0 - g$ and $\psi(a, t) = \int w \mathcal{G} F_{a,t} dq$. By (22),

$$\frac{1}{\vartheta} |\varphi_\vartheta - \varphi_0 - \vartheta \psi| \leq \int \min \left\{ \frac{\vartheta w^2}{2} p_2(|t|), 2w p_1(|t|) \right\} dq \xrightarrow{\vartheta \downarrow 0} 0$$

by dominated convergence, and $|\varphi_\vartheta - \varphi_{\vartheta'}| \leq |\vartheta - \vartheta'| \mathbb{E}_q w p_1(|t|)$. The last bound shows that $\mathcal{J}(\vartheta) = \mathbb{E}_{a,t} |\varphi_\vartheta - g|^2$ is continuous. Writing $\mathcal{J}(\vartheta) - \mathcal{J}(0) = 2 \text{Re } \mathbb{E}_{a,t} [\overline{D}(\varphi_\vartheta - \varphi_0)] + \mathbb{E}_{a,t} |\varphi_\vartheta - \varphi_0|^2$, with $|D| \leq 2$ and $\mathbb{E}_t p_1(|t|)^2 < \infty$, dominated convergence gives

$$\partial_\vartheta \mathcal{J}|_{\vartheta=0^+} = 2 \text{Re } \mathbb{E}_{a,t} [\overline{D(a, t)} \psi(a, t)]. \quad (23)$$

If w is bounded, $\partial_\vartheta^k \varphi_\vartheta = \int w^k Q_{\vartheta w} L_2^k G_{a,t}(x, x) dq$ is bounded by $\|w\|_\infty^k p_k(|t|)$, and $\mathcal{J}$ and Δ are C^∞ by dominated convergence.

(i), *spectral evaluation.* For finite signed measures μ, ν on $\mathbb{S}^{d-1}$ and fixed t let $\Phi_\nu(a) = \int F_{a,t} d\nu$ and $\Psi_\mu(a) = \int \mathcal{G} F_{a,t} d\mu$. For $Y \in \mathcal{H}_\ell$ and $x \in \mathbb{S}^{d-1}$, (15) (applied in the variable a) gives $\int F_{a,t}(x) Y(a) d\sigma(a) = i^\ell h_\ell(t) Y(x)$. Since $\mathcal{G} F_{a,t}(x) = \frac{1}{2} \nabla_{\mathbb{S}}^2 F_{a,t}(x) - \frac{d-1}{2} t \partial_t F_{a,t}(x)$ on $\mathbb{S}^{d-1}$ and all derivatives are dominated, we may differentiate under the integral:

$$\int \mathcal{G} F_{a,t}(x) Y(a) d\sigma(a) = i^\ell k_\ell(t) Y(x), \quad k_\ell(t) := -\frac{1}{2} \ell(\ell + d - 2) h_\ell(t) - \frac{d-1}{2} t h'_\ell(t).$$

By Fubini, the degree- ℓ components of the bounded functions $\Phi_\nu, \Psi_\mu \in L^2(\sigma)$ are $i^\ell h_\ell(t) \nu_\ell$ and $i^\ell k_\ell(t) \mu_\ell$, where $\mu_\ell = \sum_m (\int Y_{\ell m} d\mu) Y_{\ell m}$. Parseval's identity (h_ℓ, k_ℓ are real) gives $\mathbb{E}_a [\Phi_\nu \Psi_\mu] = \sum_{\ell \geq 0} h_\ell(t) k_\ell(t) \langle \nu_\ell, \mu_\ell \rangle$, and, by Cauchy-Schwarz in ℓ and Parseval again, $\sum_\ell |h_\ell k_\ell \langle \nu_\ell, \mu_\ell \rangle| \leq \|\Phi_\nu\|_{L^2(\sigma)} \|\Psi_\mu\|_{L^2(\sigma)} \leq |\nu|(\mathbb{S}^{d-1}) |\mu|(\mathbb{S}^{d-1}) p_1(|t|)$, which is integrable in t . Hence $\mathbb{E}_t$ and $\sum_\ell$ commute. Gaussian integration by parts, as in Step 2 of [Appendix A.13](#), gives $\mathbb{E}_t [h_\ell k_\ell] = -\frac{1}{2} (\ell(\ell + d - 2) \lambda_\ell^{\text{lift}} + \frac{d-1}{2} \mathbb{E}_t [(t^2 - 1) h_\ell^2]) = -\frac{1}{2} G_\ell$ for $\ell \geq 1$. Take $\nu = q - \sigma$ (so $\nu_0 = 0$, $\nu_\ell = u_\ell$, $\Phi_\nu = D$) and $\mu = w q$ (so $\Psi_\mu = \psi$). By (23),

$$\partial_\vartheta \mathcal{J}|_{0^+} = - \sum_{\ell \geq 1} G_\ell \langle u_\ell, \mu_\ell \rangle = - \sum_{\ell \geq 1} G_\ell \int u_\ell w dq = -\mathbb{E}_q [w \Gamma_q],$$

where $\langle u_\ell, \mu_\ell \rangle = \int u_\ell d\mu$ because $u_\ell \in \mathcal{H}_\ell$, and the last sum converges absolutely by [Lemma 8](#). Dividing by $-\Delta'(0)$ gives $g(w)$.

(ii) For $w \equiv 1$, $\mathbb{E}_q \Gamma_q = \sum_\ell G_\ell \int u_\ell dq = \sum_\ell G_\ell \|\hat{q}_\ell\|^2$, since $\int u_\ell dq = \int u_\ell d(q - \sigma) = \|u_\ell\|^2$. The expression with $\rho(q)$ is a rewriting, and $\rho(q) > 6\sqrt{3}$ by $G_\ell > \ell(\ell + d - 2) \lambda_\ell^{\text{lift}} \geq (d-1) \lambda_\ell^{\text{lift}}$.

(iii) $g(w)$ is the mean of $\Gamma_q/(d-1)$ under the probability measure $w dq/\mathbb{E}_q w$, hence at most $M/(d-1)$ with $M := \text{ess sup}_q \Gamma_q = \max_{\text{supp } q} \Gamma_q$ (continuity of Γ_q). For $\eta > 0$, $w = \mathbf{1}_{\{\Gamma_q > M - \eta\}}$ is bounded with $\mathbb{E}_q w > 0$ and

$g(w) \geq (M - \eta)/(d - 1)$. If $q(\Gamma_q = M) > 0$, $w = \mathbf{1}\{\Gamma_q = M\}$ attains $M/(d - 1)$; conversely, $g(w) = M/(d - 1)$ forces $\mathbb{E}_q[w(M - \Gamma_q)] = 0$ with a q -a.e. nonnegative integrand, so $w = 0$ q -a.e. on $\{\Gamma_q < M\}$, and $\mathbb{E}_q w > 0$ forces $q(\Gamma_q = M) > 0$.

(iv) By (21) and Cauchy–Schwarz,

$$|\Gamma_q(x)| \leq \sum_{\ell} G_{\ell} \sqrt{N_{\ell}} \|\hat{q}_{\ell}\| = \sum_{\ell} \frac{G_{\ell} \sqrt{N_{\ell}}}{\sqrt{\lambda_{\ell}^{\text{lift}}}} \cdot \sqrt{\lambda_{\ell}^{\text{lift}}} \|\hat{q}_{\ell}\| \leq K_d \sqrt{\mathcal{J}_{\text{unif}}(q)}.$$

With (iii) this gives the bound. Sharpness: fix $x_0 \in \mathbb{S}^{d-1}$ and $L \geq 1$, let $f_L(x) = \sum_{\ell=1}^L (G_{\ell}/\lambda_{\ell}^{\text{lift}}) N_{\ell} P_{\ell}(x_0^{\top} x)$ and $q = (1 + \varepsilon f_L)\sigma$ with $0 < \varepsilon \|f_L\|_{\infty} < 1$, so that $\text{supp } q = \mathbb{S}^{d-1}$. By the addition theorem, $N_{\ell} P_{\ell}(x_0^{\top} \cdot) \in \mathcal{H}_{\ell}$ has squared norm N_{ℓ} , so $u_{\ell} = \varepsilon (G_{\ell}/\lambda_{\ell}^{\text{lift}}) N_{\ell} P_{\ell}(x_0^{\top} \cdot)$ for $\ell \leq L$ and 0 otherwise. Hence $\mathcal{J}_{\text{unif}}(q) = \varepsilon^2 K_L^2$ and $\Gamma_q(x_0) = \varepsilon K_L^2$, with $K_L^2 = \sum_{\ell \leq L} N_{\ell} G_{\ell}^2 / \lambda_{\ell}^{\text{lift}}$. By (iii), $\sup_w g(w) \geq \frac{K_L}{d-1} \sqrt{\mathcal{J}_{\text{unif}}(q)}$, and $K_L \uparrow K_d$ while ε is free. $\square$

A nuisance certificate for centered channels

Let $k_d := 2 + \frac{2}{\sqrt{d/(d+2)}} < 6.5$.

Proposition 10 (Nuisance certificate for centered channels). *Let Λ be a random element, $W = W(\Lambda) \in \mathbb{S}^{d-1}$ with law q , and let $U, V \in \mathbb{S}^{d-1}$ be conditionally independent given Λ with*

$$\mathbb{E}[U \mid \Lambda] = \mathbb{E}[V \mid \Lambda] = c(\Lambda) W, \quad c(\Lambda) \in [0, 1].$$

Then, with P the law of (U, V) and $T_q = RW$,

$$\|\varphi_Z(t \cdot) - \varphi_{T_q}(t \cdot)\| \leq k_d \Delta(P), \quad \text{hence} \quad |\sqrt{\mathcal{J}(P)} - \sqrt{\mathcal{J}_{\text{unif}}(q)}| \leq k_d \Delta(P) \leq 6.5 \Delta(P),$$

and $\mathcal{J}(P) + \alpha \Delta(P) \geq \mathcal{J}_{\text{unif}}(q) + (\alpha - 2k_d \sqrt{\mathcal{J}_{\text{unif}}(q)}) \Delta(P)$ for every $\alpha \geq 0$. In particular, if $\alpha > 2k_d \sqrt{\mathcal{J}_{\text{unif}}(q)}$ (for instance $\alpha > 13 \sqrt{\mathcal{J}_{\text{unif}}(q)}$), then $\mathcal{J}_{\alpha}(P) > \mathcal{J}_{\alpha}(P_q) = \mathcal{J}_{\text{unif}}(q)$ for every such P with $\Delta(P) > 0$, where P_q is the aligned law $U = V = W$.

Proof. Fix (a, t) and put $x = tRa$; R is independent of (Λ, U, V) . Since $|e^{iy} - 1 - iy| \leq y^2/2$,

$$\varphi_Z(ta) - \varphi_{T_q}(ta) = \mathbb{E}[e^{ix^{\top} W} i x^{\top} (C - W)] + \mathbb{E}[e^{ix^{\top} W} r] =: T_1(a, t) + T_2(a, t), \quad |r| \leq \frac{1}{2} (x^{\top} (C - W))^2.$$

Conditional moments. Let $\Delta_{\Lambda} = \frac{1}{4} \mathbb{E}[\|U - V\|^2 \mid \Lambda]$, so $\mathbb{E} \Delta_{\Lambda} = \Delta$. By conditional independence, $\mathbb{E}[U^{\top} V \mid \Lambda] = c^2$, so $\Delta_{\Lambda} = \frac{1}{2} (1 - c^2)$ and, as $c \in [0, 1]$, $1 - c \leq (1 - c)(1 + c) = 2\Delta_{\Lambda}$. Moreover $\mathbb{E}[C \mid \Lambda] = cW$ and $\mathbb{E}[\|C\|^2 \mid \Lambda] = 1 - \Delta_{\Lambda}$, so

$$\mathbb{E}[\|C - W\|^2 \mid \Lambda] = 2(1 - c) - \Delta_{\Lambda} \leq 3\Delta_{\Lambda}.$$

First-order term. Conditionally on (Λ, R) , $\mathbb{E}[x^{\top} (C - W) \mid \Lambda, R] = tR(c - 1) a^{\top} W$. Hence $|T_1| \leq 2|t| \mathbb{E} R \mathbb{E}[|a^{\top} W| \Delta_{\Lambda}]$. By the Cauchy–Schwarz inequality with weights Δ_{Λ} , $(\mathbb{E}[|a^{\top} W| \Delta_{\Lambda}])^2 \leq \Delta \mathbb{E}[(a^{\top} W)^2 \Delta_{\Lambda}]$, and $\mathbb{E}_a(a^{\top} W)^2 = 1/d$. Using $(\mathbb{E} R)^2 \leq \mathbb{E} R^2 = d$ (F5) and $\mathbb{E} t^2 = 1$, we get $\|T_1\|^2 \leq 4 \mathbb{E} t^2 (\mathbb{E} R)^2 \Delta^2 / d \leq 4\Delta^2$.

Remainder. With $\Sigma = \mathbb{E}[(C - W)(C - W)^{\top}] \succeq 0$, $\text{tr } \Sigma \leq 3\Delta$, independence of R and (F5) give $|T_2| \leq \frac{d}{2} t^2 a^{\top} \Sigma a$. Exactly as in Lemma 2, (F3)–(F4) and $\text{tr}(\Sigma^2) \leq (\text{tr } \Sigma)^2$ give

$$\|T_2\|^2 \leq \frac{d^2}{4} \cdot 3 \cdot \frac{3(\text{tr } \Sigma)^2}{d(d+2)} \leq \frac{81d}{4(d+2)} \Delta^2.$$

By Minkowski’s inequality, $\|\varphi_Z - \varphi_{T_q}\| \leq (2 + \frac{2}{\sqrt{d/(d+2)}}) \Delta = k_d \Delta$. Since $\sqrt{\mathcal{J}} = \|\varphi_Z - g\|$ and $\sqrt{\mathcal{J}_{\text{unif}}(q)} = D_{\text{lift}}(q, \sigma) = \|\varphi_{T_q} - g\|$, the reverse triangle inequality gives the second claim.

Consequence. Let $s = \sqrt{\mathcal{J}_{\text{unif}}(q)}$. If $s \geq k_d \Delta$, then $\mathcal{J} \geq (s - k_d \Delta)^2 \geq s^2 - 2k_d s \Delta$. If $s < k_d \Delta$, then $s^2 - 2k_d s \Delta < -k_d s \Delta \leq 0 \leq \mathcal{J}$. In both cases $\mathcal{J} \geq \mathcal{J}_{\text{unif}}(q) - 2k_d \sqrt{\mathcal{J}_{\text{unif}}(q)} \Delta$; add $\alpha \Delta$. Finally, $\mathcal{J}(P_q) = D_{\text{lift}}^2(q, \sigma) = \mathcal{J}_{\text{unif}}(q)$ and $\Delta(P_q) = 0$. $\square$

Corollary 4 (Gains of centered channels). *Let (P_{θ}) be a family as in Proposition 6 with $F = \mathcal{J}$, such that each P_{θ} is a centered channel as in Proposition 10 with the same code law q , and $P_0 = P_q$. Then its gain satisfies $g \leq 2k_d \sqrt{\mathcal{J}_{\text{unif}}(q)} \leq 13 \sqrt{\mathcal{J}_{\text{unif}}(q)}$. Consequently, $\mathcal{J}_{\alpha}$ has the aligned code as a strict local minimizer along every such family once $\alpha > 2k_d \sqrt{\mathcal{J}_{\text{unif}}(q)}$. By Proposition 9(iv), the best constant c_d^{gain} in $g \leq c_d^{\text{gain}} \sqrt{\mathcal{J}_{\text{unif}}(q)}$ over all centered families satisfies $K_d/(d - 1) \leq c_d^{\text{gain}} \leq 2k_d$, so the order $\sqrt{\mathcal{J}_{\text{unif}}}$ cannot be improved.*

Proof. By [Proposition 10](#), $(f(\vartheta) - f(0))/\Delta(\vartheta) \geq -2k_d\sqrt{\mathcal{J}_{\text{unif}}(q)}$ for small $\vartheta > 0$ (where $\Delta(\vartheta) > 0$ because $\Delta'(0) > 0$), and the left side tends to $-g$. Heteroscedastic heat channels are centered, with $c = e^{-(d-1)\vartheta w(W)} \in (0, 1]$ and $\Lambda = W$. $\square$

Remark 1 (Scope of the centering hypothesis). Whenever the two views are conditionally independent given a latent Λ with the same conditional mean (e.g. i.i.d. per-view style given content), the hypothesis of [Proposition 10](#) holds with the *mean code* $W = \mathbb{E}[U \mid \Lambda]/\|\mathbb{E}[U \mid \Lambda]\|$ and $c = \|\mathbb{E}[U \mid \Lambda]\|$ (W arbitrary where $c = 0$). The hypothesis cannot be dropped: an aligned displacement of the code, $U = V = W'$ with $W' \neq W$, has $\Delta = 0$ and $\mathcal{J} = \mathcal{J}_{\text{unif}}(\text{law}(W'))$, which in general differs from $\mathcal{J}_{\text{unif}}(q)$.

A.16. Finite batches

We use the setting of [Section 5](#), with $B \geq 2$, $M \geq 1$, and frequencies independent of the pairs and radii.

In the notation of [Section 5](#), (10) reads

$$\hat{\mathcal{J}}_U = \frac{1}{M} \sum_{m=1}^M \left[\frac{B|\hat{\phi}_m|^2 - 1}{B-1} - 2g_m \text{Re } \hat{\phi}_m + g_m^2 \right]. \quad (24)$$

Unbiasedness averages over both the i.i.d. examples and the fresh frequencies; a fixed quadrature in (a, t) defines a different discrepancy.

Proof of Proposition 4. Condition on a frequency ω . The Z_j are i.i.d. (pairs are i.i.d. and each has its own radius), so $\mathbb{E}|\hat{\phi}|^2 = B^{-2} \sum_{j,k} \mathbb{E}e^{i\omega^\top(Z_j - Z_k)} = \frac{1}{B} + (1 - \frac{1}{B})|\varphi_Z(\omega)|^2$ and $\mathbb{E}\hat{\phi} = \varphi_Z(\omega)$. Since g is real, $\mathbb{E}|\hat{\phi} - g|^2 = \mathbb{E}|\hat{\phi}|^2 - 2g \text{Re } \varphi_Z + g^2 = |\varphi_Z - g|^2 + \frac{1}{B}(1 - |\varphi_Z|^2)$. Averaging over i.i.d. frequencies proves the identity. On the family of [Section 4.3](#), $\mathbb{E}_{a,t}|\varphi_Z|^2 = \mathbb{E}e^{-st^2} = (1 + 2s)^{-1/2}$, so by (17)

$$F_B(s) := \mathbb{E}\hat{\mathcal{J}}_V = (1 - \frac{1}{B})(1 + 2s)^{-1/2} - 2(2 + s)^{-1/2} + 3^{-1/2} + \frac{1}{B}.$$

Then $F'_B(s) = (2 + s)^{-3/2} - (1 - \frac{1}{B})(1 + 2s)^{-3/2}$, which is negative if and only if $r(s) := \frac{1+2s}{2+s} < a_B := (1 - \frac{1}{B})^{2/3}$. The map r increases from $\frac{1}{2}$ to 1 on $[0, 1]$, and $a_B \in (\frac{1}{2}, 1)$ for every $B \geq 2$ (since $a_2 = 2^{-2/3}$). So F_B decreases and then increases, with unique minimizer $s_B = r^{-1}(a_B) = (2a_B - 1)/(2 - a_B) \in (0, 1)$. Moreover $1 - s_B = 3(1 - a_B)/(2 - a_B)$ and $1 - a_B = \frac{2}{3B} + O(B^{-2})$, so $1 - s_B = \frac{2}{B} + O(B^{-2})$ and $2s_B - 1 = 1 - \frac{4}{B} + O(B^{-2})$. Every aligned uniform law has $Z \sim \mathcal{N}(0, I_d)$ and thus the value $F_B(1) > F_B(s_B)$, so it is not a minimizer over paired laws. $\square$

Proof of Proposition 5. Unbiasedness. Conditionally on ω_m , $\mathbb{E}[(B|\hat{\phi}_m|^2 - 1)/(B-1)] = |\varphi_Z(\omega_m)|^2$ by the computation above, so the bracket in (24) has conditional mean $|\varphi_Z(\omega_m) - g_m|^2$. Averaging over $\omega_m = t_m a_m$ gives $\mathcal{J}$.

U-statistic form and cost. With $\xi_j \in \mathbb{C}^M$, $(\xi_j)_m = M^{-1/2}(e^{i\omega_m^\top Z_j} - g_m)$, we have $\hat{\mathcal{J}}_U = \frac{1}{B(B-1)} \sum_{j \neq k} H(Z_j, Z_k)$ with the symmetric kernel $H(z, z') = \text{Re}\langle \xi(z), \xi(z') \rangle$. Indeed, $B^2|\hat{\phi}_m|^2 = B + \sum_{j \neq k} \cos(\omega_m^\top(Z_j - Z_k))$ and $\sum_{j \neq k} (\cos \omega_m^\top Z_j + \cos \omega_m^\top Z_k) = 2(B-1) \sum_j \cos \omega_m^\top Z_j$. Equivalently, $\hat{\mathcal{J}}_U = (\|\sum_j \xi_j\|^2 - \sum_j \|\xi_j\|^2)/(B(B-1))$. Computing $\{\omega_m^\top Z_j\}$ is one $B \times d$ by $d \times M$ product, $O(BdM)$; the rest is $O(BM)$. $\square$

B. Experimental Details

Exact families ([Fig. 1a](#)). (i) Extremal family: (17) and $\mathcal{L}_\beta - \mathcal{L}_\beta^* = 2\beta(1 - s)$. (ii) Noisy views: $U \sim \sigma$, $V = (U + \eta_n \xi)/\|U + \eta_n \xi\|$ with $\xi \sim \mathcal{N}(0, I_d)$. The pair law is rotation invariant, so $Z = R\|C\|\theta$ with $\theta \sim \sigma$ independent of $\|C\|$, and $\varphi_Z(ta) = \mathbb{E}e^{-t^2\|C\|^2/2}$. We evaluate $\mathcal{J}$ by 80-point Gauss–Hermite quadrature in t and 4×10^5 Monte Carlo samples of $\|C\|^2 = (1 + U^\top V)/2$ with common random numbers across t ; $\mathcal{U}_\beta = 0$. (iii) Aligned vMF: $U = V \sim \text{vMF}(e, \kappa_{\text{vMF}})$. Then $\|\hat{q}_\ell\|^2 = N_\ell r_\ell^2$ with $r_\ell = I_{\ell+\nu}(\kappa_{\text{vMF}})/I_\nu(\kappa_{\text{vMF}})$, $\mathcal{J} = \sum_{\ell \geq 1} \lambda_\ell^{\text{lift}} N_\ell r_\ell^2$ (truncated at $\ell = 80$), and $\mathcal{U}_\beta$ is a one-dimensional integral; $\lambda_\ell^{\text{lift}}$ is computed from [Lemma 5](#) by adaptive quadrature. (iv) Mixture path: $\mathcal{J}(P_p) = p^2 \mathcal{J}(\sigma \otimes \sigma)$ with $\mathcal{J}(\sigma \otimes \sigma) = 0.02046$ at $d = 16$.

Spectral threshold ([Fig. 2b](#)). For $d = 8$, $\beta = 5$, $\varepsilon = 0.3$ and $q = (1 + \varepsilon P_\ell(e^\top \cdot))\sigma$ with P_ℓ the Gegenbauer polynomial normalized by $P_\ell(1) = 1$, $\mathcal{U}_\beta(q) = \int \log(1 + \varepsilon \varrho_\ell P_\ell)(1 + \varepsilon P_\ell) d\sigma$ with $\varrho_\ell = \lambda_\ell^\beta / \kappa = I_{\ell+D-1}(\beta)/I_{D-1}(\beta)$, computed with 3000-point Gauss–Jacobi quadrature for the law of $e^\top U$, and $\text{MMD}_\gamma^2 = \varepsilon^2 \|P_\ell\|^2 \Gamma(D) \gamma^{1-D} e^{-2\gamma} I_{\ell+D-1}(2\gamma)$ in closed form (Funk–Hecke).

Heat channel (Fig. 1c–d). We use $q = (1 + \varepsilon f_\ell)\sigma$ with $f_\ell = \sqrt{N_\ell} P_\ell(e^\top \cdot)$, so $\|f_\ell\|_{L^2(\sigma)} = 1$, and restrict to $\varepsilon \|f_\ell\|_\infty < 1$. We evaluate $\mathcal{J}$ from the pair law using the heat-kernel Gegenbauer expansion and Gaussian quadrature, and estimate minimizers by spline interpolation in heat time. Panels (c,d) numerically illustrate the local asymptotic law of Theorem 3.

Estimator bias (Fig. 1b). $d = 16$, $M = 64$ fresh frequencies per batch; 6×10^4 ($B = 16$) and 3×10^4 ($B = 64$) batches; the same batches are reused across s (common random numbers).

Latent-variable study (Section 7). *Data.* Content $c \sim \sigma$ on $\mathbb{S}^7$. (a) *Content–style:* each view has independent style $s_i \sim \mathcal{N}(0, I_4)$, and $x_i = m(c, s_i) \in \mathbb{R}^{16}$. (b) *vMF:* $x_1 = m(c)$, $x_2 = m(\tilde{c})$ with $\tilde{c} \sim \text{vMF}(c, \kappa_{\text{vMF}} = 10)$ (Zimmermann et al., 2021). The mixing map m acts on $(c, s_i) \in \mathbb{R}^{12}$ (vMF: on $c \in \mathbb{R}^8$) and is a fixed random three-layer leaky-ReLU (slope 0.2) network with orthogonal square layers after a Gaussian input layer, hence injective almost surely. Fresh samples are drawn at every step, and the three training seeds share the mixing map. LeJEPA and VICReg hyperparameters are selected on validation samples; all reported metrics, probes and gains are computed on the frozen encoders with independently generated test samples (3000 pairs per model, shared across methods). *Encoder.* MLP $16 \rightarrow 256 \rightarrow 256 \rightarrow 256 \rightarrow 8$ with leaky-ReLU (slope 0.2), shared across views; InfoNCE, SG, SG_α and moment matching act on the normalized outputs. *Optimization.* Adam (default moments, no weight decay or clipping), one-cycle schedule with peak learning rate 10^{-3} (5% warm-up), 8000 steps, $B = 256$ unless stated, three seeds. InfoNCE→ SG_α trains InfoNCE for 4000 steps and SG_α for the remaining 4000 steps of the same schedule. *Losses.* InfoNCE: symmetric NT-Xent with $\tau = 0.2$ ($\beta = 5$). SG: $\hat{\mathcal{J}}_U$ or $\hat{\mathcal{J}}_V$ with $M = 256$ fresh directions and frequencies per step; see *Number of frequencies* below. SG_α : $\hat{\mathcal{J}}_U + \alpha \cdot \frac{1}{4} \|U - V\|^2$. LeJEPA follows the minimal reference implementation: the invariance term I is the mean over views, batch and coordinates of the squared deviation from the view average, and SIGReg is applied to each view separately with 256 random slices resampled at every step, 17 knots on $[0, 3]$ with trapezoidal weights and window $e^{-t^2/2}$, scaled by the batch size; the loss is $(1 - \lambda)I + \lambda$ SIGReg on unnormalized embeddings. VICReg follows the reference implementation on the $d = 8$ outputs, without an expander (variance hinge with target 1 and $\epsilon = 10^{-4}$, squared off-diagonal covariances divided by d). For both we searched one seed over $\lambda \in \{0.005, 0.01, 0.02, 0.05, 0.1\}$ and $(\text{sim}, \text{std}, \text{cov}) \in \{(10, 25, 1), (25, 25, 1), (50, 25, 1)\}$ and selected by validation R_c^2 : $\lambda = 0.01$ and $(10, 25, 1)$, used in both latent models. All methods share the encoder, optimizer, schedule, batch size and number of steps; the other methods use fixed settings. *Metrics* (test samples, after normalizing). $\mathcal{J}$ by the Rao–Blackwellized U-statistic with 5000 frequencies, the radius integrated by generalized Gauss–Laguerre quadrature. $\mathcal{U}_\beta$ and the InfoNCE excess at $\beta = 5$ use all other test samples as negatives, averaged over the two views. R_c^2 : linear regression of c on U , fitted on one half of the test set and evaluated on the other. CKA: linear CKA (Kornblith et al., 2019) of the view-1 embeddings to the InfoNCE encoders of the other seeds. Style: held-out R^2 of an MLP (one hidden layer of width 64) predicting s_1 from (U, c) , fitted on one half and evaluated on the other; we call style detected when $R^2 > 0.02$.

Number of frequencies. With $M \in \{16, 64, 128, 256, 1024\}$ at $B = 256$ (three seeds each, same evaluation protocol as Table 1), the CKA of $\text{SG}_{0.2}$ to InfoNCE is 0.920, 0.968, 0.970, 0.993, 0.993 on vMF and 0.818, 0.827, 0.834, 0.828, 0.825 on content–style; on content–style, the style R^2 of pure SG is 0.29, 0.17, 0.15, 0.11, 0.10. On vMF, the lower means at $M = 64$ and 128 come from one seed that converges to a different solution (CKA 0.92); the same seed reaches 0.99 at $M \geq 256$, the other two seeds reach 0.99 at every $M \geq 64$, and three further seeds at $M = 128$ reach 0.992–0.993 (three further seeds at $M = 256$: 0.994). Thus $M = B/2$ matches $M = 256$ in five of six vMF runs and on content–style.

Style probes: robustness. On the seed-0 encoders of InfoNCE, SG, InfoNCE→SG, $\text{SG}_{0.2}$, LeJEPA ($\lambda = 0.01$) and VICReg (Table 2), a larger probe (MLP 256×256, five-fold cross-validation on 10^4 fresh pairs) gives the same classification, and refits of the reported probe on permuted style targets give $R^2 \leq -0.004$, below the detection threshold 0.02. $\text{SG}_{0.2}$ is reported as the representative price because 0.2 lies well above the critical price $\alpha \approx 0.023$ of Fig. 3a; the conclusions on the price rest on the full sweep.

Measuring the gain (Fig. 3). For a trained encoder, we rescale the style input, $x_i = m(c, \eta s_i)$ with $\eta \in \{0, 0.03, 0.06, 0.1, 0.15, 0.2\}$, and evaluate the objective on 3000 fresh pairs with common frequencies across η ; we repeat this five times with independent pairs and frequencies and report the mean (bars in Fig. 3: one standard deviation over repetitions). At $\eta = 0$ the two views coincide. We fit $F - F(0) = -g\Delta + c_2\Delta^2$ by least squares on $\eta \leq 0.2$ (the comparisons of Fig. 3a,b also hold for a refit on $\eta \leq 0.1$), with $F = \hat{\mathcal{J}}_U$ and $\Delta = \frac{1}{4} \mathbb{E} \|U - V\|^2$ on the normalized embeddings; for LeJEPA, F is the per-sample SIGReg statistic (1024 slices, 8 repetitions with common random numbers) and Δ is replaced by I , which gives g_S . Since s_i and $-s_i$ have the same law, F and Δ are even in η , and $\vartheta = \eta^2$ plays the role of the parameter in Proposition 6, with $\Delta'(0) > 0$ in ϑ whenever the embedding depends on the style at first order: $\eta = 0$ gives the aligned law P_0 of the trained encoder, and the fitted slope estimates

$g = -dF/d\Delta|_{\Delta=0}$ (for LeJEPA, $g_S = -dS/dI|_{I=0}$) along this empirical channel. LeJEPA is trained with the plug-in SIGReg statistic. Applying the identity behind Proposition 4 to each slice, its batch-scaled expectation is BS plus a term that does not grow with B . Measured along the same channel, this term contributes at most 0.33 in absolute value to the gain of each of the 34 encoders, and adding it to Bg_S leaves every classification in Fig. 3b unchanged.

SG $_{\alpha}$ sweep (Fig. 3a). We trained SG_{α} for $\alpha \in \{0, 0.001, 0.003, 0.01, 0.015, 0.02, 0.025, 0.03, 0.05, 0.1, 0.2\}$ with three seeds each (two at $\alpha = 0.001$, five at $\alpha \in \{0.02, 0.025, 0.03\}$). The mean own-channel gain falls from 0.036 at $\alpha = 0$ to 0.006 at $\alpha = 0.2$, and interpolating it places the crossing $g(\alpha) = \alpha$ at $\alpha \approx 0.023$. Style is detected in every seed for $\alpha \leq 0.003$, in some seeds for $\alpha \in \{0.01, 0.015\}$, and in none for $\alpha \geq 0.02$.

LeJEPA (Fig. 3b). At $B = 256$ (22 encoders: one seed at $\lambda = 0.005$, three otherwise; gain range over seeds):

λ	0.005	0.01	0.02	0.03	0.05	0.07	0.1	0.2
price $(1 - \lambda)/\lambda$	199	99	49	32	19	13	9.0	4.0
gain Bg_S	6	5–9	13–15	19–23	22–28	28–40	37–61	73–127
style R^2 max	−0.01	−0.01	−0.02	−0.01	0.14	0.28	0.61	0.83

Batch size. At $B = 64$ and $B = 1024$:

B	λ	price	gain Bg_S	style R^2	kept
64 (2 seeds)	0.03, 0.05, 0.1	32, 19, 9	1.6–4.2	≤ -0.006	no
	0.2	4	9.6, 9.8	0.41, 0.35	yes
1024 (seed 0)	0.005, 0.01	199, 99	51, 90	≤ -0.010	no
	0.02, 0.03	49, 32	127, 221	0.27, 0.41	yes

Warm start (Fig. 3c). For $\alpha = 0, 0.001, 0.003, 0.01, 0.03, 0.2$, the style R^2 is 0.146, 0.081, 0.027, −0.014, −0.018, −0.016 and the CKA to InfoNCE encoders of the other seeds is 0.962, 0.967, 0.971, 0.975, 0.977, 0.975 (InfoNCE seed to seed: 0.982). The own-channel gains of these encoders lie in $[0.0017, 0.0083]$. The switch-point encoders (InfoNCE after 4000 steps; regenerated deterministically, with final branch weights identical to the reported runs) have style R^2 −0.003, 0.057, 0.045, $\mathcal{J} = (1.87, 0.57, 1.37) \times 10^{-3}$ and $\mathcal{J}_{0.2} = (4.3, 3.4, 3.9) \times 10^{-3}$ for the three seeds, evaluated with the same protocol.

Condition (U) in training (Fig. 4, Section 4.5). Moment matching minimizes $\|\mathbb{E}W\|^2 + \|\mathbb{E}WW^T - I/d\|_F^2 + \alpha\Delta$ over the normalized embeddings W of both views, with $\alpha = 1$ (3 seeds); its encoders reach pair cosine 0.98 and $\mathcal{U}_{\beta} = 0.29$ (InfoNCE: 0.021). The Gaussian uniformity loss is $\frac{\beta}{2}\mathbb{E}\|U - V\|^2 + \log \text{mean}_{i \neq j} e^{-\gamma\|w_i - w_j\|^2}$ over the $2B$ embeddings of a batch, with $\gamma \in \{1, 2, 2.5, 5\}$ (2 seeds). The residual $\|\hat{q}_{\ell}\|^2$ is estimated from 3000 test embeddings by the U-statistic $N_{\ell} \text{mean}_{i \neq j} P_{\ell}(w_i^T w_j)$ and averaged over seeds; for moment matching it is 7–63 times InfoNCE’s at every degree $\ell = 3, \dots, 8$. VICReg with invariance weight $a \in \{1, 2.5, 5, 10, 25, 50\}$ (variance and covariance weights 25, 1; 3 seeds, one at $a = 50$) reaches pair cosine 0.91–0.99 and $\mathcal{U}_{\beta} = 0.9$ –2.0 after normalization; style is not detected in any of the 16 encoders.

Code and compute. We will release the code, the training command of every encoder and all evaluated numbers; training is deterministic on CPU. Exact-law and controlled-latent experiments ran on a 2-core CPU; a latent-model run takes 2–13 minutes, and the heat-channel integration for $d = 8$ about 40 minutes.

Table 2 | Content–style model, all metrics (mean $\pm$ s.d. over 3 seeds; Table 1 reports a subset, together with the vMF CKA of Table 3). Here cos is the mean pair cosine $\mathbb{E} U^T V$; $\mathcal{J}$ in units of 10^{-3} ; excess is $\mathcal{L}_\beta - \mathcal{L}_\beta^* = 2\beta\Delta + \mathcal{U}_\beta$ at $\beta = 5$; R_c^2 is the held-out linear decodability of the content; CKA and style as in Table 1. $\hat{\mathcal{J}}_U$ is the unbiased estimator; InfoNCE→SG and InfoNCE→SG_{0.2} switch from InfoNCE halfway through training; λ is LeJEPa’s SIGReg weight.

Method	cos	$\mathcal{J}$	$\mathcal{U}_\beta$	excess	R_c^2	CKA	style
InfoNCE	0.985 ± 0.001	0.17 ± 0.01	$0.02 \pm <0.005$	$0.10 \pm <0.005$	0.77 ± 0.01	$0.98 \pm <0.005$	-0.02 ± 0.01
SG, $\hat{\mathcal{J}}_U$	0.802 ± 0.006	0.85 ± 0.02	0.05 ± 0.01	1.03 ± 0.02	0.57 ± 0.04	0.65 ± 0.03	0.11 ± 0.01
InfoNCE→SG	0.961 ± 0.002	0.12 ± 0.01	$0.02 \pm <0.005$	0.21 ± 0.01	0.75 ± 0.01	$0.96 \pm <0.005$	0.14 ± 0.01
InfoNCE→SG _{0.2}	0.981 ± 0.001	0.27 ± 0.01	$0.04 \pm <0.005$	$0.13 \pm <0.005$	0.76 ± 0.01	$0.98 \pm <0.005$	-0.02 ± 0.01
SG _{0.2}	0.973 ± 0.001	0.98 ± 0.04	$0.16 \pm <0.005$	0.29 ± 0.01	$0.67 \pm <0.005$	0.83 ± 0.01	$-0.02 \pm <0.005$
SG _{0.05}	0.934 ± 0.002	0.79 ± 0.04	$0.12 \pm <0.005$	0.45 ± 0.01	$0.63 \pm <0.005$	0.77 ± 0.01	$-0.02 \pm <0.005$
LeJEPa, $\lambda = 0.01$	$0.956 \pm <0.0005$	1.57 ± 0.03	0.26 ± 0.01	0.48 ± 0.01	$0.56 \pm <0.005$	$0.70 \pm <0.005$	$-0.02 \pm <0.005$
LeJEPa, $\lambda = 0.1$	0.682 ± 0.024	2.20 ± 0.36	0.04 ± 0.01	1.63 ± 0.12	0.49 ± 0.04	0.57 ± 0.03	0.52 ± 0.09
VICReg	0.979 ± 0.002	95 ± 5	1.38 ± 0.06	1.48 ± 0.06	0.47 ± 0.03	0.60 ± 0.05	-0.02 ± 0.01

Table 3 | vMF latent model ($d = 8$, $B = 256$, 8k steps; mean $\pm$ s.d. over 3 seeds). Columns as in Table 2, without style; VICReg weights (sim, std, cov) = (10, 25, 1).

Method	cos	$\mathcal{J}$	$\mathcal{U}_\beta$	excess	R_c^2	CKA
InfoNCE	$0.698 \pm <0.0005$	1.37 ± 0.01	$0.00 \pm <0.005$	$1.51 \pm <0.005$	$0.99 \pm <0.005$	$0.997 \pm <0.0005$
SG, $\hat{\mathcal{J}}_U$	0.668 ± 0.003	1.75 ± 0.08	$0.02 \pm <0.005$	1.67 ± 0.02	0.93 ± 0.01	0.938 ± 0.010
SG, $\hat{\mathcal{J}}_V$	0.667 ± 0.004	1.76 ± 0.08	$0.02 \pm <0.005$	1.68 ± 0.02	0.92 ± 0.01	0.935 ± 0.012
SG _{0.2}	0.694 ± 0.001	1.42 ± 0.03	$0.00 \pm <0.005$	$1.53 \pm <0.005$	$0.99 \pm <0.005$	0.993 ± 0.001
SG _{0.05}	0.686 ± 0.001	1.51 ± 0.03	$0.01 \pm <0.005$	$1.57 \pm <0.005$	$0.97 \pm <0.005$	0.981 ± 0.001
LeJEPa, $\lambda = 0.01$	0.687 ± 0.001	1.54 ± 0.02	$0.01 \pm <0.005$	1.57 ± 0.01	$0.97 \pm <0.005$	0.981 ± 0.003
VICReg, (10, 25, 1)	0.745 ± 0.022	57 ± 18	0.89 ± 0.18	2.16 ± 0.07	0.75 ± 0.02	0.811 ± 0.009

C. Notation

Symbols are listed with the place where they are defined; symbols used only in part of the paper are marked with the sections in which they appear.

Symbol	Meaning	Defined in
<i>Paired laws and the shared loss</i>		
$d, \mathbb{S}^{d-1}, \sigma$	dimension, unit sphere, its uniform law	Section 3
$P, (U, V), q_U, q_V, \bar{q}, q$	paired law, views, marginals, mixture and equal marginal	Section 3
$\Delta(P)$	misalignment $\frac{1}{4}\mathbb{E}\ U - V\ ^2$	(1)
C, R, Z	centroid, χ_d radius, $Z = RC$	(2)
$\varphi_Z, \mathcal{J}(P)$	characteristic function of Z , SG loss	(3)
a, t	frequency direction and scale	(3)
H, Q	$\frac{1}{2}(U - V), \ H\ ^2$	Section 4.1
j, k_0, Δ^*	alignment bound, $k_0 = 2 \cdot 3^{3/4}, \min\{1, j^{-1}\}$	Lemma 1, Theorem 1
$D_{\text{lift}}, K_{\text{lift}}$	lifted discrepancy and its zonal kernel	Section 4.3, Appendix A.2
$\mathcal{J}_\alpha (\text{SG}_\alpha)$	$\mathcal{J} + \alpha \Delta$	(12)
<i>InfoNCE</i>		
$\beta = 1/\tau, \mathcal{L}_\beta, \mathcal{L}_\beta^*$	inverse temperature, population InfoNCE, its minimum	(4)
$\kappa_d(\beta), \mathcal{U}_\beta(q)$	$\mathbb{E}_{W \sim \sigma} e^{\beta u^\top W}$, uniformity excess	(5)
k_β, D_β	kernel $e^{\beta u^\top v}$ and its MMD	Sections 4.3 and 4.5
$C_{d,\beta}^*, \bar{c}_{d,\beta}, c_{\beta,\alpha}$	kernel-comparison and certificate constants	Proposition 7, Theorem 1, Proposition 8
<i>Finite batches</i>		
$B, M, \omega_m = t_m a_m$	batch size, number of frequencies, frequencies	Section 5
$\hat{\phi}_m, g_m, \hat{\mathcal{J}}_V, \hat{\mathcal{J}}_U$	empirical and target characteristic functions, estimators	Section 5, (10)
<i>Nuisance and price</i>		
$\vartheta, \mathcal{J}_q(\vartheta), \mathcal{J}_{\text{unif}}(q)$	heat time, SG loss along the channel, its value at $\vartheta = 0$	Section 6.1
$G_\ell, \rho(f), \rho_\ell, \Delta_{\text{eq}}$	first-variation weights, equilibrium ratio (ρ_ℓ : f of pure degree ℓ), equilibrium misalignment	Theorem 3, (11)
g, α_c	gain of a channel, critical price	(11), Proposition 6, Corollary 3
(A)	alignment price above the gain, $\alpha > g$	Section 6.3, Proposition 6
$\lambda, S, I, g_S, \lambda_c$	LeJEPa weight, SIGReg and invariance terms, gain, critical weight	Proposition 6
$W, \Lambda, c(\Lambda)$	shared or mean code, latent, length of the conditional mean	Sections 6.1 and 6.3
<i>Spherical harmonics and uniformity tests (from Section 4.5 on)</i>		
$\hat{q}_\ell$	degree- ℓ harmonic component of $q - \sigma$	Section 4.5
$\mathcal{H}_\ell, N_\ell, Y_{\ell m}, P_\ell$	harmonics of degree ℓ , dimension, basis, Gegenbauer polynomial	Appendix A.8
$\lambda_\ell^\beta, \lambda_\ell^{\text{lift}}, \mu_\ell, S_K$	eigenvalues of k_β, K_{lift} , a zonal kernel K ; spectral ratio	Section 4.5, Theorem 2, Appendix A.8
$Q_\beta, \delta, c_K, (U)$	quadratic approximation of $\mathcal{U}_\beta$, density deviation, best linear constant, spectral dominance $S_K < \infty$	Theorem 2
$D = d/2, \nu = D - 1$	half-dimension, Bessel index	Appendix A.2, Appendix A.8
$g(t) = e^{-t^2/2}, T_\mu, \psi_u$	target characteristic function (not the gain g), lifted variable, feature map	Appendix A.2
k_d, K_d, Γ_q	centered-channel constant, gain constant, gain density	Appendix A.15