

Sharp Asymptotic Theory of Maximum Likelihood Estimation for Gaussian Processes with an RBF Kernel

Ameer Qaqish* and Didong Li†

Department of Biostatistics, University of North Carolina at Chapel Hill

Abstract

Gaussian processes (GPs) are widely used across machine learning, spatial statistics, time-series analysis, optimization, Bayesian statistics, and scientific applications. A central component of a GP model is its kernel, which is typically specified through a parametric family. Among the most widely used choices is the radial basis function (RBF), also known as the squared exponential or Gaussian kernel, owing to its simple form, smoothness, and flexibility. In practice, the kernel parameters are routinely estimated by the maximum likelihood estimators (MLEs), as implemented by standard GP software. Despite this widespread use, the asymptotic behavior of the MLEs remains poorly understood under fixed-domain asymptotics, even for the RBF kernel. The main difficulty arises from the increasingly strong dependence among densely sampled observations and the nonlinear dependence of the covariance matrix on the kernel parameters. In this paper, we address this gap by providing, to the best of our knowledge, the first complete asymptotic characterization of the joint MLE of the spatial variance, lengthscale, and nugget variance under fixed-domain asymptotics. We establish consistency, derive convergence rates for all three parameters, prove joint asymptotic normality, and show that these rates are minimax optimal.

Keywords: Gaussian process; RBF kernel; maximum likelihood; fixed-domain asymptotics; minimax optimal.

Contents

1	Introduction	3
2	Background and related work	5
2.1	Background	5
2.2	Related work on RBF parameter inference	6

*ameer@unc.edu

†Corresponding author: didongli@unc.edu. The authors used GPT-6 and Claude Opus 5.5 to assist with proofs and manuscript polishing, and Claude Opus 5.5 to assist with coding. The authors independently verified the results and take full responsibility for the content of this paper.

3	Theory	7
3.1	Problem setup, assumptions, and notations	7
3.2	MLE asymptotics	8
3.3	Minimax optimality	10
3.4	Some useful corollaries	11
3.4.1	Log scale	11
3.4.2	Confidence intervals	12
3.4.3	Random observation locations	13
3.5	Proof outline and key ideas	14
4	Simulation	16
4.1	Simulation setup	16
4.2	Simulation results	17
5	Discussion and future work	21
A	Lemmas	24
A.1	Gaussian affinity and elementary bounds	24
A.2	A maximal inequality for continuous processes	25
A.3	Auxiliary lemmas	26
B	Proofs of the main results	30
B.1	Proof of Theorem 3.1	30
B.2	Proof of the uniformity in θ_0 in Theorem 3.2	32
B.3	Proof of Theorem 3.3	33
B.4	Proofs of the Corollaries in Section 3.4	34
B.4.1	Proof of Theorem 3.4	34
B.4.2	Proof of Theorem 3.5	35
B.4.3	Proof of Theorem 3.6	36
C	Proofs of the lemmas in Appendix A	38
C.1	Proof of Theorem A.1	38
C.2	Proof of Theorem A.2	39
C.3	Proof of Theorem A.3	39
C.4	Proof of Theorem A.4	39
C.5	Proof of Theorem A.5	40
C.6	Proof of Theorem A.6	40
C.7	Proof of Theorem A.7	41
C.8	Proof of Theorem A.8	41
C.9	Proof of Theorem A.9	45
C.10	Proof of Theorem A.10	50
C.11	Proof of Theorem A.11	53
C.12	Proof of Theorem A.12	53
C.13	Proof of Theorem A.13	54
C.14	Proof of Theorem A.14	55

C.15 Proof of Theorem A.16	57
C.16 Proof of Theorem A.17	58
D Additional simulation details	59
D.1 Exact evaluation of the likelihood	59
D.2 Data generation	60
D.3 Computation of the MLE	61
D.4 Computation of the Fisher information	61
D.5 Monte Carlo summaries	61
D.6 Additional normality diagnostics	62
D.7 Finite-sample behavior of the nugget	62
D.8 Finite-sample behavior of the spatial parameters	62
D.9 Validation against dense computations	64
D.10 Monte Carlo implementation and software	64
D.11 Results on the original scale	64

1 Introduction

Gaussian processes (GPs) are a fundamental class of probabilistic models with broad applications across statistics, machine learning, and the sciences. In machine learning, GPs provide flexible nonparametric models for regression, classification, and nonlinear dimension-reduction methods [Rasmussen and Williams, 2006]. In spatial statistics and time-series analysis, GPs provide a natural framework for modeling spatial [Banerjee et al., 2025] and temporal dependence [Roberts et al., 2013]. They are also widely used as surrogate models in Bayesian optimization [Garnett, 2023], and, more generally, as flexible prior distributions over unknown functions in Bayesian inference [Ghosal and van der Vaart, 2017]. Applications have further extended to biomedical problems, such as spatial transcriptomics [Svensson et al., 2018, Weber et al., 2023].

A key component of a GP is its covariance function, also known as the kernel function, which determines the dependence and smoothness of the underlying random function [Rasmussen and Williams, 2006]. Among the most commonly used choices is the radial basis function (RBF) kernel, also known as the squared exponential or Gaussian kernel. Its popularity stems from its simple form and its smoothness. In this paper, we consider the RBF kernel

$$K_{\sigma^2, l}(x, x') := \sigma^2 \exp\left(-\frac{\|x - x'\|^2}{2l^2}\right), \quad (1.1)$$

where σ^2 is the spatial variance, which controls the magnitude of spatial variation, and l is the lengthscale, which determines the scale over which the process remains correlated. We observe the GP at n locations with additive, independent Gaussian measurement errors of variance τ^2 , known as the nugget; the precise observation model is given in Section 2.1. In practice, the parameter $\theta = (\sigma^2, l, \tau^2)$ is routinely estimated by maximizing the Gaussian likelihood, leading to the maximum likelihood estimator (MLE). This is the standard approach implemented in widely used GP software, including GPML [Rasmussen and Nickisch, 2010] and `scikit-learn` [Pedregosa et al., 2011] in Python and DiceKriging [Roustant et al., 2012] and GPfit [MacDonald et al., 2015] in R.

However, the theoretical understanding of the MLEs is considerably less developed than their widespread practical use would suggest [Stein, 1999]. Throughout this paper, we focus on fixed-domain asymptotics, where observations become increasingly dense within a fixed bounded domain as the sample size increases. Several recent results address related aspects of this problem. Qaqish and Li [2025] established the identifiability of all parameters of the RBF kernel, showing that consistent estimation of these parameters is at least possible in principle. Identifiability alone, however, does not establish the consistency or asymptotic distribution of any particular estimator. Moving from identifiability to estimation, Loh and Lam [2000] established consistency of sieve MLEs for smooth Gaussian random fields, including the RBF case, under regular-grid sampling and a restricted parameterization. More recently, Loh [2026] constructed consistent estimators of RBF parameters from noiseless scattered observations using local differences. These estimators, however, are not MLEs. Xu and Stein [2017] studied likelihood estimation for a smooth Gaussian random field model when the observations are generated from a deterministic function rather than from the assumed GP. From a different perspective, Karvonen and Oates [2023] showed that lengthscale MLE can be ill-posed in noiseless GP regression. Despite these developments, a general asymptotic theory for the joint MLE of the RBF spatial variance, lengthscale, and nugget variance remains unavailable.

Consequently, even for the RBF kernel, a basic statistical question remains unresolved: *what is the asymptotic behavior of the MLE that practitioners actually compute?* In particular, it is not even known in general whether the joint MLE of $\theta = (\sigma^2, l, \tau^2)$ is consistent, at what rates its three components converge, or whether it admits an asymptotically normal distribution. The difficulty is fundamentally different from standard likelihood problems based on independent or weakly dependent observations. As the sampling locations become increasingly dense, the observations arise from a single realization of the underlying GP and become strongly dependent. At the same time, the covariance matrix is a nonlinear function of the kernel parameters, and its dimension and dependence structure both change with n . Standard likelihood arguments based on sums of approximately independent contributions therefore do not apply directly.

In this paper, we resolve this problem for the RBF kernel. We establish consistency of the joint MLE of θ . We further derive the convergence rates of all three parameters, characterize the Fisher information, and establish joint asymptotic normality. In particular, we show that the MLEs of the spatial variance σ^2 , lengthscale l , and nugget variance τ^2 converge at rates

$$\left(\frac{\log n}{\log \log n}\right)^{-p/2}, \quad \left(\frac{\log n}{\log \log n}\right)^{-(p+2)/2}, \quad n^{-1/2},$$

respectively, where p is the dimension of the input domain. We further establish matching minimax lower bounds and show that the MLE attains these bounds uniformly over the parameter space. Thus, the above three convergence rates are minimax optimal. The theory accommodates irregular observation locations under a polynomial coverage condition and does not require a regular grid or a minimum separation condition. Numerical experiments complement the theoretical results and examine the convergence rates empirically.

Beyond the specific asymptotic results, a key methodological contribution of this work is a general strategy for statistical inference with strongly dependent observations. The central idea is to construct a function of the observations that is a lower triangular linear transformation of

i.i.d. standard normals, while controlling the approximation error and showing that it vanishes sufficiently fast. This converts a difficult dependent-data problem into one that can be analyzed using tools closer to those for independent observations. In the RBF setting studied here, the analyticity of the kernel allows us to construct such a transformation through a growing collection of Taylor coefficients, whose approximation errors decay rapidly with the sample size. The resulting representation makes the information carried by different kernel parameters explicit and provides the key mechanism for establishing consistency, convergence rates, and asymptotic normality of the MLEs. More broadly, this dependence-to-independence transformation provides a potentially useful strategy for asymptotic inference in other strongly dependent models whenever an appropriate latent representation can be recovered with sufficiently small error.

The remainder of the paper is organized as follows. [Section 2](#) reviews background on fixed-domain GP inference and related work on kernel parameter inference. [Section 3](#) presents the main theoretical results for the joint MLEs. [Section 4](#) reports simulation studies supporting the rates. [Section 5](#) discusses implications, limitations, and directions for future work. All proofs and additional experimental details are provided in the Appendix.

2 Background and related work

2.1 Background

Definition 1 (GP). Let $\mathcal{X} \subset \mathbb{R}^p$ be an input domain. A stochastic process $\{F(x) : x \in \mathcal{X}\}$ is a GP, denoted by $F \sim \text{GP}(m, K)$, if, for every finite collection $x_1, \dots, x_n \in \mathcal{X}$, the random vector

$$(F(x_1), \dots, F(x_n))^\top \sim N(\mu, \Sigma),$$

where $\mu = [m(x_1), \dots, m(x_n)]^\top$ and $\Sigma_{ij} = K(x_i, x_j)$. m is called the mean function while K is called the covariance function or kernel function, which together uniquely determine the GP.

Throughout this paper, we assume that the mean function m is known to be zero and focus on inference for the kernel parameters of RBF as defined in [\(1.1\)](#). We observe noisy outcomes

$$Y_{n,i} = F(x_{n,i}) + \varepsilon_{n,i}, \quad i = 1, \dots, n, \tag{2.1}$$

at deterministic locations $x_{n,1}, \dots, x_{n,n} \in \mathcal{X}$, where $\varepsilon_{n,i} \stackrel{\text{iid}}{\sim} N(0, \tau^2)$ and the errors are independent of F . We first develop the theory for deterministic designs satisfying a weak coverage condition; random designs, including independent sampling from densities bounded away from zero on a region of $\mathcal{X}$, are treated in [Theorem 3.6](#). For the deterministic-design formulation, the covariance matrix of the observations is

$$C_n(\theta) = \sigma^2 R_n(l) + \tau^2 I_n, \quad \theta = (\sigma^2, l, \tau^2),$$

where

$$[R_n(l)]_{ij} = \exp\left(-\frac{\|x_{n,i} - x_{n,j}\|^2}{2l^2}\right).$$

Thus,

$$Y_n = (Y_{n,1}, \dots, Y_{n,n})^\top \sim N_n(0, C_n(\theta)).$$

We study fixed-domain asymptotics: the domain $\mathcal{X} \subset \mathbb{R}^p$ remains fixed and bounded while the number of observation locations increases. This regime differs from increasing-domain asymptotics, in which the observation region itself expands with n [Stein, 1999]. Under fixed-domain asymptotics, increasingly many observations correspond to a single realization of F over the same bounded region, resulting in strong dependence among nearby observations. Whether kernel parameters can be consistently estimated in this regime depends critically on the kernel family, which motivates the identifiability and estimation literature reviewed next.

2.2 Related work on RBF parameter inference

Parameter estimation under fixed-domain asymptotics differs substantially across kernel families. A fundamental issue is whether different kernel parameters induce equivalent or orthogonal Gaussian measures. Parameters that can be distinguished in this sense are commonly referred to as microergodic parameters [Stein, 1999]. For the RBF kernel, Qaqish and Li [2025] showed that all RBF parameters are identifiable. This result, however, concerns the distinguishability of the induced Gaussian measures and does not establish the behavior of a particular estimator.

Several works have studied RBF parameter inference. Loh and Lam [2000] considered an RBF GP observed on a regular lattice and derived an explicit likelihood using the special structure of the resulting correlation matrix. They established strong consistency of sieve MLEs for the kernel parameters. This was an important early consistency result for likelihood-based estimation of smooth kernels but its scope is substantially different from the problem considered here. In particular, the analysis relies on a highly structured regular design and on a sieve likelihood construction tailored to that setting. In contrast, the MLE routinely used in GP software is obtained by directly maximizing the Gaussian likelihood over a prescribed parameter space, and observations in applications are rarely restricted to a regular lattice. Our goal is to characterize this exact MLE under substantially more general observation designs.

More recently, Loh [2026] constructed consistent estimators of the spatial variance and lengthscale of the RBF kernel from noiseless observations at scattered, space-filling locations. The result substantially relaxes the regular-grid requirement and establishes directly that these parameters can be consistently recovered. However, the estimators are specifically constructed for the theoretical analysis and are not obtained by maximizing the Gaussian likelihood. Moreover, the observations are noiseless, corresponding to $\tau^2 = 0$ in our notation. Measurement error is present in real data applications, and introducing an unknown nugget changes both the likelihood and the asymptotic problem. Thus, while Loh [2026] establishes the existence of consistent estimators, it does not determine whether the routinely used MLE is consistent or characterize its asymptotic behavior.

Xu and Stein [2017] studied a different question motivated by computer experiments. There, the observations are values of a deterministic function f , while an RBF GP is used as a statistical model for interpolation. Consequently, there is no true covariance parameter θ_0 generating the observations, and consistency toward a true RBF parameter is not the relevant question. For regularly spaced observations on $[0, 1]$ with the lengthscale fixed, they showed that the MLE of the spatial variance converges to zero when f is constant and diverges to infinity when f is linear; they also derived asymptotic orders for related polynomial examples. Their results therefore characterize the behavior of likelihood estimation under this deterministic-function formulation rather than estimation of the kernel parameters of a correctly specified RBF GP. Indeed, Xu and Stein [2017] explicitly noted that

establishing estimation theory when the GP model is correctly specified remained a substantially different and difficult problem.

Finally, [Karvonen and Oates \[2023\]](#) studied the well-posedness of MLE-based GP regression rather than consistency of covariance parameter estimators. They showed that, in the noiseless setting, estimating the lengthscale by maximum likelihood can make the resulting predictive distribution non-Lipschitz with respect to perturbations of the observed data. In particular, for certain data configurations the likelihood drives the estimated lengthscale to infinity. This notion of ill-posedness does not contradict the consistency result established in this paper. Their result concerns sensitivity of the fitted predictor to perturbations of a fixed noiseless data set, whereas our results concern the probabilistic behavior of the MLE when the data are generated from an RBF GP with noise as n increases. The nugget is particularly important: it keeps the covariance matrices uniformly nonsingular and excludes the noiseless setting in which their negative result occurs.

Taken together, the existing literature establishes identifiability of the RBF parameters, consistency of specially constructed estimators, consistency of sieve MLEs under highly structured designs, and both positive and negative results for MLEs in other formulations. None of these results characterizes the joint MLE of $\theta = (\sigma^2, l, \tau^2)$ under the observation model in (2.1). In particular, consistency, convergence rates, and asymptotic normality of this MLE remain unresolved. These are the questions addressed in [Section 3](#).

3 Theory

This section states the model and assumptions in [Section 3.1](#), establishes the asymptotic behavior in [Section 3.2](#) and minimax optimality of the MLE in [Section 3.3](#), with a few useful corollaries in [Section 3.4](#), and gives a sketch of the proof in [Section 3.5](#). Complete proofs are deferred to the Appendices [A](#), [B](#), and [C](#).

3.1 Problem setup, assumptions, and notations

Recall from [Section 2.1](#) that

$$Y_n \sim N_n(0, C_n(\theta)), \quad C_n(\theta) = \sigma^2 R_n(l) + \tau^2 I_n, \quad \theta = (\sigma^2, l, \tau^2).$$

For notational simplicity, in the theoretical analysis we write

$$(s, l, t) = (\sigma^2, l, \tau^2),$$

so that $\theta = (s, l, t) \in \Theta \subset (0, \infty)^3$ and $C_n(\theta) = sR_n(l) + tI_n$. We also write $K_\theta = K_{s,l}$ for the RBF kernel (1.1), which does not depend on t . Let $\theta_0 = (s_0, l_0, t_0)$ denote the true parameter. We estimate θ by maximizing the Gaussian likelihood over Θ . Ignoring an additive constant, the log likelihood is

$$l_n(\theta) = -\frac{1}{2} \log \det C_n(\theta) - \frac{1}{2} Y_n^\top C_n(\theta)^{-1} Y_n, \quad (3.1)$$

and the MLE is

$$\hat{\theta}_n \in \arg \max_{\theta \in \Theta} l_n(\theta).$$

We now state the two assumptions of our theoretical analysis.

Assumption 1 (Positive compact parameter set). The parameter space is

$$\Theta = [s_-, s_+] \times [l_-, l_+] \times [t_-, t_+] \subseteq (0, \infty)^3,$$

i.e., Θ is compactly contained in $(0, \infty)^3$, and assume that $\theta_0 \in \text{int}(\Theta)$.

For a fixed closed cube $Q \subset \mathcal{X}$, define its fill distance by

$$h_n(Q) = \sup_{x \in Q} \min_{1 \leq i \leq n} \|x - x_{n,i}\|.$$

Assumption 2 (Polynomial coverage). There exist a fixed closed cube $Q \subset \mathcal{X}$ with positive side length and constants $C_h < \infty$ and $\kappa > 0$ such that, for all sufficiently large n ,

$$h_n(Q) \leq C_h n^{-\kappa}. \quad (3.2)$$

[Assumption 2](#) requires the observation locations to become sufficiently dense on at least one fixed region of the domain, but imposes no minimum spacing condition. In particular, irregular, clustered, and repeated observation locations are allowed. Necessarily $\kappa \leq 1/p$, since n balls of radius $h_n(Q)$ must cover a set of positive p -dimensional volume.

We finally introduce the Fisher information and the normalization used in our asymptotic results. Let subscripts on C_n denote parameter derivatives. At θ_0 , define the Fisher information matrix

$$[\mathcal{I}_n(\theta_0)]_{ab} = \frac{1}{2} \text{tr} \left(C_{n,0}^{-1} C_{n,a}(\theta_0) C_{n,0}^{-1} C_{n,b}(\theta_0) \right), \quad a, b \in \{s, l, t\}, \quad (3.3)$$

where $C_{n,0} = C_n(\theta_0)$. For $n \geq 16$, let

$$b_n = \frac{\log n}{\log \log n}, \quad D_n = \text{diag}(b_n^{p/2}, b_n^{p/2+1}, \sqrt{n}), \quad J_n = D_n^{-1} \mathcal{I}_n(\theta_0) D_n^{-1}, \quad \epsilon_n = b_n^{-p/2}.$$

Thus, J_n is the normalized Fisher information, with the normalization D_n corresponding to the convergence rates of the three kernel parameters that will be established below.

Throughout this article, constants denoted by c, C may change between displays and can depend on the fixed parameter rectangle, dimension, bounded domain, cube, and coverage constants, but not on n . For symmetric matrices, $\preceq$ is the Loewner order. The symbols $\|\cdot\|_{\text{op}}$ and $\|\cdot\|_{\text{F}}$ denote operator and Frobenius norms. For a multiindex $\alpha \in \mathbb{N}_0^p$, write $|\alpha| = \sum_j \alpha_j$, $\alpha! = \prod_j \alpha_j!$, and $x^\alpha = \prod_j x_j^{\alpha_j}$.

3.2 MLE asymptotics

Our main result is the following.

Theorem 3.1 (MLE asymptotics). Under [Assumptions 1](#) and [2](#), the following assertions hold.

(i) **Fisher information.** The normalized Fisher information is uniformly nondegenerate:

$$cI_3 \preceq J_n \preceq CI_3 \quad (3.4)$$

for all sufficiently large n . Its spatial block $J_{n,\text{sp}}$ satisfies $cI_2 \preceq J_{n,\text{sp}} \preceq CI_2$, and

$$J_n = \begin{pmatrix} J_{n,\text{sp}} & o(1) \\ o(1) & (2t_0^2)^{-1} + o(1) \end{pmatrix}. \quad (3.5)$$

(ii) **Consistency and convergence rates.** The MLE $\hat{\theta}_n$ admits the expansion

$$D_n(\hat{\theta}_n - \theta_0) = J_n^{-1} D_n^{-1} \nabla l_n(\theta_0) + o_{\mathbb{P}}(1). \quad (3.6)$$

In particular, $\hat{\theta}_n$ is consistent with rates

$$\hat{s}_n - s_0 = O_{\mathbb{P}}(b_n^{-p/2}), \quad \hat{l}_n - l_0 = O_{\mathbb{P}}(b_n^{-(p+2)/2}), \quad \hat{t}_n - t_0 = O_{\mathbb{P}}(n^{-1/2}). \quad (3.7)$$

(iii) **Asymptotic normality.** Let M_n be any 3×3 matrices with $M_n^{\top} M_n = \mathcal{I}_n(\theta_0)$. Then

$$M_n(\hat{\theta}_n - \theta_0) \implies N_3(0, I_3). \quad (3.8)$$

Moreover,

$$\sqrt{n}(\hat{t}_n - t_0) \implies N(0, 2t_0^2). \quad (3.9)$$

(iv) **Exponential tail bounds.** Write $\hat{h}_n = D_n(\hat{\theta}_n - \theta_0) = (\hat{h}_{n,\text{sp}}, \hat{h}_{n,t})$, where $\hat{h}_{n,\text{sp}} \in \mathbb{R}^2$ collects the two spatial coordinates s and l . There are constants $c, C > 0$ and n_0 such that, for all $n \geq n_0$ and all $H \geq 0$,

$$\mathbb{P}_{\theta_0,n}\{\|\hat{h}_{n,\text{sp}}\| \geq H\} \leq C \exp\{-c \min(H^2, b_n^{p/2} H)\}, \quad \mathbb{P}_{\theta_0,n}\{|\hat{h}_{n,t}| \geq H\} \leq C e^{-cH^2}. \quad (3.10)$$

(v) **Moment convergence.** The normalized MLE has uniformly bounded exponential moments: for every $\lambda > 0$ there is n_λ such that

$$\sup_{n \geq n_\lambda} \mathbb{E}_{\theta_0,n} \exp\{\lambda \|D_n(\hat{\theta}_n - \theta_0)\|\} < \infty. \quad (3.11)$$

Consequently, let $Z \sim N_3(0, I_3)$, and let $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ be continuous with $|f(x)| \leq C e^{\lambda \|x\|}$ for some constants $C, \lambda > 0$. Then, for every choice of M_n in (iii),

$$\mathbb{E}_{\theta_0,n} f(D_n(\hat{\theta}_n - \theta_0)) - \mathbb{E} f(J_n^{-1/2} Z) \rightarrow 0, \quad \mathbb{E}_{\theta_0,n} f(M_n(\hat{\theta}_n - \theta_0)) \rightarrow \mathbb{E} f(Z). \quad (3.12)$$

In particular, all moments converge, and

$$D_n \mathbb{E}_{\theta_0,n}(\hat{\theta}_n - \theta_0) \rightarrow 0, \quad D_n \mathbb{E}_{\theta_0,n}[(\hat{\theta}_n - \theta_0)(\hat{\theta}_n - \theta_0)^{\top}] D_n - J_n^{-1} \rightarrow 0. \quad (3.13)$$

Theorem 3.1 provides a complete asymptotic characterization of the MLE. All three kernel parameters are consistently estimated, but at substantially different rates. The nugget variance t has the standard parametric rate $n^{-1/2}$, whereas the spatial variance s and lengthscale l converge at the slower logarithmic rates. We show in [Section 3.3](#) that these rates are minimax optimal.

The MLE is also jointly asymptotically normal after normalization by the Fisher information.

An important feature of [Theorem 3.1](#) is that it does not require the normalized Fisher information J_n to converge. If, for a particular design sequence, $J_{n,\text{sp}} \rightarrow J_{\text{sp}}$, then it also implies

$$D_n(\hat{\theta}_n - \theta_0) \implies N_3 \left(0, \begin{pmatrix} J_{\text{sp}}^{-1} & 0 \\ 0 & 2t_0^2 \end{pmatrix} \right).$$

Thus, when J_n converges, the MLE admits a conventional asymptotic normal distribution under the deterministic normalization D_n . When it does not, the Fisher-normalized result in part (iii) continues to provide a valid asymptotic normal approximation for the actual observation design.

Beyond convergence in probability and distribution, parts (iv) and (v) provide exponential tail bounds and moment convergence for the normalized MLE. In particular, the moment convergence allows the stochastic rates in part (ii) to be translated into corresponding MSE rates, which are used in [Section 3.3](#) to establish minimax optimality.

Remark 3.2 (Uniformity in θ_0). Inspection of the proofs shows that parts (i) and (iv) also hold uniformly in $\theta_0 \in \Theta$. Parts (ii), (iii), and the limits in part (v) hold uniformly over θ_0 in compact subsets K of $\text{int}(\Theta)$. For example, writing $\mathcal{L}_{\theta_0}$ for the law under θ_0 and d_{BL} for the bounded Lipschitz distance,

$$\sup_{\theta_0 \in K} d_{\text{BL}} \{ \mathcal{L}_{\theta_0} (M_n(\hat{\theta}_n - \theta_0)), N_3(0, I_3) \} \rightarrow 0.$$

A sketch of the proofs of these uniform claims is given in [Appendix B.2](#).

Uniformity is what makes these asymptotic statements usable for inference when θ_0 is unknown. It ensures that Wald confidence intervals based on $\mathcal{I}_n(\theta_0)^{-1}$ (or $\mathcal{I}_n(\hat{\theta}_n)^{-1}$, as will be shown later) attain their nominal coverage asymptotically, uniformly over $\theta_0 \in K$, rather than only at each fixed parameter value.

3.3 Minimax optimality

The convergence rates in [Theorem 3.1](#) are logarithmic for the spatial variance s and lengthscale l . In fact, the logarithmic rates also arise for the specially constructed estimators of [Loh \[2026\]](#) for noiseless observations, whose established rates are slower than those obtained here for the MLE. A natural question is therefore whether the rates in [Theorem 3.1](#) can be improved by using a different estimator. The following result shows that they cannot: the rates in [Theorem 3.1](#) are minimax optimal for all three parameters, and the MLE attains these optimal rates.

Theorem 3.3 (Minimax optimality). Assume [Assumption 1](#). Let $d_{n,j} = D_{n,jj}$ for $j \in \{s, l, t\}$. For each $j \in \{s, l, t\}$, there exist constants $a > 0$ and $c_0 > 0$ such that, for every fixed $\theta_0 \in \text{int}(\Theta)$,

$$\theta_{1,n}^{(j)} = \theta_0 + a d_{n,j}^{-1} e_j \in \Theta$$

for all sufficiently large n , and

$$\inf_{\theta_{n,j}} \max_{\theta \in \{\theta_0, \theta_{1,n}^{(j)}\}} \mathbb{P}_{\theta,n} \left\{ |\tilde{\theta}_{n,j} - \theta_j| \geq \frac{a}{2d_{n,j}} \right\} \geq c_0. \quad (3.14)$$

The other two parameters are identical under the two alternatives, so this lower bound continues to

hold even when they are known. Consequently,

$$\inf_{\tilde{\theta}_{n,j}} \sup_{\theta \in \Theta} \mathbb{E}_{\theta,n}(\tilde{\theta}_{n,j} - \theta_j)^2 \geq c d_{n,j}^{-2}. \quad (3.15)$$

If [Assumption 2](#) also holds, then the MLE satisfies

$$\sup_{\theta \in \Theta} \mathbb{E}_{\theta,n}(\hat{\theta}_{n,j} - \theta_j)^2 \leq C d_{n,j}^{-2}. \quad (3.16)$$

Therefore, for each parameter $j \in \{s, l, t\}$,

$$\inf_{\tilde{\theta}_{n,j}} \sup_{\theta \in \Theta} \mathbb{E}_{\theta,n}(\tilde{\theta}_{n,j} - \theta_j)^2 \asymp d_{n,j}^{-2}, \quad (3.17)$$

and the MLE attains the minimax rate. Equivalently, the minimax root MSE rates for (s, l, t) are

$$b_n^{-p/2}, \quad b_n^{-(p+2)/2}, \quad n^{-1/2},$$

respectively.

3.4 Some useful corollaries

The preceding results characterize the asymptotic behavior and prove minimax optimality of the MLE. We next give three corollaries that connect this theory more directly to statistical practice. They show that the theory carries over to the commonly used log parametrization ([Theorem 3.4](#)), justify confidence intervals based on estimated Fisher information rather than the Fisher information at the unknown true parameter ([Theorem 3.5](#)), and extend the deterministic-design theory to random designs ([Theorem 3.6](#)). All proofs are given in [Appendix B.4](#).

3.4.1 Log scale

As all three parameters are positive, GP implementations commonly optimize them on the log scale, which automatically enforces positivity. Our computations and simulations in [Section 4](#) therefore use $\lambda = \log \theta = (\log s, \log l, \log t)$, coordinatewise. By equivariance, the MLE of λ is $\hat{\lambda}_n = \log \hat{\theta}_n$. Let $\Lambda_0 = \text{diag}(s_0, l_0, t_0)$. Since $\partial_{\log \theta_j} = \theta_j \partial_{\theta_j}$, the score and the Fisher information for λ at λ_0 are $\Lambda_0 \nabla l_n(\theta_0)$ and

$$\mathcal{I}_n^{\log}(\theta_0) = \Lambda_0 \mathcal{I}_n(\theta_0) \Lambda_0 = D_n J_n^{\log} D_n, \quad J_n^{\log} = \Lambda_0 J_n \Lambda_0,$$

because the diagonal matrices Λ_0 and D_n commute.

Corollary 3.4 (Log scale). Under [Assumptions 1](#) and [2](#), all assertions of [Theorem 3.1](#) and [Theorem 3.3](#) hold for $\hat{\lambda}_n$, with $\hat{\theta}_n - \theta_0$ replaced by $\hat{\lambda}_n - \lambda_0$, $\mathcal{I}_n(\theta_0)$ by $\mathcal{I}_n^{\log}(\theta_0)$, J_n by $J_n^{\log}$, and $\nabla l_n(\theta_0)$ by $\Lambda_0 \nabla l_n(\theta_0)$, after adjusting the constants c and C . In particular, $M_n(\hat{\lambda}_n - \lambda_0) \Rightarrow N_3(0, I_3)$ whenever $M_n^\top M_n = \mathcal{I}_n^{\log}(\theta_0)$,

$$\sqrt{n}(\log \hat{t}_n - \log t_0) \implies N(0, 2),$$

and

$$\mathbb{E}(\log \hat{s}_n - \log s_0)^2 \asymp b_n^{-p}, \quad \mathbb{E}(\log \hat{l}_n - \log l_0)^2 \asymp b_n^{-(p+2)}, \quad n \mathbb{E}(\log \hat{t}_n - \log t_0)^2 \longrightarrow 2.$$

The uniformity statements of [Theorem 3.2](#) also carry over.

Thus, working on the log scale does not change the convergence rates or the first-order asymptotic theory. It has additional advantages of respecting positivity and yielding a parameter-free limiting variance for the nugget, and, as the simulations in [Section 4](#) illustrate, can provide a better normal approximation at moderate sample sizes.

3.4.2 Confidence intervals

Asymptotic normality is useful for practical statistical inference only when the asymptotic covariance matrix can be estimated from the data. In particular, to report standard errors or confidence intervals for the kernel parameters, the Fisher information at the unknown true parameter θ_0 needs to be replaced by an estimated quantity. The next corollary shows that both the expected and observed information evaluated at the MLE $\hat{\theta}_n$ are valid replacements.

Corollary 3.5 (Estimated information and confidence intervals). Under [Assumptions 1](#) and [2](#), let $\hat{\mathcal{I}}_n$ denote either the plug-in Fisher information $\mathcal{I}_n(\hat{\theta}_n)$ or the observed information $-\nabla^2 l_n(\hat{\theta}_n)$. Then the following assertions hold.

(i) **Consistency of the estimated Fisher information.**

$$D_n^{-1} \hat{\mathcal{I}}_n D_n^{-1} - J_n = O_{\mathbb{P}}(\epsilon_n). \quad (3.18)$$

In particular, $\hat{\mathcal{I}}_n$ is positive definite with probability tending to one, and for each $j \in \{s, l, t\}$,

$$\frac{[\hat{\mathcal{I}}_n^{-1}]_{jj}}{[\mathcal{I}_n(\theta_0)^{-1}]_{jj}} = 1 + O_{\mathbb{P}}(\epsilon_n).$$

(ii) **Coordinatewise intervals.** For each $j \in \{s, l, t\}$,

$$\frac{\hat{\theta}_{n,j} - \theta_{0,j}}{[\hat{\mathcal{I}}_n^{-1}]_{jj}^{1/2}} \implies N(0, 1).$$

Consequently, for every $\alpha \in (0, 1)$, letting $z_{1-\alpha/2}$ denote the $(1 - \alpha/2)$ quantile of the standard normal distribution, the Wald interval $\hat{\theta}_{n,j} \pm z_{1-\alpha/2} [\hat{\mathcal{I}}_n^{-1}]_{jj}^{1/2}$ contains $\theta_{0,j}$ with probability tending to $1 - \alpha$.

(iii) **Joint inference.** Let $\hat{L}_n$ be the lower-triangular Cholesky factor of $\hat{\mathcal{I}}_n$. Then

$$\hat{L}_n^\top (\hat{\theta}_n - \theta_0) \implies N_3(0, I_3), \quad (\hat{\theta}_n - \theta_0)^\top \hat{\mathcal{I}}_n (\hat{\theta}_n - \theta_0) \implies \chi_3^2.$$

Hence the ellipsoid $\{\theta : (\hat{\theta}_n - \theta)^\top \hat{\mathcal{I}}_n (\hat{\theta}_n - \theta) \leq \chi_{3,1-\alpha}^2\}$ is an asymptotic $1 - \alpha$ confidence region for θ_0 .

(iv) **Log scale.** Let $\hat{\Lambda}_n = \text{diag}(\hat{\theta}_n)$ and $\hat{\mathcal{I}}_n^{\log} = \hat{\Lambda}_n \hat{\mathcal{I}}_n \hat{\Lambda}_n$. Then (i)–(iii) hold for $\hat{\lambda}_n - \lambda_0$, with $\hat{\mathcal{I}}_n$ replaced by $\hat{\mathcal{I}}_n^{\log}$ and J_n by $J_n^{\log}$. In particular, the interval

$$\left[\hat{\theta}_{n,j} \exp\{-z_{1-\alpha/2} [(\hat{\mathcal{I}}_n^{\log})^{-1}]_{jj}^{1/2}\}, \hat{\theta}_{n,j} \exp\{z_{1-\alpha/2} [(\hat{\mathcal{I}}_n^{\log})^{-1}]_{jj}^{1/2}\} \right]$$

contains $\theta_{0,j}$ with probability tending to $1 - \alpha$. When $\hat{\mathcal{I}}_n$ is the observed information, $\hat{\mathcal{I}}_n^{\log}$ coincides with the observed information $-\nabla_{\lambda}^2 l_n$ at $\hat{\lambda}_n$ whenever $\hat{\theta}_n \in \text{int}(\Theta)$, an event of probability tending to one.

On the event that $\hat{\mathcal{I}}_n$ is not positive definite, the intervals and regions in (ii)–(iv) may be defined arbitrarily.

The estimated standard errors thus have relative error $O_{\mathbb{P}}(\epsilon_n) = O_{\mathbb{P}}(b_n^{-p/2})$. At moderate n , the estimated standard errors of the spatial parameters therefore carry nonnegligible uncertainty, which is a further reason to form intervals on the log scale and exponentiate them; this also keeps them positive. For $\hat{\mathcal{I}}_n = \mathcal{I}_n(\hat{\theta}_n)$, the proof gives the explicit bound $\|D_n^{-1}\mathcal{I}_n(\hat{\theta}_n)D_n^{-1} - J_n\|_{\text{op}} \leq C\epsilon_n \|D_n(\hat{\theta}_n - \theta_0)\|$ for every $\theta_0 \in \Theta$. Combined with [Theorem 3.2](#), the coverage statements hold uniformly over θ_0 in compact subsets of $\text{int}(\Theta)$.

3.4.3 Random observation locations

So far, all results are formulated for deterministic observation locations, which allows the likelihood analysis to condition on the design. In many applications, however, the locations themselves are naturally modeled as random. The next corollary shows that the deterministic-design formulation is not a substantive restriction: a broad class of random designs satisfies the required coverage condition ([Assumption 2](#)) almost surely, so the preceding results continue to hold.

Corollary 3.6 (Independent random sampling). Let $X_1, X_2, \dots$ be independent and identically distributed locations in a fixed bounded set $\mathcal{X}$, independent of the process and the errors. Suppose their distribution has a Lebesgue density f satisfying $f(x) \geq f_- > 0$ for almost every x in a fixed cube $Q \subset \mathcal{X}$. Write $\mathbf{X} = (X_1, X_2, \dots)$ for the design sequence. The quantities $\mathcal{I}_n(\theta_0)$, J_n , and M_n in [Theorem 3.1](#) are evaluated at the realized locations, and are therefore random.

- (a) **Conditionally on the design.** For almost every realization of $\mathbf{X}$, [Assumption 2](#) holds, and [Theorems 3.1](#) and [3.3](#) and [Theorems 3.4](#) and [3.5](#) hold conditionally on $\mathbf{X}$. The constants c and C in these results do not depend on $\mathbf{X}$; only the thresholds n_0 and n_{λ} may.
- (b) **Unconditionally.** Under the joint law of the design, the process, and the errors, the following assertions hold.
 - (i) [Theorem 3.1](#)(i) holds almost surely for all sufficiently large n .
 - (ii) [Theorem 3.1](#)(ii) and (iii) hold.
 - (iii) [Theorem 3.1](#)(iv) holds with an additional term $Ce^{-cn^{3/4}}$ on the right-hand side of each tail bound.
 - (iv) In [Theorem 3.1](#)(v), the exponential moment bound holds, and $\mathbb{E}g(M_n(\hat{\theta}_n - \theta_0)) \rightarrow \mathbb{E}g(Z)$ for every test function g as in [Theorem 3.1](#)(v). In particular, $D_n\mathbb{E}(\hat{\theta}_n - \theta_0) \rightarrow 0$ and $\mathbb{E}[M_n(\hat{\theta}_n - \theta_0)(\hat{\theta}_n - \theta_0)^{\top}M_n^{\top}] \rightarrow I_3$.
 - (v) [Theorem 3.3](#) holds, so the MLE attains the minimax rate.
 - (vi) [Theorem 3.5](#) holds, and (i)–(v) above hold on the log scale.

The remaining statements of [Theorem 3.1\(v\)](#) compare the MLE with the design-dependent matrix J_n , through $\mathbb{E}g(J_n^{-1/2}Z)$ and J_n^{-1} . They hold conditionally on the design by part (a). Unconditional versions would require integrability of J_n^{-1} over the design, which we do not pursue: near-degenerate designs, in which all locations are close together, have small but positive probability and make J_n nearly singular. Part (b)(iv) gives the standardized analogues instead. A continuous density that is bounded away from zero on a neighborhood of Q satisfies the assumptions of the corollary. Thus, the corollary covers standard random designs whose sampling density is sufficiently well behaved on some region of the domain $\mathcal{X}$.

3.5 Proof outline and key ideas

The proof of [Theorem 3.1](#) requires overcoming difficulties that make standard likelihood asymptotics inapplicable. Under fixed-domain asymptotics, the observations become increasingly strongly dependent, so the likelihood cannot easily be decomposed into a growing number of approximately independent contributions. Our main strategy is to exploit the analytic structure of the RBF kernel to recover, from the observations, a growing collection of latent Gaussian components whose conditional variance structure makes their information about the kernel parameters tractable. This leads to a global separation bound for the observation laws at the precise scales of the optimal convergence rates. After this global step is established, consistency is obtained via the method of [Ibragimov and Has'minskii \[1981\]](#), and then a Taylor expansion of the local log likelihood is used to derive the asymptotic expansion and distribution of the MLE.

The power series of the field. Writing $\|x - y\|^2 = \|x\|^2 + \|y\|^2 - 2\langle x, y \rangle$ and expanding $e^{\langle x, y \rangle / l^2}$ by the multinomial theorem gives

$$K_\theta(x, y) = s \sum_{\alpha \in \mathbb{N}_0^p} \phi_{\alpha, l}(x) \phi_{\alpha, l}(y), \quad \phi_{\alpha, l}(x) = e^{-\|x\|^2 / (2l^2)} \frac{x^\alpha}{l^{|\alpha|} \sqrt{\alpha!}}.$$

Hence F can be realized as the entire random function

$$F(x) = \sqrt{s} e^{-\|x\|^2 / (2l^2)} \sum_{\alpha \in \mathbb{N}_0^p} \xi_\alpha \frac{x^\alpha}{l^{|\alpha|} \sqrt{\alpha!}}, \quad \xi_\alpha \text{ independent } N(0, 1).$$

Its normalized Taylor coefficients at a fixed point, $\zeta_\alpha = \sqrt{\alpha!} [x^\alpha] F(x)$, are linear in the ξ_γ . Multiplication by $e^{-\|x\|^2 / (2l^2)}$ raises degrees in steps of two, so ζ_α depends only on ξ_α and on variables ξ_γ of strictly lower degree. In degree order, the map from the ξ 's to the ζ 's is therefore lower triangular with diagonal entries $\sqrt{s} l^{-|\alpha|}$. Consequently, given all normalized coefficients of lower degree, ζ_α is normal with variance $s l^{-2|\alpha|}$. On the log scale, this conditional variance is $\log s - 2|\alpha| \log l$, which is linear in the degree with intercept $\log s$ and slope $-2 \log l$.

This triangular representation is the key structural feature of the proof: it reveals that the highly dependent Gaussian field is a relatively simple function of a sequence of Gaussian variables with simple conditional variance structure. If these normalized coefficients were directly observed, the different information scales for s and l would already be visible from this linear dependence on the degree. The difficulty is that the coefficients are latent, and not all of them can be recovered accurately from the n noisy observations at the irregular locations.

Estimating the normalized coefficients. The normalized coefficients are not observed, but they can be estimated by a fixed linear statistic of the data. Recovering a growing number of them with sufficiently small error is a central technical step, particularly because we assume only polynomial coverage and allow irregular, clustered, and repeated observation locations ([Assumption 2](#)). We fit a polynomial of total degree $K_n \asymp b_n = \log n / \log \log n$ to the data by a discrete weighted least squares regression onto a Legendre basis for $L^2(Q)$, and then read off its normalized Taylor coefficients. We show that all normalized coefficients through degree $m_n \asymp b_n$ are recovered with error n^{-c} , uniformly in θ ([Theorem A.9](#)). There are on the order of b_n^p such coefficients. The estimator $\hat{\zeta}^{(m_n)} = H_n Y_n$ of the exact normalized coefficients $\zeta^{(m_n)} = (\zeta_\alpha)_{|\alpha| \leq m_n}$ does not depend on θ . The fact that H_n is independent of θ is essential: if one can distinguish parameters using the observed $\hat{\zeta}^{(m_n)} = H_n Y_n$, as we will show, then one can distinguish between parameters using the full data Y_n .

From coefficient recovery to global likelihood separation. For laws P and Q , the affinity $\text{Aff}(P, Q) = \int \sqrt{dP dQ}$ is the Bhattacharyya coefficient, and $-\log \text{Aff}(P, Q)$ is the Bhattacharyya distance [[Bhattacharyya, 1943](#)]. The lower triangular representation of $\zeta^{(m_n)}$ allows us to derive an explicit lower bound on the Bhattacharyya distance between its laws under θ and θ_0 . The uniform recovery result then transfers this bound from the latent coefficients $\zeta^{(m_n)}$ to their estimators $\hat{\zeta}^{(m_n)}$. Finally, because $\hat{\zeta}^{(m_n)} = H_n Y_n$ is a parameter-independent statistic of the observations, the data processing inequality transfers the bound to the observation laws themselves. This yields a global separation bound in terms of $(\log s - \log s_0)^2$ and $(\log l - \log l_0)^2$.

For the nugget, the factorial decay in the Taylor series also makes the kernel matrix numerically of rank $O(b_n^p)$ along a common subspace, uniformly in θ ([Theorem A.8](#)). The data component W_n orthogonal to that subspace is essentially white noise with variance t , and contributes a Bhattacharyya distance of order $n(\log t - \log t_0)^2$.

The proof therefore separates two fundamentally different sources of information: only $O(b_n^p)$ effective signal directions carry information about the spatial parameters, whereas the remaining directions provide essentially n independent pieces of information about the nugget. This distinction is what ultimately produces the logarithmic rates for (s, l) and the parametric $n^{-1/2}$ rate for t .

Putting the signal and nugget Bhattacharyya distance bounds together yields that in rescaled coordinates $h = D_n(\theta - \theta_0)$, for all $\theta \in \Theta$,

$$-\log \text{Aff}(P_{\theta,n}, P_{\theta_0,n}) \geq c\{\min(\|h_{\text{sp}}\|^2, b_n^{p/2} \|h_{\text{sp}}\|) + h_t^2\} - o(1).$$

In the case of i.i.d. data, such a lower bound comes very easily, but in this case, it is one of the hardest parts of the proof. Crucially, the bound holds uniformly over the entire parameter space, that is, the bound is uniform in θ_0 .

From global separation to the limit theorem. The global separation bound first controls the global maximizer of the likelihood. A chaining argument based on [Ibragimov and Has'minskii \[1981\]](#) converts this separation into the exponential tail bounds for $\hat{h}_n = D_n(\hat{\theta}_n - \theta_0)$ in [Theorem 3.1\(iv\)](#), thereby establishing consistency at the rates (3.7) before any local likelihood approximation is invoked. This localization is essential: without first ruling out distant likelihood maximizers, a quadratic expansion around θ_0 alone cannot establish the behavior of the global MLE.

Once the MLE has been localized, we establish a uniform quadratic expansion of the log likelihood on the relevant rescaled neighborhoods ([Theorem A.14](#)). The normalized score is a vector of Gaussian quadratic forms whose individual spectral contributions vanish, allowing a Lyapunov central limit theorem argument to establish asymptotic normality of the normalized score, despite the strong dependence of the original observations. Together, these three ingredients—global separation and localization, a uniform local quadratic approximation, and a central limit theorem for the normalized score—yield (3.6) and the joint asymptotic normality in [Theorem 3.1\(iii\)](#).

4 Simulation

We examine how closely the asymptotic theory in [Theorem 3.1](#) describes the MLE at finite sample sizes. Because the convergence rates for σ^2 and l are logarithmic, we consider sample sizes spanning several orders of magnitude.

4.1 Simulation setup

Model and designs. We simulate from (2.1) with the RBF kernel on the unit cube $\mathcal{X} = [0, 1]^p$, for $p \in \{1, 2, 3\}$, with true parameter

$$\theta_0 = (\sigma_0^2, l_0, \tau_0^2) = (1, 0.25, 0.01).$$

This gives a signal-to-noise ratio $\sigma_0^2/\tau_0^2 = 100$, and the lengthscale is one quarter of the side length of the domain. The locations form a regular midpoint grid with G points per axis,

$$x \in \{(i - \frac{1}{2})/G : i = 1, \dots, G\}^p,$$

so that $n = G^p$. This design satisfies [Assumption 2](#) with $\kappa = 1/p$. For each p , we use nine sample sizes ranging from approximately 10^2 to 10^6 :

$$\begin{aligned} G &\in \{10^2, 3 \cdot 10^2, 10^3, \dots, 3 \cdot 10^5, 10^6\}, & p = 1, \\ G &\in \{10, 18, 32, 56, 100, 178, 316, 562, 1000\}, & p = 2, \\ G &\in \{5, 7, 10, 14, 22, 32, 46, 68, 100\}, & p = 3. \end{aligned}$$

For every (p, n) , we generate 1000 independent data sets, each with a new realization of the field F and measurement errors.

Likelihood evaluation and MLE. Both data generation and likelihood evaluation exploit the power-series representation of the RBF kernel and, for $p \geq 2$, the Kronecker structure of the regular grid. This avoids forming the full $n \times n$ covariance matrix and makes likelihood optimization feasible for sample sizes up to $n = 10^6$. The MLE $\hat{\theta}_n$ is computed over

$$\Theta = [10^{-2}, 10^2] \times [0.05, 2] \times [10^{-4}, 1],$$

which contains θ_0 well inside its interior. Of the 2.7×10^4 estimates, only two lie on the boundary of Θ : one estimate of l for $p = 1$ and $n = 10^2$, and one estimate of τ^2 for $p = 3$ and $n = 125$. Details of

the likelihood computation, optimization, and numerical validation are provided in Appendix D.1.

Error summaries. For each parameter $\theta_j \in \{\sigma^2, l, \tau^2\}$, $j = 1, 2, 3$, we report three quantities as functions of n .

- (a) The *root MSE* (RMSE),

$$\text{RMSE}_j = \left\{ \frac{1}{R} \sum_{r=1}^R (\hat{\theta}_j^{(r)} - \theta_{0j})^2 \right\}^{1/2}, \quad R = 1000.$$

By Theorem 3.1(v), RMSE_j has the same order as the corresponding entry of D_n^{-1} .

- (b) The *robust standard deviation* $\text{IQR}_j/1.349$, where IQR_j is the interquartile range of the replicate errors $\hat{\theta}_j^{(r)} - \theta_{0j}$. The factor 1.349 is the interquartile range of a standard normal distribution, so $\text{IQR}/1.349$ equals the standard deviation under normality. Unlike the RMSE, this quantity is insensitive to a small number of extreme estimates.
- (c) The *asymptotic standard deviation* $[\mathcal{I}_n(\theta_0)^{-1}]_{jj}^{1/2}$, where $\mathcal{I}_n(\theta_0)$ is the Fisher information in (3.3). By Theorem 3.1(iii)–(v), the MLE is approximately normal with this standard deviation, whose order is $b_n^{-p/2}$, $b_n^{-(p+2)/2}$, and $\sqrt{2}\tau_0^2 n^{-1/2}$ for σ^2 , l , and τ^2 , respectively. This quantity is computed exactly for the actual grid design at each n .

Bootstrap 95% intervals for the first two quantities are obtained by resampling the replicates.

Log parametrization. We report the results on the log scale. As shown in Theorem 3.4, the log transformation preserves the convergence rates and yields the parameter-free nugget limit

$$\sqrt{n}(\log \hat{\tau}^2 - \log \tau_0^2) \implies N(0, 2).$$

Results on the original scale are reported in Appendix D.11 and lead to the same conclusions about the convergence rates.

4.2 Simulation results

Graphical summaries. Figures 1 to 3 show the results for $p = 1, 2, 3$. The top row compares the Monte Carlo RMSE and robust standard deviation with the asymptotic standard deviation and the theoretical rates in Theorem 3.1(ii). The bottom row shows normal Q–Q plots of the standardized errors at three representative sample sizes, providing a direct assessment of the asymptotic normality in Theorem 3.1(iii).

The rates agree with the theory. In all three dimensions and for all three parameters, the Monte Carlo errors decrease at rates consistent with Theorem 3.1 and closely track the exact asymptotic standard deviations. For $p = 3$, the RMSEs of $\log \hat{\sigma}^2$ and $\log \hat{l}$ are within 7% of their asymptotic standard deviations at every n from 125 to 10^6 . For $p = 2$, they are within 10% at every n and within 6% for $n \geq 10^3$. For $p = 1$, the RMSEs are somewhat larger but approach the asymptotic standard deviations as n increases.




Figure 1: Simulation results for $p = 1$ on the log scale, with $\theta_0 = (1, 0.25, 0.01)$ and 1000 replicates per n . Top: RMSE (filled circles), robust standard deviation IQR/1.349 (open squares), both with bootstrap 95% intervals, and the exact asymptotic standard deviation $[\mathcal{I}_n(\theta_0)^{-1}]_{jj}^{1/2}/\theta_{0j}$ (grey line). The grey region indicates sample sizes beyond the Monte Carlo range, where repeated MLE computation is computationally infeasible but the Fisher-information can still be evaluated. Dashed lines show the theoretical rates in [Theorem 3.1\(ii\)](#). Bottom: normal Q-Q plots of the standardized errors, with a 95% band for exact normality; triangles mark points outside the frame.

[Table 1](#) compares slopes fitted to the simulated errors with those of the exact asymptotic standard deviation over the same range of n , as well as with the limiting exponents in [Theorem 3.1](#). For $p = 2, 3$, the simulated RMSE slopes agree with those of the asymptotic standard deviation to within 0.1. The finite-sample slopes need not yet equal their limiting values because $b_n = \log n / \log \log n$ grows extremely slowly; this issue is examined further in [Appendix D.8](#).

The contrast between the spatial parameters and the nugget is substantial. From $n = 10^2$ to $n = 10^6$, the RMSE of $\log \hat{\sigma}^2$ decreases only by factors of 1.2, 1.5, and 2.1 for $p = 1, 2, 3$, respectively, whereas the RMSE of $\log \hat{\tau}^2$ decreases by factors between 36 and 350. This is the slow logarithmic behavior predicted by the theory.

The errors are close to normal. For $p = 2$ and $p = 3$, the standardized errors of $\log \hat{\sigma}^2$ and $\log \hat{\ell}$ lie essentially within the 95% band for exact normality at the displayed sample sizes. The RMSE and robust standard deviation are also very similar, providing further evidence for the




Figure 2: Simulation results for $p = 2$, with the layout and graphical elements as in Figure 1.

Table 1: Least-squares slopes of $\log(\text{error})$ against $\log b_n$ for $\log \sigma^2$ and $\log l$, and against $\log n$ for $\log \tau^2$, over the nine simulated sample sizes ($10^2 \lesssim n \leq 10^6$). The last column gives the theoretical rate shown in Theorem 3.1.

		RMSE	IQR/1.349	Asymptotic SD	Theory
$\log \sigma^2$	$p = 1$	-0.49	-0.49	-0.35	-0.5
	$p = 2$	-0.75	-0.72	-0.74	-1
	$p = 3$	-1.34	-1.48	-1.32	-1.5
$\log l$	$p = 1$	-2.05	-1.81	-1.66	-1.5
	$p = 2$	-2.32	-2.24	-2.22	-2
	$p = 3$	-3.11	-3.11	-3.09	-2.5
$\log \tau^2$	$p = 1$	-0.50	-0.50	-0.50	-0.5
	$p = 2$	-0.52	-0.53	-0.52	-0.5
	$p = 3$	-0.60	-0.58	-0.58	-0.5

Fisher-normalized limit in (3.8). The nugget errors show similarly good agreement except at the smallest sample size for $p = 3$. Convergence to normality is slower for $p = 1$, particularly for the lengthscale, where a heavier right tail remains visible even at the largest simulated sample sizes.




Figure 3: Simulation results for $p = 3$, with the layout and graphical elements as in Figure 1.

Additional normality diagnostics are reported in Appendix D.6.

The nugget is the easiest parameter. The nugget variance is estimated at the parametric rate. In every dimension, $\sqrt{n} \text{RMSE}(\log \hat{\tau}^2)$ is within 6% of its limit $\sqrt{2}$ for all $n \geq 3 \times 10^3$. On the original scale, for example,

$$\sqrt{n} \text{RMSE}(\hat{\tau}^2) = 0.0143$$

at $n = 10^6$ for $p = 2$, compared with the theoretical limit $\sqrt{2} \tau_0^2 = 0.01414$. The approximation is less accurate at the smallest sample sizes, especially for $p = 3$. This behavior is consistent with the low-rank structure underlying Theorem 3.1: most directions eventually contain essentially only nugget noise, allowing τ^2 to be estimated at the usual parametric rate. A more detailed finite-sample explanation is given in Appendix D.7.

Convergence rates of the spatial parameters are slow. For the spatial parameters σ^2 and ℓ , the effective amount of information grows with the number of recoverable Taylor coefficients, whose maximal degree is of order $b_n = \log n / \log \log n$. This quantity grows very slowly: over $10^2 \leq n \leq 10^6$, it increases only from approximately 3.0 to 5.3. Consequently, the finite-sample slopes in Table 1 need not closely match the limiting exponents even when the MLE already tracks the Fisher-information benchmark well. Indeed, the exact asymptotic standard deviation itself

exhibits the same finite-sample deviations from the limiting slopes. Further analysis of these effects is provided in Appendix D.8.

The different spatial rates are nevertheless already clearly visible. Since D_n contains an additional factor b_n for l , the lengthscale is estimated more accurately than the spatial variance by a factor of order b_n , in agreement with the simulations.

Higher dimensions help. The rates $b_n^{-p/2}$ and $b_n^{-(p+2)/2}$ imply that estimation of σ^2 and l becomes easier as the dimension p increases. This may initially appear counterintuitive, since all observations still come from a single realization of the field. The explanation is that a higher-dimensional RBF field contains more recoverable Taylor coefficients. The number of coefficients of degree at most m is

$$N_p(m) = \binom{m+p}{p},$$

which is asymptotically of order m , $m^2/2$, and $m^3/6$ for $p = 1, 2, 3$, respectively. Thus a single realization contains more independent information about the spatial kernel parameters as p increases. This is reflected both in the smaller estimation errors and in the improved normal approximation from $p = 1$ to $p = 3$. In contrast, the nugget information is of order n in every dimension.

5 Discussion and future work

In this article, we establish an asymptotic theory for MLEs of RBF kernel parameters under fixed-domain asymptotics. We prove consistency of the joint MLE of the spatial variance, lengthscale, and nugget variance, derive their convergence rates, establish joint asymptotic normality, and show that all three rates are minimax optimal. To the best of our knowledge, these results provide the first complete asymptotic characterization of the joint MLE for the RBF kernel in this setting. The results provide theoretical justification for the MLE routinely used in GP software. At the same time, they reveal an important limitation: while the nugget variance converges at the standard parametric rate $n^{-1/2}$, the spatial variance and lengthscale converge only at logarithmic rates. Since these rates are sharp, the slow convergence is not merely an artifact of the MLE or our analysis. In practice, substantial uncertainty in the estimated spatial variance and lengthscale may therefore remain even with a large number of densely sampled observations, and these estimates should be interpreted with caution when they are used for inference or downstream tasks. There are, however, some limitations of our study that motivate important directions for future work.

First, our results concern the exact MLE, whose computation requires $O(n^3)$ operations and becomes prohibitive for large n . A large literature has developed scalable GP approximations, including inducing-point, variational inference, nearest-neighbor, and Vecchia-type methods, which replace the exact likelihood with computationally tractable approximations [Liu et al., 2020]. Indeed, our simulation studies also rely on scalable numerical approximations because repeated exact likelihood optimization becomes computationally prohibitive at the sample sizes needed to examine the asymptotic behavior. Thus, an important next question is how these approximations affect kernel parameter inference. In particular, it remains to determine which scalable approximations yield consistent estimators, whether they preserve the convergence rates and asymptotic distributions of the exact MLE, and how the approximation accuracy or computational complexity must scale with

n to retain these statistical properties. Conversely, aggressive approximations may lose information about the kernel parameters and lead to slower rates or even inconsistency. Developing a statistical theory connecting computational approximation to kernel parameter inference would therefore provide useful guidance for scalable GP methodology.

A second limitation is that our analysis is specific to the RBF kernel. A key ingredient of our proof is its analytic structure, which allows a growing collection of normalized Taylor coefficients to be recovered from the observations with sufficiently small error. This argument does not extend directly to kernels with finite smoothness, most notably the widely used Matérn family. Establishing analogous results for such kernels will therefore require different mathematical tools. More generally, understanding how kernel smoothness determines the amount of recoverable information and the resulting behavior of likelihood-based kernel parameter estimation is an important direction for future work.

References

- Sudipto Banerjee, Alan E. Gelfand, and Bradley P. Carlin. *Hierarchical Modeling and Analysis for Spatial Data*. Chapman and Hall/CRC, Boca Raton, FL, third edition, 2025.
- A. Bhattacharyya. On a measure of divergence between two statistical populations defined by their probability distributions. *Bulletin of the Calcutta Mathematical Society*, 35:99–109, 1943.
- DLMF. NIST Digital Library of Mathematical Functions. <https://dlmf.nist.gov/>, 2026. Release 1.2.8 of 2026-09-15.
- Roman Garnett. *Bayesian Optimization*. Cambridge University Press, Cambridge, 2023.
- Subhashis Ghosal and Aad van der Vaart. *Fundamentals of Nonparametric Bayesian Inference*, volume 44 of *Cambridge Series in Statistical and Probabilistic Mathematics*. Cambridge University Press, Cambridge, 2017.
- I. A. Ibragimov and R. Z. Has’minskii. *Statistical Estimation: Asymptotic Theory*, volume 16 of *Applications of Mathematics*. Springer, New York, 1981.
- Toni Karvonen and Chris J. Oates. Maximum likelihood estimation in Gaussian process regression is ill-posed. *Journal of Machine Learning Research*, 24(120):1–47, 2023.
- Haitao Liu, Yew-Soon Ong, Xiaobo Shen, and Jianfei Cai. When Gaussian process meets big data: A review of scalable GPs. *IEEE Transactions on Neural Networks and Learning Systems*, 31(11):4405–4423, 2020.
- Wei-Liem Loh. Consistent estimation for a Gaussian random field with squared exponential covariance using scattered data under fixed-domain asymptotics. *Electronic Journal of Statistics*, 20(2):4045–4082, 2026.
- Wei-Liem Loh and Tao-Kai Lam. Estimating structured correlation matrices in smooth Gaussian random field models. *The Annals of Statistics*, 28(3):880–904, 2000.

- Blake MacDonald, Pritam Ranjan, and Hugh Chipman. GPfit: An R package for fitting a Gaussian process model to deterministic simulator outputs. *Journal of Statistical Software*, 64(12):1–23, 2015.
- Fabian Pedregosa, Gaël Varoquaux, Alexandre Gramfort, Vincent Michel, Bertrand Thirion, Olivier Grisel, Mathieu Blondel, Peter Prettenhofer, Ron Weiss, Vincent Dubourg, et al. Scikit-learn: Machine learning in Python. *Journal of Machine Learning Research*, 12:2825–2830, 2011.
- Ameer Qaqish and Didong Li. Identifiability for Gaussian processes with holomorphic kernels. In *International Conference on Learning Representations*, 2025.
- Carl Edward Rasmussen and Hannes Nickisch. Gaussian processes for machine learning (GPML) toolbox. *Journal of Machine Learning Research*, 11:3011–3015, 2010.
- Carl Edward Rasmussen and Christopher K. I. Williams. *Gaussian Processes for Machine Learning*. MIT Press, Cambridge, MA, 2006.
- Daniel Revuz and Marc Yor. *Continuous Martingales and Brownian Motion*, volume 293 of *Grundlehren der mathematischen Wissenschaften*. Springer, Berlin, third edition, 1999.
- Stephen Roberts, Michael Osborne, Mark Ebdon, Steven Reece, Neale Gibson, and Suzanne Aigrain. Gaussian processes for time-series modelling. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 371(1984):20110550, 2013.
- Olivier Roustant, David Ginsbourger, and Yves Deville. DiceKriging, DiceOptim: Two R packages for the analysis of computer experiments by kriging-based metamodeling and optimization. *Journal of Statistical Software*, 51(1):1–55, 2012.
- Michael L. Stein. *Interpolation of Spatial Data: Some Theory for Kriging*. Springer Series in Statistics. Springer, New York, 1999.
- Valentine Svensson, Sarah A. Teichmann, and Oliver Stegle. SpatialDE: identification of spatially variable genes. *Nature Methods*, 15(5):343–346, 2018.
- Michel Talagrand. *Upper and Lower Bounds for Stochastic Processes: Decomposition Theorems*, volume 60 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge*. Springer, Cham, second edition, 2021.
- Lukas M. Weber, Arkajyoti Saha, Abhirup Datta, Kasper D. Hansen, and Stephanie C. Hicks. nnSVG for the scalable identification of spatially variable genes using nearest-neighbor Gaussian processes. *Nature Communications*, 14(1):4059, 2023.
- Wanting Xu and Michael L. Stein. Maximum likelihood estimation for a smooth Gaussian random field model. *SIAM/ASA Journal on Uncertainty Quantification*, 5(1):138–175, 2017.

Appendix

In Appendix A, we state key lemmas to prove the main theorems, together with auxiliary lemmas to prove the key lemmas. In Appendix B, we prove the main theorems and their corollaries using the lemmas in Appendix A. In Appendix C, we prove all lemmas in Appendix A. Finally, in Appendix D, we present additional experimental details for simulations in Section 4.

A Lemmas

This appendix first records elementary Gaussian facts and a maximal inequality for continuous processes, and then states the auxiliary lemmas used in the proofs of Theorems 3.1 and 3.3 and their corollaries. The lemma proofs appear in Appendix C. All design matrices and linear statistics introduced below are deterministic functions of the locations.

A.1 Gaussian affinity and elementary bounds

For probability laws P, Q , define

$$\text{Aff}(P, Q) = \int \sqrt{dP dQ}, \quad \text{TV}(P, Q) = \sup_A |P(A) - Q(A)|.$$

The affinity is also called the Bhattacharyya coefficient, and $-\log \text{Aff}(P, Q)$ is the Bhattacharyya distance. For positive definite $d \times d$ covariance matrices A, B , abbreviate $\text{Aff}(A, B) = \text{Aff}(N_d(0, A), N_d(0, B))$. Direct Gaussian integration gives

$$\text{Aff}(A, B) = \frac{\det(A)^{1/4} \det(B)^{1/4}}{\det((A+B)/2)^{1/2}}. \quad (\text{A.1})$$

Applying the same statistic to both laws cannot decrease their affinity.

Lemma A.1 (Monotonicity of affinity). Let P, Q be probability laws on a measurable space $(\mathcal{Y}, \mathcal{F})$, and let $T : \mathcal{Y} \rightarrow \mathcal{Y}'$ be a measurable map into another measurable space. Then

$$\text{Aff}(P \circ T^{-1}, Q \circ T^{-1}) \geq \text{Aff}(P, Q).$$

In particular, if $Y \sim N_d(0, A)$ under one law and $Y \sim N_d(0, B)$ under the other, and H is a fixed $r \times d$ matrix of rank r , then $\text{Aff}(H A H^\top, H B H^\top) \geq \text{Aff}(A, B)$.

In the applications, T is a linear statistic of the data that does not depend on the parameter. Both uses are in the proof of Theorem A.10: the coefficient statistic of Theorem A.9, and the noise block W_n .

The next two lemmas are elementary. Theorem A.2 controls all moments of the scores and is used in Theorem A.11; Theorem A.3 is used in the proof of Theorem A.10 to show that a fixed fraction of polynomial degrees is separated.

Lemma A.2 (Moments of Gaussian quadratic forms). Let $X \sim N_d(0, I_d)$ and let B be a symmetric $d \times d$ matrix. For every $q \geq 1$ there is C_q , depending only on q , such that $\mathbb{E}|X^\top B X - \text{tr } B|^q \leq C_q \|B\|_F^q$.

Lemma A.3 (A Paley–Zygmund-type bound). Let X be a random variable with $|X| \leq M$ almost surely and $\mathbb{E}X^2 \geq \sigma^2 > 0$. Then, for every $\lambda \in [0, 1)$,

$$\mathbb{P}(|X| \geq \lambda\sigma) \geq (1 - \lambda^2) \frac{\sigma^2}{M^2}.$$

In particular, if $X = f(\iota)$ for an index ι uniformly distributed on a finite set $\mathcal{A}$, then at least $(1 - \lambda^2)\sigma^2|\mathcal{A}|/M^2$ elements $\iota \in \mathcal{A}$ satisfy $|f(\iota)| \geq \lambda\sigma$.

A.2 A maximal inequality for continuous processes

The exponential tail bounds rest on the following chaining inequality, a variant of the shell argument of Ibragimov and Has'minskii [1981, Theorem I.5.1]. Theorems A.4 and A.5 are the building blocks, Theorem A.6 is the general bound, and Theorem A.7 is the form used in Theorem A.12. Throughout, $m \geq 1$, $\|\cdot\|_{L^m}$ refers to a fixed probability measure, $|\cdot|$ is the Euclidean norm, and a supremum over the empty set is zero.

Lemma A.4 (Kolmogorov continuity criterion). Let $k \in \mathbb{N}$, $m \geq 1$, and $\alpha > k$. If a real process $(\xi(x))_{x \in [0,1]^k}$ satisfies $\|\xi(x) - \xi(y)\|_{L^m} \leq L|x - y|^{\alpha/m}$ for all x, y , then it has a continuous modification, again denoted ξ , such that

$$\left\| \sup_{x, y \in [0,1]^k, |x-y| \leq h} |\xi(x) - \xi(y)| \right\|_{L^m} \leq C_{k,m,\alpha} L h^{(\alpha-k)/m}, \quad h \in (0, 1].$$

Lemma A.5 (Supremum over a compact convex set). Let $k \in \mathbb{N}$ and $m > k$. Let $\Gamma \subseteq \mathbb{R}^k$ be a nonempty compact convex set of diameter at most $\delta > 0$, and let $(\xi(u))_{u \in \Gamma}$ be a real process with continuous paths such that $\sup_{u \in \Gamma} \|\xi(u)\|_{L^m} \leq M$ and $\|\xi(u) - \xi(v)\|_{L^m} \leq L|u - v|$ for $u, v \in \Gamma$. Then

$$\mathbb{E} \sup_{u \in \Gamma} |\xi(u)|^m \leq C_{k,m} \{M^m + (L\delta)^k M^{m-k}\}.$$

Proposition A.6 (Maximal inequality over a cover). Let $k \in \mathbb{N}$, $m > k$, $U \subseteq \mathbb{R}^k$, and let $(\xi(u))_{u \in U}$ be a real process with continuous paths. Let $(\Gamma_j)_{j \in \mathcal{J}}$ be a countable family of nonempty compact convex subsets of U of diameters at most δ_j . Suppose that $\sup_{u \in \Gamma_j} \|\xi(u)\|_{L^m} \leq M_j$ and $\|\xi(u) - \xi(v)\|_{L^m} \leq L_j|u - v|$ for $u, v \in \Gamma_j$. Then

$$\mathbb{E} \sup_{u \in \bigcup_j \Gamma_j} |\xi(u)|^m \leq C_{k,m} \sum_{j \in \mathcal{J}} \{M_j^m + (L_j \delta_j)^k M_j^{m-k}\}.$$

Only Lipschitz control inside each piece is required, so neither product structure nor convexity of U itself is needed. Taking the pieces to be annular shells gives bounds of Ibragimov–Has'minskii type. For the Bernstein-type envelopes arising here, unit cubes are more convenient.

Corollary A.7 (Bernstein-type envelopes). Let $k = k_1 + k_2$ with $k_1, k_2 \in \mathbb{N}$, let $m > k$, and let $U \subseteq \mathbb{R}^{k_1} \times \mathbb{R}^{k_2}$ be a product of compact intervals. For $\gamma \in [1, \infty]$ and $r \geq 0$ put $\psi_\gamma(r) = \min(r^2, \gamma r)$, so that $\psi_\infty(r) = r^2$. Let $(\xi(u))_{u \in U}$ have continuous paths, and suppose that, for constants $A \geq 0$,

$a > 0$, $B \geq 0$, and $\gamma_1, \gamma_2 \in [1, \infty]$,

$$\|\xi(u)\|_{L^m}^m \leq A \exp[-a\{\psi_{\gamma_1}(|u_1|) + \psi_{\gamma_2}(|u_2|)\}], \quad u \in U, \quad (\text{A.2})$$

$$\|\xi(u) - \xi(v)\|_{L^m} \leq B|u - v|, \quad u, v \in U. \quad (\text{A.3})$$

Put $a'' = a(m - k)/(32m)$. Then, for all $H_1, H_2 \geq 0$,

$$\mathbb{E} \sup_{\substack{u \in U \\ |u_1| \geq H_1, |u_2| \geq H_2}} |\xi(u)|^m \leq C(A + B^k A^{1-k/m}) \exp[-a''\{\psi_{\gamma_1}(H_1) + \psi_{\gamma_2}(H_2)\}], \quad (\text{A.4})$$

where C depends only on k_1, k_2, m , and a , and not on γ_1, γ_2 , or U .

A.3 Auxiliary lemmas

The auxiliary lemmas follow the stages of the proof outline in [Section 3.5](#).

- [Theorem A.8](#) splits the likelihood into a low-dimensional signal block and a noise block, and supplies the covariance derivative bounds used throughout.
- [Theorems A.9](#) and [A.10](#) give global identifiability. [Theorem A.9](#) recovers normalized Taylor coefficients of the field from the data, and [Theorem A.10](#) turns their triangular covariance structure into an affinity bound between the observation laws.
- [Theorem A.11](#), [Theorem A.6](#), and [Theorem A.7](#) convert that affinity bound into the exponential tail bound of [Theorem A.12](#), which is [Theorem 3.1\(iv\)](#).
- [Theorems A.13](#), [A.14](#) and [A.16](#) give the information bounds, the local quadratic expansion, and the score central limit theorem, which together prove parts (i)–(iii) of [Theorem 3.1](#). Part (v) follows from parts (iii) and (iv).
- [Theorem A.17](#) is used only for [Theorem 3.6](#).

The RBF kernel is numerically of low rank on a bounded set. Its Taylor coefficients decay factorially, so only polynomial features of degree $O(b_n)$ are visible at the noise level. The next lemma makes this precise with a single projection that works for all parameters simultaneously.

Lemma A.8 (Common projection and covariance derivatives). Assume [Assumption 1](#) and bounded observation locations. For any fixed $A > 0$, there is a parameter-independent orthogonal projection P_n of rank $r_n = O(b_n^p)$ such that, with $S_n(\theta) = sR_n(l)$,

$$\sup_{\theta \in \Theta} \|\partial_\theta^\gamma (S_n - P_n S_n P_n)\|_{\text{op}} \leq n^{-A}, \quad |\gamma| \leq 3. \quad (\text{A.5})$$

For $\tilde{C}_n = tI_n + P_n S_n P_n$ and its log likelihood,

$$\sup_{\theta \in \Theta} |\partial_\theta^\gamma (l_n - \tilde{l}_n)| = O_{\mathbb{P}}(n^{4-A}), \quad |\gamma| \leq 3. \quad (\text{A.6})$$

For a spatial derivative with a derivatives in s and c in l , $1 \leq a + c \leq 3$, a nonzero derivative

has $a \leq 1$ and satisfies

$$\sup_{\theta} \left\| C_n^{-1/2} \partial_s^a \partial_l^c C_n C_n^{-1/2} \right\|_{\text{op}} \leq C b_n^c, \quad (\text{A.7})$$

$$\sup_{\theta} \left\| C_n^{-1/2} \partial_s^a \partial_l^c C_n C_n^{-1/2} \right\|_{\text{F}} \leq C b_n^{p/2+c}. \quad (\text{A.8})$$

The same bounds hold after compression. Precisely, let U be any $n \times r$ matrix with orthonormal columns, $1 \leq r \leq n$, that does not depend on θ , and let $A_{\theta} = U^{\top} C_n(\theta) U$ be the compressed covariance. Then (A.7) and (A.8) hold with C_n replaced by A_{θ} , that is, for the compressed derivatives $\partial_s^a \partial_l^c A_{\theta} = U^{\top} (\partial_s^a \partial_l^c C_n) U$ whitened by $A_{\theta}^{-1/2}$, with constants that do not depend on U . In the coordinates $\theta = \theta_0 + D_n^{-1} h$, normalized spatial derivatives of total order $k \leq 3$, whitened either by C_n or by a compression A_{θ} , have respective bounds $C \epsilon_n^k$ in operator norm and $C \epsilon_n^{k-1}$ in Frobenius norm. The first normalized nugget derivative has operator norm $O(n^{-1/2})$ and Frobenius norm $O(1)$, and all covariance derivatives of order at least two involving t vanish.

In coordinates adapted to this projection, the reduced likelihood separates into a signal block of dimension r_n and a noise block of dimension $d_n = n - r_n$ that depends only on the nugget. Fix a projection from Theorem A.8 with $A = 20$. Choose orthonormal matrices U_n, V_n spanning its range and orthogonal complement, and write

$$Z_n = U_n^{\top} Y_n, \quad W_n = V_n^{\top} Y_n, \quad d_n = n - r_n, \quad A_{\theta} = t I_{r_n} + s U_n^{\top} R_n(l) U_n. \quad (\text{A.9})$$

The reduced negative twice log likelihood decomposes as

$$\begin{aligned} -2\tilde{l}_n(\theta) &= F_n(t) + G_n(s, l, t), \\ F_n(t) &= d_n \log t + \|W_n\|^2 / t, \\ G_n(s, l, t) &= \log |A_{\theta}| + Z_n^{\top} A_{\theta}^{-1} Z_n. \end{aligned}$$

The spatial parameters are identified through the Taylor coefficients of the field at a fixed point. For the next lemma, apply a fixed rigid motion to the spatial coordinates so that $Q = [0, L]^p$ for some $L > 0$. This preserves the covariance model. The field F has a version that is an entire function on $\mathbb{R}^p$ (see the proof of Theorem A.9). Its normalized Taylor coefficients at the origin are

$$\zeta_{\alpha} = \sqrt{\alpha!} [x^{\alpha}] F(x), \quad \alpha \in \mathbb{N}_0^p. \quad (\text{A.10})$$

Let $\zeta^{(m)} = (\zeta_{\alpha} : |\alpha| \leq m)$, $q_m = \binom{m+p}{p}$, and $\Sigma_{m,\theta} = \text{Cov}_{\theta}(\zeta^{(m)})$.

These coefficients have a triangular structure. By (C.19) in the proof of Theorem A.9, $\zeta^{(m)} = \sqrt{s} \mathcal{T}_{m,\theta} \xi^{(m)}$ with independent standard normal ξ_{α} and a matrix $\mathcal{T}_{m,\theta}$ that is lower triangular in degree order with diagonal entries $l^{-|\alpha|}$. Given the coefficients of lower degree, ζ_{α} therefore has variance $s l^{-2|\alpha|}$. The next lemma shows that a fixed linear statistic recovers these coefficients through degree $m_n \asymp b_n$, with polynomially small error uniformly in θ . The regression is fitted to a higher degree $K_n \approx 2m_n$: extracting a coefficient of degree m from the fitted polynomial amplifies errors by a factor $e^{m \log m}$, which the truncation error $e^{-K \log K}$ of the fit must overcome.

Lemma A.9 (Recovery of normalized Taylor coefficients). Under [Assumptions 1](#) and [2](#), set

$$\beta = \kappa p, \quad K_n = \left\lfloor \frac{\beta}{2} b_n \right\rfloor, \quad m_n = \left\lfloor \frac{\beta}{4} b_n \right\rfloor.$$

There is a parameter-independent linear statistic $\widehat{\zeta}^{(m_n)} = H_n Y_n$ such that

$$\sup_{\theta \in \Theta} \max_{|\alpha| \leq m_n} \mathbb{E}_\theta (\widehat{\zeta}_\alpha - \zeta_\alpha)^2 \leq n^{-\beta/4+o(1)}. \quad (\text{A.11})$$

The exact coefficient covariance satisfies

$$e^{-Cm} I_{q_m} \preceq \Sigma_{m,\theta} \preceq e^{Cm} I_{q_m}, \quad m \geq 1, \quad (\text{A.12})$$

uniformly in θ . Writing $\widehat{\Sigma}_{n,\theta} = \text{Cov}_\theta(H_n Y_n)$, we have

$$\sup_{\theta \in \Theta} \left\| \Sigma_{m_n,\theta}^{-1/2} (\widehat{\Sigma}_{n,\theta} - \Sigma_{m_n,\theta}) \Sigma_{m_n,\theta}^{-1/2} \right\|_{\text{op}} \leq n^{-\beta/8+o(1)}. \quad (\text{A.13})$$

Consequently,

$$\sup_{\theta, \theta' \in \Theta} \left| \log \text{Aff}(\widehat{\Sigma}_{n,\theta}, \widehat{\Sigma}_{n,\theta'}) - \log \text{Aff}(\Sigma_{m_n,\theta}, \Sigma_{m_n,\theta'}) \right| = o(1). \quad (\text{A.14})$$

The logarithm of the conditional variance $s l^{-2|\alpha|}$ is $\log s - 2|\alpha| \log l$, which is linear in the degree. Separating (s, l) from (s_0, l_0) therefore amounts to a regression of log-variances on degrees spread over $[0, m_n]$. With about b_n^p coefficients, this explains the information scales b_n^p for the intercept s and b_n^{p+2} for the slope l . The next lemma turns this heuristic into a global bound on the affinity between observation laws, and adds the noise block to separate the nugget.

Lemma A.10 (Affinity separation). Under [Assumptions 1](#) and [2](#), there are $c > 0$ and a deterministic sequence $e_n \rightarrow 0$ such that, for all $\theta = (s, l, t) \in \Theta$,

$$\text{Aff}(P_{\theta,n}, P_{\theta_0,n}) \leq \exp \left[e_n - c b_n^p \min \left\{ \Lambda_n(\theta)^2, \Lambda_n(\theta_0) \right\} \right], \quad (\text{A.15})$$

where $\Lambda_n(\theta) \geq 0$ is defined by $\Lambda_n(\theta)^2 = \log^2(s/s_0) + b_n^2 \log^2(l/l_0)$. Moreover, write $h = D_n(\theta - \theta_0) = (h_{\text{sp}}, h_t)$, where $h_{\text{sp}} \in \mathbb{R}^2$ collects the two spatial coordinates. After decreasing c and enlarging e_n , for all $\theta \in \Theta$,

$$\text{Aff}(P_{\theta,n}, P_{\theta_0,n}) \leq \exp [e_n - c \{ \min(\|h_{\text{sp}}\|^2, b_n^{p/2} \|h_{\text{sp}}\|) + h_t^2 \}]. \quad (\text{A.16})$$

The constant c and the sequence e_n do not depend on $\theta_0 \in \Theta$, and they depend on the design only through $\mathcal{X}$, Q , C_h , and κ . Both bounds hold at every sufficiently large n at which [\(3.2\)](#) is satisfied.

To control the global maximizer, the likelihood ratio $Z_n(h)$ enters only through the process $\xi_n = Z_n^{1/(2m)}$. The exponent $1/(2m)$ makes $\mathbb{E} \xi_n(h)^m$ exactly the affinity, and a change of measure bounds the L^m Lipschitz constant of ξ_n uniformly over the whole parameter box. This holds even where covariances at different parameters are not comparable.

Lemma A.11 (Likelihood-ratio process). Assume [Assumption 1](#) and bounded observation locations, and fix $m \geq 1$. On the box $U_n = D_n(\Theta - \theta_0)$ define

$$Z_n(h) = \exp \{ l_n(\theta_0 + D_n^{-1} h) - l_n(\theta_0) \}, \quad \xi_n(h) = Z_n(h)^{1/(2m)}.$$

Then ξ_n has continuous paths,

$$\mathbb{E}_{\theta_0,n} \xi_n(h)^m = \text{Aff}(P_{\theta_0 + D_n^{-1}h,n}, P_{\theta_0,n}), \quad h \in U_n, \quad (\text{A.17})$$

and there is a constant C_L , depending only on m, p, Θ , and $\mathcal{X}$, such that for all n and all $u, v \in U_n$,

$$\|\xi_n(u) - \xi_n(v)\|_{L^m(\mathbb{P}_{\theta_0,n})} \leq C_L \|u - v\|. \quad (\text{A.18})$$

Combining [Theorems A.10](#) and [A.11](#) with [Theorem A.7](#) gives the tail bound. The key observation is that $\xi_n(\hat{h}_n) \geq \xi_n(0) = 1$, because $\hat{\theta}_n$ maximizes the likelihood. Hence a large deviation of $\hat{h}_n$ forces a large supremum of ξ_n far from the origin.

Lemma A.12 (Exponential tails of the global maximizer). Under [Assumptions 1](#) and [2](#), write $\hat{h}_n = D_n(\hat{\theta}_n - \theta_0) = (\hat{h}_{n,\text{sp}}, \hat{h}_{n,t})$. There are constants $c, C > 0$ and n_0 such that [\(3.10\)](#) holds for all $n \geq n_0$ and $H \geq 0$. The constants do not depend on $\theta_0 \in \Theta$, and they depend on the design only through $\mathcal{X}, Q, C_h$, and κ . For $n \geq n_0$, the bounds hold at every n at which [\(3.2\)](#) is satisfied. Interiority of θ_0 is not used.

The upper information bounds follow from the derivative bounds of [Theorem A.8](#). The lower bound on $J_{n,\text{sp}}$ avoids differentiating the coefficient recovery error. Instead, it compares a local quadratic expansion of the affinity at distance one, in rescaled coordinates, with the separation bound of [Theorem A.10](#).

Lemma A.13 (Information bounds). Under [Assumption 1](#) and bounded observation locations,

$$\sup_{\theta \in \Theta} \mathcal{I}_{ss,n}(\theta) \leq C b_n^p, \quad \sup_{\theta \in \Theta} \mathcal{I}_{ll,n}(\theta) \leq C b_n^{p+2}, \quad \sup_{\theta \in \Theta} \mathcal{I}_{tt,n}(\theta) \leq C n.$$

If [Assumption 2](#) also holds, then [\(3.4\)](#) and [\(3.5\)](#) hold.

The remaining stage is local. Because [Theorem 3.1\(iv\)](#) already gives tightness, the main proof needs the quadratic expansion only on balls of fixed radius. The next lemma gives it on balls of every radius $o(b_n^{p/6})$, and [Theorem A.15](#) shows that this radius cannot be enlarged.

Lemma A.14 (Uniform local quadratic expansion). Under [Assumption 1](#) and bounded observation locations, set $\Delta_n = D_n^{-1} \nabla l_n(\theta_0)$ and $L_n(h) = l_n(\theta_0 + D_n^{-1}h)$. Then $\mathbb{E} \Delta_n = 0$, $\text{Cov}(\Delta_n) = J_n$, and

$$\nabla_h^2 L_n(0) = -J_n + O_{\mathbb{P}}(\epsilon_n). \quad (\text{A.19})$$

Let $H_n \geq 1$ be any sequence with $\epsilon_n H_n \rightarrow 0$. Then

$$\sup_{\|h\| \leq H_n} \max_{a,b,c} |\partial_{h_a h_b h_c}^3 L_n(h)| = O_{\mathbb{P}}(\epsilon_n). \quad (\text{A.20})$$

For all sufficiently large n , the ball $\{\|h\| \leq H_n\}$ lies in the rescaled parameter space $D_n(\Theta - \theta_0)$, and

$$\sup_{\|h\| \leq H_n} \left| L_n(h) - L_n(0) - h^\top \Delta_n + \frac{1}{2} h^\top J_n h \right| = O_{\mathbb{P}}(\epsilon_n H_n^3). \quad (\text{A.21})$$

In particular, if $\epsilon_n H_n^3 \rightarrow 0$, that is, $H_n = o(b_n^{p/6})$, then

$$\sup_{\|h\| \leq H_n} \left| L_n(h) - L_n(0) - h^\top \Delta_n + \frac{1}{2} h^\top J_n h \right| = o_{\mathbb{P}}(1). \quad (\text{A.22})$$

This applies, in particular, to every fixed radius.

Remark A.15 (The radius $b_n^{p/6}$ is sharp). The condition $\epsilon_n H_n^3 \rightarrow 0$ cannot be weakened. Under [Assumptions 1](#) and [2](#), there is $c > 0$ such that, for every fixed $\kappa > 0$, the point $h_n = \kappa \epsilon_n^{-1/3} e_s$ in the variance direction satisfies

$$\mathbb{P}\left\{L_n(h_n) - L_n(0) - h_n^\top \Delta_n + \frac{1}{2} h_n^\top J_n h_n \geq c\kappa^3\right\} \rightarrow 1.$$

Hence [\(A.22\)](#) fails for every sequence H_n with $\liminf_n \epsilon_n H_n^3 > 0$. The reason is that the third derivative of the log likelihood in the variance direction is of exact order ϵ_n , as for the variance of $N \asymp b_n^p$ independent normal observations.

The score is a vector of centered Gaussian quadratic forms. Each normalized whitened derivative has operator norm $O(\epsilon_n)$ but Frobenius norm of order one, so the quadratic forms are sums of many small independent contributions, and a Lyapunov argument gives the normal limit.

Lemma A.16 (Fisher-normalized score central limit theorem). Assume [Assumptions 1](#) and [2](#), and let M_n be 3×3 matrices with $M_n^\top M_n = \mathcal{I}_n(\theta_0)$. Then

$$M_n^{-\top} \nabla l_n(\theta_0) \implies N_3(0, I_3). \quad (\text{A.23})$$

The same quadratic-form argument applies to bounded sequences of linear combinations of Δ_n whose variances converge to a positive limit.

The last lemma is used only for [Theorem 3.6](#). Independent sampling with a density bounded below on Q satisfies [Assumption 2](#) almost surely, with polynomially small failure probability at each n .

Lemma A.17 (Coverage for independent sampling). Under the sampling assumptions of [Theorem 3.6](#), almost surely,

$$h_n(Q) = O((\log n/n)^{1/p}). \quad (\text{A.24})$$

More precisely, for every $q > 0$ there are constants $A_q, C_q < \infty$ such that $\mathbb{P}\{h_n(Q) > A_q(\log n/n)^{1/p}\} \leq C_q n^{-q}$ for all $n \geq 2$. Thus [Assumption 2](#) holds almost surely for every fixed $0 < \kappa < 1/p$.

B Proofs of the main results

B.1 Proof of [Theorem 3.1](#)

Proof. [Theorem A.13](#) proves **part (i)**, and [Theorem A.12](#) proves **part (iv)**. Write $\hat{h}_n = D_n(\hat{\theta}_n - \theta_0)$, recall $L_n(h) = l_n(\theta_0 + D_n^{-1}h)$ from [Theorem A.14](#), and set

$$Q_n(h) = h^\top \Delta_n - \frac{1}{2} h^\top J_n h, \quad h_n^* = J_n^{-1} \Delta_n, \quad R_{n,K} = \sup_{\|h\| \leq K} |L_n(h) - L_n(0) - Q_n(h)|.$$

Part (ii). For every fixed K , (A.22) applies with the fixed radius $H_n = \max(K, 1)$, so $R_{n,K} = o_{\mathbb{P}}(1)$; that is, $L_n(h) - L_n(0) = Q_n(h) + o_{\mathbb{P}}(1)$ uniformly on $\{\|h\| \leq K\}$. The function Q_n is quadratic, with $\nabla Q_n(h_n^*) = \Delta_n - J_n h_n^* = 0$ and $\nabla^2 Q_n = -J_n$, so its Taylor expansion about h_n^* is exact:

$$Q_n(h) = Q_n(h_n^*) - \frac{1}{2}(h - h_n^*)^\top J_n (h - h_n^*), \quad h \in \mathbb{R}^3. \quad (\text{B.1})$$

In particular, $Q_n(h) \leq Q_n(h_n^*)$ for all h .

Fix $\varepsilon > 0$. The score satisfies $\mathbb{E} \|\Delta_n\|^2 = \text{tr } J_n \leq C$, so $h_n^* = O_{\mathbb{P}}(1)$ by part (i), and $\hat{h}_n = O_{\mathbb{P}}(1)$ by part (iv). Choose K such that the event $E_n = \{\|\hat{h}_n\| \leq K, \|h_n^*\| \leq K\}$ has $\mathbb{P}(E_n^c) \leq \varepsilon$ for all large n . Because θ_0 is interior, the ball $\{\|h\| \leq K\}$ lies in $D_n(\Theta - \theta_0)$ for large n . Hence, on E_n , the point $\theta_0 + D_n^{-1} h_n^*$ belongs to Θ , and global optimality gives $L_n(\hat{h}_n) \geq L_n(h_n^*)$. Using the expansion at h_n^* , this inequality, the expansion at $\hat{h}_n$, and $Q_n \leq Q_n(h_n^*)$, we obtain on E_n

$$Q_n(h_n^*) - R_{n,K} \leq L_n(h_n^*) - L_n(0) \leq L_n(\hat{h}_n) - L_n(0) \leq Q_n(\hat{h}_n) + R_{n,K} \leq Q_n(h_n^*) + R_{n,K}.$$

Thus the maximized log likelihood ratio is approximated by the maximum of the quadratic:

$$|L_n(\hat{h}_n) - L_n(0) - Q_n(h_n^*)| \leq R_{n,K} \quad \text{on } E_n. \quad (\text{B.2})$$

By (B.1) at $h = \hat{h}_n$, (B.2), and the expansion at $\hat{h}_n$, on E_n

$$\begin{aligned} \frac{1}{2}(\hat{h}_n - h_n^*)^\top J_n (\hat{h}_n - h_n^*) &= Q_n(h_n^*) - Q_n(\hat{h}_n) \\ &\leq |L_n(\hat{h}_n) - L_n(0) - Q_n(h_n^*)| + |L_n(\hat{h}_n) - L_n(0) - Q_n(\hat{h}_n)| \leq 2R_{n,K}. \end{aligned}$$

Since $J_n \succeq cI_3$ by part (i), for every $\eta > 0$,

$$\mathbb{P}\{\|\hat{h}_n - h_n^*\| > \eta\} \leq \mathbb{P}(E_n^c) + \mathbb{P}\{R_{n,K} \geq c\eta^2/4\} \leq \varepsilon + o(1).$$

As ε is arbitrary, $\hat{h}_n - h_n^* = o_{\mathbb{P}}(1)$, which is (3.6). The three rates and consistency follow.

Part (iii). Let $T_n = D_n J_n^{1/2}$, so that $\mathcal{I}_n(\theta_0) = D_n J_n D_n = T_n T_n^\top$. Then

$$T_n^\top (\hat{\theta}_n - \theta_0) = J_n^{1/2} \hat{h}_n, \quad T_n^{-1} \nabla l_n(\theta_0) = J_n^{-1/2} D_n^{-1} \nabla l_n(\theta_0) = J_n^{-1/2} \Delta_n = J_n^{1/2} h_n^*,$$

and therefore

$$T_n^\top (\hat{\theta}_n - \theta_0) - T_n^{-1} \nabla l_n(\theta_0) = J_n^{1/2} (\hat{h}_n - h_n^*), \quad (\text{B.3})$$

whose norm is at most $\|J_n\|_{\text{op}}^{1/2} \|\hat{h}_n - h_n^*\| = o_{\mathbb{P}}(1)$ by parts (i) and (ii). Now let $M_n^\top M_n = \mathcal{I}_n(\theta_0)$. The matrix $O_n = M_n T_n^{-\top}$ is orthogonal, because $O_n O_n^\top = M_n \mathcal{I}_n(\theta_0)^{-1} M_n^\top = I_3$. Hence $M_n = O_n T_n^\top$ and $M_n^{-\top} = O_n T_n^{-1}$, and (B.3) gives

$$M_n (\hat{\theta}_n - \theta_0) = M_n^{-\top} \nabla l_n(\theta_0) + O_n J_n^{1/2} (\hat{h}_n - h_n^*) = M_n^{-\top} \nabla l_n(\theta_0) + o_{\mathbb{P}}(1).$$

Theorem A.16 and Slutsky's lemma prove (3.8).

For the nugget, write $\Delta_{n,t}$ for the third score coordinate. The block structure in (3.5) and (3.6) give

$$\sqrt{n}(\hat{t}_n - t_0) = 2t_0^2 \Delta_{n,t} + o_{\mathbb{P}}(1).$$

Its score variance tends to $(2t_0^2)^{-1}$, and the final assertion of [Theorem A.16](#) gives $\Delta_{n,t} \Rightarrow N(0, (2t_0^2)^{-1})$. This proves (3.9).

Part (v). Let $\mu > 0$. For $n \geq n_0$, part (iv) and $e^{-c \min(x,y)} \leq e^{-cx} + e^{-cy}$ give

$$\mathbb{E} e^{\mu \|\hat{h}_{n,\text{sp}}\|} = 1 + \int_0^\infty \mu e^{\mu H} \mathbb{P}\{\|\hat{h}_{n,\text{sp}}\| > H\} dH \leq 1 + C \int_0^\infty \mu e^{\mu H} (e^{-cH^2} + e^{-cb_n^{p/2}H}) dH,$$

which is bounded uniformly in n once $cb_n^{p/2} \geq 2\mu$. Similarly, $\mathbb{E} e^{\mu |\hat{h}_{n,t}|} \leq 1 + C \int_0^\infty \mu e^{\mu H - cH^2} dH$. Since $\|\hat{h}_n\| \leq \|\hat{h}_{n,\text{sp}}\| + |\hat{h}_{n,t}|$, the Cauchy–Schwarz inequality with $\mu = 2\lambda$ proves (3.11).

Let f be as in part (v). By part (iii) with $M_n = T_n^\top$, $Y_n = J_n^{1/2} \hat{h}_n \Rightarrow Z$. The family $\{f(\hat{h}_n)\}$ is uniformly integrable, because $f(\hat{h}_n)^2 \leq C^2 e^{2\lambda \|\hat{h}_n\|}$ has bounded expectation by (3.11). Suppose that the first convergence in (3.12) fails. Then there are $\delta > 0$ and a subsequence along which $|\mathbb{E} f(\hat{h}_n) - \mathbb{E} f(J_n^{-1/2} Z)| \geq \delta$. By part (i), a further subsequence has $J_n \rightarrow J_*$ with $cI_3 \preceq J_* \preceq CI_3$. Along it, $\hat{h}_n = J_n^{-1/2} Y_n \Rightarrow J_*^{-1/2} Z$ by Slutsky’s lemma, so uniform integrability gives $\mathbb{E} f(\hat{h}_n) \rightarrow \mathbb{E} f(J_*^{-1/2} Z)$. Also $\mathbb{E} f(J_n^{-1/2} Z) \rightarrow \mathbb{E} f(J_*^{-1/2} Z)$ by dominated convergence, since $|f(J_n^{-1/2} Z)| \leq C e^{\lambda c^{-1/2} \|Z\|}$. This contradicts the choice of δ . For the second convergence, the proof of part (iii) gives $M_n(\hat{\theta}_n - \theta_0) = O_n Y_n$ with O_n orthogonal, so $\|M_n(\hat{\theta}_n - \theta_0)\| = \|Y_n\| \leq C^{1/2} \|\hat{h}_n\|$. By (3.11), $f(M_n(\hat{\theta}_n - \theta_0))$ is uniformly integrable, and (3.8) gives $\mathbb{E} f(M_n(\hat{\theta}_n - \theta_0)) \rightarrow \mathbb{E} f(Z)$.

Finally, polynomials satisfy the growth condition. Applying the first convergence in (3.12) to the coordinates x_a and to the products $x_a x_b$, and using $\mathbb{E} J_n^{-1/2} Z = 0$ and $\mathbb{E} J_n^{-1/2} Z Z^\top J_n^{-1/2} = J_n^{-1}$, proves (3.13). $\square$

B.2 Proof of the uniformity in θ_0 in [Theorem 3.2](#)

We indicate why the statements in [Theorem 3.2](#) hold. The arguments are those of the preceding proofs, and we only point out where uniformity enters.

Part (i). The upper bounds in [Theorem A.13](#) are suprema over Θ . The lower bound compares a second-order expansion of the affinity on the unit sphere $\{\|h\| = 1\}$ with (A.16). The remainders in that expansion are controlled by the derivative bounds of [Theorem A.8](#), which hold uniformly over Θ , and the constants in (A.16) do not depend on θ_0 . For θ_0 near the boundary, the points $\theta_0 + D_n^{-1} h$ may leave Θ . Applying the same lemmas on a slightly larger rectangle $\Theta' \supset \Theta$, whose constants are again uniform, removes this difficulty. Finally, $J_{n,tt} \rightarrow (2t_0^2)^{-1}$ and the vanishing of the cross-information hold uniformly in $t_0 \in [t_-, t_+]$.

Parts (ii), (iii), and (v). Let $K \subset \text{int}(\Theta)$ be compact. Since $\|D_n^{-1}\|_{\text{op}} \rightarrow 0$, for every fixed radius H the ball $\{\|h\| \leq H\}$ lies in $D_n(\Theta - \theta_0)$ for all $\theta_0 \in K$ once n is large. The remainders in [Theorem A.14](#) are controlled by bounds whose laws do not depend on θ_0 . The Hessian fluctuation is a centered Gaussian quadratic form with Frobenius norm $O(\epsilon_n)$, and the third-derivative bound uses $Z_n^\top A_{\theta_0}^{-1} Z_n$, which is exactly $\chi_{r_n}^2$ under θ_0 . In [Theorem A.16](#), the Lyapunov ratio is at most $C\epsilon_n^2$ uniformly in θ_0 . Along any sequence $\theta_{0,n} \in K$, the Lindeberg–Feller theorem for triangular arrays therefore gives the normal limit of each fixed linear combination of the normalized score, and the Cramér–Wold device gives the joint limit. The tail bounds of [Theorem A.12](#) are also uniform. Running the argument of [Appendix B.1](#) along an arbitrary sequence $\theta_{0,n} \in K$ therefore yields the conclusions of parts (ii), (iii), and (v) along that sequence, which is equivalent to uniformity over K .

Near the boundary the argument breaks down. It uses global optimality in the form $l_n(\hat{\theta}_n) \geq$

$l_n(\theta_0 + D_n^{-1}h_n^*)$, which requires the unconstrained maximizer h_n^* of the quadratic approximation to be feasible, that is, $\theta_0 + D_n^{-1}h_n^* \in \Theta$. If θ_0 lies within distance of order $d_{n,j}^{-1}$ of the boundary in some coordinate j , this fails with probability bounded away from zero. The constrained maximizer $\hat{h}_n$ then approximates the maximizer of the quadratic over the feasible set, whose limit law is not normal.

B.3 Proof of Theorem 3.3

Proof. Consider a path varying only coordinate j . Differentiating its Gaussian density p_u under the integral and using Cauchy–Schwarz gives

$$\|\partial_u p_u\|_{L^1} = \mathbb{E}_u |\partial_u \log p_u| \leq \sqrt{\mathcal{I}_{jj,n}(u)}.$$

Consequently,

$$\text{TV}(P_{u_1,n}, P_{u_0,n}) \leq \frac{1}{2} \int_{u_0}^{u_1} \sqrt{\mathcal{I}_{jj,n}(u)} du. \quad (\text{B.4})$$

The uniform upper bounds in Theorem A.13 are $\mathcal{I}_{jj,n} \leq C d_{n,j}^2$. Choose $a > 0$ small enough that the right side of (B.4) is at most 1/4 when $u_1 - u_0 = a/d_{n,j}$. Both alternatives lie in Θ eventually.

We now reduce estimation to testing between the two alternatives (Le Cam’s two-point method). Write $u_0 = \theta_{0,j}$, $u_1 = u_0 + a/d_{n,j}$, $P_0 = P_{u_0,n}$, and $P_1 = P_{u_1,n}$, and let $\delta = (u_1 - u_0)/2 = a/(2d_{n,j})$ be half the separation. Let $\tilde{\theta}_{n,j}$ be any estimator of the j th coordinate, that is, any measurable function of the data vector Y_n . It defines a test of P_0 against P_1 : decide for whichever of the two candidate values u_0, u_1 is closer to the estimate,

$$\psi = \mathbf{1}\{|\tilde{\theta}_{n,j} - u_1| < |\tilde{\theta}_{n,j} - u_0|\},$$

so that $\psi = 1$ means “decide u_1 ” and $\psi = 0$ means “decide u_0 ”. Equivalently, $\psi = 1$ exactly when $\tilde{\theta}_{n,j}$ exceeds the midpoint $(u_0 + u_1)/2$.

A wrong decision forces a large estimation error. If $\psi = 1$, then $|\tilde{\theta}_{n,j} - u_0| > |\tilde{\theta}_{n,j} - u_1|$, and the triangle inequality $|\tilde{\theta}_{n,j} - u_0| + |\tilde{\theta}_{n,j} - u_1| \geq u_1 - u_0$ gives $|\tilde{\theta}_{n,j} - u_0| > \delta$. Likewise, if $\psi = 0$, then $|\tilde{\theta}_{n,j} - u_1| \geq \delta$. Hence

$$P_0\{|\tilde{\theta}_{n,j} - u_0| \geq \delta\} \geq P_0(\psi = 1), \quad P_1\{|\tilde{\theta}_{n,j} - u_1| \geq \delta\} \geq P_1(\psi = 0).$$

The total variation distance bounds how well any test can separate the two laws. For the event $E = \{\psi = 1\}$,

$$P_0(\psi = 1) + P_1(\psi = 0) = 1 - \{P_1(E) - P_0(E)\} \geq 1 - \text{TV}(P_0, P_1) \geq \frac{3}{4}.$$

Combining the last two displays, the two estimation error probabilities sum to at least 3/4, so the larger of them is at least 3/8. This proves (3.14) with $c_0 = 3/8$, since the estimator was arbitrary.

Finally, by Markov’s inequality, $\mathbb{E}_{\theta,n}(\tilde{\theta}_{n,j} - \theta_j)^2 \geq \delta^2 \mathbb{P}_{\theta,n}\{|\tilde{\theta}_{n,j} - \theta_j| \geq \delta\}$. Taking the larger of the two alternatives and using (3.14) gives $\sup_{\theta \in \Theta} \mathbb{E}_{\theta,n}(\tilde{\theta}_{n,j} - \theta_j)^2 \geq \frac{3}{8} \cdot \frac{a^2}{4d_{n,j}^2}$, which is (3.15).

For the upper bound, recall from Theorem A.12 that the constants in the tail bound (3.10) do not depend on $\theta_0 \in \Theta$, and that interiority of θ_0 is not needed. Integrating the tail bound as in the

proof of [Theorem 3.1\(v\)](#) therefore gives, for $n \geq n_0$,

$$\sup_{\theta \in \Theta} \mathbb{E}_{\theta,n} \{d_{n,j}^2(\hat{\theta}_{n,j} - \theta_j)^2\} \leq C,$$

with C independent of n and θ . Since Θ is bounded, the finitely many $n < n_0$ are covered by enlarging C . This proves (3.16). Combining this with (3.15) gives (3.17). Since

$$(d_{n,s}^{-1}, d_{n,l}^{-1}, d_{n,t}^{-1}) = (b_n^{-p/2}, b_n^{-(p+2)/2}, n^{-1/2}),$$

the stated minimax root MSE rates follow. $\square$

B.4 Proofs of the Corollaries in [Section 3.4](#)

B.4.1 Proof of [Theorem 3.4](#)

Proof. Let $\theta_- = \min(s_-, l_-, t_-)$ and $\theta_+ = \max(s_+, l_+, t_+)$, and write $\hat{h}_n = D_n(\hat{\theta}_n - \theta_0)$.

Information. The diagonal entries of Λ_0 lie in $[\theta_-, \theta_+]$. Hence $c\theta_-^2 I_3 \preceq J_n^{\log} = \Lambda_0 J_n \Lambda_0 \preceq C\theta_+^2 I_3$, and $J_n^{\log}$ inherits the block structure (3.5), with $J_{n,tt}^{\log} = t_0^2 J_{n,tt} \rightarrow 1/2$. This is part (i) of [Theorem 3.1](#) on the log scale.

Expansion and rates. A second-order Taylor expansion of $\log$, coordinatewise on $[\theta_-, \theta_+]$, gives

$$D_n(\hat{\lambda}_n - \lambda_0) = \Lambda_0^{-1} \hat{h}_n + R_n, \quad \|R_n\| \leq \frac{\|D_n^{-1}\|_{\text{op}} \|\hat{h}_n\|^2}{2\theta_-^2}. \quad (\text{B.5})$$

Since $\|D_n^{-1}\|_{\text{op}} = \epsilon_n$ for large n , (3.11) gives $R_n = O_{\mathbb{P}}(\epsilon_n)$, together with all moments. By (3.6),

$$\Lambda_0^{-1} \hat{h}_n = \Lambda_0^{-1} J_n^{-1} D_n^{-1} \nabla l_n(\theta_0) + o_{\mathbb{P}}(1) = (J_n^{\log})^{-1} D_n^{-1} \{\Lambda_0 \nabla l_n(\theta_0)\} + o_{\mathbb{P}}(1),$$

which is part (ii) on the log scale. The rates follow.

Normal limits. Let $M_n^{\top} M_n = \mathcal{I}_n^{\log}(\theta_0)$. Then $\|M_n D_n^{-1}\|_{\text{op}}^2 = \|J_n^{\log}\|_{\text{op}} \leq C$, and

$$M_n D_n^{-1} (J_n^{\log})^{-1} D_n^{-1} = M_n \mathcal{I}_n^{\log}(\theta_0)^{-1} = M_n^{-\top}.$$

Hence, by the previous step,

$$M_n(\hat{\lambda}_n - \lambda_0) = M_n^{-\top} \Lambda_0 \nabla l_n(\theta_0) + o_{\mathbb{P}}(1) = \widetilde{M}_n^{-\top} \nabla l_n(\theta_0) + o_{\mathbb{P}}(1), \quad \widetilde{M}_n = M_n \Lambda_0^{-1}.$$

Since $\widetilde{M}_n^{\top} \widetilde{M}_n = \Lambda_0^{-1} \mathcal{I}_n^{\log}(\theta_0) \Lambda_0^{-1} = \mathcal{I}_n(\theta_0)$, [Theorem A.16](#) gives $M_n(\hat{\lambda}_n - \lambda_0) \Rightarrow N_3(0, I_3)$. For the nugget, the third coordinate of (B.5) reads $\sqrt{n}(\log \hat{t}_n - \log t_0) = t_0^{-1} \sqrt{n}(\hat{t}_n - t_0) + R_{n,t}$ with $R_{n,t} = O_{\mathbb{P}}(n^{-1/2})$, and (3.9) gives the $N(0, 2)$ limit.

Tails and moments. Since $|\log x - \log y| \leq |x - y|/\theta_-$ on $[\theta_-, \theta_+]$, each coordinate satisfies $|d_{n,j}(\hat{\lambda}_{n,j} - \lambda_{0,j})| \leq |\hat{h}_{n,j}|/\theta_-$. Hence (3.10) holds for $D_n(\hat{\lambda}_n - \lambda_0)$ with H replaced by $\theta_- H$, and (3.11) holds with the constant in the exponent divided by θ_- . For the limits in part (v), $R_n \rightarrow 0$ in probability in (B.5). Along any subsequence with $J_n \rightarrow J_*$, $\Lambda_0^{-1} \hat{h}_n \Rightarrow \Lambda_0^{-1} J_*^{-1/2} Z$, which has law $N_3\{0, (\Lambda_0 J_* \Lambda_0)^{-1}\}$. The subsequence argument in the proof of [Theorem 3.1\(v\)](#) therefore applies with J_n replaced by $J_n^{\log}$. In particular, $D_n \mathbb{E}[(\hat{\lambda}_n - \lambda_0)(\hat{\lambda}_n - \lambda_0)^{\top}] D_n = (J_n^{\log})^{-1} + o(1)$. Since

$cI_3 \preceq J_n \preceq CI_3$, the diagonal entries of $(J_n^{\log})^{-1} = \Lambda_0^{-1} J_n^{-1} \Lambda_0^{-1}$ lie between $(C\theta_+^2)^{-1}$ and $(c\theta_-^2)^{-1}$, which gives the stated MSE orders for $\log s$ and $\log l$. For the nugget, the block structure (3.5) and the Schur complement formula give $[J_n^{-1}]_{tt} \rightarrow 2t_0^2$, hence $[(J_n^{\log})^{-1}]_{tt} = t_0^{-2}[J_n^{-1}]_{tt} \rightarrow 2$.

Uniformity. The additional constants above depend only on θ_- and θ_+ , so the uniformity statements of Theorem 3.2 carry over. $\square$

B.4.2 Proof of Theorem 3.5

Proof. Write $\hat{h}_n = D_n(\hat{\theta}_n - \theta_0)$ and $\hat{J}_n = D_n^{-1} \hat{\mathcal{I}}_n D_n^{-1}$. By Theorem 3.1(iv), $\hat{h}_n = O_{\mathbb{P}}(1)$.

Part (i) for the plug-in Fisher information. For $\theta \in \Theta$, put $J_n(\theta) = D_n^{-1} \mathcal{I}_n(\theta) D_n^{-1}$, so that $J_n(\theta_0) = J_n$. With

$$B_a(\theta) = d_{n,a}^{-1} C_n(\theta)^{-1/2} C_{n,a}(\theta) C_n(\theta)^{-1/2}, \quad B_{ab}(\theta) = d_{n,a}^{-1} d_{n,b}^{-1} C_n(\theta)^{-1/2} C_{n,ab}(\theta) C_n(\theta)^{-1/2},$$

we have $[J_n(\theta)]_{ab} = \frac{1}{2} \text{tr}\{B_a(\theta) B_b(\theta)\}$. Differentiating $\mathcal{I}_n(\theta)$, using $\partial_c C_n^{-1} = -C_n^{-1} C_{n,c} C_n^{-1}$, and normalizing gives, in the coordinates $h = D_n(\theta - \theta_0)$,

$$\partial_{h_c} [J_n(\theta)]_{ab} = \frac{1}{2} \text{tr}\{-B_c B_a B_b - B_a B_c B_b + B_{ac} B_b + B_a B_{bc}\},$$

with all matrices evaluated at θ . By Theorem A.8, uniformly over $\theta \in \Theta$, $\|B_a(\theta)\|_{\text{op}} \leq C\epsilon_n$ and $\|B_a(\theta)\|_{\text{F}} \leq C$ for $a \in \{s, l, t\}$, where for the nugget $\|B_t\|_{\text{op}} = O(n^{-1/2}) \leq C\epsilon_n$. Moreover, $\|B_{ac}(\theta)\|_{\text{F}} \leq C\epsilon_n$, and $B_{ac} = 0$ when a or c is t . Since $|\text{tr}(XYZ)| \leq \|X\|_{\text{op}} \|Y\|_{\text{F}} \|Z\|_{\text{F}}$ and $|\text{tr}(XY)| \leq \|X\|_{\text{F}} \|Y\|_{\text{F}}$, every entry of $\nabla_h J_n(\theta)$ is bounded by $C\epsilon_n$, uniformly over Θ . As Θ is convex, the mean value theorem along the segment from θ_0 to θ gives

$$\|J_n(\theta) - J_n\|_{\text{op}} \leq C\epsilon_n \|D_n(\theta - \theta_0)\|, \quad \theta \in \Theta. \quad (\text{B.6})$$

Taking $\theta = \hat{\theta}_n$ proves (3.18) for $\hat{\mathcal{I}}_n = \mathcal{I}_n(\hat{\theta}_n)$. This step uses neither interiority of θ_0 nor any local expansion, and it holds for every $\theta_0 \in \Theta$.

Part (i) for the observed information. Let L_n be as in Theorem A.14. Then

$$D_n^{-1} \{-\nabla^2 l_n(\hat{\theta}_n)\} D_n^{-1} = -\nabla_h^2 L_n(\hat{h}_n).$$

Fix $K \geq 1$. On the event $\{\|\hat{h}_n\| \leq K\}$, the mean value theorem and (A.20) with $H_n = K$ give

$$\left\| \nabla_h^2 L_n(\hat{h}_n) - \nabla_h^2 L_n(0) \right\|_{\text{op}} \leq CK \sup_{\|h\| \leq K} \max_{a,b,c} |\partial_{h_a h_b h_c}^3 L_n(h)| = O_{\mathbb{P}}(\epsilon_n),$$

and (A.19) gives $-\nabla_h^2 L_n(0) = J_n + O_{\mathbb{P}}(\epsilon_n)$. Since $\mathbb{P}\{\|\hat{h}_n\| > K\}$ is arbitrarily small for large K , (3.18) follows for the observed information.

Consequences of (3.18). Since $cI_3 \preceq J_n \preceq CI_3$, with probability tending to one $\frac{c}{2}I_3 \preceq \hat{J}_n \preceq 2CI_3$, so that $\hat{\mathcal{I}}_n = D_n \hat{J}_n D_n$ is positive definite. On this event,

$$\left\| \hat{J}_n^{-1} - J_n^{-1} \right\|_{\text{op}} \leq \left\| \hat{J}_n^{-1} \right\|_{\text{op}} \left\| \hat{J}_n - J_n \right\|_{\text{op}} \left\| J_n^{-1} \right\|_{\text{op}} = O_{\mathbb{P}}(\epsilon_n).$$

Since $[\widehat{\mathcal{I}}_n^{-1}]_{jj} = d_{n,j}^{-2}[\widehat{J}_n^{-1}]_{jj}$, $[\mathcal{I}_n(\theta_0)^{-1}]_{jj} = d_{n,j}^{-2}[J_n^{-1}]_{jj}$, and $[J_n^{-1}]_{jj} \geq C^{-1}$, the ratio statement in (i) follows.

Part (ii). Let $\Delta_n = D_n^{-1} \nabla l_n(\theta_0)$. By (3.6), $\widehat{h}_{n,j} = e_j^\top J_n^{-1} \Delta_n + o_{\mathbb{P}}(1)$. The variable $V_n = e_j^\top J_n^{-1} \Delta_n / [J_n^{-1}]_{jj}^{1/2}$ has variance one, since $\text{Cov}(\Delta_n) = J_n$, and it is a linear combination of Δ_n with coefficient vector $J_n^{-1} e_j / [J_n^{-1}]_{jj}^{1/2}$ of norm at most $C^{1/2}/c$. By the final assertion of Theorem A.16, $V_n \Rightarrow N(0, 1)$. As $[J_n^{-1}]_{jj} \geq C^{-1}$, also $\widehat{h}_{n,j} / [J_n^{-1}]_{jj}^{1/2} = V_n + o_{\mathbb{P}}(1)$. Finally,

$$\frac{\widehat{\theta}_{n,j} - \theta_{0,j}}{[\widehat{\mathcal{I}}_n^{-1}]_{jj}^{1/2}} = \frac{\widehat{h}_{n,j}}{[J_n^{-1}]_{jj}^{1/2}} \left(\frac{[J_n^{-1}]_{jj}}{[\widehat{J}_n^{-1}]_{jj}} \right)^{1/2},$$

and the last factor is $1 + O_{\mathbb{P}}(\epsilon_n)$ by part (i). Slutsky's lemma gives the normal limit. The interval contains $\theta_{0,j}$ exactly when the absolute value of this ratio is at most $z_{1-\alpha/2}$, which gives the coverage statement.

Part (iii). For a positive definite matrix A , write $\text{chol}(A)$ for its lower-triangular Cholesky factor with positive diagonal. Because D_n is diagonal with positive entries, $D_n \text{chol}(\widehat{J}_n)$ is lower triangular with positive diagonal and $\{D_n \text{chol}(\widehat{J}_n)\} \{D_n \text{chol}(\widehat{J}_n)\}^\top = \widehat{\mathcal{I}}_n$. By uniqueness of the Cholesky factor, $\widehat{L}_n = D_n \text{chol}(\widehat{J}_n)$. The map chol is smooth on positive definite matrices, hence Lipschitz on the compact convex set $\{A : \frac{c}{2} I_3 \preceq A \preceq 2C I_3\}$, so $\text{chol}(\widehat{J}_n) - \text{chol}(J_n) = O_{\mathbb{P}}(\epsilon_n)$. Therefore

$$\begin{aligned} \widehat{L}_n^\top (\widehat{\theta}_n - \theta_0) &= \text{chol}(\widehat{J}_n)^\top \widehat{h}_n = \text{chol}(J_n)^\top \widehat{h}_n + O_{\mathbb{P}}(\epsilon_n) \\ &= M_n (\widehat{\theta}_n - \theta_0) + O_{\mathbb{P}}(\epsilon_n), \quad M_n = \text{chol}(J_n)^\top D_n. \end{aligned}$$

Since $M_n^\top M_n = D_n \text{chol}(J_n) \text{chol}(J_n)^\top D_n = \mathcal{I}_n(\theta_0)$, Theorem 3.1(iii) gives the first limit. The second follows from $(\widehat{\theta}_n - \theta_0)^\top \widehat{\mathcal{I}}_n (\widehat{\theta}_n - \theta_0) = \left\| \widehat{L}_n^\top (\widehat{\theta}_n - \theta_0) \right\|^2$ and the continuous mapping theorem, and the ellipsoid statement is a restatement of it.

Part (iv). Since $\widehat{\Lambda}_n - \Lambda_0 = \text{diag}(\widehat{\theta}_n - \theta_0)$, $\left\| \widehat{\Lambda}_n - \Lambda_0 \right\|_{\text{op}} \leq \|D_n^{-1}\|_{\text{op}} \left\| \widehat{h}_n \right\| = O_{\mathbb{P}}(\epsilon_n)$. Because diagonal matrices commute,

$$D_n^{-1} \widehat{\mathcal{I}}_n^{\log} D_n^{-1} = \widehat{\Lambda}_n \widehat{J}_n \widehat{\Lambda}_n = \Lambda_0 J_n \Lambda_0 + O_{\mathbb{P}}(\epsilon_n) = J_n^{\log} + O_{\mathbb{P}}(\epsilon_n),$$

which is part (i) on the log scale. By Theorem 3.4, $D_n(\widehat{\lambda}_n - \lambda_0) = (J_n^{\log})^{-1} D_n^{-1} \{\Lambda_0 \nabla l_n(\theta_0)\} + o_{\mathbb{P}}(1)$, and the normal limit of Theorem 3.1(iii) holds on the log scale. The proofs of parts (ii) and (iii) therefore apply verbatim, with $\widehat{h}_n$ replaced by $D_n(\widehat{\lambda}_n - \lambda_0)$, J_n by $J_n^{\log}$, Δ_n by $\Lambda_0 \Delta_n$, and $\mathcal{I}_n(\theta_0)$ by $\mathcal{I}_n^{\log}(\theta_0)$. The displayed interval is the image under $\exp$ of the log-scale Wald interval for $\lambda_{0,j}$. Finally, the chain rule gives

$$-\nabla_{\lambda}^2 l_n = \Lambda \{-\nabla_{\theta}^2 l_n\} \Lambda - \text{diag}(\theta \circ \nabla_{\theta} l_n), \quad \Lambda = \text{diag}(\theta),$$

where $\circ$ denotes the entrywise product. At an interior maximizer the gradient vanishes. Since $\theta_0 \in \text{int}(\Theta)$ and $\widehat{\theta}_n$ is consistent, $\widehat{\theta}_n \in \text{int}(\Theta)$ with probability tending to one. $\square$

B.4.3 Proof of Theorem 3.6

Proof. Write $\widehat{h}_n = D_n(\widehat{\theta}_n - \theta_0)$.

A coverage event with stretched-exponential failure probability. Let Q have side length L . For $n \geq 1$, let $k_n = \lfloor n^{1/(4p)} \rfloor$, and partition Q into k_n^p closed subcubes of side length L/k_n . Let H_n be the event that every subcube contains at least one of $X_1, \dots, X_n$. As in the proof of [Theorem A.17](#), each subcube has sampling probability at least $f_- L^p / k_n^p$, so a union bound gives

$$\mathbb{P}(H_n^c) \leq k_n^p \exp\left(-\frac{nf_- L^p}{k_n^p}\right) \leq n^{1/4} \exp(-f_- L^p n^{3/4}), \quad (\text{B.7})$$

because $k_n^p \leq n^{1/4}$. On H_n , every point of Q lies within the diameter $\sqrt{p} L / k_n$ of an observation. Since $\lfloor y \rfloor \geq y/2$ for $y \geq 1$, this gives $h_n(Q) \leq 2\sqrt{p} L n^{-1/(4p)}$. Hence, on H_n , (3.2) holds at n with $\kappa = 1/(4p)$ and $C_h = 2\sqrt{p} L$. Fix these constants from now on. The constants in [Theorems A.8](#) to [A.10](#) and [A.12](#) depend on the design only through $\mathcal{X}$, Q , C_h , and κ , and the bounds of [Theorem A.12](#) hold at every $n \geq n_0$ at which (3.2) is satisfied. Therefore every bound of [Theorem 3.1](#)(iv) and (v) holds on H_n for $n \geq n_0$, with constants that do not depend on the design. By (B.7), $\sum_n \mathbb{P}(H_n^c) < \infty$, so by the first Borel–Cantelli lemma, almost surely H_n occurs for all sufficiently large n .

Two bounds hold for every design in the bounded set $\mathcal{X}$. The upper information bounds of [Theorem A.13](#) require only bounded locations, so $\|J_n\|_{\text{op}} \leq C$. The rescaled parameter set $D_n(\Theta - \theta_0)$ has diameter at most $C\sqrt{n}$, so $\|\hat{h}_n\| \leq C\sqrt{n}$.

Part (a). The design is independent of the process and the errors, so conditionally on $\mathbf{X}$ the observations follow the model (2.1) with deterministic locations. For almost every realization of $\mathbf{X}$, H_n occurs for all large n , so [Assumption 2](#) holds with the constants fixed above. All results proved for deterministic designs therefore apply conditionally on $\mathbf{X}$, with design-independent constants and design-dependent thresholds.

Part (b)(i). This is part (a) applied to [Theorem 3.1](#)(i), since J_n is a function of $\mathbf{X}$ alone.

Part (b)(ii), and the convergence statements in (b)(vi). Let V_n be any statistic in these statements whose limit law does not depend on the design, and let V have that limit law. This covers the standardized errors in [Theorem 3.1](#)(iii) and [Theorems 3.4](#) and [3.5](#) and the quadratic forms with χ_3^2 limits. For bounded continuous φ , part (a) gives $\mathbb{E}[\varphi(V_n) \mid \mathbf{X}] \rightarrow \mathbb{E}\varphi(V)$ almost surely, and dominated convergence gives $\mathbb{E}\varphi(V_n) \rightarrow \mathbb{E}\varphi(V)$. Coverage probabilities are treated in the same way. For statements of the form $R_n = o_{\mathbb{P}}(1)$, such as the expansion in [Theorem 3.1](#)(ii), part (a) gives $\mathbb{P}(\|R_n\| > \eta \mid \mathbf{X}) \rightarrow 0$ almost surely for every $\eta > 0$, and dominated convergence gives $\mathbb{P}(\|R_n\| > \eta) \rightarrow 0$. For statements of the form $R_n = O_{\mathbb{P}}(r_n)$, such as the rates in [Theorem 3.1](#)(ii) and [Theorem 3.5](#)(i), let $q_{n,M}(\mathbf{X}) = \mathbb{P}(\|R_n\| > Mr_n \mid \mathbf{X})$. Part (a) gives $\lim_{M \rightarrow \infty} \limsup_n q_{n,M}(\mathbf{X}) = 0$ almost surely. Since $0 \leq q_{n,M} \leq 1$, the reverse Fatou lemma gives

$$\lim_{M \rightarrow \infty} \limsup_n \mathbb{E} q_{n,M}(\mathbf{X}) \leq \lim_{M \rightarrow \infty} \mathbb{E} \left\{ \limsup_n q_{n,M}(\mathbf{X}) \right\} = 0.$$

Part (b)(iii). For $n \geq n_0$, by (B.7),

$$\mathbb{P}\{\|\hat{h}_{n,\text{sp}}\| \geq H\} \leq \mathbb{E}[\mathbb{P}\{\|\hat{h}_{n,\text{sp}}\| \geq H \mid \mathbf{X}\} \mathbf{1}_{H_n}] + \mathbb{P}(H_n^c) \leq C \exp\{-c \min(H^2, b_n^{p/2} H)\} + C e^{-cn^{3/4}},$$

and the same argument applies to the nugget coordinate.

Part (b)(iv). For $n \geq n_\lambda$, the exponential moment bound of [Theorem 3.1](#)(v) on H_n , together

with $\|\hat{h}_n\| \leq C\sqrt{n}$, gives

$$\mathbb{E} \exp\{\lambda \|\hat{h}_n\|\} \leq \mathbb{E}[\mathbb{E}\{\exp(\lambda \|\hat{h}_n\|) \mid \mathbf{X}\} \mathbf{1}_{H_n}] + e^{\lambda C\sqrt{n}} \mathbb{P}(H_n^c) \leq C_\lambda + e^{\lambda C\sqrt{n}} n^{1/4} e^{-f-L^p n^{3/4}},$$

which is bounded in n . Now let g be continuous with $|g(x)| \leq Ce^{\lambda\|x\|}$. Since $\|M_n(\hat{\theta}_n - \theta_0)\| = \|J_n^{1/2}\hat{h}_n\| \leq C\|\hat{h}_n\| \leq C\sqrt{n}$,

$$\mathbb{E}[|g(M_n(\hat{\theta}_n - \theta_0))| \mathbf{1}_{H_n^c}] \leq Ce^{\lambda C\sqrt{n}} \mathbb{P}(H_n^c) \rightarrow 0.$$

On H_n , [Theorem 3.1\(v\)](#) bounds $\mathbb{E}[\exp\{\lambda C\|\hat{h}_n\|\} \mid \mathbf{X}]$ uniformly in the design, so $\mathbb{E}[g(M_n(\hat{\theta}_n - \theta_0)) \mid \mathbf{X}] \mathbf{1}_{H_n}$ is uniformly bounded. It converges to $\mathbb{E}g(Z)$ almost surely, by part (a) and because $\mathbf{1}_{H_n} \rightarrow 1$ almost surely. Dominated convergence over the design gives $\mathbb{E}g(M_n(\hat{\theta}_n - \theta_0)) \rightarrow \mathbb{E}g(Z)$. Taking $g(x) = x_a x_b$ gives the second-moment statement. For the bias, part (a) gives $\mathbb{E}[\hat{h}_n \mid \mathbf{X}] \rightarrow 0$ almost surely; on H_n this conditional mean is bounded uniformly in the design by the exponential moment bound, and $\|\mathbb{E}[\hat{h}_n \mathbf{1}_{H_n^c}]\| \leq C\sqrt{n} \mathbb{P}(H_n^c) \rightarrow 0$. Dominated convergence gives $D_n \mathbb{E}(\hat{\theta}_n - \theta_0) = \mathbb{E}\hat{h}_n \rightarrow 0$.

Part (b)(v). The proof of the lower bounds in [Theorem 3.3](#) uses only bounded locations. For every realized design, it shows that any estimator satisfies $\mathbb{P}_{\theta_0}\{|\tilde{\theta}_{n,j} - \theta_{0,j}| \geq \delta \mid \mathbf{X}\} + \mathbb{P}_{\theta_{1,n}^{(j)}}\{|\tilde{\theta}_{n,j} - \theta_{1,n,j}^{(j)}| \geq \delta \mid \mathbf{X}\} \geq 3/4$, with $\delta = a/(2d_{n,j})$. Taking expectations over the design, the two unconditional error probabilities also sum to at least $3/4$, so the larger is at least $3/8$. This gives the lower bounds in [Theorem 3.3](#), as in its proof. For the upper bound, the constants of [Theorem A.12](#) also do not depend on $\theta_0 \in \Theta$. Hence, for $n \geq n_0$ and every $\theta \in \Theta$,

$$\mathbb{E}_\theta\{d_{n,j}^2(\hat{\theta}_{n,j} - \theta_j)^2\} \leq \mathbb{E}_\theta[\mathbb{E}_\theta\{d_{n,j}^2(\hat{\theta}_{n,j} - \theta_j)^2 \mid \mathbf{X}\} \mathbf{1}_{H_n}] + C d_{n,j}^2 \mathbb{P}(H_n^c) \leq C,$$

because Θ is bounded, $d_{n,j}^2 \leq n$, and $n \mathbb{P}(H_n^c) \rightarrow 0$ by [\(B.7\)](#). The finitely many $n < n_0$ are covered by enlarging C .

Part (b)(vi). The convergence in distribution, convergence in probability, and coverage statements of [Theorems 3.4](#) and [3.5](#) were treated above. For the plug-in Fisher information in [Theorem 3.5\(i\)](#), the Lipschitz bound for $J_n(\theta)$ in the proof of [Theorem 3.5](#) ([Appendix B.4.2](#)) gives $\|D_n^{-1} \mathcal{I}_n(\hat{\theta}_n) D_n^{-1} - J_n\|_{\text{op}} \leq C\epsilon_n \|\hat{h}_n\|$ for every bounded design, and $\hat{h}_n = O_{\mathbb{P}}(1)$ unconditionally by part (b)(ii). For the observed information, [Theorem 3.5\(i\)](#) holds conditionally by part (a), and the $O_{\mathbb{P}}$ argument above gives the unconditional version. Finally, the proofs of (b)(i)–(v) apply verbatim on the log scale with $\hat{h}_n$ replaced by $D_n(\hat{\lambda}_n - \lambda_0)$, because $|d_{n,j}(\hat{\lambda}_{n,j} - \lambda_{0,j})| \leq |\hat{h}_{n,j}|/\theta_-$, where $\theta_- = \min(s_-, l_-, t_-)$, and because the log-scale versions of the conditional statements hold by part (a). $\square$

C Proofs of the lemmas in [Appendix A](#)

C.1 Proof of [Theorem A.1](#)

Proof. Let $\rho = (P+Q)/2$, a probability law dominating P and Q , and let $p = dP/d\rho$ and $q = dQ/d\rho$. The affinity does not depend on the dominating measure, so $\text{Aff}(P, Q) = \mathbb{E}_\rho \sqrt{pq}$. Put $\rho_T = \rho \circ T^{-1}$,

which dominates $P \circ T^{-1}$ and $Q \circ T^{-1}$. For $F \in \mathcal{F}'$,

$$P\{T^{-1}(F)\} = \mathbb{E}_\rho[p \mathbf{1}\{T \in F\}] = \mathbb{E}_\rho[\mathbb{E}_\rho(p \mid T) \mathbf{1}\{T \in F\}],$$

so the density of $P \circ T^{-1}$ with respect to ρ_T , evaluated at T , is $\mathbb{E}_\rho(p \mid T)$, ρ -almost surely. The same holds for Q . Hence

$$\text{Aff}(P \circ T^{-1}, Q \circ T^{-1}) = \mathbb{E}_\rho \sqrt{\mathbb{E}_\rho(p \mid T) \mathbb{E}_\rho(q \mid T)}.$$

The conditional Cauchy–Schwarz inequality gives $\mathbb{E}_\rho(\sqrt{pq} \mid T) \leq \sqrt{\mathbb{E}_\rho(p \mid T) \mathbb{E}_\rho(q \mid T)}$ almost surely. Taking expectations yields $\text{Aff}(P, Q) = \mathbb{E}_\rho \sqrt{pq} \leq \text{Aff}(P \circ T^{-1}, Q \circ T^{-1})$. The Gaussian statement is the case $T(y) = Hy$, since HY is $N_r(0, HAH^\top)$ or $N_r(0, HBH^\top)$, and these covariances are positive definite because H has full row rank. $\square$

C.2 Proof of Theorem A.2

Proof. If $B = 0$ there is nothing to prove. Diagonalizing B , the variable $Q = X^\top BX - \text{tr } B$ has the law of $\sum_j \lambda_j(\eta_j^2 - 1)$ for independent standard normal η_j , where $\sum_j \lambda_j^2 = \|B\|_F^2$. For $|v| \leq 1/2$,

$$-\frac{1}{2} \log(1 - v) - \frac{v}{2} = \sum_{r \geq 2} \frac{v^r}{2r} \leq \frac{v^2}{4(1 - |v|)} \leq \frac{v^2}{2}.$$

Hence, for $|s| \leq 1/(4\|B\|_F)$, so that $|2s\lambda_j| \leq 1/2$ for every j ,

$$\log \mathbb{E} e^{sQ} = \sum_j \left\{ -\frac{1}{2} \log(1 - 2s\lambda_j) - s\lambda_j \right\} \leq 2s^2 \|B\|_F^2 \leq \frac{1}{8}.$$

Applying this with $s = \pm 1/(4\|B\|_F)$ gives $\mathbb{E} \exp\{|Q|/(4\|B\|_F)\} \leq 2e^{1/8}$. Since $x^q \leq C_q e^x$ for $x \geq 0$, it follows that $\mathbb{E}|Q|^q \leq 2e^{1/8} C_q (4\|B\|_F)^q$. $\square$

C.3 Proof of Theorem A.3

Proof. Splitting the second moment according to whether $|X| < \lambda\sigma$,

$$\sigma^2 \leq \mathbb{E} X^2 = \mathbb{E}[X^2 \mathbf{1}\{|X| < \lambda\sigma\}] + \mathbb{E}[X^2 \mathbf{1}\{|X| \geq \lambda\sigma\}] \leq \lambda^2 \sigma^2 + M^2 \mathbb{P}(|X| \geq \lambda\sigma).$$

Rearranging gives the first claim. For the second, $\mathbb{P}(|f(\iota)| \geq \lambda\sigma)$ is the number of $\iota \in \mathcal{A}$ with $|f(\iota)| \geq \lambda\sigma$, divided by $|\mathcal{A}|$. $\square$

C.4 Proof of Theorem A.4

Proof. Inequality (1.10) of Talagrand [2021] shows that there is a countable dense set $G \subseteq [0, 1]^k$ such that, for all $j \in \mathbb{N}_0$,

$$\left\| \sup_{x, y \in G, |x-y| \leq 2^{-j}} |\xi(x) - \xi(y)| \right\|_{L^m} \leq C_{k, m, \alpha} L 2^{-j(\alpha-k)/m}.$$

Letting $j \rightarrow \infty$ shows that $\xi|_G$ is almost surely uniformly continuous. Its continuous extension to $[0, 1]^k$ is a modification of ξ , because ξ is continuous in L^m . For $h \in (0, 1)$, choose $j \in \mathbb{N}_0$ with $2^{-j-1} \leq h < 2^{-j}$. If $|x - y| \leq h$ and $x_r \rightarrow x$, $y_r \rightarrow y$ along G , then $|x_r - y_r| < 2^{-j}$ eventually, so the supremum over $|x - y| \leq h$ is at most the supremum over G at scale 2^{-j} . Since $2^{-j} \leq 2h$, this proves the claim for $h < 1$. The case $h = 1$ follows from $h = 1/2$ by the triangle inequality through midpoints. $\square$

The classical Kolmogorov–Chentsov theorem [Revuz and Yor, 1999, Theorem I.2.1] gives the same conclusion with $h^{(\alpha-k)/m}$ replaced by h^γ for any $\gamma < (\alpha - k)/m$. This weaker form would suffice below, after replacing $m - k$ by a smaller positive number in Theorem A.5.

C.5 Proof of Theorem A.5

Proof. If $M = 0$, then $\xi = 0$ almost surely on a countable dense subset of Γ , and hence everywhere by continuity. Assume $M > 0$. Fix $a \in \Gamma$, let π be the metric projection of $\mathbb{R}^k$ onto Γ , which is 1-Lipschitz, and put $\eta(x) = \xi(\pi(a - \delta \mathbf{1} + 2\delta x))$ for $x \in [0, 1]^k$, where $\mathbf{1} = (1, \dots, 1)$. Since $\Gamma \subseteq a + [-\delta, \delta]^k$ and π is the identity on Γ , $\sup_{[0, 1]^k} |\eta| = \sup_\Gamma |\xi|$. Moreover, η has continuous paths, $\sup_x \|\eta(x)\|_{L^m} \leq M$, and $\|\eta(x) - \eta(y)\|_{L^m} \leq 2\delta L|x - y|$. By Theorem A.4 with $\alpha = m > k$, applied to η , whose continuous modification is indistinguishable from η ,

$$\left\| \sup_{|x-y| \leq h} |\eta(x) - \eta(y)| \right\|_{L^m} \leq C\delta L h^{1-k/m}, \quad h \in (0, 1].$$

Let $h \in (0, 1]$ and cover $[0, 1]^k$ by at most $\lceil 1/h \rceil^k \leq (2/h)^k$ closed cubes B of side h , choosing a point x_B in each. Every $x \in [0, 1]^k$ lies within $\sqrt{k}h$ of some x_B , so

$$\sup_x |\eta(x)| \leq \max_B |\eta(x_B)| + \sup_{|x-y| \leq \sqrt{k}h} |\eta(x) - \eta(y)|.$$

Since $\|\max_B |X_B|\|_{L^m} \leq (\sum_B \|X_B\|_{L^m}^m)^{1/m}$, the first term has L^m norm at most $(2/h)^{k/m} M$. Dividing the segment from x to y into $\lceil \sqrt{k} \rceil$ pieces of length at most h , the second term has L^m norm at most $C\delta L h^{1-k/m}$. Thus

$$\left\| \sup_\Gamma |\xi| \right\|_{L^m} \leq C\{Mh^{-k/m} + \delta L h^{1-k/m}\}, \quad 0 < h \leq 1.$$

If $M \geq \delta L$, take $h = 1$ to obtain the bound $2CM$. Otherwise take $h = M/(\delta L)$ to obtain $2CM^{1-k/m}(\delta L)^{k/m}$. Raising to the power m proves the claim. $\square$

C.6 Proof of Theorem A.6

Proof. Each $\sup_{\Gamma_j} |\xi|$ is measurable, being a supremum over a countable dense subset of Γ_j . Since $\sup_{u \in \bigcup_j \Gamma_j} |\xi(u)|^m \leq \sum_j \sup_{u \in \Gamma_j} |\xi(u)|^m$, monotone convergence and Theorem A.5 prove the claim. $\square$

C.7 Proof of Theorem A.7

Proof. We use two elementary properties of $\psi = \psi_\gamma$. First, ψ is nondecreasing, and $\psi(r/2) \geq \psi(r)/4$. Hence, for $c \geq 0$,

$$\psi((r-c)_+) \geq \psi(r)/4 - c^2, \quad r \geq 0 : \quad (\text{C.1})$$

if $r \geq 2c$ this follows from $(r-c)_+ \geq r/2$, and if $r < 2c$ the right side is negative. Second, $\psi_\gamma \geq \psi_1$ because $\gamma \geq 1$.

Let $A_H = \{u \in U : |u_1| \geq H_1, |u_2| \geq H_2\}$, which we may assume nonempty. For $z = (z_1, z_2) \in \mathbb{Z}^{k_1} \times \mathbb{Z}^{k_2}$ put $\Gamma_z = U \cap (z + [0, 1]^k)$, and let $\mathcal{Z}$ be the set of z with $\Gamma_z \cap A_H \neq \emptyset$. Each Γ_z with $z \in \mathcal{Z}$ is a nonempty product of compact intervals, hence compact and convex, of diameter at most $\sqrt{k}$, and these sets cover A_H . If $u \in \Gamma_z$, then $|u_i - z_i| \leq \sqrt{k_i}$, so $|u_i| \geq \tau_i(z_i) := (|z_i| - \sqrt{k_i})_+$. If moreover $z \in \mathcal{Z}$, then $|z_i| \geq H_i - \sqrt{k_i}$. Let $E_z = \exp[-a \sum_i \psi_{\gamma_i}(\tau_i(z_i))] \leq 1$. By (A.2), $\sup_{\Gamma_z} \|\xi\|_{L^m}^m \leq A E_z$, and $E_z \leq E_z^{1-k/m}$. Theorem A.6, applied with $L_z = B$ and $\delta_z = \sqrt{k}$, gives

$$\mathbb{E} \sup_{A_H} |\xi|^m \leq C \sum_{z \in \mathcal{Z}} \{A E_z + B^k A^{1-k/m} E_z^{1-k/m}\} \leq C(A + B^k A^{1-k/m}) \prod_{i=1}^2 S_i,$$

where, with $a' = a(m-k)/m$,

$$S_i = \sum_{z_i \in \mathbb{Z}^{k_i} : |z_i| \geq H_i - \sqrt{k_i}} \exp[-a' \psi_{\gamma_i}(\tau_i(z_i))].$$

Fix i and suppress it from the notation, writing $k' = k_i$ and $\psi = \psi_{\gamma_i}$. The number of $z \in \mathbb{Z}^{k'}$ with $j \leq |z| < j+1$ is at most $C(1+j)^{k'-1}$, and for such z , (C.1) gives $\psi(\tau(z)) \geq \psi(j)/4 - k'$. Only integers $j \geq J := (H - \sqrt{k'} - 1)_+$ occur. Since ψ is nondecreasing and $\psi \geq \psi_1$,

$$S_i \leq C e^{a'k'} \sum_{j \geq J} (1+j)^{k'-1} e^{-a' \psi(j)/4} \leq C e^{a'k'} e^{-a' \psi(J)/8} \sum_{j \geq 0} (1+j)^{k'-1} e^{-a' \min(j^2, j)/8} \leq C e^{-a' \psi(J)/8}.$$

By (C.1) again, $\psi(J) \geq \psi(H)/4 - (\sqrt{k'} + 1)^2$, so $S_i \leq C \exp\{-a' \psi_{\gamma_i}(H_i)/32\}$. Since $a'' = a'/32$, this proves (A.4). No constant depends on γ_i or on U . $\square$

C.8 Proof of Theorem A.8

Proof. A common projection. Since $\mathcal{X}$ is bounded and $l \geq l_- > 0$, the quantity $u_l(x, y) = \|x - y\|^2 / (2l^2)$ is uniformly bounded. Set

$$T_{M,l}(x, y) = \sum_{j=0}^M \frac{(-1)^j}{j!} u_l(x, y)^j.$$

For each fixed $k \leq 3$, differentiating l^{-2j} introduces a polynomial factor of degree k in j , with bounded additional powers of l^{-1} . Therefore, for a fixed $B < \infty$,

$$\sup_{x, y \in \mathcal{X}, l \in [l_-, l_+]} |\partial_l^k (R_l(x, y) - T_{M,l}(x, y))| \leq C_k \sum_{j > M} \frac{(j+1)^k B^j}{j!} \leq e^{-M \log M + C_k M}. \quad (\text{C.2})$$

The last inequality is the case $\gamma = 1$ of the following factorial tail bound. Since $e^j \geq j^j/j!$, Stirling's lower bound

$$\log j! \geq j \log j - j, \quad j \geq 1, \quad (\text{C.3})$$

holds, together with the trivial upper bound $\log j! \leq j \log j$. Fix $k \geq 0$, $B > 0$, and $\gamma \in (0, 1]$, and put $a_j = (j+1)^k B^j / (j!)^\gamma$. Then

$$\sum_{j>M} a_j \leq e^{-\gamma M \log M + CM}, \quad M \geq 1, \quad (\text{C.4})$$

where C depends only on k , B , and γ . Indeed, $a_{j+1}/a_j = \{(j+2)/(j+1)\}^k B(j+1)^{-\gamma} \leq 2^k B(j+1)^{-\gamma}$, which is at most $1/2$ once $j+1 \geq (2^{k+1}B)^{1/\gamma}$. For M beyond this threshold, the tail is at most $2a_{M+1}$. By (C.3), $\log(M+1)! \geq (M+1) \log(M+1) - (M+1) \geq M \log M - M - 1$, so

$$\log(2a_{M+1}) \leq \log 2 + k \log(M+2) + (M+1) \log^+ B - \gamma(M \log M - M - 1) \leq -\gamma M \log M + CM.$$

For the finitely many smaller M , the tail is at most $\sum_j a_j < \infty$, which is absorbed by enlarging C .

Let P_n project onto evaluations of polynomials of total degree at most $2M$. Each column and row of the matrix $T_{M,l}$ lies in that space, as do its first three lengthscale derivatives. Thus $P_n(\partial_l^k T_{M,l})P_n = \partial_l^k T_{M,l}$ and

$$\left\| \partial_l^k R_n - P_n(\partial_l^k R_n)P_n \right\|_{\text{op}} \leq 2ne^{-M \log M + C_k M}.$$

Choose $M = \lceil c_A b_n \rceil$ with a fixed $c_A > A + 1$. Then $\log M = (1 + o(1)) \log \log n$, so $M \log M = (c_A + o(1)) \log n$, while $C_k M = o(\log n)$. Hence the last bound is $n^{1-c_A+o(1)} \leq n^{-A}$ for large n . The rank is at most $\binom{2M+p}{p} = O(b_n^p)$. Derivatives in s cause only constant factors; derivatives of S_n in t vanish. This proves (A.5).

Both exact and reduced covariance matrices are bounded below by $t_- I_n$. Put $\Omega_n = C_n - \tilde{C}_n = S_n - P_n S_n P_n$, so that $\|\Omega_n\|_{\text{op}} \leq n^{-A}$ uniformly over Θ by (A.5). The resolvent identity $C_n^{-1} - \tilde{C}_n^{-1} = -C_n^{-1} \Omega_n \tilde{C}_n^{-1}$ gives

$$\left\| C_n^{-1} - \tilde{C}_n^{-1} \right\|_{\text{op}} \leq t_-^{-2} n^{-A}.$$

Every raw covariance derivative through order three has operator norm $O(n)$ uniformly over Θ . Formulas for likelihood derivatives through order three contain finitely many traces and quadratic forms with at most three covariance derivative factors. Replacing factors one at a time, using (A.5), bounds each difference by $C n^{4-A} (1 + \|Y_n\|^2/n)$. For the zeroth-order term,

$$l_n(\theta) - \tilde{l}_n(\theta) = -\frac{1}{2} \{ \log \det C_n - \log \det \tilde{C}_n \} - \frac{1}{2} Y_n^\top (C_n^{-1} - \tilde{C}_n^{-1}) Y_n.$$

The quadratic part is at most $t_-^{-2} n^{-A} \|Y_n\|^2$. For the log determinant, let $C_n(r) = \tilde{C}_n + r \Omega_n = (1-r)\tilde{C}_n + r C_n$ for $r \in [0, 1]$. As a convex combination of two matrices bounded below by $t_- I_n$, each $C_n(r)$ satisfies $C_n(r) \succeq t_- I_n$. Jacobi's formula gives $\frac{d}{dr} \log \det C_n(r) = \text{tr}\{C_n(r)^{-1} \Omega_n\}$, and

$|\text{tr}(AB)| \leq n \|A\|_{\text{op}} \|B\|_{\text{op}}$ for $n \times n$ matrices. Hence

$$|\log \det C_n - \log \det \tilde{C}_n| \leq \int_0^1 |\text{tr}\{C_n(r)^{-1} \Omega_n\}| dr \leq t^{-1} n^{1-A}.$$

Thus all differences in (A.6) are at most $Cn^{4-A}(1 + \|Y_n\|^2/n)$ uniformly over Θ . Since $\mathbb{E}_{\theta_0} \|Y_n\|^2 = \text{tr} C_n(\theta_0) = n(s_0 + t_0)$, (A.6) follows.

Feature differentiation. Define

$$\phi_{\alpha,l}(x) = e^{-\|x\|^2/(2l^2)} \frac{x^\alpha}{l^{|\alpha|} \sqrt{\alpha!}}, \quad R_l(x, y) = \sum_{\alpha \in \mathbb{N}_0^p} \phi_{\alpha,l}(x) \phi_{\alpha,l}(y).$$

This feature expansion of the kernel is derived in (C.9) in the proof of Theorem A.9. The series and its fixed-order lengthscale derivatives converge absolutely and uniformly on bounded sets. Direct differentiation gives

$$\partial_l \phi_{\alpha,l} = l^{-1} \left[-|\alpha| \phi_{\alpha,l} + \sum_{j=1}^p \sqrt{(\alpha_j + 1)(\alpha_j + 2)} \phi_{\alpha+2e_j,l} \right]. \quad (\text{C.5})$$

For $M \geq 0$, let $I_M = \{\alpha \in \mathbb{N}_0^p : |\alpha| \leq M\}$, which has $q_M = \binom{M+p}{p}$ elements, and let $B_M(l) \in \mathbb{R}^{n \times q_M}$ be the observation matrix of the features of degree at most M :

$$[B_M(l)]_{i\alpha} = \phi_{\alpha,l}(x_{n,i}), \quad 1 \leq i \leq n, \quad \alpha \in I_M.$$

Thus the column of $B_M(l)$ indexed by α is the feature $\phi_{\alpha,l}$ evaluated at the n sampling points, and

$$[B_M(l) B_M(l)^\top]_{ij} = \sum_{\alpha \in I_M} \phi_{\alpha,l}(x_{n,i}) \phi_{\alpha,l}(x_{n,j})$$

is the degree- M truncation of $R_n(l)$. For $M \leq M'$, let $E_{M,M'} \in \mathbb{R}^{q_{M'} \times q_M}$ be the coordinate embedding, $E_{M,M'} e_\alpha = e_\alpha$ for $\alpha \in I_M$. It has orthonormal columns, so $\|E_{M,M'}\|_{\text{op}} = 1$, and $B_M = B_{M'} E_{M,M'}$ because the columns of B_M are among those of $B_{M'}$.

Applying (C.5) column by column gives $\partial_l B_M(l) = B_{M+2}(l) V_M(l)$, where $V_M(l) \in \mathbb{R}^{q_{M+2} \times q_M}$ has columns

$$V_M(l) e_\alpha = l^{-1} \left[-|\alpha| e_\alpha + \sum_{j=1}^p \sqrt{(\alpha_j + 1)(\alpha_j + 2)} e_{\alpha+2e_j} \right], \quad \alpha \in I_M.$$

In particular, $V_M(l) = l^{-1} V_M^\circ$ with V_M° independent of l . Since $\sqrt{(\alpha_j + 1)(\alpha_j + 2)} \leq |\alpha| + 2 \leq M + 2$, every column has norm at most $(p+1)(M+2)/l$. A bound on column norms does not by itself bound an operator norm, but it does once the matrix is split into pieces with orthogonal columns. Write

$$l V_M(l) = -E_{M,M+2} N_M + \sum_{j=1}^p W_{M,j}, \quad N_M e_\alpha = |\alpha| e_\alpha, \quad W_{M,j} e_\alpha = \sqrt{(\alpha_j + 1)(\alpha_j + 2)} e_{\alpha+2e_j}.$$

In each of these $p+1$ matrices, distinct columns are multiples of distinct coordinate vectors, because the maps $\alpha \mapsto \alpha$ and $\alpha \mapsto \alpha + 2e_j$ are injective. The columns are therefore orthogonal. For a matrix W with orthogonal columns, $\|Wx\|^2 = \sum_{\alpha} x_{\alpha}^2 \|We_{\alpha}\|^2$, so its operator norm equals its largest column norm, here at most $M+2$. Hence

$$\|V_M(l)\|_{\text{op}} \leq l^{-1}(p+1)(M+2) \leq C(M+1). \quad (\text{C.6})$$

Equivalently, each row of $V_M(l)$ also has at most $p+1$ nonzero entries, so the Schur test applies.

We now show that, for $k \leq 3$,

$$\partial_l^k \{B_M(l)B_M(l)^{\top}\} = B_{M+2k}(l)H_{M,k}(l)B_{M+2k}(l)^{\top}, \quad \|H_{M,k}(l)\|_{\text{op}} \leq C_k(M+1)^k, \quad (\text{C.7})$$

where $H_{M,k}(l) = l^{-k}H_{M,k}^{\circ}$ is a symmetric $q_{M+2k} \times q_{M+2k}$ matrix and $H_{M,k}^{\circ}$ does not depend on l . For $k=0$, take $H_{M,0} = I_{q_M}$. Suppose that (C.7) holds for some k , and abbreviate $M_k = M+2k$ and $E = E_{M_k, M_k+2}$. Since $B_{M_k} = B_{M_k+2}E$ and $\partial_l B_{M_k} = B_{M_k+2}V_{M_k}$, the product rule gives

$$\partial_l^{k+1} \{B_M B_M^{\top}\} = B_{M_k+2} \{V_{M_k} H_{M,k} E^{\top} + E(\partial_l H_{M,k}) E^{\top} + E H_{M,k} V_{M_k}^{\top}\} B_{M_k+2}^{\top}.$$

The matrix in braces is $H_{M,k+1}$. It is symmetric, and it has the form $l^{-(k+1)}H_{M,k+1}^{\circ}$ because $V_{M_k} = l^{-1}V_{M_k}^{\circ}$ and $\partial_l H_{M,k} = -kl^{-1}H_{M,k}$. Since $\|E\|_{\text{op}} = 1$, by (C.6),

$$\|H_{M,k+1}\|_{\text{op}} \leq \{2C(M+2k+1) + kl^{-1}\} \|H_{M,k}\|_{\text{op}} \leq C_{k+1}(M+1)^{k+1}.$$

This is how all smaller feature spaces are embedded in the common space through degree $M+2k$: the embeddings E carry every intermediate term into $\mathbb{R}^{q_{M+2k}}$, so that the single feature matrix B_{M+2k} appears on both sides.

Positive feature truncation gives $sB_{M+2k}B_{M+2k}^{\top} \preceq C_n$ and therefore

$$\left\| \sqrt{s} C_n^{-1/2} B_{M+2k} \right\|_{\text{op}} \leq 1. \quad (\text{C.8})$$

The whitened truncated derivative consequently has operator norm $O(M^k)$ and rank $O(M^p)$, so its Frobenius norm is $O(M^{p/2+k})$.

To control the derivative tail, collect terms with $|\alpha| = j$. Absolute summation of the monomials bounds these terms by $B^j/j!$ on the fixed bounded set, and a k th lengthscale derivative adds at most $C_k(j+1)^k$. Hence (C.4) with $\gamma = 1$ gives the tail bound in (C.2). For M a sufficiently large multiple of b_n , its operator norm is an arbitrarily small power of n^{-1} , and its Frobenius norm is at most $\sqrt{n}$ times that bound. Whitening multiplies these errors by at most t_-^{-1} . This proves (A.7) and (A.8) for lengthscale derivatives. One s -derivative removes the factor s ; two s -derivatives give zero.

For a compression $A_{\theta} = U^{\top} C_n U$, the derivatives are $U^{\top} (\partial_l^k C_n) U$, so by (C.7) the truncated part becomes $s(U^{\top} B_{M+2k}) H_{M,k} (U^{\top} B_{M+2k})^{\top}$. Since $sB_{M+2k}B_{M+2k}^{\top} \preceq C_n$, also $sU^{\top} B_{M+2k}B_{M+2k}^{\top} U \preceq U^{\top} C_n U = A_{\theta}$, which gives the analogue $\left\| \sqrt{s} A_{\theta}^{-1/2} U^{\top} B_{M+2k} \right\|_{\text{op}} \leq 1$ of (C.8). The rank is still at most q_{M+2k} . The compressed tail $U^{\top} \{\partial_l^k (R_n - B_M B_M^{\top})\} U$ has operator norm at most that of the uncompressed tail and Frobenius norm at most $\sqrt{r} \leq \sqrt{n}$ times it, and $A_{\theta} \succeq t_- I_r$. Hence every

estimate above holds verbatim for A_θ , with the same constants.

Finally,

$$\partial_{h_s}^a \partial_{h_l}^c = b_n^{-(a+c)p/2-c} \partial_s^a \partial_l^c.$$

Combining this identity with the preceding estimates gives $O(\epsilon_n^{a+c})$ in operator norm and $O(\epsilon_n^{a+c-1})$ in Frobenius norm. Since $C_{n,t} = I_n$, its normalized relative derivative is $n^{-1/2} C_n^{-1}$, which has operator norm at most $t_-^{-1} n^{-1/2}$ and Frobenius norm at most t_-^{-1} . All higher covariance derivatives involving t vanish. $\square$

C.9 Proof of Theorem A.9

Proof. Kernel expansion. For $x, y \in \mathbb{R}^p$, $\|x - y\|^2 = \|x\|^2 + \|y\|^2 - 2\langle x, y \rangle$. By the multinomial theorem, $\langle x, y \rangle^j = \sum_{|\alpha|=j} \frac{j!}{\alpha!} x^\alpha y^\alpha$, and therefore

$$\exp\left(\frac{\langle x, y \rangle}{l^2}\right) = \sum_{j=0}^{\infty} \frac{\langle x, y \rangle^j}{l^{2j} j!} = \sum_{j=0}^{\infty} \sum_{|\alpha|=j} \frac{x^\alpha y^\alpha}{l^{2j} \alpha!} = \sum_{\alpha \in \mathbb{N}_0^p} \frac{x^\alpha y^\alpha}{l^{2|\alpha|} \alpha!}.$$

The rearrangement is justified because the same series with every term replaced by its absolute value sums to $\exp(\sum_i |x_i y_i|/l^2) < \infty$. Hence

$$K_\theta(x, y) = s e^{-\|x\|^2/(2l^2)} e^{-\|y\|^2/(2l^2)} e^{\langle x, y \rangle/l^2} = s \sum_{\alpha \in \mathbb{N}_0^p} \phi_{\alpha,l}(x) \phi_{\alpha,l}(y), \quad (\text{C.9})$$

where

$$\phi_{\alpha,l}(x) = e^{-\|x\|^2/(2l^2)} \frac{x^\alpha}{l^{|\alpha|} \sqrt{\alpha!}}.$$

In particular, $\sum_\alpha \phi_{\alpha,l}(x)^2 = K_\theta(x, x)/s = 1$.

An analytic representation. Let ξ_α , $\alpha \in \mathbb{N}_0^p$, be independent standard normal variables, and define

$$F(x) = \sqrt{s} \sum_{\alpha \in \mathbb{N}_0^p} \xi_\alpha \phi_{\alpha,l}(x) = \sqrt{s} e^{-\|x\|^2/(2l^2)} \sum_{\alpha \in \mathbb{N}_0^p} \xi_\alpha \frac{x^\alpha}{l^{|\alpha|} \sqrt{\alpha!}}. \quad (\text{C.10})$$

For $\rho > 0$ and $x \in [-\rho, \rho]^p$, $|\phi_{\alpha,l}(x)| \leq (\rho/l_-)^{|\alpha|}/\sqrt{\alpha!}$, so

$$\mathbb{E} \sum_\alpha |\xi_\alpha| \sup_{x \in [-\rho, \rho]^p} |\phi_{\alpha,l}(x)| \leq \sum_\alpha \frac{(\rho/l_-)^{|\alpha|}}{\sqrt{\alpha!}} = \left(\sum_{a=0}^{\infty} \frac{(\rho/l_-)^a}{\sqrt{a!}} \right)^p < \infty.$$

Hence, almost surely, the series in (C.10) converges absolutely and uniformly on every box, and the power series in it defines an entire function. For fixed x, y , the partial sums also converge in L^2 because $\sum_\alpha \phi_{\alpha,l}(x)^2 = 1$, and independence of the ξ_α together with (C.9) gives

$$\text{Cov}\{F(x), F(y)\} = s \sum_\alpha \phi_{\alpha,l}(x) \phi_{\alpha,l}(y) = K_\theta(x, y).$$

Thus (C.10) is an analytic version of the centered Gaussian process with covariance (1.1).

Taylor coefficients and their tails. Let

$$H(x) = e^{\|x\|^2/(2l^2)} F(x) = \sqrt{s} \sum_{\gamma \in \mathbb{N}_0^p} \xi_\gamma \frac{x^\gamma}{l^{|\gamma|} \sqrt{\gamma!}}, \quad a_l = -\frac{1}{2l^2},$$

so that $F(x) = e^{a_l \|x\|^2} H(x)$. Since $e^{a_l \|x\|^2} = \prod_j e^{a_l x_j^2} = \sum_{r \in \mathbb{N}_0^p} a_l^{|r|} x^{2r} / r!$, the Cauchy product of these absolutely convergent series gives

$$F(x) = \sqrt{s} \sum_{r \in \mathbb{N}_0^p} \sum_{\gamma \in \mathbb{N}_0^p} \frac{a_l^{|r|}}{r! l^{|\gamma|} \sqrt{\gamma!}} \xi_\gamma x^{2r+\gamma}.$$

Collecting the terms with $2r + \gamma = \alpha$, the Taylor coefficients $f_\alpha = [x^\alpha]F(x)$ are

$$f_\alpha = \sqrt{s} \sum_{2r \leq \alpha} \frac{a_l^{|r|}}{r! l^{|\alpha|-2|r|} \sqrt{(\alpha-2r)!}} \xi_{\alpha-2r}. \quad (\text{C.11})$$

The summands involve distinct independent normal variables. For each coordinate, the terms in

$$\sum_{r=0}^{\lfloor q/2 \rfloor} \frac{q!}{r! r! (q-2r)!} |a_l|^{2r} (l^{-2})^{q-2r}$$

form a subset of the nonnegative terms in the multinomial expansion of $(2|a_l| + l^{-2})^q = (2l^{-2})^q$. Therefore

$$\alpha! \text{Var}_\theta(f_\alpha) \leq s(2l^{-2})^{|\alpha|}, \quad \|f_\alpha\|_{L^2(P_\theta)} \leq \frac{C^{|\alpha|+1}}{\sqrt{\alpha!}}.$$

Let $T_K F(x) = \sum_{|\alpha| \leq K} f_\alpha x^\alpha$, and write $|x|^\alpha = \prod_i |x_i|^{\alpha_i}$. The number of $\alpha \in \mathbb{N}_0^p$ with $|\alpha| = j$ is $\binom{j+p-1}{p-1}$. Hence the Cauchy–Schwarz inequality over these indices and the multinomial theorem give

$$\begin{aligned} \left(\sum_{|\alpha|=j} \frac{|x|^\alpha}{\sqrt{\alpha!}} \right)^2 &\leq \binom{j+p-1}{p-1} \sum_{|\alpha|=j} \frac{x^{2\alpha}}{\alpha!} = \binom{j+p-1}{p-1} \frac{1}{j!} \sum_{|\alpha|=j} \frac{j!}{\alpha!} \prod_{i=1}^p (x_i^2)^{\alpha_i} \\ &= \binom{j+p-1}{p-1} \frac{\|x\|^{2j}}{j!}. \end{aligned} \quad (\text{C.12})$$

Since $|F(x) - T_K F(x)| \leq \sum_{j>K} \sum_{|\alpha|=j} |f_\alpha| |x|^\alpha$, Minkowski's inequality, the bound on $\|f_\alpha\|_{L^2(P_\theta)}$, (C.12), and $\binom{j+p-1}{p-1} \leq (j+1)^{p-1}$ give, for x in a fixed bounded set,

$$\|F(x) - T_K F(x)\|_{L^2(P_\theta)} \leq C \sum_{j>K} \frac{(j+1)^{(p-1)/2} (C\|x\|)^j}{(j!)^{1/2}}.$$

By (C.4) with $\gamma = 1/2$, which rests on Stirling's bound $\log j! \geq j \log j - j$, the right side is at most $e^{-(K \log K)/2 + CK}$. Squaring yields

$$\sup_{\theta \in \Theta} \sup_{x \in \mathcal{X}} \mathbb{E}_\theta |F(x) - T_K F(x)|^2 \leq e^{-K \log K + CK}. \quad (\text{C.13})$$

Weights and polynomial Gram matrix. Partition Q into nearest-observation cells $V_{n,i}$, with a fixed tie-breaking rule. Define $w_{n,i} = |V_{n,i}|/|Q|$. Then

$$\sum_i w_{n,i} = 1, \quad 0 \leq w_{n,i} \leq Ch_n(Q)^p \leq Cn^{-\beta}. \quad (\text{C.14})$$

Indeed, $V_{n,i} \subseteq B(x_{n,i}, h_n(Q))$. Zero weights and repeated locations are allowed.

Use the product Legendre basis on $[0, L]^p$, restricted to total degree at most K , and normalized for uniform probability measure on Q . Write its elements as ψ_ν , $|\nu| \leq K$, and set

$$\Phi_{i,\nu} = \psi_\nu(x_{n,i}), \quad W = \text{diag}(w_{n,1}, \dots, w_{n,n}), \quad M_K = \Phi^\top W \Phi.$$

The shifted Legendre polynomials have the explicit form

$$P_j(2x - 1) = \sum_{a=0}^j (-1)^{j-a} \binom{j}{a} \binom{j+a}{a} x^a. \quad (\text{C.15})$$

This follows from Rodrigues' formula $P_j(t) = (2^j j!)^{-1} \frac{d^j}{dt^j} (t^2 - 1)^j$ [DLMF, §18.5]. With $t = 2x - 1$, we have $t^2 - 1 = 4(x^2 - x)$ and $\frac{d}{dt} = \frac{1}{2} \frac{d}{dx}$, so

$$P_j(2x - 1) = \frac{1}{j!} \frac{d^j}{dx^j} \{x^j (x - 1)^j\} = \frac{1}{j!} \sum_{a=0}^j \binom{j}{a} (-1)^{j-a} \frac{d^j}{dx^j} x^{j+a} = \sum_{a=0}^j (-1)^{j-a} \binom{j}{a} \frac{(j+a)!}{j! a!} x^a,$$

which is (C.15). This identity implies bounds $e^{CK} K^C$ for these basis polynomials and their first derivatives on any fixed enlarged box. The same bounds hold for products of two basis polynomials, after changing C . Comparing the integral over each $V_{n,i}$ with its value at $x_{n,i}$ gives

$$|[M_K - I]_{\mu\nu}| \leq h_n(Q) \sup |\nabla(\psi_\mu \psi_\nu)|.$$

The supremum is taken on a fixed box containing all relevant line segments. This remains valid when the assigned node is outside Q . The number of basis elements is polynomial in K , and hence

$$\|M_K - I\|_{\text{op}} \leq h_n(Q) e^{CK} K^C. \quad (\text{C.16})$$

For $K = K_n$, this is $n^{-\kappa+o(1)}$, so M_{K_n} and its inverse are uniformly bounded for all sufficiently large n .

Weighted regression and coefficient extraction. Fit the polynomial with Legendre coefficient vector

$$\hat{c} = M_{K_n}^{-1} \Phi^\top W Y_n.$$

The basis has $q_K = \binom{K+p}{p}$ elements. For $|\alpha| \leq K$, let $L_{\alpha,K} \in \mathbb{R}^{1 \times q_K}$ be the row vector that maps the Legendre coefficient vector of a polynomial to its normalized monomial coefficient of index α :

$$L_{\alpha,K} c = \sqrt{\alpha!} [x^\alpha] \sum_{|\nu| \leq K} c_\nu \psi_\nu(x), \quad c \in \mathbb{R}^{q_K}.$$

By (C.15), individual unnormalized Legendre coefficients are at most 8^j on $[0, 1]$. Rescaling to $[0, L]$,

taking products, and allowing polynomial dimension factors gives

$$\|L_{\alpha,K}\|^2 \leq e^{m \log m + CK}, \quad |\alpha| \leq m \leq K. \quad (\text{C.17})$$

Here $\alpha! \leq |\alpha|! \leq m! \leq e^{m \log m}$ was used, by the trivial upper counterpart of (C.3). Define $\hat{\zeta}_\alpha = L_{\alpha,K_n} \hat{c}$ for $|\alpha| \leq m_n$. This is a fixed linear function of Y_n .

Write $Y_n = F_n + \bar{\varepsilon}$, where $F_n = (F(x_{n,1}), \dots, F(x_{n,n}))^\top$ and $\bar{\varepsilon} = (\varepsilon_{n,1}, \dots, \varepsilon_{n,n})^\top$. Weighted regression reproduces $T_K F$ exactly. If $r_i = F(x_{n,i}) - T_K F(x_{n,i})$, the residual coefficient error obeys

$$|L_{\alpha,K} M_K^{-1} \Phi^\top W r|^2 \leq (L_{\alpha,K} M_K^{-1} L_{\alpha,K}^\top) r^\top W r.$$

For the noise error, $\text{Cov}_\theta(\bar{\varepsilon}) = t I_n$ and $W^2 \preceq (\max_i w_{n,i}) W$ imply

$$\text{Var}_\theta(L_{\alpha,K} M_K^{-1} \Phi^\top W \bar{\varepsilon}) \leq t_+ \max_i w_{n,i} L_{\alpha,K} M_K^{-1} L_{\alpha,K}^\top.$$

The field residual and noise are independent. Using (C.13), (C.14), (C.16), and (C.17), we obtain

$$\sup_\theta \mathbb{E}_\theta(\hat{\zeta}_\alpha - \zeta_\alpha)^2 \leq e^{m_n \log m_n + CK_n} \{n^{-\beta} + e^{-K_n \log K_n + CK_n}\}. \quad (\text{C.18})$$

Since $\log m_n$ and $\log K_n$ are both $(1 + o(1)) \log \log n$, we have $m_n \log m_n = (\beta/4 + o(1)) \log n$ and $K_n \log K_n = (\beta/2 + o(1)) \log n$, while $CK_n = o(\log n)$. Hence the two contributions in (C.18) are respectively $n^{-3\beta/4 + o(1)}$ and $n^{-\beta/4 + o(1)}$. This proves (A.11).

Conditioning of the exact coefficient covariance. Let e_α denote the coordinate vector indexed by α in the space $|\alpha| \leq m$. Define

$$D_{m,l} e_\alpha = l^{-|\alpha|} e_\alpha$$

and a truncated raising matrix

$$Q_m e_\alpha = \sum_{\substack{1 \leq j \leq p \\ |\alpha|+2 \leq m}} \sqrt{(\alpha_j + 1)(\alpha_j + 2)} e_{\alpha+2e_j}.$$

Since Q_m raises the total degree by two, $Q_m^r = 0$ for $2r > m$. As for V_M in the proof of Theorem A.8, Q_m splits into p matrices with orthogonal columns of norm at most m , so $\|Q_m\|_{\text{op}} \leq pm$.

We now derive (C.19) below in four steps. For a function f given by a power series $f(x) = \sum_\alpha ([x^\alpha] f) x^\alpha$ that converges absolutely on $\mathbb{R}^p$, its normalized coefficients are

$$[\mathcal{N}f]_\alpha = \sqrt{\alpha!} [x^\alpha] f, \quad \text{so that} \quad f(x) = \sum_{\alpha \in \mathbb{N}_0^p} [\mathcal{N}f]_\alpha \frac{x^\alpha}{\sqrt{\alpha!}}.$$

Write $\mathcal{N}_m f = ([\mathcal{N}f]_\alpha : |\alpha| \leq m) \in \mathbb{R}^{q_m}$. By (A.10), $\zeta^{(m)} = \mathcal{N}_m F$.

Step 1: multiplication by $\|x\|^2$. Since $\|x\|^2 f(x) = \sum_{j=1}^p \sum_\alpha ([x^\alpha] f) x^{\alpha+2e_j}$, for every β ,

$$[x^\beta](\|x\|^2 f) = \sum_{j: \beta_j \geq 2} [x^{\beta-2e_j}] f.$$

Multiply by $\sqrt{\beta!}$ and write $\alpha = \beta - 2e_j$. Then $\beta!/\alpha! = (\alpha_j + 1)(\alpha_j + 2)$, so

$$[\mathcal{N}(\|x\|^2 f)]_\beta = \sum_{j: \beta_j \geq 2} \sqrt{(\alpha_j + 1)(\alpha_j + 2)} [\mathcal{N}f]_{\beta - 2e_j}.$$

For $|\beta| \leq m$, the right side involves only indices of degree at most $m - 2$, and it is exactly the β th entry of $Q_m \mathcal{N}_m f$. Hence $\mathcal{N}_m(\|x\|^2 f) = Q_m \mathcal{N}_m f$.

Step 2: multiplication by $e^{a\|x\|^2}$. By induction, $\mathcal{N}_m(\|x\|^{2r} f) = Q_m^r \mathcal{N}_m f$ for every $r \geq 0$. For $a \in \mathbb{R}$, $e^{a\|x\|^2} f = \sum_{r \geq 0} a^r \|x\|^{2r} f / r!$, and all these power series converge absolutely, so coefficients may be extracted term by term. Since $Q_m^r = 0$ for $2r > m$,

$$\mathcal{N}_m(e^{a\|x\|^2} f) = \sum_{r \leq m/2} \frac{a^r}{r!} Q_m^r \mathcal{N}_m f = e^{aQ_m} \mathcal{N}_m f.$$

Step 3: the coefficients of H . By the definition of H , $[x^\gamma]H = \sqrt{s} \xi_\gamma / (l^{|\gamma|} \sqrt{\gamma!})$, so $[\mathcal{N}H]_\gamma = \sqrt{s} l^{-|\gamma|} \xi_\gamma$, that is, $\mathcal{N}_m H = \sqrt{s} D_{m,l} \xi^{(m)}$.

Step 4: the coefficients of F . Since $F = e^{a_l \|x\|^2} H$ with $a_l = -1/(2l^2)$, Steps 2 and 3 give

$$\zeta^{(m)} = \mathcal{N}_m F = e^{a_l Q_m} \mathcal{N}_m H = \sqrt{s} e^{-Q_m/(2l^2)} D_{m,l} \xi^{(m)}. \quad (\text{C.19})$$

Entrywise, this is (C.11) multiplied by $\sqrt{\alpha!}$. The transform and its inverse have operator norms at most e^{Cm} : $\|e^{\pm Q_m/(2l^2)}\|_{\text{op}} \leq e^{pm/(2l^2)} \leq e^{Cm}$ and $\|D_{m,l}^{\pm 1}\|_{\text{op}} \leq e^{Cm}$. Constants can be absorbed into the exponent for $m \geq 1$. This proves (A.12).

Relative covariance error and affinity. Let $\delta_n = \widehat{\zeta}^{(m_n)} - \zeta^{(m_n)}$, a centered random vector. Since $q_{m_n} \leq (m_n + 1)^p = n^{o(1)}$ and $e^{Cm_n} = n^{o(1)}$, (A.11) and (A.12) give

$$\begin{aligned} \sup_{\theta} \mathbb{E}_{\theta} \|\delta_n\|^2 &= \sup_{\theta} \sum_{|\alpha| \leq m_n} \mathbb{E}_{\theta} (\widehat{\zeta}_{\alpha} - \zeta_{\alpha})^2 \leq n^{-\beta/4+o(1)}, \\ \sup_{\theta} \mathbb{E}_{\theta} \|\zeta^{(m_n)}\|^2 &= \sup_{\theta} \text{tr} \Sigma_{m_n, \theta} \leq q_{m_n} e^{Cm_n} = n^{o(1)}. \end{aligned}$$

Since $\widehat{\zeta}^{(m_n)} = \zeta^{(m_n)} + \delta_n$,

$$\widehat{\Sigma}_{n, \theta} - \Sigma_{m_n, \theta} = \mathbb{E}_{\theta} [\zeta^{(m_n)} \delta_n^{\top}] + \mathbb{E}_{\theta} [\delta_n \zeta^{(m_n)\top}] + \mathbb{E}_{\theta} [\delta_n \delta_n^{\top}].$$

For unit vectors u, v , the Cauchy–Schwarz inequality gives

$$|u^{\top} \mathbb{E}_{\theta} [\zeta^{(m_n)} \delta_n^{\top}] v| \leq \{\mathbb{E}_{\theta} (u^{\top} \zeta^{(m_n)})^2 \mathbb{E}_{\theta} (v^{\top} \delta_n)^2\}^{1/2} \leq \{\mathbb{E}_{\theta} \|\zeta^{(m_n)}\|^2 \mathbb{E}_{\theta} \|\delta_n\|^2\}^{1/2},$$

and similarly for the transposed term, while $\|\mathbb{E}_{\theta} [\delta_n \delta_n^{\top}]\|_{\text{op}} = \sup_{\|u\|=1} \mathbb{E}_{\theta} (u^{\top} \delta_n)^2 \leq \mathbb{E}_{\theta} \|\delta_n\|^2$. Therefore

$$\begin{aligned} \|\widehat{\Sigma}_{n, \theta} - \Sigma_{m_n, \theta}\|_{\text{op}} &\leq \mathbb{E}_{\theta} \|\delta_n\|^2 + 2\sqrt{\mathbb{E}_{\theta} \|\delta_n\|^2 \mathbb{E}_{\theta} \|\zeta^{(m_n)}\|^2} \\ &\leq n^{-\beta/4+o(1)} + n^{-\beta/8+o(1)} = n^{-\beta/8+o(1)}. \end{aligned}$$

Finally, $\left\| \Sigma^{-1/2} A \Sigma^{-1/2} \right\|_{\text{op}} \leq \left\| \Sigma^{-1} \right\|_{\text{op}} \|A\|_{\text{op}}$ and $\left\| \Sigma_{m_n, \theta}^{-1} \right\|_{\text{op}} \leq e^{Cm_n} = n^{o(1)}$ by (A.12), which proves (A.13).

Let $\eta_n = n^{-\beta/8+o(1)}$ be a common bound in that equation. For large n ,

$$(1 - \eta_n) \Sigma_{m_n, \theta} \preceq \widehat{\Sigma}_{n, \theta} \preceq (1 + \eta_n) \Sigma_{m_n, \theta}.$$

The same inequalities hold for sums of two such covariance matrices. Using (A.1) and the corresponding log determinant bounds, the absolute difference of log affinities is at most $Cq_{m_n}\eta_n = o(1)$. This proves (A.14). The comparison is multiplicative for affinity, which will be essential when the affinity itself is exponentially small. $\square$

C.10 Proof of Theorem A.10

Proof. Triangular structure. Order the indices $\alpha \in I_m = \{|\alpha| \leq m\}$ by total degree, with an arbitrary order within each degree. Write $\alpha(1), \dots, \alpha(q)$ for the ordered indices, where $q = q_m$, write ζ_k and ξ_k for the coordinates of $\zeta^{(m)}$ and $\xi^{(m)}$ with index $\alpha(k)$, and put $\zeta_{<k} = (\zeta_1, \dots, \zeta_{k-1})$ and $\xi_{<k} = (\xi_1, \dots, \xi_{k-1})$. By (C.19), $\zeta^{(m)} = \sqrt{s} \mathcal{T}_\theta \xi^{(m)}$ with $\mathcal{T}_\theta = e^{-Q_m/(2l^2)} D_{m,l}$. Since Q_m raises total degree by two, every nonzero entry of Q_m^r , $r \geq 1$, lies in a row of strictly higher degree than its column. Hence $e^{-Q_m/(2l^2)} = \sum_{r \leq m/2} (-2l^2)^{-r} Q_m^r / r!$ is unit lower triangular in this order, and $\mathcal{T}_\theta$ is lower triangular with diagonal entries $l^{-|\alpha(k)|}$:

$$\zeta_k = \sqrt{s} l^{-|\alpha(k)|} \xi_k + \sqrt{s} \sum_{i < k} (\mathcal{T}_\theta)_{ki} \xi_i, \quad k = 1, \dots, q.$$

The leading $(k-1) \times (k-1)$ block of $\mathcal{T}_\theta$ is invertible, so $\zeta_{<k}$ and $\xi_{<k}$ are invertible linear functions of each other, and ξ_k is independent of $\xi_{<k}$. Therefore, under P_θ , the conditional law of ζ_k given $\zeta_{<k}$ is $N(\mu_{\theta,k}(\zeta_{<k}), \omega_{\theta,k})$, where $\mu_{\theta,k}$ is a linear function of $\zeta_{<k}$ and

$$\omega_{\theta,k} = \text{Var}_\theta(\zeta_k \mid \zeta_{<k}) = s l^{-2|\alpha(k)|}. \quad (\text{C.20})$$

The conditional variance does not depend on $\zeta_{<k}$.

Univariate affinity. For normal densities with means μ, μ' and variances ω, ω' , the exponent of the product of the two square roots satisfies

$$-\frac{(x - \mu)^2}{4\omega} - \frac{(x - \mu')^2}{4\omega'} = -\frac{(x - \bar{\mu})^2}{2w} - \frac{(\mu - \mu')^2}{4(\omega + \omega')}, \quad w = \frac{2\omega\omega'}{\omega + \omega'},$$

for a suitable $\bar{\mu}$. Integrating in x gives

$$\text{Aff}\{N(\mu, \omega), N(\mu', \omega')\} = \frac{\sqrt{2\pi w}}{\sqrt{2\pi} (\omega\omega')^{1/4}} \exp\left\{-\frac{(\mu - \mu')^2}{4(\omega + \omega')}\right\} = \left(\frac{2\sqrt{\omega\omega'}}{\omega + \omega'}\right)^{1/2} \exp\left\{-\frac{(\mu - \mu')^2}{4(\omega + \omega')}\right\}.$$

Since $(\omega + \omega')/(2\sqrt{\omega\omega'}) = \frac{1}{2}(\sqrt{\omega/\omega'} + \sqrt{\omega'/\omega}) = \cosh\{\frac{1}{2}\log(\omega/\omega')\}$, it follows that

$$\text{Aff}\{N(\mu, \omega), N(\mu', \omega')\} \leq \cosh\left\{\frac{1}{2}\log \frac{\omega}{\omega'}\right\}^{-1/2} \quad \text{for all } \mu, \mu'. \quad (\text{C.21})$$

Successive integration. Let π_θ denote the density of $\zeta^{(m)}$ under P_θ , and $\pi_\theta(\zeta_k \mid \zeta_{<k})$ its conditional densities, so that $\pi_\theta(\zeta) = \prod_{k=1}^q \pi_\theta(\zeta_k \mid \zeta_{<k})$. For $0 \leq k \leq q$ put

$$\mathcal{A}_k = \int_{\mathbb{R}^k} \prod_{i=1}^k \sqrt{\pi_\theta(\zeta_i \mid \zeta_{<i}) \pi_{\theta_0}(\zeta_i \mid \zeta_{<i})} d\zeta_1 \cdots d\zeta_k,$$

so that $\mathcal{A}_0 = 1$ and $\mathcal{A}_q = \text{Aff}(\Sigma_{m,\theta}, \Sigma_{m,\theta_0})$. The integrand is nonnegative, so by Tonelli's theorem we may integrate over ζ_k first, with $\zeta_{<k}$ fixed. By (C.21),

$$\int_{\mathbb{R}} \sqrt{\pi_\theta(\zeta_k \mid \zeta_{<k}) \pi_{\theta_0}(\zeta_k \mid \zeta_{<k})} d\zeta_k \leq \rho_k := \cosh\left\{\frac{1}{2} \log \frac{\omega_{\theta,k}}{\omega_{\theta_0,k}}\right\}^{-1/2}.$$

This bound is the same for every value of $\zeta_{<k}$, because the conditional variances (C.20) do not depend on $\zeta_{<k}$. The conditional means, which do depend on $\zeta_{<k}$ and on the parameter, enter only through the exponential factor of the univariate affinity, which is at most one. Hence $\mathcal{A}_k \leq \rho_k \mathcal{A}_{k-1}$, and by induction $\text{Aff}(\Sigma_{m,\theta}, \Sigma_{m,\theta_0}) \leq \prod_{k=1}^q \rho_k$. Finally, with $u = \log(s/s_0)$ and $v = \log(l/l_0)$, (C.20) gives $\log(\omega_{\theta,k}/\omega_{\theta_0,k}) = u - 2|\alpha(k)|v$. Therefore

$$\text{Aff}(\Sigma_{m,\theta}, \Sigma_{m,\theta_0}) \leq \prod_{k=1}^q \cosh(u/2 - |\alpha(k)|v)^{-1/2} = \exp\left\{-\frac{1}{2} \sum_{|\alpha| \leq m} \log \cosh(u/2 - |\alpha|v)\right\}. \quad (\text{C.22})$$

A fixed fraction of separated degrees. To bound the sum, select a multiindex α uniformly from I_m , let $J = |\alpha|$, and put $Z = J/m$ for $m \geq 1$. The number of indices of degree j is $\binom{j+p-1}{p-1}$. Binomial summation gives

$$\mathbb{E}Z = \frac{p}{p+1}, \quad \mathbb{E}Z^2 = \frac{p}{p+2} + \frac{p}{(p+1)(p+2)m}.$$

The matrix $\mathbb{M} = \begin{pmatrix} 1 & -\mathbb{E}Z \\ -\mathbb{E}Z & \mathbb{E}Z^2 \end{pmatrix}$ has $\det \mathbb{M} = \text{Var } Z \geq \frac{p}{p+2} - \frac{p^2}{(p+1)^2} = \frac{p}{(p+2)(p+1)^2}$ and $\text{tr } \mathbb{M} \leq 2$, so

$$\lambda_{\min}(\mathbb{M}) \geq \frac{\det \mathbb{M}}{\text{tr } \mathbb{M}} \geq c_p := \frac{p}{2(p+2)(p+1)^2}.$$

Put $a = u/2$, $b = mv$, and $L_0 = \sqrt{a^2 + b^2}$, so that $a - bZ = u/2 - |\alpha|v$. Then

$$\mathbb{E}(a - bZ)^2 = a^2 - 2ab\mathbb{E}Z + b^2\mathbb{E}Z^2 = (a, b) \mathbb{M} (a, b)^\top \geq c_p L_0^2,$$

and, since $0 \leq Z \leq 1$, $|a - bZ| \leq |a| + |b| \leq \sqrt{2}L_0$. If $L_0 = 0$, then $u = v = 0$ and (C.23) below is trivial. Otherwise, Theorem A.3 with $X = a - bZ$, $\sigma^2 = c_p L_0^2$, $M = \sqrt{2}L_0$, and $\lambda = 1/\sqrt{2}$ shows that the set

$$\mathcal{G} = \{\alpha \in I_m : |u/2 - |\alpha|v| \geq c_0 L_0\}, \quad c_0 = (c_p/2)^{1/2},$$

contains at least $(c_p/4)q_m \geq (c_p/4)m^p/p!$ indices.

Lower bound for the sum. For all real z ,

$$\log \cosh(1) \min(z^2, |z|) \leq \log \cosh z \leq \min(z^2/2, |z|).$$

The upper bound is elementary. For the lower bound, the function $g(z) = 2 \log \cosh z - z \tanh z$ satisfies $g(0) = 0$ and $g'(z) = \operatorname{sech}^2 z (\frac{1}{2} \sinh 2z - z) \geq 0$ for $z \geq 0$, so $(\log \cosh z)/z^2$ is nonincreasing on $(0, \infty)$; this gives the bound for $|z| \leq 1$. For $|z| \geq 1$, it follows from convexity of $\log \cosh$ and $\log \cosh 0 = 0$. For $\alpha \in \mathcal{G}$, since $\log \cosh$ is increasing in $|z|$ and $c_0 \leq 1$,

$$\log \cosh(u/2 - |\alpha|v) \geq \log \cosh(c_0 L_0) \geq \log \cosh(1) c_0^2 \min(L_0^2, L_0).$$

Since $L_0^2 \geq (u^2 + m^2 v^2)/4$, we have $\min(L_0^2, L_0) \geq \frac{1}{4} \min\{u^2 + m^2 v^2, (u^2 + m^2 v^2)^{1/2}\}$. Discarding the nonnegative terms with $\alpha \notin \mathcal{G}$ and using $|\mathcal{G}| \geq (c_p/4)m^p/p!$ gives

$$\sum_{|\alpha| \leq m} \log \cosh(u/2 - |\alpha|v) \geq cm^p \min\{u^2 + m^2 v^2, (u^2 + m^2 v^2)^{1/2}\}. \quad (\text{C.23})$$

By [Theorem A.1](#), the affinity of the observation laws is at most the affinity of the laws of the parameter-independent linear statistic $H_n Y_n$. By [Theorem A.9](#), the latter log affinity differs from that of the exact coefficient vector by $o(1)$, uniformly over the parameters. Combining this fact with (C.22), (C.23), and $m_n \asymp b_n$ proves (A.15). The coefficient law has no nugget parameter, and its recovery bound is uniform in t ; this proves the claimed uniformity in the nugget as well.

We now prove (A.16). On the rectangle, the mean value theorem gives $|\log(s/s_0)| \geq |s - s_0|/s_+$ and $|\log(l/l_0)| \geq |l - l_0|/l_+$. Hence, with $c_1 = \min\{1, s_+^{-2}, l_+^{-2}\}$,

$$\Lambda_n(\theta)^2 \geq c_1 \{(s - s_0)^2 + b_n^2 (l - l_0)^2\} = c_1 b_n^{-p} \|h_{\text{sp}}\|^2.$$

Since $x \mapsto \min(x^2, x)$ is increasing on $[0, \infty)$,

$$b_n^p \min\{\Lambda_n(\theta)^2, \Lambda_n(\theta)\} \geq \min\{c_1 \|h_{\text{sp}}\|^2, c_1^{1/2} b_n^{p/2} \|h_{\text{sp}}\|\} \geq c_1 \min(\|h_{\text{sp}}\|^2, b_n^{p/2} \|h_{\text{sp}}\|),$$

and (A.15) gives $\text{Aff}(P_{\theta,n}, P_{\theta_0,n}) \leq \exp\{e_n - cc_1 \min(\|h_{\text{sp}}\|^2, b_n^{p/2} \|h_{\text{sp}}\|)\}$.

For the nugget, use the noise block $W_n = V_n^\top Y_n$ of (A.9). Since $P_n V_n = 0$, $\text{Cov}_\theta(W_n) = tI_{d_n} + E_\theta$ with $E_\theta = V_n^\top S_n(\theta)V_n = V_n^\top \{S_n(\theta) - P_n S_n(\theta)P_n\}V_n$. By (A.5), $0 \preceq E_\theta \preceq n^{-20}I_{d_n}$ for all $\theta \in \Theta$. Therefore the eigenvalues $\lambda_1, \dots, \lambda_{d_n}$ of $\text{Cov}_{\theta_0}(W_n)^{-1/2} \text{Cov}_\theta(W_n) \text{Cov}_{\theta_0}(W_n)^{-1/2}$ satisfy $|\log \lambda_k - \log(t/t_0)| \leq n^{-20}/t_-$. By (A.1),

$$-\log \text{Aff}\{\text{Cov}_\theta(W_n), \text{Cov}_{\theta_0}(W_n)\} = \frac{1}{2} \sum_{k=1}^{d_n} \log \cosh\left(\frac{1}{2} \log \lambda_k\right).$$

The function $y \mapsto \log \cosh(y/2)$ is $\frac{1}{2}$ -Lipschitz, and since its second derivative is $\frac{1}{4} \operatorname{sech}^2(y/2)$, it is at least $c_2 y^2$ for $|y| \leq \log(t_+/t_-)$, where $c_2 = \frac{1}{8} \operatorname{sech}^2\{\frac{1}{2} \log(t_+/t_-)\}$. Using $|\log(t/t_0)| \geq |t - t_0|/t_+$, $d_n \leq n$, and $d_n \geq n/2$ for large n ,

$$-\log \text{Aff}\{\text{Cov}_\theta(W_n), \text{Cov}_{\theta_0}(W_n)\} \geq \frac{c_2}{4t_+^2} n(t - t_0)^2 - \frac{n^{-19}}{t_-} = \frac{c_2}{4t_+^2} h_t^2 - \frac{n^{-19}}{t_-}.$$

The statistic $W_n = V_n^\top Y_n$ is linear and parameter independent, so [Theorem A.1](#) shows that the affinity of the observation laws is at most this affinity. Since $\min(e^{-x}, e^{-y}) \leq e^{-(x+y)/2}$, the spatial and nugget bounds together prove (A.16), with c replaced by $\frac{1}{2} \min\{cc_1, c_2/(4t_+^2)\}$ and e_n by

$\max(e_n, n^{-19}/t_-)$. The design enters only through [Theorems A.8](#) and [A.9](#). Their constants depend on $\mathcal{X}$, Q , C_h , and κ , and their proofs use (3.2) only at the current n . No constant depends on θ_0 : the bounds of [Theorem A.9](#) hold uniformly over pairs $\theta, \theta' \in \Theta$, and the remaining constants depend only on p and the endpoints of Θ . $\square$

C.11 Proof of [Theorem A.11](#)

Proof. The matrix $C_n(\theta)$ is a smooth positive definite function of θ on a neighborhood of Θ , so l_n is continuously differentiable there and ξ_n has continuous paths. Write $\theta_h = \theta_0 + D_n^{-1}h$ and $\nabla_h l_n(h) = D_n^{-1}(\nabla l_n)(\theta_h)$. Since $Z_n(h)$ is the density of $P_{\theta_h, n}$ with respect to $P_{\theta_0, n}$,

$$\mathbb{E}_{\theta_0, n} \xi_n(h)^m = \mathbb{E}_{\theta_0, n} Z_n(h)^{1/2} = \int \sqrt{dP_{\theta_h, n} dP_{\theta_0, n}},$$

which is [\(A.17\)](#).

For [\(A.18\)](#), let $u, v \in U_n$ and $w_r = v + r(u - v)$ for $r \in [0, 1]$; these points lie in the box U_n . Since $\frac{d}{dr} \xi_n(w_r) = (2m)^{-1} \xi_n(w_r) \nabla_h l_n(w_r)^\top (u - v)$, Minkowski's integral inequality gives

$$\|\xi_n(u) - \xi_n(v)\|_{L^m} \leq \frac{\|u - v\|}{2m} \sup_{w \in U_n} (\mathbb{E}_{\theta_0, n} [Z_n(w)^{1/2} \|\nabla_h l_n(w)\|^m])^{1/m}.$$

Fix w and abbreviate $\theta = \theta_w$. Changing measure and applying the Cauchy–Schwarz inequality under $P_{\theta, n}$,

$$\begin{aligned} \mathbb{E}_{\theta_0, n} [Z_n(w)^{1/2} \|\nabla_h l_n(w)\|^m] &= \mathbb{E}_{\theta, n} [Z_n(w)^{-1/2} \|\nabla_h l_n(w)\|^m] \\ &\leq (\mathbb{E}_{\theta, n} Z_n(w)^{-1})^{1/2} (\mathbb{E}_{\theta, n} \|\nabla_h l_n(w)\|^{2m})^{1/2}, \end{aligned}$$

and $\mathbb{E}_{\theta, n} Z_n(w)^{-1} = \mathbb{E}_{\theta_0, n} 1 = 1$. Under $P_{\theta, n}$, $X = C_n(\theta)^{-1/2} Y_n \sim N_n(0, I_n)$, and the coordinates of $\nabla_h l_n(w)$ are $\frac{1}{2} \{X^\top B_a(\theta) X - \text{tr } B_a(\theta)\}$ for $a \in \{s, l, t\}$, where $B_a(\theta) = D_{n, aa}^{-1} C_n(\theta)^{-1/2} C_{n, a}(\theta) C_n(\theta)^{-1/2}$. Dividing [\(A.8\)](#) by $D_{n, aa}$ gives $\|B_a(\theta)\|_F \leq C$ for $a \in \{s, l\}$, uniformly over $\theta \in \Theta$ and n , and $\|B_t(\theta)\|_F = n^{-1/2} \|C_n(\theta)^{-1}\|_F \leq t_-^{-1}$. [Theorem A.2](#) therefore gives $\mathbb{E}_{\theta, n} \|\nabla_h l_n(w)\|^{2m} \leq C$, uniformly in w and n , which proves [\(A.18\)](#). $\square$

C.12 Proof of [Theorem A.12](#)

Proof. Take $m = 4$, so that $m > k = 3$, and let ξ_n be the process of [Theorem A.11](#) on the box $U_n = D_n(\Theta - \theta_0)$. Write $u = (u_1, u_2)$, where $u_1 \in \mathbb{R}^2$ holds the spatial coordinates and $u_2 \in \mathbb{R}$ the nugget coordinate. Let n be large enough that $b_n \geq 1$. By [\(A.17\)](#) and [\(A.16\)](#), the envelope [\(A.2\)](#) holds with $A = \exp(\sup_n e_n)$, with a equal to the constant c of [\(A.16\)](#), and with $\gamma_1 = b_n^{p/2}$ and $\gamma_2 = \infty$. By [\(A.18\)](#), [\(A.3\)](#) holds with $B = C_L$. These constants do not depend on n .

Since $\hat{\theta}_n$ maximizes l_n over Θ , $\xi_n(\hat{h}_n) \geq \xi_n(0) = 1$. Hence Markov's inequality and [Theorem A.7](#) with $(H_1, H_2) = (H, 0)$ give

$$\begin{aligned} \mathbb{P}\{\|\hat{h}_{n, \text{sp}}\| \geq H\} &\leq \mathbb{P}\left\{\sup_{u \in U_n, |u_1| \geq H} \xi_n(u) \geq 1\right\} \leq \mathbb{E} \sup_{u \in U_n, |u_1| \geq H} \xi_n(u)^m \\ &\leq C \exp\{-a'' \min(H^2, b_n^{p/2} H)\}, \end{aligned}$$

with $a'' = a/128$, since $\psi_{\gamma_2}(0) = 0$. With $(H_1, H_2) = (0, H)$, the same argument gives $\mathbb{P}\{|\hat{h}_{n,t}| \geq H\} \leq C \exp(-a'' H^2)$. This proves (3.10). The constants A , a , B , and γ_1 do not depend on θ_0 , and neither do the constants of Theorem A.7, which do not depend on the box U_n . The design enters only through (A.16) and Theorem A.11. This gives the stated dependence of the constants. $\square$

C.13 Proof of Theorem A.13

Proof. The spatial upper bounds follow from (3.3) and the first derivative Frobenius bounds in Theorem A.8. The nugget upper bound follows from $C_n^{-1} \preceq t_-^{-1} I_n$.

Spatial lower bound. Write $D_{n,\text{sp}} = \text{diag}(b_n^{p/2}, b_n^{p/2+1})$. For $\|h\| \leq 1$ in $\mathbb{R}^2$, set $\theta_h = ((s_0, l_0) + D_{n,\text{sp}}^{-1} h, t_0)$ and

$$E_n(h) = C_{n,0}^{-1/2} \{C_n(\theta_h) - C_{n,0}\} C_{n,0}^{-1/2}.$$

For $a \in \{s, l\}$, let $B_{n,a} = D_{n,aa}^{-1} C_{n,0}^{-1/2} C_{n,a}(\theta_0) C_{n,0}^{-1/2}$ be the normalized whitened first derivatives at θ_0 .

Covariance comparability. Fix h with $\|h\| \leq 1$, put $C(u) = C_n(\theta_{uh})$ for $u \in [0, 1]$, and fix $x \in \mathbb{R}^n \setminus \{0\}$. The function $f(u) = x^\top C(u) x$ is positive, and

$$f'(u) = \sum_{a \in \{s, l\}} h_a D_{n,aa}^{-1} x^\top C_{n,a}(\theta_{uh}) x = \sum_{a \in \{s, l\}} h_a y^\top \tilde{B}_a(u) y,$$

where $y = C(u)^{1/2} x$ and $\tilde{B}_a(u) = D_{n,aa}^{-1} C(u)^{-1/2} C_{n,a}(\theta_{uh}) C(u)^{-1/2}$. Here the whitening is at the same parameter as the derivative, so dividing (A.7) by $D_{n,aa}$ gives $\|\tilde{B}_a(u)\|_{\text{op}} \leq C\epsilon_n$, uniformly in u and h . Since $|h_s| + |h_l| \leq \sqrt{2}$ and $\|y\|^2 = f(u)$,

$$|f'(u)| \leq \sqrt{2} C\epsilon_n f(u), \quad \text{that is,} \quad \left| \frac{d}{du} \log f(u) \right| \leq \sqrt{2} C\epsilon_n.$$

Integrating over $[0, u]$ gives $e^{-\sqrt{2} C\epsilon_n} f(0) \leq f(u) \leq e^{\sqrt{2} C\epsilon_n} f(0)$. Since x was arbitrary, after renaming the constant,

$$e^{-C\epsilon_n} C_{n,0} \preceq C_n(\theta_{uh}) \preceq e^{C\epsilon_n} C_{n,0}, \quad u \in [0, 1], \quad \|h\| \leq 1. \quad (\text{C.24})$$

Change of whitening. Let $K(u) = C_{n,0}^{-1/2} C(u)^{1/2}$. Then $\|K(u)\|_{\text{op}}^2 = \|C_{n,0}^{-1/2} C(u) C_{n,0}^{-1/2}\|_{\text{op}}$, which is at most $e^{C\epsilon_n}$ by (C.24). For any $n \times n$ matrix A ,

$$C_{n,0}^{-1/2} A C_{n,0}^{-1/2} = K(u) \{C(u)^{-1/2} A C(u)^{-1/2}\} K(u)^\top,$$

so

$$\|C_{n,0}^{-1/2} A C_{n,0}^{-1/2}\|_{\text{F}} \leq e^{C\epsilon_n} \|C(u)^{-1/2} A C(u)^{-1/2}\|_{\text{F}}. \quad (\text{C.25})$$

Taylor expansion. Taylor's formula with integral remainder, applied to $u \mapsto C(u)$, gives $E_n(h) = h_s B_{n,s} + h_l B_{n,l} + R_n(h)$ with

$$R_n(h) = \int_0^1 (1-u) C_{n,0}^{-1/2} C''(u) C_{n,0}^{-1/2} du, \quad C''(u) = \sum_{a,b \in \{s,l\}} h_a h_b D_{n,aa}^{-1} D_{n,bb}^{-1} C_{n,ab}(\theta_{uh}).$$

By [Theorem A.8](#), the normalized second derivatives whitened at θ_{uh} have Frobenius norm $O(\epsilon_n)$, so $\|C(u)^{-1/2}C''(u)C(u)^{-1/2}\|_F = O(\epsilon_n)$. By [\(C.25\)](#), the same holds after whitening at $C_{n,0}$, and therefore

$$E_n(h) = h_s B_{n,s} + h_l B_{n,l} + R_n(h), \quad \sup_{\|h\| \leq 1} \|R_n(h)\|_F = O(\epsilon_n). \quad (\text{C.26})$$

Moreover, $\|B_{n,a}\|_F \leq C$ by [Theorem A.8](#), so $\|E_n(h)\|_F = O(1)$, and [\(C.24\)](#) with $u = 1$ places every eigenvalue of $E_n(h)$ in $[e^{-C\epsilon_n} - 1, e^{C\epsilon_n} - 1]$, so $\|E_n(h)\|_{\text{op}} = O(\epsilon_n)$, uniformly on $\|h\| \leq 1$.

For any symmetric E with small operator norm,

$$\frac{1}{2} \log \det(I + E/2) - \frac{1}{4} \log \det(I + E) = \frac{1}{16} \text{tr}(E^2) + O(\|E\|_{\text{op}} \|E\|_F^2).$$

By [\(A.1\)](#) and [\(C.26\)](#),

$$-\log \text{Aff}(P_{\theta_h,n}, P_{\theta_0,n}) = \frac{1}{8} h^\top J_{n,\text{sp}} h + o(1) \quad (\text{C.27})$$

uniformly on $\|h\| \leq 1$.

For $\|h\| = 1$, the point θ_h has rescaled coordinates $(h, 0)$, with spatial part h and nugget part 0. Hence [\(A.16\)](#) gives, once $b_n \geq 1$,

$$-\log \text{Aff}(P_{\theta_h,n}, P_{\theta_0,n}) \geq c \min(1, b_n^{p/2}) - e_n = c - e_n.$$

Comparing with [\(C.27\)](#), we obtain $\frac{1}{8} h^\top J_{n,\text{sp}} h \geq c - o(1)$ uniformly on the unit circle, so $J_{n,\text{sp}} \succeq 4cI_2$ for all sufficiently large n . This argument uses a finite local shift and does not differentiate the coefficient recovery error.

Nugget block and cross information. By [Theorem A.8](#), the positive signal covariance $S_n(\theta_0)$ has at most $r_n = O(b_n^p)$ eigenvalues exceeding n^{-20} . Therefore

$$\frac{\mathcal{I}_{tt,n}(\theta_0)}{n} = \frac{1}{2n} \text{tr}(C_{n,0}^{-2}) \longrightarrow \frac{1}{2t_0^2}.$$

For $a = s, l$, approximate the normalized whitened derivative $B_{n,a}$ by a matrix $B_{n,a}^{(r)}$ of rank at most r_n , using the same projection. Its Frobenius norm is bounded and the discarded part is negligible in all traces below. The normalized nugget derivative is $B_{n,t} = C_{n,0}^{-1}/\sqrt{n}$, so

$$|J_{n,at}| \leq \frac{1}{2} \|B_{n,t}\|_{\text{op}} \|B_{n,a}^{(r)}\|_* + o(1) \leq C \sqrt{\frac{r_n}{n}} + o(1) = o(1),$$

where $\|\cdot\|_*$ is the sum of singular values and $\|B^{(r)}\|_* \leq \sqrt{r_n} \|B^{(r)}\|_F$. The spatial lower bound and the nugget limit establish [\(3.4\)](#) and [\(3.5\)](#). $\square$

C.14 Proof of [Theorem A.14](#)

Proof. Let $X_n = C_{n,0}^{-1/2} Y_n \sim N_n(0, I_n)$. Define, for $a, b \in \{s, l, t\}$,

$$B_a = \frac{C_{n,0}^{-1/2} C_{n,a}(\theta_0) C_{n,0}^{-1/2}}{D_{n,aa}}, \quad B_{ab} = \frac{C_{n,0}^{-1/2} C_{n,ab}(\theta_0) C_{n,0}^{-1/2}}{D_{n,aa} D_{n,bb}}.$$

The Gaussian score is

$$\Delta_{n,a} = \frac{1}{2} \{X_n^\top B_a X_n - \text{tr } B_a\}.$$

It has mean zero and covariance $J_{n,ab} = \frac{1}{2} \text{tr}(B_a B_b)$.

Directly differentiating the likelihood gives

$$[\nabla_h^2 L_n(0) + J_n]_{ab} = \frac{1}{2} \{X_n^\top M_{ab} X_n - \text{tr } M_{ab}\}, \quad M_{ab} = B_{ab} - B_a B_b - B_b B_a. \quad (\text{C.28})$$

[Theorem A.8](#) gives $\|B_a\|_{\text{op}} \leq C\epsilon_n$, $\|B_a\|_F \leq C$, and $\|B_{ab}\|_F \leq C\epsilon_n$; the nugget's operator bound is smaller, and second derivatives involving it are zero. Thus $\|M_{ab}\|_F \leq C\epsilon_n$. For a symmetric matrix M , $\text{Var}(X_n^\top M X_n) = 2 \|M\|_F^2$. Equation (C.28) proves (A.19).

Uniform third derivatives in the signal block. Use the fixed subspace and covariance A_θ from (A.9). Compression preserves the relative derivative bounds in [Theorem A.8](#). In particular, every nonzero covariance derivative of total order $k \leq 3$ in h has relative operator norm at most $C\epsilon_n^k$. A first nugget derivative is bounded by $Cn^{-1/2} \leq C\epsilon_n$, and higher covariance derivatives involving the nugget vanish.

For $\|h\| \leq H_n$, integration of the first relative derivative bound along the segment from 0 to h , exactly as in the proof of (C.24), gives

$$e^{-C\epsilon_n H_n} A_{\theta_0} \preceq A_{\theta_0 + D_n^{-1} h} \preceq e^{C\epsilon_n H_n} A_{\theta_0}.$$

Since $\epsilon_n H_n \rightarrow 0$ and $Z_n^\top A_{\theta_0}^{-1} Z_n \sim \chi_{r_n}^2$,

$$\sup_{\|h\| \leq H_n} Z_n^\top A_{\theta_0 + D_n^{-1} h}^{-1} Z_n = O_{\mathbb{P}}(r_n).$$

This bounds the entire neighborhood using the same random quadratic form.

Every third derivative of $-\frac{1}{2} \log \det A_\theta - \frac{1}{2} Z_n^\top A_\theta^{-1} Z_n$ is a finite sum of traces or quadratic forms formed from products of whitened covariance derivatives of total order three. Each product has operator norm at most $C\epsilon_n^3$. The traces are bounded by $C r_n \epsilon_n^3$, and the quadratic forms by $C \epsilon_n^3 Z_n^\top A_\theta^{-1} Z_n$. Therefore,

$$\sup_{\|h\| \leq H_n} \max_{a,b,c} |\partial_{h_a h_b h_c}^3 \tilde{L}_{n,\text{signal}}(h)| \leq C \epsilon_n^3 \left(r_n + \sup_{\|h\| \leq H_n} Z_n^\top A_\theta^{-1} Z_n \right) = O_{\mathbb{P}}(\epsilon_n),$$

because $r_n = O(b_n^p) = O(\epsilon_n^{-2})$.

In the complementary block, the log likelihood is $-\frac{1}{2} d_n \log t - \|W_n\|^2 / (2t)$. Only derivatives in h_t are nonzero. Its third derivative is bounded by $C n^{-3/2} (d_n + \|W_n\|^2) = O_{\mathbb{P}}(n^{-1/2})$ uniformly on the local ball. The exact likelihood derivative differs by $O_{\mathbb{P}}(n^{-16})$ from this reduced likelihood derivative, by (A.6); normalization only decreases that error. This proves (A.20).

The interior assumption and $H_n \|D_n^{-1}\|_{\text{op}} = \epsilon_n H_n \rightarrow 0$ ensure admissibility of the whole ball. Since $\nabla_h L_n(0) = \Delta_n$, Taylor's formula gives, for $\|h\| \leq H_n$,

$$L_n(h) - L_n(0) - h^\top \Delta_n + \frac{1}{2} h^\top J_n h = \frac{1}{2} h^\top \{\nabla_h^2 L_n(0) + J_n\} h + \frac{1}{6} \sum_{a,b,c} \partial_{h_a h_b h_c}^3 L_n(\bar{h}) h_a h_b h_c$$

for some $\bar{h}$ on the segment from 0 to h . By (A.19), (A.20), and $\sum_a |h_a| \leq \sqrt{3} \|h\|$, the right side is $O_{\mathbb{P}}(\epsilon_n H_n^2 + \epsilon_n H_n^3) = O_{\mathbb{P}}(\epsilon_n H_n^3)$ uniformly on the ball, because $H_n \geq 1$. This proves (A.21), and (A.22) follows.

Sharpness (Theorem A.15). Along the variance direction $h = ue_s$, the covariance is $C(\sigma) = \sigma R_n(l_0) + t_0 I_n$ with $\sigma = s_0 + b_n^{-p/2} u$, so $\partial_\sigma C = R_n(l_0)$ and $\partial_\sigma^2 C = 0$. Differentiating $l = -\frac{1}{2} \log \det C - \frac{1}{2} Y_n^\top C^{-1} Y_n$ gives

$$\partial_\sigma^3 l = -\text{tr}\{(C^{-1} R_n)^3\} + 3Y_n^\top C^{-1} (R_n C^{-1})^3 Y_n, \quad \partial_\sigma^4 l = 3\text{tr}\{(C^{-1} R_n)^4\} - 12Y_n^\top C^{-1} (R_n C^{-1})^4 Y_n,$$

with $R_n = R_n(l_0)$. At $u = 0$, in terms of $X_n = C_{n,0}^{-1/2} Y_n$ and the normalized derivative B_s , $\partial_{h_s}^3 L_n(0) = -\text{tr}(B_s^3) + 3X_n^\top B_s^3 X_n$. Its mean is $2\text{tr}(B_s^3)$, and its standard deviation is $3\sqrt{2} \|B_s^3\|_F \leq C \|B_s\|_{\text{op}}^2 \|B_s\|_F \leq C \epsilon_n^2$ by Theorem A.8. The matrix B_s is positive semidefinite. By the Cauchy-Schwarz inequality over its eigenvalues, $(\text{tr} B_s^2)^2 \leq \text{tr}(B_s) \text{tr}(B_s^3)$. Here $\text{tr} B_s^2 = 2J_{n,ss} \geq 2c$ by Theorem A.13. Moreover, $B_s = b_n^{-p/2} R_n(s_0 R_n + t_0 I_n)^{-1}$ has eigenvalues $b_n^{-p/2} \mu / (s_0 \mu + t_0)$, where μ ranges over the eigenvalues of R_n . By (A.5) and Weyl's inequality, at most r_n of these μ exceed n^{-20}/s_0 , so $\text{tr} B_s \leq b_n^{-p/2} (r_n/s_0 + n^{-19}/(s_0 t_0)) \leq C \epsilon_n^{-1}$. Hence $\text{tr}(B_s^3) \geq c \epsilon_n$, and $\partial_{h_s}^3 L_n(0) \geq c \epsilon_n$ with probability tending to one, after decreasing c .

For the fourth derivative, all matrices involved commute with R_n . In an eigenbasis of R_n , with Y_n having independent coordinates of variances $s_0 \mu_i + t_0$, each eigenvalue of $C^{-1} R_n$ is at most $C \min(1, \mu_i)$ for $|\sigma - s_0| \leq s_0/2$. Therefore $\sup_{|\sigma - s_0| \leq s_0/2} |\partial_\sigma^4 l|$ is bounded by a constant times $\sum_i \min(1, \mu_i^4) (1 + \chi_i^2)$ with χ_i independent standard normal, which is $O_{\mathbb{P}}(r_n)$ by the eigenvalue count above. After normalization, the fourth derivative in h_s is $O_{\mathbb{P}}(b_n^{-2p} r_n) = O_{\mathbb{P}}(\epsilon_n^2)$ uniformly on $|u| \leq \kappa \epsilon_n^{-1/3}$.

Now let $u = \kappa \epsilon_n^{-1/3}$, so that $\sigma - s_0 = \kappa \epsilon_n^{2/3} \rightarrow 0$. Taylor's formula in u , together with (A.19), gives

$$L_n(ue_s) - L_n(0) - u \Delta_{n,s} + \frac{1}{2} J_{n,ss} u^2 = O_{\mathbb{P}}(\epsilon_n u^2) + \frac{1}{6} \partial_{h_s}^3 L_n(0) u^3 + O_{\mathbb{P}}(\epsilon_n^2 u^4).$$

The first and last terms are $O_{\mathbb{P}}(\kappa^2 \epsilon_n^{1/3})$ and $O_{\mathbb{P}}(\kappa^4 \epsilon_n^{2/3})$, and the middle term is at least $c \kappa^3/6$ with probability tending to one. This proves Theorem A.15. $\square$

C.15 Proof of Theorem A.16

Proof. Use X_n and B_a from the preceding proof. For a fixed $v \in \mathbb{R}^3$, write

$$v^\top M_n^{-\top} \nabla l_n(\theta_0) = \frac{1}{2} \{X_n^\top B_n(v) X_n - \text{tr} B_n(v)\},$$

where

$$B_n(v) = \sum_a [D_n M_n^{-1} v]_a B_a.$$

Since $M_n^{-1} M_n^{-\top} = (M_n^\top M_n)^{-1} = \mathcal{I}_n(\theta_0)^{-1}$ and $\mathcal{I}_n(\theta_0) = D_n J_n D_n$,

$$(D_n M_n^{-1})(D_n M_n^{-1})^\top = D_n \mathcal{I}_n(\theta_0)^{-1} D_n = J_n^{-1}, \quad \|D_n M_n^{-1}\|_{\text{op}}^2 = \|J_n^{-1}\|_{\text{op}} \leq C$$

by [Theorem A.13](#), so the coefficient vector is bounded. Hence $\|B_n(v)\|_{\text{op}} \leq C_v \epsilon_n \rightarrow 0$. The variance is exactly $\frac{1}{2} \text{tr}(B_n(v)^2) = v^\top M_n^{-\top} \mathcal{I}_n(\theta_0) M_n^{-1} v = \|v\|^2$.

For $v \neq 0$, diagonalize the symmetric matrix $B_n(v)$, with eigenvalues $\lambda_{j,n}$. The quadratic form is distributed as $\frac{1}{2} \sum_j \lambda_{j,n} (\xi_j^2 - 1)$ for independent standard normal variables. Its Lyapunov ratio of order four is bounded by a constant times

$$\frac{\sum_j \lambda_{j,n}^4}{(\sum_j \lambda_{j,n}^2)^2} \leq \frac{\max_j \lambda_{j,n}^2}{\sum_j \lambda_{j,n}^2} \rightarrow 0.$$

The univariate central limit theorem and the Cramér–Wold device give [\(A.23\)](#). The same calculation applies to any bounded sequence of coefficient vectors for Δ_n whose variances converge to a positive limit. $\square$

C.16 Proof of [Theorem A.17](#)

Proof. Let Q have side length L , and let $A \geq 1$ be a fixed constant. For $n \geq 3$, put

$$k_n = \left\lfloor \left(\frac{n}{A \log n} \right)^{1/p} \right\rfloor,$$

and suppose $k_n \geq 1$. Partition Q into k_n^p closed subcubes $Q_1, \dots, Q_{k_n^p}$ of side length L/k_n , which overlap only on their boundaries. Each has volume L^p/k_n^p , so its sampling probability satisfies

$$\pi_j = \mathbb{P}(X_1 \in Q_j) = \int_{Q_j} f(x) dx \geq \frac{f_- L^p}{k_n^p}.$$

Since $X_1, \dots, X_n$ are independent and $1 - x \leq e^{-x}$,

$$\mathbb{P}(\text{no } X_i \text{ lies in } Q_j) = (1 - \pi_j)^n \leq \left(1 - \frac{f_- L^p}{k_n^p}\right)^n \leq \exp\left(-\frac{n f_- L^p}{k_n^p}\right).$$

A union bound over the k_n^p subcubes gives

$$\mathbb{P}(\text{some } Q_j \text{ is empty}) \leq k_n^p \exp\left(-\frac{n f_- L^p}{k_n^p}\right).$$

By the definition of k_n , $k_n^p \leq n/(A \log n)$. Hence $k_n^p \leq n$, because $A \log n \geq 1$, and $n/k_n^p \geq A \log n$, so that $\exp(-n f_- L^p/k_n^p) \leq \exp(-f_- L^p A \log n) = n^{-f_- L^p A}$. Therefore

$$\mathbb{P}(\text{some } Q_j \text{ is empty}) \leq n \exp(-f_- L^p A \log n) = n^{1-f_- L^p A}.$$

Given $q > 0$, choose $A \geq \max\{1, (q+1)/(f_- L^p)\}$, so that this probability is at most n^{-q} . When every subcube contains an observation, every $x \in Q$ lies in some Q_j together with an observation, so it is within the diameter $\sqrt{p} L/k_n$ of that observation. Since $\lfloor y \rfloor \geq y/2$ for $y \geq 1$,

$$h_n(Q) \leq \frac{\sqrt{p} L}{k_n} \leq 2\sqrt{p} L \left(\frac{A \log n}{n} \right)^{1/p}$$

on this event, for all $n \geq 3$ with $k_n \geq 1$. This proves the probability bound with $A_q = 2\sqrt{p} LA^{1/p}$, after enlarging C_q to cover the finitely many n with $n < 3$ or $k_n = 0$. With $q = 2$, the first Borel–Cantelli lemma proves (A.24). Since $(\log n/n)^{1/p} = o(n^{-\kappa})$ for $\kappa < 1/p$, the coverage conclusion follows. $\square$

D Additional simulation details

This appendix provides additional computational details and finite-sample results for the simulations in Section 4.

D.1 Exact evaluation of the likelihood

The sample sizes considered in Section 4 are too large for direct $O(n^3)$ evaluation of the Gaussian likelihood. We therefore exploit the analytic structure of the RBF kernel and, for $p \geq 2$, the Kronecker structure of the regular grid. These calculations evaluate the same Gaussian likelihood in (3.1), up to numerical truncation at machine precision.

Feature representation. On $[0, 1]$, let $u = (x - \frac{1}{2})/l$. We use the expansion

$$\exp\left\{-\frac{(x - x')^2}{2l^2}\right\} = \sum_{k=0}^{\infty} \phi_k(x)\phi_k(x'), \quad \phi_k(x) = e^{-u^2/2} \frac{u^k}{\sqrt{k!}}.$$

We truncate the expansion after

$$M(l) = \left\lceil \bar{u} + 10\sqrt{\bar{u}} + 25 \right\rceil, \quad \bar{u} = (2l)^{-2}.$$

Over the parameter range $l \geq 0.05$, the reconstructed kernel and its lengthscale derivative agree with direct evaluation to approximately 10^{-13} in maximum norm. The features are evaluated recursively using $\phi_k = \phi_{k-1}u/\sqrt{k}$ to avoid numerical over- and underflow.

Let Φ denote the $n \times M$ feature matrix. Then

$$R_n(l) \approx \Phi\Phi^\top,$$

with the approximation at machine precision under the truncation above.

Spectral form of the likelihood. For fixed l , let

$$R_n(l) = U\Lambda U^\top, \quad \Lambda = \text{diag}(\lambda_1, \dots, \lambda_r),$$

where eigenvalues smaller than $10^{-12}\lambda_{\max}$ are discarded. Writing $z = U^\top Y$, the Gaussian likelihood can be evaluated as

$$-2l_n(\theta) = \sum_{k=1}^r \left\{ \log(\sigma^2\lambda_k + \tau^2) + \frac{z_k^2}{\sigma^2\lambda_k + \tau^2} \right\} + (n - r) \log \tau^2 + \frac{\|Y\|^2 - \|z\|^2}{\tau^2},$$

up to an additive constant. The quantities Λ and z are obtained from the much smaller Gram matrix $\Phi^\top \Phi$ and the vector $\Phi^\top Y$.

Linear-cost Gram matrix. The feature products satisfy

$$\phi_j(x)\phi_k(x) = e^{-u^2} \frac{u^{j+k}}{\sqrt{j!k!}},$$

so $\Phi^\top \Phi$ is a scaled Hankel matrix:

$$[\Phi^\top \Phi]_{jk} = h_{j+k} \binom{j+k}{j}^{1/2}, \quad h_m = \sum_{i=1}^n e^{-u_i^2} \frac{u_i^m}{\sqrt{m!}}.$$

Thus, forming the Gram matrix requires $O(nM)$ rather than $O(nM^2)$ operations. The midpoint grid is symmetric around $1/2$, so the odd moments vanish. This symmetry further reduces the computation without changing the likelihood.

Kronecker structure for $p \geq 2$. For the regular grid used in the simulations,

$$R_n(l) = R_G(l)^{\otimes p}.$$

Consequently, its eigenvalues are products of the one-dimensional eigenvalues,

$$\lambda_{k_1} \cdots \lambda_{k_p},$$

and projections of the observations are obtained by multiplying the data array by the one-dimensional eigenvector matrix along each axis. Thus, only one-dimensional quantities involving $G = n^{1/p}$ grid points need to be explicitly constructed.

D.2 Data generation

For $p = 1$, data are generated as

$$Y = \sigma_0 \Phi \xi + \tau_0 \varepsilon,$$

where $\xi \sim N_M(0, I_M)$ and $\varepsilon \sim N_n(0, I_n)$ are independent and Φ is evaluated at l_0 . Therefore,

$$\text{Cov}(Y) = \sigma_0^2 \Phi \Phi^\top + \tau_0^2 I_n = C_n(\theta_0)$$

up to the machine-precision truncation described above.

For $p \geq 2$, the feature matrix is replaced by its Kronecker product and ξ by an array of independent standard normal variables. This generates a new independent realization of both the GP and measurement errors for each Monte Carlo replicate.

D.3 Computation of the MLE

The MLE is computed over

$$\Theta = [10^{-2}, 10^2] \times [0.05, 2] \times [10^{-4}, 1]$$

for (σ^2, l, τ^2) .

For a fixed l , we optimize over $(\log \sigma^2, \log \tau^2)$ using L-BFGS-B with analytic gradients, relative function tolerance 10^{-15} , and gradient tolerance 10^{-10} . The profile likelihood is first evaluated at 25 log-equispaced values of l in $[0.05, 2]$. We then refine the best candidate by a bounded one-dimensional Brent search in $\log l$, with tolerance 10^{-7} . If the original grid value gives a larger likelihood than the refined value, the grid value is retained.

Among the 2.7×10^4 fitted data sets, only two estimates occur on the boundary of Θ : one estimate of l for $p = 1$, $n = 10^2$, and one estimate of τ^2 at its lower bound for $p = 3$, $n = 125$.

For several $p = 1$ data sets producing extreme values of $\hat{\sigma}^2$, we additionally evaluated the profile likelihood over a fine grid of 400 lengthscales. In each case, the profile likelihood had a single maximum coinciding with the computed MLE. Thus, these extreme estimates arise from the likelihood itself rather than numerical optimization failure.

D.4 Computation of the Fisher information

For each n , the Fisher information $\mathcal{I}_n(\theta_0)$ in (3.3) is computed directly from the observation design. Let V be an orthonormal basis for the span of Φ and $\partial_l \Phi$ at l_0 . Both R_n and $\dot{R}_n$ vanish, up to the machine-precision truncation, on the orthogonal complement of this space, where C_n acts as $\tau_0^2 I$.

The traces in (3.3) therefore reduce to calculations involving $V^\top R_n V$ and $V^\top \dot{R}_n V$, together with the contribution

$$\frac{n - \dim(V)}{2\tau_0^4}$$

to the nugget information. For $p \geq 2$, the Kronecker structure again reduces these calculations to one-dimensional quantities.

Because the resulting computation depends on n primarily through one-dimensional Gram matrices, we can evaluate the Fisher-information benchmark beyond the range used for Monte Carlo simulation. We evaluate it up to $n = 10^8$ for $p = 1$ and $n = 10^{12}$ for $p = 2, 3$. These extended calculations are used below to examine how slowly the finite-sample information approaches the limiting rates in Theorem 3.1.

D.5 Monte Carlo summaries

For each parameter, the main text reports the RMSE, the robust standard deviation IQR/1.349, and the asymptotic standard deviation obtained from the Fisher information.

For a normal random variable with variance v ,

$$\text{IQR} = 2\Phi^{-1}(0.75)\sqrt{v} \approx 1.349\sqrt{v}.$$

Thus, IQR/1.349 estimates the standard deviation under normality. Unlike the RMSE, it depends only on the central half of the sampling distribution and is therefore less sensitive to a small number

of extreme estimates. Agreement between the RMSE and robust standard deviation provides an additional indication that the estimation error is approximately normal.

Bootstrap 95% intervals for the RMSE and robust standard deviation are computed using 2000 bootstrap resamples of the 1000 Monte Carlo replicates.

D.6 Additional normality diagnostics

The Q–Q plots in [Figures 1 to 3](#) provide the primary graphical assessment of asymptotic normality. Additional numerical diagnostics lead to the same conclusion.

For $p = 3$ and $n = 10^6$, the standardized errors of $\log \hat{\sigma}^2$ and $\log \hat{l}$ have skewness -0.05 and 0.00 , respectively, and their Kolmogorov–Smirnov distances from $N(0, 1)$ are below 0.02 . Consistent with these diagnostics, the RMSE and robust standard deviation are nearly identical.

Convergence is slower for $p = 1$. The ratio of the RMSE to the Fisher-information standard deviation remains above one over the simulated range. For $\log \hat{\sigma}^2$, the excess decreases from 18% at $n = 10^2$ to 10% at $n = 10^6$; for $\log \hat{l}$, it decreases from 54% to 26%. The robust standard deviation is closer to the Fisher-information benchmark, with deviations between 4% and 20%. Because these ratios continue to decrease with n , the Monte Carlo slopes for $p = 1$ over the simulated range are steeper than the corresponding slopes of the Fisher-information standard deviation.

D.7 Finite-sample behavior of the nugget

The particularly rapid convergence of the nugget estimator can be understood from the effective rank of the RBF kernel matrix. By [Theorem A.8](#), the signal component has numerical rank $O(b_n^p)$. The remaining directions are approximately pure noise with variance τ^2 . Since b_n^p grows much more slowly than n , most directions eventually contribute almost exclusively to estimation of the nugget.

This explains the limit

$$\sqrt{n}(\hat{\tau}^2 - \tau_0^2) \longrightarrow N(0, 2\tau_0^4)$$

asymptotically, the nugget behaves approximately like a variance estimated from n independent $N(0, \tau_0^2)$ observations.

The same interpretation explains the deviations at the smallest sample sizes. The approximation becomes accurate only when n is large relative to the number of signal directions. For example, for $p = 3$ and $n = 125$, 96 of the 125 eigen-directions of $R_n(l_0)$ have signal variance $\sigma_0^2 \lambda_k$ exceeding the noise variance τ_0^2 . Thus, relatively few directions behave as nearly pure noise, and the nugget estimator is farther from its asymptotic regime.

D.8 Finite-sample behavior of the spatial parameters

The limiting rates for σ^2 and l are powers of

$$b_n = \frac{\log n}{\log \log n},$$

which grows extremely slowly. Over the entire simulated range $10^2 \leq n \leq 10^6$, b_n increases only from approximately 3.0 to 5.3. As a result, finite-sample slopes need not be close to their limiting exponents even when the Monte Carlo errors already closely track the Fisher-information benchmark.

Lower-order terms in the effective dimension. The proof suggests approximately

$$N_p(m) = \binom{m+p}{p}$$

recoverable Taylor coefficients through degree $m \asymp b_n$. The limiting rate for σ^2 uses

$$N_p(m) \sim \frac{m^p}{p!},$$

but the lower-order terms are substantial when m is only around 3–5. For example,

$$\frac{d \log N_2(m)}{d \log m} = 1.55$$

at $m = 5$, rather than its limiting value 2. Since the variance of $\hat{\sigma}^2$ is of order $N_p(m)^{-1}$, this produces a shallower finite-sample slope than the limiting value $-p/2$.

The exact Fisher-information calculations show this effect directly. Near $n = 10^6$, the local slopes of the asymptotic standard deviation against b_n are -0.37 , -0.74 , and -1.13 for σ^2 when $p = 1, 2, 3$, compared with the limiting values -0.5 , -1 , and -1.5 . The convergence is extremely slow: the corresponding slopes reach only -0.39 at $n = 10^8$ for $p = 1$ and -0.81 and -1.22 at $n = 10^{12}$ for $p = 2, 3$. For $p = 2, 3$, the Monte Carlo slopes in [Table 1](#) closely reproduce these finite- n Fisher-information slopes.

The behavior of the lengthscale is somewhat different because its information is weighted toward the highest recoverable degrees. Near $n = 10^6$, the local Fisher-information slopes are -1.59 , -2.06 , and -2.58 for $p = 1, 2, 3$, already close to the limiting values $-(p+2)/2$. At smaller sample sizes they are steeper; for example, the local slopes between $n = 10^2$ and 10^3 are approximately -2.57 for $p = 2$ and -4.46 for $p = 3$. This explains why slopes fitted over the entire simulated range can be steeper than the limiting exponent.

The Fisher information is the relevant finite-sample benchmark. [Theorem 3.1](#) determines the orders of the Fisher information but does not require J_n to converge. Consequently, the limiting powers of b_n describe the eventual rate but need not provide an accurate finite- n approximation. The design-specific quantity

$$[\mathcal{I}_n(\theta_0)^{-1}]_{jj}^{1/2}$$

is therefore the appropriate benchmark for the Monte Carlo errors at a given n . The close agreement between this quantity and the simulated errors for $p = 2, 3$ shows that the MLE is already well described by the Fisher approximation even though the Fisher information itself has not yet reached its limiting power law.

The one-dimensional case. The normal approximation is slower for $p = 1$ because the effective number of signal directions remains small. Over $10^2 \leq n \leq 10^6$, only approximately 7–11 eigen-directions have signal variance exceeding the noise variance. Thus, $\hat{\sigma}^2$ behaves roughly like a variance estimator based on a small number of independent Gaussian components, leading to visible skewness on the original scale.

The log transformation removes much of this skewness. The remaining tail behavior is associated with the strong positive dependence between $\hat{\sigma}^2$ and $\hat{l}$ along the likelihood ridge. As noted in [Appendix D.3](#), direct evaluation of the profile likelihood for the extreme cases confirms that these estimates are genuine likelihood maximizers rather than numerical optimization failures.

Correspondingly, at $n = 10^6$ the asymptotic standard deviations of $\log \hat{\sigma}^2$ are 0.59, 0.26, and 0.12, and those of $\log \hat{l}$ are 0.059, 0.022, and 0.010, for $p = 1, 2, 3$, respectively. The larger number of informative directions in higher dimensions also helps explain why the normal approximation improves from $p = 1$ to $p = 3$.

D.9 Validation against dense computations

For designs small enough to permit direct $O(n^3)$ linear algebra, we compared the fast implementation with dense Cholesky-based calculations. The comparisons used n between 300 and 4000 for $p = 1$, $G = 20$ for $p = 2$, and $G = 6$ for $p = 3$.

The log likelihoods evaluated at a range of parameter values agreed to within 10^{-6} . The MLEs agreed with those obtained by direct Nelder–Mead optimization of the dense likelihood to six significant digits. The Fisher information matrices agreed to relative error approximately 10^{-13} . Finally, the empirical covariance matrices computed from up to 2×10^4 independently generated data sets agreed with $C_n(\theta_0)$ within Monte Carlo error.

These checks confirm that the feature and Kronecker calculations used in the simulations reproduce the exact Gaussian likelihood and Fisher information to numerical precision.

D.10 Monte Carlo implementation and software

Each Monte Carlo replicate uses an independent random seed generated from

```
numpy.random.SeedSequence((20261003, p)).
```

Consequently, the results do not depend on the number of parallel workers, and individual sample sizes can be recomputed independently.

Computations used Python 3.13 with NumPy 2.4 and SciPy 1.18 on an 8-core laptop. The total computation time was approximately 5 hours: about 4.5 hours for $p = 1$, dominated by the cases $n \geq 3 \times 10^5$, approximately 15 minutes for $p = 2$, and approximately 10 minutes for $p = 3$. For $p \geq 2$, a single MLE fit at $n = 10^6$ requires approximately 0.1–2 seconds.

D.11 Results on the original scale

[Figures A1 to A3](#) repeat the simulation results in [Figures 1 to 3](#) for (σ^2, l, τ^2) on the original parameter scale. The reference rates are unchanged, and the asymptotic standard deviation is $[\mathcal{I}_n(\theta_0)^{-1}]_{jj}^{1/2}$ rather than its log-scale counterpart $[\mathcal{I}_n(\theta_0)^{-1}]_{jj}^{1/2}/\theta_{0j}$.

The conclusions concerning the convergence rates are unchanged. The main visible difference is greater right skewness of $\hat{\sigma}^2$, particularly for $p = 1$, and also of $\hat{l}$ in the one-dimensional case. This skewness increases the RMSE relative to the robust standard deviation and is substantially reduced by the log transformation, consistent with the discussion in [Section 4](#).




Figure A1: Simulation results for $p = 1$ on the original parameter scale, with the layout and graphical elements as in Figure 1.




Figure A2: Simulation results for $p = 2$ on the original parameter scale, with the layout and graphical elements as in Figure 1.




Figure A3: Simulation results for $p = 3$ on the original parameter scale, with the layout and graphical elements as in Figure 1.