Stable Computation of Revised Spherical Cap Harmonics

Pan Zhang[1,2,3], Jinsong Du[2,3,4,*] and Jiaxuan Zhang[5]

[1]School of Geology and Jewelry, Lanzhou Resources & Environment Voc-Tech University, Lanzhou 730021, China
[2]School of Geophysics and Geomatics, China University of Geosciences, Wuhan 430074, China
[3]Hubei Subsurface Multi-scale Imaging Key Laboratory, China University of Geosciences, Wuhan 430074, China
[4]Key Laboratory of Geological Survey and Evaluation of Ministry of Education, China University of Geosciences (Wuhan), Wuhan 430074, China
[5]Macau Institute of Space Technology and Application, Macau University of Science and Technology, Macao 999078, China
*Corresponding author: jinsongdu@cug.edu.cn

Abstract

High-resolution regional magnetic-field modeling requires special functions that remain stable from the cap axis to oscillatory high-degree regimes. Revised spherical cap harmonic analysis (R-SCHA) couples these functions to derivatives, boundary roots and cap norms, making consistency across numerical representations essential. We develop an open C++ basis layer for its real-degree Legendre, Mehler and zero-eigenvalue families. Scaled Legendre representations carry analytic degree sensitivities; a Riccati-sensitivity identity removes the raw amplitude from Mehler normalization; and a positive integral avoids cancellation in the zero-eigenvalue norm. Family-specific normalization, residual tables and sign–logarithm records connect these algorithms across floating-point ranges. Across 40,995 precision-refined reference evaluations, the maximum scaled value or angular-derivative error is 1.11×10^{-8} and the maximum direct degree-sensitivity error is 1.06×10^{-10}. Independent tests cover Legendre degrees below 10^5 and Mehler parameters and zero-eigenvalue orders through 10^4; algebraic consistency checks extend to degree 10^8. The maximum sampled relative Newton correction for boundary roots is 2.83×10^{-14}. Peak throughput ranges from 2.80×10^4 to 5.40×10^6 evaluations s^{-1} across the three families on a 10-core workstation. Cascadia and North American Midcontinent inversions combine dense near-surface constraints with satellite differences at internal equivalent degrees near 9.1×10^3. Their common-radius B_z maps and cap spectra retain shorter-wavelength structure beyond EMM2017, WDMAMv2 and SH1050, with smoothly decaying high-degree power. The resulting basis makes high-degree three-dimensional regional modeling numerically tractable and connects its additional bandwidth to geological interpretation.

Keywords: revised spherical cap harmonic analysis; associated Legendre function; conical function; nonoscillatory phase; Riccati sensitivity; canonical-residual interpolation

1 Introduction

Lithospheric magnetic fields are spatially multiscale: satellite missions constrain long wavelengths, whereas crustal anomalies and tectonic boundaries require shorter regional structure. Global spherical-harmonic models provide a consistent reference but impose a domain-wide bandwidth through their truncation and sampling. Spherical-cap harmonic analysis (SCHA) concentrates a basis on a finite region (Haines, 1985); revised SCHA (R-SCHA) extends it to a three-dimensional spherical-conical shell and has become a practical framework for regional geomagnetic and lithospheric-field modeling (Thébault et al., 2004, 2006; Vervelidou et al., 2018; Zhang et al., 2025). The broader cap-harmonic literature is reviewed by Torta (2020). At high resolution,

however, the value of a cap expansion depends on whether its basis remains numerically consistent as the truncation increases.

High-degree R-SCHA exposes a coupled numerical problem. Regular non-integer-degree solutions can be exponentially small near the cap axis, pass through turning regions and become rapidly oscillatory. Values, angular derivatives, degree sensitivities, boundary roots and cap norms then span different scales and may be evaluated by different branches. A plausible point value can therefore carry an incompatible phase, normalization or derivative into the field kernel and inverse problem. Extended-range arithmetic, asymptotics, recurrences and phase methods provide important Legendre components (Olver and Smith, 1983; Smith et al., 1981; Fukushima, 2012; Bremer and Rokhlin, 2017; Bremer, 2018); normalized cap functions and eigenvalues have also been studied (Hwang and Chen, 1997), and general conical-function algorithms cover broader classes (Gil et al., 2009, 2012; Dunster et al., 2015). The remaining challenge is to connect all three angular families, their derivatives, roots and norms through compatible family-specific normalizations and a common validation framework.

We develop that layer for the real-degree Legendre, Mehler and zero-eigenvalue families. The Legendre evaluator combines series, recurrence, propagation and phase representations on the adopted Schmidt scale and carries analytic degree sensitivities. A Riccati-sensitivity identity removes the raw exponentially varying amplitude from Mehler normalization, while a positive integral avoids cancellation in the zero-eigenvalue norm. Residual and logarithmic-norm tables provide reusable corrections for repeated evaluations; direct fallbacks and sign–logarithm records preserve robustness outside ordinary binary64 ranges. This architecture makes the coupled quantities required by R-SCHA interoperable rather than treating each special-function call in isolation.

We validate the implementation with precision-refined references, dense identities, boundary-root residuals and cap-norm integrations. The tests cover Legendre degrees below 10^5, Mehler parameters and zero-eigenvalue orders through 10^4, and caps up to a hemisphere; wider identity checks reach degree 10^8. We then apply the same basis to Cascadia and North American Midcontinent inversions. At a common radius and grid, the regional fields retain coherent B_z structure and cap-spectrum power beyond EMM2017, WDMAMv2 and SH1050, with a smoothly decaying high-degree tail. Geological and modified horizontal-gradient comparisons organize plausible magnetic transitions without assigning unique source depths.

2 R-SCHA boundary-value formulation

2.1 Spherical-conical domain and separation of Laplace's equation

Let (r, θ, φ) denote spherical coordinates with polar axis through the cap center, and consider the source-free spherical-conical domain

$$\Omega = \{(r, \theta, \varphi) : r_1 \le r \le r_2,\ 0 \le \theta \le \theta_0,\ 0 \le \varphi < 2\pi\}\,. \tag{1}$$

A scalar potential V in Ω satisfies

$$\nabla^2 V = 0. \tag{2}$$

For a separated mode $V = R(r)f(\theta)\Phi_m(\varphi)$, with $\Phi_m = \cos m\varphi$ or $\sin m\varphi$, the radial and angular equations may be written as

$$r^2 R'' + 2rR' - \lambda R = 0, \tag{3}$$

and

$$\mathcal{A}_m f = -\frac{1}{\sin\theta}\frac{\mathrm{d}}{\mathrm{d}\theta}\left(\sin\theta\frac{\mathrm{d}f}{\mathrm{d}\theta}\right) + m^2\csc^2\theta\, f = \lambda f. \tag{4}$$

Regularity at the axis requires $f(\theta) = O(\theta^m)$ for $m > 0$ and a bounded even solution for $m = 0$. Boundary conditions on the conical side and the two spherical surfaces determine the spectral parameters and radial combinations (Thébault et al., 2004, 2006).

2.2 The three angular families

The separation constant falls into three classes,

$$\lambda = \nu(\nu+1), \qquad \lambda = -\left(\tau^2 + \tfrac{1}{4}\right), \qquad \lambda = 0. \tag{5}$$

For $\lambda > 0$, the regular angular solution is an associated Legendre function of real degree ν; the radial solutions are r^ν and $r^{-\nu-1}$. For $\lambda = -(\tau^2 + 1/4)$, the angular solution is the real regular branch proportional to $\mathsf{P}^m_{-1/2+\mathrm{i}\tau}(\cos\theta)$. For $\tau > 0$, independent real radial solutions are

$$r^{-1/2}\cos[\tau\log(r/r_*)], \qquad r^{-1/2}\sin[\tau\log(r/r_*)], \tag{6}$$

where r_* is a reference radius. At $\tau = 0$, the independent pair is $r^{-1/2}$ and $r^{-1/2}\log(r/r_*)$, obtained by taking the limit of the sine solution divided by τ. For $\lambda = 0$, the regular angular shape is $\tan^m(\theta/2)$ for $m \geq 1$ and constant for $m = 0$; the radial solution is $A + B/r$.

The resulting R-SCHA potential can therefore be written schematically as

$$V = V^{\mathrm{in}}_{\mathrm{Leg}} + V^{\mathrm{out}}_{\mathrm{Leg}} + V_{\mathrm{Meh}} + V_0. \tag{7}$$

The first two terms use the real-degree Legendre family, the third the Mehler family, and the fourth the zero-eigenvalue family. Coefficient and radial conventions depend on the model boundary conditions. All require angular values, derivatives and cap norms; the Legendre terms also require real-degree boundary roots. We develop this angular basis independently of the model drivers.

3 Angular basis and normalization identities

Let $0 < \theta_0 < \pi$, $0 \leq \theta \leq \theta_0$, and $m = 0, 1, 2, \ldots$. The angular operator and its three spectral branches are given by Eqs. (4) and (5). For a real angular function f, define

$$I_f(\theta_0) = \|f\|^2_{\mathrm{cap}} = \int_0^{\theta_0} f^2(\theta)\sin\theta\,\mathrm{d}\theta, \qquad \widehat{f}(\theta) = \frac{f(\theta)}{\sqrt{I_f(\theta_0)}}. \tag{8}$$

Thus I_f is a squared norm and $\widehat{f}$ has unit cap norm. Longitude factors are excluded: their squared integrals over $[0, 2\pi)$ are 2π for $\cos 0\varphi$ and π for the nonzero-order sine and cosine branches. Analytical angles are in radians; degree-valued compatibility interfaces are identified in Section 4.1.

For a differentiable regular family $f_{\lambda m}$, put $\mathcal{S}_{\lambda m} = \partial_\lambda f_{\lambda m}$ at fixed (m, θ). The Lagrange identity for the angular operator gives

$$\frac{\mathrm{d}}{\mathrm{d}\theta}\left\{\sin\theta\left[\mathcal{S}_{\lambda m} f'_{\lambda m} - f_{\lambda m}\mathcal{S}'_{\lambda m}\right]\right\} = f^2_{\lambda m}\sin\theta. \tag{9}$$

The regular axis condition makes the lower boundary term vanish. The cap integral can therefore be recovered from the solution and its spectral sensitivity at θ_0, provided both use a consistent amplitude convention. This identity underlies the Legendre boundary norm and Mehler Riccati norm below. The zero-eigenvalue norm instead admits a positive-integral reduction. The identities are exact; their evaluation introduces truncation, integration, interpolation and rounding errors.

3.1 Non-integer-degree Legendre family: normalization and scale

We distinguish the standard Ferrers function, the Schmidt-scaled point function used by the R-SCHA interfaces, and the cap-unit basis. Let $\mathsf{P}_\nu^m(x)$ be the Ferrers function on $-1 < x < 1$ with the Condon–Shortley convention. The point evaluator returns

$$L_{\nu m}(\theta) = N_{\nu m}^{\mathrm{Sch}} \mathsf{P}_\nu^m(\cos\theta), \qquad N_{\nu m}^{\mathrm{Sch}} = (-1)^m \sqrt{\epsilon_m \frac{\Gamma(\nu - m + 1)}{\Gamma(\nu + m + 1)}}, \tag{10}$$

where $\epsilon_0 = 1$ and $\epsilon_m = 2$ for $m \geq 1$. The factor $(-1)^m$ cancels the Condon–Shortley sign, giving a positive regular polar amplitude. Cap normalization gives

$$\widehat{L}_{\nu m}(\theta) = \frac{L_{\nu m}(\theta)}{\sqrt{I_{\nu m}(\theta_0)}}, \qquad I_{\nu m}(\theta_0) = \int_0^{\theta_0} L_{\nu m}^2(\theta) \sin\theta \, \mathrm{d}\theta. \tag{11}$$

The public function `CapNormLegacySchmidt` returns $I_{\nu m}$, not $\sqrt{I_{\nu m}}$. Hereafter $L_{\nu m}$ denotes the Schmidt-scaled function; a hat denotes unit cap norm.

Substitution of the positive eigenvalue in Eq. (5) gives

$$L''_{\nu m} + \cot\theta L'_{\nu m} + \left[\nu(\nu+1) - m^2 \csc^2\theta\right] L_{\nu m} = 0. \tag{12}$$

With $\kappa = \nu + 1/2$ and $U = \sqrt{\sin\theta}\, L_{\nu m}$, its Liouville form is

$$U'' + Q(\theta; \kappa, m) U = 0, \qquad Q = \kappa^2 - (m^2 - 1/4)\csc^2\theta. \tag{13}$$

For $m > 0$, $Q = 0$ defines the turning angle $\theta_{\mathrm{tp}} = \arcsin[\sqrt{m^2 - 1/4}/\kappa]$ when $\sqrt{m^2 - 1/4} \leq \kappa$. Within $0 < \theta \leq \pi/2$, the interval below the turning point is nonoscillatory and the interval above it is oscillatory. Dirichlet cap degrees are the simple roots of $L_{\nu m}(\theta_0) = 0$; multiplication by the nonzero Schmidt factor changes neither their location nor their multiplicity. The primary domain considered here is integer $m < \nu < 10^5$ with $0 \leq \theta \leq \theta_0 \leq 90^\circ$; the equator is handled analytically.

The degree sensitivity required by the root and normalization calculations is $S_{\nu m} = \partial_\nu L_{\nu m}$, including the derivative of the Schmidt factor. Differentiating Eq. (12) at fixed (m, θ) gives the inhomogeneous problem

$$S''_{\nu m} + \cot\theta S'_{\nu m} + \left[\nu(\nu+1) - m^2 \csc^2\theta\right] S_{\nu m} = -(2\nu+1) L_{\nu m}. \tag{14}$$

For this family, $\partial_\lambda L_{\nu m} = S_{\nu m}/(2\nu+1)$. The common spectral identity, Eq. (9), therefore specializes to

$$\frac{\mathrm{d}}{\mathrm{d}\theta}\left\{\sin\theta \left[S_{\nu m} L'_{\nu m} - L_{\nu m} S'_{\nu m}\right]\right\} = (2\nu+1) L_{\nu m}^2 \sin\theta, \tag{15}$$

and integration from the axis to the cap boundary gives

$$I_{\nu m}(\theta_0) = \frac{\sin\theta_0}{2\nu+1}\left[L'_{\nu m}(\theta_0) S_{\nu m}(\theta_0) - L_{\nu m}(\theta_0) S'_{\nu m}(\theta_0)\right]. \tag{16}$$

The regular polar behavior makes the lower boundary term vanish. At a Dirichlet root, $L_{\nu m}(\theta_0) = 0$ and only $L'_{\nu m} S_{\nu m}$ remains; its sign must be positive because $I_{\nu m} > 0$. The same sensitivity gives the Newton correction $-L_{\nu m}/S_{\nu m}$, used in the selected root-solver regime described below. It is not the sensitivity of the cap-unit function, which at fixed θ_0 is

$$\partial_\nu \widehat{L}_{\nu m} = \frac{S_{\nu m}}{\sqrt{I_{\nu m}}} - \frac{L_{\nu m}\, \partial_\nu I_{\nu m}}{2 I_{\nu m}^{3/2}}. \tag{17}$$

The code computes $S_{\nu m}$ for the boundary norm, exposes its boundary value and uses it in selected Newton steps. The cap-unit sensitivity in Eq. (17) is not a public pointwise output.

3.2 Mehler family

For the negative spectrum, write $\lambda = -\Lambda$ with $\Lambda = \tau^2 + 1/4 > 0$. Cross-order relations require a fixed raw amplitude convention. We use

$$K_\tau^{m,\mathrm{raw}}(\theta) = \sin^m\theta\, {}_2F_1\left(m + \tfrac{1}{2} - \mathrm{i}\tau, m + \tfrac{1}{2} + \mathrm{i}\tau; m+1; z\right),$$
$$z = \sin^2(\theta/2), \qquad K_\tau^{m,\mathrm{raw}}(\theta) \sim \theta^m \quad (\theta \to 0). \tag{18}$$

For $m = 0$, the last condition means $K_\tau^{0,\mathrm{raw}}(0) = 1$. This real solution is proportional to the Ferrers function $\mathsf{P}^m_{-1/2+\mathrm{i}\tau}(\cos\theta)$. Euler's transformation gives the equivalent positive-series representation

$$K_\tau^{m,\mathrm{raw}}(\theta) = [2\tan(\theta/2)]^m\, {}_2F_1\left(\tfrac{1}{2} - \mathrm{i}\tau, \tfrac{1}{2} + \mathrm{i}\tau; m+1; z\right). \tag{19}$$

Both series have positive coefficients for $0 \le z < 1$. Their fixed raw convention is required for cross-order identities, whereas a positive angle-independent multiplier cancels from cap normalization.

Writing $K = K_\tau^{m,\mathrm{raw}}$, the angular equation is

$$K_{\theta\theta} + \cot\theta K_\theta - \left[\Lambda + m^2\csc^2\theta\right] K = 0. \tag{20}$$

The half-angle coordinate $t = \log\tan(\theta/2)$, for which $\sin\theta = \operatorname{sech} t$, removes the first derivative:

$$K_{tt} = \left[m^2 + \Lambda\operatorname{sech}^2 t\right] K. \tag{21}$$

The regular polar solution has $K > 0$ and $K_t > 0$ at finite t. Since the coefficient in Eq. (21) is positive, neither can acquire a first zero. Unlike the Legendre solution, this branch has no turning point or oscillatory interval, so its logarithmic derivative stays regular on a finite cap. Tests cover $m, \tau \le 10^4$ and $0 \le \theta \le \theta_0 \le 90^\circ$.

3.3 Zero-eigenvalue constant family

For $\lambda = 0$, the regular angular shape is $\tan^m(\theta/2)$. This is distinct from the integer-degree Ferrers function P^m_0, which vanishes for integer $m \ge 1$. We denote the regularized R-SCHA branch by

$$C_0^{\mathrm{raw}}(\theta) = 1, \qquad C_m^{\mathrm{raw}}(\theta) = \Gamma(m)\tan^m\frac{\theta}{2}, \quad m \ge 1. \tag{22}$$

This is the function customarily called the Thébault constant function in the R-SCHA literature (Thébault et al., 2004, 2006). Its multiplicative gamma factor cancels after cap normalization. For $m = 0$, $\widehat{C}_0 = (1 - \cos\theta_0)^{-1/2}$; for $m \ge 1$, Section 4.4 supplies a cancellation-free norm. The tested order range is $m \le 10^4$.

4 Stable numerical realization of the angular basis

Each evaluator preserves the regular polar branch and its prescribed amplitude convention, and recovers derivatives from their own scaled states. Legendre point and sensitivity calls may use different numerical routes but return quantities on the same Schmidt scale. Mehler and zero-eigenvalue point calls include cap normalization. We first describe the family algorithms, then the shared table construction and lookup layer.

4.1 Interfaces, units, and extended-range records

Table 1 lists the public conventions. Legendre point calls return Schmidt-scaled quantities; Mehler and constant-family calls return cap-unit quantities. The basis layer can be used independently of the full inverse or forward model.

Table 1: Implemented public conventions. A prime denotes differentiation with respect to the radian argument, even when the wrapper receives an angle in degrees.

Family/interface	Point output	Input angle	Additional public quantity
Legendre `Legendre_2rank`	$L_{\nu m}, L'_{\nu m}, L''_{\nu m}$ on the legacy Schmidt scale	degrees	squared norm $I_{\nu m}$ and boundary sensitivity $S_{\nu m}(\theta_0)$ (radian cap angle)
Mehler `Mehler_1rank`, `Mehler_2rank`	$\widehat{K}, (\widehat{K})', (\widehat{K})''$	degrees for θ and θ_0	cap normalization is included
Constant `Constant_1rank`, `Constant_2rank`	$\widehat{C}, (\widehat{C})', (\widehat{C})''$	degrees for θ and θ_0	cap normalization is included

Internal records retain ordinary values, component signs and logarithmic magnitudes, derivative-to-value ratios, range flags and status codes. Compatibility wrappers return ordinary `double` values; validation accesses the full records. Each derivative is formed from its own logarithm, so an underflowed value can coexist with a normal-range derivative. Flushed or rounded subnormals do not replace the mathematical logarithmic state. A range flag thus distinguishes a valid out-of-range quantity from evaluation failure. The namespace `DegreeTrig` handles degree-valued trigonometry; coordinate rotation is separate. The two additional Legendre quantities listed in the first row are exposed by `CapNormLegacySchmidt` and `BoundaryDegreeDerivativeLegacySchmidt`, respectively.

4.2 Legendre evaluation

Every Legendre representation computes the Schmidt-scaled $L_{\nu m}$ of Eq. (10), with angular derivatives in radians. Series, recurrences, propagation and phase methods cover overlapping regimes. Their domain and consistency checks determine which representation is usable (Table 2). Atlas lookups provide residual corrections within these representations; they do not define another basis.

Table 2: Legendre representations and their numerical roles. The regions overlap; these rows describe method families rather than a universal dispatch order.

Local structure	Representation	Quantity checked
Regular polar solution	Differentiated hypergeometric series; convergent high-order polar form	Term decay, convergence, and cancellation
Recessive near-order solution	Endpoint/Laplace anchors, scaled recurrence, or carrier-removed implicit propagation	Anchor domain, recurrence consistency, or mesh refinement
Turning and oscillatory solution	Matched direct propagation; Airy, Liouville–Green, or corrected WKB carrier	Propagation success; carrier defect or atlas cell acceptance
Near-equatorial solution	Exact gamma data with short propagation; coalescent WKB phase difference	Finite analytic data, propagation domain, and phase defect

4.2.1 Regular and recessive representations

Near the pole, remove the known vanishing factor and write

$$\mathcal{L}_{\kappa m}(\theta) = \sin^m \theta \sum_{j=0}^{\infty} c_j z^j, \qquad z = \sin^2 \frac{\theta}{2}, \tag{23}$$

with $c_0 = 1$ and

$$c_j = c_{j-1} \frac{(m + j - 1/2)^2 - \kappa^2}{j(m + j)}. \tag{24}$$

The physical scale is restored by $L_{\nu m} = A_{\nu m} \mathcal{L}_{\kappa m}$, where

$$A_{\nu m} = \frac{1}{2^m \Gamma(m + 1)} \sqrt{\epsilon_m \frac{\Gamma(\nu + m + 1)}{\Gamma(\nu - m + 1)}}. \tag{25}$$

The Ferrers definition and Schmidt factor determine $A_{\nu m}$, evaluated logarithmically. An equivalent form is useful at high order:

$$L_{\nu m} = 2^m A_{\nu m} \tan^m \frac{\theta}{2} H(z), \qquad H(z) = {}_2F_1(-\nu, \nu + 1; m + 1; z). \tag{26}$$

Its term ratio is $z(j - 1 - \nu)(j + \nu)/[j(m + j)]$. In the convergence-screened high-order polar region, the implementation differentiates this series for both angular derivatives. A large degree need not force an asymptotic route if the local series is well conditioned.

When $d = \nu - m$ is small relative to m, endpoint and Laplace-moment representations remove the dominant recessive factor. The moment expansion is used only in its small-moment domain while terms decrease. Scaled order recurrence can otherwise continue from admissible anchors. A near-equatorial evanescent route writes $L = \text{const.} \sin^m \theta \, G$ and propagates $v = \partial_t \log G$:

$$v_t = 2m \tanh t \, v - v^2 - d(\nu + m + 1) \operatorname{sech}^2 t. \tag{27}$$

A two-stage singly diagonally implicit Runge–Kutta scheme uses mesh refinement and Richardson extrapolation of second-order solutions. This route is restricted to the nonoscillatory branch and does not cross zeros.

Removing a carrier is useful only if the remaining scale can be computed accurately. In the endpoint recurrence, put $b = m + 1$ and evaluate the balanced logarithm

$$\log(\sin \theta) + \log\left(2 + \frac{d - 1}{b}\right) - \log 2, \tag{28}$$

from positive factors. This avoids amplifying an endpoint subtraction error by a large order. Double-double arithmetic carries the scale through anchor normalization, recurrence rescaling, the final Schmidt value and the adjacent-order ratio. It preserves cancellation in the numerical scale without changing the normalization.

4.2.2 Turning, oscillatory, and equatorial representations

In the half-angle coordinate, the Legendre equation becomes

$$L_{tt} + \left[(\kappa^2 - 1/4) \operatorname{sech}^2 t - m^2\right] L = 0 \tag{29}$$

Within the zero-free interval before the first zero, it is propagated through its logarithmic derivative $r = L_t/L$. With $h = \partial_\kappa r$,

$$r_t = m^2 - (\kappa^2 - 1/4) \operatorname{sech}^2 t - r^2, \tag{30}$$

$$h_t = -2\kappa \operatorname{sech}^2 t - 2rh. \tag{31}$$

The polar series fixes the regular branch. Before reaching a Riccati pole, the calculation matches to a state or phase representation.

In the oscillatory sector, a nonoscillatory phase α for $U = \sqrt{\sin\theta}\, L$ satisfies Kummer's equation,

$$Q - (\alpha')^2 - \frac{1}{2}\frac{\alpha'''}{\alpha'} + \frac{3}{4}\left(\frac{\alpha''}{\alpha'}\right)^2 = 0, \tag{32}$$

and gives

$$U(\theta) = C\frac{\sin[\alpha(\theta) + \delta]}{\sqrt{\alpha'(\theta)}}. \tag{33}$$

The phase approach follows Bremer and Rokhlin (2017) and Bremer (2018). The implementation constructs Liouville–Green, Airy and corrected WKB carriers. An available atlas cell interpolates the residual correction; domain, defect or sampled-cell checks must pass. Direct phase integration, endpoint propagation and order recurrence remain available when a carrier is unsuitable.

Near the equator, exact endpoint data avoid choosing the wrong oscillatory branch or subtracting two large phases. Define

$$a_m = \frac{\nu + m + 2}{2}, \qquad b_m = \frac{1 - \nu + m}{2}. \tag{34}$$

The Schmidt-scaled equatorial value is

$$\begin{aligned}\log|L_{\nu m}(\pi/2)| = {} & \frac{1}{2}\log\epsilon_m + \frac{1}{2}[\log\Gamma(\nu + m + 1) - \log\Gamma(\nu - m + 1)] \\ & - m\log 2 + \frac{1}{2}\log\pi - \log\Gamma(a_m) - \log|\Gamma(b_m)|\,, \end{aligned} \tag{35}$$

$$\operatorname{sgn} L_{\nu m}(\pi/2) = \operatorname{sgn}\Gamma(b_m). \tag{36}$$

At a pole of $\Gamma(b_m)$, the reciprocal-gamma limit gives zero. Reflection formulas reduce the endpoint data to sine factors and positive gamma ratios. Their derivatives remain finite at value zeros. Away from those zeros, with $\psi = \Gamma'/\Gamma$,

$$q_m^{(0)} = \partial_\nu \log|L_{\nu m}(\pi/2)| = \frac{1}{2}\left[\psi(\nu + m + 1) - \psi(\nu - m + 1) - \psi(a_m) + \psi(b_m)\right], \tag{37}$$

so that $S_{\nu m}(\pi/2) = q_m^{(0)} L_{\nu m}(\pi/2)$. The product is evaluated by the differentiated reflection formula at a zero, not by multiplying a zero by a divergent logarithmic derivative.

For $m \geq 1$ and $\nu \geq m + 1$, put $\rho_{\nu m} = \sqrt{(\nu - m)(\nu + m + 1)}$. The adjacent-order identity gives

$$L'_{\nu m}(\pi/2) = -\rho_{\nu m} L_{\nu,m+1}(\pi/2), \tag{38}$$

$$S'_{\nu m}(\pi/2) = -\frac{2\nu + 1}{2\rho_{\nu m}} L_{\nu,m+1}(\pi/2) - \rho_{\nu m} S_{\nu,m+1}(\pi/2). \tag{39}$$

The $m = 0$ case uses separate exact endpoint formulas. Short propagation from the equator resolves nearby intervals. If the turning point approaches the equator, a corrected WKB route integrates the local phase density from the exact equatorial anchor; using a phase difference avoids subtracting large actions. These point-evaluation routes retain their domain and defect checks. The four-state equatorial sensitivity seed requires $\nu - m \geq 1$; smaller gaps use regular polar data. This restriction applies to the sensitivity route, not to the equatorial value formula.

4.2.3 Scale matching and degree sensitivity

Proportional solutions have different norms and degree derivatives unless their scale is fixed. The polar coefficient $c_0 = 1$ and amplitude $A_{\nu m}$ fix that scale. Matching L and L_θ at a common angle fixes the continuation by uniqueness. For a phase match, define

$$y_* = \alpha'(\theta_*)L(\theta_*), \qquad x_* = L_\theta(\theta_*) + \frac{1}{2}\left[\cot\theta_* + \frac{\alpha''(\theta_*)}{\alpha'(\theta_*)}\right] L(\theta_*). \tag{40}$$

Then

$$\alpha(\theta_*) + \delta = \operatorname{atan2}(y_*, x_*) \pmod{2\pi}. \tag{41}$$

The $\cot\theta_*$ term accounts for $U = \sqrt{\sin\theta}\, L$. The signed pair fixes the phase quadrant, and Eq. (33) then fixes C. The exact equatorial data provide an alternative anchor on the same Schmidt scale.

Degree sensitivity is propagated analytically, rather than obtained by subtracting neighboring-degree values. Differentiating the polar recurrence gives

$$\dot{c}_0 = 0, \tag{42}$$

$$\dot{c}_j = \dot{c}_{j-1}\frac{(m+j-1/2)^2 - \kappa^2}{j(m+j)} - c_{j-1}\frac{2\kappa}{j(m+j)}, \tag{43}$$

where a dot denotes $\partial_\nu = \partial_\kappa$. The scale derivative is included:

$$\partial_\nu \log A_{\nu m} = \frac{1}{2}\left[\psi(\nu+m+1) - \psi(\nu-m+1)\right]. \tag{44}$$

Thus S differentiates the Schmidt-scaled function, including its amplitude. In a zero-free interval, $q_\nu = \partial_\nu \log|L|$ satisfies $(q_\nu)_t = h$, giving

$$S = q_\nu L, \qquad S_\theta = \frac{L}{\sin\theta}(h + r q_\nu). \tag{45}$$

Across turning and oscillatory intervals, the coupled physical state satisfies

$$\frac{\mathrm{d}}{\mathrm{d}\theta}\begin{bmatrix} L \\ L' \\ S \\ S' \end{bmatrix} = \begin{bmatrix} L' \\ -\cot\theta\, L' - D_{\nu m} L \\ S' \\ -\cot\theta\, S' - D_{\nu m} S - (2\nu+1)L \end{bmatrix}, \qquad D_{\nu m} = \nu(\nu+1) - m^2\csc^2\theta. \tag{46}$$

Differentiated downward-order recurrence is used when it avoids a long or poorly conditioned propagation. It inherits the scale of the differentiated regular anchors. Temporary rescaling divides all four components by a common factor recorded separately: the Newton ratio L/S is unchanged, and restoring the squared factor recovers the boundary norm. Arbitrary-angle degree sensitivity is internal; public point calls expose angular derivatives.

4.2.4 Point evaluation, normalization, and roots

Point and sensitivity calculations share a mathematical scale but can use different numerical routes. Algorithms 1 and 2 summarize their flow.

Algorithm 1. Schmidt-scaled point evaluation. Validate the order, degree, and angle; evaluate exact axis limits when applicable. Select a locally admissible representation from Table 2, with convergent regular forms preferred where they are well conditioned. In the bounded-degree small-angle regime $\nu + 1/2 < 10^4$, $\theta \le \pi/6$, try the scaled downward order recurrence before polar propagation. If the later asymptotic paths are inadmissible, also try that recurrence for $\nu < 10^4$ before constructing a

phase initial-value solution. For a tabulated or asymptotic candidate, apply its domain and cell, defect, or recurrence check. If the candidate fails, try another admissible representation or direct propagation. Retain the differentiated series second derivative in the regular polar neighborhood; elsewhere reconstruct $L_{\theta\theta}$ from Eq. (12). Return component-wise signed logarithms and range metadata together with ordinary values. Failure of all admissible paths is reported as a numerical failure.

The downward recurrence starts from regular polar states at orders $\lfloor\nu\rfloor$ and $\lfloor\nu\rfloor - 1$, preserving a common logarithmic scale. For $\nu < 10^4$, compensated log-gamma evaluation supplies the seed normalization with work independent of order, instead of summing one term per order. The recurrence itself still requires work proportional to its order span, and the seed series must satisfy its convergence criterion. This change removes redundant normalization work without making the full point evaluator a constant-cost procedure. The differentiated WKB construction likewise retains only the Taylor coefficients needed by the remaining derivative steps; its acceptance criteria are unchanged.

Algorithm 2. Degree sensitivity, cap normalization, and root refinement. Construct $(L, L_\theta, S, S_\theta)$ from a differentiated regular anchor, exact equatorial sensitivity data in its allowed domain, or differentiated downward-order recurrence. Propagate with short-segment rescaling when needed. Recover $I_{\nu m}$ from the full boundary bilinear identity. For $\kappa\theta_0 \le 1.5$, integrate the square of the polar series with the $\sin\theta$ measure; additional integral or logarithmically scaled routes are available if the boundary calculation is unusable. Predict a boundary zero from the action described below, establish a computed sign bracket, and refine it with a safeguarded solver. Root-array interpolation supplies starting values only; each returned entry is subsequently checked with the public point evaluator.

The root predictor uses the exact Liouville turning point, distinct from the leading atlas carrier. For $m \ge 1$, define

$$A_{\mathrm{root}}(\kappa, m, \theta_0) = \int_{\arcsin(\beta_m/\kappa)}^{\theta_0} \sqrt{\kappa^2 - \beta_m^2 \csc^2 s}\, \mathrm{d}s, \qquad \beta_m = \sqrt{m^2 - \tfrac{1}{4}}. \tag{47}$$

Set the action to zero when $\kappa \le \beta_m$ or the boundary precedes the turning point. The leading predictor solves $A_{\mathrm{root}}(\kappa_{\mathrm{pred}}, m, \theta_0) = \pi(j - 1/4)$; for $m = 0$, it uses $\kappa_{\mathrm{pred}}\theta_0 = \pi(j - 1/4)$. The action already includes the degree factor. These conditions predict roots; numerical bracketing and refinement determine acceptance.

The standalone root interface normally refines a computed sign bracket with an Illinois-weighted signed-log secant and periodic bisection. A safeguarded Newton step using S is used only for $m \le 1$, $j \ge 100$, and $\theta_0 \le 2°$. When a near-zero value cannot be materialized, the signed-log solver requires a bracket width no greater than $64\epsilon_{\mathrm{mach}} \max(1, |\nu_-|, |\nu_+|)$ before accepting the result. For triangular root arrays, the first 32 roots and every subsequent 16th root provide explicitly refined anchors; a local interpolation in $1/j$, using at most eight anchors, predicts the remaining corrections. Every entry is then re-evaluated with the public evaluator and refined in a computed sign bracket, or accepted at a computed zero; increasing root order is checked before storage. These checks are floating-point consistency checks, not interval-arithmetic certification.

A computed zero from the scaled recurrence is retained when the adjacent-order state gives a finite, nonzero angular derivative. The raising identity then gives L_θ, and the ODE gives $L_{\theta\theta} = -\cot\theta\, L_\theta$. Their separate signed logarithms avoid division by the zero value. The root

solver can accept this floating-point zero, subject to the residual tests below; it is not an exact-zero certificate.

Normalization uses the full boundary bilinear form, including near a root. The term involving $L(\theta_0)$ vanishes only at an exact Dirichlet zero. Scaled products and logged amplitudes retain both terms outside the ordinary range; an accumulated integral provides an alternative where implemented. An exhausted refinement budget returns failure rather than a converged root.

4.2.5 Conditioning and error control

These checks address different error sources and do not provide a global forward-error bound. For a polar sum F with tail estimate R_J, the truncation contribution scales as $R_J/|F|$: term decay alone cannot resolve cancellation. Differentiated recurrences avoid finite-difference subtraction, while log-gamma ratios and double-double arithmetic reduce amplitude cancellation.

For logarithmic propagation, a perturbation obeys

$$(\delta r)_t = -2r\,\delta r + \delta V + O(\delta r^2), \tag{48}$$

Here δV includes coefficient and integration errors. Where $r > 0$, forward propagation damps perturbations of the regular branch. This applies only before a zero. A computed phase with Kummer residual R_K is exact for the perturbed coefficient $Q - R_K$; the defect measures a coefficient perturbation, not relative value error near a zero. There, signed phase, scaled residuals and Newton corrections are better diagnostics.

The $\csc^2\theta$ coefficient also makes unconditional ODE reconstruction of the second derivative unsuitable near the axis. Exact axis limits and differentiated series resolve that removable singularity. For a term $T_j = a_j z T_{j-1}$, the stable recurrences

$$T_{j,z} = a_j(zT_{j-1,z} + T_{j-1}), \qquad T_{j,zz} = a_j(zT_{j-1,zz} + 2T_{j-1,z}) \tag{49}$$

avoid division of underflowed terms by z or z^2. Scale preservation is also needed for matching and normalization, although cancellation can remain in their bilinear expressions. Representation checks and the tests in Section 6 assess these remaining errors.

4.3 Mehler normalization from a positive Riccati flow

The zero-free Mehler branch permits its shape and scale to be separated without crossing a pole. Define

$$q = \frac{K_t}{K}, \qquad s = \partial_\Lambda q, \qquad \ell = \log K. \tag{50}$$

Equation (21) then gives

$$q_t = m^2 + \Lambda\,\mathrm{sech}^2\, t - q^2, \tag{51}$$

$$s_t = \mathrm{sech}^2\, t - 2qs, \qquad \ell_t = q, \tag{52}$$

with $q \to m$ and $s \to 0$ as $t \to -\infty$. The scale-invariant variables q and s determine the shape and norm relative to the boundary amplitude; ℓ carries the amplitude.

Proposition 1 (Riccati-sensitivity normalization)**.** *Let K be a positive regular solution of Eq. (21), and let $s = \partial_\Lambda(K_t/K)$ satisfy the polar condition above. At every finite t, $s(t) > 0$. For $t_0 = \log\tan(\theta_0/2)$,*

$$\|K\|^2_{\mathrm{cap}} = K^2(t_0)s(t_0), \qquad \widehat{K}^m_\tau(\theta) = \frac{\exp\left[\int_{t_0}^{t(\theta)} q(\xi)\,\mathrm{d}\xi\right]}{\sqrt{s(t_0)}}. \tag{53}$$

Proof. Regularity gives $K(t) = ce^{mt}[1 + O(e^{2t})]$ for $m > 0$ and $K(t) = c[1 + O(e^{2t})]$ for $m = 0$, with $c > 0$. The positive coefficient in Eq. (21) preserves $K > 0$ and $K_t > 0$, so q is well defined. Multiplying the sensitivity equation by K^2 yields

$$\frac{\mathrm{d}}{\mathrm{d}t}(K^2 s) = K^2 \operatorname{sech}^2 t. \tag{54}$$

The regular polar expansion gives $K^2 s \to 0$ as $t \to -\infty$. Consequently,

$$K^2(t)s(t) = \int_{-\infty}^{t} K^2(\xi) \operatorname{sech}^2 \xi \, \mathrm{d}\xi > 0. \tag{55}$$

Since $\sin\theta \, \mathrm{d}\theta = \operatorname{sech}^2 t \, \mathrm{d}t$, the integral at t_0 is exactly the squared cap norm. The ratio $K(t)/K(t_0) = \exp[\int_{t_0}^{t} q \, \mathrm{d}\xi]$ proves the normalized expression and removes c. □

Equation (53) is the negative-spectrum form of the Lagrange identity. It removes the raw amplitude and the need for a separate norm quadrature.

Conditioning of the propagated state. Two exact solutions of the Riccati equation have difference $e = q_1 - q_2$ satisfying

$$e(t) = e(t_a) \exp\left[-\int_{t_a}^{t} (q_1 + q_2) \, \mathrm{d}\xi \right]. \tag{56}$$

Forward flow damps initialization error while both solutions remain positive. Discretization and interpolation errors still enter: if $\widetilde{q}$ has differential defect d, then $e_t = -(\widetilde{q} + q)e + d$. The defect accumulates through the variation-of-constants integral. To first order,

$$\delta \log \widehat{K}(t) = \delta\ell(t) - \delta\ell(t_0) - \frac{\delta s(t_0)}{2s(t_0)}. \tag{57}$$

The log-amplitude difference and boundary sensitivity therefore require their own error controls; contraction of q does not certify the normalized value.

Offline construction. The polar state can be initialized reproducibly by setting $u = e^{2t}$ and expanding

$$\operatorname{sech}^2 t = \sum_{j=1}^{\infty} d_j u^j, \qquad d_j = 4(-1)^{j-1} j. \tag{58}$$

With $q = m + \sum_{j\geq 1} a_j u^j$ and $s = \sum_{j\geq 1} b_j u^j$, substitution gives

$$a_j = \frac{\Lambda d_j - \sum_{k=1}^{j-1} a_k a_{j-k}}{2(j+m)}, \tag{59}$$

$$b_j = \frac{d_j - 2\sum_{k=1}^{j-1} a_k b_{j-k}}{2(j+m)}. \tag{60}$$

The raw series in Eq. (18) supplies the log-amplitude anchor at the same small polar angle.

To reduce the scale of the propagated quantities, let $p(t) = [m^2 + \Lambda \operatorname{sech}^2 t]^{1/2}$ and choose an analytic carrier C with $C_t = p$. The builder advances $r = q - p$, s, and $\ell - C$:

$$r_t = -2pr - r^2 - p_t, \tag{61}$$

$$s_t = \operatorname{sech}^2 t - 2(p + r)s, \qquad (\ell - C)_t = r. \tag{62}$$

Implicit midpoint steps are compared with two half steps and corrected by Richardson extrapolation. The scaled local-error estimator uses relative tolerance 3×10^{-14}, absolute tolerance 3×10^{-17} for r and s, and 3×10^{-16} for the log-amplitude residual. These are construction controls, not bounds on the final interpolated error. Only the two slowly varying fields $\ell - C$ and $\log s$ are stored in the fixed tensor atlas described in Section 4.5.

Online reconstruction. At the requested point and cap boundary, the stored fields recover $\ell(t)$, $\ell(t_0)$, and $s(t_0)$. The normalized value is formed directly in logarithmic form,

$$\log \widehat{K}(t) = \ell(t) - \ell(t_0) - \tfrac{1}{2} \log s(t_0). \tag{63}$$

Differentiating the first Chebyshev field restores $q = p + \partial_t(\ell - C)$; the angular derivatives follow from the same state and the differential equation:

$$\widehat{K}_\theta = \frac{q}{\sin\theta} \widehat{K}, \tag{64}$$

$$\widehat{K}_{\theta\theta} = -\cot\theta \widehat{K}_\theta + (\Lambda + m^2 \csc^2\theta)\widehat{K}. \tag{65}$$

Online evaluation solves neither a Riccati initial-value problem nor a norm quadrature. Section 4.5 specifies the local-series routes, endpoints and table domains.

4.4 Cancellation-free zero-eigenvalue normalization

For the zero-eigenvalue family with $m \geq 1$, the angular shape is analytic; only its cap norm needs numerical treatment. Let $u = \tan(\theta/2)$, $u_0 = \tan(\theta_0/2)$, and $\chi = u_0^2$. Removing the angle-independent raw amplitude, the squared cap norm of the boundary-scaled shape $(u/u_0)^m$ is

$$I_m^{\text{shape}} = \frac{4}{u_0^{2m}} \int_0^{u_0} \frac{u^{2m+1}}{(1+u^2)^2}\, \mathrm{d}u = 2\chi J_m(\chi), \tag{66}$$

where

$$J_m(\chi) = \int_0^1 \frac{x^m}{(1+\chi x)^2}\, \mathrm{d}x > 0, \qquad m\Phi(-\chi, 1, m) - \frac{1}{1+\chi} = \chi J_m(\chi). \tag{67}$$

Integration by parts in the defining Lerch integral gives the second identity. Its left side subtracts nearly equal terms at large m; the right side evaluates the required positive quantity directly.

Set $a = \chi/(1+\chi)$. Expanding about the positive denominator at $x = 1$ and integrating each term gives

$$J_m(\chi) = \frac{1}{(1+\chi)^2} \sum_{k=0}^{\infty} \frac{(k+1)!a^k}{(m+1)(m+2)\cdots(m+k+1)}. \tag{68}$$

The terms are generated without subtraction:

$$T_0 = \frac{1}{(1+\chi)^2(m+1)}, \qquad T_{k+1} = T_k \frac{(k+2)a}{m+k+2}. \tag{69}$$

For $m \geq 1$, $T_{k+1}/T_k \leq a$, so the tail after T_K is at most $T_{K+1}/(1-a)$. On the reported hemispheric domain, $\chi \leq 1$ and hence $a \leq 1/2$: convergence is rapid and the tail estimate remains well conditioned. Although the series converges for every finite χ, this observation does not imply uniformly bounded cost as $\theta_0 \to \pi$.

The builder sums the positive series in its `long double` arithmetic until the tail estimate is below 2×10^{-19} of the partial sum. This is a truncation criterion, not a rounding-error bound. It stores $\frac{1}{2}\log[2\chi J_m(\chi)]$. Online calls interpolate this field or, for caps below $0.01°$, use its first four

positive terms. Table domains and degrees appear in Section 4.5; an unsupported lookup returns failure.

Once the norm is known, the value and angular derivatives retain their analytic form:

$$\widehat{C}_m(\theta) = \left(\frac{u}{u_0}\right)^m [2\chi J_m(\chi)]^{-1/2}, \tag{70}$$

$$\widehat{C}'_m = \frac{m}{\sin\theta}\widehat{C}_m, \tag{71}$$

$$\widehat{C}''_m = \frac{m(m-\cos\theta)}{\sin^2\theta}\widehat{C}_m. \tag{72}$$

The value is assembled from $m(\log u - \log u_0) - \frac{1}{2}\log[2\chi J_m(\chi)]$, and each derivative is materialized from its own logarithmic magnitude. For $m = 1$, the second-derivative factor is evaluated as $1/(1+\cos\theta)$ to avoid polar cancellation. The removable axis limits are supplied explicitly: the first derivative can be nonzero only for $m = 1$, and the second derivative only for $m = 2$. The $m = 0$ mode uses the exact cap measure $1 - \cos\theta_0$ and has zero angular derivatives. Thus numerical approximation enters only through the scalar norm, while the angular dependence and endpoint limits remain analytic.

4.5 Residual compression: offline construction and online recovery

The table layer compresses residuals and logarithmic norms after removing rapid phase and amplitude variation. It does not interpolate the raw oscillatory function. Offline construction generates numerical source fields; online evaluation restores their carriers and normalization.

4.5.1 Carrier coordinates and stored fields

For the Legendre charts, set $u = \nu + 1/2$ and $\eta = m/u$. The Liouville coefficient can be written as

$$Q(\theta) = u^2 F_\eta(\theta) + \tfrac{1}{4}\csc^2\theta, \qquad F_\eta(\theta) = 1 - \eta^2\csc^2\theta.$$

Thus $\theta_c = \arcsin\eta$, for $0 < \eta < 1$, is the leading-carrier turning point; the residual includes the $\frac{1}{4}\csc^2\theta$ correction to Q. Define

$$A = uJ, \qquad J = \begin{cases} \displaystyle\int_\theta^{\theta_c} \sqrt{-F_\eta(s)}\,\mathrm{d}s, & 0 < \theta \le \theta_c, \\ \displaystyle\int_{\theta_c}^{\theta} \sqrt{F_\eta(s)}\,\mathrm{d}s, & \theta_c \le \theta \le \pi/2, \end{cases} \qquad \zeta = \operatorname{sgn}(F_\eta)(3J/2)^{2/3}. \tag{73}$$

At θ_c, $\zeta = 0$. The Airy argument $-u^{2/3}\zeta$ is negative in the oscillatory interval and positive in the recessive interval. This carrier action differs from the root predictor based on the exact Q.

On the oscillatory side, the Airy-core charts use $(u^{-1}, \eta, (3A/2)^{2/3})$ for $0 \le A \le 7.5$; the outer charts use $(u^{-1}, \eta, \log A)$ for $7.5 < A \le 400$. For a consistently scaled numerical fundamental pair $W = W_1 + iW_2$ and its Airy carrier $B = B_1 + iB_2$, the stored fields are $u^2[\mathrm{Re}(W/B) - 1]$ and $u^2\,\mathrm{Im}(W/B)$. Removing both the leading unit multiplier and its u^{-2} decay makes these fields suitable for interpolation. Separate degree slabs cover $512 \le \nu \le 10^6$ and $10^6 < \nu \le 10^8$, with additional refinement in intermediate action wedges and with cells truncated to the available angular interval. Farther into the oscillatory sector, the analytic WKB route remains available.

The near-order chart instead uses $(u^{-1}, \nu - m, \log(1 + A))$ on the recessive side. Its field is the difference between the numerical logarithm and that of the recessive Airy carrier, evaluated at the same internal scale. The double-double endpoint formula and scaled recurrence generate the numerical source; they are not the carrier subtracted from it. Candidate cells are constructed within $10^4 \le \nu \le 10^6$, $0 \le \nu - m \le 1024$, and $0 \le A \le 150$. A small direct chart in $(m, \nu - m, \theta)$

covers $m, \nu - m \le 8$; outside its domain, the low-degree evaluator retains the non-tabulated kernels.

For Mehler functions, the stored fields are $\rho = \log K_{\mathrm{raw}} - C$ and $\log s$, where C is the analytic carrier introduced in the Riccati construction. For the constant family, only $\ell_N = \frac{1}{2}\log(2\chi J_m(\chi))$ is tabulated; the angular shape and its derivatives remain analytic. The fixed grids used for these two positive families are summarized in Table 3.

Table 3: Fixed table domains for the positive families. Angular limits are given in degrees; transformed coordinates use radians.

Family	Domain	Coordinates and fields	Cells; degrees
Mehler	$0 \le m, \tau \le 10^4$; $0.01^\circ \le \theta \le 90^\circ$	$(\log(1+m), \log(1+\tau), t)$; ρ, $\log s$	2025; $(10, 14, 14)$
Constant	$1 \le m \le 10^4$; $0.01^\circ \le \theta_0 \le 90^\circ$	$(\log(1+m), \log\tan(\theta_0/2))$; ℓ_N	24; $(12, 12)$

4.5.2 Offline approximation and its acceptance scope

Within a cell, each field has the tensor-product expansion

$$R(\boldsymbol{x}) = \sum_{i=0}^{d_1}\sum_{j=0}^{d_2}\sum_{k=0}^{d_3} c_{ijk} T_i(\widehat{x}_1) T_j(\widehat{x}_2) T_k(\widehat{x}_3), \tag{74}$$

with affine local coordinates $\widehat{x}_l \in [-1, 1]$; the constant norm uses the corresponding two-dimensional form. Coefficients are computed at tensor Chebyshev–Lobatto nodes. Legendre source values are supplied by direct, phase, or asymptotic kernels in their respective domains; the positive-family sources are generated by the Riccati and positive-series constructions above. These are numerical approximations produced by the same framework, rather than independent exact references. Nested Clenshaw recurrences evaluate the stored fields and, where required, their derivatives.

The near-order section has an additional sampling screen. If E_{samp} is the maximum discrepancy between the interpolant and its generating kernel on a fixed check grid, including points on every face, edge, and corner, the stored indicator is $8E_{\mathrm{samp}} + 2 \times 10^{-11}$. A lookup is eligible only if this indicator does not exceed 4×10^{-11}. All 2880 near-order candidate cells in the present atlas exceed this threshold. This section therefore provides no interpolation acceleration in the present implementation: near-order calls use the endpoint formula and scaled recurrence, including beyond the candidate table range. The screen measures sampled compression discrepancy against the generating kernel; it neither certifies the source error nor gives a uniform bound on the reconstructed function. The independent accuracy assessment is reported in Section 6.

4.5.3 Online recovery and endpoint branches

A Mehler lookup restores $\log K_{\mathrm{raw}} = C + \rho$, $q = C_t + \rho_t$, and $s = \exp(\log s)$. Applying these fields at t and t_0 gives the Riccati-sensitivity normalization without online integration. The zero-order raw amplitude uses the fixed amplitude convention of the Mehler definition and the exact identity

$$K^0_{\mathrm{raw}}(\theta) = \cot\theta\, K^1_{\mathrm{raw}}(\theta) + \frac{\Lambda + 2}{8} K^2_{\mathrm{raw}}(\theta). \tag{75}$$

The implementation first obtains the zero-order table state, then evaluates the two adjacent orders to reconstruct its raw amplitude. No long order-recurrence chain is used.

Endpoint and convergent-series branches avoid unnecessary interpolation. For $\Lambda \le 16(m+1)$, both point and boundary data are obtained from the positive hypergeometric series in Eq. (19). Denote its hypergeometric factor by $H_M(z)$. The coefficient ratio is $z[j(j-1) + \Lambda]/[j(m+j)]$.

Differentiating the same recurrence gives $s = \frac{1}{2}\sin^2\theta\,\partial_\Lambda(H_{M,z}/H_M)$. Value and differentiated tail tests must both pass within 256 terms; failure returns an invalid status. For the remaining modes, a fixed 64-term polar expression supplies point or boundary data through 0.1°, with analytic axis limits.

For $m > 1$ and $0 < \theta < 10^{-150}$ radians, the Mehler second derivative is recovered directly from its independent logarithm,

$$\log K_{\theta\theta} = \log K - 2\log\theta + \log[m(m-1)] + \log\bigl(1 + O(\theta^2)\bigr). \tag{76}$$

The implemented leading polar expression omits a relative correction below 10^{-292} within the admitted $m, |\tau| \le 10^4$ range. It avoids materializing $m(m-1)/\sin^2\theta$, which can overflow even when $\sin^2\theta$ is a nonzero subnormal number. The normalized mode's logarithm is used, so an underflowed ordinary amplitude does not force an underflowed derivative. Orders zero and one retain their separate polar expressions and axis limits.

At a hemispheric boundary, set $a = (m + 3/2 + i\tau)/2$ and $b = (m + 1/2 + i\tau)/2$. The fixed raw convention gives

$$K_{\mathrm{raw}}(\pi/2) = \frac{\Gamma(m+1)\sqrt{\pi}}{|\Gamma(a)|^2}, \qquad R_\theta = \frac{K_\theta}{K} = 2\frac{|\Gamma(a)|^2}{|\Gamma(b)|^2}, \qquad s = \frac{R_\theta}{2\tau}\,\mathrm{Im}[\psi(b) - \psi(a)].$$

The first two expressions follow from the Ferrers special values at zero (National Institute of Standards and Technology, n.d.); the last is their spectral derivative. They are used outside the positive-series branch, so $\tau > 0$. Complex gamma and digamma values are evaluated by recurrence followed by eight-term Stirling expansions after shifting the argument to modulus at least 16; a first-order correction retains the actual binary64 radian endpoint. A polar point uses the analytic boundary amplitude. A table point retains the table-to-table logarithmic difference, which cancels a common atlas amplitude offset, while replacing the boundary sensitivity by its analytic value. Mehler evaluation uses no online Riccati propagation, WKB branch, or arbitrary-precision fallback.

Lookups remain inside their cells and apply the near-order sampling screen where relevant. Legendre evaluation uses a non-tabulated representation if no lookup is admissible. Mehler and constant calls retain their specified series and endpoint routes, returning failure when none applies. No table is extrapolated.

4.6 Algorithmic cost and storage

Within an accepted cell, lookup and nested Clenshaw evaluation use fixed polynomial degrees. Mehler and zero-eigenvalue calls use fixed tables or bounded series and endpoint branches in the tested domain. The hybrid Legendre evaluator has no such global cost bound: recurrence or adaptive propagation can grow with order distance, phase span and tolerance.

The atlas occupies 158,806,776 bytes in nine sections, loaded on first use by family. Its size is fixed for the current parameterization; extending a range or tightening a screen requires rebuilding the affected sections. Section 7 measures throughput and the remaining Legendre cost tail.

5 Validation design and error evaluation

5.1 Complementary evidence and accuracy criteria

The numerical formulation must preserve the regular branch, its derivatives and its normalization across changes of representation. Validation therefore addresses three distinct questions: whether independently computed values agree, whether the implemented identities remain consistent across the parameter domain, and whether the resulting basis satisfies its cap boundary and

norm conditions. Table 4 separates the evidence supplied by each experiment. Internal identities provide broad coverage, whereas independent references test errors that different numerical representations could share. An extended-degree grid probes algebraic consistency beyond the primary reference range.

The numerical targets are 10^{-7} for the specified identity, modal-error and norm diagnostics, and 10^{-10} for the boundary-root backward residual. Status and finiteness checks accompany these thresholds. A requested reference that does not converge under precision refinement, a missing required component, or an unsuccessful evaluation is counted as a numerical failure. The grids and thresholds are fixed for the reported experiments. Results are presented in Section 6; computational performance is assessed separately in Section 7.

Table 4: Validation layers and the scope of the conclusions they support. Coverage and independence are complementary properties; neither substitutes for the other.

Experiment	Evidence supplied	Limit of inference
Arbitrary-precision references	Point values and derivatives from separate arithmetic and reference representations	Accuracy at the requested tuples, not a uniform bound between them
Internal identities	Dense recurrence and derivative consistency across numerical regimes	A common scale or branch error can satisfy the same identities
Cap roots and norms	Boundary backward residuals and integration cross-checks of normalization	Integration reuses computed basis values or IVP states; it tests normalization consistency
Extended-degree identities	Algebraic consistency on a grid extending to degree 10^8	Identity residuals do not establish independent pointwise accuracy

5.2 Dense algebraic consistency

The primary identity grid contains 279,446 points at 17 interior colatitudes between 0.01° and 89.99°. The exact 90° branch is tested separately. Legendre evaluation uses 180 logarithmically distributed degree samples from 1.371 to 99999, with up to 121 integer orders at each degree. These are predominantly non-integer degrees; the upper endpoint is exactly 99999. Mehler evaluation uses 44 values of τ from zero through the exactly represented endpoint 10000. Its order identities call modes through $m + 2$, so center orders end at 9998 and all evaluated orders remain within $m \leq 10^4$. The constant-family identity calls the adjacent order, with center orders through 9999 and the final called order at 10000.

For an identity $\sum_i a_i = 0$, the diagnostic is

$$\rho = \frac{|\sum_i a_i|}{\sum_i |a_i|}, \tag{77}$$

when the denominator is nonzero; an identity whose terms are all exactly zero is assigned residual zero. The sum is evaluated after subtracting the largest logarithmic magnitude from all terms. This permits the same diagnostic in the ordinary and extended-range records of Section 4.1, without forming an overflowing or underflowing amplitude. The point statistic is $\rho_{\max} = \max(\rho_0, \rho_1, \rho_2)$ for the recurrence, first-derivative and second-derivative identities. The normalization is invariant under a common scale, which is both its strength and its limitation: a small ρ establishes algebraic consistency, not absolute pointwise accuracy.

5.3 Independent references and range-aware error evaluation

The primary reference design comprises 2,350 modes at 17 angles: 1,400 Legendre, 850 Mehler and 100 constant modes, giving 39,950 requested comparisons. The stratification resolves near-diagonal, turning, interior and extreme cases. High-degree Legendre references concentrate near the diagonal, where independent evaluation is especially demanding; they are complemented by

the broader identity and cap sweeps. Separate grids add 510 Legendre degree-sensitivity points, 439 endpoint and boundary points, and 96 zero-neighborhood and degree-sensitivity points. Thus the four reference grids contain 40,995 requests. The 68 analytic regressions and the reference runner's 24 representation and differentiation controls are separate checks, rather than additional primary-grid observations.

References use mpmath 1.3.0 (Johansson et al., 2023) with the gmpy2 2.2.2 backend. Precision is refined through 80, 120 and, when required, 180 decimal digits, with 40 guard digits. Successive scaled references must agree to 10^{-25}; lack of convergence fails the comparison. Exact input bit patterns are exported by C++, including the radian argument used by the degree-sensitivity path, so the two implementations are evaluated at the same represented inputs. Values and derivatives are retained as sign–logarithm pairs. Reference validity therefore does not require the mathematical amplitude to fit in binary64.

Let x_k and y_k denote the production and reference angular derivatives of order $k = 0, 1, 2$. Define the modal frequency and joint reference envelope by

$$\omega = \begin{cases} \nu + 1, & \text{Legendre,} \\ \sqrt{m^2 + \tau^2} + 1, & \text{Mehler,} \\ m + 1, & \text{constant,} \end{cases} \qquad M = \max\left(|y_0|, \frac{|y_1|}{\omega}, \frac{|y_2|}{\omega^2} \right). \tag{78}$$

For $M > 0$, the component error is

$$e_k = \frac{|x_k - y_k|}{M\omega^k}, \qquad k = 0, 1, 2. \tag{79}$$

The shared envelope remains meaningful when one component crosses zero. An exactly zero reference state is accepted only when the corresponding production components are also zero. The statistic measures error relative to a modal state, rather than claiming a relative forward-error bound for each individual component. Differences and scales are formed in sign–logarithm arithmetic, using a two-part binary64 logarithm on the production side and arbitrary-precision logarithms on the reference side. No binary64 amplitude floor is imposed. The comparator additionally checks ordinary outputs whenever the reference component has magnitude at least 2^{-1022}; these errors enter the same acceptance statistic. Ordinary subnormal components are not assigned the same accuracy requirement and may be flushed to zero, while their mathematical sign–logarithm records remain subject to comparison.

Degree sensitivity is tested independently of the production differentiated recurrences and sensitivity equations. For each Legendre reference, mpmath forms the five-point derivative

$$\partial_\nu f(\nu) \approx \frac{-f(\nu + 2h) + 8f(\nu + h) - 8f(\nu - h) + f(\nu - 2h)}{12h}, \tag{80}$$

with $h = 10^{-\lceil p/4 \rceil}$, $p = d + 40$ and $d = 80, 120, 180$. Each function evaluation uses 30 additional digits. Precision refinement uses the same 10^{-25} criterion; independent controls also compare against numerical differentiation of a separate Ferrers representation. If $S^{\mathrm{ref}} = \partial_\nu L^{\mathrm{ref}}$ and $S_\theta^{\mathrm{ref}} = \partial_\theta \partial_\nu L^{\mathrm{ref}}$, the implemented joint sensitivity scale is

$$D = \max\left(\left|S^{\mathrm{ref}}\right|, \frac{\left|S_\theta^{\mathrm{ref}}\right|}{\omega}, \frac{M}{\max(1, \nu)} \right), \qquad e_S = \frac{\left|S - S^{\mathrm{ref}}\right|}{D}, \quad e_{S_\theta} = \frac{\left|S_\theta - S_\theta^{\mathrm{ref}}\right|}{\omega D}. \tag{81}$$

The last term supplies a degree-scaled reference-state floor when the sensitivities vanish; it is not a fixed amplitude floor.

5.4 Boundary roots and cap normalization

The Legendre and constant sweeps use 18 non-hemispherical cap half-angles from 0.4° to 89.99°. Mehler additionally uses 0.0001°, 0.01°, 0.1° and 90°, covering small-cap and hemispheric limits. The Legendre sweep samples Dirichlet root indices 1, 2, 10, 50 and 200. Fixed orders through 10000 are supplemented by fractions of the cap-dependent admissible upper order, subject to $m, \nu \leq 10^5$; the resulting grid reaches order 99600. This tests high-order roots whose admissibility changes with cap angle, rather than imposing the same order list on every cap.

Legendre cap norms from Eq. (16) are compared with adaptive integration of Eq. (8) at 781 points. At 149 high-order near-diagonal points, specified by $m \geq 4096$ and $m/(\nu + 1/2) \geq 0.90$, a continuous turning-point IVP accumulates the squared regular solution across the cap. This avoids quadrature aliasing of the oscillatory tail, but uses the production IVP and is explicitly an algorithm cross-check. Mehler and constant normalized functions are checked by direct norm quadrature. Both kinds of test compare normalization with a separate integration procedure while reusing production states; the independent reference grids address the corresponding point-evaluation errors.

A computed boundary degree is assessed with

$$e_{\mathrm{root}} = \frac{|L_{\nu m}(\theta_0)|}{\max(|S_{\nu m}(\theta_0)|, u_{\min}) \max(1, |\nu|)}, \qquad u_{\min} = 2^{-1022}, \tag{82}$$

where L and S use the same legacy Schmidt convention, and $u_{\min}$ is the smallest positive normal binary64 number. When the safeguard is inactive and the root is simple, this is the relative Newton correction to first order. It is a backward-residual proxy, not a certified interval enclosing the exact root. The angular-ODE residual is retained as an interface diagnostic: away from the polar series, the second derivative is reconstructed from that same ODE, while polar derivatives share their series. It therefore supplies regression evidence rather than an independent accuracy measurement.

5.5 Extended-degree identity tests

A separate 69,156-point Legendre identity grid extends the sampled degree range to 10^8. The recurrence and derivative residuals defined above are evaluated on this grid to examine algebraic consistency beyond the primary reference range. These residuals complement the independent comparisons but do not establish pointwise accuracy throughout the extended domain.

5.6 Performance measurements

Timing uses two observables: sustained throughput of a fixed heterogeneous workload, and amortized cost for individual warmed parameter tuples. The workstation runs Windows 11 on an Intel Core i9-10900K (10 physical cores, 20 logical threads, 3.70 GHz nominal frequency and 20 MiB shared L3 cache), with 32 GiB memory in two 16 GiB modules configured at 2133 MT/s. No accelerator is used. Intel oneAPI DPC++/C++ Compiler 2025.3.2 compiles C++17 with `/O2 /fp:precise`; OpenMP builds additionally use `/Qopenmp`. The Legendre header also scopes precise floating-point semantics around its compensated and sign–logarithm arithmetic. On this Windows build, `long double` has binary64 precision; the compensated normalization does not assume extended hardware precision. Timing is performed without concurrent numerical experiments, with processes at BelowNormal priority. The throughput experiment uses `granularity=fine,scatter` affinity, and the single-thread cost experiment uses compact affinity.

A first pass records initialization separately from warmed throughput. Steady-state tests cover one to 20 OpenMP threads, with at least 27 s per family/thread combination; the heterogeneous pools contain 4,624 Legendre, 4,641 Mehler and 357 constant tuples. Each call returns the value

and first two angular derivatives. The separate single-thread grid contains 400 Legendre, 245 Mehler and 75 constant parameter tuples. Each tuple is warmed three times, its repetition count is calibrated with a batch lasting at least 50 ms, and five timed batches use that repetition count; the reported tuple cost is their median. Quantiles describe variation between parameter tuples, rather than the distribution of individual-call jitter. The timed loops include checksum instrumentation, including inverse tangents of returned components; no overhead is subtracted. The measurements therefore characterize the instrumented evaluation workload, not isolated intrinsic function latency.

6 Validation results

The independently referenced grids support the prescribed 10^{-7} modal accuracy target for the tested values and derivatives. All 40,995 point and sensitivity requests converged and were compared without exclusion. The largest point-component error is 1.11×10^{-8} across the primary and supplementary boundary grids; their individual maxima are 2.15×10^{-9} and 1.11×10^{-8}. The 930 sampled Legendre boundary degrees also meet the separate 10^{-10} root target, with a maximum relative Newton correction of 2.83×10^{-14}. These results are assessed below alongside dense identity checks and cap-norm integration tests, which have different failure modes. They provide empirical error envelopes on fixed grids, rather than a uniform error theorem for the continuous parameter domain.

6.1 Independent pointwise and degree-sensitivity accuracy

The 39,950 primary mpmath references all satisfy the predeclared 10^{-25} refinement criterion. Table 5 reports the worst component statistic among the value and first two angular derivatives under Eq. (79). The maximum errors are 6.39×10^{-10} for Legendre, 2.15×10^{-9} for Mehler, and 4.41×10^{-11} for the constant family. The comparison uses mathematical sign–logarithm records outside ordinary binary64 range and separately checks normal-range ordinary outputs. Thus agreement does not result from assigning a zero error to underflowed values. Figure 1f shows the component distributions, and panel (g) resolves their angular dependence.

Table 5: Arbitrary-precision mpmath comparison using Eq. (79). The P99 column is the largest componentwise P99; the maximum is taken over all components and points.

Family	Valid references	P99 scaled error	Maximum scaled error
Legendre	23,800	1.17×10^{-10}	6.39×10^{-10}
Mehler	14,450	8.13×10^{-10}	2.15×10^{-9}
Constant	1,700	4.35×10^{-11}	4.41×10^{-11}

The analytic Legendre sensitivity of Sections 3 and 4 is tested separately against degree-differenced arbitrary-precision references. All 510 requested pairs $(\partial_\nu L, \partial_\theta \partial_\nu L)$ converge and have both C++ logarithmic components available. Their maxima are 1.06×10^{-10} and 1.03×10^{-10} under the joint scale of Eq. (81) (Table 6). This set includes 117 points with a nonzero mathematical value but zero ordinary binary64 output; their logarithmic states remain comparable. These tests therefore address the degree-sensitive scale used by the boundary norm identity, not only the angular point evaluator. The sensitivity curves are also shown in Fig. 1g.

The supplementary grids add 439 endpoint/boundary comparisons and 96 zero-neighborhood sensitivity comparisons. All references converge and every required component remains below 10^{-7}. The endpoint grid includes normal/subnormal derivative transitions and axis cases with an underflowed amplitude but a representable derivative; its largest component error is 1.11×10^{-8}. Its coverage therefore supplies evidence beyond the primary grid. Each reference grid also passes 24 independent controls before comparison. A separate 68-case analytic regression tests polar

Table 6: Direct arbitrary-precision validation of the Legendre degree sensitivity and mixed angular–degree derivative. All 510 requested references converged and had both C++ sign–logarithm components available for paired comparison.

Quantity	Comparable points	Median error	P99 error	Maximum error
$\partial_\nu L_{\nu m}$	510	1.21×10^{-13}	1.05×10^{-10}	1.06×10^{-10}
$\partial_\theta \partial_\nu L_{\nu m}$	510	1.89×10^{-13}	8.48×10^{-11}	1.03×10^{-10}

derivatives, angular-gradient combinations, non-root small-cap norms, invalid orders, and scalar inverse problems with known solutions; its maximum diagnostic error is 4.36×10^{-14}.

6.2 Dense identity consistency

The 279,446 primary identity points return valid diagnostics with no failures (Table 7). The largest P99 residual is 4.25×10^{-10} and the largest maximum is 7.02×10^{-9}. These dense recurrence and derivative checks exercise consistency across the overlapping numerical representations of Section 4. They complement the independent references, but cannot exclude an error in scale or branch shared by both sides of an identity.

Table 7: Internal identity validation over the target parameter domains. The point statistic is $\rho_{\max} = \max(\rho_0, \rho_1, \rho_2)$ over the value and first two derivative identities.

Family	Valid points	Failures	P99 residual	Maximum residual
Legendre	252,365	0	9.81×10^{-11}	7.02×10^{-9}
Mehler	24,684	0	4.25×10^{-10}	2.02×10^{-9}
Constant	2,397	0	3.64×10^{-12}	7.28×10^{-12}

Figure 1a displays intervals from median to maximum, with circles for P99 and diamonds for maximum. The constant-family first-derivative *identity residual* is exactly zero on this grid and is indicated separately; the function derivative itself is generally nonzero. Panels (c, d) show maximum combined identity residuals in occupied parameter cells, with shared colors and 10^{-12} and 10^{-10} contours; gray cells are unsampled. Panel (e) resolves the constant-family residuals by order and colatitude. The 10^{-16} display floor has no role in acceptance.

6.3 Boundary roots and cap normalization

Table 8 tests the quantities required to turn point functions into a cap basis. The maximum Legendre relative Newton correction, Eq. (82), is 2.83×10^{-14} over 930 boundary degrees. This is a backward-error proxy using the computed sensitivity, rather than an independently known forward error in the root. It tests the boundary solution selected by the refinement workflow, including the scale-preserving sensitivity described in Section 4.

The 930 Legendre cap norms agree with their integration cross-checks to 1.28×10^{-9}: 781 use separate adaptive quadrature and 149 high-order near-diagonal cases use the production turning-point IVP integral. The latter are algorithm cross-checks, not independent accuracy references. Mehler and constant norm comparisons reach 2.74×10^{-10} and 8.42×10^{-11}, respectively, supporting the numerical realization of Proposition 1 and the positive-integral norm in Section 4.4. Because these integrations reuse production point values, their agreement establishes normalization consistency; the arbitrary-precision grids address pointwise accuracy through a separate computation.

The angular-ODE row is an internal regression diagnostic. It shares the production state, and outside the stable polar series the second derivative is reconstructed from that same ODE. Its small residual therefore supplies no additional independent pointwise-accuracy evidence. Figure 1h




Figure 1: Numerical validation of the three R-SCHA angular families. (a–e) Internal identity diagnostics and their parameter dependence. (f, g) Independent arbitrary-precision errors and angular dependence. (h, i) Cap-wide normalization, boundary-root and angular-ODE checks. Red lines mark the 10^{-7} accuracy target; the dotted line marks the 10^{-10} root target. Protocols and diagnostic limitations are described in the text.

Table 8: Cap-wide validation. Norm integration cross-checks normalization consistency; the boundary row is a root backward residual, and the angular-ODE row is an internal regression diagnostic.

Metric	Valid cases	P99 error	Maximum error
Legendre cap norm	930	8.37×10^{-10}	1.28×10^{-9}
Legendre boundary zero	930	1.28×10^{-14}	2.83×10^{-14}
Mehler cap norm	242	7.61×10^{-11}	2.74×10^{-10}
Constant cap norm	126	8.06×10^{-11}	8.42×10^{-11}
Angular ODE	3,349	7.86×10^{-13}	2.32×10^{-12}

summarizes the norm and root distributions, while panel (i) resolves maxima by cap angle. The red and dotted lines mark the 10^{-7} modal and 10^{-10} root targets, respectively.

6.4 Extended-degree algebraic consistency

The separate 69,156-point Legendre identity grid extends calls to $\nu \leq 10^8$. Among its 32,147 points above 10^5, the P99 and maximum normalized residuals are 9.40×10^{-9} and 7.73×10^{-8} (Fig. 1b). These are algebraic stress results. They demonstrate consistency on the additional grid but do not extend the independently referenced primary-domain claim to every degree or parameter tuple up to 10^8.

7 Computational performance

The two performance experiments measure complementary quantities: the sustained rate of completing a heterogeneous workload and the amortized cost of repeatedly evaluating one parameter tuple. Figure 2 separates these observables. Panels (a, b) show the OpenMP rate and its speedup relative to the same family's one-thread workload; panels (c–f) resolve variation across parameter tuples. The reciprocal of aggregate throughput includes parallel execution and scheduling and is therefore not an individual-call latency.

7.1 Sustained throughput and parallel scaling

Table 9 reports warmed rates, after the initialization pass. The best observed Legendre and Mehler rates are 2.80×10^4 and 8.99×10^5 evaluations s^{-1}, at 20 and 16 OpenMP threads, respectively. Their speedups over one thread are 7.64 and 9.77. Thread counts refer to logical threads on the 10-core workstation.

Table 9: Warmed sustained throughput. Each family uses its fixed heterogeneous parameter workload and the same timing protocol.

Family	One thread	Best throughput	Threads	Speedup
Legendre	$3.67 \times 10^3\,s^{-1}$	$2.80 \times 10^4\,s^{-1}$	20	7.64
Mehler	$9.20 \times 10^4\,s^{-1}$	$8.99 \times 10^5\,s^{-1}$	16	9.77
Constant	$3.84 \times 10^6\,s^{-1}$	$5.40 \times 10^6\,s^{-1}$	2	1.41

For a thread count p, define the throughput efficiency as $E(p) = S(p)/p$, where $S(p)$ is the rate relative to the one-thread run. At 10 threads, E is 65.3% for Legendre and 84.5% for Mehler. At their observed peaks, these efficiencies are 38.2% and 61.1%, with the denominator counting logical threads. Relative to 10 threads, the best observed rates are 16.9% and 15.6% higher. The sweep identifies the best configuration in this run rather than a hardware-independent optimum or a separately replicated uncertainty estimate.

The constant family behaves differently: its highest measured rate, 5.40×10^6 evaluations s^{-1}, occurs on 2 threads. The best measured configuration has a speedup of 1.41 over its one-thread run. The timing includes scheduling, reductions, and checksum instrumentation; it does not




Figure 2: Computational performance over the target parameter domains. (a, b) Sustained throughput and parallel speedup over 1–20 threads. (c) Distribution of single-thread parameter costs. (d, e) Legendre and Mehler cost envelopes with a shared color scale. (f) Constant-function cost versus order at five colatitudes. Costs are five-trial medians and include checksum instrumentation.

separate their contributions from implementation or hardware costs. A particular bottleneck cannot be assigned from this scaling curve alone.

7.2 Parameter-dependent evaluation cost

The single-thread measurements reveal the heterogeneity hidden by an aggregate rate. Across the 400 Legendre tuples, the median cost is 7.65 μs, whereas P99 is 10.8 ms and the maximum is 15.2 ms. P99 is approximately 1.41×10^3 times the median. The 245 Mehler tuples have a median of 10.3 μs, P99 of 52.6 μs, and maximum of 52.8 μs. For the 75 constant tuples, the corresponding costs are 244 ns, 259 ns, and 264 ns to three significant figures. Figure 2c also reports P90. These distributions summarize parameter-to-parameter variation in the five-trial median of repeated-call costs; they are not distributions of individually timed calls or measures of temporal jitter. All costs include the unsubtracted loop and inverse-tangent checksum operations.

Panels (d, e) locate this variation in parameter space. Each occupied cell contains the largest five-trial median over its sampled colatitudes, so the plotted envelope is an empirical cost summary rather than a bound on all calls within that cell. Panel (f) shows the considerably narrower constant-family profile by order and colatitude. The large Legendre cost spread is consistent with the range of series, recurrence, phase, and initial-value representations in Section 4. The Mehler and constant implementations emphasize fixed-degree residual interpolation and analytic reconstruction, with a convergent positive-series option for Mehler. These paths provide qualitative context: the benchmark does not record route frequencies or profile individual stages, so it cannot apportion the observed cost to a particular fallback or identify a cache, memory, or arithmetic bottleneck.

For basis-intensive calculations, these results favor retaining initialized table sections and grouping independent evaluations into sufficiently large mode or observation blocks. They also

show why thread counts should be selected for the actual family and workload. The measured scaling curves provide the family-specific evidence for that choice. These are implications for use of the basis layer; they do not establish a speedup over another library or predict an end-to-end model speedup.

8 High-resolution regional magnetic-field modeling

8.1 Data and regional model setup

Two regional applications connect the numerical basis to magnetic-field modeling in contrasting geological settings: the Cascadia plate margin and the North American Midcontinent. Both use a 7.5° modeling cap, with field comparisons restricted to an inner 6.5° cap. The spherical shell extends from 6356.2 to 6921.2 km, or 15 km below to 550 km above the reference radius $a = 6371.2$ km. The internal and external Legendre truncation indices are 378 and 7; the two side-family limits are 3. This parameterization stores 144,124 coefficient entries and includes internal equivalent degrees of approximately 9.1×10^3. These are cap-basis indices and degrees, rather than the degree of a global spherical-harmonic expansion.

The near-surface constraints are total-intensity anomalies from the NAMAG-CM compilation at 1-km node spacing (Bankey et al., 2002). Missing nodes are supplemented by degrees 16–185 of LCS-1 (Olsen et al., 2017); their numbers are reported separately in Table 10. The near-surface data are distributed at a nominal height of 1 km above ground. The background direction for the near-surface total-intensity operator is obtained from degrees 1–15 of CHAOS-8.4 at the nominal epoch 1 January 2003 (Kloss et al., 2026, 2025). Previously processed CHAMP (Maus, 2007; GFZ Helmholtz Centre for Geosciences, n.d.), Swarm A/B/C (Friis-Christensen et al., 2006; European Space Agency, n.d.) and MSS-1 (Zhang, 2023; Macau University of Science and Technology, n.d.) endpoint pairs supply along-track vector and total-intensity differences. Swarm C contributes vector differences only. Details of the satellite data processing are given by Zhang et al. (2025).

The three constraint types enter a weighted, damped least-squares problem,

$$J(\boldsymbol{c}) = \|W_d(G\boldsymbol{c} - \boldsymbol{d})\|_2^2 + \lambda \|\boldsymbol{c}\|_2^2, \qquad (W_d)_{ii} = \sigma_i^{-1}, \qquad \lambda = 0.01. \tag{83}$$

Here G contains the near-surface total-intensity and satellite endpoint-difference operators, and $\boldsymbol{c}$ contains the stored model coefficients. Vector endpoints are rotated into the cap frame before differencing; differences are endpoint 1 minus endpoint 2, without division by separation. The nominal near-surface uncertainty is 40 nT; satellite weights use task- and magnetic-latitude-dependent empirical difference scatter. The damping acts directly on the squared coefficient norm, with zero prior and initial coefficients; it is not a spatial-smoothness penalty. A symmetrically diagonally preconditioned BiCGSTAB iteration (van der Vorst, 1992) solves the normal equations with a limit of 100 iterations and no robust reweighting. Appendix A.3 defines the preconditioner and stopping criterion. The recovered models use the same basis for subsequent field and derivative evaluation. Shared long-wavelength information among the compilation, its gap values and the satellite products is represented only by diagonal weights, rather than a full covariance model.

8.2 Field structure and spectral bandwidth

Figure 3 compares the downward vertical component B_z with EMM2017 (NOAA National Centers for Environmental Information, 2017), WDMAMv2 (Lesur et al., 2016), and SH1050 (Thébault et al., 2021a,b). All fields are synthesized at the same spherical radius, 6371.2 km, on the same 0.02° geocentric grid. The reference models use a common lower SH-degree cutoff of 16, with native upper cutoffs of 790, 800 and 1050, respectively. This geometry removes differences in display height and sampling from the comparison. The 0.02° grid is an evaluation spacing, rather than a measured resolving scale.

Table 10: Regional inputs and agreement with fitted observations. Centers are geocentric. Near-surface counts and residual statistics include the separately identified LCS-1 gap values. Satellite counts refer to endpoint pairs. Vector-difference residuals are expressed in the cap frame used by the inversion; z is positive downward. All residuals use the saved observations and reconstructions over the full modeling caps and describe fitted-data agreement.

Quantity	Cascadia	Midcontinent
Cap center (λ_c, ϕ_c)	$(-127°, 45.5°)$	$(-89°, 45.8°)$
Near-surface total-intensity nodes	2,083,788	1,905,983
LCS-1 gap values	73,008	40
Satellite vector-difference pairs	7,911	7,802
Satellite total-intensity-difference pairs	6,595	6,556
Forward grid nodes	474,496	477,038
Near-surface residual RMS (nT)	60.8	97.5
Near-surface median absolute residual (nT)	10.1	15.3
Near-surface P95 absolute residual (nT)	107.6	124.9
Cap-frame ΔB_{x_c} residual RMS (nT)	1.47	2.23
Cap-frame ΔB_{y_c} residual RMS (nT)	1.74	1.63
ΔB_z residual RMS (nT)	0.70	1.16
Total-intensity-difference residual RMS (nT)	0.72	1.28

The regional solutions retain the principal anomaly organization while expressing finer spatial variations. In Cascadia, elongated offshore anomalies remain coherent across the regional map, and the coastal and onshore field contains substantially more local structure than the global references. In the Midcontinent, the strong arcuate anomaly around Lake Superior persists while smaller anomalies and sharper transitions become visible around its flanks. The enhanced local structure is reflected in the spectra below: the regional solution retains coherent power beyond the global cutoffs, and the anomalies remain continuous on the original 0.02° evaluation grid rather than being clipped at a global-model degree limit. Together, the maps and spectra show that the stable high-degree basis resolves shorter-wavelength regional structure while preserving the same three-dimensional potential representation for near-surface and satellite-altitude constraints.

For the spectral diagnostic, all four models are represented on the same 7.5° cap and evaluated at the comparison radius. Following the cap-spectrum framework of Vervelidou and Thébault (2015), which relates R-SCHA spectral spacing to the cap half-angle and permits comparison with global spherical-harmonic spectra, the plotted power density sums the internal and external Legendre-family contributions in equivalent-degree bins of width $\Delta\nu = 24$, divided by that width, in nT^2 per degree. Mehler, zero-eigenvalue and cross-family terms are excluded. Consequently, this diagnostic does not equal the total energy of the complete model or the variance of the displayed B_z map. Across approximately $\nu = 12$–300, the regional and global curves share the same rise and broad peak. From about $\nu = 400$–800 in Cascadia and $\nu = 400$–900 in the Midcontinent, the regional solution carries systematically more power than the localized global references. The global curves then roll off near their native upper degrees of 790, 800 and 1050, whereas the regional spectrum remains non-negligible through approximately $\nu = 5000$. Beyond the reference bandwidths, regional power decreases smoothly over several orders of magnitude, with a steepening terminal tail. This combination of enhanced intermediate-degree power and a tapered high-degree spectrum distinguishes the regional solutions from the truncated global references. The decay rate varies with the fitted interval, so a single exponential scale is not assigned to the whole tail.

The residual distribution provides a complementary measure of fitted-data agreement (Table 10). Near-surface median absolute residuals are 10.1 and 15.3 nT, compared with RMS values of 60.8 and 97.5 nT and P95 absolute residuals of 107.6 and 124.9 nT. Thus a relatively small

fraction of large residuals contributes strongly to the RMS; all nodes, including gap values, enter these statistics. Satellite vector-difference RMS values range from 0.70 to 2.23 nT in the fitted cap frame, and total-intensity-difference RMS values are 0.72 and 1.28 nT. These statistics quantify agreement with the constraints. Establishing a spatial resolving scale or an accuracy advantage over the global models requires independent observations and treatment of data correlation and regularization.




Figure 3: Regional vertical magnetic fields and common-cap Legendre-family spectra. Panels (a–d) and (e–h) show Cascadia and the Midcontinent, with EMM2017, WDMAMv2, SH1050 and the regional R-SCHA solution in successive columns. B_z is positive downward at spherical radius 6371.2 km. Reference SH bands are 16–790, 16–800 and 16–1050. Circles mark the 6.5° useful caps. All maps share a −400 to 400 nT color scale; saturation affects display only. Panels (i, j) show internal-plus-external Legendre-family power density on 7.5° caps, with $\Delta\nu = 24$. Colored vertical lines mark the reference models' native upper SH degrees; their finite-cap projections can retain spectral tails beyond these cutoffs.

8.3 Geological expression and magnetic transitions

Figure 4 places the regional fields beside geological and tectonic layers compiled from the Cascadia database, regional US geological maps and Canadian geological mapping (Staisch and Walton, 2022; Nicholson et al., 2004; Wheeler et al., 1996); the Minnesota outcrop polygons provide an additional local reference (Minnesota Geological Survey, n.d.). Offshore tectonic context follows the oceanic-crust framework of Seton et al. (2020), and coastlines follow GSHHG (Wessel and Smith, 1996). The geological panels provide surface context; they do not define a unique map of magnetic sources at depth.

We apply the modified horizontal gradient amplitude (MHGA) of Ai et al. (2024) to the modeled B_z field. With $H = \|\nabla_h B_z\|$, the plotted dimensionless response is

$$\mathrm{MHGA} = \mathrm{clip}_{[-1,1]}\left(\frac{H_z}{\sqrt{H_x^2 + H_y^2}} - \frac{\pi}{3}\right), \qquad H_z = \mathcal{F}^{-1}(|\boldsymbol{k}|\,\mathcal{F}H)\,, \tag{84}$$

where subscripts x, y denote horizontal derivatives, H_z is a Fourier pseudo-vertical derivative, and nearly zero denominators are masked. Calculations use a 2-km local projected grid, reflection extension and a 160-km computational buffer. The outer 80 km of the useful cap is excluded, and the B_z and MHGA panels share an identical display mask. This grid-based diagnostic is applied to the modeled vertical component; reduction to the pole and the additional preprocessing of the original MHGA study are not applied.

In Cascadia, A1 identifies a mapped spreading-ridge segment within the offshore banded field. The repeated, ridge-parallel magnetic texture and its corresponding MHGA structure are consistent with contrasts in oceanic-crust magnetization associated with seafloor spreading and remanence carried by oceanic crust (Vine and Wilson, 1965; Seton et al., 2020). This interpretation makes a testable prediction: lineations should remain organized along the ridge direction in B_z and should be concentrated by MHGA where their gradients are strongest, while their continuity should weaken toward the continental margin. A2 marks the mapped subduction boundary and supplies a reference for comparing anomaly texture across the forearc. A change in wavelength or gradient concentration near A2 would be compatible with juxtaposed oceanic, forearc and continental sources, but it would not by itself locate the slab interface or determine source depth. Magnetic responses in the forearc can also reflect crustal and mantle lithology, and the Coast Range includes the Siletzia igneous province (Blakely et al., 2005; Wells et al., 2014). The figure therefore supports a structured comparison of ridge fabric, subduction-margin transition and continental lithology; it does not establish a unique tectonic boundary from magnetic texture alone.

In the Midcontinent, the broad Lake Superior anomaly is consistent with the strong magnetic expression of volcanic and intrusive rocks of the Midcontinent Rift (Hinze et al., 1992; Stein et al., 2015). B1 marks an igneous unit in Minnesota and B2 a mapped fault; identical geographic targets are used across the geological, field and MHGA panels. Sharp or elongated B_z transitions and their MHGA maxima can be compared with the mapped igneous-unit and fault fabric. Coincidence is compatible with contrasts in mafic and intrusive lithology or fault-controlled offsets in magnetized crust. A mismatch may instead reflect a buried source, remanent magnetization, overlapping bodies or a contact displaced from the surface trace. Integrated interpretations of Lake Superior illustrate this non-uniqueness (Grauch et al., 2020). B1 and B2 therefore define candidate source associations. Their depths and geometry require independent geological contacts, susceptibility or remanence constraints, or altitude-dependent field tests.




Figure 4: Geological context and magnetic transitions in Cascadia (a–c) and the Midcontinent (d–f). Columns show mapped geology and tectonic features, regional B_z, and MHGA applied to B_z following Ai et al. (2024). B_z has the same radius and color scale as Fig. 3. The field and MHGA panels use identical masks, including an 80-km edge exclusion; geology retains the full contextual extent. A1/A2 mark a spreading-ridge segment and subduction-boundary position; B1/B2 mark a Minnesota igneous-unit reference and mapped fault. Targets have identical coordinates in all three columns. Enhanced MHGA responses indicate candidate magnetic transitions; spatial association with a mapped feature alone does not verify a subsurface contact. Geological sources are specified in the text.

9 Discussion

9.1 What the numerical formulation adds

The three families require different algorithms, but each benefits from removing the poorly scaled quantity before forming ordinary values. Legendre evaluation carries a scaled series, logarithmic derivative or phase while preserving the regular polar branch. Mehler normalization uses a boundary sensitivity instead of a large raw amplitude and a separate quadrature. The zero-eigenvalue norm is summed with positive terms. These transformations address conditioning; sign–logarithm records then address floating-point range. Extended range alone would not remove cancellation or select the correct branch.

Most ingredients are established. The Legendre solver uses Ferrers series, order recurrences, Liouville–Green asymptotics and the nonoscillatory-phase methods of Bremer and Rokhlin (2017)

and Bremer (2018). Parameter derivatives also have a substantial analytical literature (Cohl, 2011; Szmytkowski, 2011, 2012). Here the representation used in the root or norm calculation is differentiated on the same Schmidt scale. Table 11 identifies the resulting integration and the family-specific normalization formulas. The common Lagrange identity is a standard consequence of the angular Sturm–Liouville equation; the numerical contribution is its stable use with the selected regular branch.

Table 11: Established ingredients and their implementation for the R-SCHA angular basis.

Family	Established ingredient	Implementation in this work
Legendre	Ferrers series, order recurrence, Liouville–Green asymptotics and nonoscillatory phase	Common Schmidt scale across regimes; residual charts; analytic degree sensitivity for boundary roots and cap norms
Mehler	Conical functions, Riccati transformation and adjacent-order identities	Positive regular cap branch; Riccati-sensitivity norm; fixed residual tables through $m, \tau = 10^4$
Zero eigenvalue	Elementary angular shape and Lerch representation	Positive norm integral and series with a tail bound; two-dimensional logarithmic-norm table through $m = 10^4$

The Mehler solver is specialized to $x = \cos\theta \in [0, 1]$. It incorporates cap normalization and does not replace conical-function libraries that support broader arguments and solution classes (Gil et al., 2009, 2012; Dunster et al., 2015). Likewise, the positive-integral transformation addresses the particular zero-eigenvalue norm used by R-SCHA. For Legendre functions, the present near-order table fails its sampled acceptance threshold in every candidate cell; those calls retain the endpoint and recurrence algorithms. Residual compression is therefore an acceleration where accepted, rather than a uniform replacement for direct evaluation.

9.2 Accuracy and limits of the evidence

The 40,995 precision-refined reference requests test values, angular derivatives and Legendre degree sensitivities at specified tuples. Their largest value or angular-derivative error, 1.11×10^{-8}, is below the 10^{-7} modal target. Dense identities add coverage but cannot detect a shared scale or branch error. Norm integration reuses production values, and the root statistic is a relative Newton correction rather than an independently known root error. These distinctions matter especially for second derivatives reconstructed from the angular ODE: a small residual of that same ODE is a regression check, not an independent validation.

Range-aware comparison is essential. At 5,616 of the 14,450 primary Mehler points, the joint reference amplitude is below 10^{-280}. An imposed amplitude floor would conceal discrepancies there. The sign–logarithm comparison retains them and gives a primary-grid maximum of 2.15×10^{-9}; the supplementary Mehler boundary grid reaches 7.73×10^{-9}. Near a component zero, the modal envelope avoids an ill-conditioned component-wise relative error. None of these sampled maxima is a uniform bound between test points. In particular, the identity grid to degree 10^8 does not establish independent accuracy over that extended domain.

9.3 Cost and parallel use

The Legendre cost varies by more than three orders of magnitude between the median and P99 parameter tuples. Recurrence distance, integration span and representation selection can all vary, although the benchmark does not record their individual contributions. Mehler and zero-eigenvalue calls use fixed tables or analytic branches and have narrower measured cost distributions. Retaining initialized tables, reusing mode-dependent quantities and grouping independent evaluations can reduce repeated work in basis-intensive applications.

OpenMP throughput peaks at different thread counts for the three families. This favors workload-specific tuning of the outer evaluation loops, consistent with the observation- or mode-

block decomposition used by the MPI application drivers. The measurements include loop and checksum costs and do not establish superiority over another library or an end-to-end inversion speedup. A matched-domain accuracy and timing comparison would be needed for either claim.

9.4 Implications for regional modeling

The applications demonstrate an operational high-degree basis: each inversion combines approximately two million near-surface nodes with satellite differences and reaches internal equivalent degrees near 9.1×10^3. The common-radius comparison shows how this capacity preserves regional anomaly organization while extending shorter-wavelength field structure beyond the three global references. Geological and MHGA comparisons then relate the additional structure to ridge fabric, igneous provinces and candidate magnetic transitions. These associations guide source hypotheses; depths, dips and buried contacts require additional geological or altitude-dependent constraints.

The accompanying reproducibility supplement provides exact numerical-validation requests, reference-generation scripts and summary records, together with the plotted spectra and scripts that audit saved application residuals. These records make the numerical evidence and reported application statistics traceable. Full regional replication additionally requires the processed observational inputs and their preparation history. The next tests with the greatest scientific value are independent references in difficult Legendre regimes and regional observations withheld from the inversion.

10 Conclusions

The angular basis developed here connects the three families needed for three-dimensional R-SCHA through stable, compatible numerical representations. Legendre values and analytic degree sensitivities share the adopted Schmidt scale; Riccati sensitivity removes the raw amplitude from Mehler normalization; and a positive integral removes cancellation from the zero-eigenvalue norm. Accepted residual tables reduce repeated work, while direct algorithms cover regimes where interpolation is unsuitable.

All 40,995 precision-refined point and sensitivity requests converged. The maximum scaled value or angular-derivative error was 1.11×10^{-8}, below the 10^{-7} modal target, and the maximum direct degree-sensitivity error was 1.06×10^{-10}. The maximum sampled relative Newton correction for boundary roots was 2.83×10^{-14}. These independent references and boundary diagnostics establish the tested basis accuracy, while identity checks extend consistency evidence to degree 10^8. Measured throughput and its parameter dependence provide practical guidance for parallel use.

The Cascadia and Midcontinent inversions show the consequence for regional magnetic-field modeling: dense near-surface and satellite constraints can be represented within one three-dimensional potential expansion at internal equivalent degrees near 9.1×10^3. Common-radius maps and cap spectra express shorter-wavelength structure beyond the global references, and geological comparisons connect that structure to candidate crustal sources and boundaries. Stable high-degree evaluation therefore expands the bandwidth available to regional models and the geological questions they can address.

CRediT authorship contribution statement

Pan Zhang: Conceptualization, Methodology, Formal analysis, Validation, Visualization, Writing – original draft. **Jinsong Du:** Conceptualization, Supervision, Project administration. **Jiaxuan Zhang:** Software, Investigation, Validation.

Funding

This work is supported by the National Natural Science Foundation of China (Grant Nos. 42574077, 42250103, and 42674085), and the Natural Science Foundation of Hubei Province of China (Grant No. 2024AFE003).

Acknowledgements

We thank the teams responsible for NAMAG-CM, the CHAMP, Swarm and MSS-1 missions, the global magnetic reference models, and the geological and coastline datasets used in this study for making their products available. The individual sources are identified in Section 8 and the reference list.

Data and code availability

The R-SCHA v2.0.1 release contains the OpenMP and MPI source trees, the canonical-residual lookup table, build scripts, validation summaries and the mathematical foundations handbook. Download it from the GitHub v2.0.1 release or the project repository (R-SCHA contributors, 2026). The manuscript source package contains the reproducibility records and supporting spectra described in the text; external magnetic, geological and coastline datasets remain available from their original sources listed in Section 8.

Declaration of competing interest

The authors declare no competing financial interests or personal relationships that could have influenced the work reported in this paper.

A Overview of the R-SCHA software package

The basis algorithms described in this paper form the numerical core of a broader C++ R-SCHA package. The package separates special-function evaluation, spectral-root calculation, coordinate transformations, model kernels, inversion, and parallel communication. The function-level numerical experiments in Sections 5–6 are separate from the application drivers described below.

A.1 Numerical modules

Table 12 lists the principal numerical modules. The package separates the public basis interface from the underlying Legendre and canonical-residual algorithms, the spectral-root solver, model-level vector and tensor kernels, coordinate transformations, inversion routines, and MPI communication. Exact source-file names and their roles are given in the table.

A.2 Application drivers

Two executable drivers illustrate how the numerical core is used in the inversion and forward-evaluation workflows considered here.

1. **Model construction (`Work1_RSCHAInversion.cpp`).** The program reads the cap geometry and truncation parameters and can combine total-field intensity, vector observations, and along-track differences of either data type. Coordinates and vector components are transformed to the spherical-cap system before the corresponding R-SCHA kernels are assembled. The reference workflow solves the weighted inverse problem with a preconditioned BiCGSTAB iteration and writes the recovered coefficients, model spectrum, and reconstructed data.

Table 12: Main components of the current C++ R-SCHA program. File names are given to make the numerical layers and the model-level workflows unambiguous.

Source component	Main role
`Legendre.h`, `CanonicalResidual`	Real-degree Legendre evaluation, extended-range state, analytic degree sensitivity, asymptotic/direct fallbacks, and canonical-residual lookup.
`Legendre.h`, `CRTable.h`	
`Basis Function.h`	Public Legendre, Mehler, and zero-eigenvalue values and angular derivatives, together with the family-specific normalization wrappers.
`RSCHARoots.h`	Prediction and refinement of the real-degree Dirichlet roots required by the Legendre series.
`R_SCHA.h`	R-SCHA coefficient bookkeeping, first- and second-rank field kernels, model input/output, and spectral summaries.
`CoordinateTransformation.h`	Transformation between geographic and spherical-cap coordinates and rotation of vector and second-rank tensor components.
`Inversion.h`	Dense kernel storage and the preconditioned biconjugate gradient stabilized (BiCGSTAB) solver used by the reference inversion workflow.
`MPIParallel.h`	MPI initialization, rank/size access, and collective reductions used by the model-level routines.

2. **Forward evaluation (`Work2_RSCHAForward.cpp`).** A stored R-SCHA model is evaluated at an arbitrary list of radii and geographic coordinates. The current driver computes the scalar potential, the three first-rank field components, and the six independent second-rank tensor components. The results are rotated from the spherical-cap system to the local Cartesian frame before output.

The inversion driver reads the cap geometry, truncation parameters and observations, and writes the recovered coefficient model. The forward driver reads this model together with evaluation coordinates.

A.3 Regional inversion equations and termination

For Eq. (83), the normal equations are

$$H\boldsymbol{c} = \boldsymbol{b}, \qquad H = G^{\mathsf{T}} W_d^2 G + \lambda I, \qquad \boldsymbol{b} = G^{\mathsf{T}} W_d^2 \boldsymbol{d}. \tag{85}$$

All active model weights are unity, so the penalty is on the coefficients as stored. The scalar λ multiplies the penalty directly; it is not squared or rescaled by the number of observations. Products with H use products with G and G^{T} rather than an explicitly assembled normal matrix. With

$$F_{jj} = H_{jj}^{-1/2}, \qquad \widetilde{H} = FHF, \qquad \widetilde{\boldsymbol{b}} = F\boldsymbol{b}, \qquad \boldsymbol{c} = F\boldsymbol{z}, \tag{86}$$

BiCGSTAB operates on $\widetilde{H}\boldsymbol{z} = \widetilde{\boldsymbol{b}}$.

The implementation uses $\tau = 64\epsilon_{\mathrm{mach}}$ for its scaled algebraic residual test. At initialization or restart, with iterate $\boldsymbol{z}_0$, it sets

$$s_0 = \max\left(\left\| \widetilde{\boldsymbol{b}} \right\|_\infty , \left\| \widetilde{H}\boldsymbol{z}_0 \right\|_\infty \right). \tag{87}$$

The initial residual is accepted if its infinity norm is at most τs_0. Subsequently, a sufficiently small recursive residual triggers explicit recomputation of $\boldsymbol{r} = \widetilde{\boldsymbol{b}} - \widetilde{H}\boldsymbol{z}$, which must satisfy

$$\|\boldsymbol{r}\|_\infty \le \tau \max\left(s_0, \left\| \widetilde{H}\boldsymbol{z} \right\|_\infty \right). \tag{88}$$

The configured unweighted data-RMS threshold is active only for zero damping and therefore does not govern these runs. Both regional parameter files set the iteration limit to 100 and disable robust reweighting. Application logs were unavailable in the inspected archive, so the

actual iteration counts and termination histories cannot be recovered from the saved models. The residuals in Table 10 are recomputed from the saved reconstructions and do not imply that the algebraic stopping test was met.

A.4 Parallel execution

The application programs share a common MPI layer. Independent observations or spectral-order blocks are distributed across MPI ranks and combined by collective reductions; root-array construction also uses OpenMP across independent order blocks. This decomposition avoids synchronization within a single basis evaluation.

A.5 Supplementary reproducibility checks

A separate six-hour Bremer comparison with seed 20260827 records 24,821,889 post-warmup samples, with no production status failures. It retains 264 exact-input dispute entries (249 distinct parameter tuples) for independent 100/150-digit integral or 270/430-bit recurrence adjudication. All retained entries meet 10^{-7}, with maximum new-method error 4.71×10^{-9}. The raw Bremer comparison has 8,491,571 unavailable comparisons and a largest available discrepancy 7.55×10^{-1}. Raw discrepancies are not automatically attributed to the new method, and unavailable references are not counted as agreement. Only the retained queue is independently adjudicated; this experiment does not certify every unavailable comparison or the continuous parameter domain. The source and exact-input provenance are retained with the results.

For the inversion and forward-evaluation outputs retained from the serial and three-rank MPI regression, all seven output comparisons pass. These outputs cover 70,264 forward coordinates, 7,136 model coefficients, and 20 inversion iterations per implementation. The largest relative L_2 difference is 1.81×10^{-12}, and the largest difference scaled by $\max(1, |a|, |b|)$ is 3.56×10^{-10}. Model metadata, coordinates and observed data match exactly, and the forward tensor-trace diagnostic passes in both implementations. These are integration-consistency checks rather than independent geophysical validation.

References

Hanbing Ai, Hazel Deniz Toktay, Ahmad Alvandi, Roman Pašteka, Kejia Su, and Qiang Liu. Advancing potential field data analysis: the modified horizontal gradient amplitude method (MHGA). *Contributions to Geophysics and Geodesy*, 54(2):119–143, 2024. https://doi.org/10.31577/congeo.2024.54.2.1.

Viki Bankey, Alejandro Cuevas, David Daniels, Carol A. Finn, Israel Hernandez, Patricia Hill, Robert Kucks, Warner Miles, Mark Pilkington, Carter Roberts, Walter Roest, Victoria Rystrom, Sarah Shearer, Stephen Snyder, Ronald E. Sweeney, Julio Velez, J. D. Phillips, and D. K. A. Ravat. Digital data grids for the magnetic anomaly map of North America. Open-File Report 2002-414, U.S. Geological Survey, 2002. URL https://www.usgs.gov/publications/digital-data-grids-magnetic-anomaly-map-north-america.

Richard J. Blakely, Thomas M. Brocher, and Ray E. Wells. Subduction-zone magnetic anomalies and implications for hydrated forearc mantle. *Geology*, 33(6):445–448, 2005. https://doi.org/10.1130/G21447.1.

James Bremer. An algorithm for the numerical evaluation of the associated Legendre functions that runs in time independent of degree and order. *Journal of Computational Physics*, 360:15–38, 2018. https://doi.org/10.1016/j.jcp.2018.01.014.

James Bremer and Vladimir Rokhlin. On the nonoscillatory phase function for Legendre's differential equation. *Journal of Computational Physics*, 350:326–342, 2017. https://doi.org/10.1016/j.jcp.2017.08.041.

Howard S. Cohl. On parameter differentiation for integral representations of associated Legendre functions. *Symmetry, Integrability and Geometry: Methods and Applications*, 7:050, 2011. https://doi.org/10.3842/SIGMA.2011.050.

T. M. Dunster, A. Gil, J. Segura, and N. M. Temme. Computation of a numerically satisfactory pair of solutions of the differential equation for conical functions of non-negative integer orders. *Numerical Algorithms*, 68:497–509, 2015. https://doi.org/10.1007/s11075-014-9857-5.

European Space Agency. Swarm data access. ESA Earth Online, n.d. URL https://earth.esa.int/eogateway/missions/swarm/data. Official Swarm data archive; accessed 17 October 2026.

Eigil Friis-Christensen, Hermann Lühr, and Gauthier Hulot. Swarm: A constellation to study the Earth's magnetic field. *Earth, Planets and Space*, 58(4):351–358, 2006. https://doi.org/10.1186/BF03351933.

Toshio Fukushima. Numerical computation of spherical harmonics of arbitrary degree and order by extending exponent of floating point numbers. *Journal of Geodesy*, 86:271–285, 2012.

GFZ Helmholtz Centre for Geosciences. Access to the CHAMP data base. Information System and Data Center, n.d. URL https://isdc.gfz.de/champ-isdc/access-to-the-champ-data. Official CHAMP data archive; accessed 17 October 2026.

Amparo Gil, Javier Segura, and Nico M. Temme. Computing the conical function $p^{\mu}_{-1/2+i\tau}(x)$. *SIAM Journal on Scientific Computing*, 31(3):1716–1741, 2009. https://doi.org/10.1137/070712006.

Amparo Gil, Javier Segura, and Nico M. Temme. An improved algorithm and a Fortran 90 module for computing the conical function $p^{m}_{-1/2+i\tau}(x)$. *Computer Physics Communications*, 183:794–799, 2012. https://doi.org/10.1016/j.cpc.2011.11.025.

V. J. Grauch, Eric D. Anderson, Samuel J. Heller, Esther K. Stewart, and Laurel G. Woodruff. Integrated geophysical analysis provides an alternate interpretation of the northern margin of the North American Midcontinent Rift System, central Lake Superior. *Interpretation*, 8(4):SS63–SS85, 2020. https://doi.org/10.1190/INT-2019-0262.1.

G. V. Haines. Spherical cap harmonic analysis. *Journal of Geophysical Research: Solid Earth*, 90(B3): 2583–2591, 1985. https://doi.org/10.1029/JB090iB03p02583.

William J. Hinze, David J. Allen, Adam J. Fox, Don Sunwood, Timothy Woelk, and Alan G. Green. Geophysical investigations and crustal structure of the North American Midcontinent Rift system. *Tectonophysics*, 213(1–2):17–32, 1992. https://doi.org/10.1016/0040-1951(92)90248-5.

Cheinway Hwang and Shin-Kuen Chen. Fully normalized spherical cap harmonics: Application to the analysis of sea-level data from TOPEX/POSEIDON and ERS-1. *Geophysical Journal International*, 129(2):450–460, 1997. https://doi.org/10.1111/j.1365-246X.1997.tb01595.x.

Fredrik Johansson et al. mpmath: a python library for arbitrary-precision floating-point arithmetic, version 1.3.0, 2023. URL https://github.com/mpmath/mpmath/tree/1.3.0.

Clemens Kloss, Christopher C. Finlay, Nils Olsen, Lars Tøffner-Clausen, Nicolas Gillet, and Alexander Grayver. CHAOS-8 geomagnetic field model, September 2025. URL https://zenodo.org/records/17225475. Version 8.4, published 29 September 2025.

Clemens Kloss, Christopher C. Finlay, Nils Olsen, Lars Tøffner-Clausen, Nicolas Gillet, and Alexander Grayver. The CHAOS-8 geomagnetic field model. *Earth, Planets and Space*, 78(1):21, 2026. https://doi.org/10.1186/s40623-025-02352-0.

Vincent Lesur, Mohamed Hamoudi, Yujin Choi, Jérôme Dyment, and Erwan Thébault. Building the second version of the world digital magnetic anomaly map (WDMAM). *Earth, Planets and Space*, 68: 27, 2016. https://doi.org/10.1186/s40623-016-0404-6.

Macau University of Science and Technology. Macau Science Satellite-1 data. MSS-1 mission data portal, n.d. URL https://mss.must.edu.mo/data.html. Official MSS-1 data portal; accessed 17 October 2026.

Stefan Maus. CHAMP. In David Gubbins and Emilio Herrero-Bervera, editors, *Encyclopedia of Geomagnetism and Paleomagnetism*, pages 59–60. Springer, Dordrecht, 2007. https://doi.org/10.1007/978-1-4020-4423-6_25.

Minnesota Geological Survey. D-02, Minnesota bedrock outcrop database. Official ArcGIS feature service, n.d. URL https://mngs-umn.opendata.arcgis.com/datasets/UMN::d-02-minnesota-bedrock-outcrop-database/about. Accessed 17 October 2026. Source polygons indicate bedrock exposed at or near the land surface.

National Institute of Standards and Technology. NIST Digital Library of Mathematical Functions: Section 14.5, special values. https://dlmf.nist.gov/14.5, n.d. Equations 14.5.1 and 14.5.2; accessed 7 October 2026.

Suzanne W. Nicholson, Connie L. Dicken, Michael P. Foose, and Julia A. L. Mueller. Preliminary integrated geologic map databases for the united states: Minnesota, wisconsin, michigan, illinois, and indiana. Open-File Report 2004-1355, U.S. Geological Survey, 2004. URL https://pubs.usgs.gov/of/2004/1355/. Version 1.1, updated December 2007.

NOAA National Centers for Environmental Information. Enhanced magnetic model (EMM2017). Model coefficients and software, released 5 July 2017, 2017. URL https://www.ncei.noaa.gov/products/enh

anced-magnetic-model. Accessed 17 October 2026. Lithospheric part used here: spherical-harmonic degrees 16–790.

Nils Olsen, Dhananjay Ravat, Christopher C. Finlay, and Livia K. Kother. LCS-1: a high-resolution global model of the lithospheric magnetic field derived from CHAMP and Swarm satellite observations. *Geophysical Journal International*, 211(3):1461–1477, 2017. https://doi.org/10.1093/gji/ggx381.

F. W. J. Olver and J. M. Smith. Associated Legendre functions on the cut. *Journal of Computational Physics*, 51:502–518, 1983.

R-SCHA contributors. R-SCHA v2.0.1 OpenMP and MPI software. GitHub, versioned software release, 2026. URL https://github.com/ZhangPan-666/RSCHA-v2.0/releases/tag/v2.0.1. Source repository: https://github.com/ZhangPan-666/RSCHA-v2.0. Software under MIT; mathematical handbook under CC BY 4.0.

Maria Seton, R. Dietmar Müller, Sabin Zahirovic, Simon Williams, Nicky M. Wright, John Cannon, Joanne M. Whittaker, Kara J. Matthews, and Rebecca McGirr. A global data set of present-day oceanic crustal age and seafloor spreading parameters. *Geochemistry, Geophysics, Geosystems*, 21(10): e2020GC009214, 2020. https://doi.org/10.1029/2020GC009214.

J. M. Smith, F. W. J. Olver, and D. W. Lozier. Extended-range arithmetic and normalized Legendre polynomials. *ACM Transactions on Mathematical Software*, 7(1):93–105, 1981.

L. M. Staisch and M. A. Walton. Cascadia subduction zone database: compilation of published datasets relevant to Cascadia subduction zone earthquake hazards and tectonics. U.S. Geological Survey data release, 2022. URL https://www.usgs.gov/special-topics/subduction-zone-science/science/cascadia-subduction-zone-database.

Carol A. Stein, Jonas Kley, Seth Stein, David Hindle, and G. Randy Keller. North America's Midcontinent Rift: When rift met LIP. *Geosphere*, 11(5):1607–1616, 2015. https://doi.org/10.1130/GES01183.1.

Radosław Szmytkowski. On the derivative of the associated Legendre function of the first kind of integer order with respect to its degree. *Journal of Mathematical Chemistry*, 49(7):1436–1477, 2011. https://doi.org/10.1007/s10910-011-9826-3.

Radosław Szmytkowski. On parameter derivatives of the associated Legendre function of the first kind. *Journal of Mathematical Analysis and Applications*, 386(1):332–342, 2012. https://doi.org/10.1016/j.jmaa.2011.08.003.

E. Thébault, J. J. Schott, M. Mandea, and J. P. Hoffbeck. A new proposal for spherical cap harmonic modelling. *Geophysical Journal International*, 159(1):83–103, 2004. https://doi.org/10.1111/j.1365-246X.2004.02361.x.

E. Thébault, J. J. Schott, and M. Mandea. Revised spherical cap harmonic analysis (R-SCHA): Validation and properties. *Journal of Geophysical Research: Solid Earth*, 111(B1):B01102, 2006. https://doi.org/10.1029/2005JB003836.

Erwan Thébault, Gauthier Hulot, Benoit Langlais, and Pierre Vigneron. A spherical harmonic model of Earth's lithospheric magnetic field up to degree 1050. *Geophysical Research Letters*, 48(21): e2021GL095147, 2021a. https://doi.org/10.1029/2021GL095147.

Erwan Thébault, Gauthier Hulot, Benoit Langlais, and Pierre Vigneron. A spherical harmonic model of Earth's lithospheric magnetic field up to degree 1050. Zenodo, coefficient dataset, version 1, 2021b. URL https://zenodo.org/records/5546528.

J. Miquel Torta. Modelling by spherical cap harmonic analysis: A literature review. *Surveys in Geophysics*, 41(2):201–247, 2020. https://doi.org/10.1007/s10712-019-09576-2.

H. A. van der Vorst. Bi-CGSTAB: A fast and smoothly converging variant of Bi-CG for the solution of nonsymmetric linear systems. *SIAM Journal on Scientific and Statistical Computing*, 13(2):631–644, 1992. https://doi.org/10.1137/0913035.

F. Vervelidou, E. Thébault, and M. Korte. A high-resolution lithospheric magnetic field model over Southern Africa based on a joint inversion of CHAMP, Swarm, WDMAM, and ground magnetic field data. *Solid Earth*, 9(4):897–910, 2018. https://doi.org/10.5194/se-9-897-2018.

Foteini Vervelidou and Erwan Thébault. Global maps of the magnetic thickness and magnetization of the Earth's lithosphere. *Earth, Planets and Space*, 67:173, 2015. https://doi.org/10.1186/s40623-015-0329-5.

F. J. Vine and J. T. Wilson. Magnetic anomalies over a young oceanic ridge off Vancouver Island. *Science*, 150(3695):485–489, 1965. https://doi.org/10.1126/science.150.3695.485.

Ray Wells, David Bukry, Richard Friedman, Douglas Pyle, Robert Duncan, Peter J. Haeussler, and Joe Wooden. Geologic history of Siletzia, a large igneous province in the Oregon and Washington coast range: Correlation to the geomagnetic polarity time scale and implications for a long-lived yellowstone hotspot. *Geosphere*, 10(4):692–719, 2014. https://doi.org/10.1130/GES01018.1.

Pål Wessel and Walter H. F. Smith. A global, self-consistent, hierarchical, high-resolution shoreline database. *Journal of Geophysical Research: Solid Earth*, 101(B4):8741–8743, 1996. https://doi.org/10.1029/96JB00104.

J. O. Wheeler, P. F. Hoffman, K. D. Card, A. Davidson, B. V. Sanford, A. V. Okulitch, and W. R. Roest. Geological map of Canada. Geological Survey of Canada, A Series Map 1860A, scale 1:5,000,000, 1996. URL https://open.canada.ca/data/en/dataset/4c9f59d6-667f-8285-a75d-25ac73d61e9b.

Keke Zhang. A novel geomagnetic satellite constellation: Science and applications. *Earth and Planetary Physics*, 7(1):4–21, 2023. https://doi.org/10.26464/epp2023019.

Pan Zhang, Jinsong Du, Liang Yin, Qing Yan, Jiaxuan Zhang, Houpu Li, Keke Zhang, and Chao Chen. A high-resolution lithospheric magnetic field model of China and surroundings by using CHAMP, Swarm, CSES-1, and MSS-1 satellite magnetic measurements and aeromagnetic anomaly data. *Earth and Planetary Physics*, 9(3):1–11, 2025. https://doi.org/10.26464/epp2025057.