

Recitation 1

Maximum Likelihood Estimation

Review and Practice Problems

Xuanxi Zhang

1 Maximum likelihood estimation: the basic setup

Suppose $x_1, \dots, x_n$ are observations from a model indexed by a parameter $\theta \in \Theta$. Under an i.i.d. assumption, the **likelihood** is

$$L(\theta; x_1, \dots, x_n) = \prod_{i=1}^n p_\theta(x_i) \quad \text{or} \quad L(\theta; x_1, \dots, x_n) = \prod_{i=1}^n f_\theta(x_i),$$

depending on whether the model is discrete or continuous. Once the data are observed, L is a function of θ ; it is not a pmf or pdf in θ .

Definition. *The maximum likelihood estimator (MLE) is*

$$\hat{\theta}_{\text{ML}} \in \arg \max_{\theta \in \Theta} L(\theta; x_1, \dots, x_n).$$

Because log is strictly increasing, maximizing L is equivalent to maximizing the log-likelihood

$$\ell(\theta) = \log L(\theta) = \sum_{i=1}^n \log p_\theta(x_i) \quad \text{or} \quad \ell(\theta) = \sum_{i=1}^n \log f_\theta(x_i).$$

A reliable workflow is:

1. state the parameter space and the model assumptions;
2. write the likelihood, including any parameter-dependent support;
3. take logs and remove only terms that do not depend on the parameter;
4. find candidate maximizers and check the boundary or support;
5. report the estimate in the units of the original parameter.

2 Poisson MLE

Let $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} \text{Poisson}(\lambda)$, where $\lambda \geq 0$. For observed counts $x_1, \dots, x_n$,

$$L(\lambda) = \prod_{i=1}^n e^{-\lambda} \frac{\lambda^{x_i}}{x_i!},$$
$$\ell(\lambda) = -n\lambda + \left(\sum_{i=1}^n x_i \right) \log \lambda - \sum_{i=1}^n \log(x_i!).$$

If $\sum_i x_i > 0$, then

$$\ell'(\lambda) = -n + \frac{\sum_i x_i}{\lambda}, \quad \ell''(\lambda) = -\frac{\sum_i x_i}{\lambda^2} < 0.$$

Therefore

$$\hat{\lambda}_{\text{ML}} = \frac{1}{n} \sum_{i=1}^n X_i = \bar{X}.$$

The estimator has a natural interpretation: the fitted expected count equals the sample average. If every observed count is zero, the likelihood decreases as λ increases, so the maximizer is the boundary value $\hat{\lambda} = 0$.

3 Gaussian MLE

Let $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} \mathcal{N}(\mu, \sigma^2)$, where $\mu \in \mathbb{R}$ and $\sigma^2 > 0$. Write $v = \sigma^2$. The log-likelihood is

$$\ell(\mu, v) = -\frac{n}{2} \log(2\pi) - \frac{n}{2} \log v - \frac{1}{2v} \sum_{i=1}^n (x_i - \mu)^2.$$

For fixed v ,

$$\frac{\partial \ell}{\partial \mu} = \frac{1}{v} \sum_{i=1}^n (x_i - \mu) = 0 \quad \implies \quad \boxed{\hat{\mu}_{\text{ML}} = \bar{X}}.$$

The identity

$$\sum_{i=1}^n (x_i - \mu)^2 = \sum_{i=1}^n (x_i - \bar{x})^2 + n(\mu - \bar{x})^2$$

also shows directly that $\mu = \bar{x}$ gives the global maximum for fixed v . Substitute $\hat{\mu} = \bar{x}$ and let $S = \sum_i (x_i - \bar{x})^2$. Then

$$\ell(\bar{x}, v) = C - \frac{n}{2} \log v - \frac{S}{2v}, \quad \frac{d\ell}{dv} = -\frac{n}{2v} + \frac{S}{2v^2}.$$

Thus, when $S > 0$,

$$\boxed{\hat{\sigma}_{\text{ML}}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2}, \quad \boxed{\hat{\sigma}_{\text{ML}} = \sqrt{\frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2}}.$$

MLE variance versus the usual sample variance. The Gaussian MLE divides by n . The unbiased sample variance divides by $n - 1$. Maximum likelihood targets the parameter value that best fits the observed data; it does not guarantee an unbiased estimator.

Degenerate data. If all observations are equal, then $S = 0$. With parameter space $v > 0$, the likelihood grows without bound as $v \downarrow 0$, so no MLE for v exists in that open parameter space. This is a useful reminder to check the boundary.

Practice problems

Problem 1: A two-point model

[7 minutes]

Suppose X is a discrete random variable with

$$\mathbb{P}(X = 1) = \theta, \quad \mathbb{P}(X = 2) = 1 - \theta, \quad 0 \leq \theta \leq 1.$$

We draw n independent observations and find that n_1 of them equal 1 and n_2 equal 2, where $n_1 + n_2 = n$.

- (a) Write the likelihood function.
- (b) Find the maximum likelihood estimator of θ , including the cases $n_1 = 0$ or $n_2 = 0$.
- (c) Rewrite the MLE in terms of the sample mean $\bar{X}$.

Problem 2: Calls to a help desk

[7 minutes]

The numbers of calls received in eight comparable 15-minute periods are

$$3, 0, 2, 1, 4, 2, 0, 4.$$

Assume the counts are independent $\text{Poisson}(\lambda)$ random variables.

- (a) Find the MLE of λ .
- (b) Using the fitted model, estimate the probability that a future 15-minute period has at least three calls.
- (c) Give one modeling assumption that would make the calculation in part (b) questionable.

Problem 3: Repeated measurements**[8 minutes]**

A device records the same reference quantity five times:

9.7, 10.1, 10.2, 9.9, 10.1.

Assume the measurements are i.i.d. $\mathcal{N}(\mu, \sigma^2)$.

- (a) Compute $\hat{\mu}_{\text{ML}}$.
- (b) Compute $\hat{\sigma}_{\text{ML}}^2$ and $\hat{\sigma}_{\text{ML}}$.
- (c) Compute the usual unbiased sample variance s^2 . Why does its denominator differ from the MLE denominator?
- (d) If 2 is added to every observation, how do the two MLEs change?

Problem 4 (backup): True or false?

Optional if time allows. Suggested working time: 5–8 minutes.

For each statement, decide whether it is true or false and give a brief reason.

- (a) MLEs are always unbiased.
- (b) If $\hat{\theta}_n \xrightarrow{P} \theta$, then $\hat{\theta}_n^2 \xrightarrow{P} \theta^2$.
- (c) Every solution of the score equation $\ell'(\theta) = 0$ is an MLE.
- (d) A likelihood must satisfy $\int_{-\infty}^{\infty} L(\theta) d\theta = 1$.

Takeaways

- For i.i.d. data, likelihoods multiply and log-likelihoods add.
- The Poisson MLE matches the sample mean count.
- The Gaussian MLE matches the sample mean and uses divisor n for the variance.
- A stationary point is not enough: check the parameter space, boundary, and parameter-dependent support.

Sources

The review follows the course handouts [Comparing Parametric and Nonparametric Models](#) and [Maximum Likelihood Estimation for Continuous Models](#). Problem 1 and Problem 4 are adapted from the 2025 Fall statistics recitation materials (Recitation 5).