


A


Exported cross-feeding metabolites (pooled L0–L9; * documented in culture, † contradicted; Table S11)
Probiotic
Exported metabolite
Metabolite class
Akkermansia
Lactobacillus
Formate* 25%
L-Malate 20%
Succinate* 14%
Propionate 12%
Acetate* 11%
Fumarate 10%
Methanol* 47%
(R)-1,2-Propanediol* 30%
1,2-Propanediol* 23%
D-Ribose* 23%
Cytidine 10%
Methanethiol 34%
Putrescine 28%
Pantothenate* 26%
Indole 17%
L-Malate 40%
Succinate* 17%
Methanol* 15%
D-Ribose* 23%
Inosine† 43%
Glycine 12%
L-Glutamine 12%
Putrescine 15%
Organic acid
Alcohol / diol
Sugar
Nucleoside
Amino acid
Other

B


Akkermansia
Methanol
D-Fructose
Methanethiol
D-Ribose
(R)-1,2-Propanediol
Putrescine
L-Malate
Pantothenate
Formate
1,2-Propanediol
Acetate
Indole
L0 L1 L2 L3 L4 L5 L6 L7 L8 L9
Prevalence (%)
50
40
30
20
10
Lactobacillus
L-Malate
Inosine
D-Ribose
Succinate
Putrescine
Methanol
Glycine
L-Glutamine
Cytidine
1,2-Propanediol
Methanethiol
Xanthosine
L0 L1 L2 L3 L4 L5 L6 L7 L8 L9
Prevalence (%)
40
30
20
10

C


Currency class composition
Share of currencies
100%
75%
50%
25%
0%
27%
7%
29%
32%
32%
24%
9%
12%
11%
12%
Akkermansia
Lactobacillus
Metabolite class
organic acid
nucleoside
sugar
amino acid
alcohol/diol
other