

Asymptotic Diameter of Preferential Attachment Model

Shuyang Gong and Zhangsong Li

School of Mathematical Sciences, Peking University

May 29, 2025


Joint work with Hang Du (MIT) and Haodong Zhu (TU/E)

YMSC Probability Seminar

Preferential attachment model (PAM)

At each time t , a new vertex labeled t arrives and forms m edges, one at a time, to existing nodes $v \in [t-1]$:


$$\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta \text{ where } \delta > -m.$$



Preferential attachment model (PAM)

At each time t , a new vertex labeled t arrives and forms m edges, one at a time, to existing nodes $v \in [t-1]$:


$$\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta \text{ where } \delta > -m.$$



Preferential attachment model (PAM)

At each time t , a new vertex labeled t arrives and forms m edges, one at a time, to existing nodes $v \in [t-1]$:


$$\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta \text{ where } \delta > -m.$$



Preferential attachment model (PAM)

At each time t , a new vertex labeled t arrives and forms m edges, one at a time, to existing nodes $v \in [t-1]$:


$$\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta \text{ where } \delta > -m.$$



Preferential attachment model (PAM)

At each time t , a new vertex labeled t arrives and forms m edges, one at a time, to existing nodes $v \in [t-1]$:

$$\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta \text{ where } \delta > -m.$$



Preferential attachment model (PAM)

At each time t , a new vertex labeled t arrives and forms m edges, one at a time, to existing nodes $v \in [t - 1]$:

$$\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta \text{ where } \delta > -m.$$

- $\deg(v)$ is updated after each edge is added

Preferential attachment model (PAM)

At each time t , a new vertex labeled t arrives and forms m edges, one at a time, to existing nodes $v \in [t - 1]$:

$$\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta \text{ where } \delta > -m.$$

- $\deg(v)$ is updated after each edge is added
- $\delta = \infty$: uniform-attachment (no degree preference)

Preferential attachment model (PAM)

At each time t , a new vertex labeled t arrives and forms m edges, one at a time, to existing nodes $v \in [t - 1]$:

$$\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta \text{ where } \delta > -m.$$

- $\deg(v)$ is updated after each edge is added
- $\delta = \infty$: uniform-attachment (no degree preference)
- $\delta = 0$: Barabási-Albert model [Barabási-Albert'99]

Preferential attachment model (PAM)

At each time t , a new vertex labeled t arrives and forms m edges, one at a time, to existing nodes $v \in [t - 1]$:

$$\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta \text{ where } \delta > -m.$$

- $\deg(v)$ is updated after each edge is added
- $\delta = \infty$: uniform-attachment (no degree preference)
- $\delta = 0$: Barabási-Albert model [Barabási-Albert'99]
- The smaller δ , the stronger preference for high-degree vertices

Preferential attachment model (PAM)

At each time t , a new vertex labeled t arrives and forms m edges, one at a time, to existing nodes $v \in [t - 1]$:

$$\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta \text{ where } \delta > -m.$$

- $\deg(v)$ is updated after each edge is added
- $\delta = \infty$: uniform-attachment (no degree preference)
- $\delta = 0$: Barabási-Albert model [Barabási-Albert'99]
- The **smaller** δ , the **stronger** preference for high-degree vertices
- A popular **dynamical** model that shares many similar features as in empirically studied real-world networks.

Features of PAM: Power-law degree distribution

Theorem (Bollobás-Riordan-Spencer-Tusnády'01, Deijfen-van den Esker-van der Hofstad-Hooghiemstra'09)

PAM with parameter m, δ yields power-law degree sequence with exponent $\tau = 3 + \delta/m > 2$.

Features of PAM: Power-law degree distribution

Theorem (Bollobás-Riordan-Spencer-Tusnády'01, Deijfen-van den Esker-van der Hofstad-Hooghiemstra'09)

PAM with parameter m, δ yields power-law degree sequence with exponent $\tau = 3 + \delta/m > 2$.




Figure: degree sequences in PAM with $m = 2, \delta = 0, \tau = 3, n = 10^6$ (picture courtesy of Remco van der Hofstad)




Figure: degree sequences in Internet Movie Data Base 2007
[Britton-Deijfen-Löf'2007]

Features of PAM: Small world phenomenon



Figure: Six degrees of separation:
“Everybody on this planet is separated
only by six other people”.




Figure: Distances in social networks
Livejournal [Backstrom-Boldi-Rosa-
Ugander-Vigna'2010]

Features of PAM: Small world phenomenon



Figure: Six degrees of separation:
“Everybody on this planet is separated
only by six other people”.

- Question: Can we rigorously justify the small world phenomenon in PAM?




Figure: Distances in social networks
Livejournal [Backstrom-Boldi-Rosa-
Ugander-Vigna'2010]

Features of PAM: Small world phenomenon



Figure: Six degrees of separation:
“Everybody on this planet is separated
only by six other people”.




Figure: Distances in social networks
Livejournal [Backstrom-Boldi-Rosa-
Ugander-Vigna'2010]

- Question: Can we rigorously justify the small world phenomenon in PAM?
- Equivalently, does PAM have small **diameters**?

Previous results on diameters of PAM

average degree: $2m$; $\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta$;

Previous results on diameters of PAM

average degree: $2m$; $\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta$;

- [Pittel'94]: the diameter of PAM with $m = 1, \delta > -1$ is typically

$$(1 + o(1))^{\frac{2(1+\delta)\log n}{(2+\delta)\theta}},$$

where $\theta \in (0, 1)$ is the solution to $\theta + (1 + \delta)(1 + \log \theta) = 0$.

Previous results on diameters of PAM

average degree: $2m$; $\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta$;

- [Pittel'94]: the diameter of PAM with $m = 1, \delta > -1$ is typically

$$(1 + o(1))^{\frac{2(1+\delta)\log n}{(2+\delta)\theta}},$$

where $\theta \in (0, 1)$ is the solution to $\theta + (1 + \delta)(1 + \log \theta) = 0$.

- [Bollobás-Riordan'09]: the diameter of PAM with $m \geq 2, \delta = 0$ is typically $(1 + o(1))^{\frac{\log n}{\log \log n}}$.

Previous results on diameters of PAM

average degree: $2m$; $\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta$;

- [Pittel'94]: the diameter of PAM with $m = 1, \delta > -1$ is typically

$$(1 + o(1))^{\frac{2(1+\delta) \log n}{(2+\delta)\theta}},$$

where $\theta \in (0, 1)$ is the solution to $\theta + (1 + \delta)(1 + \log \theta) = 0$.

- [Bollobás-Riordan'09]: the diameter of PAM with $m \geq 2, \delta = 0$ is typically $(1 + o(1))^{\frac{\log n}{\log \log n}}$.
- [Caravenna-Garavaglia-van der Hofstad'19]: the diameter of PAM with $m \geq 2, -m < \delta < 0$ is typically

$$(1 + o(1))^{\left(\frac{4}{|\log(1+\delta/m)|} + \frac{2}{\log m} \right) \log \log n}.$$

Previous results on diameters of PAM

average degree: $2m$; $\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta$;

- [Pittel'94]: the diameter of PAM with $m = 1, \delta > -1$ is typically

$$(1 + o(1))^{\frac{2(1+\delta)\log n}{(2+\delta)\theta}},$$

where $\theta \in (0, 1)$ is the solution to $\theta + (1 + \delta)(1 + \log \theta) = 0$.

- [Bollobás-Riordan'09]: the diameter of PAM with $m \geq 2, \delta = 0$ is typically $(1 + o(1))^{\frac{\log n}{\log \log n}}$.
- [Caravenna-Garavaglia-van der Hofstad'19]: the diameter of PAM with $m \geq 2, -m < \delta < 0$ is typically

$$(1 + o(1))^{\left(\frac{4}{|\log(1+\delta/m)|} + \frac{2}{\log m}\right) \log \log n}.$$

- Remaining case: PAM with $m \geq 2, \delta > 0$.

PAM with $m \geq 2, \delta > 0$

average degree: $2m$; $\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta$;

PAM with $m \geq 2, \delta > 0$

average degree: $2m$; $\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta$;

- Difficulties for extending previous argument:

PAM with $m \geq 2, \delta > 0$

average degree: $2m$; $\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta$;

- Difficulties for extending previous argument:
 - $m \geq 2 \implies$ no tree structure;

PAM with $m \geq 2, \delta > 0$

average degree: $2m$; $\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta$;

- Difficulties for extending previous argument:
 - $m \geq 2 \implies$ no tree structure;
 - $\delta > 0 \implies$ no hub structure.

PAM with $m \geq 2, \delta > 0$

average degree: $2m$; $\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta$;

- Difficulties for extending previous argument:
 - $m \geq 2 \implies$ no tree structure;
 - $\delta > 0 \implies$ no hub structure.
- Difficulties in the model:

PAM with $m \geq 2, \delta > 0$

average degree: $2m$; $\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta$;

- Difficulties for extending previous argument:
 - $m \geq 2 \implies$ no tree structure;
 - $\delta > 0 \implies$ no hub structure.
- Difficulties in the model:
 - Lack of independence;

PAM with $m \geq 2, \delta > 0$

average degree: $2m$; $\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta$;

- Difficulties for extending previous argument:

- $m \geq 2 \implies$ no tree structure;
- $\delta > 0 \implies$ no hub structure.

- Difficulties in the model:

- Lack of independence;
- Harder to couple to the local limit.

PAM with $m \geq 2, \delta > 0$

average degree: $2m$; $\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta$;

- Difficulties for extending previous argument:
 - $m \geq 2 \implies$ no tree structure;
 - $\delta > 0 \implies$ no hub structure.
- Difficulties in the model:
 - Lack of independence;
 - Harder to couple to the local limit.
- [Dommers-van der Hofstad-Hooghiemstra'10]: the diameter of PAM with $m \geq 2, \delta > 0$ is typically $O(\log n)$.

Typical distance of PAM with $m \geq 2, \delta > 0$

average degree: $2m$; $\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta$; $\text{PA}_n^{(m,\delta)}$: law of PAM

Theorem (van der Hofstad-Zhu'25+)

Let ν to be the exponential growth parameter of the local limit of the preferential attachment model, then

$$\mathbb{P}_{G \sim \text{PA}_n^{(m,\delta)}} \mathbb{P}_{u,v \sim \text{unif}(V(G))} (\text{dist}_G(u, v) = (1 + o(1)) \log_\nu n) = 1 - o(1),$$

Typical distance of PAM with $m \geq 2, \delta > 0$

average degree: $2m$; $\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta$; $\text{PA}_n^{(m,\delta)}$: law of PAM

Theorem (van der Hofstad-Zhu'25+)

Let ν to be the exponential growth parameter of the local limit of the preferential attachment model, then

$$\mathbb{P}_{G \sim \text{PA}_n^{(m,\delta)}} \mathbb{P}_{u,v \sim \text{unif}(V(G))} (\text{dist}_G(u, v) = (1 + o(1)) \log_\nu n) = 1 - o(1),$$

- Implies that typically we have $\text{dist}_G(u, v) = (1 + o(1)) \log_\nu n$ for $\geq 99\%$ vertex pairs (thus typically $\text{diam}(G) \geq (1 + o(1)) \log_\nu n$).

Typical distance of PAM with $m \geq 2, \delta > 0$

average degree: $2m$; $\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta$; $\text{PA}_n^{(m,\delta)}$: law of PAM

Theorem (van der Hofstad-Zhu'25+)

Let ν to be the exponential growth parameter of the local limit of the preferential attachment model, then

$$\mathbb{P}_{G \sim \text{PA}_n^{(m,\delta)}} \mathbb{P}_{u,v \sim \text{unif}(V(G))} (\text{dist}_G(u, v) = (1 + o(1)) \log_\nu n) = 1 - o(1),$$

- Implies that typically we have $\text{dist}_G(u, v) = (1 + o(1)) \log_\nu n$ for $\geq 99\%$ vertex pairs (thus typically $\text{diam}(G) \geq (1 + o(1)) \log_\nu n$).
- Relies on first/second moment method + path counting technique.

Typical distance of PAM with $m \geq 2, \delta > 0$

average degree: $2m$; $\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta$; $\text{PA}_n^{(m,\delta)}$: law of PAM

Theorem (van der Hofstad-Zhu'25+)

Let ν to be the exponential growth parameter of the local limit of the preferential attachment model, then

$$\mathbb{P}_{G \sim \text{PA}_n^{(m,\delta)}} \mathbb{P}_{u,v \sim \text{unif}(V(G))} (\text{dist}_G(u, v) = (1 + o(1)) \log_\nu n) = 1 - o(1),$$

- Implies that typically we have $\text{dist}_G(u, v) = (1 + o(1)) \log_\nu n$ for $\geq 99\%$ vertex pairs (thus typically $\text{diam}(G) \geq (1 + o(1)) \log_\nu n$).
- Relies on first/second moment method + path counting technique.
- Conjecture in [van der Hofstad-Zhu'25+]: typically the diameter of PAM with $m \geq 2, \delta > 0$ is also $(1 + o(1)) \log_\nu n$.

Our result: from typical distance to diameter

average degree: $2m$; $\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta$; $\text{PA}_n^{(m,\delta)}$: law of PAM

Theorem (Du-G.-L.-Zhu'25+)

- Let $M_n = M_n(G)$ be the median of pairwise vertex distances of $G \sim \text{PA}_n^{(m,\delta)}$.
- Let $R_n = R_n(G)$ satisfying $\#\{R_n\text{-neighborhood of } u\} \geq (\log n)^2$ for all $u \in V(G)$.

Then we have $\mathbb{P}_{G \sim \text{PA}_n^{(m,\delta)}}(\text{diam}(G) \leq M_n + O(1) \cdot R_n) = 1 - o(1)$.

Our result: from typical distance to diameter

average degree: $2m$; $\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta$; $\text{PA}_n^{(m,\delta)}$: law of PAM

Theorem (Du-G.-L.-Zhu'25+)

- Let $M_n = M_n(G)$ be the median of pairwise vertex distances of $G \sim \text{PA}_n^{(m,\delta)}$.
- Let $R_n = R_n(G)$ satisfying $\#\{R_n\text{-neighborhood of } u\} \geq (\log n)^2$ for all $u \in V(G)$.

Then we have $\mathbb{P}_{G \sim \text{PA}_n^{(m,\delta)}}(\text{diam}(G) \leq M_n + O(1) \cdot R_n) = 1 - o(1)$.

- Note that [van der Hofstad-Zhu'25+] implies that typically $M_n(G) = (1 + o(1)) \log_\nu n$.

Our result: from typical distance to diameter

average degree: $2m$; $\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta$; $\text{PA}_n^{(m,\delta)}$: law of PAM

Theorem (Du-G.-L.-Zhu'25+)

- Let $M_n = M_n(G)$ be the median of pairwise vertex distances of $G \sim \text{PA}_n^{(m,\delta)}$.
- Let $R_n = R_n(G)$ satisfying $\#\{R_n\text{-neighborhood of } u\} \geq (\log n)^2$ for all $u \in V(G)$.

Then we have $\mathbb{P}_{G \sim \text{PA}_n^{(m,\delta)}}(\text{diam}(G) \leq M_n + O(1) \cdot R_n) = 1 - o(1)$.

- Note that [van der Hofstad-Zhu'25+] implies that typically $M_n(G) = (1 + o(1)) \log_\nu n$.
- [Du-G.-L.-Zhu'25+]: typically $R_n(G) \leq (\log n)^{\frac{2}{3}}$. (expected to be far from tight).

Our result: from typical distance to diameter

average degree: $2m$; $\mathbb{P}(t \rightarrow v) \propto \deg(v) + \delta$; $\text{PA}_n^{(m,\delta)}$: law of PAM

Theorem (Du-G.-L.-Zhu'25+)

- Let $M_n = M_n(G)$ be the median of pairwise vertex distances of $G \sim \text{PA}_n^{(m,\delta)}$.
- Let $R_n = R_n(G)$ satisfying $\#\{R_n\text{-neighborhood of } u\} \geq (\log n)^2$ for all $u \in V(G)$.

Then we have $\mathbb{P}_{G \sim \text{PA}_n^{(m,\delta)}}(\text{diam}(G) \leq M_n + O(1) \cdot R_n) = 1 - o(1)$.

- Note that [van der Hofstad-Zhu'25+] implies that typically $M_n(G) = (1 + o(1)) \log_\nu n$.
- [Du-G.-L.-Zhu'25+]: typically $R_n(G) \leq (\log n)^{\frac{2}{3}}$. (expected to be far from tight).
- Conclusion: typically $\text{diam}(G) \leq (1 + o(1)) \log_\nu n$.

Discussions: from typical distance to diameter

It seems that our result

$$\text{diam}(G) \leq M_n(G)^{\leftarrow \text{average distance}} + O(1) \cdot R_n(G)^{\leftarrow \text{depth for large neighborhood}}$$

holds for many interesting cases beyond the scope of PAM, e.g.

Discussions: from typical distance to diameter

It seems that our result

$$\text{diam}(G) \leq M_n(G)^{\leftarrow \text{average distance}} + O(1) \cdot R_n(G)^{\leftarrow \text{depth for large neighborhood}}$$

holds for many interesting cases beyond the scope of PAM, e.g.

- Random d -regular graph ($d \geq 3$):

Discussions: from typical distance to diameter

It seems that our result

$$\text{diam}(G) \leq M_n(G)^{\leftarrow \text{average distance}} + O(1) \cdot R_n(G)^{\leftarrow \text{depth for large neighborhood}}$$

holds for many interesting cases beyond the scope of PAM, e.g.

- Random d -regular graph ($d \geq 3$):
 - [Chung-Lu'02]: $M_n = \log_{d-1} n + O(\log \log n)$;

Discussions: from typical distance to diameter

It seems that our result

$$\text{diam}(G) \leq M_n(G)^{\leftarrow \text{average distance}} + O(1) \cdot R_n(G)^{\leftarrow \text{depth for large neighborhood}}$$

holds for many interesting cases beyond the scope of PAM, e.g.

- Random d -regular graph ($d \geq 3$):
 - [Chung-Lu'02]: $M_n = \log_{d-1} n + O(\log \log n)$;
 - $R_n = O(\log \log n)$;

Discussions: from typical distance to diameter

It seems that our result

$$\text{diam}(G) \leq M_n(G)^{\leftarrow \text{average distance}} + O(1) \cdot R_n(G)^{\leftarrow \text{depth for large neighborhood}}$$

holds for many interesting cases beyond the scope of PAM, e.g.

- Random d -regular graph ($d \geq 3$):
 - [Chung-Lu'02]: $M_n = \log_{d-1} n + O(\log \log n)$;
 - $R_n = O(\log \log n)$;
 - [Bollobás-Fernandez De La Vega'81]:
diameter $= \log_{d-1} n + O(\log \log n)$.

Discussions: from typical distance to diameter

It seems that our result

$$\text{diam}(G) \leq M_n(G)^{\leftarrow \text{average distance}} + O(1) \cdot R_n(G)^{\leftarrow \text{depth for large neighborhood}}$$

holds for many interesting cases beyond the scope of PAM, e.g.

- Random d -regular graph ($d \geq 3$):
 - [Chung-Lu'02]: $M_n = \log_{d-1} n + O(\log \log n)$;
 - $R_n = O(\log \log n)$;
 - [Bollobás-Fernandez De La Vega'81]:
diameter $= \log_{d-1} n + O(\log \log n)$.
- Giant component of Erdős-Rényi graph with average degree $\lambda = 1 + \Omega(1)$:

Discussions: from typical distance to diameter

It seems that our result

$$\text{diam}(G) \leq M_n(G)^{\leftarrow \text{average distance}} + O(1) \cdot R_n(G)^{\leftarrow \text{depth for large neighborhood}}$$

holds for many interesting cases beyond the scope of PAM, e.g.

- Random d -regular graph ($d \geq 3$):
 - [Chung-Lu'02]: $M_n = \log_{d-1} n + O(\log \log n)$;
 - $R_n = O(\log \log n)$;
 - [Bollobás-Fernandez De La Vega'81]:
diameter $= \log_{d-1} n + O(\log \log n)$.
- Giant component of Erdős-Rényi graph with average degree $\lambda = 1 + \Omega(1)$:
 - [Riordan-Wormald'08]: $M_n = c(\lambda) \log n$;

Discussions: from typical distance to diameter

It seems that our result

$$\text{diam}(G) \leq M_n(G)^{\leftarrow \text{average distance}} + O(1) \cdot R_n(G)^{\leftarrow \text{depth for large neighborhood}}$$

holds for many interesting cases beyond the scope of PAM, e.g.

- Random d -regular graph ($d \geq 3$):
 - [Chung-Lu'02]: $M_n = \log_{d-1} n + O(\log \log n)$;
 - $R_n = O(\log \log n)$;
 - [Bollobás-Fernandez De La Vega'81]:
diameter $= \log_{d-1} n + O(\log \log n)$.
- Giant component of Erdős-Rényi graph with average degree $\lambda = 1 + \Omega(1)$:
 - [Riordan-Wormald'08]: $M_n = c(\lambda) \log n$;
 - $R_n = \Theta(1) \cdot \log n$;

Discussions: from typical distance to diameter

It seems that our result

$$\text{diam}(G) \leq M_n(G)^{\leftarrow \text{average distance}} + O(1) \cdot R_n(G)^{\leftarrow \text{depth for large neighborhood}}$$

holds for many interesting cases beyond the scope of PAM, e.g.

- Random d -regular graph ($d \geq 3$):
 - [Chung-Lu'02]: $M_n = \log_{d-1} n + O(\log \log n)$;
 - $R_n = O(\log \log n)$;
 - [Bollobás-Fernandez De La Vega'81]:
diameter $= \log_{d-1} n + O(\log \log n)$.
- Giant component of Erdős-Rényi graph with average degree $\lambda = 1 + \Omega(1)$:
 - [Riordan-Wormald'08]: $M_n = c(\lambda) \log n$;
 - $R_n = \Theta(1) \cdot \log n$;
 - [Fernholz-Ramachandran'07] (see also [Ding-Kim-Lubetzky-Peres'10] for more general λ): diameter $= (1 + \Theta(1)) \cdot \text{average distance}$.

Proof idea

Proof idea

- **Two random sources:** denote $\text{PA}(\cdot)$ the distribution of G_n and $\mathbb{P}_{u,v}$ the uniform selection.

Proof idea

- **Two random sources:** denote $\text{PA}(\cdot)$ the distribution of G_n and $\mathbb{P}_{u,v}$ the uniform selection.
- Let M_n be the upper bound of “**typical**” **median distance**: with prob. $1 - o(1)$ over PA on G_n

$$\mathbb{P}_{u,v \sim \text{UNIF}(V(G_n))} [\text{dist}(u, v) \leq M_n \mid G_n] \geq 1/2.$$


- By [van der Hofstad and Zhu 25], we can take $M_n = (1 + o(1)) \log_v n$.

Proof idea

- **Two random sources:** denote $\text{PA}(\cdot)$ the distribution of G_n and $\mathbb{P}_{u,v}$ the uniform selection.
- Let M_n be the upper bound of “typical” median distance: with prob. $1 - o(1)$ over PA on G_n

$$\mathbb{P}_{u,v \sim \text{UNIF}(V(G_n))} [\text{dist}(u, v) \leq M_n \mid G_n] \geq 1/2.$$

- By [van der Hofstad and Zhu 25], we can take $M_n = (1 + o(1)) \log_v n$.
- **High level idea:** $\forall u, v$, with probability $1 - o(1/n^2)$, there exists two vertices in their respective R_n -neighborhoods with distance at most M_n .




Proof idea

- **Two random sources:** denote $\text{PA}(\cdot)$ the distribution of G_n and $\mathbb{P}_{u,v}$ the uniform selection.
- Let M_n be the upper bound of “typical” median distance: with prob. $1 - o(1)$ over PA on G_n

$$\mathbb{P}_{u,v \sim \text{UNIF}(V(G_n))} [\text{dist}(u, v) \leq M_n \mid G_n] \geq 1/2.$$

- By [van der Hofstad and Zhu 25], we can take $M_n = (1 + o(1)) \log_{\nu} n$.
- **High level idea:** $\forall u, v$, with probability $1 - o(1/n^2)$, there exists two vertices in their respective R_n -neighborhoods with distance at most M_n .



- Diameter at most $M_n + 2R_n$. $M_n = \log_{\nu} n$, $R_n = o(\log n)$.

Uniform growth of neighborhood size

Lemma

Taking $R_n = (\log n)^{2/3}$,

$$\text{PA} \left[|N_{R_n}(u)| \geq (\log n)^4, \forall u \in V(G_n) \right] = 1 - o(1).$$

Uniform growth of neighborhood size

Lemma

Taking $R_n = (\log n)^{2/3}$,

$$\text{PA} \left[|N_{R_n}(u)| \geq (\log n)^4, \forall u \in V(G_n) \right] = 1 - o(1).$$

- **Far from tight.** Expected to be $R_n = O(\log \log n)$ for $|N_{R_n}(u)| \geq \text{polylog}(n)$.

Uniform growth of neighborhood size

Lemma

Taking $R_n = (\log n)^{2/3}$,

$$\text{PA} \left[|N_{R_n}(u)| \geq (\log n)^4, \forall u \in V(G_n) \right] = 1 - o(1).$$

- **Far from tight.** Expected to be $R_n = O(\log \log n)$ for $|N_{R_n}(u)| \geq \text{polylog}(n)$.
 - $m \geq 2$ condition \Rightarrow exponential growth in neighborhood size.

Uniform growth of neighborhood size

Lemma

Taking $R_n = (\log n)^{2/3}$,

$$\text{PA} \left[|N_{R_n}(u)| \geq (\log n)^4, \forall u \in V(G_n) \right] = 1 - o(1).$$

- **Far from tight.** Expected to be $R_n = O(\log \log n)$ for $|N_{R_n}(u)| \geq \text{polylog}(n)$.
 - $m \geq 2$ condition \Rightarrow exponential growth in neighborhood size.
- **Major Challenge:** dealing with dependence issue.

Lemma (Conditional attachment lemma)

Let E be a set of potential edges in $G_n \sim \text{PA}$ and A be a set of vertices. Assume that $A \subset [s, n]$, then

$$\text{PA}[u \rightarrow A \mid E \subset E(G_n)] \leq \frac{|A|(m + \delta + 1) + |E|}{(2s - 2)m + s\delta}.$$

Key ingredient: typical vertices

Key ingredient: typical vertices

- **Typical vertices**: for any u , let $\mathcal{A}(u)$ denote the set of vertices w with $\text{dist}(u, w) \leq M_n$. u is called **typical** if $|\mathcal{A}(u)| \geq n/10$.

Key ingredient: typical vertices

- **Typical vertices**: for any u , let $\mathcal{A}(u)$ denote the set of vertices w with $\text{dist}(u, w) \leq M_n$. u is called **typical** if $|\mathcal{A}(u)| \geq n/10$.
- With prob. $1 - o(1)$, $\#\{\text{typical vertices}\} \geq n/10$. Denoted by \mathcal{G}_1 .

Key ingredient: typical vertices

- **Typical vertices**: for any u , let $\mathcal{A}(u)$ denote the set of vertices w with $\text{dist}(u, w) \leq M_n$. u is called **typical** if $|\mathcal{A}(u)| \geq n/10$.
- With prob. $1 - o(1)$, $\#\{\text{typical vertices}\} \geq n/10$. Denoted by \mathcal{G}_1 .
 - [van der Hofstad-Zhu'25] implies

$$\tilde{\mathcal{G}}_1 \triangleq \left\{ G_n : \mathbb{P}_{u,v \sim \text{unif}^{\otimes 2}}[\text{dist}_{G_n}(u, v) \leq M_n \mid G_n] \geq 1/2 \right\}.$$

holds w.p. $1 - o(1)$.

Key ingredient: typical vertices

- **Typical vertices**: for any u , let $\mathcal{A}(u)$ denote the set of vertices w with $\text{dist}(u, w) \leq M_n$. u is called **typical** if $|\mathcal{A}(u)| \geq n/10$.
- With prob. $1 - o(1)$, $\#\{\text{typical vertices}\} \geq n/10$. Denoted by \mathcal{G}_1 .
 - [van der Hofstad-Zhu'25] implies

$$\tilde{\mathcal{G}}_1 \triangleq \left\{ G_n : \mathbb{P}_{u,v \sim \text{unif}^{\otimes 2}}[\text{dist}_{G_n}(u, v) \leq M_n \mid G_n] \geq 1/2 \right\}.$$

holds w.p. $1 - o(1)$.

- We have $\tilde{\mathcal{G}}_1 \subset \mathcal{G}_1$. Assuming \mathcal{G}_1^c ,

$$\begin{aligned} & \mathbb{P}_{u,v}[\text{dist}_{G_n}(u, v) \leq M_n \mid G_n] \\ & \leq \mathbb{P}_u[u \text{ is typical} \mid G_n] + \mathbb{P}_{u,v}[u \text{ is not typical, } \text{dist}(u, v) \leq M_n \mid G_n] \\ & \leq \frac{1}{10} + \frac{1}{10} < 1/2. \end{aligned}$$

Key ingredient: typical vertices

- **Typical vertices**: for any u , let $\mathcal{A}(u)$ denote the set of vertices w with $\text{dist}(u, w) \leq M_n$. u is called **typical** if $|\mathcal{A}(u)| \geq n/10$.
- With prob. $1 - o(1)$, $\#\{\text{typical vertices}\} \geq n/10$. Denoted by \mathcal{G}_1 .
 - [van der Hofstad-Zhu'25] implies

$$\tilde{\mathcal{G}}_1 \triangleq \left\{ G_n : \mathbb{P}_{u,v \sim \text{unif}^{\otimes 2}}[\text{dist}_{G_n}(u, v) \leq M_n \mid G_n] \geq 1/2 \right\}.$$

holds w.p. $1 - o(1)$.


- We have $\tilde{\mathcal{G}}_1 \subset \mathcal{G}_1$. Assuming \mathcal{G}_1^c ,

$$\begin{aligned} & \mathbb{P}_{u,v}[\text{dist}_{G_n}(u, v) \leq M_n \mid G_n] \\ & \leq \mathbb{P}_u[u \text{ is typical} \mid G_n] + \mathbb{P}_{u,v}[u \text{ is not typical, } \text{dist}(u, v) \leq M_n \mid G_n] \\ & \leq \frac{1}{10} + \frac{1}{10} < 1/2. \end{aligned}$$

- $\Rightarrow \text{PA}(\mathcal{G}_1) = 1 - o(1)$ under PA.


Bridging the gap by sprinkling

- Breaking $[1, n]$ into three sets: $G_{n-2K_n} \triangleq [1, n - 2K_n]$, $I_1 \triangleq [n - 2K_n, n - K_n]$ and $I_2 \triangleq [n - K_n, n]$ where $K_n = n / \log n$.
- There exists a w_1 in I_1 , such that $w_1 \rightarrow$ a typical vertex and $w_1 \rightarrow N_{R_n}(u)$, with probability $1 - (1 - O((\log n)^4/n))^{K_n} = 1 - \exp(-O((\log n)^3))$.
- For any $u, v \in [1, n - 2K_n]$, $\text{dist}_{G_n}(u, v) \leq M_n + 2R_n + 4$ with prob. $1 - o(1/n^2)$.




Bridging the gap by sprinkling

- Breaking $[1, n]$ into three sets: $G_{n-2K_n} \triangleq [1, n - 2K_n]$, $I_1 \triangleq [n - 2K_n, n - K_n]$ and $I_2 \triangleq [n - K_n, n]$ where $K_n = n / \log n$.
- There exists a w_1 in I_1 , such that $w_1 \rightarrow$ a typical vertex and $w_1 \rightarrow N_{R_n}(u)$, with probability $1 - (1 - O((\log n)^4/n))^{K_n} = 1 - \exp(-O((\log n)^3))$.
- For any $u, v \in [1, n - 2K_n]$, $\text{dist}_{G_n}(u, v) \leq M_n + 2R_n + 4$ with prob. $1 - o(1/n^2)$.




Bridging the gap by sprinkling

- Breaking $[1, n]$ into three sets: $G_{n-2K_n} \triangleq [1, n - 2K_n]$, $I_1 \triangleq [n - 2K_n, n - K_n]$ and $I_2 \triangleq [n - K_n, n]$ where $K_n = n / \log n$.
- There exists a w_1 in I_1 , such that $w_1 \rightarrow$ a typical vertex and $w_1 \rightarrow N_{R_n}(u)$, with probability $1 - (1 - O((\log n)^4/n))^{K_n} = 1 - \exp(-O((\log n)^3))$.
- For any $u, v \in [1, n - 2K_n]$, $\text{dist}_{G_n}(u, v) \leq M_n + 2R_n + 4$ with prob. $1 - o(1/n^2)$.




Bridging the gap by sprinkling

- Breaking $[1, n]$ into three sets: $G_{n-2K_n} \triangleq [1, n - 2K_n]$, $I_1 \triangleq [n - 2K_n, n - K_n]$ and $I_2 \triangleq [n - K_n, n]$ where $K_n = n / \log n$.
- There exists a w_1 in I_1 , such that $w_1 \rightarrow$ a typical vertex and $w_1 \rightarrow N_{R_n}(u)$, with probability $1 - (1 - O((\log n)^4/n))^{K_n} = 1 - \exp(-O((\log n)^3))$.
- For any $u, v \in [1, n - 2K_n]$, $\text{dist}_{G_n}(u, v) \leq M_n + 2R_n + 4$ with prob. $1 - o(1/n^2)$.




Bringing the gap by sprinkling

- Breaking $[1, n]$ into three sets: $G_{n-2K_n} \triangleq [1, n - 2K_n]$, $I_1 \triangleq [n - 2K_n, n - K_n]$ and $I_2 \triangleq [n - K_n, n]$ where $K_n = n / \log n$.
- There exists a w_1 in I_1 , such that $w_1 \rightarrow$ a typical vertex and $w_1 \rightarrow N_{R_n}(u)$, with probability $1 - (1 - O((\log n)^4/n))^{K_n} = 1 - \exp(-O((\log n)^3))$.
- For any $u, v \in [1, n - 2K_n]$, $\text{dist}_{G_n}(u, v) \leq M_n + 2R_n + 4$ with prob. $1 - o(1/n^2)$.



Bringing the gap by sprinkling

- Breaking $[1, n]$ into three sets: $G_{n-2K_n} \triangleq [1, n - 2K_n]$, $I_1 \triangleq [n - 2K_n, n - K_n]$ and $I_2 \triangleq [n - K_n, n]$ where $K_n = n / \log n$.
- There exists a w_1 in I_1 , such that $w_1 \rightarrow$ a typical vertex and $w_1 \rightarrow N_{R_n}(u)$, with probability $1 - (1 - O((\log n)^4/n))^{K_n} = 1 - \exp(-O((\log n)^3))$.
- For any $u, v \in [1, n - 2K_n]$, $\text{dist}_{G_n}(u, v) \leq M_n + 2R_n + 4$ with prob. $1 - o(1/n^2)$.




Tackling the last $2K_n$ vertices

- Show for any $u \in [n - 2K_n, n]$, $\text{dist}_{G_n}(u, [1, n - 2K_n]) \leq R_n + 2$.
- **BFS** (breadth-first search) in $N_{R_n}(u)$. \mathcal{F}_{k-1} as the attachment of v_1, \dots, v_{k-1} .
- Applying the conditional attachment lemma,

$$\mathbb{P}[v_k \not\rightarrow [1, n - 2K_n] \mid \mathcal{F}_{k-1}] = O\left(\frac{1}{\log n}\right).$$

- $\mathbb{P}[N_{R_n}(u) \cap [1, n - 2K_n] = \emptyset] \leq (1/\log n)^{(\log n)^3} = o(1/n^3)$, by **iterative conditioning**.




Tackling the last $2K_n$ vertices

- Show for any $u \in [n - 2K_n, n]$, $\text{dist}_{G_n}(u, [1, n - 2K_n]) \leq R_n + 2$.
- **BFS** (breadth-first search) in $N_{R_n}(u)$. \mathcal{F}_{k-1} as the attachment of v_1, \dots, v_{k-1} .
- Applying the conditional attachment lemma,

$$\mathbb{P}[v_k \not\rightarrow [1, n - 2K_n] \mid \mathcal{F}_{k-1}] = O\left(\frac{1}{\log n}\right).$$

- $\mathbb{P}[N_{R_n}(u) \cap [1, n - 2K_n] = \emptyset] \leq (1/\log n)^{(\log n)^3} = o(1/n^3)$, by **iterative conditioning**.




Tackling the last $2K_n$ vertices

- Show for any $u \in [n - 2K_n, n]$, $\text{dist}_{G_n}(u, [1, n - 2K_n]) \leq R_n + 2$.
- **BFS** (breadth-first search) in $N_{R_n}(u)$. \mathcal{F}_{k-1} as the attachment of v_1, \dots, v_{k-1} .
- Applying the conditional attachment lemma,

$$\mathbb{P}[v_k \not\rightarrow [1, n - 2K_n] | \mathcal{F}_{k-1}] = O\left(\frac{1}{\log n}\right).$$

- $\mathbb{P}[N_{R_n}(u) \cap [1, n - 2K_n] = \emptyset] \leq (1/\log n)^{(\log n)^3} = o(1/n^3)$, by **iterative conditioning**.




Tackling the last $2K_n$ vertices

- Show for any $u \in [n - 2K_n, n]$, $\text{dist}_{G_n}(u, [1, n - 2K_n]) \leq R_n + 2$.
- **BFS** (breadth-first search) in $N_{R_n}(u)$. \mathcal{F}_{k-1} as the attachment of v_1, \dots, v_{k-1} .
- Applying the conditional attachment lemma,

$$\mathbb{P}[v_k \not\rightarrow [1, n - 2K_n] \mid \mathcal{F}_{k-1}] = O\left(\frac{1}{\log n}\right).$$

- $\mathbb{P}[N_{R_n}(u) \cap [1, n - 2K_n] = \emptyset] \leq (1/\log n)^{(\log n)^3} = o(1/n^3)$, by **iterative conditioning**.



Outlook and discussions

- We prove the asymptotic diameter of the PA model is $\log_{\nu} n$ when $m \geq 2, \delta > 0$.
- **End of the story?** We hope the proof technique can be applied to other graph models.
- **Open question:**
 - (1) Conditional on diameter being $C \log_{\nu} n$ with $C > 1$, what is the graph structure?
 - (2) Pinpointing the second order of the diameter of PA model. **Conjecture:** $\log_{\nu} n + O(\log \log n)$.

Thank you!